



UNIVERSITI PUTRA MALAYSIA

**THE SYNTHESIS AND BIOACTIVITY STUDY OF
ANTHRAQUINONES, AND THE ISOLATION OF BIOACTIVE
COMPOUNDS FROM LEEA INDICA (BURM.F.) MERR.**

KOUSHIK SAHA.

IB 2005 7

**THE SYNTHESIS AND BIOACTIVITY STUDY OF
ANTHRAQUINONES, AND THE ISOLATION OF BIOACTIVE
COMPOUNDS FROM *LEEA INDICA* (BURM.F.) MERR.**

By

KOUSHIK SAHA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

May 2005



*Dedicated to
My Dearest Parents;
Loving Wife, Jhumu and
Dearest Son, Ani.*



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirements for the degree of Doctor of Philosophy

**THE SYNTHESIS AND BIOACTIVITY STUDY OF
ANTHRAQUINONES, AND THE ISOLATION OF BIOACTIVE
COMPOUNDS FROM *LEEA INDICA* (BURM.F.) MERR.**

By

KOUSHIK SAHA

May 2005

Chairman: Professor Nordin Hj. Lajis, PhD

Institute: Bioscience

Damnacanthal and nordamnacanthal were synthesized due to their interesting property as cytotoxic agents against several cancer cell-lines. Friedel-Craft condensation reaction has been used to prepare anthraquinone back-bone in view of its less stringent requirements with regards to reaction conditions (such as humidity), highly reactive reagents and lower cost. Various other synthetic methods were used to prepare their derivatives. A total 43 derivatives of anthraquinone have been successfully synthesized including damnacanthal and nordamnacanthal. A new rearrangement was observed when phthalic anhydride reacted with different halophenols leading to a few rearrangement products depending on the types of halophenol used. All the synthesized compounds were subjected to cytotoxicity assay against different cancer cell lines and antioxidant activities. A total 22 anthraquinone derivatives showed the cytotoxicity against the cell-lines used. Among the compounds, 2-bromomethyl-1,3-dimethoxyanthraquinone (A-34) was found as most cytotoxic against all the cell-lines. 3-Acetoxy-2-bromomethyl-1-methoxyanthraquinone (A-42) and damnacanthal (A-46) also showed strong cytotoxicity against different cancer cell-lines. On the other hand, only two



synthesized anthraquinones, damnacanthal (A-46) and nordamnacanthal (A-37) showed strong antioxidant activity comparable with vitamin E.

A preliminary antioxidant and nitric oxide inhibition activities screening of seven medicinal plants including *Lasianthus oblongus*, *Psychotria rostrata*, *Spermacoce exilis*, *Chasalia chartacea*, *Hedyotis verticillata*, *Spermacoce articularis* and *Leea indica* have been conducted properly. *Leea indica* (family: Leeaceae) has been selected for further investigation. Phytochemical study on this plant has resulted the isolation of a novel carotenoid, leeatene (L-7) and nine other known compounds including squalene (L-1), hexadecanoyl- β -amyrin (L-2), vitamin E (L-3), 1-tetratriacontanol (L-4), β -amyrin (L-5), 3-hydroxy-12-oleanen-28-oic acid (L-6), β -sitosteryl- β -D-glucopyranoside (L-8), 2 α ,3 α ,23-trihydroxy-12-oleanen-28-oic acid (L-9) and phloridzin (L-10). The chemical structure of the compounds was established based on spectral studies including ultraviolet-visible, infrared, one and two dimensional nuclear magnetic resonance and mass spectroscopies. All isolated compounds were tested for their antioxidant and cytotoxic activities. Three compounds including vitamin E (L-3), leeatene (L-7) and phloridzin (L-10) showed antioxidant activity whereas squalene (L-1), hexadecanoyl- β -amyrin (L-2), 1-tetratriacontanol (L-4), 3-hydroxy-12-oleanen-28-oic acid (L-6), leeatene (L-7), β -sitosteryl- β -D-glucopyranoside (L-8), 2 α ,3 α ,23-trihydroxy-12-oleanen-28-oic acid (L-9) were found cytotoxic against different cancer cell lines.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**SINTESIS DAN KEAKTIFAN BIOLOGI ANTRAKUINON, DAN
PEMENCILAN SEBATIAN AKTIF DARIPADA *LEEA INDICA* (BURM.F.)
MERR.**

Oleh

KOUSHIK SAHA

Mei 2005

Pengerusi: Profesor Nordin Hj. Lajis, PhD

Institut: Biosains

Damnakantal dan nordamnakantal telah disintesis kerana didapati mempunyai sifat keaktifan biologi yang menarik sebagai agen sitotoksik terhadap beberapa sel kanser. Kondensasi Fridel-Craft telah digunakan untuk menghasilkan gelang antrakuinon berdasarkan keperluan yang kurang ketat termasuk keadaan tindakbalas (seperti kelembapan) dan kereaktifan reagen yang tinggi. Sejumlah 43 terbitan antrakuinon telah berjaya disintesis termasuk damnakantal dan nordamnakantal. Satu kaedah baru penyusunan semula telah dijalankan apabila ftalik anhidrida ditindak balas dengan halofenol yang berbeza mengarah kepada beberapa hasil penyusunan semula bergantung kepada jenis halofenol yang digunakan. Kesemua sebatian yang telah disintesis diuji kesan sitotoksik terhadap beberapa talian sel kanser dan aktiviti antioksidan. Dua puluh dua terbitan antrakuinon didapati sitotoksik terhadap beberapa talian sel yang digunakan. Daripada semua sebatian, 2-bromometil-1,3-dimetoksiantrakuinon (A-34) didapati paling sitotoksik diikuti 3-asitoksi-2-bromometil-1-metoksiantrakuinon (A-42) dan damnakantal (A-46). Selain itu, hanya 2 antrakuinon yang talah disintesis iaitu damnakantal (A-46) dan

nordamnakantal (A-37) menunjukkan aktiviti antioksidan yang tinggi berbanding vitamin E.

Kajian awal aktiviti antioksidan dan perencatan nitrik oksida (NO) daripada 7 tumbuhan ubatan termasuk *Lasianthus oblongus*, *Psychotria rostrata*, *Spermacoce exilis*, *Chasalia chartacea*, *Hedyotis verticillata*, *Spermacoce articularis* and *Leea indica* telah dijalankan. *Leea indica* (keluarga: Leeaceae) telah dipilih untuk kajian lebih lanjut. Kajian fitokimia ke atas tumbuhan ini telah berakhir dengan pemencilan satu novel karotenoid, leeatene (L-7) serta sembilan sebatian yang telah diketahui termasuk skualena (L-1), heksadekanoil- β -amirin (L-2), vitamin E (L-3), 1-tetratriakontanol (L-4), β -amirin (L-5), asid 3-hidroksi-12-oleanen-28-oik (L-6), β -sitosteril- β -D-glukopiranosida (L-8), asid 2 α ,3 α ,23-trihidroksi-12-oleanen-28-oik (L-9) dan floridzin (L-10). Struktur kimia sebatian-sebatian tersebut telah dikenalpasti berdasarkan kajian spektroskopi termasuk spektroskopi ultra lembayung-nampak, spektroskopi inframerah, spektroskopi resonan magnetik nuclear satu dan dua dimensi dan spektroskopi jisim. Kesemua sebatian telah diuji aktiviti antioksidan dan sitotoksik menggunakan kaedah yang berbeza. Tiga sebatian termasuk vitamin E (L-3), leeatene (L-7) dan floridzin (L-10) telah menunjukkan aktiviti antioksidan manakala skualena (L-1), heksadekanoil- β -amirin (L-2), 1-tetratriakontanol (L-4), asid 3-hidroksi-12-oleanen-28-oik (L-6), leeatene (L-7), β -sitosteril- β -D-glukopiranosida (L-8), asid 2 α ,3 α ,23-trihidroksi-12-oleanen-28-oik (L-9) didapati sitotoksik terhadap beberapa talian sel kanser.

ACKNOWLEDGEMENTS

I would like to express my deepest sense of gratitude and sincere thanks to my supervisor, Prof. Dr. Nordin Hj. Lajis for his constant encouragement, keen interest and thoughtful suggestions throughout my research work. I am also delighted to express my heartiest gratitude to the members of the supervisory committee, Prof. Dr. Ahmad Sazali Hamzah, Assoc. Prof. Dr. Daud Ahmad Israf Ali and Assoc. Prof. Dr. Khozirah Shaari for their thoughtful suggestions and unfailing co-operations throughout the study.

I would like to offer my sincere thanks to Dr. Johnson Stanslas for his co-operation to get the cytotoxic assay results. I wish to offer my warm thanks to all of my lab mates, friends and well wishers for their positive attitude and co-operations.

I wish to express my hearty gratitude to my parents, brother and sisters for their constant encouragement and sacrifice throughout this study.

Finally, I am grateful to my wife, Jhumu, for her loving support and inspiration during the course of this higher study.



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF SCHEMES	xxii
LIST OF ABBREVIATIONS	xxvii
 CHAPTER	
 I INTRODUCTION	1
General	1
Plant Secondary Metabolites	3
Natural Product Synthesis	6
Anthraquinones in <i>Morinda elliptica</i>	8
Objective of Research	9
 II LITERATURE REVIEW	11
Synthesis of Anthraquinones	11
Biosynthesis of Anthraquinones	45
Biological Activity of Anthraquinones	48
Botany, Distribution and Ethnobotany of <i>Leea</i> Species	59
Previous Work on Genus <i>Leea</i>	62
 III RESEARCH EXPERIMENTAL	63
General Instrumentations	63
Chromatographic methods	63
Solvents	65
Synthesis of Anthraquinones	65
Friedel-Craft Condensation of Phthalic Anhydride and	
Benzene Derivatives	65
Halogen Rearrangement in Friedel-Craft	
Condensation	77
Debromination of Bromoanthraquinones	77
Methylation of Hydroxyanthraquinones	78
Oxidation of Methylanthraquinone	88
Hydrolysis of Geminal Dibromomethyl or	
Ethoxybromomethylanthraquinone	89
Bromination of Methylanthraquinone	90
Demethylation of Methoxyanthraquinone	95
Substitution of Bromine	97
Acetylation of Hydroxyanthraquinone	98
Plant Materials	101

	Extraction and Fractionation of Crude Sample from <i>Leea indica</i>	101
	Isolation of Compounds from <i>Leea indica</i>	103
	Compounds from Chloroform Extracts	103
	Compounds from Methanol Extracts	104
	Spectral Data of Isolated Compounds	105
	Bioassay Procedures	111
	Cytotoxic Assay	111
	Nitric Oxide Inhibitory Assay	113
	Antioxidant Assay	115
IV	CHEMISTRY AND BIOLOGICAL ACTIVITIES OF ANTHRAQUINONES	119
	Synthesis of Anthraquinones	119
	Synthesis of Anthraquinone Skeleton	119
	Debromination of Bromoanthraquinone	138
	Methylation of Hydroxyanthraquinone	139
	Formylation of Anthraquinone	143
	Oxidation of Benzylic Methyl of Anthraquinone	144
	Bromination of Benzylic Methyl of Anthraquinone	145
	Demethylation of Methoxyanthraquinone	148
	Synthesis of Nordamnacanthal	149
	Alternative Route for the Synthesis of Nordamnacanthal	151
	Synthesis of Damnacanthal	153
	Biological Activities of Anthraquinones	157
	Evaluation of Cytotoxicity	157
	Evaluation of Antioxidant Activity	162
	Structure-Activity Relationship of Anthraquinones	165
V	CHEMISTRY AND BIOLOGICAL ACTIVITIES OF ISOLATED COMPOUNDS FROM <i>LEEA INDICA</i>	168
	Isolation of Compounds	168
	Characterization of Isolated Compounds	170
	Squalene (L-1)	170
	O-Hexadecanoyl- β -amyrin (L-2)	179
	Vitamin E (L-3)	189
	1-Tetratriacontanol (L-4)	199
	β -Amyrin (L-5)	204
	3-Hydroxy-12-oleanen-28-oic acid (L-6)	215
	Leeatene (L-7)	221
	β -Sitosteryl- β -D-glucopyranoside (L-8)	249
	2 α ,3 α ,23-Trihydroxy-12-oleanen-28-oic acid (L-9)	260
	4,4',6'-Trihydroxydihydrochalcone-2'- β - D-glucopyranoside (L-10)	274
	Screening of Some Malaysian Medicinal Plants	286
	Biological Activity of the Crude Extracts of <i>Leea indica</i>	294
	Biological Activity of the Fractions from the Column	
	Chromatography of Crude Extracts of <i>Leea indica</i>	296
	Biological Activity of the Isolated Compounds	297

VI	CONCLUSION	301
BIBLIOGRAPHY		308
BIODATA OF THE AUTHOR		326



LIST OF TABLES

Table	Page
1.1 Some Pharmaceuticals of Plant or Microorganism Origin	2
4.1 Cytotoxic Activity of Anthraquinones (IC_{50} in μM)	161
4.2 Absorbance Value of Control, Standard and Samples on the Last Day (8 th Day) of FTC and TBA Methods	162
5.1 1H NMR, ^{13}C NMR and HMBC Spectral Data	224
5.2 List of Plant Species, the Percentage of Yields and Their Traditional Uses	287
5.3 DPPH Free Radical Scavenging Activity of Plant Samples and Standards (Vit C, Quercetin & BHT)	290
5.4 Percent Inhibition of H_2O_2 Production in Macrophages by Plant Extracts	291
5.5 Percent Inhibition of Nitric Oxide Accumulation in Cell Culture Supernatants of Macrophages by Plant Extracts	293
5.6 Cytotoxic Activity of Chloroform and Methanol Extracts Against Breast Cancer Cell Line (Mcf7) Using MTT Assay	296
5.7 DPPH Free Radical Scavenging Activity of Isolated Compounds from <i>Leea indica</i> Including Standards (Quercetin & BHT)	300
5.8 Cytotoxic Activity of Isolated Compounds from <i>Leea indica</i>	300



LIST OF FIGURES

Figure		Page
1.1 Main Stream of Secondary Metabolism		4
2.1 Some Structural Resemblance of Substituted Anthraquinones to Anthracyclinones / Anthracyclines		26
2.2 Biogenetic Numbering for Anthraquinones		46
2.3 Anthraquinones Found as Mutagenic <i>in Vitro</i> Study (Westendorf <i>et al.</i> , 1990)		49
2.4 Structure of 1,8-Dihydroxyanthraquinone Derivatives Investigated For Genotoxicity (Mueller <i>et al.</i> , 1998)		50
2.5 Antileukemic Anthraquinones from <i>Morinda parvifolia</i> (Chang <i>et al.</i> , 1982 & 1984)		51
2.6 Anthraquinones from <i>Morinda elliptica</i> Found as Cytotoxic (Ali <i>et al.</i> , 2000)		52
2.7 Structure of Cytotoxic BQ-I and BQ-II (Inbaraj <i>et al.</i> , 1999)		52
2.8 Structure of Mitoxantrone, a Clinically Useful Antineoplastic Agent		53
2.9 Derivatives of 1,3-Dihydroxyanthraquinone as Cytotoxic Compounds (Wei <i>et al.</i> , 2000)		54
2.10 Synthetic Esters of 2-Substituted 1,4-Dihydroxyanthraquinones as Multifunctional Anticancer Agents (Jin <i>et al.</i> , 2001)		55
2.11 Structure of Two New Naturally Occurring Bioactive Anthraquinones (Onogi <i>et al.</i> , 2002; Babu <i>et al.</i> , 2003)		56
2.12 Anthraquinones That Inhibit Lipid Peroxidation in Rat Heart Mitochondria (Huang <i>et al.</i> , 1995)		57
2.13 Plant <i>Leea indica</i>		61
2.14 Leaves and Flowers of <i>Leea indica</i>		61
2.15 Favanoids in <i>L. guineensis</i>		62
4.1 Reaction Intermediate of Demethylation According to Preston <i>et al.</i> , 1983		149

4.2	Antioxidant Activity Using FTC Method	164
4.3	Antioxidant Activity Using TBA Method	164
5.1	Structure of Squalene (L-1)	171
5.2	Infrared Spectrum of Squalene (L-1)	172
5.3	Mass Spectrum of Squalene (L-1)	173
5.4	Proton NMR Spectrum of Squalene (L-1)	174
5.5	Carbon-13 NMR Spectrum of Squalene (L-1)	175
5.6	DEPT Spectrum of Squalene (L-1)	176
5.7	HSQC Spectrum of Squalene (L-1)	177
5.8	HMBC Spectrum of Squalene (L-1)	178
5.9	Structure and Numbering of <i>O</i> -Hexadecanoyl- β -amyrin (L-2)	180
5.10	Selected HMBC Correlations of <i>O</i> -Hexadecanoyl- β -amyrin (L-2)	181
5.11	Infrared Spectrum of <i>O</i> -Hexadecanoyl- β -amyrin (L-2)	182
5.12	Mass Spectrum of <i>O</i> -Hexadecanoyl- β -amyrin (L-2)	183
5.13	Proton NMR Spectrum of <i>O</i> -Hexadecanoyl- β -amyrin (L-2)	184
5.14	Carbon-13 NMR Spectrum of <i>O</i> -Hexadecanoyl- β -amyrin (L-2)	185
5.15	DEPT Spectrum of <i>O</i> -Hexadecanoyl- β -amyrin (L-2)	186
5.16	HSQC Spectrum of <i>O</i> -Hexadecanoyl- β -amyrin (L-2)	187
5.17	HMBC Spectrum of <i>O</i> -Hexadecanoyl- β -amyrin (L-2)	188
5.18	Structure and Numbering of Vitamin E (L-3)	190
5.19	Selected ^1H - ^{13}C Long Range Correlations in HMBC Spectrum of Vitamin E (L-3)	190
5.20	Infrared Spectrum of Vitamin E (L-3)	191
5.21	Mass Spectrum of Vitamin E (L-3)	192
5.22	Proton NMR Spectrum of Vitamin E (L-3)	193

5.23	Carbon-13 NMR Spectrum of Vitamin E (L-3)	194
5.24	DEPT Spectrum of Vitamin E (L-3)	195
5.25	COSY Spectrum of Vitamin E (L-3)	196
5.26	HSQC Spectrum of Vitamin E (L-3)	197
5.27	HMBC Spectrum of Vitamin E (L-3)	198
5.28	Structure of 1-Tetratriacontanol (L-4)	199
5.29	Infrared Spectrum of 1-Tetratriacontanol (L-4)	200
5.30	Mass Spectrum of 1-Tetratriacontanol (L-4)	201
5.31	Proton NMR Spectrum of 1-Tetratriacontanol (L-4)	202
5.32	Carbon-13 NMR Spectrum of 1-Tetratriacontanol (L-4)	203
5.33	Structure and Numbering of β -Amyrin (L-5)	205
5.34	Selected ^1H - ^{13}C Long Range Correlations in HMBC Spectrum of β -Amyrin (L-5)	206
5.35	Infrared Spectrum of β -Amyrin (L-5)	207
5.36	Mass Spectrum of β -Amyrin (L-5)	208
5.37	Proton NMR Spectrum of β -Amyrin (L-5)	209
5.38	Carbon-13 NMR Spectrum of β -Amyrin (L-5)	210
5.39	DEPT Spectrum of β -Amyrin (L-5)	211
5.40	COSY Spectrum of β -Amyrin (L-5)	212
5.41	HSQC Spectrum of β -Amyrin (L-5)	213
5.42	HMBC Spectrum of β -Amyrin (L-5)	214
5.43	Structure and Numbering of 3-Hydroxy-12-oleanen-28-oic acid (L-6)	216
5.44	Infrared Spectrum of 3-Hydroxy-12-oleanen-28-oic acid (L-6)	217
5.45	Mass Spectrum of 3-Hydroxy-12-oleanen-28-oic acid (L-6)	218
5.46	Proton NMR Spectrum of 3-Hydroxy-12-oleanen-28-oic acid (L-6)	219



5.47	Carbon-13 NMR Spectrum of 3-Hydroxy-12-oleanen-28-oic acid (L-6)	220
5.48	Structure and Numbering of Leeatene (L-7)	223
5.49	Selected NOESY Interactions and Configuration of Leeatene (L-7)	224
5.50	Infrared Spectrum of Leeatene (L-7)	227
5.51	Mass Spectrum of Leeatene (L-7)	228
5.52	Proton NMR Spectrum of Leeatene (L-7)	229
5.53	Carbon-13 NMR Spectrum of Leeatene (L-7)	230
5.54	DEPT Spectrum of Leeatene (L-7)	231
5.55	COSY Spectrum of Leeatene (L-7)	232
5.56	COSY Spectrum of Leeatene (Expansion 1)	233
5.57	COSY Spectrum of Leeatene (Expansion 2)	234
5.58	HSQC Spectrum of Leeatene (L-7)	235
5.59	HMBC Spectrum of Leeatene (L-7)	236
5.60	HMBC Spectrum of Leeatene (Expansion 1)	237
5.61	HMBC Spectrum of Leeatene (Expansion 2)	238
5.62	HMBC Spectrum of Leeatene (Expansion 3)	239
5.63	HMBC Spectrum of Leeatene (Expansion 4)	240
5.64	HMBC Spectrum of Leeatene (Expansion 5)	241
5.65	1D TOCSY Spectrum of Leeatene (L-7)	242
5.66	1D TOCSY Spectrum of Leeatene (L-7)	243
5.67	1D TOCSY Spectrum of Leeatene (L-7)	244
5.68	1D TOCSY Spectrum of Leeatene (L-7)	245
5.69	CD Spectrum of Leeatene (L-7)	246
5.70	CD Spectrum of Leeatene (L-7)	247
5.71	NOESY Spectrum of Leeatene (L-7)	248

5.72	Structure and Numbering of β -Sitosteryl- β -D-glucopyranoside (L-8)	250
5.73	Selected ^1H - ^{13}C Long Range Correlations in HMBC Spectrum of β -Sitosteryl- β -D-glucopyranoside (L-8)	251
5.74	Infrared Spectrum of β -Sitosteryl- β -D-glucopyranoside (L-8)	252
5.75	Mass Spectrum of β -Sitosteryl- β -D-glucopyranoside (L-8)	253
5.76	Proton NMR Spectrum of β -Sitosteryl- β -D-glucopyranoside (L-8)	254
5.77	Carbon-13 NMR Spectrum of β -Sitosteryl- β -D-glucopyranoside (L-8)	255
5.78	DEPT Spectrum of β -Sitosteryl- β -D-glucopyranoside (L-8)	256
5.79	COSY Spectrum of β -Sitosteryl- β -D-glucopyranoside (L-8)	257
5.80	HSQC Spectrum of β -Sitosteryl- β -D-glucopyranoside (L-8)	258
5.81	HMBC Spectrum of β -Sitosteryl- β -D-glucopyranoside (L-8)	259
5.82	Structure and Numbering of $2\alpha,3\alpha,23$ -Trihydroxy-12-oleanen-28-oic acid (L-9)	262
5.83	Selected ^1H - ^{13}C Long Range Correlations in HMBC Spectrum of $2\alpha,3\alpha,23$ -Trihydroxy-12-oleanen-28-oic acid (L-9)	262
5.84	Selected ^1H - ^1H NOESY Correlations of $2\alpha,3\alpha,23$ -Trihydroxy-12-oleanen-28-oic acid (L-9)	263
5.85	Infrared Spectrum of $2\alpha,3\alpha,23$ -Trihydroxy-12-oleanen-28-oic acid (L-9)	264
5.86	Mass Spectrum of $2\alpha,3\alpha,23$ -Trihydroxy-12-oleanen-28-oic acid (L-9)	265
5.87	Proton NMR Spectrum of $2\alpha,3\alpha,23$ -Trihydroxy-12-oleanen-28-oic acid (L-9)	266
5.88	Carbon-13 NMR Spectrum of $2\alpha,3\alpha,23$ -Trihydroxy-12-oleanen-28-oic acid (L-9)	267
5.89	DEPT Spectrum of $2\alpha,3\alpha,23$ -Trihydroxy-12-oleanen-28-oic acid (L-9)	268
5.90	COSY Spectrum of $2\alpha,3\alpha,23$ -Trihydroxy-12-oleanen-28-oic acid (L-9)	269

5.91	HSQC Spectrum of 2 α ,3 α ,23-Trihydroxy-12-oleanen-28-oic acid (L-9)	270
5.92	HMBC Spectrum of 2 α ,3 α ,23-Trihydroxy-12-oleanen-28-oic acid (L-9)	271
5.93	NOESY Spectrum of 2 α ,3 α ,23-Trihydroxy-12-oleanen-28-oic acid (L-9)	272
5.94	NOESY Spectrum of 2 α ,3 α ,23-Trihydroxy-12-oleanen-28-oic acid (Expansion)	273
5.95	Structure and Numbering of Phloridzin (L-10)	276
5.96	Selected ^1H - ^{13}C Long Range Correlations in HMBC Spectrum of Phloridzin (L-10)	276
5.97	Infrared Spectrum of Phloridzin (L-10)	277
5.98	Mass Spectrum of Phloridzin (L-10)	278
5.99	Proton NMR Spectrum of Phloridzin (L-10)	279
5.100	Proton NMR Spectrum of Phloridzin (L-10) in Acetone-d ₆	280
5.101	Carbon-13 NMR Spectrum of Phloridzin (L-10)	281
5.102	DEPT Spectrum of Phloridzin (L-10)	282
5.103	COSY Spectrum of Phloridzin (L-10)	283
5.104	HSQC Spectrum of Phloridzin (L-10)	284
5.105	HMBC Spectrum of Phloridzin (L-10)	285
5.106	Antioxidant Activity of Crude Methanol Extracts Using FTC Method (Data Indicate Mean of 3 Replicate Determination, n = 3)	288
5.107	Antioxidant Activity of Crude Methanol Extracts Using TBA Method (Data Indicate Mean of 3 Replicate Determination, n = 3)	289
5.108	Antioxidant Activity of Chloroform and Methanol Extracts of <i>Leea indica</i> Using FTC Method (Data Indicate Mean of 3 Replicate Determination, n = 3)	295
5.109	Antioxidant Activity of Chloroform and Methanol Extracts of <i>Leea indica</i> Using TBA Method (Data Indicate Mean of 3 Replicate Determination, n = 3)	295

5.110	Antioxidant Activity of Isolated Compounds from <i>Leea indica</i> Using FTC Method (Data Indicate Mean of 3 Replicate Determination, n = 3), Note: Q = Quercetin	299
5.111	Antioxidant Activity of Isolated Compounds from <i>Leea indica</i> Using TBA Method (Data Indicate Mean of 3 Replicate Determination, n = 3), Note: Q = Quercetin	299
6.1	Structure of Prepared Anthraquinones	302
6.2	Structure of Isolated Compounds from <i>Leea indica</i>	305



LIST OF SCHEMES

Scheme	Page
2.1 General Friedel-Craft Condensation to Anthraquinone	11
2.2 Preparation of Anthraquinones Using Phthalic Anhydride and Benzene Derivatives as Starting Materials (Hirose, 1960)	12
2.3 Synthesis of Polyhydroxyanthraquinone Using Friedel-Craft Acylation (Cameron <i>et al.</i> , 1981)	13
2.4 Synthesis of 4-Bromo-1-hydroxyanthraquinone from Phthalic Anhydride and 4-Bromophenol (Aleksandrzak and Gronowska 1987)	13
2.5 Synthesis of Hydroxyanthraquinone-2-carboxylic Acid (Smith <i>et al.</i> , 1993)	14
2.6 Preparation of Naturally Occurring 2-Hydroxy-1-methoxy-3-methylanthraquinone (Zhang <i>et al.</i> , 1996)	15
2.7 Synthesis of Aza Anthraquinones, Tolypocladin and Isotolypocladin by Using Friedel-Craft Reaction (Werner <i>et al.</i> , 1997)	16
2.8 Synthesis of Anthraquinone from Phthalic Anhydride and Aryl Magnesium Bromide (Braun, 1979)	16
2.9 Synthesis of 9,10-Anthraquinone from Naphthoquinone and 1,4-Diacetoxybutadiene Using Diels-Alder Reaction (Hill and Carlson, 1964)	17
2.10 Condensation of Keten Acetals and Halonaphthoquinones: Regiospecific Synthesis of Anthraquinone (Banville <i>et al.</i> , 1974)	18
2.11 Synthesis of 1,3,8-Trihydroxyanthraquinone Using 5-Hydroxy-1,4-naphthoquinone and 1,1-Diethoxyethene (Cameron <i>et al.</i> , 1978)	19
2.12 Anthraquinones from Naphthoquinonoid Dienophiles and 1-Methoxy-1-trimethylsilyloxybutadienes (Cameron <i>et al.</i> , 1986)	19
2.13 Synthesis of 2,3-Dimethoxy-6-methylanthraquinone Based on Diels-Alder Reaction (Zhang <i>et al.</i> , 1996)	20
2.14 Synthesis of Anthraquinones from 6-Methoxy-4-methyl-2-pyrone and Naphthoquinones by Diels-Alder Reaction (Jung <i>et al.</i> , 1984)	22

2.15	Reaction of 2-Prop-2-enylidene-1,3-dioxolan with Naphthoquinones Finally Afforded Hydroxyanthraquinones (Ley <i>et al.</i> , 1981)	23
2.16	Synthesis of Quinone Aldehyde by Cycloaddition Followed by Oxidation (Cormier <i>et al.</i> , 1992)	23
2.17	Synthesis of Anthraquinone Glycosides by Regio- and Stereoselective Cycloadditions (Beagley <i>et al.</i> , 1992)	24
2.18	Synthesis of Naturally Occurring 1-Hydroxyanthraquinones Based on 1,4-Addition Elimination Reaction of Phenylsulfonyl Stabilized Anion (Hauser and Prasanna, 1982)	28
2.19	Reaction of Lithium Salts of Phthalide with Arynes (Sammes and Dodsworth, 1979)	29
2.20	Reaction Between Aryne and Lithium Cyanophthalide (Khanapure <i>et al.</i> , 1987)	30
2.21	Anthraquinone Synthesis Involves the Reaction of the Anion of 3-Phenylsulphonylphthalides with Substituted Quinone Monoacetals (Russell and Warrener, 1981)	31
2.22	Synthesis of Anthraquinone Using Cyclohexadienone as Precursor (Mitchell and Russell, 1997)	32
2.23	Anthraquinone Synthesis by Using 3-Bromophthalide and Aromatic Substrate as Starting Materials (Kim <i>et al.</i> , 1979)	33
2.24	Synthesis of Anthraquinones Using Cyanophthalides and Quinone Monoketals (Freskos <i>et al.</i> , 1985)	34
2.25	Formation of Phthalide from Organometallic Compound as a Precursor of Anthraquinone Synthesis (Baldwin and Bair, 1978)	34
2.26	Preparation of Phthalide from Silylated or Brominated Benzamide for Anthraquinone Synthesis (Mills and Snieckus, 1984)	36
2.27	Synthesis of <i>O</i> -Methylchrysophanol from Cyanophthalide and Cyclohexenone (Kraus <i>et al.</i> , 1983)	36
2.28	Preparation of Methoxyanthraquinone from Cyanomethyl Methoxybenzophenone (Davies <i>et al.</i> , 1969)	37
2.29	Methoxy Substituted Benzyl Magnesium Chlorides Displace the <i>Ortho</i> -Methoxy Group from <i>o</i> -Methoxyaryl-dihydro-oxazole, Resultant <i>o</i> -Benzylbenzoic Acids Converted to Anthraquinones (Nicoletti <i>et al.</i> , 1988)	38

2.30	Synthesis of Anthraquinones Using Methoxycarbonylbenzoic Acid and Benzene Derivative as Starting Materials (Gleim <i>et al.</i> , 1978)	39
2.31	Arylacetonitrile and Cyclohexene Ester Used as Starting Materials for Anthraquinone Synthesis (Parker and Kallmerten, 1980)	39
2.32	Synthesis of Anthraquinones by Using Bromobenzaldehydes and Phenyllithium Derivatives as Starting Materials (Almeida and Costa, 1996)	40
2.33	Reactions of 2-Acetyl-1,4-naphthoquinone with Enamines (Kobayashi <i>et al.</i> , 2000)	41
2.34	Conversion of Methyl to Formyl by Using NBS Followed by Hydrolysis (Roberts <i>et al.</i> , 1977)	42
2.35	β -Formylation of α -Hydroxyanthraquinone by the Rearrangement of Allyl Group Followed by Ozonolysis (Roberts and Rutledge, 1977)	42
2.36	Selective Demethylation of Methoxyanthraquinone via Aryloxydifluoroboron Chelates (Preston <i>et al.</i> , 1983)	43
2.37	Conversion of 1,4-Anthraquinones to 9,10-Anthraquinones (Wiseman <i>et al.</i> , 1980)	43
2.38	Synthesis of Anthra[1,2- <i>b</i>]pyrans by Marschalk Alkylation Followed by Baker-Venkataraman Chain Elongation and Acid Catalyzed Cyclization (Krohn and Vitz, 2004)	44
2.39	Biosynthetic Route of Anthraquinones via Shikimate-mevalonate Pathway (Wijnsma and Verpoorte, 1986)	47
2.40	Biosynthesis of Anthraquinones via Polyketide Pathway (Torsell, 1983)	48
4.1	Basic Equation for Anthraquinone Synthesis	120
4.2	Synthesis of Anthraquinones (Hirose, 1960)	120
4.3	Friedel-Craft Condensation Between Phthalic Anhydride and 4-Bromophenol	122
4.4	Reaction Between Phthalic Anhydride and 4-Bromoanisole	123
4.5	Condensation Between Phthalic Anhydride and 3-Bromophenol	124
4.6	Reaction Between Phthalic Anhydride and 2-Bromophenol	125
4.7	Friedel-Craft Acylation of Phthalic Anhydride with 4-Chlorophenol or 4-Iodophenol	126