



UNIVERSITI PUTRA MALAYSIA

***CLIMATE RISK MAPPING OF DENGUE AND MALARIA CASES IN
KUALA LUMPUR AND SELANGOR, MALAYSIA***

HAFIZ HASSAN

FPAS 2013 18



**CLIMATE RISK MAPPING OF DENGUE AND MALARIA CASES IN KUALA
LUMPUR AND SELANGOR, MALAYSIA**

By

HAFIZ BIN HASSAN

Thesis Submitted to the School of Graduate Studies, Universiti Putra

Malaysia, in Fulfillment of the Requirements for the Degree of Masters of Science

JUNE 2013

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement of the degree of Masters of Science

CLIMATE RISK MAPPING OF DENGUE AND MALARIA CASES IN KUALA LUMPUR AND SELANGOR, MALAYSIA

By

HAFIZ BIN HASSAN

June 2013

Chair: Shamarina Shohaimi, PhD

Faculty: Faculty of Science

Local health enforcement activities regarding mosquito-borne disease control and eradication seldom takes climate factors into consideration. The objectives of this study are to find out the historical trend of dengue fever (DF) and malaria fever (MF) in Peninsular Malaysia, to find out the spatial relationship between the climate factors and cases of DF and MF in Selangor and Kuala Lumpur and to prove the feasibility of climate factors to be used in predictive risk mapping for dengue and malaria in Selangor and Kuala Lumpur. For objective one, simple choropleth maps are used with secondary data to visualize the trend and change of DF and MF cases from 1980 to 2010 among states in Peninsular Malaysia supplemented with simple correlation analysis with land uses of forest and urban area percentage. The next objective is carried out by using the geospatial analysis of Standard Deviation Ellipse (SDE) and cluster analysis based on the Getis-Ord General G (G_i^*) Hotspot Analysis spatial statistics. Lastly, the risk map based on climate

factors is formulated through Co-Kriging method which will then be validated against real data. In the initial mapping of Peninsular Malaysia's DF and MF cases of 1980 to 2010, dengue has a rising trend while malaria decreases with Malaysia's population and urbanization growth. Statistical analysis has shown positive correlations for urban areas and dengue ($r = 0.49$), population density and dengue ($r = 0.48$); and forested areas and malaria ($r = 0.74$). Next, SDEs visualized the distribution of climate factors against dengue and malaria cases in Selangor and Kuala Lumpur and through overlay, showed that the mean distributions of dengue and malaria cases were situated within the same focal hotspot as the climate factors. The DF, MF, and rainfall values are found to be clustered with z-values of 2.72, 3.64 and 2.77. A risk map was produced using multiple co-kriging method to predict the cases of DF and MF in Selangor and Kuala Lumpur. Validation using scaled choropleth comparisons showed that the risk map's predicted cases have a difference of 0.2 levels against dengue cases and malaria cases. In conclusion, Peninsular Malaysia's DF trend is rising while MF is decreasing between 1990 and 2010. The climate factors, which are spatially correlated with the distribution of DF and MF cases, can be used to predict of the distribution of future DF and MF cases in Selangor and Kuala Lumpur.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah sarjana.

PEMETAAN RISIKO IKLIM KES DEMAM DENGGI DAN MALARIA KUALA LUMPUR DAN SELANGOR, MALAYSIA

Oleh

HAFIZ BIN HASSAN

Jun 2013

Pengerusi: Shamarina Shohaimi, PhD

Fakulti: Fakulti Sains

Aktiviti penguatkuasaan kesihatan tempatan berkenaan kawalan dan pembasmian penyakit bawaan nyamuk jarang mengambil kira faktor-faktor iklim. Objektif kajian ini adalah untuk mengetahui tren sejarah demam denggi (DF) dan demam malaria (MF) di Semenanjung Malaysia, untuk mengetahui hubungan ruang antara faktor-faktor iklim dan kes-kes DF dan MF di Selangor dan Kuala Lumpur dan untuk membuktikan kebolegunaan faktor-faktor iklim untuk digunakan dalam pemetaan risiko ramalan untuk DF dan MF di Selangor dan Kuala Lumpur. Untuk objektif pertama, peta 'choropleth' yang ringkas dibina menggunakan data sekunder untuk menggambarkan trend dan perubahan kes DF dan MF antara tahun 1980-2010 di antara negeri-negeri di Semenanjung Malaysia. Ia disokong dengan analisis korelasi DF dan MF dengan guna tanah hutan dan peratusan kawasan bandar. Objektif seterusnya dijalankan dengan menggunakan analisis ruang daripada Elips Sisihan piawai (SDE) dan analisis kluster berdasarkan statistik ruang Getis-Ord G_i^* Hotspot Analisis. Akhir sekali, peta risiko berdasarkan faktor-faktor iklim dirumuskan melalui kaedah 'Co-kriging' yang

kemudiannya akan disahkan dengan data sebenar. Dalam pemetaan awal kes DF dan MF dari 1980 hingga 2010 bagi Semenanjung Malaysia, denggi mempunyai tren yang semakin meningkat sementara malaria berkurangan seiring dengan pertumbuhan populasi dan perbandaran Malaysia. Analisis statistik telah menunjukkan korelasi positif bagi kawasan bandar dan denggi ($r = 0.49$), kepadatan penduduk dan denggi ($r = 0.48$), dan kawasan hutan dan malaria ($r = 0.74$). Seterusnya, SDE membayangkan taburan faktor-faktor iklim dan kes-kes DF dan MF di Selangor dan Kuala Lumpur dan melalui tindihan. Dari tindihan ini, ia menunjukkan bahawa taburan min kes DF dan MF yang terletak di dalam fokus hotspot yang sama dengan faktor-faktor iklim. DF, MF, dan taburan hujan purata didapati berkelompok dengan nilai-Z sebanyak 2.72, 3.64 dan 2.77. Sebuah peta risiko telah dihasilkan menggunakan kaedah 'Multiple co-kriging' untuk meramalkan kes DF dan MF di Selangor dan Kuala Lumpur. Pengesahan menggunakan perbandingan choropleth berskala menunjukkan bahawa kes yang diramalkan peta risiko mempunyai perbezaan tahap 0.2 berbanding kes denggi dan malaria sebenar. Kesimpulannya, trend DF Semenanjung Malaysia semakin meningkat manakala MF semakin berkurangan dari tahun 1990-2010. Faktor-faktor iklim, yang berkait rapat dengan taburan reruang kes DF dan MF, boleh digunakan untuk meramal taburan kes-kes DF dan MF di Selangor dan Kuala Lumpur.

ACKNOWLEDGEMENTS

In the making of this thesis, I would like to first and foremost thank almighty God Allah s.w.t. and his Messenger Muhammad (Peace Be Upon Him) for giving me the opportunity and strength to finish my research in UPM. To my parents, Norlidza Abdul and Dr Hassan Mohd Daud whose support is priceless, I sincerely extend my gratitude knowing that it will never be enough to be repaid. Not forgotten to UPM, which GRF allowance provided funds to make this study possible.

I wish to also express my gratitude to the following parties: Dr Nor Rasidah Hashim my supervisor who continually helped me along the way in form of guidance and positive criticism; Dr Shamarina Shohaimi my co-supervisor who provided valuable advice in biostatistics and mosquito epidemiology researcher Dr Nazni who kindly gave valuable input while in the International Public Health Symposium. I would also like to express my gratitude to the governmental agencies including Air Quality Division Department of Environment (DOE), Malaysian Meteorological Department (JMM), Department of Irrigation and Drainage (DID), the National Archives, and lastly the staff at the Kuala Lumpur Territory Health Office especially Deputy Director of Health (Public Health) Dr Zainol Ariffin Pawanchee, Health Officer Muhammad Shafizi B Abdul Wahab (Inspectorate Unit) and Deputy Health Officer Amirul Hisham Muhamad Isa (Vector Diseases Unit) for the input and access to case data. I thank all of the parties above for their precious support and help, because without them I would not have finished all my work in this study.

I certify that a Thesis Examination Committee has met on 18 June 2013 to conduct the final examination of Hafiz Bin Hassan on his thesis entitled "Climate Risk Mapping Of Dengue And Malaria Cases" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Examination Committee were as follows:

Zelina Zaiton Ibrahim, PhD

Associate Professor
Faculty of Environmental Studies
Universiti Putra Malaysia
(Chairman)

Malina Osman, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Internal Examiner)

Hasmadi Ismail, PhD

Associate Professor
Faculty of Forestry
Universiti Putra Malaysia
(Internal Examiner)

NORITAH OMAR, PhD

Associate Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia
Date: 20 November 2013

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Masters of Science. The members of the Supervisory Committee were as follows:

Nor Rasidah Hashim, PhD

Senior Lecturer
Faculty of Environmental Studies
Universiti Putra Malaysia
(Chairman)

Shamarina Shohaimi, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: _____

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of
Chairman of
Supervisory

Committee: Shamarina Shohaimi (PhD)

Signature: _____

Name of
Member of
Supervisory

Committee: Nor Rasidah Hashim (PhD)



TABLE OF CONTENTS

ABSTRACT	Page
ABSTRAK	ii
ACKNOWLEDGEMENTS	iv
APPROVAL	vi
DECLARATION	vii
LIST OF TABLES	ix
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xiv
	xvi

CHAPTER

1	INTRODUCTION	
	1.1 General Introduction	1
	1.2 Problem Statement	3
	1.3 Objectives	5
	1.4 Limitations of Study	6
	1.5 Thesis Outline	7
2	LITERATURE REVIEW	
	2.1 Mosquito-borne diseases	9
	2.1.1 Dengue	10
	2.1.2 Malaria	11
	2.2 The vectors	13
	2.2.1 Aedes	13
	2.2.2 Anopheles	14
	2.3 History of Dengue and Malaria in Malaysia	15
	2.3.1 History of Dengue	16
	2.3.2 History of Malaria	17
	2.4 Climatic Factors Influencing Dengue and Malaria Cases	20
	2.4.1 Rainfall	20
	2.4.2 Temperature	21
	2.4.3 Relative Humidity	22
	2.5 Application of Mapping in Disease Surveillance	23
3	MATERIALS AND METHODS	
	3.1 Conceptual Framework of Study	25
	3.2 Data Types	26
	3.2.1 Spatial Data	27
	3.2.2 Case Data	27
	3.2.3 Population Data	30
	3.2.4 Climate Data	31

	3.3 Analysis Methods	35
	3.3.1 Statistical Analysis	34
	3.3.2 Geostatistical Analysis	35
	3.4 Study Area	36
	3.4.1 Peninsular Malaysia	36
	3.4.2 Selangor and Kuala Lumpur	38
4	THE DISTRIBUTION OF DENGUE AND MALARIA IN PENINSULAR MALAYSIA	
	4.1 Introduction	41
	4.2 Materials and Methods	42
	4.2.1 Choropleth	43
	4.2.2 Statistical Analysis	44
	4.3 Results and Discussion	45
	4.3.1 Dengue Rate	45
	4.3.2 Malaria Rate	48
	4.3.3 Statistical analysis	50
	4.4 Conclusion	52
5	SPATIAL DISTRIBUTION OF DENGUE, MALARIA AND EFFECTS OF CLIMATE FACTORS IN SELANGOR AND KUALA LUMPUR	
	5.1 Introduction	53
	5.2 Materials and Methods	53
	5.2.1 Climate Data	54
	5.2.2 Spatial Analyses	56
	5.3 Results and Discussion	61
	5.3.1 Standard Deviation Ellipse	61
	5.3.2 Cluster Analysis	68
	5.4 Conclusion	69
6	CLIMATE FACTOR MAPPING FOR DENGUE AND MALARIA IN SELANGOR AND KUALA LUMPUR	
	6.1 Introduction	71
	6.2 Materials and Methods	72
	6.3 Results and Discussion	76
	6.3.1 Climate risk prediction	77
	6.3.2 Risk map validation	78
	6.4. Conclusion	82

7	SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	
	7.1 Thesis summary and conclusion	84
	7.2 Recommendations for future research	86
	REFERENCES	88
	APPENDICES	94
	BIODATA OF THE STUDENT	108
	LIST OF PUBLICATIONS	110-

LIST OF TABLES

		Page
Table 3.1	List of stations of climate data in Selangor and Kuala Lumpur	32
Table 3.2	List of unavailable data within MET's stations	34
Table 5.1	SDE data input prompt	59
Table 6.1(a)	Calculation of difference between climate risk map and cases of dengue 2010	79
Table 6.1(b)	Scatterplot between climate risk map and cases of dengue 2010.	79
Table 6.2 (a)	Calculation of difference between climate risk map and cases of malaria 2010	81
Table 6.2 (b)	Scatterplot between climate risk map and cases of malaria 2010	81

LIST OF FIGURES

		Page
Figure 3.1	Conceptual Framework of Study	25
Figure 3.2	Location of Peninsular Malaysia in South East Asia	36
Figure 3.3	The states in Peninsular Malaysia	37
Figure 3.4	Districts and main towns of Selangor	38
Figure 3.5	Administrative districts of Kuala Lumpur	39
Figure 4.1	Flowchart of analysis steps	42
Figure 4.2a-f	The prevalence rate of dengue in Peninsular Malaysia 1980, 1985, 1990, 1995, 2000 and 2010	45
Figure 4.3a-d	The prevalence rate of malaria in Peninsular Malaysia 1998, 2000, 2005 and 2010	48
Figure 5.1	Flowchart of data analysis	53
Figure 5.2	Cluster analysis output, in graphical form	61
Figure 5.3	Standard Deviation Ellipse of dengue	62
Figure 5.4	Standard Deviation Ellipse of malaria	63
Figure 5.5	Standard Deviation Ellipse of rainfall.	64
Figure 5.6	Standard Deviation Ellipse of temperature	65
Figure 5.7	Standard Deviation Ellipse of rain days.	66
Figure 5.8	Overlay of Standard Deviation Ellipses (with town names).	67
Figure 6.1	Risk map generated from climatic factors, in its original raster form.	76

Figure 6.2	Choropleth map based on predicted values.	77
Figure 6.3	Comparison of climate risk map and cases of dengue 2010	78
Figure 6.4	Comparison of climate risk map and cases of malaria 2010.	80



LIST OF ABBREVIATIONS

DF Dengue Fever

GIS Geographical Information Systems

KL Kuala Lumpur

JUPEM Jabatan Ukur dan Pemetaan Malaysia

MET Meteorological Department

ASMA Alam Sekitar Malaysia

JAS Jabatan Alam Sekitar

JPS Jabatan Pengairan dan Saliran

MOH Ministry of Health, Malaysia

CHAPTER 1

INTRODUCTION

1.1 General Introduction

Dengue and malaria are the most persistent insect-borne diseases in the Third world. Both of these diseases thrive in poor and developing countries alike, within their endemic zone near the equator with favorable environmental conditions. The diseases claim millions of lives living in such endemic areas, including children. The problem regarding dengue and malaria also concerns the socio-economic conditions of the endemic countries, which hamper the eradication and control efforts. Malaysia is fortunate to have adequate health service funding, but it is still insufficient to eradicate the diseases although the overall number of deaths and cases has decreased (Statistics Department of Malaysia, 2011).

Malaysia has been dealing with dengue fever since its first epidemic in the 1970s. Between the years 1962-1972, an average of 19.4 dengue cases were reported per year in Peninsular Malaysia with majority of the cases in Johor, Melaka and Selangor (Ministry of Health Malaysia, 1986). In the 1980s, a biannual pattern is seen, where case numbers and death tolls peak every two years before going irregular at the end of the decade. Besides the humid and rainy climate being suitable for the vector (*Aedes sp.*) to thrive, the unsanitary conditions of citizens' homes provide plenty of breeding sites, which account for 70% of mosquito larvae found during inspections (Merican, M.I., 2010).

Malaria fever has been endemic in Malaya since before independence with the local name 'Demam Kura'. Compared to dengue's painful symptoms, the common symptoms of malaria is a period of chills and cold sensations phase alternating with a hot, feverish phase. The disease is related to the conditions of colonial Malaya, with lush vegetation and orchards in place before rapid urbanization (Mohamed Hassan, 2008). The clearing of jungles for economic agricultural activities in support of British's Industrial Revolution coupled with migrant labor saw the dawn of an emerging threat of malaria. In the 1960s, the Malayan government started its Malaria Eradication Program in which free Quinine tablets were given to alleviate the sickness (WHO, 2001). The national situation improved, with lesser infections and deaths. Three decades later, malaria saw a resurgence in disease rates, with a possibility of imported cases caused by migrant workers (Dewi, 2009).

The highly urbanized areas in Selangor and Kuala Lumpur have high population density, and are the most populated states in the whole Peninsula. This is due to the various developments and economic opportunities in the region, which attracts many people to work and live in the area, dubbed the Klang Valley. Dengue fever, which vector is the anthropophilic the *Aedes sp.*, is closely monitored as they present immediate risk to the human populations (Ahmad Zahawir, 2009; Shaharuddin et al. 2001). Related to places with high human population in urban areas is the factor of rapid and far-reaching transportation networks such as roads, trains, airplanes and ships which can bridge the gap between places easily in a short time. Such conveniences are built for commerce and travel alike, but it also helps facilitate the spread of infectious diseases (Ferreira, 2005).

Control of dengue and malaria fever has been carried since the first instance of its emergence by the Ministry of Health, Malaysia. For a common locality in Malaysia, the pre-emptive

activity done by the local health office includes breeding source reduction (sanitation), usage of larvicidal powder ABATE, distribution of treated mosquito nets, preventive fogging and mass blood screening (Ministry of Health Malaysia, 1986). If a case does occur and was confirmed, vector control activities were carried out to eliminate mosquitoes and prevent further spread of the disease by fogging and mass sanitation programs (Ministry of Health Malaysia, 1986).

Despite the efforts of controlling, prevention and education by the Ministry of Health, the number of cases still continues to rise after instances of heavy rain and dry spells (Muhammad Harawi, Deputy Environmental Health Officer, Vector Department, Hulu Langat Health Office, pers comm. August 2008). Although the population has done everything it could to control their surroundings, but nature prevailed at last and favored the rapid breeding and growth of the vectors.

1.2 Problem Statement

The extent of mosquito control activities in Malaysian local health authorities' heavily depends on the occurrence of cases (incidences) and the density of mosquito adult and larvae population (Aziz, 2011) in terms of Aedes Index and Breteau Index (Ministry of Health Malaysia, 1986). However, in literature by Kolivras (2006), Richards et al. (2010), and Troyo et al. (2009) environmental and demographic factors rank high in influencing the breeding of mosquitoes and subsequently the incidence and spread of dengue cases in tropical countries.

This study aims to utilize the available climatic data for determining the susceptibility of

dengue and malaria in Selangor and Kuala Lumpur without relying on reactive input such as incidences or mosquito population density.

Mapping has been used by the local health authorities, but only retroactively. The extent of the usage of mapping only involves plotting the point coordinate of an event of interest. While useful in determining locations of cases or causal factors, the mapping method employed only records events which have already happened. Such is the case with the dengue enforcement units, which cases pop up rapidly. The modus operandi for the dengue eradication programme's call for action also depends on received case reports reactively (Aziz, 2011).

Modeling serves to fill in the gap of knowledge encountered by decision-makers when complete data is unavailable (Lloyd and Yu, 1994). Due to breakdowns in communication or inefficiency in relaying information, empirical data could be unavailable for use. As an example, some districts may have an abundance of case data while others do not. At times, the case data collected by field staff are incomplete and compromised, leaving it unusable to be recorded and used. Rather than being discarded, the data can be compared with the modeled ones to gain approximation to the real values.

From the past local studies (Che Dom et al. 2010, Aziz et al. 2012), there are some applications of GIS in the sector of health, especially dengue monitoring. However, none of them involved a state-wide scale with multiple environmental factors including population density using the method of co-kriging. There is no malaria risk mapping in Selangor and Kuala Lumpur as of yet. Being a factor that influences the cases of dengue (and probably malaria too), this study has found that climate factors can be mapped as a risk factor for dengue and malaria can be done to characterize the susceptibility for the area of Selangor and Kuala Lumpur. Thus, the

mapping and modeling in this study can prove useful not only in demonstrating climate factors' influence on the cases, but also predict the areas which will have the risk of having high or low potential for outbreaks.

1.3 Objectives

The main objective is to create a climate-based risk map of dengue and malaria fever-susceptible areas in Kuala Lumpur and Selangor. The high, medium and low risk areas are identified and displayed in a map, and compared to the actual data. The map also needs to be verified to be as close as it can be to the real data, to become a predictive tool. The resulting risk map will function as a map of the susceptibility of areas in Selangor and Kuala Lumpur to dengue, and the probability map for any location in the area to have the highest future incidence. The secondary objectives for this study are to provide a historical overview of the dengue and malaria situation in Peninsular Malaysia and to determine the spatial correlation between climate factors and dengue and malaria cases in Selangor and Kuala Lumpur.

1.4 Limitations of Study

This study has delimitations and limitations in terms of factors, scope and breadth given its time and resources. The limitations of this study are time, cost and manpower. A lot of time was taken for gathering, organizing and processing the data from its raw form for it to be ready for spatial analysis. Budget constraints have limited the amount of data bought from the Meteorological Department to only two selected parameters (rainfall and rain days) for limited number of years and stations.

This study's delimitations are related to its scale and extent of study. The factors used in this risk mapping only involve environmental factors i.e. climatic and population but does not include the factors of vector population. Vector populations (in this case, mosquitoes) consist of emerging populations (larval and egg survey) and infective populations (adult survey). In field surveys, field staff is instructed to eradicate the discovered larvae on sight to prevent future infections. Thus, the emerging population of mosquito larvae is removed from becoming a future risk factor. The presumptive larval surveys are not done continually to all districts under the jurisdiction of the local health office, so the records from larval surveys are not comprehensive. It is impossible to cover all districts all the time with limited resources and time. As for adult populations, the surveys are done after control measures have been taken. This time consuming process of bare leg catching is done in confirmed high risk areas or focal points to evaluate the effectiveness of control measures. The records collected do not represent the true mosquito population, and its timing is too late to be considered into a risk map as the events (dengue and malaria cases) have already taken place.

In terms of scope, the risk map focuses on the states of Selangor and Kuala Lumpur only, and the district-level data used for comparison consist of data from years 2009 and 2010 only as per its availability. The model which uses a smaller spatial scale will ensure a more accurate rendering of risk compared to a nationwide modeling with the same amount of input. In terms of population data, it does not take into account unregistered immigrants, who were also included in the incidence of malaria cases. The census done by the Statistics Department of Malaysia collected their census data in residential areas but often could not overlook shanties and slum areas resided by immigrants who are often inaccessible and hidden. As a result, the population count is not 100% representative of Malaysia's real population.

1.5 Thesis Outline

This thesis contains seven chapters including the current chapter, Introduction, which contains the problem statement and objectives of the study. Chapter 2 is the review of literature related to the problem of mosquito-borne diseases in Malaysia, the influence of weather and climate to the distribution, and application of risk mapping in mosquito-borne diseases. The following chapter, Chapter 3, contains information about the study area including the weather stations from which climatic data was extracted in this study. Chapter 4 presents the historical overview of Peninsular Malaysia's dengue and malaria pre-2010, while Chapter 5 visualizes the spatial distribution of various climatic factors with dengue and malaria cases as Standard Deviation Ellipses (SDE) and cluster analysis in the conurbation of Selangor and KL (Kuala Lumpur). Next, Chapter 6 applies the climate factors to create a model a climate-based of

dengue and malaria risk in for Selangor and KL which is verified against real data. Lastly, the conclusions of the study are presented in the final chapter with suggestions for future research.



REFERENCES

- Abu Bakar, S. and Shafee, N. (2002). Outlook of dengue in Malaysia: a century later. *Malaysian Journal of Pathology* 2002; 24(1) : 23 – 27
- Aggrey, N. and O. Douglason, 2010. Environmental and Socio-economic determinants of malaria prevalence in Uganda. *Earth Science and the Environment, Reprint*, 2(4): 194.
- Ahmad Zahawir, M.M.H (2009). *Applications Of Remote Sensing And GIS For Dengue Epidemic Surveillance In Petaling Jaya, Malaysia*. Masters Thesis. Serdang: Universiti Putra Malaysia. August 2008.
- Ali, M., Wagatsuma, Y., Emch, M. and Breiman, R.F.(2003). Use of geographic information system for defining spatial risk for dengue transmission in Bangladesh: role for *Aedes albopictus* in an urban outbreak. *American Journal of Tropical Medicine and Hygiene* 2003;69: 634–40.
- Alto, B.W. and Juliano, S.A. (2001). Temperature effects on the dynamics of *Aedes albopictus* (Diptera: Culicidae) populations in the laboratory. *Journal of Medical Entomology*. 2001.
- Aziz, S., Ngui, R., Lim, Y.A.L., Sholehah, I, Nur Farhana, J., Azizan, A.S. and Wan Yusoff, W.S. (2012). Spatial pattern of 2009 dengue distribution in Kuala Lumpur using GIS application. *Tropical Biomedicine* 29 (1): 113–120.
- Aziz, S. (2011). Evaluation of the Spatial Risk Factors for High Incidence of Dengue Fever and Dengue Hemorrhagic Fever Using GIS Application. *Sains Malaysiana* 40(8)(2011): 937–943.
- Bledsoe, G.H. (2005). Malaria primer for clinicians in the United States. *Southern Medical Journal* 98 (12): 1197–204.
- Center of Disease Control USA (2010). About Malaria: Disease. Accessible at: <http://www.cdc.gov/malaria/about/disease.html>, Retrieved on February 2 2012.
- Chan, K. L., Y. C. Chan and B. C. Ho. 1971. *Aedes aegypti* (L.) and *Aedes albopictus* (Skuse) in Singapore City. 5. Observations in relation to Dengue hemorrhagic fever. *Bull. W.H.O* 44:651-658.
- Che Dom, N., Abu Hassan, A., Rabiatul, A. and Ismail, R. (2010). Spatial Mapping of Temporal Risk Characteristic of Dengue Cases in Subang Jaya. Proceedings of the International Conference of Science and Social Research 2010. Research Managemant Institute, Universiti Teknologi MARA, 361-366.
- Cockings, S., Dunn, C.E., Bhopal, R.S. and Walker, D.R. (2004). Users' perspectives on epidemiological, GIS and point pattern approaches to analyzing environment and health data. *Health Place* 2004;10:169–82.

Cushing, J.M. (1986). Oscillatory Population Growth in Periodic Environments. *Theoretical Population Biology*. 30: 289–308.

Dawson, W. R. (1992). *Physiological responses of animals to higher temperatures*, pp. 138-158. In R. L. Peters and T. E. Lovejoy (eds.), *Global Warming and Biological Diversity*. Yale University, New Haven, CT.

Department of Statistics Malaysia (2011). Compendium of Environmental Statistics 2010. Retrieved 10 January 2012 from :
http://www.statistics.gov.my/portal/index.php?option=com_content&view=article&id=1464&Itemid=169&lang=en

Dewi, K.K., (2009, January 11). Penang concerned about ‘alarming’ malaria rate. *The Star Online*. Retrieved 5 March 2011 from:
<http://www.thestar.com.my/story.aspx?file=%2f2009%2f1%2f11%2fnation%2f20090111132809&sec=nation>

Dow, R.P. and Gerrish, G.M. (1970). Day-to-day change in relative humidity and the activity of *Culex nigripalpus* (Diptera: Culicidae). *Annals of the Entomological Society of America*. 63: 995–999.

Estrada-Franco, R.G. and Craig, G.B. (1995). *Biology, disease relationship and control of Aedes albopictus*. Pan American Health Organization, Washington DC: Technical Paper No. 42.

Evenhuis, N.L. and Gon III, S. M. (2007). 22. *Family Culicidae*. In Neal L. Evenhuis Catalog of the Diptera of the Australasian and Oceanian Regions. Bishop Museum.

Fang, R., Lo, E. and Lim, T.W. (1983). *The 1982 dengue epidemic in Malaysia*. Proceedings of the International Conference on Dengue/Dengue Haemorrhagic fever September 1-3, 1982, Kuala Lumpur; T. Pang and R. Pathmanathan (eds.); University of Malaya, Kuala Lumpur, 519 p; 96-102.

Fernando, R.L., Fernando, S.S.E. and Siew-Yin Leong, A. (2001). Tropical Infectious Diseases: Epidemiology, Investigation, Diagnosis and Management. *Cambridge University Press*. p. 26.

Ferreira, M. (2005). *Spatial Diffusion Maps Of Dengue Fever Epidemics Occurring In Southeastern Brazil: A Methodology For Cartographic Modeling In GIS*. Proceeding of the International Cartographic Conference 2005, 9-16 July 2005, Coruna, Spain.

Frazier, M. R., Huey, R. B. and Berrigan, D. (2006). Population growth rates: “Warmer is better.” *American Naturalist*. 168: 512-520.

Gubler, D.J. (2002). Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic problem in the 21st century. *Trends in Microbiology* 2002. (10):100–3.

Goovaerts, P. (1997). *Geostatistics for Natural Resources Evaluation*, Oxford University Press. 483pp.

- Harbach, R. (2008). Family Culicidae Meigen, 1818. Mosquito Taxonomic Inventory. Available online at : <http://mosquito-taxonomic-inventory.info/family-culicidae-meigen-1818>. Last updated November 2, 2008.
- Hii, J.L., Chee, K.C., Vun, Y.S., Awang, J., Chin, K.H. and Kan, S.K. (1996). Sustainability of a successful malaria surveillance and treatment program in a Runggus community in Sabah, East Malaysia. *Southeast Asian Journal of Tropical Medicine and Public Health* 1996;27:512-21.
- Jamaiah, I., Anuar, A.K., Najib, N.A.R.N. and Zurainee, M.N. (1998). Imported malaria: a retrospective study in University Hospital, Kuala Lumpur, a ten-year experience. *Medical Journal of Malaysia* 1998;53:6-9.
- Jekel, J.F., Katz, D.L., Wild, D. and Elmore, J.G. (2001) *Epidemiology, Biostatistics, and Preventive Medicine* (2nd edition). Philadelphia, WB Saunders.
- Khormi, H.M., Kumar, L. and Elzahrany, R.A (2011). Modeling spatio-temporal risk changes in the incidence of dengue fever in Saudi Arabia: a Geographical Information System case study. *Geospatial Health* – 6(1): 77-84.
- Kolivras, K.N. (2006). Mosquito habitat and dengue risk potential in Hawaii: a conceptual framework and GIS application. *Professional Geographer* 58: 139-154.
- Last, J.M., Spasoff, R.A. and Harris, S.G. (2001) *A Dictionary of Epidemiology* (4th edition). Oxford, Oxford University Press.
- Lawson, A.B. and Williams, F.L.R. (2001). *An Introductory Guide to Disease Mapping*. England: John Wiley & Son, Ltd.
- Lloyd, O. and Yu, T. (1994). Disease Mapping: A valuable technique for environmental medicine. *Journal of Hong Kong Medical Association*, 46 (1): 3-15.
- McMichael, A.J., Haines, A., Slooff, R. and Kovats, S. (1996). *Climate change and human health: an assessment prepared by a task group on behalf of the World Health Organization, the World Meteorological Organization and the United Nations Environment Programme*. Redhill: Horley Studios. 333p.
- Merican, M.I., (2011). *Press Release of Director-General, Ministry of Health Malaysia for week 52 2011*. Retrieved February 2 2012 from: http://www.moh.gov.my/press_releases/94.
- Mohamed Hassan, N. (2008). *Penyakit Tropika di Negeri-negeri Melayu Bersekutu 1896-1914*. Penerbit Universiti Malaya. 144pp.
- Moore C.S. and Cheong, I.(1995). Audit of imported and domestic malaria cases at Kuala Lumpur Hospital. *British Journal of Clinical Practitioners*. 49:304-7.
- Morrison, A.C., Getis, A., Santiago, M., Rigau-Perez, J. and Reiter, P. (1998). Exploratory space–time analysis of reported dengue cases during an outbreak in Florida, Puerto Rico, 1991–1992. *American Journal of Tropical Medicine and Hygiene*. 1998;58:287–98.

Ministry of Health Malaysia (1986). *Handbook of Prevention and Control Guide for Dengue Fever/ Dengue Hemorrhagic Fever*. Kuala Lumpur: Vector-borne Disease Unit, Ministry of Health Malaysia.

Nawrocki, S.J.. and Hawley, W.A. (1987). Estimation of the northern limits of distribution of *Aedes albopictus* in North America. *Journal of American Mosquito Control Association* 1987. 3:314–317.

O’Gower, A.K. (1963). Environmental Stimuli And The Oviposition Behaviour Of *Aedes aegypti* var. *queenslandis* Theobald. *Animal Behaviour*.11(1):189–197

Palmer, K. (2002). Combating communicable diseases. Geneva: World Health Organization 2002. Retrieved 3 March 2010 from http://www.wpro.who.int/themes_focuses/theme1/focus2/tlf2malaysia.asp

Pathirana, S., Kawabata, M. and Goonetilake, R. (2009), 'Study of potential risk of dengue disease outbreak in Sri Lanka using GIS and statistical modelling', *Journal of Rural and Tropical Public Health*, vol. 8, pp. 8-17.

Poovaneswari, S. (1993). Dengue situation in Malaysia. *Malaysian Journal of Pathology* 1993. 3-7.

Provost, M.W. (1973). Mosquito flight and night relative humidity in Florida. *Florida Scientist*. 36: 217–225.

Rahman, W.A., Adanan, C.R. and Abu Hassan, A. (2002). Species composition of adult *Anopheles* populations and their breeding habitats in Hulu Perak district, Peninsular Malaysia. *Southeast Asian Journal of Tropical Medicine and Public Health*, 33 (3), 547-550.

Reid, J.A. (1954). A Preliminary *Aedes aegypti* survey. *Medical Journal of Malaya*, 9:161-168.

Richards, E.E., Masuoka, P., Major, D.B., Smith, M., Klein, T.A., Kim, H.C., Anyamba, A. and Grieco, J. (2010). The relationship between mosquito abundance and rice field density in the Republic of Korea. *International Journal of Health Geographic* 9: 32-42.

Richardson, K., Hoffmann, A.A., Johnson, P., Ritchie, S. and Kearney, M.R. (2011). Thermal Sensitivity of *Aedes aegypti* from Australia: Empirical Data and Prediction of Effects on Distribution. *Journal of Medical Entomology*, 48(4):914-923. 2011.

Rudnick, A. (1978). Ecology of dengue virus. *Asian Journal of Infectious Diseases*. 2:156-160.

Rudolfs, W. (1925). Relation between Temperature, Humidity and Activity of House Mosquitoes. *Journal of the New York Entomological Society*, 33(3): 163-169.

Selangor State Investment Centre (2005). Opportunities of investment in Selangor. Retrieved 3 March 2010 from <http://www.ssic.com.my>

Shaharudin, I., Shamsul, A., Tahir, A., Mariam, M., Azah, D. and Nik Shamsidah, N. (2001). Geographical Information Systems (GIS) and the Public Health Sector: Dengue Fever Studies in Bandar Baru Bangi and Kajang. *Malaysian Journal of Community Health, Special Issue*: 34-42.

Sidhu, P.S. and Ng, S.C. (1991). A retrospective study on malaria cases admitted to the University Hospital, Kuala Lumpur, 1984-1988. *Medical Journal Malaysia*. 46:177-82.

Skae, F.M.T. (1901). Dengue Fever in Penang. *British Medical Journal*, 2:1581-1582.

Smith, C.E.G. (1956). A localized outbreak of dengue fever in Kuala Lumpur: epidemiological and clinical aspects. *Medical Journal of Malaya*, 10(4):289-303.

Stanton, A.T. 1919. The mosquitoes of Far Eastern ports with special reference to the prevalence of *Stegomyia fasciata*, *Bulletin of Entomological Research*. 10:333-344.

Teng H.J. and Apperson, C.S. (2000). Development and survival of immature *Aedes albopictus* and *Aedes triseriatus* (Diptera: Culicidae) in the laboratory: Effects of density, food, and competition in response to temperature. *Journal of Medical Entomology* 2000;37:40-52.

Tran, A., Deparis, X., Dussart, P., Morvan, J., Rabarison, P. and Remy, F. (2004). Dengue spatial and temporal patterns, French Guiana. *Emerging Infectious Diseases* 2004;10:615-21.

Tropical Disease Research, WHO Special Programme for Research and Training in Tropical Diseases (2005). *Scaling up home-based management of malaria: From research to implementation*. WHO, Geneva. 96pp.

Troyo, A., Fuller, D.O., Arguedas, O.C., Solano, M.E. and Beier, J.C. (2009). Urban structure and dengue fever in Puntarenas, Costa Rica. *Singapore Journal of Tropical Geography* 30(2): 265-282.

Wan Fairos, W. Y., Wan Azaki, W. H., Mohamad Alias, L. and Y. Bee Wah (2010). Modeling Dengue Fever (DF) and Dengue Haemorrhagic Fever (DHF) Outbreak Using

Poisson and Negative Binomial Model. *World Academy of Science, Engineering and Technology* 38 2010.

Watson, M. (1903) *The Prevention of Malaria in the Federated Malay States*. 381pp.

Whitehorn, J. and Farrar, J. (2010). "Dengue". *British Medical Bulletin*. 95: 161-73

World Health Organization (1993). Global Alert and Responses for Dengue Fever. Retrieved from <http://www.who.int/csr/disease/dengue/en/index.html> on 3 March 2010

World Health Organization (2001). Review of Roll-Back Malaria Strategies in the South-East Asia Region. Proceedings from Report of an Intercountry Consultative Meeting Yogyakarta, Indonesia, 6-10 November 2001.

World Health Organization (2002). Dengue/DHF—global public health burden. *Weekly Epidemiology Records* 2002;77:300–4. Retrieved 3 March 2010 from <http://www.who.int/denguenet>.

World Health Organization (2008). *World Malaria Report 2008* (Report). ISBN 978-92-4-156369-7.

Yap, H.H. and Thiruvengadam, V. (1979). Relative abundance of *Aedes aegypti* (Linnaeus) and *Aedes albopictus* (Skuse) in different habitats. *Medical Journal of Malaysia*. 34: 76-79.



© COPYRIGHT UPM

LIST OF PUBLICATIONS

1. Hafiz, H; Hashim, N.R., Shohaimi, S. (2011). *Spatial and temporal distribution of dengue fever in Selangor and Kuala Lumpur 2009-2010*. Journal of Community Health, Volume 1 (Supplement 1) 2011.
2. Hafiz, H; Hashim, N.R., Shohaimi, S. (2012). *Risk Mapping of Dengue Fever in Selangor and Kuala Lumpur, Malaysia*. Geospatial Health 7(1), 2012, pp. 21-25.
3. Hafiz, H; Hashim, N.R. (2012). *Spatial and Temporal Distribution of Malaria in Peninsular Malaysia from 1998-2010*. Health and The Environment Journal, 3(3), 2012.
4. Hafiz, H; Hashim, N.R., Shohaimi, S. (2013). *Centrography of Dengue and Malaria in Selangor and Kuala Lumpur*. Poster presentation. Fundamental Science Congress 2013, Universiti Putra Malaysia.