

UNIVERSITI PUTRA MALAYSIA

ESTIMATING CARBON STORAGE OF FOREST FLOOR COMPONENTS AT VARYING ALTITUDES IN TROPICAL FOREST OF PAHANG, MALAYSIA

V. JEYANNY VIJAYANATHAN

FP 2015 91



ESTIMATING CARBON STORAGE OF FOREST FLOOR COMPONENTS AT VARYING ALTITUDES IN TROPICAL FOREST OF PAHANG, MALAYSIA

By

V. JEYANNY VIJAYANATHAN

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfillment of the Requirement for the Degree of Doctor of Philosophy

December 2014


All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

Specially dedicated to all my pillars of strength,
my beloved family, Mr. Gerard Felix, Ms. Juliyana and Michael Andrew
For their constant encouragement and inspiration.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

ESTIMATING CARBON STORAGE OF FOREST FLOOR COMPONENTS AT VARYING ALTITUDES IN TROPICAL FOREST OF PAHANG, MALAYSIA

By

V. JEYANNY VIJAYANATHAN

December 2014

Chairman : Associate Professor Ahmad Husni Mohd. Hanif, PhD

Faculty : Agriculture

Increasing atmospheric carbon dioxide (CO₂) concentrations at alarming rates has triggered the need to conserve and monitor carbon (C) stocks for climate change mitigation. Tropical forests are important carbon sinks which are dynamic due to topographic variations, biomass components, forest floor quality, decomposition processes and spatial variation. Precise and reliable estimation of C stocks and its confounding processes that release/store C are still absent in tropical montane and lowland forests in Malaysia. The objectives of this study were i) to quantify above and belowground biomass C stocks in a lowland forest and montane forest with varying topography; ii) to determine the potential indicators (i.e. litter and duff decomposition rate, forest floor component and properties, soil CO2 fluxes and C:N ratios) of soil C storage and iii) to determine the spatial variability of litter, soil C, C:N, and forest floor component depths of a tropical lowland forest and tropical montane forest with varying topography. A systematic design of 10 m x 10 m plots was established for soil (0- 15 cm depth), litter and aboveground biomass sampling along three slope positions at the montane forest and one plot in the lowland forest. Basic soil characteristics and botanical distribution were determined. A litter bag study and soil CO₂ flux measurements were conducted for 480 days in the montane and lowland forest.

Forest floor materials were carbon dated and segregated for precise bulk density and carbon fraction measurements. Soil C, C:N, litter depth and various decomposing layers were explored using geostatistics to determine spatial variability. Litter and soil carbon stocks were significantly higher (3 and 5-fold) in the montane forest compared to the lowlands. The aboveground biomass ranged from 100 to 120 Mg C ha⁻¹ and was the most dominant pool (> 40%) for all sites. The decomposition decay rate constant, k ranged from -0.002 to -0.004 day⁻¹ for the tropical montane and lowland forest. Lowlands showed increased mass loss and significant linear regression relationships between mass loss and litter quality except for C, lignin and cellulose. Soil CO₂ fluxes were higher in the lowlands and positively correlated with decomposition and water filled pore space (WFPS). Duff (hemic + sapric) segregation resulted higher bulk density values (0.2) compared to litter (0.04) and revealed a more precise carbon fraction for litter (0.43) and duff (0.55) to be used for forest floor C stocks predictions utilizing significant linear regressions. Duff may reside up to 60 years. Duff decomposition is impeded even at higher temperatures in Forest Research Institute Malaysia campus and retained most of its carbon, nitrogen and lignin. Relationship between soil C:N and C were strong for all plots. Soil total C, C:N, and litter depth exhibited spatial variability at both forest types. Similarly, the litter, hemic and the total forest floor depth fractions confirmed spatial variations. Most variables exhibited a strong spatial dependence with the exception of C:N at the sideslope, litter depth at Jengka VJR and hemic depth at the summit (moderate). Surface maps for total C, C:N, litter depths, hemic and total forest floor depth showed distinct spatial clustering and displayed acceptable accuracy of interpolated values. Forest floor in the montane forest acts as an important C stock in the tropical forest which needs to accounted precisely for national C accounting. Research and developments in monitoring C stocks in montane forests within national and regional areas must be explored to avoid undestimation.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

ANGGARAN STOK KARBON DALAM KOMPONEN LAPISAN ORGANIK PADA KEDUDUKAN CERUN BERLAINAN DI HUTAN TROPIKA PAHANG, MALAYSIA

Oleh

V. JEYANNY VIJAYANATHAN

Disember 2014

Pengerusi : Profesor Madya Ahmad Husni Mohd. Hanif, PhD

Fakulti : Pertanian

Peningkatan drastik dalam kepekatan gas karbon dioksida di atmosfera telah mencetuskan keperluan untuk memelihara dan memantau stok karbon (C) untuk mitigasi perubahan iklim. Hutan tropika merupakan stok karbon yang penting dan dinamik kerana variasi topografi, komponen biomas, kualiti lapisan organik, proses penguraian, dan variasi ruang. Penganggaran stok karbon yang tepat dan persis serta proses-proses yang berkaitan dengan pembebasan dan penyimpanan karbon masih tidak wujud di hutan tanah gunung dan hutan tanah pamah di Malaysia. Objektif kajian projek ini adalah i) untuk menentukan stok biomas karbon diatas dan dibawah tanah di hutan tanah pamah dan hutan gunung dengan kecerunanan berlainan; ii) menentukan indikator-indikator yang berpotensi (kadar penguraian daun luruhan organik dan 'duff'; ciri-ciri lapisan organik, fluks CO2 tanah, nisbah C:N) dalam penyimpanan stok karbon dan iii) menentukan variasi ruang ketebalan daun luruhan organik, nilai C tanah, nisbah C:N dan ketebalan lapisan organik hutan gunung dengan kecerunan berlainan dan hutan tanah pamah. Kuadrat bersaiz 10 x 10 m persegi telah ditubuhkan secara sistematik untuk persampelan tanah (kedalaman 0 hingga 15 cm), daun luruhan organik dan biomas di atas tanah di tiga kedudukan cerun di hutan gunung dan satu plot di hutan tanah pamah.

Ciri-ciri tanah dan taburan am vegetasi telah ditentukan. Kajian 'litterbag' dan penentuan fluks CO2 tanah telah dijalankan untuk 480 hari di hutan gunung dan hutan tanah pamah. Bahan lapisan organik telah ditentukan usia karbon dan dipisahkan untuk mendapatkan ketumpatan pukal dan pecahan karbon. Nilai karbon tanah, C:N dan ketebalan daun luruhan serta lapisan organik pada tahap penguraian yang berbeza telah dianalisis menggunakan kaedah geostatistik untuk penentuan variasi ruang. Hasil kajian mendapati stok karbon daun luruhan organik dan tanah adalah 3 dan 5 kali lebih tinggi secara signifikan di hutan gunung berbanding hutan tanah pamah. Kandungan stok karbon biomas pokok adalah dalam lingkungan 100 -120 Mg C ha⁻¹ di hutan gunung dan merupakan komponen paling dominan (> 40 %) berbanding plot lain. Pemboleh ubah malar kadar penguraian bahan organik, k, adalah dalam lingkungan -0.002 – 0.004 day⁻¹ untuk hutan gunung dan hutan tanah pamah. Hutan tanah pamah menunjukkan kehilangan berat yang tinggi dan hubungan regresi linear yang signifikan di antara kehilangan berat dan kualiti daun luruhan organik kecuali C, kandungan lignin dan sellulosa daun. Fluks CO2 tanah adalah lebih tinggi di hutan tanah pamah dan mempunyai korelasi yang positif dengan penguraian daun luruhan organik dan 'water filled pore space' (WFPS). Pengasingan duff (hemic + sapric) menunjukkan ketumpatan pukal yang lebih tinggi (0.2) berbanding daun luruhan organik (0.04) serta menunjukkan pecahan karbon yang lebih persis iaitu 0.43 untuk daun luruhan organik dan 0.55 untuk duff bagi penentuan ramalan stok C dalam lapisan organik permukaan tanah menggunakan hubungan regresi linear yang signifikan. Duff dapat bertahan selama 60 tahun. Penguraian duff adalah lambat walaupun dengan suhu tinggi di kampus Institut Penyelidikan Perhutanan Malaysia (FRIM) dan menyimpan kebanyakan unsur karbon, nitrogen dan lignin. Hubungan regresi yang signifikan didapati antara nisbah C:N tanah dan C untuk semua plot kajian. Analisis geostatistik menunjukkan variasi struktur ruang (spatial) untuk pemboleh ubah nilai C tanah, nisbah C:N dan ketebalan daun luruhan di kedua-dua jenis hutan. Pengasingan pecahan daun luruhan, hemic dan keseluruhan lapisan organik juga mengesahkan variasi ruang yang signifikan. Kebanyakan pemboleh ubah menunjukkan ianya bergantung secara kuat kepada variasi ruang, kecuali nisbah C:N di permukaan landai bukit, ketebalan daun luruhan di Jengka VJR dan ketebalan hemic di puncak bukit yang bergantung secara sederhana sahaja. Peta permukaan menunjukkan kelompok ruang yang berbeza dan memaparkan nilai interpolasi yang tepat untuk C, C:N, ketebalan daun luruhan, *hemic* dan ketebalan keseluruhan lapisan organik. Lapisan organik di hutan gunung merupakan komponen stok C yang penting di hutan tropika yang perlu dikaji secara tepat untuk penentuan stok karbon negara. Usaha penyelidikan dan pembangunan dalam pemantauan stok karbon hutan gunung di negara dan rantau Asia perlu diterokai untuk mengelakkan kesalahan dalam pengganggaran.

ACKNOWLEDGEMENTS

The author would first extend her grace and gratitude to the Lord Almighty, who gave the courage and endurance to make all things possible.

The author is also greatly indebted to Associate Professor, Dr. Ahmad Husni Mohd. Hanif, the Chairman of the Supervisory Committee, for his tireless guidance and constant support throughout the course of her research and the preparation of this thesis. The willing assistance provided by the members of the Supervisory Committee, Dr. Siva Kumar Balasundram, Dr. Wan Rasidah Kadir and Dr. Arifin Abdu cannot be forgotten. Their constructive criticisms were very crucial in her completion of the thesis.

The author also wishes to express her gratitude to the Director General of Forest Research Insitute Malaysia (FRIM), Dato' Dr. Abdul Latif Bin Mohmod for granting permission and rendering support throughout the course of her PhD, by providing infrastructure, manpower and leave. She is also grateful to the Department of Forestry, Pahang which provided permission and assistance during the research plot selection.

The author is greatly indebted to Ministry of Science, Technology and Innovation (MOSTI) for granting study leave and providing financial allowances during her PhD training. The financial contribution provided through the E- Science fund by the Ministry of Agriculture (MOA) is also acknowledged for running the project.

Her special thanks goes to Mr. Muhammad Firdaus Sulaiman and Mr. Kamarul Hisham Mustafa for assistance provided in field and laboratory during soil CO₂ flux measurements and botanical description, respectively. Her sincere appreciation is also addressed to the entire staff of the Soil Management Department of FRIM, with special reference to Mr. Suhaimi Wan Chik and Miss. Hasnidarul Ahmad for their help during the running of the field experimentation and data collection. The author also acknowledges the support and advice given by Miss Law Mei Ching, Miss Lee Yit Leng, and Mr. Sudharsan Selvarajah during laboratory instrumentations and geostatistical analysis troubleshooting. Their timely aid during the research is deeply appreciated.

Last but not least, her sincere appreciation is reserved for her family's kindness, Mr. Gerard Felix Louis, her husband, Ms. Juliyana Savarimuthu Lazarus, her mother in law and her son, Master Michael Andrew Gerard for their faith and patience throughout her responsibility as a PhD student. Their undivided love and sacrifice cannot be measured.

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Ahmad Husni Mohd. Hanif, PhD

Associate Professor Faculty of Agriculture Universiti Putra Malaysia (Chairman)

Siva Kumar Balasundram, PhD

Associate Professor Faculty of Agriculture Universiti Putra Malaysia (Member)

Wan Rasidah Kadir, PhD

Senior Research Officer Soil Management Branch Forest Research Institute Malaysia (Member)

Arifin Abdu, PhD

Associate Professor Faculty of Forestry Universiti Putra Malaysia (Member)

BUJANG KIM HUAT, PhD

Professor/Dean School of Graduate Studies Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature:	Date:	
<i></i>		

Name and Matric No.: V. Jeyanny Vijayanathan (GS 27229)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature:	Signature:
Name of	Name of
Chairman of	Member of
Supervisory	Supervisory
Committee:	Committee:
Signature:	Signature:
Name of	Name of
Member of	Member of
Supervisory	Supervisory
Committee:	Committee:

TABLE OF CONTENTS

ARST	ГКАСТ	Page
ABST		iii
	NOWLEDGEMENTS	v
	ROVAL	vii
	LARATION	ix
	OF TABLES	xiv
	OF FIGURES	xvi
	OF ABBREVIATIONS	xix
OTT 1		
CHA	PTER	
I.	INTRODUCTION	1
II.	LITERATURE REVIEW	5
	Climate change and carbon in tropical forest biomass	5
	Causes of climate change	5
	Impacts of climate change	6
	Approaches for climate change mitigation	8
	Role of C in carbon balance management	8
	World forest biomass statistics and carbon	9
	Malaysia forest statistics	10
	Carbon sequestration	12
	Carbon sequestration in various components	12
	Role of soil in carbon sequestration	13
	Factors controlling soil carbon sequestration	14
	Indicators for soil carbon sequestration	17
	Soil carbon and nitrogen	17
	C:N	17
	Forest floor component	17
	Spatial variability	20
	Precision forestry (PF)	20
	Precision forestry tools	20
	Mechanics of geostatistics	21
	Application of geostatistics in CS	25
	Summary	26

26

III.	QUANTIFICATION OF SOIL, LITTER, ABOVE AND BELOW	
	GROUND BIOMASS CARBON STOCKS	
	Introduction	27
	Material and Methods	28
	Exp. 1: Determination of litter, soil, roots and aboveground	
	biomass C stocks in a lowland forest and a montane forest	
	with varying topography.	
	Results	36
	Discussion	41
	Conclusions and Recommendations	42
TX 7		
IV.	SOIL AND FOREST FLOOR PROPERTIES AS POTENTIAL	12
	INDICATORS FOR CARBON SEQUESTRATION Introduction	43 43
	Materials and Methods	45
		40
	Exp. 2(i): Litter decomposition and soil respiration of tropical lowland and montane forest	
	Exp. 2(ii): Contribution of duff properties and decomposition	
	at lowland temperatures towards C management	
	Exp.2(iii): Determination of soil C:N of 0-15 cm soil depth	
	Results	56
	Litter decomposition and soil respiration in	50
	Sungai Kial FR and Jengka VJR	56
	Duff dynamics	67
	Soil C:N	84
	Discussion	86
	Litter decomposition and soil respiration in	
	Sungai Kial FR and Jengka VJR	86
	Duff dynamics	89
	Soil C:N	95
	Conclusion	96

٧.	SPATIAL VARIABILITY OF SOIL CAND FOREST FLOOR	
	COMPONENTS DEPTH FOR CARBON SEQUESTRATION	
	MANAGEMENT	97
	Introduction	97
	Material and Methods	98
	Results	101
	Comparison of C, C:N and forest floor depth	
	at different sites	101
	Comparison of different components of forest	
	floor depth according to slope types	113
	Discussion	125
	Comparison of C, C:N and forest floor depth	
	at different sites	125
	Comparison of different components of forest floor depth	120
	according to slope types	127
	Conclusion	130
	Conclusion	130
VI.	SUMMARY, GENERAL CONCLUSION AND	
V 1.		
	RECOMMENDATIONS FOR FUTURE	121
	RESEARCH	131
	FERENCES	133
	PENDICES	156
	DATA OF STUDENT	173
LIS	T OF PUBLICATIONS	174

LIST OF TABLES

Table		Page
2.1	Major forest types and soil regimes in Peninsular Malaysia in 1990	11
2.2	Common allometric equations for above ground biomass of tropical forest	13
2.3	Organic Carbon (C) loss in soil erosion from tropical lands	15
3.1	Basic information on environmental conditions at study sites	31
3.2	Soil physio-chemical properties of the soil profile at study sites	37
3.3	Brief botanical description of vegetation in Sungai Kial FR and Jengka FR	38
3.4	Soil bulk density and carbon stocks estimates for Sungai Kial FR and Jengka VJR	39
4.1	Crossvalidation of litter decay models	50
4.2	Initial nutrient concentration of leaf litter analysis	58
4.3	Changes in litter nutrient content during decomposition, 120-480 days	61
4.4	Pearson linear correlation coefficients (r) between mass loss and litter quality from 0 to 480 days	63
4.5	Soil bulk density, soil temperature and relative humidity of study area, $n = 6$	64
4.6	Descriptive statistics for bulk density (BD, g cm ⁻³) of segregated forest floor layers	68
4.7	Descriptive statistics for carbon fraction (CF, C%/100) of segregated forest floor layers	69
4.8	Carbon dating results of hemic and sapric from the summit plot,	75

4.9	Initial concentration of litter and duff analysis	77
4.10	Crossvalidation of forest floor decay models in FRIM campus	78
4.11	Changes in tissue content during decomposition, 60 to 330 days	81
4.12	Pearson linear correlation coefficients (r) between mass loss and forest floor quality from 0 to 330 days	83
5.1	Classification of spatial dependence	100
5.2	Total C, C:N and forest floor depth of Sungai Kial FR and Jengka VJR	101
5.3	Descriptive statistics for selected soil properties in Sungai Kial FR and Jengka VJR	103
5.4	Cross validation statistics of kriged values for variables	110
5.5	Litter, hemic, sapric, and TFF of Sungai Kial FR	113
5.6	Descriptive statistics for selected forest floor components in Sungai Kial FR	115
5.7	Cross validation statistics of kriged values at	120

LIST OF FIGURES

Figure		Page
2.1	Projection of increased temperature anomaly since 1850	6
2.2	Principal global C pools and fluxes between them	9
2.3	The generalized semivariogram model	22
3.1	Location of study sites in the state on Pahang, Malaysia	29
3.2	10 m x 10 m plot design	30
3.3	Annual mean air temperature and rainfall at Sungai Kial Forest Reserve from 2009 to 2012	32
3.4	Annual mean air temperature and rainfall at Jengka VJR from 2009 to 2012	33
3.5	Carbon stocks in various components of tropical montane and lowland forest (% Mg ha ⁻¹)	40
4.1	Processes involved in AMS dating for sapric and hemic material	54
4.2	Changes of mean temperature and rainfall from July 2011 to November 2012 in Sg. Kial FR	56
4.3	Changes of mean temperature and rainfall from July 2011 to November 2012 in Jengka VJR	57
4.4	Trend of litter remaining (%) against time (days) at Sungai Kial FR and Jengka VJR	60
4.5	Soil CO ₂ flux measurements at summit and toeslope of Sungai Kial FR and Jengka VJR.	65
4.6	Soil water filled pore space measurements at summit and toeslope of Sungai Kial FR and Jengka VIR	66

4.7	Soil air filled pore space measurements at summit and toeslope of Sungai Kial FR and Jengka VJR	66
4.8	Organic material profile at toeslope, Sungai Kial FR	67
4.9	Regression between bulk density (g cm ⁻³) against litter (a) and hemic (b) C stocks (Mg C ha ⁻¹) and their respective predicted versus calculated C stocks (c,d)	70
4.10	Regression between bulk density (g cm ⁻³) against sapric (a) and duff (b) C stocks (Mg C ha ⁻¹) and their respective predicted versus calculated C stocks (c,d)	71
4.11	Regression between bulk density (g cm ⁻³) against total and forest floor C stocks (Mg C ha ⁻¹) and their respective predicted versus calculated C stocks	72
4.12	Mycorrhizal root infection (%) of hemic in the toeslope and summit of Sungai Kial FR	73
4.13	Mycorrhizal spore count (per 10 g material) of hemic in the toeslope and summit of Sungai Kial FR	74
4.14	Meteorological data of rainfall and mean temperature for FRIM	76
4.15	Trend of forest floor remaining (%) against time (days) at FRIM campus, Selangor	79
4.16	Pearson correlation coefficient of Total C against C:N in Sungai Kial FR and Jengka VJR	85

5.1	Spatial structure and attributes of selected soil variables at the summit and sideslope of Sungai Kial FR	105
5.2	Spatial structure and attributes of selected soil variables at the sideslope and toeslope of Sungai Kial FR	106
5.3	Spatial structure and attributes of selected soil variables at Jengka VJR.	107
5.4	Spatial variability of C and C:N at the summit and sideslope of Sungai Kial FR	109
5.5	Spatial variability of litter, C and C:N at the sideslope and toeslope of Sungai Kial FR	111
5.6	Spatial variability of C and C:N at Jengka VJR	112
5.7	Spatial structure and attributes of litter, hemic and TFF depth at the summit, Sungai Kial FR	115
5.8	Spatial structure and attributes of litter, hemic and TFF depth at the sideslope, Sungai Kial FR	116
5.9	Spatial structure and attributes of litter, hemic and TFF depth at the toeslope, Sungai Kial FR	118
5.10	Spatial variability of litter, hemic and TFF at the summit of Sungai Kial FR	119
5.11	Spatial variability of litter, hemic and TFF at the sideslope of Sungai Kial FR	123
5.12	Spatial variability of litter, hemic and TFF at the	124

LIST OF ABBREVIATIONS

ABD Above Ground Biomass

AFPC Air Filled Pore Space

AGB Above Ground Biomass

AMF Arbuscular Mycorrhizal Fungi

AMS Accelerated Mass Spectrometry

ANOVA Analysis of Variance

ATP Adenosine Triphosphate

BD Bulk Density

C Carbon

C:N Carbon: Nitrogen

C:P Carbon: Phosphorus

Ca Calcium

CDM Clean Development Mechanism

CF Carbon Fraction

CNS Carbon: Nitrogen: Sulfur

CS Carbon Sequestration

CTFS Centre for Tropical Forest Science

CV Coefficient of Variation

CWD Coarse Woody Debris

DBH Diameter at Breast Height

DNA Deoxyribonucleic acid

DWD Downed Woody Debris

EDA Exploratory Data Analysis

ENSO El Nino Southern Oscillation

ER Effective Range

ESD Extreme Studentized Deviated

FAO Food Aid Organization

FFC Forest Floor Carbon

FR Forest Reserve

FRIM Forest Research Institute Malaysia

GCM Global Circulation Model

GHG Greenhouse Gases

GIS Geographic Information System

GPG Good Practice Guidance

GPS Global Positioning System

Gt Giga tonnes

HCL Hydrochloric Acid

IDW Inverse Distance Weighting

IPCC Intergovernmental Panel on Climate Change

K Kalium/Potassium

LULUCF Land Use and Land Use Chang and Forestry

ME Mean Error

Mg Mega gram (10⁶)

Mn Manganese

MOA Ministry of Agriculture

MRT Mean Residence Time

MSE Mean Square Error

N Nitrogen

NaOH Sodium Hydroxide NH₄OAc Ammonium Acetate

NH₄OAc Ammonium Acetate

NPP Net Primary Production

NSCCC National Steering Committee on Climate Change

NW North West

Oa Sapric (organic)

Oe Hemic (organic)

Oi Fibric (organic)

OM Organic Matter

P Phosphorus

PA Precision Agriculture

Pg C Pentagram (10¹⁵)

PMC Percent Modern Carbon

PRECIS Providing Regional Climates for Impacts Studies

PVC Poly Vinyl Chloride

REDD Reducing Emissions from Deforestation and Forest

Degradation in Developing Countries

RNA Ribonucleic acid

SE South East

SIC Soil Inorganic Carbon

SOC Soil Organic Carbon

SMSE Standardized Mean Squared Error

SOM Soil Organic Matter

SOS Soil Organic Carbon

SW South West

TFF Total Forest Floor

UNFCCC United National Framework Convention On Climate Change

UPM Universiti Putra Malaysia

VJR Virgin Jungle Reserve

WFPS Water Filled Pore Space



CHAPTER I

INTRODUCTION

The progressive global industrial and agricultural development has caused increased carbon dioxide (CO_2) emissions. The concentration of CO_2 in the atmosphere has been reported to soar up to 391 ppm in 2011 (IPCC, 2013) compared to 280 ppm in the 1800 (Pitelka & Rojas 2001). According to the Intergovernmental Panel on Climate Change (IPCC), this will simultaneously increases Earth's temperature between 0.6 to 1.1° C over the period of 1880 to 2012. In order to reduce the Greenhouse Gases (GHG) in the atmosphere, two key activities are relevant: decrease the anthropogenic emissions of CO_2 and create or promote carbon (C) sinks in the biosphere.

Tropical forests clearly dominate the role of forests both on C flux, and the volume of C stored. Tropical soils constitute 26% of the soil organic carbon (SOC) in the world (Batjes, 1996). Worldwide, SOC in the top one meter of soil comprises about 3/4 of the earth's terrestrial carbon and has tremendous potential to sequester additional carbon in soil (Soil Science Society of America, 2009). Malaysia had voluntarily agreed to CO₂ emission reductions, increased carbon sinks and promoting research on carbon cycling in the Kyoto Protocol (Soderstrom, 2002). During the Bali Convention on Climate Change (2007), Malaysia being part of the 80% of the world's tropical rainforest has further reaffirmed its stand by showing interest in obtaining funds to reduce emissions from deforestation. At present, the total forest cover of Malaysia stands at 60% (excluding rubber) [FAO, 2010].

However, the amount of C which can be sequestered in native ecosystem types are limited. Regional estimation which is currently used may not reflect the actual values for Malaysia in the Land Use and Land Use Change and Forestry (LULUCF). Moreover, definitive quantification of various carbon stocks, especially in aboveground and belowground biomass has not be delineated successfully due to dynamic ecological systems, budget constraints, and lack of collaboration in the research and development of climate change related issues.

Soil organic matter (SOM) accumulation influences the SOC stocks in soil. Other indicators that may also influence the C stock in forests are C:N ratios, moisture, temperature, topography, elevation, vegetation, forest floor components, litter decomposition and spatial variability. Changes in SOM decomposition rates at different altitudes was influenced by temperature (Wang et al., 2009) and soil moisture (Reynolds, 2000). Therefore, it is believed that increase in soil C stocks was significantly associated with elevation and temperature (Leifeld et al., 2005). In subtropical and temperate forests, the stocks of SOC were found to be decreasing with altitude, from 185.6 (1600-1800 m) to 160.8 t C ha⁻¹ (2000-2200 m) and from 141.6 (600-800 m) to 124.8 tC ha⁻¹ (1000- 1200 m) respectively (Sheikh et al., 2009).

Research in quantifying carbon stocks in the Malaysian region according to topographical variations is scarce. Earlier work by Kitayama and Aiba (2002) on sedimentary substrate interspersed with ultrabasic rock of Mount Kinabalu, Sabah, and presented evidence that SOC concentrations were increased with increasing elevations of 700 – 3100 masl. Lim (2002) whereas estimated the highest amount of SOC in Malaysia were found in the highlands (300-1200 masl), followed by agricultural land between 20-100 masl. His values were derived from different soil profiles of agricultural areas in Peninsular Malaysia. However, both studies did not account the landform segmentation (i.e summit, sideslope and toeslope) variations and the influence of parent material type which were present in the tropical forest of the Peninsular Malaysia region.

Generally, most research agrees that temperature, precipitation, litter chemistry (Adair et al., 2008), soil nutrient content and microbial communities (Attignon et al., 2004) strongly control rates of litter decomposition. However, how these factors independently or interactively influence decomposition across large spatial and temporal scales remains unclear. Wang et al., (2009) reported that mean mass loss of litter was positively correlated with mean annual temperature along elevational gradient in the Wuyi mountains, China. In another study, temperature effects were evident when comparing the decomposition rates of pine needle litter and the forest floor (Prescott et al., 2004). In tropical forests, the decomposition coefficient (k) of leaf litter ranges from 0.1 to 0.5 per month (He et al., 2009).

C:N influences the decomposition rate of organic matter, which partly controls the mineralisation or immobilization of soil nitrogen (N). Carbon: nitrogen ratio were significantly influenced by altitude, parent material, and their interactions (Wagai, 2008).

The thickness of forest floor is usually influenced by climate, vegetation, time and faunal communities (Bens et al., 2006). Although litter decomposition studies in the Malaysian tropical forest is quite common, segregation of various components of the forest floor (i.e duff, litter, and DWD) in providing accurate estimates of carbon densities for tropical forests, especially in a tropical montane forest has not been initiated. Separated data acquired via this exercise would provide accurate duff and litter bulk density values for use in calculating mass and carbon, and the influence of different components in CS in mineral soil. Understanding SOC and its associations (C:N and forest floor) will provide useful insights in forest CS and quantification of C stocks at varying topography will be vital.

Geostatistics, a branch from geographic information system (GIS) is a tool that has been successfully used in tropical soil studies (Couto et al., 1997). Soil organic carbon (SOC) is unevenly distributed in forest and agricultural soils (Grove et al., 2008). Spatial patterns of SOC have been computed using geostatistics (Van Meirvenne et al., 1996; Frogbrook and Oliver, 2001). Besides SOC, it will be vital to quantify spatial variability of C:N and forest floor depth variations to determine the C stocks in soil. These innovative methods will greatly reduce human resource requirements, costs, and allow forest managers to strategize management zoning in order to avoid areas which are significantly potential in sequestering carbon. Based on the current knowledge gaps and scarce information available on the various factors as mentioned above, the dissertation mainly investigates the various components of C stocks in the tropical lowland and montane forest in Chapter II. Chapter III identifies potential indicators that are related to soil C storage whereas Chapter IV determines the spatial variability of forest floor material and its depths, soil C and C:N in a tropical lowland and montane forest with varying topography.

Main objectives

To quantify soil C, C:N, aboveground biomass, forest floor components (depths and layers of decomposing materials) and decomposition at varying topographic positions in a tropical montane forest and in a lowland forest that could assist in quantifying C stocks related to C management.

Sub objectives

Experiment 1: To quantify litter, soil (0-15 cm depth), roots and

aboveground biomass C stocks in a tropical lowland forest

and tropical montane forest with varying topography.

Experiment 2: To determine the potential indicators (i.e. litter and duff

decomposition rate, forest floor component and properties,

soil CO₂ fluxes, C:N ratios) of soil C storage.

Experiment 3: To determine the spatial variability of litter and soil C, C:N,

and various depths of forest floor component of a tropical lowland forest and tropical montane forest with varying

topography.

REFERENCES

- Adachi, M., Bekku, Y.S., Rashidah, W., Okuda, T., and Koizumi, H. 2006. Differences in soil respiration between different tropical ecosystems. *Applied Soil Ecology* 34(2): 258-265.
- Adair, E.C., Parton, W.J., Del Grosso, S.J., Silver, W.L., Harmon, M.E., Hall, S.A., Burke, I. C.and Hart, S.C. 2008. Simple three-pool model accurately describes patterns of long-term litter decomposition in diverse climates. *Global Change Biology* 14(11): 2636-2660
- Adams, A., Harrison, R., Sletten, R., Strahm, B., Turnblom, E., and Jensen, C. 2005. Nitrogen-fertilization impacts on carbon sequestration and flux in managed coastal Douglas-fir stands of the Pacific Northwest. *Forest Ecology and Management* 220(1): 313-325.
- Ahmed, O.H., Hasbullah, N.A. and Muhamad, A. 2010. Accumulation of soil carbon and phosphorus contents of a rehabilitated forest. *The Scientific World Journal* 10: 1988-1995
- Alexander, I. and Lee, S.S. 2005. Mycorrhizas and ecosystem processes in tropical rain forest:Implications for diversity.In *Biotic interactions in the tropics:Their role in the maintenance of species diversity*, ed. D.F.R. Burslem, P. Pinard, M. A. Hartley, pp. 165-203. Cambridge: Cambridge University Press
- Andriesse, J. 1988. *Nature and management of tropical peat soils*: FAO Soils Bulletin 59. Rome: Food and Agriculture Organization of the United Nations.
- Arnold, J., Corre, M.D., and Veldkamp, E. 2009. Soil N cycling in old-growth forests across an Andosol toposequence in Ecuador. *Forest Ecology and Management* 257(10):2079-2087.
- Attignon, S.E., Weibel, D., Lachat, T., Sinsin, B., Nagel, P. and Peveling, R. 2004. Leaf litter breakdown in natural and plantation forests of the Lama forest reserve in Benin. *Applied Soil Ecology* 27(2): 109-124.
- Austin A.T. and Vitousek P.M. 2000. Precipitation, decomposition and litter decomposability of *Metrosideros polymorpha* in native forests on Hawai'i. *Journal of Ecology* 88:129–138.

- Austin, A.T. and Ballare, C.L. 2010. Dual role of lignin in plant litter decomposition in terrestrial ecosystems. *Proceedings of the National Academy of Sciences* 107(10): 4618-4622.
- Averill, C., Turner, B.L., Finzi, A.C. 2014. Mycorrhiza-mediated competition between plants and decomposers drives soil carbon storage. *Nature* 505:543–545
- Aznar, J.C., Paucar-Munoz, H., Richer-Lafleche, M., and Begin, Y. 2010. Field litter thickness assessed by gamma-ray spectrometry. *Forest Ecology and Management* 260(10): 1640-1645.
- Baggie, I., Rowell, D., Robinson, J., and Warren, G. 2005. Decomposition and phosphorus release from organic residues as affected by residue quality and added inorganic phosphorus. *Agroforestry systems* 63(2): 125-131.
- Balasundram, S., Robert, P., Mulla, D., and Allan, D. 2006. Spatial variability of soil fertility variables influencing yield in oil palm (*Elaeis guineensis Jacq.*). *Asian Journal of Plant Sciences* 5(2): 397-408
- Balasundram, S.K., Mulla, D., and Robert, P. 2007. Spatial data calibration for site-specific phosphorus management. *International Journal of Agricultural Research* 2: 888-889.
- Balasundram, S.K., M.H.A., Husni and Ahmed, O.H. 2008. Application of geostatistical tools to quantify spatial variability of selected soil chemical properties from a cultivated tropical peat. *Journal of Agronomy* 7(1): 82-82.
- Baldock, J.A., and Skjemstad, J.O. 2000. Role of the soil matrix and minerals in protecting natural organic materials against biological attack. *Organic Geochemistry* 31:697-710
- Barrow, C.J. 1991. Land degradation development and breakdown of terrestrial environment. Cambridge: Cambridge University Press
- Basile-Doelsch, I., Amundson, R., Stone, W., Borschneck, D., Bottero, J.Y., Moustier, S., Masin, F., and Colin, F. 2005. Mineral control of carbon pools in a volcanic soil horizon. *Geoderma* 137(3): 477-489.
- Basuki, T.M., van Laake, P. E., Skidmore, A. K., and Hussin, Y. A. 2009. Allometric equations for estimating the above-ground biomass in tropical lowland Dipterocarp forests. *Forest Ecology and Management* 257(8): 1684-1694.
- Bationo, A., Kihara, J., Vanlauwe, B., Waswa, B., and Kimetu, J. 2007. Soil organic carbon dynamics, functions and management in West African agroecosystems. *Agricultural Systems* 94(1): 13-25.

- Batjes, N. H. 1996 Total C and N in soils of the world. *European Journal of Soil Science* 47:151–163.
- Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N.,Rodenbeck, C.,Altaf Arain, M., Baldocchi, D., Bonan, G.B., Bondeau, A., Cescatti, A., Lasslop, G.,Lindroth, A., Lomas, M., Luyssaert, S., Margolis, H., Oleson, K.W., Roupsard, O.,Veenendaal, E., Viovy, N., Williams, C., Woodward, F.I.,Papale, D.2010. Terrestrial gross carbon dioxide uptake: global distribution and covariation with climate. *Science* 329:834–838.
- Bens, O., Buczko, U., Sieber, S., and Huttle, R.F. 2006. Spatial variability of O layer thickness and humus forms under different pine beech-forest transformation stages in NE Germany. *Journal of Plant Nutrition and Soil Science* 169(1): 5-15.
- Berg, B. and Laskowski, R. 2006. Litter decomposition: A guide to carbon and nutrient turnover. New York: Academic Press
- Berg, B., and McClaugherty, C.2008. *Plant litter: Decomposition, humus formation, carbon sequestration.* Berlin: Springer Verlag
- Berger, T.W., Inselsbacher, E., and Zechmeister-Boltenstern, S. 2010. Carbon dioxide emissions of soils under pure and mixed stands of beech and spruce, affected by decomposing foliage litter mixtures. *Soil Biology and Biochemistry* 42(6): 986-997.
- Bernhard-Reversat, F. and Schwartz, D. 1997. Change in lignin content during litter decomposition in tropical forest soils (Congo): Comparison of exotic plantations and native stands. *Earth and Planetary Science* 325(6): 427-432.
- Beta Analytic Inc. 2013. Introduction to Radiocarbon Determination by the Accelerator Mass Spectrometry Method. Retrieved 20 February 2013 from http://www.radiocarbon.com/PDF/Beta-AMS-Methodology.pdf
- Blais, A.M., Lorrain, S.P., Plourde, Y. and Varfalvy, L., 2005. Organic carbon densities of soils and vegetation of tropical, temperate and boreal forests.In *Greenhouse gas emissions-fluxes and processes* ed. A., Tremblay, L., Varfalvy, C., Roehm, and M. Garneau, pp155-185. Berlin: Springer-Verlag
- Bohlman, S.A., Matelson, T.J., and Nadkarni, N.M. 1995. Moisture and temperature patterns of canopy humus and forest floor soil of a montane cloud forest, Costa Rica. *Biotropica* 27(1): 13-19.

- Borders, B.D., Pushnik, J.C., and Wood, D.M. 2006. Comparison of leaf litter decomposition rates in restored and mature riparian forests on the Sacramento River, California. *Restoration Ecology* 14(2): 308-315.
- Bot, A., and Benites, J. 2005. *The importance of soil organic matter: Key to drought-resistant soil and sustained food production*. FAO Soils Bulletin 80. Rome: Food & Agriculture Organization.
- Bouwman, A., and Leemans, R. 1995. The role of forest soils in the global carbon cycle. In *Carbon forms and functions in forest soils*. ed. W., McFee, and J.M. Kelly, pp. 503-525. Wisconsin: Soil Science Society of America Inc.
- Boyd, H.A., Flanders, W.D., Addiss, D.G. and Waller, L.A. 2005. Residual spatial correlation between geographically referenced observations: A Bayesian hierarchical modeling approach. *Epidemilogy* 16(4): 532-541.
- Brady, N.C. and R.R. Weil. 2008. *The nature and properties of soils* (14th Ed). New Jersey: Prentice Hall.
- Bray, R.H. and Kurtz, L.T. 1945. Determination of total, organic, and available forms of phosphorus in soils. *Soil Science* 59:39-45.
- Bridson, D. and Forman, L. 1992. *The herbarium handbook*, ed. 2. Richmond Kew: Royal Botanic Gardens.
- Brown, S., Hall, C.A.S., Knabe, W., Raich, J., Trexler, M.C., Woomer, P. 1993. Tropical forests: Their past, present, and potential future role in the terrestrial carbon budget. *Water, Air & Soil Pollution* 70(1-4):71-94.
- Brown, S., Gillespie, A.J.R., and Lugo, A.E. 1989. Biomass estimation methods for tropical forests with applications to forest inventory data. *Forest Science* 35(4): 881-902.
- Brown, S., Lugo, A.E., Iverson, L.R. 1992. Processes and lands to sequestering carbon in the tropical forest landscape. *Water, Air and Soil Pollution* 64:139-155.
- Bruun, T.B., Elberling, B. and Christensen, B.T. 2010. Lability of soil organic carbon in tropical soils with different clay minerals. *Soil Biology and Biochemistry* 42(6): 888-895.
- Brundrett, M., Dell, B., Malajczuk, N., and Gong Mingqin. 1995. *Mycorrhizal Research for Forestry in Asia*. Canberra: Australian Centre for International Agricultural Research (ACIAR)
- Burnham, C. 1974. Altitudinal changes in soils on granite in Malaysia. *Proceedings* of the 10th International Congress of Soil Science 10(6): 290–296

- Cairns, M.A., Brown, S., Helmer, E.H., and Baumgardner, G.A. 1997. Root biomass allocation in the world's upland forests. *Oecologia* 111(1): 1-11.
- Cambardella, C.A., Moorman, T.B., Novak, J.M., Parkin, T.B., Karlen, D.L., Turco, R.F., and Konopka, A.E. 1994. Field-scale variability of soil properties in central Iowa soils. *Soil Science Society of American Journal* 58: 1501-1511.
- Chave, J., Andalo, C., Brown, S., Cairns, M., Chambers, J., Eamus, D., Folster, H., Fromard, F., Higuchi, N., and Kira, T. 2005. Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia*, 145(1): 87-99.
- Chimner, R. and Karberg, J. 2008. Long-term carbon accumulation in two tropical mountain peatlands, Andes Mountains, Ecuador. *Mires & Peat* 3:1-10.
- Chiti, T., Neubert, R., Janssens, I., Certini, G., Curiel Yuste, J., and Sirignano, C. 2009. Radiocarbon dating reveals different past managements of adjacent forest soils in the Campine region, Belgium. *Geoderma* 149(1): 137-142.
- Choi, D., Quoreshi, A., Maruyama, Y., Jin, H., and Koike T. 2005. Effect of ectomycorrhizal infection on growth and photosynthetic characteristics of Pinus densiflora seedlings grown under elevated CO₂ concentrations. *Photosynthetica* 43(2): 223-229.
- Chojnacky, D.C., Mickler, R.A., Heath, L.S., and Woodall, C.W. 2004. Estimates of down woody materials in eastern US forests. *Environmental Management* 33(1): S44-S55.
- Chojnacky, D., Amacher, M and Gavazzi. M. 2009. Separating duff and litter for improved mass and carbon estimates. *Southern Journal of Applied Forestry* 33(1): 29-34.
- Currie, W.S, Yanai, R.D., Piatek, K.B., Prescott, C.E. and Goodale, C.L. 2003. Processes affecting carbon storage in the forest floor and in downed woody debris. In *The potential of US forest soil to sequester carbon and mitigate the greenhouse effect, ed.* J.M., Kimble, R., Lal, R., Birdsey, and L.S. Heath, pp 135-157.Boca Raton: CRC Press
- Couto, E.G., Stein, A., and Klamt, E. 1997. Large area spatial variability of soil chemical properties in central Brazil. *Agriculture*, *Ecosystems & Environment* 66(2):139-152.
- Couteaux, M.M., Bottner, P. and Berg, B.R. 1995. Litter decomposition, climate and liter quality. *Trends in Ecology & Evolution* 10(2): 63-66.
- Dent, D.H., Bagchi, R., Robinson, D., Majalap-Lee, N. and Burslem, D.F. 2006. Nutrient fluxes via litterfall and leaf litter decomposition vary across a gradient of soil nutrient supply in a lowland tropical rain forest. *Plant and Soil* 288(1-2):197-215.

- Department of Agriculture. 1989. Soil correlation of steepland:Soil in Cameron Highlands: 20-21 February 1989. Kuala Lumpur: Soil Management Branch, Department of Agriculture.
- Department of Agriculture. 2000. Agricultural development plan of Cameron Highlands. . Kuala Lumpur: Soil Management Branch, Department of Agriculture.
- Djomo, A.N., Knohl, A., and Gravenhorst, G. 2011. Estimations of total ecosystem carbon pools distribution and carbon biomass current annual increment of a moist tropical forest. *Forest Ecology and Management* 261:1448–1459.
- Dixon, R.K., Solomon, A.M., Brown, S., Houghton, R.A., Trexler, M.C., Wisniewski, J. 1994. Carbon pools and flux of global forest ecosystems. *Science* 263: 185–190.
- Dof Sotta, E., Meir, P., Malhi, Y., Hodnett, M., and Grace, J. 2004. Soil CO₂ efflux in a tropical forest in the central Amazon. *Global Change Biology* 10(5): 601-617.
- Domisch, T., Finer, L., Laine, J. and Laiho, R. 2006. Decomposition and nitrogen dynamics of litter in peat soils from two climatic regions under different temperature regimes. *European Journal of Soil Biology* 42(2): 74-81.
- Duchaufour, P. 1982. *Pedology: Pedogenesis and classification*. Translated by T.R. Paton. London: George Allen & Unwin.
- ESRI, 1994. Understanding GIS:the ARC/INFO method, version 7 for UNIX and Open VMS. Redlands: ESRI
- Eswaran, H., Berg, E., and van den Reich, P. 1993. Organic carbon in soils of the world. *Soil Science Society of American Journal* 57:192–194.
- Fioretto, A., Di Nardo, C., Papa, S., and Fuggi, A. 2005. Lignin and cellulose degradation and nitrogen dynamics during decomposition of three leaf litter species in a Mediterranean ecosystem. *Soil Biology and Biochemistry* 37(6): 1083-1091.
- Food and Agriculture Organization, 2010. *Global forest resources assessment 2010*. FAO Forestry Paper 163 Rome: FAO of the United Nations
- Frogbrook, Z. and Oliver, M. 2001. Comparing the spatial predictions of soil organic matter determined by two laboratory methods. *Soil Use and Management* 17(4): 235-244.
- Gamma Design Software, 2004. *GS+: Geostatistics for environmental scientists. GS + User guide version 7.* Michigan: Gamma Design Software.

- Garten Jr., C.T.2005. Forest soil carbon inventories and dynamics along an elevation gradient in the southern Appalachian mountains. Retrieved 12 June 2013 from http://www.esd.ornl.gov/research/terrestrial_ecology/soil_carbon.shtml.
- Garten, C.T., and Hanson, P.J. 2006. Measured forest soil C stocks and estimated turnover times along an elevation gradient. *Geoderma* 136(1): 342-352.
- Gershenson, A., Bader, N.E., and Cheng, W. 2008. Effects of substrate availability on the temperature sensitivity of soil organic matter decomposition. *Global Change Biology* 15(1): 176-183.
- Gibbs, H.K., Brown, S., Niles, J.O., and Foley, J.A. 2007. Monitoring and estimating tropical forest carbon stocks: making REDD a reality. *Environmental Research Letters* 2(4): 045023.
- Goetz, S., Baccini, A., Laporte, N., Johns, T., Walker, W., Kellndorfer, J., Houghton, R. and Sun, M. 2009. Mapping and monitoring carbon stocks with satellite observations: A comparison of methods. *Carbon Balance and Management* 4(1):2.
- Goslar, T., Van der Knaap, W., Hicks, S., Andric, M., Czernik, J., Goslar, E., Rasanen, S., and Hyotyla, H. 2005. Radiocarbon dating of modern peat profiles: pre-and post-bomb 14C variations in the construction of age-depth models. *Radiocarbon* 47(1): 115-134.
- Gosz, J. R., Likens, G. E., and Bormann, F.H. 1976. Organic matter and nutrient dynamics of the forest and forest floor in the Hubbard Brook Forest. *Oecologia* 22(4): 305-320.
- Gourbiere, F., and Debouzie, D. 1995. Spatial distribution and estimation of forest floor components in a 37-year-old *Casuarina equisetifolia* (forst.) plantation in coastal Senegal. *Soil Biology and Biochemistry* 27(3): 297-304.
- Greenpeace, 2008. Greenpeace Briefing: Tropical deforestation and the Kyoto Protocol.Retrieved 20 February 2013 from http://www.greenpeace.org/international/press/reports/briefing-deforestation-climate-NY-meet.
- Griffiths, R.P., Madritch, M.D., and Swanson, A.K. 2009. The effects of topography on forest soil characteristics in the Oregon Cascade Mountains (USA): Implications for the effects of climate change on soil properties. *Forest Ecology and Management* 257(1): 1-7.
- He, X., Zhang, P., Lin, Y., Li, A., Tian, X., and Zhang, Q.H. 2009. Responses of litter decomposition to temperature along a chronosequence of tropical montane rainforest in a microcosm experiment. *Ecological Research* 24(4): 781-789.

- Heath, L.S., Smith, J., Skog, K., Nowak, D., Woodall, C.W. 2011. Managed forest carbon stock and stock-change estimates for the U.S. greenhouse gas inventory,1990–2008. *Journal of Forestry* 109:167–173.
- Heath, L.S., Smith, J.E., and Birdsey, R.A. 2003. Carbon trend in U.S. forestland: A context for the role of soils in forest carbon sequestration. In *The potential of U.S. forest soils to sequester carbon and mitigate the greenhouse effect*, ed. J.M., Kimble, L.S. Heath, R.A. Birdsey, and R. Lal. pp.35-45. Boca Raton: CRC Press
- Hengl, T.,2007. A practical guide to geostatistical ,mapping of environmental variables. JRC scientific and technical reports. Luxemburg: European Commission JRC, Institut for Environment and Sustainability. Retrieved 15 February 2013 from http://eusoils.jrc.ec.europa.eu/esdb_archive/eusoils_docs/other/eur22904en.p df
- Hertel, D. and Leuschner, C. 2010. Fine root mass and fine root production in tropical moist forests as dependent on soil, climate, and elevation. In *Tropical Montane Cloud Forests: Science for Conservation and Management*, ed. L.A., Bruijnzeel, F.N., Scatena, and L.S. Hamilton, pp. 428-442 Cambridge: Cambridge University Press.
- Hilli, S., Stark, S. and Derome, J. 2010. Litter decomposition rates in relation to litter stocks in boreal coniferous forests along climatic and soil fertility gradients. *Applied Soil Ecology* 46(2): 200-208.
- Hugget, R. and Cheeseman, J. 2002. *Topography and the Environment*. Harlow: Prentice-Hall.
- Ho, C.C., Newbery, D.M., and Poore, M.E.D. 1987. Forest composition and inferred dynamics in Jengka Forest Reserve, Malaysia. *Journal of Tropical Ecology* 3(1): 25-56.
- Hoover, C.M. 2011. Management impacts on forest floor and soil organic carbon in northern temperate forests of the US. *Carbon Balance and Management* 6(1): 1-8.
- IPCC, 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Prepared by the National Greenhouse Gas Inventories Programme ed. H.S. Eggleston, L. Buendia, K., Miwa, T., Ngara and K. Tanabe. Japan: Institute For Global Environmental Strategies
- IPCC, 2007: Climate Change 2007: Synthesis Report. *In Contribution of Working Groups I,II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* ed. R.K., Pachauri, A. Reisinger, et al.. Geneva: International Panel for Climate Change

- Isaaks, E.H. and Srivastava, R.M.1989. *An Introduction to Applied Geostatistics (1st edition)*. New York :Oxford University Press.
- Itoh, M., Kosugi, M., Takanashi, S., Hayashi, Y., Kanemitsu, S., Osaka, K., Tani, M., and NIk, A.R. 2010. Temporal and spatial variations of soil carbon dioxide, methane, and nitrous oxide fluxes in a Southeast Asian tropical rainforest. *Biogeosciences Discussions* 7: 6847-6887.
- Jackson, R.B., Mooney, H., and Schulze, E.D. 1997.A global budget for fine root biomass, surface area and nutrient contents. *Proceedings of the National Academy of Sciences* 94(14):7362-7366.
- Jacob, M., Weland, N., Platner, C., Schaefer, M., Leuschner, C., and Thomas, F.M. 2009. Nutrient release from decomposing leaf litter of temperate deciduous forest trees along a gradient of increasing tree species diversity. *Soil Biology and Biochemistry* 41(10): 2122-2130.
- Jastrow J.D. and Miller R.M. 1997. Soil aggregate stabilization and carbon sequestration: Feedbacks through organomineral associations. *In Soil Processes and the Carbon Cycle*. ed. R., Lal, J.M., Kimble, R.F., Follett and B.A. Stewart. pp. 207-223. Boca Raton:CRC Press.
- Journel, A. G. and Huijbregts, C.J. 1978. *Mining Geostatistics*. London: Academic Press.
- Kalbitz, K., Meyer, A., Yang, R., and Gerstberger, P. 2007. Response of dissolved organic matter in the forest floor to long-term manipulation of litter and throughfall inputs. *Biogeochemistry* 86(3): 301-318.
- Kane, E., Valentine, D., Schuur, E.A.G., and Dutta, K. 2005. Soil carbon stabilization along climate and stand productivity gradients in black spruce forests of interior Alaska. *Canadian Journal of Forest Research* 35(9): 2118-2129.
- Keith D.M., Johnson E.A., and Valeo C. 2010. A hillslope forest floor (duff) water budget and the transition to local control. *Hydrology Processes* 24: 2738–2751.
- Ketterings, Q.M., Coe, R., van Noordwijk, M., Ambagau, Y., and Palm, C.A. 2001. Reducing uncertainty in the use of allometric biomass equations for predicting above-ground tree biomass in mixed secondary forests. *Forest Ecology and Management* 146(1): 199-209.
- Kirkby, C., Kirkegaard J., Richardson A., Wade L., Blanchard C., and Batten G. 2013. Nutrients-the real constraint to sequestering carbon in soil organic matter? Retreived 5 January 2013 from http://aciar.gov.au/files/node/14068/nutrients_the_real_constraint_to_sequest ering_ca_61722.pdf

- Kitayama, K. and Aiba, S.I. 2002. Ecosystem structure and productivity of tropical rain forests along altitudinal gradients with contrasting soil phosphorus pools on Mount Kinabalu, Borneo. *Journal of Ecology* 90(1):37-51.
- Kogel-Knabner, I. 2002. The macromolecular organic composition of plant and microbial residues as inputs to soil organic matter. *Soil Biology and Biochemistry* 34(2): 139-162.
- Kohl, M. and Gertner G.1997. Geostatistics in evaluating forest damage surveys: considerations on methods for describing spatial distributions. *Forest Ecology and Management* 95(2): 131-140.
- Kosugi, Y., Mitani, T., Itoh, M., Noguchi, S., Tani, M., Matsuo, N., Takanashi, S., Ohkubo, S., and Rahim Nik, A. 2007. Spatial and temporal variation in soil respiration in a Southeast Asian tropical rainforest. *Agricultural and Forest Meteorology* 147(1-2): 35-47.
- Kravchenko, A.N., Robertson, G., Snap, S., and Smucker, A.J.M. 2006. Using information about spatial variability to improve estimates of total soil carbon. *Agronomy Journal* 98(3): 823-829.
- Krige, D.G. 1951. A statistical approach to some basic mine valuation problems on the Witwatersrand. *Journal of the Chemical, Metallurgical and Mining Society of South Africa* 52 (6):119–139.
- Laidlaw, R.K. 2011. Effects of habitat disturbance and protected areas on mammals of Peninsular Malaysia. *Conservation Biology* 14(6):1639-1648.
- Lal, R. 1995. Global soil erosion by water and carbon dynamics. In *Soils and global change* ed. R., Lal, J., Kimble, E., Levine, and B.A. Stewart, pp.131-142. Boca Raton: CRC Press Inc.
- Lal, R., 2008. Carbon sequestration. *Philosophical Transactions of the Royal Society B: Biological Sciences* 363(1492):815-830.
- Lal, R., J.M., Kimble, R.F., Follett, B.A. Stewart. 1998. Soil processes and the carbon cycle. Boca Raton:CRC Press
- Lal, R. 2004. Soil carbon sequestration impacts on global climate change and food security *Science* 304: 1623-1627.
- Lal, R. and Follett, F. 2009. *Soil carbon sequestration and the greenhouse effect*. Special Publ. 57, Second Edition. Madison: Soil Science Society of America.
- Lasco, R.D., Guillermo, I.Q., Cruz, R.V.O., Bantayan, N.C., and Pulhin, F.B.2004. Carbon stocks assessment of a secondary forest in Mount Makiling Forest Reserve, Philippines. *Journal of Tropical Forest Science* 16:35–45.

- Law, M.C., Balasundram, S.K., and Husni, M.H.A. 2009. Spatial variability of soil organic carbon in oil palm. *International Journal of Soil Science* 4(4): 93-103.
- Leifeld, J., Bassin, S., and Fuhrer, J. 2005. Carbon stocks in Swiss agricultural soils predicted by land-use, soil characteristics, and altitude. *Agriculture, Ecosystems & Environment* 105(1-2): 255-266.
- Li, P., Wang, Q., Endo, T., Zhao, X. and Kakubari, Y. 2010, Soil organic carbon stock is closely related to aboveground vegetation properties in cold-temperate mountainous forests. *Geoderma* 154(3-4): 407-415.
- Liaghat, S., and Balasundram, S.K. 2010. A Review: The Role of Remote Sensing in Precision Agriculture. *American Journal of Agricultural and Biological Sciences* 5(1): 50-55.
- Lichter, J., Billings, S. A., Ziegler, S. E., Gaindh, D., Ryals, R., Finzi, A.C., Jackson, R.B., Stemmler, E.A., and Schlesinger W.H. 2008. Soil carbon sequestration in a pine forest after 9 years of atmospheric CO₂ enrichment. *Global Change Biology* 14(12):2910-2922.
- Lim, J.S. 2002. *Estimating organic carbon in soils of Peninsular Malaysia*. Paper presented at the Seminar on Climate Change Carbon Accounting, SIRIM, Shah Alam. 8 January 2002.
- Liski, J. and Westman, C. J. 1995. Density of organic carbon in soil at coniferous forest sites in southern Finland. *Biogeochemistry* 29(3): 183-197.
- Liu, D., Wang, Z., Zhang, B., Song, K., Li, X., Li, J., Li, F., and Duan, H. 2006. Spatial distribution of soil organic carbon and analysis of related factors in croplands of the black soil region, Northeast China. *Agriculture, Ecosystems & Environment* 113(1): 73-81.
- Lorenz, K. and Lal, R. 2010. *Carbon sequestration in forest ecosystems*. New York: Springer.
- Luo, Y and Zhuo, X. 2006. Soil respiration and the environment. Burlington: Academic Press.
- Luyssaert, S., Inglima, I., Jung, M., Richardson, A., Reichstein, M., Papale, D., et al. 2007. CO₂ balance of boreal, temperate, and tropical forests derived from a global database. *Global Change Biology* 13(12): 2509-2537.
- Maden, K. 2004. Plant collection and herbarium techniques. *Our Nature* 2(1): 53-57.

- Majila, B.S., Joshi, G.C., and Kala, C.P. 2005. Patterns in litter fall and litter decomposition along an altitudinal gradient in the Binsar Wildlife Sanctuary, Central Himalaya. *International Journal of Sustainable Development & World Ecology* 12(2): 205 212.
- Malaysian Standard 678. 1980. Recommended methods for soil chemical analysis: Part I V. Selangor: Standards & Industrial Research Institute of Malaysia.
- Malhi, Y., T.R., Baker, O.L., Phillips, et al. 2004. The above-ground coarse wood productivity of 104 Neotropical forest plots. *Global Change Biology* 10: 563–591.
- Manton, M.J., Della-Marta, P.M., Haylock M.R., Hennessy K.J., Nicholls N., Chambers L.E., Collins D.A., Daw G., Finet A., Gunawan D., et.al. 2001. Trends in extreme daily rainfall and temperature in Southeast Asia and the South Pacific: 1961-1998. *International Journal of Climatology* 21: 269-284.
- Manton, M.J., Heath L., Salinger, J., and Stevenson L.A. 2011. *Climate in Asia and the Pacific: A Synthesis of APN Activities*. Japan: Asia-Pacific Network for Global Change Research.
- Marchetti, A., Piccini, C., Francaviglia, R., and Mabit, L. 2012. Spatial distribution of soil organic matter using geostatistics: A key indicator to assess soil degradation status in central Italy. *Pedosphere* 22(2): 230-242.
- Martin, W.K.E. and Timmer, V.R. 2006. Capturing spatial variability of soil and litter properties in a forest stand by landform segmentation procedures. *Geoderma* 132(1-2): 169-181.
- Martin, D., Lal, T., Sachdev, C. B., and Sharma, J. P. 2010. Soil organic carbon storage changes with climate change, landform and land use conditions in Garhwal hills of the Indian Himalayan mountains. *Agriculture, Ecosystems & Environment*, 138: 64-73.
- Martin, Y.E., Johnson, E.A., Gallaway, J.M., and Chaikina, O. 2011. Negligible soil erosion in a burned mountain watershed, Canadian Rockies: field and modelling investigations considering the role of duff. *Earth Surface Processes and Landforms* 36(15): 2097-2113.
 - Mason, I.G., McLachlan, R.I., and Gerard, D.T. 2006. A double exponential model for biochemical oxygen demand. *Bioresource Technology* 97(2): 273-282.
- McGroddy, M. and Silver, W.L. 2000. Variations in belowground carbon storage and soil CO₂ flux rates along a wet tropical climate gradient. *Biotropica* 32(4a): 614-624.

- McKenzie, N., Ryan, P., Fogarty, P. and Wood, J. 2000. Sampling, measurement and analytical protocols for carbon estimation in soil, litter and coarse woody debris: National Carbon accounting technical report 14; Canberra: Australian Greenhouse Office.
- Melling, L., Hatano, R., and Goh, K.J. 2005. Soil CO₂ flux from three ecosystems in tropical peatland of Sarawak, Malaysia. *Tellus B* 57(1): 1-11.
- Melvin, K.K.K., Japar, S.B., Ahmed, O.H., Majid, N.M., Roland, K.J.H., and Silvester, J. 2011. Litter production, decomposition and standing forest litter of rehabilitated forests at Bintulu, Sarawak, Malaysia. *In* Proceedings of the Rehabilitation of Tropical Rainforest Ecosystems, ed. S.B., Japar, O.H., Ahmed, N.M., Majid, K.J.H. Roland.pp 287-291. 24 25 October 2011, Kuala Lumpur: Universiti Putra Malaysia
- Ministry of Natural Resources and Environment, Malaysia. 2011. Second national communication to the UNFCCC. Putrajaya: NRE.
- Moore, T.R., Trofymov J.A., Taylor B., Prescott C., Camire C., Duschene L., Fyles J., Kozak L., Kranabetter M., Morrison I., Siltanen M., Smith S., Titus B, Visser S, Wein R, and Zoltai S., 1999. Litter decomposition rates in Canadian forests. *Global Change Biology* 5: 75–82.
- Moser, G., M. Roderstein, N., Soethe, D., Hertel, and C.H. Leuschner. 2008. Altitudinal changes in stand structure and biomass allocation of tropical mountain forest in relation to microclimate and soil chemistry. In *Gradients in a tropical mountain ecosystem of Ecuador*, ed. E. Beck, J., Bendix, I., Kottke, F., Makeschin, and R. Mosandl, pp. 229–242. Berlin: Springer-Verlag.
- Moyersoen, B., Alexander, I.J., and Fitter, A.H. 1998. Phosphorus nutrition of ectomycorrhizal and arbuscular mycorrhizal tree seedlings from a lowland tropical rain forest in Korup National Park, Cameroon. *Journal of Tropical Ecology*: 47-61.
- Mulla, D.I. and A.B. McBratney, 1999. Soil Spatial Variability. In *Handbook of Soil Science*, ed. M., Sumner, pp. A321-A352. Boca Raton: CRC Press.
- Muller, R.N. 2003. Landscape patterns of change in coarse woody debris accumulation in an old-growth deciduous forest on the Cumberland Plateau, southeastern Kentucky. *Canadian Journal of Forestry Research* 33: 763–769.

- Mulligan, M. 2010. Modelling the tropics-wide extent and distribution of cloud forest and cloud forest loss, with implications for conservation priority. In *Tropical Montane Cloud Forests: Science for Conservation and Management*, ed. L.A., Bruijnzeel, F.N., Scatena, and L.S. Hamilton, pp. 14-38 Cambridge: Cambridge University Press.
- Murtedza, M., Padmanabhan, E., B.L.H. Mei and W.B. Siong. 2002. *The peat soil of Sarawak*. Kuching: Universiti Malaysia Sarawak
- Myneni, R.B., Keeling, C.D., Tucker, C.J., Asrar, G., and Nemani R.R. 1997. Increased plant growth in the northern high latitudes from 1981 to 1991. *Nature* 386: 698 –702.
- Nahrawi, H., Husni, M., Othman, R., and Bah, A. 2011. Decomposition of Leaf and Fine Root Residues of Three Different Crop Species in Tropical Peat under Controlled Condition. *Malaysian Journal of Soil Science* 15: 63-74.
- Nakashizuka, T., Yusop, Z. and Nik, A.R., 1992. Altitudinal zonation of forest communities in Selangor, Peninsular Malaysia. *Journal of Tropical Forest Science* 4(3): 233-244.
- Ngo, K.M., Turner, B.L., Muller-Landau, H.C., Davies, S.J., Larjavaara, M., Nik Hassan, N.F.B. and Lum, S. 2013. Carbon stocks in primary and secondary tropical forests in Singapore. *Forest Ecology and Management* 296: 81-89.
- Nicholls, N., Baek H.J., Gosai, A., Chambers L.E., Choi, Y., Collins, D., Della-Marta, P.M., Griffiths, G.M., Haylock, M.R., Iga N, et.al. 2005. The El Nino Southern Oscillation and daily temperature extremes in East Asia and the West Pacific. *Geophysics Research Letters* 32: L16714
- Niiyama, K., Kajimoto, T., Matsuura, Y., Yamashita, T., Matsuo, N., Yashiro, Y., Ripin, A., Kassim, A.R., and Noor, N.S. 2010. Estimation of root biomass based on excavation of individual root systems in a primary dipterocarp forest in Pasoh Forest Reserve, Peninsular Malaysia. *Journal of Tropical Ecology* 26: 71-284.
- Nizam, M.S. and D. Rohaiza. 2011. Species composition, diversity and biomass of tree communities at two forest reserves at Cameron Highlands. *In Hutan pergunungan Cameron Highlands: Pengurusan hutan, persekitaran fizikal dan kepelbagaian biologi* (Vol.14) ed. A.R., Abdul Rahman, H.L, Koh, K., Mohd. Paiz, A., Muhammad and Latiff A. pp. 97-107. Kuala Lumpur: Forestry Department of Peninsular Malaysia.
- Nogueira, F., Couto, E., and Bernardi, C. 2002. Geostatistics as a tool to improve sampling and statistical analysis in wetlands: A case sudy on dynamics of organic matter distribution in the Pantanal of Mato Grosso, Brazil. *Brazilian Journal of Biology* 62(4b): 861-870.

- Norini, H., Abdul Razak, M.A., Phillips E., and Samsudin, M. 2007. Opportunities and Challenges of the Forestry Clean Development Mechanism (CDM) for the Timber Industry in Malaysia. *Sarawak Developmental Journal* 8:16-29
- Nuccio, E. E., Hodge, A., Pett-Ridge, J., Herman, D. J., Weber, P. K., and Firestone, M.K. 2013. An arbuscular mycorrhizal fungus significantly modifies the soil bacterial community and nitrogen cycling during litter decomposition. *Environmental Microbiology* 15(6): 1870-1881.
- Olsen, S.R. and Sommers, L.E. 1982. Phosphorus. In *Methods of soil analysis, Part* 2, *Chemical and Microbiological Properties*, ed. A.L., Page, R.H., Miller, and D.R. Keeney, (Eds.). pp. 403-430. Madison: ASA and SSSA
- Olson, J. S. 1963. Energy storage and the balance of producers and decomposers in ecological systems. *Ecology* 44(2): 322-331.
- Ottmar, R., and Andreu, A. 2007. Litter and duff bulk densities in the Southern United States. Final report, JFSP project, 04-2-1-49. USDA Forest Service. Retrieved on 13 March 2013 from https://www.firescience.gov/projects/04-2-1-49/project/04-2-1-49_final_report.pdf
- Ovington, J.D. 1959. The circulation of minerals in plantations of *Pinus sylvestris* L. *Annals of Botany* 23:229–239.
- Pachauri, R.K., and Reisinger, A., 2007. *Climate change 2007: Synthesis Report*. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.Geneva: IPCC
- Paramanathan, 1977. Soil genesis on igneous and metamorphic rocks in Malaysia. PhD Thesis, University of Ghent, Belgium.
- Parsons, S.A., and Congdon, R.A. 2008. Plant litter decomposition and nutrient cycling in north Queensland tropical rain-forest communities of differing successional status. *Journal of Tropical Ecology* 24(3): 317-327.
- Paul, E. 1984. Dynamics of organic matter in soils. *Plant Soil* 76: 75-285.
- Pearson, T., Walker, S. and Brown, S. 2005. Sourcebook for land use, land-use change and forestry projects. Winrock International and the BioCarbon Fund of the World Bank: WinRock International
- Peh, K.S.H., Soh, M.C.K., Sodhi N.S., Laurance, W.F., Ong D.J., and Clements, R. 2011. Up in the clouds: Is sustainable use of tropical montane cloud forests possible in Malaysia. *Bioscience* 61: 27-38.
- Penne, C., Ahrends, B., Deurer, M., and Böttcher, J. 2010. The impact of the canopy structure on the spatial variability in forest floor carbon stocks. *Geoderma* 158(3-4): 282-297.

- Perez, J. and Jeffries, T.W. 1992. Roles of manganese and organic acid chelators in regulating lignin degradation and biosynthesis of peroxidases by *Phanerochaete chrysosporium*. *Applied and Environmental Microbiology* 58(8): 2402-2409.
- Perez-Corona, M.E., Hernandez, M.C.P. and de Castro, F.B. 2006. Decomposition of alder, ash, and poplar litter in a Mediterranean riverine area. *Communications in Soil Science and Plant Analysis* 37(7-8): 1111-1125.
- Persson, H. 1980. Spatial distribution of fine-root growth, mortality and decomposition in a young Scots pine stand in Central Sweden. *Oikos* 34: 77-87.
- Pfeffer, P.E., Bago, B. and Shachar-Hill, Y. 2001. Exploring mycorrhizal function with NMR spectroscopy. *New Phytologist* 150(3): 543-553.
- Pingintha, N., Leclerc, M.Y., Beasley Jr, J.P., Zhang, G., and Senthong, C. 2010. Assessment of the soil CO₂ gradient method for soil CO₂ efflux measurements: Comparison of six models in the calculation of the relative gas diffusion coefficient. *Tellus B* 62(1): 47-58.
- Piotrowska, N., Blaauw, M., Mauquoy, D., and Chambers, F. 2011. Constructing deposition chronologies for peat deposits using radiocarbon dating. *Mires and Peat* 7: 1-14.
- Pitelka, L. and A.R. Rojas. 2001. IPCC WGI Third Assessment Report, Chapter 3: The carbon cycle & atmospheric carbon dioxide. Retrieved 1 June 2013 fromhttp://www.ipcc.ch/ipccreports/tar/wg1/558.htm.
- Podwojewski, P., Poulenard, J.R.M., Nguyet, M.L., De Rouw, A., Nguyen, V.T., Pham, Q.H. and Tran, D.T. Climate and vegetation determine soil organic matter status in an alpine inner-tropical soil catena in the Fan Si Pan Mountain, Vietnam. *Catena* 87(2): 226-239.
- Poore, M. 1968. Studies in Malaysian Rain Forest: I. The Forest on Triassic Sediments in Jengka Forest Reserve. *The Journal of Ecology* 56(1): 143-196.
- Pregitzer, K.S., and Euskirchen, E.S. 2004. Carbon cycling and storage in world forests: biome patterns related to forest age. *Global Change Biology* 10(12): 2052-2077.
- Prescott, C.E., Blevins, L.L., and Staley, C. 2004. Litter decomposition in British Columbia forests: controlling factors and influences of forestry activities. *BC Journal of Ecosystem and Management* 5(2): 44-57.

- Quadir, D.A., Shrestha, M.L., Khan, T.M.A., Ferdousi, N., Rahman, M., and Mannan, A. 2004. Variations of surface air temperature over the land areas in and around the Bay of Bengal. *Natural Hazards* 31(2): 561-584.
- Raich, J.W., Potter, C.S., and Bhagawati, D. 2002. Interannual variability in global soil respiration, 1980-94. *Global Change Biology* 8(8):800-812.
- Rillig, M.C., Wright, S.F., Nichols, K.A., Schmidt, W.F., and Torn, M.S. 2001. Large contribution of arbuscular mycorrhizal fungi to soil carbon pools in tropical forest soils. *Plant and Soil* 233(2): 167-177.
- Robinson, D. 2007. Implications of a large global root biomass for carbon sink estimates and for soil carbon dynamics. *Proceedings of the Royal Society B: Biological Sciences* 274(1626): 2753-2759.
- Roman, L., Scatena, F.N. and Bruijnzeel, L.A. 2010. Global and local variations in tropical mounane forest soils. *Tropical Montane Cloud Forests: Science for Conservation and Management*, ed. L.A., Bruijnzeel, F.N., Scatena, and L.S. Hamilton, pp. 77-89 Cambridge: Cambridge University Press.
- Rossi, J., Govaerts, A., De Vos, B., Verbist, B., Vervoort, A., Poesen, J., Muys, B. and Deckers, J. 2009. Spatial structures of soil organic carbon in tropical forests-A case study of Southeastern Tanzania. *Catena* 77(1): 19-27.
- Rossi, R.E., Mulla, D.J., Journel, A.G. and Franz, E.H. 1992. Geostatistical tools for modeling and interpreting ecological spatial dependence. *Ecological Monographs* 62(2): 277-314.
- Sammathuria, M.K., Ling L.K., and Wan Hassan, W.A. 2010. Extreme climate change scenarios over Malaysia for 2001 2099. JMM Research Publication No.12/2010.Retrieved 20 February 2013 from http://www.met.gov.my/images/pdf/research/researchpaper_201013.pdf
- Samonil, P., Kral, K., Douda, J., and Sebkova, B. 2008. Variability in forest floor at different spatial scales in a natural forest in the Carpathians: Effect of windthrows and mesorelief. *Canadian Journal of Forest Research* 38(10): 2596-2606.
- Saner, P., Loh, Y.Y., Ong, R.C. and Hector, A. 2012. Carbon stocks and fluxes in tropical lowland Dipterocarp rainforests in Sabah, Malaysian Borneo. *PloS one* 7(1): e29642.
- Satrio, A.E., Gandaseca, S., Ahmed, O.H., and Majid, N.M.A. 2009. Influence of chemical properties on soil carbon storage of a tropical peat swamp forest. *American Journal of Applied Sciences* 6 (11): 1970-1973.

- Sauer, T.J., Cambardella, C.A., and Brandle, J.R., 2007. Soil carbon and tree litter dynamics in a red cedar and scotch pine shelterbelt. *Agroforestry systems* 71(3):163-174.
- Saw, L.G. 2010. Vegetation of Peninsular Malaysia. In *Malayan Forest Records No.* 49. Series II. Seed Plant, Vol. 1. Flora of Peninsular Malaysia, ed. R., Kiew, R.C.K, Chung, L.G., Saw, E. Soepadmo, and P.C. Boyce.
- Scheffer R.A., Van Logtestijn, R.S.P. and Verhoeven J.T.A. 2001.Decomposition of Carex and *Sphagnum* litter in two mesotrophic fens differing in dominant plant species. *Oikos* 92:44–54.
- Scherer-Lorenzen, M., Luis Bonilla, J., and Potvin, C. 2007. Tree species richness affects litter production and decomposition rates in a tropical biodiversity experiment. *Oikos* 116(12):2108-2124.
- Schiermeier, Q. Increased flood risk linked to global warming. *Nature* 470(7334): 316-316.
- Schlesinger, W.H. 1977. Carbon balance in terrestrial detritus. *Annual review of ecology and systematics* 8: 51-81.
- Schnitzer, M. 1991. Soil organic matter-the next 75 years. Soil Science 151(1): 41-58.
- Schiffman, P.M., Johnson W.C. 1989. Phytomass and detrital carbon storage during forest regrowth in the southeastern United States Piedmont. *Canadian Journal of Foest Research* 19:67–78.
- Schrag, D. P. 2007. Preparing to capture carbon. *Science* 315(5813): 812-813.
- Schuur, E. A. and Matson, P. A. 2001. Net primary productivity and nutrient cycling across a mesic to wet precipitation gradient in Hawaiian montane forest. *Oecologia*, 128(3):431-442.
- Selvaraju, R. 2003. Impact of El Niño-Southern Oscillation on Indian foodgrain production. *International Journal of Climatology* 23:187-206.
- Sheikh, M.A., Kumar, M., and Bussmann, R.W. 2009. Altitudinal variation in soil organic carbon stock in coniferous subtropical and broadleaf temperate forests in Garhwal Himalaya. *Carbon Balance and Management* 4:6
- Sigurdsson, B.D., Roberntz P., Freeman M., Naess M., Saxe H., Thorgeirsson H. and Linder S. 2002. Impact studies on Nordic forests: Effects of elevated CO₂ and fertilization on gas exchange. *Canadian Journal of Forest Research* 32: 779–788.

- Silveira, M., Comerford, N., Reddy, K., Prenger, J. and Debusk, W. 2009. *Soil properties as indicators of disturbance in forest ecosystems of Georgia, USA*. Ecological Indicators 9: 740–747.
- Singh. K.D. 1993. The 1990 tropical forest resources assessment. In Forest resources assessment. Unasylva No.174.Vol.44 1993/3. Retrieved 1 June 2013 from http://www.fao.org/docrep/v0290E/v0290e04.
- Soderstrom, B. 2002. Challenges for mycorrhizal research into the new millennium. *Plant and Soil* 244: 1-7
- Sodhi N.S., and Brook B.W. 2006. *Southeast Asian Biodiversity in Crisis*. Cambridge: Cambridge University Press
- Soil Science Society of America. 2009. Carbon Sequestration: Position of the Soil Science Society of America, Retrieved 10 January 2013 from https://www.soils.org/pdf/pos_paper_carb_seq.pdf
- Soil Survey Staff, 1993. *Soil survey manual*. USDA Agricultural Handbook No. 18. Washington: U.S. Government Printing Office.
- Soil Survey Staff, 2010. *Keys to Soil Taxonomy*, 11th edition. Washington: USDA Natural Resources Conservation Service.
- Solomon, S., D., Qin, M., Manning, Z., Chen, M., Marquis, K.B., Averyt, M., Tignor and H.L. Miller. 2007. *Climate change 2007: The physical science basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC) 2007. Cambridge: Cambridge University Press
- Stafford, J.V. 2005. *Precision Agriculture 2005*. The Netherlands: Wagenigen Academic Publishers
- Stevenson, F. 1982. *Humus Chemistry: Genesis, composition and reactions*. New York: John Wiley and Sons.
- Strickland, M.S., Osburn, E., Lauber, C., Fierer, N., and M.A. Bradford .2009. Litter quality is in the eye of the beholder: initial decomposition rates as a function of inoculum characteristics. *Functional Ecology* 23(3):627-636.
- Symington, C.F. 1943. *Foresters Manual on Dipterocarps*. Malayan Forest Records No. 16. Kuala Lumpur: University of Malaya Press.
- Talbot, J.M. and Treseder, K.K. 2012. Interactions among lignin, cellulose, and nitrogen drive litter chemistry-decay relationships. *Ecology* 93(2): 345-354.

- Tang, C.Q. 2006. Forest vegetation as related to climate and soil conditions at varying altitudes on a humid subtropical mountain, Mount Emei, Sichuan, China. *Ecological Research* 21(2): 174-180.
- Tangang, F., Juneng, L., and Ahmad, S. 2007. Trend and interannual variability of temperature in Malaysia: 1961-2002. *Theoretical and Applied Climatology*, 89(3): 127-141.
- Tangang, F. 2007. Climate change and global warming: Malaysia perspective: Malaysia perspective and challenges. Paper presented at the Syarahan Umum, Universiti Kebangsaan Malaysia, Kuala Lumpur.23 March 2007.
- Thi, B.K., Lee, S.S., Hattori, T., Ota, Y., and Lee, S.L. 2012. Assessment of growth rate and enzymatic activities of *Microporus* spp.: A preliminary data. Malaysian Science and Technology Congress: Transforming the S&T Resources for Sustainable High Income Society (MSTC 2012). 19-21 November 2012. Sime Darby Convention Centre, Kuala Lumpur. (CD publ.)
- Thomas, G.W. 1982. Exchangeable cations. In: *Methods of soil analysis Part 2*, *Chemical and Microbiological Properties.* ed. A.L., Page, R.H., Miller, and D.R. Keeney, (Eds.). pp. 159-165. Madison: ASA and SSSA
- Thwaites, R.N. 2000. From biodiversity to geodiversity and soil diversity. A spatial understanding of soil in ecological studies of the forest landscape. *Journal of Tropical Forest Science* 12(2): 388-405.
- Tian G., Kang, B.T., and Brussard L.1992. Biological effects of plant residues with contrasting chemical composition under tropical conditions decomposition and nutrient release. *Soil Biology and Biochemistry* 25(6): 731-737.
- Tinker, P.B. and Ineson, P. 1990. Soil organic matter and biology in relation to climate change. In *Soils on a warmer earth-effects of expected climate change on soil processes, with emphasis on the tropics and sub-tropics*, Developments in Soil Science 20.ed. Scharpenseel, H.W., Schomaker, M., and A.Ayoub, pp.71-88. Amsterdam: Elsevier
- Titus B, Visser S, Wein R, and Zoltai S. 1999. Litter decomposition rates in Canadian forests. *Global Change Biology* 5: 75–82.
- Torreta, N.K. and Takeda, H.1999. Carbon and nitrogen dynamics of decomposing leaf litter in a tropical hill evergreen forest. *European Journal of Soil Biology* 35:57–63.
- Trum, F., Titeux, H., Delvaux, B., and Gilkes, R. 2010. Effects of manganese concentration on beech leaf litter decomposition: Results from field and laboratory experiments. In: *Proceedings of the 19th World Congress on Soil Science: Solutions for a changing world* ed. R.J., Gilkes, N. Prakongkep, N.

- ISBN 978-0-646-53783-2; Published on DVD; http://www.iuss.org; Symposium 4.2.1, 1-6 August 2010, Brisbane Australian IUSS.
- Turner B.L. 2010. Soil carbon in tropical forests: An inventory in CTFS forest dynamics plots. Paper presented in Soil Carbon Seminar, 26 March 2010, Forest Research Institute Malaysia.
- Updegraff, K., Zimmerman, P.R., Price, M., and Capehart, W.J., 2005. C-Lock: An online system to standardize the estimation of agricultural carbon sequestration credits. *Fuel Processing Technology* 86 (14-15): 1695-1704.
- Valentini, C.M.A., Sanches, L., de Paula, S.R.R., Vourlitis, G.L., de Souza Nogueira, J., Pinto, O. B., and de Almeida Lobo, F. 2008. Soil respiration and aboveground litter dynamics of a tropical transitional forest in northwest Mato Grosso, Brazil. *Journal of Geophysical Research: Biogeosciences* 113(G1).
- Van Breugel, M., Ransijn, J., Craven, D., Bongers, F., and Hall, J.S. 2011. Estimating carbon stock in secondary forests: Decisions and uncertainties associated with allometric biomass models. *Forest Ecology and Management* 262(8): 1648-1657.
- Van Cleve, K. and Powers, R.F.1995. Soil carbon, soil formation and ecosystem development. In *Carbon forms and functions in soils*, ed. W.W., McFee, and M. Kelly. pp.155-200. Wisconsin: Soil Science Society of America.
- Van Meirvenne, M., Pannier, J., Hofman, G., and Louwagie, G. 1996. Regional characterization of the long-term change in soil organic carbon under intensive agriculture. *Soil Use and Management* 12(2): 86-94.
- Van Soest, P.J. 1963. Use of detergents in analysis of fibrous feeds: a rapid method for the determination of fiber and lignin. *J. Assoc. Off. Agric. Chem*, 46: 829-835.
- Varallyay, G. 1990. Influence of climatic change on soil moisture regime, texture, structure and erosion. In *Soils on a warmer earth-effects of expected climate change on soil processes, with emphasis on the tropics and sub-tropics, Developments in Soil Science 20.*ed. H.W., Scharpenseel, M., Schomaker, and A. Ayoub, pp.39-49. Amsterdam: Elsevier.
- Vitousek, P.M. 1994. Beyond global warming: ecology and global change. *Ecology* 75(7): 1861-1876.

- Vitousek, P.M., Turner, D.R., Parton, W.J., and Sanford, R.L. 1994. Litter decomposition on the Mauna Loa environmental matrix, Hawai'i: patterns, mechanisms, and models. *Ecology*: 418-429.
- Vitousek, P., Chadwick, O., Matson, P., Allison, S., Derry, L., Kettley, L., Luers, A., Mecking, E., Monastra, V., and Porder, S. 2003. Erosion and the Rejuvenation of Weathering-derived Nutrient Supply in an Old Tropical Landscape. *Ecosystems*, 6(8): 762-772.
- Vogt, K.A., Vogt, D.J., and Bloomfield, J. 1998. Analysis of some direct and indirect methods for estimating root biomass and production of forests at an ecosystem level. *Plant and Soil* 200: 71-89.
- Wagai, R., Mayer, L.M., Kitayama, K., and Knicker, H. 2008. Climate and parent material controls on organic matter storage in surface soils: A three-pool, density-separation approach. *Geoderma* 147(1-2):23-33.
- Wardle D.A., Zachrisson, O, Hörnberg, G., and Gallet C. 1997. The influence of island area on ecosystem properties. *Science* 277:1296–1299.
- Wang, S., Ruan, H. and Wang, B. 2009. Effects of soil microarthropods on plant litter decomposition across an elevation gradient in the Wuyi Mountains. *Soil Biology and Biochemistry* 41(5): 891-897.
- Wang, Z., Zhang, B., Song, K., Liu, D., Li, F., Guo, Z. and Zhang, S. 2008. Soil organic carbon under different landscape attributes in croplands of Northeast China. *Plant, Soil and Environment* 10: 420-427.
- Webster, R. and Oliver, M.A., 2007. Geostatistics for environmental scientists. Wiley & Sons Inc: John West Sussex
- Wilcke, W., Hess, T., Bengel, C., Homeier, J., Valarezo, C., and Zech, W. 2005. Coarse woody debris in a montane forest in Ecuador: Mass, C and nutrient stock, and turnover. *Forest Ecology and Management* 205(1): 139-147.
- Wiseman, C.L.S. and Püttmann, W. 2006. Interactions between mineral phases in the preservation of soil organic matter. *Geoderma* 134:109-118.
- Woodall, C.W., Perry, C.H., and Westfall, J.A. 2012. An empirical assessment of forest floor carbon stock components across the United States. *Forest Ecology and Management* 269: 1-9.
- Wopereis. M.C., Gascuel-Odoux, C., Bourrie. G., So and Net, G., 1988. Spatial variability of heavy metals in soil on a one hectare scale. *Soil Science* 146 (2):113-118.
- Wright S.F. and Upadhyaya, A. 1996. Extraction of an abundant and unusual protein from soil and comparison with hyphal protein of arbuscular mycorrhizal fungi. *Soil Science* 161(9): 575-586.

- Wyatt-Smith, J. 1964. *Manual of Malayan Silviculture for Inland Forest. Part III, Natural Regeneration*. Malayan Forest Records 23. Kuala Lumpur: Forest Research Institute Malaysia.
- Xu, M., Qi, Y., 2001. Soil-surface CO₂ efflux and its spatial and temporal variation in a young ponderosa pine plantation in northern California. *Global Change Biology* 7: 667–677.
- Yanai, R.D., Currie, W.S., and Goodale, C.L. 2003. Soil carbon dynamics afterforest harvest: an ecosystem paradigm reconsidered. *Ecosystems* 6(3): 197-212.
- Yimer, F., Ledin, S. and Abdelkadir, A. 2006. Soil organic carbon and total nitrogen stocks as affected by topographic aspect and vegetation in the Bale Mountains, Ethiopia. *Geoderma* 135: 335-344.
- Yohannes, Y., Shibistova, O., Abate, A., Fetene, M. and Guggenberger, G. 2011. Soil CO₂ efflux in an Afromontane forest of Ethiopia as driven by seasonality and tree species. *Forest Ecology and Management* 261:1090-1098.
- Zhang, C. and McGrath, D. 2004. Geostatistical and GIS analyses on soil organic carbon concentrations in grassland of southeastern Ireland from two different periods. *Geoderma* 119(3-4): 261-275.