



UNIVERSITI PUTRA MALAYSIA

***INFLUENCE OF DIFFERENT FAT REPLACERS AND DRYING
TECHNIQUES ON PHYSICOCHEMICAL CHARACTERISTICS AND
SENSORY ATTRIBUTES OF REGULAR AND INSTANT REDUCED-FAT
COFFEE CREAMER***

SIMIN HEDAYATNIA



**INFLUENCE OF DIFFERENT FAT REPLACERS AND DRYING
TECHNIQUES ON PHYSICOCHEMICAL CHARACTERISTICS
AND SENSORY ATTRIBUTES OF REGULAR AND INSTANT
REDUCED-FAT COFFEE CREAMER**

By

SIMIN HEDAYATNIA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia
in Fulfilment of the Requirement for the Degree of Master of Science**

January 2015

COPYRIGHT

All materials contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This thesis is dedicated to my beloved father and mother who are always giving me their unlimited support, love, patience and understanding.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement of the degree of Master of Science

INFLUENCE OF DIFFERENT FAT REPLACERS AND DRYING TECHNIQUES ON PHYSICOCHEMICAL CHARACTERISTICS AND SENSORY ATTRIBUTES OF REGULAR AND INSTANT REDUCED-FAT COFFEE CREAMER

By

SIMIN HEDAYAT NIA

January 2015

Chairman : Assoc. Prof. Seyed Hamed Mirhosseini

Faculty : Food Science and Technology

Coffee is one of the most popular soft drinks all around the world. Most of coffee drinkers prefer to add creamer and/or whitener to their coffee before consumption. Coffee creamers usually contain high amount of the saturated fat (15-40%). Therefore, the frequent consumption of the whitened coffee can induce many health issues (e.g. cardiovascular and chronic diseases) for coffee drinkers. In recent years, the demand for low- and reduced-fat products has been extensively increased. The aim of the present study was to formulate and characterize the reduced-fat coffee creamer with the most desirable characteristics comparable with commercial creamers.

The main objective of the present study was to investigate the effects of different type and content of fat replacer (i.e. inulin, 0, 2.5, 5 and 7.5%; maltodextrin, 0, 15, 20 and 25%, w/w) as well as different drying techniques (i.e. spray drying, drum drying and fluidized-bed drying) on physicochemical properties, microstructures, and sensory attributes of the regular-and instant reduced-fat creamers. The regular coffee creamers were produced by a single-stage drying (either spray drying or drum drying only); while the instant reduced-fat coffee creamers were produced by a double-stage drying (i.e. spray drying or drum drying along with fluidized-bed drying). Physicochemical properties of all formulated creamers were compared with the control (as a negative control) and commercial creamers (as a positive control).

The current study revealed that the physicochemical characteristics, microstructures, and sensory attributes of both regular-and instant reduced-fat creamers were significantly ($p \leq 0.05$) influenced by both fat replacers and drying techniques. Moisture content, water activity of regular-and instant creamers were notably decreased by increasing the concentration of maltodextrin and inulin. This could be due to significant ($p \leq 0.05$) increase in solid content of samples. The bulk density of regular-and instant creamers was dropped by increasing the content of target fat replacers and enlarging the particle size. The current study revealed that the

wettability, solubility, viscosity and glass transition temperature of the formulated creamer were significantly ($p \leq 0.05$) improved as the contents of inulin or maltodextrin in the creamer formulation were increased.

The present study revealed that spray dried reduced-fat creamer had smaller spherical or oval shape particles than the drum dried creamers; while drum dried samples had much bigger particles with irregular shape. In this study, the drum-dried creamers had darker colour (or lower lightness) than the spray-dried samples. This might be because of its higher drying temperature and longer residence time. The drum-dried creamers with markedly bigger particle size and lower moisture content had considerably lower bulk density than the spray-dried creamer.

The current study revealed that the instant reduced-fat creamer had higher glass transition temperature than the regular reduced-fat creamer. This could be explained by the fact that the instant reduced-fat creamer had markedly lower moisture content than the regular creamer because the application of fluidized-bed drying led to decrease the moisture content, water activity, bulk density and stickiness. The agglomeration induced by fluidized bed drying significantly increased the reconstitution properties (wettability and solubility), viscosity and glass transition temperature of the reduced fat creamer. The morphology analysis revealed that agglomeration caused by fluidized-bed drying resulted in bigger particles with more porous structure than the regular creamer. Finally, the current study revealed that the instant spray-dried creamer (containing 25% maltodextrin and 7.5% inulin) had better quality comparable with commercial creamer than the instant drum-dried sample with similar formulation. The current study showed that instant spray-and drum dried reduced-fat creamers containing high amounts of maltodextrin (25%, w/w) and inulin (7.5%, w/w) had the most desirable characteristics among all formulated creamers comparable with the commercial creamer.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**KESAN PELBAGAI JENIS LEMAK PEGANTI DAN TEKNIK
PENGERINGAN KE ATAS SIFAT FIZIOKIMIA DAN DERIA RASA KOPI
KRIMER KURANG LEMAK BIASA DAN SEGERA**

Oleh

SIMIN HEDAYATNIA

Januari 2015

Pengerusi : Profesor Madya. Seyed Hamed Mirhosseini

Fakulti : Sains dan Teknologi Makanan

Kopi merupakan salah satu minuman paling digemari di seluruh dunia. Majoriti peminum kopi biasanya memilih untuk menambah krimer dan/atau pemutih di dalam kopi. Namun begitu, kopi krimmer kebiasaanya mengandungi kandungan lemak tepu yang tinggi iaitu 15-40%. Pengambilan yang kerap akan menyebabkan pelbagai isu kesihatan akan timbul seperti penyakit kardiovaskular dan penyakit kronik. Justeru itu, permintaan terhadap produk kurang lemak semakin meningkat sejak kebelakangan ini. Kajian ini bertujuan untuk menghasilkan formulasi dan ciri-ciri kopi krimmer rendah lemak yang paling standing dengan krimmer komersial. Objektif utama kajian ini adalah untuk mengkaji kesan pelbagai jenis dan kandungan pengganti lemak (seperti inulin, 0, 2.5, 5 and 7.5%; maltodextrin, 0, 15, 20 dan 25%, w/w) selain penggunaan teknik pengeringan (seperti pengeringan semburan pengeringan dram dan pengeringan terbendalir katil) ke atas sifat fizikimia, struktur mikro dan deria rasa krimer rendah lemak biasa dan segera. Krimer kopi yang biasa dihasilkan dengan menggunakan pengeringan peringkat tunggal (sama ada pengeringan semburan atau pengeringan dram), manakala kopi krimer segera dihasilkan menggunakan pengeringan peringkat berganda (seperti pengeringan semburan atau pengeringan dram bersama terbendalir katil). Sifat fizikimia semua krimer berformulasi akan dibandingkan dengan krimer kawalan (sebagai kawalan negatif) dan krimer komersial (sebagai kawalan positif).

Kajian ini telah menunjukkan ciri fizikimia, struktur mikro dan deria rasa kedua-dua krimer biasa dan segera menghasilkan perbezaan yang ketara setelah teknik pengeringan digunakan ($p \leq 0.05$). Kandungan kelembapan, iaitu aktiviti air krimer biasa dan segera menunjukkan penurunan dengan peningkatan kepekatan maltodekstrin dan inulin. Ini mungkin disebabkan perbezaan yang ketara ($p \leq 0.05$) meningkat dalam kandungan sampel pepejal. Ketumpatan pukal krimer biasa dan segera menurun dengan peningkatan sasaran kandungan pengganti lemak dan pembesaran saiz zarah. Kajian ini juga menunjukkan bahawa kebolehbasaan, kelarutan, kelikatan dan suhu peralihan kaca krimer berformulasi menunjukkan

perbezaan yang ketara ($p \leq 0.05$) meningkat setelah kandungan inulin atau maltodekstrin ditingkatkan.

Kajian ini juga menunjukkan bahawa pengeringan semburan krimer kurang lemak mempunyai zarah berbentuk sfera atau bujur yang lebih kecil berbanding penggunaan pengeringan dram yang mempunyai zarah lebih besar dengan bentuk yang tidak teratur. Dalam kajian ini, krimer dari pengeringan dram mempunyai warna lebih gelap (atau kecerahan yang lebih rendah) daripada sampel krimer daripada pengeringan semburan. Ini mungkin kerana suhu pengeringan yang lebih tinggi dan masa pengeringan lama. Krimer daripada pengeringan dram dengan saiz zarah yang lebih besar dan kandungan lembapan yang lebih rendah mempunyai ketumpatan pukal lebih rendah daripada krimer pengeringan semburan.

Kajian ini juga menunjukkan krimer rendah lemak mempunyai suhu peralihan kaca yang lebih tinggi (T_g) daripada krimer yang biasa. Ini dapat dijelaskan oleh fakta yang mengatakan bahawa krimer segera rendah lemak mempunyai kandungan kelembapan yang rendah berbanding krimer biasa kerana penggunaan pengeringan terbendalir katil mengurangkan kandungan kelembapan, aktiviti air, ketumpatan pukal dan kelekitan. Aglomerasi disebabkan oleh pengeringan terbendalir katil meningkat dengan ketara sifat pelarut (kebolehasahan dan kelarutan), kelikatan dan suhu peralihan kaca krimer kurang lemak. Analisis morfologi mendedahkan aglomerasi disebabkan oleh pengeringan terbendalir katil menghasilkan zarah yang lebih besar dengan struktur lebih porous daripada krimer biasa.

Kesimpulannya, kajian ini menunjukkan bahawa krimer segera daripada pengeringan semburan (yang mengandungi 25% maltodekstrin dan 7.5% inulin) mempunyai kualiti yang lebih baik setanding dengan krimer komersial daripada kopi segera daripada pengeringan dram dengan formulasi yang sama. Kajian ini juga menunjukkan bahawa krimer segera pengeringan semburan dan krimer rendah lemak pengeringan dram yang mengandungi jumlah maltodekstrin yang tinggi (25%, w / w) dan inulin (7.5%, w / w) mempunyai ciri-ciri yang paling dikehendaki di kalangan semua Krimer berformulasi setanding dengan krimer komersial.

ACKNOWLEDGEMENTS

First and foremost, I give my greatest thanks to God for giving me the wisdom, ability, health and endurance to complete this degree.

I would like to express the deepest appreciation to my supervisor **Assoc. Prof. Dr. Seyed Hamed Mirhosseini** for his encouragement, ever-lasting support, guidance and supervision of this thesis. You have been a tremendous mentor for me. Without his supervision and constant help this dissertation would not have been possible.

I would also like to appreciate my other committee members: **Prof. Dr. Yazid Abd Manap** and **Assoc. Prof. Dr. Roselina Karim** for their encouragement, useful comments, and challenging questions. I would also like to express thanks to the technical staff of Faculty of Food Science and Technology in Food Biochemistry and Engineering Laboratories. A very special thank you goes to **Mr. Amran** and **Madam Rosmawati Othman** for being so supportive and warm welcome.

The most importantly, I take pleasure in expressing my gratitude to my father, **Saeed** and my mother **Marzieh**, who always encouraged me to obtain higher education. They bore me, raised me, supported me, taught me, and loved me. To them I dedicate this thesis. I'm so lucky to have them be my parents.

I want to thank my beloved sisters **Sima** and **Mahshad** who have always stimulated me to stay open-minded and to keep on doing the best I could. I would also like to thank **Azad** and **Rozhin** for all emotional support. Thanks for being with me.

A special thanks to my caring, loving, and supportive fiancé **Rasa**. Words cannot express how grateful I am to my fiancé for all of the sacrifices that you've made on my behalf. Himself, Rasa has been my best friend and a great companion, loved, supported, encouraged, entertained, and helped me get through this agonizing period in the most positive way. My heartfelt thanks.

Lastly, my special thanks to my best friends **Maryam, Elham, Mahdokht, Homa, Hanie** and **Adila** for helping me get through the difficult times, and for all the emotional support, camaraderie, entertainment, and caring they provided.

I certify that a Thesis Examination Committee has met on 15 January 2015 to conduct the final examination of Simin Hedayatnia on her thesis entitled "Influence of Different Fat Replacers and Drying Techniques on Physicochemical Characteristics and Sensory Attributes of Regular and Instant Reduced-Fat Coffee Creamer" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Yaya Rukayadi, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Jinap bt Selamat, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Internal Examiner)

Russly b Abdul Rahman, PhD

Professor
Halal Products Research Institute
Universiti Putra Malaysia
(Internal Examiner)

Kamariah Long, PhD

Senior Lecturer
Center for Biotechnology
MARDI
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 19 March 2015

This Thesis was submitted to the Senate of University Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the supervisory committee were as follows:

Seyed Hamed Mirhosseini, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairperson)

Mohd Yazid Abdul Manap, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Roselina Karim, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

BUJANG KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Simin Hedayatnia GS31758

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____

Name of

Chairman of

Supervisory

Committee: _____

Signature: _____

Name of

Member of

Supervisory

Committee: PROF. DATO' DR. MOHD. YAZID ABD. MANAP, D.S.I.S.
Pengarah
Institut Penyelidikan Produk Halal
Universiti Putra Malaysia
43400 UPM Serdang, Selangor Darul Ehsan, Malaysia

Signature: _____

Name of

Member of

Supervisory

Committee: PROF. MARYA DR. ROSELINA KARIM
Pensyarah
Jabatan Teknologi Makanan
Fakulti Sains & Teknologi Makanan
Universiti Putra Malaysia,
43400 UPM Serdang, Selangor,
Malaysia

TABLE OF CONTENTS

ABSTRACT	Page
ABSTRAK	i
ACKNOWLEDGEMENTS	iii
APPROVAL	v
DECLARATION	vi
LIST OF TABLES	viii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xv
	xviii
 CHAPTER	
 1 INTRODUCTION	 1
 2 LITERATURE REVIEW	 4
2.1 Coffee	4
2.2 Coffee Creamer	5
2.2.1 History	8
2.2.2 Consumption of Coffee Creamer	8
2.2.3 Classification of Coffee Creamers	9
2.2.3.1 Coffee Creamer in Powder Form	9
2.2.3.2 Coffee Creamer in Liquid Form	9
2.3 Coffee Creamer Composition	10
2.3.1 Fat	10
2.3.2 Sweetener	10
2.3.3 Emulsifier and Stabilizer	11
2.3.4 Protein	11
2.3.5 Other Ingredients	12
2.3.5.1 Fat Replacers	12
2.4 Drying Techniques	16
2.5 Instantization and Agglomeration	18
 3 METHODOLOGY	 20
3.1 Materials and Methods	20
3.1.1 Materials and Chemicals	20
3.1.2 Methodology	20
3.1.3 Production of Reduced-Fat Creamer	22
3.1.3.1 Preparation of Creamer Emulsion	22
3.1.3.1.1 Commercial Sample (ControlPositive)	23
3.1.3.1.2 Control Sample (Control Negative)	23
3.1.3.2 Homogenization Process	24
3.1.3.3 Spray Dryer Operation	25
3.1.3.4 Drum Dryer Operation	26
3.1.3.5 Fluidized-Bed Dryer Procedure	26
3.2 Analytical Tests	27
3.2.1 Moisture Content and Water Activity Analysis	27
3.2.2 Measurement of Bulk Density	28
3.2.3 Morphology Structure Analysis	28

3.2.4	Analysis of Droplet Size Distribution	29
3.2.5	Determination of Wettability	29
3.2.6	Solubility Index Measurement	29
3.2.7	Measurement of Apparent Viscosity	29
3.2.8	Glass transition temperature Analysis	30
3.2.9	Lightness (L*) Measurement	30
3.2.10	Sensory Evaluation	30
3.3	Statistical Design and Data Analysis	31
4	RESULTS AND DISCUSSION	34
4.1	Preliminary Study	34
4.1.1	Application of Protein in Creamer Formulation	34
4.1.2	Application of Emulsifier and Stabilizer in Creamer Formulation	35
4.1.3	Establishment of Desirable Spray Drying Condition	35
4.1.4	Establishment of Desirable Drum Drying Condition	36
4.1.5	Establishment of Desirable Fluidized-Bed Drying Condition	37
4.2	Effects of Different Compositions and Drying Techniques on Physicochemical Characteristics of Regular Reduced-Fat Creamers	37
4.2.1	Moisture Content and Water of Regular Reduced-Fat Creamers	37
4.2.2	Bulk Density of Regular Reduced-Fat Creamers	41
4.2.3	Morphology Structure of Regular Reduced-Fat Creamers	43
4.2.4	Volume-Weighted Mean (or Particle Size) of Regular Reduced-Fat Creamers	45
4.2.5	Wettability of Regular Reduced-Fat Creamers	49
4.2.6	Solubility of Regular Reduced-Fat Creamers	51
4.2.7	Apparent Viscosity of Regular Reduced-Fat Creamer	54
4.2.8	Glass transition temperature of Regular Reduced-Fat Creamers	56
4.2.9	Colour of Regular Reduced-Fat Creamers	60
4.2.10	Single and Interaction Effects of Drying Variables on Characteristics of Regular Creamers	64
4.3	Effects of Different Compositions and Fluidized Bed Drying on Physicochemical Characteristics of Instant Reduced-Fat Creamer	67
4.3.1	Moisture Content and Water Activity of Instant Reduced-Fat Creamers	67
4.3.2	Bulk Density of Instant Reduced-Fat Creamers	70
4.3.3	Morphology Structure of Instant Reduced-Fat Creamers	72
4.3.4	Volume-Weighted Mean (or Particle Size) of Instant Reduced-Fat Creamers	74
4.3.5	Wettability of Instant Reduced-Fat Creamers	77
4.3.6	Solubility of Instant Reduced-Fat Creamers	80
4.3.7	Apparent Viscosity of Instant Reduced-Fat Creamers	82
4.3.8	Glass transition temperature of Instant Reduced-Fat Creamers	83
4.3.9	Colour of Instant Reduced-Fat Creamers	86
4.3.10	Sensorial Analysis	88
4.3.11	Single and Interaction Effects of Drying Variables on Characteristics of Regular-Creamers	91

5 SUMMARY, CONCLUSION AND RECOMMENDATIONS	93
5.1 Summary and conclusion	93
5.2 Recommendations	94
REFERENCES	95
APPENDICES	111
BIODATA OF STUDENT	120
LIST OF PUBLICATIONS	121



LIST OF TABLES

Table	Page
1.1	1
2.1	6
2.2	8
2.3	14
2.4	15
2.5	15
2.6	16
2.7	17
2.8	18
3.1	22
3.2	23
3.3	24
3.4	32
3.5	33
4.1	34
4.2	35
4.3	35
4.4	36
4.5	36
4.6	37
4.7	41
4.8	65
4.9	65
4.10	69
4.11	72
4.12	76

4.13	<i>t-test</i> analysis on wettability of regular-and-instant spray-and drum dried creamers	79
4.14	<i>t-test</i> analysis on solubility of regular-and-instant spray-and-drum dried creamers	81
4.15	<i>t-test</i> analysis on viscosity of regular-and-instant spray-and-drum dried creamers	83
4.16	<i>t-test</i> analysis on glass transition temperature of regular-and-instant spray-and-drum dried creamers	85
4.17	<i>t-test</i> analysis on lightness (L*) of regular-and instant spray-and drum dried creamers	87
4.18	Sensory analysis of tested instant creamers	91
4.19	Two-way ANOVA for different analysis of instant spray-dried creamers	91
4.20	Two-way ANOVA for different analysis of instant drum-dried creamers	91

LIST OF FIGURES

Figure		Page
2.1	Average annual coffee consumption in some Asian countries	5
2.2	Classification of coffee creamer	9
2.3	Obesity in international perspective	11
2.4	Four drying process steps	16
2.5	(A) Schematic mechanism of agglomeration process, (B) Spray-dried creamer agglomeration process	19
3.1	General methodology	21
3.2	The drum-dried samples containing 0% fat replacers (maltodextrin and inulin)	24
3.3	Laboratory scale spray drying operation	25
3.4	Aluminium packed spray-dried creamer	25
3.5	A) A Schematic diagram of Simon drum dryer B) Laboratory scale Simon drum dryer	26
3.6	(A) Schematic diagram of fluidised-bed dryer and (B) Laboratory scale fluidised-bed dryer	27
3.7	Scanning Electron Microscope (SEM) instrument	28
4.1	Moisture content of regular spray-dried reduced-fat creamers	38
4.2	Water activity of regular spray-dried reduced-fat creamers	38
4.3	Moisture content of regular drum-dried reduced-fat creamers	39
4.4	Water activity of regular drum-dried reduced-fat creamers	40
4.5	Bulk density of regular spray-dried reduced-fat creamers	42
4.6	Bulk density of regular drum-dried reduced-fat creamers	43
4.7	Scanning electron micrographs (SEM) of creamers produced by spray drying (a and b) and drum drying (c and d) (magnification of 500)	44
4.8	Volume-weighted mean of regular Spray-dried reduced-fat creamers	46
4.9	Size enlargements induced with increasing maltodextrin content for spray-dried creamers containing 0% inulin and different maltodextrin content A (0%), B (15%), and C (25%)	47
4.10	Volume-weighted mean of regular drum-dried reduced-fat creamers	48
4.11	Drum-dried creamers with different volume weighed mean, (A: 15% maltodextrin and 0% inulin; B: 25% maltodextrin and 0% inulin)	49
4.12	Wettability of regular spray-dried reduced-fat creamers	50
4.13	Wettability of regular drum-dried reduced-fat creamers	51
4.14	Solubility of regular spray-dried reduced-fat creamers	52
4.15	Solubility of regular drum-dried reduced-fat creamers	53
4.16	Solubility improvement of regular drum-dried creamer A) 0% maltodextrin , B) 15% maltodextrin and C) 25% maltodextrin creamers by increasing the maltodextrin concentration (inulin 0%)	53
4.17	Apparent viscosity of regular spray-dried reduced-fat creamers	55
4.18	Apparent viscosity of regular drum-dried reduced-fat creamers	56
4.19	Glass transition temperature of regular spray-dried creamers	57

4.20	DSC Thermogram analyzed for T_g of regular spray-dried powders	58
4.21	Glass transition temperature of regular spray-dried creamers	58
4.22	DSC Thermogram analyzed for T_g of regular drum-dried powders	59
4.23	DSC Thermogram of glass transition temperature of the regular spray-dried powders	60
4.24	Lightness (L^*) of regular spray-dried reduced-fat creamers	61
4.25	Lightness (L^*) of regular drum-dried reduced-fat creamers	62
4.26	Images of spray-dried creamers (a: commercial creamer; c: control creamer; e: creamer) and drum dried creamers (b: commercial creamer; d: control creamer; f: creamer)	63
4.27	Moisture content of instant spray-dried reduced-fat creamers	67
4.28	Water activity of instant spray-dried reduced-fat creamers	68
4.29	Moisture content of instant drum-dried reduced-fat creamers	68
4.30	Water activity of instant drum-dried reduced-fat creamers	69
4.31	Effects of agglomeration process on the moisture content of the regular-and instant spray-dried creamers (15% Maltodextrin and 5% inulin)	70
4.32	Bulk density of instant spray-dried reduced-fat creamers	70
4.33	Bulk density of instant drum-dried reduced-fat creamers	71
4.34	Microstructure of regular-and-instant spray-dried creamers (a, b) and regular-and-instant drum-dried creamers (c, d)	73
4.35	Scanning electron micrographs (SEM) of individual instant reduced-fat creamer particles dried using spray drying followed by fluidized-bed drying (a), drum drying followed by fluidized-bed drying (b)	74
4.36	Volume-weighted mean of instant spray-dried reduced-fat creamers	75
4.37	Volume-weighted mean of instant drum-dried reduced-fat creamers	75
4.38	Particle size of A) regular and B) instant spray-dried reduced-fat creamer containing 25% maltodextrin and 0% inulin	76
4.39	Particle size of A) regular and B) instant drum-dried reduced-fat creamer containing 25% maltodextrin/ 0% inulin	77
4.40	Wettability of instant spray-dried reduced-fat creamers	78
4.41	Wettability of instant drum-dried reduced-fat creamers	79
4.42	Solubility of instant spray-dried reduced-fat creamers	80
4.43	Solubility of instant drum-dried reduced-fat creamers	81
4.44	Apparent viscosity of instant spray-dried reduced-fat creamers	82
4.45	Apparent viscosity of instant drum-dried reduced-fat creamers	83
4.46	Glass transition temperature of instant reduced-fat spray-dried creamers	84
4.47	Glass transition temperature of instant reduced-fat drum-dried creamers	85
4.48	DSC thermograms of the drum-dried creamer a) before agglomeration process (regular creamer) b) after agglomeration process (instant creamer)	86
4.49	Lightness (L^*) of instant reduced-fat spray-dried creamers	86
4.50	Lightness (L^*) of instant reduced-fat drum-dried creamers	87

4.51	Significant effect of agglomeration process on colour of drum-dried creamers	88
4.52	The effect of different chosen-creamers on taste evaluation	89
4.53	The effect of different chosen-creamers on aroma evaluation	89
4.54	The effect of different chosen-creamers on colour evaluation	90
4.55	The effect of different chosen-creamers on overall accetibity evaluation	90



LIST OF ABBREVIATIONS

ANOVA	Analysis of Variance
DE	Dextrose equivalent
DPHP	Di-potassium hydrogen phosphate
e.g	Exempli gratia
etc	Et cetera
et al	Et alibi
g	Gram
pH	Hydrogen ion exponent
IN	Inulin
Kcal/g	Kilocalories per gram
Kg	Kilogram
Kj/g	kilojoules per gram
Kpa	kilopascals
kWh.tonne-1	kilowatt hour per tonne
MA	Maltodextrin
μL	Microliter
mL	Milliliter
mm	Millimeter
mg	Milligram
mL/min	Milliliters per minute
min	Minute
MPa	Mega Pascal
rpm	Revolution per minute
DRFC	Drum-dried reduced-fat creamer
SRFC	Spray-dried reduced-fat creamer
RS	Rotation speed
SEM	Scanning electron microscopy
SMP	Skim-milk powder
SP	Steam pressure
T	Temperature
h	Time (Hour)
s	Time (Second)
US	United States
D _{3,4}	Volume-weighted mean particle size, μm
H ₂ O	Water
w/w	Weight/weight
WPNI	Whey Protein Nitrogen Index
°C	Degree centigrade
≤	Equal or less
%	Percentage

CHAPTER 1

INTRODUCTION

The recent changes in the lifestyles of numerous people worldwide have led to an increase in demand for convenient health food products, along with healthier foods in general, such as food products containing low-fat content and high fibre level, which produce lower energy level in the human's body (Nishinari, 2009). The recommended daily intakes of total fibre for adults are 38 and 35 g for men and women, respectively (Trumbo et al., 2002). Fat is the most concentrated source of energy in the diet program that is providing 9 kcal/g energy as compared to 4 kcal/g for proteins and carbohydrates (American Heart Association, 1996). However, consumers prefer to consume foods with minimal or reduced-fat irrespective of the food taste. As shown by the Calorie Control Council (CCC, 1996), 88% of American adults prefer to consume the low- or reduced-fat or even fat-free foods and beverages because high fat daily intake is always associated with high risk for the obesity, cancer, chronic and cardiovascular diseases. Table 1.1 summarized some of the health issues that describing the consumer preferences for low-fat products. One of the most common strategies for low fat products is to use fat replacers to compensate for the shortcomings in the sensory attributes and textural properties of the product (Sandrou and Arvantoyannis, 2000). However, this strategy does not guarantee the textural properties (such as creaminess) and consumer acceptability (Szczeniak, 2002). Food scientists have conducted extensive research to develop an "ideal fat replacer" that could provide similar taste and functional properties comparable to conventional fat, but without inducing any negative side effects on the human health (Akoh, 1998).

Table 1.1. Main consumer preference to have reduced-fat products

Why people use reduced-fat food products?	Percentages%
To stay in better overall health	77
To eat or drink healthier food and beverages	71
To reduce fat intake	68
To reduce cholesterol	61
To maintain current weight	57
To reduce calories	56
To maintain an attractive physical appearance	52
To reduce weight	43
For refreshment or taste	39
To help with a medical condition	31

Source: Calorie Control Council (CCC) 1996, natural survey

Coffee creamer, also known as "coffee whitener" or "coffee sweetener" are liquid or granular substances intended to substitute for milk or cream as an additive to coffee or other beverages. Coffee additives are dried milk concentrates, evaporated

milk, coffee cream, liquid milk, and coffee whiteners or creamer (Kelly et al., 1999). A desired or preferred coffee creamer is supposed to have certain characteristics in terms of solubility, stability, whitening ability and viscosity (Golde and Schmidt, 2005; Tuot et al., 2014). Coffee creamer should remain physically stable during storage and its viscosity should be constant over the time of storage. Coffee creamer should be dissolved rapidly in the hot water without separation of its components. In addition, it should provide a good whitening effect after adding to hot coffee or similar hot beverages (Oldfield and Singh, 2005).

One of the main health issues for coffee drinkers is the presence of high percentage of fat in creamer formulation. In this regard, the reduced-fat creamer can be alternatively produced by the partial replacement of fat portion with fat replacer components (i.e. fat replacers such as maltodextrin and inulin). Maltodextrin is one of the most popular polysaccharide-based fat replacers. It has many industrial applications based on the degree of starch hydrolysis. Furthermore, it is a white powder with low bulk density and soluble in water which is widely used as a texture modifier, gelling agent, fat replacer, volume enhancer, and encapsulation agent (Kiessling and Zeller, 2005). Inulin is another type of water soluble carbohydrate with a neutral taste and minimal side effects on organoleptic attributes of the food product (El-Nagar et al., 2002). In the current study, the effects of different concentrations of maltodextrin and inulin on physiochemical characteristics and organoleptic attributes of the regular-and instant reduced-fat creamers were investigated.

The characteristics of coffee creamer are also highly influenced by the processing conditions. Drying is the most important processing step as it has different effect on the characteristics of powder products. Spray drying is one of the most common techniques applied for different food products (Chegini and Ghobadian, 2005; Chavez and Ledebor, 2007) such as, creamer powder (Kiessling and Zeller, 2005; Beeson and Erickson, 2001), milk powder (Yazdanpanah and Langrish, 2011) and yogurt (Koc et al., 2010). However, it has several technical disadvantages such as high energy consumption, thermal degradation and production of the amorphous particles (White and Cakebread, 1966). Drum-drying is another drying technique that is widely used in bakery goods, beverages, cereal and dairy foods (Pua et al., 2010). The main advantages of drum drying are high drying rate and low production cost compared to other drying techniques (Vega et al., 2001). Moreover, further agglomeration is highly recommended to improve the quality and reconstitution properties of spray-and drum dried powders. In this regards, fluidized-bed drying is widely used for agglomeration purpose after drying process especially for spray drying. This may be possibly lead to induce further crystallization (Yazdanpanah and Lngrish, 2011). The main research questions were as follows:

- Whether different drying techniques and conditions can significantly affect the physicochemical characteristics and overall acceptability of the regular-and instant reduced-fat creamer?

- Is there any significant difference among all formulated creamers before and after fluidized-bed drying? Or is there any significant difference between physicochemical characteristics and overall acceptability of the regular-and instant reduced-fat creamers?
- Which one of fat replacers and drying techniques can provide creamer with more desirable characteristics and overall acceptability?
- Is there any significant difference between commercial creamers and newly formulated reduced-fat creamer in terms of overall quality and acceptability?

In this study, the effects of different drying techniques (i.e. drum-drying, spray-drying and fluidized-bed drying) and type and concentration of the fat replacers (maltodextrin and inulin) on physicochemical characteristics, functional properties and overall acceptability of the regular-and instant reduced-fat creamer were all investigated. The efficiency of different drying techniques and creamer composition were determined by assessing moisture content, water activity, bulk density, morphology structure, particle size distribution, wettability, solubility, viscosity, and glass transition temperature, colour intensity, and sensory attributes of various formulated creamers. The main goal of the present work was to produce the regular-and instant reduced-fat creamers with desirable physicochemical and functional characteristics comparable with commercial coffee creamers. In the current study, the specific objectives were as follows:

- To investigate the effect of type and content of fat replacers on physicochemical characteristics and overall acceptability of regular-and instant reduced-fat coffee creamer.
- To evaluate the effect of drying techniques on physicochemical characteristics and overall acceptability of regular-and instant reduced-fat coffee creamer.

REFERENCES

- Abadio, F. D. B., Domingues, A. M., Borges, S. V., & Oliveira, V. M. (2004). Physical properties of powdered pineapple (*Ananas comosus*) juice effect of maltedextrin concentration and atomization speed. *Journal of Food Engineering*, 64(3), 285-287.
- Abdenouri, N., Idlimam, A., & kouhila, M. (2010). Sorption Isotherms and Thermodynamic Properties of Powdered Milk. *Chemical Engineering. Comminucation*, 197(2), 1109–1125.
- Abdullah, A., Malundo, T. M. M., Resurreccion, A. V. A., & Beuchat, L. R. (1993). Descriptive sensory profiling for optimizing the formula of a peanut milk-based liquid coffee whitener. *Journal of Food Science*, 58(1), 120-123.
- Abiad, M. G., Gonzalez, D. C., Mert, B., Campanella, O. H., & Carvajal, M. T. (2010). A novel method to measure the glass and melting transitions of pharmaceutical powders. *International Journal of Pharmaceutics*, 396(1), 23-29.
- Aguilar, C. A., & Ziegler, G. R. (1994). Physical and microscopic characterization of dry whole milk with altered lactose content. 2. Effect of lactose crystallization. *Journal of Dairy Science*, 77(5), 1198-1204.
- Akoh, C. C. (1998). Fat replacers. *Food Technology*, 52(3), 47-53.
- American Dairy Products Institute. (1992). Standards for grades of dry milk including method of analysis. Bull. 916. Chicago, IL, USA: ADPI.
- American Heart Association. (1996). *Dietary guidelines for healthy American Adults: A statement for physicians and health professionals*. American Heart Association (pp.1795-1800).
- Amid, B. T., & Mirhosseini, H. (2012). Effect of different purification techniques on the characteristics of heteropolysaccharide-protein biopolymer from durian (*Durio zibethinus*) seed. *Molecules*, 17(9), 10875-10892.
- Anandaraman, S., & Reineccius, G. A. (1986). Stability of encapsulated orange peel oil. *Food technology (USA)*, 40, 88–93.
- Anastasiades, A., Thanou, S., Loulis, D., Stapatoris, A., & Karapantsios, T. D. (2002). Rheological and physical characterization of pregelatinized maize starches. *Journal of Food Engineering*, 52(1), 57-66.
- Anema, S.G., Pinder, D.N., Hunter, R.J., & Hemar, Y. (2006). Effects of storage temperature on the solubility of milk protein concentrate (MPC85). *Food Hydrocolloids*, 20, 386–393.

- Aravind, N., Sissons, M. J., Fellowes, C. M., Blazek, J., Gilbert, E. P. (2012). Effect of inulin soluble dietary fibre addition on technological, sensory, and structure properties of durum wheat spaghetti. *Food Chemistry*, 993(6)-1002.
- Arcia, P. L., Costell, E., & Tárrega, A. (2011). Inulin blend as prebiotic and fat replacer in dairy desserts: optimization by response surface methodology. *Journal of Dairy Science*, 94(5), 2192-2200.
- A/S Niro Atomizer. Analytical Methods for Dry Milk Products, 4th Ed; Niro: Copenhagen, Denmark, 1978.
- Astrup, A., Dyerberg, J., Elwood, P., Hermansen, K., Hu, F. B., Jakobsen, M. U., & Willett, W. C. (2011). The role of reducing intakes of saturated fat in the prevention of cardiovascular disease: where does the evidence stand in 2010?. *The American Journal of Clinical Nutrition*, 93(4), 684-688.
- Barbosa-Canovas, G. V., Malave-Lopez, J., & Peleg, M. (1987). Density and compressibility of selected food powders mixture. *Journal of Food Process Engineering*, 10(1), 1-19.
- Barbosa-Canovas, G.V., Juliano, P., & Yan, H. (2005). Physical and chemical properties of food powders. In: *Encapsulated and Powdered Foods*, pp. 40-71. Onwulata, C., Ed., pp.40-71. CRC Press, Boca Roton, FL.
- Barbosa-Canovas, G.V., Rivas, E. O., Juliano, P., & Yan, H. (2005). Size enlargement. In: *Food powders: Physical properties, processing and functionality*, pp. 175-198. Kluwer Academic/Plenum, New York.
- Baruah, B. K., Das, B., Medhi, C., & Misra, A. K. (2012). Fertility status of soil in the tea garden belts of golaghat district, Assam, India. *Journal of Chemistry*.
- Bayarri, S., Costell, E., McClements, D. J., & Decker, E. A. (2009). Optimising the flavour of low-fat foods. *Designing functional foods: measuring and controlling food structure breakdown and nutrient absorption*, 431-452.
- Beeson, C. A., & Erickson, L. (2001). *U.S. Patent No. 6,287,616*. Washington, DC: U.S. Patent and Trademark Office.
- Bhandari, B. R., & Howes, T. (1999). Implication of glass transition for the drying and stability of dried foods. *Journal of Food Engineering*. 40(1), 71-79.
- Bhandari, B. R., Datta, N., & Howes, T. (1997a). Problem associated with spray drying of sugar- rich foods. *Drying Technology*, 15(2), 671-684.
- Bhandari, B. R., Datta, N., Crooks, R., Howes, T., & Rigby, S. (1997b). A semi empirical approach to optimise the quantity of drying aids required to spray dry sugar-rich foods. *Drying Technology*, 15(10), 2509-2525.
- Bhandari, B., & Howes, T. (2005). Relating the stickiness property of foods undergoing drying and dried products to their surface energetics. *Drying Technology*, 23(4), 781-797.

- Bhupathiraju, S. N., Pan, A., Malik, V. S., Manson, J. E., Willett, W. C., & van Dam, R. M. (2013). Caffeinated and caffeine-free beverages and risk of type 2 diabetes. *The American Journal of Clinical Nutrition*, 97(1), 155–166.
- Blecker, C., Fougny, C., Van Herck, J. C., Chevalier, J. P., & Paquot, M. (2002). Kinetic study of the acid hydrolysis of various oligofructose samples. *Journal of Agricultural and Food Chemistry*, 50(6), 1602-1607.
- Bonazzi, C., & Bimbenet, J. J., (2008). Sechage des produits alimentaires – Mat ériels et applications, in Techniques de l'ingénieur- 18j 1 Quality Changes in Food Materials as Influenced by Drying Processes Traité Agroalimentaire F3. Editions T. I., Paris, France, p. 3002, 1–17.
- Bonazzi, C., & Dumoulin, E. (2011). Quality changes in food materials as influenced by drying processes . In *Modern Drying Technology* ; Tsotsas , E. ; Mujumdar , A.S. , Eds.; Wiley-VCH Verlag: Weinheim , Germany, 1 – 20.
- Borderías, A. J., Sánchez-Alonso, I., & Pérez-Mateos, M. (2005). New applications of fibres in foods: addition to fishery products. *Trends in Food Science & Technology*, 16(10), 458-465.
- Brennan, C. S., & Tudorica, C. M. (2008). Carbohydrate-based fat replacers in the modification of the rheological, textural and sensory quality of yoghurt: comparative study of the utilisation of barley beta-glucan, guar gum and inulin. *International Journal of Food Science & Technology*, 43(5), 824-833.
- Bröckel, U., Wahl, M., Kirsch, R., & Feise, H. J. (2006). Formation and growth of crystal bridges in bulk solids. *Chemical Engineering & Technology*, 29(6), 691-695.
- Buffo, R. A., Probst, K., Zehentbauer, G., Luo, Z., & Reineccius, G. A. (2002). Effects of agglomeration on the properties of spray-dried encapsulated flavours. *Flavour and Fragrance Journal*, 17(4), 292-299.
- Calorie Control Council. (1996). Fat reduction in food. (pp.111). Calorie Control Council. Atlanta, GA.
- Calorie Control Council. (2004). Fat replacers: Food ingredients for healthy eating, [cited:26January2006]. Available from: <http://www.caloriecontrol.org/fatreprint.htm>.
- Caparino, O. A., Tang, J., Nindo, C. I., Sablani, S. S., Powers, J. R., & Fellman, J. K. (2012). Effect of drying methods on the physical properties and microstructures of mango (Philippine ‘Carabao’ var.) powder. *Journal of Food Engineering* 111(1), 135–148.
- Caric ´, M. (2003). Milk powders: Types and manufacture and physical and functional properties of milk powders. In H. Roginski, J. W. Fuquay, & P. F. Fox (Eds.), *Encyclopedia of Dairy Sciences* (pp. 1869–1880). New York: Academic Press.

- Chávez, B. E., & Ledebøer, A. M. (2007). Drying of probiotics: Optimization of formulation and process to enhance storage survival. *Drying Technology*, 25(7-8), 1193–1201.
- Chegini, G. R., & Ghobadian, B. (2005). Effect of Spray-drying Conditions on Physical Properties of Orange Juice Powder. *Drying Technology*, 23(3), 657–668.
- Chen, X. D., & Özkan, N. (2007). Stickiness, functionality, and microstructure of food powders. *Drying Technology*, 25(6), 959-969.
- Chen, X. D., & Patel, K. C. (2008). Manufacturing better quality food powders from spray drying and subsequent treatments. *Drying Technology*, 26(11), 1313-1318.
- Chiou, D., & Langrish, T. A. G. (2007). Crystallization of amorphous components in spray-dried powders. *Drying Technology*, 25(9), 1427-1435.
- Cho, I. C., & Kolar Jr, C. W. (1977). *U.S. Patent No. 4,025,659*. Washington, DC: U.S. Patent and Trademark Office.
- Chronakis, I. S. (1998). On the molecular characteristics, compositional properties, and structural-functional mechanisms of maltodextrins: a review. *Critical Reviews in Food Science and Nutrition*, 38(7), 599-637.
- Chua, K. J., Mujumdar, A. S., Chou, S. K., Hawlader, M. N. A., & Ho, C. (2000). Convective drying of banana, guava and potato pieces: Effect of cyclical variations of air temperature on drying kinetics and color change. *Drying Technology*, 18(4-5), 907–936.
- Chung, M. S., Ruan, R. R., Chen, P., Chung, S. H., Ahn, T. H., & Lee, K. H. (2000). Study of caking in powdered foods using nuclear magnetic resonance spectroscopy. *Journal of Food Science*, 65(1), 134-138.
- Coussemment, P. A. (1999). Inulin and oligofructose: Safe intakes and legal status. *The Journal of Nutrition*, 129(7), 1412S-1417S.
- Cruz, M. A. A., Passos, M. L., & Ferreira, W. R. (2005). Final drying of whole milk powder in vibrated-fluidized beds. *Drying Technology*, 23(9-11), 2021-2037.
- Cunha, P. L., Paula, R., & Feitosa, J. (2007). Purification of guar gum for biological applications. *International Journal of Biological Macromolecules*, 41(3), 324-331.
- Dacanal, G. C., & Menegalli, F. C. (2009). Experimental study and optimization of the agglomeration of acerola powder in a conical fluid bed. *Drying Technology*, 188(3), 187-194.
- Dacanal, G. C., & Menegalli, F. C. (2010). Selection of operational parameters for the production of instant soy protein isolate by pulsed fluid bed agglomeration. *Powder Technology*, 203(3), 565-573.

- De Castro, F. P., Cunha, T. M., Barreto, P. L., Amboni, R. D. D., & Prudencio, E. S. (2009). Effect of oligofructose incorporation on the properties of fermented probiotic lactic beverages. *International Journal of Dairy Technology*, 62(1), 68-74.
- De Vito, F., Veytsmanb, B., Painter, P., & Kokini, J. (2015). Simulation of the effect of hydrogen bonds on water activity of glucose and dextran using the Veytsman model. *Carbohydrate Polymers*, (117), 236–246.
- De Vriese, S., De Backer, G., De Henauw, S., Huybrechts, I., Kornitzer, K., Levêque, A., & Van Oyen, H. (2005). The Belgian food consumption survey: aims, design and methods. *Archives of Public Health*, 63(1), 1-16.
- Debon, J., Prudêncio, E. S., & Cunha Petrus, J. C. (2010). Rheological and physico-chemical characterization of prebiotic microfiltered fermented milk. *Journal of Food Engineering*, 99(2), 128-135.
- Desai, K. G., & Park, H. G. (2004). Solubility studies of valdecoxib in presence of carriers, co-solvent and surfactants. *Drug Development Research*, 62, 41-48.
- Descamps, N., Palzer, S., Roos, Y. H., & Fitzpatrick, J. J. (2013). Glass transition and flowability/caking behaviour of maltodextrin DE 21. *Journal of Food Engineering*, 119(4), 809-813.
- Desobry, S. A., Netto, F. M., & Labuza, T. P. (1997). Comparison of spray-drying, drum-drying and freeze-drying for β -carotene encapsulation and preservation. *Journal of Food Science*, 62(6), 1158–1162.
- Dhanalakshmi, K., Bhattacharya, S. (2014). Agglomeration of turmeric powder and its effect on physico-chemical and microstructural characteristics. *Journal of Food Engineering*, 120, 124–134.
- Dhanalakshmi, K., Ghosal, S., & Bhattacharya, S. (2011). Agglomeration of food powder and applications. *Critical Reviews in Food Science and Nutrition*, 51(5), 432-441.
- Dijkgraaf, J. T., Westerbeek, J. M., & Zijlmans, J. B. (1995). *U.S. Patent No. 5,462,759*. Washington, DC: U.S. Patent and Trademark Office.
- Domagala, J., Sady, M., Grega, T., & Bonczar, G. (2006). Rheological properties and texture of yogurts when oat-maltodextrin is used as a fat substitute. *International Journal of Food Properties*, 9(1), 1–11.
- Dziezak, J. D. (1988). Microencapsulation and encapsulated ingredients. *Food Technology*, 42(4), 136–148.
- Eapen, K. E., & Kahn, M. L. (1980). *U.S. Patent No. 4,199,604*. Washington, DC: U.S. Patent and Trademark Office.
- Eissens, A. C., Bolhuis, G. K., Hinrichs, W. L., & Frijlink, H. W. (2002). Inulin as filler-binder for tablets prepared by direct compaction. *European Journal of Pharmaceutical Sciences*, 15(1), 31-38.

- Elleuch, M., Bedigian, D., Roiseux, O., Besbes, S., Blecker, C., & Attia, H. (2011). Dietary fibre and fibre-rich by-products of food processing: Characterisation, technological functionality and commercial applications: A review. *Food Chemistry*, 124(2), 411-421.
- Fang, Y., Selomulya, C., & Chen, X. D. (2007). On measurement of food powder reconstitution properties. *Drying Technology*, 26(1), 3-14.
- Fazio, S. C. (1994). *U.S. Patent No. 5,284,674*. Washington, DC: U.S. Patent and Trademark Office.
- Ferrari, C. C., Germer, S. P. M., & de Aguirre, J. M. (2012). Effects of spray-drying conditions on the physicochemical properties of blackberry powder. *Drying Technology*, 30(2), 154-163.
- Filkova, I., & Mujumdar, A. S. (1995). Industrial spray drying systems. *Handbook of Industrial Drying*, 1, 263-308.
- Fitzpatrick, J. J., Iqbal, T., Delaney, C., Twomey, T., & Keogh, M. K. (2004). Effect of powder properties and storage conditions on the flowability of milk powders with different fat contents. *Journal of Food Engineering*. 64(4), 435-444.
- Flaczyk, E., Gorecka, D., Kobus, J., & Szymandera-Buszk, K. (2009). The influence of inulin addition as fat substitute on reducing energy value and consumer acceptance of model pork meatballs. *Zywnosc. Nauka. Technologia. Jakosc*, 4(65), 41-46.
- Franck, A. (2002). Technological functionality of inulin and oligofructose. *British Journal of Nutrition*, 87(S2), S287-S291.
- Frary, C. D., Johnson, R. K., & Wang, M. Q. (2005). Food sources and intakes of caffeine in the diets of persons in the United States. *Journal of the American Dietetic Association*, 105, 110-113.
- Gaiani, C., Ehrhardt, J. J., Scher, J., Hardy, J., Desobry, S., & Banon, S. (2006). Surface composition of dairy powders observed by X-ray photoelectron spectroscopy and effects on their rehydration properties. *Colloids and Surfaces B: Biointerfaces*, 49, 71-78.
- Gardiner, D. S. (1977). *U.S. Patent No. 4,046,926*. Washington, DC: U.S. Patent and Trademark Office.
- Gianfrancesco, A., Turchiuli, C., Dumoulin, E., & Palzer, S. (2009). Prediction of powder stickiness along spray drying process in relation to agglomeration. *Particulate Science and Technology*, 27(5), 415-427.
- Gibson, G. R., Probert, H. M., Van Loo, J., Rastall, R. A., & Roberfroid, M. B. (2004). Dietary modulation of the human colonic microbiota: updating the concept of prebiotics. *Nutrition Research Reviews*, 17(2), 259-275.

- Giese, J. (1996). Fats and fat replacers: balancing the health benefits. *Food Technology*.
- Gilmore, C., & Miller, D. E. (1980). *U.S. Patent No. 4,239,786*. Washington, DC: U.S. Patent and Trademark Office.
- Glibowski, P., & Bukowska, A. (2011). The effect of pH, temperature and heating time on inulin chemical stability. *Acta Scientiarum Polonorum. Seria: Technologia Alimentaria*, 10(2), 189-196.
- Golde, A. E., & Schmidt, K. A. (2005). Quality of coffee creamers as a function of protein source. *Journal of Food Quality*, 28(1), 46–61
- Gong, Z., Zhang, M., Mujumdar, A. S., & Sun, J. (2007). Spray drying and agglomeration of instant bayberry powder. *Drying Technology*, 26(1), 116-121.
- Gonzalez-Tomás, L., Coll-Marqués, J., & Costell, E. (2008). Viscoelasticity of inulin–starch-based dairy systems. Influence of inulin average chain length. *Food Hydrocolloids*, 22(7), 1372-1380.
- Goula, A. M., & Adamopoulos, K. G. (2005). Spray drying of tomato pulp in dehumidified air: II. The effect on powder properties. *Journal of Food Engineering*, 66(1), 35-42.
- Goula, A. M., & Adamopoulos, K. G. (2008). Effect of maltodextrin addition during spray drying of tomato pulp in dehumidified air: II. Powder properties. *Drying Technology*, 26(6), 726–737.
- Goula, A. M., & Adamopoulos, K. G. (2010). A new technique for spray drying orange juice concentrate. *Innovative Food Science & Emerging Technologies*, 11(2), 342-351.
- Grabowski, J. A., Truong, V. D., & Daubert, C. R. (2008). Nutritional and rheological characterization of spray dried sweetpotato powder. *LWT-Food Science and Technology*, 41(2), 206-216.
- Groenewold, H., & Tsotsas, E. (2001). Experimental Investigation and Modeling of the Influence of Indirect Heating on Fluidized Bed Drying. *Drying Technology*, 19(8), 1739-1754.
- Hager, A. S., Ryan, L. A., Schwab, C., Gänzle, M. G., O'Doherty, J. V., & Arendt, E. K. (2011). Influence of the soluble fibres inulin and oat β -glucan on quality of dough and bread. *European Food Research and Technology*, 232(3), 405-413.
- Hall, C. W., Hedrick, T. I. (1975). *Dairy of milk powders (2nd ed.)*. The Avi publishing company: Westport, CT.
- Haugaard Sorensen, I., Krag, J., Pisecky, J., & Westergaard, V. (1978). Méthodes d'analyses des produits laitiers déshydratés. *Niro A/S, Copenhagen, Denmark*.
- Hennelly, P. J., Dunne, P. G., O'Sullivan, M., & O'Riordan, D. (2005). Increasing the moisture content of imitation cheese: effects on texture, rheology and

- microstructure. *European Food Research and Technology*, 220(3-4), 415-420.
- Hennigs, C., Kockel, T. K., & Langrish, T. A. G. (2001). New measurements of the sticky behavior of skim milk powder. *Drying Technology*, 19(3-4), 471-484.
- Higdon, J. V., & Frei, B. (2006). Coffee and health: a review of recent human research. *Critical Review Food Science and Nutrition*, 46, 101–23.
- Hooda, S., & Jood, S. (2005). Organoleptic and nutritional evaluation of wheat biscuits supplemented with untreated and treated fenugreek flour. *Food Chemistry*, 90(3), 427-435.
- Hunter, R. S. (1987). *The measurement of appearance*. John Wiley & Sons.
- Hursh, H., & Martin, J. (2005). Low-carb and beyond: The health benefits of inulin. *Cereal Foods World*, 50(2), 57-60.
- Icoz, D. Z., & Kokini, J. L. (2007a). Examination of the validity of the Flory–Huggins solution theory in terms of miscibility in dextran systems. *Carbohydrate Polymers*, 68(1), 59–67.
- Jakubczyk, E., Ostrowska-Ligeza, E., & Gondek, E. (2010). Moisture sorption characteristics and glass transition temperature of apple puree powder. *International Journal of Food Science & Technology*, 45(12), 2515-2523.
- Jaya, S. (2009). Microstructure analysis of dried yogurt: Effect of different drying methods. *International Journal of Food Properties*, 12(3), 469-481.
- Jaya, S., & Das, H. (2003). A vacuum drying model for mango pulp. *Drying Technology*, 21(7), 1215-1234.
- Jaya, S., Das, H., & Mani, S. (2006). Optimization of maltodextrin and tricalcium phosphate for producing vacuum dried mango powder. *International Journal of Food Properties*, 9(1), 13-24.
- Jayasundera, M., Adhikari, B., Adhikari, R., & Aldred, P. (2011). The effects of proteins and low molecular weight surfactants on spray drying of model sugar-rich foods: Powder production and characterisation. *Journal of food Engineering*, 104(2), 259-271.
- Jimenez-Flores, R., & Kosikowski, F. V. (1987). Low-fat dairy coffee creamer. *U.S. Patent No. 4,689,245*. Washington, DC: U.S. Patent and Trademark Office.
- Jinapong, N., Supphantharika, M., & Jamnong, J. (2008). Production of instant soymilk powders by ultrafiltration, spray drying and fluidized bed agglomeration. *Journal of Food Engineering*, 84, 194–205.

- Johari, G. P., Hallbrucker, A., & Mayer, E. (1987). The glass–liquid transition of hyperquenched water. *Nature*, 330(6148), 552-553.
- Kage, H., Nishihara, K., Ishimatsu, H., Ogura, H., & Matsuno, Y. (2001). Effect of drying of powder coating efficiency and agglomeration in vibro-fluidized bed. *Drying Technology*, 19, 359–373.
- Kage, M., Yang, Q., Sato, H., Matsumoto, S., Kaji, R., Akiguchi, I., & Tooyama, I. (2001). Acidic fibroblast growth factor (FGF-1) in the anterior horn cells of ALS and control cases. *Neuroreport*, 12(17), 3799-3803.
- Kalogiannia, E. P., Xynogalos, V. A., Karapantsios, T. D., & Kostoglou, M. (2002). Effect of feed concentration on the production of pregelatinized starch in a double drum dryer. *LWT- Food Science and Technology*, 35(8), 703-711.
- Kelly, P. M., Oldfield, D. J., & O’Kennedy, B.T. (1999). The thermostability of spray dried imitation coffee whiteners. *International Journal of Dairy Technology*, 52(3), 107–113.
- Kha, T. C., Nguyen, M. H., & Roach, P. D. (2010). Effects of spray drying conditions on the physicochemical and antioxidant properties of the Gac (*Momordica cochinchinensis*) fruit aril powder. *Journal of Food Engineering*, 98(3), 385-392.
- Kiessling, T. R., & Zeller, B. L. (2005). *U.S. Patent No. 6,129,943*. Washington, DC: U.S. Patent and Trademark Office.
- Kim, E. H. J., Chen, X. D., & Pearce, D. (2009). Surface composition of industrial spray-dried milk powders. 2. Effects of spray drying conditions on the surface composition. *Journal of Food Engineering*, 94(2), 169-181.
- Kim, E.H.J., Chen, X.D., & Pearce, D. (2002). Surface characterization of four industrial spray-dried dairy powders in relation to chemical composition, structure and wetting property. *Colloids and Surfaces B: Biointerfaces*, 26, 197–212.
- Kim, Y., Faqih, M. N., & Wang, S. S. (2001). Factors affecting gel formation of inulin. *Carbohydrate Polymers*, 46(2), 135-145.
- Knight, P. C. (2001). Structuring agglomerated products for improved performance. *Powder Technology*, 119(1), 14-25.
- Koc, B., Yilmazer, M. S., Balkır, P., & Ertekin, F. K. (2010). Spray drying of yogurt: Optimization of process conditions for improving viability and other quality attributes. *Drying Technology*, 28(4), 495-507.
- Kumar, P., & Mishra, H. N. (2004). Yoghurt powder—a review of process technology, storage and utilization. *Food and Bioproducts Processing*, 82(2), 133-142.

- Lopez-Garcia, E., van Dam, R. M., Rajpathak, S., Willett, W. C., Manson, J. E., & Hu, F. B. (2006). Changes in caffeine intake and long-term weight change in men and women. *The American Journal of Clinical Nutrition*, 83(3), 674–680.
- Maa, Y. F., Nguyen, P. A., Sit, K., & Hsu, C. C. (1998). Spray-drying performance of a bench-top spray dryer for protein aerosol powder preparation. *Biotechnology and Bioengineering*, 60(3), 301-309.
- Machado, V. G., Hirata, T. A. M., & Menegalli, F. C. (2014). Agglomeration of soy protein isolate in a pulsed fluidized bed: Experimental study and process optimization. *Powder Technology*, 254, 248-255.
- Malvern Instruments (1999). Operators guide: Man.0247 Issue 2.0. Worcestershire, United Kingdom: Malvern Instruments Ltd.
- Mann, J. I., & Cummings, J. H. (2009). Possible implications for health of the different definitions of dietary fibre. *Nutrition, Metabolism and Cardiovascular Diseases*, 19(3), 226-229.
- Masters, K. (1985). Spray drying handbook. *Spray drying handbook*, (Ed.4). Longman Scientific and Technical: Essex, UK.
- Masters, K. (1992). Spray drying handbook. *Spray drying handbook*, (Ed.5). Ed; Longman Scientific and Technical: Essex, UK.
- Menshutima, N. V., Gordienko, M. G., Vionovskiy, A. A., & Zbicinski, I. (2010). Spray drying of prebiotics: process development and scale-up. *Drying Technology*, 28 (3), 1170-1177.
- Meyer, D., Bayarri, S., Tárrega, A., & Costell, E. (2011). Inulin as texture modifier in dairy products. *Food Hydrocolloids*, 25(8), 1881-1890.
- Miller, D. E., & Werstak, C. E. (1983). *U.S. Patent No. 4,415,600*. Washington, DC: U.S. Patent and Trademark Office.
- Millqvist-Fureby, A., & Smith, P. (2007). In-situ lecithination of dairy powders in spray-drying for confectionery applications. *Food Hydrocolloids*, 21(5), 920-927.
- Mirhosseini, H., Tan, C. P., Hamid, N. S., & Yusof, S. (2008). Effect of Arabic gum, xanthan gum and orange oil contents ζ -potential, conductivity, stability, size index and pH of orange beverage emulsion. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 315(1), 47-56.
- Mirhosseini, H., Tan, C. P., Hamid, N. S., Yusof, S., & Chern, B. H. (2009). Characterization of the influence of main emulsion components on the physicochemical properties of orange beverage emulsion using response surface methodology. *Food Hydrocolloids*, 23(2), 271-280.

- Mishra, P., Srivastava, V., Verma, D., Chauhan, O. P., & Rai, G. K. (2009). Physicochemical properties of chakiya variety of amla (*emblica officinalis*) and effect of different dehydration methods on quality of powder. *African Journal of Food Science*, 3(10), 303-306.
- Monteiro, C. A., Mondini, L., & Costa, R. B. L. (2000). Mudanças na composição e ~ adequação nutricional da dieta familiar nas ~ áreas metropolitanas do Brasil (1988e1996). *Revista de Saúde Pública*. Sao Paulo, 34, 251-258.
- Moore, J. G. (1995). Drum dryer, in *Handbook of Industrial Drying*, 2nd edn., Mujumdar, A. S., ed., Marcel Dekker, New york, p.249.
- Mounir, S., & Allaf, K. (2008). Three-stage spray drying: new process involving instant controlled pressure drop. *Drying Technology*, 26(4), 452-463.
- Mujumdar, A. S. (Ed.). (2014). *Handbook of industrial drying*. CRC Press
- Mukherjee, S., & Bhattacharya, S. (2006). Characterization of agglomeration process as a function of moisture content using a model food powder. *Journal of Texture Studies*, 37(1), 35-48.
- Musielak, G., & Mierzwa, D. (2009). Permanent strains in clay-like material during drying. *Drying Technology*, 27(7-8), 894-902.
- National Center for Health Statistics (US). (1994). *Plan and operation of the third National Health and Nutrition Examination Survey, 1988-94* (No. 32). Natl Ctr for Health Statistics.
- Nep, E. I., & Conway, B. R. (2011). Physicochemical characterization of grewia polysaccharide gum: Effect of drying method. *Carbohydrate Polymers*, 84(1), 446-453.
- Nijdam, J., & Langrish, T. A., (2005). An investigation of milk powders produced by a laboratory-scale spray dryer. *Drying Technology*, 23(5), 1049-1056.
- Nijdam, J., Ibach, A., & Kind, M. (2008). Fluidisation of whey powders above the glass-transition temperature. *Powder Technology*, 187(1), 53-61.
- Nindo, C. I., & Tang, J. (2007). Refractance Window dehydration technology: a novel contact drying method. *Drying Technology*, 25(1), 37-48.
- Niness, K. R. (1999). Inulin and oligofructose: What are they?. *The Journal of Nutrition*. 129(7), 1402s-1406s.
- Nishinari, K. (2009). Texture and rheology in food and health. *Food Science and Technology Research*, 15(2), 99-106.
- Non-polyol low-digestible carbohydrates: food applications and functional benefits. *British Journal of Nutrition*, 85(S1), S47-S53.
- O'Neil, E. (1993). Low-fat products. *Meat Focus International*, 2, 70-75.

- Oldfield, D., & Singh, H. (2005). Functional properties of milk powders. *Food Science and Technology-New York-Marcel Dekker*, 146, 365.
- Papadakis, S. E., Gardeli, C., & Tzia, C. (1998, August). Raisin extract powder: Production, physical and sensory properties. In *Proc. 11th International Drying Symposium IDS* (Vol. 98, pp. 1207-1213).
- Paseephol, T., Small, D. M., & Sherkat, F. (2008). Rheology and texture of set yogurt as affected by inulin addition. *Journal of Texture Studies*, 39(6), 617-634.
- Peleg, M., & Hollenbech, A. M. (1984). Flow conditioners and anti-caking agents. *Food Technology*, 91-100.
- Peleg, M., & Mannheim, C. H. (1977). The mechanism of caking of powdered onion. *Journal of Food Processing and Preservation*, 1(1), 3-11.
- Pietsch, W. (1999). Readily engineer agglomerates with special properties from micro-and nanosized particles. *Chemical Engineering Progress*, 95(8), 67-81.
- Popat, R. A., Van Den Eeden, S. K., Tanner, C.M., Kamel, F., Umbach, D.M., & Marder, K. (2011). Coffee, ADORA2A, and CYP1A2: The caffeine connection in Parkinson's disease. *European Journal of Neurology*, 18(5), 756-765.
- Pordy, W. T. (1996). *U.S. Patent No. 5,480,670*. Washington, DC: U.S. Patent and Trademark Office.
- Potter, N. N. (1968). *Food Science*. AVI Publishing Company: Westport, CT.
- Pua, C. K., Hamid, N. S. A., Tan, C. P., Mirhosseini, H., Rahman, R. B. A., & Rusul, G. (2010). Optimization of drum drying processing parameters for production of jackfruit (*Artocarpus heterophyllus*) powder using response surface methodology. *LWT-Food Science and Technology*, 43(2), 343-349.
- Pua, C. K., Hamid, N., Rusul, G., & Rahman, R. A. (2007). Production of drum-dried jackfruit (*Artocarpus heterophyllus*) powder with different concentration of soy lecithin and gum arabic. *Journal of Food Engineering*, 78(2), 630-636.
- Ren, G. Y., Li, D., Wang, L. J., Özkan, N., & Mao, Z. H. (2010). Morphological properties and thermoanalysis of micronized cassava starch. *Carbohydrate Polymers*, 79(1), 101-105.
- Rennie, P. R., Chen, X. D., Hargreaves, C., & Mackereth, A. R. (1999). A study of the cohesion of dairy powders. *Journal of Food Engineering*, 39(3), 277-284.
- Ronkart, S. N., Blecker, C. S., Fourmanoir, H., Fougnes, C., Deroanne, C., Van Herck, J. C., & Paquot, M. (2007). Isolation and identification of inulooligosaccharides resulting from inulin hydrolysis. *analytica chimica acta*, 604(1), 81-87.

- Roos, Y. (1995). Characterization of food polymers using state diagrams. *Journal of Food Engineering*, 24, 339–360.
- Roos, Y. H. (1995). Glass transition-related physicochemical changes in foods: Chemical and rheological changes during phase transition in food. *Food Technology*, 49(10), 97-102.
- Roos, Y. H., & Karel, M. (1991). Water and molecular weight effect in glass transition in amorphous carbohydrates and carbohydrate solution. *Journal of Food Science*, 56(6), 1676-1681.
- Roustapour, O. R., Hosseinalipour, M., & Ghobadian, B. (2006). An experimental investigation of lime juice drying in a pilot plant spray dryer. *Drying Technology*, 24(2), 181-188.
- Sandrou, D. K., & Arvantoyannis, I. S. (2000). Low fat/calorie foods: current state and perspectives. *Critical Reviews in Food Science and Nutrition*, 40(5), 427-447.
- Sang, L. X., Chang, B., Li, X. H., & Jiang, M. (2013). Consumption of coffee associated with reduced risk of liver cancer: A meta-analysis. *BMC Gastroenterology*, 13(1), 34.
- Schaefer, B. (2004). Coffee consumption and type 2 diabetes mellitus. *Ann International Medicine* 141:321. author reply 3–4.
- Schaller-Povolny, L. A., & Smith, D. E. (2001). Original papers-Viscosity and freezing point of a reduced fat ice cream mix as related to inulin content. *Milchwissenschaft*, 56(1), 25-28.
- Schein, C.H. (1990). Solubility as a function of protein structure and solvent components. *Nature Biotechnology*, 8, 308–317.
- Schuck, P., Jeantet, R., & Dolivet, A. (2012). *Analytical methods for food and dairy powders*. John Wiley & Sons.
- Schuck, P., le Floch-Fouere, C., & Jeantet, R. (2013). Changes in functional properties of milk protein powders: effects of vacuum concentration and drying. *Drying Technology*, 31(13-14), 1578-1591.
- Shittu, T. A., & Lawal, M. O. (2007). Factors affecting instant properties of powdered cocoa beverages. *Food Chemistry*, 100 (1), 91-98.
- Shrestha, A. K., Howes, T., Adhikari, B. P., & Bhandari, B. R. (2007). Water sorption and glass transition properties of spray dried lactose hydrolysed skim milk powder. *LWT-Food Science and Technology*, 40(9), 1593-1600.
- Singh, A. K., Selvam, R. P., & Sivakumar, T. (2010). Isolation, characterisation and formulation properties of a new plant gum obtained from mangifera indica. *Int J Pharm Biomed Res*, 1, 35.

- Soukoulis, C., Lebesi, D., & Tzia, C. (2009). Enrichment of ice cream with dietary fibre: Effects on rheological properties, ice crystallisation and glass transition phenomena. *Food Chemistry*, 115(2), 665–671.
- Srinivasan, M., Singh, H., & Munro, P. A. (2002). Formation and stability of sodium caseinate emulsions: influence of retorting (121°C for 15 min) before or after emulsification. *Food Hydrocolloids*, 16(2), 153–160.
- Stone, H., & Sidel, J. L. (1993). Sensory evaluation practices (2nd ed.). San Diego, CA: Academic Press.
- Sudha, M. L., Srivastava, A. K., Vetrmani, R., & Leelavathi, K. (2007). Fat replacement in soft dough biscuits: Its implications on dough rheology and biscuit quality. *Journal of Food Engineering*, 80(3), 922–930.
- Sudhagar, M. (2000). *Spray drying of fruit juices*. (Unpublished master dissertation). Department of Agriculture and Food Engineering, Indian Institute of Technology: Kharagpur, India.
- Sundaram, J., & Durance, T. D. (2008). Water sorption and physical properties of locust bean gum–pectin–starch composite gel dried using different drying methods. *Food Hydrocolloids*, 22(7), 1352–1361.
- Sunooj, K. V., Radhakrishna, K., George, J., & Bawa, A. S. (2009). Factors influencing the calorimetric determination of glass transition temperature in foods: A case study using chicken and mutton. *Journal of Food Engineering*, 91(2), 347–352.
- Szczesniak, A. S. (2002). Texture is a sensory property. *Food Quality and Preference*, 13(4), 215–225.
- Szulc, K., & Lenart, A. (2013). Surface modification of dairy powders: Effects of fluid-bed agglomeration and coating. *International Dairy Journal*, 33(1), 55–61.
- Takeiti, C. Y., Kieckbusch, T. G., & Collares-Queiroz, F. P. (2010). Morphological and physicochemical characterization of commercial maltodextrins with different degrees of dextrose-equivalent. *International Journal of Food Properties*, 13(2), 411–425.
- Tamime, A. Y., & Robinson, R. K. (1999). *Yoghurt: science and technology*. Wood head Publishing.
- Thomas, M., Scher, J., Desobry-Banon, S., & Desobry, S. (2004). Milk powders ageing: Effect on physical and functional properties. *Critical Reviews in Food Science and Nutrition*, 44, 297–322.
- Toneli, J., Park, K., Negreiros, A., & Murr, F. (2010). Spray-drying process optimization of chicory root inulin. *Drying Technology*, 28(3), 369–379.

- Tonon, R. V., Brabet, C., & Hubinger, M. D. (2008). Influence of process conditions on the physicochemical properties of açai (*Euterpe oleraceae* Mart.) powder produced by spray drying. *Journal of Food Engineering*, 88(3), 411-418.
- Trumbo, P., Schlicker, S., Yates, A., & Poos, M. (2002). Dietary reference intakes for energy, carbohydrate, fibre, fat, fatty acids, cholesterol, protein, and amino acids. *Journal of the American Dietetic Association*, 102(11), 1621–1630.
- Tuot, J., Octavia, W., & Sher, A. A. (2014). *U.S. Patent No. 8,658,230*. Washington, DC: U.S. Patent and Trademark Office.
- Turchiuli, C., Eloualia, Z., El Mansouri, N., & Dumoulin, E. (2005). Fluidised bed agglomeration: Agglomerates shape and end-use properties. *Powder Technology*, 157(1), 168-175.
- United States Department of Agriculture Foreign Agricultural Service. Global agricultural information network report. Coffee market brief update_Seoul ATO_Korea–Republic of 2010-12-09. United States Department of Agriculture Foreign Agricultural Service; 2011.
- United States. Department of Health. (1996). *Physical activity and health and Human Services: a report of the Surgeon General*. DIANE Publishing.
- Uthumporn, U., Zaidul, I. S., & Karim, A. A. (2010). Hydrolysis of granular starch at sub-gelatinization temperature using a mixture of amylolytic enzymes. *Food and Bioproducts Processing*, 88(1), 47-54.
- Valous, N. A., Gavrielidou, M. A., Karapantsios, T. D., & Kostoglou, M. (2002). Performance of a double drum dryer for producing pregelatinized maize starches. *Journal of Food Engineering*, 51(3), 171-183.
- Van Loo, J. (2007). How chicory fructans contribute to zootechnical performance and well-being in livestock and companion animals. *The Journal of Nutrition*, 137(11), 2594S-2597S.
- Vega-Mercado, H., Gongora-Nieto, M., & Barbosa- Cánovas, G. V. (2001). Advances in dehydration of food. *Journal of Food Engineering*, 49(4), 271-289.
- Villegas, B., Carbonell, I., & Costell, E. (2007). Inulin milk beverages: sensory differences in thickness and creaminess using R-index analysis of the ranking data. *Journal of Sensory Studies*, 22(4), 377-393.
- Wang, Q., & Cui, S. W. (2005). Understanding the physical properties of food polysaccharides. In S. W. Cui (Ed.), *Food carbohydrate: chemistry, physical properties, and applications*. (pp. 162-214). Boca Raton, Florida, Taylor and Fraiss.
- Wang, S. W. (2000). A new ingredient for non-fat yogurt. *Milk Industry International, Technical & Research Supplement*, September. 6-7.

- White, G. W., & Cakebread, S. H. (1966). The glassy state in certain sugar-containing food products. *International Journal of Food Science & Technology*, 1(1), 73-82.
- Williams, S. ur. (1984) Official methods of analysis. Arlington, VA: Association of Official Analytical Chemists/AOAC.
- Word health organization (WHO), (2012). Bulk density and tapped density of powders. Document QAS/11.450.
- World Resources Institute. (2011). Earth trends data tables: Energy and resources. Retrieved from:
http://www.earthtrends.wri.org/searchable_db/index.php?theme=8&variable_ID=1677&action=select_countries. Accessed 31.10.11.
- Wouters, E. (1998). The benefits of inulin and oligofructose in ice cream. *Word of Ingredients*. September 44-45.
- Yazdanpanah, N., & Langrish, T. A. (2011). Crystallization and drying of milk powder in a multiple-stage fluidized bed dryer. *Drying Technology*, 29 (9), 1046-1057.
- Zimon, A. D. (1982). *Adhesion of dust and powder* (2nd ed). Consultants Bureau: New York.

LIST OF PUBLICATIONS

Hedayat Nia, S., Mirhosseini, H., Abd Manap, M. Y., Karim, R. Anew formulation and process for manufacturing of low-fat, enriched coffee creamer. Patent (Pending NO.38/2014).

Hedayatnia, S., Mirhosseini, H., Tabatabaee Amid, B., Islam Sarker, Z., Karim, R., Abd Manap, Y. (2015). Effect of Different Fat Replacers and Drying Methods on Thermal Behaviour, Glass Transition, Morphology and Flowability of Reduced-Fat Coffee Creamer. *LWT - Food Science and Technology* (under riview).





UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : _____

TITLE OF THESIS / PROJECT REPORT :

Influence of different fat replacers and drying techniques
on physicochemical characteristics and sensory attributes of
Regular and instant reduced-fat coffee creamer.

NAME OF STUDENT : _____

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (✓)

☐

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

☐

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

☐

OPEN ACCESS


I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

☐

PATENT

Embargo from _____ until _____
(date) (date)



(Signature of Student)
New IC No/ Passport No.:

M27760546

Date :

21/4/2015

Approved by:



(Signature of Chairman of Supervisory Committee)
Name:

Department of Food Technology
Faculty of Food Science & Technology
Universiti Putra Malaysia

Date :

21/4/15

43400 UPM Serdang, SELANGOR
Tel No: 603-8946 8390 Fax No: 603-8942 3552