



**UNIVERSITI PUTRA MALAYSIA**

***EFFECTS OF FERMENTATION AND DE-BITTERING METHODS ON  
PRODUCTION OF ALMOND-LIKE RAMBUTAN SEED  
(*Nephelium lappaceum* L.) SNACK***

**SHABNAM MEHDIZADEH**

**FSTM 2015 43**



**EFFECTS OF FERMENTATION AND DE-BITTERING METHODS ON  
PRODUCTION OF ALMOND-LIKE RAMBUTAN SEED  
(*Nephelium lappaceum* L.) SNACK**

By

**SHABNAM MEHDIZADEH**


**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirement for the Degree of Master of Science**

**December 2014**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia





Dedicated to My Family

© COPYRIGHT UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Degree of Master of Science.

**EFFECTS OF FERMENTATION AND DE-BITTERING METHODS ON  
PRODUCTION OF ALMOND-LIKE RAMBUTAN SEED  
(*Nephelium lappaceum* L.) SNACK**

By

**SHABNAM MEHDIZADEH**

**December 2014**

**Chairman: Associate Professor Lasekan Olusegun Olaniyi, PhD**

**Faculty: Food Science Technology**

There are so many current studies going on in the field of waste products. One of the instances is seeds of rambutan, which is discarded yearly because of its slightly bitter taste despite the fact that it contains valuable source of oleic acid, arachidic acid and vitamins. The first part of this study determined the effect of fermentation on variability of fermentation index, polyphenol constituents and amino acids of rambutan seeds (anak sekolah). Results of fermentation showed that time of fermentation had a significant decrease in pH, whereas a significant increase was evident in the fermentation index ( $p < 0.05$ ). Findings also showed that microbial activities in the pulp produce alcohols to some extent and acids such as acetic acid and lactic acid, and raises the temperature and humidity and these changes inside the seed resulted in the death of the embryo. Moreover, it was found that fermentation significantly reduced the levels of total polyphenols (59%), tannin (60%) and saponins (33%). However, fermentation appeared to have a moderate effect on geraniin, corilagin and much stronger effect on ellagic and gallic acids. During fermentation, variability in gallic acid, geraniin, corilagin and ellagic acid did not show consistent trend. In contrast, amino acids that elicited bitter taste (phenylalanine, tyrosine and leucine) showed reduction of 20%, 30% and 40% respectively after 10 days of fermentation. Furthermore, it was discovered that sucrose, which was the only sugar present in significant amount, in unfermented seed was remarkably reduced during fermentation. The second part of this study concentrated on the effect of different debittering methods on the seed of rambutan. There were 14 different methods for reducing bitterness with different times and concentrations and in different soaking medium such as soaking in water and debittering through impregnation processes by sugar and salt as well as lye treatment. Three methods were selected based on the substantial reduction of bitterness compounds. These three methods were soaking in water for 7 days, treatment of seeds with 22% sodium chloride for 8 h and treatment of seeds with 1 N NaOH for 8 h, accordingly. The three most efficient methods indicated that tannin content declined by 55.4%, 48.2% and 40.9%. Similarly, polyphenol contents were reduced to 92.7%, 84.4% and 85.1%. Saponin, gallic acid and corilagin, which are known as

the prime causes of bitterness, were completely removed by these different treatments. However, geraniin was reduced by 20% during debittering methods. Tyrosine, leucine and phenylalanine were the three amino acids, which contain bitter taste in rambutan seeds were also reduced by the different debittering methods. The third part of this study focused on the effect of roasting on the texture, color and flavor of debittered seeds of rambutan and almond. Debittered rambutan seeds and almonds were roasted at 125 °C for 15 minutes, the volatile extract of rambutan seeds produced 13 odors and roasted almond produced 122 odors-active compounds by GC-MS. Findings showed that the highest peaks were associated to acids and ester groups. The results also showed that roasted debittered by 22% NaCl for 8 h and 7 days of soaking treatment were more similar to roasted almond than the other two treatments. In general, therefore, not only do the current findings add to a growing body of literature on the different debittering methods of rambutan seed, but also some efficient approaches were identified for debittering rambutan seed in which this may contribute significantly to produce a new almond-like snack.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Master Sains.

**KAJIAN KESAN PENAPAIAN DAN KAEDAH NYAH PAHIT TERHADAP  
PENGHASILAN SNEK BIJI RAMBUTAN (*Nephelium lappaceum* L.) KE  
SERUPA ALMOND**

Oleh

**SHABNAM MEHDIZADEH**

**Disember 2014**

**Pengerusi: Professor. Madya Lasekan Olusegun Olaniyi, PhD**

**Fakulti: Sains dan Teknologi Makanan**

Pada masa ini, beberapa kajian telah dijalankan ke atas bahan buangan. Salah satu contoh adalah biji rambutan yang dibuang setiap tahun kerana ia rasanya pahit walaupun ia mengandungi sumber yang berharga seperti asid oleik, asid arakidik dan vitamin. Bahagian pertama kajian ini telah menentukan kesan penapaian pada kebolehubahan indeks penapaian, jujuk polifenol dan asid amino rambutan. Keputusan penapaian menunjukkan bahawa masa penapaian mempunyai penurunan ketara dalam pH, manakala peningkatan yang ketara jelas didapati pada indeks penapaian ( $p < 0.05$ ). Dapatan kajian juga menunjukkan bahawa aktiviti mikrob pada pulpa menghasilkan sedikit sebanyak alkohol dan asid seperti asid asetik dan asid laktik, dan meningkatkan suhu dan kelembapan serta perubahan beginimengakibatkan kematian embrio pada benih. Selain itu, didapati bahawa penapaian mengurangkan aras jumlah polifenol (59%), tanin (60%) dan saponin (33%). Walau bagaimanapun, penapaian mempunyai kesan sederhana pada geraniin, korilagin dan kesan lebih kukuh pada elagik dan asid galik. Semasa penapaian, kepelbagaian asid galik, geraniin, korilagin dan asid elagik tidak menunjukkan tren yang konsisten. Sebaliknya, asid amino yang memberi rasa pahit (fenylalanina, tirosina dan leusina) menunjukkan pengurangan masing-masing 20%, 30% dan 40% selepas 10 hari penapaian. Tambahan pula, didapati bahawa sukrosa, yang merupakan satu-satunya sumber gula yang hadir dengan jumlah yang besar, di dalam biji yang tidak beralkohol telah berkurangan dengan ketara akibat penapaian. Bahagian kedua kajian ini tertumpu kepada kesan kaedah penyahpahitan berbeza pada rambutan. Terdapat 14 kaedah yang berbeza untuk mengurangkan kepahitan dalam pada masa yang berbeza dan melalui konsentrasi yang berbeza biji seperti rendaman di dalam air selama 3 hari dan 7 hari dan penyahpahitan melalui proses gula dan garam serta rawatan 'lye'. Tiga kaedah telah dipilih berdasarkan pengurangan sebatian kepahitan yang berpotensi. Tiga kaedah ini adalah dengan merendamnya di dalam air selama 7 hari, rawatan biji rambutan dengan 22% natrium klorida selama 8 jam dan rawatan biji dengan 1 N NaOH selama 8 jam. Dengan menggunakan tiga kaedah yang berkesan, hasil telah menunjukkan bahawa kandungan tanin masing-masing berkurangan menjadi 55.4%, 48.2% dan 40.9%.

Selain itu, kandungan polifenol masing-masing dikurangkan sebanyak 92.7%, 84.4% dan 85.1%. Saponin, asid galik dan korilagin yang diketahui merupakan punca utama kepahitan telah dibuang sepenuhnya oleh rawatan-rawatan yang berbeza. Walau bagaimanapun geraniin dikurangkan sebanyak 20% dengan kaedah penyahpahitan. Tirosina, leusina dan fenilalanina merupakan tiga asid amino, yang mengandungi rasa pahit dalam biji rambutan, juga dikurangkan dengan kaedah penyahpahitan berbeza dengan nilai-nilai yang berbeza. Bahagian ketiga kajian ini memberi tumpuan kepada kesan memanggang terhadap tekstur, warna dan rasa biji rambutan dan badam yang dinyahpahit. Biji rambutan yang dinyahpahit dan badam telah dipanggang pada 125°C selama 15 minit, ekstrak volatile biji rambutan menghasilkan 13 bau dan badam panggang menghasilkan 122 sebatian bau-aktif oleh GC-MS. Dapatan kajian menunjukkan bahawa puncak tertinggi adalah berkaitan dengan asid dan kumpulan ester. Keputusan juga menunjukkan bahawa hasil panggang yang dinyahpahit oleh NaCl 22% selama 8 jam dan 7 hari rawatan rendaman lebih serupa dengan badam panggang berbanding dengan dua rawatan yang lain. Secara umum, oleh kerana itu, kajian ini bukan sahaja menghasilkan penemuan yang menambahkan bahan literatur kepada badan literatur yang semakin berkembang tentang kaedah penyahpahitan yang berbeza pada biji rambutan, tetapi juga kepada beberapa pendekatan yang cekap yang telah dikenalpasti untuk penyahpahitan biji rambutan di mana ini boleh menyumbang dengan ketara untuk menghasilkan makanan ringan ala badam yang baharu.



## ACKNOWLEDGEMENTS

I would like to express my gratitude to a number of people for their help, support, and cooperation in preparing this dissertation.

First of all, I would like to thank my supervisor, Associate Professor Dr. Lasekan Olusegun Olaniyiwho took his precious time and rendered me valuable advice and constructive comments in writing this thesis. He was a great help in shaping my study. A profound gratitude is also owed to the feedback of the committee members, Associate Prof. Dr. Sharifah Kharidah Syed Muhammad and Assoc. Prof. Badlishah Sham Bahrain for their valuable recommendations, worthwhile suggestions and encouragement.

Besides, I would like to thank all lecturers and staff in the faculty of Food Science and Technology for their invaluable help, suggestions and constructive comments in my thesis.

Special thanks go to my husband and my family for their constant support and encouraging me all through my life when I needed it most.

Last but not least, I am truly grateful to Allah for giving me patience, strength, and wisdom to complete this thesis.

I certify that a Thesis Examination Committee has met on 30 December 2014 to conduct the final examination of Shabnam Mehdizadeh on her thesis entitled "Effects of Fermentation and De-Bittering Methods on Production of Almond-Like Rambutan Seed (*Nephelium lappaceum* L.) Snack" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U. (A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

**Chong Gun Hean, PhD**

Senior Lecturer  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Roselina binti Karim, PhD**

Associate Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Internal Examiner)

**Zaidul Islam Sarker, PhD**

Professor  
Kulliyyah of Pharmacy, Kuantan Campus  
International Islamic University Malaysia  
(External Examiner)



---

**ZULKARNAIN ZAINAL, PhD**  
Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 26 February 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

**Lasekan Olusegun Olaniyi, PhD**

Associate Professor  
Faculty of Food Science & Technology  
Universiti Putra Malaysia  
(Chairman)

**Sharifah Kharidah Syed Muhammad, PhD**

Associate Professor  
Faculty of Food Science & Technology  
Universiti Putra Malaysia  
(Member)

**Badlishah Sham Bahrain, PhD**

Associate Professor  
Faculty of Food Science & Technology  
Universiti Putra Malaysia  
(Member)

---

**BUJANG BIN KIM HUAT, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work
- quotations, illustrations and citations have been duly referenced
- the thesis has not been submitted previously or concurrently for any other degree at any institutions
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be owned from supervisor and deputy vice –chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No: Shabnam Mehdizadeh GS31860

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the University Putra Malaysia(Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_  
Name of  
Chairman of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xiv
<b>LIST OF FIGURES</b>	xv
<b>LIST OF ABBREVIATIONS</b>	xvi
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	<b>3</b>
2.1 Characteristics of rambutan fruit (morphology and taxonomy)	3
2.2 Characteristics of rambutan Seed	4
2.2.1 Practical usage of rambutan seed	4
2.2.2 Nutritional value of rambutan seed	5
2.2.3 Health benefit	7
2.3 Characteristics of almond	7
2.3.1 Comparison the characteristics of rambutan seed and almond	8
2.4 Definition of bitter taste	9
2.4.1 Detection of bitter taste	9
2.4.2 Bitterness of phenols and tannins	10
2.4.3 Bitterness and consumer acceptability	11
2.4.4 The debittering process of plant foods	12
2.5 Bitterness in almond	14
2.6 Causes of bitterness in plants	15
2.6.1 Poly phenol	15
2.6.2 Tannin	15
2.6.3 Traces of alkaloid	16
2.6.4 Saponin	16
2.6.5 Amino acid	16
2.7 Definition of fermented food	16
2.7.1 Traditional fermentation in cocoa	17
2.7.2 Importance of fermentation	17
2.7.3 Fermentation and bitterness compounds	18
2.8 Reduction of bitterness by different processing methods	19
2.8.1 Chemical substances (lye)	19
2.8.2 Osmotic dehydration and impregnation processes	20
2.8.3 Soaking and washing as a natural methods	22
2.9 Role of roasting in food/ nut processing	22



<b>3</b>	<b>VARIABILITY IN THE FERMENTATION INDEX, POLYPHENOLS AND AMINO ACIDS OF SEEDS OF RAMBUTAN (<i>Nephelium lappaceum</i> L.) DURING FERMENTATION</b>	<b>23</b>
3.1	Introduction	23
3.2	Methods and materials	24
3.2.1	Raw material (plant material)	24
3.2.2	Chemicals and reagent	24
3.2.3	Rambutan seed fermentation process	24
3.2.4	Determination of pH, titratable acidity, and fermentation index	24
3.2.5	Lipid and protein contents determination	24
3.2.6	Color measurement	25
3.2.7	Sample preparation and determination of total saponin	25
3.2.8	Preparation of solvent extraction and determination of Tannin content	25
3.2.9	Preparation of seed extracts and phenolic content determination	26
3.2.10	Quantification of phenolic compounds	26
3.2.11	Determination of amino acid	26
3.2.12	Determination of sugar content	27
3.2.13	Statistical analysis	27
3.3	Results and discussion	27
3.3.1	Effect of fermentation on fermentation index, acids and pH	27
3.3.2	Effect of fermentation on lipid and protein contents	30
3.3.3	Color changes during fermentation	30
3.3.4	Changes in sugar content	33
3.3.5	Total polyphenols, tannins and saponins	33
3.3.6	Quantification of phenolic constituents in seeds of rambutan	35
3.3.7	Changes in free amino acid during fermentation process	36
3.4	Conclusion	38
<b>4</b>	<b>EFFECT OF DIFFERENT DE-BITTERING METHODS ON THE PHYSICOCHEMICAL PROPERTIES OF RAMBUTAN SEED (<i>Nephelium lappaceum</i> L.)</b>	<b>39</b>
4.1	Introduction	39
4.2	Materials and Methods	40
4.2.1	Sample preparation	40
4.2.2	Debitting processes	40
4.2.2.1	Soaking in water	40
4.2.2.2	Lye treatment (alkaline treatment)	41
4.2.2.3	Osmotic dehydration and impregnation process	41
4.2.2.4	Laboratory procedure for calculation of solid gain, water loss and weight reduction	42
4.2.3	Chemicals and Reagent	42
4.3	Statistical analysis	43
4.4	Results and Discussion	43
4.4.1	Lipid content and protein content	43
4.4.2	Changes of color during debittering methods	46
4.4.3	Water Loss and solid gain and weight reduction during osmotic dehydration process	47

4.4.4	Effect of different debittering methods on the total phenolic, tannin and saponin contents and polyphenol constituent in rambutan seed	49
4.4.4.1	Effect of soaking in water on the total phenolic, tannin and saponin contents	49
4.4.4.2	Effect of alkaline solution on the total phenolic, tannin and saponin contents	52
4.4.4.3	Effect of osmotic dehydration and impregnation process on the total phenolic, tannin and saponin contents	54
4.4.4.4	Quantification of phenolic constituents in different debittering methods	57
4.4.5	Changes of free amino acids during different debittering methods	61
4.5	Conclusion	65
<b>5</b>	<b>EFFECT OF DEBITTERING AND ROASTING ON THE VOLATILE COMPOUNDS OF RAMBUTAN SEEDS</b>	<b>66</b>
5.1	Introduction	66
5.2	Materials and Methods	67
5.2.1	Sample preparation	67
5.2.2	Roasting Conditions	67
5.2.3	Chemicals	67
5.2.4	Extraction	67
5.2.5	Gas chromatography –Mass spectrometry (GC-MS)	67
5.2.6	Color measurement	68
5.2.7	Textural analysis	68
5.3	Statistical Analysis	68
5.4	Results and discussion	68
5.4.1	Effect of roasting on the color values and hardness of debittered treatments	68
5.4.2	Identification of volatile compounds in unroasted rambutan seeds	69
5.4.3	Identification of volatile compounds in fermented and roasted rambutan seeds	74
5.4.4	Identification of volatile compounds of roasted rambutan seed treated with 22% NaCl, 8 h	76
5.4.5	Identification of volatile constituents of roasted debittered rambutan seed treated with 1 N NaOH for 8 h	78
5.4.6	Identification of volatile compounds of roasted seeds subjected to soaking treatment (144 h)	80
5.4.7	Effect of roasting on the volatile compounds of roasted almond	81
5.4.8	Identification of volatile compounds in roasted almond	82
5.5	Conclusion	89
<b>6</b>	<b>GENERAL CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	<b>91</b>



<b>REFERENCES</b>	<b>92</b>
<b>APPENDICES</b>	<b>116</b>
Appendix A: Standards of Polyphenolic compounds in HPLC	116
Appendix B: Polyphenol constituents of rambutan seed in day 0 and day 10 of fermentation	118
Appendix C: Standards of Amino acid and Amino acids in seeds of rambutan	119
Appendix D: Rambutan seed debittered by 7days soaking, 1N NaoH, 8h and 22% NaCl, 8h	120
Appendix E: Results of two-way ANOVA of bitterness components	121
Appendix F: GC-MS volatile of raw, roasted debittered rambutan seeds, and roasted almond	122
<b>BIODATA OF STUDENT</b>	<b>125</b>
<b>LIST OF PUBLICATIONS</b>	<b>126</b>



## LIST OF TABLES

<b>Table</b>	<b>Page</b>
2. 1 Fatty acids composition of rambutan seed	5
2. 2 Amino acid composition (g per 100 g protein) of rambutan seeds	6
2. 3 Chemical and mineral contents of rambutan ( <i>Nephellium lappaceum L.</i> ) seed and fruit	7
2.4 Comparison of the fatty acid in rambutan seeds and almonds	8
2. 5 Free amino acid in rambutan seeds and almonds	9
2. 6 Previous literature on different debittering methods	13
3. 1 Correlation between Variables and day of fermentation 29	
3. 2 Effect of fermentation duration on color and sugar variations in seed rambutan	32
3. 3 Effect of fermentation on changes of Total poly phenol (mg/g gallic acid), tannin (mg/g catechin), saponin (mg/g soya saponin) and polyphenol constituents	34
3. 4 Free amino acids during fermentation process of rambutan seed (mg g-1)	37
4. 1 Effect of different debittering methods on the color values and lipid and protein contents of rambutan seeds	45
4. 2 Effect of time and concentration of solute of impregnation and osmotic dehydration process on the	48
4. 3 Effect of different debittering methods on total phenolic content, tannin and saponin of rambutan seeds	51
4. 4 Effect of different debittering methods on polyphenol constituents of rambutan seeds	58
4. 5 Variation in free amino acid during osmotic dehydration process	64
5. 1 Effect of roasting on color values and hardness of roasted rambutan seeds	69
5. 2 Volatile constituents of unroasted rambutan seeds	71
5. 3 Volatile constituents of fermented roasted seeds of rambutan	75
5. 4 Volatile constituents of roasted rambutan seeds treated with 22%NaCl, 8 h	77
5. 5 Volatile constituents of roasted debittered rambutan seed treated with 1 N NaOH for 8 h	79
5. 6 Volatile constituents of roasted debittered rambutan seed subjected to soaking (144 h)	81
5. 7 Volatile constituents of roasted almond	84
5. 8 Summary of comparison of volatile compounds between samples	89

## LIST OF FIGURES

Figure	Page
3. 1A Effect of fermentation time on changes in lipid and protein contents [means with different letters are significantly different, $P < 0.05$ ]	28
3. 1B Effect of fermentation time on changes in pH and fermentation index (FI) [means with different letters are significantly different, $P < 0.05$ ]	28
3. 1C Acetic acid and lactic acid changes during fermentation [mean $\pm$ SD]	29
3. 2 Scatter plot matrix of the different responses: Red color within the box represents day 0, green color = days 1-5 and blue color days 6-10 of fermentation	30
3. 3 Phenolic constituents in seed rambutan	35
4. 1A TPC (mg/g)	53
4. 1B Tannin (mg/g)	53
4. 1C Saponin content (mg/g)	53
4. 2 Osmotic dehydration of rambutan seeds by salt (left) and sugar (right)	55
4. 3A TPC (mg/g)	56
4. 3B Tannin (mg/g)	56
4. 3C Saponin (mg/g)	56
4. 4A Gallic acid (mg/kg)	59
4. 4B Geraniin (mg/kg)	59
4. 4C Gallic acid (mg/kg)	60
4. 4D Geraniin (mg/kg)	60

## LIST OF ABBREVIATIONS

ANOVA	Analysis of variance
C	Corilagin
°C	Celsius
EA	Ellagic acid
FI	Fermentation Index
G	Geraniin
GA	Gallic Acid
GC-MS	Gas Chromatography- Mass Spectrometry
g	Gram
hr	Hour
h°	hue angles
HCl	Hydrochloric acid
HPLC	High Performance Liquid Chromatography
KCl	Potassium chloride
Kg	Kilogram
mg	Milligram
µg	Microgram
µl	Microliter
Min	Minute
N	Normality
NaCl	Sodium Chloride
NaOH	Sodium Hydroxide
Na <sub>2</sub> CO <sub>3</sub>	Sodium Carbonate
R <sup>2</sup>	Coefficient of determination
SG	Solid gain

TPC	Total Phenolic Content
tR	Retention time
WL	Water loss
WR	Weight reduction
w/w	Weight per weight
w/v	Weight/volume
3D	Three-dimensional
%	Percentage



## CHAPTER 1

### INTRODUCTION

Rambutan (*Nephelium lappaceum* L.) is one of the famous fruit that typically can be found in Southeast Asia belongs to the family of Sapindaceae (Marisa, 2006). Rambutan comes from the Malay word 'rambut' meaning hair. In Malaysia the rambutan bear its fruit two times in a year, the first, being in June and a second one in December. The rambutan is well adapted to humid climate and tropical area. It is widely cultivated in Malaysia with an annual production rate of (80, 000 tons/year), Indonesia (148, 000 tons/year), Thailand (430, 000 tons/year), and the Philippines (20, 000 tons/year). A warm climate is ideal for its growth and high yield. Rambutan tree is about 10-20 feet (3.05-6.1 m) high. The color of the skin can be red, yellow or green and the shape is round to oval, size is about 2" to 3" inch. The surface of fruit is made of spiny peel while inside of the fruit is a white and juicy acid pulp that is rich in vitamin C (Dadshani, 2002). The pulp is sweet, succulent and adheres to the seed (Nakasone; Paull, 1998; Wall, 2006). Among many appealing and desirable attributes of fruits from the tropics and subtropics is their characteristic flavor, which is the most remarkable to consumers. While rambutan fruit is still little known in many parts of the world, it is an important commercial crop in Asia (Morton 1987; Ong, Acree & Lavin, 1998).

The edible portion of rambutan (*Nephelium lappaceum* L.) is sweet and juicy; the light brown oval seed is generally remains as a waste material annually in large amount in Southeast Asia because it is slightly bitter (Kamel & Kakuda, 1992). The bitterness of the seeds of rambutan has been linked to traces of alkaloid, tannin, saponin and phenolic content like ellagic acid corillagin, geraniin and gallic acid (Kondo, Tsuda, Muto, Ueda, 2002; Singh, Murthy, Jayaprakasha, 2002). Another study has also shown that some Amino acids (AA) could have contributed to the bitter taste (Kano & Goto, 2003). As far as our literature survey could establish, there are little or no studies on the identification of phenolic compounds in the seed of *Nephelium*. Although a few polyphenolic components of longan seeds (same family with rambutan) have been characterized (Zheng et al., 2009). Although roasted rambutan seed are considered edible in some Asian countries, for more safety the seed must be evaluated before used (Gray, 2011). Furthermore, seeds are highly rich in vitamins such as vitamin C. They are also a valuable source of zinc, iron, Mg and Mn. Some experiments have revealed that seed of rambutan contains almost great quantity of fat with values between 14 g/100 g and 41 g/100 g (Solís- Fuentes, Camey-Ortiz, Hernández-Medel, Pérez-Mendoza & Durán-de-Bazúa, 2010) that is highly valued for manufacturing of food and production of soap and candle. The seed oil provides enough minerals necessary for human requirements (Standard from Institute of Medicine 2001). Main fatty acids in seed of rambutan (*Nephelium lappaceum* L.) are oleic acid (40.3%) and arachidic acid (34.5%). The processing of rambutan seeds for the reduction of bitterness (Supplied by MARDI) involves two major stages namely fermentation and application of debittering methods on rambutan seeds. Fermentation is used as the first stage in the bitterness reduction.

Fermentation, cause several chemical reactions and a deep adjustment of the biochemical feature of seeds (Timbie, Sechrist & Keeney, 1978; Gill, Macleod & Moreau, 1984). Certainly, sugars in the mucilage are changed into alcohol by micro-organism, during fermentation, heat is generated and many organic substances are produced such as ethanol, lactic, acetic acids and other organic acids leading to the death of the seed embryo (Kostinek, Ban-Koffi, Ottah-Atikpo, Teniola, Schillinger, Holzapfel & Franz, 2008). The mass then becomes runny and drops off from the beans. Enzymatic reactions occur, leading to the creation of flavor and changes in the color of cotyledons. These biochemical variations inside the beans cause the decrease of astringency and bitterness and the production of flavor precursors (Lagunes-Gálvez et al., 2007). Fermentation is one of the most widely used methods for food preparation before humans recognized the benefits. Firstly, fermentation was utilized to preserve food, modify the flavor, color as well as producing foods that are more digestible, and it could reduce undesirable and toxic constituents in foods, which is another significant aspect of the fermentation process (Jinap, Dimick & Hollander, 1995; Afoakwa, Quao, Takrama, Budu & Saalia, 2012). Fermentation is an earliest method for improving the shelf life, nutritional and organoleptic quality of foods (Doblado, Frias, Muñoz & Vidal-Valverde, 2003). Lately, fermentation as a kind of bioprocess system has been applied for the creation and removal of bioactive constituents in foodstuff, and in chemical and pharmaceutical productions (Martins, Mussatto, Martínez-Avila, Montañez-Saenz, Aguilar & Teixeira, 2011). Apart from fermentation, other debittering methods that have been used are soaking in water, blanching with sodium hydroxide and impregnation processes by sugar and salt. Roasting is an important stage in coffee bean and nuts processing (Ozdemir & Devres, 2000; Pittia, Rosa & Lerici, 2001; Saklar, Katnas & Urgan, 2001). Roasting is performed in order to promote flavor, color and desirable textural changes that eventually promote the complete accessibility of nuts. It is becoming increasingly difficult to ignore the enormous capacities of agricultural-food biomass waste discarded yearly all around the world. Moreover, disposal of agricultural-food waste can have a serious ecological impact on the environment. In recent years, there have been increased interests in numerous investigations on waste reuse (Singh, Chidambara Murthy & Jayaprakasha, 2002). Nowadays, large amounts of fruit most especially rambutan seeds (at least ~8000 metric tons of seed in Malaysia and ~60000 tons in other countries) are discarded annually at processing plants. Kamel and Kakuda, (1992) reported that to be economically viable, however, both oil and meal from these fruit seeds must be used.

Almond (*Prunus dulcis*) is one of the most reputable nuts with a high nutritional value, which have been used as a snack all over the world. In this study almond was used as control for roasting since the texture, volatile compounds and some nutritional values are similar to rambutan seed.

The main objectives of this study are:

- 1) To examine the effect of fermentation on fermentation index, poly phenol and amino acids profile of rambutan
- 2) To evaluate the effect of different debittering methods on the reduction of bitterness in rambutan seeds and
- 3) To determine the effect of roasting conditions on the aroma profile of debittered seeds.



## REFERENCES

- Abers, J. E., & Wrolstad, R. E. (1979). Causative factors of color deterioration in strawberry preserves during processing and storage. *Journal of Food Science*, 44(1), 75-81.
- Abu, O. A., Tewe, O. O., Losel, D. M., & Onifade, A. A. (2000). Changes in lipid, fatty acids and protein composition of sweet potato (*Ipomoea batatas*) after solid-state fungal fermentation. *Bioresource Technology*, 72(2), 189-192.
- Adams, M. R. (1990). Topical aspects of fermented foods. *Trends in Food Science & Technology*, 1, 140-144.
- Adeniran, H. A., Farinde, E. O., & Obatolu, V. A. (2013). Effect of Heat Treatment and Fermentation on Anti-Nutrients Content of Lima Bean (*Phaseolus lunatus*) during production of Daddawa Analogue. *Annual Review & Research in Biology*, 3 (3), 256-266.
- Adeyemi, O. T., Muhammad, N. O., & Oladiji, A. T. (2012). Biochemical assessment of the mineral and some anti-nutritional constituents of *Aspergillus niger* fermented *Chrysophyllum albidum* seed meal. *African Journal of Food Science*, 6(1), 20-28.
- Adler, E., Hoon, M. A., Mueller, K. L., Chandrashekar, J., Ryba, N. J., & Zuker, C. S. (2000). A novel family of mammalian taste receptors. *Cell*, 100(6), 693-702.
- Afoakwa, E. O., Paterson, A., & Fowler, M. (2007). Factors influencing rheological and textural qualities in chocolate—a review. *Trends in Food Science & Technology*, 18(6), 290-298.
- Afoakwa, E. O., Paterson, A., Fowler, M., & Ryan, A. (2008). Flavor formation and character in cocoa and chocolate: a critical review. *Critical Reviews in Food Science and Nutrition*, 48(9), 840-857.
- Afoakwa, E. O., Quao, J., Takrama, F. S., Budu, A. S., & Saalia, F. K. (2012). Changes in total polyphenols, o-diphenols and anthocyanin concentrations during fermentation of pulp pre-conditioned cocoa (*Theobroma cacao*) beans. *International Food Research Journal*, 19(3), 1071-1077.
- Afoakwa, E. O., Budu, A. S., Mensah-Brown, H., Felix Takrama, J., & Ofori-Ansah, E. (2014). Effect of Roasting Conditions on the Browning Index and Appearance Properties of Pulp Pre-Conditioned and Fermented Cocoa (*Theobroma Cacao*) Beans. *J Nutrition Health Food Sci*, 1(1), 5.
- Agila, A. (2012). Volatile Profile of Cashews (*Anacardium occidentale L.*), Almonds, and Honeys from Different Origins by Selected Ion Flow Tube Mass Spectrometry (Doctoral dissertation, The Ohio State University).
- Agila, A., & Barringer, S. (2012). Effect of roasting conditions on color and volatile profile including HMF level in sweet almonds (*Prunus dulcis*). *Journal of Food Science*, 77(4), C461-C468.
- Ahmad, N., & Mukhtar, H. (1999). Green tea polyphenols and cancer: biologic mechanisms and practical implications. *Nutrition Reviews*, 57(3), 78-83.
- Akasha, A., Haas, G., Meibach, R., & West, L. (2002). US20030203089 A1. Washington, DC: U.S.
- Alasalvar, C., & Shahidi, F. (Eds.). (2008). *Tree nuts: composition, phytochemicals, and health effects*. FL, CRC Press.
- Aliya, S., & Geervani, P. (1981). An assessment of the protein quality and vitamin B content of commonly used fermented products of legumes and millets. *Journal of the Science of Food and Agriculture*, 32(8), 837-842.



- Allen, J. G. (1998). Toxins and lupinosis. *Lupines as a crop plant biology, production and utilization*, 411-428.
- Alonso-Salces, R. M., Barranco, A., Abad, B., Berrueta, L. A., Gallo, B., & Vicente, F. (2004). Polyphenolic profiles of Basque cider apple cultivars and their technological properties. *Journal of Agricultural and Food Chemistry*, 52(10), 2938-2952.
- Ames, B. N., Profet, M., & Gold, L. S. (1990). Dietary pesticides (99.99% all natural). *Proceedings of the National Academy of Sciences*, 87(19), 7777-7781.
- Antony, U., & Chandra, T. S. (1998). Antinutrient reduction and enhancement in protein, starch, and mineral availability in fermented flour of finger millet (*Eleusine coracana*). *Journal of Agricultural and Food Chemistry*, 46(7), 2578-2582.
- AOAC. (1980). In Official Methods of Analysis. Association of Official Analytical Chemists, 13<sup>th</sup> Edition, Washington, D.C.
- AOAC, (1995). pH of acidified foods. Section 42.1.04, method 981.12. In: Cunniff, P.A. (Ed.), Official Methods of Analysis of AOAC International, 16th ed. AOAC International, Gaithersburg, MD, pp.2-4.
- Ardhana, M. M., & Fleet, G. H. (2003). The microbial ecology of cocoa bean fermentations in Indonesia. *International Journal of Food Microbiology*, 86(1), 87-99.
- Arrázola Paternina, G. (2002). Análisis de glucósidos cianogénicos en variedades de almendro: implicaciones en la mejora genética.
- Augustin, M. A., & Chua, B. C. (1988). Composition of rambutan seeds. *Pertanika*, 11(2), 211-215.
- Augustin, M. A., & Ling, E. T. (1987). Composition of mango seed kernel. *Pertanika (Malaysia)*.
- Authority, A. N. Z. F. (2001). Lupin alkaloids in food. *A Toxicological Review and Risk Assessment*.
- Auw, J. M., Blanco, V., O'keefe, S. F., & Sims, C. A. (1996). Effect of processing on the phenolic and color of Cabernet Sauvignon, Chambourcin, and Noble wines and juices. *American Journal of Enology and Viticulture*, 47(3), 279-286.
- Avila, I. M. L. B., & Silva, C. L. M. (1999). Modelling kinetics of thermal degradation of colour in peach puree. *Journal of Food Engineering*, 39(2), 161-166.
- Awada, S. H., Hady, A., Hassan, A. B., Ali, I., & Babiker, E. E. (2005). Antinutritional factors content and availability of protein, starch and mineral of maize (*Zeamays linnaus*) and lentil (*Lens culinaris*) as influenced by domestic processing. *Journal of Food Technology*, 3(4), 523-528.
- Axena, A. K., Chadha, M., & Sharma, S. (2003). Nutrients and antinutrients in chickpea (*Cicer arietinum* L.) cultivars after soaking and pressure cooking. *Journal of Food Science and Technology*, 40(5), 493-497.
- Bakare, M. A., Ameh, D. A., & Agbaji. A. S. (2011). Biochemical Assessment of Daddawa Food Seasoning Produced by Fermentation of Pawpaw (*Carica papaya*) Seeds. *Journal of Nutrition*, 10(3), 220-223.
- Baker, D. M., Tomlins, K. I., & Gay, C. (1994). Survey of Ghanaian cocoa farmer fermentation practices and their influence on cocoa flavour. *Food Chemistry*, 51(4), 425-431.

- Barampama, Z., & Simard, R. E. (1995). Effects of soaking, cooking and fermentation on composition, in-vitro starch digestibility and nutritive value of common beans. *Plant Foods for Human Nutrition*, 48(4), 349-365.
- Baxter, N. J., Lilley, T. H., Haslam, E., & Williamson, M. P. (1997). Multiple interactions between polyphenols and a salivary proline-rich protein repeat result in complexation and precipitation. *Biochemistry*, 36(18), 5566-5577.
- Bayram, M., Öner, M. D., & Kaya, A. (2004). Influence of soaking on the dimensions and colour of soybean for bulgur production. *Journal of Food Engineering*, 61(3), 331-339.
- Bazzano, L. A., He, J., Ogden, L. G., Loria, C. M., Vupputuri, S., Myers, L., & Whelton, P. K. (2002). Fruit and vegetable intake and risk of cardiovascular disease in US adults: the first National Health and Nutrition Examination Survey Epidemiologic Follow-up Study. *The American Journal of Clinical Nutrition*, 76(1), 93-99.
- Bchir, B., Besbes, S., Attia, H., & Blecker, C. (2009). Osmotic dehydration of pomegranate seeds: mass transfer kinetics and differential scanning calorimetry characterization. *International Journal of Food Science & Technology*, 44(11), 2208-2217.
- Beck, J. J., Higbee, B. S., Gee, W. S., & Dragull, K. (2011a). Ambient orchard volatiles from California almonds. *Phytochemistry Letters*, 4(2), 199-202.
- Beck, J. J., Mahoney, N. E., Cook, D., & Gee, W. S. (2011b). Volatile analysis of ground almonds contaminated with naturally occurring fungi. *Journal of Agricultural and Food Chemistry*, 59(11), 6180-6187.
- Beecher, G. R. (1999). Phytonutrients' role in metabolism: Effects on resistance to degenerative processes. *Nutrition Reviews*, 57(9), 3-6.
- Behrens, M., & Meyerhof, W. (2006). Bitter taste receptors and human bitter taste perception. *Cell Mol Life Sci*, 63(13), 1501-1509.
- Belitz, H. D., Grosch, W., & Schieberle, P. (2009). 5 Aroma Compounds. In Belitz, Grosch, & Schieberle (Eds.) *Food Chemistry* (pp 340-402), Berlin, Germany.
- Belitz H D, Grosch W. (1999). Cocoa and Chocolate. In *Food Chemistry 2nd ed.*, Springer-Verlag Berlin, 894-903.
- Belitz, H. D., & Wieser, H. (1985). Bitter compounds: Occurrence and structure-activity relationships. *Food Reviews International*, 1(2), 271-354.
- Belitz, H. D., Grosch, W., & Schieberle, P. (2001). *Lehrbuch der lebensmittelchemie* (Vol. 6). Berlin: Springer.
- Benz, K. (2002). Debacterization of cocoa beans gains increasing acceptance. Available from: <<http://www.buhlergroup.com/sc/en/publikationen.asp>>
- Breslin, P. A. S., & Beauchamp, G. K. (1995). Suppression of bitterness by sodium: variation among bitter taste stimuli. *Chemical senses*, 20(6), 609-623.
- Biehl, B., Meyer, B., Said, M. B., & Samarakoddy, R. J. (1990). Bean spreading: a method for pulp preconditioning to impair strong nib acidification during cocoa fermentation in Malaysia. *Journal of the Science of Food and Agriculture*, 51(1), 35-45.
- Birch, G. G., Cowell, N. D., & Young, R. H. (1972). Structural basis of and interaction between sweetness and bitterness in sugars. *Journal of the Science of Food and Agriculture*, 23(10), 1207-1212.
- Bishnoi, S., Khetarpaul, N., & Yadav, R. K. (1994). Effect of domestic processing and cooking methods on phytic acid and polyphenol contents of pea cultivars (*Pisum sativum*). *Plant Foods for Human Nutrition*, 45(4), 381-388.

- Boateng, J., Verghese, M., Walker, L. T., & Ogutu, S. (2008). Effect of processing on antioxidant contents in selected dry beans (*Phaseolus spp. L.*). *LWT-Food Science and Technology*, 41(9), 1541-1547.
- Bodenhofer, U., Kothmeier, A., & Hochreiter, S. (2011). AP Cluster: An R package for affinity propagation clustering. *Bioinformatics*, 27(17), 2463-2464.
- Bonvehí, J. S., & Coll, F. V. (2000). Evaluation of purine alkaloids and diketopiperazines contents in processed cocoa powder. *European Food Research and Technology*, 210(3), 189-195.
- Bosland, P. W., Votava, E. J., & Votava, E. M. (2012). Peppers: Vegetable and Spice Capsicums (Vol. 22). Cabi.
- Bravo, L. (1998). Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutrition Reviews*, 56(11), 317-333.
- Bressani, R. (1993). Grain quality of common beans. *Food Reviews International*, 9(2), 237-297.
- Broadhurst, R. B., & Jones, W. T. (1978). Analysis of condensed tannins using acidified vanillin. *Journal of the Science of Food and Agriculture*, 29(9), 788-794.
- Brockhoff, A., Behrens, M., Massarotti, A., Appendino, G., & Meyerhof, W. (2007). Broad tuning of the human bitter taste receptor hTAS2R46 to various sesquiterpene lactones, clerodane and labdane diterpenoids, strychnine, and denatonium. *Journal of Agricultural and Food Chemistry*, 55(15), 6236-6243.
- Buchaillet, A., Caffin, N., & Bhandari, B. (2009). Drying of lemon myrtle (*Backhousia citriodora*) leaves: Retention of volatiles and color. *Drying Technology*, 27(3), 445-450.
- Burdack-Freitag, A., & Schieberle, P. (2010). Changes in the key odorants of Italian hazelnuts (*Coryllus avellana L. var. Tonda Romana*) induced by roasting. *Journal of Agricultural and Food Chemistry*, 58(10), 6351-6359.
- Burns, D. J. W., & Noble, A. C. (1985). Evaluation of the Separate Contributions of Viscosity and Sweetness of Sucrose to Perceived Viscosity, Sweetness and Bitterness of Vermouth. *Journal of Texture Studies*, 16(4), 365-380.
- Cai, Y. Z., Sun, M., Xing, J., Luo, Q., & Corke, H. (2006). Structure-radical scavenging activity relationships of phenolic compounds from traditional Chinese medicinal plants. *Life Sciences*, 78(25), 2872-2888.
- Calvino, A. M., García-Medina, M. R., & Cometto-Muniz, J. E. (1990). Interactions in caffeine-sucrose and coffee-sucrose mixtures: evidence of taste and flavor suppression. *Chemical Senses*, 15(5), 505-519.
- Campbell-Platt, G. (1987). Fermented foods of the world. *A dictionary and Guide*. Butterworths.
- Camu, N., De Winter, T., Addo, S.K., Takrama, J.S., Bernaert, H. and De Vuyst, L. (2008). Fermentation of cocoa beans: Influence of microbial and polyphenol concentrations on flavor of cocoa. *Journal of the Science of Food and Agriculture*, 88, 2288-2297.
- Camu, N., De Winter, T., Verbrugghe, K., Cleenwerck, I., Vandamme, P., Takrama, J. S., & De Vuyst, L. (2007). Dynamics and biodiversity of populations of lactic acid bacteria and acetic acid bacteria involved in spontaneous heap fermentation of cocoa beans in Ghana. *Applied and Environmental Microbiology*, 73(6), 1809-1824.

- Candela, M., Astiasaran, I., & Bello, J. (1997). Cooking and warm-holding: effect on general composition and amino acids of kidney beans (*Phaseolus vulgaris*), chickpeas (*Cicer arietinum*), and lentils (*Lens culinaris*). *Journal of Agricultural and Food Chemistry*, 45(12), 4763-4767.
- Casado, F. J., Sánchez, A. H., Rejano, L., & Montaña, A. (2007). D-amino acid formation in sterilized alkali-treated olives. *Journal of Agricultural and Food Chemistry*, 55(9), 3503-3507.
- Chandrashekar, J., Hoon, M. A., Ryba, N. J., & Zuker, C. S. (2006). The receptors and cells for mammalian taste. *Nature*, 444(7117), 288-294.
- Chandrashekar, J., Mueller, K. L., Hoon, M. A., Adler, E., Feng, L., Guo, W., & Ryba, N. J. (2000). T2Rs function as bitter taste receptors. *Cell*, 100(6), 703-711.
- Chen, C. Y., & Blumberg, J. B. (2008). Phytochemical composition of nuts. *Asia Pac J Clin Nutr*, 17(Suppl 1), 329-332.
- Chen, C. Y., Milbury, P. E., Chung, S. K., & Blumberg, J. (2007). Effect of almond skin polyphenolics and quercetin on human LDL and apolipoprotein B-100 oxidation and conformation. *The Journal of Nutritional Biochemistry*, 18(12), 785-794.
- Chenlo, F., Moreira, R., Fernandez-Herrero, C., & Vázquez, G. (2006). Mass transfer during osmotic dehydration of chestnut using sodium chloride solutions. *Journal of Food Engineering*, 73(2), 164-173.
- Chenlo, F., Moreira, R., Fernández-Herrero, C., & Vázquez, G. (2007). Osmotic dehydration of chestnut with sucrose: Mass transfer processes and global kinetics modelling. *Journal of Food Engineering*, 78(3), 765-774.
- Chu, F. L., & Yaylayan, V. A. (2008). Model studies on the oxygen-induced formation of benzaldehyde from phenylacetaldehyde using pyrolysis GC-MS and FTIR. *Journal of Agricultural and Food Chemistry*, 56(22), 10697-10704.
- Chung, K. T., Wong, T. Y., Wei, C. I., Huang, Y. W., & Lin, Y. (1998). Tannins and human health: a review. *Critical Reviews in Food Science and Nutrition*, 38(6), 421-464.
- Clapperton, J., Yow, S., Chan, J., Lim, D., Lockwood, R., Romanczyk, L., & Hammerstone, J. (1994). The contribution of genotype to cocoa (*Theobroma cacao* L.) flavour. *Tropical Agriculture*, 71(4), 303-308.
- Coleman III, W. M., White Jr, J. L., & Perfetti, T. A. (1994). Characteristics of heat-treated aqueous extracts of peanuts and cashews. *Journal of Agricultural and Food Chemistry*, 42(1), 190-194.
- Committee Report. World Wide Web: <http://www.health.gov/dietaryguidelines/dga2005/report>.
- Contreras, J. E., & Smyrl, T. G. (1981). An evaluation of osmotic concentration of apple rings using corn syrup solids solutions. *Canadian Institute of Food Science and Technology Journal*, 14(4), 310-314.
- Dadshani, S.W. (2002). *Nephelium lappaceum* L. (Rambutan). *ARTS*, 3, 71-79.
- De Brito, E. S., García, N. H. P., Gallão, M. I., Cortelazzo, A. L., Fevereiro, P. S., & Braga, M. R. (2001). Structural and chemical changes in cocoa (*Theobroma cacao* L) during fermentation, drying and roasting. *Journal of the Science of Food and Agriculture*, 81(2), 281-288.
- De Vries, R. P., & Visser, J. (2001). Aspergillus enzymes involved in degradation of plant cell wall polysaccharides. *Microbiology and Molecular Biology Reviews*, 65(4), 497-522.



- Del Río, J. A., Arcas, M. C., Benavente, O., Sabater, F., & Ortuño, A. (1998). Changes of Polymethoxylated Flavones Levels during Development of EM EMTYPE. *Planta Medica*, 64(06), 575-576.
- Delahunty, C. M., & Piggott, J. R. (1995). Current methods to evaluate contribution and interactions of components to flavour of solid foods using hard cheese as an example. *International journal of food science & technology*, 30(5), 555-570.
- Deshpande, S. S., Sathe, S. K., Salunkhe, D. K., & Cornforth, D. P. (1982). Effects of dehulling on phytic acid, polyphenols, and enzyme inhibitors of dry beans (*Phaseolus vulgaris L.*). *Journal of Food Science*, 47(6), 1846-1850.
- Devic, E., Guyot, S., Daudin, J. D., & Bonazzi, C. (2009). Effect of temperature and cultivar on polyphenol retention and mass transfer during osmotic dehydration of apples. *Journal of Agricultural and Food Chemistry*, 58(1), 606-614.
- Devic, E., Guyot, S., Daudin, J. D., & Bonazzi, C. (2010). Kinetics of polyphenol losses during soaking and drying of cider apples. *Food and Bioprocess Technology*, 3(6), 867-877.
- Dicenta, F., Martínez-Gómez, P., Grané, N., Martín, M. L., León, A., Cánovas, J. A., & Berenguer, V. (2002). Relationship between cyanogenic compounds in kernels, leaves, and roots of sweet and bitter kernelled almonds. *Journal of Agricultural and Food Chemistry*, 50(7), 2149-2152.
- Diczbalis, Y., Eamus, D. and Menzel, C.M. (1996). Environmental factors influencing the growth and yield of rambutan grown in the wet/dry tropics of northern Australia. In 'Proceedings of the International Conference on Tropical fruits, 23-26 July 1996, KL, Malaysia.' Volume 2 (Compiled by S. Vijaysegran, m. Pauziah, M.S. Mohamed and S. Ahmad Tarmizi). Malaysian Agricultural Research and Development institute (MARDI), Serdang, Selangor.
- Dixon, G. M., & Jen, J. J. (1977). Changes Of Sugars And Acids Of Osmovac-Dried Apple Slices. *Journal of Food Science*, 42(4), 1126-1127.
- Djendoubi Mrad, N., Boudhrioua Mihoubi, N., Kechaou, N., Courtois, F., & Bonazzi, C. (2013). Effect of Osmo-dehydration Conditions on the Quality Attributes of Pears. *Journal of Food Process Technol*, 4(256), 2.
- Doblado, R., Frias, J., Muñoz, R., & Vidal-Valverde, C. (2003). Fermentation of (*Vigna sinensis var.*) carilla flours by natural microflora and Lactobacillus species. *Journal of Food Protection*®, 66(12), 2313-2320.
- Doyle, B., & Griffiths, L. A. (1980). The metabolism of ellagic acid in the rat. *Xenobiotica*, 10(4), 247-256.
- Drewnowski, A. (1996). From asparagus to zucchini: mapping cognitive space for vegetable names. *Journal of the American College of Nutrition*, 15(2), 147-153.
- Drewnowski, A. (1997). Taste preferences and food intake. *Annual Review of Nutrition*, 17(1), 237-253.
- Drewnowski, A. (2001). The science and complexity of bitter taste. *Nutrition reviews*, 59(6), 163-169.
- Drewnowski, A., & Gomez-Carneros, C. (2000). Bitter taste, phytonutrients, and the consumer: a review. *The American Journal of Clinical Nutrition*, 72(6), 1424-1435.
- DRIs (2001). Dietary Reference Intakes: Estimated Mineral Average. Food & Nutrition Board, Institute of Medicine, the National Academic Press, Washington, DC.

- Drumm, T. D., Gray, J. I., Hosfield, G. L., & Uebersax, M. A. (1990). Lipid, saccharide, protein, phenolic acid and saponin contents of four market classes of edible dry beans as influenced by soaking and canning. *Journal of the Science of Food and Agriculture*, 51(4), 425-435.
- Eckert, M., & Riker, P. (2007). Overcoming challenges in functional beverages. *Food technology*.
- Eka, O. U. (1980). Effect of fermentation on the nutrient status of locust beans. *Food chemistry*, 5(4), 303-308.
- El Alaily, H. A., Anwar, A., & El Banna, I. (1976). Mango seed kernels as an energy source for chicks 1. *British Poultry Science*, 17(2), 129-133.
- El Hag, M. E., El Tinay, A. H., & Yousif, N. E. (2002). Effect of fermentation and dehulling on starch, total polyphenols, phytic acid content and in vitro protein digestibility of pearl millet. *Food Chemistry*, 77(2), 193-196.
- El Nahry, F., Darwish, N. M., & Tharwat, S. (1977). Effect of preparation and cooking on the nutritive value of local kidney bean. *Qualitas Plantarum*, 27(2), 141-150.
- ElMaki, H. B., AbdelRahaman, S. M., Idris, W. H., Hassan, A. B., Babiker, E. E., & El Tinay, A. H. (2007). Content of antinutritional factors and HCl-extractability of minerals from white bean (*Phaseolus vulgaris*) cultivars: Influence of soaking and/or cooking. *Food Chemistry*, 100(1), 362-368.
- Engel, E., Nicklaus, S., Septier, C., Salles, C., & Le Quéré, J. L. (2000). Taste active compounds in a goat cheese water-soluble extract. 2. Determination of the relative impact of water-soluble extract components on its taste using omission tests. *Journal of Agricultural and Food Chemistry*, 48(9), 4260-4267.
- Erbas, M. (2010). The effects of different debittering methods on the production of lupin bean snack from bitter *Lupinus albus L.* seeds. *Journal of Food Quality*, 33(6), 742-757.
- Espinosa Diaz, M. (2004). Comparison between orthonasal and retronasal flavour perception at different concentrations. *Flavour and Fragrance Journal*, 19(6), 499-504.
- European Union, Off. J. Eur. Commun., L182, European Community, Brussels, 1999.
- European Union, Off. J. Eur. Commun., L312/3, European Community, Brussels, 2008.
- Fahey, J. W., Stephenson, K. K., Talalay, P., Shibamoto, T., Terao, J., & Osawa, T. (1998). Glucosinolates, myrosinase, and isothiocyanates: three reasons for eating brassica vegetables. In Functional foods for disease prevention I. Fruits, vegetables, and teas. Symposium sponsored by the Division of Agricultural and Food Chemistry at the 213th National Meeting of the American Chemical Society, San Francisco, California, USA, April 13-17, 1997.(pp. 16-22). *American Chemical Society*.
- Fan, X. (2005). Formation of furan from carbohydrates and ascorbic acid following exposure to ionizing radiation and thermal processing. *Journal of Agricultural and Food Chemistry*, 53(20), 7826-7831.
- FAO (1970). Amino acid content of foods and biological data on proteins, FAO Nutrition Studies, Rome.
- Fatemi, H. (2010). *Textbook of Food Chemistry*. The Enteshar Company, Tehran, Iran.
- Fenwick, D. E. & Oakenfull, D. (1983). Saponin content of food products and prepared foods. *Journal of the Science of Food and Agriculture* 34, 186-91.

- Fenwick, G. R., Griffiths, N. M., & Heaney, R. K. (1983). Bitterness in Brussels sprouts (*Brassica oleracea* L. var. *gemmifera*): the role of glucosinolates and their breakdown products. *Journal of the Science of Food and Agriculture*, 34(1), 73-80.
- Fernandes, A. C., Nishida, W., & da Costa Proença, R. P. (2010). Influence of soaking on the nutritional quality of common beans (*Phaseolus vulgaris* L.) cooked with or without the soaking water: a review. *International Journal of Food Science & Technology*, 45(11), 2209-2218.
- Fernández, A. G., Fernandez-Diez, M. J., & Adams, M. R. (1997). Table olives: production and processing. *Springer*.
- Fernández, M., López-Jurado, M., Aranda, P., & Urbano, G. (1996). Nutritional assessment of raw and processed faba bean (*Vicia faba* L.) cultivar major in growing rats. *Journal of Agricultural and Food Chemistry*, 44(9), 2766-2772.
- Ferreira, D., Guyot, S., Marnet, N., Delgadillo, I., Renard, C. M., & Coimbra, M. A. (2002). Composition of phenolic compounds in a Portuguese pear (*Pyrus communis* L. var. *S. Bartolomeu*) and changes after sun-drying. *Journal of Agricultural and Food Chemistry*, 50(16), 4537-4544.
- Folin, O., & Ciocalteu, V. (1927). On tyrosine and tryptophane determinations in proteins. *J. biol. Chem*, 73(2), 627-650.
- Foo, L. Y. (1993). Amariin, a di-dehydrohexahydroxydiphenoyl hydrolysable tannin from *Phyllanthus amarus*. *Phytochemistry*, 33(2), 487-491.
- Forsyth, W. G., & Quesnel, V. C. (1963). The mechanism of cacao curing. *Adv Enzymol Relat Areas Mol Biol*, 25, 457-492.
- Fowler, M. S. (2009). Cocoa beans: from tree to factory. *Industrial Chocolate Manufacture and Use, Fourth Edition*, 10-47.
- Frankel, E. N., Waterhouse, A. L., & Teissedre, P. L. (1995). Principal phenolic phytochemicals in selected California wines and their antioxidant activity in inhibiting oxidation of human low-density lipoproteins. *Journal of Agricultural and Food chemistry*, 43(4), 890-894.
- Fraser, G. E., Sabate, J., Beeson, W. L., & Strahan, T. M. (1992). A possible protective effect of nut consumption on risk of coronary heart disease: the Adventist Health Study. *Archives of Internal Medicine*, 152(7), 1416-1424.
- Frehner, M., Scalet, M., & Conn, E. E. (1990). Pattern of the Cyanide-Potential in Developing Fruits Implications for Plants Accumulating Cyanogenic Monoglucosides (*Phaseolus lunatus*) or Cyanogenic Diglucosides in Their Seeds (*Linum usitatissimum*, *Prunus amygdalus*). *Plant physiology*, 94(1), 28-34.
- Duffus, C. M., & Duffus, J. H. (Eds.). (1991). Toxic substances in crop plants. *Elsevier*.
- Gálvez, S., Loiseau, G., Paredes, J. L., Barel, M., & Guiraud, J. P. (2007). Study on the microflora and biochemistry of cocoa fermentation in the Dominican Republic. *International Journal of Food Microbiology*, 114(1), 124-130.
- García-Viguera, C., Zafrilla, P., Romero, F., Abellán, P., Artés, F., & Tomás-Barberán, F. A. (1999). Color stability of strawberry jam as affected by cultivar and storage temperature. *Journal of Food Science*, 64(2), 243-247.
- Garrido, I., Monagas, M., Gómez-Cordovés, C., & Bartolomé, B. (2008). Polyphenols and antioxidant properties of almond skins: influence of industrial processing. *Journal of Food Science*, 73(2), C106-C115.



- Gayte-sorbier, A., Airaudo, C. B., & Armand, P. (1985). Stability of glutamic acid and monosodium glutamate under model system conditions: influence of physical and technological factors. *Journal of Food Science*, 50(2), 350-352.
- Gill, M. S., Macleod, A. J., & Moreau, M. (1984). Volatile components of cocoa with particular reference to glucosinolate products. *Phytochemistry*, 23(9), 1937-1942.
- Gillette, M. (1985). Flavor effects of sodium chloride. *Food technology (USA)*.
- Glanz, K., Basil, M., Maibach, E., Goldberg, J., & Snyder, D. A. N. (1998). Why Americans eat what they do: taste, nutrition, cost, convenience, and weight control concerns as influences on food consumption. *Journal of the American Dietetic Association*, 98(10), 1118-1126.
- Glendinning, J. I. (1994). Is the bitter rejection response always adaptive. *Physiology & Behavior*, 56(6), 1217-1227.
- Goldman, I. L., Kader, A. A., & Heintz, C. (1999). Influence of production, handling, and storage on phytonutrient content of foods. *Nutrition Reviews*, 57(9), 46-52.
- Gorinstein, S., Zemser, M., Haruenkit, R., Chuthakorn, R., Grauer, F., Martin-Belloso, O., & Trakhtenberg, S. (1999). Comparative content of total polyphenols and dietary fiber in tropical fruits and persimmon. *The Journal of Nutritional Biochemistry*, 10(6), 367-371.
- Gourieva, K. B., & Tserevitinov, O. B. (1979). Method of evaluating the degree of fermentation of cocoa beans. *USSR patent*, 646.
- Granito, M., Frias, J., Doblado, R., Guerra, M., Champ, M., & Vidal-Valverde, C. (2002). Nutritional improvement of beans (*Phaseolus vulgaris*) by natural fermentation. *European Food Research and Technology*, 214(3), 226-231.
- Gray, N. (2011). Seed waste may be source of new fats. Retrieved April 29, 2012 from <http://www.foodnavigator.com/Science-Nutrition/Seed-waste-may-be-source-of-new-fats-Study>.
- Grundhöfer, P., Niemetz, R., Schilling, G., & Gross, G. G. (2001). Biosynthesis and subcellular distribution of hydrolysable tannins. *Phytochemistry*, 57(6), 915-927.
- Guadagni, D. G., Maier, V. P., & Turnbaugh, J. H. (1974). Some factors affecting sensory thresholds and relative bitterness of limonin and naringin. *Journal of the Science of Food and Agriculture*, 25(10), 1199-1205.
- HA, E. Y., Morr, C. V., & SEO, A. (1992). Isoflavone aglucones and volatile organic compounds in soybeans; effects of soaking treatments. *Journal of Food Science*, 57(2), 414-417.
- Hammami, C., Rene, F., & Marin, M. (1999). Process-quality optimization of the vacuum freeze-drying of apple slices by the response surface method. *International journal of food science & technology*, 34(2), 145-160.
- Hansen, C. E., Del Olmo, M., & Burri, C. (1998). Enzyme activities in cocoa beans during fermentation. *Journal of the Science of Food and Agriculture*, 77, 273-281.
- Harnly, J. M., Doherty, R. F., Beecher, G. R., Holden, J. M., Haytowitz, D. B., Bhagwat, S., & Gebhardt, S. (2006). Flavonoid content of US fruits, vegetables, and nuts. *Journal of Agricultural and Food Chemistry*, 54(26), 9966-9977.
- Harwood, J. L., & Stumft, P. K. (1970) Fat metabolism in higher plants. XL. Synthesis of fatty acids in the initial stage of seed germination. *Plant Physiol* 46:500.



- Hawladar, M. N. A., Perera, C. O., & Tian, M. (2006). Properties of modified atmosphere heat pump dried foods. *Journal of Food Engineering*, 74(3), 392-401.
- Heaney, R. K., & Fenwick, G. R. (1980). Glucosinolates in Brassica vegetables. Analysis of 22 varieties of Brussels sprout (*Brassica oleracea* var. *gemmifera*). *Journal of the Science of Food and Agriculture*, 31(8), 785-793.
- Heaney, R. K., & Fenwick, G. R. (1980). The analysis of glucosinolates in *Brassica* species using gas chromatography. Direct determination of the thiocyanate ion precursors, glucobrassicin and neoglucobrassicin. *Journal of the Science of Food and Agriculture*, 31(6), 593-599.
- Hellekant G., Gopal V., Breslin P. A., Beauchamp G. K. (1995). *Chem. Senses.*, 20, 609-623.
- Hemingway, R. W., Laks, P. E., & Branham, S. J. (Eds.). (1992). Plant polyphenols: synthesis, properties, significance (Vol. 59). *Springer*.
- Henney, J. E., Taylor, C. L., & Boon, C. S. (2010). Taste and flavor roles of sodium In foods: a unique challenge to reducing sodium intake.
- Hertog, M. G., Hollman, P. C., & Katan, M. B. (1992). Content of potentially anticarcinogenic flavonoids of 28 vegetables and 9 fruits commonly consumed in the Netherlands. *Journal of Agricultural and Food Chemistry*, 40(12), 2379-2383.
- Hiai, S., Oura, H., & Nakajima, T. (1976). Color Reaction of Some Sapogenins and Saponins with Vanillin and Sulfuric Acid. *Planta Medica*, 29(02), 116-122.
- Hii, C. L., Law, C. L., Cloke, M., & Suzannah, S. (2009). Thin layer drying kinetics of cocoa and dried product quality. *Biosystems Engineering*, 102(2), 153-161.
- Hladik, C. M., & Simmen, B. (1996). Taste perception and feeding behavior in nonhuman primates and human populations. *Evolutionary Anthropology: Issues, News, and Reviews*, 5(2), 58-71.
- Hooda, S., & Jood, S. (2003). Effect of soaking and germination on nutrient and antinutrient contents of fenugreek (*Trigonella foenum graecum* L.). *Journal of Food Biochemistry*, 27(2), 165-176.
- Hostettmann, K., & Marston, A. (2005). Saponins. Cambridge University Press. html.
- Huang, A. L., Chen, X., Hoon, M. A., Chandrashekar, J., Guo, W., Tränkner, D., ... & Zuker, C. S. (2006). The cells and logic for mammalian sour taste detection. *Nature*, 442(7105), 934-938.
- Huma, N., Anjum, M., Sehar, S., Khan, M. I., & Hussain, S. (2008). Effect of soaking and cooking on nutritional quality and safety of legumes. *Nutrition & Food Science*, 38(6), 570-577.
- Hurst, W. J., Martin, R. A., & Zoumas, B. L. (1979). Application of HPLC to characterization of individual carbohydrates in foods. *Journal of Food Science*, 44(3), 892-895.
- Hutton, T. (2002). Sodium technological functions of salt in the manufacturing of food and drink products. *British Food Journal*, 104(2), 126-152.
- Igbedioh, S. O., Shaire, S., & Aderiyi, B. J. I. (1995). Effects of processing on total phenols and proximate composition of pigeon pea (*Cajanus cajan*) and climbing bean (*Vigna umbellata*). *Journal of Food Science and Technology*, 32(6), 497-500.
- Ikan, R. (1999). Naturally occurring glycosides. John Wiley & Sons.

- In G. Reed (ed.), *Biotechnology, a comprehensive treatise*, vol. 5. Verlag in wine after the end of fermentation by *Saccharomyces cerevisiae* strains. *Journal of the Science of Food and Agriculture*, 83(8), 830-835.
- Ishikawa, T., & Noble, A. C. (1995). Temporal perception of astringency and sweetness in red wine. *Food Quality and Preference*, 6(1), 27-33.
- Islam, M. N., & Flink, J. N. (1982). Dehydration of potato. *International Journal of Food Science & Technology*, 17(3), 387-403.
- Ito, H., Iguchi, A., & Hatano, T. (2007). Identification of urinary and intestinal bacterial metabolites of ellagitannin geraniin in rats. *Journal of Agricultural and Food Chemistry*, 56(2), 393-400.
- Jadhav, S. J., Sharma, R. P., & Salunkhe, D. K. (1981). Naturally occurring toxic alkaloids in foods. *CRC Critical Reviews in Toxicology*, 9(1), 21-104.
- Janovitz-Klapp, A. H., Richard, F. C., Goupy, P. M., & Nicolas, J. J. (1990). Inhibition studies on apple polyphenol oxidase. *Journal of Agricultural and Food Chemistry*, 38(4), 926-931.
- Jenkins, D. J., Kendall, C. W., Marchie, A., Josse, A. R., Nguyen, T. H., Faulkner, D. A., & Singer, W. (2008). Effect of almonds on insulin secretion and insulin resistance in nondiabetic hyperlipidemic subjects: a randomized controlled crossover trial. *Metabolism*, 57(7), 882-887.
- Jinap, S., & Dimick, P. S. (1991). Effect of roasting on acidic characteristics of cocoa beans. *Journal of the Science of Food and Agriculture*, 54(2), 317-321.
- Jinap, S., Dimick, P. S., & Hollender, R. (1995). Flavour evaluation of chocolate formulated from cocoa beans from different countries. *Food Control*, 6(2), 105-110.
- Jinap, S., Jamilah, B., & Nazamid, S. (2003). Effects of incubation and polyphenol oxidase enrichment on colour, fermentation index, procyanidins and astringency of unfermented and partly fermented cocoa beans. *International Journal of Food Science & Technology*, 38(3), 285-295.
- Jinap, S., Rosli, W. I., Russly, A. R., & Nordin, L. M. (1998). Effect of roasting time and temperature on volatile component profiles during nib roasting of cocoa beans (*Theobroma cacao*). *Journal of the Science of Food and Agriculture*, 77(4), 441-448.
- Jinap, Thien & Yap (1994). Effect of drying on acidity and volatile fatty acids content of cocoa beans. *Journal of the Science of Food and Agriculture*, 65, 67-75.
- Jinap, M., Nazamid, S., & Jamilah, B. (2002). Activation of remaining key enzymes in dried under-fermented cocoa beans and its effect on aroma precursor formation. *Food chemistry*, 78(4), 407-417.
- Jood, S., Bishnoi, S., & Sehgal, S. (1998). Effect of processing on nutritional and antinutritional factors of moongbean cultivars. *Journal of Food Biochemistry*, 22(3), 245-257.
- Jood, S., Chauhan, B. M., & Kapoor, A. C. (1987). Polyphenols of chickpea and blackgram as affected by domestic processing and cooking methods. *Journal of the Science of Food and Agriculture*, 39(2), 145-149.
- Kahyaoglu, T., & Kava, S. (2006). Modeling of moisture, color and texture changes in sesame seeds during the conventional roasting. *Journal of Food Engineering*, 75(2), 167-177.
- Kallithraka, S., Bakker, J., & Clifford, M. N. (1997). Evaluation Of Bitterness and Astringency Of (+)-Catechin and (-)-Epicatechin in Red Wine and in Model Solution. *Journal of Sensory Studies*, 12(1), 25-37.

- Kamel, B. S., & Kakuda, Y. (1992). Characterization of the seed oil and meal from apricot, cherry, nectarine, peach and plum. *Journal of the American Oil Chemists' Society*, 69(5), 492-494.
- Kamen, J. M., Pilgrim, F. J., Gutman, N. J., & Kroll, B. J. (1961). Interactions of suprathreshold taste stimuli. *Journal of Experimental Psychology*, 62(4), 348. Kroeze J. H., Bartoshuk L. M., *Physiol. Behav.*, 35, 779—83 (1985).
- Kano, Y., & Goto, H. (2003). Relationship between the occurrences of bitter fruit in cucumber (*Cucumis sativus* L.) and the contents of total nitrogen, amino acid nitrogen, protein and HMG-CoA reductase activity. *Scientia Horticulturae*, 98, 1–8.
- Kantha, S. S., & Erdman, J. W. (1984). The winged bean as an oil and protein source: A review. *Journal of the American Oil Chemists' Society*, 61(3), 515-525.
- Kataria, A., Chauhan, B. M., & Gandhi, S. (1988). Effect of domestic processing and cooking on the antinutrients of black gram. *Food Chemistry*, 30(2), 149-156.
- Kaya, S. (2002). Effect of salt on hardness and whiteness of Gaziantep cheese during short-term brining. *Journal of Food Engineering*, 52(2), 155-159.
- Keast, R. S., Breslin, P. A., & Beauchamp, G. K. (2001). Suppression of bitterness using sodium salts. *Chimia International Journal for Chemistry*, 55(5), 441-447.
- Keast, R. S., Canty, T. M., & Breslin, P. A. (2004). The influence of sodium salts on binary mixtures of bitter-tasting compounds. *Chemical Senses*, 29(5), 431-439.
- Keurentjes, J. T. F., Janssen, A. E. M., Broek, A. P., Van der Padt, A., Wesselingh, J. A., & Van't Riet, K. (1992). Multicomponent diffusion in dialysis membranes. *Chemical Engineering Science*, 47(8), 1963-1971.
- Khetarpaul, N., & Chauhan, B. M. (1989). Effect of fermentation on protein, fat, minerals and thiamine content of pearl millet. *Plant Foods for Human Nutrition*, 39(2), 169-177.
- Khan, M. R. (2012). Osmotic dehydration technique for fruits preservation—a review. *Pakistan Journal of Food Sciences*, 22(2), 71-85.
- Khandelwal, S., Udipi, S. A., & Ghugre, P. (2010). Polyphenols and tannins in Indian pulses: Effect of soaking, germination and pressure cooking. *Food Research International*, 43(2), 526-530.
- Kheiri, M. S. A., & bin Mohammad Som, M. N. (1979). Physico-chemical characteristics of rambutan (*Nephelium lappaceum*) kernel fat [Peninsular Malaysia]. *Agricultural Product Utilisation Report (Malaysia)*. no. 186.
- Kiefl, J., Cordero, C., Nicolotti, L., Schieberle, P., Reichenbach, S. E., & Bicchi, C. (2012). Performance evaluation of non-targeted peak-based cross-sample analysis for comprehensive two-dimensional gas chromatography–mass spectrometry data and application to processed hazelnut profiling. *Journal of Chromatography A*, 1243, 81-90.
- Kim, J. S., & Lee, Y. S. (2009). Enolization and racemization reactions of glucose and fructose on heating with amino-acid enantiomers and the formation of melanoidins as a result of the Maillard reaction. *Amino acids*, 36(3), 465-474.
- Klein, L. B., & Mondy, N. I. (1981). Comparison of microwave and conventional baking of potatoes in relation to nitrogenous constituents and mineral composition. *Journal of Food Science*, 46(6), 1874-1877.
- Köksal, A. İ., Artık, N., Şimşek, A., & Güneş, N. (2006). Nutrient composition of hazelnut (*Corylus avellana* L.) varieties cultivated in Turkey. *Food Chemistry*, 99(3), 509-515.

- Kondo, S., Tsuda, K., Muto, N., Ueda, J.-E., (2002). Antioxidative activity of apple skin or flesh extracts associated with fruit development on selected apple cultivars. *Sci. Hort.* 96, 177–185.
- Kong, F., & Singh, R. P. (2009). Digestion of raw and roasted almonds in simulated gastric environment. *Food Biophysics*, 4(4), 365-377.
- Kostinek, M., Ban-Koffi, L., Ottah-Atikpo, M., Teniola, D., Schillinger, U., Holzappel, W. H., & Franz, C. M. (2008). Diversity of predominant lactic acid bacteria associated with cocoa fermentation in Nigeria. *Current microbiology*, 56(4), 306-314.
- Kowalska, H., & Lenart, A. (2001). Mass exchange during osmotic pretreatment of vegetables. *Journal of Food Engineering*, 49(2), 137-140.
- Kratzer, U., Frank, R., Kalbacher, H., Biehl, B., Wöstemeyer, J., & Voigt, J. (2009). Subunit structure of the vicilin-like globular storage protein of cocoa seeds and the origin of cocoa-and chocolate-specific aroma precursors. *Food Chemistry*, 113(4), 903-913.
- Kreutzmann, S., Christensen, L. P., & Edelenbos, M. (2008). Investigation of bitterness in carrots (*Daucus carota*L.) based on quantitative chemical and sensory analyses. *LWT-Food Science and Technology*, 41(2), 193-205.
- Kyi, T. M., Daud, W. R. W., Mohammad, A. B., Wahid Samsudin, M., Kadhum, A. A. H., & Talib, M. Z. M. (2005). The kinetics of polyphenol degradation during the drying of Malaysian cocoa beans. *International Journal of Food Science & Technology*, 40(3), 323-331.
- La Londe, R. T.; Perakyla, H.; Hayes, M. P. (1990). Potentially mutagenic, chlorine-substituted 2 (5H)-furanones: studies of their synthesis and NMR properties. *J. Org. Chem.* , 55, 2847-2855.
- Lagunes Gálvez, S., Loiseau, G., Paredes, J. L., Barel, M., & Guiraud, J. P. (2007). Study on the microflora and biochemistry of cocoa fermentation in the Dominican Republic. *International Journal of Food Microbiology*, 114(1), 124-130.
- Lamberts, L., Brijs, K., Mohamed, R., Verhelst, N., & Delcour, J. A. (2006). Impact of browning reactions and bran pigments on color of parboiled rice. *Journal of Agricultural and Food Chemistry*, 54(26), 9924-9929.
- Land DG. (1996). Perspectives On the Effect on Interactions of Flavour Perception: An Overview. In: McGorin RJ, Leland JV , editors. *Flavor-Food Interactions. ACS Symposium Series 633. Washington DC: Am Chem Soc.* p 2-11.
- Laohakunjit, N., Kerchoechuen, O., Matta, F. B., Silva, J. L., & Holmes, W. E. (2007). Postharvest survey of volatile compounds in five tropical fruits using headspace-solid phase microextraction (HS-SPME). *HortScience*, 42(2), 309-314.
- Lasekan, O., & Abbas, K. (2010). Analysis of volatile flavour compounds and acrylamide in roasted Malaysian tropical almond (*Terminalia catappa*) nuts using supercritical fluid extraction. *Food and Chemical Toxicology*, 48(8), 2212-2216.
- Lazarides, H. N., Katsanidis, E., & Nickolaidis, A. (1995). Mass transfer kinetics during osmotic preconcentration aiming at minimal solid uptake. *Journal of Food Engineering*, 25(2), 151-166.
- Le Bourvellec, C., Le Ouéré, J. M., Sanoner, P., Drilleau, J. F., & Guvot, S. (2004). Inhibition of apple polyphenol oxidase activity by procyanidins and polyphenol oxidation products. *Journal of Agricultural and Food Chemistry*, 52(1), 122-130.



- Lee, S. Y., Yoo, S. S., Lee, M. J., Kwon, I. B., & Pyun, Y. R. (2001). Optimization of nibs roasting in cocoa bean processing with lotte-better taste and color process. *Food Science and Biotechnology*, 10(3), 286-293.
- Lehrian, D. W. and Patterson, G. R. (1983). Cocoa fermentation, In: *Biotechnology, a Comprehensive Treatise*, vol. 5, ed. by Reed G. Verlag Chemie, Basel, pp. 529–575.
- Lehrian, D. W., and G. R. Patterson. (1983). Cocoa fermentation. In: Reed, G. (Ed.), *Biotechnology*, vol. 5. Verlag Chemie, Basel, Switzerland, pp. 529–575.
- Lehrian, D.W, Patterson, G.R. (1983). Cocoa fermentation. In *Biotechnology*, vol. 5, ed.by Rehm, H.J. & Reed, G. Verlag Chemie, Weinheim, PP.529-575.
- Leong, L. P., & Shui, G. (2002). An investigation of antioxidant capacity of fruits in Singapore markets. *Food Chemistry*, 76(1), 69-75.
- Lerici, C. R., Pinnavaia, G., Rosa, M. D., & Bartolucci, L. (1985). Osmotic dehydration of fruit: Influence of osmotic agents on drying behavior and product quality. *Journal of Food Science*, 50(5), 1217-1219.
- Ley, J. P. (2008). Masking bitter taste by molecules. *Chemosensory Perception*, 1(1), 58-77.
- Lim, T. K., & Diczbalis, Y. (1998). Rambutan. *The new rural industries—a handbook for farmers and investors*. Rural Industries Research & Development Corporation, Canberra, 306-313.
- Liu, D., Su, Z., Wang, C., Gu, M., & Xing, S. (2010). Separation and purification of hydrolyzable tannin from *Geranium wilfordii* Maxim by reversed-phase and normal-phase high-speed counter-current chromatography. *Journal of Separation Science*, 33(15), 2266-2271.
- López de Lerma, N., Peinado, J., Moreno, J., & Peinado, R. A. (2010). Antioxidant activity, browning and volatile Maillard compounds in Pedro Ximénez sweet wines under accelerated oxidative aging. *LWT-Food Science and Technology*, 43(10), 1557-1563.
- López, A. (1983). Factors associated with cacao bean acidity and the possibility of its reduction by improved fermentation. *Revista Theobroma*, 13(3), 233-248.
- Lopez, A. S. (1986). Chemical changes occurring during the processing of cacao.
- Lopez, A. S., and P. S. Dimick. (1995). Cocoa fermentation, p. 561–577. In G. Reed and T. W. Nagodawithana (ed.), *Enzymes, biomass, food and feed*, 2<sup>nd</sup>ed. VCH, Weinheim, Germany.
- Lopez, A.S.F. & Dimick, P.S. (1995). Cocoa Fermentation. In: Reed, G. and T.W. Nagodawithana (Eds.), *Enzymes, Biomass, Food and Feed*. 2nd E dn., *Biotechnology, Weinheim, Germany, VCH*, 9: 561-577.
- Luger, P., Weber, M., Kashino, S., Amakura, Y., Yoshida, T., Okuda, T., & Dauter, Z. (1998). Structure of the tannin geraniin based on conventional X-ray data at 295 K and on synchrotron data at 293 and 120 K. *Acta Crystallographica Section B: Structural Science*, 54(5), 687-694.
- Madrau, M. A., Piscopo, A., Sanguinetti, A. M., Del Caro, A., Poiana, M., Romeo, F. V., & Piga, A. (2009). Effect of drying temperature on polyphenolic content and antioxidant activity of apricots. *European Food Research and Technology*, 228(3), 441-448.
- Maga, J. A. (1990). Compound structure versus bitter taste. *Developments in Food Science*, 25, 35-48.

- Maignial, L., Pibarot, P., Bonetti, G., Chaintreau, A., & Marion, J. P. (1992). Simultaneous distillation—extraction under static vacuum: isolation of volatile compounds at room temperature. *Journal of Chromatography A*, 606(1), 87-94.
- Makkar, H. P., Siddhuraju, P., & Becker, K. (2007). Plant secondary metabolites. Humana Press.
- Manan, D. M. A., Nadiah, W. A. W., & Ng, Y. M. (2012). Changes in the concentration of carbohydrates, organic acids and amino acids of soaked peanuts with lactic acid bacteria.
- Mancilla-Margalli, N. A., & López, M. G. (2002). Generation of Maillard compounds from inulin during the thermal processing of Agave tequilana Weber Var. azul. *Journal of Agricultural and Food Chemistry*, 50(4), 806-812.
- Marisa, M. W. (2006). Ascorbic acid and mineral composition of longan (*Dimocarpus longan*), lychee (*Litchi chinensis*) and rambutan (*Nephelium lappaceum*) cultivars grown in Hawaii. *Journal of Food Composition & Analysis*, 19, 655-663.
- Martins, S., Mussatto, S. I., Martínez-Avila, G., Montañez-Saenz, J., Aguilar, C. N., & Teixeira, J. A. (2011). Bioactive phenolic compounds: production and extraction by solid-state fermentation. A review. *Biotechnology Advances*, 29(3), 365-373.
- Maskan, A., Kaya, S., & Maskan, M. (2002). Effect of concentration and drying processes on color change of grape juice and leather (*pestil*). *Journal of Food Engineering*, 54(1), 75-80.
- Maskan, M. (2001). Kinetics of colour change of kiwifruits during hot air and microwave drying. *Journal of Food Engineering*, 48(2), 169-175.
- McBurney, D. H. (1978). Psychological dimensions and perceptual analyses of taste. *Handbook of Perception*. Vol. VIA: Tasting and Smelling. EC Carterette and MP Freidman, editors. Academic Press, Inc., New York, 125-155.
- Medoua, G. N., Mbome, I. L., Agbor-Egbe, T., & Mbofung, C. M. F. (2008). Influence of fermentation on some quality characteristics of trifoliate yam (*Dioscorea dumetorum*) hardened tubers. *Food Chemistry*, 107, 1180-1186.
- Meibach, R. L., Akashe, A., Haas, G. W., & West, L. G. (2006). *U.S. Patent No. 7,097,874*. Washington, DC: U.S. Patent and Trademark Office.
- Meyer, B., Biehl, B., Said, M. B., & Samarakoddy, R. J. (1989). Post-harvest pod storage: A method for pulp preconditioning to impair strong nib acidification during cocoa fermentation in Malaysia. *Journal of the Science of Food and Agriculture*, 48(3), 285-304.
- Miliauskas, G., Venskutonis, P. R., & Van Beek, T. A. (2004). Screening of radical scavenging activity of some medicinal and aromatic plant extracts. *Food Chemistry*, 85, 231-237.
- Min, D. B., & Smouse, T. H. (Eds.). (1985). Flavor chemistry of fats and oils (Vol. 15). *The American Oil Chemists Society*.
- Minifie, B. W. (1989). *Chocolate, cocoa and confectionery: Science and Technology*. Springer. Berlin, Germany
- Mohibbe Azam, M., Waris, A., & Nahar, N. M. (2005). Prospects and potential of fatty acid methyl esters of some non-traditional seed oils for use as biodiesel in India. *Biomass and Bioenergy*, 29(4), 293-302.
- Møller, B. L., & Seigler, D. S. (1999). Biosynthesis of cyanogenic glucosides, cyanolipids, and related compounds. *Plant Amino Acids, Biochemistry and Biotechnology*. Marcel Dekker, New York, 563-609.

- Montano, A., Sanchez, A. H., Lopez-Lopez, A., De Castro, A., & Rejano, L. (2010). Chemical composition of fermented green olives: acidity, salt, moisture, fat, protein, ash, fiber, sugar and polyphenol. *Olives and Olive Oil in Health and Disease*, 291-296.
- Montgomery, D. C. (2013). *Applied Statistics and Probability for Engineers* 6th edition. Wiley.
- Morton, J. F. (1987). *Fruits of warm climates*. Miami, USA, JF Morton.
- Mousavi, Z. E., Mousavi, S. M., Razavi, S. H., Hadinejad, M., Emam-Djomeh, Z., & Mirzapour, M. (2013). Effect of Fermentation of Pomegranate Juice by *Lactobacillus plantarum* and *Lactobacillus acidophilus* on the Antioxidant Activity and Metabolism of Sugars, Organic Acids and Phenolic Compounds. *Food Biotechnology*, 27, 1-13.
- Mubarak, A. E. (2005). Nutritional composition and antinutritional factors of mung bean seeds (*Phaseolus aureus*) as affected by some home traditional processes. *Food Chemistry*, 89(4), 489-495.
- Mulimani, V. H., & Supriya, D. (1994). Tannic acid content in sorghum (*Sorghum bicolor* M.): Effects of processing. *Plant Foods for Human Nutrition*, 46(3), 195-200.
- Murray, N. J., & Williamson, M. P. (1994). Conformational study of a salivary proline-rich protein repeat sequence. *European Journal of Biochemistry*, 219(3), 915-921.
- Murray, N. J., Williamson, M. P., Lilley, T. H., & Haslam, E. (1994). Study of the interaction between salivary proline-rich proteins and a polyphenol by 1H-NMR spectroscopy. *European Journal of Biochemistry*, 219(3), 923-935.
- Nahrstedt, A. (1985). Cyanogenic compounds as protecting agents for organisms. *Plant Systematics and Evolution*, 150(1-2), 35-47.
- Nakamura, T., Tanigake, A., Miyanaga, Y., Ogawa, T., Akiyoshi, T., Matsuyama, K., & Uchida, T. (2002). The effect of various substances on the suppression of the bitterness of quinine-human gustatory sensation, binding, and taste sensor studies. *Chemical and Pharmaceutical Bulletin*, 50(12), 1589-1593.
- Nazaruddin, R., Seng, L. K., Hassan, O., & Said, M. (2006). Effect of pulp preconditioning on the content of polyphenols in cocoa beans (*Theobroma Cacao*) during fermentation. *Industrial Crops and Products*, 24(1), 87-94.
- Nebesny, E., & Rutkowski, J. (1998). Effect of roasting and secondary fermentation on cocoa bean enrichment. *Polish Journal of Food and Nutrition Sciences*, 7(3), 437-445.
- Nelson, S. L. (1998). *U.S. Patent No. 5,766,622*. Washington, DC: U.S. Patent and Trademark Office.
- Noble, A. C. (1994). Bitterness in wine. *Physiology & behavior*, 56(6), 1251-1255.
- Norlia, M. I., Roshazita, C. A., Nuraiti, T. I. T., Salwa, M. Z. M., & Fatimah, M. S. S. (2011). Preparation and Characterisation of Activated Carbon from Rambutan Seed (*Nephelium Lappaceum*) by Chemical Activation. *Empowering Science, Technology and Innovation Towards a Better Tomorrow*, 184-191.
- Notka, F., Meier, G., & Wagner, R. (2004). Concerted inhibitory activities of *Phyllanthus amarus* on HIV replication in vitro and ex vivo. *Antiviral Research*, 64(2), 93-102.
- Obizoba, I. C., & Atii, J. V. (1991). Effect of soaking, sprouting, fermentation and cooking on nutrient composition and some anti-nutritional factors of sorghum (*Guinesia*) seeds. *Plant Foods for Human Nutrition*, 41(3), 203-212.



- Oboh, G., Raddatz, H., & Henle, T. (2008). Antioxidant properties of polar and non-polar extracts of some tropical green leafy vegetables. *Journal of the Science of Food and Agriculture*, 88(14), 2486-2492.
- Ohene, Afoakwa, E., & Alistair, P. (2010). Cocoa Fermentation: Chocolate Flavor Quality. Available from <http://ugspace.ug.edu.gh/handle/123456789/2521>
- Okabe, S., Suganuma, M., Imayoshi, Y., Taniguchi, S., Yoshida, T., & Fujiki, H. (2001). Osman, M. A. (2007). Effect of different processing methods, on nutrient composition, antinutritional factors, and in vitro protein digestibility of Dolichos lablab bean (*Lablab purpureus* (L) Sweet). *Pakistan Journal of Nutrition*, 6(4), 299-303.
- Okuda, T., Yoshida, T., & Hatano, T. (1995). Hydrolyzable tannins and related polyphenols. In Fortschritte der Chemie organischer Naturstoffe/Progress in the Chemistry of Organic Natural Products (pp. 1-117). Springer Vienna.
- Omafuvbe, B. O., Shonukan, O. O., & Abiose, S. H. (2000). Microbiological and biochemical changes in the traditional fermentation of soybean for 'soy-daddawa' Nigerian food condiment. *Food Microbiology*, 17(5), 469-474.
- Ong, P. K., Acree, T. E., & Lavin, E. H. (1998). Characterization of volatiles in rambutan fruit (*Nephelium lappaceum* L.). *Journal of Agricultural and Food Chemistry*, 46(2), 611-615.
- Onimawo, I. A., Oteno, F., Orokpo, G., & Akubor, P. I. (2003). Physicochemical and nutrient evaluation of African bush mango (*Irvingia gabonensis*) seeds and pulp. *Plant Foods for Human Nutrition*, 58(3), 1-6.
- Oreopoulou, V., & Russ, W. (Eds.). (2007). Utilization of by-products and treatment of waste in the food industry. New York: Springer.
- Osman, H., Nasarudin, R., & Lee, S. L. (2004). Extracts of cocoa (*Theobroma cacao* L.) leaves and their antioxidation potential. *Food Chemistry*, 86(1), 41-46.
- Osman, M. A. (2007). Effect of different processing methods, on nutrient composition, antinutritional factors, and in vitro protein digestibility of Dolichos lablab bean (*Lablab purpureus* (L) Sweet). *Pakistan Journal of Nutrition*, 6(4), 299-303.
- Osorio, C., Franco, M. S., Castaño, M. P., González-Miret, M. L., Heredia, F. J., & Morales, A. L. (2007). Colour and flavour changes during osmotic dehydration of fruits. *Innovative Food Science & Emerging Technologies*, 8(3), 353-359.
- Oszmiański, J., Wolniak, M., Woidyło, A., & Wawer, I. (2008). Influence of apple purée preparation and storage on polyphenol contents and antioxidant activity. *Food Chemistry*, 107(4), 1473-1484.
- Ouattara, H.G., Koffi, B.L., Karou, G.T., Sangare A, Niamke, S.L., Diopoh, J.K. (2008). Implication of Bacillus spp. in the production of pectinolytic enzymes during cocoa fermentation. *World Journal of Microbiology and Biotechnology*. 24, 1753–1760.
- Owusu, M., Petersen, M. A., & Heimdal, H. (2012). Effect of fermentation method, roasting and conching conditions on the aroma volatiles of dark chocolate. *Journal of Food Processing and Preservation*, 36(5), 446-456.
- Özdemir, M., & Devres, O. (2000). Kinetics of color changes of hazelnuts during roasting. *Journal of Food Engineering*, 44(1), 31-38.
- Palanisamy, U., Cheng, H. M., Masilamani, T., Subramaniam, T., Ling, L. T., & Radhakrishnan, A. K. (2008). Rind of the rambutan, *Nephelium lappaceum*, a potential source of natural antioxidants. *Food Chemistry*, 109(1), 54-63.
- Pangborn, R. M., & Dunklev, W. L. (1964). Difference-Preference Evaluation of Milk by Trained Judge's. *Journal of Dairy Science*, 47(12), 1414-1416.



- Paredes-López, O., Harry, G. I., & Murray, E. D. (1988). Food biotechnology review: Traditional solid-state fermentations of plant raw materials—application, nutritional significance, and future prospects. *Critical Reviews in Food Science & Nutrition*, 27(3), 159-187.
- Pérez, A. G., Sanz, C., Olías, R., & Olías, J. M. (1999). Lipoxygenase and hydroperoxide lyase activities in ripening strawberry fruits. *Journal of Agricultural and Food Chemistry*, 47(1), 249-253.
- Perren, R., & Escher, F. E. (1997). Investigations on the hot air roasting of nuts. *Manufacturing Confectioner*, 77, 123-127.
- Peterson, J., & Dwyer, J. (1998). Flavonoids: dietary occurrence and biochemical activity. *Nutrition Research*, 18(12), 1995-2018.
- Pettipher, G.L. (1986). An improved method for the extraction and quantification of anthocyanin in cocoa beans and its use as an index of the degree of fermentation. *Journal of the Science of Food and Agriculture*, 37, 289-296.
- Piga, A., Del Caro, A., & Corda, G. (2003). From plums to prunes: influence of drying parameters on polyphenols and antioxidant activity. *Journal of Agricultural and Food Chemistry*, 51(12), 3675-3681.
- Pittia, P., Dalla Rosa, M., & Lerici, C. R. (2001). Textural changes of coffee beans as affected by roasting conditions. *LWT-Food Science and Technology*, 34(3), 168-175.
- Porres, J. M., Urbano, G., Fernández-Figares, I., Prieto, C., Perez, L., & Aguilera, J. F. (2002). Digestive utilisation of protein and amino acids from raw and heated lentils by growing rats. *Journal of the Science of Food and Agriculture*, 82(14), 1740-1747.
- Price, M. L., Butler, L. G., Rogler, J. C., & Featherston, W. R. (1979). Overcoming the nutritionally harmful effects of tannin in sorghum grain by treatment with inexpensive chemicals. *Journal of Agricultural and Food Chemistry*, 27(2), 441-445.
- Prodanov, M., Garrido, I., Vacas, V., Lebrón-Aguilar, R., Dueñas, M., Gómez-Cordovés, C., & Bartolomé, B. (2008). Ultrafiltration as alternative purification procedure for the characterization of low and high molecular-mass phenolics from almond skins. *Analytica Chimica Acta*, 609(2), 241-251.
- Pujolà, M., Farreras, A., & Casañas, F. (2007). Protein and starch content of raw, soaked and cooked beans (*Phaseolus vulgaris* L.). *Food Chemistry*, 102(4), 1034-1041.
- Puziah, H., Jinap, S., Muhammad, K. & Ali, A. (1998). Changes in free amino acid, peptide- N, sugar and pyrazine concentration during cocoa fermentation. *Journal of the Science of Food and Agriculture*, 78(4), 535-542.
- Quesnel, V. C., & Jugmohunsingh, K. (1970). Browning reaction in drying cacao. *Journal of the Science of Food and Agriculture*, 21(10), 537-541.
- Ragasa, C. Y., De Luna, R. D., Cruz, W. C., & Rideout, J. A. (2005). Monoterpene Lactones from the Seeds of *Nephelium lappaceum*. *Journal of natural products*, 68(9), 1394-1396.
- Rahma, E. H., & Rao, M. N. (1984). Effect of debittering treatment on the composition and protein components of lupine seed (*Lupinus termis*) flour. *Journal of Agricultural and Food Chemistry*, 32(5), 1026-1030.
- Rangkadilok, N., Worasuttayangkurn, L., Bennett, R.N., Satayavivad, J., (2005) Identification and quantification of polyphenolic compounds in longan (*Euphoria longana* Lam.) fruit. *Journal of Agricultural and Food Chemistry*, 53, 1387-1392.

- Rao, P. U., & Deosthale, Y. G. (1982). Tannin content of pulses: varietal differences and effects of germination and cooking. *Journal of the Science of Food and Agriculture*, 33(10), 1013-1016.
- Raoult-Wack, A. L., Guilbert, S., Maguer, M. L., & Rios, G. (1991). Simultaneous water and solute transport in shrinking media-part 1: application to dewatering and impregnation soaking process analysis (osmotic dehydration). *Drying Technology*, 9(3), 589-612.
- Rastogi, N. K., Raghavarao, K. S. M. S., & Niranjani, K. (1997). Mass transfer during osmotic dehydration of banana: Fickian diffusion in cylindrical configuration. *Journal of Food Engineering*, 31(4), 423-432.
- Reddy, N. R., & Pierson, M. D. (1994). Reduction in anti-nutritional and toxic components in plant foods by fermentation. *Food Research International*, 27, 281-290.
- Reddy, N. R.; Salunkhe, D. K. Fermentation. In Handbook of World Food Legumes: *Nutritional Chemistry, Processing Technology*,
- Redgwell, R. J., Trovato, V., & Curti, D. (2003). Cocoa bean carbohydrates: roasting-induced changes and polymer interactions. *Food Chemistry*, 80(4), 511-516.
- Renard, C. M. (2005). Effects of conventional boiling on the polyphenols and cell walls of pears. *Journal of the Science of Food and Agriculture*, 85(2), 310-318.
- Review (10/18/06). [http://www.nass.usda.gov/Statistics\\_by\\_State/California/](http://www.nass.usda.gov/Statistics_by_State/California/)
- Riboli, E., & Norat, T. (2003). Epidemiologic evidence of the protective effect of fruit and vegetables on cancer risk. *The American Journal of Clinical Nutrition*, 78(3), 559S-569S.
- Rocha, A. M. C. N., & Morais, A. M. (2001). Influence of controlled atmosphere storage on polyphenoloxidase activity in relation to colour changes of minimally processed 'Jonagored' apple. *International journal of food science & technology*, 36(4), 425-432.
- Rodriguez-Campos, J., Escalona-Buendía, H. B., Orozco-Avila, I., Lugo-Cervantes, E., & Jaramillo-Flores, M. E. (2011). Dynamics of volatile and non-volatile compounds in cocoa (*Theobroma cacao L.*) during fermentation and drying processes using principal components analysis. *Food Research International*, 44(1), 250-258.
- Rodríguez-Gómez, F., Bautista-Gallego, J., Romero-Gil, V., Arroyo-López, F. N., Garrido-Fernández, A., & García-García, P. (2012). Effects of salt mixtures on Spanish green table olive fermentation performance. *LWT-Food Science and Technology*, 46(1), 56-63.
- Rouseff, R. L. (1990). Bitterness in food products: an overview. *Developments in food science*, 25, 1-14.
- Rozan, P., Kuo, Y.H., & Lambein, F. (2000). Free amino acids present in commercially available seedlings sold for human consumption. A potential hazard for consumers. *Journal of Agricultural and Food Chemistry*, 48, 716-723.
- Rufián-Henares, J. A., & Morales, F. J. (2007). Functional properties of melanoidins: In vitro antioxidant, antimicrobial and antihypertensive activities. *Food Research International*, 40(8), 995-1002.
- Sabatini, N., & Marsilio, V. (2008). Volatile compounds in table olives (*Olea Europaea L.*, *Nocellara del Belice cultivar*). *Food chemistry*, 107(4), 1522-1528.

- Saharan, K., Khetarpaul, N., & Bishnoi, S. (2002). Antinutrients and protein digestibility of Faba bean and Rice bean as affected by soaking, dehulling and germination. *Journal of Food Science and Technology*, 39(4), 418–422
- Saklar, S., Katnas, S., & Urgan, S. (2001). Determination of optimum hazelnut roasting conditions. *International Journal of Food Science & Technology*, 36(3), 271-281.
- Saklar, S., Urgan, S., & Katnas, S. (1999). Instrumental crispness and crunchiness of roasted hazelnuts and correlations with sensory assessment. *Journal of Food Science*, 64(6), 1015-1019.
- Salakpetch, S. (2003). Rambutan production in Thailand. In *II International Symposium on Lychee, Longan, Rambutan and other Sapindaceae Plants* 665 (pp. 67-72).
- Satwadhar, P. N., Kadam, S. S. and Salunkhe, D. K. (1981) Effect of germination and cooking on polyphenols and in vitro protein digestibility of horse gram and moth bean. *Quality Plant Foods Hum. Nutrition* 36, 71-76.
- Saurel, R., Raoult-Wack, A. L., Rios, G., & Guilbert, S. (1994). Mass transfer phenomena during osmotic dehydration of apple I. Fresh plant tissue. *International Journal of Food Science & Technology*, 29(5), 531-542.
- Savage, G. P., McNeil, D. L., & Dutta, P. C. (1997). Lipid composition and oxidative stability of oils in hazelnuts (*Corylus avellana L.*) grown in New Zealand. *Journal of the American Oil Chemists' Society*, 74(6), 755-759.
- Savaş, E., & Uylaser, V. (2013). Quality improvement of green table olive cv.'Domat'(*Olea europaea L.*) grown in Turkey using different de-bittering methods. *Notulae Botanicae, Horti Agrobotanici, Cluj-Napoca*, 41(1), 269-275.
- Saxena, A. K., Chadha, M., & Sharma, S. (2003). Nutrients and antinutrients in chickpea (*Cicer arietinum L.*) cultivars after soaking and pressure cooking. *Journal of Food Science and Technology*, 40(5), 493-497.
- Scalbert, A. (1991). Antimicrobial properties of tannins. *Phytochemistry*, 30(12), 3875-3883.
- Schiffman, S. S., Booth, B. J., Losee, M. L., Pecore, S. D., & Warwick, Z. S. (1995). Bitterness of sweeteners as a function of concentration. *Brain Research Bulletin*, 36(5), 505-513.
- Schiffman, S. S., & Dackis, C. (1975). Taste of nutrients: amino acids, vitamins, and fatty acids. *Perception & Psychophysics*, 17(2), 140-146.
- Schwan, R. F., & Wheals, A. E. (2004). The microbiology of cocoa fermentation and its role in chocolate quality. *Critical Reviews in Food Science and Nutrition*, 44(4), 205-221.
- Severini, C., Gomes, T., De Pilli, T., Romani, S., & Massini, R. (2000). Autoxidation of packed almonds as affected by Maillard reaction volatile compounds derived from roasting. *Journal of Agricultural and Food Chemistry*, 48(10), 4635-4640.
- Shakerardekani, A. (2011). Effect of roasting conditions on hardness, moisture content and colour of pistachio kernels. *International Food Research Journal*, (18), 704-710.
- Shakerardekani, A., Karim, R., Ghazali, H. M., & Chin, N. L. (2013). Textural, rheological and sensory properties and oxidative stability of nut spreads: a review. *International journal of molecular sciences*, 14(2), 4223-4241.
- Shi, J., & Le Maguer, M. (2002). Osmotic dehydration of foods: mass transfer and modeling aspects. *Food Reviews International*, 18(4), 305-335.

- Shibuva, M., Hoshino, M., Katsube, Y., Havashi, H., Kushiro, T., & Ebizuka, Y. (2006). Identification of  $\beta$ -amyrin and sophoradiol 24-hydroxylase by expressed sequence tag mining and functional expression assay. *FEBS journal*, 273(5), 948-959.
- Silva, F. M., & Silva, C. L. (1999). Colour changes in thermally processed cupuacu (*Theobroma grandiflorum*) puree: critical times and kinetics modelling. *International journal of food science & technology*, 34(1), 87-94.
- Singh, B., Panesar, P. S., & Nanda, V. (2008). Optimization of osmotic dehydration process of carrot cubes in sucrose solution. *Journal of Food Process Engineering*, 31(1), 1-20.
- Singh, R. P., Chidambara Murthy, K. N., & Jayaprakasha, G. K. (2002). Studies on the antioxidant activity of pomegranate (*Punica granatum*) peel and seed extracts using in vitro models. *Journal of Agricultural and Food Chemistry*, 50, 81-86.
- Singh, R.P., Murthy, K.N.C., Jayaprakasha, G.K., (2002). Studies on the antioxidant activity of pomegranate (*Punica granatum*) peel and seed extracts using in vitro models. *J. Agric. Food Chem.* 50, 81–86.
- Singh, U. (1988). Antinutritional factors of chickpea and pigeonpea and their removal by processing. *Plant Foods for Human Nutrition*, 38(3), 251-261.
- Singleton, V. L. (1992). Tannins and the qualities of wines. In *Plant Polyphenols* (pp. 859-880). Springer US.
- Sirisompong, W., Jirapakkul, W., & Klinkesorn, U. (2011). Response surface optimization and characteristics of rambutan (*Nephelium lappaceum* L.) kernel fat by hexane extraction. *LWT-Food Science and Technology*, 44(9), 1946-1951.
- Skrede, G. (1985). Color quality of blackcurrant syrups during storage evaluated by hunter L', a', b' values. *Journal of Food Science*, 50(2), 514-517.
- Smith, A. K., June, H., & Noble, A. C. (1996). Effects of viscosity on the bitterness and astringency of grape seed tannin. *Food Quality and Preference*, 7(3), 161-166.
- Solis-Fuentes, J. A., Camey-Ortiz, G., Hernández-Medel, M. D. R., Pérez-Mendoza, F., & Durán-de-Bazúa, C. (2010). Composition, phase behavior and thermal stability of natural edible fat from rambutan (*Nephelium lappaceum* L.) seed. *Bioresource Technology*, 101(2), 799-803.
- Solms, J. (1969). Taste of amino acids, peptides, and proteins. *Journal of Agricultural and Food Chemistry*, 17(4), 686-688.
- Steingarten, J. (2011). *The Man Who Ate Everything*. Random House LLC.
- Stephan, A., & Steinhart, H. (2000). Bitter taste of unsaturated free fatty acids in emulsions: contribution to the off-flavour of soybean lecithins. *European Food Research and Technology*, 212(1), 17-25.
- Štoudková, E. V. B. L. H., & Zemanová, J. (2007). Application of SPME-GC method for analysis of the aroma of white surface mould cheeses. *Journal of Food and Nutrition Research*, 46(2), 84-90.
- Su, G., Zheng, L., Cui, C., Yang, B., Ren, J., & Zhao, M. (2011). Characterization of antioxidant activity and volatile compounds of Maillard reaction products derived from different peptide fractions of peanut hydrolysate. *Food Research International*, 44(10), 3250-3258.



- Suzuki, H., Kaiimoto, Y., & Kumagai, H. (2002). Improvement of the bitter taste of amino acids through the transpeptidation reaction of bacterial  $\gamma$ -glutamyltranspeptidase. *Journal of Agricultural and Food Chemistry*, 50(2), 313-318.
- Takei, Y., & Yamanishi, T. (1974). Flavor components of roasted almond. *Agricultural and Biological Chemistry* (Japan).
- Tang, T. S., & Teoh, P. K. (1985). Palm kernel oil extraction-The Malaysian experience. *Journal of the American Oil Chemists' Society*, 62(2), 254-258.
- Taylor, A. J., & Roozen, J. P. (1996). Volatile flavor release from foods during eating. *Critical Reviews in Food Science & Nutrition*, 36(8), 765-784.
- Teguia, A., & Fru, S. F. (2007). The growth performances of broiler chickens as affected by diets containing common bean (*Phaseolus vulgaris*) treated by different methods. *Tropical Animal Health and Production*, 39(6), 405-410.
- Thitilertdecha, N., & Rakariyatham, N. (2011). Phenolic content and free radical scavenging activities in rambutan during fruit maturation. *Scientia Horticulturae*, 129(2), 247-252.
- Thitilertdecha, N., Teerawutgulrag, A., & Rakariyatham, N. (2008). Antioxidant and antibacterial activities of *Nephelium lappaceum* L. extracts. *LWT-Food Science and Technology*, 41(10), 2029-2035.
- Thitilertdecha, N., Teerawutgulrag, A., Kilburn, J. D., & Rakariyatham, N. (2010). Identification of major phenolic compounds from *Nephelium lappaceum* L. and their antioxidant activities. *Molecules*, 15(3), 1453-1465.
- Thompson, S.S., Miller, K.B., Lopez, A.S. (2007). Cocoa and coffee. In: Doyle MP, Beuchat LR, Montville TJ, editors. *Food Microbiology Fundamentals and Frontiers*. Washington.
- Timbie, D. J., Sechrist, L., & Keeney, P. G. (1978). Application of high-pressure liquid chromatography to the study of variables affecting theobromine and caffeine concentrations in cocoa beans. *Journal of Food Science*, 43(2), 560-565.
- Tindall, H. D. (1994). *Rambutan cultivation* (No. 121). Food & Agriculture Org. Tropical Crops Division. <http://www.fas.usda.gov/htp/horticulture/nuts>.
- Toelstede, S., & Hofmann, T. (2008). Sensomics mapping and identification of the key bitter metabolites in Gouda cheese. *Journal of Agricultural and Food Chemistry*, 56(8), 2795-2804.
- Toledo, T. C. F. D., & Canniatti-Brazaca, S. G. (2008). Chemical and nutritional evaluation of Carioca beans (*Phaseolus vulgaris* L.) cooked by different methods. *Food Science and Technology (Campinas)*, 28(2), 355-360.
- Troszyńska, A. (2004). Non-nutrient bioactive substances in food of plant origin causing bitterness and astringency. *Polish Journal of Food and Nutrition Sciences*, 13(Spec. Issue 1).
- Tunçel, G., Nout, M. J. R., Brimer, L., & Goektan, D. (1990). Toxicological, nutritional and microbiological evaluation of tempe fermentation with *Rhizopus oligosporus* of bitter and sweet apricot seeds. *International Journal of Food Microbiology*, 11(3), 337-344.
- Uylaser, V., Tamer, C. E., İncedayı, B., Vural, H., & Çopur, Ö. U. (2008). The quantitative analysis of some quality criteria of Gemlik variety olives. *Journal of Food, Agriculture & Environment*, 6(3&4), 26-30.
- Valero, E., Millán, C., Ortega, J. M., & Mauricio, J. C. (2003). Concentration of amino acids in wine after the end of fermentation by *Saccharomyces cerevisiae* strains. *Journal of the Science of Food and Agriculture*, 83(8), 830-835.

- Van Boekel, M. A. J. S. (2001). Kinetic aspects of the Maillard reaction: a critical review. *Food/Nahrung*, 45(3), 150-159.
- Vanzani, P., Rossetto, M., Rigo, A., Vrhovsek, U., Mattivi, F., D'Amato, E., & Scarpa, M. (2005). Major phytochemicals in apple cultivars: contribution to peroxy radical trapping efficiency. *Journal of Agricultural and Food Chemistry*, 53, 3377-3382.
- Vázquez-Araújo, L., Enguix, L., Verdú, A., García-García, E., & Carbonell-Barrachina, A. A. (2008). Investigation of aromatic compounds in toasted almonds used for the manufacture of turrón. *European Food Research and Technology*, 227(1), 243-254.
- Venkatachalam, M., & Sathe, S. K. (2006). Chemical composition of selected edible nut seeds. *Journal of Agricultural and Food Chemistry*, 54(13), 4705-4714.
- Vijayakumari, K., Siddhuraju, P. and Janardhanan, K. (1995) Effects of various water or hydrothermal treatments on certain antinutritional compounds in the seeds of the tribal pulse, *Dolichos lablab* var. *vulgaris* L. *Plant Foods Hum. Nutrition* 48, 17-29.
- Vijayakumari, K., Siddhuraju, P., Pugalenti, M., & Janardhanan, K. (1998). Effect of soaking and heat processing on the levels of antinutrients and digestible proteins in seeds of *Vigna aconitifolia* and *Vigna sinensis*. *Food Chemistry*, 63(2), 259-264.
- Vinci, G., Botrè, F., Mele, G., & Ruggieri, G. (1995). Ascorbic acid in exotic fruits: a liquid chromatographic investigation. *Food Chemistry*, 53(2), 211-214.
- Vranová, J., & Ciesarová, Z. (2009). Furan in food—a review. *Czech J Food Sci*, 27(1), 1-10.
- Voigt, J., Biehl, B., Heinrichs, H., Kamaruddin, S., Marsoner, G. G., & Hugl, A. (1994). In-vitro formation of cocoa-specific aroma precursors: aroma-related peptides generated from cocoa-seed protein by co-operation of an aspartic endo-protease and a carboxypeptidase. *Food Chemistry*, 49, 173-180.
- Wall, M. M. (2006). Ascorbic acid and mineral composition of longan (*Dimocarpus longan*), lychee (*Litchi chinensis*) and rambutan (*Nephelium lappaceum*) cultivars grown in Hawaii. *Journal of Food Composition and Analysis*, 19(6), 655-663.
- Watson, B. J. (1988). Rambutan cultivars in north Queensland. *Queensland Agricultural Journal*, 1, 37-42.
- Watson, B.J., Cunningham, I.C., Walduck, G.D., Wait, A.I. and Goebel, R.L., (1988). Rambutan culture north Queensland. Queensland Department of Primary Industries, AGDEX 238/00, 21 pp.
- Wehrli, C. (2008). US20120232163 A1. Washington, DC: U.S.
- Wijekoon, M. M., Bhat, R., & Karim, A. A. (2011). Effect of extraction solvents on the phenolic compounds and antioxidant activities of bunga kantan (*Etilingera elatior* Jack.) inflorescence. *Journal of Food Composition and Analysis*, 24(4), 615-619.
- Wirthensohn, M. G., Chin, W. L., Franks, T. K., Baldock, G., Ford, C. M., & Sedgley, M. (2008). Characterising the flavour phenotypes of almond (*Prunus dulcis* Mill.) kernels. *Journal of Horticultural Science and Biotechnology*, 83(4), 462-468.
- Wollgast, J., & Anklam, E. (2000). Review on polyphenols in *Theobroma cacao*: changes in composition during the manufacture of chocolate and methodology for identification and quantification. *Food Research International*, 33(6), 423-447.

- Wu, Y. V., & Wall, J. S. (1980). Lysine content of protein increased by germination of normal and high-lysine sorghums. *Journal of agricultural and food chemistry*, 28(2), 455-458.
- Xu, B., & Chang, S. K. (2008). Effect of soaking, boiling, and steaming on total phenolic content and antioxidant activities of cool season food legumes. *Food Chemistry*, 110(1), 1-13.
- Xu, B., & Chang, S. K. (2009). Phytochemical profiles and health-promoting effects of cool-season food legumes as influenced by thermal processing. *Journal of Agricultural and Food Chemistry*, 57, 10718-10731.
- Yadav, A. K., & Singh, S. V. (2012). Osmotic dehydration of fruits and vegetables: a review. *Journal of Food Science and Technology*, 1-20.
- Yakubu, N., Amuzat, A. O., & Hamza, R. U. (2012). Effect of processing methods on the nutritional contents of bitter leaf (*Vernonia amygdalina*). *Am. J. Food. Nutr.*, 2(1), 26-30.
- Yang, C. M., Cheng, H. Y., Lin, T. C., Chiang, L. C., & Lin, C. C. (2007). The in vitro activity of geraniin and 1, 3, 4, 6-tetra-O-galloyl- $\beta$ -D-glucose isolated from *Phyllanthus urinaria* against herpes simplex virus type 1 and type 2 infection. *Journal of Ethnopharmacology*, 110(3), 555-558.
- Zee, F., 1993. Rambutan and pili nuts: potential crops for Hawaii. In: Janick, J., Simon, M. J. E. (Eds.), *New Crops*. J. Wiley and Sons Inc., New York, pp. 461-465.
- Zhao, Y., & Xie, J. (2004). Practical applications of vacuum impregnation in fruit and vegetable processing. *Trends in Food Science & Technology*, 15(9), 434-451.
- Zheng, G., Xu, L., Wu, P., Xie, H., Jiang, Y., Chen, F., & Wei, X. (2009). Polyphenols from longan seeds and their radical-scavenging activity. *Food Chemistry*, 116, 433-436.
- Ziena, H. M. S. (1989). Hard-to-cook phenomenon in relation to physical, chemical and biological properties of faba beans (*Vicia Faba L.*) (Doctoral dissertation, PhD Thesis, Faculty of Agriculture, Alexandria University, Alexandria, Egypt).
- Zumbé, A. (1998). Polyphenols in cocoa: are there health benefits. *Nutrition Bulletin*, 23(1), 94-102.



## LIST OF PUBLICATIONS

- Mehdizadeh, S., Lasekan, O., Kharidah, M., Badlishah, S. (2014). Variability in the fermentation index, polyphenols and amino acids of seeds of rambutan (*Nephelium lappaceum* L.) during fermentation. *Journal of Food Composition and Analysis* (in press).
- Lasekan,O.,& Mehdizadeh, S. (2013). Effect of Traditional Fermentation as a Pretreatment to Decrease the Antinutritional Properties of Rambutan Seed (*Nephelium lappaceum* L.). *International Proceedings of Chemical, Biological & Environmental Engineering*, 55.





UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : 2014-2015

TITLE OF THESIS / PROJECT REPORT :

Effects of fermentation and debittering methods on  
production of almond-like *Sambutan* seed  
(*Cnephelium lappaceum* L.) snack

NAME OF STUDENT : Shabnam Mehdizadeh

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

\*Please tick (v )

**CONFIDENTIAL**

(Contain confidential information under Official Secret Act 1972).

**RESTRICTED**

(Contains restricted information as specified by the organization/institution where research was done).

**OPEN ACCESS**

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

**PATENT**

Embargo from \_\_\_\_\_ until \_\_\_\_\_  
(date) (date)

(Signature of Student)  
New IC No/ Passport No.:

Date : 20/3/2015

**Approved by:**

\_\_\_\_\_  
(Signature of Chairman of Supervisory Committee)  
Name:

Date :

[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted. ]