



UNIVERSITI PUTRA MALAYSIA

***EFFECTS OF SILENCING DISHEVELLED 1, 2 and 3 mRNA
EXPRESSION ON MDA-MB-231 CELL MIGRATION AND INVASION
USING siRNAs***

SAMUEL KHOO LEON JUAN

IB 2015 38



**EFFECTS OF SILENCING DISHEVELLED 1, 2 AND 3 mRNA
EXPRESSION ON MDA-MB-231 CELL MIGRATION AND INVASION
USING siRNAs**

By

SAMUEL KHOO LEON JUAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Master of Science**

May 2015

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia

Copyright © Universiti Putra Malaysia



DEDICATION

Specially dedicated to,

My loving parents, my awesome friends, dear supervisors,

For their invaluable love, understanding, encouragement and patience.



COPYRIGHT

UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Degree of Master of Science

**EFFECTS OF SILENCING DISHEVELLED 1, 2 AND 3 mRNA
EXPRESSION ON MDA-MB-231 CELL MIGRATION AND INVASION
USING siRNAs**

By

SAMUEL KHOO LEON JUAN

May 2015

Chairman : Professor Seow Heng Fong, PhD
Institute : Bioscience

The Wnt signalling pathway plays an important role in embryonic development, generation of cell polarity and specification. Previous studies have shown that both canonical and non-canonical Wnt signaling pathways are involved in breast cancer metastasis. Dishevelled, one of the components of both canonical and non-canonical Wnt signaling pathway is involved in proliferation, polarity, terminal differentiation and self-renewal of cells. It is well known that Dishevelled interacting with Axin plays a role in dissociating the β -catenin destruction complex in the canonical Wnt signaling pathway. Although the involvement of Dishevelled in the non-canonical Wnt signaling pathway is less well studied compared to the canonical pathway, the role of Dishevelled is also important for cytoskeletal reorganization in planar cell polarity pathway and promoting tissue separation as well as ventral axis identity in *Xenopus* via the Wnt Calcium pathway. Our hypothesis is that silencing of human Dishevelled 1,2 and 3 isoforms results in inhibition of both canonical and non-canonical Wnt signaling pathway leading to reduced β -catenin interaction with TCF (T-cell Factor) and LEF (Lymphoid Enhancer Factor), reduced cell proliferation, migration and also invasion regardless of canonical or non-canonical Wnt signaling pathway. MDA-MB-231 is the breast cancer cell line used in this study. The objectives in this study include (1) to confirm the silencing of Dishevelled 1,2 and 3 by reverse-transcription quantitative PCR and Western Blotting. (2) to determine the changes in cell viability, migration, invasion, TCF-LEF mediated transcriptional activation after gene silencing using siRNA targeting Dishevelled isoforms. (3) to identify the changes in β -catenin, Glycogen Synthase Kinase-3 β (GSK3 β) protein, Epidermal Growth Factor Receptor (EGFR) and Extracellular signal-Regulated Kinases (ERK) expression in cells transfected with siRNAs targeting Dishevelled 1, 2 and 3. This study resulted in a few interesting findings: (1) A significant reduction of Dishevelled 1,2 and 3 mRNA levels in cells transfected with siRNA based on reverse transcription quantitative PCR was observed. However, we were not able to detect Dishevelled 1 isoform at the protein

level using the commercially available anti-Dishevelled 1 antibody; (2) different phenotypical changes were observed in cells transfected with siRNA targeting Dishevelled 1 and Dishevelled 2. This occurred with siRNA D1_9 or siRNA D2_4 where cells transfected with both siRNAs showed reduced cell viability. However, there was no significant change in cell viability, migration and invasion with other siRNAs for Dishevelled 1 and Dishevelled 2 siRNA (D1_6 and D2_8, respectively) as compared to control. Further studies showed that both siRNAs D1_9 and D2_4 resulted in reduced cell invasion and migration that were statistically significant. This indicates that there is possibility of target non-specificity by both siRNAs and for siRNA D2_4 or it could also be due to a higher degree of Dishevelled 2 knockdown. (3) Dishevelled 3 plays a role in mediating Wnt3a-stimulated TCF-LEF transcriptional activity. Reduced Wnt3a-stimulated TCF-LEF transcriptional activation was observed in MDA-MB-231 cell line transfected with siRNAs targeting Dishevelled 3 especially on cells transfected with siRNA D3_7 which was found to be statistically significant. (4) both densitometric results of phospho-GSK3 β - Serine 9 expression (relative to total GSK3 β) and phospho- β -catenin expression (relative to total β -catenin) for cells transfected with Dishevelled 3 siRNAs shows that a higher expression of phospho-GSK3 β -Serine 9, which is indicative of inactive GSK3 β , can result in reduced β -catenin degradation (phospho- β -catenin). However, the level of β -catenin degradation (relative to total β -catenin) expression in Dishevelled 3 siRNA transfected cells was comparable to Control. The basal phospho- β -catenin in Dishevelled 3 siRNA transfected cells was not related to GSK3 β activity in this study. (5) by relating both phospho- β -catenin expression (relative to total β -catenin) on densitometric analysis and Luciferase assay, reduced β -catenin degradation results in a higher TCF-LEF activity for cells transfected with siRNA targeting Dishevelled 1 and 2. However, reduced TCF/LEF mediated transcriptional activity in cells transfected with Dishevelled 3 siRNAs did not show an increase in phospho- β -catenin expression (relative to total β -catenin) as compared to control. Silencing of Dishevelled 3 expression maintains expression of basal phospho- β -catenin, which is sufficient for degradation or dissociation of β -catenin destruction complex resulting in the reduced TCF/LEF mediated transcriptional activity. It is possible that disruption of the interaction between Dishevelled 3 but not Dishevelled 1 and 2 with Axin interferes with formation of β -catenin destruction complex formation resulting in reduced TCF/LEF-sensitive transcriptional activity by Wnt3a in MDA-MB-231 cells or interference at the promoter binding site for β -catenin/ TCF-LEF complex. (6) densitometric analysis of Western Blot involving components along EGFR-ERK pathway shows that Wnt3a stimulation does not result in activation of EGFR-ERK pathway and knockdown of Dishevelled isoforms do not result in reduction of EGFR-ERK pathway under Wnt3a stimulation. In conclusion, Dishevelled 3 is involved in canonical Wnt β -catenin pathway in MDA-MB-231 breast cancer cells, playing a role in inhibiting proteosomal degradation of β -catenin. It is possible that knockdown of Dishevelled 3 expression disrupted the interaction with Axin, reduced β -catenin accumulation and ultimately reduced Wnt3a stimulated TCF/LEF-mediated transcriptional activity. It is possible that silencing of Dishevelled 3 expression interfered with β -catenin/TCF-LEF-mediated transcriptional activity. On the contrary, non-reduction in β -catenin/ TCF-LEF-mediated transcriptional activity and low phospho- β -catenin expression relative to total β -catenin in Dishevelled 1 and Dishevelled 2 siRNA transfected cells suggest that Dishevelled 1 and 2 isoforms are not involved in canonical Wnt β -catenin pathway on MDA-MB-231 cell line.

Abstrak tesis yang dikemukakan kepada Senat Univerisiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Master Sains

**KESAN PENYEPAPAN PENYATAAN mRNA DISHEVELLED 1,2 DAN 3
KE ATAS PENGHIJRAHAN DAN SERANGAN SEL MDA-MB-231 DENGAN
MENGUNAKAN siRNA**

Oleh

SAMUEL KHOO LEON JUAN

Mei 2015

Pengerusi : Profesor Seow Heng Fong, PhD
Instititute : Biosains

Isyarat laluan Wnt memainkan peranan penting dalam perkembangan embrio, penjanaan kekutuban sel dan juga penspesiesan sel. Kajian sebelum ini telah menunjukkan penglibatan isyarat laluan Wnt, sama ada yang berkanun atau isyarat laluan Wnt tidak berkanun dalam perebakan sel. Dishevelled, salah satu komponen yang terkandung dalam isyarat laluan Wnt berkanun dan tidak berkanun, memainkan peranan penting dalam pembahagian, kekutuban, pembezaan terminal dan pembaharuan sendiri dalam sel. Interaksi antara Dishevelled dan Axin memainkan peranan penting dalam pemisahan kompleks pemusnahan β -catenin dalam isyarat laluan Wnt berkanun. Walaupun kajian penglibatan Dishevelled dalam isyarat laluan Wnt tidak berkanun adalah lebih kurang berbanding dengan laluan berkanun, Dishevelled juga memainkan peranan yang penting bagi penyusunan semula perangkaan sel dalam laluan kekutuban satah sel, mendorong pemisahan tisu dan juga mendorong identiti paksi ventral dalam *Xenopus* melalui isyarat kalsium Wnt. Hipotesis kami adalah penyenyapan isoform Dishevelled 1,2 dan 3 mengakibatkan perencatan kedua-dua isyarat laluan Wnt berkanun dan tidak berkanun berikutan dengan pengurangan interaksi antara β -catenin dengan TCF dan LEF, pengurangan pembahagian sel, penghijrahan dan juga serangan dalam isyarat laluan Wnt berkanun dan tidak berkanun. Sel MDA-MB-231, modal sel kanser bagi kanser payu dara telah digunakan dalam kajian ini. Tujuan kajian ini termasuk (1) untuk mengesahkan kesan penyenyapan Dishevelled 1,2 dan 3 melalui reverse transkripsi PCR berkuantitatif dan Western Blot. (2) untuk menentukan perubahan dalam kesan kemandirian, penghijrahan, serangan, dan pengaktifan transkripsi yang diantarakan oleh TCF-LEF selepas penyenyapan dengan menggunakan siRNA. (3) Mengenal pasti perubahan β -catenin, Glycogen Synthase Kinase 3 β , Faktor Pertumbuhan Epiderma (EGFR) dan Pengawalan Isyarat Luar selular Kinase (ERK), dalam cell yang dijangkit oleh siRNA yang menyasar Dishevelled 1,2 dan 3. Terdapat beberapa keputusan yang menarik yang dalam kajian ini: (1) terdapat pengurangan yang ketara dalam aras mRNA Dishevelled 1,2 dan 3 berdasarkan keputusan reverse transkripsi quantitative PCR.

Namun, kami tidak dapat mengesan pernyataan isoform Dishevelled 1 dalam aras protein dengan menggunakan antibodi anti-Dishevelled 1. (2) terdapat kesan perubahan fenotipik yang dapat diperhatikan di antara siRNA Dishevelled 1 dan Dishevelled 2. Perkara ini berlaku di antara siRNA D1_9 dan siRNA D2_4 di mana terdapat pengurangan dari segi kemandirian sel yang dijangkiti oleh kedua-dua siRNA tersebut. Namun, tiada perubahan yang ketara dari segi kemandirian, penghijrahan dan serangan sel oleh siRNA yang lain yang juga menyasar Dishevelled 1 dan Dishevelled 2 (siRNA D1_6 dan siRNA D2_8). Kajian yang seterusnya menunjukkan bahawa siRNA D1_9 dan D2_4 mengakibatkan pengurangan serangan dan penghijrahan sel yang ketara dari analisa statistik. Ini menunjukkan bahawa terdapat kemungkinan dari segi ketidak-penghususannya dari segi penyasaran bagi kedua-dua siRNA tersebut dan juga berkemungkinan bahawa kesan pernyataan yang lebih tinggi bagi menjatuhkan pernyataan Dishevelled 2 bagi siRNA D2_4. (3) Dishevelled 3 memainkan peranan yang penting dalam aktiviti transkripsi TCF-LEF yang diantar oleh Wnt3a. Pengurangan aktiviti transkripsi TCF-LEF yang diantar oleh Wnt3a yang diperhatikan dalam sel MDA-MB-231 yang dijangkiti oleh siRNA yang menyasar Dishevelled 3 terutamanya bagi sel yang dijangkiti oleh siRNA D3_7 yang mengakibatkan pengurangan yang ketara secara statistik. (4) Keputusan densitometrik oleh fosfo-GSK3 β -Serine9 (relative dengan jumlah GSK3 β) dan fosfo- β -catenin (relative kepada jumlah β -catenin) bagi sel yang dijangkiti oleh siRNA Dishevelled 3 menunjukkan pernyataan fosfo-GSK3 β -Serine9 yang lebih tinggi, ini menandakan kecenderungan GSK3 β yang tidak aktif. Pernyataan fosfo-GSK3 β -Serine9 boleh mengakibatkan pengurangan kecenderungan untuk penguraian β -catenin (fosfo- β -catenin). Namun, pernyataan aras penguraian β -catenin (relative kepada jumlah β -catenin) oleh sel yang dijangkiti oleh siRNA Dishevelled 3 hanya menunjukkan aras yang sama berbanding dengan kawalan. Aras fosfo- β -catenin (relative kepada jumlah β -catenin) yang asas sebagaimana yang ditunjukkan oleh sel yang dijangkiti oleh siRNA Dishevelled 3 tidak menunjukkan sebarang kaitan dengan aktiviti GSK3 β dalam kajian ini. (5) Dengan mengaitkan pernyataan fosfo- β -catenin (relative kepada jumlah β -catenin) dalam analisis densitometrik dengan keputusan Luciferase assay, kurangnya kecenderungan dalam penguraian β -catenin (relative kepada jumlah β -catenin) mengakibatkan aras aktiviti TCF-LEF yang lebih tinggi bagi sel yang dijangkiti oleh Dishevelled 1 dan Dishevelled 2. Namun, bagi sel yang dijangkiti oleh siRNA Dishevelled 3, tiada pertambahan pernyataan fosfo- β -catenin (relative kepada jumlah β -catenin), meskipun terdapat kekurangan dalam aktiviti transkripsi TCF-LEF. Penyenyapan pernyataan Dishevelled 3 hanya menunjukkan pernyataan asas fosfo- β -catenin jika dibandingkan dengan kawalan. Penyenyapan pernyataan Dishevelled 3 mengekalkan pernyataan aras yang asas bagi fosfo- β -catenin dan pengekalannya pernyataan ini adalah memadai untuk penguraian dan pemisahan kompleks pemusnahan β -catenin yang mengakibatkan pengurangan aktiviti transkripsi yang diantar oleh TCF-LEF. Keputusan ini menyatakan kemungkinan bahawa gangguan interaksi antara Dishevelled 3 dengan kompleks pemusnah β -catenin dan bukannya Dishevelled 1 dan Dishevelled 2 yang mengakibatkan kurangnya aktiviti transkripsi TCF-LEF yang diantar oleh Wnt3a. Ada juga kemungkinan bahawa terdapat gangguan oleh tapak promoter di antara β -catenin dan TCF-LEF oleh sel yang dijangkiti oleh Dishevelled 3 siRNA. (6) Analisa densitometrik oleh Western Blot yang mengaitkan komponen laluan EGFR-ERK menunjukkan bahawa tiada sebarang pengaktifan laluan EGFR-ERK berlaku semasa stimulasi Wnt3a dan kejatuhan pernyataan isoform Dishevelled 3 tidak mengakibatkan kurangnya aktiviti EGFR-ERK semasa stimulasi Wnt3a. Kesimpulannya, Dishevelled 3 terlibat dalam isyarat laluan Wnt berkanun dalam sel

kanser payu dara MDA-MB-231 dan memainkan peranan yang penting dalam merencat penguraian β -catenin. Terdapat kemungkinan bahawa jatuhnya penyataan Dishevelled 3 mengganggu interaksi dengan Axin, dan mengakibatkan kurangnya pengumpulan β -catenin dan akhirnya mengurangkan aktiviti transkripsi TCF-LEF. Terdapat juga kemungkinan bahawa penyenyapan Dishevelled 3 mengganggu aktiviti transkripsi TCF –LEF secara langsung. Sebaliknya, tiadanya kesan pengurangan aktiviti transkripsi TCF-LEF yang diantara oleh β -catenin dan juga kurangnya penyataan fosfo- β -catenin relatif kepada jumlah β -catenin bagi sel yang dijangkiti oleh siRNA Dishevelled 1 dan Dishevelled 2. Ini menunjukkan bahawa tiadanya penglibatan isoform Dishevelled 1 dan Dishevelled 2 dalam isyarat laluan Wnt berkanun dalam sel MDA-MB-231.



ACKNOWLEDGEMENTS

First of all, I would like to thank Prof Dr Seow Heng Fong for taking me in to be involved in this study. Also, I would like to thank Dr Yip Wai Kien for the guidance in experiments. I am grateful that both supervisors provide a space and opportunity to grow, to learn and to experience this whole process.

I would like to thank my family for the support and patience given. I really appreciate their understanding of the hardships and struggles that I had been through along the process. In the end, they made me realize that family plays an important part of my life.

Also, my dear comrades along the journey of this study. I am grateful to get to know Mr Goh Tse Hau (Andrew), Miss Ayunadirah Binti Ayub (Ayu), Mr Ng Chin Tat (Jack), Miss Iris Goh Wen Li, lovely twins Miss Chai Boon Yean and Miss Chai Boon Lee and Mr He Pei Yuan. Along the process I am grateful for the presence of all of you in the journey of my studies. Indirectly all of you made me into a better person.

Also, I would like to thank my dear comrades in the lab. Miss Yiau Kang Xian (Stephnie), Mr Tan Jun Hao (James), Miss Catherine Chieng Cheing Yun (Cat), Mr Lee Le Jie, Miss Nuramziah (Ramziah), Miss Tan Shi Wei (Shi Wei), Mr Tong Chih Kong (aka Tong Gor), Mr Haw Tatt Yhew (Randy aka Dai Lou), Mr Shinsmon Jose (Shins), for their company at the workplace. I must say that all of you are the nicest people I have ever met.

It is undeniable that I had struggled along the process and I am grateful that God lead the way and let me realize things that are important for me. I am grateful that he has given me strength to go through hard time and also allow me to rest when I was in my lowest point of my life. I am grateful that I have become a better person than before.

I certify that a Thesis Examination Committee has met on 25 May 2015 to conduct the final examination of Samuel Khoo Leon Juan on his thesis entitled "Effects of Silencing Dishevelled 1, 2 and 3 mRNA Expression on MDA-MB-231 Cell Migration and Invasion using siRNAs" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Asmah binti Rahmat, PhD

Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Cheah Yoke Kqueen, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Internal Examiner)

Judith Helen Harmey, PhD

Associate Professor
Perdana University
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 12 August 2015


This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Seow Heng Fong, PhD

Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Yip Wai Kien, PhD

Postdoctoral Researcher
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Member)



ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No: Samuel Khoo Leon Juan, GS30265

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____

Name of Chairman
of Supervisory
Committee:

Professor Dr. Seow Heng Fong

Signature: _____

Name of Member
of Supervisory
Committee:

Dr. Yip Wai Kien

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	vi
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xvii
 CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 Breast Cancer Pathogenesis	4
2.2 Breast cancer Subtypes:	4
2.3 Breast Cancer Risk Factors	5
2.4 Breast Cancer treatment:	6
2.4.1 Breast Cancer Treatment: Surgical removal	6
2.4.2 Breast Cancer Treatment: Radiation therapy	6
2.4.3 Breast Cancer Treatment: Chemotherapy	7
2.4.4 Breast Cancer Treatment: Hormone Therapy	7
2.4.5 Breast Cancer Treatment: monoclonal antibody Therapy	8
2.5 Application of siRNA in Cancer Therapy Development	8
2.6 Wnt Signaling Pathway	10
2.6.1 Dishevelled and its domains	11
2.6.2 Canonical Wnt signaling pathway and the role of Dishevelled in canonical Wnt signaling pathway	12
2.6.3 Canonical Wnt signaling pathway in Breast Cancer	12
2.6.4 Non-canonical Signalling Pathway and role of Dishevelled in Non-Canonical Signaling pathway	13
2.6.5 Non- Canonical Wnt Signaling pathway in Breast Cancer	14
2.6.6 Role of Dishevelled in Development and Cancer	15
2.6.7 Studies related to Dishevelled using siRNA silencing	17
2.7 MDA-MB-231 Breast Cancer Cell Line	17
2.8 Wnt Inhibitor XAV-939	18
2.9 Wnt inhibitor ICG-001	18
3 MATERIALS AND METHODS	20
3.1 Cell Culture	20
3.1.1 Subculture of adherent cells	20
3.1.2 Wnt Inhibitors	20
3.1.3 Cell counting	21
3.2 Preparation of siRNA	21

	3.2.1	Knockdown of Dishevelled 1,2, 3 by transfection with siRNA	23
	3.3	RNA extraction	24
	3.3.1	Reverse transcription	24
	3.3.2	Quantitative Real Time PCR	25
	3.4	Western Blot and quantitation of respective target protein of interest	26
	3.5	TCF/LEF- sensitive luciferase reporter gene assay	28
	3.6	MTS Cell Viability Assay	29
	3.7	MTT Cell Viability Assay	29
	3.8	Cell Invasion Assay	30
	3.9	Wound Healing Assay	32
	3.10	Immunocytochemistry on siRNA transfected MDA-MB-231 cell line stimulated with Wnt3a	33
	3.11	Statistical Analysis	34
4	RESULTS		35
	4.1	Effect of Human Dishevelled mRNA levels after transfection with siRNAs targeting respective Dishevelled proteins.	35
	4.2	Effect of human Dishevelled protein levels after transfection with siRNAs targeting respective Dishevelled proteins	42
	4.3	Effects of cell viability after transfection with siRNAs targeting respective Dishevelled proteins.	42
	4.4	Effects of cell invasion after transfection with siRNAs targeting respective Dishevelled proteins	47
	4.5	Effects of Cell Migration with siRNAs targeting respective Dishevelled proteins.	50
	4.6	Effects of TCF-LEF transcriptional activation with siRNAs targeting respective Dishevelled proteins.	54
	4.7	Effects of β -catenin expression in Wnt β -catenin Signaling Pathway for MDA- MB-231 cell line after transfection with Dishevelled siRNAs	55
	4.8	Effects of molecules involved in Wnt β -catenin Signaling Pathway after transfection with Dishevelled siRNAs	57
	4.9	Effects of cell viability after treatment with Wnt Inhibitors XAV939 and ICG001	65
	4.10	Effects of Cell Migration after treatment with Wnt Inhibitors XAV939 and ICG001	69
	4.11	Effects on Molecules involving in EGFR pathway after Dishevelled knockdown under Wnt3a stimulation	71
5	DISCUSSION		78
	5.1	Effects of siRNA in mRNA and protein knockdown of Dishevelled 1,2 and 3 isoforms	78
	5.2	Different effects on cell viability, cell invasion and cell migration after transfection of Dishevelled siRNAs	78
	5.3	Knockdown of Dishevelled 3 isoforms results in reduced TCF-LEF transcriptional activation	80
	5.4	Different expression levels on components involving Wnt β -catenin Pathway	82

5.5	Different effects on cell biological effects after treatment with Wnt Inhibitors with target site of Wnt β -catenin pathway.	85
5.6	Different effects on between Dishevelled siRNAs towards expression of molecules involving in EGFR- ERK pathway activation	86
6	CONCLUSION AND FUTURE RECOMMENDATIONS	87
6.1	Conclusion	87
6.2	Future Recommendations	93
	REFERENCES	94
	APPENDICES	106
	BIODATA OF STUDENT	115



LIST OF TABLES

Table		Page
3.2	Sequence of purchased siRNA targeting mRNA of respective Dishevelled isoforms	22
3.2.1	Concentration siRNA transfected among cell lines	24
3.3.2	Steps and condition involved in RT-qPCR for primers targeting Human Dishevelled 1, 2, 3 and Human β -actin.	26
3.4	Primary Antibodies used for Western Blot Analysis	28
4.4	Percentage of Invaded cells after siRNA transfection targeting Dishevelled 1,2 and 3	48
4.8(a)	Fold changes in expression of total GSK3 β and p-GSK3 β -Ser9 with normalization to α -tubulin..	61
4.8(b)	Fold changes in expression of p-GSK3 β -Ser9 against total GSK3 β	62
4.8(c)	Fold changes in expression of total β -catenin and p- β -catenin with normalization to α -tubulin	63
4.8(d)	Fold changes in expression of p- β -catenin against total β -catenin	64
4.9	IC75 and IC50 generated on respective cell lines with Wnt Inhibitors ICG001 and XAV939 based on results from MTT assay.	69
4.11(a)	Fold Changes in expression of total EGFR and p-EGFR relative to α -tubulin	74
4.11(b)	Fold Changes in expression of p-EGFR against total EGFR	75
4.11(c)	Fold Changes in expression of total ERK and p-ERK after normalization with α -tubulin	76
4.11(d)	Fold Changes in expression of p-ERK against total ERK	77
6	Overview of single or combination siRNA Dishevelled isoforms in MDA-MB-231 cell line.	90
A.1	Laemmli resolving gel, 29:1 ratio	108
D	Primer pairs targeting Dishevelled1, 2 and 3 mRNAs with Ct Values Generated Based on Gradient RT-qPCR	112

LIST OF FIGURES

Figure	Page
1	2
2.1	12
2.2	14
3.8(a)	31
3.8(b)	32
4.1.1	36
4.1.2	37
4.1.3	38
4.1.4	39
4.1.5	40
4.1.6	41
4.2	42
4.3(a)	44
4.3(b)	44
4.3(c)	45
4.3(d)	45
4.3(e)	46
4.3(f)	46
4.3(g)	47

4.4	Effect of siRNAs targeting Dishevelled 1, 2 and 3 on MDA-MB-231 cell invasion.	50
4.5.1	Effect of siRNAs Dishevelled 1,2 and 3 on MDA-MB-231 cell migration under 10% FBS. Gap Recovery (%) Against Time for MDA-MB-231 cell lines transfected with (a) Dishevelled 1, (b) Dishevelled 2, and (c) Dishevelled 3 siRNAs under 10% FBS	52
4.5.2	Effect of siRNAs to Dishevelled 1,2,and 3 on MDA-MB-231 cell migration under 1% FBS. Gap Recovery (%) Against Time for MDA-MB-231 cell lines transfected with (a) Dishevelled 1, (b) Dishevelled 2, and (c) Dishevelled 3 siRNAs under 1% FBS.	53
4.6	Effect of siRNAs to Dishevelled 1,2 and 3 on MDA-MB-231 transcriptional activation.	55
4.7	Expression of β -catenin on Wnt3a stimulated, Dishevelled siRNA transfected MDA-MB-231	57
4.8	Effect of changes in protein expression involving Wnt3a stimulated Wnt β -catenin signaling on Dishevelled 1,2 and 3 silenced, Wnt3a stimulated MDA-MB-231.	60
4.9.1	Effects of cell viability using Wnt Inhibitor treatment on MDA-MB-231 with 1% FBS.	65
4.9.2	Effects of cell viability using Wnt Inhibitor treatment on HCT116 with 1% FBS.	66
4.9.3	Effects of cell viability using Wnt Inhibitor treatment on PC3 with 1% FBS.	66
4.9.4	Effects of cell viability using Wnt Inhibitor treatment on DU145 with 1% FBS.	67
4.9.5	Effects of cell viability using Wnt Inhibitor treatment on T47D with 1% FBS.	67
4.9.6	Effects of cell viability using Wnt Inhibitor treatment on MCF7 with 1% FBS.	68
4.9.7	Effects of cell viability using Wnt Inhibitor treatment on SiHa with 1% FBS.	68
4.10	Effect of Wnt Inhibitors XAV939 and ICG001 on MDA-MB-231, PC3 and DU145 cell lines.	70
4.11	Effect of changes in protein expression involving EGFR pathway on Dishevelled 1,2 and 3 silenced, Wnt3a stimulated MDA-MB-231.	73

LIST OF ABBREVIATIONS

%	Percentages
°C	Degree Celsius
Akt	V-akt murine thymoma viral oncogene homolog 1
ANOVA	Analysis of Variance
APC	Adenomatous Polyposis Coli
BCL	B-cell lymphoma
b.p	base pairs
BRCA1	Breast Cancer 1, early onset
BRCA2	Breast Cancer 2, early onset
BSA	Bovine Serum Albumin
CAMKII	Ca ²⁺ /calmodulin-dependent protein kinase
CBP	CREB Binding Protein
cDNA	complimentary deoxyribonucleic acid
CO ₂	Carbon dioxide
Daam1	Dishevelled associate activator of morphogenesis 1
DAPI	4',6-diamidino-2-phenylindole
DEP	<i>Dishevelled</i> , Egl-10 and Pleckstrin domain
DEPC	Diethylpyrocarbonate
DIX	Dishevelled/Axin domain
DMSO	Dimethylsulfoxide
DNA	Deoxyribonucleic acid.
dNTP	Deoxy-nucleotide-triphosphates
DSH	Dishevelled (short form for <i>Drosophila sp.</i>)
DTT	Dithiothreitol

DVL	Dishevelled (short form for <i>homo sapiens</i>).
EGF	Epidermal Growth Factor
EGFR	Epidermal Growth Factor Receptor
ER	Estrogen Receptor
ERK	Extracellular signal-regulated kinase
<i>et. al</i>	And all.
FBS	Fetal Bovine Serum
FZD	Frizzled
GSK3 β	Glycogen Synthase Kinase 3 β
HCl	Hydrochloric acid
HER2	Human Epidermal Growth Receptor 2
HMGA2	High-mobility group AT-hook 2
HRP	Horse Radish Peroxidase
HRT	Hormone Replacement Therapy
IC	Inhibitory Concentration
IDC	Invasive Ductal Carcinoma
IGF	Insulin Growth Factor
JNK	c-Jun N-terminal Kinase
KCl	Potassium Chloride
LEF	Lymphoid Enhancer Factor
LiCl	Lithium Chloride
M	mol dm ⁻³
MAPK	Mitogen Activated Protein Kinase
mRNA	Messenger ribonucleic acid
MTS	3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4 sulfophenyl)-2H-tetrazolium

MTT	3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
NLK	Nemo-like kinase
OD	Optical density
PAGE	Polyacrylamide gel electrophoresis
PAI	Plasminogen activator inhibitor
PARP	Poly ADP Ribose Polymerase
PBS	Phosphate buffered saline
PCP	Planar cell polarity
PCR	Polymerase chain reaction
PDZ	Postsynaptic Sensity as, disc large, zonula occludens
PEC	Polyelectrolyte complex
p-EGFR	Phosphorylated epidermal growth factor receptor
p-ERK	Phosphorylated extracellular signal-regulated kinase
p-GSK3 β -Ser9	Phosphorylate glycogen synthase Kinase 3 beta serine 9
PI	Propidium iodide
PI3K	Phosphatidylinositol 3-kinase
PKC	Protein kinase C
PR	Progesterone receptor
PVDF	Polyvinylidene difluoride
qPCR	Quantitative Polymerase Chain Reaction.
S.D	Standard Deviation
SH3	Src homology 3
siRNA	Small interfering ribonucleic acid.
TCF	T-cell factor
TGF- β	Transforming growth factor beta

TKNS	Tankyrase
TNBC	Triple negative breast cancer
VEGF	Vaso-endothelial growth factor
Wnt	Wingless-type MMTV integration site family



© COPYRIGHT UPM

CHAPTER 1

INTRODUCTION

Introduction Breast cancer is the most common cancer amongst women which can result in death (Greenlee *et al.*, 2000). In the year 2013, 32,340 new cases of invasive breast cancer and 39,620 breast cancer deaths were reported in United States (DeSantis *et al.*, 2014). Current treatment options for breast cancer include mastectomy, radiotherapy and chemotherapy (Maughan *et al.*, 2010)

Wnt signaling pathway is a signal transduction pathway which plays an important role in embryonic development, generation of cell polarity and cell specification (Logan *et al.*, 2004). Aberrant activation of Wnt pathway leads to tumorigenesis (Giles *et al.*, 2003). The Wnt signaling pathway consists of canonical signaling pathway and non-canonical signaling pathway. The Wnt non-canonical signaling pathway can be further divided into Wnt planar cell polarity pathway (Wang, 2009) and Wnt calcium pathway (Kühl, 2004).

Wnt was discovered as an oncogene which contributes to mouse mammary tumorigenesis in mouse models of breast cancer. Aberrant activation of Wnt signaling pathway is also linked to tumorigenesis on human cancers (Brown, 2001) which caused by mutations of certain key components that are involved in Wnt signaling pathway. Mutation of certain components can be seen in colorectal cancer, namely APC (Adenomatous Polyposis Coli), a component of the canonical Wnt β -Catenin signaling pathway which plays an important role in β -catenin degradation (Axelrod *et al.*, 1998). Mutations of APC result in resistance of β -catenin to undergo the process of ubiquitinylation and phosphorylation (Bienz *et al.*, 2000). This leads to β -catenin accumulation in the cytoplasm which further results in activation of downstream pathways after T cell Factor (TCF) and Lymphoid Enhancer Factor (LEF) transcriptional activation in the nucleus (Morin *et al.*, 1997). Mutation of other components such as β -catenin itself and Axin also contributes in aberrant activation of canonical Wnt β -catenin pathway (Lammi *et al.*, 2004).




Figure 1 : Mutation of APC in Wnt Signaling Pathway. Mutation of APC resulted in the ability of APC to be involved in β -catenin degradation complex (Narayan *et al.*, 2003). Increased stabilized β -catenin resulted in accumulation of β -catenin and translocation into the nucleus, where β -catenin forms a complex with TCF and LEF to activate TCF-LEF mediated transcriptional activation of downstream of Wnt target genes such as *c-myc* and *cyclin D1*.

Both canonical and non-canonical Wnt signaling pathways play a role in the pathogenesis of breast cancer metastasis (Dey *et al.*, 2013; Jiang *et al.*, 2013; Zhu *et al.*, 2012). Increased cytoplasmic and nuclear β -catenin levels contributes to an aberrant activation of canonical Wnt Signaling Pathway. Increased expression of cytoplasmic and nuclear β -catenin levels were found in breast cancer samples which indicate an aberrant canonical wnt signaling pathway activation (Prasad *et al.*, 2007). Further studies also showed that upregulation of Wnt β -catenin pathway correlates with elevated expression of Cyclin D1, which is one of the downstream targets of Wnt β -catenin pathway, contributing towards increased cell proliferation (Prasad *et al.*, 2007). On the other hand, β -catenin levels were not observed in normal breast tissues (Prasad *et al.*, 2007). Thus, this provides evidence that stabilized β -catenin also contributes in Wnt β -catenin pathway which results in tumorigenesis of breast cancer.

There are different ways to study the role of Wnt Signaling in breast cancer tumorigenesis by blocking different components that are involved in the pathway. One of the approaches used in previous studies is to block Wnt/FZD (Frizzled) interaction, which is located upstream of the Wnt signaling pathway (Matsuda *et al.*, 2009; Yang *et al.*, 2011). Blocking of this interaction caused a reduction in cell invasion and also in cell proliferation (Matsuda *et al.*, 2009; Yang *et al.*, 2011). Also, blockade of Wnt

signaling is also performed by increasing the expression of certain components involved in β -catenin degradation complex, which can facilitate in reduction of β -catenin levels.

Dishevelled is one of the components which is normally located in the cytoplasm. It is involved in both canonical and non-canonical Wnt signaling pathway. Dishevelled has been shown to play a role in the canonical and non-canonical Wnt signaling pathway in breast cancer cells by influencing cell proliferation, polarity, terminal differentiation and self renewal ability of stem cells (Zhao *et al.*, 2010; Zhu *et al.*, 2012). The expression of Dishevelled isoforms has been found in the cytoplasm and also in the nucleus of invasive ductal carcinomas (Prasad *et al.*, 2007).

In this study, human Dishevelled 1, 2 and 3 isoforms expression was silenced by using siRNA (small interfering RNA) in order to block the canonical or non-canonical Wnt signaling pathway at the cytoplasmic level. This ribonucleic acid which contains 21-23 nucleotide bases plays a role in RNA interference by binding specifically to mRNA (messenger RNA) in order to cleave a specific mRNA. The cleaved mRNA product results in further degradation by cytoplasmic nucleases. This reduces the mRNA levels in siRNA transfected cells which ultimately leads to reduced translation of target protein.

Our hypothesis is that silencing of human Dishevelled 1,2 and 3 mRNA expression leads to blockade in activation of canonical Wnt signaling pathway leading to reduced β -catenin interaction with TCF and LEF, reduced cell proliferation, migration and also in invasion regardless of canonical or non-canonical Wnt signaling pathway.

The overall objective of this study is to determine the role of Dishevelled 1, 2 and 3 in canonical Wnt signaling pathway in breast cancer using the gene silencing approach. The specific objectives are as follows:

1. To confirm the silencing of Dishevelled 1, 2 and 3 by reverse-transcription Quantitative PCR and Western Blotting.
2. To determine the changes in cell viability, migration, invasion, TCF-LEF mediated transcriptional activation after gene silencing using siRNA targeting Dishevelled isoforms.
3. To identify the changes in β -catenin, Glycogen Synthase Kinase 3 β (GSK3 β), Epidermal Growth Factor Receptor (EGFR) and Extracellular signal-Regulated Kinases (ERK) expression in cells transfected with siRNAs targeting Dishevelled 1, 2 and 3.

REFERENCES

- Adams, J., & Cory, S. (2007). The Bcl-2 apoptotic switch in cancer development and therapy. *Oncogene*, 26(9), 1324-1337.
- Allred, C. D., Ju, Y. H., Allred, K. F., Chang, J., & Helferich, W. G. (2001). Dietary genistein stimulates growth of estrogen-dependent breast cancer tumors similar to that observed with genistein. *Carcinogenesis*, 22(10), 1667-1673.
- Axelrod, J. D., Miller, J. R., Shulman, J. M., Moon, R. T., & Perrimon, N. (1998). Differential recruitment of Dishevelled provides signaling specificity in the planar cell polarity and Wingless signaling pathways. *Genes & Development*, 12(16), 2610-2622.
- Bao, R., Christova, T., Song, S., Angers, S., Yan, X., & Attisano, L. (2012). Inhibition of tankyrases induces Axin stabilization and blocks Wnt signalling in breast cancer cells. *PLoS One*, 7(11).
- Bienz, M., & Clevers, H. (2000). Linking colorectal cancer to Wnt signaling. *Cell*, 103(2), 311-320.
- Bikkavilli, R. K., Feigin, M. E., & Malbon, C. C. (2008). Gao mediates WNT-JNK signaling through Dishevelled 1 and 3, RhoA family members, and MEKK 1 and 4 in mammalian cells. *Journal of Cell Science*, 121(2), 234-245.
- Bikkavilli, R. K., Feigin, M. E., & Malbon, C. C. (2008). p38 mitogen-activated protein kinase regulates canonical Wnt- β -catenin signaling by inactivation of GSK3 β . *Journal of Cell Science*, 121(21), 3598-3607.
- Boutros, M., & Mlodzik, M. (1999). Dishevelled: at the crossroads of divergent intracellular signaling pathways. *Mechanisms of Development*, 83(1-2), 27-37.
- Boutros, M., Paricio, N., Strutt, D. I., & Mlodzik, M. (1998). Dishevelled activates JNK and discriminates between JNK pathways in planar polarity and wingless signaling. *Cell*, 94(1), 109-118.
- Brown, A. (2001). Wnt signaling in breast cancer: have we come full circle? *Breast Cancer Research*, 3(6), 351-355.
- Bui, T. D., Beier, D. R., Jonssen, M., Smith, K., Dorrington, S. M., Kaklamanis, L., Kearney, L., Regan, R., Sussman, D. J., & Harris, A. L. (1997). cDNA Cloning of a Human Dishevelled DVL-3 Gene, Mapping to 3q27, and Expression in Human Breast and Colon Carcinomas. *Biochemical and Biophysical Research Communications*, 239(2), 510-516.
- Cailleau, R., Young, R., Olivé, M., & Reeves, W. J. (1974). Breast Tumor Cell Lines From Pleural Effusions. *Journal of the National Cancer Institute*, 53(3), 661-674.

- Carey, L. A., Perou, C. M., Livasy, C. A., Dressler, L. G., Cowan, D., Conway, K., Karaca, G., Troester, M. A., Tse, C. K., & Edmiston, S. (2006). Race, breast cancer subtypes, and survival in the Carolina Breast Cancer Study. *JAMA*, 295(21), 2492-2502.
- Carthew, R. W., & Sontheimer, E. J. (2009). Origins and Mechanisms of miRNAs and siRNAs. *Cell*, 136(4), 642-655.
- Chen, Y., & Bodles, A. M. (2007). Amyloid precursor protein modulates beta-catenin degradation. *Journal of Neuroinflammation*, 4(1), 29.
- Cianfrocca, M., & Gradishar, W. J. (2005). Controversies in the therapy of early stage breast cancer. *The Oncologist*, 10(10), 766-779.
- Clemons, M., & Goss, P. (2001). Estrogen and the risk of breast cancer. *New England Journal of Medicine*, 344(4), 276-285.
- Consonni, S. V., Maurice, M. M., & Bos, J. L. (2014). DEP domains: structurally similar but functionally different. *Nature Reviews Molecular Cell Biology*, 15(5), 357-362.
- CuboCube. (2013). Experimental Techniques for Investigating Metastasis Retrieved 20 January, 2015, from <http://cubocube.com/dashboard.php?a=378&b=1605&c=1>
- Cullen, B. R. (2006). Enhancing and confirming the specificity of RNAi experiments. *Nature Methods*, 3(9), 677-681.
- Curtis, R. E., Boice, J. D., Shriner, D. A., Hankey, B. F., & Fraumeni, J. F. (1996). Second cancers after adjuvant tamoxifen therapy for breast cancer. *Journal of the National Cancer Institute*, 88(12), 832-835.
- DeSantis, C., Ma, J., Bryan, L., & Jemal, A. (2014). Breast cancer statistics, 2013. *CA: A Cancer Journal for Clinicians*, 64(1), 52-62.
- Dey, N., Barwick, B., Moreno, C., Ordanic-Kodani, M., Chen, Z., Oprea-Illies, G., Tang, W., Catzavelos, C., Kerstann, K., Sledge, G., Abramovitz, M., Bouzyk, M., De, P., & Leyland-Jones, B. (2013). Wnt signaling in triple negative breast cancer is associated with metastasis. *BMC Cancer*, 13(1), 537.
- Dillman, A. R., Minor, P. J., & Sternberg, P. W. (2013). Origin and evolution of dishevelled. *G3: Genes/ Genomes/ Genetics*, 3(2), 251-262.
- Echeverri, C. J., Beachy, P. A., Baum, B., Boutros, M., Buchholz, F., Chanda, S. K., Downward, J., Ellenberg, J., Fraser, A. G., & Hacohen, N. (2006). Minimizing the risk of reporting false positives in large-scale RNAi screens. *Nature Methods*, 3(10), 777-779.

- Ewer, M. S., & Lippman, S. M. (2005). Type II chemotherapy-related cardiac dysfunction: time to recognize a new entity. *Journal of Clinical Oncology*, 23(13), 2900-2902.
- Ewer, S. M., & Ewer, M. S. (2008). Cardiotoxicity profile of trastuzumab. *Drug Safety*, 31(6), 459-467.
- Ford, D., Easton, D., Stratton, M., Narod, S., Goldgar, D., Devilee, P., Bishop, D., Weber, B., Lenoir, G., & Chang-Claude, J. (1998). Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families. *The American Journal of Human Genetics*, 62(3), 676-689.
- Gan, X.-q., Wang, J.-y., Xi, Y., Wu, Z.-l., Li, Y.-p., & Li, L. (2008). Nuclear Dvl, c-Jun, β -catenin, and TCF form a complex leading to stabilization of β -catenin–TCF interaction. *The Journal of Cell Biology*, 180(6), 1087-1100.
- Gao, C., & Chen, Y.-G. (2010). Dishevelled: The hub of Wnt signaling. *Cellular Signalling*, 22(5), 717-727.
- Giles, R. H., van Es, J. H., & Clevers, H. (2003). Caught up in a Wnt storm: Wnt signaling in cancer. *Biochim Biophys Acta*, 1653(1), 1-24.
- Gnerlich, J., Jeffe, D. B., Deshpande, A. D., Beers, C., Zander, C., & Margenthaler, J. A. (2007). Surgical removal of the primary tumor increases overall survival in patients with metastatic breast cancer: analysis of the 1988–2003 SEER data. *Annals of Surgical Oncology*, 14(8), 2187-2194.
- Goldhirsch, A., Glick, J., Gelber, R., Coates, A., Thürlimann, B., & Senn, H.-J. (2005). Meeting highlights: international expert consensus on the primary therapy of early breast cancer 2005. *Annals of Oncology*, 16(10), 1569-1583.
- Greenlee, R. T., Murray, T., Bolden, S., & Wingo, P. A. (2000). Cancer statistics, 2000. *CA: A Cancer Journal for Clinicians*, 50(1), 7-33.
- Ha, N.-C., Tonzuka, T., Stamos, J. L., Choi, H.-J., & Weis, W. I. (2004). Mechanism of phosphorylation-dependent binding of APC to β -catenin and its role in β -catenin degradation. *Molecular Cell*, 15(4), 511-521.
- Habas, R., Kato, Y., & He, X. (2001). Wnt/Frizzled activation of Rho regulates vertebrate gastrulation and requires a novel Formin homology protein Daam1. *Cell*, 107(7), 843-854.
- Hamblet, N. S., Lijam, N., Ruiz-Lozano, P., Wang, J., Yang, Y., Luo, Z., Mei, L., Chien, K. R., Sussman, D. J., & Wynshaw-Boris, A. (2002). Dishevelled 2 is essential for cardiac outflow tract development, somite segmentation and neural tube closure. *Development*, 129(24), 5827-5838.
- Hanahan, D., & Folkman, J. (1996). Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis. *Cell*, 86(3), 353-364.

- Hanahan, D., & Weinberg, R. A. (2011). Hallmarks of cancer: the next generation. *Cell*, 144(5), 646-674.
- Hao, S., He, W., Li, Y., Ding, H., Hou, Y., Nie, J., Hou, F. F., Kahn, M., & Liu, Y. (2011). Targeted inhibition of β -catenin/CBP signaling ameliorates renal interstitial fibrosis. *Journal of the American Society of Nephrology*, 22(9), 1642-1653.
- Hayes, D. F., Thor, A. D., Dressler, L. G., Weaver, D., Edgerton, S., Cowan, D., Broadwater, G., Goldstein, L. J., Martino, S., & Ingle, J. N. (2007). HER2 and response to paclitaxel in node-positive breast cancer. *New England Journal of Medicine*, 357(15), 1496-1506.
- Hitt. (2013). Wnt Signaling Inhibition: Will Decades of Effort Be Fruitful at Last? Retrieved 20 January 2015, from <http://www.onclive.com/publications/oncology-live/2012/december-2012/wnt-signaling-inhibition-will-decades-of-effort-be-fruitful-at-last/1>.
- Hitt, E. (2013). Wnt Signaling Inhibition : Will Decades of Effort Be Fruitful at Last? *OncLive, Intellisphere LLC*.
- Huang, S.-M. A., Mishina, Y. M., Liu, S., Cheung, A., Stegmeier, F., Michaud, G. A., Charlat, O., Wiellette, E., Zhang, Y., & Wiessner, S. (2009). Tankyrase inhibition stabilizes axin and antagonizes Wnt signalling. *Nature*, 461(7264), 614-620.
- Ireland, M., English, C., Cross, I., Houlsby, W., & Burn, J. (1991). A de novo translocation t (3; 17)(q26. 3; q23. 1) in a child with Cornelia de Lange syndrome. *Journal of Medical Genetics*, 28(9), 639-640.
- Ismail, S. (1994). Pathology of endometrium treated with tamoxifen. *Journal of Clinical Pathology*, 47(9), 827-833.
- Iwami, Y., Masuda, H., & Asahara, T. (2004). Endothelial progenitor cells: past, state of the art, and future. *Journal of Cellular and Molecular Medicine*, 8(4), 488-497.
- Jackson, A. L., Bartz, S. R., Schelter, J., Kobayashi, S. V., Burchard, J., Mao, M., Li, B., Cavet, G., & Linsley, P. S. (2003). Expression profiling reveals off-target gene regulation by RNAi. *Nature Biotechnology*, 21(6), 635-637.
- Jackson, A. L., Burchard, J., Leake, D., Reynolds, A., Schelter, J., Guo, J., Johnson, J. M., Lim, L., Karpilow, J., & Nichols, K. (2006). Position-specific chemical modification of siRNAs reduces “off-target” transcript silencing. *RNA*, 12(7), 1197-1205.
- Jackson, A. L., & Linsley, P. S. (2010). Recognizing and avoiding siRNA off-target effects for target identification and therapeutic application. *Nature Reviews Drug Discovery*, 9(1), 57-67.

- Jiang, W., Crossman, D. K., Mitchell, E. H., Sohn, P., Crowley, M. R., & Serra, R. (2013). WNT5A inhibits metastasis and alters splicing of Cd44 in breast cancer cells. *PLoS One*, 8(3), e58329.
- Ju, Y. H., Doerge, D. R., Allred, K. F., Allred, C. D., & Helferich, W. G. (2002). Dietary genistein negates the inhibitory effect of tamoxifen on growth of estrogen-dependent human breast cancer (MCF-7) cells implanted in athymic mice. *Cancer Research*, 62(9), 2474-2477.
- Julius, M., Schelbert, B., Hsu, W., Fitzpatrick, E., Jho, E., Fagotto, F., Costantini, F., & Kitajewski, J. (2000). Domains of axin and disheveled required for interaction and function in wnt signaling. *Biochemical and Biophysical Research Communications*, 276(3), 1162-1169.
- Kafka, A., Bašić-Kinda, S., & Pećina-Šlaus, N. (2014). The cellular story of dishevelleds. *Croatian Medical Journal*, 55(5), 459-467.
- Kafka, A., Tomas, D., Beroš, V., Pećina, H. I., Zeljko, M., & Pećina-Šlaus, N. (2014). Brain Metastases from Lung Cancer Show Increased Expression of DVL1, DVL3 and Beta-Catenin and Down-Regulation of E-Cadherin. *International Journal of Molecular Sciences*, 15(6), 10635-10651.
- Katoh, M. (2005). WNT/PCP signaling pathway and human cancer (Review). *Oncology Reports*, 14(6), 1583.
- Katso, R., Okkenhaug, K., Ahmadi, K., White, S., Timms, J., & Waterfield, M. D. (2001). Cellular function of phosphoinositide 3-kinases: implications for development, immunity, homeostasis, and cancer. *Annual Review of Cell and Developmental Biology*, 17(1), 615-675.
- Khan, S. A., Rogers, M. A., Obando, J. A., & Tamsen, A. (1994). Estrogen receptor expression of benign breast epithelium and its association with breast cancer. *Cancer Research*, 54(4), 993-997.
- Kim, S.-E., & Choi, K.-Y. (2007). EGF receptor is involved in WNT3a-mediated proliferation and motility of NIH3T3 cells via ERK pathway activation. *Cellular Signalling*, 19(7), 1554-1564.
- Kim, S. H., Jeong, J. H., Lee, S. H., Kim, S. W., & Park, T. G. (2008). Local and systemic delivery of VEGF siRNA using polyelectrolyte complex micelles for effective treatment of cancer. *Journal of Controlled Release*, 129(2), 107-116.
- Kishida, S., Yamamoto, H., Hino, S.-i., Ikeda, S., Kishida, M., & Kikuchi, A. (1999). DIX domains of Dvl and Axin are necessary for protein interactions and their ability to regulate β -catenin stability. *Molecular and Cellular Biology*, 19(6), 4414-4422.

- Klapholz-Brown, Z., Walmsley, G. G., Nusse, Y. M., Nusse, R., & Brown, P. O. (2007). Transcriptional program induced by Wnt protein in human fibroblasts suggests mechanisms for cell cooperativity in defining tissue microenvironments. *PLoS One*, 2(9), e945.
- Kühl, M. (2004). The WNT/calcium pathway: biochemical mediators, tools and future requirements. *Frontiers of Science: A Journal and Virtual Library*, 9(1), 967-974.
- Kühl, M., Sheldahl, L. C., Park, M., Miller, J. R., & Moon, R. T. (2000). The Wnt/Ca²⁺ pathway: a new vertebrate Wnt signaling pathway takes shape. *Trends in Genetics*, 16(7), 279-283.
- Kwan, H. T., Chan, D. W., Cai, P. C., Mak, C. S., Yung, M. M., Leung, T. H., Wong, O. G., Cheung, A. N., & Ngan, H. Y. (2013). AMPK activators suppress cervical cancer cell growth through inhibition of DVL3 mediated Wnt/beta-catenin signaling activity. *PLoS One*, 8(1), e53597.
- Lako, M., Strachan, T., Bullen, P., Wilson, D. I., Robson, S. C., & Lindsay, S. (1998). Isolation, characterisation and embryonic expression of WNT11, a gene which maps to 11q13.5 and has possible roles in the development of skeleton, kidney and lung. *Gene*, 219(1-2), 101-110. doi: 10.1016/s0378-1119(98)00393-x
- Lammi, L., Arte, S., Somer, M., Järvinen, H., Lahermo, P., Thesleff, I., Pirinen, S., & Nieminen, P. (2004). Mutations in Axin2 cause familial tooth genesis and predispose to colorectal cancer. *The American Journal of Human Genetics*, 74(5), 1043-1050.
- Lee, H. Z., JJ. (2011). DVL1 (dishevelled, dsh homolog 1 (Drosophila)). *Atlas of Genetics and Cytogenetics in Oncology and Haematology*, :998-1004.(12).
- Lee, Y.-N., Gao, Y., & Wang, H.-y. (2008). Differential mediation of the Wnt canonical pathway by mammalian Dishevelleds-1,-2, and-3. *Cellular Signalling*, 20(2), 443-452.
- Li, L., Yuan, H., Weaver, C. D., Mao, J., Farr, G. H., 3rd, Sussman, D. J., Jonkers, J., Kimelman, D., & Wu, D. (1999). Axin and Frat1 interact with dvl and GSK, bridging Dvl to GSK in Wnt-mediated regulation of LEF-1. *EMBO J*, 18(15), 4233-4240. doi: 10.1093/emboj/18.15.4233
- Liu, C., Li, Y., Semenov, M., Han, C., Baeg, G.-H., Tan, Y., Zhang, Z., Lin, X., & He, X. (2002). Control of β -catenin phosphorylation/degradation by a dual-kinase mechanism. *Cell*, 108(6), 837-847.
- Liu, R., Page, C., Beidler, D. R., Wicha, M. S., & Núñez, G. (1999). Overexpression of Bcl-x_L Promotes Chemotherapy Resistance of Mammary Tumors in a Syngeneic Mouse Model. *The American Journal of Pathology*, 155(6), 1861-1867.

- Logan, C. Y., & Nusse, R. (2004). The Wnt signaling pathway in development and disease. *Annual Review of Cell and Developmental Biology*, 20, 781-810.
- Ma, H., Nguyen, C., Lee, K.-S., & Kahn, M. (2005). Differential roles for the coactivators CBP and p300 on TCF/ β -catenin-mediated survivin gene expression. *Oncogene*, 24(22), 3619-3631.
- MacDonald, B. T., Tamai, K., & He, X. (2009). Wnt/ β -catenin signaling: components, mechanisms, and diseases. *Developmental Cell*, 17(1), 9-26.
- Matsuda, Y., Schlange, T., Oakeley, E. J., Boulay, A., & Hynes, N. E. (2009). WNT signaling enhances breast cancer cell motility and blockade of the WNT pathway by sFRP1 suppresses MDA-MB-231 xenograft growth. *Breast Cancer Research*, 11(3), R32.
- Maughan, K. L., Lutterbie, M. A., & Ham, P. S. (2010). Treatment of breast cancer. *American Family Physician*, 81(11), 1339-1346.
- McDonald, S., & Silver, A. (2009). The opposing roles of Wnt-5a in cancer. *British Journal of Cancer*, 101(2), 209-214.
- McPherson, K., Steel, C., & Dixon, J. (2000). ABC of breast diseases: breast cancer—epidemiology, risk factors, and genetics. *BMJ: British Medical Journal*, 321(7261), 624.
- Metcalfé, C., Ibrahim, A. E., Graeb, M., de la Roche, M., Schwarz-Romond, T., Fiedler, M., Winton, D. J., Corfield, A., & Bienz, M. (2010). Dvl2 promotes intestinal length and neoplasia in the ApcMin mouse model for colorectal cancer. *Cancer Research*, 70(16), 6629-6638.
- Miki, Y., Swensen, J., Shattuck-Eidens, D., Futreal, P. A., Harshman, K., Tavtigian, S., Liu, Q., Cochran, C., Bennett, L. M., & Ding, W. (1994). A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science*, 266(5182), 66-71.
- Miyake, H., HARA, I., YAMANAKA, K., ARAKAWA, S., & KAMIDONO, S. (1999). Synergistic enhancement of resistance to cisplatin in human bladder cancer cells by overexpression of mutant-type p53 and Bcl-2. *The Journal of Urology*, 162(6), 2176-2181.
- Miyashita, T., & Reed, J. C. (1993). Bcl-2 oncoprotein blocks chemotherapy-induced apoptosis in a human leukemia cell line. *Blood*, 81(1), 151-157.
- Mizutani, K., Miyamoto, S., Nagahata, T., Konishi, N., Emi, M., & Onda, M. (2005). Upregulation and overexpression of DVL1, the human counterpart of the Drosophila dishevelled gene, in prostate cancer. *Tumorigenesis*, 91(6), 546.
- Morin, P. J., Sparks, A. B., Korinek, V., Barker, N., Clevers, H., Vogelstein, B., & Kinzler, K. W. (1997). Activation of β -catenin-Tcf signaling in colon cancer by mutations in β -catenin or APC. *Science*, 275(5307), 1787-1790.

- Morrone, S., Cheng, Z., Moon, R. T., Cong, F., & Xu, W. (2012). Crystal structure of a Tankyrase-Axin complex and its implications for Axin turnover and Tankyrase substrate recruitment. *Proceedings of the National Academy of Sciences*, *109*(5), 1500-1505.
- Mourits, M. J., De Vries, E. G., Willemse, P. H., Ten Hoor, K. A., Hollema, H., & Van der Zee, A. G. (2001). Tamoxifen treatment and gynecologic side effects: a review. *Obstetrics & Gynecology*, *97*(5), 855-866.
- Mu, P., Nagahara, S., Makita, N., Tarumi, Y., Kadomatsu, K., & Takei, Y. (2009). Systemic delivery of siRNA specific to tumor mediated by atelocollagen: Combined therapy using siRNA targeting Bcl-xL and cisplatin against prostate cancer. *International Journal of Cancer*, *125*(12), 2978-2990.
- Nagahata, T., Shimada, T., Harada, A., Nagai, H., Onda, M., Yokoyama, S., Shiba, T., Jin, E., Kawanami, O., & Emi, M. (2003). Amplification, up-regulation and over-expression of DVL-1, the human counterpart of the *Drosophila* dishevelled gene, in primary breast cancers. *Cancer Science*, *94*(6), 515-518.
- Narayan, S., & Roy, D. (2003). Role of APC and DNA mismatch repair genes in the development of colorectal cancers. *Molecular Cancer*, *2*(1), 41.
- Negro, A., Brar, B. K., & Lee, K.-F. (2004). Essential roles of Her2/erbB2 in cardiac development and function. *Recent Progress in Hormone Research*, *59*(1), 1-12.
- Nicole, S., Hamida, C. B., Beighton, P., Bakouri, S., Belal, S., Romero, N., Viljoen, D., Ponsot, G., Sammoud, A., & Weissenbach, J. (1995). Localization of the Schwartz-Jampel syndrome (SJS) locus to chromosome 1p34-p36. 1 by homozygosity mapping. *Human Molecular Genetics*, *4*(9), 1633-1636.
- Orban, T. I., & Izaurralde, E. (2005). Decay of mRNAs targeted by RISC requires XRN1, the Ski complex, and the exosome. *RNA*, *11*(4), 459-469.
- Pàez-Ribes, M., Allen, E., Hudock, J., Takeda, T., Okuyama, H., Viñals, F., Inoue, M., Bergers, G., Hanahan, D., & Casanovas, O. (2009). Antiangiogenic therapy elicits malignant progression of tumors to increased local invasion and distant metastasis. *Cancer Cell*, *15*(3), 220-231.
- Penton, A., Wodarz, A., & Nusse, R. (2002). A mutational analysis of dishevelled in *Drosophila* defines novel domains in the dishevelled protein as well as novel suppressing alleles of axin. *Genetics*, *161*(2), 747-762.
- Persengiev, S. P., Zhu, X., & Green, M. R. (2004). Nonspecific, concentration-dependent stimulation and repression of mammalian gene expression by small interfering RNAs (siRNAs). *RNA*, *10*(1), 12-18.

- Prasad, C. P., Gupta, S. D., Rath, G., & Ralhan, R. (2007). Wnt signaling pathway in invasive ductal carcinoma of the breast: relationship between beta-catenin, dishevelled and cyclin D1 expression. *Oncology*, 73(1-2), 112-117.
- Pulvirenti, T., Van Der Heijden, M., Droms, L. A., Huse, J. T., Tabar, V., & Hall, A. (2011). Dishevelled 2 signaling promotes self-renewal and tumorigenicity in human gliomas. *Cancer Research*, 71(23), 7280-7290.
- Rakha, E. A., Reis-Filho, J. S., & Ellis, I. O. (2008). Basal-like breast cancer: a critical review. *Journal of Clinical Oncology*, 26(15), 2568-2581.
- Resnier, P., Montier, T., Mathieu, V., Benoit, J.-P., & Passirani, C. (2013). A review of the current status of siRNA nanomedicines in the treatment of cancer. *Biomaterials*, 34(27), 6429-6443.
- Rundhaug, J. E. (2005). Matrix metalloproteinases and angiogenesis. *Journal of Cellular and Molecular Medicine*, 9(2), 267.
- Sasaki, T., Nakashiro, K.-i., Tanaka, H., Azuma, K., Goda, H., Hara, S., Onodera, J., Fujimoto, I., Tanji, N., & Yokoyama, M. (2010). Knockdown of Akt isoforms by RNA silencing suppresses the growth of human prostate cancer cells *in vitro* and *in vivo*. *Biochemical and Biophysical Research Communications*, 399(1), 79-83.
- Sato, A., Khadka, D. K., Liu, W., Bharti, R., Runnels, L. W., Dawid, I. B., & Habas, R. (2006). Profilin is an effector for Daam1 in non-canonical Wnt signaling and is required for vertebrate gastrulation. *Development*, 133(21), 4219-4231.
- Schlange, T., Matsuda, Y., Lienhard, S., Huber, A., & Hynes, N. E. (2007). Autocrine WNT signaling contributes to breast cancer cell proliferation via the canonical WNT pathway and EGFR transactivation. *Breast Cancer Research*, 9(5), R63.
- Schnitt, S. J. (2010). Classification and prognosis of invasive breast cancer: from morphology to molecular taxonomy. *Modern Pathology*, 23, S60-S64.
- Semizarov, D., Frost, L., Sarthy, A., Kroeger, P., Halbert, D. N., & Fesik, S. W. (2003). Specificity of short interfering RNA determined through gene expression signatures. *Proceedings of the National Academy of Sciences*, 100(11), 6347-6352.
- Sheldahl, L. C., Slusarski, D. C., Pandur, P., Miller, J. R., Kuhl, M., & Moon, R. T. (2003). Dishevelled activates Ca²⁺ flux, PKC, and CamKII in vertebrate embryos. *Journal of Cell Biology*, 161(4), 769-777.
- Shelke, G., Lässer, C., Gho, Y., & Lötvall, J. (2014). Importance of exosome depletion protocols to eliminate functional and RNA-containing extracellular vesicles from fetal bovine serum. *Journal of Extracellular Vesicles*, 3.

- Singletary, S. E., Walsh, G., Vauthey, J.-N., Curley, S., Sawaya, R., Weber, K. L., Meric, F., & Hortobagyi, G. N. (2003). A role for curative surgery in the treatment of selected patients with metastatic breast cancer. *The Oncologist*, 8(3), 241-251.
- Slamon, D. J., Godolphin, W., Jones, L. A., Holt, J. A., Wong, S. G., Keith, D. E., Levin, W. J., Stuart, S. G., Udove, J., & Ullrich, A. (1989). Studies of the HER-2/neu proto-oncogene in human breast and ovarian cancer. *Science*, 244(4905), 707-712.
- Society, A. C. (2015). Hormone Therapy for Breast Cancer Retrieved 26 June 2015, 2015
- Stevens, C. A., & Qumsiyeh, M. B. (1995). Syndromal frontonasal dysostosis in a child with a complex translocation involving chromosomes 3, 7, and 11. *American Journal of Medical Genetics*, 55(4), 494-497.
- Sugimura, R., He, X. C., Venkatraman, A., Arai, F., Box, A., Semerad, C., Haug, J. S., Peng, L., Zhong, X. B., Suda, T., & Li, L. (2012). Noncanonical wnt signaling maintains hematopoietic Stem Cells in the niche. *Cell*, 150(2), 351-365.
- Sutton, L. P., & Rushlow, W. J. (2012). The dopamine D2 receptor regulates Akt and GSK-3 via Dvl-3. *International Journal of Neuropsychopharmacology*, 15(7), 965-979.
- Tauriello, D. V., Jordens, I., Kirchner, K., Slootstra, J. W., Kruitwagen, T., Bouwman, B. A., Noutsou, M., Rudiger, S. G., Schwamborn, K., Schambony, A., & Maurice, M. M. (2012). Wnt/beta-catenin signaling requires interaction of the Dishevelled DEP domain and C terminus with a discontinuous motif in Frizzled. *Proceedings of the National Academy of Sciences USA*, 109(14), E812-820. doi: 10.1073/pnas.1114802109
- Tian, X.-H., Hou, W.-J., Fang, Y., Fan, J., Tong, H., Bai, S.-L., Chen, Q., Xu, H., & Li, Y. (2013). XAV939, a tankyrase 1 inhibitor, promotes cell apoptosis in neuroblastoma cell lines by inhibiting Wnt/ β -catenin signaling pathway. *Journal of Experimental & Clinical Cancer Research*, 32(1), 1-9.
- Veeman, M. T., Slusarski, D. C., Kaykas, A., Louie, S. H., & Moon, R. T. (2003). Zebrafish prickles, a modulator of noncanonical Wnt/Fz signaling, regulates gastrulation movements. *Current Biology*, 13(8), 680-685.
- Vogel, C. L., Cobleigh, M. A., Tripathy, D., Gutheil, J. C., Harris, L. N., Fehrenbacher, L., Slamon, D. J., Murphy, M., Novotny, W. F., & Burchmore, M. (2002). Efficacy and safety of trastuzumab as a single agent in first-line treatment of HER2-overexpressing metastatic breast cancer. *Journal of Clinical Oncology*, 20(3), 719-726.

- Wallingford, J. B., & Habas, R. (2005). The developmental biology of Dishevelled: an enigmatic protein governing cell fate and cell polarity. *Development*, 132(20), 4421-4436.
- Wang, Y. (2009). Wnt/Planar cell polarity signaling: a new paradigm for cancer therapy. *Molecular Cancer Therapeutics*, 8(8), 2103-2109.
- Watanabe, J., Kushihata, F., Honda, K., Sugita, A., Tateishi, N., Mominoki, K., Matsuda, S., & Kobayashi, N. (2004). Prognostic significance of Bcl-xL in human hepatocellular carcinoma. *Surgery*, 135(6), 604-612.
- Wend, P., Runke, S., Wend, K., Anchondo, B., Yesayan, M., Jardon, M., Hardie, N., Loddenkemper, C., Ulasov, I., & Lesniak, M. S. (2013). WNT10B/ β -catenin signalling induces HMGA2 and proliferation in metastatic triple-negative breast cancer. *EMBO Molecular Medicine*, 5(2), 264-279.
- Wharton, K. A., Jr. (2003). Runnin' with the Dvl: proteins that associate with Dsh/Dvl and their significance to Wnt signal transduction. *Developmental Biology*, 253(1), 1-17.
- Winklbauer, R., Medina, A., Swain, R. K., & Steinbeisser, H. (2001). Frizzled-7 signalling controls tissue separation during *Xenopus* gastrulation. *Nature*, 413(6858), 856-860.
- Winter, C. G., Wang, B., Ballew, A., Royou, A., Karess, R., Axelrod, J. D., & Luo, L. (2001). *Drosophila* Rho-associated kinase (Drok) links Frizzled-mediated planar cell polarity signaling to the actin cytoskeleton. *Cell*, 105(1), 81-91.
- Wong, H.-C., Bourdelas, A., Krauss, A., Lee, H.-J., Shao, Y., Wu, D., Mlodzik, M., Shi, D.-L., & Zheng, J. (2003). Direct binding of the PDZ domain of Dishevelled to a conserved internal sequence in the C-terminal region of Frizzled. *Molecular Cell*, 12(5), 1251-1260.
- Wood, A. J., Smith, I. E., & Dowsett, M. (2003). Aromatase inhibitors in breast cancer. *New England Journal of Medicine*, 348(24), 2431-2442.
- Xu, W., & He, X. (2010). DEEP Insights through the DEP Domain. *Structure*, 18(10), 1223-1225.
- Yang, L., Wu, X., Wang, Y., Zhang, K., Wu, J., Yuan, Y., Deng, X., Chen, L., Kim, C., & Lau, S. (2011). FZD7 has a critical role in cell proliferation in triple negative breast cancer. *Oncogene*, 30(43), 4437-4446.
- Yang, Y., Jiao, L., Hou, J., Xu, C., Wang, L., Yu, Y., Li, Y., Yang, C., Wang, X., & Sun, Y. (2013). Dishevelled-2 silencing reduces androgen-dependent prostate tumor cell proliferation and migration and expression of Wnt-3a and matrix metalloproteinases. *Molecular Biology Reports*, 40(7), 4241-4250.

- Yin, X., Xiang, T., Li, L., Su, X., Shu, X., Luo, X., Huang, J., Yuan, Y., Peng, W., & Oberst, M. (2013). DACT1, an antagonist to Wnt/b-catenin signaling, suppresses tumor cell growth and is frequently silenced in breast cancer. *Breast Cancer*, 4, 5.
- Yokoyama, N., Yin, D., & Malbon, C. C. (2007). Abundance, complexation, and trafficking of Wnt/beta-catenin signaling elements in response to Wnt3a. *Journal of Molecular Signaling*, 2(11).
- Zhao, Y., Yang, Z. Q., Wang, Y., Miao, Y., Liu, Y., Dai, S. D., Han, Y., & Wang, E. H. (2010). Dishevelled-1 and dishevelled-3 affect cell invasion mainly through canonical and noncanonical Wnt pathway, respectively, and associate with poor prognosis in nonsmall cell lung cancer. *Molecular Carcinogenesis*, 49(8), 760-770.
- Zhu, Y., Tian, Y., Du, J., Hu, Z., Yang, L., Liu, J., & Gu, L. (2012). Dvl2-Dependent Activation of Daam1 and RhoA Regulates Wnt5a-Induced Breast Cancer Cell Migration. *PLoS One*, 7(5).



UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : Second Semester 2016/2017

TITLE OF THESIS / PROJECT REPORT :

EFFECTS OF SILENCING DISHEVELLED 1, 2 AND 3 mRNA EXPRESSION ON MDA-MB-231 CELL MIGRATION AND INVASION USING siRNAs

NAME OF STUDENT: SAMUEL KHOO LEON JUAN

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (✓)

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

OPEN ACCESS

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

PATENT

Embargo from _____ until _____
(date) (date)

Approved by:

(Signature of Student)
New IC No/ Passport No.:

Date :

(Signature of Chairman of Supervisory Committee)
Name:

Date :

[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted.]