



UNIVERSITI PUTRA MALAYSIA

***MOLECULAR IDENTIFICATION, PROBIOTIC CHARACTERIZATION,
ANTICANCER ASSESSMENT, AND ENCAPSULATION OF LACTIC AND
ACETIC ACID BACTERIA ISOLATED FROM IRANIAN TRADITIONAL
DAIRY PRODUCTS***

BABAK HAGHSHENAS

IB 2015 41



**MOLECULAR IDENTIFICATION, PROBIOTIC
CHARACTERIZATION, ANTICANCER ASSESSMENT, AND
ENCAPSULATION OF LACTIC AND ACETIC ACID
BACTERIA ISOLATED FROM IRANIAN TRADITIONAL
DAIRY PRODUCTS**

By

BABAK HAGHSHENAS

**Thesis Submitted to the School of Graduate Studies, Universiti
Putra Malaysia, in Fulfilment of the Requirements for the Degree of
Doctor of Philosophy**

March 2015

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATIONS

I wish to dedicate this work to my father who passed away during my study.

God bless his soul.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia, in fulfilment of the requirements for the degree of Doctor of Philosophy

**MOLECULAR IDENTIFICATION, PROBIOTIC CHARACTERIZATION,
ANTICANCER ASSESSMENT, AND ENCAPSULATION OF LACTIC AND
ACETIC ACID BACTERIA ISOLATED FROM IRANIAN TRADITIONAL
DAIRY PRODUCTS**

By

BABAK HAGHSHENAS

March 2015

Chairman: Associate Professor Norhafizah Abdullah, PhD

Faculty: Institute of Bioscience

Lactic and acetic acid bacteria (LAB and AAB) are generally recognized as safe microorganisms. Their increasing demand in food industry has motivate this present study to perform molecular identification, *in vitro* characterization, anticancer assessment and microencapsulation of probiotics from traditional dairy products produced in the west of Iran region. Cancer epidemiology showed lower cases recorded in this region compared with other areas in Iran which can be associated with the daily dietary intake of people living in these areas. Dairy products made from the west of Iran employs traditional methods involving no antibiotics in the animal feed and often contained higher acidity compared with commercial dairy products. Probiotic bioactive compounds have fewer side effects with low-cost when use in chemotherapy. Thus, the identification of probiotic bacteria exhibiting anticancer property isolated from dairy products for medicinal purposes is highly desirable.

The isolated bacteria from 5 types of traditional dairy products were identified and clustered into 5 genera (*Lactococcus*, *Leuconostoc*, *Lactobacillus*, *Enterococcus*, and *Acetobacter*) with 12 species, 4 subspecies, and 27 strains using combination of different molecular techniques (namely 16S rRNA sequencing, (GTG)₅-PCR, and ARDRA). *In vitro* characterization showed 11 isolates displaying high anti-pathogenic activity and antibiotic susceptibility. Their survival rates in the harsh gastrointestinal conditions were also high, calculated at >70%.

Cytotoxic assays indicated that extracted metabolites from *Acetobactersyzygii*38Lac, *Enterococcus durans* 39C, and *Lactococcus lactis subsp. lactis* 44Lac displayed high anticancer activity against AGS, MCF-7, HT29, and HeLa cancer cell lines but not on normal cell lines. Pronase assay revealed that proteins contained in the metabolites play a key role in anticancer properties of these isolates. Incidence of apoptosis was detected

in the isolates on the basis of the morphological assessments via fluorescent microscope and flow cytometry.

To improve delivery and prolong the survival rate of probiotic in the gastrointestinal tract, encapsulation approach was investigated using a simple extrusion method. Nine formulations using different blend of alginate with psyllium or fenugreek were designed and investigated aiming at improving the encapsulation efficiency. Results showed 2% (w/v) alginate, 1.5% (w/v) alginate +0.5% (w/v) psyllium, and 1.5% (w/v) alginate + 0.5% (w/v) fenugreek formulations, with >98% encapsulation efficiency, high stability during 28 days of storage and displayed high survival rates in harsh gastrointestinal conditions (>80 %), which were significantly ($P < 0.01$) higher than the non-encapsulated bacteria (>62%). Incorporation of 0.5% (w/v) fenugreek and psyllium into 1.5% (w/v) alginate gel resulted in enhanced encapsulation properties. For psyllium-alginate formulation, additional properties such as high growth-stimulating effects of probiotic and smaller bead sizes were observed.

Acetobacter strains were isolated for the first time from traditional dairy products, which is a new finding of this study. Their extracted metabolites displayed high cytotoxicity effects on various cancer cell lines comparable to commercial cancer drug, Taxol. Development of a new encapsulated probiotic formulation using fenugreek-alginate blend was also a success. The formulation showed significant improvement in the delivery of probiotic in gastrointestinal system.

In conclusion, probiotic bacteria were successfully isolated and characterized from Iranian traditional dairy products. These probiotics exhibited antimicrobial activity, antibiotic resistance and anticancer activity thus these make bacteria potential candidates for the development of nutraceutical products. In addition, the study demonstrates the improved survival and delivery of probiotics in the gastrointestinal conditions system by using encapsulation technique.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**Pengenalan Molekul, Pencirian Probiotik, Penilaian Anti
Kanser, dan Pengkapsulan Bakteria Asid Laktik dan Asid
Asetik yang Diasingkan Daripada Produk Tenusu
Tradisional Iran**

Oleh

BABAK HAGHSHENAS

Mac 2015

Pengerusi: Profesor Madya Norhafizah Abdullah, PhD

Fakulti: Institut Biosains

Bakteria asid laktik dan asid asetik (LAB dan AAB) secara umumnya dikenali sebagai mikroorganisma yang selamat. Permintaan yang semakin meningkat dalam industri makanan mendorong penyelidikan ini dijalankan bagi mengenalpasti secara molekul, pencirian *in vitro*, penilaian anti kanser dan pemikrokapsulan probiotik daripada produk tenusu tradisional yang dihasilkan di kawasan barat Iran. Epidemiologi penyakit kanser menunjukkan jumlah kes yang lebih rendah telah dicatatkan di rantau ini berbanding kawasan lain di Iran, di mana ia boleh berhubungkaitkan dengan diet pemakanan harian penduduk di kawasan ini. Produk tenusu dari kawasan ini dihasilkan dengan menggunakan kaedah tradisional yang tidak melibatkan penggunaan antibiotik dalam makanan haiwan dan mempunyai keasidan yang lebih tinggi berbanding produk-produk tenusu komersial. Sebatian bioaktif probiotik mempunyai kesan sampingan yang kurang dengan kos rendah apabila digunakan dalam kemoterapi. Oleh itu, mengenalpasti bakteria probiotik mempamerkan sifat antikanser yang dipencil dari produk tenusu untuk tujuan perubatan adalah sangat diperlukan.

Bakteria yang dipencilkan daripada lima jenis hasil tenusu tradisional telah berjaya dikenal pasti dan dikelompokkan kepada lima genera (*Lactococcus*, *Leuconostoc*, *Lactobacillus*, *Enterococcus*, dan *Acetobacter*) dengan 12 spesis, 4 subspesis dan 27 strain dengan menggunakan kombinasi teknik molekul yang berbeza (penjujukan 16S rRNA, (GTA)₅-PCR, dan ARDRA). Pencirian *in vitro* telah menunjukkan 11 pencilan yang mempunyai sifat anti patogen yang tinggi dan rentangan terhadap antibiotik.

Pengasaan sitotoksik terhadap metabolit yang diekstrak dari *Acetobacter syzygii* 38Lac, *Enterococcus durans* 39C dan *Lactococcus lactis subsp. lactis* 44Lac menunjukkan aktiviti anti kanser yang tinggi terhadap sel-sel kanser AGS, MCF-7, HT29 dan HeLa tetapi tiada kesan ke atas sel normal. Pengasaan pronase telah menunjukkan bahawa mekanisme anti-kanser adalah berpunca daripada kandungan protin yang terkandung dalam metabolit yang dihasilkan oleh pencilan-pencilan

tersebut. Kejadian apoptosis telah dikenal pasti berlaku di dalam pencilan-pencilan ini, dengan berdasarkan ke atas penilaian morfologi melalui mikroskop pendarfluor dan teknik pengalir sitometri.

Untuk meningkatkan penyampaian dan kelangsungan hidup probiotik dalam alam persekitaran saluran pencernaan, teknik pengkapsulan telah dijalankan melalui kaedah penyemperitan yang mudah. Sebanyak 9 jenis formulasi gel menggunakan kepekatan yang berbeza antara alginat dengan psyllium atau halba telah direka dan dinilai demi menghasilkan peningkatan pengkapsulan yang efisien. Keputusan menunjukkan 2% (w/v) alginat, 1.5% (w/v) alginat + 0.5 (w/v) psyllium, dan 1.5% (w/v) alginate + 0.5% (w/v) halba telah menghasilkan pengkapsulan yang efisien sebanyak >98%, kestabilan yang tinggi semasa 28 hari penyimpanan dan memaparkan kadar hidup yang tinggi dalam alam persekitaran saluran pencernaan yang mencabar (>80%), dengan signifikan yang tinggi ($P < 0.01$) berbanding dengan probiotik yang tidak dikapsulkan (<62%). Pengabungan 0.5% (w/v) halba dan psyllium ke dalam 1.5% (w/v) gel alginat menghasilkan penambahbaikan terhadap ciri-ciri pengkapsulan. Bagi formulasi psyllium-alginat, ciri tambahan iaitu penambahan kesan stimulasi terhadap pertumbuhan probiotik dan saiz kapsul yang kecil telah terhasil.

Penemuan baharu daripada kajian ini ialah strain *Acetobacter* telah berjaya diasingkan buat kali pertama daripada produk tenusu tradisional. Metabolit yang diekstrak daripada strain ini menunjukkan sifat sitotoksiti tinggi terhadap pelbagai sel kanser setanding dengan ubat kanser komersial, Taxol. Pengkapsulan dengan gabungan halba-alginat sebagai matriks baru telah diformulasi dengan jayanya. Formulasi produk ini menunjukkan peningkatan yang ketara dalam penghantaran probiotik di dalam sistem pencernaan.

Kesimpulannya, bakteria probiotik telah berjaya dipencilkan dan dicirikan dari produk tenusu tradisional dari negara Iran. Probiotik yang dipencilkan mempunyai sifat antimikrob, rintangan antibiotik dan anti-kanser, seterusnya menjadikan mereka calon yang berpotensi untuk dibangunkan sebagai produk nutraseutikal. Di samping itu, kajian ini telah menunjukkan penambahbaikan kelangsungan hidup dan penghantaran probiotik dalam sistem saluran pencernaan dengan mengadaptasi teknik pengkapsulan.

ACKNOWLEDGEMENTS

At the end of this step of my graduate period has allowed for a fragment of reflection, and the many dears who have contributed to both, my work, and my life during of this period of time, because this research project would not have been possible without the support of many people. First and foremost, I wish to express my full thanks and sincere gratitude to my dear supervisor, Dr. Norhafizah Abdullah for their invaluable suggestions, beneficial advices and their endless supports. My sincere gratitude also goes to Prof. Dr. Rozita Rosli and Dr. Dayang Radiah Awang Biak, who served on my thesis committee and kindly provided valuable advice and suggestions for this work and my thesis. I would like to acknowledge and have my best regards to Dr. Ahmad Yari Khosroushahi for their friendly collaboration during of my experimental research. I also would like to acknowledge Mr. Abolfazl Barzegari in Tabriz University and Mr. Yousef Nami in Universiti Putra Malaysia or their invaluable helps. Last but not least I wish to avail myself of this opportunity, express a sense of gratitude and love to my friends and my beloved parents for their support, strength, and help for everything during of my life. I wish to acknowledge my gratitude to all lecturers, officials and other staff members in Universiti Putra Malaysia.

Babak Haghshenas
March 2015

This thesis was submitted to the senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Norhafizah Abdullah, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Rozita Rosli, PhD

Professor
Institute of Bioscience
Universiti Putra Malaysia
(Member)

Dayang Radiah Awang Biak, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Ahmad Yari Khosroushahi, PhD

Assistant Professor
Faculty of Pharmaceutical
Tabriz University of Medical Science of Iran
(Member)

BUJANGBIN KIMHUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- This thesis is my original work;
- Quotations, illustrations and citations have been duly referenced;
- This thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- Intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- Written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- There is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No: Babak Haghshenas, GS26631

Declaration by Members of Supervisory Committee

This is to confirm that:

- The research conducted and the writing of this thesis was under our supervision;
- Supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to

Name of Chairman of Supervisory Committee:
Associate Professor Norhafizah Abdullah

Signature: _____

Name of Member of Supervisory Committee:
Professor Rozita Rosli

Signature: _____

Name of Member of Supervisory Committee:
Dr. Dayang Radiah Awang Biak

Signature: _____

Name of Member of Supervisory Committee:
Dr. Ahmad Yari Khosroushahi

Signature: _____

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF APPENDICES	xix
LIST OF ABBREVIATIONS	xx
 CHAPTER	
1 INTRODUCTION	1
1.1 Background	1
1.2 Research Problems	2
1.3 Objectives	3
1.4 Scope of Work	3
1.5 Thesis Layout	6
 2 LITERATURE REVIEW	 7
2.1 Human Gastrointestinal Microbiota (Gut Flora)	7
2.2 Probiotic	7
2.2.1 Diversity of Probiotics	8
2.2.2 Dosage and Formulations of Probiotic Products in Market	8
2.2.3 The Health Effects of Probiotics	9
2.3 Probiotic Belongs to <i>Lactobacillus</i> Genus	10
2.3.1 Classification of <i>Lactobacillus</i>	10
2.3.2 <i>Lactobacillus</i> Strains in Dairy Products	11
2.3.3 Antibiotic Resistant Species of <i>Lactobacillus</i>	11
2.3.4 Anti Pathogen Activity of <i>Lactobacillus</i> Strains and its Mechanisms	12
2.4 <i>Enterococcus</i> Genus	13
2.4.1 History and Characteristics	13
2.4.2 Products and Habitats	13
2.4.3 Antibiotic-Resistant Species of <i>Enterococcus</i> Genus	13
2.4.4 Clinical Health Promoting Effects of <i>Enterococcus</i>	14
2.4.5 <i>Enterococcus</i> Anti pathogen Mechanisms (Bacteriocins)	14
2.4.6 <i>Enterococcus</i> as Probiotic	15
2.5 Other Probiotic Associated Genus	15
2.5.1 <i>Leuconostoc</i> Genus	15
2.5.1.1 Habitat	15
2.5.1.2 The Antibiotic Resistant Strains	16
2.5.1.3 The Anti-Microbial Mechanisms	16
2.5.1.4 <i>Leuconostoc</i> as Probiotic	16
2.5.2 <i>Lactococcus</i> Genus	17

2.5.3	<i>Acetobacter</i> Genus	17
2.6	Molecular Identification Methods	18
2.6.1	Introduction	18
2.6.2	Genetic Probe	18
2.6.3	Fluorescent In Situ Hybridization (FISH)	19
2.6.4	Ribotyping	19
2.6.5	Pulsed-Field Gel Electrophoresis (PFGE)	20
2.6.6	Species Specific PCR and Sequencing	21
2.6.7	Sequences of Specific Genes	22
2.6.8	Randomly Amplified Polymorphic DNA (RAPD)	22
2.6.9	Repetitive Bacterial DNA Elements PCR (REP-PCR)	22
2.6.10	Amplified Fragment Length Polymorphism (AFLP)	23
2.6.11	Amplified Ribosomal DNA Restriction Analysis (ARDRA)	23
2.6.12	Temperature and Denaturing Gradient Gel Electrophoresis (TGGE and DGGE)	23
2.7	Probiotic and Cancer	23
2.7.1	<i>In Vivo</i> Anti-Cancer Evidences	23
2.7.2	<i>In Vitro</i> Anti-Cancer Evidences	24
2.7.3	Colorectal Cancer (CRC)	24
2.7.4	Anti-Cancer Mechanisms	25
2.8	Microencapsulation of Probiotics	25
2.8.1	Microencapsulation Methods	26
2.8.1.1	Encapsulation by Freeze Drying and Spray Drying Methods	26
2.8.1.2	Encapsulation by Emulsion Method	27
2.8.1.3	Encapsulation by Extrusion Method	27
2.8.1.4	Advantages and Disadvantages of Emulsion and Extrusion	27
2.8.2	Supporting Materials	28
2.9	Traditional Dairy Products of Kermanshah	28
2.9.1	Curd	29
2.9.2	Tarkhineh	30
2.9.3	Shiraz	31
2.9.4	Yogurt	32
2.9.5	Cheese	32
2.10	Concluding Remarks	33

3 ISOLATION, MOLECULAR IDENTIFICATION AND CLUSTERING OF ISOLATED LACTIC AND ACETIC ACID BACTERIA 35

3.1	Introduction	35
3.2	Materials and Methods	35
3.2.1	Sampling of Dairy Products	35
3.2.2	Isolation of Harsh Condition-Tolerant Strains	36
3.2.3	Morphological Assay	37
3.2.4	Biochemical Assay	37
3.2.5	DNA Extraction	37
3.2.6	Amplification and Sequencing of 16S rRNA	38
3.2.7	ARDRA Analysis	38

	3.2.8	(GTG) ₅ -PCR fingerprinting	38
	3.2.9	Data Analysis	39
3.3		Results and Discussion	39
	3.3.1	Sampling and Isolation of Harsh Condition-Tolerant Strains	39
	3.3.2	16S-rRNA Pattern Analysis	41
	3.3.3	Identification by 16S rRNA Sequencing	43
	3.3.4	Clustering by ARDRA and Data Analysis	46
	3.3.5	(GTG) ₅ -PCR fingerprinting and data analysis	50
3.4		Conclusion	55
4		GASTROINTESTINAL CHARACTERIZATION OF LACTIC AND ACETIC ACIDBACTERIA ISOLATED FROM IRANIAN FERMENTEDDAIRY PRODUCTS	57
	4.1	Introduction	57
	4.2	Materials and Methods	57
	4.2.1	Bacterial Strains and Growth Conditions	57
	4.2.2	Low pH Tolerance Assessment	58
	4.2.3	Tolerance to Bile Salt	58
	4.2.4	Antimicrobial Activity Assay	59
	4.2.5	Antibiotic Susceptibility Assay	59
	4.3	Results and Discussion	60
	4.3.1	Low pH Tolerance Assessment	60
	4.3.2	Tolerance to Bile Salt	62
	4.3.3	Antimicrobial Activity against Various Pathogens	64
	4.3.4	Antibiotic Susceptibility Assays against Clinically Important Antibiotics	71
	4.4	Conclusion	75
5		CYTOTOXIC ASSESSMENT AND DETERMINATION OF CELL DEATH MODES INDUCED BY THE EXTRACTED METABOLITES OF ISOLATED PROBIOTICBACTERIA ON VARIOUS CANCER AND NORMAL CELL LINES	76
	5.1	Introduction	76
	5.2	Materials and Methods	77
	5.2.1	Bacterial Growth Condition	77
	5.2.2	Preparation of cell free culture supernatant (CFCS)	77
	5.2.3	The Growth Condition of Cell Lines	77
	5.2.4	MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyl Tetrazolium Bromide) Assay	78
	5.2.5	Apoptosis Assessment by DAPI Staining and Fluorescent Microscopy	78
	5.2.6	Flow Cytometric Quantification of Apoptotic Cells	79
	5.2.7	Statistical Analysis	79
	5.3	Results and Discussion	80
	5.3.1	Pre Screening Cytotoxic Test on Various Cancer Cell Lines	80
	5.3.2	Screening Cytotoxic Test on Various Cancer Cell Lines	89
	5.3.3	Cytotoxic Test on Normal Cell Line	93
	5.3.4	Pronase Assessment	94

5.3.5	Apoptosis Assessment by Fluorescent Microscopy	96
5.3.6	Flow Cytometric Quantification of Apoptotic Cells	98
5.4	Conclusion	102
6	MICROENCAPSULATION OF PROBIOTIC BACTERIA <i>L. plantarum</i> 15HN AND <i>E. durans</i> 39C USING HERBAL-BASED POLYMERS BLENDS	103
6.1	Introduction	103
6.2	Materials and Methods	104
6.2.1	Materials	104
6.2.2	Preparation and Amplification of Bacteria	104
6.2.3	Extraction of Psyllium Gel	104
6.2.4	Extraction of Fenugreek Gel	105
6.2.5	Probiotic Microencapsulation Using Extrusion Technique	105
6.2.6	Morphological Analysis of the Encapsulated Probiotic Microcapsules	105
6.2.7	Encapsulation Efficiency (EE)	105
6.2.8	Probiotic Cell Viability at the Low pH Condition	106
6.2.9	Probiotic Cell Viability at the High Bile Salt Condition	106
6.2.10	Storage Stability	106
6.2.11	Release of Bacteria	107
6.2.12	Statistical Analyses	107
6.3	Results and Discussion	107
6.3.1	Morphological Analysis of Microencapsulated Probiotic Beads	107
6.3.2	Encapsulation Efficiency (EE) of Prepared Beads	112
6.3.3	Viability at Low pH Condition	113
6.3.4	Viability at High Bile Salt Condition	116
6.3.5	Storage Stability	119
6.3.6	Release of Microencapsulated Probiotic Beads in Colonic Solution	121
6.4	Conclusion	123
7	CONCLUSION AND RECOMMENDATIONS FOR FUTURE STUDY	125
7.1	Conclusion	125
7.2	Future study	126
	REFERENCES	127
	APPENDICES	168
	BIODATA OF STUDENT	175
	LIST OF PUBLICATIONS	176

LIST OF TABLES

Table	Page
2.1 Microorganisms used as probiotics	8
2.2 Beneficial effects of probiotics on health	9
3.1 Biochemical analysis (gram staining and catalase test) of isolated bacteria	40
3.2 Isolated LAB and AAB and their traditional dairy origins according to 16S-rRNA pattern analysis	42
3.3 Sequencing results of 27 isolated LAB and AAB and their dairy origin	45
3.4 Clustering of isolates after comparison the experimental ARDRA band patterns with the virtually ARDRA results	50
3.5 The various attributes of isolated bacteria	55
4.1 Survival rates of isolated LAB and AAB after 3 h incubation at pH 2.5	61
4.2 Survival rates of isolated LAB and AAB after 3 h incubation at 0.3% bile	63
4.3 Origin, incubation condition, and infectious diseases of tested pathogens	65
4.4 The inhibitory effect of isolated LAB strains against pathogens	67
4.5 The inhibitory effect of isolated AAB strains against pathogens	69
4.6 Antibiotic susceptibility profiles of isolated LAB and AAB	73
5.1 Probiotic strains and their dairy origin	80
5.2 The IC ₅₀ values of isolated probiotic secretions on various cancer cell lines for 12 hours	87
5.3 The IC ₅₀ values of isolated probiotic secretions on various cancer cell lines for 24 hours	88
5.4 The IC ₅₀ values of isolated probiotic secretions on various cancer cell lines for 48 hours	88
6.1 Compositions of study gels	108
6.2 Size, shape and number of protective layers in the prepared formulations	111
6.3 Log CFU/mL before and after encapsulation process and encapsulation	113

efficiency (%) of *E. durans* 39C and *L. plantarum* 15HN in various gel formulations

- 6.4 Mean counts (log CFU/g) for *E. durans* 39C and *L. plantarum* 15HN 114
before and after low pH (pH 1.8) treatments and the bacterial survival
rates in different prepared formulations
- 6.5 Mean counts (log CFU/g) for *E. durans* 39C and *L. plantarum* 15HN 118
before and after bile salt (0.5% (w/v) oxgall) treatments and the
bacterial survival rates in different prepared formulations



LIST OF FIGURES

Figure	Page
1.1 Schematic diagram represented the flowchart of the study in this thesis	5
2.1 Diagrammatic representation of genetic probe method	19
2.2 Diagrammatic representation of ribotyping	20
2.3 Diagrammatic representation of PFGE	21
2.4 Mountain climate and forest in Kermanshah province	29
2.5 Diagrammatic figure of dried curd preparation	30
2.6 Diagrammatic figure of tarkhineh preparation	31
2.7 Diagrammatic figure of shiraz preparation	32
2.8 Diagrammatic figure of traditional dairy product preparation	33
3.1 Topographic map of Kermanshah province with five divided sections	36
3.2 Low pH and high bile salt resistant single colonies	40
3.3 Morphological evaluation of isolated single colonies by fluorescent microscopy (Olympus BX61)	41
3.4 Agarose gel electrophoresis of amplified 16S rRNA products with approximately 1500bp length	42
3.5 Agarose gel electrophoresis of amplified 16S rRNA products with approximately 1500 bp length belonged to the representative LAB and AAB isolates	43
3.6 The percentage distribution of isolated LAB and AAB strains for dairy products	45
3.7 Agarose gel electrophoresis of amplified 16S rRNA products after restriction with <i>Pst</i> I restriction enzyme	47
3.8 Generated dendrogram by ARDRA band patterns of isolated strains restricted with <i>Pst</i> I enzyme	48
3.9 Virtual ARDRA patterns for 16S rRNA of selected strains in Gene bank after digesting with <i>Pst</i> I restriction enzyme by ApE software	49
3.10 (GTG) ₅ -PCR banding patterns of a total of 18 reference LAB and AAB isolates, representing all validly described LAB and AAB species (27 strains)	52

3.11	The combination results of (GTG) ₅ -PCR with 16S rRNA sequencing and ARDRA band patterns of a total of 18 reference strains and 27 LAB and AAB isolates	54
4.1	Antagonistic activity of isolated strains against the pathogenic bacteria by formation of inhibition zones	66
4.2	The susceptibility spectrum of routinely used antibiotics against the isolated LAB and AAB	72
5.1	The cytotoxic time-dependent graphs for the secreted metabolites of eleven Isolated LAB and AAB on AGS cancer cell line	81
5.2	The cytotoxic effects of isolated <i>L. lactis subsp. Lactis</i> 44 Lac secretions on different cancer cell lines at three time points 12, 24, and 48 h	83
5.3	The cytotoxic effects of isolated <i>E. durans</i> 39C secretions on different cancer cell lines at three time points 12, 24, and 48 h	84
5.4	The cytotoxic effects of isolated <i>A. syzygii</i> 38Lac secretions on different cancer cell lines at three time points 12, 24, and 48 h	85
5.5	The cytotoxic effects of isolated <i>E. faecalis</i> 13C secretions on different cancer cell lines at three time points 12, 24, and 48 h	86
5.6	The cytotoxic effects of <i>L. lactis subsp. Lactis</i> 44Lac secretions (40 µg/mL) and Taxol on several cancer cell lines for 48 h.	90
5.7	The cytotoxic effects of <i>E. durans</i> 39C secretions (40 µg/mL) and Taxol on several cell lines for 48 h	91
5.8	The cytotoxic effects of <i>A. syzygii</i> 38Lac secretions (40 µg/mL) and Taxol on several cancer cell lines for 48 h	92
5.9	The cytotoxic effects of <i>A. syzygii</i> 38Lac, <i>E. durans</i> 39C, and <i>L. lactis subsp. lactis</i> 44Lac secretions (40 µg/mL) and Taxol on the HUVEC normal cell line for 48 h	94
5.10	The cytotoxic effects of <i>A. syzygii</i> 38Lac, <i>E. durans</i> 39C, <i>L. lactis subsp. lactis</i> 44Lac, and their pronase treated secretions (40 µg/mL) on the AGS cell line for 48 h	95
5.11	Fluorescent photomicrographs after incubation of <i>A. syzygii</i> 38Lac secreted metabolite (40 µg/mL) on AGS cell line	96
5.12	Fluorescent photomicrographs after incubation of <i>E. durans</i> 39C secreted metabolite (40 µg/mL) on AGS cell line	97
5.13	Fluorescent photomicrographs after incubation of <i>L. lactis subsp. lactis</i> 44Lac secreted metabolite (40 µg/mL) on AGS cell line	98

5.14	Flow cytometry assessment of AGS cancer cell line after 24 hours incubation with the extracted metabolites (40 µg/mL) of isolated bacteria	100
5.15	Flow cytometry assessment of AGS cancer cell line after 48 hours incubation with the extracted metabolites (40 µg/mL) of isolated bacteria	101
6.1	Morphological image of alginate-psyllium encapsulated beads by optical microscopy (Olympus BX61)	109
6.2	Morphological image of alginate encapsulated beads by optical Microscopy (Olympus BX61)	109
6.3	Morphological image of alginate-fenugreek encapsulated beads by optical microscopy (Olympus BX61)	110
6.4	Morphological image of encapsulated beads containing entrapped probiotic bacteria	110
6.5	Survival rates of <i>E. durans</i> 39C and <i>L. plantarum</i> 15HN encapsulated with various gel formulations as compared with free cells after incubation in low pH condition (pH1.8)	116
6.6	Survival rates of <i>E. durans</i> 39C and <i>L. plantarum</i> 15HN encapsulated with various gel formulations as compared with free cells after incubation in bile salt condition (0.5% w/v oxgall)	119
6.7	Storage stability of free and encapsulated <i>E. durans</i> 39C with different gel formulations during 28 days storage at 4°C	120
6.8	Storage stability of free and encapsulated <i>L. plantarum</i> 15HN with different gel formulations during 28 days storage at 4°C.	120
6.9	Releasing rates of encapsulated <i>E. durans</i> 39C with different gel formulations at colonic pH solution in every one hour and for 12 hours	122
6.10	Releasing rates of encapsulated <i>L. plantarum</i> 15HN with different gel formulations at colonic pH solution in every one hour and for 12 hours	122

LIST OF APPENDICES

Appendix	Page
A.1 Glucose -Tris-EDTA (GTE) buffer	168
A.2 Proteinase K (20 mg/mL)	169
A.3 Phenol: chloroform: isoamyl alcohol (25:24:1)	169
A.4 Sodium acetate 3 M (pH 5.2)	169
A.5 RNase (10 mg/mL)	169
A.6 Tris-Acetic acid-EDTA (TAE) buffer	169
B.1 Morphological evaluation of isolated bacteria by fluorescent microscope (Olympus BX61)	170
B.2 Cytotoxic time-dependent graphs for the secreted metabolites of eleven isolated LAB and AAB on various cancer cell line	172
C.1 Gene Ruler 1 kb DNA Ladder 250 to 10,000 bp (Thermo Scientific, USA)	174

LIST OF ABBREVIATIONS

°C	Degree Celsius
%	Percent
4T1	Mouse breast cancer cell line
AAB	Acetic acid bacteria
AFLP	Amplified Fragment Length Polymorphism
AGS	Human gastric epithelial cancer cell line
ARDRA	Amplified Ribosomal DNA Restriction Analysis
ARE	Antibiotic-Resistant Enterococci
ATCC	America Type Culture Collection
BLAST	Basic Local Alignment Search Tool
bp	Base pair
BSA	Bovine serum albumin
CAP	Cellulose Acetate Phthalate
CFU	Colony forming unit
Co ₂	Carbon dioxide
DAPI	4',6-diamidino-2-phenylindole
DGGE	Denaturing Gradient Gel Electrophoresis
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleic acid
EDTA	Ethylene diaminetetraacetic acid
EFSA	European Food Safety Authority
ERIC-PCR	Enterobacterial Repetitive Intergenic Consensus PCR
FAO	Food and Agricultural Organization of the United States
FBS	Fetal bovine serum
FISH	Fluorescent In Situ Hybridization
G+C	Guanine plus cytosine
GTG-PCR	Polymerase chain reaction by GTG primer
HeLa	Human cervix cancer cell line
HT29	Human colon cancer cell line
g	Gravity force
g	Gram
GTE	Glucose-Tris-EDTA
h	Hour
HCL	Hydrochloric acid
ITS	Intergenic spacer region
kb	Kilo base pair
kDa	Kilo Dalton
L929	Mouse fibroblast cell line
LAB	Lactic acid bacteria
Lb	Lactobacillus
Lc	Lactococcus
LDH	Lactate dehydrogenase
LDL	Low-density lipoprotein
M	Molar
MgCl ₂	Magnesium chloride
min	Minute
Mm	Milimolar
mg	Miligram
mL	Mililiter

mm	Milimeter
MRS	Man Rogosa and Sharpe
MRSA	Methicillin-Resistant <i>Staphylococcus aureus</i>
MTT	3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyl Tetrazolium Bromide
MYP	Mannitol-Egg-yolk-Polymyxine
N	Normality
Na	Sodium
NaCl	Sodium chloride
NaOH	Sodium hydroxide
NCBI	National Center for Biotechnology Information
nm	Nanometer
NTSYS-PC	Numerical Taxonomy Analysis Program Package
PBS	Potassium buffered saline
PCR	polymerase Polymerase chain reaction
PFGE	Pulsed-Field Gel Electrophoresis
RAPD	Randomly amplified polymorphic DNA
rDNA	Ribosomal deoxyribonucleic acid
REA	Restriction enzyme analysis
Rep-PCR	Repetitive Extragenic Palindromes-PCR
rRNA	Ribosomal ribonucleic acid
RFLP	Restriction fragment length polymorphism
RNA	Ribonucleic acid
RNase	Ribonuclease
rpm	Round per minute
sdH ₂ O	Sterile distilled water
S	Streptococcus
SE	Standard error
sp.	Species
subsp.	Subspecies
TAE	Tris-acetate-EDTA
TGGE	Temperature Gradient Gel Electrophoresis
tRNA	Transfer RNA
TUNEL	Terminal deoxynucleotidyltransferase mediated dUTP
UPGMA	Unweighted Pair- Group Method by Arithmetic Average
V	Volt
v/v	Volume per volume
VRE	Vancomycin-Resistant Enterococci
w/v	Weight per volume
µg	Microgram
µL	Microliter
µg/mL	Microgram per milliliter
µM	Micromolar
XG	Xanthan Gum

CHAPTER 1

INTRODUCTION

1.1 Background

The term of “probiotics” was first introduced by Lilly and Stillwell as the microorganism’s active factors stimulating other microorganism’s growth and possessing the beneficial impacts on the host health when they are consumed in the sufficient amounts (Guarner *et al.*, 2008; Lilly and Stillwell, 1965). Probiotics significantly affect the bioavailability of nutrients in the human body (Young and Huffman, 2003) by facilitating the absorption of magnesium and calcium from milk proteins, digesting lactose, and producing folate and B vitamins. Probiotics also prevent irritable bowel syndrome (Barouei *et al.*, 2009; Parkes, 2010) and inflammatory effects on bowel disease (Geier *et al.*, 2007). These microorganisms also control antibiotic-related diarrhea (Friedman, 2012), elicit protective effects on vaginal infections (Reid and Bocking, 2003), and decrease cholesterol levels (Guarner *et al.*, 2008). As anti-cancer agents, probiotics can inactivate pro-carcinogenic compounds, particularly in colon cancer (Serban, 2013; Zhu *et al.*, 2011).

The majority of probiotics include lactic acid bacteria (LAB), (Guarner *et al.*, 2008) but isolated acetic acid bacteria (AAB) from safe sources, such as fermented dairy products, also can be introduced as probiotics because of their strong fermentation and acidification activities and vitamin C production (González and Mas, 2011). The LAB including the genera such as *Oenococcus*, *Lactococcus*, *Streptococcus*, *Enterococcus*, *Leuconostoc*, *Lactobacillus* and *Pediococcus*. Among the LAB, the *Lactobacillus* and *Enterococcus* species are commonly consumed as probiotics (Biradar *et al.*, 2005; Giraffa *et al.*, 2010; Guarner *et al.*, 2008). On the other hand, *Acetobacter* is one of the most common genera with commercial importance in the AAB group (Sengun and Karabiyikli, 2011). *Acetobacter* species, particularly *A. aceti* and *A. syzygii*, show the industrial importance and can be introduced as probiotic (Cleenwerck and De Vos, 2008; De Vuyst *et al.*, 2008; Gullo *et al.*, 2006; Lefeber *et al.*, 2011; Sengun and Karabiyikli, 2011).

The microbiota of traditional dairy products is supposed to be dominated by *Lactobacillus* and *Lactococcus* species (Kore *et al.*, 2012; Tabatabaee *et al.*, 2012; Tafvizi and Tajabadi, 2012; Veera Jothi *et al.*, 2012). But, the biodiversity of LAB species in fermented dairy products is variable and region specific. For example, in Armada cheese (Spanish goat's cheese); the predominant *Lactobacilli* are *L. casei* ssp. *casei*, *L. plantarum* and, *L. brevis* (Herreros *et al.*, 2003). But, in Batzos cheese (traditional Greek goat cheese), *L. paracasei* spp. *tolerans*, *L. plantarum*, and *L. sake* are the dominant species (Psoni *et al.*, 2003). Meanwhile, in Minas Frescal cheese (Brazilian fresh cheese) the predominant species is *L. acidophilus* (Lollo *et al.*, 2012).

Probiotics are an important part of the complex world of health-promoting food. The best-known example of which is yogurt. Bacteria found in traditional dairy products, such as yogurt, kefir, and fermented milk, are usually not found in the human intestines. These bacteria do not colonize the intestine, but are flushed and eliminated quickly from the body. Therefore, the introduction of potential probiotic cells from

native food sources is a crucial step to determine their efficacy. The diverse climate in Kermanshah Province, Iran particularly in rural areas, allows the villagers to produce different traditional dairy products. These products can be the ideal dairy source for the selection of new potential probiotic bacteria because of their production methods in semi-wild conditions and use of raw milk from cows and sheep.

1.2 Research Problems

Fermented dairy products such as curd, tarkhineh, shiraz, yogurt and cheese in the west of Iran (Kermanshah Province) can be explored for the ideal dairy source in the selection of new potential probiotic bacteria. The different production methods in semi-wild conditions and the use of raw milk from cows or sheep provide the advantage on the selection. The drying time and fermented processes are different among these dairy products. Moreover, they have various acidity values. Therefore, new strains with high probiotic capability can be isolated from these products.

The beneficial properties of probiotic bacteria are strain specific, and a majority of them belong to the lactic acid bacteria (LAB) group, which requires same growth conditions and nutrition (Dubernet *et al.*, 2002). Identification and differentiation up to the genus level between these closely related bacteria through traditional phenotypic and biochemical methods, such as sugar fermentation, is impossible and does not provide clear results (Gevers *et al.*, 2001). Recently, rapid, accurate, and practical molecular identification techniques, such as repetitive sequence-based PCR ((GTG)₅-PCR) (Gevers *et al.*, 2001), specific gene sequencing (Vaugien *et al.*, 2002), 16S rRNA sequencing (Gueimonde *et al.*, 2004), ribotyping with specific probes (Singh *et al.*, 2009), random amplified polymorphic DNA (RAPD) (Singh *et al.*, 2009), and amplified ribosomal DNA restriction analysis (ARDRA) (Satokari *et al.*, 2003), have been designed (Singh *et al.*, 2009; Ward and Roy, 2005). However, these methods individually do not create strong results for differentiation up to the strain levels. Thus, an effective combination must be considered.

A probiotic must have a number of characteristics to be effective in improving host health, such as resistance to gastrointestinal conditions, antipathogen activities, and high susceptibility to antibiotics (Biradar *et al.*, 2005; Surawicz, 2003). Most probiotics are sensitive to low acid or high bile salt conditions in the digestive system and are eliminated after consumption. Probiotics also carry antibiotic-resistant genes, which if transferred to other probiotics or pathogenic bacteria, can create high resistance to a wide variety of antibiotics (Mathur and Singh, 2005). Traditional dairy products from the west of Iran are prepared based on the traditional methods employed in this region, wherein no antibiotics are used in animal food and higher acidity of products compared with commercial dairy products. In this regard, new strains with high probiotic capability can be isolated from these products.

Although, several effects that are beneficial to health have been attributed to probiotic bacteria, the most interesting and controversial property remains to be anticancer activity (de LeBlanc *et al.*, 2010). The vast majority of studies in this area deal with protective effects against colon cancer (Rafter, 2003). A wealth of indirect evidence based largely on laboratory studies is available regarding the anticancer effects of probiotic bacteria, which may be divided into the following categories: *in vitro* studies, animal studies, epidemiological studies, and human dietary intervention studies. The

mechanisms by which probiotics may inhibit different cancer cell lines are still poorly understood. However, several potential mechanisms will be discussed in the literature. Cancer epidemiology showed lower cases recorded in rural areas of this region (Kermanshah Province), compared with other areas in Iran (Mirmomeni *et al.*, 2009; Najafi *et al.*, 2011). This difference could be related to the daily dietary intake of the areas. Through *in vitro* cytotoxic and apoptotic studies in different human cancer cell lines, effective strains can be isolated, and their anticancer mechanisms can be investigated.

Probiotic microorganisms show low survival rates in dairy products and the gastrointestinal tract (Vidhyalakshmi *et al.*, 2009). A vast variety of coating materials (e.g., starch, vegetable gums, gelatins, fats, and glyceride derivatives) can be applied using different microencapsulation methods (e.g., freeze drying (Heidebach *et al.*, 2010; Otero *et al.*, 2007), spray drying (Avila-Reyes *et al.*, 2013; Salar-Behzadi *et al.*, 2013), gel formation (Vidhyalakshmi *et al.*, 2009), co-extrusion (Shinde *et al.*, 2013), extrusion (Khan *et al.*, 2013; Lotfipour *et al.*, 2012), emulsion (Mantzouridou *et al.*, 2012), fluid bed processing (Stummer *et al.*, 2010; Stummer *et al.*, 2012), and matrix entrapment (Reyed, 2006) to increase the stability of probiotic bacteria. In addition, microencapsulation is now developed to lend protection against unwanted conditions, as well as to increase tolerance against strong acids and undesirable reactions upon storage. Therefore, this method is a novel and interesting technology in the probiotics market (Sohail *et al.*, 2011). Herbal gels, such as fenugreek and psyllium, which are cheaper than alginate and have been used for the treatment of gastrointestinal disease, have been demonstrated to be suitable candidates for encapsulation of probiotic bacteria.

1.3 Objectives

The objectives of this study are:

- 1) To isolate, identify and cluster the probiotic strains from five types of traditional Iranian dairy products using combination of 3 molecular methods, which are 16S rRNA sequencing, (GTG)₅-PCR and ARDRA.
- 2) To characterize and assess the acid and bile tolerance, anti-microbial potential and antibiotic susceptibility of the isolated probiotic strains.
- 3) To investigate the *in vitro* cytotoxic assay and morphological apoptosis assessment of probiotic metabolites on cancer and normal cell lines by MTT assay, fluorescent microscopy and flow cytometry.
- 4) To develop a delivery system on the best candidate of probiotic strains using microencapsulation method with alginate, psyllium, and fenugreek gels by extrusion technique and characterize its probiotics performance in acid, bile and storage time.

1.4 Scope of Work

The use of probiotic products as medical supplements has successfully spread throughout the world especially in Europe and Asia. *Lactobacillus* species are the most

famous group of probiotics which are isolated from fermented dairy products. Other probiotics isolated from dairy products are *Bifidobacteria*, *Bacillus*, *Streptococci*, some strains of *E. coli*, and yeast (*Saccharomyces cerevisiae*) (Biradar *et al.*, 2005; Guarner *et al.*, 2008). Recent discoveries on the benefits of probiotic populations on human health motivate further interest on probiotic communities and their functions (Moore *et al.*, 2011). For this study, 200 samples were collected randomly from 5 traditional dairy products, including cheese, yogurt, curd, shiraz, and tarkhineh, from retailers in different parts of Kermanshah Province (west of Iran). Samples of traditional dairy products were obtained across five wide area of Kermanshah province, Iran. Random clustering was chosen as a suitable method for their selection. According to this method, the area was divided to five sections (North, South, East, West and center). In each section, 2 villages and in each village 4 rancher families were chosen randomly for sampling the products.

The lactic and acetic acid bacteria were isolated from traditional dairy products for the first time in the west of Iran (Kermanshah province) by the combination of three molecular techniques ((GTG)₅-PCR, 16S rRNA sequencing and ARDRA). *Acetobacter* species normally are isolated from ethanol rich sources such as wine and beers. But, for the first time, *Acetobacter* strains were isolated and identified from dairy products.

The most of researches and studies on probiotics have been focused on their anti-colorectal properties (Commane *et al.*, 2005; de LeBlanc *et al.*, 2010; Rafter, 2003). But, the extracted metabolites of isolated bacteria displayed the high cytotoxic effects on various cancer cell lines same as Taxol.

The fenugreek polymer exhibits several prebiotic effects and has been utilized in animal husbandry and nutritional supplements (Brummer *et al.*, 2003). This gum has health-promoting effects and has been prescribed by many practitioners for the treatment of diabetes and for the regulation of cholesterol level (Brummer *et al.*, 2003). In addition, its stability and emulsifying activity (Chang *et al.*, 2011; Işıklı and Karababa, 2005) make this gel a suitable stabilizer for the food industry. In this study, fenugreek for the first time was blended with the alginate polymer as a matrix for probiotic formulation. This polymer offers added advantages of being prebiotic for the enhancement of probiotic bacterial growth in the gastrointestinal environment. Hence, its incorporation with alginate can be introduced as new and unique delivery carriers for the oral administration of probiotics. The flowchart of the study is presented in Figure 1.1.




Figure1.1. Schematic diagram represented the flowchart of the study in this thesis.

1.5 Thesis Layout

In the first chapter, samples from traditional dairy products were collected to isolate probiotic bacteria belonging to LAB and AAB groups. To discover new strains, identification and discrimination of isolated bacteria were performed up to the strain levels by combining three molecular methods (i.e., 16S rRNA sequence analysis, ARDRA, and (GTG)₅-PCR banding patterns comparison).

In the second chapter, some probiotic characteristics of the isolated strains, such as tolerance to extra bile salts and low acidic conditions, antibiotic susceptibility, and antimicrobial activity against the human pathogens, were investigated by *in vitro* assessments.

In the third chapter, MTT assay was employed to assess the anticancer property of the extracted metabolites from candidate probiotic strains by using different human cancer and normal cell lines. The results were compared with untreated (negative control) and Taxol-treated cells lines (positive controls). The anticancer mechanisms of the probiotics were characterized by comparing the normal treated with pronase-treated, untreated, and Taxol-treated cells. In addition, some morphological apoptosis characteristics, such as membrane blebbing, disruption of membrane integrity, and DNA fragmentation, were assessed using fluorescent microscopy and flow cytometry.

In the last chapter, two effective probiotic strains (*E. durans* 39C and *L. plantarum* 15HN), which displayed low viability in harsh gastric conditions, were encapsulated through extrusion method with supporting materials including alginate, psyllium, and fenugreek. Morphological characteristics and encapsulation properties were assessed and analyzed using PSWE Statistics 18 software.

REFERENCES

- Agerholm-Larsen, L., Bell, M. L., Grunwald, G. K., & Astrup, A. (2000). The effect of a probiotic milk product on plasma cholesterol: a meta-analysis of short-term intervention studies. *European Journal of Clinical Nutrition*, 54(11), 856-860.
- Agrawal, A., Houghton, L. A., Morris, J., Guyonnet, D., Goupil Feuillerat, N., Schlumberger, A., Jakob, S., & Whorwell, P. J. (2008). T1395 fermented milk containing the probiotic *Bifidobacterium animalis*, DN-173 010 (FM) improves abdominal distension, bloating and transit in irritable bowel syndrome with constipation (IBS-C). *Gastroenterology*, 134(4, Supplement 1), A-546.
- Ahmad, I., Sharma, J., & Ahmad, F. (2004). Isolation and characterization of resistance traits of indigenous strains of *Acetobacter diazotrophicus* associated with sugarcane. *Sugar Tech*, 6(1), 41-46.
- Ahmadova, A., Todorov, S. D., Choiset, Y., Rabesona, H., Mirhadi Zadi, T., Kuliye, A., Franco, B. D. G. M., Chobert, J.M., & Haertlé, T. (2013). Evaluation of antimicrobial activity, probiotic properties and safety of wild strain *Enterococcus faecium* AQ71 isolated from Azerbaijani Motal cheese. *Food Control*, 30(2), 631-641.
- Akagi, C. L., & Lesser, D. (2000). A review of chemotherapeutic agents and their oral manifestaions. *The California Dental Hygienists Association Journal*, 16(2), 9-14.
- Al Jarousha, A. M. K., El Qouqa, I. A., El Jadba, A. H. N., & Al Afifi, A. S. (2008). An outbreak of *Serratia marcescens* septicaemia in neonatal intensive care unit in Gaza City, Palestine. *Journal of Hospital Infection*, 70(2), 119-126.
- Albufera, U., Bhugaloo-Vial, P., Issack, M. I., & Jaufeerally-Fakim, Y. (2009). Molecular characterization of *Salmonella* isolates by REP-PCR and RAPD analysis. *Infection, Genetics and Evolution*, 9(3), 322-327.
- Alegria, Á., Delgado, S., Rocés, C., López, B., & Mayo, B. (2010). Bacteriocins produced by wild *Lactococcus lactis* strains isolated from traditional, starter-free cheeses made of raw milk. *International Journal of Food Microbiology*, 143(1-2), 61-66.
- Ali, A. M., Ismail, N. H., Mackeen, M. M., Yazan, L. S., Mohamed, S. M., Ho, A. S., & Lajis, N. H. (2000). Antiviral, cytototoxic and antimicrobial activities of anthraquinones isolated from the roots of morinda elliptica. *Pharmaceutical Biology*, 38(4), 298-301.
- Alonso, R., Picon, A., Gaya, P., Fernández-García, E., & Nuñez, M. (2012). Effect of high-pressure treatment of ewe raw milk curd at 200 and 300 MPa on characteristics of Hispánico cheese. *Journal of Dairy Science*, 95(7), 3501-3513.
- Altonsy, M. O., Andrews, S. C., & Tuohy, K. M. (2010). Differential induction of apoptosis in human colonic carcinoma cells (Caco-2) by atopobium, and commensal, probiotic and enteropathogenic bacteria: Mediation by the

- mitochondrial pathway. *International Journal of Food Microbiology*, 137(2–3), 190-203.
- Amann, R., & Ludwig, W. (2000). Ribosomal RNA-targeted nucleic acid probes for studies in microbial ecology. *FEMS Microbiology Reviews*, 24(5), 555-565.
- Amann, R. I., Ludwig, W., & Schleifer, K. H. (1995). Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiological Review*, 59, 143-169.
- Anal, A. K., & Singh, H. (2007). Recent advances in microencapsulation of probiotics for industrial applications and targeted delivery. *Trends in Food Science & Technology*, 18(5), 240-251.
- Anandan, S., Dey, A., Deb, S. M., Kumar, S., & Harbola, P. C. (1999). Effect of curds as probiotic supplement on performance of Cheghu crossbred kids. *Small Ruminant Research*, 32(1), 93-96.
- Anderson, H. A., Englert, R., Gursel, I., & Shacter, E. (2002). Oxidative stress inhibits the phagocytosis of apoptotic cells that have externalized phosphatidylserine. *Cell Death and Differentiation*, 9, 616-625.
- Andiç, S., Tunçtürk, Y., & Boran, G. (2015). Chapter 28 - Changes in volatile compounds of cheese. In V. Preedy (Ed.), *Processing and impact on active components in food* (pp. 231-239). San Diego: Academic Press.
- Annan, N. T., Borza, A. D., & Hansen, L. T. (2008). Encapsulation in alginate-coated gelatin microspheres improves survival of the probiotic *Bifidobacterium adolescentis* 15703T during exposure to simulated gastro-intestinal conditions. *Food Research International*, 41(2), 184-193.
- Anonymous. (1992). Yoghurt and probiotics. *Choice*, 32(1), 32–35.
- Antoine, M. J. (2009). Micro-encapsulation of probiotics. *Prebiotics and Probiotics Science and Technology*, 805-823. 10.1007/978-0-387-79058-9_20.
- Aplin, A. E., Howe, A. K., & Juliano, R. L. (1999). Cell adhesion molecules, signal transduction and cell growth. *Current Opinion in Cell Biology*, 11(6), 737-744.
- Aquino Esperanza, J. A., Aguirre, M. V., Aispuru, G. R., Lettieri, C. N., Juaristi, J. A., Alvarez, M. A., & Brandan, N. C. (2008). *In vivo* 5-fluorouracil-induced apoptosis on murine thymocytes: involvement of FAS, Bax and Caspase3. *Cell Biology and Toxicology*, 24(5), 411-422.
- Aragón, F., Carino, S., Perdigón, G., & LeBlanc, A. d. M. d. (2014). The administration of milk fermented by the probiotic *Lactobacillus casei* CRL 431 exerts an immunomodulatory effect against a breast tumor in a mouse model. *Immunobiology*, 219(6), 457-464.
- Argyri, A. A., Zoumpopoulou, G., Karatzas, K. A. G., Tsakalidou, E., Nychas, G. J. E., Panagou, E. Z., & Tassou, C. C. (2013). Selection of potential probiotic lactic acid

- bacteria from fermented olives by *in vitro* tests. Food Microbiology, 33(2), 282-291.
- Aroutcheva, A., Gariti, D., Simon, M., Shott, S., Faro, J., Simoes, J. A., Gurguis, A., & Faro, S. (2001). Defense factors of vaginal *Lactobacilli*. American Journal of Obstetrics and Gynecology, 185(2), 375-379.
- Asbahani, A. E., Miladi, K., Badri, W., Sala, M., Addi, E. H. A., Casabianca, H., Mousadik, A. E., Hartmann, D., Jilale, A., Renaud, F. N. R., & Elaissari, A. (2015). Essential oils: From extraction to encapsulation. International Journal of Pharmaceutics, 483(1–2), 220-243.
- Aslim, B., & Beyatli, Y. (2004). Antibiotic resistance and plasmid DNA contents of *Streptococcus thermophilus* strains isolated from Turkish yogurts. Turkish Journal of Veterinary and Animal Sciences, 28(2), 257-263.
- Aslim, B., Yuksekdog, Z. N., Sarikaya, E., & Beyatli, Y. (2005). Determination of the bacteriocin-like substances produced by some lactic acid bacteria isolated from Turkish dairy products. LWT - Food Science and Technology, 38(6), 691-694.
- Avila-Reyes, S. V., Garcia-Suarez, F. J., Jiménez, M. T., Martín-Gonzalez, M. F. S., & Bello-Perez, L. A. (2013). Protection of *L. rhamnosus* by spray-drying using two prebiotics colloids to enhance the viability. Carbohydrate Polymers, 4(102), 423-430.
- Babich, H., & Borenfreund, E. (1991). Cytotoxicity of T-2 toxin and its metabolites determined with the neutral red cell viability assay. Applied and Environmental Microbiology, 57(7), 2101-2103.
- Baele, M., Vanechoutte, M., Verhelst, R., Vancanneyt, M., Devriese, L. A., & Haesebrouck, F. (2002). Identification of *Lactobacillus* species using tDNA-PCR. Journal of Microbiological Methods, 50(3), 263-271.
- Baker, M. A., Cerniglia, G. J., & Zaman, A. (1990). Microtitre plate assay for the measurement of glutathione and glutathione disulfide in large numbers of biological samples. Analytical Biochemistry, 190 (2), 360-365.
- Bakry, M. H., Bahga, t. M. R., Hassaan, A. E., Hala, A. A., & A, A. E. (2013). Structure analysis and antitumor activity of the exopolysaccharide from probiotic *Lactobacillus plantarum* NRRL B- 4496 *in vitro* and *in vivo*. Journal of Applied Sciences Research, 9(1), 425-434.
- Barouei, J., Adams, M. C., & Hodgson, D. M. (2009). Prophylactic role of maternal administration of probiotics in the prevention of irritable bowel syndrome. Medical Hypotheses, 73(5), 764-767.
- Baskić, D., Popović, S., Ristić, P., & Arsenijević, N. N. (2006). Analysis of cycloheximide-induced apoptosis in human leukocytes: Fluorescence microscopy using annexin V/propidium iodide versus acridin orange/ethidium bromide. Cell Biology International, 30(11), 924-932.

- Bauer, A. W., Kirby, W. M., Sherris, J. C., & Turck, M. (1966). Antibiotic susceptibility testing by a standardized single disk method. *American Journal of Clinical Pathology*, 45(4), 493-496.
- Bautista-Gallego, J., Arroyo-López, F. N., Rantsiou, K., Jiménez-Díaz, R., Garrido-Fernández, A., & Cocolin, L. (2013). Screening of lactic acid bacteria isolated from fermented table olives with probiotic potential. *Food Research International*, 50(1), 135-142.
- Begley, M., Gahan, C. G. M., & Hill, C. (2005). The interaction between bacteria and bile. *FEMS Microbiology Reviews*, 29(4), 625-651.
- Ben Amor, K., Vaughan, E. E., & de Vos, W. M. (2007). Advanced molecular tools for the identification of lactic acid bacteria. *The Journal of Nutrition*, 137(3), 741-747.
- Ben Salah, R., Trabelsi, I., Ben Mansour, R., Lassoued, S., Chouayekh, H., & Bejar, S. (2012). A new *Lactobacillus plantarum* strain, TN8, from the gastro intestinal tract of poultry induces high cytokine production. *Anaerobe*, 18(4), 436-444.
- Bengmark, S. (1998). Immunonutrition: role of biosurfactants, fiber, and probiotic bacteria. *Nutrition*, 14(7-8), 585-594.
- Beresford, T. P., Fitzsimons, N. A., Brennan, N. L., & Cogan, T. M. (2001). Recent advances in cheese microbiology. *International Dairy Journal*, 11(4-7), 259-274.
- Bernardeau, M., Vernoux, J. P., Henri-Dubernet, S., & Guéguen, M. (2008). Safety assessment of dairy microorganisms: The *Lactobacillus* genus. *International Journal of Food Microbiology*, 126(3), 278-285.
- Berthier, F., Beuvier, E., Dasen, A., & Grappin, R. (2001). Origin and diversity of mesophilic *lactobacilli* in Comté cheese, as revealed by PCR with repetitive and species-specific primers. *International Dairy Journal*, 11(4-7), 293-305.
- Bhardwaj, A., Gupta, H., Kapila, S., Kaur, G., Vij, S., & Malik, R. K. (2010). Safety assessment and evaluation of probiotic potential of bacteriocinogenic *Enterococcus faecium* KH 24 strain under in vitro and in vivo conditions. *International Journal of Food Microbiology*, 141(3), 156-164.
- Biasucci, G., Benenati, B., Morelli, L., Bessi, E., & Boehm, G. n. (2008). Cesarean delivery may affect the early biodiversity of intestinal bacteria. *The Journal of Nutrition*, 138, 1796-1800.
- Bibe, R., & Sandine, W. E. (1992). Acetic, propionic, and lactic acids of starter culture bacteria as biopreservatives. In *Food Biopreservatives of Microbial Origin*. B. Ray and M. Daeschel (eds.), pp 103-136. Boca Raton, FL: CRC Press.
- Biffi, A., Coradini, D., Larsen, R., Riva, L., & Di Fronzo, G. (1997). Antiproliferative effect of fermented milk on the growth of a human breast cancer cell line. *Nutrition and Cancer*, 28(1), 93-99.

- Bigwood, T., Hudson, J. A., Cooney, J., McIntyre, L., Billington, C., Heinemann, J. A., & Wall, F. (2012). Inhibition of *Listeria monocytogenes* by *Enterococcus mundtii* isolated from soil. *Food Microbiology*, 32(2), 354-360.
- Binns, C., & Lee, M. K. (2010). The use of probiotics to prevent diarrhea in young children attending child care centers: A review. *Journal of Experimental & Clinical Medicine*, 2(6), 269-273.
- Biradar, S. S., Bahagvati, S. T., & Shegunshi, B. (2005). Probiotics and antibiotics: A brief overview. *The Internet Journal of Nutrition and Wellness*, 2(1), 1-5.
- Borgia, M., Sepe, N., Brancato, V., & Borgia, R. (1982). A controlled clinical study on *Streptococcus faecium* preparation for the prevention of side reactions during longterm antibiotic treatments. *Current Therapeutic Research*, 31, 265-271.
- Bouchier-Hayes, L., Munoz-Pinedo, C., Connel, S., & Green, D. R. (2008). Measuring apoptosis at the single cell level. *Methods*, 44, 222-228.
- Bourget, N., Simonet, J.-M., & Decaris, B. (1993). Analysis of the genome of the five *Bifidobacterium breve* strains: plasmid content, pulsed-field gel electrophoresis genome size estimation and rrn loci number. *FEMS Microbiology Letters*, 110(1), 11-20.
- Bravo, D., Rodríguez, E., & Medina, M. (2009). Nisin and lactacin 481 coproduction by *Lactococcus lactis* strains isolated from raw ewes' milk. *Journal of Dairy Science*, 92(10), 4805-4811.
- Brown, J. M., & Attardi, L. D. (2005). The role of apoptosis in cancer development and treatment response. *Nature Reviews Cancer*, 5(3), 231-237.
- Brown, T., Hipps, N. A., Easteal, S., Parry, A., & Evans, J. A. (2014). Reducing domestic food waste by lowering home refrigerator temperatures. *International Journal of Refrigeration*, 40(0), 246-253.
- Brummer, Y., Cui, W., & Wang, Q. (2003). Extraction, purification and physicochemical characterization of fenugreek gum. *Food Hydrocolloids*, 17(3), 229-236.
- Bunselmeyer, B., & Buddendick, K. (2010). Chapter 19 - Probiotics and prebiotics—Prevention and therapy in atopic eczema. In R. R. Watson & V. R. Preedy (Eds.), *Bioactive Foods in Promoting Health* (pp. 279-292). Boston: Academic Press.
- Çakır, I., Coşkun, H., Akoğlu, I., İşleyen, M., Kıralan, M., & Bayrak, A. (2009). Introducing a traditional dairy product Keş: Chemical, microbiological, and sensorial properties and fatty acid composition. *Journal of Food, Agriculture & Environment*, 7(3&4), 116-119.
- Capurso, G., Marignani, M., & Fave, G. D. (2006). Probiotics and the incidence of colorectal cancer: when evidence is not evident. *Digestive and Liver Disease*, 38(2), 277-282.

- Carmen Collado, M., & Hernández, M. (2007). Identification and differentiation of *Lactobacillus*, *Streptococcus* and *Bifidobacterium* species in fermented milk products with *bifidobacteria*. *Microbiological Research*, 162(1), 86-92.
- Casado Muñoz, M. d. C., Benomar, N., Lerma, L. L., Gálvez, A., & Abriouel, H. (2014). Antibiotic resistance of *Lactobacillus pentosus* and *Leuconostoc pseudomesenteroides* isolated from naturally-fermented Aloreña table olives throughout fermentation process. *International Journal of Food Microbiology*, 172, 110-118.
- Casalta, E., & Montel, M.-C. (2008). Safety assessment of dairy microorganisms: The *Lactococcus* genus. *International Journal of Food Microbiology*, 126(3), 271-273.
- Casarotti, S. N., Monteiro, D. A., Moretti, M. M. S., & Penna, A. L. B. (2014). Influence of the combination of probiotic cultures during fermentation and storage of fermented milk. *Food Research International*, 59, 67-75.
- Cebrián, R., Baños, A., Valdivia, E., Pérez-Pulido, R., Martínez-Bueno, M., & Maqueda, M. (2012). Characterization of functional, safety, and probiotic properties of *Enterococcus faecalis* UGRA10, a new AS-48-producer strain. *Food Microbiology*, 30(1), 59-67.
- Chagnaud, P., Machinis, K., Coutte, L. A., Marecat, A., & Mercenier, A. (2001). Rapid PCR-based procedure to identify lactic acid bacteria, application to six common *Lactobacillus* species. *Journal of Microbiological Methods*, 44, 139-148.
- Champagne, C. P., & Fustier, P. (2007). Microencapsulation for the improved delivery of bioactive compounds into foods. *Current Opinion in Biotechnology*, 18(2), 184-190.
- Chan, E. S. (2011). Preparation of Ca-alginate beads containing high oil content: Influence of process variables on encapsulation efficiency and bead properties. *Carbohydrate Polymers*, 84(4), 1267-1275.
- Chandramouli, V., Kailasapathy, K., Peiris, P., & Jones, M. (2004). An improved method of microencapsulation and its evaluation to protect *Lactobacillus* spp. in simulated gastric conditions. *Journal of Microbiological Methods*, 56(1), 27-35.
- Chang, Y. C., Tsai, C. Y., Lin, C. F., Wang, Y. C., Wang, I. K., & Chung, T. C. (2011). Characterization of tetracycline resistance *lactobacilli* isolated from swine intestines at western area of Taiwan. *Anaerobe*, 17(5), 239-245.
- Chang, Y. H., Cui, S. W., Roberts, K. T., Ng, P. K. W., & Wang, Q. (2011). Evaluation of extrusion-modified fenugreek gum. *Food Hydrocolloids*, 25(5), 1296-1301.
- Charlier, C., Even, S., Gautier, M., & Le Loir, Y. (2008). Acidification is not involved in the early inhibition of *Staphylococcus aureus* growth by *Lactococcus lactis* in milk. *International Dairy Journal*, 18(2), 197-203.
- Charteris, W. P., Kelly, P. M., Morelli, L., & Collins, J. K. (1998). Development and application of an *in vitro* methodology to determine the transit tolerance of

- potentially probiotic *Lactobacillus* and *Bifidobacterium* species in the upper human gastrointestinal tract. *Journal of Applied Microbiology*, 84, 759-768.
- Chávarri, M., Marañón, I., Ares, R., Ibáñez, F. C., Marzo, F., & Villarán, M. d. C. (2010). Microencapsulation of a probiotic and prebiotic in alginate-chitosan capsules improves survival in simulated gastro-intestinal conditions. *International Journal of Food Microbiology*, 142(1-2), 185-189.
- Chen, M. J., & Chen, K. N. (2008). Encapsulation of probiotics in alginate systems. *Controlled Release Society*, 25(2), 7-9.
- Cibik, R., Lepage, E., & Tailliez, P. (2000). Molecular diversity of *Leuconostoc mesenteroides* and *Leuconostoc citreum* Isolated from traditional French cheeses as revealed by RAPD fingerprinting, 16S rDNA sequencing and 16S rDNA fragment amplification. *Systematic and Applied Microbiology*, 23(2), 267-278.
- Cizeikiene, D., Juodeikiene, G., Paskevicius, A., & Bartkiene, E. (2013). Antimicrobial activity of lactic acid bacteria against pathogenic and spoilage microorganism isolated from food and their control in wheat bread. *Food Control*, 31(2), 539-545.
- Clark, P. A., & Martin, J. H. x. (1994). Selection of *bifidobacteria* for use as dietary adjuncts in cultured dairy foods: III – Tolerance to simulated bile of human stomachs. *Cultured Dairy Products Journal*, 29, 18-21.
- Cleenwerck, I., & De Vos, P. (2008). Polyphasic taxonomy of acetic acid bacteria: An overview of the currently applied methodology. *International Journal of Food Microbiology*, 125(1), 2-14.
- CLSI. (2007). Performance standards for antimicrobial susceptibility testing; Seventeenth informational supplement (Vol. 27). Wayne, PA: Clinical and laboratory standards institute.
- Coeuret, V., Gueguen, M., & Vernoux, J. P. (2004). Numbers and strains of *lactobacilli* in some probiotic products. *International Journal of Food Microbiology*, 97(2), 147-156.
- Commane, D., Hughes, R., Shortt, C., & Rowland, I. (2005). The potential mechanisms involved in the anti-carcinogenic action of probiotics. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 591(1-2), 276-289.
- Condon, S. (1987). Responses of lactic acid bacteria to oxygen. *FEMS Microbiology Letters*, 46(3), 269-280.
- Connor, M. O. (Ed.). (2011). *Ulcerative colitis - treatments, special populations and the future*: InTech.
- Conrad, P. B., Miller, D. P., Cielenski, P. R., & de Pablo, J. J. (2000). Stabilization and preservation of *Lactobacillus acidophilus* in saccharide matrices. *Cryobiology*, 41(1), 17-24.

- Conway, P. L., Gorbach, S. L., & Goldin, B. R. (1987). Survival of lactic acid bacteria in the human stomach and adhesion to intestinal cells. *Journal of Dairy Science*, 70(1), 1-12.
- Corsetti, A., Lavermicocca, P., Morea, M., Baruzzi, F., Tosti, N., & Gobbetti, M. (2001). Phenotypic and molecular identification and clustering of lactic acid bacteria and yeasts from wheat (species *Triticum durum* and *Triticum aestivum*) sourdoughs of Southern Italy. *International Journal of Food Microbiology*, 64(1-2), 95-104.
- Corsetti, A., & Valmorri, S. (2011). Lactic acid bacteria | *Lactobacillus* spp.: *Lactobacillus plantarum*. In W. F. Editor-in-Chief: John (Ed.), *Encyclopedia of Dairy Sciences* (Second Edition) (pp. 111-118). San Diego: Academic Press.
- Daglioglu, O. (2000). Tarhana as a traditional Turkish fermented cereal food. Its recipe, production and composition. *Nahrung/Food* 44(2), 85-88.
- Danielsen, M., & Wind, A. (2003). Susceptibility of *Lactobacillus* spp. to antimicrobial agents. *International Journal of Food Microbiology*, 82(1), 1-11.
- Dartsch, D. C., Schaefer, A., Boldt, S., Kolch, W., & Marquardt, H. (2002). Comparison of anthracycline-induced death of human leukemia cells: Programmed cell death versus necrosis. *Apoptosis*, 7, 537-548.
- Davies, E. A., & Delves-Broughton, J. (1999). Bacteriocins | Nisin. In K. R. Editor-in-Chief: Richard (Ed.), *Encyclopedia of Food Microbiology* (pp. 191-198). Oxford: Elsevier.
- De Kwaadsteniet, M., Todorov, S. D., Knoetze, H., & Dicks, L. M. T. (2005). Characterization of a 3944 Da bacteriocin, produced by *Enterococcus mundtii* ST15, with activity against Gram-positive and Gram-negative bacteria. *International Journal of Food Microbiology*, 105(3), 433-444.
- de LeBlanc, A. d. M., Bibas Bonet, M. E., LeBlanc, J. G., Sesma, F., & Perdígón, G. (2010). Chapter 29 - Probiotics in cancer prevention. In W. Ronald Ross & R. P. Victor (Eds.), *Bioactive Foods in Promoting Health* (pp. 497-511). Boston: Academic Press.
- De Moreno De Leblanc, A., & Perdígón, G. (2010). The application of probiotic fermented milks in cancer and intestinal inflammation. *Proceedings of the Nutrition Society*, 69(3), 421-428.
- De, S., & Robinson, D. (2003). Polymer relationships during preparation of chitosan-alginate and poly-l-lysine-alginate nanospheres. *Journal of Controlled Release*, 89(1), 101-112.
- De Vero, L., & Giudici, P. (2008). Genus-specific profile of acetic acid bacteria by 16S rDNA PCR-DGGE. *International Journal of Food Microbiology*, 125(1), 96-101.
- De Vuyst, L., Camu, N., De Winter, T., Vandemeulebroecke, K., Van de Perre, V., Vancanneyt, M., De Vos, P., & Cleenwerck, I. (2008). Validation of the (GTG)₅-

- rep-PCR fingerprinting technique for rapid classification and identification of acetic acid bacteria, with a focus on isolates from Ghanaian fermented cocoa beans. *International Journal of Food Microbiology*, 125(1), 79-90.
- De Vuyst, L., Foulquié Moreno, M. R., & Revets, H. (2003). Screening for enterocins and detection of hemolysin and vancomycin resistance in *enterococci* of different origins. *International Journal of Food Microbiology*, 84(3), 299-318.
- de Waard, R., Garssen, J., Bokken, G. C. A. M., & Vos, J. G. (2002). Antagonistic activity of *Lactobacillus casei* strain Shirota against gastrointestinal *Listeria monocytogenes* infection in rats. *International Journal of Food Microbiology*, 73(1), 93-100.
- Del Piano, M., Morelli, L., Strozzi, G. P., Allesina, S., Barba, M., Deidda, F., Lorenzini, P., Ballaré, M., Montino, F., Orsello, M., Sartori, M., Garelo, E., Carmagnola, S., Pagliarulo, M., & Capurso, L. (2006). Probiotics: from research to consumer. *Digestive and Liver Disease*, 38(2), 248-255.
- Dellaglio, F., Torriani, S., & Felis, G. E. (2004). Reclassification of *Lactobacillus cellobiosus* Rogosa, et al. 1953 as *Lactobacillus fermentum* Beijerinck 1901. *International Journal of Systematic and Evolutionary Microbiology*, 54, 809-812.
- Deng, W., Xi, D., Mao, H., & Wanapat, M. (2008). The use of molecular techniques based on ribosomal RNA and DNA for rumen microbial ecosystem studies: A review. *Molecular Biology Report*, 35, 265-274.
- Devirgiliis, C., Barile, S., Caravelli, A., Coppola, D., & Perozzi, G. (2010). Identification of tetracycline- and erythromycin-resistant Gram-positive cocci within the fermenting microflora of an Italian dairy food product. *Journal of Applied Microbiology*, 109(1), 313-323.
- Deye, G., Lewis, J., Patterson, J., & Jorgensen, J. (2003). A case of *Leuconostoc ventriculitis* with resistance to carbapenem antibiotics. *Oxford Journals*, 37(6), 869-870.
- Dijkshoorn, L., & Towner, K. (2001). 1 - An introduction to the generation and analysis of microbial typing data. In D. Lenie, J. T. Kevin, K. J. T. Marc StruelensA2 - Lenie Dijkshoorn & S. Marc (Eds.), *New Approaches for the Generation and Analysis of Microbial Typing Data* (pp. 1-30). Amsterdam: Elsevier Science B.V.
- Doleyres, Y., & Lacroix, C. (2005). Technologies with free and immobilised cells for probiotic bifidobacteria production and protection. *International Dairy Journal*, 15(10), 973-988.
- Domig, K. J., Mayrhofer, S., Zitz, U., Mair, C., Petersson, A., Amtmann, E., Mayer, H. K., & Kneifel, W. (2007). Antibiotic susceptibility testing of *Bifidobacterium thermophilum* and *Bifidobacterium pseudolongum* strains: Broth microdilution vs. agar disc diffusion assay. *International Journal of Food Microbiology*, 120(1-2), 191-195.

- Doulgeraki, A. I., Pramateftaki, P., Argyri, A. A., Nychas, G.-J. E., Tassou, C. C., & Panagou, E. Z. (2013). Molecular characterization of lactic acid bacteria isolated from industrially fermented Greek table olives. *LWT - Food Science and Technology*, 50(1), 353-356.
- Du Toit, W. J., & Lambrechts, M. G. (2002). The enumeration and identification of acetic acid bacteria from South African red wine fermentations. *International Journal of Food Microbiology*, 74(1-2), 57-64.
- Dubernet, S., Desmasures, N., & Guéguen, M. (2002). A PCR-based method for identification of *Lactobacilli* at the genus level. *FEMS Microbiology Letters*, 214(2), 271-275.
- Dubey, D., Rath, S., Sahu, M. C., Rout, S., Debata, N. K., & Padhy, R. N. (2013). A report on infection dynamics of inducible clindamycin resistance of *Staphylococcus aureus* isolated from a teaching hospital in India. *Asian Pacific Journal of Tropical Biomedicine*, 3(2), 148-153.
- Edwards, U., Rogall, T., Blocker, H., Emde, M., & Bottger, E. C. (1989). Isolation and direct complete nucleotide determination of entire genes. Characterization of a gene coding for 16S ribosomal RNA. *Nucleic Acids Research*, 17, 7843-7853.
- Ehrmann, M. A., & Vogel, R. F. (2005). Taxonomic note '*Lactobacillus pastorianus*' (Van Laer, 1892) a former synonym for *Lactobacillus paracollinoides*. *Systematic and Applied Microbiology*, 28, 54-56.
- Ejtahed, H. S., Mohtadi-Nia, J., Homayouni-Rad, A., Niafar, M., Asghari-Jafarabadi, M., Mofid, V., & Akbarian-Moghari, A. (2011). Effect of probiotic yogurt containing *Lactobacillus acidophilus* and *Bifidobacterium lactis* on lipid profile in individuals with type 2 diabetes mellitus. *Journal of Dairy Science*, 94(7), 3288-3294.
- El-Nezami, H. S., Polychronaki, N. N., Ma, J., Zhu, H., Ling, W., Salminen, E. K., Juvonen, R. O., Salminen, S. J., Poussa, T., & Mykka'nen, H. M. (2006). Probiotic supplementation reduces a biomarker for increased risk of liver cancer in young men from Southern China. *American Society for Nutrition*, 83, 1199-1203.
- Elegado, F. B., Guerra, M. A. R. V., Macayan, R. A., Mendoza, H. A., & Lirazan, M. B. (2004). Spectrum of bacteriocin activity of *Lactobacillus plantarum* BS and fingerprinting by RAPD-PCR. *International Journal of Food Microbiology*, 95(1), 11-18.
- Elliott, D. A., Kim, W. S., Jans, D. A., & Garner, B. (2007). Apoptosis induces neuronal apolipoprotein-E synthesis and localization in apoptotic bodies. *Neuroscience Letters*, 416(2), 206-210.
- Elmer, G. W. (2001). Probiotics: 'Living Drugs'. *American Society of Health-System Pharmacists*, 58(12), 1101-1109.
- Enck, P., Zimmermann, K., Menke, G., Müller-Lissner, S., Martens, U., & Klosterhalfen, S. (2008). A mixture of *Escherichia coli* (DSM 17252) and

- Enterococcus faecalis* (DSM16440) for treatment of the irritable bowel syndrome - a randomized controlled trial with primary care physicians. *Neurogastroenterology & Motility*, 20(10), 1103-1109.
- Endo, A., Futagawa-Endo, Y., & Dicks, L. M. T. (2010). Diversity of *Lactobacillus* and *Bifidobacterium* in feces of herbivores, omnivores and carnivores. *Anaerobe*, 16(6), 590-596.
- Ennahar, S., Sashihara, T., Sonomoto, K., & Ishizaki, A. (2000). Class Iia bacteriocins: biosynthesis, structure and activity. *FEMS Microbiology Reviews*, 24(1), 85-106.
- Ewaschuk, J.B., Walker, J.W., Diaz, H., & Madsen, K.L. (2006). Bioproduction of conjugated linoleic acid by probiotic bacteria occurs *in vitro* and *in vivo* in mice. *Nutrition*, 136, 1483-1487.
- Fadok, V. A., Bratton, D. L., Konowal, A., Freed, P. W., Westcott, J. Y., & Henson, P. M. (1998). Macrophages that have ingested apoptotic cells *in vitro* inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF-beta, PGE2, and PAF. *Journal of Clinical Investigation*, 101, 890-899.
- Fan, Y. j., Chen, S. j., Yu, Y. c., Si, J. m., & Liu, B. (2006). A probiotic treatment containing *Lactobacillus*, *Bifidobacterium* and *Enterococcus* improves IBS symptoms in an open label trial. *Journal of Zhejiang University Science B*, 7(12), 987-991.
- FAO/WHO. (2002). Guidelines for the evaluation of probiotics in food. FAO/WHO Working Group 1-11.
- Finegold, S. M., Martin, W. J., & Scott, E. G. (1978). Reagents and tests, in Bailey & Scott's diagnostic microbiology: CV Mosby Co.
- Fischer, M. H., Yu, N., Gray, G. R., Ralph, J., Anderson, L., & Marlett, J. A. (2004). The gel-forming polysaccharide of psyllium husk (*Plantago ovata* Forsk). *Carbohydrate Research*, 339(11), 2009-2017.
- Flórez, A. B., Tosi, L., Danielsen, M., von Wright, A., Bardowski, J., Morelli, L., & Mayo, B. (2008). Resistance-susceptibility profiles of *Lactococcus lactis* and *Streptococcus thermophilus* strains to eight antibiotics and proposition of new cut-offs. *International Journal of Probiotics and Prebiotics*, 3(4), 249-256.
- Foo, H. L., Loh, T. C., Lai, P. W., Lim, Y. Z., KufliC.N., & Rusul, G. (2003). Effects of adding *Lactobacillus plantarum* I-UL4 metabolites in drinking water of rats. *Pakistan Journal of Nutrition*, 2(5), 283-288.
- Forestier, C., De Champs, C., Vatoux, C., & Joly, B. (2001). Probiotic activities of *Lactobacillus casei* rhamnosus: *in vitro* adherence to intestinal cells and antimicrobial properties. *Research in Microbiology*, 152(2), 167-173.

- Foulquié Moreno, M. R., Sarantinopoulos, P., Tsakalidou, E., & De Vuyst, L. (2006). The role and application of *enterococci* in food and health. *International Journal of Food Microbiology*, 106(1), 1-24.
- Franciosi, E., Settanni, L., Cavazza, A., & Poznanski, E. (2009). Biodiversity and technological potential of wild lactic acid bacteria from raw cows' milk. *International Dairy Journal*, 19(1), 3-11.
- Franks, A. H., Harmsen, H. J. M., Raangs, G. C., Jansen, G. J., Schut, F., & Welling, G. W. (1998). Variations of bacterial populations in human feces measured by fluorescent in situ hybridization with group-specific 16S rRNA-targeted oligonucleotide probes. *Applied and Environmental Microbiology*, 64, 3336-3345.
- Franz, C. M. A. P., Holzapfel, W. H., & Stiles, M. E. (1999). *Enterococci* at the crossroads of food safety. *International Journal of Food Microbiology*, 47(1-2), 1-24.
- Franz, C. M. A. P., Huch, M., Abriouel, H., Holzapfel, W., & Gálvez, A. (2011). *Enterococci* as probiotics and their implications in food safety. *International Journal of Food Microbiology*, 151(2), 125-140.
- Franz, C. M. A. P., Stiles, M. E., Schleifer, K. H., & Holzapfel, W. H. (2003). *Enterococci* in foods—a conundrum for food safety. *International Journal of Food Microbiology*, 88(2-3), 105-122.
- Freese, E., Sheu, C. W., & Galliers, E. (1973). Function of lipophilic acids as antimicrobial food additives. *Nature*, 2(241), 321-325.
- Friedman, G. (2012). The role of probiotics in the prevention and treatment of antibiotic-associated diarrhea and *Clostridium difficile* colitis. *Gastroenterology Clinics of North America*, 41(4), 763-779.
- Fujimori, S., Gudis, K., Mitsui, K., Seo, T., Yonezawa, M., Tanaka, S., Tatsuguchi, A., & Sakamoto, C. (2009). A randomized controlled trial on the efficacy of synbiotic versus probiotic or prebiotic treatment to improve the quality of life in patients with ulcerative colitis. *Nutrition*, 25(5), 520-525.
- Gade, J., & Thorn, P. (1989). Paraghurt for patients with irritable bowel syndrome. A controlled clinical investigation from general practice. *Scandinavian Journal of Primary Health Care*, 7(1), 23-26.
- Gaggia, F., Mattarelli, P., & Biavati, B. (2010). Probiotics and prebiotics in animal feeding for safe food production. *International Journal of Food Microbiology*, 141, 15-28.
- Gala, E., Landi, S., Solieri, L., Nocetti, M., Pulvirenti, A., & Giudici, P. (2008). Diversity of lactic acid bacteria population in ripened Parmigiano Reggiano cheese. *International Journal of Food Microbiology*, 125(3), 347-351.

- Galluzzi, L., Maiuri, M. C., Vitale, I., Zischka, H., Zitvogel, L., & Kroemer, G. (2007). Cell death modalities: Classification and pathophysiological implications. *Cell Death and Differentiation*, 14, 1237-1266.
- Gancheva, A., Pot, B., Vanhonacker, K., Hoste, B., & Kersters, K. (1999). A polyphasic approach towards the identification of strains belonging to *Lactobacillus acidophilus* and related species. *Systematic and Applied Microbiology*, 22(4), 573-585.
- Gaspar, P., Carvalho, A. L., Vinga, S., Santos, H., & Neves, A. R. (2013). From physiology to systems metabolic engineering for the production of biochemicals by lactic acid bacteria. *Biotechnology Advances*. 31(6), 764-788.
- Geier, M. S., Butler, R. N., & Howarth, G. S. (2007). Inflammatory bowel disease: Current insights into pathogenesis and new therapeutic options; probiotics, prebiotics and synbiotics. *International Journal of Food Microbiology*, 115(1), 1-11.
- Gendrot, F., Ferchichi, M., Winters, D. A., & Hemme, D. (2000). Valine transport and biodiversity of *Leuconostoc* wild strains from French raw milk cheeses. *Systematic and Applied Microbiology*, 23(4), 461-468.
- Gerez, C. L., Torres, M. J., Font de Valdez, G., & Rollán, G. (2013). Control of spoilage fungi by lactic acid bacteria. *Biological Control*, 64(3), 231-237.
- Gevers, D., Huys, G., & Swings, J. (2001). Applicability of rep-PCR fingerprinting for identification of *Lactobacillus* species. *FEMS Microbiology Letters*, 205(1), 31-36.
- Gionchetti, P., Amadini, C., Rizzello, F., Venturi, A., Palmonari, V., Morselli, C., Romagnoli, R., & Campieri, M. (2002). Probiotics — Role in inflammatory bowel disease. *Digestive and Liver Disease*, 34(2), 58-62.
- Giraffa, G. (2002). *Enterococci* from foods. *FEMS Microbiology Reviews*, 26(2), 163-171.
- Giraffa, G., Chanishvili, N., & Widyastuti, Y. (2010). Importance of *lactobacilli* in food and feed biotechnology. *Research in Microbiology*, 161(6), 480-487.
- Glück, S. (2014). nab-Paclitaxel for the treatment of aggressive metastatic breast cancer. *Clinical Breast Cancer*, 14(4), 221-227.
- Goderska, K., & Czarnecki, Z. (2008). Influence of microencapsulation and spray drying on the viability of *Lactobacillus* and *Bifidobacterium* strains. *Polish Journal of Microbiology*, 57, 135-140.
- Goel, P. K. (2009). Patent EP2285821A1: A novel process for extraction of furostanolic saponins from fenugreek seeds.

- Goldin, B. R., & Gorbach, S. L. (1980). Effect of *Lactobacillus acidophilus* dietary supplements on 1,2-dimethylhydrazine dihydrochloride-induced intestinal cancer in rats. *Journal of the National Cancer Institute*, 64(2), 263-265.
- Goldin, B. R., Gualtieri, L. J., & Moore, R. P. (1996). The effect of *Lactobacillus* GG on the initiation and promotion of DMH- induced intestinal tumors in the rat. *Nutrition and Cancer*, 25(2), 197-204.
- Gómez Zavaglia, A., de Urza, P., & De Antoni, G. (2000). Characterization of *Bifidobacterium* strains using Box primers. *Anaerobe*, 6(3), 169-177.
- González, Á., Guillamón, J. M., Mas, A., & Poblet, M. (2006). Application of molecular methods for routine identification of acetic acid bacteria. *International Journal of Food Microbiology*, 108(1), 141-146.
- González, Á., Hierro, N., Poblet, M., Mas, A., & Guillamón, J. M. (2005). Application of molecular methods to demonstrate species and strain evolution of acetic acid bacteria population during wine production. *International Journal of Food Microbiology*, 102(3), 295-304.
- González, Á., & Mas, A. (2011). Differentiation of acetic acid bacteria based on sequence analysis of 16S–23S rRNA gene internal transcribed spacer sequences. *International Journal of Food Microbiology*, 147(3), 217-222.
- González Ferreiro, M., Tillman, L., Hardee, G., & Bodmeier, R. (2002). Characterization of alginate/poly-l-lysine particles as antisense oligonucleotide carriers. *International Journal of Pharmaceutics*, 239(1–2), 47-59.
- Gonzalez, C. J., Encinas, J. P., Garcia-Lopez, M. L., & Otero, A. (2000). Characterization and identification of lactic acid bacteria from freshwater fishes. *Food Microbiology*, 17, 383–391.
- Gopal, P. K., Prasad, J., & Gill, H. S. (2003). Effects of the consumption of *Bifidobacterium lactis* HN019 (DR10TM) and galacto-oligosaccharides on the microflora of the gastrointestinal tract in human subjects. *Nutrition Research*, 23(10), 1313-1328.
- Gornstein, E., & Schwarz, T. L. (2014). The paradox of paclitaxel neurotoxicity: Mechanisms and unanswered questions. *Neuropharmacology*, 76, Part A, 175-183.
- Grojean, S., Koziel, V., Vert, P., & Luc Daval, J. (2000). Apoptosis induced by calcein acetoxymethyl ester in human histiocytic lymphoma cell line. *Biochemical Pharmacology*, 60, 1751-1759.
- Guarner, F., Khan, A. G., Garisch, J., Eliakim, R., Gangl, A., Thomson, A., Krabshuis, J., & Le Mair, T. (2008). Probiotics and prebiotics. *World Gastroenterology Organisation Practice Guideline*.
- Gudkov, A. V., & Komarova, E. A. (2003). The role of p53 in determining sensitivity to radiotherapy. *Nature Reviews Cancer*, 3, 117-129.

- Gueimonde, M., Delgado, S., Mayo, B., Ruas-Madiedo, P., Margolles, A., & de los Reyes-Gavilán, C. G. (2004). Viability and diversity of probiotic *Lactobacillus* and *Bifidobacterium* populations included in commercial fermented milks. *Food Research International*, 37(9), 839-850.
- Guérin, D., Vuillemand, J. C., & Subirade, M. (2003). Protection of *bifidobacteria* encapsulated in polysaccharide–protein gel beads against gastric juice and bile. *Journal of Food Protection*, 66(11), 2076-2084.
- Gullo, M., Caggia, C., De Vero, L., & Giudici, P. (2006). Characterization of acetic acid bacteria in “traditional balsamic vinegar”. *International Journal of Food Microbiology*, 106(2), 209-212.
- Guo, Q., Cui, S. W., Wang, Q., & Christopher Young, J. (2008). Fractionation and physicochemical characterization of psyllium gum. *Carbohydrate Polymers*, 73(1), 35-43.
- Gupta, P. K., Mital, B. K., & Garg, S. K. (1996). Characterization of *Lactobacillus acidophilus* strains for use as dietary adjunct. *International Journal of Food Microbiology*, 29(1), 105-109.
- Gupta, R., & Srivastava, S. (2014). Antifungal effect of antimicrobial peptides (AMPs LR14) derived from *Lactobacillus plantarum* strain LR/14 and their applications in prevention of grain spoilage. *Food Microbiology*, 42, 1-7.
- Gurtler, V., Grando, D., Mayall, B. C., Wang, J., & Ghaly-Derias, S. (2012). A novel method for simultaneous *Enterococcus* species identification/typing and van genotyping by high resolution melt analysis. *Journal of Microbiological Methods*, 90(3), 167-181.
- Hadji-Sfaki, I., El-Ghaish, S., Ahmadova, A., Batdorj, B., Le Blay-Laliberté, G., Barbier, G., Haertlé, T., & Chobert, J.-M. (2011). Antimicrobial activity and safety of use of *Enterococcus faecium* PC4.1 isolated from Mongol yogurt. *Food Control*, 22(12), 2020-2027.
- Halicka, H. D., Bedner, E., & Darzynkiewicz, Z. (2000). Segregation of RNA and separate packaging of DNA and RNA in apoptotic bodies during apoptosis. *Experimental Cell Research*, 260, 248-255.
- Hamid, R., Rotshteyn, Y., Rabadi, L., Parikh, R., & Bullock, P. (2004). Comparison of alamar blue and MTT assays for high through-put screening. *Toxicology in vitro*, 18, 703-710.
- Handwerker, S., Pucci, M. J., Volk, K. J., Liu, J., & Lee, M. S. (1994). Vancomycin-resistant *Leuconostoc mesenteroides* and *Lactobacillus casei* synthesize cytoplasmic peptidoglycan precursors that terminate in lactate. *Journal of Bacteriology*, 176(1), 260–264.
- Hatakka, K., Ahola, A. J., Yli-Knuuttila, H., Richardson, M., Poussa, T., Meurman, J. H., & Korpela, R. (2007). Probiotics reduce the prevalence of oral *Candida* in the elderly a randomized controlled trial. *Journal of Dental Research*, 86(2), 125-130.

- Hayatsu, H., & Hayatsu, T. (1993). Suppressing effect of *Lactobacillus casei* administration on the urinary mutagenicity arising from ingestion of fried ground beef in the human. *Cancer Letters*, 73(2–3), 173-179.
- Heidebach, T., Först, P., & Kulozik, U. (2009). Microencapsulation of probiotic cells by means of rennet-gelation of milk proteins. *Food Hydrocolloids*, 23(7), 1670-1677.
- Heidebach, T., Först, P., & Kulozik, U. (2010). Influence of casein-based microencapsulation on freeze-drying and storage of probiotic cells. *Journal of Food Engineering*, 98(3), 309-316.
- Hemme, D., & Foucaud-Scheunemann, C. (2004). *Leuconostoc*, characteristics, use in dairy technology and prospects in functional foods. *International Dairy Journal*, 14(6), 467-494.
- Henry, C. M., Hollville, E., & Martin, S. J. (2013). Measuring apoptosis by microscopy and flow cytometry. *Methods*, 61(2), 90-97.
- Hernández-Rodríguez, L., Lobato-Calleros, C., Pimentel-González, D. J., & Vernon-Carter, E. J. (2014). *Lactobacillus plantarum* protection by entrapment in whey protein isolate: κ -carrageenan complex coacervates. *Food Hydrocolloids*, 36(0), 181-188.
- Herreros, M. A., Fresno, J. M., González Prieto, M. J., & Tornadijo, M. E. (2003). Technological characterization of lactic acid bacteria isolated from Armada cheese (a Spanish goats' milk cheese). *International Dairy Journal*, 13(6), 469-479.
- Hill, D. A., Hoffmann, C., Abt, M. C., Du, Y., Kobuley, D., Kirn, T., Bushman, F.D., & Artis, D. (2010). Metagenomic analyses reveal antibiotic-induced temporal and spatial changes in intestinal microbiota with associated alterations in immune cell homeostasis. *Mucosal Immunology*, 3(2), 148-158.
- Hirayama, K., & Rafter, J. (2000). The role of probiotic bacteria in cancer prevention. *Microbes and Infection*, 2(6), 681-686.
- Hlivak, P., Odraska, J., Ferencik, M., Ebringer, L., Jahnova, E., & Mikes, Z. (2005). One-year application of probiotic strain *Enterococcus faecium* M-74 decreases serum cholesterol levels. *Bratislava Medical Journal*, 106(2), 67-72.
- Holzapfel, W. H., Haberer, P., Snel, J., Schillinger, U., & Huis in't Veld, J. H. J. (1998). Overview of gut flora and probiotics. *International Journal of Food Microbiology*, 41(2), 85-101.
- Hong, S. I., Kim, Y. J., & Pyun, Y. R. (1999). Acid tolerance of *Lactobacillus plantarum* from Kimchi. *Lebensm.-Wiss. u.-Technol*, 32(3), 142-148.
- Hsu, Y. L., Kuo, P. L., Lin, L. T., & Lin, C. C. (2005). Asiatic acid, a triterpene, induces apoptosis and cell cycle arrest through activation of extracellular signal-regulated kinase and p38 mitogen-activated protein kinase pathways in human

- breast cancer cells. *Journal of Pharmacology and Experimental Therapeutics*, 313(1), 333-344.
- Huot, E., Meghrou, J., Barrena-Gonzalez, C., & Petitdemange, H. (1996). Bacteriocin J46, a new bacteriocin produced by *Lactococcus lactis* subsp.cremorisJ46: Isolation and characterization of the protein and its gene. *Anaerobe*, 2(3), 137-145.
- Hussein, K., Raz-Pasteur, A., Finkelstein, R., Neuberger, A., Shachor-Meyouhas, Y., Oren, I., & Kassis, I. (2013). Impact of carbapenem resistance on the outcome of patients' hospital-acquired bacteraemia caused by *Klebsiella pneumoniae*. *Journal of Hospital Infection*, 83(4), 307-313.
- Hwang, S. W., Kim, N., Kim, J. M., Huh, C. S., Ahn, Y. T., Park, S. H., Shin, C. M., Park, J.H., Lee, M.K., Nam, R. H., Lee, H. S., Kim, J. S., Jung, H. C., & Song, I. S. (2012). Probiotic suppression of the *H. pylori*-induced responses by conjugated linoleic acids in a gastric epithelial cell line. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 86(6), 225-231.
- Ibrahim, F., Ruvio, S., Granlund, L., Salminen, S., Viitanen, M., & Ouwehand, A. C. (2010). Probiotics and immunosenescence: Cheese as a carrier. *FEMS Immunology and Medical Microbiology*, 59(1), 53-59.
- Iran. (2009). Project Document for a Regional Standard for Doogh. Paper presented at the FAO/WHO COORDINATING COMMITTEE FOR THE NEAR EAST, Tunis, Tunisia.
- Irvine, S. L., Hummelen, R., & Hekmat, S. (2011). Probiotic yogurt consumption may improve gastrointestinal symptoms, productivity, and nutritional intake of people living with human immunodeficiency virus in Mwanza, Tanzania. *Nutrition Research*, 31(12), 875-881.
- Işıklı, N. D., & Karababa, E. (2005). Rheological characterization of fenugreek paste (çemen). *Journal of Food Engineering*, 69(2), 185-190.
- Isolauri, E., Salminen, S., & Ouwehand, A. C. (2004). Probiotics. *Best Practice & Research Clinical Gastroenterology*, 18(2), 299-313.
- Izquierdo, E., Bednarczyk, A., Schaeffer, C., Cai, Y., Marchioni, E., Dorsselaer, A. V., & Ennahar, S. (2008). Production of enterocins L50A, L50B, and IT, a new enterocin, by *Enterococcus faecium* IT62, a strain isolated from Italian ryegrass in Japan. *Antimicrobial Agents and Chemotherapy*, 52(6), 1917–1923.
- Janapatla, R. P., Yan, J. J., Huang, A. H., Chen, H. M., Wu, H. M., & Wu, J. J. (2007). Inducible clindamycin resistance in *Staphylococcus aureus* isolates causing bacteremia at a university hospital in southern Taiwan. *Diagnostic Microbiology and Infectious Disease*, 58(2), 203-209.
- Jang, J., Kim, B., Lee, J., & Han, H. (2003). A rapid method for identification of typical *Leuconostoc* species by 16S rDNA PCR-RFLP analysis. *Journal of Microbiological Methods*, 55(1), 295-302.

- Jasniewski, J., Cailliez-Grimal, C., Younsi, M., Millière, J. B., & Revol-Junelles, A. M. (2008). Fluorescence anisotropy analysis of the mechanism of action of mesentericin 52A: speculations on antimicrobial mechanism. *Applied Microbiology and Biotechnology*, 81(2), 339-247.
- Jatkauskas, J., & Vrotniakienė, V. (2010). Effects of probiotic dietary supplementation on diarrhoea patterns, faecal microbiota and performance of early weaned calves. *Veterinarija Medicina*, 55(10), 494–503.
- Jing, X., Deng, L., Gao, B., Xiao, L., Zhang, Y., Ke, X., Lian, J., Zhao, Q., Ma, L., Yao, J., & Chen, J. (2014). A novel polyethylene glycol mediated lipid nanoemulsion as drug delivery carrier for paclitaxel. *Nanomedicine: Nanotechnology, Biology and Medicine*, 10(2), 371-380.
- Johnson, A. P., Uttley, A. H. C., Woodford, N., & George, R. C. (1990). Resistance to vancomycin and teicoplanin: an emerging clinical problem. *Clinical Microbiology Reviews.*, 3, 280-291.
- Jokovic, N., Nikolic, M., Begovic, J., Jovcic, B., Savic, D., & Topisirovic, L. (2008). A survey of the lactic acid bacteria isolated from Serbian artisanal dairy product kajmak. *International Journal of Food Microbiology*, 127(3), 305-311.
- Jose, J. E., Padmanabhan, S., & Chitharanjan, A. B. (2013). Systemic consumption of probiotic curd and use of probiotic toothpaste to reduce *Streptococcus mutans* in plaque around orthodontic brackets. *American Journal of Orthodontics and Dentofacial Orthopedics*, 144(1), 67-72.
- Jung, J. Y., Lee, S. H., Lee, H. J., Seo, H. Y., Park, W. S., & Jeon, C. O. (2012). Effects of *Leuconostoc mesenteroides* starter cultures on microbial communities and metabolites during kimchi fermentation. *International Journal of Food Microbiology*, 153(3), 378-387.
- Kadere, T. T., Miyamoto, T., Oniang'o, R. K., Kutima, P. M., & Njoroge, S. M. (2008). Isolation and identification of the genera *Acetobacter* and *Gluconobacter* in coconut toddy (mnazi). *African Journal of Biotechnology*, 7(16), 2963-2971.
- Kailasapathy, K. (2002). Microencapsulation of probiotic bacteria: technology and potential applications. *Current Issues in Intestinal Microbiology*, 3(2), 39-48.
- Kailasapathy, K. (2006). Survival of free and encapsulated probiotic bacteria and their effect on the sensory properties of yoghurt. *LWT - Food Science and Technology*, 39(10), 1221-1227.
- Kailasapathy, K., & Chin, J. (2000). Survival and therapeutic potential of probiotic organisms with reference to *Lactobacillus acidophilus* and *Bifidobacterium* spp. *Immunology and Cell Biology*, 78(1), 80-88.
- Kailasapathy, K., & Masondole, L. (2005). Survival of free and microencapsulated *Lactobacillus acidophilus* and *Bifidobacterium lactis* and their effect on texture of feta cheese. *Australian Journal of Dairy Technology*, 60(3), 252-258.

- Kastner, S., Perreten, V., Bleuler, H., Hugenschmidt, G., Lacroix, C., & Meile, L. (2006). Antibiotic susceptibility patterns and resistance genes of starter cultures and probiotic bacteria used in food. *Systematic and Applied Microbiology*, 29(2), 145-155.
- Katla, A.-K., Kruse, H., Johnsen, G., & Herikstad, H. (2001). Antimicrobial susceptibility of starter culture bacteria used in Norwegian dairy products. *International Journal of Food Microbiology*, 67(1-2), 147-152.
- Kemp, C. J., Sun, S., & Gurley, K. E. (2001). p53 induction and apoptosis in response to radio- and chemotherapy *in vivo* is tumor-type-dependent. *Cancer Research*, 61(1), 327-332.
- Khan, N. H., Korber, D. R., Low, N. H., & Nickerson, M. T. (2013). Development of extrusion-based legume protein isolate–alginate capsules for the protection and delivery of the acid sensitive probiotic, *Bifidobacterium adolescentis*. *Food Research International*, 54(1), 730-737.
- Kiess, W., & Gallaher, B. (1998). Hormonal control of programmed cell death/apoptosis. *European Journal of Endocrinology*, 138, 482-491.
- Kim, J. Y., Woo, H. J., Kim, Y. S., & Lee, H. J. (2002). Screening for antiproliferative effects of cellular components from lactic acid bacteria against human cancer cell lines. *Biotechnology Letters*, 24(17), 1431-1436.
- Kimoto-Nira, H., Suzuki, C., Sasaki, K., Kobayashi, M., & Mizumachi, K. (2010). Survival of a *Lactococcus lactis* strain varies with its carbohydrate preference under in vitro conditions simulated gastrointestinal tract. *International Journal of Food Microbiology*, 143(3), 226-229.
- Kiwaki, M., & Sato, T. (2009). Antimicrobial susceptibility of *Bifidobacterium breve* strains and genetic analysis of streptomycin resistance of probiotic *B. breve* strain Yakult. *International Journal of Food Microbiology*, 134(3), 211-215.
- Klaenhammer, T. R., & Kullen, M. J. (1999). Selection and design of probiotics. *International Journal of Food Microbiology*, 50(1-2), 45-57.
- Klein, G., Hallmann, C., Casas, I. A., Abad, J., Louwers, J., & Reuter, G. (2000). Exclusion of vanA, vanB and vanC type glycopeptide resistance in strains of *Lactobacillus reuteri* and *Lactobacillus rhamnosus* used as probiotics by polymerase chain reaction and hybridization methods. *Journal of Applied Microbiology*, 89(5), 815-824.
- Klibi, N., Said, L. B., Jouini, A., Slama, K. B., López, M., Sallem, R. B., Boudabous, A., & Torres, C. (2013). Species distribution, antibiotic resistance and virulence traits in *enterococci* from meat in Tunisia. *Meat Science*, 93(3), 675-680.
- Knoetze, H., Todorov, S. D., & Dicks, L. M. T. (2008). A class IIa peptide from *Enterococcus mundtii* inhibits bacteria associated with otitis media. *International Journal of Antimicrobial Agents*, 31(3), 228-234.

- Komarova, E. A., Chernov, M. V., Franks, R., Wang, K., Armin, G., Zelnick, C. R., Chin, D. M., Bacus, S. S., Stark, G. R., & Gudkov, A. V. (1997). Transgenic mice with p53-responsive lacZ: p53 activity varies dramatically during normal development and determines radiation and drug sensitivity in vivo. *The EMBO Journal*, 16(6), 391-1400.
- Kore, K. B., Pattanaik, A. K., K, S., & Mirajkar, P. P. (2012). Effect of feeding traditionally prepared fermented milk dahi (curd) as a probiotics on nutritional status, hindgut health and haematology in dogs. *Indian Journal of Traditional Knowledge*, 11(1).
- Korzeniewski, C., & Callewaert, D. M. (1983). An enzyme-release assay for natural cytotoxicity. *Journal of Immunological Methods*, 64, 313-320.
- Kos, B., Suskovic, J., Goreta, J., & Matosic, S. (2000). Effect of protectors on the viability of *Lactobacillus acidophilus* M92 in simulated gastrointestinal conditions. *Food Technology and Biotechnology*, 38, 121-127.
- Kostinek, M., Pukall, R., Rooney, A. P., Schillinger, U., Hertel, C., & Holzapfel, W. H. (2005). *Lactobacillus arizonensis* is a later heterotypic synonym of *Lactobacillus plantarum*. *International Journal of Systematic and Evolutionary Microbiology*, 55, 2485-2489.
- Krasaekoopt, W., Bhandari, B., & Deeth, H. (2003). Evaluation of encapsulation techniques of probiotics for yoghurt. *International Dairy Journal*, 13(1), 3-13.
- Krasaekoopt, W., Bhandari, B., & Deeth, H. (2004). The influence of coating materials on some properties of alginate beads and survivability of microencapsulated probiotic bacteria. *International Dairy Journal*, 14(8), 737-743.
- Krasaekoopt, W., Bhandari, B., & Deeth, H. C. (2006). Survival of probiotics encapsulated in chitosan-coated alginate beads in yoghurt from UHT- and conventionally treated milk during storage. *LWT - Food Science and Technology*, 39(2), 177-183.
- Křížová, J., Španová, A., & Rittich, B. (2006). Evaluation of amplified ribosomal DNA restriction analysis (ARDRA) and species-specific PCR for identification of *Bifidobacterium* species. *Systematic and Applied Microbiology*, 29(1), 36-44.
- Kullen, M. J., Brady, L. J., & O'Sullivan, D. J. (1997). Evaluation of using a short region of the recA gene for rapid and sensitive speciation of dominant *bifidobacteria* in the human large intestine. *FEMS Microbiology Letters*, 154(2), 377-383.
- Lahtinen, S. J., Forssten, S., Aakko, J., Granlund, L., Rautonen, N., Salminen, S., Viitanen, M., & Ouwehand, A. C. (2012). Probiotic cheese containing *Lactobacillus rhamnosus* HN001 and *Lactobacillus acidophilus* NCFM® modifies subpopulations of fecal *lactobacilli* and *Clostridium difficile* in the elderly. *Age*, 34(1), 133-143.

- Langer-Safer, P. R., Levine, M., & Ward, D. C. (1982). Immunological method for mapping genes on *Drosophila polytene* chromosomes. *Proceedings of the National Academy of Sciences*, 79(14), 4381–4385.
- Le, M. G., Moulton, L. H., Hill, C., & Kramar, A. (1986). Consumption of dairy produce and alcohol in a case-control study of breast cancer. *Journal of the National Cancer Institute*, 77(3), 633-636.
- Leblond-Bouget, N., Philippe, H., Mangin, I., & Decaris, B. (1996). 16s rRNA and 16s to 23s internal transcribed spacer sequence analyses reveal inter- and intraspecific *Bifidobacterium* phylogeny. *International Journal of Systematic Bacteriology*, 46(1), 102-111.
- Lee, B. R., Hwang, J. W., Choi, Y. Y., Wong, S. F., Hwang, Y. H., Lee, D. Y., & Lee, S. H. (2012). *In situ* formation and collagen-alginate composite encapsulation of pancreatic islet spheroids. *Biomaterials*, 33(3), 837-845.
- Lee, D. K., Jang, S., Kim, M. J., Kim, J. H., Chung, M. J., Kim, K. J., & Ha, N. J. (2008). Anti-proliferative effects of *Bifidobacterium adolescentis* SPM0212 extract on human colon cancer cell lines. *BMC Cancer*, 8(310), 1-8.
- Lee, J., Jang, J., Kim, B., Kim, J., Jeong, G., & Han, H. (2004). Identification of *Lactobacillus sakei* and *Lactobacillus curvatus* by multiplex PCR-based restriction enzyme analysis. *Journal of Microbiological Methods*, 59(1), 1-6.
- Lee, J. H., Nam, S. H., Seo, W. T., Yun, H. D., Hong, S. Y., Kim, M. K., & Cho, K. M. (2012). The production of surfactin during the fermentation of cheonggukjang by potential probiotic *Bacillus subtilis* CSY191 and the resultant growth suppression of MCF-7 human breast cancer cells. *Food Chemistry*, 131(4), 1347-1354.
- Lee, S. M., & Lee, W. K. (2000). Inhibitory effects of lactic acid bacteria (LAB) on the azoxymethane-induced colonic preneoplastic lesions. *The Journal of Microbiology*, 38(3), 169-175.
- Leenhouts, K. J., Kok, J., & Venema, G. (1990). Stability of integrated plasmids in the chromosome of *Lactococcus lactis*. *Applied and Environmental Microbiology*, 56(9), 2726-2735.
- Lefeber, T., Gobert, W., Vrancken, G., Camu, N., & De Vuyst, L. (2011). Dynamics and species diversity of communities of lactic acid bacteria and acetic acid bacteria during spontaneous cocoa bean fermentation in vessels. *Food Microbiology*, 28(3), 457-464.
- Li-on, C. (2010). Cytotoxic and antiproliferative properties of metabolites produced by six strains of *Lactobacillus plantarum* on human cancer cells. Accepted thesis, Faculty of biotechnology and biomolecular sciences, Universiti Putra Malaysia.
- Li, Y., Ahmed, F., Ali, S., Philip, P. A., Kucuk, O., & Sarkar, F. H. (2005). Inactivation of nuclear factor kappaB by soy isoflavone genistein contributes to increased apoptosis induced by chemotherapeutic agent in human cancer cells. *Cancer Research*, 65(15), 6934-6942.

- Lilly, D. M., & Stillwell, R. H. (1965). Probiotics: Growth-promoting factors produced by microorganisms. *Science*, 147(3659), 747-748.
- Lim, Y. S. (2010). Probiotic characteristics of bacteriocinogenic *Lactobacillus plantarum* strains isolated from Malaysian food. Accepted thesis, Faculty of biotechnology and biomolecular sciences, Universiti Putra Malaysia.
- Lin, W. H., Hwang, C.-F., Chen, L. W., & Tsen, H. Y. (2006). Viable counts, characteristic evaluation for commercial lactic acid bacteria products. *Food Microbiology*, 23(1), 74-81.
- Liong, M. T., & Shah, N. P. (2005). Acid and bile tolerance and cholesterol removal ability of *Lactobacilli* strains. *Journal of Dairy Science*, 88(1), 55-66.
- Liu, S. N., Han, Y., & Zhou, Z. J. (2011). Lactic acid bacteria in traditional fermented Chinese foods. *Food Research International*, 44(3), 643-651.
- Liu, W., Bao, Q., Jirimutu, Qing, M., Siriguleng, Chen, X., Sun, T., Li, M., Zhang, J., Yu, J., Bilige, M., Sun, T., & Zhang, H. (2012). Isolation and identification of lactic acid bacteria from Tarag in Eastern Inner Mongolia of China by 16S rRNA sequences and DGGE analysis. *Microbiological Research*, 167(2), 110-115.
- Lollo, P. C. B., Cruz, A. G., Morato, P. N., Moura, C. S., Carvalho-Silva, L. B., Oliveira, C. A. F., Faria, J. A. F., & Amaya-Farfan, J. (2012). Probiotic cheese attenuates exercise-induced immune suppression in Wistar rats. *Journal of Dairy Science*, 95(7), 3549-3558.
- Lollo, P. C. B., de Moura, C. S., Morato, P. N., Cruz, A. G., Castro, W. d. F., Betim, C. B., Nisishima, L., Faria, J. A. F., Maróstica Junior, M., Fernandes, C. O., & Amaya-Farfan, J. (2013). Probiotic yogurt offers higher immune-protection than probiotic whey beverage. *Food Research International*, 54(1), 118-124.
- Looft, T., & Allen, H. K. (2012). Collateral effects of antibiotics on mammalian gut microbiomes. *Gut Microbes*, 3(5), 1-5.
- López-Díaz, T. M., Alonso, C., Román, C., García-López, M. L., & Moreno, B. (2000). Lactic acid bacteria isolated from a hand-made blue cheese. *Food Microbiology*, 17(1), 23-32.
- Lotfipour, F., Mirzaeei, S., & Maghsoodi, M. (2012). Preparation and characterization of alginate and psyllium beads containing *Lactobacillus acidophilus*. *The Scientific World Journal*, 1-8.
- Lourens-Hattingh, A., & Viljoen, B. C. (2001). Yogurt as probiotic carrier food. *International Dairy Journal*, 11(1-2), 1-17.
- Ly, N. P., Litonjua, A., Gold, D. R., & Celedon, J. C. (2011). Gut microbiota, probiotics, and vitamin D: Interrelated exposures influencing allergy, asthma, and obesity. *American Academy of Allergy, Asthma & Immunology*, 127(5), 1087-1094.

- Ma, Y., Pacan, J. C., Wang, Q., Sabour, P. M., Huang, X., & Xu, Y. (2012). Enhanced alginate microspheres as means of oral delivery of bacteriophage for reducing *Staphylococcus aureus* intestinal carriage. *Food Hydrocolloids*, 26(2), 434-440.
- Ma, Y. S., Pacan, J. C., Wang, Q., Xu, Y. P., Huang, X. Q., Korenevsky, A., & Sabour, P. M. (2008). Microencapsulation of bacteriophage Felix O1 into chitosan–alginate microspheres for oral delivery. *Applied and Environmental Microbiology*, 74, 4799-4805.
- Maal, B. K., & Shafiee, R. (2010). Characterization of an *Acetobacter* strain isolated from Iranian peach that tolerates high temperatures and ethanol concentrations. *World Academy of Science, Engineering and Technology*, 62, 141-145.
- Maal, K. B., & Shafiee, R. (2009). Isolation and identification of an *Acetobacter* strain from Iranian white-red cherry with high acetic acid productivity as a potential strain for cherry vinegar production in food and agriculture biotechnology. *World Academy of Science, Engineering and Technology*, 54, 201-203.
- Macedo, A. C., Costa, M. L., & Malcata, F. X. (1996). Changes in the microflora of Serra cheese: Evolution throughout ripening time, lactation period and axial location. *International Dairy Journal*, 6(1), 79-94.
- Malek, R., El-Attar, A., Mohamed, M., Anwar, S., El-Soda, M., & Béal, C. (2012). Technological and safety properties display biodiversity among *enterococci* isolated from two Egyptian cheeses, “Ras” and “Domiat”. *International Journal of Food Microbiology*, 153(3), 314-322.
- Mandal, S., Puniya, A. K., & Singh, K. (2006). Effect of alginate concentrations on survival of microencapsulated *Lactobacillus casei* NCDC-298. *International Dairy Journal*, 16(10), 1190-1195.
- Manichanh, C., Reeder, J., & Gibert, P. (2010). Reshaping the gut microbiome with bacterial transplantation and antibiotic intake. *Genome Research*, 20, 1411-1419.
- Mannu, L., Paba, A., Daga, E., Comunian, R., Zanetti, S., Duprè, I., & Sechi, L. A. (2003). Comparison of the incidence of virulence determinants and antibiotic resistance between *Enterococcus faecium* strains of dairy, animal and clinical origin. *International Journal of Food Microbiology*, 88(2–3), 291-304.
- Mantzouridou, F., Spanou, A., & Kiosseoglou, V. (2012). An inulin-based dressing emulsion as a potential probiotic food carrier. *Food Research International*, 46(1), 260-269.
- Maroof, H., Hassan, Z. M., Mobarez, A. M., & Mohamadabadi, M. A. (2012). *Lactobacillus acidophilus* could modulate the immune response against breast cancer in murine model. *Journal of Clinical Immunology*, 32(6), 1353-1359.
- Marques, A. P., Zé-Zé, L., San-Romão, M. V., & Tenreiro, R. (2010). A novel molecular method for identification of *Oenococcus oeni* and its specific detection in wine. *International Journal of Food Microbiology*, 142(1–2), 251-255.

- Martinez, R. C. R., Wachsmann, M., Torres, N. I., LeBlanc, J. G., Todorov, S. D., & Franco, B. D. G. d. M. (2013). Biochemical, antimicrobial and molecular characterization of a noncytotoxic bacteriocin produced by *Lactobacillus plantarum* ST71KS. *Food Microbiology*, 34(2), 376-381.
- Martley, F. G., & Crow, V. L. (1993). Interactions between non-starter microorganisms during cheese manufacture and ripening. *International Dairy Journal*, 3(4-6), 461-483.
- Maruo, T., Sakamoto, M., Toda, T., & Benno, Y. (2006). Monitoring the cell number of *Lactococcus lactis* subsp. *cremoris* FC in human feces by real-time PCR with strain-specific primers designed using the RAPD technique. *International Journal of Food Microbiology*, 110(1), 69-76.
- Masco, L., Huys, G., Gevers, D., Verbruggen, L., & Swings, J. (2003). Identification of *Bifidobacterium* species using rep-PCR fingerprinting. *Systematic and Applied Microbiology*, 26(4), 557-563.
- Massi, M., Vitali, B., Federici, F., Matteuzzi, D., & Brigidi, P. (2004). Identification method based on PCR combined with automated ribotyping for tracking probiotic *Lactobacillus* strains colonizing the human gut and vagina. *Journal of Applied Microbiology*, 96, 777-786.
- Masuda, Y., Ono, H., Kitagawa, H., Ito, H., Mu, F., Sawa, N., Zendo, T., & Sonomoto, K. (2011). Identification and characterization of leucocyclicin Q, a novel cyclic bacteriocin produced by *Leuconostoc mesenteroides* TK41401. *Applied and Environmental Microbiology*, 77(22), 8164-8170.
- Mathara, J. M., Schillinger, U., Kutima, P. M., Mbugua, S. K., Guigas, C., Franz, C., & Holzapfel, W. H. (2008). Functional properties of *Lactobacillus plantarum* strains isolated from Maasai traditional fermented milk products in Kenya. *Current Microbiology*, 56(4), 315-321.
- Mathur, S., & Singh, R. (2005). Antibiotic resistance in food lactic acid bacteria—a review. *International Journal of Food Microbiology*, 105(3), 281-295.
- Matsuki, T., Watanabe, K., Fujimoto, J., Miyamoto, Y., Takada, T., Matsumoto, K., Oyaizu, H., & Tanaka, R. (2002). Development of 16S rRNA-gene-targeted group-specific primers for the detection and identification of predominant bacteria in human feces. *Applied and Environmental Microbiology*, 68, 5445-5451.
- Matsuki, T., Watanabe, K., & Tanaka, R. (2003). Genus- and species-specific PCR primers for the detection and identification of *Bifidobacteria*. *Current Issues in Intestinal Microbiology*, 4, 61-69.
- Matsushita, K., Toyama, H., & Adachi, O. (1994). Respiratory chains and bioenergetics of acetic acid bacteria. *Advances in Microbial Physiology*, 36, 247-301.
- McAuley, C. M., Gobius, K. S., Britz, M. L., & Craven, H. M. (2012). Heat resistance of thermotolerant *enterococci* isolated from milk. *International Journal of Food Microbiology*, 154(3), 162-168.

- McFaddin, J. F. (1980). Biochemical tests for identification of medical bacteria: The Williams and Wilkins Co.
- McMahon, D. J., Oberg, C. J., Drake, M. A., Farkye, N., Moyes, L. V., Arnold, M. R., Ganesan, B., Steele, J., & Broadbent, J. R. (2014). Effect of sodium, potassium, magnesium, and calcium salt cations on pH, proteolysis, organic acids, and microbial populations during storage of full-fat Cheddar cheese. *Journal of Dairy Science*, 97(8), 4780-4798.
- Medici, M., Vinderola, C. G., & Perdigón, G. (2004). Gut mucosal immunomodulation by probiotic fresh cheese. *International Dairy Journal*, 14(7), 611-618.
- Medina, L., & Jordano, R. (1994). Survival of constitutive microflora in commercially fermented milk containing *bifidobacteria* during refrigerated storage. *Journal of Food Protection*, 55, 731-733.
- Mehrabian, S., Tajabadi Ebrahimi, M., Abbas Ahmadi, M., & Bahrami, H. (2012). Study of antimutagenic and anticancer effect of lactic acid bacteria isolated from Tarkhineh by Ames Test. *Arak Medical University Journal*, 15(66), 72-79.
- Merk, K., Borelli, C., & Korting, H. C. (2005). *Lactobacilli* – bacteria–host interactions with special regard to the urogenital tract. *International Journal of Medical Microbiology*, 295(1), 9-18.
- Micheline, S., Modesto, M., Oki, K., Stenico, V., Stefanini, I., Biavati, B., Watanabe, K., Ferrara, A., & Mattarelli, P. (2015). Isolation and identification of cultivable *Bifidobacterium* spp. from the faeces of 5 baby common marmosets (*Callithrix jacchus* L.). *Anaerobe*, 33(0), 101-104.
- Millette, M., Luquet, F., & Lacroix, M. (2007). In vitro growth control of selected pathogens by *Lactobacillus acidophilus*- and *Lactobacillus casei*-fermented milk. *Letters in Applied Microbiology*, 44(3), 314-319.
- Mirmomeni, M. H., Mohammadi, F., Sisakhtnezhad, S., Hashemi, R., & Nazari, G. (2009). Statistical analysis of different cancers in Kermanshah province. *Journal of Biological Sciences*, 9(3), 200-216.
- Mirzaei, H., & Barzgar, A. (2012). Isolation and molecular study of potentially probiotic *Lactobacilli* in traditional white cheese of Tabriz in Iran. *Annals of Biological Research*, 3(5), 2213-2216.
- Modler, H. W., McKellar, R. C., & Yaguchi, M. (1990). *Bifidobacteria* and bifidogenic factors. *Canadian Institute of Food Science and Technology Journal*, 23(1), 29-41.
- Modzelewska-Kapituła, M., Kobukowski, J. A. N., Kłebukowska, L., & Wiśniewska-Pantak, D. (2010). The influence of feeding diets containing white cheese, produced with prebiotics and the potentially probiotic *lactobacillus plantarum* strain, on the gastrointestinal microflora of rats. *Czech Journal of Food Sciences*, 28(2), 139-145.

- Moghadam, M. S., Foo, H. L., Leow, T. C., Abdul Rahim, R., & Loh, T. C. (2010). Novel bacteriocinogenic *Lactobacillus plantarum* strains and their differentiation by sequence analysis of 16S rDNA, 16S-23S and 23S-5S intergenic spacer regions and randomly amplified polymorphic DNA analysis. *Food Technology and Biotechnology*, 48(4), 476–483.
- Mokarram, R. R., Mortazavi, S. A., Najafi, M. B. H., & Shahidi, F. (2009). The influence of multi stage alginate coating on survivability of potential probiotic bacteria in simulated gastric and intestinal juice. *Food Research International*, 42(8), 1040-1045.
- Moonmangmee, S., Kawabata, K., Tanaka, S., Toyama, H., Adachi, O., & Matsushita, K. (2002). A novel polysaccharide involved in the pellicle formation of *Acetobacter aceti*. *Journal of Bioscience and Bioengineering*, 93(2), 192-200.
- Moore, A. M., Munck, C., Sommer, M. O. A., & Dantas, G. (2011). Functional metagenomic investigations of the human intestinal microbiota. *Frontiers in Microbiology*, 2(188), 1-8.
- Moraes, P. M., Perin, L. M., Tassinari Ortolani, M. B., Yamazi, A. K., Viçosa, G. N., & Nero, L. A. (2010). Protocols for the isolation and detection of lactic acid bacteria with bacteriocinogenic potential. *LWT - Food Science and Technology*, 43(9), 1320-1324.
- Morelli, L. (2007). *In vitro* assessment of probiotic bacteria: From survival to functionality. *International Dairy Journal*, 17(11), 1278-1283.
- Morgan, C. A., Herman, N., White, P. A., & Vesey, G. (2006). Preservation of micro-organisms by drying; A review. *Journal of Microbiological Methods*, 66(2), 183-193.
- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*, 16, 55-63.
- Mowade, S., & Bhattacharyya, P. (2000). Resistance of P-solubilizing *Acetobacter diazotrophicus* to antibiotics, *Current Science*, 79(11), 1591-1594.
- Musikasang, H., Tani, A., H-kittikun, A., & Maneerat, S. (2009). Probiotic potential of lactic acid bacteria isolated from chicken gastrointestinal digestive tract. *World Journal of Microbiology and Biotechnology*, 25(8), 1337-1345.
- Muyanja, C. M. B. K., Narvhus, J. A., Treimo, J., & Langsrud, T. (2003). Isolation, characterisation and identification of lactic acid bacteria from bushera: A Ugandan traditional fermented beverage. *International Journal of Food Microbiology*, 80(3), 201-210.
- Najafi, F., Mozaffari, H. R., Karami, M., Izadi, B., Tavvafzadeh, R., & Pashar, Y. (2011). Trends in incidence of gastrointestinal tract cancers in western Iran, 1993-2007. *Iranian Red Crescent Medical Journal*, 13(11), 805–810.

- Nakai, S. A., & Siebert, K. J. (2003). Validation of bacterial growth inhibition models based on molecular properties of organic acids. *International Journal of Food Microbiology*, 86(3), 249-255.
- Nawaz, M., Wang, J., Zhou, A., Ma, C., Wu, X., Moore, J. E., Cherie Millar, B., & Xu, J. (2011). Characterization and transfer of antibiotic resistance in lactic acid bacteria from fermented food products. *Current Microbiology*, 62(3), 1081-1089.
- Ndoye, B., Weekers, F., Diawara, B., Guiro, A. T., & Thonart, P. (2007). Survival and preservation after freeze-drying process of thermoresistant acetic acid bacteria isolated from tropical products of Subsaharan Africa. *Journal of Food Engineering*, 79(4), 1374-1382.
- Noori, A., Keshavarzian, F., Mahmoudi, S., Yousefi, M., & Nateghi, L. (2013). Comparison of traditional Doogh (yogurt drinking) and Kashk characteristics (Two traditional Iranian dairy products). *European Journal of Experimental Biology*, 3(6), 252-255.
- Ogawa, M., Shimizu, K., Nomoto, K., Tanaka, R., Hamabata, T., Yamasaki, S., Takeda, T., & Takeda, Y. (2001). Inhibition of *in vitro* growth of Shiga toxin-producing *Escherichia coli* O157:H7 by probiotic *Lactobacillus* strains due to production of lactic acid. *International Journal of Food Microbiology*, 68(1-2), 135-140.
- Ogier, J. C., Casalta, E., Farrokh, C., & Saïhi, A. (2008). Safety assessment of dairy microorganisms: The *Leuconostoc* genus. *International Journal of Food Microbiology*, 126(3), 286-290.
- Oliveira, P. M., Zannini, E., & Arendt, E. K. (2013). Cereal fungal infection, mycotoxins, and lactic acid bacteria mediated bioprotection: from crop farming to cereal products. *Food Microbiology*, 1(18).
- Omar, N. B., Castro, A., Lucas, R., Abriouel, H., Yousif, N. M. K., Franz, C. M. A. P., Holzapfel, W. H., Rubén, P. P., Martínez-Canãmero, M., & Gálvez, A. (2004). Functional and safety aspects of *Enterococci* isolated from different Spanish foods. *Systematic and Applied Microbiology*, 27(1), 118-130.
- Otero, M. C., Espeche, M. C., & Nader-Macías, M. E. (2007). Optimization of the freeze-drying media and survival throughout storage of freeze-dried *Lactobacillus gasseri* and *Lactobacillus delbrueckii* subsp. *delbrueckii* for veterinarian probiotic applications. *Process Biochemistry*, 42(10), 1406-1411.
- Ouoba, L. I. I., Lei, V., & Jensen, L. B. (2008). Resistance of potential probiotic lactic acid bacteria and *bifidobacteria* of African and European origin to antimicrobials: Determination and transferability of the resistance genes to other bacteria. *International Journal of Food Microbiology*, 121(2), 217-224.
- Ouwehand, A. C., Kirjavainen, P. V., Grönlund, M. M., Isolauri, E., & Salminen, S. J. (1999). Adhesion of probiotic micro-organisms to intestinal mucus. *International Dairy Journal*, 9(9), 623-630.

- PAAG. (2009). World gastroenterology organisation practice guideline: Probiotics and prebiotics. Arab Journal of Gastroenterology, 10(1), 33-42.
- Paik, G. (1980). Reagents, stains and miscellaneous test procedures, Manual of Clinical Microbiology. Washington, DC: American Society for Microbiology.
- Pan, X., Chen, F., Wu, T., Tang, H., & Zhao, Z. (2009). The acid, bile tolerance and antimicrobial property of *Lactobacillus acidophilus* NIT. Food Control, 20(6), 598-602.
- Paolillo, R., Romano Carratelli, C., Sorrentino, S., Mazzola, N., & Rizzo, A. (2009). Immunomodulatory effects of *Lactobacillus plantarum* on human colon cancer cells. International Immunopharmacology, 9(11), 1265-1271.
- Papagianni, M., & Anastasiadou, S. (2009). Encapsulation of *Pediococcus acidilactici* cells in corn and olive oil microcapsules emulsified by peptides and stabilized with xanthan in oil-in-water emulsions: Studies on cell viability under gastro-intestinal simulating conditions. Enzyme and Microbial Technology, 45(6-7), 514-522.
- Park, S. C., Hwang, M. H., Kim, Y. H., Kim, J. C., Song, J. C., Lee, K. W., Jeong, K. S., Rhee, M. H., Kim, K. S., & Kim, T. W. (2006). Comparison of pH and bile resistance of *Lactobacillus acidophilus* strains isolated from rat, pig, chicken, and human sources. World Journal of Microbiology and Biotechnology, 22(1), 35-37.
- Park, Y. K., Row, K. H., & Chung, S. T. (2000). Adsorption characteristics and separation of taxol from yew tree by NP-HPLC. Separation and Purification Technology, 19(1-2), 27-37.
- Parkes, G. C. (2010). Chapter 30 - The role of probiotics in the treatment of irritable bowel syndrome. In W. Ronald Ross & R. P. Victor (Eds.), Bioactive Foods in Promoting Health (pp. 513-528). Boston: Academic Press.
- Parkin, D. M. (2001). Global cancer statistics in the year 2000. The Lancet Oncology, 2(9), 533-543.
- Paseephol, T., & Sherkat, F. (2009). Probiotic stability of yoghurts containing Jerusalem artichoke inulins during refrigerated storage. Journal of Functional Foods, 1(3), 311-318.
- Patel, R. (2000). Enterococcal-type glycopeptide resistance genes in non-enterococcal organisms. FEMS Microbiology Letters, 185(1), 1-7.
- Patterson, J.A. & Burkholder, K.M. (2003). Application of prebiotics and probiotics in poultry production. Poultry Science, 82(4), 627-631.
- Peniche, C., Howland, I., Carrillo, O., Zaldívar, C., & Argüelles-Monal, W. (2004). Formation and stability of shark liver oil loaded chitosan/calcium alginate capsules. Food Hydrocolloids, 18(5), 865-871.
- Pepe, O., Blajotta, G., Anastasio, M., Moschetti, G., Ercolini, D., & Villani, F. (2004). Technological and molecular diversity of *Lactobacillus plantarum* strains isolated

- from naturally fermented sourdoughs. *Systematic and Applied Microbiology*, 27(4), 443-453.
- Pérez Elortondo, F. J., Aldámiz Echobarria, P., Albisu, M., & Barcina, Y. (1998). Indigenous lactic acid bacteria in Idiazábal Ewes' milk cheese. *International Dairy Journal*, 8(8), 725-732.
- Perreten, V., Kollöffel, B., & Teuber, M. (1997). Conjugal transfer of the Tn916-like transposon TnFO1 from *Enterococcus faecalis* isolated from cheese to other gram-positive bacteria. *Systematic and Applied Microbiology*, 20(1), 27-38.
- Peters, R. K., Pike, M. C., Garabrant, D., & Mack, T. M. (1992). Diet and colon cancer in Los Angeles County, California. *Cancer Causes and Control*, 3(5), 457-473.
- Picot, A., & Lacroix, C. (2003). Effects of micronization on viability and thermotolerance of probiotic freeze-dried cultures. *International Dairy Journal*, 13(6), 455-462.
- Pieper, R., Janczyk, P., Urubschurov, V., Hou, Z., Korn, U., Pieper, B., & Souffrant, W. B. (2010). Effect of *Lactobacillus plantarum* on intestinal microbial community composition and response to enterotoxigenic *Escherichia coli* challenge in weaning piglets. *Livestock Science*, 133(1-3), 98-100.
- Pogačić, T., Mancini, A., Santarelli, M., Bottari, B., Lazzi, C., Neviani, E., & Gatti, M. (2013). Diversity and dynamic of lactic acid bacteria strains during aging of a long ripened hard cheese produced from raw milk and undefined natural starter. *Food Microbiology*, 36(2), 207-215.
- Pontes, D. S., Lima-Bittencourt, C. I., Chartone-Souza, E., & Amaral Nascimento, A. M. (2007). Molecular approaches: advantages and artifacts in assessing bacterial diversity. *Journal of Industrial Microbiology and Biotechnology*, 34(7), 463-473.
- Pot, B., Devriese, L. A., Ursi, D., Vandamme, P., Haesebrouck, F., & Kersters, K. (1996). Phenotypic identification and differentiation of *Lactococcus* strains isolated from animals. *Systematic and Applied Microbiology*, 19(2), 213-222.
- Potten, C. S., & Grant, H. K. (1998). The relationship between ionizing radiation-induced apoptosis and stem cells in the small and large intestine. *British Journal of Cancer*, 78(8), 993-1003.
- Prinsloo, S., Pieters, R., & Bezuidenhout, C.C. (2013). A cell viability assay to determine the cytotoxic effects of water contaminated by microbes. *South African Journal of Science*, 109(7/8), 1-4.
- Psoni, L., Kotzamanidis, C., Yiangou, M., Tzanetakis, N., & Litopoulou-Tzanetaki, E. (2007). Genotypic and phenotypic diversity of *Lactococcus lactis* isolates from Batzos, a Greek PDO raw goat milk cheese. *International Journal of Food Microbiology*, 114(2), 211-220.

- Psoni, L., Tzanetakis, N., & Litopoulou-Tzanetaki, E. (2003). Microbiological characteristics of Batzos, a traditional Greek cheese from raw goat's milk. *Food Microbiology*, 20(5), 575-582.
- Rafter, J. (2002). Lactic acid bacteria and cancer: mechanistic perspective. *British Journal of Nutrition*, 88(1), 89-94.
- Rafter, J. (2003). Probiotics and colon cancer. *Best Practice & Research Clinical Gastroenterology*, 17(5), 849-859.
- Rahmati, M., Bannazadeh, H., Fazli, D., Tahmasebzadeh, F., Farrokhi, A., & Rasmi, Y. (2011). The apoptotic and cytotoxic effects of *Polygonum aviculare* extract on HeLa-S cervical cancer cell line. *African Journal of Biochemistry Research*, 5(14), 373-378.
- Rašić, J. L., & Kurmann, J. A. (1983). *Bifidobacteria* and their role: microbiological, nutritional-physiological, medical, and technological aspects and bibliography. *Experientia. Supplementum*, 39, 1-295.
- Reddy, B. S., Narisawa, T., Wright, P., Vukusich, D., Weisburger, J. H., & Wynder, E. L. (1975). Colon carcinogenesis with azoxymethane and dimethylhydrazine in germ-free rats. *Cancer Research*, 35(2), 287-290.
- Reddy, B. S., & Rivenson, A. (1993). Inhibitory effect of *Bifidobacterium longum* on colon, mammary, and liver carcinogenesis induced by 2-amino-3-methylimidazo[4,5-f]quinoline, a food mutagen. *Cancer Research*, 53(17), 3914-3918.
- Reid, G. (2002). The potential role of probiotics in pediatric urology. *The Journal of Urology*, 168(4), 1512-1517.
- Reid, G., & Bocking, A. (2003). The potential for probiotics to prevent bacterial vaginosis and preterm labor. *American Journal of Obstetrics and Gynecology*, 189(4), 1202-1208.
- Reid, G., & Burton, J. (2002). Use of *Lactobacillus* to prevent infection by pathogenic bacteria. *Microbes and Infection*, 4(3), 319-324.
- Renner, H. W., & Münzner, R. (1991). The possible role of probiotics as dietary antimutagens. *Mutation Research Letters*, 262(4), 239-245.
- Reyed, R. (2006). Novel hybrid entrapment approach for probiotic cultures and its application during lyophilization. *The Internet Journal of Microbiology*, 3(2).
- Ringo, E., & Gatesoupe, F. J. (1998). Lactic acid bacteria in fish: a review. *Aquaculture*, 160(3-4), 177-203.
- Rishniw, M., & Wynn, S. G. (2011). Azodyl, a synbiotic, fails to alter azotemia in cats with chronic kidney disease when sprinkled onto food. *Journal of Feline Medicine & Surgery*, 13(6), 405-409.

- Rivas, F. P., Castro, M. P., Vallejo, M., Marguet, E., & Campos, C. A. (2012). Antibacterial potential of *Enterococcus faecium* strains isolated from ewes' milk and cheese. *LWT - Food Science and Technology*, 46(2), 428-436.
- Rivkin, I., Cohen, K., Bod, T., Argov, M., & Margalit, R. (2014). Cancer cell sensitization and improved treatment efficacy by combined sodium butyrate and paclitaxel formulations is cancer-type specific. *International Journal of Pharmaceutics*, 461(1–2), 437-447.
- Rizzo, A., Losacco, A., Carratelli, C. R., Domenico, M. D., & Bevilacqua, N. (2013). *Lactobacillus plantarum* reduces *Streptococcus pyogenes* virulence by modulating the IL-17, IL-23 and Toll-like receptor 2/4 expressions in human epithelial cells. *International Immunopharmacology*, 17(2), 453-461.
- Robinson, R. K. (1987). Survival of *Lactobacillus acidophilus* in fermented products. *South African Society of Dairy Technology*, 19(1), 10725-10727.
- Rodas, A. M., Ferrer, S., & Pardo, I. (2003). 16S-ARDRA, a tool for identification of lactic acid bacteria isolated from grape must and wine. *Systematic and Applied Microbiology*, 26(3), 412-422.
- Rodrigues, D., Sousa, S., Rocha-Santos, T., Silva, J. P., Sousa Lobo, J. M., Costa, P., Amaral, M. H., Pintado, M. M., Gomes, A. M., Malcata, F. X., & Freitas, A. C. (2011). Influence of l-cysteine, oxygen and relative humidity upon survival throughout storage of probiotic bacteria in whey protein-based microcapsules. *International Dairy Journal*, 21(11), 869-876.
- Rönkä, E., Malinen, E., Saarela, M., Rinta-Koski, M., Aarnikunnas, J., & Palva, A. (2003). Probiotic and milk technological properties of *Lactobacillus brevis*. *International Journal of Food Microbiology*, 83(1), 63-74.
- Roy, D., & Sirois, S. (2000). Molecular differentiation of *Bifidobacterium* species with amplified ribosomal DNA restriction analysis and alignment of short regions of the *ldh* gene. *FEMS Microbiology Letters*, 191(1), 17-24.
- Rubio, R., Bover-Cid, S., Martin, B., Garriga, M., & Aymerich, T. (2013). Assessment of safe *enterococci* as bioprotective cultures in low-acid fermented sausages combined with high hydrostatic pressure. *Food Microbiology*, 33(2), 158-165.
- Ryu, E. H., Yang, E. J., Woo, E. R., & Chang, H. C. (2014). Purification and characterization of antifungal compounds from *Lactobacillus plantarum* HD1 isolated from kimchi. *Food Microbiology*, 41, 19-26.
- Saarela, M., Maukonen, J., von Wright, A., Vilpponen-Salmela, T., Patterson, A. J., Scott, K. P., Hämynen, Heikki., & Mättö, J. (2007). Tetracycline susceptibility of the ingested *Lactobacillus acidophilus* LaCH-5 and *Bifidobacterium animalis* subsp. *lactis* Bb-12 strains during antibiotic/probiotic intervention. *International Journal of Antimicrobial Agents*, 29(3), 271-280.
- Saavedra, L., Taranto, M., Sesma, F., & de Valdez, G. F. (2003). Homemade traditional cheeses for the isolation of probiotic *Enterococcus faecium* strains. *International Journal of Food Microbiology*, 88(2–3), 241-245.

- Sahadeva, R. P. K., Leong, S. F., Chua, K. H., Tan, C. H., Chan, H. Y., Tong, E. V., Wong, S.Y.W., & Chan, H. K. (2011). Survival of commercial probiotic strains to pH and bile. *International Food Research Journal*, 18(4), 1515-1522.
- Sakai, S., Ono, T., Ijima, H., Kawakami, K. (2002). Permeability of alginate/sol-gel synthesized aminopropyl-silicate/alginate membrane templated by calcium-alginate gel. *Journal of Membrane Science*, 205(1-2), 183-189.
- Salar-Behzadi, S., Wu, S., Toegel, S., Hofrichter, M., Altenburger, I., Unger, F. M., Wirth, M., & Viernstein, H. (2013). Impact of heat treatment and spray drying on cellular properties and culturability of *Bifidobacterium bifidum* BB-12. *Food Research International*, 54(1), 93-101.
- Salveti, E., Fondi, M., Fani, R., Torriani, S., & Felis, G. E. (2013). Evolution of lactic acid bacteria in the order *Lactobacillales* as depicted by analysis of glycolysis and pentose phosphate pathways. *Systematic and Applied Microbiology*, 36(5), 291-305.
- Samarzija, D., Sikora, S., Redzepovic, S., Antunac, N., & Havranek, J. (2002). Application of RAPD analysis for identification of *Lactococcus lactis* subsp. *cremoris* strains isolated from artisanal cultures. *Microbiological Research*, 157(1), 13-17.
- Sanchez, I., Sesenã, S., & Palop, L. L. (2004). Polyphasic study of the genetic diversity of *lactobacilli* associated with 'Almagro' eggplants spontaneous fermentation based on combined numerical analysis of randomly amplified polymorphic DNA and pulse-field gel electrophoresis patterns. *Journal of Applied Microbiology*, 97, 446-458.
- Sandoval-Castilla, O., Lobato-Calleros, C., García-Galindo, H. S., Alvarez-Ramírez, J., & Vernon-Carter, E. J. (2010). Textural properties of alginate-pectin beads and survivability of entrapped *Lb. casei* in simulated gastrointestinal conditions and in yoghurt. *Food Research International*, 43(1), 111-117.
- Sarantinopoulos, P., Andrighetto, C., Georgalaki, M. D., Rea, M. C., Lombardi, A., Cogan, T. M., Kalantzopoulos, G., & Tsakalidou, E. (2001). Biochemical properties of *enterococci* relevant to their technological performance. *International Dairy Journal*, 11(8), 621-647.
- Sathishkumar, G., Gobinath, G., Wilson, A., Sivaramakrishnan, S. (2014). *Dendrophthoe falcata* (L.f) Ettingsh (Neem mistletoe): A potent bioresource to fabricate silver nanoparticles for anticancer effect against human breast cancer cells (MCF-7). *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 128, 285-290.
- Sathyabama, S., Ranjith kumar, M., Bruntha devi, P., Vijayabharathi, R., & Brindha priyadharisini, V. (2014). Co-encapsulation of probiotics with prebiotics on alginate matrix and its effect on viability in simulated gastric environment. *LWT - Food Science and Technology*. 57(1), 419-425.

- Satokari, R. M., Vaughan, E. E., Smidt, H., Saarela, M., Mättö, J., & de Vos, W. M. (2003). Molecular approaches for the detection and identification of *Bifidobacteria* and *Lactobacilli* in the human gastrointestinal tract. *Systematic and Applied Microbiology*, 26(4), 572-584.
- Scheinbach, S. (1998). Probiotics: Functionality and commercial status. *Biotechnology Advances*, 16(3), 581-608.
- Schleifer, K. H., & Kilpper-Bälz, R. (1987). Molecular and chemotaxonomic approaches to the classification of *Streptococci*, *Enterococci* and *Lactococci*: A Review. *Systematic and Applied Microbiology*, 10(1), 1-19.
- Schleifer, K. H., Kraus, J., Dvorak, C., Kilpper-Bälz, R., Collins, M. D., & Fischer, W. (1985). Transfer of *Streptococcus lactis* and related *Streptococci* to the genus *Lactococcus* gen. nov. *Systematic and Applied Microbiology*, 6(2), 183-195.
- Schoug, Å., Olsson, J., Carlfors, J., Schnürer, J., & Håkansson, S. (2006). Freeze-drying of *Lactobacillus coryniformis* Si3—effects of sucrose concentration, cell density, and freezing rate on cell survival and thermophysical properties. *Cryobiology*, 53(1), 119-127.
- Sengun, I. Y., & Karabiyikli, S. (2011). Importance of acetic acid bacteria in food industry. *Food Control*, 22(5), 647-656.
- Sengun, I. Y., Nielsen, D. S., Karapinar, M., & Jakobsen, M. (2009). Identification of lactic acid bacteria isolated from Tarhana, a traditional Turkish fermented food. *International Journal of Food Microbiology*, 135(2), 105-111.
- Senok, A. C., Ismaeel, A. Y., & Botta, G. A. (2005). Probiotics: facts and myths. *Clinical Microbiology and Infection*, 11(12), 958-966.
- Seo, B. J., Rather, I. A., Kumar, V. J. R., Choi, U. H., Moon, M. R., Lim, J. H., & Park, Y. H. (2012). Evaluation of *Leuconostoc mesenteroides* YML003 as a probiotic against low-pathogenic avian influenza (H9N2) virus in chickens. *Journal of Applied Microbiology*, 113(1), 163-171.
- Serban, D. E. (2013). Gastrointestinal cancers: Influence of gut microbiota, probiotics and prebiotics. *Cancer Letters*, 345(2), 258-270.
- Servin, A. L. (2004). Antagonistic activities of *lactobacilli* and *bifidobacteria* against microbial pathogens. *FEMS Microbiology Reviews*, 28(4), 405-440.
- Settanni, L., Guarcello, R., Gaglio, R., Francesca, N., Aleo, A., Felis, G. E., & Moschetti, G. (2014). Production, stability, gene sequencing and *in situ* anti-*Listeria* activity of mundticin KS expressed by three *Enterococcus mundtii* strains. *Food Control*, 35(1), 311-322.
- Sghir, A., Gramet, G., Suau, A., Rochet, V., Pochart, P., & Dore, J. (2000). Quantification of bacterial groups within human fecal flora by oligonucleotide probe hybridization. *Applied and Environmental Microbiology*, 66, 2263-2266.

- Shah, N. P. (2000). Probiotic bacteria: Selective enumeration and survival in dairy foods. *Journal of Dairy Science*, 83(4), 894-907.
- Shahani, K. M., & Ayebo, A. D. (1980). Role of dietary *lactobacilli* in gastrointestinal microecology. *American Journal of Clinical Nutrition*, 33(11), 2448-2457.
- Shi, L. E., Li, Z. H., Li, D. T., Xu, M., Chen, H. Y., Zhang, Z. L., & Tang, Z. X. (2013a). Encapsulation of probiotic *Lactobacillus bulgaricus* in alginate-milk microspheres and evaluation of the survival in simulated gastrointestinal conditions. *Journal of Food Engineering*, 117(1), 99-104.
- Shi, L. E., Li, Z. H., Zhang, Z. L., Zhang, T. T., Yu, W. M., Zhou, M. L., & Tang, Z. X. (2013b). Encapsulation of *Lactobacillus bulgaricus* in carrageenan-locust bean gum coated milk microspheres with double layer structure. *LWT - Food Science and Technology*, 54(1), 147-151.
- Shida, K., Kiyoshima-Shibata, J., Nagaoka, M., Watanabe, K., & Nanno, M. (2006). Induction of interleukin-12 by *Lactobacillus* strains having a rigid cell wall resistant to intracellular digestion. *Journal of Dairy Science*, 89(9), 3306-3317.
- Shinde, T., Sun-Waterhouse, D., & Brooks, J. (2014). Co-extrusion encapsulation of probiotic *Lactobacillus acidophilus* alone or together with apple skin polyphenols: An aqueous and value-added delivery system using alginate. *Food and Bioprocess Technology*, 7(6), 1581-1596.
- Shoma, S., Kamruzzaman, M., Ginn, A. N., Iredell, J. R., & Partridge, S. R. (2014). Characterization of multidrug-resistant *Klebsiella pneumoniae* from Australia carrying blaNDM-1. *Diagnostic Microbiology and Infectious Disease*, 78(1), 93-97.
- Shu, Q., & Gill, H. S. (2002). Immune protection mediated by the probiotic *Lactobacillus rhamnosus* HN001 (DR20™) against *Escherichia coli* O157:H7 infection in mice. *FEMS Immunology and Medical Microbiology*, 34(1), 59-64.
- Shukla, R., Iliev, I., & Goyal, A. (2014). *Leuconostoc mesenteroides* NRRL B-1149 as probiotic and its dextran with anticancer properties. *Journal of Bioscience and Biotechnology*, 3(1), 79-87.
- Sievers, M., Gaberthüel, C., Boesch, C., Ludwig, W., & Teuber, M. (1995). Phylogenetic position of *Gluconobacter* species as a coherent cluster separated from all *Acetobacter* species on the basis of 16S ribosomal RNA sequences. *FEMS Microbiology Letters*, 126(2), 123-126.
- Simon, O., Jadamus, A. & Vahjen, W. (2001). Probiotic feed additives-effectiveness and expected modes of action. *Journal of Animal Feed Science*, 10(1), 51-67.
- Simpson, P. J., Fitzgerald, G. F., Stanton, C., & Ross, R. P. (2006). Enumeration and identification of *Pediococci* in powder-based products using selective media and rapid PFGE. *Journal of Microbiological Methods*, 64(1), 120-125.

- Singh, B. (2007). Psyllium as therapeutic and drug delivery agent. *International Journal of Pharmaceutics*, 334(1–2), 1-14.
- Singh, J., Rivenson, A., Tomita, M., Shimamura, S., Ishibashi, N., & Reddy, B. S. (1997). *Bifidobacterium longum*, a lactic acid-producing intestinal bacterium inhibits colon cancer and modulates the intermediate biomarkers of colon carcinogenesis. *Carcinogenesis*, 18(4), 833-841.
- Singh, N. P., Tiwari, A., Bansal, A., Thakur, S., Sharma, G., & Gabrani, R. (2015). Genome level analysis of bacteriocins of lactic acid bacteria. *Computational Biology and Chemistry*, 56(0), 1-6.
- Singh, S., & Goswami, P. (2009). Application of molecular identification tools for *Lactobacillus*, with a focus on discrimination between closely related species. *LWT - Food Science and Technology*, 42(2), 448-457.
- Singh, S., Goswami, P., Singh, R., & Heller, K. J. (2009). Application of molecular identification tools for *Lactobacillus*, with a focus on discrimination between closely related species: A review. *LWT - Food Science and Technology*, 42(2), 448-457.
- Siragusa, S., De Angelis, M., Calasso, M., Campanella, D., Minervini, F., Di Cagno, R., & Gobbetti, M. (2014). Fermentation and proteome profiles of *Lactobacillus plantarum* strains during growth under food-like conditions. *Journal of Proteomics*, 96, 366-380.
- Smit, G., Smit, B. A., & Engels, W. J. M. (2005). Flavour formation by lactic acid bacteria and biochemical flavour profiling of cheese products. *FEMS Microbiology Reviews*, 29(3), 591-610.
- Sohail, A., Turner, M. S., Coombes, A., Bostrom, T., & Bhandari, B. (2011). Survivability of probiotics encapsulated in alginate gel microbeads using a novel impinging aerosols method. *International Journal of Food Microbiology*, 145(1), 162-168.
- Solanki, H. K., Pawar, D. D., Shah, D. A., Prajapati, V. D., Jani, G. K., Mulla, A. M., & Thakar, P. M. (2013). Development of microencapsulation delivery system for long-term preservation of probiotics as biotherapeutics agent. *BioMed Research International*, 2013, 1-21.
- Solieri, L., Bianchi, A., & Giudici, P. (2012). Inventory of non starter lactic acid bacteria from ripened Parmigiano Reggiano cheese as assessed by a culture dependent multiphasic approach. *Systematic and Applied Microbiology*, 35, 270-277.
- Song, S., & Lambert, P. (1999). Different responses of epidermal and hair follicular cells to radiation correlate with distinct patterns of p53 and p21 induction. *The American Journal of Pathology*, 155(4), 1121-1127.
- Song, Y.-L., Kato, N., Liu, C.-X., Matsumiya, Y., Kato, H., & Watanabe, K. (2000). Rapid identification of 11 human intestinal *Lactobacillus* species by multiplex

- PCR assays using group- and species-specific primers derived from the 16S–23S rRNA intergenic spacer region and its flanking 23S rRNA. *FEMS Microbiology Letters*, 187(2), 167-173.
- Sorrells, K. M., & Speck, M. L. (1970). Inhibition of *Salmonella gallinarum* by culture filtrates of *Leuconostoc citrovorum*. *Journal of Dairy Science*, 53(2), 239-241.
- Sousa, S., Gomes, A. M., Pintado, M. M., Silva, J. P., Costa, P., Amaral, M. H., Duarte, A. C., Rodrigues, D., Rocha-Santos, T. A. P., & Freitas, A. C. (2013). Characterization of freezing effect upon stability of, probiotic loaded, calcium-alginate microparticles. *Food and Bioproducts Processing*.
- Spinler, J. K., Taweechotipatr, M., Rognerud, C. L., Ou, C. N., Tumwasorn, S., & Versalovic, J. (2008). Human-derived probiotic *Lactobacillus reuteri* demonstrate antimicrobial activities targeting diverse enteric bacterial pathogens. *Anaerobe*, 14(3), 166-171.
- Stierle, A. A., & D. B, S. (2000). Bioactive compounds from four endophytic *Penicillium* sp. of a northwest pacific yew tree. *Studies in Natural Products Chemistry*, 24 (Part E), 933-977.
- Strobel, G. A., Stierle, A., & Hess, W. M. (1993). Taxol formation in yew — *Taxus*. *Plant Science*, 92(1), 1-12.
- Strompfová, V., Lauková, A., & Ouwehand, A. C. (2004). Selection of *enterococci* for potential canine probiotic additives. *Veterinary Microbiology*, 100(1–2), 107-114.
- Stummer, S., Salar-Behzadi, S., Unger, F. M., Oelzant, S., Penning, M., & Viernstein, H. (2010). Application of shellac for the development of probiotic formulations. *Food Research International*, 43(5), 1312-1320.
- Stummer, S., Toegel, S., Rabenreither, M.-C., Unger, F. M., Wirth, M., Viernstein, H., & Salar-Behzadi, S. (2012). Fluidized-bed drying as a feasible method for dehydration of *Enterococcus faecium* M74. *Journal of Food Engineering*, 111(1), 156-165.
- Su, J. M., Wang, L. Y., Liang, Y. L., & Zha, X. L. (2005). Role of cell adhesion signal molecules in hepatocellular carcinoma cell apoptosis. *World Journal of Gastroenterology*, 11(30), 4667-4673.
- Sultana, K., Godward, G., Reynolds, N., Arumugaswamy, R., Peiris, P., & Kailasapathy, K. (2000). Encapsulation of probiotic bacteria with alginate-starch and evaluation of survival in simulated gastrointestinal conditions and in yogurt. *International Journal of Food Microbiology*, 62(1-2), 47-55.
- Surawicz, C. M. (2003). Probiotics, antibiotic-associated diarrhoea and *Clostridium difficile* diarrhoea in humans. *Best Practice & Research Clinical Gastroenterology*, 17(5), 775-783.
- Švec, P., Pantůček, R., Petráš, P., Sedláček, I., & Nováková, D. (2010). Identification of *Staphylococcus* spp. using (GTG) 5-PCR fingerprinting. *Systematic and Applied Microbiology*, 33(8), 451-456.

- Švec, P., Vancanneyt, M., Seman, M., Snauwaert, C., Lefebvre, K., Sedláček, I., & Swings, J. (2005). Evaluation of (GTG)₅-PCR for identification of *Enterococcus* spp. FEMS Microbiology Letters, 247(1), 59-63.
- Swenson, J. M., Facklan, R. R., & Thornsberry, C. (1990). Antimicrobial susceptibility of vancomycin-resistant *Leuconostoc*, *Pediococcus*, and *Lactobacillus* species. Antimicrobial Agents and Chemotherapy, 34(4), 543-549.
- Tabatabaee, F., Alizadeh Behbahani, B., Mohebbi, M., Mortazavi, S. A., & Ghaitaranpour, A. (2012). Identification of lactic acid bacteria isolated from Tarkhineh, a traditional Iranian fermented food. Scientific Journal of Microbiology, 1(7), 152-159.
- Tabatabaei Yazdi, F., Alizadeh Behbahani, B., & Mortazavi, A. (2013). Effect of temperature and salt concentration on microbial changes during Tarkhineh fermentation. Scientific Journal of Biological Sciences, 2(1), 8-16.
- Tafvizi, F., & Tajabadi Ebrahimi, M. (2012). Detection of genetic diversity and classification of *Lactobacillus* species isolated from Iranian traditional dairy products by RAPD fingerprinting and POPGENE analysis Annals of Biological Research, 3(10), 4904-4911.
- Tahara, N., Yano, H., & Yoshinaga, F. (1997). Two types of cellulase activity produced by a cellulose-producing *Acetobacter* strain. Journal of Fermentation and Bioengineering, 83(4), 389-392.
- Temmerman, R., Huys, G., & Swings, J. (2004). Identification of lactic acid bacteria: culture-dependent and culture-independent methods. Trends in Food Science & Technology, 15(7-8), 348-359.
- Temmerman, R., Pot, B., Huys, G., & Swings, J. (2003). Identification and antibiotic susceptibility of bacterial isolates from probiotic products. International Journal of Food Microbiology, 81(1), 1-10.
- Theunissen, J., Britz, T. J., Torriani, S., & Witthuhn, R. C. (2005). Identification of probiotic microorganisms in South African products using PCR-based DGGE analysis. International Journal of Food Microbiology, 98(1), 11-21.
- Todorov, S., Onno, B., Sorokine, O., Chobert, J. M., Ivanova, I., & Dousset, X. (1999). Detection and characterization of a novel antibacterial substance produced by *Lactobacillus plantarum* ST 31 isolated from sourdough. International Journal of Food Microbiology, 48(3), 167-177.
- Todorov, S. D., Wachsmann, M. B., Knoetze, H., Meincken, M., & Dicks, L. M. T. (2005). An antibacterial and antiviral peptide produced by *Enterococcus mundtii* ST4V isolated from soya beans. International Journal of Antimicrobial Agents, 25(6), 508-513.
- Trabelsi, I., Bejar, W., Ayadi, D., Chouayekh, H., Kammoun, R., Bejar, S., & Ben Salah, R. (2013). Encapsulation in alginate and alginate coated-chitosan improved

- the survival of newly probiotic in oxgall and gastric juice. *International Journal of Biological Macromolecules*, 61, 36-42.
- Trias, R., Badosa, E., Montesinos, E., & Bañeras, L. (2008). Bioprotective *Leuconostoc* strains against *Listeria monocytogenes* in fresh fruits and vegetables. *International Journal of Food Microbiology*, 127(1-2), 91-98.
- Truelstrup-Hansen, L., Allan-wojtas, P. M., Jin, Y. L., & Paulson, A. T. (2002). Survival of free and calcium-alginate microencapsulated *Bifidobacterium* spp. in stimulated gastro-intestinal conditions. *Food Microbiology*, 19, 35-45.
- Tulini, F. L., Winkelströter, L. K., & De Martinis, E. C. P. (2013). Identification and evaluation of the probiotic potential of *Lactobacillus paraplantarum* FT259, a bacteriocinogenic strain isolated from Brazilian semi-hard artisanal cheese. *Anaerobe*, 22, 57-63.
- Tuohy, K. M., Probert, H. M., Smejkal, C. W., & Gibson, G. R. (2003). Using probiotics and prebiotics to improve gut health. *Drug Discovery Today*, 8(15), 692-700.
- Tynkkynen, S., Satokari, R., Saarela, M., Mattila Sandholm, T., & Saxelin, M. (1999). Comparison of ribotyping, randomly amplified polymorphic DNA analysis, and pulsedfield gel electrophoresis in typing of *Lactobacillus rhamnosus* and *L. casei* strains. *Applied and Environmental Microbiology*, 65, 3908-3914.
- Van't Veer, P., Dekker, J. M., Lamers, J. W. J., Kok, F. J., Schouten, E. G., Brants, H. A. M., Sturmans, F., & Hermus, R. J. J. (1989). Consumption of fermented milk products and breast cancer: A case-control study in The Netherlands. *Cancer Research*, 49(14), 4020-4023.
- Van Hoorde, K., Verstraete, T., Vandamme, P., & Huys, G. (2008). Diversity of lactic acid bacteria in two Flemish artisan raw milk Gouda-type cheeses. *Food Microbiology*, 25(7), 929-935.
- Vaugien, L., Prevots, F., & Roques, C. (2002). *Bifidobacteria* identification based on 16S rRNA and pyruvate kinase partial gene sequence analysis. *Anaerobe*, 8(6), 341-344.
- Vedamuthu, E. R. (1994). The Dairy *Leuconostoc*: Use in dairy products. *Journal of Dairy Science*, 77(9), 2725-2737.
- Veera Jothi, V., Anandapandian, K. T. K., & Shankar, T. (2012). Bacteriocin production by probiotic bacteria from curd and its field application to poultry. *Archives of Applied Science Research*, 4(1), 336-347.
- Velicanski, A. S., Cvetkovic, D. D., Markov, S. L., Tumbas, V. T., & Savatovi, S. M. (2007). Antimicrobial and antioxidant activity of lemon balm kombucha. *Original Scientific Paper*, 38, 165-172.
- Ventura, M., Casas, I. A., Morelli, L., & Callegari, M. L. (2000). Rapid amplified ribosomal DNA restriction analysis (ARDRA) identification of *Lactobacillus* spp.

- isolated from fecal and vaginal samples. *Systematic and Applied Microbiology*, 23(4), 504-509.
- Ventura, M., Meylan, V., & Zink, R. (2003). Identification and tracing of *Bifidobacterium* species by use of enterobacterial repetitive intergenic consensus sequences. *Applied and Environmental Microbiology*, 69(7), 4296-4301.
- Ventura, M., & Zink, R. (2002). Specific identification and molecular typing analysis of *Lactobacillus johnsonii* by using PCR-based methods and pulsed-field gel electrophoresis. *FEMS Microbiology Letters*, 217(2), 141-154.
- Vera Pingitore, E., Todorov, S. D., Sesma, F., & Gombossy de Melo Franco, B. D. (2012). Application of bacteriocinogenic *Enterococcus mundtii* CRL35 and *Enterococcus faecium* ST88Ch in the control of *Listeria monocytogenes* in fresh Minas cheese. *Food Microbiology*, 32(1), 38-47.
- Vermes, I., Haanen, C., Steffens-Nakken, H., & Reutelingsperger, C. (1995). A novel assay for apoptosis Flow cytometric detection of phosphatidylserine early apoptotic cells using fluorescein labelled expression on Annexin V. *Journal of Immunological Methods*, 184, 39-51.
- Vidhyalakshmi, R., Bhakayaraj, R., & Subhasree, R. S. (2009). Encapsulation "The future of probiotics"-A Review. *Advances in Biological Research*, 3(3-4), 96-103.
- Vincent, D., Roy, D., Mondou, F., & Déry, C. (1998). Characterization of *bifidobacteria* by random DNA amplification. *International Journal of Food Microbiology*, 43(3), 185-193.
- Vinh, D. C., Nichol, K. A., Rand, F., & Embil, J. M. (2006). Native-valve bacterial endocarditis caused by *Lactococcus garvieae*. *Diagnostic Microbiology and Infectious Disease*, 56(1), 91-94.
- Voo, W. P., Ravindra, P., Tey, B. T., & Chan, E. S. (2011). Comparison of alginate and pectin based beads for production of poultry probiotic cells. *Journal of Bioscience and Bioengineering*, 111(3), 294-299.
- Vrancken, G., Rimaux, T., De Vuyst, L., & Leroy, F. (2008). Kinetic analysis of growth and sugar consumption by *Lactobacillus fermentum* IMDO 130101 reveals adaptation to the acidic sourdough ecosystem. *International Journal of Food Microbiology*, 128(1), 58-66.
- Walker, D. K., & Gilliland, S. E. (1993). Relationships among bile tolerance, bile salt deconjugation, and assimilation of cholesterol by *Lactobacillus acidophilus*. *Journal of Dairy Science*, 76(4), 956-961.
- Wang, W., Waterhouse, G. I. N., & Sun-Waterhouse, D. (2013). Co-extrusion encapsulation of canola oil with alginate: Effect of quercetin addition to oil core and pectin addition to alginate shell on oil stability. *Food Research International*, 54(1), 837-851.

- Ward, P., & Roy, D. (2005). Review of molecular methods for identification, characterization and detection of *bifidobacteria*. *Dairy Science and Technology*, 85, 23-32.
- Weinbreck, F., Bodnár, I., & Marco, M. L. (2010). Can encapsulation lengthen the shelf-life of probiotic bacteria in dry products? *International Journal of Food Microbiology*, 136(3), 364-367.
- Werner, G., Coque, T. M., Franz, C. M. A. P., Grohmann, E., Hegstad, K., Jensen, L., Van Schaik, W., & Weaver, K. (2013). Antibiotic resistant enterococci—Tales of a drug resistance gene trafficker. *International Journal of Medical Microbiology*, 303(6-7), 360-379.
- Wickens, K., Black, P. N., Stanley, T. V., Mitchell, E., Fitzharris, P., Tannock, G. W., Purdie, G., & Crane, J. (2008). A differential effect of 2 probiotics in the prevention of eczema and atopy: A double-blind, randomized, placebo-controlled trial. *Journal of Allergy and Clinical Immunology*, 122(4), 788-794.
- Wijtzes, T., Bruggeman, M. R., Nout, M. J. R., & Zwietering, M. H. (1997). A computerised system for the identification of lactic acid bacteria. *International Journal of Food Microbiology*, 38(1), 65-70.
- Williams, A. M., Fryer, J. L., & Collins, M. D. (1990). *Lactococcus piscium* sp. nov. a new *Lactococcus* species from salmonid fish. *FEMS Microbiology Letters*, 68(1-2), 109-113.
- Woodford, N., Johnson, A. P., Morrison, D., & Speller, D. C. E. (1995). Current perspective on glycopeptide resistance. *Clinical Microbiology Reviews*, 8, 585-615.
- Wyllie, A. H., Kerr, J. F., & Currie, A. R. (1980). Cell death: the significance of apoptosis. *International Review of Cytology*, 68, 251-306.
- Xanthopoulos, V., Ipsilandis, C. G., & Tzanetakis, N. (2012). Use of a selected multi-strain potential probiotic culture for the manufacture of set-type yogurt from caprine milk. *Small Ruminant Research*, 106(2-3), 145-153.
- Xue, R., Han, N., Xia, M., Ye, C., Hao, Z., Wang, L., Wang, Y., Yang, J., Saiki, I., & Yin, J. (2015). TXA9, a cardiac glycoside from *Streptocaulon juvenas*, exerts a potent anti-tumor activity against human non-small cell lung cancer cells *invitro* and *invivo*. *Steroids*, 94(0), 51-59.
- Yamada, Y., & Yukphan, P. (2008). Genera and species in acetic acid bacteria. *International Journal of Food Microbiology*, 125(1), 15-24.
- Yang, E. J., & Chang, H. C. (2010). Purification of a new antifungal compound produced by *Lactobacillus plantarum* AF1 isolated from kimchi. *International Journal of Food Microbiology*, 139(1-2), 56-63.
- Yazdi, M. H., Mahdavi, M., Kheradmand, E., & Shahverdi, A. R. (2012). The preventive oral supplementation of a selenium nanoparticle-enriched probiotic

- increases the immune response and lifespan of 4T1 breast cancer bearing mice. *Arzneimittel-Forschung/Drug Research*, 62(11), 525-531.
- Young, R. J., & Huffman, S. (2003). Probiotic use in children. *Journal of Pediatric Health Care*, 17(6), 277-283.
- Young, T. B., & Wolf, D. A. (1988). Case-control study of proximal and distal colon cancer and diet in Wisconsin. *International Journal of Cancer*, 42(2), 167-175.
- Zacconi, C., V. B., Rebecchi, A., Bosi, E., Sarra, P. G., & Tagliaferri, L. (1992). Serum cholesterol levels in axenic mice colonized with *Enterococcus faecium* and *Lactobacillus acidophilus*. *Microbiologica*, 15(4), 413-417.
- Zamfir, M., Callewaert, R., Cornea, P. C., & De Vuyst, L. (2000). Production kinetics of acidophilin 801, a bacteriocin produced by *Lactobacillus acidophilus* IBB 801. *FEMS Microbiology Letters*, 190(2), 305-308.
- Zhou, J. S., Pillidge, C. J., Gopal, P. K., & Gill, H. S. (2005). Antibiotic susceptibility profiles of new probiotic *Lactobacillus* and *Bifidobacterium* strains. *International Journal of Food Microbiology*, 98(2), 211-217.
- Zhu, L. J., Zheng, F. Y., Zhao, Y. Z., Xing, X. N., Li, Q., & Gu, G. X. (2006). Application of Rep-PCR fingerprint in rapid identification of beer-spoilage bacteria. *Chinese Journal of Biotechnology*, 22(6), 1013-1021.
- Zhu, Y., Michelle Luo, T., Jobin, C., & Young, H. A. (2011). Gut microbiota and probiotics in colon tumorigenesis. *Cancer Letters*, 309(2), 119-127.
- Zouain, M. G., & Araj, G. F. (2001). Antimicrobial resistance of *Enterococci* in Lebanon. *International Journal of Antimicrobial Agents*, 17(3), 209-213.

LIST OF PUBLICATIONS

1. Babak Haghshenas, Yousef Nami, Norhafizah Abdullah, Dayang Radiah, Rozita Rosli, Ahmad Yari Khosroushahi (2014). Anti-proliferative effects of *Enterococcus* strains isolated from fermented dairy products on different cancer cell lines. *Journal of Functional Foods*, 11, 363–374.
2. Babak Haghshenas, Yousef Nami, Norhafizah Abdullah, Dayang Radiah, Rozita Rosli, Ahmad Yari Khosroushahi (2015). Anticancer impacts of potentially probiotic acetic acid bacteria isolated from traditional dairy microbiota. *LWT - Food Science and Technology*, 60, 690-697.
3. Babak Haghshenas, Yousef Nami, Norhafizah Abdullah, Dayang Radiah, Rozita Rosli, Ahmad Yari Khosroushahi (2014). Potentially probiotic acetic acid bacteria isolation and identification from traditional dairies microbiota. *International Journal of Food Science and Technology*, 50(4), 1056-1064.
4. Babak Haghshenas, Norhafizah Abdullah, Yousef Nami, Dayang Radiah, Rozita Rosli, Ahmad Yari Khosroushahi (2014). Different effects of two newly-isolated probiotic *Lactobacillus plantarum* 15HN and *Lactococcus lactis* subsp. *Lactis* 44Lac strains from traditional dairy products on cancer cell lines. *Anaerobe*, 30, 51-59.
5. Babak Haghshenas, Norhafizah Abdullah, Yousef Nami, Dayang Radiah, Rozita Rosli, Ahmad Yari Khosroushahi (2015). Microencapsulation of probiotic bacteria *Lactobacillus plantarum* 15HN using alginate-psyllium-fenugreek polymeric blends. *Journal of Applied Microbiology*, 118(4), 1048-1057.
6. Yousef Nami, Babak Haghshenas, Norhafizah Abdullah, Dayang Radiah, Rozita Rosli, Ahmad Yari Khosroushahi (2014). Probiotics or antibiotics: future challenges in medicine. *Journal of Medical Microbiology*, 64(2), 137-146.