



**UNIVERSITI PUTRA MALAYSIA**

***MOLECULAR IDENTIFICATION, PROBIOTIC CHARACTERIZATION,  
ANTICANCER ASSESSMENT AND ENCAPSULATION OF LACTIC ACID  
BACTERIA ISOLATED FROM VAGINAS OF IRANIAN FERTILE WOMEN***

**Yousef Nami**

**IB 2015 43**



**MOLECULAR IDENTIFICATION, PROBIOTIC  
CHARACTERIZATION, ANTICANCER ASSESSMENT AND  
ENCAPSULATION OF LACTIC ACID BACTERIA ISOLATED  
FROM VAGINAS OF IRANIAN FERTILE WOMEN**

By

**YOUSEF NAMI**

**Thesis Submitted to the School of Graduate Studies, Universiti  
Putra Malaysia, in Fulfilment of the Requirements for the Degree of  
Doctor of Philosophy**

**February 2015**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



**This thesis dedicates to:**

*My family for their love, endless support  
and encouragement...*



Abstract of thesis presented to the Senate of Universiti Putra Malaysia, in fulfilment of the requirements for the degree of Doctor of Philosophy

**MOLECULAR IDENTIFICATION, PROBIOTIC CHARACTERIZATION,  
ANTICANCER ASSESSMENT AND ENCAPSULATION OF LACTIC ACID  
BACTERIA ISOLATED FROM VAGINAS OF IRANIAN FERTILE WOMAEN**

By

**YOUSEF NAMI**

**February 2015**

**Chairman: Associate Professor Norhafizah Abdullah, Ph.D**  
**Faculty: Institute of Bioscience**

Probiotics are non-pathogenic microorganisms that positively influence their hosts when ingested in adequate amounts. Research on probiotics has increased over the past years. This trend has resulted in the commercialization of probiotic supplements and functional food. Numerous investigations have evaluated the potential of probiotic bacteria for cancer prevention or treatment. Most previous works isolated probiotic bacteria from dairy products or other fermented foods. This study isolated and characterized lactic acid bacteria (LAB) from a healthy vaginal ecosystem. LAB that exhibited probiotic properties were screened for their antimicrobial activity, antibiotic resistance, and anticancer activity. Isolation was performed using preliminary phenotypic screening based on Gram-staining and catalase reaction, followed by 16S rRNA gene sequencing, amplified ribosomal DNA restriction analysis (ARDRA), and repetitive sequence-based PCR fingerprinting. The bacteria isolated from 40 samples were classified into three genera with 13 species, three subspecies, and 45 strains. The combined use of molecular techniques was proven effective for the identification and classification of vaginal LAB, particularly *Lactobacillus*, *Lactococcus*, and *Enterococcus*.

The 45 vaginal LAB strains were characterized for their probiotic and therapeutic potentials. Probiotic characterization included in vitro assays of acid and bile salt tolerance, antimicrobial activity, and antibiotic resistance. Results showed that the survival rate of the isolates ranged from 18 to 88% and from 31 to 93% under acidic (pH 3.0) and 0.3% bile salt conditions in a simulated gastrointestinal environment, respectively. The strains from each of the 13 species that demonstrated the highest acid and bile salt tolerance were further evaluated for their antibiotic susceptibility and antimicrobial activity. Results showed that the survival rates of these 13 isolates under acidic and bile salt conditions were higher than 58 and 78%, respectively, qualifying them to be probiotic candidates. All 13 isolates exhibited antimicrobial activity against different pathogenic bacteria. Among the LAB, *Lactococcus lactis* 2BL showed the strongest antimicrobial activity against 15 pathogenic bacteria. The 13 isolates showed

different levels of antibiotic resistance depending on antibiotic used. Among the LAB, *Enterococcus avium* 7BL and *Enterococcus durans* 6HL were resistant and sensitive to all nine antibiotics used, respectively.

The anticancer effects of the 13 isolates on some human cancer cell lines, such as HeLa, AGS, HT-29, and MCF-7, were also assessed. The human normal cell line HUVEC was used as a control. The cytotoxicity of these strains and the occurrence of apoptotic cells were evaluated using DNA fragmentation, fluorescent microscopy, and flow cytometry. The metabolites produced by *Lactobacillus plantarum* 5BL, *Lactobacillus acidophilus* 36YL, *Enterococcus faecalis* 16H, and *Enterococcus lactis* 2BL exhibited remarkable anticancer activity against the tested human carcinoma cell lines with no significant cytotoxicity on the HUVEC normal cells. Apoptotic cells were also observed in the cancer cell lines.

Finally, the isolates were subjected to encapsulation to improve their bioavailability and survival rate under harsh gastrointestinal conditions. Herbal-based biopolymer matrices, such as alginate, gum arabic, and psyllium, were prepared through extrusion. Three gel formulations, namely, 1, 1.5 and 2% (w/v) alginate, 2% (w/v) alginate + 0.1 and 0.3% (w/v) gum arabic + psyllium, 1.5% (w/v) alginate + 0.3 and 0.5% (w/v) gum arabic + psyllium, and 1% (w/v) alginate + 0.4 and 0.6% (w/v) gum arabic/psyllium were prepared to improve the encapsulation efficiency, gastrointestinal survival, and colonic release rates of the isolates. All encapsulated bacteria exhibited significantly improved ( $P < 0.05$ ) encapsulation efficiency (>98%) and survival rate (>75%) compared with the un-encapsulated bacteria (>45%) under harsh gastrointestinal conditions. The incorporation of gum arabic and psyllium into the alginate gel enhanced encapsulation properties. The use of combination gels for encapsulation improved the survival rate and storage stability of the probiotic strains under gastrointestinal conditions.

In conclusion, probiotic bacteria were successfully isolated and characterized from a healthy vaginal ecosystem. These probiotics exhibited antimicrobial activity, antibiotic resistance, and anticancer activity, making these bacteria potential candidates for the development of nutraceutical products.

Abstrak tesis yang dikemukakan kepada senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**Pengenalan Molekul, Probiotik Pencirian, Antikanser  
Penilaian dan Pengkapsulan Asid Laktik Bakteria Ribut  
dari Vagina Iran Wanita Subur**

Oleh

**YOUSEF NAMI**

**Februari 2015**

**Pengerusi: Profesor Madya Norhafizah Abdullah, Ph.D**  
**Fakulti: Institut Biosains**

Probiotik adalah mikroorganisma bukan patogen yang boleh memberi kesan positif terhadap hos mereka apabila dimakan dalam jumlah yang mencukupi. Penyelidikan probiotik meningkat sejak beberapa tahun kebelakangan ini. Kecenderungan ini telah menghasilkan pengkomersialan makanan tambahan probiotik dan makanan berfungsi. Selain dari itu, semakin banyak kajian dilakukan bagi mengkaji potensi bakteria probiotik untuk pencegahan atau rawatan kanser. Kebanyakan kajian sebelum ini telah menggunakan probiotik yang dipencil daripada produk tenusu atau makanan diperam yang lain. Pada kajian ini, kelainan sumber bakteria asid laktik (LAB) yang dipencil daripada ekosistem faraj yang sihat dan subur telah diperkenalkan. LAB yang mempamerkan sifat probiotik seterusnya disaring untuk aktiviti anti-mikrob, rintangan antibiotik, dan aktiviti anti kanser. Pengasingan dilakukan dengan menggunakan saringan awal fenotip berdasarkan teknik pewarnaan-Gram dan reaksi katalase, diikuti pula dengan teknik penjujukan gen 16S rRNA, teknik penguatan sekatan ribosoma DNA (ARDRA), dan analisa pengecaman '*finger-printing*' turutan berasaskan PCR. Bakteria yang diasingkan daripada 40 sampel telah diklasifikasikan kepada tiga genus dengan 13 spesies, tiga subspesies, dan 45 jenis. Penggunaan gabungan teknik molekul telah terbukti berkesan bagi mengenalpasti dan mengklasifikasi LAB yang dipencil dari ekosistem faraj, terutamanya *Lactobacillus*, *Lactococcus*, dan *Enterococcus*.

45 strain LAB faraj telah dicirikan mempunyai potensi probiotik dan terapeutik. Pencirian probiotik termasuk ujian toleransi '*in vitro*' terhadap asid dan garam hempedu, aktiviti antimikrob, dan rintangan antibiotik. Hasil kajian mendapati kadar kelangsungan hidup daripada strain yang dipencil adalah dari 18 kepada 88% dan dari 31% kepada 93% di dalam persekitaran berasid (pH 3.0) dan 0.3% garam hempedu yang mana masing-masing adalah mewakili persekitaran sistem pencernaan. Strain dari setiap satu daripada 13 spesies yang menunjukkan toleransi terhadap asid dan garam hempedu selanjutnya dinilai untuk kerentanan terhadap antibiotik dan aktiviti anti-mikrob. Keputusan menunjukkan bahawa kadar kelangsungan hidup dari 13 penciran LAB di bawah keadaan berasid dan garam hempedu adalah lebih tinggi daripada 58%

dan 78% masing-masing, melayakkan mereka untuk menjadi calon probiotik. Kesemua 13 pencilan menunjukkan aktiviti antimikrob terhadap bakteria patogenik yang berbeza. Antara LAB itu, *Lactococcus lactis* 2BL menunjukkan aktiviti antimikrob yang kuat terhadap 15 bakteria patogenik. 13 pencilan menunjukkan tahap yang berbeza terhadap rintangan antibiotik, bergantung kepada jenis antibiotik digunakan. Antara LAB itu, *Enterococcus avium* 7BL dan *Enterococcus durans* 6HL adalah tahan dan sensitif kepada kesemua sembilan antibiotik digunakan, masing-masing.

Kesan anti-kanser daripada 13 pencilan pada beberapa jenis sel kanser manusia, seperti HeLa, AGS, HT-29, dan MCF-7, juga dinilai. Sel normal manusia HUVEC digunakan sebagai kawalan. Sitotoksiti strain ini menunjukkan berlakunya 'apoptosis' sel, seperti yang dinilai oleh teknik pemecahan DNA, mikroskop pendarfluor, dan pengalir sitometri. Metabolit yang dihasilkan oleh *Lactobacillus Plantarum* 5BL, *Lactobacillus acidophilus* 36YL, *Enterococcus faecalis* 16H, dan *Enterococcus lactis* 2BL menunjukkan aktiviti anti-kanser yang luar biasa terhadap sel-sel karsinoma manusia yang diuji, tetapi tiada kesan sitotoksiti yang ditunjukkan ke atas sel-sel normal HUVEC.

Akhirnya, pencilan tertakluk kepada pengkapsulan bagi meningkatkan "bioavailability" mereka dan kadar kelangsungan hidup di bawah persekitaran sistem pencernaan. Matriks biopolimer berasaskan herba, seperti alginat, gam arabik dan helba telah disediakan melalui teknik penyemperitan. Tiga formulasi gel, iaitu, 1, 1.5 dan 2% (w/v) alginat, 2% (w/v) alginat + 0.1 dan 0.3% (w/v) gam arabik / helba, 1.5% (w/v) alginat + 0.3 dan 0.5% (w/v) gam arabik / helba, dan 1% (w/v) alginat + 0.4 dan 0.6% (w/v) gam arabik / helba telah disediakan untuk meningkatkan kecekapan pengkapsulan ini, kelangsungan hidup dalam sistem pencernaan, dan kadar pembebasan kolon daripada pencilan. Semua bakteria yang terkandung mempamerkan penambahbaikan yang ketara ( $P < 0.05$ ) dengan kecekapan pengkapsulan yang tinggi (>98%) dan kadar kelangsungan hidup (>75%) berbanding dengan bacteria yang tidak dikandung dalam gel matrik (>45%). Pemerbadanan polimer gam arab dan helba ke dalam pengkapsulan gel alginat turut juga dipertingkatkan. Penggunaan kombinasi gel-gel ini menghasilkan pengkapsulan baik dengan kadar kelangsungan hidup dan kestabilan penyimpanan strain probiotik dalam keadaan sistem pencernaan yang memuaskan.

Kesimpulannya, bakteria probiotik telah berjaya dipencilkan dan dicirikan dari ekosistem vagina yang sihat. Probiotik ini mempunyai sifat antimikrob, rintangan antibiotik dan anti-kanser, menjadikan mereka calon yang berpotensi untuk dibangunkan sebagai produk nutrasetikal.



## ACKNOWLEDGEMENTS

At the end of this step of my graduate period has allowed for a fragment of reflection, and the many dears who have contributed to both, my work, and my life during this period of time, because this research project would not have been possible without the support of many people.

First and foremost, I would like to express my utmost gratitude to my supervisor, AP. Dr. Norhafizah Abdullah and my consultants Prof. Dr. Rozita Rosli, Dr. Dayang Radiah Awang Biak and Dr. Ahmad Yari Khosroushahi; for their invaluable suggestions, beneficial advices, their extended long-term supports and vast reserve of patience and knowledge. I owe my deepest gratitude to my best friends, Babak Haghshenas and Abolfazl Barzegari, who have always helped me and believed that I could do it.

Finally, I would like to offer my sincere thanks to my dear wife for her encouragements, her emotional supports and her fortitude. Last but not least I wish to avail myself of this opportunity, express a sense of gratitude and love to my friends and my beloved parents for their support, strength, and help for everything during my life. I wish to acknowledge my gratitude to all lecturers, officials and other staff members in Universiti Putra Malaysia and Pharmacy College of Tabriz Universiti.

**Yousef Nami**  
**February 2015**

I certify that a Thesis Examination Committee has met on 26 February 2015 to conduct the final examination of Yousef Nami on his PhD thesis entitled "Molecular Identification, Probiotic Properties, Anticancer Assessment and Encapsulation of Lactic Acid Bacteria Isolated from Vaginas of Iranian Healthy Women" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U. (A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

**Luqman Chuah Abdullah, PhD**

Professor  
Faculty of Engineering  
Universiti Putra Malaysia  
(Chairman)

**Rosfarizan binti Mohamad, PhD**

Associate Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**Tan Wen Siang, PhD**

Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**William Chen Wei Ning, PhD**

Professor  
Nanyang Technological University  
Singapore  
(External Examiner)



---

**ZULKARNAIN ZAINAL, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 15 April 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Norhafizah Abdullah, PhD**

Associate Professor  
Faculty of Engineering  
Universiti Putra Malaysia  
(Chairman)

**Rozita Rosli, PhD**

Professor  
Institute of Bioscience  
Universiti Putra Malaysia  
(Member)

**Dayang Radiah Awang Biak, PhD**

Senior Lecturer  
Faculty of Engineering  
Universiti Putra Malaysia  
(Member)

**Ahmad Yari Khosroushahi, PhD**

Assistant Professor  
Faculty of Pharmaceutical  
Tabriz University of Medical Science of Iran  
(Member)

---

**BUJANG BIN KIM HUAT, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## CHAPTER 1

### INTRODUCTION

Lactic acid bacteria (LAB) are heterogeneous, Gram-positive, and non-spore-forming bacteria that display fermentative metabolism. These organisms naturally thrive on humans, animals, and plants. LAB are commonly utilized in the dairy and food industries because of their health-promoting and well-documented fermentative properties (Temmerman *et al.*, 2004). LAB are normal flora in the mouth, intestine, and other parts of mammals. LAB also dominate the vaginal microbiota in normal post-pubertal and pre-menopausal women (Nam *et al.*, 2007; Ehrstrom *et al.*, 2010). They share a mutual relationship with their hosts by promoting health and preventing diseases (Turnbaugh *et al.*, 2007; Ma *et al.*, 2012a). Most LAB can inhibit the growth of other bacteria by producing different metabolites, such as lactic acid, acetic acid, H<sub>2</sub>O<sub>2</sub>, bacteriocin, and nitric oxide (Ehrstrom *et al.*, 2010). These metabolites help prevent bacterial vaginosis, yeast infections, urinary tract infections, and sexually transmitted diseases (Ma *et al.*, 2012a).

However, a major health concern regarding LAB supplementation is the transmission of antibiotic resistance genes from LAB to unrelated pathogenic or potentially pathogenic bacteria in the gut. LAB strains that harbor antibiotic resistance plasmids are unsuitable for use as human or animal probiotics (Hart and Powell, 1990; Palmer *et al.*, 2010). Some strains of *Lactobacillus fermentum*, *Lactobacillus plantarum*, and *Lactobacillus reuteri* carry potentially transmissible plasmid-encoded antibiotic resistance genes (Begovic *et al.*, 2009).

Among the LAB, lactobacilli are important members of vaginal bacterial communities in women of reproductive age (Ma *et al.*, 2012a). Lactobacilli maintain the vaginal ecological equilibrium by protecting against pathogenic microorganisms. They also assist in changing the physiology of the urogenital tract by modifying the normal microbial flora (Vintini *et al.*, 2004; Pascual *et al.*, 2010). They are considered non-pathogenic because of their long-standing reputation as health-promoting organisms in many fermented food products. Lactobacilli are probiotics that have been used to treat different bacterial vaginal infections (Mastromarino *et al.*, 2009). The ecological functions and effects of other LAB on the overall dynamics and functions of vaginal bacterial communities remain unclear. Recent evolutions in biotechnological and biochemical techniques have been directed to the use of LAB in the production of bulk enzymes and biopolymers or as oral delivery vehicles (Steidler, 2002; Nam *et al.*, 2007).

Probiotics are live non-pathogenic microorganisms that confer health benefits to the host when consumed in adequate amounts (Barrons and Tassone, 2008b). Probiotics, which exist in the human body as microflora, promote the intestinal microbial balance

(Zhang *et al.*, 2012). Probiotics can adhere to epithelial cells and tolerate high pH and bile levels in the gastrointestinal tract. Acid tolerance is one of the desirable properties used during the screening of potential probiotic strains. The competition of commensal and probiotic bacteria with pathogens to adhere and colonize is an important mechanism in resisting intestinal pathogens and preventing diseases. Probiotic bacteria produce various compounds or metabolites that inhibit pathogen growth. These metabolites include organic acids (lactic and acetic acids), bacteriocin, and reuterin (Alvarez-Olmos and Oberhelman, 2001). Organic acids affect pathogen growth and elicit toxic effects on microbes by reducing pH. Not all LAB have probiotic properties that control or prevent infectious diseases (Colodner *et al.*, 2003; Reid and Bruce, 2003). Disruption and damage of the normal genital microflora, particularly *Lactobacillus* species, lead to the increase of genital infections (Reid, 2002; Pascual *et al.*, 2008a; Pascual *et al.*, 2008b). The *in vitro* attachment of pathogenic bacteria can be blocked by the normal vaginal flora of healthy females (Pascual *et al.*, 2010). Lactobacilli produce bacteriocins, such as acidophilin, acidolin, lactocidin, bulgarican, lactolin, lactobacillin, and lactobrevin (Dimitonova *et al.*, 2007; Pan *et al.*, 2009). Many *Lactobacillus*, *Bifidobacterium*, and *Streptococcus* species are safe to use for the prevention or treatment of various infectious diseases (Lee *et al.*, 2003).

Probiotic strains should not carry transmissible antibiotic resistance genes. Probiotics may become hosts of antibiotic resistance genes that can be transferred to pathogenic bacteria. Ingestion of bacteria carrying such genes is undesirable because horizontal gene transfer to recipient bacteria in the gut can lead to the development of new antibiotic-resistant pathogens (Zhou *et al.*, 2005).

Cancer is one of the main causes of human deaths. Chemotherapy and chemoprevention are applied to control metastasis and decrease human mortality (Miranda *et al.*, 1999). Probiotics also exhibit anticancer activities against different carcinoma cells, such as colon, bladder, and breast (de Blanc and Bonet, 2010). Most studies focused on the anticancer effect of probiotics on colon carcinoma cells (Rafter, 2004). Studies have discovered that some specific strains of LAB can induce the production of pro-inflammatory and anti-inflammatory cytokines in animal or human body. Evaluation of the toxicity of bacterial cytotoxic agents to different cell types is performed by *in vitro* cytotoxicity tests and by analyzing the mechanism of apoptosis in treated cancer cells (Hsu *et al.*, 2005). Induction of apoptosis, which can be applied to control cancer development, involves complex anticancer activity of several therapeutic substances (Li *et al.*, 2005). Although various methods are available for identification, quantification, and characterization of apoptosis, flow cytometry is the most common choice to study apoptosis because of its applicability to an extensive range of cell types, stimulants, and time. Evaluating apoptosis by flow cytometry provides information on symptoms and incidence of apoptosis that is interpreted as a sign. Besides, cellular morphological characteristic is another method to determine the cell death mode (apoptosis or necrosis).

In the present study, effective probiotic strains were isolated from the vaginas of Iranian women. The anticancer effects of the isolates on different human cancer cell lines were evaluated through *in vitro* cytotoxic and apoptotic analyses.

Encapsulation is the process of entrapping bioactive molecules (e.g., vitamins, minerals, antioxidants, fatty acid, lutein, and lycopene) and living cells (e.g., probiotics) within carrier materials (Nedovic *et al.* 2011). Encapsulation is the most effective technique to protect probiotic bacteria during processing and storage (Kanmani *et al.* 2011). Improving the delivery of these active agents into foods and medicines is important and many substances can be used for encapsulation. Encapsulation materials are selected based on the following criteria: functionality, stability, type of release, encapsulates concentration, and cost. In addition, carrier materials must be biodegradable, biocompatible, food grade, and capable of barrier formation (Nedovic *et al.* 2011).

Encapsulation causes the production of a physical barrier between the internal phase and its surrounding to protect them against pH alterations, moisture variations, and oxidation; thus, this process controls the release of active molecules and increases their bioavailability (Dubey *et al.* 2009). The most significant aim for the encapsulation of active agents is to make available improved stability in final products and during processing.

The most common encapsulation material is sodium alginate because of its simplicity, biocompatibility, non-toxicity and low cost (Krasaekoopt *et al.*, 2003). In addition, alginate microspheres with porous structure allow diffusion of acid in and out of microspheres easily. These disadvantages can be effectively overcome by blending alginate with other polymers or coating one polymer layer on alginate microspheres (Krasaekoopt *et al.*, 2003; Burgain *et al.*, 2011). New biopolymers for encapsulation proposes have been recently reported (Dickinson 2003; Mahfoudhi *et al.* 2014). Gum exudates are predominantly composed of polysaccharides that function as stabilizing and emulsifying agents (Orozco-Villafuerte *et al.* 2003). Gum Arabic (GA) has the highest commercial value among the gum exudates because of its extensive application in the pharmaceutical, food, and cosmetic industries (Mahfoudhi *et al.* 2014).

Psyllium (PSY), an arabinoxylan herbal-based biopolymer, extracts from *Plantago* species. PSY can stimulate the growth of probiotic bacteria in the gastrointestinal tract (GIT) and treat several gut disorders, including ulcerative colitis, chronic kidney, constipation, and diarrhea (Guo *et al.* 2008; Rishniw and Wynn, 2011).

Studies on the isolation of probiotic bacteria from the vaginas of healthy Iranian women and on the evaluation of their efficacy on pathogenic bacteria and different cancers are lacking. In addition, the probiotic activities of LAB isolated from the vagina have yet to be assessed. Accordingly, the principal objectives of this study were:

1. To isolate and identify LAB from the vaginas of healthy Iranian women using different molecular fingerprinting methods and evaluation of the isolated LAB for probiotic characteristics: low pH resistance, high bile salt tolerance, antimicrobial activity and antibiotic susceptibility.
2. To evaluate the *in vitro* cytotoxic assay and morphological apoptosis assessment of probiotic secretions on different cancerous and normal cell lines using MTT, fluorescent microscopy, DAPI staining and flow cytometry assays.
3. To enhance the survival and bioavailability of probiotic bacteria under gastrointestinal conditions by using microencapsulation with alginate, psyllium and Gum Arabic by extrusion method.

## REFERENCES

- Aga, Z.B. and Ramanan, S.R. (2014). Effect of doping on the structural parameters of pure and doped barium titanate nano powders synthesised by sol-gel emulsion technique. *International Journal of Nanoparticles* 7(3): 331-349.
- Ahmadova, A. and Todorov, S.D. (2013). Evaluation of antimicrobial activity, probiotic properties and safety of wild strain *Enterococcus faecium* AQ71 isolated from Azerbaijani Motal cheese. *Food Control* 30(2): 631-641.
- Albertini, B., Passerini, N., Pattarino, F. and Rodriguez, L. (2008). New spray-congealing atomizer for the microencapsulation of highly concentrated solid and liquid substances. *European Journal of Pharmaceutics and Biopharmaceutics* 69: 348-357.
- Albertini, B., Vitali, B., Passerini, N., Cruciani, F., Di Sabatino, M., Rodriguez, L. and Brigidi, P. (2010). Development of microparticulate systems for intestinal delivery of *Lactobacillus acidophilus* and *Bifidobacterium lactis*. *European Journal of Pharmaceutical Sciences* 40: 359-366.
- Ali, B.H., Ziada, A. and Blunden, G. (2009). Biological effects of gum Arabic: A review of some recent research. *Food and Chemical Toxicology* 47: 18-23.
- Almuslet, N.A., Hassan, E.A., Al-Sherbini, A.S.A. and Muhgoub, M.G.A. (2012). Diode Laser (532 nm) Induced Grafting of Polyacrylamide onto Gum Arabic. *Journal of Physical Science* 23(2): 43-53.
- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D.J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search Programs. *Nucleic Acids Research* 25: 3389-3402.
- Alvarez-Olmos, M.I. and Oberhelman, R.A. (2001). Probiotic agents and infectious diseases: a modern perspective on a traditional therapy. *Clinical Infectious Diseases* 32(11): 1567-1576.
- American Cancer Society. Cancer Facts and Figures 2013. Atlanta, Ga: American Cancer Society; 2013.
- American Cancer Society. Global Cancer Facts & Figures 2007.
- Amine, K.M., Champagne, C.P., Raymond, Y., St-Gelais, D., Britten, M., Fustier, P., Salmieri, S. and Lacroix, M. (2014). Survival of microencapsulated *Bifidobacterium longum* in Cheddar cheese during production and storage. *Food Control* 37: 193-199.
- Anal, K.A. and Singh, H. (2007). Recent advances in microencapsulation of probiotics for industrial applications and targeted delivery. *Trends in Food Science & Technology* 18, 240-251.



- Anderson, H.A., Englert, R., Gursel, I. and Shacter, E. (2002). Oxidative stress inhibits the phagocytosis of apoptotic cells that have externalized phosphatidylserine. *Cell Death and Differentiation* 9: 616-625.
- Andersson, H., Asp, N.G., Bruce, A., Roos, A., Wadstrom, T. and Wold, A.E. (2001). Health effects of probiotics and prebiotics. A literature review on human studies. *Scandinavian Journal of Nutrition* 45: 58-75.
- Andreu, A. (2004). Lactobacillus as a probiotic for preventing urogenital infections. *Reviews in Medical Microbiology* 15: 1-6.
- Axelsson, L. (1998). Lactic acid bacteria: Classification and Physiology. In Lactic acid bacteria: Microbiology and Functional Aspects. Marcel Dekker, Inc., New York, pp. 1-72.
- Axelsson, L.T., Salminen, S. and Wright, A.V. (1993). Lactic acid bacteria: Classification and Physiology: Lactic Acid Bacteria. *Marcel Dekker, INC., New York*: 1-52.
- Babbar, N., Dejonghe, W., Gatti, M., Sforza, S., and Kathy, E. (2015). Pectic oligosaccharides from agricultural by-products: production, characterization and health benefits. *Critical reviews in biotechnology*, (0): 1-13.
- Babich, H. and Borenfreund, E. (1991). Cytotoxicity of T-2 toxin and its metabolites determined with the neutral red cell viability assay. *Applied and Environmental Microbiology* 57(7): 2101-2103.
- Baker, M.A., Cerniglia, G.J. and Zaman, A. (1990). Microtitre plate assay for the measurement of glutathione and glutathione disulfide in large numbers of biological samples. *Analytical Biochemistry* 190: 360-365.
- Baldwin, C., Millette, M., Oth, D., Ruiz, M.T., Luquet, F.M. and Lacroix, M. (2010). Probiotic *Lactobacillus acidophilus* and *L. casei* mix sensitize colorectal tumoral cells to 5-fluorouracil-induced apoptosis. *Nutrition Cancer* 62: 371-378.
- Barrons, R. and Tassone, D. (2008). Use of Lactobacillus probiotics for bacterial genitourinary infections in women: a review. *Clin Ther* 30: 453-468.
- Barrons, R. and Tassone, D. (2008). Use of *Lactobacillus* Probiotics for Bacterial Genitourinary Infections in Women. *Clinical Therapeutics* 30(3): 453-468.
- Bassaganya-Riera, J. (2012). Immunoregulatory mechanisms underlying prevention of colitis-associated colorectal cancer by probiotic bacteria. *PLoS ONE* 7(4): p.e34676.
- Bassaganya-Riera, J., Viladomiu, M., Pedragosa, M., De Simone, C. and Hontecillas, R. (2012). Immunoregulatory Mechanisms Underlying Prevention of Colitis-Associated Colorectal Cancer by Probiotic Bacteria. *PLoS ONE* 7(4).
- Beardsley, T. (1994). Trends in cancer epidemiology: a war not won. *Science American* 270: 118-126.

- Begley, M., Hill, C. and Gahan, C.G.M. (2006). Bile Salt Hydrolase Activity in Probiotics. *Applied Environmental Microbiology* 72: 1729-1738.
- Begovic, J., Huys, G., Mayo, B., D'Haene, K., Florez, A.B., Lozo, J., Kojic, M., Strahinic, I. and Topisirovic, L. (2009). Human vaginal *Lactobacillus rhamnosus* harbor mutation in 23S rRNA associated with erythromycin resistance. *Research in Microbiology* 160(6): 421-426.
- Bell, L.N. (2001). Stability testing of nutraceuticals and functional foods. In: Wildman R.E.C. (Eds.). Handbook of nutraceuticals and functional foods, CRC Press, New York. 501-516.
- Bernardeau, M., Vernoux, J.P., Henri-Dubernet, S. and Gueguen, M. (2008). Safety assessment of dairy microorganisms: The *lactobacillus* genus. *International Journal of Food Microbiology* 126: 278-285.
- Bigwood, T. and Hudson, J.A. (2012). Inhibition of *Listeria monocytogenes* by *Enterococcus mundtii* isolated from soil. *Food Microbiology* 32(2): 354-360.
- Bingham, S. and Riboli, E. (2004). Diet and cancer- The European prospective investigation into cancer and nutrition. *Nature Reviews Cancer* 4: 206-215.
- Blum, S., Reniero, R., Schiffrin, E.J., Crittenden, R., Mattila-Sandholm, T., von Wright, A., Saarela, M., Saxelin, M., Collins, K. and Morelli, L. (1999). Adhesion studies for probiotics: need for validation and refinement. *Trends in Food Science and Technology* 10: 405-410.
- Boever, D.P., Wouters, R., Verschaeve, L., Berckmans, P., Schoeters, G. and Verstraete, W. (2000). Protective effect of the bile salt hydrolase-active *Lactobacillus reuteri* against bile salt cytotoxicity. *Applied Microbiology and Biotechnology* 53: 709-714.
- Bosnea, L.A., Moschakis, T., and Biliaderis, C.G. (2014). Complex Coacervation as a Novel Microencapsulation Technique to Improve Viability of Probiotics Under Different Stresses. *Food and Bioprocess Technology*, 7(10): 2767-2781.
- Bouchier-Hayes, L., Munoz-Pinedo, C., Connel, S. and Green, D.R. (2008). Measuring apoptosis at the single cell level. *Methods* 44: 222-228.
- Bouton, Y., Guyot, P., Beuvier, E., Tailliez, P. and Grappin, R. (2002). Use of PCR-based methods and PFGE for typing and monitoring homofermentative lactobacilli during Comte cheese ripening. *International Journal of Food Microbiology* 76: 27-38.
- Brenner, D.M., Moeller, M.J., Chey, W.D. and Schoenfeld, P.S. (2009). The utility of probiotics in the treatment of irritable bowel syndrome: a systematic review. *American Journal of Gastroenterology* 104(4): 1033-1049.
- Bulut, C. 2003. Isolation and molecular characterization of lactic acid bacteria from cheese. IYTE thesis of Master.

- Burgain, J., Gaiani, C., Linder, M. and Scher, J. (2011). Encapsulation of probiotic living cells: from laboratory scale to industrial application. *Journal of Food Engineering* 104: 467-483.
- Burton, J.P., Cadieux, P.A. and Reid, G. (2003). Improved understanding of the bacterial vaginal microbiota of women before and after probiotic instillation. *Applied Environmental Microbiology* 69(1): 97-101.
- Cakir, I. 2003. Determination of some probiotic properties on Lactobacilli and Bifidobacteria. Ankara University Thesis of Ph.D.
- Cancer Research UK. Stomach cancer symptoms. <http://www.cancerhelp.org.uk/help/default.asp?page=3897> Last accessed 1 August 2011.
- Capela, P., Hay, T.K.C. and Shah, N.P. (2006). Effect of cryoprotectants, prebiotics and microencapsulation on survival of probiotic organisms in yoghurt and freeze-dried yoghurt. *Food Research International* 39: 203-211.
- Cardinal, M.J., Meghrou, J., Lacroix, C. and Simard, R.E. (1997). Isolation of Lactococcus lactis strain producing inhibitory activity against *Listeria*. *Food Biotechnology* 11(129).
- Carmen, C.M. and Hernandez, M. (2007). Identification and differentiation of *Lactobacillus*, *Streptococcus* and *Bifidobacterium* species in fermented milk products with *bifidobacteria*. *Microbiological Research* 162(1): 86-92.
- Carvalho, A.S., Silva, J., Ho, P., Malcata, F.X. and Gibbs, P. (2003). Effect of various growth media upon survival during storage of freeze-dried *Enterococcus faecalis* and *Enterococcus durans*. *Journal of Applied Microbiology* 94: 947-952.
- Cebrian, R., Banos, A., Valdivia, E., Perez-Pulido, R., Martinez-Bueno, M. and Maqueda, M. (2012). Characterization of functional, safety, and probiotic properties of *Enterococcus faecalis* UGRA10, a new AS-48-producer strain. *Food Microbiology* 30(1): 59-67.
- Cebrian, R., Banos, A., Valdivia, E., Perez-Pulido, R., Martinez-Bueno, M. and Maqueda, M. (2012b). Characterization of functional, safety, and probiotic properties of *Enterococcus faecalis* UGRA10, a new AS-48-producer strain. *Food Microbiology* 30: 59-67.
- Chandramouli, V., Kailasapathy, K., Peiris, P. and Jones, M. (2004). An improved method of microencapsulation and its evaluation to protect *Lactobacillus spp.* in simulated gastric conditions. *Journal of Microbiological Methods* 56(1): 27-35.
- Chatterjee, S., and Judeh, Z.M. (2015). Encapsulation of fish oil with N-stearoyl O-butylglyceryl chitosan using membrane and ultrasonic emulsification processes. *Carbohydrate Polymers* 123: 432-442.
- Chen, P., Zhang, Q., Dang, H., Liu, X., Tian, F., Zhao, J., Chen, Y., Zhang, H. and Chen, W. (2014). Screening for potential new probiotic based on probiotic properties and  $\alpha$ -glucosidase inhibitory activity. *Food Control* 35(1): 65-72.

- Cheng, X., Liu, R. and He, Y. (2010). A simple method for the preparation of monodisperse protein-loaded microspheres with high encapsulation efficiencies. *European Journal of Pharmaceutics and Biopharmaceutics* 76: 336-341.
- Choi, S.Y., Chang, C.E., Kim, S.C. and So, J.S. (2003). Antimicrobial susceptibility and strain prevalence of Korean vaginal *Lactobacillus* spp. *Anaerobe* 9: 277-280.
- Cizeikiene, D. and Juodeikiene, G. (2013). Antimicrobial activity of lactic acid bacteria against pathogenic and spoilage microorganism isolated from food and their control in wheat bread. *Food Control* 31(2): 539-545.
- Claire, L., Glenn, V., Gibson, R. and Robert, A. (2006). Carbohydrate preference, acid tolerance and bile tolerance in five strains of Bifidobacterium. *Journal of Applied Microbiology* 100: 846-853.
- Clark, P.A. and Martin, J.H.x. (1994). Selection of bifidobacteria for use as dietary adjuncts in cultured dairy foods: III – Tolerance to simulated bile of human stomachs. *Cultured Dairy Products Journal* 29: 18-21.
- Collins, E.B. (1977). Influence of medium and temperature on and products and growth. *Journal Dairy Science* 60: 799-804.
- Colodner, R., Edelstein, H., Chazan, B. and Raz, R. (2003). Vaginal colonization by orally administered *Lactobacillus rhamnosus* GG. *Israel Medical Association Journal* 5(11): 767-769.
- Coolen, M.J., Post, E., Davis, C.C. and L.J., F. (2005). Characterization of microbial communities found in the human vagina by analysis of terminal restriction fragment length polymorphisms of 16S rRNA genes. *Applied Environmental Microbiology* 71: 8729-8737.
- Corcoran, B.M., Ross, R.P., Fitzgerald, G.F. and Stanton, C. (2004). Comparative survival of probiotic lactobacilli spray-dried in the presence of prebiotic substances. *Journal of Applied Microbiology* 96: 1024-1039.
- Damaskos, D. and Kolios, G. (2008). Probiotics and prebiotics in inflammatory bowel disease: microflora 'on the scope'. *British journal of clinical pharmacology* 65(4): 453-467.
- Davey, G.P. and Heap, H.A. (1993). Appearance of the arginine phenotype *Lactococcus lactis* subsp. cremoris 2204 following phage transduction. *Canadian Journal of Microbiology* 39: 745-758.
- de Blanc, A.M. and Bonet, M.E.B. (2010). Chapter 29 - Probiotics in Cancer Prevention. *Bioactive Foods in Promoting Health*. W. Ronald Ross and R. P. Victor. Boston, Academic Press: 497-511.
- de Vuyst, L., Camu, N., De Winter, T., Vandemeulebroecke, K., De Perre, V.V., Vancanneyt, M., De Vos, P. and Cleenwerck, I. (2008). Validation of the (GTG)<sub>5</sub>-rep-PCR fingerprinting technique for rapid classification and identification of acetic acid bacteria, with a focus on isolates from Ghanaian

- fermented cocoa beans. *International Journal of Food Microbiology* 125: 79-90.
- Delcenserie, V., Martel, D., Lamoureux, M., Amiot, J., Boutin, Y. and Roy, D. (2009). Immunomodulatory Effects of Probiotics in the Intestinal Tract. *Current Issues in Molecular Biology* 10: 37-54.
- Dickinson, E. (2003). Hydrocolloids at interfaces and the influence on the properties of dispersed systems. *Food Hydrocolloids* 17: 25-39.
- Dimitonova, S.P., Danova, S.T., Serkedjieva, J.P. and Bakalov, B.V. (2007). Antimicrobial activity and protective properties of vaginal lactobacilli from healthy Bulgarian women. *Anaerobe* 13: 178-184.
- Dinakar, P. and Mistry, V.V. (1994). Growth and viability of *Bifidobacterium bifidum* in cheddar cheese. *Journal of Dairy Science* 77(10): 2854-2864.
- Diosma, G., Romanin, D.E., Rey-Burusco, M.F., Londero, A., and Garrote, G.L. (2014). Yeasts from kefir grains: isolation, identification, and probiotic characterization. *World Journal of Microbiology and Biotechnology*, 30(1): 43-53.
- Doleyres, Y. and Lacroix, C. (2005). Technologies with free and immobilised cells for probiotic *bifidobacteria* production and protection. *International Dairy Journal* 15: 973-988.
- Domig, K. (2003). Methods used for the isolation, enumeration, characterisation and identification of *Enterococcus spp.* 1. Media for isolation and enumeration. *International Journal of Food Microbiology* 88(2-3): 147-164.
- Donaldson, M.S. (2004). Nutrition and cancer: A review of the evidence for an anti-cancer diet. *Nutrition Journal*: 15-21.
- Donaldson, M.S. (2004). Nutrition and cancer: A review of the evidence for an anti-cancer diet. *Nutrition Journal* 3: 19-39.
- Donders, G., Van Calsteren, K. and Bellen, G. (2009). Predictive value for preterm birth of abnormal vaginal flora, bacterial vaginosis and aerobic vaginitis during the first trimester of pregnancy. *International Journal of Obstetrics and Gynaecology* 116: 1315-1324.
- Doulgeraki, A.I. and Pramateftaki, P. (2013). Molecular characterization of lactic acid bacteria isolated from industrially fermented Greek table olives. *LWT - Food Science and Technology* 50(1): 353-356.
- Dubernet, S., Desmasures, N. and Gueguen, M. (2002). A PCR-based method for identification of lactobacilli at the genus level. *FEMS Microbiology Letters* 214: 271-275.
- Dubey, R., Shami, T.C. and Bhasker, R.K.U. (2009). Microencapsulation technology and application. *Defence Science Journal* 59(1): 82-95.

- Dugas, B., Mercenier, A., Lenoir-Wijnkoop, I., Arnaud, C., Dugas, N. and Postaire, E. (1999). Immunity and probiotics. *Trends Immunology Today* 20(9): 387-390.
- Dunne, C., O'Mahony, L., Murphy, L., Thornton, G., Morrissey, D., O'Halloran, S., Feeney, M., Flynn, S., Fitzgerald, G., Daly, C., Kiely, B., O'Sullivan, G.C., Shanahan, F. and Collins, J.K. (2001). *In vitro* selection criteria for probiotic bacteria of human origin: correlation with *in vivo* findings. *Clinical Nutrition* 73: 386-392.
- Ehrstrom, S., Daroczy, K., Rylander, E., Samuelsson, C., Johannesson, U., Anzen, B. and Pahlson, C. (2010). Lactic acid bacteria colonization and clinical outcome after probiotic supplementation in conventionally treated bacterial vaginosis and vulvovaginal candidiasis. *Microbes and Infection* 12(10): 691-699.
- Elli, M., Cattivelli, D., Soldi, S., Bonatti, M. and Morelli, L. (2008). Evaluation of prebiotic potential of refined psyllium (*Plantago ovata*) fiber in healthy women. *Journal of Clinical Gastroenterology* 42: 174-176.
- Endo, A. and Okada, S. (2005). Monitoring the lactic acid bacterial diversity during Shochu fermentation by PCR-denaturing gradient gel electrophoresis. *Journal of Bioscience and Bioengineering* 3: 216-221.
- Ennahar, S. and Sashihara, T. (2000). Class IIa bacteriocins: biosynthesis, structure and activity. *FEMS Microbiology Reviews* 24(1): 85-106.
- Ercolini, D., Moschetti, G., Blaiotta, G. and Coppola, S. (2001). Behavior of variable V3 region from 16S rDNA of lactic acid bacteria in denaturing gradient gel electrophoresis. *Current Microbiology* 42: 199-202.
- Erkkila, S. and Petaja, E. (2000). Screening of commercial meat starter cultures at low pH and in the presence of bile salts for potential probiotic use. *Meat Science* 55: 297-300.
- Eschenbach, D.A., Davick, P.R., Williams, B.L., Klebanoff, S.J., Young-Smith, K., Critchlow, C. and Holmes, K. (1989). Prevalence of Hydrogen Peroxide-Producing *Lactobacillus* Species in Normal Women and Women with Bacterial Vaginosis. *Journal of Clinical Microbiology*: 251-256.
- Ewaschuk, J.B., Walker, J.W., Diaz, H. and Madsen, K.L. (2006). Bioproduction of conjugated linoleic acid by probiotic bacteria occurs *in vitro* and *in vivo* in mice. *Nutrition* 136: 1483-1487.
- Fadok, V.A., Bratton, D.L., Konowal, A., Freed, P.W., Westcott, J.Y. and Henson, P.M. (1998). Macrophages that have ingested apoptotic cells *in vitro* inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF-beta, PGE2, and PAF. *Journal of Clinical Investigation* 101: 890-899.
- Fadok, V.A., Voelker, D.R., Campbell, P.A., Cohen, J.J., Bratton, D.L. and Henson, P.M. (1992). Exposure of phosphatidylserine on the surface of apoptotic lymphocytes triggers specific recognition and removal by macrophages. *Immunology* 148: 2207.

- FAO/WHO. (2001). Food and agriculture organization of the United Nations, World Health Organization. Report of a joint FAO-WHO expert consultation on evaluation of health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria. Cordoba, 2001.
- Fischer, M.H., Yu, N., Gray, G.R., Ralph, J., Anderson, L. and Marlett, J.A. (2004). The gel-forming polysaccharide of psyllium husk (*Plantago ovata* Forsk). *Carbohydrate Research* 339(11): 2009-2017.
- Fong, F.L., Kirjavainen, P., Wong, V.H., and El-Nezami, H. (2015). Immunomodulatory effects of *Lactobacillus rhamnosus* GG on dendritic cells, macrophages and monocytes from healthy donors. *Journal of Functional Foods*, 13: 71-79.
- Forsyth, C.B., Farhadi, A., Jakate, S.M., Tang, Y. and Keshavarzian, S.M.A. (2009). *Lactobacillus* GG treatment ameliorates alcohol-induced intestinal oxidative stress, gut leakiness, and liver injury in a rat model of alcoholic steatohepatitis. *Alcohol* 43: 163-172.
- Franz, C.M.A.P., Huch, M., Abriouel, H., Holzapfel, W. and Galvez, A. (2011). Enterococci as probiotics and their implications in food safety. *International Journal of Food Microbiology* 151(2): 125-140.
- Franz, C.M.A.P., Specht, I., Haberer, P. and Holzapfel, W.H. (2001). Bile salt hydrolase activity of Enterococci isolated from food: screening and quantitative determination. *Journal of Food Protection* 64(725-729).
- Franz, C.M.A.P., Stiles, M.E., Schleifer, K.H. and Holzapfel, W.H. (2003). Enterococci in foods-a conundrum for food safety. *International Journal of Food Microbiology* 88: 105-122.
- Fujimori, S., Gudis, K. and Mitsui, K. (2009). A randomized controlled trial on the efficacy of synbiotic versus probiotic or prebiotic treatment to improve the quality of life in patients with ulcerative colitis. *Nutrition* 25(5): 520-525.
- Galluzzi, L., Maiuri, M.C., Vitale, I., Zischka, H., Zitvogel, L. and Kroemer, G. (2007). Cell death modalities: Classification and pathophysiological implications. *Cell Death and Differentiation* 14: 1237-1266.
- Galvez, A., Lopez, R.L., Abriouel, H., Valdivia, E. and Bem Omar, N. (2008). Application of bacteriocins in the control of foodborne pathogenic and spoilage bacteria. *Critical Reviews in Biotechnology* 28: 125-152.
- Gardiner, G.E., Bouchier, P., OSullivan, E., Kelly, J., Collins, J.K., Fitzgerald, G., Ross, R.P. and Stanton, C. (2002). A spray-dried culture for probiotic cheddar cheese manufacture. *International Dairy Journal* 12(9): 749-756.
- Gevers, D., Huys, G. and Swings, J. (2001). Applicability of rep-PCR fingerprinting for identification of *Lactobacillus* species. *FEMS Microbiology Letters* 205: 31-36.

- Ghosh, C., Baker, J.A. and Moysich, K.B. (2008). Dietary intakes of selected nutrients and food groups and risk of cervical cancer. *Nutrition Cancer* 60(3): 331-341.
- Gillian, E.G., Christine, H., Andrew, W.B., Dee, B. and Gregor, R. (2002). Persistence of *Lactobacillus fermentum* RC-14 and *Lactobacillus rhamnosus* GR-1 but not *L. rhamnosus* GG in the human vagina as demonstrated by randomly amplified polymorphic DNA. *Clinical and Diagnostic Laboratory Immunology* 9: 92-96.
- Giorgi, A., S., Torriani, F., Dellaglio, G.B., Stola, E. and Bernuzzi, L. (1987). Identification of vaginal lactobacilli from asymptomatic women. *Microbiologica* 10: 377-384.
- Giraffa, G. (2002). Enterococci from foods. *FEMS Microbiology Reviews* 26: 163-171.
- Gismondo, M.R., Drago, L. and Lombardi, A. (1999). Review of probiotics available to modify gastrointestinal flora. *International Journal of Antimicrobial Agents* 12: 287-292.
- Gobetti, M., Corsetti, A., Smacchi, E., Zochetti, A. and De Angelis, M. (1998). Production of Crescenza cheese by incorporation of bifidobacteria. *Journal of Dairy Science* 81(1): 37-47.
- Godward, G. and Kailasapathy, K. (2003). Viability and survival of free and encapsulated probiotic bacteria in Cheddar cheese. *Milchwissenschaft* 58(11-12): 624-627.
- Gopal, A., Shah, N.P. and Roginski, H. (1996). Bile tolerance, taurocholate deconjugation and cholesterol removal by *Lactobacillus acidophilus* and *Bifidobacterium spp.* *Milchwissenschaft* 51: 619-623.
- Gorbach, S.I. (2002). Probiotics in the Third Millennium. *Digestive and Liver Disease* 34(2): 2-7.
- Grill, J.P., Perrin, S. and Schneider, F. (2000a). Bile salt toxicity to some *Bifidobacteria* strains: role of conjugated bile salt hydrolase and pH. *Canadian Journal of Microbiology* 46: 878-884.
- Grojean, S., Koziel, V., Vert, P. and Luc Daval, J. (2000). Apoptosis induced by calcein acetoxymethyl ester in human histiocytic lymphoma cell line. *Biochemical Pharmacology* 60: 1751-1759.
- Guarner, F., Perdigon, G., Corthier, G., Salminen, S., Koletzko, B. and Morelli, L. (2005). Should yoghurt cultures be considered probiotic? *British Journal of Nutrition* 93: 783-786.
- Gunay-Esiyok, O.Z.L.E.M., Akcelik, N., and Akcelik, M.U.S. (2014). Identification of Genomic Heterogeneity among *Lactococcus lactis* Strains by Plasmid Profiling, PFGE and 16S rDNA Sequence Analysis. *Polish Journal of Microbiology*, 63(2): 157-166.



- Guo, Q., Cui, S.W., Wang, Q. and Christopher Young, J. (2008). Fractionation and physicochemical characterization of psyllium gum. *Carbohydrate Polymers* 73(1): 35-43.
- Halicka, H.D., Bedner, E. and Darzynkiewicz, Z. (2000). Segregation of RNA and separate packaging of DNA and RNA in apoptotic bodies during apoptosis. *Experimental Cell Research* 260: 248-255.
- Hamid, R., Rotshteyn, Y., Rabadi, L., Parikh, R. and Bullock, P. (2004). Comparison of alamar blue and MTT assays for high through-put screening. *Toxicology in vitro* 18: 703-710.
- Hammes, W.E. and Vogel, R.F. (1995). The genus *Lactobacillus*: The lactic acid bacteria Vol. 2. The general of lactic acid bacteria. J.B. Wood Brian and W.H. Holzapel. Blackie academic & Professional, London, United Kingdom.
- Handwerger, S., Pucci, M.J., Volk, K.J., Liu, J.P. and Lee, M.S. (1994). Vancomycin-resistant *Leuconostoc mesenteroides* and *Lactobacillus casei* synthesize cytoplasmic peptidoglycan precursors that terminate in lactate. *Journal of bacteriology* 176: 260-264.
- Hart, B.L. and Powell, K.L. (1990). Antibacterial Properties of Saliva: Role in Maternal Periparturient Grooming and in Licking Wounds. *Physiology and Behavior* 48: 383-386.
- Hatch, E.E., Herbst, A.L. and Hoover, R.N. (2001). Incidence of squamous neoplasia of the cervix and vagina in women exposed prenatally to diethylstilbestrol (United States). *Cancer Causes Control* 12(9): 837-845.
- Heidebach, T., Forst, P. and Kulozik, U. (2009). Microencapsulation of probiotic cells by means of rennet-gelation of milk proteins. *Food Hydrocolloids* 23: 1670-1677.
- Heidebach, T., Forst, P. and Kulozik, U. (2012). Microencapsulation of probiotic cells for food applications. *Critical Reviews in Food Science and Technology* 52: 291-311.
- Herzog, T.J., Armstrong, D.K., Brady, M.F., Coleman, R.L., Einstein, M.H., Monk, B.J., and Alvarez, R.D. (2014). Ovarian cancer clinical trial endpoints: Society of Gynecologic Oncology white paper. *Gynecologic oncology*, 132(1): 8-17.
- Hirayama, K. and Rafter, J. (2000). The role of probiotic bacteria in cancer prevention. *Microbes and Infection* 2: 671-686.
- Hoesl, C.E. and Altwein, J.E. (2005). The probiotic approach: an alternative treatment option in urology. *European Urology* 47(288-296).
- Holzapel, W.H., Haberer, P., Geisen, R., Bjorkroth, J. and Schillinger, U. (2001). Taxonomy and important features of probiotic microorganisms in food nutrition. *American Journal of Clinical Nutrition* 73(365-373).

- Hou, R.C., M.Y., L., Wang, M.M. and Tzen, J.T. (2003). Increase of viability of entrapped cells of *Lactobacillus delbrueckii* spp. *Bulgarius* in artificial sesame oil emulsions. *Journal of Dairy Science* 86: 424-428.
- Hoveyda, N., Heneghan, C., Mahtani, K.R., Perera, R., Roberts, N. and Glasziou, P. (2009). A systematic review and meta-analysis: probiotics in the treatment of irritable bowel syndrome. *BMC Gastroenterology* 9(1): 15-23.
- Howarth, G.S. and Wang, H. (2013). Role of Endogenous Microbiota, Probiotics and Their Biological Products in Human Health. *Nutrients* 5: 58-81.
- Hsu, Y.L., Kuo, P.L., Lin, L.T. and Lin, C.C. (2005). Asiatic acid, a triterpene, induces apoptosis and cell cycle arrest through activation of extracellular signal-regulated kinase and p38 mitogen-activated protein kinase pathways in human breast cancer cells. *Journal of Pharmacology and Experimental Therapeutics* 313(1): 333-344.
- Hyman, R.W., Fukushima, M., Diamond, L., Kumm, J., Giudice, L.C. and Davis, R.W. (2005). Microbes on the human vaginal epithelium. *PNAS, Proceedings of the National Academy of Sciences* 102(7952-7957).
- Iannitti, T. and Palmier, B. (2010). Therapeutical use of probiotic formulations in clinical practice. *Clinical Nutrition* 29: 701-725.
- Inaba, H., Husain, S., Horibe, A. and Haruki, N. (2006). Heat and Mass Transfer Analysis of Fluidized Bed Grain Drying. *Memoirs of the Faculty of Engineering, Okayama University* 41: 52-62.
- Jacobsen, C.N., Nielsen, V.R., Hayford, A.E., Moller, P.L., Michaelsen, K.F., Paerregaard, A., Sandström, B., Tvede, M. and Jacobsen, M. (1999). Screening of probiotic activities of forty seven strains of *Lactobacillus* spp. by *in vitro* techniques and evaluation of the colonization ability of five selected strains in humans. *Applied and Environmental Microbiology* 65(11): 4949-4956.
- Jemal, A., Siegal, R., Ward, E., Hao, Y., Xu, J., Murray, T. and Thun, M.J. (2008). Cancer statistics. *CAA Cancer Journal for Clinicians* 58: 71-96.
- Jemal, A., Siegal, R., Ward, E., Murray, T., Xu, J. and Thun, M.J. (2007). Cancer statistics. *CAA Cancer Journal for Clinicians* 57: 43-66.
- Jett, B.D., Huycke, M.M. and Gilmore, M.S. (1994). Virulence of enterococci. *Clinical Microbiology Reviews* 7: 462-278.
- Johnson, A.P., Uttley, A.H.C., Woodford, N. and George, R.C. (1990). Resistance to vancomycin and teicoplanin: an emerging clinical problem. *Clinical Microbiology Reviews* 3: 280-291.
- Johnson, W. (2005). Final report of the safety assessment of Acacia catechu Gum, Acacia concinna fruit extract, Acacia dealbata leaf extract, Acacia dealbata leaf wax, Acacia decurrens extract, Acacia farnesiana flower wax, Acacia farnesiana gum, Acacia senegal extract, Acacia senegal gum, and Acacia senegal gum extract. *International Journal Toxicology* 24: 75-118.

- Jokovic, N. and Nikolic, M. (2008). A survey of the lactic acid bacteria isolated from Serbian artisanal dairy product kajmak. *International Journal of Food Microbiology* 127(3): 305-311.
- Kailasapathy, K. (2002). Microencapsulation of Probiotic Bacteria: Technology and Potential Applications. *Current Issues Intestinal Microbiology* 3: 39-48.
- Kailasapathy, K. (2002). Microencapsulation of probiotic bacteria: technology and potential applications. *Current Issues in Intestinal Microbiology* 3(2): 39-48.
- Kailasapathy, K. and Masondole, L. (2005). Survival of free and microencapsulated *Lactobacillus acidophilus* and *Bifidobacterium lactis* and their effect on texture of feta cheese. *Australian Journal of Dairy Technology* 60(3): 252-258.
- Kalai, M., Lamkanfi, M., Denecker, G., Boogmans, M., Lippens, S. and Meeus, A. (2003). Regulation of the expression and processing of caspase-12. *Journal of Cell Biology* 162: 457-467.
- Kanmani, P., Kumar, R.S., Yuvaraj, N., Paari, K.A., Pattikumar, V. and Arul, V. (2011). Cryopreservation and microencapsulation of a probiotic in alginate-chitosan capsules improves survival in simulated gastrointestinal conditions. *Biotechnology and Bioprocess Engineering* 16: 1106-1114.
- Karaiskou, S., Blekas, G. and Paraskevopoulou, A. (2008). Aroma release from gum Arabic or egg yolk/xanthan-stabilized oil-in-water emulsions. *Food Research International* 41: 637-645.
- Kashlet, E.R. (1987). Bioenergetics of Lactic acid bacteria: cytoplasmic pH osmotolerance. *FEMS Microbiology Review* 46: 233-244.
- Kaufmann, S. and Hengartner, M. (2001). Programmed cell death: alive and well in the new millennium. *Trends in Cell Biology* 262: 526-534.
- Kaur, I.P., Chopra, K. and Saini, A. (2002). Probiotics: Potential pharmaceutical applications. *European Journal of Pharmaceutical Sciences* 15: 1-9.
- Kawamoto, K., Horibe, I. and Uchida, K. (1989). Purification and characterization of a new hydrolase for conjugated bile acids, chenodeoxy- cholytaurine hydrolase, from *Bacteroides vulgatus*. *Journal of Biochemistry* 106: 1049-1053.
- Kaysner, F.H. (2003). Safety aspects of enterococci from the medical point of view. *International Journal of Food Microbiology* 88: 255-262.
- Kebary, K.M.K., Hussein, S.A. and Badawi, R.M. (1998). Improving viability of *Bifidobacteria* and their effect on frozen ice milk. *Egyptian Journal of Dairy Science* 26: 319-337.
- Kelly, J.R. and Duggan, J.M. (2003). Gastric cancer epidemiology and risk factors. *Journal of Clinical Epidemiology* 56: 1-9.

- Kiess, W. and Gallaher, B. (1998). Hormonal control of programmed cell death/apoptosis. *European Journal of Endocrinology* 138: 482-491.
- Kim, S.J., Cho, S.Y., Kim, S.H., Song, O.J., Shin, S., Cha, D.S. and Park, H.J. (2008a). Effect of microencapsulation on viability and other characteristics in *Lactobacillus acidophilus* ATCC 43121. *LWT—Food Science and Technology* 41(3): 493-500.
- Kim, Y., Lee, D., Cho, J., Yang, J., Chung, M., Kim, K. and Ha, N. (2008b). Inhibition of proliferation in colon cancer cell lines and harmful enzyme activity of colon bacteria by *Bifidobacterium adolescentis* SPM0212. *Archives of Pharmacal Research* 31(4): 468-473.
- Kim, W. (2014). The genus *Lactococcus*. *Lactic Acid Bacteria: Biodiversity and Taxonomy*, 429-443.
- Kim, S., Jo, A., and Ahn, J. (2015). Application of chitosan–alginate microspheres for the sustained release of bacteriophage in simulated gastrointestinal conditions. *International Journal of Food Science & Technology*.
- Klein, G., Pack, A., Bonaparte, C. and Reuter, G. (1998). Taxonomy and physiology of probiotic lactic acid bacteria. *International Journal of Food Microbiology* 41: 103-105.
- Klemmer, K.J., Korber, D.R., Low, N.H. and Nickerson, M.T. (2011). Pea protein-based capsules for probiotic and prebiotic delivery. *International Journal of Food Science and Technology* 46: 2248-2256.
- Korzeniewski, C. and Callewaert, D.M. (1983). An enzyme-release assay for natural cytotoxicity. *Journal of Immunological Methods* 64: 313-320.
- Kos, B., Suskovic, J., Goreta, J. and Matosic, S. (2000). Effect of protectors on the viability of *Lactobacillus acidophilus* M92 in simulated gastrointestinal conditions. *Food Technology and Biotechnology* 38: 121-127.
- Koskey, A.M., Fisher, J.C., Traudt, M.F., Newton, R.J., and McLellan, S.L. (2014). Analysis of the gull fecal microbial community reveals the dominance of *Catelicoccus marimammalium* in relation to culturable Enterococci. *Applied and environmental microbiology*, 80(2): 757-765.
- Krasaekoopt, W., Bhandari, B. and Deeth, H. (2003). Evaluation of encapsulation techniques of probiotics for yoghurt. *International Dairy Journal* 13(1): 3-13.
- Krasaekoopt, W., Bhandari, B. and Deeth, H. (2004). The influence of coating materials on some properties of alginate beads and survivability of microencapsulated probiotic bacteria. *International Dairy Journal* 14(8): 737-743.
- Kuitunen, M., Kukkonen, A. and Savilahti, E. (2012). Impact of maternal allergy and use of probiotics during pregnancy on breast milk cytokines and food antibodies and development of allergy in children until 5 years. *International Archives of Allergy and Immunology* 159(2): 162-170.

- Kumar, R. (2003). Another tie that binds the MTA family to breast cancer. *Cell* 113: 142-143.
- Kurdi, P., Veen, H.W.V., Tanaka, H., Mierau, I., Konings, W.N., Tannock, G.W., Tomita, F. and Yokota, A. (2000). Cholic acid is accumulated spontaneously, driven by membrane delta pH, in many lactobacilli. *Journal of Bacteriology* 182: 6525-6528.
- Kwon, H.S., Yang, E.H., Yeon, S.W., Kang, B.H. and Kim, T.Y. (2004). Rapid identification of probiotic *Lactobacillus* species by multiplex pcr using species-specific primers based on the region extending from 16s rRNA through 23s rRNA. *FEMS Microbiology Letters* 239: 267-275.
- Lachlak, N., Ageron, E., Zampatti, O., Michel, G. and Grimont, P.A. (1996). Composition of the *Lactobacillus acidophilus* complex isolated from vaginal flora. *New Microbiol.* 19: 123-132.
- Lachlak, N., E., Ageron, O., Zampatti, G.M. and P.A., G. (1996b). Composition of the *Lactobacillus acidophilus* complex isolated from vaginal flora. *New Microbiology* 19: 123-132.
- Lambert, J.M., Weinbreck, F. and Kleerebezem, M. (2008). *In vitro* analysis of protection of the enzyme bile salt hydrolase against enteric conditions by whey protein-gumarabic microencapsulation. *Journal of Agricultural and Food Chemistry* 56: 8360-8364.
- Lane, D.J., Pace, B., Olsen, G.J., Stahl, D.A., Sogint, M.L. and Pace, N. (1986). Rapid determination of 16S ribosomal RNA sequences for phylogenetic analyses. *PNAS, Proceedings of the National Academy of Sciences* 82: 6955-6959.
- Larena, I., Melgarejo, P. and De Cal, A. (2003). Drying of conidia of *Penicillium oxalicum*, a biological control agent against fusarium wilt of tomato. *Journal of Pytopathology* 151: 600-606.
- Lee, E., Kwon, B. and Song, G. (2004). Heyneanol A induces apoptosis via cytochrome c release and caspase activation in human leukemia U937 cells. *Life Science* 74: 2313-2326.
- Lee, H., Yoon, H., Ji, Y., Kim, H., Park, H., Lee, J., Shin, H. and Holzapfel, W. (2011). Functional properties of *Lactobacillus* strains isolated from kimchi. *International Journal of Food Microbiology* 145: 155-161.
- Lee, J.W., Shin, J.G., Kim, E.H., Kang, H.E., Yim, I.B., Kim, J.Y., Joo, H.G. and Woo, H.J. (2004). Immunomodulatory and antitumor effects in vivo by the cytoplasmic fraction of *Lactobacillus casei* and *Bifidobacterium longum*. *Journal of Veterinary Science* 5: 41-48.
- Lee, Y.J., Yu, W.K. and Heo, T.R. (2003). Identification and screening for antimicrobial activity against *Clostridium difficile* of *Bifidobacterium* and *Lactobacillus* species isolated from healthy infant faeces. *International Journal of Antimicrobial Agents* 21: 340-346.

- Li, Y., Ahmed, F., Ali, S., Philip, P.A., Kucuk, O. and Sarkar, F.H. (2005). Inactivation of nuclear factor kappaB by soy isoflavone genistein contributes to increased apoptosis induced by chemotherapeutic agent in human cancer cells. *Cancer Research* 65(15): 6934-6942.
- Lin, P.W., Nasr, T.R., Berardinelli, A.J., Kumar, A. and Neish, A.S. (2008). The probiotic *Lactobacillus* GG may augment intestinal host defense by regulating apoptosis and promoting cytoprotective responses in the developing murine gut. *Pediatric Research* 64: 511-516.
- Liong, M.T. (2008). Safety of probiotics: translocation and infection. *Nutrition Research Reviews* 66(4): 192-202.
- Liu, W. and Bao, Q. (2012). Isolation and identification of lactic acid bacteria from Tarag in Eastern Inner Mongolia of China by 16S rRNA sequences and DGGE analysis. *Microbiological Research* 167(2): 110-115.
- Lodish, H., Berk, A., Zipursky, L.S., Matsudaira, P., Baltimore, D. and Darnell, J. (2000). *Cancer. Molecular Cell Biology*. New York: W.H. Freeman and Company.
- Loreti, G., Maroni, A., Del Curto, M.D., Melocchi, A., Gazzaniga, A., and Zema, L. (2014). Evaluation of hot-melt extrusion technique in the preparation of HPC matrices for prolonged release. *European Journal of Pharmaceutical Sciences* 52: 77-85.
- Lopez-Cordoba, A., Deladino, L., Agudelo-Mesa, L., and Martino, M. (2014). Yerba mate antioxidant powders obtained by co-crystallization: Stability during storage. *Journal of Food Engineering* 124: 158-165.
- Lotfipour, F., Mirzaeei, S. and Maghsoodi, M. (2012). Preparation and Characterization of Alginate and Psyllium Beads Containing *Lactobacillus acidophilus*. *The Scientific World Journal*: 8 pages.
- Lundeen, S.G. and Savage, D.C. (1990). Characterization and purification of bile salt hydrolase from *Lactobacillus* spp. strain 100-100. *Journal of Bacteriology* 172: 4171-4177.
- Ma, B., Forney, L.J. and Ravel, J. (2012). Vaginal Microbiome: Rethinking Health and Disease. *Annual Review Microbiology* 66: 371-389.
- Ma, Y., Pacan, J.C., Wang, Q., Sabour, P.M., Huang, X.Q. and Xu, Y. (2012b). Enhanced alginate microspheres as means of oral delivery of bacteriophage for reducing *Staphylococcus aureus* intestinal carriage. *Food Hydrocolloids* 26: 434-440.
- Ma, Y., Pacan, J.C., Wang, Q., Xu, Y., Huang, X. and Korenevsky, A. (2008). Microencapsulation of bacteriophage Felix O1 into chitosan-alginate microspheres for oral delivery. *Applied and Environmental Microbiology* 74: 4799-4805.

- MacPhee, R., Hummelen, R., Bisanz, J., Miller, W. and Reid, G. (2010). Probiotic strategies for the treatment and prevention of bacterial vaginosis. *Expert Opinion on Pharmacotherapy* 11: 2985-2995.
- Mahfoudhi, N., Sessa, M., Chouaibi, M., Ferrari, G., Donsi, F. and Hamdi, S. (2014). Assessment of emulsifying ability of almond gum in comparison with gum arabic using response surface methodology. *Food Hydrocolloids* 37: 49-59.
- Maldonado, N.C., de Ruiz, C.S., Otero, M.C., Sesma, F. and Nader-Macías, M.E. (2012). Lactic acid bacteria isolated from young calves – Characterization and potential as probiotics. *Research in Veterinary Science* 92(2): 342-349.
- Maldonado Galdeano, C., Novotny Nunez, I., Carmuega, E., de Moreno de LeBlanc, A., and Perdigon, G. (2015). Role of Probiotics and Functional Foods in Health: Gut Immune Stimulation by Two Probiotic Strains and a Potential Probiotic Yoghurt. *Endocrine, Metabolic & Immune Disorders-Drug Targets (Formerly Current Drug Targets-Immune, Endocrine & Metabolic Disorders)*, 15(1): 37-45.
- Mandal, S. and Puniya, A.K. (2006). Effect of alginate concentrations on survival of microencapsulated *Lactobacillus casei* NCDC-298. *International Dairy Journal* 16(10): 1190-1195.
- Martin, H.L., Richardson, B.A., Nyange, P.M., Lavreys, L., Hillier, S.L., Chohan, B., Mandaliya, K., Ndinya-Achola, J.O., Bwayo, J. and Kreiss, J. (1999). Vaginal lactobacilli, microbial flora, and risk of human immunodeficiency virus type 1 and sexually transmitted disease acquisition. *The Journal of Infectious Diseases* 180: 1863-1868.
- Massad, L.S., Einstein, M.H. and Huh, W.K. (2013). Updated Consensus Guidelines for the Management of Abnormal Cervical Cancer Screening Tests and Cancer Precursors. *Journal of Lower Genital Tract Disease* 17(5): 1-27.
- Mastromarino, P., Macchia, S., Meggiorini, L., Trinchieri, V., Mosca, L., Perluigi, M. and Midulla, C. (2009). Effectiveness of *Lactobacillus*-containing vaginal tablets in the treatment of symptomatic bacterial vaginosis. *Clinical Microbiology and Infection* 15(1): 67-74.
- Mehra, N., Majumdar, R.S., kumar, S. and Dhewa, T. (2012). Probiotics: preventive and clinical applications. *Biotechnology Research Bulletin* 1: 15-20.
- Metchnikoff, E., 1907. (1907). The prolongation of life, London: William Heinemann.
- Meurman, J.H. (2005). Probiotics: do they have a role in oral medicine and dentistry? *European Journal Oral Science* 113: 188-196.
- Milanovic, J., Manojlovic, V., Levic, S., Rajic, N., Nedovic, V. and Bugarski, B. (2010). Microencapsulation of Flavors in Carnauba Wax. *Sensors* 10: 901-912.

- Millqvist-Fureby, A., Malmsten, M. and Bergenstahl, B. (2000). An aqueous polymer two-phase system as carrier in the spray-drying of biological material. *Journal of Colloid and Interface Science* 225: 54-61.
- Miranda, C.L., Stevens, J.F., Hemlrich, A., Henderson, M.C., Rodriguez, R.J., Yang, Y.H., Deinzer, M.L., Barnes, D.W. and Buhler, D.R. (1999). Antiproliferative and cytotoxic effects of prenylated flavonoids from hops (*Human lupulus*) in human cancer cell lines. *Food and Chemical Toxicology* 37: 271-285.
- Mirzaei, H. and Barzgar, A. (2012). Isolation and Molecular Study of Potentially Probiotic Lactobacilli in Traditional White Cheese of Tabriz in Iran. *Annals of Biological Research* 3(4): 2019-2022.
- Moghadam, M.S., Foo, H.L., Leow, T.C., Abdul Rahman, R. and Loh, T.C. (2010). Novel bacteriocinogenic *Lactobacillus plantarum* strains and their differentiation by sequence analysis of 16S rDNA, 16S-23S AND 23S-5S intergenic spacer regions and randomly amplified polymorphic DNA analysis. *Food Technology Biotechnology* 48: 476-483.
- Mokarram, R.R., Mortazavi, S.A., Najafi, M.B.H. and Shahidi, F. (2009). The influence of multi stage alginate coating on survivability of potential probiotic bacteria in simulated gastric and intestinal juice. *Food Research International* 42(8): 1040-1045.
- Mombelli, B. and Gismondo, M.R. (2000). The use of probiotics in medical practice. *International Journal of Antimicrobial Agents* 16: 531-536.
- Moreira, J.L.S., Mota, R.M., Horta, M.F., Teixeira, S.M., Neuman E., Nicoli J. R. and A.C., N. (2005). Identification to the species level of lactobacillus isolated in probiotic prospecting studies of human, animal or food origin by 16S-23S rRNA restriction profiling. *BMC Microbiology* 5(15): 1-9.
- Morgan, C.A., Herman, N., Gauci, M., White, P.A. and Vesey, G. (2006). Preservation of micro-organisms by drying; A review. *Journal of Microbiological Methods* 66: 183-193.
- Mortazavian, A., Razavi, S.H., Ehsani, M.R. and Sohrabvandi, S. (2007). Principles and methods of microencapsulation of probiotic microorganisms. *Iranian Journal of Biotechnology* 5(1): 1-17.
- Moser, S.A. and Savage, D.C. (2001). Bile salt hydrolase activity and resistance to toxicity of conjugated bile salts are unrelated properties in Lactobacilli. *Applied Environmental Microbiology* 67: 3476-3480.
- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunology Methods* 16: 55-63.
- Mundt, J.O., Sneath, P.H.A., Mair, N.S., Sharpe, M.E. and Holt, J.G. (1986). Enterococci: Bergey's Manual of Systematic Bacteriology. *Williams and Wilkins, Baltimore*. 2: 1063-1065.



- Muthukumarasamy, P., Allan-Wojtas, P. and Holley, R.A. (2006). Stability of *Lactobacillus* microcapsules and survival in simulated gastrointestinal conditions and in yoghurt. *International Dairy Journal* 14: 505-515.
- Nam, H., Whang, K. and Lee, Y. (2007). Analysis of Vaginal Lactic Acid Producing Bacteria in Healthy Women. *Journal of Microbiology* 45(6): 515-520.
- NCCLS, 2007. Performance standards for antimicrobial susceptibility testing, seventeenth informational supplement. M100-S17. Vol. 27 No. 1.
- Nedovic, V., Kalusevic, A., Manojlovic, V., Levic, S. and Bugarski, B. (2011). An overview of encapsulation technologies for food applications. *Procedia Food Science* 1: 1806-1815.
- Nesterenko, A., Alric, I., Violleau, F., Silvestre, F. and Durrieu, V. (2013). A new way of valorizing biomaterials: The use of sunflower protein for  $\alpha$ -tocopherol microencapsulation. *Food Research International* 53: 115-124.
- Nick, G., Jussila, M., Hoste, B., Niemi, R.M., Kaijalainen, S., De Lajudie, P., Gillis, M., De Bruijn, F.J. and Lindstrom, K. (1999). Rhizobia isolated from root nodules of tropical leguminous trees characterized using DNA-DNA dot-blot hybridisation and rep-PCR genomic fingerprinting. *Systematic and Applied Microbiology* 22: 287-299.
- Nishi, K.K. and Jayakrishnan, A. (2007). Self-gelling primaquine-gum Arabic conjugate: 501 an injectable controlled delivery system for primaquine. *Biomacromolecules* 8(502): 84-90.
- Nowroozi, J., Mirzaei, M. and Norouzi, M. (2004). Study of *Lactobacillus* as Probiotic Bacteria. *Iranian Journal of Public Health* 33(2): 1-7.
- O'Bryan, C.A., Crandall, P.G., Ricke, S.C., and Ndahetuye, J.B. (2014). Lactic acid bacteria (LAB) as antimicrobials in food products: types and mechanisms of action. *Handbook of Natural Antimicrobials for Food Safety and Quality*, 117.
- Onsaard, E., Putthanimon, J., Singthong, J. and Thammarutwasik, P. (2014). Influence of maltodextrin and environmental stresses on stability of whey protein concentrate/ $\kappa$ -carrageenan stabilized sesame oil-in-water emulsions. *Food Science and Technology International* 20(8): 617-628.
- O'Shea, E.F., Cotter, P.D., Stanton, C., Ross, R.P. and Hill, C. (2012). Production of bioactive substances by intestinal bacteria as a basis for explaining probiotic mechanisms: Bacteriocins and conjugated linoleic acid. *International Journal of Food Microbiology* 152(3): 189-205.
- Oelschlaeger, T.A. (2010). Mechanisms of probiotic actions – A review. *International Journal of Medical Microbiology* 300(1): 57-62.
- Onderdonk, A.B. and Wissemann, K.W. (1993). Normal vaginal microbiota, p. 285-304. In P. Elsner and J. Martins (ed.), *Vulvovaginitis*. Marcel Dekker Inc., New York, N.Y., USA.

- Orla-Jensen, S. 1919. The lactic acid bacteria. in Koeniglicher Hof Boghandel. Copenhagen.
- Orozco-Villafuerte, J., Cruz-Sosa, F., Ponce-Alquicira, E. and Vernon-Carter, E.J. (2003). Mesquite gum: fractionation and characterization of the gum exuded from *Prosopis laevigata* obtained from plant tissue culture and from wild trees. *Carbohydrate Polymers* 54: 327-333.
- Otero, M.C., Morelli, L. and Nader-Macias, M.E. (2006). Probiotic properties of vaginal lactic acid bacteria to prevent metritis in cattle. *Letters in Applied Microbiology* 43: 91-97.
- Ozer, B., Kirmaci, H.A., Senel, E., Atamer, M. and Hayaloglu, A. (2009). Improving the viability of *Bifidobacterium bifidum* BB-12 and *Lactobacillus acidophilus* LA-5 in white-brined cheese by microencapsulation. *International Dairy Journal* 19(1): 22-29.
- Ozer, B., Uzun, Y.S. and Kirmaci, H.A. (2008). Effect of microencapsulation on viability of *Lactobacillus acidophilus* LA-5 and *Bifidobacterium bifidum* BB-12 during Kasar cheese ripening. *International Journal of Dairy Technology* 61(3): 237-244.
- Palmer, K., Kos, V. and Gilmore, M. (2010). Horizontal gene transfer and the genomics of enterococcal antibiotic resistance. *Current opinion in microbiology* 13(5): 632-639.
- Pan, X., Chen, F., Wu, T., Tang, H. and Zhao, Z. (2009). The acid, bile tolerance and antimicrobial property of *Lactobacillus acidophilus* NIT. *Food Control* 20: 598-602.
- Paolillo, R., Romano Carratelli, C., Sorrentino, S., Mazzola, N. and Rizzo, A. (2009). Immunomodulatory effects of *Lactobacillus plantarum* on human colon cancer cells. *International Immunopharmacology* 9(11): 1265-1271.
- Parker, R.B. (1974). Probiotics: The other half of the antibiotics story. *Animal Nutrition Health* 29: 4-8.
- Parkin, D.M. (2005). Global Cancer Statistics. *CAA Cancer Journal for Clinicians* 55: 74-108.
- Parkin, D.M., Bray, F., Ferlay, J. and Pisani, P. (2005). Global Cancer Statistics. *CAA Cancer Journal for Clinicians* 55: 74-108.
- Pascual, L., Daniele, M., Giordano, W., Pajaro, M. and Barberis, I. (2008). Purification and partial characterization of novel bacteriocin L23 produced by *Lactobacillus fermentum* L23. *Current Microbiology* 56: 397-402.
- Pascual, L., Daniele, M.B., Ruiz, F., Giordano, W., Pajaro, C. and Barberis, L. (2008). *Lactobacillus rhamnosus* L60, a potential probiotic isolated from the human vagina. *Journal of General and Applied Microbiology* 54(3): 141-148.

- Pascual, L., Ruiz, F., Giordano, W. and Barberis, I.L. (2010). Vaginal colonization and activity of the probiotic bacterium *Lactobacillus fermentum* L23 in a murine model of vaginal tract infection. *Journal of Medical Microbiology* 59: 360-364.
- Pascual, L.M., Daniele, M.B., Pajaro, C. and Barberis, L. (2006). *Lactobacillus* species isolated from the vagina: identification, hydrogen peroxide production and noxynol-9 resistance. *Contraception* 73(1): 78-81.
- Paster, B.J., Olsen, I., Aas, J.A. and Dewhirst, F.E. (2006). The breadth of bacterial diversity in the human periodontal pocket and other oral sites. *Journal of Periodontology* 42: 80-87.
- Patterson, J.A. and Burkholder, K.M. (2003). Application of prebiotics and probiotics in poultry production. *Poultry Science* 82: 627-631.
- Pereira, D.I.A. and Gibson, G. (2002). Cholesterol assimilation by lactic acid bacteria and bifidobacteria isolated from the human gut. *Applied and Environmental Microbiology* 68: 4689-4693.
- Pessi, T., Sutas, Y., Saxelin, M., Kallioinen, H. and Isolauri, E. (1999). Antiproliferative effects of homogenates derived from five strains of candidate probiotic bacteria. *Applied Environmental Microbiology* 65: 4725-4728.
- Petterson, H.E. (1988). Starters for fermented milks. Sec.2: Mesophylic starter cultures. *IDF Bull.* 227: 19-26.
- Picot, A. and Lacroix, C. (2003). Effect of micronization on viability and thermotolerance of probiotic freeze-dried cultures. *International Dairy Journal* 13: 455-462.
- Picot, A. and Lacroix, C. (2004). Encapsulation of *bifidobacteria* in whey protein-based microcapsules and survival in simulated gastrointestinal conditions and in yoghurt. *International Dairy Journal* 14: 505-515.
- Pollak, M.N. and Foulkes, W.D. (2003). Challenges to cancer control by screening. *Nature Reviews Cancer* 3: 297-303.
- Praagman, J., Dalmeijer, G.W., van der Schouw, Y.T., Soedamah-Muthu, S.S., Monique Verschuren, W.M., Bas Bueno-de-Mesquita, H., and Beulens, J.W. (2015). The relationship between fermented food intake and mortality risk in the European Prospective Investigation into Cancer and Nutrition-Netherlands cohort. *British Journal of Nutrition*, 1-9.
- Prakash, S. and Jones, M.L. (2005). Artificial cell therapy: New strategies for the therapeutic delivery of live bacteria. *Journal of Biomedicine and Biotechnology* 1: 44-56.
- Prasad, J., Gill, H., Smart, J. and Gopal, P.K. (1998). Selection and Characterization of *Lactobacillus* and *Bifidobacterium* strains for use as probiotic. *International Dairy Journal* 8: 993-1002.

- Prevost, H. and Divies, C. (1987). Fresh fermented cheese production with continuous pre-fermented milk by a mixed culture of mesophilic lactic streptococci entrapped in Ca-alginate. *Biotechnology Letters* 9(11): 789-794.
- Prinsloo, S., Pieters, R. and Bezuidenhout, C.C. (2013). A cell viability assay to determine the cytotoxic effects of water contaminated by microbes. *South African Journal of Science* 109(7/8): 1-4.
- Prisciandaro, L.D., Geier, M.S., Butler, R.N., Cummins, A.G. and Howarth, G.S. (2011). Evidence supporting the use of probiotics for the prevention and treatment of chemotherapy-induced intestinal mucositis. *Critical Reviews in Food Science and Nutrition* 51: 239-247.
- Pothakos, V., Snauwaert, C., De Vos, P., Huys, G., and Devlieghere, F. (2014). Psychrotrophic members of *Leuconostoc gasicomitatum*, *Leuconostoc gelidum* and *Lactococcus piscium* dominate at the end of shelf-life in packaged and chilled-stored food products in Belgium. *Food microbiology*, 39: 61-67.
- Prusse, U., Bilancetti, L., Bucko, M., Bugarski, B., Bukowski, J. and Gemeiner, P. (2008). Comparison of different technologies for alginate beads production. *Chemical Papers* 62(4): 364-374.
- Quigley, E.M. (2008). Probiotics in functional gastrointestinal disorders: what are the facts? *Current Opinion in Pharmacology* 8(6): 704-708.
- Quwehand, A.C. and Vesterlund, S. (2004). Antimicrobial components from lactic acid bacteria. *Lactic Acid Bacteria Microbiological and Functional Aspects*. New York: Marcel Dekker Inc.
- Rafter, J. (2003). Probiotics and colon cancer. *Best Practice and Research Clinical Gastroenterology* 17(5): 849-859.
- Rafter, J. (2004). the effects of probiotic on colon cancer development. *Nutrition Reviews* 17: 277-284.
- Rahmati, M., Bannazadeh, H., Fazli, D., Tahmasebzadeh, F., Farrokhi, A. and Rasmi, Y. (2011). The apoptotic and cytotoxic effects of Polygonum aviculare extract on Hela-S cervical cancer cell line. *African Journal of Biochemistry Research* 5(14): 373-378.
- Rajam, R., Karthik, P., Parthasarathi, S., Joseph, G.S. and Anandharamakrishnan, C. (2012). Effect of whey protein-alginate wall systems on survival of microencapsulated *Lactobacillus plantarum* in simulated gastrointestinal conditions. *Journal of Functional Foods* 4: 891-898.
- Ravel, J., Gajer, P. and Abdo, Z. (2011). Vaginal microbiome of reproductive-age women. *PNAS, Proceedings of the National Academy of Sciences* 108: 4680-4687.

- Ray, D., Alpini, G., and Glaser, S. (2014). Probiotic *Bifidobacterium* species: potential beneficial effects in diarrheal disorders. Focus on “Probiotic *Bifidobacterium* species stimulate human SLC26A3 gene function and expression in intestinal epithelial cells”. *American Journal of Physiology-Cell Physiology*, 307(12): 1081-1083.
- Redon, R., Mulle, D., Caulee, K., Wanherdrick, K., Abecassis, J. and Manoir, S. (2001). A simple specific pattern of chromosomal aberrations at early stages of head and neck squamous cell carcinomas. *Cancer Research* 61: 63-68.
- Redondo-Lopez, V., Cook, R.L. and Sobel, J.D. (1990). Emerging role of lactobacilli in the control and maintenance of the vaginal bacterial microflora. *Reviews of infectious diseases* 12: 856-872.
- Reed, J. (2001). Apoptosis-regulating proteins as targets for drug discovery. *Trends in Molecular Medicine* 7: 314-319.
- Reed, J. (2003). Apoptosis-targeted therapies for cancer. *Cancer Cells* 3: 17-25.
- Reid, G. (2002). Probiotics for urogenital health. *Nutrition in Clinical Care* 5(1): 3-8.
- Reid, G. and Bruce, A.W. (2003). Urogenital infections in women: can probiotics help? *Postgraduate Medical Journal* 79(934): 428-432.
- Reid, G., Dols, J. and Miller, W. (2009). Targeting the vaginal microbiota with probiotics as a means to counteract infections. *Current Opinion in Clinical Nutrition and Metabolic Care* 12(6): 583-587.
- Reid, G., McGroarty, J.A., Tomeczek, L. and Bruce, A.W. (1996). Identification and plasmid profiles of *Lactobacillus* species from the vagina of 100 healthy women. *FEMS Immunology and Medical Microbiology* 15: 23-26.
- Rishniw, M. and Wynn, S.G. (2011). Azodyl, a synbiotic, fails to alter azotemia in cats with chronic kidney disease when sprinkled onto food. *Journal of Feline Medicine and Surgery* 13(6): 405-409.
- Rolfe, R.D. (2000). The Role of probiotic cultures in the control of gastrointestinal health. *Jornal of Nutrition* 130: 396-402.
- Roukos, D.H., Agnantis, N.J., Fatouros, M. and Kappas, A.M. (2002). Gastric Cancer: Introduction, Pathology, Epidemiology. *Gastric Breast Cancer* 1(1): 1-3.
- Roy, D., Sirios, S. and Vincent, D. (2001). Molecular discrimination of Lactobacilli used as starter and probiotic cultures by amplified ribosomal DNA restriction analysis. . *Current Microbiology an International Journal* 42: 282-289.
- Russell, D.A., Ross, R.P., Fitzgerald, G.F. and Stanton, C. (2011). Metabolic activities and probiotic potential of bifidobacteria. *International Journal of Food Microbiology* 149: 88-105.

- Saad, N., Delattre, C., Urdaci, M., Schmitter, J.M. and Bressollier, P. (2013). An overview of the last advances in probiotic and prebiotic field. *LWT - Food Science and Technology* 50(1): 1-16.
- Saarela, M., Mogensen, G., Fonden, R., Matto, J. and Mattila-Sandholm, T. (2000). Probiotic bacteria: safety, functional and technological properties. *Journal of Biotechnology* 84: 197-215.
- Sabikhi, L., Babu, R., Thompkinson, D.K. and Kapila, S. (2010). Resistance of microencapsulated *Lactobacillus acidophilus* LA1 to processing treatments and simulated gut conditions. *Food Bioprocess Technology* 3: 586-593.
- Sainsbury, J.R.C., Anderson, T.J. and Morgan, D.A.L. (2000). Breast cancer. *British Medical Journal* 321: 745-751.
- Sah, B.N., Vasiljevic, T., McKechnie, S., and Donkor, O.N. (2015). Identification of Anticancer Peptides from Bovine Milk Proteins and Their Potential Roles in Management of Cancer: A Critical Review. *Comprehensive Reviews in Food Science and Food Safety*, 14(2): 123-138.
- Sahoo, T.K., Jena, P.K., Nagar, N., Patel, A.K., and Seshadri, S. (2015). *In Vitro* Evaluation of Probiotic Properties of Lactic Acid Bacteria from the Gut of *Labeo rohita* and *Catla catla*. *Probiotics and antimicrobial proteins*, 1-11.
- Salminen, S., Gorbach, S., Lee, Y.K. and Benno, Y. (2004). Human studies on probiotics: What is scientifically proven today? *Lactic Acid Bacteria Microbiological and Functional Aspects*, New York: Marcel Dekker Inc.
- Salminen, S. and Isolauri, E. (2008). Opportunities for improving the health and nutrition of the human infant by probiotics. *Nestle Nutr Workshop Ser Pediatr Program* 62: 223-233
- Salminen, S., Wright, A.V., Morelli, L., Marteau, P., Brassart, D., de Vos, W.M., Fonden, R., Saxelin, M., Collins, K., Mogensen, G., Birkeland, S.-E. and Mattila-Sandholm, T. (1998). Demonstration of safety of probiotics. *International Journal of Food Microbiology* 44: 93-106.
- Santiago-Lopez, L., Hernandez-Mendoza, A., Garcia, H.S., Mata-Haro, V., Vallejo-Cordoba, B., and Gonzalez-Cordova, A.F. (2015). The effects of consuming probiotic-fermented milk on the immune system: A review of scientific evidence. *International Journal of Dairy Technology*.
- Sasco, A.J. (2001). Epidemiology of breast cancer: an environmental disease? *Acta Pathologica, Microbiologica et Immunologica Scandinavica* 109: 321-332.
- Sasco, A.J. (2008). Cancer and globalization. *Biomedicine and Pharmacotherapy* 62: 110-121.
- Schiffman, M., Castle, P.E., Jeronimo, J., Rodriguez, A.C. and Wacholder, S. (2007). Human papillomavirus and cervical cancerx. *Lancet* 370(9590): 890-907.

- Schillinger, U. and Lucke, F.K. (1987). Identification of lactobacilli from meat and meat products. *Food Microbiology* 4: 199-208.
- Schoug, A., Olsson, J., Carlfors, J., Schnurer, J. and Hakansson, S. (2006). Freeze-drying of *Lactobacillus coryniformis* S13- effects of sucrose concentration, cell density, and freezing rate on cell survival and thermophysical properties. *Cryobiology* 53: 119-127.
- Servin, A.L. (2004). Antagonistic activities of lactobacilli and *bifidobacteria* against microbial pathogens. *FEMS Microbiology Review* 28: 405-440.
- Sim, I., Koh, J.H., Kim, D.J., Gu, S.H., Park, A., and Lim, Y.H. (2015). *In vitro* assessment of the gastrointestinal tolerance and immunomodulatory function of *Bacillus methylotrophicus* isolated from a traditional Korean fermented soybean food. *Journal of applied microbiology*.
- Shah, N.P. (2000). Probiotic bacteria: Selective enumeration and survival in dairy foods. *Journal of Dairy Science* 83: 894-907.
- Shah, N.P. (2004). Probiotics and Prebiotics. *Agro-Food Industry Hi-Tech* 15: 13-16.
- Shah, N.P. (2007). Functional cultures and health benefits. *International Dairy Journal* 17(11): 1261-1275.
- Sharma, A.K., Mohan, P. and Nayak, B.B. (2005). Probiotics: Making a comeback. *Indian Journal of pharmacology* 37(6): 358-385.
- Shi, L.E., Li, Z.H., Li, D.T., Xu, M., Chen, H.Y., Zhang, Z.L. and Tang, Z.X. (2013). Encapsulation of probiotic *Lactobacillus bulgaricus* in alginate-milk microspheres and evaluation of the survival in simulated gastrointestinal conditions. *Journal of Food Engineering* 117: 99-104.
- Shi, L.E., Li, Z.H., Zhang, Z.L., Zhang, T.T., Yu, W.M., Zhou, M.L. and Tang, Z.X. (2013). Encapsulation of *Lactobacillus bulgaricus* in carrageenan-locust bean gum coated milk microspheres with double layer structure. *LWT - Food Science and Technology* 54: 147-151.
- Shi, Y., Chen, L., Tong, J. and Xu, C. (2009). Preliminary characterization of vaginal microbiota in healthy Chinese women using cultivation-independent methods. *Journal of Obstetrics and Gynaecology Research* 35(3): 525-532.
- Silvester, M.E. and Dicks, L.M. (2003). Identification of lactic acid bacteria isolated from human vaginal secretions. *Antonie Van Leeuwenhoek* 83: 117-123.
- Simon, O., Jadamus, A. and Vahjen, W. (2001). Probiotic feed additives-effectiveness and expected modes of action. *Journal of Animal Feed Science* 10: 51-67.
- Singh, B. (2007). Psyllium as therapeutic and drug delivery agent. *International Journal of Pharmaceutics* 334(1-2): 1-14.

- Singh, S. and Goswami, P. (2009). Application of molecular identification tools for *Lactobacillus*, with a focus on discrimination between closely related species. *LWT - Food Science and Technology* 42(2): 448-457.
- Smet, D., Hoorde, L.V., Woestyne, M.V., Christiaens, H. and Verstraete, V. (1995). Significance of bile salt hydrolytic activities of lactobacilli. *Journal of Applied Bacteriology* 79: 292-3011.
- Sohail, A., Turner, M.S., Coombes, A., Bostrom, T. and Bhandari, B. (2011). Survivability of probiotics encapsulated in alginate gel microbeads using a novel impinging aerosols method. *International Journal of Food Microbiology* 145(1): 162-168.
- Sohail, A., Turner, M.S., Coombes, A., Bostrom, T. and Bhandari, B. (2011). Survivability of probiotics encapsulated in alginate gel microbeads using a novel impinging aerosols method. *International Journal of Food Microbiology* 145: 162-168.
- Solieri, L., Bianchi, A. and Giudici, P. (2012). Inventory of non starter lactic acid bacteria from ripened Parmigiano Reggiano cheese as assessed by a culture dependent multiphasic approach. *Systematic and Applied Microbiology* 35: 270-277.
- Song, Y.L., Kato, N., Matsumiya, Y., Liu, C.X., Kato, H. and Watanabe, K. (1999). Identification of and hydrogen peroxide production by fecal and vaginal lactobacilli isolated from Japanese women and newborn infants. *Journal of Clinical Microbiology* 37: 3062-3064.
- Steidler, L. (2002). In situ delivery of cytokines by genetically engineered *Lactococcus lactis*. *International Journal of General and Molecular Microbiology* 82: 323-331.
- Steyn, P.L. and Holzaphel, W.H. (1991). Identity and antibiotic sensitivity of human vaginal lactic acid bacteria from the population of northern Namibia. *S-Afr. Tydskr. Wet.* 87: 68-69.
- Stiles, M.A. and Holzaphel, W.H. (1997). Lactic acid bacteria of food and their current taxonomy. *International Journal Food Microbiology* 36: 1-29.
- Sultana, K., Godward, G., Reynold, N., Arumugaswamy, R., Peiris, P. and Kailasapathy, K. (2000a). Encapsulation of probiotic bacteria with alginate-starch and evaluation of survival in simulated gastrointestinal conditions and in yogurt. *International Journal of Food Microbiology* 62: 47-55.
- Sultana, K., Godward, G., Reynolds, N., Arumugaswamy, R., Peiris, P. and Kailasapathy, K. (2000b). Encapsulation of probiotic bacteria with alginate-starch and evaluation of survival in simulated gastrointestinal conditions and in yoghurt. *International Journal of Food Microbiology* 62: 47-55.
- Sultana, K., Godward, G., Reynolds, N., Arumugaswamy, R., Peiris, P. and Kailasapathy, K. (2000c). Encapsulation of probiotic bacteria with alginate-



- starch and evaluation of survival in simulated gastrointestinal conditions and in yoghurt. *International Journal of Food Microbiology* 62(1-2): 47-55.
- Svec, P., Pantucek, R., Petrás, P., Sedlacek, I. and Novakova, D. (2010). Identification of *Staphylococcus spp.* using (GTG)5-PCR fingerprinting. *Systematic and applied microbiology* 33: 451-456.
- Svec, P., Vancanneyt, M., Seman, M., Snauwaert, C., Lefebvre, K., Sedlacek, I. and Swings, J. (2005). Evaluation of (GTG)5-PCR for identification of *Enterococcus spp.* *FEMS Microbiology Letters* 247: 59-63.
- Swenson, J.M., Facklamand, R.R. and Thornsberry, C. (1990). Antimicrobial susceptibility of vancomycin-resistant *Leuconostoc*, *Pediococcus*, and *Lactobacillus* species. *Antimicrobial Agents and Chemotherapy* 34: 543-549.
- Tambekar, D.H. and Bhutada, S.A. (2010). An evaluation of probiotic potential of *Lactobacillus sp.* from milk of domestic animals and commercial available probiotic preparations in prevention of enteric bacteria infections. *Recent Research in Science and Technology* 2(10): 82-88.
- Tannock, G.W., Dashkevicz, M.P. and Feighner, S.D. (1989). Lactobacilli and bile salt hydrolase in the murine intestinal tract. *Appl. Environ. Microbiol.* 55: 1848-1851.
- Tejero-Sarinena, S., Barlow, J., Costabile, A., Gibson, G.R. and Rowland, I. (2012). In vitro evaluation of the antimicrobial activity of a range of probiotics against pathogens: Evidence for the effects of organic acids. *Anaerobe* 18(5): 530-538. doi: <http://dx.doi.org/10.1016/j.anaerobe.2012.08.004>.
- Temmerman, R., Huys, G. and Swings, J. (2004). Identification of lactic acid bacteria: culture-dependent and culture-independent methods. *Trends in Food Science & Technology* 15: 348-359. doi: 10.1016/j.tifs.2003.12.007.
- Thakker, S.P., Rokhade, A.P., Abbigerimath, S.S., Iliger, S.R., Kulkarni, V.H., More, U. A., and Aminabhavi, T.M. (2014). Inter-polymer complex microspheres of chitosan and cellulose acetate phthalate for oral delivery of 5-fluorouracil. *Polymer bulletin* 71(8): 2113-2131.
- Thokchom, S. and Joshi, S.R. (2012). Probiotic and Bacteriocin Efficacy of lactic Acid Bacteria from Traditionally Fermented Foods: A Review. *Assam University Journal of Science & Technology* 10: 142-155.
- Todorov, S., Botes, M., Danova, S. and Dicks, L. (2007). Probiotic properties of *Lactococcus lactis ssp. lactis* HV219, isolated from human vaginal secretions. *Journal of Applied Microbiology* 103: 629-639.
- Tokudome, S., Suzuki, S. and Ichikawa, H. (2004). Condom use promotes regression of cervical intraepithelial neoplasia and clearance of human papillomavirus: a randomized clinical trial. *International Journal Cancer* 112(1): 164.

- Troisi, R., Hatch, E.E. and Titus-Ernstoff, L. (2007). Cancer risk in women prenatally exposed to diethylstilbestrol. *International Journal Cancer* 121(2): 356-360.
- Truelstrup-Hansen, L., Allan-wojtas, P.M., Jin, Y.L. and Paulson, A.T. (2002). Survival of free and calcium-alginate microencapsulated Bifidobacterium spp. in stimulated gastro-intestinal conditions. *Food Microbiology* 19: 35-45.
- Turnbaugh, P., Ley, R., Hamady, M., Fraser-Liggett, C., Knight, R. and Gordon, J. (2007). The Human Microbiome Project. *Nature* 449: 804-810.
- Uccello, M. (2012). Potential role of probiotics on colorectal cancer prevention. *BMC Surg* 12(1): S35.
- Van den Nieuwboer, M., Brummer, R.J., Guarner, F., Morelli, L., Cabana, M., and Claasen, E. (2015). The administration of probiotics and synbiotics in immune compromised adults: is it safe?. *Beneficial microbes*, 6(1): 3-17.
- Vermes, I., Haanen, C., Steffens-Nakken, H. and Reutelingsperger, C. (1995). A novel assay for apoptosis Flow cytometric detection of phosphatidylserine early apoptotic cells using fluorescein labelled expression on Annexin V. *Journal of Immunological Methods* 184: 39-51.
- Vintini, E., Ocana, V. and Elena Nader-Macias, M. (2004). Effect of lactobacilli administration in the vaginal tract of mice: evaluation of side effects and local immune response by local administration of selected strains. *Methods Mol Biol* 268: 401-410.
- Vivek, K.B. (2013). Use of encapsulation probiotics in dairy based foods. *International Journal of Food, Agriculture and Veterinary Sciences* 3(1): 188-199.
- Voo, W., Ravindra, P., Tey, B. and Chan, E. (2011). Comparison of alginate and pectin based beads for production of poultry probiotic cells. *Journal of Bioscience and Bioengineering* 111: 294-299.
- Vos, P., Faas, M.M., Spasojevic, M. and Sikkema, J. (2010). Encapsulation for preservation of functionality and targeted delivery of bioactive food components. *International Dairy Journal* 20: 292-302.
- Wanga, C.L., Ng, T.B., Yuan, F., Liu, Z.K. and Liu, F. (2007). Induction of apoptosis in human leukemia K562 cells by cyclic lipopeptide from Bacillus subtilis natto T-2. *Peptides* 28: 1344-1350.
- Wilks, M., Wiggins, R., Whiley, A., Hennessy, E., Warwick, S., Porter, H., Corfield, A. and Millar, M. (2004). Identification and H<sub>2</sub>O<sub>2</sub> production of vaginal Lactobacilli from pregnant women at high risk of preterm birth and relation with outcome. *J. Clin. Microbiol.* 42: 713-717.
- Woodford, N., Johnson, A.P., Morrison, D. and Speller, D.C.E. (1995). Current perspective on glycopeptide resistance. *Clinical Microbiology Reviews* 8: 585-615.

- Yaghoobi, M. (2004). Hereditary risk factors for the development of gastric cancer in younger patients. *BMC Gastroenterology* 4: 28.
- Yang, X., Hang, X., Zhang, M., Liu, X., and Yang, H. (2015). Relationship between acid tolerance and cell membrane in *Bifidobacterium*, revealed by comparative analysis of acid-resistant derivatives and their parental strains grown in medium with and without Tween 80. *Applied microbiology and biotechnology*, 1-10.
- Yarbro, C., Frogga, M. and Goodman, M. (2005). *Cancer nursing: Principles and practice. Jones and Bartlett Publishers, Boston, MA.*
- Zeeb, B., Saberi, A.H., Weiss, J., and McClements, D.J. (2015). Retention and release of oil-in-water emulsions from filled hydrogel beads composed of calcium alginate: impact of emulsifier type and pH. *Soft matter* 11(11): 2228-2236.
- Zhang, Y., Zhang, L., Ma, W., Yi, H.X., Yang, X., Du, M., Shan, Y.J., Han, X. and Zhang, L.L. (2012). Screening of probiotic lactobacilli for inhibition of *Shigella sonnei* and the macromolecules involved in inhibition. *Anaerobe*: 1-6. doi: 10.1016/j.anaerobe.2012.08.007
- Zhou, J.S., Pillidge, C.J., Gopal, P.K. and Gill, H.S. (2005). Antibiotic susceptibility profiles of new probiotic *Lactobacillus* and *Bifidobacterium* strains. *International Journal of Food Microbiology* 98: 211-217. doi: 10.1016/j.ijfoodmicro.2004.05.011
- Zhou, X., Brown, C.J. and Abdo, Z. (2007). Differences in the composition of vaginal microbial communities found in healthy Caucasian and black women. *Isme J* 1: 121-133.
- Zuidam, N.J. and Shimoni, E. (2009). Overview of Microencapsulates for Use in Food Products or Processes and Methods to Make Them. In: Zuidam, N.J., Nedovic, V.A., (Eds.). *Encapsulation Technologies for Food Active Ingredients and Food Processing*, Springer: Dordrecht, The Netherlands; 2009, p. 3-31.

## LIST OF PUBLICATIONS

- Nami, Y., Abdullah, N., Haghshenas, B., Radiah, D., Rosli, R., and Yari Khosroushahi, A., (2014). Assessment of probiotic potential and anticancer activity of newly isolated vaginal bacterium *Lactobacillus plantarum* 5BL. ***Microbiology and Immunology***, 58(9): 492-502.
- Nami, Y., Abdullah, N., Haghshenas, B., Radiah, D., Rosli, R., and Yari Khosroushahi, A., (2014). Probiotic potential and biotherapeutic effects of newly isolated vaginal *Lactobacillus acidophilus* 36YL strain on cancer cells. ***Anaerobe***, 28: 29-36.
- Nami, Y., Abdullah, N., Haghshenas, B., Radiah, D., Rosli, R., and Yari Khosroushahi, A., (2014). Probiotic assessment of *Enterococcus durans* 6HL and *Lactococcus lactis* 2HL isolated from vaginal microflora. ***Journal of Medical Microbiology***, 63(8): 1044-1051.
- Nami, Y., Abdullah, N., Haghshenas, B., Radiah, D., Rosli, R., and Yari Khosroushahi, A., (2014). A newly isolated probiotic *Enterococcus faecalis* strain from vagina microflora enhances apoptosis of human cancer cells. ***Journal of Applied Microbiology***, 117 (2): 498-508.
- Nami, Y., Haghshenas, B., Abdullah, N., Barzegari, A., Radiah, D., Rosli, R., and Yari Khosroushahi, A., (2015). Probiotics or antibiotics: future challenges in medicine. ***Journal of Medical Microbiology***, 64: 137-146.



**UNIVERSITI PUTRA MALAYSIA**

**STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT**

**ACADEMIC SESSION :** \_\_\_\_\_

**TITLE OF THESIS / PROJECT REPORT :**

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

**NAME OF STUDENT :** \_\_\_\_\_

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

\*Please tick (✓)

**CONFIDENTIAL**

(Contain confidential information under Official Secret Act 1972).

**RESTRICTED**

(Contains restricted information as specified by the organization/institution where research was done).

**OPEN ACCESS**

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

**PATENT**

Embargo from \_\_\_\_\_ until \_\_\_\_\_  
(date) (date)

**Approved by:**

\_\_\_\_\_  
(Signature of Student)  
New IC No/ Passport No.:

\_\_\_\_\_  
(Signature of Chairman of Supervisory Committee)  
Name:

Date :

Date :

[Note : If the thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted. ]