UNIVERSITI PUTRA MALAYSIA

EFFECTS OF EMPTY FRUIT BUNCH BIOCHAR AND NITROGEN-FIXING BACTERIA ON SOIL QUALITY AND GROWTH OF SWEET CORN

DIYAR KAREEM ABDULRAHMAN

FP 2015 64

EFFECTS OF EMPTY FRUIT BUNCH BIOCHAR AND NITROGEN-FIXING BACTERIA ON SOIL QUALITY AND GROWTH OF SWEET CORN

By

DIYAR KAREEM ABDULRAHMAN

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Master of Science.

April 2015

.

COPYRIGHT

DEDICATION

This research is dedicated to all the scientists, researchers and environmentalists who are soldiering to keep the sanctity of our Earth's atmosphere preserved. For my great supervisor who cared so much about my work and who intelligence, support and guidance played a vital role in completing this study. For my beloved father (Kareem Abdulrahman Qadir) and mother (Amira Ali Raza) whom I wish could be here to see me become Master of Science, degree. To my dearest and beloved wife Ngin Aziz Mohammed and finally, to my beloved country Kurdistan- Iraq.

Abstract of Thesis Presented to the Senate of Universiti Putra Malaysia in Fulfillment of the Requirement for the Degree of Master of Science.

EFFECTS OF EMPTY FRUIT BUNCH BIOCHAR AND NITROGEN-FIXING BACTERIA ON SOIL QUALITY AND GROWTH OF SWEET CORN

By

DIYAR KAREEM ABDULRAHMAN

April 2015

Chairman : Associate Professor Radziah Othman, PhD

Faculty : Agriculture

There has been an increasing interest in biochar research as soil amendments to improve soil properties leading to improve most of plant growth in a sustainable agriculture. Tropical soil is generally unfertile unit, low organic matter, plant nutrients, acidic and low microorganisms that affect crop production. Addition of biochar such as empty fruit bunch (EFB) could improve the soil fertility. Laboratory and glasshouse studies were conducted with the following objectives, i) to determine the effect of soil sterilization, EFB biochar and N_2-fixing bacteria *Stenotrophomonas sp.* (Sb16) on indigenous soil microbial population, enzyme activity and soil chemical properties, and ii) to determine the effect of EFB biochar and N_2-fixing bacteria Sb16 on growth and nutrient uptake of sweet corn. Five rates of EFB biochar (0, 0.25, 0.5, 0.75 and 1%) were applied to sterilized and non-sterilized soil either with or without N_2-fixing bacteria Sb16 and incubated for 40 days under laboratory condition. The factorial study was arranged in a Complete Randomized Design (CRD) with 3 replications. Microbial population, enzyme activity and chemical properties of soil were determined at the end of incubation. Sweet corn was grown in pots containing 6 kg soil and applied with the 5 levels of EFB biochar (0, 5, 10, 15 and 20 t/ha) either with or without bacteria Sb16. Plants were sampled at tasseling stage and analyzed for dry biomass, chlorophyll content and nutrient uptake. Soil samples were analyzed for soil biochemical and microbiological properties. Results of laboratory study showed that application of EFB biochar at 0.5% without inoculation and 0.25% with N_2-fixing bacteria Sb16 in both soils significantly increased population of soil bacteria, fungi, actinomycetes and N_2-fixing bacteria (NFB), enzymes (urease, acid phosphatase and fluorescein diacetate hydrolysis (FDA) activity), and soil chemical properties (pH, organic C, total N, available P and exchangeable K, Ca and Mg. A glasshouse experiment showed that application of EFB biochar at 5 t/ha and N_2-fixing bacteria Sb16 significantly ($P<0.05$) improved growth of corn (shoot and root biomass, root length, root volume, plant height and leaf chlorophyll content, nutrient uptake), soil microbial populations, FDA and selected soil chemical properties. The EFB biochar at 5 t/ha and N_2-fixing bacteria Sb16 stimulated soil quality and plant growth. Addition of high EFB rates (15 and 20

t/ha) to soil negatively affected all the observed parameters. Addition of EFB biochar to soil with N_2-fixing bacteria may be an alternative solution in improving nutrients, enzymes and diversity of microorganisms in soil and thus led to improve plant growth. The studies showed that application of EFB biochar at 5 t/ha or 0.25% with N_2-fixing bacteria Sb16 and 10 t/ha or 0.5% without bacteria inoculated improved corn growth and the quality of soil for sustainable corn production.

Abstrak tesis dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains.

KESAN TANDAN KOSONG BIOCAR (EFB) DAN BAKTERI A PENGIKAT NITROGEN TERHADAP KUALITI TANAH DAN PERTUMBUHAN JAGUNG MANIS.

Oleh

DIYAR KAREEM ABDULRAHMAN

April 2015

Pengerusi : Profesor Madya Radziah Othman, PhD

Fakulti : Pertanian

Terdapat peningkatan terhadap kajian biocar sebagai pembaik-pulih tanah untuk mengingkatkan ciri-ciri tanah ke arah meningkatkan pertumbuhan tanaman lestari. Ciri biokimia tanah seperti aktiviti enzim mempunyai pernanan utama dalam kitaran nutrien dan boleh dianggap sebagai pengukur kualiti tanah. Tanah tropika mempunyai kandungan yang rendah dalam bahan organik, nutrient tumbuhan, kebolehan pertukaran kation (CEC), pH, larut lesap yang tinggi, dan kekurangan mikroorganisma hidup yang memberi kesan terhadap hasil tanaman. Kajian makmal dan rumah kaca telah dijalankan berdasarkan objektif berikut, i) Untuk mengenalpasti kesan tanah steril, tandan kosong biocar (EFB) dan bakteria pengikat nitrogen *Stenotrophomonas sp.* Sb16 terhadap populasi asal mikroorganisma tanah, aktiviti enzim dan ciri-ciri kimia tanah, dan ii) Untuk mengenalpasti kesan tandan kosong biocar (EFB) dan bakteria pengikat nitrogen Sb16 terhadap pertumbuhan dan penyerapan nutrient oleh jagung manis. Lima kadar tandan kosong kelapa sawit biocar (EFB) (0, 0.25, 0.5, 0.75 and 1 %) telah digunakan untuk yang steril dan tidak steril dengan samaada mengandungi atau tidak mengandungi bakteria pengikat nitrogen Sb16 dan inkubasi selama 40 hari di bawah keadaan makmal. Kajian faktorial adalan mengikut rekabentuk rawak lengkap (CRD) dengan tiga replikasi. Populasi mikrob, aktiviti enzim dan ciri-ciri kimia telah dikenalpasti diakhir proses inkubasi. Jagung manis telah ditanam dalam 6 kg tanah/pasu dan ditambah dengan lima kadar tandan kosong biocar (EFB) (0, 5, 10, 15, dan 20 t/ha) samaada mengandungi atau tidak mengandungi bakteria Sb16. Tanaman dan tanah telah diambil sampel pada peringkat teasseling dan telah dianalisis untuk berat kering biojisim, pengambilan nutrien, biokimia tanah dan ciri-ciri mikrobiologi. Hasil kajian makmal menunjukkan penggunaan tandan kosong biocar (EFB) pada 0.5% tanpa inokulasi dan 0.25% dengan bakteria pengikat nitrogen Sb16 dalam kedua dua tanah dengan nyata menginggikan populasi bakteria tanah, kulat, aktinomiset, dan bakteria pengikat nitrogen (NFB), enzim (urea, asid fosfotas, aktiviti hidrolisis fluoresin diasetat (FDA) aktiviti, dan ciri- ciri kimia tanah (pH, C organic, jumlah N, P tersedia dan pertukaran K, Ca dan Mg). Kajian rumah kaca menunjukkan bahawa penggunaan biocar EFB pada 5 t/ha dan bakteria pengikat Sb16 dengan nyata (P

<0.05) meningkatkan pertumbuhan jagung dengan biojisim pucuk dan akar, panjang akar, jumlah akar, tinggi pokok dan kandungan klorofil daun), pengambilan nutrient, populasi mikrob tanah, FDA dan ciri-ciri kimia tanah yang dipilih. Biocar EFB pada 5 t/ha dan bakteria pengikat nitrogen Sb16 merangsang kualiti tanah dan pertumbuhan pokok. Penambahan kadar biocar EFB yang tinggi (15 dan 20 t/ha) kepada tanah tidak memberi kesan kepada semua parameter yang diperhatikan. Penambahan biocar EFB kepada tanah dan bakteria pengikat nitrogen mungkin menjadi penyelesaian alternatif dalam meningkatkan nutrien, enzim dan kepelbagaian mikroorganisma tanah dan seterusnya menjadi tunjang untuk meningkatkan pertumbuhan tanaman. Kajian menunjukkan penggunaan biocar EFB pada 10 t/ha or 0.5% tanpa bakteria Sb16 dan kadar yang rendah 5 t/ha or 0.25% dengan bakteria pengikat nitrogen menigkatkan pertumbuhan jagung dan kualiti tanah untuk penghasilan jagung yang mampan.

ACKNOWLEDGEMENTS

Alhamdullilah

First of all, my deepest gratitude goes to the most merciful almighty Allah for giving me good health, strength and perseverance. It has given me a lot of pleasure and pride to work under the supervision of Assoc. Prof Dr. Radziah Othman. Her continuous support and encouragement over the past two years and her intelligence, support and guidance played a vital role in completing this study. I am greatly indebted for her constructive criticism and patience. I thank her for her excellent teaching as well as for being a mother figure to me. I am grateful to her for giving me this opportunity. I am indebted to Assoc. Prof Dr. Halimi b Mohd Saud for his continued technical assistance and guidance in every part of this work. I appreciate the help of Prof Dr. Rosenani bt Abu Bakar for introducing the present topic and for her inspiring guidance, constructive criticism and valuable suggestion throughout this project work. My special thanks go to Prof. Dr. Zulkifli Shamsuddin who was always available to provide valuable ideas, valuable suggestion, editing, input and extremely supportive and for being the wind beneath my wings. His immense help and guidance has been instrumental in accelerating the progress of my research. I extend my gratitude to all my friends who helped made my stay comfortable and gave me a good moral support. I extend my gratitude to Mr. Zul for his advice and assistance, friendly attitude during the study and other staff in the Department of Land Management and Department of Crop Science for their extensive cooperation and friendship. I acknowledge the help and support provided by our department graduate secretaries in making all the paper work timely and efficiently. My heartiest thanks to my government Kurdistan Regional Government and my amazing family for their unwavering encouragement and the unconditional support offered by them both morally and financially.Their love and blessings are reflected in all my endeavors. Last but not the least, my sincere thanks to all my friends who have patiently extended all help for accomplishing this work.

I certify that a Thesis Examination Committee has met on 21 April 2015 to conduct the final examination of Diyar Kareem Abdulrahman on his thesis entitled "Effects of Empty Fruit Bunch Biochar and Nitrogen-Fixing Bacteria on Soil Quality and Growth of Sweet Corn" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Shamshuddin bin Jusop, PhD
Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Samsuri bin Abd. Wahid, PhD
Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Datin Ainon Hamzah, PhD
Professor
Universiti Kebangsaan Malaysia
Malaysia
(External Examiner)

ZULKARNAIN ZAINAL, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 17 June 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Radziah Othman, PhD
Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Halimi b Mohd Saud, PhD
Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Rosenani bt Abu Bakar, PhD
Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD
Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date :

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work
- quotations, illustrations and citations have been duly referenced
- the thesis has not been submitted previously or comcurrently for any other degree at any institutions
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be owned from supervisor and deputy vice –chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No: Diyar Kareem Abdulrahman

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman of Supervisory Committee: Prof. Madya Dr. Radziah Othman
Ketua
Jabatan Pengurusan Tanah
Fakulti Pertanian
Universiti Putra Malaysia

Signature: _____
Name of Member of Supervisory Committee: DR HALIMI MOHD SAUD

PROF MADYA DR. HALIMI MOHD SAUD
Timbalan Dekan (Pembangunan Jaringan Industri dan Masyarakat)
Fakulti Pertanian
Universiti Putra Malaysia
43400 UPM Serdang, Selangor

Signature: _____
Name of Member of Supervisory Committee: PROF. DR. ROSENANI ABU BAKAR
Professor
Jabatan Pengurusan Tanah
Fakulti Pertanian
Universiti Putra Malaysia
43400 Serdang, Malaysia

TABLE OF CONTENTS

LIST OF TABLES

LIST OF FIGURES

LIST OF PLATES

LIST OF ABBREVIATIONS

A.A	Actinobacteria Isolation Agar
AAS	Atomic Absorption Spectrophotometer
ANOVA	Analysis of Variance
BET	Brunauer, Emmett and Teller
BNF	Biological Nitrogen Fixation
CEC	Cation Exchange Capacity
CFU	Colony Forming Unit
C:N	Carbon to Nitrogen Ratio
CRBD	Completely Randomized Block Design
DNA	Deoxy Ribonucleic Acid
EFB	Empty Fruit Bunch
FDA	Fluorescein Diacetate Hydrolysis
GLM	General Linear Model
MUB	Modified Universal Buffer
NA	Nutrient Agar
NFB	N_2- Fixing Bacteria
NFB	N- Free Broth
OD	Optical Density
OM	Organic Matter
PAWC	Plant Available Water Content
P-NPP	Aldisodium P- nitrophenyl Phosphate Tetrahydrate
RBSA	Rose Bengal Isolation Agar
SAS	Statistical Analysis System
SDW	Sterile Distilled Water
SOC	Soil Organic Carbon
THAM BUFFER	Tris (hydroxyl methyl) Amino Methan
TSP	Triple Super Phospha

CHAPTER 1

INTRODUCTION

Tropical soils are mostly infertile soil that affect soil quality and plant growth. The main characteristics of tropical soils are acidic soils and have low fertility (Shamshuddin and Fauziah, 2010). Aluminium (Al) and manganese (Mn) is toxic elements to plants, while Ca and Mg often are deficient in acidic soil. In addition, this soil has low pH, cation exchange capacity (CEC), poor water holding capacity, low microbial population, poor soil enzyme activity and high leaching of basic cations that are essential for soil properties and plant growth (Aini and Vimala, 2002). The amount of N present in acidic soil is extremely small to provide the productivity of agricultural systems and responsible for the decomposition of organic matter. The beneficial soil free living N_2 fixing bacteria that supply some nutrients in a form that plants can absorb easily from the soil and improve plant growth can die of nutrient deficiency in tropical soil (Topoliantz, 2005).

In tropical soil, the size of most crops is restricted and yield is low. Plants suffering from element toxicity and poor nutrient will show a restricting root system, which may affect the capacity for mineral nutrient acquisition and resulting in reducing plant growth (Rout et al., 2001). These acidic soils are constraints to crop production. Optimum growth of sweet corn occurs when essential nutrients are available, but growth will not satisfy when sweet corn requires a continuous supply of chemical properties in soil (Fageria and Baliger, 2008). In Malaysia, the production of corn is unsatisfactory and its imports for a amounting of feed to millions of Ringgit (USDA, 2014). To meet these requirements, yield corn production should be increased by way of using proper fertilizer and other inputs.

However, this soil infertility can be ameliorated effectively by applying liming material, organic matter, bio-fertilizer, biochar and other amendments (Topoliantz, 2005). In order to control infertile soil, soil microbial activity and soil pH need to be substantially improved. Application of oil palm waste into soil can be an alternative to improve soil fertility. Empty fruit bunch (EFB) has been turned to EFB biochar roughly 20 tonnes/day by a Nasmeck company in Malaysia (USDA, 2014). EFB biochar is produced from the burning of oil palm EFB at high temperatures under low oxygen conditions with the intention to sequester carbon, mitigate CO_2 emission and at the same time to be used as a soil amendment (Norazlina et al., 2014).

A few researches have been conducted to produce biochar from oil palm EFB and used as soil amendments for its role as a nutrient source for crop production (Nair and Lawson, 2012). Sahara and Lim (2000) showed that application of biochar increased soil mineral N, pH, exchangeable K, Ca, Mg, soil moisture and organic matter leading to improved plant growth. Biochar has been known to affect the soil chemical, biological and enzyme activities and plant growth (Bailey et al., 2010; Lehmann et al., 2011). The ameliorating effect of biochars on acidic soil was assumed to be consistent with their composition and properties which depend on the biomass feedstock type and pyrolytic conditions.

Addition of biochar with inoculation of beneficial microbes can be important for improving soil fertility and plant physiological activity through the transfer of nutrients into forms that can be uptake easily from the soil (Arnoldus et al., 2011; Jin, 2010; Blackwell et al., 2009). Inoculation of rice plant with *Stenotrophomonas sp* strain (Sb16), a free living N_2 fixing bacteria has improved rice growth isolated from rice crop (*Oryza sativa* L.) (Radziah et al., 2013). In Malaysian soil, Studies showed that the inoculation significantly increased plant growth, yield and nutrient uptake of rice and can supplement about 40% of total N to the rice (Naher et al., 2011). Total world biological nitrogen fixation is estimated at 139-172 million t/year, which is three times higher than industrially fixed nitrogen and contributes significantly to biological nitrogen fixation in agriculture and natural N cycle (Ishizuka, 1992).

Previous studies showed that biochar addition, affected bacterial population due to its effect on soil pH, temperature, nutrients, water, oxygen, and metabolic compounds (Jeffery et al., 2011). Infertile soil can be ameliorated effectively by applying liming material, organic matter and beneficial microorganisms. Little information is available on the potential of biochars with free living N_2-fixing bacteria and their interactions to reduce soil acidity. Studies suggest that crop yields and soil field capacity can increase as a result of applying biochar as an amendment to the soil (Christopher et al., 2012). Insufficient information is available on the application of EFB biochar with N_2-fixing bacteria on plant growth and soil quality. The following studies were conducted with the following objectives;

1- To determine the effect of soil sterilization, EFB biochar and N_2-fixing bacteria *Stenotrophomonas sp*. (Sb16) on indigenous soil microbial population, enzyme activity and soil chemical properties.

2- To determine the effect of EFB biochar and N_2-fixing bacteria on growth and nutrient uptake of sweet corn.

REFERENCES

Abebe, N., Endalkachew, K., Mastawesha, M., & Gebermedihin, A. (2012) Effect of biochar application on soil properties and nutrient uptake of lettuces(Lactuca sativa) grown in chromium polluted soils. American-Eurasian Journal of Agricultural & Environmental Sciences, 12: 369-376.

Afeng, Z., Rongjun, B., Genxing, P., Liqiang, C., Qaiser, H., Lianqing, L., Jinwei,Z., Jufeng, Z., Xuhui, Z., Xiaojun, H., & Xinyan, Yua. (2012) Effects of biochar amendment on soil quality, crop yield and greenhouse gas emission in a Chinese rice paddy: A field study of 2 consecutive rice growing cycles. Field Crops Research An International Journal, 127:153–160.

Aini, Z., Sivaprasagam, A., Vimala, P., & Mohammad, R. (2005) Organic vegetablecultivation in Malaysia: 1-70. MARDI (Malaysian Agricultural Research and Development Institute, Malaysia.

Aini, Z., & Vimala, P. (2002) Research of Organic Crop Production in Malaysia. Paper presented at workshop report of FAO Expert Group Workshop on the Preparation of Technical Guidelines on the Organic Cultivationof Tropical and Subtropical Fruits, 22-26, Kuala Lumpur.

Alan, L., Wright, L., Provin., Frank, M., Hons., David, A., Zuberer., & Richard, H. W. (2007) Compost source and rate effects on soil macronutrientavailability under saint augustine grass and bermuda grass turf. Compost Science & Utilization, 15: 22-32.

Albiach, R., Canet, R., Pomares, F., & Ingelmo, F. (2000) Microbial biomasscontent and enzymatic activities after the application of organic amendments to a horticultural soil. Bioresource Technology, 75: 43-48.

Amonette, J., & Joseph, S. (2009) Characteristics of biochar: Microchemical properties." Chapter 3 in Biochar for Environmental Management: Science and Technology, ed. J Lehmann & S Joseph. 33-52. Earthscan, London,United Kingdom.

Anderson, C.R., Condron, L.M., Clough, T.J., Fiers, M., Stewart, A., Hill, R.A.,&Sherlock, R.R. (2011) Biochar induced soil microbial community change: Implications for biogeochemical cycling of carbon, nitrogen and phosphorus. Pedobiologia, 54:309-320.

Antonious, G. F. (2003) Impact of Soil Management and two Botanical Insecticides on Urease and Invertase Activity. Journal of Environmental Science and Health, Part B: Pesticides, Food Contaminants, and Agricultural Wastes,38: 479-488.

Aon, M.A., & Colaneri, A.C. (2001) Temporal and Spatial Evolution of Enzymatic Activities and Physico-Chemical Properties in an Agricultural soil. Applied Soil Ecology, 18: 255–270.

Arnoldus, K. B., Nguyen, H., & Amjad, A. (2011) Beneficial use of biochar to correct soil acidity. www.ctahr.hawaii.edu.

Asai, H., Samson, B.K., Stephan, H.M., Songyikhangsuthor, K., Homma, K., Kiyono, Y., Inoue,Y., Shiraiwa,T., & Horie, T. (2009) Biochar amendment techniques for upland rice production in Northern Laos 1. Soil physicalproperties, leaf SPAD and grain yield. Field Crops Research, 111: 81–84

Atkinson, C. J., Fitzgerald, J. D., & Hipps, N.A. (2010) Potential mechanisms for achieving agricultural benefits from biochar application to temperate soils: A review. Plant and Soil, 337: 1-18.

Bailey, V.L., Fansler, S.J., Smith, J.L., & Bolton, Jr., H. (2010) Reconciling apparent variability in effects of biochar amendment on soil enzyme activities by assay optimization. Soil Biology and Biochemistry, 43 (9): 296-301.

Baldock, J.A., & Smernik, R.J. (2002) Chemical composition and bio-availability ofthermally altered Pinus resinosa (Red pine) wood. Organic Geochemistry, 33: 1093-1109.

Basso, Andres Santiago. (2012) Effect of fast pyrolysis biochar on physical andchemical properties of a sandy soil. Graduate Theses and Dissertations. Paper 12586.

Biederman, L.A., & Harpole, W.S. (2013) Biochar and its effects on plant productivity and nutrient cycling: a meta-analysis. Global Change Biology–Bioenergy, 5: 202–214.

Birk, J. J., Steiner, C., Teixiera, W. C., Zech, W., & Glaser, B. (2009) Microbialresponse to charcoal amendments and fertilization of a highly weathered tropical soil and Amazonian Dark Earths in Central Amazonia – Preliminary Results. Institute of Soil Science, 46: 309-324.

Blackwell, P., Riethmuller, G & Collins, M. (2009) Biochar Application to Soil. In:Lehmann, J. & Joseph, S. (eds.) Biochar for Environmental Management, Science & Technology, 207-226.

Boguslaw, U., Mateusz, L., & Jerzy, L. (2014) Thermal properties of soils: effect of biochar application. Geophysical Research Abstracts, 16: 9533.

Bolan, N. S., Kunhikrishnan, A., Choppala, G. K., Thangarajan, R., & Chung, J. W. (2012) Stabilization of carbon in composts and biochars in relation to carbon sequestration and soil fertility. Science of the Total Environment, 424: 264-270.

Brady, N.C & Weil, R.R. (2004). Elements of the Nature and Properties of Soils, 2nd ed. Pearson Prentice Hall: Upper Saddle River, NJ: 111-112.

Bray, R.H., & Kurtz, L.T. (1945) Determination of total organic, and available forms of phosphorus in soils. Soil Science, 59: 39-45.

Bremner, J. M., & Edwards, AP. (1965) Determination and isotoperatio analysis of different forms of nitrogen in soils. Apparatus and procedure for distillation and determination of ammonium. Soil Science Society of America, 29: 504–507.

Bremner, J.M., & Mulvaney, C.S. (1982) Nitrogen-Total N. In Methods of Soil nalysis. Part 2. Chemical and Microbiological Properties Agronomy 9. Arctic, Antarctic, and Alpine Research, 42:139-151.

Brockhoff, S. R., Christians, N. E., Killorn, R. J., Horton, R., & Davis, D. (2010) Physical and Mineral-Nutrition Properties of Sand-Based Turfgrass Root Zones Amended with Biochar. Agronomy Journal, 102: 1627-1631.

Brown, T. R., Wright, M. M., & Brown, R. C. (2011) Estimating profitability of two biochar production scenarios: Slow pyrolysis vs. fast pyrolysis. iofuels, Bioproducts and Biorefining 5: 54–68.

Brussaard, L., de Ruiter, P.C. & Brown, G.G. (2007) Soil biodiversity for agricultural sustainability. Agriculture, Ecosystems and Environment, 121: 233-244.

Bruun, E.S., Ambus, P., Egsgaard, H & Huggaard-N, H. (2012) Effects of slow and fast pyrolysis biochar on soil C and N dynamics. Soil Biology and Biochemistry, 46: 73–79.

Burgmann, H., F. Widmer, W., Von Sigler, & J. Zeyer. (2004) New molecular creening tools for analysis of free-living diazotrophs in soil. Applied Environmental Microbiology, 70: 240-247.

Burris, R.H. (1991) Nitrogenases. The Journal of Biological Chemistry, 226: 9339-9342.

Caires, E.F., Corrêa, J.C.L., Churka, S., Barth, G. & Garbuio, F. (2006) Surfaceapplication of lime ameliorates subsoil acidity and improves root growth and yield of wheat in an acid soil under no-till system. Scientia Agricola, 63: 502–509.

Caliman, J.P., Martha, B., & Salètes, S. (2001) Dynamics of nutrient release from empty fruit bunches in field conditions and soil characteristic changes In: Cutting-edge technologies for sustained competitiveness. Agriculture conference: Proceedings of the 2001 PIPOC International palm oil congress, 20 - 22, Kuala Lumpur, Malaysia: 550-556.

Carson, J. K., Rooney, D., Gleeson, D. B., & Clipson, N. (2007) Altering the mineral composition of soil causes a shift in microbial community structure. FEMS Microbiology Ecology, 61: 414–423.

Castaldi, S., Riondino, M., Baronti, S., Esposito, F.R., Marzaioli, R & Rutigliano, F.A. (2012) Impact of biochar application to a mediterranean wheat crop on soil microbial activity and greenhouse gas fluxes. Chemosphere, 85: 1464–1471.

Chan, K.Y. & Z. Xu. (2009) Biochar:Nutrient properties and their enhancement. In: Lehmann, J. and S. Joseph (Eds.), Biochar for Environmental Management: Science and Technology. Earthscan, London, 67-84.

Chan, K.Y., Van Zwieten, L., Meszaros, I., Downie, A & Joseph, S. (2008) Using poultry litter biochars as soil amendments. Australian Journal of Soil Research, 46: 437-444.

Chang, E.H., Chung, R.S. & Tsai, Y.H. (2007) Effect of different application rates of organic fertilizer on soil enzyme activity and microbial population. Soil and Plant Nutrition, 53: 132-140.

Chantigny, M.H., D.A. Angers, & P. Rochette. (2002) Fate of carbon and nitrogenfrom animal manure and crop residues in wet and cold soils. Soil Biology and Biochemistry, 34: 509-517.

Cheng, C-H., Lehmann, J., Thies, J.E, Burton, S.D, & Engelhard, M.H. (2006) Oxidation of black carbon by biotic and abiotic processes. OrganicGeochemistry, 37: 1477–1488.

Christopher, J., Ennis, A., Garry Evans, Meez Islam, T., Komang R.S. & Eric Senior. (2012) Biochar: Carbon Sequestration, Land Remediation, and Impacts on Soil Microbiology, Critical Reviews in Environmental Science and Technology, 42: 2311-2364.

Clough, T.J., & Condron, L.M. (2010) Biochar and the nitrogen cycle. Journal of Environmental Quality, 39: 1218–1223.

Cross, A., & Sohi, S.P. (2011) The priming potential of biochar products in relation to labile carbon contents and soil organic matter status. Soil Biology and Biochemistry, 43: 2127–2134.

Cui, H.J., Wang, M.K., Fu, M.L., & Ci, E. (2011) Enhancing phosphorus availability inphosphorus-fertilized zones by reducing phosphate adsorbed on ferrihydrite using rice straw-derived biochar. Journal of Soils and Sediments, 11: 1135–1141.

Daniel, N., Dempster, A., Deirdre, B., Gleeson, A., Zakaria M., Solaiman, A., David, L., Jones, B., Daniel, V., Murphy, A. (2010) Biochar addition to soil changed microbial Community structure and decreased microbial biomass carbon and net inorganic nitrogen mineralised. School of the Environment and Natural Resources, University of Wales, Bangor, Gwynedd LL57, 2UW, UK.

DeLuca, T. H., MacKenzie, M.D., Gundale, M. J., & Holben,W. E. (2006) Wildfire-produced charcoal directly influences nitrogen cycling in forest ecosystems . Soil Science Society America Journal, 70: 448–453.

Domene, X., Stefania, M., Kelly, H., Akio Enders & Lehmann, J. (2014) Medium-term effects of corn biochar addition on soil biota activities and functions in a emperate soil cropped to corn. Soil Biology and Biochemistry, 72: 152-162.

Downie, A., Crosky, A., & Munroe, P (2009) Physical properties of biochar. In Biochar for Environmental Management. Science and Technology, Lehmann, J., Joseph, S., Eds., Earthscan: London, UK.13–32.

Fageria, N. K., & Baligar, V. C. (2008) Ameliorating soil acidity of tropical Oxisols by liming for sustainable crop production. Advances in agronomy, 99: 345-399.

Forster, P., V. Ramaswamy, P. Artaxo, T. Berntsen, R. Betts, D.W., Fahey, J., Haywood, J., Lean, D.C., Lowe, G., Myhre, J., Nganga, R., Prinn, G., Raga, M., Schulz, & R. Van Dorland. (2007). Changes in Atmospheric Constituents and in Radiative Forcing. In: Climate Change 2007: The Physical Science Basis. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Gaskin, JW., Steiner, C., Harris, K., Das, KC., & Bibens, B. (2008) Effect of low-temperature pyrolysis conditions on biochar for agricultural use. American Society of Agricultural and Biological Engineers, 51: 2061–2069.

Gaspar, M., Cabello, M., Pollero, R., & Aon, M. (2001) Fluorescein diacetate hydrolysis as a measure of fungal biomass in soil. Current Microbiology, 42: 339-344.

Giller, KE. (2001) Nitrogen fixation in tropical cropping systems, 2nd edn. CAB International, Wallingford.

Glaser, B. (2007) Prehistorically modified soils of central Amazonia: a model for ustainable agriculture in the twenty-first century. Philosophical Transactions of the Royal Society B: Biological Sciences, 362: 187-196.

Glaser, B., Lehmann J., & Zech.W. (2002) Amelorating physical and chemichal properties of highly weathered soils in the tropics with charcoal-A review. Biofertilizers, 35: 219-230.

Harris, RF., & Sommers, LE. (1968) Plate-dilution frequency technique for assay of microbial ecology. Applied Microbiology, 16: 330–334.

Haug, A., & Foy, C. E.(1984) Molecular aspects of aluminum toxicity. Critical Reviews in Plant Sciences, 1: 345-373.

Herath, H.M.S.K., Marta Camps-Arbestain, & Mike Hedley. (2013) Effect of biochar on soil physical properties in two contrasting soils: An Alfisol and an Andisol. Journal of Geoderma, 209–210:188–197.

Hina, K., Bishop, P., Arbestain, M.C.,Calvelo-Pereira, R., Macia-Agullo, J.A., Hindmarsh,J., Hanly, J.A., Macias, F., & Hedley, M.J. (2010) Producing biochars with enhanced surface activity through alkaline pretreatment of feedstocks. Australian Journal of Soil Research, 48: 606–617.

Hoshi, T. (2001) Growth promotion of tea trees by putting bamboo charcoal in soil, In Proceedings of International Conference on O-cha (Tea) Culture and Science, Tokyo, Japan, 147-150.

Ingram, C. W., Coyne, M. S., & Williams, D. W. (2005) Effects of commercial diazinon and imidacloprid on microbial urease activity in soil. Journal of Environmental Quality, 34: 1573–1580.

Ishizuka, J. (1992) Trends in biological nitrogen fixation research and application. Plant and Soil, 141: 197-209.

Islami, T., Bambang, G., Nur, B., & Agus, S. (2011) Maize yield and associated soil quality changes in cassava , Maize intercropping system After 3 Years of Biochar Application. Journal of Agriculture and Food Technology 1: 112-15.

Ivan, R., Kennedy, & Yao-Tseng Tchan. (2006) Biological nitrogen fixation in non-leguminous field crops: Recent advances. Plant and Soil, 141: 93-118.

Jeffery, S., Verheijen, A., vander, V. M., & Bastos, A. C. (2011) A quantitativereview of the effects of biochar application to soils on crop productivity using meta-analysis. Agriculture, Ecosystems and Environment, 144: 175–187.

Jensen, H L. (1951) Notes on the biology of Azotobacter. Proceedings of the Society of Applied Bacteriology, 14: 89–94.

Jin, H. (2010) Characterization of microbial life colorizing biochar and biochar-amended soils. PhD Dissertation, Cornell University,Ithaca, NY.

Jindo, K., Sánchez-Monedero, M. A., Hernández, T., García, C., Furukawa, T., Matsumoto, K., & Bastida, F. (2012) Biochar influences the microbial community structure during manure composting with agricultural wastes. Science of the Total Environment, 416: 476-481.

Johnson, D.W., & P.S. Curtis. (2001) Effects of forest management on soil C and N storage: meta analysis. Forest Ecology and Management 140: 227-238.

Jones, J.B. (1985) Soil testing and plant analysis: guides to the fertilization of horticultural crops. HortTechnology, 7:1–68.

Joseph, Stephen, Camps-Arbestain, M., Lin, Y., Munroe, P., Chia, C. H., Hook, J & van Zwieten, L. (2010) An investigation into the reactions of biochar in soil. Australian Journal of Soil Research, 48: 501-515.

Josiah, H., Michael, D., Dwight, S., & Andrew, K. (2010) The Basics of biochar : A natural soil amendment. College of Tropical Agriculture and HumanResources, SCM-30.

Kamprath, E. J. (1984) Crop response to lime on soils in the tropics. In soil acidity and liming. 2nd ed. F Adams, Agronomy Monograph, 12: 349-368.

Khalid, E., Mary, S., James, I., Kenneth, B., & Matthew, W. (2014) Biochar effectson soil microbial communities and resistance of enzymes to stress. MSc Dissertation. Colorado State University. UMI Number: 1550608.

Khalil, M.I., Hossain, M.B., & Schmidhalter, U. (2005) Carbon and nitrogenmineralization in different upland soils of the subtropics treated with organic materials. Soil Biology and Biochemistry, 37: 1507-1518.

Kim, J.-S., Sparovek, S., Longo, R.M., De Melo, W.J., & Crowley, D. (2007) Bacterial diversity of terra preta and pristine forest soil from the Western Amazon. Soil Biology and Biochemistry, 39: 648-690.

Kimetu, Joseph, M., J. Lehmann, Solomon, O., Ngoze, Daniel, N., Mugendi, James, M., Kinyangi, Susan Riha, Lou Verchot, John, W., Recha, & Alice, N. Pell. (2008) Reversibility of Soil Productivity Decline with Organic Matter of Differing Quality Along a Degradation Gradient. Ecosystems 11: 726-39.

Krull, E.S., J. Lehmann, J. Skjemstad, & J. Baldock. (2008) The global extent of black C in soil; is it everywhere? In: Hans G. Schroder (ed.), Grasslands; ecology, management and restoration. New York: Nova Science Publishers, 4: 13–17.

Kumari, K. G. I. D., PerMoldrup., Marcos, P., & Lis W. de J. (2014) Phenanthrene Sorption on Biochar-Amended Soils: Application Rate, Aging, and Physicochemical Properties of Soil. Water Air and Soil Pollution. 25:2105.

Laird, D., Fleming, P., Wang, B., Horton, R., & Karlen, D. (2010) Biochar impact on nutrient leaching from a Midwestern agricultural soil. Geoderma, 158: 436–442.

Lammirato, C., Miltner, A., & Kaestner, M. (2011) Effects of wood char andactivated carbon on the hydrolysis of cellobiose by [beta]-glucosidase from Aspergillus niger. Soil Biology and Biochemistry, 43:1936-1942.

Lehmann, J., J. P. da Silva Jr., C. Steiner, T. Nehls, W. Zech, and B.Glaser. 2003. Nutrient availability and leaching in an archaeological anthrosol and a ferralsol of the central Amazon basin: Fertilizer, manure and charcoal amendments. Plant & Soil, 249: 343-357.

Lehmann, J., & Joseph, S. (2009) Biochar for environmental management: An introduction. In Biochar for Environmental Management, Science and Technology, Lehmann, J., Joseph, S., Eds., Earthscan: London, UK. 17: 416.

Lehmann, J. (2007) Bio-energy in the black. Frontiers in Ecology and the Environment, 5: 381- 387.

Lehmann, J., Rillig, M.C.,Thies, J., Masiello, C.A., Hockaday,W.C & Crowley, D. (2011) Biochar effects on soil biota—A review. Soil Biology and Biochemistry, 43: 1812–1836.

Lehmann, J., & Rondon, M. (2006) Bio-char soil management on highly weathered soils in the humid tropics. Biological approaches to sustainable soil systems. CRC Press Boca Raton, Florida, 20: 517-530.

Liang, B., Lehmann, J., Solomon, D., Kinyangi, J., Grossman, J., O'Neill, B., Skjemstad, J. O.,Thies, J., Luizao, F. J., Petersen, J., & Neves, E. G. (2006) Black carbon increases cation exchange capacity in soils. Soil Science Society of America Journal, 70:1719–1730.

Major, J., M. Rondon, D. Molina, S.J. Riha., & J. Lehmann. (2010) Maize yield and nutrition after 4 years of doing biochar application to a Colombian savanna oxisol. Plant and Soil, 333:117–128.

Major, J., Steiner, C., Downie, A., & Lehmann, J. (2009) Biochar effects on nutrient leaching. In: Lehmann J, Joseph S (eds) Biochar for Environmental Management. Earthscan, London, 271-287.

Marco, A., Rondon., Johannes Lehmann., Juan, R., & Maria, H. (2007) Biological nitrogen fixation by common beans (*Phaseolus vulgaris L.*) increases with bio-char additions. Biology and Fertility in Soils, 43: 699-708.

Martha, W., Gilliland. (1987) Nitrogen fixing biotechnologies for corn in Mexico and central American, Impact Assessment For International Development, 5: 71-94.

Martin, J.P. (1950) Use of acid, rose bengal, and streptomycin in the plate method for estimating soil fungi. Soil Science, 69: 215-232.

Milla, O., Wang, H., & Huang, W. (2013) Feasibility Study using Municipal Solid Waste Incineration Bottom Ash and Biochar from Binary Mixtures of Organic Waste as Agronomic Materials. J. Hazard. Toxic Radioact. Waste, 17: 187–195.

Monkiedje, A., Ilori, M. O., & Spiteller, M. (2002) Soil quality changes resulting from the application of the fungicides mefenoxam and metalaxyl to a sandy loam soil. Soil Biology and Biochemistry, 34: 1939–1948.

MPOB. (2012) Malaysian oil palm statistics 2011. MPOB, Bangi, Malaysia.

Mukhlis, M. (2006) Composting of oil palm empty fruit bunches with Trichorderma and its effects on growth of tomato and corn. Phd Thesis, Universiti Putra Malaysia.

Naher, UA., Radziah, O., Shamsuddin, ZH., Halimi, MS., & Mohd Razi. (2011) Effect of root exuded specific sugars on biological nitrogen fixation and growth promotion in rice (Oryza sativa). Australian Journal of Crop Science, 5: 1210–1217.

Nair, Ajay., & Lawson,Vincent. (2012) Effect of Biochar on Sweet Corn Production. Agricultural Science Commons, Agriculture Commons, and the Horticulture Commons, 368: 12-20.

Nannipieri, P. L., Giagnoni, G., Renella, E., Puglisi, B., Ceccanti, G., Masciandaro, F., Fornasier, M. C., Moscatelli & S. Marinari. (2012) Soil enzymology: classical and molecular approaches. Biology and Fertility in Soils, 47:863–874

Nelson, D.W., & Sommers L.E. (1982) Total Carbon, Organic Carbon, and Organic Matter. In Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties Agronomy. 9: 539-577.

Nelson, N.O., Agudelo, S.C., Yuan, W., & Gan, J. (2011) Nitrogen and phosphorus avail-ability in biochar-amended soils. Soil Science, 176: 218-226.

Nguyen, B., Lehmann, J., Kinyangi, J., Smernik, R., Riha, S.J & Engelhard, M.H. (2008) Long-term black carbon dynamics in cultivated soil. Bio-geochemistry 89: 295-308.

Norazlina A. S, Che Fauziah Ishak & Rosenani A. B. (2014) Characterization of oil palm empty fruit bunch and rice husk biochars and their potential to adsorb arsenic and cadmium. American Journal of Agricultural and Biological Sciences, 9: 450-456.

Nowak, A., Nowak, J., Klodka, D., Pryzbulewska, K., Telesinski, A., & Szopa, E. (2004) Changes in the microflora and biological activity of the soil during the degradation of isoproturon. Journal of Plant Diseases and Protection,19: 1003-1016.

Novak, J. M., W. J. Busscher, D. L. Laird, M. Ahmedna, D. W. Watts, & M. A. S. Niandou. (2009) Impact of Biochar Amendment on Fertility of a Southeastern Coastal Plain Soil. Soil Science, 174: 105-112.

Oguntunde, P.G., Fosu, M., Ajayi, A.E & Van De Giesen., N.D. (2004) Effects of charcoal production on maize yield, chemical properties and texture of soil. Biology and Fertility of Soils, 39: 295-299.

O'Neill, B., J. Grossman, M.T. Tsai, J.E. Gomes, J. Lehmann, J. Peterson, E. Neves, & J.E. Thies. (2009) Bacterial community composition in Brazilian Anthrosols and adjacent soils characterized using culturing and molecular identification. Microbial Ecology 58: 23–35.

Palm, CA., Gachengo, CN., Delve, RJ., Cadisch, G., & Giller, K. (2001) Organic inputs for soil fertility management in tropical agroecosystems: application of an organic resource database. Agriculture Ecosystems Environment Journal, 83: 27–42.

Parkinson, D., Gray, T.R.G., & Williams, S.T. (1971) Methods for studying the ecology of soil microorganisms. IBP Handbook No.19.

Peoples, M. B., & E. T. Craswell. (1992) Biological nitrogen fixation: investments, expectations and actual contributions to agriculture. Plant and Soil, 141: 13-39.

Raymond, J., Siefert, J.L., Staples, C.R., & Blankenship, R.E. (2004) The naturalhistory of nitrogen fixation. Molecular Biology and Evolution, 21: 541-554.

Rachel, U., & Randy, K. (2011) Effect of Three Different Qualities of Biochar on Selected Soil Properties. Soil Science and Plant Analysis, 42: 2274–2283,

Radziah, O., Umme, A. N., & Siti, Z. Y. (2013) Effect of urea-N on growth andindoleacetic acid of urea-N on growth and indoleacetic acid production of *Stenotrophomonas maltophilia* (Sb16) isolated from rice growing soils in Malaysia. Chilean Journal of Agricultural Research, 73: 187-192.

Rajesh, C., Javier, M., Thomas, E., Schumacher, Douglas, D., Malo, & James, L., Julson. (2014) Effect of biochar on chemical properties of acidic soil, Archives Agronomy and Soil Science, 60: 393-404.

Renner, R. (2007) Rethinking biochar. Environmental Science and Technology, 41: 5932-5933.

Rillig, M. C., Wagner, M., Salem, M., Antunes, P.M., George, C., Ramke, H.G., Titirici, M.M., & Antonietti, M. (2010) Material derived fromhydrothermal carbonization: Effects on plant growth and arbuscular mycorrhiza. Applied Soil Ecology, 45: 238-242.

Rondon, M. A., J. Lehmann, J. Ramirez, & M. Hurtado. (2007) Biological nitrogen fixation by common beans (Phaseolus vulgaris L.) increases with bio-char additions. Biology and Fertility of Soils, 43: 699-708.

Rout, G. R., Samantaray, S., & Das, P. (2001) Aluminium toxicity in plants: a review. Agronomie, 21: 3-21.

Rutigliano, F. A., Romano, M., Marzaioli, R., Baglivo, I., Baronti, S., Miglietta, F & Castaldi, S. (2014) Effect of biochar addition on soil microbial community in a wheat crop. European Journal of Soil Biology, 60: 9-15.

Samsuri, A.W., F. Sadegh-Zadeh & B.J. She-Bardan. (2013) Adsorption of As (III) and As (V) by Fe coated biochars and biochars produced from empty fruit bunch and rice husk. Journal of Environmental Chemical Engineering 1: 981-988,

Samuel, T., Partey, Richard, F., Preziosi, & Geoffrey, D., Robson. (2014) Short-Term Interactive Effects of Biochar, Green Manure, and Inorganic Fertilizer on Soil Properties and Agronomic Characteristics of Maize.Journal of Agricultural Research, 2: 124-128.

Sanchez- M., M C. Mondini, M. Cayuela, A. Roig, M. Contin., & M. De Nobili. (2008) Fluorescein diacetate hydrolysis, respiration and microbial biomass in freshly amended soils. Biology and Fertility of Soils, 44: 885-895

Sander Bruun, T El-Zahery., & L. Jensen.(2010) Carbon sequestration with biochar stability and effect on decomposition of soil organic matter. Earth and Environmental Science, 6: 242-256.

Sarah Carter, Simon, S., Saran Sohi, Tan Boun, S., & Stephan, H. (2013) The Impact of Biochar Application on Soil Properties and Plant Growth of Pot Growth Lettuce (Lactuca sativa) and Cabbage (Brassica chinensis). Agronomy, 3: 404-418.

Saran Sohi, & Miranda P-M. (2013) Report on the effect of fresh and weatheredbiochar on the architecture of crop roots and soil mineral nitrogenin the rhizosphere. The Interreg IVB project Biochar.
http://archive.northsearegion.eu.

Schnurer, J., & Rosswall, T. (1982) Fluorescein Diacetate Hydrolysis as a Measure of Total Microbial Activity in Soil and Litter, Applied and Environmental Microbiology, 43: 1256-1261.

Schulz, H., & Glaser, B. (2012) Effects of biochar compared to organic and inorganic fertilizers on soil quality and plant growth in a greenhouse experiment. Journal of Plant Nutrition and Soil Science, 175: 410–422.

Shackley, S., Sohi, S., Ibarrola, R., Hammond, J., Mašek, O., Brownsort, P., & Haszeldine, S. (2012) Biochar as a Tool for Climate Change Mitigation and Soil Management. In Encyclopedia of Sustainability Science and Technology, 183: 947–966.

Shao, X., & Zheng Jian-wei. (2014) Soil Organic Carbon, Black Carbon, and Enzyme Activity Under Long-Term Fertilization. Journal of Integrative Agriculture,13: 517-524.

Shamshuddin, J., & Fauziah, C.I. (2010) Weathered Tropical Soils:The Ultisols and Oxisols. UPM Press, Serdang, Malaysia.

Shelby, R., & J. Lehmann. (2010) Biochar as an amendment to improve soil fertility. Thesis of College of Agriculture of Cornell University.

Shukla, G., & A. Varma. (2011) Role of enzymes in maintaining soil health. Soil Enzymology, Soil Biology, 22: 25-42.

Silva, J. A., & Uchida, R. (2000) Biological nitrogen fixation, nature's partnership for sustainable agricultural production. College of Tropical Agriculture and Human Resources, 13: 121-126.

Singer, J.W., Logsdon S.D., & D.W. Meek. (2007) Tillage and compost effect on orn growth, nutrient accumulation and grain yield. Agronomy Journal, 99: 80-87.

Singer, M., & Munn, D. N. (1999) Soils an introduction. Four edition. Prentice Hall Upper Saddle River, 1: 07458. 527.

Singh, B.P., Hatton, B.J., Singh, B., Cowie, A., & Kathuria, A. (2010) Influence of biochars on nitrous oxide emission and nitrogen leaching from two contrasting soils. Journal of Environmental Quality, 39:1224-35.

Sohi, S. P., E. Krull, E. Lopez-Capel, & R. Bol. (2010) A Review of Biochar and its Use and Function in Soil. Advances in Agronomy, 105: 47-82.

Solomon, D., J. Lehmann, J. Thies, T. Schafer, B. Liang, J. Kinyangi, E. Neves, J. Petersen, F. Luizo & J. Skjemstad. (2007) Molecular signature and sources of biochemical recalcitrance of organic C in Amazonian dark earths. Geochimica et cosmochimica Acta, 71: 2285-2298.

Somasegaran, P., & Hoben, H.J. (1985) Methods in Legume-Rhizobium Technology. University of Hawaii NifTAL. http://www.ctahr.hawaii.edu.

Sophie, M., & Bruno T. (2003) Plant growth-promoting bacteria and nitrate availability: impacts on root development and nitrate uptake. Journal of Experimental Botany, 55: 394-419.

Steinbeiss, S., Gleixner, G., & Antonietti, M. (2009) Effect of biochar amendment on soil carbon balance and soil microbial activity. Soil Biology and Biochemistry, 41: 1301-1310.

Steiner, C., Glaser, B., Teixeira, W.G., Lehmann, J., Blum, W.E. & Zech, W. (2008) Nitrogen retention and plant uptake on a highly weathered central Amazonian Ferralsol amended with compost and charcoal. Journal of Plant Nutrition and Soil Science, 171: 893-899.

Steinweg, J. M., Dukes, J. S., Paul, E. A., & Wallenstein, M. D. (2013) Microbial responses to multi-factor climate change: effects on soil enzymes. Frontiers in Microbiology, 4: 218-230.

Streubel, J.D., Collins, H.P., Garcia-Perez, M., Tarara, J., Granatstein, D & Kruger, C.E. (2011) Influence of contrasting biochar types on five soils at increasing rates of application. Soil Biology and Biochemistry, 75: 1402–1413.

Tabatabai, M.A. & Bremner, J.M. (1969) Use of p-Nitrophenyl phosphate for assay of soil phosphatase activity. Soil Biology & Biochemistry, 1: 301–307.

Tabatabai, M.A & Bremner, J.M. (1972) Assay of urease activity in soils Soil Biology and Biochemistry, 4: 479–487.

Tejada, M., Garcia, C., Gonzalez, J. L., & Hernandez, M. T. (2006) Use of organic amendment as a strategy for saline soil remediation: influence on the physical, chemical and biological properties of soil. Soil Biology and Biochemistry, 38: 1413–1421.

Thies, J. E., & Rillig, M. C. (2009) Characteristics of biochar: biological properties. Biochar for environmental management: Science and Technology, 10: 85-105.

Thomas, G.W. (1982) Exchangeable cations. In A.L. Page (ed.). Methods of soil analysis, Part 2, Second Edition, Agronomy Monograph 9. American Society of Agronomy.

Thomas, R.L, Sheard, R.W., & Moyer, J.R. (1967) Comparison of conventional and automated procedures for nitrogen, phosphorus and potassium analysis of plant materials using a single digestion. Agronomy, 59: 240-243.

Tim, J., Clough, Leo, M., Condron, Claudia Kammann, & Christoph Muller. (2013) A Review of Biochar and Soil Nitrogen Dynamics. Agronomy, 3: 275-293.

Topoliantz, S., J.-F. Ponge, & S. Ballof. (2005) Manioc peel and charcoal: a potential organic amendment for sustainable soil fertility in the tropics. Biology and Fertility in Soils, 41: 15-21.

Unger, R., & Killorn, R. (2012) Effect of the application of biochar on selected soil chemical properties, corn grain, and biomass yields in Iowa. Commun. Soil Science and Plant Analysis, 42: 2441–2451.

USDA Foreign Agriculture Service. (2014) Assessments of commodity and trade issues. GAIN Report Number: MY 4001. http://gain.fas.usda.gov.

Van Zwieten, L., Kimber, S., Downie, A., Chan, K.Y., Cowie, A., Wainberg, R & Morris, S. (2007) Papermill char: Benefits to soil health and plant production in Proceedings of the Conference of the International Agrichar Initiative, Terrigal, NSW, Australia.

Van Zwieten, L., S. Kimber, S. Morris, K. Y. Chan, A. Downie, J. Rust, S. Joseph & A. Cowie. (2010) Effects of biochar from slow pyrolysis of papermill waste on agronomic performance and soil fertility. Plant and Soil, 327: 235–246.

Verheijen, F.G.A., Jeffery, S., Bastos, A.C., van der Velde, M., & Diafas, I. (2009) Biochar Application to Soils—A Critical Scientific Review of Effects on Soil Properties, Processes and Functions. Office for the Official Publications of the European Communities, 21: 61-77.

Volker, R., & Gabriel, S. (2004) Iron transport in plants: Future research in view of a plant nutritionist and a molecular biologist, Soil Science and Plant Nutrition, 50: 1003-1012.

Warnock, D.D., J. Lehmann, T.W. Kuyper, & M.C. Rillig. (2007) Mycorrhizal responses to biochar in soil-concepts and mechanisms. Plant and Soil, 300: 9-20.

Watson, M. E., & Brown, J. R. (2011) pH and Lime Requirement. Chemical Soil Test Procedures for the North Central Region. North Central Regional Research Publication, 221: 13-16.

Winsley, P. (2007) Biochar and bioenergy production for climate change mitigation.New Zealand Science. Review, 64: 5-10.

Woods,W. I., & Denevan, W. M. (2009) Amazonian dark earths: the first centuryoF reports. In Amazonian Dark Earths: Wim Sombroek's Vision, Springer Netherlands, 1-14.

Woods, W. I., & McCann, J. M. (1999) The anthropogenic origin and persistence of Amazonian dark earths. In Yearbook. Conference of Latin Americanist Geographers, 7-14. Conference of Latin Americanist Geographers.

Wu, Y., G. Xu & H. B. Shao. (2014) Furfural and its biochar improve the general properties of a saline soil. Solid Earth, 5: 665–671.

Xu, G., Shao, H.B., & Sun, J.N. (2013) What is more important for enhancing nutrient bio-availability with biochar application into a sandy soil: direct or indirect mechanism? Journal of Ecological Engineering, 52: 119–124.

Yang, Z, Liu S, Zheng D & Feng, S. (2006) Effects of cadmium, zinc and lead on soil enzyme activities. Journal of Environmental Science, 18:1135–1141.

Yamato, M., Okimori, Y., Wibowo, I.F., Anshori, S & Ogawa, M. (2006) Effects of the application of charred bark of Acacia mangium on the yield of maize, cowpea and peanut, and soil chemical properties in South Sumatra, Indonesia. Soil Science and Plant Nutrition, 52: 489-495.

Yuan, J.H., & Xu, R.K. (2011) The amelioration effects of low temperature biochar generated from nine crop residues on an acidic Ultisol. Soil Use and Management, 27: 110–115.

Zaharah, A.R., & Lim, K.C. (2000) Oil palm empty fruit bunches a source of nutrients and soil ameliorant in oil palm plantations. Malays. Journal of Soil Science, 4:51-66.

Zurich, ETH., Gattinger, A., & Scheifele, M. (2014) Effects of biochar amendments on plant growth, microbial communities and biochar decomposition in agricultural soils. Official Project number, 315230_138238 / 1.