



UNIVERSITI PUTRA MALAYSIA

***IMIDACLOPRID RESIDUE IN PADDY, UNPOLISHED AND POLISHED
RICE, AND HEALTH RISK ASSESSMENT OF RICE CONSUMPTION
IN TANJUNG KARANG, MALAYSIA***

RAIHANAH BINTI CHOKELI

FPSK(m) 2015 58



**IMIDACLOPRID RESIDUE IN PADDY, UNPOLISHED AND POLISHED
RICE, AND HEALTH RISK ASSESSMENT OF RICE CONSUMPTION
IN TANJUNG KARANG, MALAYSIA**

By

RAIHANAH BINTI CHOKELI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

October 2015

All materials contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Degree of Master of Science

IMIDACLOPRID RESIDUE IN PADDY, UNPOLISHED AND POLISHED RICE, AND HEALTH RISK ASSESSMENT OF RICE CONSUMPTION IN TANJUNG KARANG, MALAYSIA

By

RAIHANAH BINTI CHOKELI

October 2015

Chair : Professor Zailina Hashim, PhD
Faculty : Medicine and Health Sciences

Background: Rice (*Oryza sativa* L.) by large is staple food consumed and produced by most of the world population including Malaysia. In order to combat from brown planthoppers attack, farmers have used imidacloprid in the paddy fields. However, the used of imidacloprid might give some effect to the population that consume rice.

Objective: The objective of this study was to evaluate the non-carcinogenic health risks on Tanjung Karang's population (n=552) through rice consumption.

Methodology: Fifty samples of paddy, unpolished rice and polished rice were collected from paddy fields and milling factory in Tanjung Karang, Selangor and were analysed for imidacloprid residues. The population was divided into 3 age groups which are adult, teenagers and children. Imidacloprid residue analysis in paddy and rice samples was conducted using the simplified and validated QuEChERS (quick, easy, cheap, effective, rugged, and safe) preparation method. Samples were extracted with acetonitrile (ACN) and salts anhydrous magnesium sulphate ($MgSO_4$) and sodium chloride (NaCl) while cleaning was by treating with primary secondary amine (PSA). Imidacloprid residue was quantified with Ultra High Performance Liquid Chromatography (UHPLC) using C18 column and UV detector. The method was validated in term of linearity, range, limit of detection (LOD), limit of quantification (LOQ), precision (repeatability and intermediate precision), and accuracy (recovery). LOD and LOQ of the method was 0.001 mg/L and 0.003 mg/L, respectively.

Result: The validated method provides good analytical results wherein the linearity r^2 was 0.9996, recoveries 80–120 % and RSDs <20%. Through of this method, imidacloprid residue was detected in 20%, 14% and 10% of paddy samples, unpolished rice and polished rice, respectively with 4% of the total samples exceeded the MRL. The mean imidacloprid concentration was 0.14 mg/kg in the paddy sample, 0.04 mg/kg in unpolished rice sample and 0.03 mg/kg in polished rice sample. The EDI for adult, adolescent and children were 0.08 mg/kg, 0.03 mg/kg and 0.0002 mg/kg, respectively. HQ value of adult was 1.32 showed that this group was exposed to non-carcinogenic health risk through rice consumption. **Conclusion:** The result of analysis showed that imidacloprid residues were present in paddy and rice samples originate from Tanjung Karang paddy cultivation area. Imidacloprid was detected in mostly in paddy samples compared to brown rice and white rice, respectively with 4% of the total samples exceeded the Malaysian Residue Level (MRL). This indicates that routine monitoring of

this contaminants in food items is required to prevent, control and reduce the contamination and to minimize health risks. This research has provided important information on imidacloprid residues contamination on local paddy and rice for the first time.

Keywords: Imidacloprid, QuEChERS, Health Risk Assessment, UHPLC-UV



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Master Sains

**IMIDACLOPRID DALAM PADI, BERAS YANG TIDAK DIGILAP DAN
BERAS YANG DIGILAP SERTA PENILAIAN RISIKO KESIHATAN
PENGUNAAN BERAS PADA PENDUDUK TANJUNG KARANG,
MALAYSIA**

Oleh

RAIHANAH BINTI CHOKELI

Oktober 2015

Pengerusi : Profesor Zailina Hashim, PhD
Fakulti : Perubatan dan Sains Kesihatan

Pendahuluan: Beras (*Oryza sativa* L.) adalah makanan ruji yang digunakan dan dihasilkan oleh kebanyakan penduduk di dunia termasuk Malaysia. Dalam usaha untuk memerangi daripada serangan bena perang ke atas padi, petani telah menggunakan imidacloprid di sawah padi. Walau bagaimanapun, penggunaan imidacloprid mungkin dapat memberi sedikit kesan kepada mereka yang makan nasi. **Objektif:** Tujuan kajian ini adalah untuk menilai risiko kesihatan bukan karsinogen terhadap penduduk di Tanjung Karang (n = 552) melalui penggunaan beras. **Metodologi:** Lima puluh sampel padi, beras tidak digilap dan beras digilap telah dikumpulkan dari sawah padi dan kilang di Tanjung Karang, Selangor. Sampel tersebut telah dianalisis untuk mengetahui kandungan sisa imidacloprid. Populasi kajian telah dibahagikan kepada 3 kumpulan umur iaitu dewasa, remaja dan kanak-kanak. Analisis sisa imidacloprid dalam sampel padi dan beras telah dijalankan menggunakan kaedah QuEChERS (cepat, mudah, murah, berkesan, lasak, dan selamat). Sampel telah diekstrak dengan menggunakan asetonitril (ACN), garam magnesium sulfat kontang (MgSO_4) dan natrium klorida (NaCl) manakala pembersihan dilakukan menggunakan primary secondary amina (PSA). Sisa imidacloprid telah diukur dengan Ultra High Performance Liquid Chromatography (UHPLC) menggunakan kolum C18 dan pengesan UV. Kaedah ini telah disahkan dalam tempoh kelinearan, pelbagai, had pengesanan (LOD), had kuantifikasi (LOQ), ketepatan (kebolehulungan dan ketepatan perantaraan), dan ketepatan (pemulihan). Bacaan LOD dan LOQ adalah 0.001 mg / L dan 0.003 mg / L. **Hasil kajian:** Kaedah di dalam kajian ini disahkan memberikan hasil analisis yang baik di mana kelinearan r^2 adalah 0.9996, pemulihan 80-120% dan RSDs <20%. Melalui kaedah ini, sisa imidacloprid dikesan dalam 20%, 14% dan 10% daripada sampel padi, beras tidak digilap dan beras digilap dengan 4% daripada jumlah sampel melebihi MRL. Kepekatan imidacloprid min adalah 0.14 mg / kg dalam sampel padi, 0.04 mg / kg dalam sampel beras tidak digilap dan 0.03 mg / kg dalam sampel beras digilap. EDI bagi populasi orang dewasa, remaja dan kanak-kanak adalah 0.08 mg / kg, 0.03 mg / kg dan 0.0002 mg / kg. Nilai HQ dewasa adalah 1.32 menunjukkan bahawa

kumpulan ini telah terdedah kepada risiko kesihatan bukan karsinogen melalui penggunaan beras. **Kesimpulan:** Hasil analisis menunjukkan bahawa terdapat sisa imidacloprid di dalam padi dan beras sampel yang diperoleh dari kawasan penanaman padi Tanjung Karang. Imidacloprid dikesan di dalam kebanyakan sampel padi berbanding beras perang dan beras putih, masing-masing dengan 4% daripada jumlah sampel melebihi Tahap Sisa Baki Malaysia (MRL). Ini menunjukkan bahawa pemantauan rutin bahan cemar ini dalam barangan makanan diperlukan untuk mencegah, mengawal dan mengurangkan pencemaran dan untuk mengurangkan risiko kesihatan. Kajian ini telah menyediakan maklumat penting mengenai pencemaran sisa imidacloprid pada padi tempatan dan beras buat kali pertama.

Kata Kunci: Imidacloprid, QuEChERS, Penilaian Risiko Kesihatan, UHPLC-UV

ACKNOWLEDGEMENT

In the name of Allah, the Most Gracious and the Most Merciful

Alhamdulillah, all praises to Allah for the strengths and His blessing in completing this thesis. Special appreciation goes to my dedicated supervisor, Prof. Dr. Zailina Hashim, for her supervision and never ending support. Her meaningful words and invaluable help of constructive comments and suggestions throughout the research and thesis works have contributed a lot to the success of this research. Not forgotten, my appreciation to my co-supervisors, Dr. Ho Yu Bin, Dr. Saliza Mohd Elias and Dr. Norida Mazlan for their support and knowledge pertaining to this research.

I would like to express my deepest gratitude to my beloved and supportive parents, Mr. Chokeli Bin Isa and Mrs. Zahariah Binti Ariffin and also my siblings, for their endless love, prayers and encouragement. My acknowledgement also goes to all the laboratory staffs of Environmental Health Laboratory, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia for their guidance and assistance throughout the sample analysis. And not forgotten to all office staffs of Department of Occupational and Environmental Health, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia for their co-operations.

I would like also to dedicate my appreciation to Department Of Agriculture, Kuala Selangor and Jawatankuasa dan Keselamatan Kampung (JKKK) Kampung Sawah Sempadan for the permission to conduct study in the paddy field agricultural area. Special thanks to all the farmers for their cooperation and participation. Last but not least, sincere thanks to all my adorable friends especially Nurul Syazani, Fauzudin Mislan Aziemah, Najihah, Nur Aida, Noreen Adila and others for their kindness and moral support during my study.

Thanks for the friendship and memories.

Thank you

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Zailina Hashim, PhD

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Ho Yu Bin, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Saliza Mohd Elias, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Norida Mazlan, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations, and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials, as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Raihanah Binti Chokel, GS34867

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: Prof . Zailina Hashim

Signature: _____
Name of
Member of
Supervisory
Committee: Dr Ho Yu Bin

Signature: _____
Name of
Member of
Supervisory
Committee: Dr Saliza Mohd Elias

Signature: _____
Name of
Member of
Supervisory
Committee: Dr Norida Mazlan

TABLE OF CONTENTS

ABSTRACT	Page
<i>ABSTRAK</i>	i
ACKNOWLEDGEMENT	iii
APPROVAL	v
DECLARATION	vi
LIST OF TABLES	viii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xiv
	xv

CHAPTER

1	INTRODUCTION	1
1.1	Introduction	1
1.2	Problem Statement	3
1.3	Study Justification	4
1.4	Study Contribution	4
1.5	Conceptual Framework	5
1.6	Definition Of Term	6
1.6.1	Conceptual Definition	6
1.6.2	Operational definition	6
1.7	Research Objectives	7
1.7.1	General Objective	7
1.7.2	Specific Objectives	7
1.7.3	Hypothesis	7
2	LITERATURE REVIEW	8
2.1	Agriculture sector in Malaysia	8
2.2	Rice production in Malaysia	9
2.3	Rice Consumption Pattern in Malaysia	11
2.4	Pesticides Usage Worldwide	12
2.5	Pesticides usage in agricultural setting in Malaysia	14
2.6	Pesticide	14
2.7	Imidacloprid	16
2.7.1	Background	16
2.7.2	Uptake, Translocation and Metabolism of Imidacloprid in Plants	17
2.7.3	Half lives of imidacloprid	18
2.7.4	Mode of action on insects	19
2.7.5	Imidacloprid fate in animal body	20
2.7.6	Toxicological study on animals (in vivo)	20
2.7.7	The toxicity of imidacloprid on human	22
2.7.8	Pesticides residue in food crop	23
2.7.9	Existing regulatory authorities to monitoring Maximum Residue Level (MRLs)	24
2.8	Method of analysis for imidacloprid residue in paddy and rice	25
2.8.1	Analysis of imidacloprid using QuEChERS Method	26

2.9	Health risk assessment	28
2.10	Health risk assessment of imidacloprid through dietary exposure	32
3	METHODOLOGY	34
3.1	Study Location	34
3.2	Study Design	36
3.3	Sampling	36
3.3.1	Paddy and rice sampling point	36
3.3.2	Community survey	36
3.3.3	Study population	37
3.3.3.1	Inclusion criteria for study population	37
3.3.4	Sampling frame	37
3.3.4.1	Sampling unit	37
3.3.5	Sample size	38
3.3.5.1	Estimation of respondents for community survey	38
3.3.5.2	Collection of paddy and rice sample	39
3.3.6	Sampling techniques	40
3.3.7	Targeted pesticides	40
3.4	Sampling Materials	40
3.4.1	Paddy and rice samples	40
3.4.2	Community Survey	40
3.5	Sample Analysis	41
3.5.1	Chemicals and Standards	41
3.5.2	Method Validation	42
3.5.3	Linearity	42
3.5.4	Recovery	42
3.5.5	Precision	42
3.5.6	Limit of Detection (LOD) and Limit of Quantification (LOQ)	43
3.5.7	Recovery studies and quality control	43
3.6	Extraction Procedure	43
3.6.1	UHPLC-UV analysis	43
3.7	Strategy for Health Risk Assessment used in this study	44
3.8	Ethical Issues	46
4	RESULTS AND DISCUSSION	47
4.1	Community Survey	47
4.2	Laboratory analysis	49
4.2.1	UHPLC-UV peak area	49
4.2.2	Linearity	50
4.2.3	Recovery and precision	51
4.2.4	Limit of Detection (LOD) and Limit of Quantification (LOQ)	51
4.3	Analysis of imidacloprid in different type of samples	52
4.4	Health Risk Assessment	54
4.5	Uncertainties of Health Risk Assessment	56
5	CONCLUSION AND RECOMMENDATIONS	58

REFERENCES	60
APPENDICES	73
BIODATA OF STUDENT	75
PUBLICATION	76



LIST OF TABLES

Table	Page
2.1 The studies of uptake, translocation and metabolism of imidacloprid in plants	18
2.2 Half life of imidacloprid	19
2.3 Bodies of regulation establishing the MRL of imidacloprid in rice	25
2.4 Example of studies that used QuEChERS methods to analyze Imidacloprid residue in rice	27
3.1 Tanjung Karang total population	39
3.2 Technique of Sample Collection	40
3.3 Details of questionnaire for community survey	41
3.4 The risk acceptability for non-carcinogenic health effect (USEPA, 2007)	46
4.1 Background information of respondent	48
4.2 Amount of rice intake annually (kg) per each age group	48
4.3 Frequency of rice intake for each group	48
4.4 Portion size of rice intake (per day)	49
4.5 LOD and LOQ of imidacloprid in rice sample	51
4.6 The imidacloprid concentration in paddy, brown rice and white rice samples	52
4.7 EDI and HQ of imidacloprid residue in white rice samples	52
4.8 EDI and HQ of imidacloprid residue in white rice samples	55

LIST OF FIGURES

Figure	Page
1.1 Worldwide consumption of pesticides	1
1.2 Imidacloprid – chemical structure	2
2.1 Land areas of major crops in Malaysia (x1000ha)	8
2.2 Map of eight main granaries in Malaysia	9
2.3 Average yield of paddy by state in Malaysia (2010/2011)	10
2.4 Average yield of paddy by granary area in Malaysia (2010/2011)	10
2.5 Annual per capita consumption of rice and wheat in Malaysia, 1960-2003	11
2.6 Worldwide consumption of pesticides	13
2.7 Consumption of pesticides in the Indian scenario	13
2.8 Imidacloprid chemical structure	16
2.9 Imidacloprid	19
2.10 Nicotine	19
2.11 The 4 steps of risk assessment process	28
2.12 Illustration of typical dose response curve	31
3.1 Paddy sampling point in Kampung Sawah Sempadan, Tanjung Karang	35
3.2 Map of location of blocks in Kampung Sawah Sempadan	38
3.3 Health Risk Assessment process in this study	44
3.4 The risk acceptability for non-carcinogenic health effect (USEPA, 2007)	46
4.1 Chromatogram of a standard solution of imidacloprid (0.5 ppm)	50
4.2 Calibration curve of imidacloprid	50
4.3 Occurrences of imidacloprid residue in paddy and rice samples	54
4.4 Non cancer risk (HQ) of imidacloprid in various groups of respondents	56

LIST OF ABBREVIATIONS

a.i.	Active Ingredient
ACN	Acetonitrile
ADI	Acceptable Daily Intake
AOAC	Association Of Official Agricultural Chemists
ARD	Acute Reference Dose
BDL	Below Detection Limit
BERNAS	Beras Nasional
CDC	Centers For Disease Control
CODEX	Codex Alimentarius Commission
DOA	Department Of Agriculture
EDI	Estimated Daily Intake
EFSA	European Food Safety Authority
EU	European Union
FAO	Food And Agriculture Organization
FDA	Food And Drug Administration
FFQ	Food Frequency Questionnaires
GAP	Good Agricultural Practice
GC	Gas Chromatography
GDP	Gross Domestic Product
HQ	Hazard Quotient
IADA	Integrated Agricultural Development Area
IUPAC	International Union Of Pure And Applied Chemistry
kg/ha	Kilogram Per Hectare
LC	Liquid Chromatography
LC ₅₀	Lethal Concentration Of Half Population
LC-MS/MS	Liquid Chromatography-Tandem Mass Spectrometry
LD ₅₀	Lethal Dose
LOAEL	(Lowest Observed Adverse Effect Level)
LOD	Limit Of Detection
LOQ	Limit Of Quantification
MANS	Malaysian Adult Nutrition Survey

mg/kg	Milligram per kilogram
MgSO ₄	Magnesium Sulfate
MOH	Ministry Of Health
MRL	Maximum Residue Limit
nAChR	Nicotinic Acetylcholine Receptors
NaCl	Sodium Chloride
NOAEL	No-Observed Adverse Effect Level
OC	Organochlorines
Organophosphates	(Op)
PPM	Part Per Million
PSA	Primary Secondary Amine
QuEChERS	Quick, Easy, Cheap, Effective, Rugged, And Safe)
RfD	Reference Doses
RSD	Relative Standard Deviation
S/N	Signal-To-Noise
SF	Safety Factors
UF	Uncertainty Factors
UHPLC-UV	Ultra High Performance Liquid Chromatography (UHPLC) With Ultra Violet Detector
USDA	United States Department Of Agriculture
USEPA	United State Environmental Protection Agency
WHO/FAO	World Health Organization/ Food And Agriculture Organization

CHAPTER 1

INTRODUCTION

Rice (*Oryza sativa L.*) by large is staple food consumed and produced by most of the world population ranking third after wheat and maize (Akinbile et al., 2011). China and India yield an approximate of 332 million tonnes of rice amounting to almost 50% and 89% of the world's cereal and harvested rice production respectively (Laborte et al., 2012). In Malaysia, the rice production is in the 25th place with the total production of 2.4 million tonnes. The national average rice production in the country is only 3.0 tons per hectare (Akinbile et al., 2011). Rice are grown mainly in eight granaries of the Peninsular Malaysia making it the most valuable plantation produced nationally after rubber and palm oil. About 300 500 hectares in Malaysia Peninsular is devoted to rice production (Food and Agriculture Organisation, 2002).

About one-third of the agricultural productions are maximized using various types of pesticides (Liu et al., 2002). Chemical use is the most common pest control method currently. According to Sondhia, (2014), two million metric tons of pesticides used by 1.8 billion people engage in agriculture annually onto only 25% of the land area worldwide (Alavanja, 2009). From Figure 1.1, herbicides (47.5%) is a highly used pesticides compared to insecticides (29.5%), fungicides (17.5%) and others (5.5%).




Figure 1.1: Worldwide consumption of pesticides
(Source: De et al., 2013)

Furthermore, the study conducted by Cai, (2008) found that without the usage of agrichemical like pesticides, the production of fruits, vegetables and cereals would reduced to 78%, 54% and 32%, respectively. Zhang et al., (2011) stated four main concerns of pesticide usage around the world: (1) a lot of countries manufacture and utilized illegal harmful pesticides to date; (2) high exposure of pesticides to produces such as rice, cotton and vegetables to fulfill the food demand; (3) lack of enforcement to effectively regulate pesticides application which results in the consumption of unqualified pesticides, and lastly (4) the pesticide residue standard were not implemented effectively. Pesticide residues data in most of the Asian country shows that agriculture products namely cereals, fruits and vegetables are the major dietary source of consumers contaminated with pesticide residues (Wanwimolruk et al, 2012). The increasing rate in food demand from the ever-growing trend of the world population growth can also contribute to the excessive and uncontrollable usage of pesticides in agricultural setting. Human exposure towards pesticide residues through the food, among other things, is of significant concern to everyone from the local authorities, researchers, producers, as well as consumers (Curbelo et al., 2012).

Oryza sativa, the variety of rice that is a staple with highest production worldwide after maize (Harrison et al., 2002) is consumed in particular at the West Indies and Asian regions. In Malaysia, rice is a crucial part of the nation's diet. Although wheat and livestock are slowly gaining influence in Malaysian food consumption pattern, rural people especially still favors rice. Per capita consumption of rice constituted about 11 percent (82 kilograms) nationwide over the total estimation around 2.3 million tons annually (Shamsudin et al., 2008). The use of pesticides is essential in crop protection during growth, storage and transportation which leads to the improvement in agricultural activities in twentieth century (Mohan et al., 2010). In consequence, food was contaminated extensively due to the arbitrary and imprudent use of pesticides that have begun to receive much attention from many organizations because these residues are found in food commodities that may be hazardous to human health (Jayashree and Vasudevan, 2007).




Figure 1.2: Imidacloprid – chemical structure

Imidacloprid, 1-(6-chloro-3-pyridinyl) methyl)-N-nitro-2-Imidazolidinimine), with Malaysia commercial name Confidor is a systemic insecticide and categorized under neonicotinoids (Figure 1.2). Neonicotinoids (a class of chemicals) insecticides are human-made derived from an alkaloid component, nicotine occurring naturally in tobacco plant (Tomizawa and Casida, 2005). Worldwide, it is used in massive amount as insecticides. It acts differently from the current pesticides by adhering to nicotinic receptors as an agonist on the postsynaptic membrane (Fishel, 2005). Therefore, disturbing the stimulus delivery in the insect's body mechanism and obstructing the above-mentioned neuronal tract in specific. Acetylcholine, a crucial neurotransmitter accumulates leading to paralysis and eventually death of the pest's. It works through the exposure by touch as well as via stomach activity. Imidacloprid is classified as a toxicity class II, which from the scale of I to IV, I is being classified as the top-level, by the Environmental Protection Agency (EPA) and labeled with the "Warning" signal word. Expected symptoms after high level exposure are lethargy, muscle twitch, cramps and weakness. The Lethal Dose (LD₅₀) is 131 and 450mg/kg for mice and rats, respectively. Airborne concentration of more than 69 and 5323 mg/meters cubed air in aerosol and dust forms respectively are fatal to half of the tested animal (LC₅₀). As supported by study done by Fishel (2005) that imidacloprid has been considered as safe and non-sensitive to eyes and skin.

In farming, imidacloprid is commonly utilized for insect's monitoring. It is powerful for managing dangerous pest species including aphids, whiteflies, thrips, scales, psyllids, plant bugs, leafhoppers, planthoppers. A study done by Jaun, (2008) reported that 50.7% of paddy farmers in (Department of Irrigation and Drainage Malaysia) IADA Barat Laut Kuala Selangor including Kampung Sawah Sempadan is using insecticide imidacloprid (Confidor) as one of their pesticides mixture in spraying activities; also lambda-cylothrins (Karate), Lufenuron (Match) and Chlorpyrifos + cypermethrin (Nurelle) with 45.8%, 43.8% and 33.3% concentration respectively.

In this study, it is more focusing on insecticide residue namely Imidacloprid in paddy and rice, planted in Selangor Darul Ehsan. Monitoring of imidacloprid residue in paddy and rice sample was combined with data of rice consumption of Tanjung Karang residential to estimate the potential non-carcinogenic health risk.

1.2 Problem Statements

Although pesticides is effective in crop protection, its usage affect the environment as well as consumers in the case of the presence of pesticides residues in edible rice products (Darko and Akoto, 2008). Exposure to pesticides can always occur during the paddy cultivation process. Basically, some of the pesticides are used to keep pests from attacking food in order to preserve the crops. These compounds have been shown to be toxic to both humans and environment because they are hormone mimickers which are carcinogenic, mutagenic, teratogenic and besides having the ability to persist in the soil (Hamilton et al., 2004). Depending on their chemical characteristics, these compounds may migrate into the grain particularly rice.

The contamination of rice by chemical hazards like pesticides residues has become an alarming worldwide public health concern as people demand for food that are high quality, nutritious and safe for consumption, high-quality. It is indispensable to control the use of pesticides and minimize contamination because of its high consumption by

people (Munshi et al., 2011). According to US EPA findings, pesticides residue should only be present on the outer layer and eliminated during grain milling, mostly in by products such as bran and husk. However, a number of prior studies have been executed analyzing the distribution of contaminants in irrigated rice milling, especially in the inner part of this grain (Kapoor et al., 2012; Chen et al., 2009; Kaushik et al., 2009).

Monitoring of pesticide contamination in food products such as rice is very important. Being an essential alimentary source throughout the world, the quality of rice should not be compromised by the presence of any chemical contamination like pesticide residues. Sometimes, contamination in the rice is due to illegal or non-scientific use of pesticide at the time of early cultivation process until the harvest. Pesticide residues although in negligible amount or in traces are harmful to human (Munshi et al., 2011).

1.3 Study Justification


Basically, an estimated half of the human population represented by almost 3.0 billion Asians feed on the rice. Thus this daily consumption which provides up to 80 percent calories needs to be of healthy and standard quality. According to the Malaysian Dietary Guidelines, National Coordinating Committee on Food and Nutrition (2010), it states that we, the Asian population must to take 6 to 11 servings a day from carbohydrate sources such as in the bread, cereal, rice and pasta group. Recently, through the Malaysian Adult Nutrition Survey (MANS) in 2012, it is reported that 97% of the nation have an average of 2½ plates of *nasi putih* (cooked rice) twice daily (Norimah et al., 2008).

It clearly shows that most of the Malaysian population consumes rice in their daily life. Thus, this study is expected to prove that our rice contain permissible level of pesticide residue and do not pose any risk to the consumer. The health risk assessments were evaluated through the Hazard Quotient (HQ) for non carcinogenic risk.

1.4 Study contribution

There are still a limited number of studies referring to neonicotinoids in terms of potential hazard to human health especially in Malaysia. Therefore, this dissertation is aimed to provide data and fill the knowledge gap about the pesticides residue level in Selangor's rice production. Furthermore, its impact on human health in term of non carcinogenic risks for long term consumption of local rice would be estimated through the study. Nonetheless, the residue data is only focused on imidacloprid insecticides as the popular usage among paddy farmers. In addition, findings from this study would also provide a baseline value for pesticide contamination in our rice production and its impacts to human health. This type of data also can be used as the recommendations for the responsible management or authorities in order to control the usage of pesticides in the field. Moreover, in Malaysia, there is no data of health risk assessment on imidacloprid residue in paddy and rice yet. In view of this, this study has contributed the publication of this issue in order to understand more on health risk assessment through the local rice consumption among three different populations.

1.5 Conceptual Framework



1.6 Definition of Term

1.6.1 Conceptual definition

Imidacloprid

A pesticide is any substance or mixture of substances intended for preventing, destroying, repelling, or mitigating any pest. Pests can be insects and insect-like organisms, mice and other vertebrate animals, unwanted plants (weeds), or fungi, bacteria and viruses that cause plant diseases. Though often misunderstood to refer only to insecticides, the term pesticide also applies to herbicides, fungicides, and various other substances used to control pests (Lehotay et al., 2005).

Method validation

The process used to confirm the analytical procedure employed for a specific test is suitable for its intended use. Results from method validation can be used to judge the quality, reliability and consistency of analytical results; it is an integral part of any good analytical practice (Kalra, 2011).

Health Risk Assessment

According to the Report of a Joint FAO/WHO Consultation (2005), health risk assessment through dietary intake was calculated using food consumption data on the concentration of chemical residue found in food. The resulting dietary exposure estimate was then compared with the relevant toxicological or nutritional reference value for the food chemical of concern.

1.6.2 Operational definition

Imidacloprid

Imidacloprid insecticide came from neonicotinoid class found in the paddy, brown and white rice samples has been selected to be analyzed in this study is extracted using QuEChERS method and analyzed through UHPLC-UV.

Method Validation

Method validation is a process that verifies if an analytical procedure matches the laboratory and instrument condition for the intended analysis. To ensure data credibility in the quantitative residue analyses, sample preparation was thoroughly validated considering parameters like recovery, precision (relative standard deviation), determination coefficient (R^2), linearity, detection limit and quantification limit. In this study, method was validated following the established European Union guidelines on quality control procedures for pesticide analysis (European Union, 2007).

Health Risk Assessment

In this study, the risk assessment through dietary intake was calculated on the imidacloprid exposure through rice consumption. The assessment was evaluated

through the Hazard Quotient (HQ) for non carcinogenic effects to estimate the health risk of those who were exposed to imidacloprid from rice ingestion in their daily meal.

1.7 Research Objectives

1.7.1 General Objective

To determine the imidacloprid residue concentration in paddy and rice as well as the Dietary Health Risk Assessment among the Tanjung Karang's population, Selangor, Malaysia.

1.7.2 Specific Objectives

- 1) To optimize and validate the QuEChERS method prior the analysis of sample using UHPLC - UV.
- 2) To determine and compare the concentration of imidacloprid residue in samples from 3 different stages of rice processes (paddy, brown rice and white rice).
- 3) To determine the rice consumption pattern (ingestion rate) among three different age group (children, adolescent and adult).
- 4) To estimate and compare the Estimated Daily Intake (EDI) of imidacloprid residue through the white rice consumption among the three groups of Tanjung Karang's population
- 5) To determine and compare the Hazard Quotient (HQ) for imidacloprid residue in white rice consumptions among the three groups of Tanjung Karang's population

1.7.3 Hypothesis

- 1) There is a significant difference between the imidacloprid residue concentration in three different types of rice samples
- 2) There is a significant difference in the Estimated Daily Intake (EDI) for imidacloprid residue through the white rice consumption among the three groups of Tanjung Karang's population
- 3) There is a significant difference in Hazard Quotient (HQ) for imidacloprid residue in white rice sample between the three groups of Tanjung Karang population.

REFERENCES

- Akinbile, C. O., El-Latif, K. M., Abdullah, R., and Yusoff, M.S. (2011). Rice Production and Water use Efficiency for Self-Sufficiency in Malaysia: A Review. *Trends in Applied Sciences Research*, 6: 1127-1140.
- Akoijam, R., and Singh, B. (2014). Persistence and Metabolism of Imidacloprid in Rice. *Bulletin of Environmental Contamination and Toxicology*, 92: 609-615.
- Akoijam, R., Singh, B., and Mandal, K. (2014). Development and Validation of a Quick, Easy, Cheap, Effective, Rugged and Safe Method for the Determination of Imidacloprid and Its Metabolites in Soil. *Journal of chromatographic science*, 53: 542-547.
- Alavanja, M. C., and Bonner, M. R. (2012). Occupational Pesticide Exposures and Cancer Risk: A Review. *Journal of Toxicology and Environmental Health, Part B*, 15: 238-263.
- Alavanja, M.C. (2009). Introduction: Pesticides Use and Exposure, Extensive Worldwide. *Reviews on Environmental Health*, 24:303-310.
- Al-Rimawi, F. (2014). A HPLC-UV Method for Determination of Three Pesticides in Water. *International Journal of Advances in Chemistry*, 2: 1-8.
- Arredondo, M., Martinez, R., Nunez, M. T., Ruz, M. and Olivares, M. (2006). Inhibition of iron and copper uptake by iron, copper and zinc. *Biological Research*, 39: 95-102.
- Anastassiades, M., Lehotay, S.J., Štajnbaher, D., and Schenck, F.J. (2003). Fast and Easy Multiresidue Method Employing Acetonitrile Extraction/Partitioning and Dispersive Solid-Phase Extraction for the Determination of Pesticide Residues in Produce. *Journal AOAC International*, 86: 412-43.
- Anatra-Cordone, M., and Durkin, P. (2005). *Imidacloprid-Human Health and Ecological Risk Assessment-Final Report*. US Department of Agriculture, Arlington, VA.
- Anyusheva, M., Lamers, M., La, N., Nguyen, V. V., and Streck, T. (2012). Fate of Pesticides in Combined Paddy Rice–Fish Pond Farming Systems In Northern Vietnam. *Journal of Environmental Quality*, 41: 515-525.
- Aponso, G. L. M., Manuweera, G. K., Anderson, K., and Tinsley, I. J. (2002). Exposure and Risk Assessment for Farmers Occupationally Exposed to Chlorpyrifos. *Annals of the Sri Lanka Department of Agriculture*, 4: 233-244.
- Bagri, P., Kumar, V., and Sikka, A. K. (2014). An *In Vivo* Assay of The Mutagenic Potential of Imidacloprid Using Sperm Head Abnormality Test and Dominant Lethal Test. *Drug and Chemical Toxicology*, (38): 342-348.

- Banerjee, K., Oulkar, D. P., Dasgupta, S., Patil, S. B., Patil, S. H., Savant, R., and Adsule, P. G. (2007). Validation and Uncertainty Analysis of A Multi-Residue Method For Pesticides in Grapes using Ethyl Acetate Extraction and Liquid Chromatography–Tandem Mass Spectrometry. *Journal of Chromatography A*, 1173: 98-109.
- Barbash, J.E. (2006). *The Geochemistry of Pesticides US Geological Survey, Tacoma*. US Geological Survey, Tacoma, WA, USA.
- Bhardwaj, S., Srivastava, M. K., Kapoor, U. and Srivastava, L. P. (2010). A 90 Days Oral Toxicity of Imidacloprid in Female Rats: Morphological, Biochemical and Histopathological Evaluations. *Food and Chemical Toxicology*, 48:1185-1190.
- Blasco, C., Font, G., and Pico, Y. (2002). Comparison of Microextraction Procedures to Determine Pesticides in Oranges by Liquid Chromatography–Mass Spectrometry. *Journal of Chromatography A*, 970: 201-212.
- Boobis, A. R., Ossendorp, B. C., Banasiak, U., Hamey, P. Y., Sebestyen, I., and Moretto, A. (2008). Cumulative Risk Assessment of Pesticide Residues in Food. *Toxicology Letters*, 180: 137-150.
- Boobis, A. R., Ossendorp, B. C., Banasiak, U., Hamey, P. Y., Sebestyen, I., and Moretto, A. (2008). Cumulative Risk Assessment of Pesticide Residues in Food. *Toxicology Letters*, 180: 137-150.
- Cai, D.W. (2008). Understand The Role of Chemical Pesticides and Prevent Misuses of Pesticides. *Bulletin of Agriculture Science Technology*, 1: 36-38.
- Carrere, M., Demaria, F., and Drogue, S. (2014). *Are consumers over-protected? A world picture of regulations on pesticide MRL in fruits and vegetables* (No. 264683).
- Cattani, M., Cena, K., Edwards, J., and Pisaniello, D. (2001). Potential Dermal and Inhalation Exposure to Chlorpyrifos in Australian Pesticide Workers. *Annals of Occupational Hygiene*, 45: 299-308.
- Champagne, E.T., Wood, D.F., Juliano, B.O., and Bechtel, D.B. (2004). The Rice Grain and Its Gross Composition. *Rice Chemistry and Technology*, 3:77-100.
- Chen, C., Li, Y., Chen, M., Chen, Z., and Qian, Y. (2009). Organophosphorus Pesticide Residues in Milled Rice (*Oryza Sativa*) on the Chinese Market and Dietary Risk Assessment. *Food Additives and Contaminants*, 26: 340-347.
- Chen, C., Qian, Y., Chen, Q., Tao, C., Li, C., and Li, Y. (2011). Evaluation of Pesticide Residues in Fruits and Vegetables From Xiamen, China. *Food Control*, 22: 1114-1120.

- Chen, L., and Li, B. (2012). Determination of Imidacloprid in Rice By Molecularly Imprinted-Matrix Solid-Phase Dispersion With Liquid Chromatography Tandem Mass Spectrometry. *Journal of Chromatography B*, 897: 32-36.
- Chourasiya, S., Khillare, P. S., and Jyethi, D. S. (2015). Health Risk Assessment of Organochlorine Pesticide Exposure through Dietary Intake of Vegetables Grown In the Periurban Sites of Delhi, India. *Environmental Science and Pollution Research*, 22: 5793-5806.
- Connell, D. W. (2005). *Basic Concepts of Environmental Chemistry*. CRC Press, Taylor and Francis Group.
- Cox, C. (2001). Insecticide Factsheet: Imidacloprid. *Journal of Pesticide Reform*, 21: 15-21.
- Darko, G., and Akoto, O. (2008). Dietary Intake of Organophosphorus Pesticide Residues through Vegetables from Kumasi, Ghana. *Food and Chemical Toxicology*, 46: 3703-3706
- De, A., Bose, R., Kumar, A., and Mozumdar, S. (2013). *Targeted Delivery of Pesticides Using Biodegradable Polymeric Nanoparticles*. Springer, India, Private.
- Debby Christy Jaun (2008). *Intergrated Pest Management's Practice among Paddy Farmers in IADA Barat Laut Selangor, Tanjung Karang, Selangor*. Degree Thesis. Universiti Putra Malaysia.
- Deka, S.C. and Barman, N. (2004). Pesticide Residues in Unpolished, Polished and Parboiled Rice in Assam, India. *Pesticide Residue Journal*, 16: 71-74.
- Department of Agriculture, (2011). List of Registered Pesticides. Retrieved on 2 December 2014 at <http://www.doa.gov.my/senarai-racun-makhluk-perosak-berdaftar>.
- Di Muccio, A., Fidente, P., Barbini, D. A., Dommarco, R., Seccia, S., and Morrica, P. (2006). Application of Solid-Phase Extraction and Liquid Chromatography–Mass Spectrometry to the Determination of Neonicotinoid Pesticide Residues in Fruit and Vegetables. *Journal of Chromatography A*, 1108: 1-6.
- Donnarumma, L., Pulcini, P., Pochi, D., Rosati, S., Lusco, L., and Conte, E. (2011). Preliminary Study on Persistence in Soil and Residues in Maize of Imidacloprid. *Journal of Environmental Science and Health, Part B*, 46: 469-472.
- Dors, G.C. E.G. Primel, C.A. Fagundes, C.H. Mariot, and Badiale-Furlong, E. (2011). Distribution of Pesticide Residues in Rice Grain and in its Co-products. *Journal Brazilian Chemistry Society*, 22: 1921-1930.

- Duzguner, V., and Erdogan, S. (2010). Acute Oxidant and Inflammatory Effects of Imidacloprid on the Mammalian Central Nervous System and Liver in Rats. *Pesticide Biochemistry and Physiology*, 97: 13-18.
- EFSA PPR Panel (EFSA Panel on Plant Protection Products and their Residues), (2013). Scientific Opinion on the developmental neurotoxicity potential of acetamiprid and imidacloprid. *EFSA Journal*, 11: 3471.
- Elbert, A., Haas, M., Springer, B., Thielert, W., and Nauen, R. (2008). Applied Aspects of Neonicotinoid Uses in Crop Protection. *Pest Management Science*, 64: 1099-1105.
- Faustman, E. M., and Omenn, G. S. (2001). *Risk Assessment*. Cassarett and Doull's Toxicology (Klaassen C, ed). 6th ed. San Francisco, CA: McGraw-Hill, 83-104.
- Fernández-Alba, A. R., Tejedor, A., Agüera, A., Contreras, M., and Garrido, J. (2000). Determination of Imidacloprid and Benzimidazole Residues in Fruits and Vegetables by Liquid Chromatography–Mass Spectrometry after Ethyl Acetate Multiresidue Extraction. *Journal of AOAC International*, 83: 748-755.
- Fidente, P., Seccia, S., Vanni, F., and Morrica, P. (2005). Analysis of Nicotinoid Insecticides Residues in Honey by Solid Matrix Partition Clean-Up and Liquid Chromatography–Electrospray Mass Spectrometry. *Journal of Chromatography A*, 1094: 175-178.
- Fishel, F. M. (2005). *Pesticide Toxicity Profile: Neonicotinoid Pesticides*. University of Florida, IFAS.
- Fossen, M. (2006). *Environmental Fate of Imidacloprid*. California Department of Pesticide Regulation, 1-16: 2006.
- Fuad, M., Junaidi, A. B., Habibah, A., Hamzah, J., Toriman, M.E., Lyndon and A.M. Azima (2012). The Impact of Pesticides on Paddy Farmers and Ecosystem. *Advances in Natural and Applied Sciences*, 6: 65-70.
- Gao, N., Guo, X., Zhang, K., and Hu, D. (2014). High-Performance Liquid Chromatography and Gas Chromatography-Mass Spectrometry Methods for the Determination of Imidacloprid, Chlorpyrifos, and Bifenthrin Residues in Tea Leaves. *Instrumentation Science and Technology*, 42: 267-277.
- Gels, J. A., Held, D. W., and Potter, D. A. (2002). Hazards of insecticides to the bumble bees *Bombus impatiens* (Hymenoptera: Apidae) foraging on flowering white clover in turf. *Journal of Economic Entomology*. 95(4):722-728.
- Gerd, W.E and Piao, Y. (2005). *Regional Overview, Pesticides Policy and Monitoring Guidelines*. Proceedings of the Asia Regional Workshop on the Implementation, Monitoring and Observance International Code of Conduct on the Distribution and Use of Pesticides. Regional Office for Asia and the Pacific.

- González-Curbelo, M. Á., Herrera-Herrera, A. V., Ravelo-Pérez, L. M., and Hernández-Borges, J. (2012). Sample Preparation Methods for Pesticide-Residue Analysis in Cereals and Derivatives. *Trends in Analytical Chemistry*, 38: 32-51.
- Goulson, D. (2013). Review: An Overview of the Environmental Risks Posed by Neonicotinoid Insecticides. *Journal of Applied Ecology*, 50: 977-987.
- Gupta, P. K. (2004). Pesticide Exposure-Indian Scene. *Toxicology*, 198: 83-90.
- Gupta, V. K., Ali, I., and Saini, V. K. (2006). Adsorption of 2, 4-D and Carbofuran Pesticides using Fertilizer and Steel Industry Wastes. *Journal of Colloid and Interface Science*, 299: 556-563.
- Guzsvány, V., Madžgalj, A., Trebše, P., Gaál, F., and Franko, M. (2007). Determination of Selected Neonicotinoid Insecticides by Liquid Chromatography with Thermal Lens Spectrometric Detection. *Environmental Chemistry Letters*, 5: 203-208.
- Hamidin, N., Yu, Q. J., and Connell, D. W. (2008). Human Health Risk Assessment of Chlorinated Disinfection By-Products in Drinking Water Using A Probabilistic Approach. *Water Research*, 42: 3263-3274.
- Harrison, P., Bruinsma, J., de Haen, H., Alexandratos, N., Schmidhuber, J., Bödeker, G., and Ottaviani, M. G. (2002). World Agriculture: Towards 2015/2030. Online, <http://www.fao.org/documents>.
- Hashmi, I., and Khan, A. D. (2011). Adverse Health Effects of Pesticide Exposure in Agricultural and Industrial Workers of Developing Country. *Pesticides-The Impacts of Pesticides Exposure, In Tech*. <http://dx.doi.org/10.5772/13835>.
- Hill, R. D. (2012). *Rice in Malaya: A Study in Historical Geography*. NUS Press.
- Hill, R. D. (2013). *Agriculture in The Malaysian Region*. NUS Press.
- Hines, C. J., and Deddens, J. A. (2001). Determinants of Chlorpyrifos Exposures and Urinary 3, 5, 6-Trichloro-2-Pyridinol Levels Among Termiticide Applicators. *Annals of Occupational Hygiene*, 45: 309-321.
- Iqbal, S., Uddin, R., Saied, S., Ahmed, M., Abbas, M., and Aman, S. (2012). Extraction, Cleanup, and Chromatographic Determination of Imidacloprid Residues in Wheat. *Bulletin Of Environmental Contamination And Toxicology*, 88: 555-558.
- Ishii, Y., Kobori, I., Araki, Y., Kuroguchi, S., Iwaya, K., and Kagabu, S. (1994). HPLC Determination of the New Insecticide Imidacloprid and its Behavior in Rice and Cucumber. *Journal of Agricultural and Food Chemistry*, 42: 2917-2921.

- Jansson, C., Pihlström, T., Österdahl, B. G., and Markides, K. E. (2004). A New Multi-Residue Method for Analysis of Pesticide Residues in Fruit and Vegetables Using Liquid Chromatography with Tandem Mass Spectrometric Detection. *Journal of Chromatography A*, 1023: 93-104.
- Jayashree, R., and Vasudevan, N. (2007). Organochlorine Pesticide Residues in Ground Water of Thiruvallur District, India. *Environmental Monitoring and Assessment*, 128: 209-215.
- Jeschke, P., Nauen, R., and Beck, M. E. (2013). Nicotinic Acetylcholine Receptor Agonists: A Milestone for Modern Crop Protection. *Angewandte Chemie International Edition*, 52: 9464-9485.
- Jeschke, P., Nauen, R., Schindler, M., and Elbert, A. (2010). Overview of The Status and Global Strategy for Neonicotinoids. *Journal of Agricultural and Food Chemistry*, 59: 2897-2908.
- Juraske, R., Mutel, C. L., Stoessel, F., and Hellweg, S. (2009). Life Cycle Human Toxicity Assessment of Pesticides: Comparing Fruit and Vegetable Diets in Switzerland and The United States. *Chemosphere*, 77: 939-945.
- Kalra, K. (2011). *Method Development and Validation of Analytical Procedures*. Quality Control of Herbal Medicines and Related Areas, InTech.
- Kamel, F., and Hoppin, J. A. (2004). Association of Pesticide Exposure with Neurologic Dysfunction and Disease. *Environmental Health Perspectives*, 112: 950-958.
- Kanrar, B., Ghosh, T., Pramanik, S. K., Dutta, S., Bhattacharyya, A., and Dhuri, A. V. (2006). Degradation Dynamics and Persistence of Imidacloprid in a Rice Ecosystem Under West Bengal Climatic Conditions. *Bulletin of Environmental Contamination and Toxicology*, 77: 631-637.
- Kapoor, U., Srivastava, M. K., Srivastava, A. K., Patel, D. K., Garg, V., and Srivastava, L. P. (2013). Analysis of Imidacloprid Residues in Fruits, Vegetables, Cereals, Fruit Juices, and Baby Foods, and Daily Intake Estimation in and Around Lucknow, India. *Environmental Toxicology and Chemistry*, 32: 723-727.
- Kaushik, G., Satya, S., and Naik, S. N. (2009). Food Processing a Tool to Pesticide Residue Dissipation-A Review. *Food Research International*, 42: 26-40.
- Kimura-Kuroda, J., Komuta, Y., Kuroda, Y., Hayashi, M., and Kawano, H. (2012). Nicotine-Like Effects of the Neonicotinoid Insecticides Acetamiprid and Imidacloprid on Cerebellar Neurons from Neonatal Rats. *PLoS One*, 7: e32432.
- Köck, M., Farré, M., Martínez, E., Gajda-Schranz, K., Ginebreda, A., Navarro, A., and Barceló, D. (2010). Integrated Ecotoxicological and Chemical Approach for the Assessment of Pesticide Pollution in the Ebro River Delta (Spain). *Journal of Hydrology*, 383: 73-82.

- Koesukwiwat, U., Lehotay, S. J., Miao, S., and Leepipatpiboon, N. (2010). High Throughput Analysis of 150 Pesticides in Fruits and Vegetables Using Quechers and Low-Pressure Gas Chromatography–Time-Of-Flight Mass Spectrometry. *Journal of Chromatography A*, 1217: 6692-6703.
- Krueve, A., Künnapas, A., Herodes, K., and Leito, I. (2008). Matrix Effects in Pesticide Multi-Residue Analysis by Liquid Chromatography–Mass Spectrometry. *Journal of Chromatography A*, 1187: 58-66.
- Kumar, A., Verma, A., and Kumar, A. (2013). Accidental Human Poisoning with a Neonicotinoid Insecticide, Imidacloprid: A Rare Case Report From Rural India With A Brief Review of Literature. *Egyptian Journal of Forensic Sciences*, 3: 123-126.
- Kwadwo, A.O., Davis, K., Aredo D. (2008). Advancing Agriculture in Developing Countries Through Knowledge and Innovation. International Food Policy Institute. Available at: <http://www.ifpri.org/sites/default/files/publications/oc59.pdf>.
- Laborte, A. G., de Bie, K. C., Smaling, E., Moya, P. F., Boling, A. A., and Van Ittersum, M. K. (2012). Rice Yields and Yield Gaps in Southeast Asia: Past Trends and Future Outlook. *European Journal of Agronomy*, 36: 9-20.
- Land Areas of Major Crops in Malaysia, (2012). ICID – Irrigation and Drainage in the World – A Global Review. Retrieved at http://www.icid.org/i_d_malaysia.pdf
- Laycock, I., Lenthall, K. M., Barratt, A. T., and Cresswell, J. E. (2012). Effects of Imidacloprid, A Neonicotinoid Pesticide, on Reproduction in Worker Bumble Bees (*Bombus Terrestris*). *Ecotoxicology*, 21: 1937-1945.
- Lehotay, S. J. (2011). QuEChERS Sample Preparation Approach for Mass Spectrometric Analysis of Pesticide Residues in Foods. *Mass Spectrometry in Food Safety*, 747: 65-91.
- Lehotay, S. J., Maštovská, K., and Yun, S. J. (2005). Evaluation of Two Fast and Easy Methods for Pesticide Residue Analysis in Fatty Food Matrixes. *Journal of AOAC International*, 88: 630-638.
- Lehotay, S. J., Son, K. A., Kwon, H., Koesukwiwat, U., Fu, W., Mastovska, K., and Leepipatpiboon, N. (2010). Comparison of QuEChERS Sample Preparation Methods for the Analysis of Pesticide Residues in Fruits and Vegetables. *Journal of Chromatography A*, 1217: 2548-2560.
- Liu, C. J., Men, W. J., and Liu, Y.J. (2002). The Pollution of Pesticides in Soils and Its Bioremediation. *System Sciences and Comprehensive Studies in Agriculture*, 18: 295-297.

- Liu, S., Zheng, Z., Wei, F., Ren, Y., Gui, W., Wu, H., and Zhu, G. (2010). Simultaneous Determination of Seven Neonicotinoid Pesticide Residues In Food By Ultra Performance Liquid Chromatography Tandem Mass Spectrometry. *Journal of Agricultural and Food Chemistry*, 58: 3271-3278.
- Liu, Y., Pan, X., and Li, J. (2015). A 1961–2010 Record of Fertilizer Use, Pesticide Application and Cereal Yields: A Review. *Agronomy for Sustainable Development*, 35: 83-93.
- Matsumura, F. (2012). *Toxicology of Insecticides*. Springer Science and Business Media.
- Miró, G., Gálvez, R., Mateo, M., Montoya, A., Descalzo, M. A., and Molina, R. (2007). Evaluation of the Efficacy of a Topically Administered Combination of Imidacloprid and Permethrin against *Phlebotomus Perniciosus* in Dog. *Veterinary Parasitology*, 143: 375-379.
- Mohamed, F., Gawarammana, I., Robertson, T. A., Roberts, M. S., Palangasinghe, C., Zawahir, S., and Roberts, D. M. (2009). Acute Human Self-Poisoning With Imidacloprid Compound: A Neonicotinoid Insecticide. *PLoS One*, 4:5127.
- Mohan, C., Kumar, Y., Madan, J., and Saxena, N. (2010). Multiresidue Analysis of Neonicotinoids By Solid-Phase Extraction Technique Using High-Performance Liquid Chromatography. *Environmental Monitoring and Assessment*, 165: 573-576.
- Mohd Shariff, Z., Mohd Yusof, S., and MY, K. Z. (2008). Energy and Nutrient Intakes: Findings from the Malaysian Adult Nutrition Survey (MANS). *Malaysian Journal of Nutrition*, 14: 1-24.
- Moniruzzaman, M., Chowdhury, M. A. Z., Rahman, M. A., Sulaiman, S. A., and Gan, S. H. (2014). Determination of Mineral, Trace Element, and Pesticide Levels in Honey Samples Originating from Different Regions of Malaysia Compared to Manuka Honey. *Biomed Research International*, 1: 1-10.
- Munshi, A. B., Siddiqi, H. A., Ansari, F. A., Begum, R., and Usmani, T. H. (2011). Multiresidue Analysis of Pesticides by Gas Chromatography with Electron-capture in Rice Samples from Different Geographical Regions of Pakistan. *Pakistan Journal of Nutrition*, 10: 851-859.
- National Coordinating Committee on Food and Nutrition (NCCFFN) (2010). *Recommended Nutrient Intakes for Malaysia*. Ministry of Health Malaysia, Kuala Lumpur.
- Nauen, R., and Denholm, I. (2005). Resistance of Insect Pests to Neonicotinoid Insecticides: Current Status and Future Prospects. *Archives of Insect Biochemistry and Physiology*, 58: 200-215.
- Newman, W.L. (1997). *Social Research Methods: Qualitative and Quantitative Approaches*. Needham Heights: Allyn and Bacon.

- Nolan, R. J., Rick, D. L., Freshour, N. L., and Saunders, J. H. (1984). Chlorpyrifos: Pharmacokinetics in Human Volunteers. *Toxicology and Applied Pharmacology*, 73: 8-15.
- Norimah, J., Safiah, A. K., Jamal, M., Haslinda, K., Zuhaida S., Rohida, H., and Azmi, M. Y. (2008). Food Consumption Patterns: Findings from the Malaysian Adult Nutrition Survey (MANS). *Malaysian Journal of Nutrition*, 14: 25-39.
- Ntow, W. J., Gijzen, H. J., Kelderman, P., and Drechsel, P. (2006). Farmer Perceptions and Pesticide Use Practices in Vegetable Production in Ghana. *Pest Management Science*, 62: 356-365.
- Obana, H., Okimashi, M., Akutsu, K., Kitagawa, Y., and Hori, S. (2003). Determination of Neonicotinoid Pesticide Residues in Vegetables and Fruits with Solid Phase Extraction and Liquid Chromatography Mass Spectrometry. *Journal of Agricultural and Food Chemistry*, 51: 2501-2505.
- Ong, H. C., Mahlia, T. M. I., and Masjuki, H. H. (2011). A Review on Energy Scenario and Sustainable Energy in Malaysia. *Renewable and Sustainable Energy Reviews*, 15: 639-647.
- Pareja, L., Colazzo, M., Pérez-Parada, A., Besil, N., Heinzen, H., Böcking and Fernández-Alba, A.R. (2012). Occurrence and Distribution Study of Residues from Pesticides Applied Under Controlled Conditions in The Field During Rice Processing. *Journal of Agricultural and Food Chemistry*, 60: 4440-4448.
- Pareja, L., Fernández-Alba, A.R., Cesio, V., and H. Heinzen. (2011). Analytical Methods for Pesticide Residues in Rice. *Trends in Analytical Chemistry*, 30: 270-291.
- Payá, P., Anastassiades, M., Mack, D., Sigalova, I., Tasdelen, B., Oliva, J., and Barba, A. (2007). Analysis of Pesticide Residues using the Quick Easy Cheap Effective Rugged and Safe (QuEChERS) Pesticide Multiresidue Method in Combination with Gas and Liquid Chromatography and Tandem Mass Spectrometric Detection. *Analytical and Bioanalytical Chemistry*, 389: 1697-1714.
- Pimentel, D. (2009a). *Environmental and Economic Costs of the Application of Pesticides Primarily in the United States*. In Integrated pest management: innovation-development process (pp 89-111), Springer Netherlands.
- Poh, B. K., Ng, B. K., Siti Haslinda, M. D., Nik Shanita, S., Wong, J. E., Budin, S. B., and Norimah, A. K. (2013). Nutritional status and dietary intakes of children aged 6 months to 12 years: findings of the Nutrition Survey of Malaysian Children (SEANUTS Malaysia). *British Journal of Nutrition*, 110: S21-S35.
- Pohanish, R. P. (2014). *Sittig's Handbook of Pesticides and Agricultural Chemicals*. William Andrew.

- Pous, X., Ruíz, M., Picó, Y., and Font, G. (2001). Determination of imidacloprid, metalaxyl, myclobutanil, prothiofos, and thiamethoxam in fruits and vegetables by liquid chromatography–atmospheric pressure chemical ionization–mass spectrometry. *Fresenius' Journal of Analytical Chemistry*, 371: 182-189.
- Rees, N., and Watson, D. (2000). *International Standards for Food Safety*. Springer Science and Business Media.
- Reza, M. A., Firdaus, M., and Novianti, T. (2012) Analyses of Food Price Stabilization in Indonesia. *ASEAN Journal of Economics, Management and Accounting* 2, 1&2: 48-63.
- Sahoo, S.K., Chahil, G.S., Mandal, K., Battu, R.S. and Singh, B. (2012). Estimation of β -Cyfluthrin and Imidacloprid in Okra Fruits and Soil by Chromatography Techniques. *Journal of Environmental Science and Health, Part B: Pesticides, Food Contaminants, and Agricultural Wastes*, 47: 42-50.
- Saka, M., Iijima, K., Nishida, M., Koma, Y., Hasegawa, N., Sato, K., and Kato, Y. (2008). Effects of Processing and Cooking On the Levels of Pesticide Residues in Rice Samples. *Journal of the Food Hygienic Society of Japan*, 49: 141-149.
- Schoning, R., and Schmuck, R. (2003). Analytical Determination of Imidacloprid and Relevant Metabolite Residues by LC MS/MS. *Bulletin of Insectology*, 56: 41-50.
- Schwedler, D. A., Thomas, A. D., and Yeh, L. T. (2000). Determination of Spinosad and Its Metabolites in Food and Environmental Matrices. Liquid Chromatography-Mass Spectrometry. *Journal of Agricultural and Food Chemistry*, 48: 5138-5145.
- Seccia, S., Albrizio, S., Fidente, P., and Montesano, D. (2011). Development and Validation of a Solid-Phase Extraction Method Coupled To High-Performance Liquid Chromatography With Ultraviolet-Diode Array Detection For The Determination Of Sulfonylurea Herbicide Residues In Bovine Milk Samples. *Journal of Chromatography A*, 1218: 1253-1259.
- Shamsudin, M. N., Mohamed, Z., Abdullah, A. M., and Radam, A. (2008). *Demand Analyses of Rice in Malaysia*. University Library of Munich, Germany.
- Sondhia, S. (2014). Herbicides Residues in Soil, Water, Plants and Non-Targeted Organisms and Human Health Implications: An Indian Perspective. *Indian Journal of Weed Science*, 46: 66-85.
- Soon, K., and Wee, K. (2011). *Demand and Supply of Rice in Malaysia*. Available at SSRN 1882764.

- Štajnbaher, D., and Zupančič-Kralj, L. (2003). Multiresidue Method for Determination of 90 Pesticides in Fresh Fruits and Vegetables Using Solid-Phase Extraction and Gas Chromatography-Mass Spectrometry. *Journal of Chromatography A*, 1015: 185-198.
- Sudo, N., Sekiyama, M., Watanabe, C., Mozammel Haque Bokul, A. T. M., and Ohtsuka, R. (2004). Gender Differences in Food and Energy Intake Among Adult Villagers in Northwestern Bangladesh: A Food Frequency Questionnaire Survey. *International Journal of Food Sciences and Nutrition*, 55: 499-509.
- Tan, V. M. H., Wu, T., Henry, C. J., and Lee, Y. S. (2015). Glycaemic and Insulin Responses, Glycaemic Index and Insulinaemic Index Values of Rice Between Three Asian Ethnic Groups. *British Journal of Nutrition*, 113: 1228-1236.
- Tan, Y., Biondi, A., Desneux, N., and Gao, X. W. (2012). Assessment of Physiological Sublethal Effects of Imidacloprid on the Mirid Bug *Apolygus Lucorum* (Meyer-Dür). *Ecotoxicology*, 21: 1989-1997.
- Teló, G. M., Senseman, S. A., Marchesan, E., Camargo, E. R., Jones, T., and McCauley, G. (2015). Residues of Thiamethoxam and Chlorantraniliprole in Rice Grain. *Journal of Agricultural and Food Chemistry*, 63: 2119-2126.
- Tomizawa, M., and Casida, J. E. (2005). Neonicotinoid Insecticide Toxicology: Mechanisms of Selective Action. *Annual Review on Pharmacology and Toxicology*, 45: 247-268.
- Tomlin, C. D. S. (2000). *Imidacloprid*. The Pesticide Manual, 537-538.
- Törnkvist, A., Glynn, A., Aune, M., Darnerud, P. O., and Ankarberg, E. H. (2011). PCDD/F, PCB, PBDE, HBCD and Chlorinated Pesticides in A Swedish Market Basket From 2005—Levels and Dietary Intake Estimations. *Chemosphere*, 83: 193-199.
- Tsochatzis, E. D., Tzimou-Tsitouridou, R., Menkissoglu-Spirodi, U., Karpouzas, D. G., and Katsantonis, D. (2013). Laboratory and Field Dissipation of Penoxsulam, Tricyclazole and Profoxydim in Rice Paddy Systems. *Chemosphere*, 91: 1049-1057.
- Tucker, K. L. (2010). Dietary Patterns, Approaches, and Multicultural Perspective. *Applied Physiology, Nutrition, and Metabolism*, 35: 211-218.
- United State Environment Protection Agency (US EPA), (1993). *Reference Dose (RfD): Description and use in health risk assessments- Background Document 1A*. Integrated Risk Information System, U.S. Environmental Protection Agency.
- USEPA Method 1694, (2007). *Pharmaceuticals and Personal Care Products in Water, Soil, Sediment, and Biosolids by HPLC/MS/MS*. Washington, D.C. EPA-821-R-08-002, 2011.

- USEPA, *Human Health Risk Assessment*, available at http://www.epa.gov/risk_assessment/health-risk.htm Retrieved on April 2014
- Wang, P., Yang, X., Wang, J., Cui, J., Dong, A. J., Zhao, H. T., and Jing, J. (2012). Multi-Residue Method for Determination of Seven Neonicotinoid Insecticides in Grains Using Dispersive Solid-Phase Extraction and Dispersive Liquid-Liquid Micro-Extraction by High Performance Liquid Chromatography. *Food Chemistry*, 134: 1691-1698.
- Wanwimolruk, S., Kanchanamayoon, O., Phopin, K., and Prachayasittikul, V. (2015). Food Safety in Thailand 2: Pesticide Residues Found in Chinese Kale (Brassica Oleracea), A Commonly Consumed Vegetable in Asian Countries. *Science of the Total Environment*, 532: 447-455.
- Watanabe, E., Baba, K., and Eun, H. (2007). Simultaneous Determination of Neonicotinoid Insecticides in Agricultural Samples by Solid-Phase Extraction Cleanup and Liquid Chromatography Equipped With Diode-Array Detection. *Journal of Agricultural and Food Chemistry*, 55: 3798-3804.
- Watts, M. (2010). *Pesticides: Sowing poison, growing hunger, reaping sorrow*. Pesticide Action Network Asia and the Pacific.
- Weeds, G. R. (2005). Glyphosate-Resistant Weeds. *Review Article*, 1: 183-187.
- Williams, P. N., Raab, A., Feldmann, J., and Meharg, A. A. (2007). Market Basket Survey Shows Elevated Levels of As in South Central US Processed Rice Compared to California: Consequences for Human Dietary Exposure. *Environmental Science and Technology*, 41: 2178-2183.
- Winter, C. K. (1992). Dietary Pesticide Risk Assessment. In *Reviews of environmental Contamination and Toxicology*, 1: 23-67.
- World Health Organization. (1997). *Guidelines for Predicting Dietary Intake of Pesticide Residues, revised*. Global Environment monitoring System-Food Contamination and Assessment Programme (GEMS/Food) in collaboration with Codex Committee on Pesticide Residues, Geneva, Switzerland.
- Xie, W., Qian, Y., Ding, H. Y., Chen, X. M., Xi, J. Y., and Jiang, X. Y. (2009). Determination of Six Neonicotinoid Pesticides Residues in Tea Samples using High Performance Liquid Chromatography Tandem Mass Spectrometry. *China Journal Analytical Chemistry*, 37: 495-499.
- Yeoh, C. B., and Chong, C. L. (2012). LC-MSMS Analysis of Acetamiprid Residue in Crude Palm Oil. *European Journal of Lipid Science and Technology*, 114: 1358-1361.
- Ying, G. G., and Kookana, R. S. (2004). Simultaneous Determination of Imidacloprid, Thiacloprid, and Thiamethoxam in Soil and Water by High-Performance Liquid Chromatography with Diode-Array Detection. *Journal of Environmental Science and Health, Part B*, 39: 737-746.

Yoo, J. K., Lee, S. W., Ahn, Y. J., Nagata, T., and Shono, T. (2002). Altered Acetylcholinesterase as A Resistance Mechanism in the Brown Planthopper (Homoptera: Delphacidae), Nilaparvata Lugens Stal. *Applied Entomology and Zoology*, 37: 37-41.

