

# **UNIVERSITI PUTRA MALAYSIA**

## APPLICATION OF DIFFERENT INSTRUMENTAL TECHNIQUES FOR ADULTERATION ANALYSIS OF BUTTER WITH LARD

# NURRULHIDAYAH AHMAD FADZLILLAH

**IPPH 2015 10** 



### APPLICATION OF DIFFERENT INSTRUMENTAL TECHNIQUES FOR ADULTERATION ANALYSIS OF BUTTER WITH LARD

By

### NURRULHIDAYAH BINTI AHMAD FADZLILLAH

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of Philosophy

February 2015

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia

 $\int$ 



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

### APPLICATION OF DIFFERENT INSTRUMENTAL TECHNIQUES FOR ADULTERATION ANALYSIS OF BUTTER WITH LARD

By

#### NURRULHIDAYAH BINTI AHMAD FADZLILLAH

February 2015

Chair: Professor Amin Ismail, PhD

#### Faculty: Halal Products Research Institute

Determination of food authenticity and detection of adulteration are major concerns not only to consumers, but also to industries and policy makers at all levels of the production process. The major authenticity issues from the food products are the alteration of food ingredients labeling whereby the high value materials are substituted with the cheaper ones without declaring it on the label. Butter is a fairly expensive raw material from an economic point of view, and thus the manufacturers are tempted to look for cheaper substitutes animal fats such as lard. Lard in food are serious matters in view of religious concerns, biological complication and health risk associated with daily intake. In the view of the risk associated with consumption of lard, it is essential to find the most effective, rapid, accurate and sensitive method to identifying the adulterated butter with lard. Therefore, the overall objective of this study is to apply different instrumental techniques for the analysis adulterated butter with lard.

In the first phase, butter is prepared through the conventional method, while lard was rendered from adipose tissue of pig using microwave oven. Subsequently, samples of butter and lard were mixed in different concentration (v/v). Each sample admixtures were subjected to analysis of Fourier Transform Infrared (FTIR) spectroscopy, Differential Scanning Calorimetry (DSC), Gas Chromatography Mass Spectrometry (GCMS), High Performance Liquid Chromatography (HPLC) and Nuclear Magnetic Resonance (NMR) spectroscopy.

Partial Least Square (PLS) from Fourier Transform Infrared Spectroscopy (FTIR) was successful used to predict lard adulteration in butter with the

equation y = 0.998x + 0.0617; was obtained with R<sup>2</sup> and RMSEC values were 0.998 and 1.42% (v/v), respectively using 2 PCs. Meanwhile, differential scanning calorimetry (DSC) analysis showed the cooling enthalpy and melting enthalpy for butter, which were -54.06±8.87 J/g and 35.97±4.18 J/g respectively, compared to the cooling and melting enthalpy of lard -10.72 ± 0.89 J/g and 13.95 ± 0.07 J/g, respectively. The cooling enthalpy supports their comparative FA and TAG saturation levels, as butter has a higher saturation FA and TAG, thus leading to a larger cooling enthalpy. In addition, fatty acid (FA) and triacylglycerol (TAG) analysis were also conducted using GCMS and HPLC. Furthermore, the <sup>1</sup>H-NMR peak signals at  $\delta$ 2.63 showed a specific characteristic of lard. Heteronuclear multiple bond correlation (HMBC). reconfirmed the assignments of the carbon atom of the palmitoyloleoylinoleic. The results from this study showed the complimentary of various instrumental techniques such as FTIR, DSC, HPLC, GCMS and NMR spectroscopy had adequate sensitivity and selectivity which would be an efficient tool for halal authentication.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

### PENGGUNAAN PELBAGAI INSTRUMEN UNTUK MENGESAN PENCAMPURAN LEMAK KHINZIR DALAM MENTEGA

Oleh

### NURRULHIDAYAH BINTI AHMAD FADZLILLAH

Februari 2015

Pengerusi: Professor Amin Ismail, PhD

#### Fakulti: Institut Penyelidikan Produk Halal

Penentuan ketulenan makanan dan pengesanan pencemaran merupakan kebimbangan utama bukan sahaja kepada pengguna, tetapi juga kepada industri dan penggubal dasar pada semua peringkat dalam proses pengeluaran makanan. Salah satu isu ketulenan utama daripada produk makanan adalah pemindaan pelabelan ramuan makanan, di mana bahanbahan bernilai tinggi digantikan dengan bahan-bahan bernilai lebih rendah tanpa mengisytiharkan pada label. Dari sudut pandangan ekonomi, mentega merupakan bahan mentah yang mahal dan menyebabkan pengeluar makanan mencari pengganti yang lebih murah daripada lemak haiwan seperti lemak babi. Lemak babi dalam produk makanan merupakan perkara yang serius dari sudut agama, komplikasi biologi dan risiko. Berdasarkan risiko yang dikaitkan dengan penggunaan lemak babi, maka adalah penting untuk membangunkan kaedah yang paling berkesan, cepat, tepat dan sensitif untuk mengenal pasti mentega yang telah dicemari oleh lemak babi. Oleh itu, objektif keseluruhan dalam kajian ini adalah untuk menggunakan pelbagai kaedah instrumentasi untuk menganalisis mentega yang telah dicemari oleh lemak babi.

Dalam fasa pertama, mentega telah disediakan dengan menggunakan kaedah konvensional manakala lemak babi daripada adipos tisu di ekstrak dengan menggunakan ketuhar gelombang mikro. Kedua-dua sampel telah dicampurkan dengan kepekatan yang berbeza (v/v). Setiap campuran telah di analisis menggunakan Spektroskopi Pengubah Fourier Inframerah (FTIR), Pembezaan Pengimbasan Kalorimeter (DSC), Kromatografi Gas Spektrometri Jisim (GCMS), Kromatografi Cecair

Berprestatsi Tinggi (HPLC) dan Spektroskopi Magnetik Resonans Nuklear (NMR).

Kombinasi spektroskopi FTIR dan kalibrasi "partial least square" (PLS) telah dapat digunakan untuk mengesan dan menentukan lemak babi dalam mentega dengan regresi linear yang bagus, iaitu y = 0.998x + 0.0617, dan nilai R<sup>2</sup> dan nilai kesalahan standard rata-rata pada kalibrasi (RMSEC) masing-masing 0.998 dan 1.42% (v/v) diperoleh menggunakan Manakala, Pembezaan Pengimbsan Kalorimeter (DSC) PCs. 2 menunjukkan profil entalpi penyejukan dan pemanasan mentega, -54.06 ± 8.87 J / g dan 35.97 ± 4.18 J / g, berbanding profil entalpi penyejukan dan pemanasan lemak babi, - 10.72 ± 0.89 J / g dan 13.95 ± 0.07 J / g, masing-masing. Entalpi penyejukan telah menyokong perbezaan ketepuan di antara asid lemak (FA) dan triasilgliserol (TAG), seperti mentega yang mempunyai tahap ketepuan FA dan TAG yang lebih tinggi membawa kepada entalpi penyejukan yang lebih besar. Di samping itu, asid lemak (FA) dan triasigliserol (TAG) analisis juga telah dijalankan dengan menggunakan GCMS dan HPLC. Tambahan pula, Nuklear magnetik resonans spektroskopi memberi puncak pada isyarat δ2.63 ppm menunjukkan ciri-ciri tertentu lemak babi. Ikatan heteronukleus berganda (HMBC), ini mengesahkan semula pemberian atom karbon bagi palmitoyloleoylinoleic. Dapatan daripada kajian ini menunjukkan pelbagai teknik instrumental yang saling melengkapi seperti FTIR, DSC, HPLC, GCMS dan NMR spektroskopi mempunyai kepekaan dan kepilihan yang mencukupi, dan boleh menjadi alat yang efisien bagi pengesahan halal yang pantas dan boleh dipercayai

#### ACKNOWLEDGEMENTS

Alhamdulillah thank you to Almighty God Allah for His mercies and blessings, thank you for always there to listen my prayers and open the door when I least expected.

At this moment of accomplishment, first of all I would like to express my deepest appreciation to my late supervisor, **Prof Dato' Dr Yaakob bin Che Man**. This work would not have been possible without his guidance, support and encouragement. Under his guidance I successfully overcame many difficulties and learned a lot. His unflinching courage and conviction will always inspire me.

Special appreciation and immense thanks to my current supervisor **Prof Dr Amin bin Ismail**, whose support and guidance from the initial phase till the end has enabled me to learn something new and valuable, as well as giving me insight to the understanding of the subject. I am also greatly indebt to my co-supervisor, **Prof Dr Shuhaimi Mustafa** and **Assoc Prof Dr Alfi Khatib** for his unremitting assistance, consideration and thoughtfulness towards myself. I would also like to thank some people from early days of my research tenure, **Prof Dr Abdul Rohman**, **Prof Dr Sugeng Triwahyono**, **Prof Dr Russly Ab Rahman**, **Assoc Prof Dzulkifly Mat Hashim** were among those who kept me going at the beginning.

Special thanks to my colleagues and staff from Halal Product Research Institute, who would always be mentors to me and for being there with me through thick and thin. I greatly appreciate **Ministry of Higher Education** and **Universiti Putra Malaysia** for providing the funds for the study and to give me the chance to study here.

It's my fortune to gratefully acknowledge the support of some special individuals. Word fail me to express my appreciation to my beloved husband **Ade Erawan** for his support, patience and love. I can see the good shape of my thesis because of his help and suggestions in formatting the entire thesis. It is also a pleasure to mention my son my great hero **Aleef Haseef**. I feel so proud you are a part of me.

Last but not least, I would like to pay high regards to my parents Ahmad Fadzlillah and Raihanah, my siblings Ahmad Mahyuddin, Nurrulashikin, Noramalina and Ahmad Hafizuddin for their sincere encouragement and inspiration throughout my journey and lifting me uphill this phase of life. I owe everything to them. Besides this, several people have knowingly and unknowingly helped me in the successful completion of this research.

Thank you all.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of **Doctor of Philosophy**. The members of the Supervisory Committee were as follows:

### Amin Bin Ismail, PhD

Professor Department of Nutrition and Dietetics Faculty of Medicine and Health Science Universiti Putra Malaysia (Chairman)

### Shuhaimi bin Mustafa, PhD

Professor Halal Products Research Institute Universiti Putra Malaysia (Member)

### Alfi Khatib, PhD

Associate Professor Department of Pharmaceutical Chemistry Kuliyyah of Pharmacy International Islamic University Malaysia (Member)

### **BUJANG BIN KIM HUAT, PhD**

Professor and Dean School of Graduate Studies Universiti Putra Malaysia

Date:

### TABLE OF CONTENTS

			Page		
ABSTRACT iii ABSTRAK v ACKNOWLEDGEMENTS vii APPROVAL viii DECLARATION x LIST OF TABLES xvi LIST OF FIGURES xvi LIST OF ABBREVIATIONS xix					
CHAPTE	:R				
1	<b>GEN</b> 1.1 1.2 1.3	ERAL INTRODUCTIONProblem statementsSignificance of the studyObjectives1.3.1General objective1.3.2Specific objectives	1 1 3 4 4 4		
2	LITE 2.1 2.2 2.3 2.4	<b>RATURE REVIEW</b> Edible fats and oils Food authentication Recent analytical techniques for food authentication Recent development in butter	7 7 8 9		
	2.5 2.6	adulterations Butter 2.5.1 Fatty acid composition of butter 2.5.2 Health benefits of butter Lard	10 11 12 12 13		
	2.7 2.8	<ul> <li>2.6.1 Physicochemical characteristic or lard</li> <li>2.6.2 Application of lard</li> <li>Recent issues in Halal food adulteration</li> <li>Recent technology in Halal</li> </ul>	14 15 16		
	2.9	authentication Validation method of analytical chemistry	17 18		
3	USE (FTI CHE ADU	OF FOURIER TRANSFORM INFRARED R) SPECTROSCOPY AND MOMETRIC DATA TO ANALYSIS LARD LTERATED WITH BUTTER	21		

 $\bigcirc$ 

	3.1	Introdu	ction	21
	3.2	Materia	als and Methods	23
		3.2.1	Sample preparation	23
		3.2.2	Fatty acid analysis	24
		3.2.3	Discriminant analysis	24
		3.2.4	Calibration and validation	
			standard	24
		3.2.5	FTIR instrumental analysis	25
		3.2.6	Statistical and chemometrics	
			analysis	25
	3.3	Result	s and discussion	25
		3.3.1	FTIR spectra	25
		3.3.2	Fatty acid composition of butter	
			and lard	29
		3.3.3	Discriminant analysis	29
		3.3.4	Quantitative analysis of lard in	
			butter	31
	3.4	Conclu	sion	34
_				
4	MON	ITORIN	G ADULTERATION OF BUTTER	
	WITI	H LARD	USING DIFFERENTIAL	
	SCA	NNING	CALORIMETRY	36
	4.1	Introdu		36
	4.2	Materia	als and Methods	37
		4.2.1	Preparation of blends	37
		4.2.2	Analysis of the composition of	27
		400	TAG USING HPLC	37
		4.2.3	Coopping Colorimetry	27
		121	Statistical analysis	ى ەر
	12	4.2.4 Docult	Statistical analysis	20
	4.3	A 2 4		30 20
		4.3.1	Heating thermogram	30
		4.3.2	Cooling thermogram	39
	11	4.3.3 Conclu		43
	4.4	COncio	3011	50
5	DIFF	FRENT	IATION OF FATTY ACID	
U	CON	IPOSITI	ON OF BUTTER ADULTERATED	
	WITI	HLARD	USING GAS	
	CHR	OMATO	OGRAPHY MASS	
	SPE	CTROM	ETRY COMBINED WITH	
	PRIN		COMPONENT ANALYSIS	51
	5.1	Introdu	iction	51
	5.2	Materia	als and Methods	52
		5.2.1	Sample preparation	52
		5.2.2	Calibration and validation	52
		523	Fatty acid analysis	53

### xiii

		5.2.4	Statistical and chemometric	53
	53	Result	s and discussion	53
	0.0	5.3.1	Fatty acid composition	53
		5.3.2	Principal component analysis of	
			butter and butter adulterated with	1
			lard	58
		5.3.3	Analysis butter adulteration with	
			lard using fatty acid composition	58
	5.4	Conclu	ision	60
6	IDE	NTIFICA	TION OF TRIACYLGLYCEROL	
	CON	<b>IPOSIT</b>	ION OF LARD AS A FATTY	
	ACII	DS MAR	KER FOR HALAL	
	AUT	HENTIC	CATION USING HIGH	
	RES	OLUTIC	ON OF NMR SPECTROSCOPY	61
	6.1	Introdu	iction	61
	6.2	Materia	als and Methods	62
		6.2.1	Sample preparation	63
		6.2.2	NMP measurement	63
		624	Bucketing of <sup>1</sup> H NMP spectra	03
		625	Analysis of TAG compositional	05
		0.2.0	by HPLC	63
		6.2.6	Determination of fatty acid	
			composition	64
	6. <mark>3</mark>	Result	s and discussion	64
		6.3.1	Spectral analysis	64
		6.3.2	Fatty acid profile	69
		6.3.3	Triacylglycerol (TAG)	
			composition	70
		6.3.4	C-NMR	75
		6.3.5	Heteronuclear Multiple Bond	
	6.4	Canali		//
	0.4	Concil	ISION	80
7	DISC	CRIMIN	ATION OF BUTTER AND	
	ADU	ILTERA	TED BUTTER BY GC-MS AND	
	NMF	R SPEC	TROSCOPY COMBINED WITH	81
	PRI	NCIPAL		
	7.1	Introdu	ICTION	81
	1.2	iviateria	als and Methods	83
		1.Z.I	Sample preparation	83 02
		1.2.2 723	Fally ablu analysis Statistical and chemometrics	63
		1.2.5	analysis	83
	73	Result	s and discussion	84
	1.0	, count		07

xiv

		7.3.1 7.3.2	Fatty acid composition Discriminant analysis	84 87
	7.4	Conclus	market sample of butter compared with lard sion	90 93
8	SUM REC 8.1 8.2 8.3	IMARY, COMMEN Summa Genera Recom	GENERAL CONCLUSION AND DATIONS ary I conclusion mendations	<b>95</b> 95 96 96
REFEREN BIODATA LIST OF F	ICES OF ST PUBLIC			98 113 114

 $\bigcirc$ 

### LIST OF TABLES

Table		Page
1	Fatty acid composition of lard	14
2	Functional groups and modes of vibration in lard and butter	28
3	Fatty acid compositions of butter and lard determined using GC with flame ionization	
	detector	29
4	Triacylglycerol (TAG) composition of butter adulterated with different lard concentrations	
	(V/V)	40
5	Cooling and melting partial enthalpy	50
6	Fatty acid compositions of butter added with	
	different levels of lard (v/v)	56
7	Assignments of signals of <sup>1</sup> H-NMR spectra	
	from samples	69
8	Fatty acid composition (%, peak area) of	
	butter and lard	71
9	Triacylglycerol composition of butter and	
	other animal fats (lard, chicken, beef and	
4.0	mutton)	71
10	Fatty acid compositions of butter, lard,	
	chicken fat, beef fat and mutton fat	85
11	Tentative identification of functional groups of	0.0
10	List of the according acording to a second acording	88
12	of buttor available in Malaysia	01
		91

 $\mathbf{G}$ 

### LIST OF FIGURES

F	igure		Page
	1	FTIR-ATR spectra of pure butter and lard	-
		peaks shown at wavenumber of 4000-650	
		cm <sup>-1</sup>	27
	2	Coomans plot of pure butter, butter	
		containing lard and pure lard	32
	3	Partial least square (PLS) model for the	
		relationship between actual and FTIR	
		predicted values (%v/v) of lard as adulterant	
		in butter in calibration samples.	33
	4	PLS model for the relationship between	
		actual and FTIR estimated values (%v/v) of	
		lard as adulterant in butter in	
		prediction/validation samples	34
	5	Differential Scanning Calorimetry (DSC)	
		heating thermogram of pure butter and pure	40
	~	lard	42
	6	Melting curves of lard, butter and lard-butter	45
	7	blends containing 1–80 % butter	45
	1	blende containing 1, 80 % butter and the	
		oplarged molting profile ground 0°C	46
	0	Differential Scapping Calorimetry (DSC)	40
	0	cooling thermogram of pure butter and pure	
		lard	47
	٥	Cooling curves of lard, butter and lard butter	4/
	9	blends containing 1–80 % butter	18
	10	Cooling curves of lard, butter and lard-butter	-0
	10	blends containing 1–80 % butter and the	
		enlarged cooling profile around 0 °C known	
		as peak B	49
	11	Eatty acids composition of pure butter (A) and	
		pure lard (B)	55
	12	PCA classification for butter and lard	
		projected by PC1 and PC2; a) score plot and	
		b) loading plot	59
	13	Composition changes of fatty acids (FAs) in	
		butter adulterated with different levels of lard:	
		(a) FA with decreased level and (b) FA with	
		increased level with increasing concentration	
		of lard	60
	14	NMR spectra of pure animal fats including	
		beef fat, chicken fat, mutton fat and lard	65
	15	NMR spectra of butter and lard mixtures	

6

	showing changes in the height ratio at 2.60-	
40	2.84ppm	66
10	butter (B)	68
17	The chromatogram of TAG of butter as	
	determined using HPLC with refractive	
	detector.	72
18	Chromatogram of TAG of lard as determined	
4.0	using HPLC with refractive detector	73
19	observed in the NMR spectra, with the	
	chemical shifts of the most important signals	75
20	<sup>13</sup> C Nuclear Magnetic Resonance (NMR)	
	spectrum of lard	76
21	Structural correlation between proton and	
	carbon in lard	78
22	Heteronuclear multiple bonds correlation	
~~	(HMBC) spectrum of lard	79
23	Score plot of PCA of butter derived from	
~ 4	animal fats	86
24	Loading plot of PCA of butter derived from	07
25	Compose plot of buttor and buttor adultorated	87
20	with animal fate $( - )$ pure butter: $( A )$ butter	
	sample adulterated with animal fats	80
26	Coomans plot of butter and those adulterated	03
20	with three nimal fats: $(\Box)$ pure butter: $(\Lambda)$	
	butter sample adulterated with three animal	
	fats	90
27	Score plot of PCA market sample of butter	
	and lard based on <sup>1</sup> H-NMR spectra	92
28	Loading plot of PCA market sample of butter	
	and lard based on <sup>1</sup> H-NMR spectra	92

 $(\mathbf{C})$ 

### LIST OF ABBREVIATIONS

AOAC	Association of Official Analytical Chemists
AOCS	American Oil Chemists' Society
ATR	Attenuated total reflectance
BF	Beef fat
CF	Chicken fat
cm⁻¹	Reciprocal centimetre
DA	Discriminant analysis
FAME	Fatty acid methyl ester
FTIR	Fourier transform infrared spectroscopy
FID	Flame ionization detector
GC	Gas chromatography
HPLC	High performance liquid chromatography
NMR	Nuclear Magnetic Resonance Spectroscopy
IR	Infrared
MF	Mutton fat
MUFA	Monounsaturated fatty acid
MIR	Mid infrared
NIR	Near infrared
PC	Principal component
PCA	Principal component analysis
PLS	Partial least square
PRESS	Predicted residual error sum of square
PUFA	Polyunsaturated fatty acid
$R^2$	Coefficient of determination
RMSEC	C Root mean standard error of calibration
RMSEC	CV Root mean standard error of cross validation
RMSEF	P Root mean standard error of prediction
rpm	Revolution per minute
TG	Triglyceride
v/v	Volume/volume

 $(\mathbf{C})$ 

### CHAPTER 1

### **GENERAL INTRODUCTION**

Butter, a type of edible fat, is one of the emerging dairy products in the market because of its nutritional benefits to human. Butter contains protein, calcium and phosphorus. The main vitamins in butter include A, D, E and vitamin K (Van Ruth *et al.*, 2008). Meanwhile, additional nutrients in butter include fatty acids, lactones, methyl ketones as well as dimethyl and diacetyl sulfide. Butter is rich in the most easily absorbable form of Vitamin A necessary for thyroid and adrenal health (Ecologist, 2015; Icattoreb, 2013; Tokach, 2013)

The denomination "butter" is reserved, for law, to the product got only and exclusively from the cream drawn from the milk of cow. It is one of the most expensive fats that have standard of quality. In this connection, the European Community has developed special regulation, strict standards for identifying butter and has established that butter must be obtained exclusively from cow's milk or cream (Commission of the European Communities, 1990; Coni *et al.*, 1994). According to (*Commission of the European Communities*, 1990; Coni, *et al.*, 1994), it has been stated under essential composition and quality factors section that commercial butter should consist of 80% milk fat, 16% water content and 2% milk solids other than fat, including salt.

Food Safety Information System Malaysia under Food Act 1983 and Food Regulation 1985 law has stated that butter is the solid product derived exclusively from milk or cream, shall contain not less than 80% of milk fat, 16% of water and may contain salt. Confirmation of the authenticity of a food or food ingredient is an increasing challenge for food scientist. This is especially the case when a substitute value claim, such as one relating to geographic origin or a particular processing history, is made on food label (Commission, 2011). Criteria and procedures for assessing the authenticity of these high-quality products need to be developed.

### 1.1 **Problem statements**

Adulteration is a serious problem for both producers and consumers in the food manufacturing industry. The driving force behind food adulteration is when the producer uses partially or totally low-cost ingredients rather than

more expensive ones to increase the revenue (Downey and Kelly, 2006). The growing interest in the authenticity of food products requires reliable verification methods because the properties of foods from different origins can be also differentiated.

Butter is a common raw material for many food products. It contains partially saturated animal milk fat. Butter is sometimes illegally mixed with cheaper animal fats such as lard for use in adulteration. This occurrence of adulteration is to gain economic benefit due to more expensive price of butter than lard and other animal fats (Koca *et al.*, 2010). However, in the case of adulteration using animal fats, this would affect in terms of consumer beliefs and practice such as for Hindus, Jews and Muslims (Didar *et al.*, 2013). For Muslims especially in Malaysia, adulteration using lard or improperly slaughtered animal fats is a big concern.

From the above considerations, the detection of other animal fats in butter is emphasized today more than ever, and a number of investigations have been carried out by several research groups to develop analytical methods for this purpose (Kamm *et al.*, 2001). When the added fat is of vegetable origin, it is usually easy for the analyst to detect its presence (Burros, 1986; Coni, *et al.*, 1994; Grob *et al.*, 1992; Hammond *et al.*, 1988; Plattner and J, 1981). In contrast, if the added fat is of animal origin, few analytical methods are available, and most of the methods are either difficult to be performed or time consuming ((Coni, *et al.*, 1994; Grob *et al.*, 1989).

Lard is one of the cheapest fats that can be found in some countries. Consequently, some manufacturers prefer to mix lard with other oils to reduce the production cost (Geeraert and Sandra, 1985; Luf *et al.*, 1987; Precht, 1991a; 1991b; Precht, 1992a; 1992b; Shukla *et al.*, 1983; Timms, 1980). It is widely applied as dough shortening and frying medium (Marikkar *et al.*, 2005). Medical reports on the effect of lard consumption in promoting carcinogenesis have alerted people to be more vigilant about pork and lard in food products (Marikkar, *et al.*, 2005; Matz, 1992). It is important to detect adulteration of foods with porcine-based products especially in monitoring and implementation of Halal regulations.

There is an increasing trend in some countries to mix pork and lard in their food products for the purposes of gaining extra economic profit. Methods have been developed for detection of lard in food product formulations, namely cake (Al-Rashood *et al.*, 1996), chocolate (Syahariza *et al.*, 2005) and biscuits (Che Man *et al.*, 2005), meat (Che Man *et al.*, 2011) ghee

and butter (Al-Jowder *et al.*, 1997), and vegetable oils (Farag *et al.*, 1983; Kowalski, 1989).

In last several years, there have been some issues associated with the adulteration of butter in Malaysia. Among the issues are pig DNA in Golden Churn Butter and in Kluang coffee. These issues have given a great impact to the producer and consumers since the majority of the population in Malaysia are Muslim. These are some examples of how important it is to develop a reliable technique to ensure the Halal status of food products, which is crucial for Muslims and Halal authentication (Marikkar, et al., 2005). Some published articles related to food authentication are mainly on lard or pork. Such reports are as described in (Yahya Ishmael, 2005) for detecting lard in some vegetable oils and (Rohman and Man, 2010a) for analysis of pork in pork meatball. However, from an extensive literature search, there is no report regarding rapid detection of butter adulterated with lard using Nuclear Magnetic Resonance (NMR) spectroscopy. The application of NMR in this study is to identify selected TAG and FA composition as potential chemical marker for lard as an adulterant and its adulteration in butter Therefore, this study was carried out, and the general objective was to apply different instrumental techniques for adulteration analysis of butter with lard.

### 1.2 Significance of the study

Testing of food products for the purpose of labeling and authentication is necessary to avoid unfair competition and assure consumer's protection against fraudulent practices in the food industry. The major issues concerning authenticity are where, the potential financial rewards for substitution of cheaper ingredients are relatively high (Rohman and Sismindari, 2011).

The advance in food technology has resulted in the issues getting more complicated where ingredients used in foods are more difficult to understand by consumers unless they are directly involved in the related field. In Halal authentication, one cannot rely solely on physical inspection and documentation, but will need complementary evidence of the latest, high technology of analytical instrumentation. The analytical method used for the analysis of lard is based on the differences in the nature and the composition of minor and major components present or absent in lard and those in foods. These methods typically rely on physical-chemical constants or based on chemical and biological measurements (Lai *et al.*, 1994). Ideally, the methods developed should be rapid, accurate, reliable, easy to operate and low cost. As a result, high-resolution NMR spectroscopy and physicochemical methods have been developed for

analysis of lard in butter. The developed method can be used in food industries to authenticate butter from lard. This finding will also help the Department of Islamic Development Malaysia (JAKIM) as the authorized body responsible for the standardization of Halal certifications.

Global Halal industry has consistently been on the rise in many parts of the world at least for a decade already. The market for Halal food around the globe is getting bigger by years. There is an emerging interest among Muslim consumers everywhere to consume Halal products, and this becomes a good sign for global Halal market. The research conducted by the World Halal Secretariat estimated that the global Halal products market is estimated at an enormous USD 2.3 trillion (not including banking), which USD1.4 trillion (67 per cent) of this market represents food and beverages. Pharmaceuticals on the other hand make up USD506 billion (22 per cent), with cosmetics and personal care amounting to USD230 billion (Kowalski, 1989). Therefore, detection of lard in Halal food becomes more demanding.

The advancement of sophisticated technology in Halal industry will lead to the innovation outcomes that could give alternatives way to authenticate butter from which types of animal fat sources and would overcome the 'Halal' issues regarding animal fats. More complex techniques such as NMR have led to greater accuracy of the analytical results. There have been important efforts made to develop new applications of existing analytical techniques for detection and quantification of lard as an adulterant and its adulteration with butter.

### 1.3 Objectives

The objectives of the study were as follows:

### 1.3.1 General objective

The general objective of this research was to apply different instrumental techniques for adulteration analysis of butter with lard.

#### 1.3.2 Specific objectives

i. To identify and quantify the functional groups in lard and adulterated butter using FTIR and PLS method.

- ii. To identify the pure butter from the adulterated butter using thermal properties and triacylglycerol profile analysis
- iii. To examine the fatty acid composition of butter adulterated with lard using gas chromatography mass spectrometry and multivariate data analysis.
- iv. To determine the triacylglycerol marker profile of lard and butter adulterated with lard using high-resolution NMR spectroscopy.
- v. To investigate the potential of NMR spectroscopy and principal component analysis (PCA) to detect the adulteration of butter with lard.



#### REFERENCES

- Abbas, O., Fernández Pierna, J. A., Codony, R., von Holst, C. and Baeten, V. (2009). Assessment of the discrimination of animal fat by FT-Raman spectroscopy. *Journal of Molecular Structure*, 924-926, 294-300.
- Abu-Taha, M., Halasa, M. and Abu-Samreh, M. (2013). On the Usage of the Faraday Effect as an Authentication Technique for Vegetable Oils.
- Aida, A., Che Man, Y., Wong, C., Raha, A. and Son, R. (2005). Analysis of raw meats and fats of pigs using polymerase chain reaction for Halal authentication. *Meat science*, 69(1), 47-52.
- Aktaş, N. and Kaya, M. (2001a). Detection of beef body fat and margarine in butterfat by differential scanning calorimetry. *Journal of Thermal Analysis and Calorimetry*, 66(3), 795-801.
- Aktaş, N. and Kaya, M. (2001b). The influence of marinating with weak organic acids and salts on the intramuscular connective tissue and sensory properties of beef. *European Food Research and Technology*, 213(2), 88-94.
- Aktaş, N. and Kaya, M. (2001c). Influence of weak organic acids and salts on the denaturation characteristics of intramuscular connective tissue. A differential scanning calorimetry study. *Meat science*, 58(4), 413-419.
- Al-Jowder, O., Kemsley, E. and Wilson, R. (1997). Mid-infrared spectroscopy and authenticity problems in selected meats: a feasibility study. *Food chemistry*, 59(2), 195-201.
- Al-Rashood, K. A., Abdel-Moety, E. M., Rauf, A., Abou-Shaaban, R. R. and Al-Khamis, K. I. (1995). Triacylglycerols-profiling by high performance liquid chromatography: A tool for detection of pork fat (lard) in processed foods. *Journal of Liquid Chromatography & Related Technologies*, 18(13), 2661-2673.
- Al-Rashood, K. A., Abou-Shaaban, R. R., Abdel-Moety, E. M. and Rauf, A. (1996). Compositional and thermal characterization of genuine and randomized lard: a comparative study. *Journal of the American Oil Chemists' Society*, 73(3), 303-309.
- Ali, M., Hashim, U., Mustafa, S., Che Man, Y., Adam, T. and Humayun, Q. (2014). Nanobiosensor for the detection and quantification of pork adulteration in meatball formulation. *Journal of Experimental Nanoscience*, 9(2), 152-160.
- Ando, Y., Satake, M. and Takahashi, Y. (2000). Reinvestigation of positional distribution of fatty acids in docosahexaenoic acid-rich fish oil triacylsn-glycerols. *Lipids*, 35(5), 579-582.
- Apetrei, C., Apetrei, I. M., Villanueva, S., de Saja, J. A., Gutierrez-Rosales, F. and Rodriguez-Mendez, M. L. (2010). Combination of an e-nose, an etongue and an e-eye for the characterisation of olive oils with different degree of bitterness. *Analytica Chimica Acta*, 663(1), 91-97.
- Arlorio, M., Coisson, J. D., Bordiga, M., Travaglia, F., Garino, C., Zuidmeer, L., et al. (2010). Olive oil adulterated with hazelnut oils: simulation to identify possible risks to allergic consumers. *Food Additives and Contaminants*, 27(1), 11-18.

- Arvanitoyannis, I. S., Varzakas, T. H., Kiokias, S. and Labropoulos, A. E. (2009). 5 Lipids, Fats, and Oils. *Advances in Food Biochemistry*, 131.
- Aryee, A. N. A., van de Voort, F. R. and Simpson, B. K. (2009). FTIR determination of free fatty acids in fish oils intended for biodiesel production. *Process Biochemistry (Amsterdam, Netherlands)*, 44(4), 401-405.
- Asensio, L., González, I., García, T. and Martin, R. (2008). Determination of food authenticity by enzyme-linked immunosorbent assay (ELISA). *Food Control*, 19(1), 1-8.
- Aursand, M., Jørgensen, L. and Grasdalen, H. (1995). Positional distribution of ω3 Fatty acids in marine lipid triacylglycerols by high-resolution13C nuclear magnetic resonance spectroscopy. *Journal of the American Oil Chemists' Society*, 72(3), 293-297.
- Bailey, N. J., Sampson, J., Hylands, P. J., Nicholson, J. K. and Holmes, E. (2002). Multi-Component Metabolic Classification of Commercial Feverfew Preparations via high-field 1H-NMR spectroscopy and chemometrics. *Planta medica*, 68(08), 734-738.
- Baker, S., Nicklin, J. and Griffiths, C. (2011). *BIOS instant notes in microbiology*: Taylor & Francis.
- Ballabio, D. and Todeschini, R. (2009). Multivariate classification for qualitative analysis (pp. 83-104): Elsevier: Burlington, MA, USA.
- Barron, L. J., Hierro, M. T. G. and Santa-María, G. (1990). HPLC and GLC analysis of the triglyceride composition of bovine, ovine and caprine milk fat. *Journal of Dairy Research*, 57(04), 517-526.
- Belitz, H.-D., Grosch, W. and Schieberle, P. (2009). Edible fats and oils. *Food chemistry*, 640-669.
- Belton, P. S., Colquhoun, I. J., Kemsley, E. K., Delgadillo, I., Roma, P., Dennis, M. J., et al. (1998). Application of chemometrics to the <sup>1</sup>H NMR spectra of apple juices: discrimination between apple varieties. *Food chemistry*, 61(1), 207-213.
- Bendini, A., Cerretani, L., Carrasco-Pancorbo, A., Gómez-Caravaca, A. M., Segura-Carretero, A., Fernández-Gutiérrez, A., et al. (2007a). Phenolic molecules in virgin olive oils: A survey of their sensory properties, health effects, antioxidant activity and analytical methods. An overview of the last decade Alessandra. *Molecules*, 12(8), 1679-1719.
- Bendini, A., Cerretani, L., Di Virgilio, F., Belloni, P., BONOLI-CARBOGNIN, M. and Lercker, G. (2007b). Preliminary evaluation of the application of the FTIR spectroscopy to control the geographic origin and quality of virgin olive oils. *Journal of Food Quality*, 30(4), 424-437.
- Bernuy, B., Meurens, M., Mignolet, E., Turu, C. and Larondelle, Y. (2009). Determination by fourier transform Raman spectroscopy of conjugated linoleic acid in i2-photoisomerized soybean oil. *Journal of Agricultural* and Food Chemistry, 57(15), 6524-6527.
- Biliaderis, C., Maurice, T. and Vose, J. (1980). Starch gelatinization phenomena studied by differential scanning calorimetry. *Journal of food science*, 45(6), 1669-1674.
- Burros, B. C. (1986). International Laboratory 4:18.

- Cajka, T., Riddellova, K., Tomaniova, M. and Hajslova, J. (2010). Recognition of beer brand based on multivariate analysis of volatile fingerprint. *Journal* of Chromatography A, 1217(25), 4195-4203.
- Camin, F., Larcher, R., Perini, M., Bontempo, L., Bertoldi, D., Gagliano, G., et al. (2010). Characterisation of authentic Italian extra-virgin olive oils by stable isotope ratios of C, O and H and mineral composition. *Food Chemistry*, 118(4), 901-909.
- Cebula, D. J. and Smith, K. W. (1991). Differential scanning calorimetry of confectionery fats. Pure triglycerides: effects of cooling and heating rate variation. *Journal of the American Oil Chemists Society*, 68(8), 591-595.
- Cebula, D. J. and Smith, K. W. (1992). Differential scanning calorimetry of confectionery fats: Part II—Effects of blends and minor components. *Journal of the American Oil Chemists' Society*, 69(10), 992-998.
- Che Man, Y., Nurrulhidayah, A., Rohman, A., Amin, I., Shuhaimi, M. and Khatib, A. (2013). Authentication analysis of butter from beef fat using Fourier Transform Infrared (FTIR) spectroscopy coupled with chemometrics. *International Food Research Journal*, 20(3), 1383-1388.
- Che Man, Y., Syahariza, Z., Mirghani, M. E. S., Jinap, S. and Bakar, J. (2005). Analysis of potential lard adulteration in chocolate and chocolate products using Fourier transform infrared spectroscopy. *Food chemistry*, 90(4), 815-819.
- Che Man, Y. B. (2012). The Contribution of Science and Technology in Determining the Permissibility (Halalness) of Food Products. *Revelation and Science*, 2(01).
- Che Man, Y. B., Abidin, S. Z. and Rohman, A. (2011). Discriminant analysis of selected edible fats and oils and those in biscuit formulation using FTIR spectroscopy. *Food Analytical Methods*, 4(3), 404-409.
- Cheong, L.-Z., Zhang, H., Xu, Y. and Xu, X. (2009). Physical characterization of lard partial acylglycerols and their effects on melting and crystallization properties of blends with rapeseed oil. *Journal of Agricultural and Food Chemistry*, 57(11), 5020-5027.
- Chiavaro, E., Vittadini, E., Rodriguez-Estrada, M. T., Cerretani, L. and Bendini, A. (2008). Differential scanning calorimeter application to the detection frefined hazelnut oil in extra virgin olive oil. *Food Chemistry*, 110(1), 248-256.
- Chowdhry, B. Z. and Cole, S. C. (1989). Differential scanning calorimetry: applications in.
- Christie, W. and Moore, J. (1969). The effect of dietary copper on the structure and physical properties of adipose tissue triglycerides in pigs. *Lipids*, 4(5), 345-349.
- Christy, A. A. and Egeberg, P. K. (2006). Quantitative determination of saturated and unsaturated fatty acids in edible oils by infrared spectroscopy and chemometrics. *Chemometrics and intelligent laboratory systems*, 82(1), 130-136.
- Collomb, M., Bütikofer, U., Spahni, M., Jeangros, B. and Bosset, J. (1999). Fatty acid and glyceride composition of cow's milk fat in high-and lowland regions. *Sciences des Aliments (France)*.

- Collomb, M. and Spahni, M. (1991). Adulteration of dairy products. Review of analytical methods for the determination of vegetable and animal fats in butterfat. *Mitteilungen aus dem Gebiete der Lebensmitteluntersuchung und Hygiene*, 82, 615-662.
- Commission, C. A. (2011). Milk and Milk Products, Rome: Food and Agriculture Organization of The United Nations. (2<sup>nd</sup> Edition).
- Commission of the European Communities. (1990). Doc. VI/6366/90, Brussels.
- Coni, E., Di Pasquale, M., Coppolelli, P. and Bocca, A. (1994). Detection of animal fats in butter by differential scanning calorimetry: A pilot study. *Journal of the American Oil Chemists' Society*, 71(8), 807-810.
- Contarini, G., Povolo, M. and Bonfitto, E. (1999). The present and future for controlling the authenticity of butter. *Latte*, 24, 60-69.
- Cosio, M. S., Ballabio, D., Benedetti, S. and Gigliotti, C. (2006). Geographical origin and authentication of extra virgin olive oils by an electronic nose in combination with artificial neural networks. *Analytica Chimica Acta*, 567(2), 202-210.
- Cuesta, M. J. R. (2007). *Limit of detection for second-order calibration methods*: Universitat Rovira i Virgili.
- De La Fuente, M. A. and Juárez, M. (2005). Authenticity assessment of dairy products. Critical reviews in food science and nutrition, 45(7-8), 563-585.
- Defernez, M., Kemsley, E. K. and Wilson, R. H. (1995). Use of infrared spectroscopy and chemometrics for the authentication of fruit purees. *Journal of Agricultural and Food Chemistry*, 43(1), 109-113.
- Derewiaka, D., Sosińska, E., Obiedziński, M., Krogulec, A. and Czaplicki, S. (2011). Determination of the adulteration of butter. *European journal of lipid science and technology*, 113(8), 1005-1011.
- Didar, Ü., İlaslan, K., Boyacı, İ. H. and Özay, D. S. (2013). Rapid detection of fat adulteration in bakery products using Raman and near-infrared spectroscopies. *European Food Research and Technology*, 237(5), 703-710.
- Downey, G. and Kelly, J. D. (2006). Food authentication using infrared spectroscopic methods: Teagasc.
- Duce, S., Amin, M., Horsfield, M., Tyszka, M. and Hall, L. (1995). Nuclear Magnetic Resonance Imaging of Dairy Products in Two and Three Dimensions. *International Dairy Journal*, 5(4), 311-319.
- Dupuy, N., Duponchel, L., Huvenne, J., Sombret, B. and Legrand, P. (1996). Classification of edible fats and oils by principal component analysis of Fourier transform infrared spectra. *Food chemistry*, 57(2), 245-251.
- Echizen, A. and Diki, M. (1975). Determination of butterfat in mixed fats. Relation between butyric acid and butterfat content in mixed fats. *Rep. Cent. Customs Lab*, 15, 111.
- Ecologist, C. B. (Producer). (2015). The 20 Health Benefits of Real Butter. Retrieved from

http://bodyecology.com/articles/benefits\_of\_real\_butter.php#.VQmNG4 7LdcQ

- Encinar, J., González, J. and Rodríguez-Reinares, A. (2007). Ethanolysis of used frying oil. Biodiesel preparation and characterization. *Fuel Processing Technology*, 88(5), 513-522.
- Enser, M. (1995). Meat lipids Developments in oils and fats (pp. 1-31): Springer.
- Fadzlillah, N. A., Rohman, A., Ismail, A., Mustafa, S. and Khatib, A. (2013). Application of FTIR-ATR Spectroscopy Coupled with Multivariate Analysis for Rapid Estimation of Butter Adulteration. *Journal of oleo science*, 62(8), 555-562.
- Fan, Y., Chen, M., Shentu, C., El-Sepai, F., Wang, K., Zhu, Y., et al. (2009). Ionic liquids extraction of Para Red and Sudan dyes from chilli powder, chilli oil and food additive combined with high performance liquid chromatography. *Analytica Chimica Acta*, 650(1), 65-69.
- Farag, R., Abo-Raya, S., Ahmed, F., Hewedi, F. and Khalifa, H. (1983). Fractional crystallization and gas chromatographie analysis of fatty acids as a means of detecting butterfat adulteration. *Journal of the American Oil Chemists Society*, 60(9), 1665-1669.
- Fasina, O., Craig-Schmidt, M., Colley, Z. and Hallman, H. (2008). Predicting melting characteristics of vegetable oils from fatty acid composition. *LWT-Food Science and Technology*, 41(8), 1501-1505.
- Fernandez, C., Astier, C., Rock, E., Coulon, J. B. and Berdagué, J. L. (2003). Characterization of milk by analysis of its terpene fractions. *International journal of food science & technology*, 38(4), 445-451.
- Foglia, T. A., Nelson, L. A., Dunn, R. O. and Marmer, W. N. (1997). Lowtemperature properties of alkyl esters of tallow and grease. *Journal of* the American Oil Chemists' Society, 74(8), 951-955.
- Forssell, P., Parovuori, P., Linko, P. and Poutanen, K. (1993). Enzymatic transesterification of rapeseed oil and lauric acid in a continuous reactor. *Journal of the American Oil Chemists Society*, 70(11), 1105-1109.
- Freire, E. (1995). Differential scanning calorimetry *Protein stability and folding* (pp. 191-218): Springer.
- Fuente, M. D. L. and Juarez, M. (1999). ARTICLES-Review: Application of chromatographic techniques to the study of triglycerides and sterols of milk fat. *Food Science and Technology International-Frederick*, 5(2), 103-120.
- García-González, D. L. and Aparicio, R. (2010). Coupling MOS sensors and gas chromatography to interpret the sensor responses to complex food aroma: Application to virgin olive oil. *Food Chemistry*, 120(2), 572-579.
- Gaydou, E. M., Bianchini, J. P. and Ratovohery, J. V. (1983). Triterpene alcohols, methylsterols, sterols, and fatty acids in five Malagasy legume seed oils. *Journal of Agricultural and Food Chemistry*, 31(4), 833-836.
- Geeraert, E. and Sandra, P. (1985). Capillary GC of triglycerides in fats and oils using a high temperature phenylmethylsilicone stationary phase, part I. *Journal of High Resolution Chromatography*, 8(8), 415-422.
- Gibson, R. A., German, J. B., Krauss, R. M., Nestel, P., Lamarche, B., Van Staveren, W. A., et al. (2009). A reappraisal of the impact of dairy foods and milk fat on cardiovascular disease risk. *European journal of nutrition*, 48(4), 191-203.

- Gonzalvez, A., Armenta, S. and de la Guardia, M. (2009). Trace-element composition and stable-isotope ratio for discrimination of foods with Protected Designation of Origin. *TrAC Trends in Analytical Chemistry*, 28(11), 1295-1311.
- Grob, K., Lanfranchi, M. and Mariani, C. (1989). Determination of free and esterified sterols and of wax esters in oils and fats by coupled liquid chromatography-gas chromatography. *Journal of Chromatography A*, 471, 397-405.
- Grob, K., Mariani, C., Lanfranchi, M., Artho, A. and Biedermann, M. (1992). CRC Handbook of Chromatography: Analysis and Characterization of Steroids (Vol. III): CRC Press.
- Guillén, M. a. D. and Ruiz, A. (2001). High resolution <sup>1</sup>H nuclear magnetic resonance in the study of edible oils and fats. *Trends in Food Science & Technology*, 12(9), 328-338.
- Guillén, M. D. and Cabo, N. (1997). Infrared spectroscopy in the study of edible oils and fats. *Journal of the Science of Food and Agriculture*, 75(1), 1-11.
- Gunstone, F. (2009). *Oils and fats in the food industry* (Vol. 6): John Wiley & Sons.
- Gunstone, F. and Seth, S. (1994). A study of the distribution of eicosapentaenoic acid and docosahexaenoic acid between the  $\alpha$  and  $\beta$  glycerol chains in fish oils by 13 C-NMR spectroscopy. *Chemistry and physics of lipids*, 72(2), 119-126.
- Gurdeniz, G. and Ozen, B. (2009). Detection of adulteration of extra-virgin olive oil by chemometric analysis of mid-infrared spectral data. *Food chemistry*, 116(2), 519-525.
- Guyot, A. (2002). Recent progress in reactive surfactants in emulsion polymerisation. Proceedings of the 2002 Macromolecular Symposia, 105-132.
- Hammond, E., Irwin, J. and Macrae, R. (1988). HPLC in Food Analysis: Academic Press, Ltd.
- Harker, M., Holmberg, N., Clayton, J. C., Gibbard, C. L., Wallace, A. D., Rawlins, S., et al. (2003). Enhancement of seed phytosterol levels by expression of an N-terminal truncated Hevea brasiliensis (rubber tree) 3-hydroxy-3-methylglutaryl-CoA reductase. *Plant biotechnology journal*, 1(2), 113-121.
- Hatzakis, E., Dagounakis, G., Agiomyrgianaki, A. and Dais, P. (2010). A facile NMR method for the quantification of total, free and esterified sterols in virgin olive oil. *Food Chemistry*, 122(1), 346-352.
- Hoffman, L. and Mellett, F. (2003). Quality characteristics of low fat ostrich meat patties formulated with either pork lard or modified corn starch, soya isolate and water. *Meat science*, 65(2), 869-875.
- Hofman, P., Vickers, M., Gluckman, P., Coveny, A., Cutfield, W., Gertler, A., et al. (2005). Neonatal leptin treatment reverses developmental programming. *Endocrinology*, 146(10), 4211-4216.
- Holmberg, K. (2001). Natural surfactants. Current Opinion in Colloid & Interface Science, 6(2), 148-159.

- Hubbard, W., Sheppard, A., Newkirk, D., Prosser, A. and Osgood, T. (1977). Comparison of various methods for the extraction of total lipids, fatty acids, cholesterol, and other sterols from food products. *Journal of the American Oil Chemists' Society*, 54(2), 81-83.
- Icattoreb, C. A.-G. (2013). Growth outcome: nutritionist perspective. *Nutrition* and Growth, 106, 12-18.
- Indrasti, D., Man, Y. B. C., Mustafa, S. and Hashim, D. M. (2010). Lard detection based on fatty acids profile using comprehensive gas chromatography hyphenated with time-of-flight mass spectrometry. *Food chemistry*, 122(4), 1273-1277.
- Iwadia, M. (1979). Y. Ito, I. Tongai, M. Toyoda, T. Watanabe. Detection of adulterated milkfats by gas-liquid chromatographic determination of butyrate and caproate, J. Food Hyg. Soc. Jpn, 20, 328.
- Jee, M. (2002a). Milk fat and other animal fats. *Oils and Fats Authentication*, 115.
- Jee, M. (2002b). Oils and fats authentication: CRC Press.
- Jensen, R. G. (2002). The composition of bovine milk lipids: January 1995 to December 2000. *Journal of Dairy Science*, 85(2), 295-350.
- Jensen, R. G., Ferris, A. M. and Lammi-Keefe, C. J. (1991). The composition of milk fat. *Journal of Dairy Science*, 74(9), 3228-3243.
- Juárez, M. (1990). Test for fraudulent identification. Proceedings of the 1990 Proceedings of the XXIII International Dairy Congress, 1285-1300.
- Juárez, P., Chase, J. and Blomquist, G. J. (1992). A microsomal fatty acid synthetase from the integument of Blattella germanica synthesizes methyl-branched fatty acids, precursors to hydrocarbon and contact sex pheromone. *Archives of biochemistry and biophysics*, 293(2), 333-341.
- Kagawa, Y., Takaoka, T. and Katsuta, H. (1970). Absence of essential fatty acids in mammalian cell strains cultured in lipid-and protein-free chemically defined synthetic media. *The Journal of Biochemistry*, 68(1), 133-136.
- Kallio, H., Yli-Jokipii, K., Kurvinen, J.-P., Sjövall, O. and Tahvonen, R. (2001). Regioisomerism of triacylglycerols in lard, tallow, yolk, chicken skin, palm oil, palm olein, palm stearin, and a transesterified blend of palm stearin and coconut oil analyzed by tandem mass spectrometry. *Journal* of Agricultural and Food Chemistry, 49(7), 3363-3369.
- Kamm, W., Dionisi, F., Hischenhuber, C. and Engel, K.-H. (2001). Authenticity assessment of fats and oils. *Food Reviews International*, 17(3), 249-290.
- Khalek, A. A. (2014). Young Consumers' Attitude towards Halal Food Outlets and JAKIM's Halal Certification in Malaysia. *Procedia-Social and Behavioral Sciences*, 121, 26-34.
- Kittur, F., Prashanth, K. H., Sankar, K. U. and Tharanathan, R. (2002). Characterization of chitin, chitosan and their carboxymethyl derivatives by differential scanning calorimetry. *Carbohydrate polymers*, 49(2), 185-193.
- Koca, N., Kocaoglu-Vurma, N., Harper, W. and Rodriguez-Saona, L. (2010). Application of temperature-controlled attenuated total reflectance-midinfrared (ATR-MIR) spectroscopy for rapid estimation of butter adulteration. *Food chemistry*, 121(3), 778-782.

- Kowalski, B. (1989). Sub-ambient differential scanning calorimetry of lard and lard contaminated by tallow. *International journal of food science & technology*, 24(4), 415-420.
- Kuksis, A. and McCarthy, M. (1962). Gas-liquid chromatographic fractionation of natural triglyceride mixtures by carbon number. *Canadian journal of biochemistry and physiology*, 40(5), 679-686.
- Kumosinski, T. F. and Farrell Jr, H. M. (1993). Determination of the global secondary structure of proteins by Fourier transform infrared (FTIR) spectroscopy. *Trends in Food Science & Technology*, 4(6), 169-175.
- Lai, Y. W., Kemsley, E. K. and Wilson, R. H. (1994). Potential of Fourier transform infrared spectroscopy for the authentication of vegetable oils. *Journal of Agricultural and Food Chemistry*, 42(5), 1154-1159.
- Lavine, B. K., Moores, A. and Helfend, L. K. (1999). A genetic algorithm for pattern recognition analysis of pyrolysis gas chromatographic data. *Journal of Analytical and Applied Pyrolysis*, 50(1), 47-62.
- Lee, J. H., Jones, K. C., Foglia, T. A., Lee, C.-R., Akoh, C. C. and Lee, K.-T. (2008). High performance liquid chromatographic separation of interesterified palm oil with tributyrin. *LWT Food Science and Technology*, 41(8), 1446-1451.
- Lees, M. (2003). Food authenticity and traceability: Elsevier.
- Lerma-García, M., Ramis-Ramos, G., Herrero-Martínez, J. and Simó-Alfonso, E. (2010a). Authentication of extra virgin olive oils by Fourier-transform infrared spectroscopy. *Food chemistry*, 118(1), 78-83.
- Lerma-García, M. J., Cerretani, L., Cevoli, C., Simó-Alfonso, E. F., Bendini, A. and Toschi, T. G. (2010b). Use of electronic nose to determine defect percentage in oils. Comparison with sensory panel results. *Sensors and Actuators B: Chemical*, In Press, Corrected Proof.
- Lim, H. K., Tan, C. P., Karim, R., Ariffin, A. A. and Bakar, J. (2010). Chemical composition and DSC thermal properties of two species of Hylocereus cacti seed oil: Hylocereus undatus and Hylocereus polyrhizus. *Food Chemistry*, 119(4), 1326-1331.
- Lipp, M. (1995). Review of methods for the analysis of triglycerides in milk fat: application for studies of milk quality and adulteration. *Food chemistry*, 54(2), 213-221.
- Lipp, M. (1996). Determination of the adulteration of butter fat by its triglyceride composition obtained by GC. A comparison of the suitability of PLS and neural networks. *Food chemistry*, 55(4), 389-395.
- Lísa, M., Holcapek, M. and Sovová, H. (2009). Comparison of various types of stationary phases in non-aqueous reversed-phase high-performance liquid chromatography-mass spectrometry of glycerolipids in blackcurrant oil and its enzymatic hydrolysis mixture. *Journal of Chromatography A*, 1216(47), 8371-8378.
- Liu, F., Yusuf, B. L., Zhong, J., Feng, L., He, Y. and Wang, L. (2011). Variety identification of rice vinegars using visible and near infrared spectroscopy and multivariate calibrations. *International Journal of Food Properties*, 14(6), 1264-1276.

- Lorenz-Meyer, H., Bauer, P., Nicolay, C., Schulz, B., Purrmann, J., Fleig, W., et al. (1996). Omega-3 fatty acids and low carbohydrate diet for maintenance of remission in Crohn's disease: a randomized controlled multicenter trial. *Scandinavian journal of gastroenterology*, 31(8), 778-785.
- Luf, W., Stock, A. and Brandl, E. (1987). Detection of Foreign Fat in Milkfat by Means of Triglyceride Analysis. I. Triglyceride-Composition of Austrian Milkfat—Seasonal and Regional Fluctuation. *Oesterr. Milchwirtsch*, 42, 23-28.
- Maggio, R. M., Cerretani, L., Chiavaro, E., Kaufman, T. S. and Bendini, A. (2010). A novel chemometric strategy for the estimation of extra virgin olive oil adulteration with edible oils. *Food Control*, 21(6), 890-895.
- Maggio, R. M., Kaufman, T. S., Carlo, M. D., Cerretani, L., Bendini, A., Cichelli, A., et al. (2009). Monitoring of fatty acid composition in virgin olive oil by Fourier transformed infrared spectroscopy coupled with partial least squares. *Food chemistry*, 114(4), 1549-1554.
- Man, Y. C., Aida, A., Raha, A. and Son, R. (2007). Identification of pork derivatives in food products by species-specific polymerase chain reaction (PCR) for halal verification. *Food Control*, 18(7), 885-889.
- Man, Y. C. and Mirghani, M. (2001). Detection of lard mixed with body fats of chicken, lamb, and cow by fourier transform infrared spectroscopy. *Journal of the American Oil Chemists' Society*, 78(7), 753-761.
- Manaf, M. A., Man, Y. B. C., Hamid, N. S. A., Ismail, A. and Abidin, S. Z. (2007). Analysis of adulteration of virgin coconut oil by palm kernel olein using Fourier transform infrared spectroscopy. *Journal of Food Lipids*, 14(2), 111-121.
- Mannina, L., Luchinat, C., Emanuele, M. C. and Segre, A. (1999). Acyl positional distribution of glycerol tri-esters in vegetable oils: a <sup>13</sup>C NMR study. *Chemistry and physics of lipids*, 103(1), 47-55.
- Mansor, T. S. T., Che Man, Y. B. and Shuhaimi, M. (2012). Employment of Differential Scanning Calorimetry in Detecting Lard Adulteration in Virgin Coconut Oil. Journal of the American Oil Chemists' Society, 89(3), 485-496.
- Mansor, T. S. T., Man, Y. B. C. and Rohman, A. (2011). Application of fast gas chromatography and Fourier transform infrared spectroscopy for analysis of lard adulteration in virgin coconut oil. *Food Analytical Methods*, 4(3), 365-372.
- Marikkar, J., Dzulkifly, M., Nadiha, M. N. and Man, Y. C. (2012). Detection of animal fat contaminations in sunflower oil by differential scanning calorimetry. *International Journal of Food Properties*, 15(3), 683-690.
- Marikkar, J., Ghazali, H., Che Man, Y., Peiris, T. and Lai, O. M. (2005). Distinguishing lard from other animal fats in admixtures of some vegetable oils using liquid chromatographic data coupled with multivariate data analysis. *Food chemistry*, 91(1), 5-14.
- Marikkar, J., Ghazali, H., Long, K. and Lai, O. (2003a). Lard uptake and its detection in selected food products deep-fried in lard. *Food research international*, 36(9), 1047-1060.

- Marikkar, J., Ghazali, H., Man, Y. C. and Lai, O. (2003b). Differential scanning calorimetric analysis for determination of some animal fats as adulterants in palm olein. *Journal of Food Lipids*, 10(1), 63-79.
- Marikkar, J., Lai, O., Ghazali, H. and Man, Y. C. (2001). Detection of lard and randomized lard as adulterants in refined-bleached-deodorized palm oil by differential scanning calorimetry. *Journal of the American Oil Chemists' Society*, 78(11), 1113-1119.
- Marikkar, J., Lai, O., Ghazali, H. and Man, Y. C. (2002). Compositional and thermal analysis of RBD palm oil adulterated with lipase-catalyzed interesterified lard. *Food chemistry*, 76(2), 249-258.
- Matz, S. A. (1992). *Cookie and cracker technology*: AVI Publishing Company, Inc.
- Mayer, K., Fegbeutel, C., Hattar, K., Sibelius, U., Krämer, H.-J., Heuer, K.-U., et al. (2003). ω-3 vs. ω-6 lipid emulsions exert differential influence on neutrophils in septic shock patients: impact on plasma fatty acids and lipid mediator generation. *Intensive care medicine*, 29(9), 1472-1481.
- McClements, D. J., Waraho, T., Cardenia, V., Rodriguez-Estrada, M. T. and Decker, E. A. (2009). Prooxidant mechanisms of free fatty acids in stripped soybean oil-in-water emulsions. *Journal of Agricultural and Food Chemistry*, 57(15), 7112-7117.
- Miklos, R., Zhang, H., Lametsch, R. and Xu, X. (2013). Physicochemical properties of lard-based diacylglycerols in blends with lard. *Food chemistry*, 138(1), 608-614.
- Miyake, T., Petek, H., Takeda, K. and Hinode, K. (1997). Atomic hydrogen enhanced reflow of copper. *Applied Physics Letters*, 70(10), 1239-1241.
- Moses, J. M., Weiss, A. H., Matusek, K. and Guczi, L. (1984). The effect of catalyst treatment on the selective hydrogenation of acetylene over palladium/alumina. *Journal of Catalysis*, 86(2), 417-426.
- Murugaiah, C., Noor, Z. M., Mastakim, M., Bilung, L. M., Selamat, J. and Radu, S. (2009). Meat species identification and Halal authentication analysis using mitochondrial DNA. *Meat Science*, 83(1), 57-61.
- Muuse, B. and Martens, R. (1993). Mixtures of milkfat with non-milkfat: determination of the milkfat content. *Bulletin-International Dairy Federation*, (285), 65-69.
- Nelson, D. and Cox, M. (2005). *I. ehninger principles of biochemistry*: New York: WH Freeman.
- Nicholson, J. K., Lindon, J. C. and Holmes, E. (1999). 'Metabonomics': understanding the metabolic responses of living systems to pathophysiological stimuli via multivariate statistical analysis of biological NMR spectroscopic data. *Xenobiotica*, 29(11), 1181-1189.
- Nor Hayati, I., Man, C., Yaakob, B., Tan, C. P. and Nor Aini, I. (2009). Physicochemical characteristics of soybean oil, palm kernel olein, and their binary blends. *International journal of food science & technology*, 44(1), 152-161.
- Ntakatsane, M., Liu, X. and Zhou, P. (2013). Short communication: Rapid detection of milk fat adulteration with vegetable oil by fluorescence spectroscopy. *Journal of Dairy Science*, 96(4), 2130-2136.

- Ntakatsane, M. P., Liu, X., Zhou, P., Mothibe, K. J., Adegoke, G. O. and Odenya, W. O. (2014). Characterization of fatty acid profile by FFFS. *Journal of Food Measurement and Characterization*, 8(1), 1-8.
- Nurjuliana, M., Man, Y. C. and Hashim, D. M. (2011). Analysis of lard's aroma by an electronic nose for rapid Halal authentication. *Journal of the American Oil Chemists' Society*, 88(1), 75-82.
- O'brien, R. D. (2008). Fats and oils: formulating and processing for applications: CRC press.
- O'brien, R. D. (2010). *Fats and oils: formulating and processing for applications:* CRC press.
- O'Brien, R. D. (2005). Shortenings: types and formulations. *Bailey's Industrial Oil and Fat Products*.
- Ogunniyi, D. (2006). Castor oil: A vital industrial raw material. *Bioresource Technology*, 97(9), 1086-1091.
- Oliveira, F. C. C., Brandal fo, C. R. R., Ramalho, H. F., da Costa, L. A. F., Suarez, P. A. Z. and Rubim, J. C. (2007). Adulteration of diesel/biodiesel blends by vegetable oil as determined by Fourier transform (FT) near infrared spectrometry and FT-Raman spectroscopy. *Analytica Chimica Acta*, 587(2), 194-199.
- Picariello, G., Sacchi, R., Fierro, O., Melck, D., Romano, R., Paduano, A., et al. (2013). High resolution 13C NMR detection of short-and medium-chain synthetic triacylglycerols used in butterfat adulteration. *European journal of lipid science and technology*, 115(8), 858-864.
- Plattner and J, R. D. (1981). Oil Chem. Soc. 58.
- Poulli, K. I., Mousdis, G. A. and Georgiou, C. A. (2007). Rapid synchronous fluorescence method for virgin olive oil adulteration assessment. *Food chemistry*, 105(1), 369-375.
- Precht, D. (1991a). Control of milk fat purity by gas chromatographic triglyceride analysis. *Kieler Milchwirtschaftliche Forschungsberichte*, 43(3), 219-242.
- Precht, D. (1991b). Detection of Adulterated Milk Fat by Fatty Acid an Triglyceride Analysis. *Lipid/Fett*, 93(S4), 538-544.
- Precht, D. (1992a). Detection of foreign fat in milk fat I. Qualitative detection by triacylglycerol formulae. Zeitschrift für Lebensmittel-Untersuchung und Forschung, 194(1), 1-8.
- Precht, D. (1992b). Detection of foreign fat in milk fat II. Quantitative evaluation of foreign fat mixtures. *Zeitschrift für Lebensmittel-Untersuchung und Forschung*, 194(2), 107-114.
- Precht, D. and Molkentin, J. (2000). Identification and quantitation of cis/trans C16: 1 and C17: 1 fatty acid positional isomers in German human milk lipids by thin-layer chromatography and gas chromatography/mass spectrometry. *European journal of lipid science and technology*, (102), 102-113.
- Precht, D., Molkentin, J., Destaillats, F. and Wolff, R. L. (2001). Comparative studies on individual isomeric 18: 1 acids in cow, goat, and ewe milk fats by low-temperature high-resolution capillary gas-liquid chromatography. *Lipids*, 36(8), 827-832.

- Ramos, M. J., Fernández, C. M., Casas, A., Rodríguez, L. and Pérez, Á. (2009). Influence of fatty acid composition of raw materials on biodiesel properties. *Bioresource Technology*, 100(1), 261-268.
- Regenstein, J., Chaudry, M. and Regenstein, C. (2003). The kosher and halal food laws. *Comprehensive Reviews in Food Science and Food Safety*, 2(3), 111-127.
- Reyhan, Boyaci, İ. H., Temiz, T., Shendi, E. G., Yadegari, R. J., Rishkan, M. M., et al. (2014). A rapid method for determination of the origin of meat and meat products based on the extracted fat spectra by using of Raman spectroscopy and chemometric method. *European Food Research and Technology*, 238(5), 845-852.
- Riaz, M. N. and Chaudry, M. M. (2004). The value of Halal food production-Mian N. Riaz and Muhammad M. Chaudry define what Halal and kosher foods are, describe why they are not the same thing, and what is required of processors and. *Inform-International News on Fats Oils and Related Materials*, 15(11), 698-701.
- Robinson, D. J. and Rajah, K. K. (2002). 6 Spreadable products. Fats in Food Technology, 10, 192.
- Rohman, A. and Che Man, Y. (2009). Analysis of Cod-Liver Oil Adulteration Using Fourier Transform Infrared (FTIR) Spectroscopy. *Journal of the American Oil Chemists' Society*, 86(12), 1149-1153.
- Rohman, A., Man, C., Yaakob, B. and Riyanto, S. (2011). Authentication analysis of red fruit (Pandanus Conoideus Lam) oil using FTIR spectroscopy in combination with chemometrics. *Phytochemical Analysis*, 22(5), 462-467.
- Rohman, A. and Man, Y. (2010a). Fourier transform infrared (FTIR) spectroscopy for analysis of extra virgin olive oil adulterated with palm oil. *Food research international*, 43(3), 886-892.
- Rohman, A. and Man, Y. (2012a). Chemical Methods for Determination Oof Lard in Food Products for Halal Authentication Study. *Jurnal Agritech Fakultas Teknologi Pertanian UGM*, 28(04).
- Rohman, A. and Man, Y. B. C. (2010b). Fourier transform infrared (FTIR) spectroscopy for analysis of extra virgin olive oil adulterated with palm oil. *Food Research International*, 43(3), 886-892.
- Rohman, A. and Man, Y. B. C. (2012b). The chemometrics approach applied to FTIR spectral data for the analysis of rice bran oil in extra virgin olive oil. *Chemometrics and intelligent laboratory systems*, 110(1), 129-134.
- Rohman, A. and Sismindari, E. (2011). Analysis of pork adulteration in beef meatball using Fourier transform infrared (FTIR) spectroscopy. *Meat science*, 88(1), 91-95.
- Romía, M. and Bernàrdez, M. (2009). Multivariate calibration for quantitative analysis. *Infrared Spectroscopy for Food Quality Analysis and Control*, 51-82.
- Rutan, S. C. (1989). Fast on-line digital filtering. *Chemometrics and intelligent laboratory systems*, 6(3), 191-201.
- Sacchi, M. C., Barsties, E., Tritto, I., Locatelli, P., Brintzinger, H.-H. and Stehling, U. (1997). Stereochemistry of first monomer insertion into

metal-methyl bond: a tool for evaluating ligand-monomer interactions in propene polymerization with metallocene catalysts. *Macromolecules*, 30(14), 3955-3957.

- Sadło, S., Duda, M., Piechowicz, B. and Jaźwa, A. (2013). Comparative study on disappearance trends of captan and trifloxystrobin residues on fruit and apple tree leaves using internal normalisation method. *Food Additives & Contaminants: Part A*, 30(5), 826-832.
- Saeed, T., Ali, S., Rahman, H. A. and Sawaya, W. (1988). Detection of pork and lard as adulterants in processed meat: liquid chromatographic analysis of derivatized triglycerides. *Journal-Association of Official Analytical Chemists*, 72(6), 921-925.
- Safar, M., Bertrand, D., Robert, P., Devaux, M. and Genot, C. (1994). Characterization of edible oils, butters and margarines by Fourier transform infrared spectroscopy with attenuated total reflectance. *Journal of the American Oil Chemists' Society*, 71(4), 371-377.
- Safwan, Khanfar, M. S. and Obeidat, W. M. (2009). Classification of edible oils and uncovering adulteration of virgin olive oil using FTIR with the aid of chemometrics. *Australian Journal of Basic and Applied Sciences*, 3(3), 2048-2053.
- Schieber, A. (2008). Introduction to Food authentication in Modern techniques for food authentication. *Elsevier Inc, New York, USA*, 1-23.
- Schripsema, J. and Dagnino, D. (2002). Complete assignment of the NMR spectra of [D-Leu1]-microcystin-LR and analysis of its solution structure. *Magnetic Resonance in Chemistry*, 40(9), 614-617.
- Schripsema, J., Vianna, M., Rodrigues, P., de Oliveira, J. and Franco, R. (2008). Metabolomic investigation of fruit flesh gelling of papaya fruit (Carica papaya L.'Golden') by nuclear magnetic resonance and principle component analysis. Proceedings of the 2008 *II International Symposium on Papaya 851*, 505-512.
- Seriburi, V. and Akoh, C. C. (1998). Enzymatic interesterification of lard and high-oleic sunflower oil with Candida antarctica lipase to produce plastic fats. *Journal of the American Oil Chemists' Society*, 75(10), 1339-1345.
- Shin, E.-C., Craft, B. D., Pegg, R. B., Phillips, R. D. and Eitenmiller, R. R. (2010). Chemometric approach to fatty acid profiles in Runner-type peanut cultivars by principal component analysis (PCA). *Food chemistry*, 119(3), 1262-1270.
- Shukla, V., Schiøtz Nielsen, W. and Batsberg, W. (1983). A Simple and Direct Procedure for the Evaluation of Triglyceride Composition of Cocoa Butters by High Performance Liquid Chromatography-A Comparison with the Existing TLC-GC Method. *Fette, Seifen, Anstrichmittel*, 85(7), 274-278.
- Sim, C., Ahmad, M., Ismail, Z., Othman, A., Noor, N. and Zaihidee, E. (2003). Chemometric Classification of Herba Orthosiphon stamineus According to Its Geographical Origin Using Virtual Chemical Sensor Based Upon Fast GC. Sensors, 3(10), 458-471.

- Simoneau, C., Lipp, M., Ulberth, F. and Anklam, E. (2000). Quantification of cocoa butter equivalents in mixtures with cocoa butter by chromatographic methods and multivariate data evaluation. *European Food Research and Technology*, 211(2), 147-152.
- Smejkalová, D. and Piccolo, A. (2010). High-power gradient diffusion NMR spectroscopy for the rapid assessment of extra-virgin olive oil adulteration. *Food Chemistry*, 118(1), 153-158.
- Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E., Johansen-Berg, H., et al. (2004). Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage*, 23, S208-S219.
- Sonntag, N. (1979). Fat splitting. *Journal of the American Oil Chemists' Society*, 56(11), 729A-732A.
- Sonntag, N. O. (1982). Glycerolysis of fats and methyl esters—status, review and critique. *Journal of the American Oil Chemists Society*, 59(10), 795A-802A.
- Sorouraddin, M.-H., Rashidi, M.-R., Ghorbani-Kalhor, E. and Asadpour-Zeynali, K. (2005). Simultaneous spectrofluorimetric and spectrophotometric determination of melatonin and pyridoxine in pharmaceutical preparations by multivariate calibration methods. *Il Farmaco*, 60(5), 451-458.
- Stewart. (2002). Oil and Fats Authentication, Blackwell Publishing, pp. 181–205.
- Svenstrup, G., Brüggemann, D., Kristensen, L., Risbo, J. and Skibsted, L. H. (2005). The influence of pretreatment on pork fat crystallization. *European journal of lipid science and technology*, 107(9), 607-615.
- Syahariza, Z., Man, Y., Selamat, J. and Bakar, J. (2005). Detection of lard adulteration in cake formulation by Fourier transform infrared (FTIR) spectroscopy. *Food chemistry*, 92(2), 365-371.
- Tan, C. and Man, Y. C. (2000). Differential scanning calorimetric analysis of edible oils: comparison of thermal properties and chemical composition. *Journal of the American Oil Chemists' Society*, 77(2), 143-155.
- Timms, R. E. (1980). Detection and quantification of non-milk fat in mixtures of milk and non-milk fats. *Journal of Dairy Research*, 47(03), 295-303.
- Todeschini, R. and Consonni, V. (2009). Molecular Descriptors for Chemoinformatics, Volume 41 (2 Volume Set) (Vol. 41): John Wiley & Sons.
- Tokach, M. D. (2013). What Factors Need To Be Considered When Doing Fat Quality Research-Perspective of A Nutritionist. Proceedings of the 2013 2013 Midwest Meeting (March 11-13, 2013),
- Toppino, P., Contarini, G., Amelotti, G., Traversi, A. and Gargano, A. (1982). Parametri gascromatografici di valutazione della genuinita del burro. *Rivista Italiana delle Sostanze Grasse*.
- Tunick, M. H., Basch, J. J., Maleeff, B. E., Flanagan, J. F. and Holsinger, V. (1989). Characterization of natural and imitation Mozzarella cheeses by differential scanning calorimetry. *Journal of Dairy Science*, 72(8), 1976-1980.

- Ulberth, F. (1995). Quantitation of foreign fat in foreign fat/milkfat mixtures by multivariate regression analysis of fatty acid data. *Journal of Agricultural and Food Chemistry*, 43(6), 1556-1560.
- Ulberth, F. (2003). Testing the authenticity of milk and milk products. *Dairy Processing. Improving Quality*, 208.
- Ulberth, F. and Buchgraber, M. (2000). Authenticity of fats and oils. *European journal of lipid science and technology*, 102(11), 687-694.
- Vaclavik, V. A. and Christian, E. W. (2014). Fat and oil products *Essentials of Food Science* (pp. 233-261): Springer.
- Van de Voort, F., Sedman, J. and Ismail, A. (1993). A rapid FTIR quality-control method for determining fat and moisture in high-fat products. *Food chemistry*, 48(2), 213-221.
- Van Ruth, S., Koot, A., Akkermans, W., Araghipour, N., Rozijn, M., Baltussen, M., et al. (2008). Butter and butter oil classification by PTR-MS. *European Food Research and Technology*, 227(1), 307-317.
- Vogels, J. T., Terwel, L., Tas, A. C., van den Berg, F., Dukel, F. and van der Greef, J. (1996). Detection of adulteration in orange juices by a new screening method using proton NMR spectroscopy in combination with pattern recognition techniques. *Journal of Agricultural and Food Chemistry*, 44(1), 175-180.
- Wang, J., Kliks, M. M., Jun, S., Jackson, M. and Li, Q. X. (2010). Rapid analysis of glucose, fructose, sucrose, and maltose in honeys from different geographic regions using Fourier transform infrared spectroscopy and multivariate analysis. *Journal of food science*, 75(2), C208-C214.
- Watkins, S. M., Carter, L. C., Mak, J., Tsau, J., Yamamoto, S. and German, J. B. (1999). Butyric acid and tributyrin induce apoptosis in human hepatic tumour cells. *Journal of Dairy Research*, 66(04), 559-567.
- Weiss, A. H., Sarkany, A. and Guczi, L. (1984). On the aging phenomenon in palladium catalysed acetylene hydrogenation. *Applied catalysis*, 10(3), 369-388.
- Wootton, M. and Bamunuarachchi, A. (1980). Application of differential scanning calorimetry to starch gelatinization. III. Effect of sucrose and sodium chloride. *Starch-Stärke*, 32(4), 126-129.
- Yahya Ishmael, R. (2005). When ignorance is not bliss. *The Halal Journal*, 28-29.
- Yakovlev, V. (2003). Long flying component produced by protons at energies above 12 TeV in the ionization calorimeter. *Nuclear Physics B*-*Proceedings Supplements*, 122, 417-421.
- Yang, H., Irudayaraj, J. and Paradkar, M. M. (2005). Discriminant analysis of edible oils and fats by FTIR, FT-NIR and FT-Raman spectroscopy. *Food chemistry*, 93(1), 25-32.
- Yang, L. (Producer). (2008). China's Tainted Milk Scandal of 2008. Retrieved from http://content.time.com/time/world/article/0,8599,1844750,00.html
- Yanty, N., Marikkar, J., Man, Y. C. and Long, K. (2011). Composition and thermal analysis of lard stearin and lard olein. *Journal of oleo science*, 60(7), 333-338.

### LIST OF PUBLICATIONS

### Journals:-

- 1. **Nurrulhidayah A.F.**, Y.B Che Man, Amin I., Shuhaimi M., Alfi Khatib."FTIR-ATR spectroscopy based metabolite fingerprinting as a direct determination of butter adulterated with lard. International Journal of Food Properties. DOI: 10.1080/10942912.2012.692224.
- Nurrulhidayah A.F., A. Rohman, Y.B Che Man, Amin I., Shuhaimi M., Alfi Khatib.2012 "Authentication analysis of butter from beef fat using FTIR spectroscopy coupled with chemometrics. International Food Research Journal 20(3): 1383-1388.
- Nurrulhidayah A.F., A. Rohman, Y.B Che Man, Amin I., Shuhaimi M., Alfi Khatib. 2013 "Application of FTIR-ATR spectroscopy coupled with multivariate analysis of rapid estimation of butter adulteration. Journal of Oleo Science 62 (8): 555-562.
- 4. **Nurrulhidayah A.F.**, A. Rohman, Y.B Che Man, Amin I., Shuhaimi M., Alfi Khatib. 2013 "Analysis of chicken fat as adulterant in butter using Fourier Transform Infrared Spectroscopy and chemometrics. Grasias Y Aceitas 64 (4): 349-355.

### Awards:-

- Nurrulhidayah A.F.,Y.B Che Man, Amin I., Shuhaimi M., Alfi Khatib. FTIR-ATR spectroscopy based metabolite fingerprinting as a direct determination of butter adulterated with lard. Pameran Rekacipta, Penyelidikan dan Inovasi 2012, Universiti Putra Malaysia (Gold medal).
- 2. **Nurrulhidayah A.F.**,Y.B Che Man, Amin I., Shuhaimi M., Alfi Khatib, A.Rohman. Analysis butter adulterated with lard using FTIR-<sup>1</sup>H-NMR spectroscopy. **World Halal Research 2012 (TOP 5 Winner-Gold).**
- Nurrulhidayah A.F., Y.B Che Man, Amin I., Shuhaimi M., Alfi Khatib, A. 2012. Analysis butter adulterated with lard using FTIR and <sup>1</sup>H-NMR spectroscopy for Halal verification. Research Innovation, Invention Design 2012. (Gold medal).
- 4. Nurrulhidayah A.F.,Y.B Che Man, Amin I., Shuhaimi M., Alfi Khatib. 2013. Application of FTIR and proton NMR spectroscopy for analysis butter adulterated with lard for Halal verification. Malaysia Technology Expo 2013. (Silver Medal)