



**UNIVERSITI PUTRA MALAYSIA**

***FIELD EVALUATION OF PHAGE FERTILIZER ON CONTROLLING  
Ralstonia solanacearum IN TOMATO AND Erwinia mallotivora IN  
PAPAYA***

**MOHD ZAFRUL ARIF RADHI**

**FP 2016 75**



**FIELD EVALUATION OF PHAGE FERTILIZER ON CONTROLLING  
*Ralstonia solanacearum* IN TOMATO AND *Erwinia mallotivora* IN PAPAYA**

By

**MOHD ZAFRUL ARIF RADHI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra  
Malaysia, in Fulfilment of the Requirements for the Degree of Master of  
Science**

**December 2016**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirement for the degree of Master of Science

**FIELD EVALUATION OF PHAGE FERTILIZER ON CONTROLLING  
*Ralstonia solanacearum* IN TOMATO AND *Erwinia mallotivora* IN PAPAYA**

By

**MOHD ZAFRUL ARIF RADHI**

**December 2016**

**Chairman : Tan Geok Hun, PhD**  
**Faculty : Agriculture**

The results of some previous studies have shown that bacteriophage, a virus that is 'eating' bacteria, by infecting and replicating in the bacterial cell-specific manner have potential to control plant pathogens. Bacterial wilt disease caused by *Ralstonia solanacearum* is one of the major threats to the cultivation of tomatoes in Malaysia. In addition, papaya dieback caused by *Erwinia mallotivora* also brings huge losses to the papaya growers across the country since 2005 until now. In this study, field evaluation was conducted to determine the effectiveness of phage fertilizer for controlling bacterial wilt disease in tomato and papaya dieback in papaya. This fertilizer plus bacteriophage is the one with specific crop nutrient requirements for both crops that has been incorporated with bacteriophage. The attempts on tomato plants showed T1 (ACI (All Cosmos Industries) fertilizer + bacteriophage) recorded the lowest score of 0.5 compared to T2 (ACI fertilizer only) of 1.56 and T3 (control) of 3.02. There were significant differences between T1 and T3 at  $p < 0.05$ . Field study on papaya plants also recorded high percentage of plant mortality for all three treatments, T1, T2 and T3, three months after being inoculated with the pathogens *E. mallotivora*. T1 (ACI fertilizer + bacteriophage) recorded a percentage of 44.74% mortality, T2 (ACI fertilizer only) with 27.5% while T3 (control) with 69.77%. There was no significant difference between the three treatments recorded at  $p < 0.05$ . This proves the specificity of bacteriophage to the host (bacterial pathogens) plays an important role in determining the success of disease control using phage.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Master Sains

**PENILAIAN LAPANGAN BAJA FAJ UNTUK KAWALAN  
*Ralstonia solanacearum* PADA POKOK TOMATO DAN  
*Erwinia mallotivora* PADA POKOK BETIK**

Oleh

**MOHD ZAFRUL ARIF RADHI**

**Disember 2016**

**Pengerusi : Tan Geok Hun, PhD**  
**Fakulti : Pertanian**

Hasil daripada beberapa kajian sebelum ini mendapati bakteriofaj, iaitu sejenis virus 'pemakan' bakteria, berpotensi sebagai kawalan terhadap patogen tumbuhan dengan cara menginfeksi dan mereplikasi di dalam sel bakteria secara spesifik. Penyakit layu bakteria yang disebabkan oleh *Ralstonia solanacearum* merupakan salah satu ancaman utama terhadap penanaman tomato di negara ini. Selain itu, penyakit mati rosot betik yang berpunca daripada *Erwinia mallotivora* juga mendatangkan kerugian besar kepada penanam betik seluruh negara sejak tahun 2005 hingga kini. Dalam kajian ini, penilaian lapangan dilakukan untuk menentukan keberkesanan baja faj dalam mengawal penyakit layu bakteria pada tomato dan penyakit mati rosot pada betik. Baja faj ini merupakan kombinasi baja dengan formulasi khusus mengikut keperluan nutrien untuk kedua-dua tanaman tersebut yang telah dikombinasikan dengan bakteriofaj di dalamnya. Percubaan terhadap tanaman tomato menunjukkan T1 (baja ACI + bakteriofaj) mencatatkan skor penyakit terendah iaitu 0.5 berbanding T2 (baja ACI sahaja) iaitu 1.56 dan T3 (kawalan) iaitu 3.02. Terdapat perbezaan signifikan antara T1 dengan T3 pada  $p < 0.05$ . Kajian lapangan terhadap tanaman betik pula mencatatkan hasil peratus kematian pokok yang tinggi untuk ketiga-tiga rawatan, T1, T2 dan T3 selepas tiga bulan diinokulasikan dengan patogen *E. mallotivora*. T1 (baja ACI + bakteriofaj) mencatatkan peratus kematian sebanyak 44.74%, T2 (baja ACI sahaja) sebanyak 27.5% manakala T3 (kawalan) sebanyak 69.77%. Tiada perbezaan signifikan antara ketiga-tiga rawatan dicatatkan pada  $p < 0.05$ . Ini membuktikan kespesifikan bakteriofaj terhadap perumahnya (bakteria patogen) memainkan peranan penting dalam penentuan kejayaan kawalan penyakit menggunakan faj.

## ACKNOWLEDGEMENTS

First of all, I am heartily thankful to my supervisor, Dr. Tan Geok Hun for her guidance, support and advice for my study. I appreciate her encouragement, efforts, scientific knowledge and understanding. With her excellent guidance and supervision, I am able to carry out the study smoothly. Again, million thanks to Dr. Tan Geok Hun.

Secondly, I wish to show my gratitude to my co-supervisors, Associate Professor Dr. Halimi Mohd Saud and Dr. Mohd Norowi Hamid. My study would not be complete without their crucial contribution, guidance and advice. I really appreciate their patience and encouragement to guide me throughout the study. Besides that, I would like to thanks the students and technical staff from Department of Agriculture Technology and Institute of Bioscience for their guidance, patience and assistance. Thank you for those who were willing to lend their hands to me when I carried out my work.

Nevertheless, I convey my gratitude to my father, mother, beloved wife (Khairunnisa Izzati) and daughter (Aufa Humaira), family members and friends for their understanding and support during the completion of my study. Without their encouragement and support, this study would not have been completed. This study was funded by Ministry of Science, Technology and Innovation Grant Scheme (TechnoFund) (TF0512B074).

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Tan Geok Hun, PhD**

Senior Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Halimi Mohd Saud, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Mohd Norowi Hamid, PhD**

Director  
Biodiversity and Environment Centre  
Malaysian Agricultural Research and Development Institute  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Mohd Zafrul Arif Radhi, GS39770



## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_

Name of Chairman  
of Supervisory  
Committee:

\_\_\_\_\_  
Tan Geok Hun, PhD

Signature: \_\_\_\_\_

Name of Member of  
Supervisory  
Committee:

\_\_\_\_\_  
Halimi Mohd Saud, PhD

Signature: \_\_\_\_\_

Name of Member of  
Supervisory  
Committee:

\_\_\_\_\_  
Mohd Norowi Hamid, PhD

## TABLE OF CONTENTS

	Page
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	ii
<b>ACKNOWLEDGEMENTS</b>	iii
<b>APPROVAL</b>	iv
<b>DECLARATION</b>	vi
<b>LIST OF TABLES</b>	x
<b>LIST OF FIGURES</b>	xi
<b>LIST OF ABBREVIATIONS</b>	xiv
 <b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	<b>3</b>
2.1 History of diseases	3
2.2 <i>Ralstonia solanacearum</i>	3
2.3 <i>Erwinia mallotivora</i>	7
2.4 Bacteriophage	10
2.4.1 History of bacteriophage	10
2.4.2 Application of bacteriophage	10
2.5 Fertilizer	11
2.6 Fermentation	14
2.6.1 Bioreactors	15
2.6.1.1 Principle of bioreactor	16
<b>3 PRODUCTION OF HOST <i>E. coli</i> BY FERMENTATION</b>	<b>19</b>
3.1 Introduction	19
3.2 Materials and methods	19
3.2.1 Fermentation	19
3.2.1.1 Biomass production of phages	19
3.2.1.2 Purification of phages	22
3.2.1.3 Survival rate study	22
3.2.1.4 Plaque assay method for phage titre determination	22
3.3 Results and discussion	22
3.3.1 Host cells production using fermenter	22
3.3.1.1 Biomass count	22
3.3.1.2 Colony forming unit (cfu/mL)	23
3.3.1.3 Optical density (OD)	24
3.3.2 Survival study of bacteriophages	25
3.4 Conclusion	27

<b>4</b>	<b>FIELD EVALUATION OF PHAGE FERTILIZER AGAINST <i>Ralstonia solanacearum</i> IN TOMATO</b>	<b>28</b>
4.1	Introduction	28
4.2	Materials and methods	29
4.2.1	Field trial	29
4.2.1.1	Experimental period	29
4.2.1.2	Layout and plot size	30
4.2.1.3	Treatments	30
4.2.1.4	Fertilization program	30
4.2.1.5	Data recording	30
4.2.1.6	Statistical analysis	30
4.3	Results and discussion	30
4.3.1	Soil analysis before experiment	30
4.3.2	Tomato field trial	31
4.3.2.1	Tomato plant growth	31
4.3.2.2	Disease scoring	33
4.3.2.3	Fruit yield	35
4.4	Conclusion	36
<b>5</b>	<b>FIELD EVALUATION OF PHAGE FERTILIZER AGAINST <i>Erwinia mallotivora</i> IN PAPAYA</b>	<b>37</b>
5.1	Introduction	37
5.2	Materials and methods	37
5.2.1	Preparation of inoculums	37
5.2.2	Field trial	37
5.2.2.1	Experimental period	38
5.2.2.2	Layout and plot size	38
5.2.2.3	Treatments	38
5.2.2.4	Fertilization program	39
5.2.2.5	Data recording	39
5.2.2.6	Statistical analysis	39
5.3	Results and discussion	39
5.3.1	Soil analysis before experiment	39
5.3.2	Papaya field trial	40
5.3.2.1	Papaya plant growth	40
5.3.2.2	Disease scoring	46
5.4	Conclusion	46
<b>6</b>	<b>SUMMARY, GENERAL CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	<b>47</b>
	<b>REFERENCES</b>	<b>48</b>
	<b>APPENDICES</b>	<b>53</b>
	<b>BIODATA OF STUDENT</b>	<b>54</b>
	<b>LIST OF PUBLICATIONS</b>	<b>55</b>

## LIST OF TABLES

Table	Page
1 Different types of fertilizers	11
2 Main parts and their functions in a bioreactor	18
3 Fermentation medium composition for F1	20
4 Fermentation medium composition for F2	20
5 Fermentation parameters for F1 and F2	20
6 Chemical content for pot soil before tomato planting	31
7 The ANOVA result for tomato plant girth week 4 for T3 resulted in no significant difference	33
8 The Kruskal-Wallis rank sum test for tomato plant disease scoring between T1 and T3	34
9 The Kruskal-Wallis rank sum test for tomato plant disease scoring between T1 and T2	34
10 The Kruskal-Wallis rank sum test for tomato plant disease scoring between T2 and T3	35
11 Initial papaya soil analysis	40
12 The Kruskal-Wallis rank sum test for papaya plant height at week 21 between T1 and T3	41
13 The Kruskal-Wallis rank sum test for papaya plant height at week 21 between T1 and T2	41
14 The Kruskal-Wallis rank sum test for papaya plant height at week 21 between T2 and T3	42
15 The Kruskal-Wallis rank sum test for papaya plant height at week 36 between T1 and T3	42
16 The Kruskal-Wallis rank sum test for papaya plant height at week 36 between T1 and T2	42
17 The Kruskal-Wallis rank sum test for papaya plant height at week 36 between T2 and T3	42

18	The Kruskal-Wallis rank sum test for papaya plant girth at week 21 between T1 and T3	44
19	The Kruskal-Wallis rank sum test for papaya plant girth at week 21 between T1 and T2	44
20	The Kruskal-Wallis rank sum test for papaya plant girth at week 21 between T2 and T3	44
21	The Kruskal-Wallis rank sum test for papaya plant girth at week 36 between T1 and T3	45
22	The Kruskal-Wallis rank sum test for papaya plant girth at week 36 between T2 and T3	45
23	The Kruskal-Wallis rank sum test for papaya plant girth at week 36 between T1 and T2	46

## LIST OF FIGURES

Figure	Page
1 Virulent colonies of <i>R. solanacearum</i> cultured on tetrazolium chloride (TZC) agar, indicated by pink dot in the middle of the white colony which differs from the non virulent one which only formed white colonies	5
2 An Image of Gram staining of <i>R. solanacearum</i> under 100X magnification, showing red coloured cells, confirming its Gram negative traits	5
3 Tomato plant infected with <i>R. solanacearum</i>	6
4 Potato plant showing wilting of one stem	6
5 Initial wilt symptoms of the lower leaves of tobacco plant	7
6 Papaya dieback on papaya plant caused by <i>E. mallotivora</i>	8
7 <i>E. mallotivora</i> cultured on NA	9
8 Transmission electron microscope image of the strain of <i>E. mallotivora</i> isolated from dieback-infected papaya tree (Noriha <i>et al.</i> , 2011).	9
9 Smart fertilizer produced in pellet form by All Cosmos Industries Sdn. Bhd.	13
10 Smart fertilizer produced in granule form by Smart Fert Sdn. Bhd	13
11 Smart fertilizer produced in liquid form by Nosktahco Holdings Sdn. Bhd.	14
12 A five litre working volume bioreactor in operation	17
13 General structure of a typical bioreactor	18
14 The 5 L bio reactor used for mass propagation of <i>E. coli</i> host cells, Infors, Switzerland	21
15 <i>E. coli</i> culture aliquoted and added with 1/10 volume phage stock	21
16 Dry cell weight of <i>E. coli</i> ER2738 (g/mL) obtained from two batches of fermentation using LB broth as growth medium (F1) and Terrific Broth as a growth medium (F2)	23

17	Population count for <i>E.coli</i> ER2738 (cfu/mL) obtained from two batches of fermentation, a) LB as growth medium (F1); b) Terrific Broth as growth medium (F2)	24
18	Optical density (OD <sub>600</sub> ) of <i>E. coli</i> strain ER2738 results from two batches of fermentation, a) LB as growth medium (F1); b) Terrific Broth as growth Medium (F2)	25
19	Survival study of bacteriophage P631 in fertilizer. Data was collected from Day 1 until eight months	26
20	Survival study of bacteriophage P45 and P47 in fertilizer. Data was collected from Day 1 until eight months	26
21	Survival study of bacteriophage P459 and P483 in fertilizer. Data was collected from Day 1 until eight months	27
22	Plot layout of the tomato trial in Ladang 2 Glasshouse, UPM	29
23	The height of tomato plants at week 4, week 8 and week 12 under different treatments. The error bars represent the standard deviation from the arithmetic mean	32
24	The stem diameter (girth) of tomato plants at week 4, week 7 and week 12. The error bars represent the standard deviation from the arithmetic mean	32
25	The height of tomato plants at week 4 under different treatments. Bar plot generated using RStudio Statistical Software	33
26	Disease scoring of tomato plants at week 4, week 8 and week 12	35
27	Total fruit yield for different treatments. The error bars represent the standard deviation from the arithmetic mean	36
28	Plot layout of the papaya trial in Ladang 2 open field, UPM	38
29	Papaya plant height at week 21 and week 36. The error bars represent the standard deviation from the arithmetic mean.	41
30	Papaya plant height at week 21. Bar plot generated using RStudio Statistical Software	43
31	Papaya plant girth at week 21 and week 36. The error bars represent the standard deviation from the arithmetic mean.	43
32	Papaya plant girth at week 36. Bar plot generated using RStudio Statistical Software	45

## LIST OF ABBREVIATIONS

ACI	All Cosmos Industries Sdn. Bhd.
ATCC	American Type Culture Collection
CRD	Complete Randomized Design
DNA	Deoxyribonucleic acid
dsRNA	Double stranded ribonucleic acid
NA	Nutrient agar
PBS	Phosphate-buffered saline
PEG	Polyethylene glycol
TBS	Tris-buffered saline



## CHAPTER 1

### INTRODUCTION

Agriculture remains an important sector of Malaysia's economy. In the year of 2014, Malaysia has a total of 7,839,000 ha of agricultural land from its total land area of 32,855,000 ha. Malaysia's Gross Domestic Product (GDP) was RM 1062.8 billion in 2015, which RM 94.1 billion (8.9%) from total GDP contributed by the agriculture sector. Total vegetable export value in 2013 was RM887.59 million (4.02% from total Malaysia food product export). The export value of tomato in 2013 was RM100.7 million (0.46% from total Malaysia food product export). Total fruit export value in 2013 was RM675.73 million (3.06% from total Malaysia food product export). The export value of pineapple (fresh and processed) in 2013 was RM38.61 million (0.17% from total Malaysia food product export) (Department of Agriculture, 2014).

Tomato production in Malaysia is commonly for domestic consumption. Tomato growing areas are usually in the highlands such as Cameron Highlands and Kundasang whereas lowland cultivation can be found in Kelantan, Johor and Pahang. Although there is no published data on the exact losses in Malaysia, bacterial wilt which brought about by a bacterial pathogen which is *Ralstonia solanacearum* dependably be the most wrecking causing agent in tomato and yield misfortunes up to 90 percent. Bacterial wilt is a critical disease, particularly in the lowlands. It was initially written about tomato and potato in 1910 (Bancroft, 1910). The pathogen had been accounted for to taint more than 200 plant species from 53 diverse plant families (Álvarez *et al.*, 2010). The most widely recognized hosts are tomato, banana, eggplant, potato, groundnut and ginger. In fresh market tomato in Taiwan, malady occurrences of 15 to 55% have been accounted for, bringing about misfortunes surpassing 12 million U.S. dollars yearly (Lin *et al.*, 2014).

Papaya or *Carica papaya* is another monetarily noteworthy fruit crop grown in Malaysia with an export value of USD 8.42 million in 2011 (FAOSTAT, 2015). In any case, papaya dieback disease has turned out to be one of the significant threats of this industry in Malaysia as of late. Symptoms commonly recognized include greasy, water soaked lesions and spots on leaves, as well as foliar and angular lesions. These lesions can prompt to optional contamination, which can in the end cause the demise of the papaya plant. This infection has been an issue for papaya producers for just about 10 years, devastating more than one million plants. The infection was initially distinguished in Malaysia close Batu Pahat, Johor in late 2003. Another occurrence was later announced in Bidor, Perak, in October 2004. All the more as of late, Maktar *et al.*, (2008) detailed *E. papayae* as creating papaya dieback in Malaysia. Notwithstanding, the most recent review by Noriha *et al.*, (2011) affirmed that the causal agent for papaya dieback in Peninsular Malaysia is *Erwinia mallotivora*.

## REFERENCES

- Agrios, G.N. (2005). Plant pathology, 5th ed. *Elsevier Academic, London, United Kingdom*. p 398–401.
- Akhtar, M. and Malik, A. (2000). Roles of organic soil amendments and soil organisms in the biological control of plant-parasitic nematodes: a review. *Bioresource Technology*, **74(1)**: 35-47.
- Álvarez, B., Biosca, E. G., and López, M. M. (2010). On the life of *Ralstonia solanacearum*, a destructive bacterial plant pathogen. *Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology*, **1**: 267–279.
- Bailey, K.L. and Lazarovitz, G. (2003). Suppressing soil-borne diseases with residue management and organic amendments. *Soil and Tillage Research*, **72**, 169-180.
- Bancroft, C. K. (1910). Researches on the Life-history of Parasitic Fungi. *Annals of Botany*, **24(94)**: 359-372.
- Cavalieri, D., McGovern, P. E., Hartl, D. L., Mortimer, R., & Polsinelli, M. (2003). Evidence for *S. cerevisiae* fermentation in ancient wine. *Journal of Molecular Evolution*, **57(1)**: S226-S232.
- Department of Agriculture Malaysia (2014). Fruit Crops Statistic 2014.
- Department of Agriculture Malaysia (2014). Vegetables and Cash Crops Statistic 2014.
- d'Hérelle, F. (1949). "The bacteriophage" (PDF). *Science News* **14**: 44–59. Retrieved 5 September 2015.
- Denny, T. P. and Baek, S. R. (1991). Genetic evidence that extracellular polysaccharide is a virulence factor of *Pseudomonas solanacearum*. *Molecular Plant-Microbe Interaction*, **4**: 198-206.
- Dirar, H. A. (1993). *The indigenous fermented foods of the Sudan: a study in African food and nutrition*. CAB International. pp.xvii + 552 pp. ref.43
- Dittmar, H., Drach, M., Vosskamp, R., Trenkel, M.E., Gutser, R. and Steffens, G. (2009). "Ullmann's Encyclopedia of Industrial Chemistry". doi:10.1002/14356007.n10\_n01. ISBN 3527306730
- FAO (2007). State of the World's Forests 2007. Rome. [www.fao.org/docrep/009/a0773e/a0773e00.htm](http://www.fao.org/docrep/009/a0773e/a0773e00.htm).

- Faostat, F. (2015). Agriculture Organization of the United Nations, 2011.FAO, Retrieved am from <http://faostat3.fao.org/faostat-gateway/go/to/download/Q/QC/S>. Acceso, 20.
- Fox, J.L. (2000). Phage treatments yield healthier tomato, pepper plants. *ASM News*, **66**, 455–456.
- Genin, S., and Boucher, C. (2002). *Ralstonia solanacearum*: secrets of a major pathogen unveiled by analysis of its genome. *Molecular plant pathology*, **3(3)**: 111-118.
- González, E. T., & Allen, C. (2003). Characterization of a *Ralstonia solanacearum* operon required for polygalacturonate degradation and uptake of galacturonic acid. *Molecular plant-microbe interactions*, **16(6)**: 536-544.
- Gough, L. P. (1993). *Understanding our fragile environment; lessons from geochemical studies*. Report (No. 1105). pp. 1-34, USGPO; USGS Map Distribution.
- Grath, S. and Van Sinderen, D. (editors) (2007). *Bacteriophage: Genetics and Molecular Biology* (1st ed.). Caister Academic Press. ISBN 978-1-904455-14-1
- Guest, J. R., Green, J., Irvine, A. S., and Spiro, S. (1996). Regulation of Gene Expression in *Escherichia coli* (Lin, E. C. C., and Lynch, A. S., eds) pp. 317–342, R. G. Landes & Co, Austin, TX
- Hamidah, S. and Lum, K. Y. (1993). Bacterial wilt of groundnut in Malaysia. In *Groundnut bacterial wilt: proceedings of the Second Working Group Meeting, 2 Nov 1992, Asian Vegetable Research and Development Center, Tainan, Taiwan* (Vol. 502, p. 6)
- Hayward, A. C. (1991). Biology and epidemiology of bacterial wilt caused by *Pseudomonas solanacearum*. *Annual review of phytopathology*, **29(1)**: 65-87.
- Hayward, A.C. (2000). *Ralstonia solanacearum*. In: Encyclopedia of microbiology, Vol. 4, (Lederberg, L., ed.), p. 32– 42. Don Mills, Ont.: Academic Press Canada
- Kiiski, H., Dittmar, H., Drach, M., Vosskamp, R., Trenkel, M. E., Gutser, R. and Steffens, G. (2016). Fertilizers, 2. Types. *Ullmann's Encyclopedia of Industrial Chemistry*. 1–53.
- IUPAC, *Compendium of Chemical Terminology*, 2nd ed. (the "Gold Book") (1997). Online corrected version: (2006–) "bioreactor"

- Jones, J.B., Lacy, G.H., Bouzar, H., Stall, R.E., Schaad, N.W. (2004). Reclassification of the Xanthomonads Associated with bacterial spot disease of tomato and pepper, *System. & Appl Microbiol.*, **27**:755–62
- Jones, J. B., Vallad, G. E., Iriarte, F. B., Obradović, A., Wernsing, M. H., Jackson, L. E., Momol, M. T. (2012). Considerations for using bacteriophages for plant disease control. *Bacteriophage*, **2(4)**, 208–214.
- Kang, Y., Liu, H., Genin, S., Schell, M. A. and Denny, T. P. (2002). *Ralstonia solanacearum* requires type 4 pili to adhere to multiple surfaces and for natural transformation and virulence. *Molecular microbiology*, **46(2)**: 427-437.
- Keen, E. C. (2012). "Phage Therapy: Concept to Cure". *Frontiers in Microbiology* **3**. doi:10.3389/fmicb.2012.00238.PMC3400130.PMID 22833738
- Klein, Donald W.; Lansing M.; Harley, John (2006). *Microbiology* (6th ed.). New York:McGraw-Hill.ISBN 978-0-07-255678-0
- Lin, C.H., Tsai, K.C., Prior, P. and Wang, J.F. (2014). Phylogenetic relationships and population structure of *Ralstonia solanacearum* isolated from diverse origins in Taiwan. *Plant Pathology*, **63**: 1395–1403
- Lopez A, Lazaro N, Marques AM. (1997). The interphase technique: a simple method of cell immobilization in gel-beads. *Journal Microbiology Methods*, **30**: 231-234.
- Maktar, N.H., Kamis, S., Mohd Yusof, F.Z. and Hussain, N.H., (2008). *Erwinia papayae* causing papaya dieback in Malaysia. *Plant Pathology* 2008, **57**: 774.
- Masyitah (2004). Development of disease suppressive compost and potting mix for control of bacterial wilt of tomato. Master's Thesis, Universiti Putra Malaysia
- McGovern, P. E., Zhang, J., Tang, J., Zhang, Z., Hall, G. R., Moreau, R. A. and Cheng, G. (2004). Fermented beverages of pre-and proto-historic China. *Proceedings of the National Academy of Sciences of the United States of America*, **101(51)**: 17593-17598.
- Noriha, M.A., Hamidun B., Rohaiza A. R. and Indu Bala S. J., (2011). *Erwinia mallotivora* sp., a New Pathogen of Papaya (*Carica papaya*) in Peninsular Malaysia. *International Journal of Molecular Science* 2011, **12(1)**: 39-45
- Obradovic, A., Jones, J.B., Momol, M.T., Balogh, B. and Olson, S.M. (2004). Management of tomato bacterial spot in the field by foliar applications of bacteriophages and SAR inducers. *Plant Dis*, **88**, 736–740.

- Oyedun, O.S., Kufu, F.O. and Nwanguma, E.I. (1997). Bacterial wilt in the tomato cropping systems of Nigeria: Its prevalence and yield loss. *Proceedings of the 2nd International Bacterial Wilt Symposium*, Gosier, Guadeloupe, France, June 22-27
- Peinado, R. A., Moreno, J. J., Villalba, J. M., González-Reyes, J. A., Ortega, J. M. and Mauricio, J. C. (2006). Yeast biocapsules: a new immobilization method and their applications. *Enzyme and Microbial Technology*, **40(1)**: 79-84.
- Peng, S.H., Wan-Azha, W.M., Wong, W.Z., Go, W.Z., Chai, E.W., Chin, K.L. and H'ng, P.S. (2013). Effect of Using Agro-fertilizers and N-fixing *Azotobacter* Enhanced Biofertilizers on the Growth and Yield of Corn. *Journal of Applied Sciences*, **13**: 508-512.
- Prescott, L. (1993). Microbiology, Wm. C. Brown Publishers, ISBN 0-697-01372-3
- Rabu, M.R. and Mat Lin, R., (2005). Prospect of papaya in the world market: Malaysia perspective. *Proceeding of First International Symposium on Papaya, Genting Highlands, Malaysia*.
- Ratlidge, C., Kanagachandran, K., Anderson, A. J., Grantham, D. J. and Stephenson, J. C. (2001). Production of docosahexaenoic acid by *Cryptocodinium cohnii* grown in a pH-auxostat culture with acetic acid as principal carbon source. *Lipids*, **36(11)**: 1241-1246.
- Salanoubat, M., Genin, S., Artiguenave, F., Gouzy, J., Mangelot, S., Arlat, M. and Chandler, M. (2002). Genome sequence of the plant pathogen *Ralstonia solanacearum*. *Nature*, **415(6871)**: 497-502.
- Sartain, J.B. University of Florida (2011). "Food for turf: Slow-release nitrogen". *Grounds Maintenance*.
- Schell, M. A. (2000). Control of virulence and pathogenicity genes of *Ralstonia solanacearum* by an elaborate sensory network. *Annual review of phytopathology*, **38(1)**: 263-292.
- Schnabel, E.L. and Jones, A.L. (2001). Isolation and characterization of five *Erwinia amylovora* bacteriophages and assessment of phage resistance in strains of *Erwinia amylovora*. *Appl Environ Microbiol*, **67**, 59–64.
- Smith, G.P. and Scott, J.K. (1993). Libraries of peptides and proteins displayed on filamentous phage. *Methods in Enzymology*. **217**: 228-257.
- Tahat, M. M., & Sijam, K. (2010). *Ralstonia solanacearum*: The Bacterial Wilt Causal Agent. *Asian Journal of Plant Sciences*, **9(7)**, 385-393.



- Tahat, M. M., Sijam, K., & Othman, R. (2012). The potential of endomycorrhizal fungi in controlling tomato bacterial wilt *Ralstonia solanacearum* under glasshouse conditions. *African Journal of Biotechnology*, **11**(67): 13085-13094.
- Tan, G.H., Nordin, M.S. and Napsiah, A.B., (2008). Isolation and characterization of lytic bacteriophages from sewage water. *Journal of Tropical Agriculture and Food Science*, **36** (1): 287-291
- Tan, G.H., Nordin, M.S., Kert, T.L., Napsiah, A.B. and Jeffrey, L.S.H., (2009). Isolation of beneficial microbes from biofertilizer products. *Journal of Tropical Agriculture and Food Science*, **37**(1): 103-109
- Tan, G.H., Nordin, M.S., Napsiah, A.B. and Rosnah, H., (2009). The Lysis Activity of Bacteriophages Isolated from Sewage against *Ralstonia solanacearum* and *Erwinia chrysanthemi*. *Journal of Tropical Agriculture and Food Science*, **7**(2): 203-209
- Tan, G.H., Nordin, M.S. and Napsiah, A.B., (2010). The effect of phage infection on pathogenic activity of *Ralstonia solanacearum* in tomato. *Journal of Tropical Agriculture and Food Science*, **38**(1): 123-130
- Tans-Kersten, J., Huang, H. and Allen, C. (2001). *Ralstonia solanacearum* needs motility for invasive virulence on tomato. *Journal of Bacteriology*, **183**(12): 3597-3605.
- Vouillamoz, J. F., McGovern, P. E., Ergul, A., Söylemezoğlu, G., Tevzadze, G., Meredith, C. P. and Grando, M. S. (2006). Genetic characterization and relationships of traditional grape cultivars from Transcaucasia and Anatolia. *Plant Genetic Resources: characterization and utilization*, **4**(2): 144-158.
- Williams, J. A. (2002). Keys to bioreactor selections. *Chemical engineering progress*, **98**(3): 34-41.
- Wommack, K. E. and Colwell, R. R. (2000). "Virioplankton: Viruses in Aquatic Ecosystem". *Microbiology and Molecular Biology Reviews*. **64** (1): 69–114.
- Yabuuchi, E., Kosako, V., Yano, I., Hotta, H. and Nishiuchi, Y. (1995). Transfer of two *Burkholderia* and an *Alcaligenes* species to *Ralstonia* gen. nov.: proposal of *Ralstonia pickettii* (Ralston, Palleroni and Doudoroff 1973) comb. nov., *Ralstonia solanacearum* (Smith 1896) comb. nov. and *Ralstonia eutropha* (Davis 1969) comb. nov. *Microbiology Immunology*, **39**: 897–904.
- Yamada, T., Kawasaki, T., Nagata, S., Fujiwara, A., Usami, S. and Fujie, M. (2007). New bacteriophages that infect the phytopathogen *Ralstonia solanacearum*. *Microbiology* **153**: 2630–2639

## BIODATA OF STUDENT

Mohd Zafrul Arif was born in Johor Bahru, Johor, Malaysia on 11<sup>th</sup> September 1984. He obtained his early education in Sekolah Kebangsaan Taman Tun Aminah, Skudai in 1991 and Sekolah Kebangsaan Sungai Tua Baru, Batu Caves from 1992 to 1996. He continued secondary education in Sekolah Alam Shah, Kuala Lumpur (SBP) from 1997 to 2001 and later enrolled in Kolej Matrikulasi Negeri Sembilan, Kuala Pilah for one year in 2002. He then pursued his tertiary education in Universiti Kebangsaan Malaysia, Bangi in year 2003 and did receive his Bachelor of Science (Hons.) in Microbiology in year 2006, under the Jabatan Perkhidmatan Awam (JPA) scholarship. After graduated, he joined Malaysian Agricultural Research and Development Institute (MARDI) as a research officer in the Strategic Resource Research Centre from 2007 until now. During his service with MARDI, he led few researches such as MARDI's Short Term Grant (indigenous microorganism research) and MARDI's RMKe-10 Mega Projects (development of biofertilizer standards and organic goat and chicken research) and MOSTI's Science Fund Project (vermicompost research). He was also an active collaborator in few projects such as the Development of Taman Tropika Kenyir in Terengganu, Integrated Organic Farming and Research on Climate Change. In 2014, he was offered to further his study by doing Master Degree in Agriculture Technology in Universiti Putra Malaysia under MARDI's scholarship.

## LIST OF PUBLICATIONS

- Mohd Zafrul Arif, R.**, Tan, G.H., Wan Azha, W.M. and Tony Peng, S.H. (2015). Field evaluation of Eksotika II (*Carica papaya* L.) against *Erwinia mallotivora* sp., causal agent for papaya dieback in Peninsular Malaysia. *Proceeding of International Congress of the Malaysian Society for Microbiology 2015* (ICMSM 2015), Bayview Beach Resort, Penang, Malaysia, 7-10 December 2015, P:68-70
- M. Z. A. Radhi**, M. B. Adam, H.M. Saud, M.N. Hamid, P.S.H. Tony and G.H. Tan (2016). Efficacy of Smart Fertilizer for Combating Bacterial Wilt Disease in *Solanum lycopersicum*. *Direct Research Journal of Agriculture and Food Science*, **4(7)**, pp. 137-143
- Radhi, M.Z.A.**, Malik, Z., Ibrahim, I.Z., Wan Othman, W.Y.F. and Tan, G.H. (2014). Evaluation of Different Diets for Organically Reared Kampung Chicken, *Proceeding of International Agriculture Congress*, 25-27th November 2014, Pullman Putrajaya Lakeside, Putrajaya, Malaysia.





UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : \_\_\_\_\_

TITLE OF THESIS / PROJECT REPORT :

FIELD EVALUATION OF PHAGE FERTILIZER ON CONTROLLING *Ralstonia solanacearum*  
IN TOMATO AND *Erwinia mallotivora* IN PAPAYA

NAME OF STUDENT: MOHD ZAFRUL ARIF RADHI

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

\*Please tick (✓)

☐

**CONFIDENTIAL**

(Contain confidential information under Official Secret Act 1972).

☐

**RESTRICTED**

(Contains restricted information as specified by the organization/institution where research was done).

☐

**OPEN ACCESS**

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

☐

**PATENT**

Embargo from \_\_\_\_\_ until \_\_\_\_\_  
(date) (date)

**Approved by:**

\_\_\_\_\_  
(Signature of Student)  
New IC No/ Passport No.:

Date :

\_\_\_\_\_  
(Signature of Chairman of Supervisory Committee)  
Name:

Date :

[Note : If the thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach with the letter from the organization/institution with period and reasons for confidentiality or restricted. ]