



UNIVERSITI PUTRA MALAYSIA

***HEAVY METALS IN SOIL FROM URBAN AREAS OF KLANG,
MALAYSIA, AND THEIR HEALTH RISK ASSESSMENT USING IN
VITRO DIGESTION METHOD***

NURUL SYAZANI YUSWIR

FPSK(m) 2015 56



**HEAVY METALS IN SOIL FROM URBAN AREAS OF KLANG, MALAYSIA,
AND THEIR HEALTH RISK ASSESSMENT USING *IN VITRO* DIGESTION
METHOD**

By

NURUL SYAZANI BINTI YUSWIR

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of
Master of Science**

October 2015

All materials contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Degree of Master of Science

HEAVY METALS IN SOIL FROM URBAN AREAS OF KLANG, MALAYSIA, AND THEIR HEALTH RISK ASSESSMENT USING *IN VITRO* DIGESTION METHOD

By

NURUL SYAZANI BINTI YUSWIR

October 2015

Chair : Sarva Mangala Praveena, PhD
Faculty : Medicine and Health Sciences

Background: Klang District is considered as one of the area that undergoes a rapid urbanization and surrounded by land and sea based anthropogenic activities. These anthropogenic activities have given various impacts on environmental and human health. **Objective:** Thus, the aim of this study was to determine the total and bioavailable of heavy metal concentrations (Al, Fe, Zn, Cu, Co, Cd, Pb and Cr) in urban surface soil samples. In addition, this study also aimed to determine possible sources in Klang District urban surface soil using multivariate analysis. Besides, this study also aimed to determine health risks posed by bioavailable of heavy metal in urban soil on adult and children. **Methodology:** A total of 76 urban soils in Klang District have been sampled and analyzed for total and bioavailable of heavy metal concentrations (Al, Fe, Zn, Cu, Co, Cd, Pb and Cr) using Inductively Coupled Plasma-Optical Emission Spectrometry. Total heavy metal concentration was determined using *aqua regia* digestion method while bioavailable of heavy metal concentration was determined using Physiologically Based Extraction Test (PBET) *in vitro* digestion method. **Result:** Results showed that the bioavailable of heavy metal concentrations were in the order of Al (25 mg/kg), Fe (6.7 mg/kg), Zn (5.6 mg/kg), Cu (3.0 mg/kg), Co (0.22 mg/kg), Cd (0.14 mg/kg), Pb (0.11 mg/kg) and Cr (0.10 mg/kg), while concentration of total heavy metal were in the order of Fe (9090 mg/kg), Al (6171 mg/kg), Cu (294 mg/kg), Zn (276 mg/kg), Pb (53 mg/kg), Cr (16 mg/kg), Co (1.2 mg/kg) and Cd (0.71 mg/kg). From the Spearman correlation coefficient (r) value, significant correlation were observed for Al-Fe (r = 0.681), Cd-Co (r = 0.495), Cu-Zn (r = 0.232), Fe-Pb (r = 0.260), Fe-Zn (r = 0.239), Al-Cu (r = -0.503), Co-Pb (r = -0.241) and Cu-Fe (r = -0.492). Spearman correlation output showed that heavy metal such as Al and Fe may come from natural sources while heavy metal such as Cd, Co, Cr and Cu may come from anthropogenic sources. Cluster Analysis output showed that these heavy metals can be classified into four clusters namely Cluster 1 which consisted of Cd, Cr, Co and Pb, Cluster 2 consisted of Zn and Cu, Cluster 3 consisted of Fe and Cluster 4 consisted of Al. For Clusters 1 and 2, anthropogenic activities were believed as the sources while for Clusters 3 and 4, the heavy metals originated from natural

sources. For the health risks, the results of Hazard Quotient and Total Risk showed that heavy metal contamination via soil ingestion pathway in Klang District may not pose carcinogenic and non-carcinogenic risk to human adult, but it may pose carcinogenic and non-carcinogenic risks to children. **Conclusion:** Output from this study can be used to fill the knowledge gap on effect of heavy metal and potential health risks due to heavy metal exposure in urban area especially in Malaysia.

Keywords: Urban soil, heavy metal, total, bioavailable, health risks



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Master Sains

**LOGAM BERAT DALAM TANAH DARI KAWASAN BANDAR KLANG,
MALAYSIA, DAN PENILAIAN RISIKO KESIHATAN MENGGUNAKAN
KAEDAH PENGHADAMAN *IN VITRO***

Oleh

NURUL SYAZANI BINTI YUSWIR

Oktober 2015

Pengerusi : Sarva Mangala Praveena, PhD
Fakulti : Perubatan dan Sains Kesihatan

Pendahuluan: Daerah Klang dianggap sebagai salah satu kawasan yang mengalami pembangunan pambangunan yang pesat dan dikelilingi oleh aktiviti antropogenik berdasarkan tanah dan laut. Aktiviti-aktiviti antropogenik ini telah memberikan pelbagai kesan terhadap alam sekitar dan juga kesihatan manusia. **Objektif:** Oleh itu, tujuan kajian ini adalah untuk menentukan kepekatan total dan kebolehdapatan logam berat (Al, Fe, Zn, Cu, Co, Cd, Pb dan Cr) di dalam permukaan sampel tanah di kawasan bandar. Di samping itu, kajian ini juga bertujuan untuk mengenal pasti hubungan antara kebolehdapatan logam berat, dan sumbernya di dalam permukaan tanah di bandar Klang dengan menggunakan analisis multivariat. Selain itu, kajian ini juga bertujuan untuk menentukan risiko kesihatan pada orang dewasa dan kanak-kanak di bandar Klang melalui kaedah kebolehdapatan logam berat di dalam tanah. **Metodologi:** Sebanyak 76 sampel tanah di kawasan bandar di Daerah Klang telah disampel dan dianalisis untuk menilai kandungan kepekatan total dan kebolehdapatan logam berat (Al, Fe, Zn, Cu, Co, Cd, Pb dan Cr) menggunakan Inductively Coupled Plasma-Optical Emission Spectrometry. Total kepekatan logam berat telah ditentukan dengan menggunakan kaedah “*aqua regia*” manakala kebolehdapatan kepekatan logam berat telah ditentukan dengan menggunakan kaedah Physiologically Based Extraction Test (PBET). **Hasil kajian:** Keputusan menunjukkan bahawa kebolehdapatan logam berat adalah dalam susunan Al (25 mg / kg), Fe (6.7 mg / kg), Zn (5.6 mg / kg), Cu (3 mg / kg), Co (0.22 mg / kg), Cd (0.14 mg / kg), Pb (0.11 mg / kg) dan Cr (0.10 mg / kg), manakala kepekatan total logam berat adalah dalam susunan Fe (9090 mg / kg), Al (6171 mg / kg), Cu (294 mg / kg), Zn (276 mg / kg), Pb (53 mg / kg), Cr (16 mg / kg), Co (1.2 mg / kg) dan Cd (0.71 mg / kg). Daripada nilai pekali korelasi Spearman (r), hubungan yang signifikan adalah Al-Fe (r = 0.681), Cd-Co (r = 0.495), Cu-Zn (r = 0.232), Fe-Pb (r = 0.260), Fe-Zn (r = 0.239), Al-Cu (r = -0.503), Co-Pb (r = -0.241) dan Cu-Fe (r = -0.492). Nilai pekali korelasi Spearman (r) di antara 0.681 - 0.503 menunjukkan bahawa logam berat seperti Al dan Fe mungkin datang daripada sumber semula jadi manakala logam berat seperti Cd, Co, Cr dan Cu mungkin datang daripada sumber antropogenik. Dapatan daripada Analisis Kluster menunjukkan bahawa logam berat ini boleh dikelaskan kepada empat kelompok di

mana Kluster 1 terdiri daripada Cd, Cr, Co dan Pb, Kluster 2 terdiri daripada Zn dan Cu, Kluster 3 terdiri daripada Fe dan Kluster 4 terdiri daripada Al. Untuk Kluster 1 dan 2, aktiviti antropogenik dipercayai sebagai sumber kepada kandungan logam berat tersebut manakala bagi Kluster 3 dan 4, logam berat tersebut mungkin berasal daripada sumber semula jadi. Bagi Penilaian Risiko Kesihatan, keputusan Hazard Quotient dan Total Risk menunjukkan bahawa pencemaran logam berat melalui tanah di daerah Klang tidak mungkin menimbulkan risiko karsinogenik dan bukan karsinogenik kepada golongan dewasa, tetapi ia boleh menimbulkan risiko karsinogenik dan bukan karsinogenik kepada kanak-kanak. **Kesimpulan:** Hasil kajian ini boleh digunakan bagi mengisi jurang pengetahuan tentang kesan pencemaran tanah dan risiko kesihatan yang berpotensi di hadapi oleh populasi hasil daripada pendedahan logam berat di kawasan bandar terutamanya di Malaysia.

Kata Kunci: Tanah bandar, logam berat, total, kebolehdapatan, risiko kesihatan

ACKNOWLEDGEMENT

In the name of Allah, the Most Gracious and the Most Merciful

Alhamdulillah, all praises to Allah for the strengths and His blessing in completing this thesis. Special appreciation goes to my dedicated supervisor, Dr Sarva Mangala Praveena, for her supervision and never ending support. Her meaningful words and invaluable help of constructive comments and suggestions throughout the research and thesis works have contributed a lot to the success of this research. Not forgotten, my appreciation to my co-supervisors, Prof. Dr. Zailina Hashim and Associate Prof Dr. Ahmad Zaharin Aris for their support and knowledge pertaining to this research.

I would like to express my deepest gratitude to my beloved and supportive parents, Mr. Yuswir Bin Yunos and Mrs. Azizah Binti Sabidin and also my siblings for their endless love, prayers and encouragement. My acknowledgement also goes to all the laboratory staffs of Environmental Health Laboratory, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia for their assistance throughout the sample analysis. And not forgotten to all office staffs of Department of Occupational and Environmental Health, Faculty of Medicine and Health Sciences, University Putra Malaysia for their co-operations. I also would like to thank to Kementerian Pengajian Tinggi Malaysia (My Brain 15), Graduate Research Fellowship UPM, Research University Grant Scheme (RUGS) vote number 9335100 and Negeri Sembilan Government for their financial support through this study.

Last but not least, sincere thanks to all my adorable friends especially Raihanah Gojes, Aida, Najihah, Aziemah, Noreen, Madiha, Hafiza and others for their kindness and moral support during my study. Thanks for the friendship and memories.

Thank you

I certify that a Thesis Examination Committee has met on 22 October 2015 to conduct the final examination of Nurul Syazani binti Yuswir on her thesis entitled "Heavy Metals in Soil from Urban Areas of Klang, Malaysia, and their Health Risk Assessment Using *In Vitro* Digestion Method" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Huda binti Zainuddin, PhD, M.CM (OH)

Medical Lecturer
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Samsuri bin Abd. Wahid, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Kamaruzzaman bin Hj. Yunus, PhD

Associate Professor
International Islamic University Malaysia
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 15 December 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Masters of Science .The members of the Supervisory Committee were as follows:

Sarva Mangala Praveena, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Zailina Hashim, PhD

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Ahmad Zaharin Aris, PhD

Associate Professor
Faculty of Environmental Sciences
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations, and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials, as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Nurul Syazani Binti Yuswir GS34965

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of

Chairman of

Supervisory

Committee : Dr Sarva Mangala Praveena

Signature: _____

Name of

Member of

Supervisory

Committee : Professor Zailina Hashim

Signature: _____

Name of

Member of

Supervisory

Committee : Associate Professor Ahmad Zaharin Aris

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENT	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xiv
CHAPTER	
1 INTRODUCTION	1
1.1 Introduction	1
1.2 Problem Statement	4
1.3 Study Justification	5
1.4 Study Contribution	5
1.5 Conceptual Framework	6
1.6 Definition	7
1.6.1 Conceptual Definition	7
1.6.1.1 Heavy Metal	7
1.6.1.2 Bioavailable	7
1.6.1.3 Health Risk Assessment	7
1.6.2 Operational Definition	7
1.6.2.1 Heavy Metal	7
1.6.2.2 Bioavailable	8
1.6.2.3 Total	8
1.6.2.2 Health Risk Assessment	8
1.7 Research Objectives	10
1.7.1 General Objective	10
1.7.2 Specific Objectives	10
1.7.3 Hypothesis	10
2 LITERATURE REVIEW	11
2.1 Soil Profile	11
2.2 Heavy Metal in Urban Soil Worldwide	13
2.3 Heavy Metal Studies in Malaysia Urban Soil	17
2.4 Method of Heavy Metal Analysis in Soil Analysis	19
2.5 Pathway of Heavy Metal Exposure From Soil	21
2.6 <i>In Vitro</i> Digestion Model	22
2.7 Studies With <i>In Vitro</i> Digestion Model Application	27
2.8 Health Risk Assessment	28
2.9 Health Risk Assessment Application Using <i>In Vitro</i> Digestion Model Output	29
2.10 Health Effects of Heavy Metal	31
2.11 Cluster Analysis Application in Soil Studies	32

3	METHODOLOGY	33
	3.1 Study Location	33
	3.2 Study Design	35
	3.3 Data Collection	35
	3.3.1 Soil Sampling	35
	3.4 Instrumentation	38
	3.5 Sample Analysis	38
	3.5.1 Total Heavy Metal Analysis	38
	3.5.2 Bioavailable Heavy Metal Analysis	40
	3.6 Determination of Heavy Metal Concentration using ICP-OES Pelkin Elmer Optima 8300	42
	3.7 Quality Control	42
	3.8 Statistical Analysis and Health Risk on Data Collected	44
	3.8.1 Normality Test For Various Heavy Metal Concentration	44
	3.8.2 Various Statistical Test Used	44
	3.8.3 Health Risk Assessment Calculation	46
4	RESULTS AND DISCUSSION	
	4.1 Concentration of Total Heavy Metal Concentration in Urban Surface Soil	50
	4.2 Concentration of Bioavailable Heavy Metal in Urban Surface Soil	54
	4.3 Comparison Between Total and Bioavailable of Heavy Metal	57
	4.4 Source Determination of Heavy Metal in the Soil	59
	4.4.1 Spearman correlation	59
	4.4.2 Cluster analysis	60
	4.5 Health Risk Assessment of Bioavailable Heavy Metal Concentration	64
	4.5.1 Health Risk Assessment based on element	64
	4.5.2 Health Risk Assessment based on population	66
	4.5.3 Pattern distribution of carcinogenic and non carcinogenic risk in Klang district	68
	4.6 Uncertainty of Health Risk Assessment	70
5	CONCLUSION	72
	REFERENCES	73
	APPENDICES	90
	BIODATA OF STUDENT	100
	LIST OF PUBLICATIONS	101

LIST OF TABLES

Table		Page
1.1	The risk acceptability for non-carcinogenic health effect (U.S. EPA, 2007)	9
1.2	The risk acceptability for carcinogenic health effects (USEPA., 2007)	9
2.1	Heavy metal concentration (mg/kg) in urban soils worldwide	15
2.2	Heavy metal studies of urban soil in Malaysia	17
2.3	Types of in vitro digestion model	23
2.4	Analytical technique for each model	24
2.5	Studies using in vitro digestion model to access heavy metal bioavailable (mg/kg) in soil	27
2.6	Health risk assessment application in heavy metal studies on urban soil	30
3.1	Recovery rate for the standard reference soil	43
3.2	Normality test for each heavy metal	44
3.3	Value for parameter used in Equation 1 -4	46
3.4	Reference Dose value	48
3.5	Cancer Slope Factor value	49
4.1	Descriptive statistic analysis of total heavy metal concentration (mg/kg)	50
4.2	Comparison of mean total heavy metal concentrations (mg/kg) from Klang urban	52
4.3	Comparison of mean total heavy metal concentrations (mg/kg) from Klang urban soil samples with other studies	53
4.4	Descriptive statistic analysis of bioavailable of heavy metal concentration (mg/kg)	54
4.5	Comparison of mean bioavailable heavy metal concentrations (mg/kg) from Klang urban soil samples with other studies	56
4.6	Comparison of total and bioavailable concentration using Kruskal Wallis test (n = 76)	57
4.7	Correlation of Various Heavy Metals In Klang Urban Soil	60

LIST OF FIGURES

Figure		Page
1.1	Percentage of urbanization in Malaysia	1
1.2	Distribution of urban centres in Peninsular Malaysia, 1957 and 2000	2
2.1	Soil Horizon	12
2.2	Method involved in heavy metal analysis in soil	19
2.3	In vitro digestion model process in human body	26
2.4	Steps in HRA	29
3.1	Location of Klang District bounded by Kuala Selangor District, Petaling District and Kuala Langat District	34
3.2	Study area and sampling site locations in Klang District, Selangor	37
3.3	Process of the <i>aqua regia</i> digestion method	39
3.4	Process of in vitro digestion model	41
4.1	Dendrogram derived from hierarchical cluster analysis of heavy metal content in analysed urban soils	62
4.2	Hazard Quotient value of each heavy metal in adult and children	65
4.3	Total Risk value of each heavy metal in adult and children	65
4.4	Hazard Quotient _{sum} value of all heavy metal in adult and children	67
4.5	Total Risk _{sum} value of all heavy metal in adult and children	67
4.6	Non carcinogenic risks of Klang district	68
4.7	Carcinogenic risks of Klang district	70

LIST OF ABBREVIATIONS

μm	Micrometer
Ag	Silver
Al	Aluminium
As	Arsenic
AT	Averaging Time
$A_{t_{\text{cancer}}}$	Averaging Time For Cancer
$A_{t_{\text{non Cancer}}}$	Averaging Time For Non Cancer
ATSDR	Agency for Toxic Substances and Disease Registry
BW	Body Weight
CA	Cluster Analysis
Cd	Cadmium
CDI	Chronic Daily Intake
Co	Cobalt
Cr	Chromium
CSF	Cancer Slope Factors
Cu	Copper
ED	Exposure Duration
EF	Exposure Frequency
HCl	Hydrochloric Acid
Hg	Mercury
HNO_3	Nitric Acid
HQ	Hazard Quotient
HRA	Health Risk Assessment
IARC	International Agency For Research On Cancer
ICP-OES	Inductively Coupled Plasma-Optical Emission Spectrometry
Ingr	Ingestion Rate
IRIS	Integrated Risk Information System
LCR	Lifetime Cancer Risk
NaCl	Sodium Chloride
NaOH	Sodium Hydroxide
Ni	Nickel
Pb	Lead
PBET	Physiologically Based Extraction Test
RAIS	Risk Assessment Information System
Rfd	Reference Dose
SBET	Simplified Bioaccessibility Extraction Test
SBRC	Solubility/Bioavailable Research Consortium
Se	Selenium
SF	Slope Factor
SHIME	Simulator Of Human Intestinal Microbial Ecosystems Of Infants
SPSS	Statistical Package For Social Science
TR	Total Risk
USDA	United States Department Of Agriculture
USDOE	United States Department Of Environment
USEPA	United States Environmental Protection Administration
Zn	Zinc

CHAPTER 1

INTRODUCTION

1.1 Introduction

Malaysia is one of the countries in Southeast Asian that is currently undergoing a rapid urbanization. Urbanization in Malaysia rose dramatically from 26.8 % to 61.8 % between 1970 to 2000 (Jaafar, 2004). In addition, Malaysia is also projected to be an urban society with the majority (over 70 %) of the country's total population living in its cities (Hassan, 2009). Figure 1.1 shows the level of urbanization in Malaysia by Department of Statistics Malaysia from 1980 to 2010 while Figure 1.2 shows the distribution of urban centres in Peninsular Malaysia in 1957 and 2000. Over 60% of the Malaysian population are concentrated in the west part of Peninsular Malaysia. Thus, this area has most of the development activities (Bin, 2011). However, this rapid urbanization has produced environmental pollutions such as water and air pollution that have given impacts on vegetation, river and urban soil (Zarcinas et al., 2004; Henderson and Wang, 2007). According to Shazili et al., (2006), manufacturing along with port and shipping activities are the major contributors of pollution in the environment of west Peninsular Malaysia.




Figure 1.1: Percentage of urbanization in Malaysia
Source: Department of Statistics Malaysia, (2014)




Figure 1.2: Distribution of urban centres in Peninsular Malaysia, 1957 and 2000

Source: Masron et al., (2012)

According to Wei and Yang (2010) and Foo et al., (2008), urban soils have become polluted due to the rapidly urbanization. Rapid urbanization has polluted the urban soil due to the land use activity by human. In urbanization process, raw land is converted and covered with pavement. The impervious surfaces created by buildings and pavement causes rainwater and snowmelt that carries environmental pollutants to flow quickly over the landscape, rather than soaking naturally into the urban soil or being absorbed by plants (Foster, 1990). There are many pollutants that can cause pollution in

urban soil such as dioxin, persistent organic pollutants (POPs) and heavy metal (Habe et al., 2001; Foo et al., 2008; Wang et al., 2012). Among these pollutants in urban soil, heavy metal is one of the major environmental problems throughout the world because heavy metal has significant toxic effects and bioaccumulate in human, animals, microorganisms and plants (Mehdi et al., 2003; Majid et al., 2012). According to Coen et al., (2001) heavy metals can accumulate in human tissue as a result of rapid urbanization pollution and give toxic effect. In addition, bioaccumulation of heavy metal over prolonged time can become hazardous to human health after entering in their body system (Mehdi et al., 2003).

Studies done by researchers have stated that urban soils worldwide have high concentration of heavy metal (Wong et al., 2006; Thornton et al., 2008; Luo et al., 2012). In urban soil, heavy metal can be present in both natural and anthropogenic forms. Natural forms of heavy metal due to weathering of rock minerals are present at relative low concentrations. Mineral weathering of rocks is the primary source of background heavy metal in urban soil (Dakane, 2012). However, in recent years a number of anthropogenic sources have implied notable contributions to the increase of the heavy metal concentrations (Luo et al., 2011). This is supported by Wei and Yang (2010) in China, stating that the pollution sources of heavy metal in urban soil are mainly derived from anthropogenic activities. In urban soils, the anthropogenic sources of heavy metals include traffic emissions, industrial emissions, domestic emission, weathering of building and pavement surface and atmospheric deposition. Traffic emissions include vehicle exhaust particles, tire wear particles, weathered street surface particles and brake lining wear particles meanwhile industrial emissions includes the power plants, coal combustion, metallurgical industry, auto repair shop and chemical plants (Foo et al., 2008). A review done by Afroz et al., (2003) in Malaysia stated that the general major sources of pollution in Malaysia are from motor vehicles, open burning and industrial emissions. Additionally, Foo et al., (2008) in Kuala Terengganu (Terengganu) stated that vehicular emission is the most commonly known significant and increasing source of soil pollution in urban environment.

Elevation of heavy metal in urban soil due to anthropogenic sources can pose significant health risks to human. Heavy metal in urban soil can enter into human body from three different exposure pathways which are by accidental ingestion (Luo et al., 2011; Okorie et al., 2011), inhalation (Laidlaw and Filippelli, 2008; Schmidt, 2010) and dermal contact (Siciliano et al., 2009). Accidental ingestion of heavy metal in urban soil can occur due to mouthing behaviour, contacting with dirty hands or eating dropped food (USEPA, 2011). Moreover, according to Doyle et al., (2012) stated that accidental ingestion may occur through the inadvertent ingestion of soil or dust particles that adhere to food, objects and hands, or the deliberate ingestion of soil such as soil pica and geophagy, which is considered to be relatively uncommon in the general population. However, geophagy can be a relatively common practice in indigenous peoples on all continents (Doyle, 2012). Meanwhile, inhalation of heavy metal in urban soil requires the heavy metal to be present in air and then be inhaled whereas dermal absorption requires contact between heavy metal and skin during contact with urban soil (WHO, 2010). However, accidental ingestion is considered as an important pathway for heavy metal to enter the human body and is the largest area of concern in exposure route pathway in soil followed by dermal and inhalation (Paustenbach, 2000). Moreover, study by Intawongse et al., (2006) also suggest that

accidental ingestion is an important exposure routes to humans from potentially harmful pollutants in soil including heavy metal.

1.2 Problem Statement

In Malaysia, Klang is located in west part of Peninsular Malaysia which is considered as one of the area that undergoes rapid urbanization (Bin, 2011). However, this rapid urbanization in Klang has given impacts on environment such as river and urban soil and produced environmental pollutions such as water and soil pollution (Zarcinas et al., 2004). Beside rapid urbanization, Klang is also an area that is surrounded by land and sea based anthropogenic activities. According to Shazili et al., (2006), anthropogenic activities in West Malaysia including Klang area has become one of the major contributor to environmental pollution in that area including heavy metal. This is supported by Yap et al. (2003), who stated that the major input sources of heavy metal in Klang District can directly contribute to high level of heavy metal concentrations in Klang waters, sediment, soil and biota. Other than that, studies by Ismail et al., (1993) and Yap et al., (2003) showed high concentrations of Cd, Pb, Cu and Zn detected in Port Klang sediments. In addition, studies done in Klang area by Lihan and Rahim, (2006) and Dayang et al., (2013) have also stated that the total heavy metal concentrations in urban soils sampled from Klang were highest for Cu, Zn, Ni, As, Cr and Cd concentrations.

So far, studies only employed total heavy metal concentration and explained about the pollution status, spatial behavior and speciation of heavy metal in Klang areas (Yap et al., 2003; Lihan and Rahim, 2006; Sany et al., 2012; Naji and Ismail., 2012; Dayang et al., 2013). However, these studies only measure the total heavy metal concentration in their soil sample and focusing on heavy metal pollution status of the soil. However, there are no any studies related to bioavailable of heavy metal in Klang urban soil which can be used to indicate human health.

1.3 Study Justification


Rapid urbanization and anthropogenic activities in Klang area have polluted the urban soil. Studies done by Ismail et al., (1993), Yap et al., (2003), Lihan and Rahim, (2006) and Dayang et al., (2013) found a high concentration of heavy metal in Klang waters, sediment, soil and biota. From time to time, there are an increment in rapid urbanization and anthropogenic activities in Klang area. This situation will increase the concentration of heavy metal in Klang urban surface soil. Thus, there is a need for further research that evaluates the recent concentrations of heavy metal in Klang area and determines the possible sources of heavy metal pollution in Klang urban surface soil.

So far, heavy metal studies on urban soil in Klang have only focused on pollution status, spatial behaviour and speciation of heavy metal using total heavy metal concentration. Thus, there was a need for further research on Klang urban soil that evaluates impacts of heavy metal on human health by assessment of bioavailable form and incorporate health risk assessment. Bioavailable of heavy metal measures heavy metal concentration that is absorbed in gastrointestinal tract of human (Oomen et al., 2003). Bioavailable of heavy metal concentration would give accurate results on human health impacts via health risk assessment compared to total heavy metal.

1.4 Study Contribution

This study output would fill the gap about Klang area which includes information on the impact of heavy metal toward human health. Furthermore, findings from this study would also provide a baseline value for heavy metal contamination in urban soil and its impacts to human health that can be used in the recommendation for the management of heavy metals released to the environment. Findings from this study will also help in the policy making on development of housing area by providing a location for the type of land that are safe to be placed.

1.5 Conceptual Framework



1.6 Definition Of Term

1.6.1 Conceptual definition

Heavy metal

Any metallic element that has a relatively high density and is toxic or poisonous even at low concentration (Duruibe et al., 2007). Of the 92 naturally occurring elements, approximately 30 heavy metal are potentially toxic to humans such as Aluminium (Al), (Chromium) Cr, Cobalt (Co), Nickel (Ni), Copper (Cu), Arsenic (As), Selenium (Se), Argentum (Ag), Cadmium (Cd), Mercury (Hg) and Lead (Pb) (Morais et al., 2012).

Bioavailable

The fraction of an orally administered dose that reaches the systemic circulation (Versantvoort et al., 2004). There are 5 analytical methods used to determine the bioavailable of heavy metal which are Physiologically Bioavailable Extraction Test (PBET) method (British Geological Survey, UK), E DIN method (Ruhr Universitat Bochum, Germany), RIVM method (National Institute of Public Health and the Environment, The Netherlands), Simulator of Human Intestinal Microbial Ecosystems of Infants (SHIME) method (LabMET, Belgium) and TIM method (TNO Nutrition, The Netherlands) (Oomen et al., 2002).

Health Risk Assessment

Health risk assessment is a scientific process to estimate the probability of adverse health effects in humans who may be exposed to pollutants in contaminated environmental media. This process utilizes the tools of science and statistics to identify and measure a hazard, determine possible routes of exposure, and finally use that information to calculate a numerical value to represent the potential risk. A human health risk assessment consists of four steps. These steps include hazard identification, dose-response assessment, exposure assessment, and risk characterization (USEPA, 2002).

1.6.2 Operational definition

Heavy metal

Heavy metal (Zn, Cu, Fe, Cd, Co, Al, Pb, Cr) concentration in urban soil that are selected in this study was digested using *aqua regia* and Physiologically Based

Extraction Test method and then the concentration was determined and analyzed using Inductively Coupled Plasma-Optical Emission Spectrometry (ICP-OES).

Bioavailable

Bioavailable of heavy metal in this study was measured using Physiologically Based Extraction Test (PBET) method. There are 5 analytical methods used in an *in vitro* digestion model to determine the bioavailable of heavy metal which are Physiologically Bioavailable Extraction Test (PBET) method (British Geological Survey, UK), E DIN method (Ruhr Universitat Bochum, Germany), RIVM method (National Institute of Public Health and the Environment, The Netherlands), Simulator of Human Intestinal Microbial Ecosystems of Infants (SHIME) method (LabMET, Belgium) and TIM method (TNO Nutrition, The Netherlands) (Oomen et al., 2002). Among 5 of the analytical methods used in an *in vitro* digestion model, PBET is one of the most widely used models.

Total

Total heavy metal in this study was measured using *aqua regia* method which is this method used the combination of hydrochloric acid and nitric acid.

Health Risk Assessment

In this study, health risk assessment was defined and done specifically on heavy metal exposure through the bioavailable of ingested urban soil. The assessment was evaluated through Hazard Quotient (HQ) for non-carcinogenic risks and Total Risk (TR) for carcinogenic risks to estimate the health risks of adult and children who are exposed to heavy metal through ingestion of urban soil.

Non carcinogenic risks were quantified by calculation of Hazard Quotient (HQ) as below: The oral route was the route of exposure observed in this study, therefore, oral RfD values from the IRIS database was used.

$$\text{Hazard Quotient (HQ)} = \frac{\text{CDI}}{\text{RfD}}$$

Where:

HQ = Non cancer Hazard Index of a health effect from intake of heavy metal

CDI = Chronic daily intake (mg/kg/day)

Oral RfD = Oral Reference dose of heavy metal (mg/kg/day)

Then, the hazard quotient value was compared with the following values of risk acceptability for non-carcinogenic health effects. In cases where the non-cancer HQ does not exceed unity ($HI < 1$), it is assumed that no chronic risks are likely to occur at the site.

Table 1.1: The risk acceptability for non-carcinogenic health effect (U.S. EPA, 2002)

Hazard Quotient (HQ)	
>1	Unacceptable
<1	Acceptable

As expressed earlier, carcinogens were assumed to not have an effective or safe threshold. Carcinogenic risk is expressed as slope factor (SF) and the following equation was used to quantify lifetime risk of cancer:

$$\text{Total Risk (TR)} = \text{CDI} \times \text{SF}$$

Where:

CDI = Chronic daily intake (mg/kg/day)

SF = Slope factor (mg/kg/day)

Cancer slope factor values can be found on EPA's (2009) Integrated Risk Information System (IRIS). EPA guidelines specified that an acceptable risk is a lifetime cancer risk of no greater than 1 in 1,000,000. Thus, lifetime cancer risk (LCR) value would be referred to the following table to access the risk acceptability of carcinogenic health effect.

Table 1.2: The risk acceptability for carcinogenic health effects (USEPA, 2002)

Total Risk (TR)	
$<10^{-6}$	Clearly acceptable
10^{-6} to 10^{-4}	Acceptable
$> 10^{-4}$	Clearly unacceptable

1.7 Objectives

1.7.1 General Objective

To determine health risk assessment for bioavailable of heavy metal concentration (Al, Fe, Zn, Cu, Co, Cd, Pb and Cr) in urban soil of Klang District.

1.7.2 Specific Objectives

- To determine and compare the total and bioavailable of heavy metal concentration (Al, Fe, Zn, Cu, Co, Cd, Pb and Cr) in urban soil.
- To determine the correlations among bioavailable heavy metal concentrations.
- To identify possible sources of bioavailable heavy metal concentration in urban soil.
- To determine health risks through accidental ingestion exposure pathway of heavy metal exposure in urban soil based on element and population.

1.7.3 Hypotheses

- There was a significant difference between total and bioavailable of heavy metal concentration in urban soil.
- There were significant correlations between bioavailable heavy metal concentrations.

REFERENCES

- Abdi, H., & Williams, L. J. (2010). Principal component analysis. *Wiley Interdisciplinary Reviews: Computational Statistics*, 2: 433-459.
- Abdullah, A., R., Tahir, N. M., Tong, S. L., Hoque, T. M., & Sulaiman, A. H. (1999). The GEF/UNDP/IMO Malacca Straits Demonstration Project: Sources of pollution. *Marine Pollution Bulletin*, 39: 229-233.
- Afroz, R., Hassan, M. N., & Ibrahim, N. A. (2003). Review of air pollution and health impacts in Malaysia. *Environmental research*, 92: 71-77.
- Ahmed, F., & Ishiga, H. (2006). Trace metal concentrations in street dusts of Dhaka city, Bangladesh. *Atmospheric Environment*, 40: 3835-3844.
- Ajorlo, M., Abdullah, R. B., Hanif, A. H. M., Halim, R. A., & Yusoff, M. K. (2010). How cattle grazing influences heavy metal concentrations in tropical pasture soils. *Polish Journal of Environmental Studies*, 19: 895-902.
- Alexander, G. (2000). Health risk appraisal. *International Electronic Journal Health Education*, 3: 133-137.
- Al-Khashman, O. A. (2007). Determination of metal accumulation in deposited street dusts in Amman, Jordan. *Environmental Geochemical Health*, 29:1-10.
- Alloway, B.J. (1995). *Heavy metals in Soil*. Blackie Academic & Professional, Glasgow, pp 206-210.
- Amato, F., Pandolfi, M., Viana, M., Querol, X., Alastuey, A. & Moreno, T. (2009). Spatial and chemical patterns of PM10 in road dust deposited in urban environment. *Atmospheric Environment*, 43: 1650-1659.
- Antoniadis, V. & Alloway, B. J. (2003). Evidence of heavy metal movement down the profile of heavily sludged soil. *Communication in Soil Sciences and Plant Analysis*, 34: 1225-1231, 2003.
- ATSDR. (2002). Public Health Statements – Toxicological Profiles (Al, As, Cd, Co, Cr, Cu, Hg, Mn, Ni, Pb, Zn). <http://www.atsdr.cdc.gov/phshome.html>. [12 August 2013].
- Baco, S., Chik, A., Musta, B., Yassin, F. M., & Halim, S. A. (2010). Study On Magnetic Properties, Mineralogy And Heavy Metals Content Of Soil And Concretion. *Borneo Science*, 27: 23-34.
- Baize, D., & Sterckeman, T. (2001). Of the necessity of knowledge of the natural pedo-geochemical background content in the evaluation of the contamination of soils by trace elements. *Sciences of Total Environment*, 264: 127-139.

- Baralkiewicz, D., & Siepak, J. (1999). Chromium, nickel and cobalt in environmental samples and existing legal norms. *Polish Journal of Environmental Studies*, 8: 201-208.
- Bermejo, P., Pena, E. M., Dominguez, R., Bermejo, A., Cocho, J., Fraga, J. M. (2002). Iron and zinc in hydrolised fractions of human milk and infant formulas using an in vitro method. *Food Chemistry*, 77: 361-369.
- Białk-Bielińska, A., Kumirska, J., Palavinskas, R., & Stepnowski, P. (2009). Optimization of multiple reaction monitoring mode for the trace analysis of veterinary sulfonamides by LC–MS/MS. *Talanta*, 80: 947-953.
- Biasioli, M., Barberi, R., Ajmone-Marsan, F. (2005). The influence of a large city on some soil properties and metals content. *Sciences of Total Environment*, 356: 154–164.
- Bin, T. J. (2011). *UTAR Seri Kembangan new village community project report*. Undergraduate thesis, Universiti Tunku Abdul Rahman.
- Birke, M. & Rauch, U. (2000). Urban geochemistry: Investigations in the Berlin metropolitan area. *Environmental Geochemistry and Health*, 22: 233-248.
- Brady, N. C., & Weil, R. R. (1996). *The nature and properties of soil*. Prentice- Hall, International, Inc. London.
- Brandon, E. F., Oomen, A. G., Rompelberg, C. J., Versantvoort, C. H., Van Engelen, J. G. & Sips, A. J. (2006). Consumer product in vitro digestion model: Bioavailable of contaminants and its application in risk assessment. *Regulatory Toxicology and Pharmacology*, 44:161-171.
- Bremmer, J. (2015). *Update and Upgrade the current quality assurance in the global supply chain of a heavy truck manufacturing company: managing product quality in the North Bound Flow (NBF) at Scania Production Zwolle*. University of Twente, Netherlands.
- Brinkmann, R. (1994). Lead pollution in soils adjacent to homes in Tampa, Florida. *Environmental Geochemistry and Health*, 16: 59-64.
- Broadway, A., Cave, M. R., Wragg, J., Fordyce, F. M., Bewley, R. J. F., Graham, M. C, et al. (2010). Determination of the bioaccessibility of chromium in Glasgow soil and the implications for human health risk assessment. *Science of the Total Environment*, 409: 267-277.
- Brown, G. E., Foster, A. L. & Ostergren, J. D. (1999). Mineral surfaces and bioavailable of heavy metals: a molecular-scale perspective. *Proceedings of the National Academy of Sciences, USA*, 96: 3388.
- Cabanero, A. I., Madrid, Y. & Camara, C. (2004). Selenium and mercury bioavailable in fish sample: an in vitro digestion method. *Analytica Chimica Acta*, 526: 51-61.

- Cai, L., Xu, Z., Ren, M., Guo, Q., Hu, X., Hu, G. & Peng, P. (2012). Source identification of eight hazardous heavy metals in agricultural soils of Huizhou, Guangdong Province, China. *Ecotoxicology Environmental Safety*, 78: 2-8.
- Carlou, C., D'Alessandro, M., & Swartjes, F. (2007). *Derivation methods of soil screening values in Europe. A review and evaluation of national procedures towards harmonization*. JRC Scientific and Technical Report EUR, 22805: 1-320.
- Centeno, J. A. (2006). Global impacts of geogenic arsenic – a medical geology research case. *International Symposium on Medical Geology*, 18-19 May 2006, Stockholm, Sweden.
- Centeno, J. A., Mullick, F. G., Ishak, K. G., Franks, T. J., Burke, A. P., Koss, M. N., Perl, D. P., Tchounwou, P. B., & Pestaner, J. P. (2006). *Environmental pathology*. In: Selinus, O., Alloway, B., Centeno, J.A., Finkelman, R.B., Fuge, R., Lindh, U. & Smedley, P. (eds) *Essentials of Medical Geology*. Elsevier Academic Press. Amsterdam, the Netherlands. p 563-594.
- Chabukdhara, M., & Nema, A. K. (2013). Heavy metals assessment in urban soil around industrial clusters in Ghaziabad, India: probabilistic health risk approach. *Ecotoxicology and environmental safety*, 87: 57-64.
- Chan, Y. H. 2003. Basic statistics for Doctors. *Singapore Medical Journal*, 44: 614-619.
- Chen, T. B., Zheng, Y. M., Lei, M., Huang, Z. C., Wu, H. T., Chen, H., & Tian, Q. Z. (2005). Assessment of heavy metal pollution in surface soils of urban parks in Beijing, China. *Chemosphere*, 60: 542-551.
- Chon, H. T., Kim, K. W., & Kim, J. Y. (1995). Metal contamination of soils and dusts in Seoul metropolitan city, Korea. *Environmental Geochemistry and Health*, 17: 139-146.
- Coen, N., Mothersill, C., Kadhim, M., & Wright, E. G. (2001). Heavy metals of relevance to human health induce genomic instability. *The Journal of pathology*, 195: 293-299.
- Crews, H. M., Burrell, J. A., & McWeeny, D. J. (1985). Trace element solubility from food following enzymolysis. *Z. Lebensm Unters Forsch*, 180: 221-226.
- Crews, H. M., Burrell, J. A., & McWeeny, D. J. (1983). Preliminary enzymolysis studies on trace element extractability from food. *Journal of Sciences Food Agriculture*, 34: 997-1004.
- Dakane, A. (2012). *Bioaccessibility of metals in Toronto city parks*. Doctoral dissertation, Royal Roads University.

- Davies, B. E., Bowman, C., Davies, T. C., & Selinus, O. (2005). *Medical geology: Perspectives and prospects*. In: Selinus, O., Alloway, B., Centeno, J.A., Finkelman, R.B., Fuge, R., Lindh, U. & Smedley, P. (eds) *Essentials of Medical Geology*. Elsevier Academic Press. Amsterdam, the Netherlands. p 563-594.
- Dayang, S. N., & Che Fauziah, I. (2013). Soil factors influencing heavy metal concentrations in medicinal plants. *Pertanika Journal of Tropical Agriculture Sciences*, 36: 161-177.
- De Miguel, E., De Grado, M. J., Llamas, J. F., Martín-Dorado, A., & Mazadiego, L. F. (1998). The overlooked contribution of compost application to the trace element load in the urban soil of Madrid (Spain). *Science of the Total Environment*, 215: 113-122.
- De Moura, M. C. S., Moita, G. C., & Neto, J. M. M. (2010). Analysis and assessment of heavy metals in urban surface soils of Teresina, Piauí State, Brazil: a study based on multivariate analysis. *Comunicata Scientiae*, 1: 120.
- Debnárová, A., & Weissmannová, H. (2010). Assessment of Heavy Metal Pollution (Cd, Cu, Pb, Hg) in Urban Soils of Roadsides in Brno. *Transport Sciences*, 3: 147-156.
- Department of Statistics Malaysia, (2014). *Distribution of urbanization in Malaysia*. Malaysia.
- Doyle, J. R., Blais, J. M., Holmes, R. D., & White, P. A. (2012). A soil ingestion pilot study of a population following a traditional lifestyle typical of rural or wilderness areas. *Science of the Total Environment*, 424: 110-120.
- Duruibe, J. O., Ogwuegbu, M. O. C., & Egwurugwu, J. N. (2007). Heavy metal pollution and human biotoxic effects. *International Journal Physical Sciences*, 2: 112-118.
- Ehi-Eromosele, C. O., Adaramodu, A. A., Anake, W. U., Ajanaku, C. O., & Edobor-Osoh, A. (2012). Comparison of Three Methods of Digestion for Trace Metal Analysis in Surface Dust Collected from an Ewaste Recycling Site. *Nature and Science*, 10: 1-6.
- Elless, M. P., Bray, C. A., Blaylock, & M. J. (2007). Chemical behavior of residential lead in urban yards in the United States. *Environmental Pollution*, 148: 291-300.
- Ellickson, K. M., Meeker, R. J., Gallo, M. A., Buckley, B. T., & Liroy, P. J. (2001). Oral bioavailable of lead and arsenic from a NIST standard reference soil material. *Environmental Contamination Toxicology*, 40: 128-135.
- Escribano, S. T., Ruiz, A., Barrios, L., Veleza, D., & Montoroa, R. (2010). Influence of mercury bioavailable on exposure assessment associated with consumption of cooked predatory fish in Spain. *Journal of Sciences Food Agriculture*, 91: 981-986.

- Esu, I. E. (2010). *Soil characterization, classification and survey*. HEBN Publishers.
- Facchinelli, A., Sacchi, E., & Mallen, L. (2001). Multivariate statistical and GIS-based approach to identify heavy metal sources in soils. *Environmental Pollution*, 114: 313-324.
- Foo, T. F., Poh, S. C., Asrul, A. M. & Mohd Tahir, N. (2008). Possible source and pattern distribution of heavy metals content in urban soil at Kuala Terengganu town center. *The Malaysian Journal of Analytical Sciences*, 12: 458 – 467.
- Foster, S. S. (1990). Impacts of urbanization on groundwater. Hydrological processes and water management in urban areas. Wallingford, UK, *International Association of Hydrological Sciences–Association Internationale des Sciences Hydrologiques* (IAHS–AISH Pub. No. 198).
- Garba, S. T., Ahmed, I., Akan, J. C., & Dauda, B. A. (2014). Distribution Pattern of the Heavy Metals: Pb, Zn, Cd and Cu in Roadside Soils of Maiduguri Metropolis, Borno State Nigeria. *International Research Journal of Pure and Applied Chemistry*, 4: 486.
- Garret, D. A., Failla, M. L., & Sarama, R. J. (1999). Development of an in vitro digestion method to assess carotenoid bioavailable from meals. *Journal Agriculture Food Chemistry*, 47: 4301-4309.
- Gaw, S., Kim, N., Northcott, G., Wilkins, A., & Robinson, G. (2008). *Developing Site-Specific Guidelines for Orchard Soils Based on Bioaccessibility – Can It Be Done?*. Proceedings, Waste Management Institute of New Zealand Annual Conference.
- Glahn, R. P., Wien, E. M., van Campen, D. R., & Miller, D. D. (1996). Caco-2 cell iron uptake from meat and casein digests parallels in vivo studies: use of a novel in vitro method for rapid estimation of iron bioavailable. *Journal of Nutritional*, 126: 332-339.
- Greim, H., & Snyder, R. (Eds.). (2008). *Toxicology and risk assessment: a comprehensive introduction*. John Wiley & Sons.
- Grzetic, I., & Ghariani, R. H. (2008). Potential health risk assessment for soil heavy metal contamination in the central zone of Belgrade (Serbia). *Journal of Serbia Chemistry Society*, 73: 923–934.
- Guerra, A., Etienne-Mesmin, L., Livrelli, V., Denis, S., Blanquet-Diot, S., & Alric, M. (2012). Relevance and challenges in modeling human gastric and small intestinal digestion. *Trends Biotechnology*, 30: 591-600.
- Guerra, F., Trevizam, A. R., Muraoka, T., Marcante, N. C., & Canniatti-Brazaca, S. G. (2012). Heavy metals in vegetables and potential risk for human health. *Sciences Agriculture*, 69: 54-60.

- Gunawardena, J., Egodawatta, P., Ayoko, G. A., & Goonetilleke, A. (2013). Atmospheric deposition as a source of heavy metals in urban stormwater. *Atmospheric Environment*, 68: 235-242.
- Guney, M., & Zagury, G. J. (2012). Heavy metals in toys and low-cost jewelry critical review of U.S. and Canadian legislations and recommendations for testing. *Environmental Sciences Technology*, 46: 4265-4274.
- Habe, H., Chung, J. S., Lee, J. H., Kasuga, K., Yoshida, T., Nojiri, H., & Omori, T. (2001). Degradation of chlorinated dibenzofurans and dibenzo-p-dioxins by two types of bacteria having angular dioxygenases with different features. *Applied and Environmental Microbiology*, 67: 3610-3617.
- Hafiza, Z. (2001). *Kandungan logam surih dalam tanah di sekitar Bandar Kuala Terengganu*. Bsc Environmental Analytical Chemistry, FST, KUSTEM.
- Haiyan, W., & Stuanes, A. (2003). Heavy metal pollution in air-water-soil-plant system of Zhuzhou city, Hunan province, china. *Water, Air, and Soil Pollution*, 147: 79-107.
- Hassan N. (2009). *Issues and challenges of sustainable urban development in Malaysia*. In: Nurhaslina Hassan (ed) Sustainable urban development issues in Malaysia, pp 1 – 22. Dewana Sdn. Bhd., Selangor.
- Henderson, J. V., & Wang, H. G. (2007). Urbanization and city growth: The role of institutions. *Regional Science and Urban Economics*, 37: 283-313.
- Hseu, Z.Y., Chen, Z.S., Tsai, C.C., Tsui, C.C., Cheng, S.F., Liu, C.L., Lin, H.T. (2002). Digestion methods for total heavy metals in sediments and soils. *Water Air and Soil Pollution*, 141: 189-205.
- Hu, Y., Liu, X., Bai, J., Shih, K., Zeng, E. Y., & Cheng, H. (2013). Assessing heavy metal pollution in the surface soils of a region that had undergone three decades of intense industrialization and urbanization. *Environmental Science and Pollution Research*, 20: 6150-6159.
- Hur, S. J., Lim, B. O., Decker, E. A., & McClements, D. J. (2011). In vitro human digestion models for food applications. *Food Chemistry*, 125: 1-12.
- Imperato, M., Adamo, P., Naimo, D., Arienzo, M., Stanzione, D. & Violante, P. (2003). Spatial distribution of heavy metals in urban soils of Naples city (Italy). *Environmental Pollution*, 124: 247-256.
- Intawongse, M., & Dean, J. R. (2006). In-vitro testing for assessing oral bioavailable of trace metals in soil and food samples. *Trends in Analytical Chemistry*, 25: 876-886.
- Ismail, A., Badri, M. A., & Noor Ramlan, M. (1993). The background levels of heavy metal concentration in sediments of the west coast of Peninsular Malaysia. *Sciences of Total Environment*, 134: 315-323.

- Jaafar, J. (2004). Emerging trends of urbanisation in Malaysia. *Journal of the Department of Statistics, Malaysia*, 1: 43-60.
- Jacobs, A., & Greenman, D. A. (1969). Availability of Food Iron. *British Medical Journal*, 1(5645): 673–676.
- Jarup, L. (2003). Hazards of heavy metal contamination. *British Medical Bulletin*, 68: 167-182.
- Juhasz, A. L., Smith, E., Weber, J., Rees, M., Rofe, A., Kuchel, T., & Sansom, L. (2007). In vitro assessment of arsenic bioavailable in contaminated (anthropogenic and geogenic) soils. *Chemosphere*, 69: 69-78.
- Kachenko, A. G. & Singh, B. (2006). Heavy metals contamination in vegetables grown in urban and metal smelter contaminated sites in Australia. *Water, Air & Soil Pollution*, 169: 101–123.
- Kalbitz, K., & Wennrich, R. (1998). Mobilization of heavy metals and arsenic in polluted wetland soils and its dependence on dissolved organic matter. *Sciences of Total Environment*, 209: 27-39.
- Kamaruzzaman, B. Y., Ong, M. C., Zaleha, K., & Shahbudin, S. (2008). Levels of heavy metals in green-lipped mussel *Perna veridis* (Linnaeus) from Muar Estuary, Johore, Malaysia. *Pakistan Journal of Biology Sciences*, 11: 2249-2253.
- Karim, Z. & Qureshi, B. A. (2013). Health risk assessment of heavy metals in urban soil of Karachi, Pakistan. *Human Ecological Risk Assessment*, 20: 658-667.
- Kelly, J., Thornton, I., & Simpson, P. R. (1996). Urban geochemistry: A study of the influence of anthropogenic activity on the heavy metal content of soils in traditionally industrial and non-industrial areas of Britain. *Applied Geochemistry*, 11: 363–370.
- Khairiah, J., Habibah, H. J., Anizan, I., Maimon, A., Aminah, A. & Ismail, B. S. (2009). Content of heavy metals in soil collected from selected paddy cultivation areas in Kedah and Perlis, Malaysia. *Journal of Applied Sciences Research*, 5: 2179-2188.
- Klaassen, C. D. (1996). Heavy metals and heavy-metal antagonists. *The pharmacological basis of therapeutics*, 12: 1851-75.
- Klang Municipal Council. 2008. *Draf Rancangan Tempatan Majlis Perbandaran Klang* (Pengubahan 1) 2020.
- Kouji, A., & Yoshiaki, T. (2004). Characterization of heavy metal particles in tire dust. *Journal of Environmental International*, 8: 1009-1017.
- Laidlaw, M. A., & Filippelli, G. M. (2008). Resuspension of urban soils as a persistent source of lead poisoning in children: a review and new directions. *Applied Geochemistry*, 23: 2021-2039.

- Lam, K. L. (2003). *Kajian taburan logam surih dalam tanah di sekitar Daerah Seberang Perai Tengah, Pulau Pinang*. Bsc Environmental Analytical Chemistry, FST, KUSTEM.
- Lee, G. F. & Jones-Lee, A. (1996). Evaluation of the Water Quality Significance of Eroded Suspended Sediment-Associated Constituents. *Land and Water*, 40: 19-23.
- Lee, S. W., Lee, B. T., Kim, J. Y., Kim, K. W. & Lee, J. S. (2006). Human risk assessment for heavy metals and As concentration in the abandoned metal mine areas, Korea. *Environmental Monitoring Assessment*, 119: 233–244.
- Li, X., Liu, L., Wang, Y., Luo, G., Chen, X., Yang, X. & He, X. (2013). Heavy metal contamination of urban soil in an old industrial city (Shenyang) in Northeast China. *Geoderma*, 192: 50-58.
- Li, X., Poon, C. & Liu, P. S. (2001). Heavy metal contamination of urban soil and street dusts in Hong Kong. *Applied Geochemical*, 16: 1361-1368.
- Li, X. D., Lee, S. L., Wong, S. C., Shi, W. Z. & Thornton, I. (2004). The study of metal contamination in urban soils of Hong Kong using a GIS-based approach. *Environmental Pollution*, 129: 113–124.
- Lihan, T., Rahim, S. A., & Kadir, A. A. (2006). Physico-chemical Characteristics of Soil at Western Part of Pulau Indah, Klang, Selangor. *Sains Malaysiana*, 35: 23–30.
- Linde, M., Bengtsson, H., & Öborn, I. (2001). Concentrations and pools of heavy metals in urban soils in Stockholm, Sweden. *Water Air & Soil Pollution*, 1: 83-101.
- Ljung, K. (2006). *Metals in Urban Playground Soils (Distribution and Bioavailable)*. Doctoral thesis, Swedish University of Agricultural Sciences.
- Lock, S. & Bender, A. E. (1980). Measurement of chemically-available iron in foods by incubation with human gastric juice in vitro. *British Journal of Nutrition*, 43: 413-420.
- Lopez, F. F., Cabrera, C., Lorenzo, M. L., & Lopez, M. C. (2002). Aluminium levels in convenience and fast foods: in vitro study of the absorbable fraction. *Sciences of Total Environment*, 300: 69-79.
- Luo, X. S., Ding, J., Xu, B., Wang, Y. J., Li, H. B. & Yu, S. (2012). Incorporating bioavailable into human health risk assessments of heavy metals in urban park soils. *Sciences of Total Environment*, 424: 88–96.
- Ma, A., Wang, J., & Zhang, K. (2012). *Sampling Survey of Heavy Metal in Soil Using SSSI*. In *Advances in Spatial Data Handling and GIS* (pp. 15-26). Springer Berlin Heidelberg.

- Ma, L., & Rao, G. N. (1997). Chemical fraction of cadmium, copper, nickel and zinc in contaminated soil. *Journal of Environmental Quality*, 26: 259-264.
- Madrid, L., Díaz-Barrientos, E., & Madrid, F. (2002). Distribution of heavy metal contents of urban soils in parks of Seville. *Chemosphere*, 49: 1301-1308.
- Majid, N. M., Islam, M. M. & Riasmi, Y. (2012). Heavy metal uptake and translocation by *Jatropha curcas* L. in sawdust sludge contaminated soils. *Australian Journal of Crop Sciences*, 6: 891-898.
- Mallory, A. L. (2010). *A risk Assessment for Ingestion of Toxic Chemical in Fish from Imperial Beach, California*. San Diego State University.
- Man, Y. B., Sun, X. L., Zhao, Y. G., Lopez, B. N., Chung, S. S., Wu, S. C. & Wong, M. H. (2010). Health risk assessment of abandoned agricultural soils based on heavy metal contents in Hong Kong, the world's most populated city. *Environment International*, 36: 570-576.
- Manta, D. S., Angelone, M., Bellanca, A., Neri, R., & Sprovieri, M. (2002). Heavy metals in urban soils: a case study from the city of Palermo (Sicily), Italy. *Science of the Total Environment*, 300: 229-243.
- Manz, M., Weissflog, L., Kühne, R., & Schüürmann, G. (1999). Ecotoxicological hazard and risk assessment of heavy metal contents in agricultural soils of central Germany. *Ecotoxicology and environmental safety*, 42: 191-201.
- Marchand, C., Lallier-Verges, E., Baltzer, F., Albéric, P., Cossa, D., & Baillif, P. (2006). Heavy metals distribution in mangrove sediments along the mobile coastline of French Guiana. *Marine chemistry*, 98: 1-17.
- Marchand, C., Fernandez, J. M., Moreton, B., Landi, L., Lallier-Vergès, E., & Baltzer, F. (2012). The partitioning of transitional metals (Fe, Mn, Ni, Cr) in mangrove sediments downstream of a ferralitized ultramafic watershed (New Caledonia). *Chemical Geology*, 300: 70-80.
- Martin, S., & Griswold, W. (2009). Human health effects of heavy metals. *Environmental Science and Technology Briefs for Citizens*, 15: 1-6.
- Masron, T., Yaakob, U., Ayob, N. M., & Mokhtar, A. S. (2012). Population and spatial distribution of urbanisation in Peninsular Malaysia 1957-2000. *Geografia: Malaysian Journal of Society and Space*, 8: 20-29.
- Mehdi, S. M., Abbas, G., Sarfraz, M. & Hassan, G. (2003). Effect of industrial effluents on mineral nutrition of rice and soil health. *Pakistan Journal of Applied Sciences*, 3: 462-473.
- Mendoza, C. A., Cortes, G., & Munoz, D. (1996). Heavy metal pollution in soils and sediments of rural developing district 063, Mexico. *Environmental Toxicology and Water Quality*, 11: 327-333.

- Mielke, H.W., Wang, G., Gonzales, C.R., Powell, E.T., Le. B. & Quach, N. (2004). PAHs and metals in soils of inner-city and suburban New Orleans, Louisiana, USA. *Environmental Toxicology and Pharmacology*, 18: 243-247.
- Miller, D. D., Schrickler, B. R., Rasmussen, R. R., & Campen, D. V. (1981). An in vitro method for estimation of iron availability from meals. *American Journal Clinical Nutritional*, 34: 2248-2256.
- Minekus, M., Marteau, P., Havenaar, R., Huis in 't Veld, J. H. J. (1995). A multi compartmental dynamic computer controlled model simulating the stomach and small intestine. *Atlanta*, 23: 197-209.
- Mitra, S., & Kebbekus, B. B. (1997). *Environmental Chemical Analysis*. CRC Press.
- Mmolawa, K. B., Likuku, A. S., & Gaboutloeloe, G. K. (2011). Assessment of heavy metal pollution in soils along major roadside areas in Botswana. *African Journal of Environmental Science and Technology*, 5: 186-196.
- Mohamad, J. (2005). Urban transport and growth management strategies: A tale of two Southeast Asian Cities at the dawn of the new millennium. *Malaysian Journal Society Space*, 1: 11-22.
- Mohd Din, M.A., Yusof, S. & Wan Jaafar, W.Z. (2008). *An integration of the GIS and multi criteria decision making method for landfill siting*. MSc Thesis, University Malaya.
- Möller, A., Muller, H. W., Abdullah, A., Abdelgawad, G., & Utermann, J. (2005). Urban soil pollution in Damascus, Syria: Concentrations and patterns of heavy metals in the soils of the Damascus Ghouta. *Geoderma*, 124: 63-71.
- Montagne, D., Cornu, S., Bourennane, H., Baize, D., Ratié, C. & King, D. (2007). Effect of agricultural practices on trace-element distribution in soil. *Communications in Soil Science and Plant Analysis*, 38: 473-491.
- Mounicou, S., Szpunar, J., Andrey, D., Blake, C., Lobinski, R. (2002). Development of a sequential enzymolysis approach for the evaluation of the bioavailable of Cd and Pb from cocoa. *Analyst (Cambridge, UK)*, 127: 1638-1641.
- Morais, S., e Costa, F. G., & de Lourdes Pereira, M. (2012). *Heavy metals and human health*. INTECH Open Access Publisher.
- Naji, A. & Ismail, A. (2010). Chemical speciation and contamination assessment of Zn and Cd by sequential extraction in surface sediment of Klang River, Malaysia. *Microchemical Journal*, 95: 285-292.
- Naji, A. & Ismail, A. (2011). Risk assessment of mercury contamination in surface sediment of the Klang River, Malaysia. *Australian Journal of Basic and Applied Sciences*, 5: 215-221.

- Naji, A. A & Ismail, A. (2012). Metal fractionation and evaluation of their risk connected with urban and industrial influx in the Klang River surface sediments, Malaysia. *Environmental Asia*, 5: 17-25.
- Najib, N. W. A.Z., Mohammed, S. A., Ismail, S. A., Amiza, W., Ahmad, W. A. (2012). Assessment of Heavy Metals in Soil due to Human Activities in Kangar, Perlis, Malaysia. *International Journal of Civil Environmental Engineering*, 12: 28-33.
- Nordberg, M. & Cherian, M.G. (2005). *Biological responses of elements*. In: Selinus, O., Alloway, B., Centeno, J.A., Finkelman, R.B., Fuge, R., Lindh, U. & Smedley, P. 2005 (eds). *Essentials of Medical Geology*. Elsevier Academic Press. Amsterdam, the Netherlands. 179-200.
- Okorie, A., Entwistle, J., & Dean, J. R. (2011). The application of in vitro gastrointestinal extraction to assess oral bioaccessibility of potentially toxic elements from an urban recreational site. *Applied Geochemistry*, 26: 789-796.
- Okoro, H. K., Fatoki, O. S., Adekola, F. A., Ximba, B. J., & Snyman, R. G. (2012). A review of sequential extraction procedures for heavy metals speciation in soil and sediments. *Open Access Scientific Reports*, 181(1).
- Olawoyin, R., Oyewole, S. A., & Grayson, R. L. (2012). Potential risk effect from elevated levels of soil heavy metals on human health in the Niger delta. *Ecotoxicology Environmental Safety*, 85: 120-130.
- Oliver, D. P., McLaughlin, M. J., Naidu, R., Smith, L. H., Maynard, E. J., Calder, I. C. (1999). Measuring Pb Bioavailable from Household Dusts Using an in vitro Model. *Environmental Sciences Technology*, 33: 4434-4439.
- Oomen, A. G., Hack, A., Minekus, M., Zeijdner, E., Schoeters, G., Verstraete, W., Wiele, T. V. D., Wragg, J., Rompelberg, C. J. M., Sips, A. J. A. M. & Wijnen, J. H. V. (2002). Comparison of Five in vitro Digestion Models to Study the Bioavailable of Soil Contaminants. *Environmental Sciences Technology*, 36: 3326.
- Oomen, A. G., Rompelberg, C. J. M., Bruil, M. A., Dobbe, C. J. G., Pereboom, D. P. K. H. & Sips, A. J. A. M. (2003). Development of an in vitro digestion model for estimation of bioavailable of soil contaminants. *Achieve Environmental Contamination Toxicology*, 44: 281-287.
- Oomen, A. G., Van Twillert, K., Hofhuis, M. F. A., Rompelberg, C. J. M. & Versantvoort, C. H. M. (2003). *Development and suitability of in vitro digestion models in assessing bioavailable of lead from toy matrices*. Report No. 320102001. National Institute for Public Health and the Environment.
- Osman, K. (1998). *Health effects of environmental lead exposure in children*. Doctoral thesis, Karolinska Institute, Institute of Environmental Medicine, Stockholm.

- Paustenbach, D. J. (2000). The practice of exposure assessment: A state-of-the-art review (Reprinted from Principles and Methods of Toxicology, 4th edition, (2001). *Journal of Toxicology Environmental Health B Critical Review*, 3: 179-291.
- Paustenbach, D. J. (2002). *Human and ecological risk assessment: Theory and practice*. New York, Wiley.
- Peijnenburg, W. J., Zablotskaja, M. & Vijver, & M. G. (2007). Monitoring metals in terrestrial environments within a bioavailable framework and a focus on soil extraction. *Ecotoxicology Environmental Safety*, 67: 163-179.
- Peltola, P. & Åström, M. (2003). Urban geochemistry: A multimedia and multielement survey of a small town in northern Europe. *Environmental Geochemistry and Health*, 25: 397-419.
- Peltoniemi, M., Heikkinen, J., & Makipaa, R. (2007). Stratification of regional sampling by model-predicted changes of carbon stocks in forested mineral soils. *Silva Fennica*, 41: 527.
- Petersson, L. (2005). Traffic related metals in soil and sediment in Mauritius.
- Petruzzelli, G., Szymura, I., Lubrano, L. & Pezzarossa, B. (1989). Chemical speciation of heavy metals in different size fractions of compost from solid urban wastes. *Environmental Technology Letter*, 10: 521-526.
- Plaster, E. (2013). *Soil science and management*. Cengage learning.
- Poggio, L., Vrscaj, B., Schulin, R., Hepperle, E., & Ajmone Marsan, F. (2009). Metal pollution and human bioaccessibility of topsoils in Grugliasco (Italy). *Environmental Pollution*, 157: 680-689.
- Poveromo, J. J. (1999). *Iron ore: Chapter 8 in the making, shaping and treating of steel*. 11th edition. Edited by DH Wakelin Warrendele, PA: Association for iron and steel technology, pp 547-642.
- Praveena, S. M., Aris, A. Z. & Radojevic, M. (2010). Heavy metals dynamics and source in intertidal mangrove sediment of Sabah, Borneo Island. *Environmental Asia*, 3: 72–81.
- Praveena, S. M., Yuswir, N. S., Aris, A. Z. & Hashim, Z. (2014). Potential Health Risk Assessment of Urban Soil on Heavy Metal Content in Seri Kembangan. *In From Sources to Solution pp 77-81. Springer Singapore*.
- Radojevic, M., & Bashkin, V. N. (1999). *Practical environmental analysis*. Royal Society of Chemistry, pp 340-348.
- Rao, B. S. N., Prabhavathi, T. (1978). An in vitro method for predicting the bioavailable of iron from foods. *American Journal of Clinical Nutritional*, 31: 169–175.

- Rayment, G. E., & Lyons, D. J. (2011). *Soil chemical methods: Australasia (Vol. 3)*. CSIRO publishing.
- Rodriguez, R. R., Basta, N. T., Casteel, S. W. & Pace, L. W. (1999). An in vitro gastrointestinal method to estimate bioavailable arsenic in contaminated soils and solid media. *Environmental Sciences Technology*, 33: 642-649.
- Ruby, M. V., Davis, A., Link, T. E., Schoof, R., Chaney, R. L., Freeman, G. B. & Bergstrom, P. (1993). Development of an in vitro screening test to evaluate the in vivo bioavailable of ingested mine-waste lead. *Environmental Sciences Technology*, 27: 2870-2877.
- Ruby, M. V., Davis, A., Schoof, R., Eberle, S. & Sellstone, C. M. (1996). Estimation of lead and arsenic bioavailable using a physiologically based extraction test. *Environmental Sciences Technology*, 39: 422-430.
- Salonen, V. P., & Korkka-Niemi, K. (2007). Influence of parent sediments on the concentration of heavy metals in urban and suburban soils in Turku, Finland. *Applied Geochemistry*, 22: 906-918.
- Samat, N., Hasni, R. & Elhadary, Y. A. E. (2011). Modelling land use changes at the peri-urban areas using geographic information systems and cellular automata model. *Journal of Sustainable Development*, 4: 72.
- Sany, S. B. T., Salleh, A., Sulaiman, A. H., Sasekumar, A., Rezayi, M., & Tehrani, G. M. (2012). Heavy metal contamination in water and sediment of the Port Klang coastal area, Selangor, Malaysia. *Environmental Earth Sciences*, 69: 2013-2025.
- Sapari, N., Idris, A., Hisham, N. & Hamid, A (1996). Total removal of heavy metal from mixed plating rinse wastewater. *Desalination*, 106: 419-422.
- Sasirekha, K., & Baby, P. (2013). Agglomerative Hierarchical Clustering Algorithm-A. *International Journal of Scientific and Research Publications*, 3: 1-3.
- Scancar, J., Milacic, R. & Hoorvat, M. (2000). Comparison of various digestion and extraction procedures in analysis of heavy metals in sediments. *Water, Air Soil Pollution*, 118: 87-99.
- Schmidt, C. W. (2010). Lead in Air: Adjusting to a new standard. *Environmental Health Perspectives*, 118: A76-A79.
- Schroeder, J. L., Basta, N. T., Casteel, S. W., Evans, T., Payton, T. J. & Si, J. (2004). Validation of the in vitro intestinal (IVG) method to estimate relative bioavailable lead in contaminated soils. *Journal of Environmental Quality*, 33: 513-521.
- Schroeder, J. L., Basta, N. T., Si, J., Casteel, S. W., Evans, T. & Payton, M. (2003). In vitro gastrointestinal method to estimate relative bioavailable cadmium in contaminated soil. *Environmental Sciences Technology*, 37: 1365-1370.

- Schütz, A., Barregård, L., Sällsten, G., Wilske, J., Manay, N., Pereira, L. & Cousillas, Z.A. (1997). Blood lead in Uruguayan children and possible sources of exposure. *Environmental Research*, 74: 17-23.
- Science Communication Unit, University of the West of England, (2013). *Soil Contamination: Impacts on Human Health*. European Commission.
- Shefsky, S. (1997). *Sample handling strategies for accurate lead-in-soil measurements in the field and laboratory*. Field Analytical Methods for Hazardous Wastes and Toxic Chemicals (A&WMA), Las Vegas, 29-31.
- Selim, H.M, Sparks, & D.L., (2001). *Heavy Metals Release in Soils*. CRC Press, Boca Raton, FL.
- Sezgin, N., Ozcan, H. K., Demir, G., Nemlioglu, S. & Bayat, C. (2003). Determination of heavy metal concentrations in street dusts in Istanbul E-5 highway. *Environment International*, 29: 979–985.
- Shamshuddin, J., & Anda, M. (2008). Charge properties of soils in Malaysia dominated by Kaolinite, Gibbsite, Goethite, and Hematite. *Bulletin Geological Society Malaysia*, 54: 27-31.
- Shazili, N. A. M., Yunus, K., Ahmad, A. S., Abdullah, N. & Rashid, M. K. A. (2006). Heavy metal pollution status in the Malaysia aquatic environment. *Aquatic Ecosystem Health Management*, 9: 137–145.
- Shi, G., Chen, Z., Xu, S., Zhang, J., Wang, L., Bi, C., & Teng, J. (2008). Potentially toxic metal contamination of urban soils and roadside dust in Shanghai, China. *Environmental Pollution*, 156: 251-260.
- Siciliano, S. D., James, K., Zhang, G., Schafer, A. N., & Peak, J. D. (2009). Adhesion and enrichment of metals on human hands from contaminated soil at an Arctic urban brownfield. *Environmental Science & Technology*, 43: 6385-6390.
- Singh, R., Gautam, N., Mishra, A. & Gupta, R. (2011). Heavy metals and living systems: an overview. *Indian Journal Pharmacology*, 43: 246.
- Soriano, A., Pallarés, S., Pardo, F., Vicente, A.B., Sanfeliu, T. & Bech, J. (2012). Deposition of heavy metals from particulate settleable matter in soils of an industrialised area. *Journal of Geochemical Exploration*, 113: 36-44.
- Sun, Y., Zhou, Q., Xie, X., & Liu, R. (2010). Spatial, sources and risk assessment of heavy metal contamination of urban soils in typical regions of Shenyang, China. *Journal of Hazardous Materials*, 174: 455-462.
- Swartjes, F. A. (Ed.). (2011). *Dealing with contaminated sites: from theory towards practical application*. Springer Science & Business Media.
- Tchounwou, P. B., Yedjou, C. G., Patlolla, A. K., and Sutton, D. J. (2012). *Heavy metal toxicity and the environment*. In Molecular, Clinical and Environmental Toxicology (pp. 133-164). Springer Basel.

- Theodore, L., & Dupont, R. R. (2012). *Environmental health and hazard risk assessment: principles and calculations*. CRC Press.
- Thornton, I., Farago, M. E., Thums, C. R., Parrish, R. R., McGill, R. A., Breward, N., ... & Watt, J. (2008). Urban geochemistry: research strategies to assist risk assessment and remediation of brownfield sites in urban areas. *Environmental Geochemistry And Health*, 30: 565-576.
- Trochim, W. M., & Donnelly, J. P. (2001). *Research methods knowledge base*. Cornell University.
- Tume, P., Bech, J., Sepulveda, B., Tume, L., & Bech, J. (2008). Concentrations of heavy metals in urban soils of Talcahuano (Chile): a preliminary study. *Environmental Monitoring And Assessment*, 140: 91-98.
- United States Department of Agriculture, (1993). *From the Surface Down An Introduction to Soil Surveys for Agronomic Use*. Second Edition.
- USDOE, (2011) *The Risk Assessment Information System (RAIS)*. U.S. Department of Energy's Oak Ridge Operations Office (ORO).
- USEPA, (2002) *Supplemental guidance for developing soil screening levels for superfund sites*. U.S. Environmental Protection Agency pp 4-24.
- USEPA, (2011) *Integrated Risk Information System (IRIS)*. U.S. Environmental Protection Agency.
- Versantvoort, C. H. M., Van de Kamp, E. & Rompelberg, C. J. M. (2004). *Development and applicability of an in vitro digestion model in assessing the bioavailable of contaminants from food*. Report No 320102002. National Institute for Public.
- Wang, X. P., Sheng, J. J., Gong, P., Xue, Y. G., Yao, T. D., & Jones, K. C. (2012). Persistent organic pollutants in the Tibetan surface soil: Spatial distribution, air-soil exchange and implications for global cycling. *Environmental Pollution*, 170: 145-151.
- Wasay, S.A., Barrington, S., & Tokunaga, S. (2001). Organic acids for the in situ remediation of soils polluted by heavy metals: soil flushing in columns. *Water, Air, and Soil Pollution*, 127: 301-314.
- Wei, B., & Yang, L. (2010). A review of heavy metal contaminations in urban soils, urban road dusts and agricultural soils from China. *Microchemical Journal*, 94: 99-107.
- Wilcke, W., Müller, S., Kanchanakool, N., & Zech, W. (1998). Urban soil contamination in Bangkok: heavy metal and aluminium partitioning in topsoils. *Geoderma*, 86: 211-228.

- Williams, T. M., Rawlins, B. G., Smith, B., & Breward, N. (1998). In-vitro determination of arsenic bioavailable in contaminated soil and mineral beneficiation waste from Ron Phibun, Southern Thailand: A basis for improved human risk assessment. *Environmental Geochemical Health*, 20: 169-177.
- Wong, C. S. C., Li, X. D. & Thornton, I. (2006). Urban environmental geochemistry of trace metals. *Environmental Pollution*, 142: 1-16.
- World Health Organization, (1993). *Guidelines for drinking-water quality*. World Health Organization Volume 1.
- Wuana, R. A., & Okieimen, F. E. (2011). *Heavy metals in contaminated soils: a review of sources, chemistry, risks and best available strategies for remediation*. ISRN Ecology, 2011.
- Wust, R. A., Ward, C. R., Bustin, R. M. & Hawke, M. I. (2002). Characterization and quantification of inorganic constituents of tropical peats and organic rich deposits from Tasek Bera (Peninsular Malaysia): implications for coals. *International Journal of Coal Geology*, 49: 215-249.
- Xia, X., Chen, X., Liu, R., & Liu, H. (2011). Heavy metals in urban soils with various types of land use in Beijing, China. *Journal of Hazardous Materials*, 186: 2043-2050.
- Yafa, C., & Farmer, J. G. (2006). A comparative study of acid-extractable and total digestion methods for the determination of inorganic elements in peat material by inductively coupled plasma-optical emission spectrometry. *Analytica Chimica Acta*, 557: 296-303.
- Yang, Z., Lu, W., Long, Y., Bao, X. & Yang, Q. (2011). Assessment of heavy metals contamination in urban topsoil from Changchun City, China. *Journal of Geochemical Explore*, 108: 27-38.
- Yap, C. K., Ismail, A. & Tan, S. G. (2003). Cd and Zn concentrations in the straits of Malacca and intertidal sediments of the west coast of Peninsular Malaysia. *Marine Pollution Bulletin*, 46: 1341-1358.
- Yongming, H., Peixuan, D., Junji, C. & Posmentier, E. S. (2006). Multivariate analysis of heavy metal contamination in urban dusts of Xi'an, Central China. *Sciences Total Environment*, 355: 176-186.
- Yu, C. H., Yin, L. H., & Paul, J. L. (2006). The Bioavailable of Lead (Pb) from Vacuumed House Dust on Carpets in Urban Residences. *Risk Analysis*, 26: 125-134.
- Yuswir, N. S., Praveena, S. M., Aris, A. Z., Hashim, Z. (2013). Bioavailable of heavy metals using in vitro digestion model: a state of present knowledge. *Review on Environmental Health*, 128: 181-187.

Zarcinas, B. A., Ishak, C. F., McLaughlin, M. J. & Cozens, G. (2004). Heavy metals in soils and crops in Southeast Asia. *Environmental Geochemistry and Health*, 26: 343-357.

Zhitkovich, A., Song, Y., Quievryn, G. & Voitkun, V. (2001). Non-oxidative mechanisms are responsible for the induction of mutagenesis by reduction of Cr(VI) with cysteine: role of ternary DNA adducts in Cr(III)-dependent mutagenesis. *Biochemistry*, 40: 549-560.



LIST OF PUBLICATIONS

- Yuswir, N. S., Praveena, S. M., Aris, A. Z., & Hashim, Z. (2013). Bioavailability of heavy metals using in vitro digestion model: a state of present knowledge. *Reviews on environmental health*, 28(4), 181-187.
- Yuswir, N. S., Praveena, S. M., Aris, A. Z., Hashim, Z., & Ismail, S. N. S. (2014). Health Risk Assessment on Bioavailability of Heavy Metals in Klang District Urban Surface Soil. *Iranian Journal of Public Health*, 43(3), 167-171.
- Yuswir, N. S., Praveena, S. M., Aris, A. Z., Hashim, Z., & Ismail, S. N. S. (2015). Health risk assessment of heavy metal in urban surface soil of Klang district (Malaysia). *Bulletin of Environmental Contamination and Toxicology*, 1-10
- Yuswir, N. S., Praveena, S. M., Aris, A. Z., Syed Ismail, S. N., De Burbure, C., & Hashim, Z. (2015). Heavy metal contamination in urban surface soil of Klang District (Malaysia). *Soil and Sediment Contamination: An International Journal*, (just-accepted).



UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : _____

TITLE OF THESIS / PROJECT REPORT :

HEAVY METALS IN SOIL FROM URBAN AREAS OF KLANG, MALAYSIA, AND THEIR
HEALTH RISK ASSESSMENT USING *IN VITRO* DIGESTION METHOD

NAME OF STUDENT : NURUL SYAZANI BINTI YUSWIR

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (✓)

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

OPEN ACCESS

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

PATENT

Embargo from _____ until _____
(date) (date)

Approved by:

(Signature of Student)
New IC No/ Passport No.:

Date :

(Signature of Chairman of Supervisory Committee)
Name:

Date :

[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted.]