



**UNIVERSITI PUTRA MALAYSIA**

***MODIFICATION OF KOSARIC'S MEDIUM WITH NITROGEN AND  
PHOSPHORUS FOR *Spirulina* (*Arthrospira platensis*) IN SHELTERED  
OPEN CULTURE SYSTEM***

**NURSHAZREEN MASHOR**

**FS 2015 50**



**MODIFICATION OF KOSARIC'S MEDIUM WITH NITROGEN AND  
PHOSPHORUS FOR *Spirulina*  
(*Arthrospira platensis*) IN SHELTERED OPEN CULTURE SYSTEM**

By

**NURSHAZREEN BINTI MASHOR**

**Thesis Submitted to the School of Graduate Studies,  
Universiti Putra Malaysia, in Fulfillment of the  
Requirements for the Degree of Master of Science**

**August 2015**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia  
fulfillment of the requirement for the degree of Master of Science

**MODIFICATION OF KOSARIC'S MEDIUM WITH COMMERCIAL NITROGEN  
AND PHOSPHORUS FOR SPIRULINA (*Arthrospira platensis*) IN  
SHELTERED OPEN CULTURE SYSTEM**

By

**NURSHAZREEN BINTI MASHOR**

**August 2015**

**Chairman: Hishamuddin Bin Omar, PhD**  
**Faculty: Science**

Presently the world is experiencing serious environmental crisis such as global warming that effect the source of human nutrition hence, increasing starvation, and malnutrition. Demand for microalgae as supplemental food had increased steadily in recent years. However, in Malaysia, cultivation of microalgae is scanty due to lack of knowledge in microalgae production in variable weather conditions. One of the fundamental problems is lack of cheap and effective fertilizer.

The aim of this research was to develop economical and inexpensive fertilizer for a commercial production and give higher productivity of *A. platensis*. The objective was implemented by substituting all the nitrogen and phosphorus sources present in Kosaric Medium (SM) with locally available commercial fertilizers. The *A. platensis* was treated with 19 treatments include SM as control with three different nitrogen sources (ammonium nitrate, ammonium sulphate, and urea) and three different phosphorus sources (phosphoric acids, triple superphosphate (TSP), and diammonium phosphate (DAP)). Two grades of nitrogen concentration representing 50% (high concentration) and 25% (low concentration) of SM nitrogen concentration were compared in this study. 30L *A. platensis* was grown in each polyethylene (PE) bag in outdoor condition for seven days in first cultivation, ten days in second cultivation and 8 days in third cultivation. The growth parameters (optical density, biomass and chlorophyll *a* concentration) and biochemical content (proteins, carbohydrate and lipid) were recorded and compared with *A. platensis* cultivated in SM.

Significant differences of the growth parameters and biochemical composition were recorded for the different nitrogen and phosphorus sources and concentration. The result of this study revealed that *A. platensis* cultivation during dry season had better performance and *A. platensis* could utilize ammonium nitrate (ammonium nitrate + triple superphosphate in low concentration) most efficiently and gave the highest productivity on day 8 with

biomass, chlorophyll, and protein yield of  $1.24 \pm 0.004\text{g/L}$ ,  $11.41 \pm 0.21\text{mg/L}$ , and 62.5%, respectively with average of pH,  $10.12 \pm 0.08$  which was significantly different ( $p < 0.05$ ) with other treatments. This was better than that of SM ( $0.73 \pm 0.01\text{g/L}$ ,  $8.64 \pm 0.13\text{mg/L}$ , and 52%, respectively). Further increased in nitrogen concentration will limit growth.

This study indicated that the utilization of ammonium nitrate and TSP as nitrogen and phosphorus has the potential to replace the expensive Kosaric media. The modified medium can be used cost effectively for large scale mass production of protein rich *A. platensis* and yields similar performances to Kosaric medium.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk Ijazah Master Sains

**PENGUBAHSUAIAN MEDIA KOSARIC DENGAN NITROGEN DAN  
PHOSPHORUS UNTUK SPIRULINA (*Arthrospira platensis*) DALAM  
SISTEM PENGKULTURAN DI LUAR MAKMAL YANG TERLINDUNG**

Oleh

**NURSHAZREEN BINTI MASHOR**

**Ogos 2015**

**Pengerusi: Hishamuddin Omar, PhD**  
**Fakulti: Sains**

Perubahan iklim yang ekstrim juga dikenali pemanasan global semakin kritikal pada masa ini memberi kesan ke atas sumber makanan dan nutrisi manusia terjejas menyebabkan kebuluran dan kekurangan zat makanan. Permintaan alga sebagai makanan tambahan semakin meningkat pada abad ini. Walau bagaimanapun, pengkulturan mikroalga di Malaysia tidak dilakukan secara besar-besaran dan masih dijalankan di dalam makmal disebabkan kekurangan kajian dan pengetahuan dalam pengkulturan mikroalga yang dijalankan dalam keadaan cuaca yang berubah-ubah.

Oleh itu, tujuan kajian ini dijalankan adalah untuk menghasilkan media kultur yang efektif dengan kos media yang berpatutan untuk pengkulturan *A. platensis* berskala besar serta memberikan produktiviti yang lebih tinggi. Objektif kajian telah dilaksanakan dengan menggantikan sumber nitrogen dan phosphorus dalam media Kosaric (SM) dengan baja nitrogen dan phosphorus komersial tempatan yang sedia ada. Pertumbuhan *A. platensis* diuji dengan 19 rawatan termasuk standard media (SM) sebagai tanda aras. Tiga jenis sumber nitrogen komersial (ammonium nitrat, ammonium sulfat, dan urea) dan phosphorus (asid phosphoric, triple superphosphate (TSP), and diammonium phosphate (DAP)) diuji dengan kepekatan tinggi (50%) dan rendah (25%) daripada kepekatan asal sumber nitrogen dan phosphorus dalam SM.

30L *A. platensis* telah dikulturkan dalam beg polyethylene (PE) diluar makmal selama tujuh hari pada pengkulturan pertama, 10 hari pengkulturan kedua dan lapan hari pada pengkulturan ketiga. Parameter pertumbuhan (ketumpatan optik, berat bersih dan kepekatan chlorophyll *a*) dan kandungan biokimia (protein, karbohidrat dan lipid) telah direkodkan dan dibandingkan dengan kesemua rawatan.

Pertumbuhan *A. platensis* dan komposisi biokimia dalam T16 telah menunjukkan perbezaan yang ketara secara statistik antara kesemua rawatan yang diuji. Hasil kajian menunjukkan pertumbuhan *A. platensis* adalah tinggi pada pengkulturan pada musim panas dan menunjukkan bahawa *A. platensis*

dapat menggunakan ammonium nitrat (ammonium nitrat + triple superphosphat, kepekatan rendah) secara optimum dan menghasilkan produktiviti tertinggi pada hari ke lapan dengan berat bersih, klorofil, dan protein  $1.24 \pm 0.004\text{g/L}$ ,  $11.41 \pm 0.21\text{mg/l}$ , and 62.5%, masing-masing dengan purata pH,  $10.12 \pm 0.08$ . Ini ada lebih tinggi berbanding *A. platensis* yang dikulturkan dalam SM dengan masing-masing ( $0.73 \pm 0.01\text{g/L}$ ,  $8.64 \pm 0.13\text{mg/L}$ , and 52%). Peningkatan kepekatan nitrogen telah menyebabkan pertumbuhan *A. platensis* menjadi terhad.

Kajian ini menunjukkan bahawa penggunaan ammonium nitrat dan tsp sebagai sumber nitrogen dan phosphorus mempunyai potensi untuk menggantikan media Kosaric yang mahal. Medium yang diubahsuai boleh digunakan untuk pengeluaran *A. platensis* yang kaya protein secara berskala besar dan menghasilkan pertumbuhan yang lebih tinggi dengan kos yang efektif dibandingkan dengan media Kosaric.

## ACKNOWLEDGEMENTS

Alhamdulillah and praise to Allah S.W.T for all the blessing and ease throughout the project. The completion of this undertaking could not have been possible without the participation and assistance of so many people whose name may not all be enumerated.

First of all, I would like to express my greatest appreciation to my supervisor, Dr. Hishamuddin bin Omar for his guidance to make this project a success. Thanks for the extensive comments, and assistance given throughout this project.

Special thanks to all lecturers and lab assistant from Plant Biochemistry Lab, Biology Department, Faculty of Science for the facilities in the lab and allowed me to conduct the proximate analysis. Not forget to my co-supervisor Dr. Shahrizim B. Zulkifly for the ideas and supports during the research works. My appreciation is extend towards the cooperation of the staff of Biology Department for the assistance and kindness support.

My deepest gratitude to my lab mates and friends, Mohd. Shahmen, Noraaini Mastor, Siti Norhidayah Othman, Adibah Shakri and Aqlili Riana for their support, encouragement and understanding spirit throughout our Master Degree journey. All the best to all of you in your future undertaking.

Finally, I would like to extend my deepest gratitude to my parents and family for their endless support either morally, financially and physically.



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Hishamuddin Bin Omar, PhD**

Senior Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

**Shahrizim Bin Zulkifly, PhD**

Senior Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Member)

---

**BUJANG BIN KIM HUAT, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in theUniversiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Nurshazreen Binti Mashor, GS30153

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_

Name of Chairman  
of Supervisory  
Committee:

Dr. Hishamuddin Bin Omar

Signature: \_\_\_\_\_

Name of Member of  
Supervisory  
Committee:

Dr. Shahrizim Bin Zulkifly

## TABLE OF CONTENT

	Page
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xiii
<b>LIST OF FIGURES</b>	xvi
<b>LIST OF ABBREVIATIONS</b>	xviii
 <b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
1.1 Problem statement	1
1.2 Justification of study	2
 <b>2 LITERATURE REVIEW</b>	<b>4</b>
2.1 Global warming: definition, causes and the effect	4
2.2 How microalgae can mitigate global warming	5
2.3 Present status of microalgae research	5
2.4 Taxonomy and Morphology of <i>A. platensis</i>	6
2.5 The potential of <i>A. platensis</i> as source of food	7
2.6 Current studies of <i>A. platensis</i>	7
2.6.1 Mass outdoor cultivation strategies	7
2.6.2 Challenges (unstable weather and frequent cloud cover)	8
2.6.3 Culture media	9
2.6.4 Fertilizer	10
2.7 Factors affecting the growth and productivity of <i>A. platensis</i>	11
2.7.1 Light	11
2.7.2 Temperature	11
2.7.3 pH	12
2.7.4 Salinity	12
2.7.5 Agitation and Aeration	12
2.8 Biochemical composition of <i>A. platensis</i>	13
2.8.1 Chlorophyll	13
2.8.2 Carbohydrate	13
2.8.3 Lipid	13
2.8.4 Protein	14
 <b>3 MATERIALS AND METHODS</b>	<b>16</b>
3.1 Site of study	16
3.2 Study duration	16
3.3 Source of <i>A. platensis</i>	16
3.4 Maintenance of stock culture	16
3.5 Culture media	17

	3.5.1	Kosaric's medium	17
	3.5.2	Formulation of test fertilizer	17
	3.6	Experimental design	18
	3.7	Outdoor cultivation setup	19
	3.8	Weather condition	20
	3.9	Growth monitoring of <i>A. platensis</i>	20
	3.9.1	Growth parameters and environment factors	20
	3.9.2	Cell dry weight determination	20
	3.9.3	Specific growth rate	21
	3.9.4	Chlorophyll a content	21
	3.10	Harvesting and sample handling	22
	3.11	Samples preservation	22
	3.12	Biochemical analysis	23
	3.12.1	Total protein	23
	3.12.2	Total lipid	23
	3.12.3	Total carbohydrate	24
	3.13	Statistical analysis	25
4		<b>RESULT</b>	26
	4.1	Preliminary result	26
	4.2	Physical parameter of culture media (Actual experiment)	28
	4.2.1	pH	28
	4.2.2	Temperature	33
	4.2.3	Light intensity during first and second cultivation cycle	37
	4.3	Growth parameter of <i>A. platensis</i> in different concentration of nitrogen and phosphorus sources in two different cultivation cycles	40
	4.3.1	Optical density	40
	4.3.2	Chlorophyll a content	46
	4.3.3	Dry weight	52
	4.4	Cultivation of <i>A. platensis</i> using optimum concentration of urea and triple super phosphate (low concentration), ammonium sulphate and phosphoric acid (high concentration), ammonium nitrate and triple super phosphate (low concentration)	58
	4.4.1	Physical properties of culture media	59
	4.4.1.1	pH	59
	4.4.1.2	Temperature	60
	4.4.2	Growth performance of <i>A. platensis</i>	60
	4.4.2.1	Optical density of <i>A. platensis</i>	60
	4.4.2.2	Chlorophyll a content of <i>A. platensis</i>	62

	4.4.2.3	Dry Weight of <i>A. platensis</i>	63
4.4.3		Biochemical properties of <i>A. platensis</i>	66
	4.4.3.1	Total protein content of <i>A. platensis</i>	66
	4.4.3.2	Total carbohydrate content of <i>A. platensis</i>	67
	4.4.3.3	Total lipid content of <i>A. platensis</i>	68
<b>5</b>		<b>DISCUSSION</b>	<b>70</b>
	5.1	The importance of acclimatization	70
	5.1.1	Preliminary study	70
	5.2	Environmental parameter	71
	5.2.1	Effect of light	71
	5.2.2	Effect of temperature	72
	5.2.3	Effect of Ph	73
	5.3	Effect of different concentration of nitrogen and phosphorus sources	74
	5.4	Biochemical composition	78
<b>6</b>		<b>SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	<b>80</b>
		<b>REFERENCES</b>	<b>81</b>
		<b>APPENDICES</b>	<b>92</b>
		<b>BIODATA OF STUDENT</b>	<b>103</b>
		<b>LIST OF PUBLICATION</b>	<b>104</b>

## LIST OF TABLES

Table	Page
2.1 Outdoor Mass Cultivation of <i>A. platensis</i>	8
2.2 General composition of different algae (% of dry matter)	15
3.1 List of nitrogen and phosphorus fertilizer with different concentration level	18
4.1 Average dry weight in seven days cultivation and specific growth rate of <i>A. platensis</i>	27
4.2 Average pH in first and second cultivation of <i>A. platensis</i>	32
4.3 Light intensity ( $\mu\text{molm}^{-2}\text{s}^{-1}$ ) at different time interval of <i>A. platensis</i> cultivation cycle	39
4.4ai Optical density of <i>A. platensis</i> supplemented with different concentration of urea and phosphorus sources in first cultivation	41
4.4aii Optical density of <i>A. platensis</i> supplemented with different concentration of urea and phosphorus sources in second cultivation	41
4.5bi Optical density of <i>A. platensis</i> supplemented with different concentration of ammonium sulphate and phosphorus sources in first cultivation	43
4.5bii Optical density of <i>A. platensis</i> supplemented with different concentration of ammonium sulphate and phosphorus sources in second cultivation	43
4.6ci Optical density of <i>A. platensis</i> supplemented with different concentration of ammonium nitrate and phosphorus sources in first cultivation	45
4.6cii Optical density of <i>A. platensis</i> supplemented with different concentration of ammonium nitrate and phosphorus sources in second cultivation	45
4.7ai Chlorophylla of <i>A. platensis</i> supplemented with different concentration of urea and phosphorus sources in first cultivation	47
4.7aii Chlorophylla of <i>A. platensis</i> supplemented with different concentration of urea and phosphorus sources in second	47

	cultivation	
4.8bi	Chlorophyll a of <i>A. platensis</i> supplemented with different concentration of ammonium sulphate and phosphorus sources in first cultivation	49
4.8bii	Chlorophyll a of <i>A. platensis</i> supplemented with different concentration of ammonium sulphate and phosphorus sources in second cultivation	49
4.9ci	Chlorophyll a of <i>A. platensis</i> supplemented with different concentration of ammonium nitrate and phosphorus sources in first cultivation	51
4.9cii	Chlorophyll a of <i>A. platensis</i> supplemented with different concentration of ammonium nitrate and phosphorus sources in second cultivation	51
4.10ai	Dry weight of <i>A. platensis</i> supplemented with different concentration of urea and phosphorus sources in first cultivation	53
4.10aii	Dry weight of <i>A. platensis</i> supplemented with different concentration of urea and phosphorus sources in second cultivation	53
4.11bi	Dry weight of <i>A. platensis</i> supplemented with different concentration of ammonium sulphate and phosphorus sources in first cultivation	55
4.11bii	Dry weight of <i>Arthrospira platensis</i> supplemented with different concentration of ammonium sulphate and phosphorus sources in second cultivation	55
4.12ci	Dry weight of <i>A. platensis</i> supplemented with different concentration of ammonium nitrate and phosphorus sources in first cultivation	57
4.12cii	Dry weight of <i>A. platensis</i> supplemented with different concentration of ammonium nitrate and phosphorus sources in second cultivation	57
4.13	The average of pH in all treatments for 8 days	59
4.14	Optical density and specific growth rate of <i>A. platensis</i>	61
4.15	The average of chlorophyll content of <i>A. platensis</i> (mg/l)	62
4.16	Average dry weight and specific growth rate of <i>A. platensis</i>	65
5.1	Comparison of <i>A. platensis</i> productivity using urea and	76



ammonium as a nitrogen source

- 5.2 Protein, carbohydrate, and lipid contents of the *A. platensis* biomass recorded using different sources of nitrogen 78



## LIST OF FIGURES

Figure		Page
3.1	The arrangement of <i>A. platensis</i> culture bags in outdoor culture system	19
4.1	Biomass (mean dry weight $\pm$ standard error) of <i>A. platensis</i> cultured in the media containing different sources of nitrogen and phosphorus as fertilizers.	26
4.2a	pH of culture media of <i>A. platensis</i> supplemented by different concentration of urea and phosphorus sources	29
4.2b	pH of culture media of <i>A. platensis</i> supplemented by different concentration of ammonium sulphate and phosphorus sources	30
4.2c	pH of culture media of <i>A. platensis</i> supplemented by different concentration of ammonium nitrate and phosphorus sources	31
4.3a	Temperature of air and culture media of <i>A. platensis</i> supplemented by different concentration of urea and phosphorus sources	34
4.3b	Temperature of air and culture media of <i>A. platensis</i> supplemented by different concentration of ammonium sulphate and phosphorus sources	35
4.3c	Temperature of air and culture media of <i>A. platensis</i> supplemented by different concentration of ammonium nitrate and phosphorus sources	36
4.4	Light intensity and air temperature during <i>A. platensis</i> cultivation. A: first cycle, B: second cycle.	38
4.5	pH content of <i>A. platensis</i> as cultured in Kosaric media (control), T4 (urea + tsp, low concentration), T7 (ammonium sulphate + phosphoric acid, high concentration), T16 (ammonium nitrate + tsp, low concentration)	59
4.6	Air and culture media temperature of <i>A. platensis</i> as cultured in Kosaric's media (control), T4 (urea + tsp, low concentration), T7 (ammonium sulphate + phosphoric acid, high concentration), T16 (ammonium nitrate + tsp, low concentration)	60
4.7	Optical density of <i>A. platensis</i> as cultured in Kosaric's (control), T4 (urea + tsp, low concentration), T7 (ammonium sulphate + phosphoric acid, high concentration), T16 (ammonium nitrate + tsp, low concentration)	62

4.8	Chlorophyll content of <i>A. platensis</i> as cultured in Kosaric's media (control), T4 (urea + tsp, low concentration), T7 (ammonium sulphate + phosphoric acid, high concentration), T16 (ammonium nitrate + tsp, low concentration)	63
4.9	Dry weight of <i>A. platensis</i> as cultured in Kosaric's media (control), T4 (urea + tsp, low concentration), T7 (ammonium sulphate + phosphoric acid, high concentration), T16 (ammonium nitrate + tsp, low concentration)	64
4.10	Total protein content of <i>A. platensis</i> as supplemented with Kosaric's media (control), T4 (urea + tsp, low concentration), T7 (ammonium sulphate + phosphoric acid, high concentration), T16 (ammonium nitrate + tsp, low concentration)	66
4.11	Total carbohydrate content of <i>A. platensis</i> as supplemented with Kosaric's media (control), T4 (urea + tsp, low concentration), T7 (ammonium sulphate + phosphoric acid, high concentration), T16 (ammonium nitrate + tsp, low concentration)	67
4.12	Total lipid content of <i>A. platensis</i> as supplemented with Kosaric's media (control), T4 (urea + tsp, low concentration), T7 (ammonium sulphate + phosphoric acid, high concentration), T16 (ammonium nitrate + tsp, low concentration)	68
4.13	Total protein, carbohydrate, and lipid content of <i>A. platensis</i> in Kosaric's medium (control), T4 (urea + tsp, low concentration), T7 (ammonium sulphate + phosphoric acid, high concentration), T16 (ammonium nitrate + tsp, low concentration)	69

## LIST OF ABBREVIATIONS

%	Percentage
μl	Microlitre
μm	Micrometer
μmolm <sup>-2</sup> s <sup>-1</sup>	Micromoles per square meter per second
CH <sub>4</sub> N <sub>2</sub> O	Urea
cm	Centimetre
CO <sub>2</sub>	Carbon Dioxide
DAP	Diammonium Phosphate
DNA	Deoxyribo Nucleic Acid
g	Gram
g/l	Gram per litre
H <sub>3</sub> PO <sub>4</sub>	Phosphoric Acid
HCL	Hydrochloric
HDPE	High Density Polyethylene
m	Meter
mg/l	Miligram per litre
MgCl <sub>2</sub>	Magnesium Chloride
ml	Mililitre
NaCl	Sodium chloride
NaNO <sub>3</sub>	Sodium Nitrate
NaOH	Sodium Hydroxide
NH <sub>3</sub>	Ammonia
NH <sub>4</sub>	Ammonium
NO <sub>2</sub>	Nitrite
PE	Polyethylene
rpm	Rotation per minute
SM	Standard Media
TSP	Triple Super Phosphate
UV	Ultraviolet
α	Alpha
β	Beta
°C	Centrugade Celcius

## CHAPTER 1

### INTRODUCTION

#### 1.1 Problem statement

Presently the world is experiencing serious environmental crisis. The quest for affluent lifestyle and economic dominance has resulted in global warming (Liz, 2014; Xue *et al.*, 2011; Soletto *et al.*, 2008). Accumulation of methane and carbon dioxide (CO<sub>2</sub>) in the atmosphere trap solar energy and heat generated from human activities in the earth atmosphere resulting in present global warming. Global warming not only causing high atmospheric temperature but the effect is more far ranging. Global warming caused climate change which results in climate change altering weather pattern: atmospheric air circulation, oceanic water circulation causing too much rain in some part of the world, too little rain in other part of the world, typhoon, hurricane, tornado, and acid rain (Peter *et al.*, 2011). The collective effects cause havoc to the environment, and well being of the earth population. Diseases, rising of the sea water level and crop failure is threatening the world population particularly the poorer countries. As a consequence, the source of human nutrition is reduced, increasing starvation and malnutrition.

So much has been debated about global warming at international forum but the action has been very slow. This is because developed countries is unwilling to make the unnecessary action to reduce carbon foot print but instead accusing poorer countries of cutting the rainforest. United Nation has identified that bulk of the carbon emission originate from North America, Europe and China (Liz, 2014). Scientist has recommended various measures to reduce green house gas such as adopting green technology, reducing fossil energy consumption, recycling and microalgae cultivation (Christopher, 2014; Raja *et al.*, 2008; Converti *et al.*, 2006; Binaghi *et al.*, 2003).

Recent studies have shown that microalgae are the potential candidate for renewable biofuel (Parmar *et al.*, 2011; Xin *et al.*, 2010; Chen *et al.*, 2009;). This is of no surprise because microalgae are one of the pioneer organisms living on earth and changing earth into habitable planet. Microalgae has the potential to reduce global warming because as stated by Garcia (2000), microalgae act as primary producer of oxygen where it uses CO<sub>2</sub> and produce oxygen (O<sub>2</sub>), and functional in carbon and nitrogen fixation while at the same time produce valuable biomass high in protein and lipid. The advantage of microalgae cultivation is that it can be carried out on land or sea. Due to the limited land area to accommodate a growing population and agricultural activity, microalgae culture is very flexible and can be done at sea or freshwater areas. It does not require a large area for high productivity. Regarding to this, this biomass can be used for food, feed, and biofuel.

Unfortunately most of the researches are carried out in the laboratories focusing on growth studies where the environment and growth factors are being

controlled (Anaga and Abu, 1996), finding the fast growing species (Day *et al.*, 2012) and high lipid content (Jiang *et al.*, 2011; Pruvost *et al.*, 2011). However very few of the studies is applicable in the real field conditions. This is because microalgae are very sensitive to the environment they are growing. Slight changes to the culture conditions will inhibit their growth. Therefore more effort should be devoted to microalgae culture in outdoor conditions. One of the microalgae adapted well in outdoor conditions is *Arthrospira* sp. commonly known as Spirulina.

*A. platensis* is a blue green algae which can be found in wide geographical distribution (Uday and Ahluwalia, 2013). It is very hardy species and can live where other microalgae facing difficulties to establish, therefore it is almost free from contaminating microalgae. *A. platensis* has been cultured in many parts of the world, good growth, and high in nutritional values where commercial cultivation of *A. platensis* are well established on a few countries including China, India, Bangladesh, America and Africa. *A. platensis* has been known as animal and human food in countries such as Asia and Africa. In addition, *A. platensis* has a high market demand and rich in pigments such as phycobiliproteins, beta carotene and luteina for pharmaceutical and nutraceutical industries (Raoof *et al.*, 2006).

## 1.2 Justification of study

As *A. platensis* is known as a potential solution to the problem of world protein source, various studies conducted and found that it has higher protein content than vegetables (Becker, 2007; FAO, 2001). *A. platensis* gives high productivity in a short time and easily adapted to a variety of climates.

Despite its potential and versatility, commercial *A. platensis* cultivation is almost nonexistent in Malaysia. This is because Malaysia lies in the tropic, with frequent rain and cloud covers and the condition is not conducive for outdoor cultivation. Most of the available references pertaining to *A. platensis* culture are from subtropical and arid region where the weather is fairly stable and clear sky (Isichei, 1990 and Hernandez *et al.*, 2009). Although Malaysia's weather is not conducive for outdoor cultivation, preliminary studies have shown that *A. platensis* can be grown in outdoor condition under a rain shelter (Puganeswary *et al.*, 2014). This study was conducted outside the laboratory to identify the potential of *A. platensis* growth and adaptation in the tropical rain forest climate with a variety of weather conditions.

In order for outdoor cultivation of *A. platensis* to be successful, the effect of weather conditions and nutrient requirement should be look into. Accordingly, management measures should be taken to ensure good microalgae cultivation is suitable for the environment in Malaysia in line with the development of the aquaculture industry. Since *A. platensis* growth is a function of environmental conditions such as light, temperature, and nutrient, it is necessary to study those interactions in Malaysia. In addition, to ensure the successful of *A. platensis* cultivation in large scale is dependent on effective development and economics culture. Study conducted by (Rodrigues *et al.*, 2010) found that over

25% of the total productivity of microalgae was related to the culture media that acts as a fertilizer to supply nutrients, particularly nitrogen and phosphorus.

Presently there is no cheap fertilizer available for commercial *A. platensis* cultivation in Malaysia. Therefore it is very important to develop cheap fertilizer from easily available chemicals to reduce production cost. There are several aspects that need to be improved in current fertilizer formula that will be used towards production of microalgae in a large scale in the future. Several factors have been studied throughout the study to compare the relationship between productivity and environmental factors change (Tomaselli *et al.*, 1997 and Prabuthas *et al.*, 2011). Hence the objectives of this study are:

1. To develop and improve the current fertilizer formula using cheap and effective nitrogen and phosphorus source for large scale production of microalgae especially in Malaysia
2. To determine the effect of different nitrogen and phosphorus source on *A. platensis* biochemical properties
3. To determine the effect of variable weather on *A. platensis* production and its biochemical composition.



## REFERENCES

- Abid, O., Sellami-Kammoun, A., Ayadi, H., Drira, Z., Bouain, A., & Aleya, L. (2008). Biochemical adaptation of phytoplankton to salinity and nutrient gradients in a coastal solar saltern, *Tunisia. Estuarine, Coastal And Shelf Science*. 391-400
- Ajayan, K.V. & Selvaraju, M. (2011). Reflector based chlorophyll production by *Spirulina platensis* through energy safe mode. *Bioresource Technology*. 7591–7594
- Ajayan, K.V., Selvaraju, M., & Thirugnanamoorthy, K. (2012). Enrichment of chlorophyll and phycobiliproteins in *Spirulina platensis* by the use of reflector light and nitrogen sources: An in-vitro study. *Biomass and Bioenergy*. 436-441
- Anaga, A. & Abu, G.O. (1996). A laboratory-scale cultivation of *Chlorella* and *Spirulina* using waste effluent from a fertilizer company in Nigeria. *Bioresource Technology*. 93–95
- AOAC International, & Latimer, G. W. (2012). Official methods of analysis of AOAC International. AOAC International.
- Avigad, V. & Richmond, A. (1988). Mass Production of the Blue-green Alga *Spirulina*: *Biomass*. 233-247
- Avila-Leon, M., Matsudo, C., Sato, S. & Carvalho, J.C.M. (2012). *Arthrospira platensis* biomass with high protein content cultivated in continuous process using urea as nitrogen source. *Journal of Applied Microbiology*. 1086–1094
- Becker, E. (2007). Micro-algae as a source of protein. *Biotechnology Advances*. 207-210
- Belay, A., Kato, T., & Ota, Y. (1996). *Spirulina* (*Arthrospira*): potential application as an animal feed supplement. *Journal of Applied Phycology*. 303-311
- Belkin, S. & Boussiba, S. (1991). High internal pH conveys ammonia resistance in *Spirulina platensis*. *Bioresource Technology*. 167-169
- Berberoglu, H., Gomez, P.S., & Pilon, L. (2009). Radiation characteristics of *Botryococcus braunii*, *Chlorococcum littorale*, and *Chlorella* sp. used for CO<sub>2</sub> fixation and biofuel production. *Journal of Quantitative and Spectroscopy*. 1879–93
- Bezerra, R.P., Matsudo, M.C., Converti, A., Sato, S., & Carvalho, J.C.M. (2008). Influence of Ammonium Chloride Feeding Time and Light Intensity on



the Cultivation of *Spirulina (Arthrospira) platensis*. *Biotechnology and Bioengineering*. 297–305

- Bezerra, R.P., Montoya, E.Y.O, Sato, S., Perego, P., Carvalho, J.C.M., & Converti, A. (2011). Effects of light intensity and dilution rate on semicontinuous cultivation of *Arthrospira (Spirulina) platensis*. A kinetic monod-type approach. *Bioresource Technology*. 3215-3219
- Binaghi, L., Del Borghi, A., Lodi, A., Converti, A., & Del Borghi, M. (2003). Batch and fed-batch uptake of carbon dioxide by *Spirulina platensis*. *Process Biochemistry*. 1341-1346
- Borowitzka, M. A. (1999). Commercial production of microalgae: ponds, tanks, tubes and fermenters. *Journal Of Biotechnology*. 313-321
- Borowitzka, M. A., Huisman, J. M., & Osborn, A. (1991). Culture of the astaxanthin-producing green alga *Haematococcus pluvialis* 1. Effects of nutrients on growth and cell type. *Journal of Applied Phycology*. 295-304
- Campo, J.A.D., González, M.G., & Guerrero, M.G. (2007). Outdoor cultivation of microalgae for carotenoid production: current state and perspectives. *Applied Microbiology and Biotechnology*. 1163-1174
- Carvalho, J.C.M., Francisco, F.R., Almeida, K.A., Sato S., & Converti A. (2004). Cultivation of *Arthrospira platensis* by fed batch addition of ammonium chloride a exponentially increasing feeding rates. *Journal Phycology*. 589-597
- Castenholz, R. W. (2001). General characteristics of the cyanobacteria. *Bergey's manual of systematic bacteriology*. 474-487
- Celekli, A., & Yavuzatmaca, M. (2009). Predictive modeling of biomass production by *Spirulina platensis* as function of nitrate and NaCl concentrations. *Bioresource Technology*. 1847-1851
- Chaiklahan, R., Chirasuwan, N., Loha, V., & Bunnag, B. (2008). Lipid and fatty acids extraction from the cyanobacterium *Spirulina*. *Science Asia*. 299-305
- Chang, Y., Wu, Z., Bian, L., Feng, D., & Leung, D.Y.C. (2013). Cultivation of *Spirulina platensis* for biomass production and nutrient removal from synthetic human urine. *Applied Energy*. 427-431
- Chen, W., Zhang, C., Song, L., Sommerfeld, M., & Hu, Q. (2009). A high throughput Nile red method for quantitative measurement of neutral lipids in microalgae. *Journal of Microbiological Methods*. 41-47
- Chojnacka, K. & Noworyta, A. (2004). Evaluation of *Spirulina* sp. growth in photoautotrophic, heterotrophic and mixotrophic cultures. *Enzyme and Microbial Technology*. 461- 465.

- Christopher, B. (2014). *Climate Change 2014: Impacts, Adaptation, and Vulnerability*. IPCC WGII AR5 Summary for Policymakers.
- Chronakis, I. S., Galatanu, A. N., Nylander, T., & Lindman, B. (2000). The behaviour of protein preparations from blue-green algae (*Spirulina platensis* strain Pacifica) at the air/water interface. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*. 181-192
- Cohen, E., Koren, A., & Arad, S.M. (1991). A closed system for outdoor cultivation of microalgae. *Biomass and Bioenergy*. 83-88
- Collet, P., Hélias, A., Lardon, L., Ras, M., Goy, R. A., & Steyer, J. P. (2011). Life-cycle assessment of microalgae culture coupled to biogas production. *Bioresource Technology*. 207-214
- Converti, A., Scapazzoni, S., Lodi, A., & Carvalho, J.C.M. (2006). Ammonia and urea removal by *Spirulina platensis*. *Journal Industry Microbial Biotechnology*. 8-16
- Converti, A., Lodi, A., Del Borghi, A., & Solisio, C. (2006). Cultivation of *Spirulina platensis* in a combined airlift-tubular reactor system. *Biochemical Engineering Journal*. 13-18
- Costa, J. A. V., Colla, L. M., Duarte Filho, P., Kabke, K., & Weber, A. (2002). Modelling of *Spirulina platensis* growth in fresh water using response surface methodology. *World Journal of Microbiology and Biotechnology*. 603-607
- Danesi, E.D.G., Rangel, C.O., de Carvalho, J.C.M., & Sato, S. (2002). An investigation of effect of replacing nitrate by urea in the growth and production of chlorophyll by *Spirulina platensis*. *Biomass and Bioenergy*. 261-269
- Day, J.G., Slocombe, S.P., & Stanley, M.S. (2012). Overcoming biological constraints to enable the exploitation of microalgae for biofuels. *Bioresource Technology*. 245-251
- Doucha, J., Straka, F., & Lívansky, K. (2005). Utilization of flue gas for cultivation of microalgae *Chlorella* sp. in an outdoor open thin-layer
- Evans, R. W., Kates, M., Ginzburg, M., & Ginzburg, B. Z. (1982). Lipid composition of halotolerant algae, *Dunaliella parva* Lerche and *Dunaliella tertiolecta*. *Biochimica et Biophysica Acta (BBA)-Lipids and Lipid Metabolism*. 186-195
- Elsayed, E. A., & Boucher, T. O. (1994). Analysis and control of production systems. Prentice Hall
- Ferreira, L.S., Rodrigues, M.S., Converti, A., & Carvalho, J.C.M. (2012). *Arthrospira platensis* cultivation in tubular photobioreactor: use on no-cost CO<sub>2</sub> from ethanol fermentation. *Applied Energy*. 379-385

- Food and Agriculture Organization of the United Nations (FAO). (2008). The state of food and agriculture 2008. Viale delle Terme di Caracalla, 00153 Rome, Italy: Food and Agriculture Organization of the United Nations
- Food and Agriculture Organization of the United Nations (FAO). (2001). The State of Food Insecurity in the World 2001. Viale delle Terme di Caracalla, 00100 Rome, Italy: Food and Agriculture Organization of the United Nations
- Gad, A. S., Khadrawy, Y. A., El-Nekeety, A. A., Mohamed, S. R., Hassan, N. S., & Abdel-Wahhab, M. A. (2011). Antioxidant activity and hepatoprotective effects of whey protein and Spirulina in rats. *Nutrition*. 582-589
- General Climate of Malaysia*. (MMD) (n.d.). Retrieved February 16, 2014, from Official Portal Malaysian Meteorological Department: [http://www.met.gov.my/index.php?option=com\\_content&task=view&id=75&Itemid=1089&limit=1&limitstart=0](http://www.met.gov.my/index.php?option=com_content&task=view&id=75&Itemid=1089&limit=1&limitstart=0)
- Habib, M.A.B.; Parvin, M.; Huntington, T.C., & Hasan, M.R. (2008). *A review on culture, production and use of spirulina as food for humans and feeds for domestic animals and fish*. Rome: FAO Fisheries and Aquaculture Circular.
- Hasan, M.R. & Chakrabarti, R. (2009). *Use of algae and aquatic macrophytes as feed in small-scale aquaculture: a review*. Rome: FAO Fisheries and Aquaculture Technical Paper
- Hase, R., Oikawa, H., Sasso, C., Morito, M. & Watabe, Y. (2000). Photosynthetic production of microalgal biomass in a race way system under greenhouse conditions in Sendi City. *Journal of Bioscience Bioengineering*. 157–163
- Henrikson, R. (1989). Earth food Spirulina. Laguna Beach, CA: Ronore Enterprises, Inc.
- Hernandez, J.P., De-Bashan, L.E., Rodriguez, D.J., Rodriguez, Y., and Bashan, Y. (2009). Growth promotion of the freshwater microalga *Chlorella vulgaris* by the nitrogen-fixing, plant growth-promoting bacterium *Bacillus pumilus* from arid zone soils. *European Journal of Soil Biology*. 88-93
- Ishikawa, T., Fujiyama, Y., Igarashi, O., Morino, M., Tada, N., Kagami, A & Nakamura, H. (1989). Effects of gammalinolenic acid on plasma lipoproteins and apolipoproteins. *Atherosclerosis*. 95-104
- Isichei, A. O. (1990). The role of algae and cyanobacteria in arid lands. A review. *Arid Soil Research and Rehabilitation*. 1-17

- Jeong, M.J., Gillis, J.M., & Hwang, J.Y. (2003). Carbon dioxide mitigation by microalgal photosynthesis. *Chemical Society*. 1763–1766
- Jiang, L., Luo, S., Fan, X., Yang, Z. & Guo, R. (2011). Biomass and lipid production of marine microalgae using municipal wastewater and high concentration of CO<sub>2</sub>. *Applied Energy*. 3336-3341
- Jimenez, C., Cossi, B. R., & Niell, F. X. (2003). Relationship between physicochemical variables and productivity in open ponds for the production of *Spirulina*: a predictive model of algal yield. *Aquaculture*. 331-345
- Jorquera, O., Kiperstok, A., Sales, E. A., Embiruçu, M., & Ghirardi, M. L. (2010). Comparative energy life-cycle analyses of microalgal biomass production in open ponds and photobioreactors. *Bioresource Technology*. 1406-1413
- Kadir, M.Z.A.A & Rafeeu, Y. (2010). A review on factors for maximizing solar fraction under wet climate environment in Malaysia. *Renewable and Sustainable Energy Reviews*. 2243–2248
- Katarzyna, C. & Noworyta, A. (2004). Evaluation of *Spirulina* sp. growth in photoautotrophic, heterotrophic and mixotrophic cultures. *Enzyme and Microbial Technology*. 461–465
- Komarek, J. & Kastovsky, J. (2003). Coincidences of structural and molecular characters in evolutionary lines of cyanobacteria. *Algological Studies*. 305-325
- Kosaric, N., Nguyen, H.T. & M. A. Bergougnou. (1974). Growth of *Spirulina maxima* algae in effluents from secondary waste-water treatment plants. *Biotechnology and Bioengineering*. 881–896
- Lee, Y. (2001). Microalgal mass culture systems and methods: Their limitation and potential. *Journal of Applied Phycology*. 307-315
- Lichtenthaler, H.K. (1987). Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. *Methodology Enzymology*. 350-382
- Liz, K. (2014, March 31). 'We're All Sitting Ducks': Latest U.N. Global Warming Report Paints Dark Picture. Retrieved from <http://www.theblaze.com/stories/2014/03/31/were-all-sitting-ducks-latest-u-n-global-warming-report-paints-dark-picture/>
- Madkour, F.F., Kamil, A.W., & Nasr, H.S. (2012). Production and nutritive value of *Spirulina platensis* in reduced cost media. *Egyptian Journal of Aquatic Research*. 51-57
- Mary Leema, J.T., Kirubakaran, R., Vinithkumar, N.V., Dheenan, P.S., & Karthikayulu, S. (2010). High value pigment production from

*Arthrospira platensis* cultured in seawater. *Bioresource Technology*. 9221-9227

- Markou, G., Chatzipavlidis, I., & Georgakakis, D. (2012). Effects of phosphorus concentration and light intensity on the biomass composition of *Arthrospira (Spirulina) platensis*. *World Journal Microbiology Biotechnology*. 2661-2670
- Matsudo, M.C., Bezerra, R.P., Sato, S., Perego, P., Converi, A., & Carvalho, J.C.M. (2009). Repeated fed-batch cultivation of *Arthrospira (Spirulina) platensis* using urea as nitrogen source. *Biochemical Engineering Journal*. 52-57
- Matsudo, M.C., Bezerra, R.P., Converi, A., Sato, S., & Carvalho, J.C.M. (2011). Carbon dioxide from alcoholic fermentation for continuous cultivation of *Arthrospira (Spirulina) platensis* in tubular photobioreactor using urea as nitrogen source. *Bioechnology Progress*. 650-656
- Molitor, V., Erber, W., & Peschek, G. A. (1986). Increased levels of cytochrome oxidase and sodium-proton antiporter in the plasma membrane of *Anacystis nidulans* after growth in sodium-enriched media. *FEBS letters*. 251-256
- Mosulishvili, L. M., Frontasyeva, M. V., Pavlov, S. S., Belokobylsky, A. I., Kirkesali, E. I. & Khizanishvili, A. I. (2004). Epithermal neutron activation analysis of *Spirulina platensis* biomass and extracted C-phycocyanin and DNA. *Journal Of Radioanalytical And Nuclear Chemistry*. 41-45
- Nakamura, M. T., & Nara, T. Y. (2003). Essential fatty acid synthesis and its regulation in mammals. *Prostaglandins, Leukotrienes And Essential Fatty Acids*. 145-150
- Ogbonda, K.H., Aminigo, R.E., & Abu, G.O. (2007). Influence of temperature and pH on biomass production and protein biosynthesis in a putative *Spirulina sp.* *Bioresource Technology*. 2207-2211
- Olguin, E. J., Galicia, S., Mercado, G., & Perez, T. (2003). Annual productivity of *Spirulina (Arthrospira)* and nutrient removal in a pig wastewater recycling process under tropical conditions. *Journal of Applied Phycology*. 249-257
- Oliveira, M.A.C.L.D, Monteiro, M.P.C., Robbs, P.G. & Leite, S.G.F. (1999). Growth and chemical composition of *Spirulina maxima* and *Spirulina platensis* biomass at different temperatures. *Aquaculture International*. 261-275
- Parmar, A., Singh, N.K., Pandey, A., Gnansounou, E., & Madamwar, D. (2011). Cyanobacteria and microalgae: A positive prospect for biofuels. *Bioresource Technology*. 10163-10172



- Parsons, T. R., & Strickland, J. D. (1963). Discussion of spectrophotometric determination of marine-plant pigments, with revised equations for ascertaining chlorophylls and carotenoids. *Journal of Marine Research*. 155-163
- Peter, H. P., Vincent, A.B. & Richard, N. (2011). Sustainability of algae derived biodiesel: A mass balance approach. *Bioresource Technology*. 1185–1193
- Pfromm, P.H., Amanor-Boadu, V., & Nelson, R. (2011). Sustainability of algae derived biodiesel: a mass balance approach. *Bioresource Technology*. 1185-1193
- Phang, S.M., Miah, M.S., Yeoh, B.G. & Hashim, M.A. (2000). Spirulina cultivation in digested sago starch factory wastewater. *Journal of Applied Phycology*. 395-400
- Piorreck, M., Baasch, K. H., & Pohl, P. (1984). Biomass production, total protein, chlorophylls, lipids and fatty acids of freshwater green and blue-green algae under different nitrogen regimes. *Phytochemistry*. 207-216
- Prabuthas, P., Srivastav, P.P. & Mishra, H.N. (2011). Optimization of environmental factors using RSM for *Spirulina platensis* cultivation. *Nutrition & Food Science*. 175 – 182
- Pruvost, J., Van Vooren, G., Le Gouic, B., Couzinet-Mossion, A., & Legrand, J. (2011). Systematic investigation of biomass and lipid productivity by microalgae in photobioreactors for biodiesel application. *Bioresource Technology*. 150-158
- Puganeswary, S., Rosimah, N., Sharizim, Z., Normala, H., Hishamuddin, O., & Ahmad, I. (2014). Potential of fresh POME as a growth medium in a mass production of *Arthrospira platensis*. *International Journal of Current Microbiology and Applied Sciences*. 235-253
- Rafiqul, I.M., Jalal, K.C.A., & Alam, M.Z., (2005). Environmental factors for optimization of *Spirulina* biomass in laboratory culture. *Biotechnology*. 19-22
- Raja, R., Hemaiswarya, S., Kumar, N.A., Sridhar, S., & Rengasamy, R. (2008). A perspective on the biotechnological potential of microalgae. *Critical Reviews in Biotechnology*. 77-88
- Raoof, B., Kaushik, B. D. & Prasanna, R. (2006). Formulation of a low-cost medium for mass production of *Spirulina*. *Biomass and Bioenergy*. 537-542
- Rangel, C.O., Danesi, E.D.G., Carvalho, J.C.M. & Sato, S. (2004). Chlorophyll production from *Spirulina platensis*: cultivation with urea addition by fed-batch process. *Bioresource Technology*. 133-141

- Ravelonandrob, P.H., Ratianarivoa, D.H., & Cassand, C.J. (2011). Improvement of the growth of *Arthrospira (Spirulina) platensis* from Toliara (Madagascar): Effect of agitation, salinity and CO<sub>2</sub> addition. *Food And Bioproducts Processing*. 209–216
- Richmond, A. & Qiang, H. (1997). Principles for efficient utilization of light for mass production of photoautotrophic microorganisms. *Applied Biochemistry And Biotechnology*. 649-658
- Richmond, A., Lichtenberg, E., Stahl, B. & Vonshak, A. (1990). Quantitative assessment of the major limitations on productivity of *Spirulina platensis* in open raceways. *Journal of Applied Phycology*. 195–206
- Rodolfi, L., Chini Zittelli, G., Bassi, N., Padovani, G., Biondi, N., Bonini, G., & Tredici, M. R. (2009). Microalgae for oil: Strain selection, induction of lipid synthesis and outdoor mass cultivation in a low-cost photobioreactor. *Biotechnology and Bioengineering*. 100-112
- Rodrigues, M. S., Ferreira, L. S., Converti, A., Sato, S. & Carvalho, J. C. M. (2010). Fed-batch cultivation of *Arthrospira (Spirulina) platensis*: potassium nitrate and ammonium chloride as simultaneous nitrogen sources. *Bioresource Technology*. 4491-4498
- Rodrigues, M.S., Ferreira, L.S., Converti, A., Sato S., & Carvalho J.C.M. (2011). Influence of ammonium sulphate feeding time on fed-batch *Arthrospira platensis* cultivation and biomass composition with and without pH control. *Bioresource Technology*. 6587-6592
- Sajilata, M. G., Singhal, R. S., & Kamat, M. Y. (2008). Fractionation of lipids and purification of  $\gamma$ -linolenic acid (GLA) from *Spirulina platensis*. *Food Chemistry*. 580-586
- Sanchez-Luna, L. D., Converti, A., Tonini, G. C., Sato, S., & De Carvalho, J. C. (2004). Continuous and pulse feedings of urea as a nitrogen source in fed-batch cultivation of *Spirulina platensis*. *Aquacultural Engineering*. 237-245
- Selmi, C., Leung, P. S., Fischer, L., German, B., Yang, C. Y., Kenny, T. P. & Gershwin, M. E. (2011). The effects of *Spirulina* on anemia and immune function in senior citizens. *Cellular & Molecular Immunology*. 248-254
- Sharma, M. K., Sharma, A., Kumar, A., & Kumar, M. (2007). *Spirulina fusiformis* provides protection against mercuric chloride induced oxidative stress in Swiss albino mice. *Food and Chemical Toxicology*. 2412-2419
- Shen, Z., Zhou, J., Zhou, X., & Zhang, Y. (2011). The production of acetic acid from microalgae under hydrothermal conditions. *Applied Energy*. 3444-3447

- Simsek, N., Karadeniz, A., Kalkan, Y., Keles, O. N., & Unal, B. (2009). *Spirulina platensis* feeding inhibited the anemia-and leucopenia-induced lead and cadmium in rats. *Journal of Hazardous Materials*. 1304-1309
- Singh, R.K. & Kundu, S. (2014). Review on Changing Natural Nitrogen Cycle: Special Reference to Kingdom of Saudi Arabia. *International Journal of Engineering Science Invention Research & Development*. 1-9
- Soletto, D., Binaghi, L., Lodi, A., Carvalho, J.C.M., & Converti, A. (2005). Batch and fed-batch cultivations of *Spirulina platensis* using ammonium sulphate and urea as nitrogen sources. *Aquaculture*. 217-224
- Soletto, D., Binaghi, L., Ferrari, L., Lodi, A., Carvalho, J.C.M., Zilli, M., & Converti, A. (2008). Effects of carbon dioxide feeding rate and light intensity on the fed-batch pulse-feeding cultivation of *Spirulina platensis* in helical photobioreactor. *Biochemical Engineering Journal*. 369-375
- Spolaore, P., Joannis-Cassan, C., Duran, E., & Isambert, A. (2006). Commercial application of microalgae. *Journal of Bioscience and Bioengineering*. 87-96
- Stewart, K. (2009). Physical Properties of Water. *Reference Module in Earth Systems and Environmental Sciences, Encyclopedia of Inland Waters*. 148–154
- Swanson, D., Block, R., & Mousa, S. A. (2012). Omega-3 fatty acids EPA and DHA: health benefits throughout life. *Advances in Nutrition: An International Review Journal*. 1-7
- Tomaselli, L., Boldrini, G., & Margheri, M.C. . (1997). Physiological behaviour of *Arthrospira (Spirulina) maxima* during acclimation to changes in irradiance. *Journal of Applied Phycology*. 37-43
- Tompkins, J.M.M., Day, J.G., & Turner M.F. (1995). Culture collection of algae and protozoa. In: De Ville, (Eds.), *Catalogue of Strains: Natural Environment Research Council*. Titus Wilson and Sons Ltd., Kendal, U.K. pp. 144-173
- Toyub, M.A., Uddin, M.Z., Miah, M.I, & Habib, M.A.B. (2011). Growth performance and nutritional analysis of *Spirulina platensis* growth performance and nutritional analysis of *Spirulina platensis* in different concentrations of papaya skin powder media . *Bangladesh Journal Of Scientific And Industrial Research*. 333-338
- Uday, B.S. & Ahluwalia, A. S. (2013). Microalgae: a promising tool for carbon sequestration. *Mitigation and Adaptation Strategies for Global Change* . 73–95
- Ugwu, C. U., Aoyagi, H., & Uchiyama, H. (2008). Photobioreactors for mass cultivation of algae. *Bioresource Technology*. 4021-4028



- United Nations(UNSCN). (November 29th - December 10th, 2010). Climate Change and Nutrition Security. *16th United Nations Conference of the Parties (COP16)* (pp. United Nations System, Standing Committee on Nutrition). Cancun: World Health Organization
- Usmanova, A., Astier, C., Mejean, C., Hubert, F., Feinberg, J., Benyamin, Y., & Roustan, C. (1998). Coevolution of actin and associated proteins: an  $\alpha$ -actinin-like protein in a cyanobacterium (*Spirulina platensis*). *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*. 693-700
- Vardon, D.R., Sharma, B.K., Scott, J., Yu, G., Wang, Z., Schideman, L., Zhang, Y., & Strathmann, T.J. (2011). Chemical properties of biocrude oil from the hydrothermal liquefaction of *Spirulina* algae, swine manure, and digested anaerobic sludge. *Bioresource Technology*. 8295-8303
- Vonshak, A. (1997). *Spirulina platensis (Arthrospira)*. In A. Vonshak, *Physiology, Cell Biology and Biotechnology* (pp. 213-226). London: Taylor and Francis
- Vonshak, A. & Richmond, A. (1988). Mass production of the blue-green alga *Spirulina*: an overview. *Biomass*. 233-247
- Vonshak, A., Abeliovich, A., Boussiba, S., Arad, S., & Richmond, A. (1982). Production of *Spirulina* biomass: effects of environmental factors and population density. *Biomass*. 175-185
- Wang, B., Li, Y.Q., Wu, N., & Lan, C.Q. (2008). CO<sub>2</sub> bio-mitigation using microalgae. *Applied Microbiology Biotechnology*, 707–180
- World Food Programme (WFP). (2013). *Hunger statistics*. Retrieved April 28, 2014, from World Food Programme, fighting hunger worldwide: <http://www.wfp.org/hunger/stats>
- Xin, L., Hongying, Hu., Ke, G., & Yingxue, S. (2010). Effects of different nitrogen and phosphorus concentrations on the growth, nutrient uptake, and lipid accumulation of a freshwater microalgae *Scenedesmus sp.* *Bioresource Technology*. 5494-5500
- Xue, S., Su, Z., & Cong, W. (2011). Growth of *Spirulina platensis* enhanced under intermittent illumination. *Journal of Biotechnology*. 271-277
- Yang, C., Liu, H., Li, M., & Yu, C. (2008). Treating urine by *Spirulina platensis*. *Acta Astronautica*. 1049-1054
- Zeng, M. T., & Vonshak, A. (1998). Adaptation of *Spirulina platensis* to salinity-stress. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 113-118

Zhang, H. Q., Lin, A. P., Sun, Y., & Deng, Y. M. (2001). Chemoand radioprotective effects of polysaccharide of *Spirulina platensis* on hemopoietic system of mice and dogs. *Acta Pharmacologica Sinica*. 1121-1124.



## PUBLICATION

Mashor, N., Yazam, M.S.M., Naquiuddin, M.A., Omar, H., & Ismail, A. (2016). Different nitrogen sources effects on the growth and productivity of *Spirulina* grown in outdoor conditions. *Acta Biologica Malaysiana*. 16-26





**UNIVERSITI PUTRA MALAYSIA**

**STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT**

**ACADEMIC SESSION :** \_\_\_\_\_

**TITLE OF THESIS / PROJECT REPORT :**

MODIFICATION OF KOSARIC'S MEDIUM WITH NITROGEN AND PHOSPHORUS FOR  
Spirulina (*Arthrospira platensis*) IN SHELTERED OPEN CULTURE SYSTEM

**NAME OF STUDENT :** NURSHAZREEN BINTI MASHOR

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

\*Please tick (✓)

☐

**CONFIDENTIAL**

(Contain confidential information under Official Secret Act 1972).

☐

**RESTRICTED**

(Contains restricted information as specified by the organization/institution where research was done).

☐

**OPEN ACCESS**

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

☐

**PATENT**

Embargo from \_\_\_\_\_ until \_\_\_\_\_  
(date) (date)

**Approved by:**

\_\_\_\_\_  
(Signature of Student)  
New IC No/ Passport No.:

Date :

\_\_\_\_\_  
(Signature of Chairman of Supervisory Committee)  
Name:

Date :

[Note : If the thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach with the letter from the organization/institution with period and reasons for confidentiality or restricted. ]