



UNIVERSITI PUTRA MALAYSIA

***IMPOSITION OF BENEFICIAL WATER STRESS FOR IMPROVEMENT OF
POSTHARVEST QUALITY OF LOWLAND TOMATO FRUITS
(*Lycopersicon esculentum* Mill.)***

MOHAMMED HASSAN NAMA

FP 2018 9



**IMPOSITION OF BENEFICIAL WATER STRESS FOR IMPROVEMENT
OF POSTHARVEST QUALITY OF LOWLAND TOMATO FRUITS
(*Lycopersicon esculentum* Mill.)**

By

MOHAMMED HASSAN NAMA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Master of Science**

January 2018

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**IMPOSITION OF BENEFICIAL WATER STRESS FOR IMPROVEMENT
OF POSTHARVEST QUALITY OF LOWLAND TOMATO FRUIT
(*Lycopersicon esculentum* Mill.)**

By

MOHAMMED HASSAN NAMA

January 2018

Chairman : Mahmud Tengku Muda Mohamed, PhD
Faculty : Agriculture

Water stress affects crop performance by influencing nutrient availability and crop functionality. There is a lack of information on utilization of deficit irrigation strategies in manipulating the growth rates, yield and quality of low land tomato plant in Malaysia. Present study was indicated to investigate the effects of different level of water stress on plant growth, yield and postharvest qualities. Two greenhouse experiments were conducted at Field 15, Faculty of Agriculture, Universiti Putra Malaysia. Experiment 1 was conducted to describe the effect of limiting strategically the water supply during plant development on plant growth, yield and postharvest quality, with the aim of identifying the best deficit schedule for the plants under low land tropical conditions. Three- week old MT1 tomato seedlings from the trays were transplanted into polybags filled with mixture of coco peat and paddy husk (2:1 V/V). After 40 days, the seedlings were treated with T1 control (daily watering to field capacity), T2 (restoring water supply to field capacity every two days) and T3 (restoring water supply to field capacity every four days). At harvest, plant height, leaves number, stem diameter, leaf area, dry shoot and root fresh and dry weight, fruit weight and fruit number were measured. In addition, data were collected on the following fruit quality parameters: firmness, total soluble solids, titratable acidity, pH, ascorbic acid and lycopene. The deficit irrigation applications increased soluble solids concentration significantly at T3 in the first experiment and at T4 (fruiting growth stage) in the second experiment. However, the rates of increment was not significantly different in titratable acidity, ascorbic acid, lycopene content, pH and firmness. The T2 (restoring water supply to field capacity every two days) promoted total fruit weight. Water stress treatments decreased plant heights, leaves number, leaf area, trusses number and both fresh and dry shoot and root weight. The leaf relative water content was reduced by 22.2% in the most stressed plants T3 (four days deficit irrigation) compared to the control. Whilst experiment 2 was conducted to identify the

most critical phenological (plant growth stages) and fruit maturity stages to impose deficit irrigation and their effects on growth, yield and postharvest quality of tomato. Three weeks old seedlings were transplanted into polybags. Then, the seedlings were exposed to four water stress treatments: T1 (control), T2 (deficit irrigation every four days at the vegetative stage), T3 (deficit irrigation every four days at the flowering stage) and T4 (deficit irrigation every four days at the fruiting stage). All growth, yield and postharvest parameters were determined as in Experiment 1. The plants that subjected to deficit irrigation levels produced similar plant height and number of leaves to the control plants (full irrigation). However, fruit weight and number of fruit increased significantly under T3 (flowering stage) but not significantly different from those in control plants. In addition, water deficit irrigation at T4 (deficit imposed during fruiting stage) significantly reduced fresh and dry weight of shoots and root compared to the other treated and control plants. In conclusion, the optimum yield of tomato could be obtained at T3 (deficit imposed at flowering stage). The vegetative and flowering growth stages could be considered as the most tolerant to deficit irrigation, and the fruiting growth stage could be considered the most critical stage. Imposition water stress at flowering growth stage on tomato produced better plants condition, while using the water stress during fruiting stage retards plant growth by decreased plant growth and rate of yield. Fruits quality such as fruit firmness, pH of fruit, SSC, TA, AA and lycopene were affected significantly by deficit irrigation treatments. T4 (fruiting stage) fruits had the highest SSC in those harvested at the turning maturity fruit stage; high SSC improves both paste yield per unit of fresh fruit and overall processing efficiency. While the highest lycopene content was observed with those treated during vegetative stage harvested at the red maturity fruit stage. On the other hand, fruits harvested from T2 (vegetative stage) plants at turning maturity fruit stage gave the highest firmness. Highest pH fruit was obtained with T3 plants at red fruit maturity index. However, the results demonstrated that different deficit irrigation regimes did not affect TA and AA contents of tomato fruits, indicating that the results from of this study can enhance and maintaining post-harvest quality, also deficit irrigation strategy can help in the development of water management system for tomato production in the scenario of reduced water availability and enable the tomato growers to produce tomato with optimum yield by allowing little water stress without substantial yield reduction.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGENAAN TEGASAN AIR YANG BERMANFAAT BAGI
PENAMBAHBAIKAN KUALITI PASCA TUAI BUAH TOMATO TANAH
RENDAH (*Lycopersicon esculentum* Mill.)**

Oleh

MOHAMMED HASSAN NAMA

Januari 2018

Pengerusi : Profesor Mahmud Tengku Muda Mohamed, PhD
Fakulti : Pertanian

Tegasan air memberi kesan terhadap prestasi tanaman dengan mempengaruhi ketersediaan nutrien dan fungsi tanaman. Terdapat kekurangan maklumat mengenai penggunaan sistem pengairan defisit dalam memanipulasikan kadar pertumbuhan, hasil tuai dan kualiti tanaman tomato tanah rendah di Malaysia. Kajian ini dijalankan bagi mengkaji kesan perbezaan tahap ketegasan air terhadap pertumbuhan pokok, hasil dan kualiti pasca tuai. Dua kajian dalam rumah hijau telah dijalankan di ladang 15, Fakulti Pertanian, Universiti Putra Malaysia. Eksperimen 1 telah dijalankan untuk menerangkan kesan mengehadkan bekalan air secara strategik semasa pertumbuhan tanaman terhadap tumbesaran, hasil tuai dan kualiti pasca tuai tanaman, dengan sasaran untuk mengenalpasti jadual defisit air terbaik untuk tanaman pada kondisi tanah rendah tropika. Anak benih tomato MT1 yang berusia tiga minggu telah dipindahkan dari dulang semaian ke dalam polibeg berisi sabut kelapa dan sekam padi (2:1 V/V). Selepas 40 hari, anak benih dirawat dengan T1, kawalan (pengairan harian ke kapasiti lapangan), T2 (memulihkan air kepada kapasiti lapangan setiap dua hari) dan T3 (memulihkan air kepada kapasiti lapangan setiap empat hari). Pada peringkat penuaian, tinggi pokok, bilangan daun, diameter batang, luas daun, berat segar dan kering pucuk dan akar, berat buah dan bilangan buah diukur. Di samping itu, data parameter kualiti buah: ketegasan, jumlah pepejal terlarut, asid tertitrat, pH, asid askorbik dan kandungan lycopene turut dikumpulkan. Pengairan defisit pada T3 dalam eksperimen pertama dan T4 (peringkat pemuahan) dalam eksperimen kedua meningkatkan kepekatan pepejal terlarut secara bererti, tetapi kadar peningkatan asid tertitrat, asid askorbik, kandungan lycopene, pH dan ketegasan adalah tidak berbeza bererti di antara rawatan. T2 (pemulihan sumber air kepada kapasiti lapangan setiap dua hari) menggalakkan jumlah berat buah. Rawatan tegasan air mengurangkan ketinggian tumbuhan, bilangan daun, luas daun, bilangan tangkai dan kedua-dua berat segar dan kering pucuk dan akar. Kandungan relatif air pada daun berkurang kepada

22.2% pada tanaman paling tinggi tegasan, T3 (pengairan defisit setiap empat hari) berbanding rawatan kawalan. Manakala eksperimen 2 telah dijalankan untuk mengenalpasti peringkat fonologikal dan kematangan buah paling kritikal untuk mengenakan pengairan defisit dan kesannya terhadap tumbesaran, hasil dan kualiti pasca tuai tomato. Anak benih berusia tiga minggu dipindahkan ke dalam polibeg. Kemudian, anak benih ini didedahkan kepada empat rawatan tegasan air: T1 (kawalan), T2 (pengairan defisit setiap empat hari pada peringkat pertumbuhan vegetatif), T3 (pengairan defisit setiap empat hari pada peringkat pembungaan) dan T4 (pengairan defisit setiap empat hari pada peringkat pemuahan). Semua parameter tumbesaran, termasuklah hasil dan pasca tuai telah dikenalpasti seperti dalam Eksperimen 1. Tanaman dengan tegasan air menunjukkan tinggi pokok dan bilangan daun yang sama dengan tanaman kawalan (pengairan penuh). Walau bagaimanapun, berat dan bilangan buah meningkat secara bererti pada T3 (peringkat pembungaan) tetapi tidak bererti berbanding tanaman kawalan. Di samping itu, pengairan defisit air mengurangkan berat segar dan kering pucuk dan akar pada T4 (defisit pada peringkat pemuahan) berbanding tanaman kawalan dan rawatan lain. Kesimpulannya, hasil tuai tomato yang optimum boleh dicapai pada T3 (pengairan defisit pada peringkat pembungaan). Peringkat tumbesaran vegetatif dan pembungaan boleh dianggap sebagai peringkat paling toleran kepada pengairan defisit air, dan peringkat tumbesaran buah boleh dianggap sebagai peringkat paling kritikal. Pengenaan tegasan air pada peringkat pembungaan tomato menghasilkan tanaman yang baik, manakala pengenaan tegasan air pada peringkat pemuahan merencat pertumbuhan pokok dengan mengurangkan tumbesaran pokok dan kadar hasil tuai. Kualiti buah-buahan seperti ketegasan buah, pH, kepekatan pepejal terlarut, asid tertitrat, asid askorbik dan lycopene terjejas dengan ketara terhadap rawatan pengairan defisit. Buah-buahan T4 (peringkat pemuahan) mengandungi kepekatan pepejal terlarut tertinggi berbanding hasil lain yang dituai pada peringkat pertukaran kematangan buah; tinggi kepekatan pepejal terlarut meningkatkan kedua-dua hasil pes per unit buah segar dan kecekapan pemprosesan keseluruhan. Manakala kandungan lycopene tertinggi pada hasil tuai dengan rawatan di peringkat vegetatif dituai pada peringkat kematangan buah merah. Sebaliknya, buah-buahan yang dituai dari tanaman T2 (peringkat vegetatif) memberikan kepejalan buah tertinggi pada peringkat pertukaran kematangan. Selain itu, pH buah tertinggi telah dicapai melalui tanaman T3 pada indeks kematangan buah merah. Walau bagaimanapun, hasil kajian menunjukkan perbezaan rejim pengairan defisit tidak menjejaskan kandungan asid tertitrat dan asid askorbik buah-buahan tomato, menunjukkan bahawa keputusan dari kajian ini dapat menambah dan mengekalkan kualiti pasca tuai, strategi pengairan defisit juga dapat membantu dalam pembangunan sisem pengurusan air untuk pengeluaran tomato dalam senario pengurangan sumber air dan membolehkan pengusaha tomato untuk menghasilkan tomato dengan hasil yang optimum dengan membenarkan sedikit ketegasan air tanpa mengalami pengurangan hasil.

ACKNOWLEDGEMENTS

In the Name of Allah, the most Beneficent and the most Compassionate

First and foremost, my gratitude is to Allah SWT for giving me the strength to complete this thesis. I wish to express my deepest gratitude to Professor Dr. Mahmud T. Muda Mohamed, the chairman of my supervisory committee and Dr. Puteri Edaroyati Megat Wahab, the member of my supervisory committee, for their invaluable advice, guidance, patience and encouragement throughout the research and the completion of this thesis.

My special appreciation and gratitude goes to Universiti Putra Malaysia for their support. I also would like to express my sincere gratitude to all lecturers and the staff in the Department of Crop Science, Faculty of Agriculture, Universiti Putra Malaysia, especially the Postharvest Laboratory staff, Mr. Azahar, Mr. Yusoff, Mr. Mazlan, and Mr. Yaacob and all the staff in greenhouse for their kindness and help in the project and my study. Furthermore, my deepest thanks to all my friends; Omar Ali, Qusay A. Alhamza, Stev Agada and Mr. Fared for the help and unforgettable moments.

My special appreciation and gratitude goes to my beloved father and mother, I pray to Allah to bless my father who guided me to achieve the highest level of education. And also wish to express my sincere gratitude to my brother, Islam, and my uncles, Dr. Samee, Dr. Hussam and Dhiya, and my cousin Mohammed for their constant encouragement, love and ever ready to give helping hands.

Completion of this research work was also done by the continuous encouragement of my wife and my beloved sons, Ali and Hussein, and my beloved daughter, Zahraa who provided the comfort and for being the constant source of inspiration for me to complete this research.

Last but not least, I would like to express my deepest thanks to the Malaysian government and people represented by Universiti Putra Malaysia, for giving me this opportunity to study in their prestigious and reputed college.

I certify that a Thesis Examination Committee has met on 4 January 2018 to conduct the final examination of Mohammed Hassan Nama on his thesis entitled "Imposition of Beneficial Water Stress for Improvement of Postharvest Quality of Lowland Tomato Fruits (*Lycopersicon esculentum* Mill.)" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Dato' Mohd Fauzi bin Hj. Ramlan, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Phebe Ding, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Normaniza Osman, PhD

Associate Professor
University of Malaya
Malaysia
(External Examiner)



NOR AINI AB. SHUKOR, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 29 January 2018

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Mahmud Tengku Muda Mohamed, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Puteri Edaroyati Megat Wahab, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____

Date: _____

Name and Matric No: Mohammed Hassan Nama GS42237

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Professor Dr. Mahmud Tengku Muda Mohamed

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Puteri Edaroyati Megat Wahab

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xvi
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 Tomato	4
2.1.1 Tomato Growth Conditions	4
2.1.2 Tomato Growth and Fruit Maturity Stages	4
2.2 Global Water Scarcity and Crop Production	5
2.3 Plant Response to Water Stress	7
2.4 Deficit Irrigation and the Improvement of Water Use Efficiency	7
2.5 Beneficial Effect of Deficit Irrigation Strategies	8
2.5.1 Effect of Deficit Irrigation on Crop Growth and Development	9
2.5.2 Effect of Deficit Irrigation on the Physico-Chemical Qualities of Fruits	10
2.5.3 Deficit Irrigation and The Maintenance of Fruit Quality After Harvest	12
2.5.4 Effect of Deficit Irrigation Timing and Fruit Quality	13
2.5.5 Effect of Deficit Irrigation on Different Fruit Maturity Stage	15
3 GENERAL MATERIALS AND METHODS	16
3.1 Glasshouse Experiments	16
3.2 Growing Condition	16
3.3 Preparation of Plant Materials	16
3.4 Imposition of Water Stress Treatments and Design	17
3.5 Measurement of Soil Water Content	17
3.6 Measurement of Leaf Water Potential	17
3.7 Plant Growth Parameter	18
3.7.1 Plant height, stem diameter (mm) and leaves number	18
3.7.2 Fresh and dry weight of shoots and roots (g) and root: shoots ratio	18
3.7.3 Determination of Fruit Number and Fruit Weight	18
3.7.3.1 Total Fruit	18

	3.7.3.2 Marketable Fruit	19
	3.7.3.3 Unmarketable Fruit	19
3.8	Physico-Chemical Characteristics of Fruits	19
	3.8.1 Fruit Sampling	19
	3.8.2 Fruit Firmness	19
	3.8.3 Soluble Solids Concentration (SSC)	19
	3.8.4 Titratable Acidity (TA)	20
	3.8.4.1 Preparation of Samples	20
	3.8.4.2 Preparation of Reagents	20
	3.8.5 Fruit pH	20
	3.8.6 Ascorbic Acid	20
	3.8.7 <i>Lycopene</i>	21
3.9	Statistical Analysis	21
4	EFFECT OF DEFICIT IRRIGATION ON GROWTH, YIELD AND POSTHARVEST QUALITIES OF LOWLAND TOMATO FRUITS	22
4.1	Introduction	22
4.2	Materials and Methods	23
	4.2.1 Planting Materials and Treatments	23
	4.2.2 Plant Growth Parameter	23
	4.2.2.1 Plant Height, Stem Diameter (mm) and Leaves Number	23
	4.2.2.2 Total Fruit Weight and Fruit Number	23
	4.2.2.3 Truss and Flowers Numbers	23
	4.2.2.4 Leaf Area Measurement (LA)	23
	4.2.2.5 Root Length, Root Surface Area, Root Volume and Average Root Diameter Determination	24
	4.2.3 Leaf Relative Water Content (LRWC %)	24
	4.2.4 Physico-Chemical Characteristics Fruit Quality Determination	24
	4.2.5 Firmness, Soluble Solids Concentration (SSC), Titratable Acidity, pH, Ascorbic Acid and <i>Lycopene</i>	24
4.3	Results and Discussion	25
	4.3.1 Leaf Water Potential	25
	4.3.2 Growth Parameter	26
	4.3.2.1 Plant Height and Number of Leaf	26
	4.3.2.2 Leaf Area	27
	4.3.2.3 Stem Diameter	28
	4.3.2.4 Flower and Trusses Number	29
	4.3.2.5 Fresh and Dry Weight of Shoots and Root	30
	4.3.2.6 Root Characteristics	35
	4.3.3 Percentage of Leaf Relative Water Content (RWC)	37
	4.3.4 Fruit Weight Per Plant	38
	4.3.5 Fruit Number Per Plant	41
	4.3.6 Physico-Chemical Parameters	44
	4.3.6.1 Firmness	44
	4.3.6.2 Total Soluble Solids Concentration	44

	4.3.6.3 Titratable Acid (TA)	45
	4.3.6.4 Fruit pH	46
	4.3.6.5 Ascorbic Acid	47
	4.3.6.6 <i>Lycopene</i>	48
4.4	Conclusion	49
5	IMPOSITION OF BENEFICIAL WATER STRESS AT DIFFERENT GROWTH STAGES FOR IMPROVEMENT OF POSTHARVEST QUALITY OF LOWLAND TOMATO FRUITS AT DIFFERENT MATURITY STAGES	50
5.1	Introduction	50
5.2	Materials and Methods	51
	5.2.1 Preparation of Planting Materials	51
	5.2.2 Growing conditions	51
	5.2.3 Growth Parameters	51
	5.2.3.1 Plant Height (cm) and Number of Leaves	51
	5.2.3.2 Total Fruit Weight (g/plant) and Number	52
	5.2.3.3 Fresh and dry weight of shoots and root (g/plant)	52
	5.2.4 Analysis of Physicochemical Parameters	52
	5.2.4.1 Firmness (N), SSC (°Brix), Titratable Acidity (%) citric acid, pH , Ascorbic Acid (mg/100g) and <i>Lycopene</i> (mg/kg)	52
	5.2.5 Statistical analysis	52
5.3	Results and Discussion	52
	5.3.1 Growth Parameter	52
	5.3.1.1 Plant Height and Number of Leaves	52
	5.3.1.2 Fresh and Dry weight measurements	53
	5.3.2 Yield and Fruit Number	57
	5.3.3 Physicochemical Parameter	59
	5.3.3.1 Firmness	59
	5.3.3.2 Soluble Solids Content (SSC)	60
	5.3.3.3 Titratable Acidity (TA)	61
	5.3.3.4 Fruit pH	63
	5.3.3.5 Ascorbic Acid (AA)	63
	5.3.3.6 <i>Lycopene</i> Content	65
5.4	Conclusion	65
6	CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	67
	REFERENCES	68
	APPENDICES	80
	BIODATA OF STUDENT	96

LIST OF FIGURES

Figure		Page
2.1	USDA tomato color scheme	5
2.2	Map showing future global water supply status	6
4.1	Effect of DI levels on leaf water potential of tomato plants	25
4.2	Effect of DI levels on height of tomato plants	27
4.3	Effect of DI levels on leaves number of tomato plants	27
4.4	Effect of DI levels on leave area of tomato plants	28
4.5	Effect of DI levels on stem diameter of tomato plants	29
4.6	Effect of DI levels on flower number of tomato plants	30
4.7	Effect of DI levels on trusses of tomato plants	30
4.8	Effect of DI levels on fresh shoots of tomato plants	31
4.9	Effect of DI levels on fresh root of tomato plants	32
4.10	Effect of DI levels on dry biomass of tomato plants	32
4.11	Effect of DI levels on dry root of tomato plants	33
4.12	Effect of DI levels on dry shoots of tomato plants	34
4.13	Effect of DI levels on shoots:root ratio of tomato plants	35
4.14	Effect of DI levels on root length of tomato plants	36
4.15	Effect of DI levels on root surface area of tomato plants	36
4.16	Effect of DI levels on root volume of tomato plants	37
4.17	Effect of DI levels on root diameter of tomato plants	37
4.18	Effect of DI levels on LRWC% of tomato plants	38
4.19	Effect of DI on total fruit weight of tomato	39
4.20	Effect of DI on marketable fruit weight of tomato	40

4.21	Effect of DI on unmarketable fruit weight of tomato	41
4.22	Effect of DI on total fruit number of tomato	42
4.23	Effect of DI on unmarketable fruit number of tomato	43
4.24	Effect of DI on marketable fruit number of tomato	43
4.25	Effect of DI levels on firmness of tomato fruit	44
4.26	Effect of DI levels on SSC of tomato fruit	45
4.27	Effect of DI levels on TA content of tomato fruit	46
4.28	Effect of DI levels on pH juice value of tomato fruit	47
4.29	Effect of DI levels on AA concentration of tomato fruit	48
4.30	Effect of DI levels on lycopene content of tomato fruit	49
5.1A	Effect of DI levels on plant height of tomato	53
5.1B	Effect of DI levels on leaves number of tomato plant	53
5.2A	Effect of DI levels on fresh shoots weight of tomato plant	54
5.2B	Effect of DI levels on fresh root weight of tomato plant	55
5.3	Effect of DI levels on dry shoots weight of tomato plant	55
5.4	Effect of DI levels on dry root weight of tomato plant	56
5.5	Effect of DI levels on shoots to root ratio of tomato plant	56
5.6	Effect of DI levels on biomass of tomato plant	57
5.7	Effect of DI levels on fruit yield of tomato plant	58
5.8	Effect of DI levels on fruit number of tomato plant	58
5.9	Interaction between DI levels and Maturity fruit stages on firmness	60
5.10	Interaction between DI levels and Maturity fruit stages on SSC	61
5.11A	Effect of DI levels on titratable acidity TA of tomato fruit	62
5.11B	Effect of maturity stages on TA of tomato fruit	62

5.12	Interaction between DI levels and Maturity fruit stages on pH	63
5.13A	Effect of DI levels on Ascorbic Acid AA of tomato fruit	64
5.13B	Effect of maturity stages on AA of tomato fruit	64
5.14	Interaction between DI levels and Maturity fruit stages on Lycopene	65



LIST OF ABBREVIATIONS

$^{\circ}\text{C}$	Degree celcius
AA	Ascorbic acid
Anova	Analysis of variance
cm	Centimetre
DAT	Day after treatment
DI	Deficit Irrigation
Dw	Dry weight
ET	Evapotranspiration
ETc	Crop Evapotranspiration
FC	Field Capacity
Fw	Fresh weight
g	Gram
ha	Hectare
kg	Kilogram
LRWC	Leaf Relative Water Content
LSD	Least Significant Difference
LWP	Leaf water potential
M	Molarity
m	Meter
MARDI	Malaysian Agricultural Research and Development Institute
MC	Moisture content
mg	Milligram
mm	Millimetre
Ms	Fruit maturity stage

PRD	Partial root zone drying
RCBD	Randomized Complete Block Design
RDI	Regulated deficit irrigation
SAS	Statistical Analysis System
SSC	Soluble Solids Concentration
TA	Titrateable Acidity
USDA	United State Department of Agriculture
WUE	Water Use Efficiency



CHAPTER I

INTRODUCTION

Tomato (*Lycopersicon esculentum* Mill.) is a member of the night shade family, Solanaceae (Costa and Heuvelink, 2005). It is a highly versatile crop and is among the most widely consumed vegetables. The fruits are rich in vitamins and minerals and contains the valuable carotenoid, lycopene, which is a vital factor in cardiovascular protection and relief of oxidative stress (Abete et al., 2013). Along with total soluble solids (TSS), a valuable attribute for the fruit processing industries, the aforementioned properties constitute part of tomato's important quality parameters that are usually the target of enhancement or maintenance strategies, either at the pre- or postharvest end of the value chain (Ilahy et al., 2011).

According to Arpaia (1994), a postharvest physiologist is interested in maximizing the postharvest quality of horticultural commodities. However, it has been suggested that optimum postharvest quality of vegetables is intrinsically tied to pre-harvest processes or factors. According to Meaza et al., (2007) cited in Sibomana et al. (2015), pre-harvest factors such as the crop genetic status, cultural practices and environmental conditions influence the post-harvest quality of crops. Workneh et al. (2012), reported improved postharvest quality and storability when tomato plants were subjected to pre-harvest treatments that included natural growth enhancers. However, Silva, (2011) reported that the effects of pre-harvest factors are not usually factored into the planning of post-harvest programmes, thereby overlooking a very significant aspect. Whereas post-harvest programmes aim to maintain the quality of produce until they are utilized by consumers, a major determinant of those qualities is the conditions and management practices to which the crop was subjected while in the field. Such factors as irrigation scheduling, including its adequacy and timeliness, fertilization, cultural practices such as weeding, pest control, plant population etc, which influence crop growth and development invariably affect the development and quality of the harvest. It must be noted that nothing can be added to the quality attained at harvest during postharvest treatment. It is therefore pertinent that pre-harvest factors are carefully planned and manipulated with a view to influencing postharvest quality and shelf life (Silva, 2011).

Two major pre-harvest factors that influence post-harvest qualities are watering and maturity fruit harvesting stages (Sibomana et al. (2015); Kader, (1997). According to Sibomana et al. (2015), water supply is a critical determinant of fruit yield and the crop is sensitive to soil moisture level during growth and development. It significantly affects fruit weight, firmness, total soluble solids and titratable acidity. According to Boamah et al. (2010), water deficit reduced tomato growth, fruit yield and quality while Cantore et al. (2012) held that too much water may result in root hypoxia which invariably reduces yield. Mpelasoka et al. (2001) on the other hand enthused that deficit irrigation (DI) enhanced fruit total soluble solid and firmness as well as their maintenance during storage at 0°C. Dorji et al. (2005) also reported enhanced fruit

quality and postharvest quality with DI treatment. They found that the total soluble solid of pepper was higher than control by 8 %. The DI treatment also modified the performance of different quality parameters at different maturity stages. For example, Dorji et al. (2005) found that TSS was 10.2 % for DI at the firm red stage as against 8.4 % for control and 5.0 % at the matured green stage as against 4.5 % in control. From the foregoing, it can be concluded that proper water management for crop growth and yield is not just an important crop management factor but also an important strategy for maintaining post-harvest quality. It would therefore be beneficial to determine the level of water supply that would result in better quality and storability. In this regard a paradigm shift from a principle of adequate water supply to optimum water supply becomes necessary.

This becomes even more pertinent when taken together with the fact that meeting agricultural water demand is increasingly been complicated by its growing scarcity. Globally, water resources have been observed to be declining at an alarming rate, leading to fear of future wide spread scarcity. According to Escobar (2010), by 2025 about half of the population of the world would be facing water scarcity. Water deficit or drought is, globally, the most common stress condition and it is increasingly of concern worldwide (Reddy et al., 2004; Mahajan & Tuteja, 2005). Absolute water stress is found most notably in arid and semi-arid regions with high population densities such as parts of India, China and the Middle East/North Africa (MENA) region. The MENA region is increasingly unable to produce the food required locally due to increasing water stress from a combination of population increase, economic development and climate change, and will have to rely more and more on food (and virtual water) imports. In arid and semi-arid regions, water availability is often a key limiting input.

Tomatoes are very sensitive to drought stress, initially during vegetative development and, later, when the tomato is in the reproductive stage (Wudiri & Henderson, 1985). Poor management is, in most cases the real culprit of water was related to low productivity in agriculture and also a major factor in the growing scarcity of water. In line with this opinion, Boutraa, (2010a) reported that only about 50 % of all water extracted for agricultural purposes are utilized. What is required then is deft management of water resources in such a way as to enhance crop productivity. Water management practices are the tools which can serve to protect our natural capital in water resources and avoid the critical situation for the survival and sustainability of agriculture and economic activities which would ensue from their decline (Postel, 2000). Although the development of irrigation has contributed greatly to increased crop productivity as well as improvement in overall agricultural performance (Hussain and Wijerathna. 2004), it is not without its cost, including negative environmental and health consequences such as increased water logging, scarcity, salinization and water-borne diseases. Some of these problems could be remedied by better management. One management strategy in use is deficit irrigation. Geerts & Raes, (2009) had recommended it as a water saving technology in arid regions and other water scarcity prone areas.

This study, will describe the effects of limiting water supply to emulate water stress during plant development on postharvest qualities of tomato fruits with the aim of identifying the best deficit irrigation program and also the effects of water stress on tomato fruits at different phenological stages of tomato plant development on yield and quality of different fruit maturity stages. This will enable us to look at the benefits that can be derived with controlled stress imposed as a management practice. Even numerous studies were performed on beneficial DI effects on tomato, but with the current variety used, this study was deemed with merit.

The objectives of this study were to examine the soil moisture depletion that can be allowed in irrigating tomatoes with a view to investigate the effects of different level of water stress on plant growth, yield and postharvest qualities as well as determined water stress effects on lycopene content and tomato fruit quality in the lowland under tropical conditions. The specific objectives of this study were to:

- i. Determine the best water stress level for optimum tomato productivity.
- ii. Determine the effects of water stress at different phenological stages on plant growth and yield of tomato plant development, and quality of fruits at different fruit stages.

REFERENCES

- Abbott, J. A. (1999). Quality measurement of fruits and vegetables. *Postharvest Biology and Technology*, 15(3), 207–225.
- Abdel-Razzak, H., Wahb-Allah, M., Ibrahim, A., Alenazi, M., & Alsadon, A. (2016). Response of cherry tomato to irrigation levels and fruit Pruning under greenhouse conditions. *Journal of Agricultural Science and Technology*, 18(4), 1091–1103.
- Abete, I., Perez-Cornago, A., Navas-Carretero, S., Bondia-Pons, I., Zulet, M. A., & Martinez, J. A. (2013). A regular lycopene enriched tomato sauce consumption influences antioxidant status of healthy young-subjects: A crossover study. *Journal of Functional Foods*, 5(1), 28–35.
- Agbemafle, R., Owusu-sekyere, J., Bart-plange, A., & Otchere, J. (2014). Effect of deficit irrigation and storage on physicochemical quality of tomato (*Lycopersicon esculentum* Mill . Var . Pechtomech). *Journal of Food Science and Quality Management*, (34), 113-120.
- Ahmed, A. F., Yu, H., Yang, X., & Jiang, W. (2014). Deficit irrigation affects growth, yield, vitamin C content, and irrigation water use efficiency of hot pepper grown in soilless culture. *HortScience*, 49(6), 722–728.
- Amor, M. A., & Amor, F. M. (2007). Response of tomato plants to deficit irrigation under surface or subsurface drip irrigation. *Journal of Applied Horticulture*, 9(2): 97-100.
- Anthon, G., & Barrett, D. M. (2006). Standardization of a rapid spectrophotometric method for lycopene analysis. In *X International Symposium on the Processing Tomato 758* (pp. 111–128).
- Arpaia, M. L. (1994). Preharvest factors influencing postharvest quality of tropical and subtropical fruit, 29(9), 982–985.
- Barr, H. D., & Weatherley, P. E. (1962). A re-examination of the relative turgidity technique for estimating water deficits in leaves. *Australian Journal of Biological Sciences*, 15(3), 413-428.
- Barry, C. S., & Giovannoni, J. J. (2007). Ethylene and fruit ripening. *Journal of Plant Growth Regulation*, 26(2), 143.
- Behboudian, M. H., Clothier, B. E., & Zegbe, J. A. (2007). Response of tomato to partial rootzone drying and deficit irrigation. *Revista Fitotecnia Mexicana*, 30 (2), 125 - 131.

- Bekhradi, F., Luna, M. C., Delshad, M., Jordan, M. J., Sotomayor, J. A., Mart'inez-Conesa, C., & Gil, M. I. (2015). Effect of deficit irrigation on the postharvest quality of different genotypes of basil including purple and green Iranian cultivars and a Genovese variety. *Postharvest Biology and Technology*, *100*, 127–135.
- Bian, S., & Jiang, Y. (2009). Reactive oxygen species, antioxidant enzyme activities and gene expression patterns in leaves and roots of Kentucky bluegrass in response to drought stress and recovery. *Scientia Horticulturae*, *120* (2), 264–270.
- Birhanu, K., & Tilahun, K. (2010). Fruit yield and quality of drip-irrigated tomato under deficit irrigation. *African Journal of Food, Agriculture, Nutrition and Development*, *10*(2), 2136-2151.
- Blokhina, O., Virolainen, E., & Fagerstedt, K. V. (2003). Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Annals of Botany*, *91*(2), 179–194.
- Boamah, P. O., Sam-Amoah, L. K., & Owusu-Sekyere, J. D. (2010). Effect of irrigation interval on growth and development of tomato under sprinkler. *Asian Journal of Agricultural Research*, *4*(4), 196–203.
- Boutraa, T. (2010a). Growth performance and biomass partitioning of the desert shrub *Calotropis procera* under water stress conditions. *Research Journal of Agriculture and Biological Sciences*, *6*(1), 20–26.
- Boutraa, T. (2010b). Improvement of water use efficiency in irrigated agriculture: a review. *Journal of Agronomy*, *9*(1), 1–8.
- Bower, J. P. (1984). Effect of fruit water stress and irrigation regime in the ripening of stored avocado fruit, cultivar Fuerte. *South African Avocado Growers Association Yearbook*, *7*, 55–56.
- Boyer, J. S. (1968). Relationship of water potential to growth of leaves. *Plant Physiology*, *43*(7), 1056–1062.
- Bradford, K. J., & Hsiao, T. C. (1982). Physiological responses to moderate water stress. In *Physiological plant ecology II*. Springer (pp. 263–324)..
- Calzadilla, A., Rehdanz, K., & Tol, R. S. J. (2010). The economic impact of more sustainable water use in agriculture: A computable general equilibrium analysis. *Journal of Hydrology*, *384*(3), 292–305.
- Camelo, A. F. L. (2004). Manual for the preparation and sale of fruits and vegetables: from field to market. (Vol. 151). Food & Agriculture Org.
- Cantore, V., Pace, B., Todorović, M., De Palma, E., & Boari, F. (2012). Influence of salinity and water regime on tomato for processing. *Italian Journal of Agronomy*, *7*(1), 64-70.

- Chartzoulakis, K., Michelakis, N., & Stefanoudaki, E. (1999). Water use, growth, yield and fruit quality of Bonanza oranges under different soil water regimes. *Advances in Horticultural Science*, 6–11.
- Chen, J., Kang, S., Du, T., Guo, P., Qiu, R., Chen, R., & Gu, F. (2014). Modeling relations of tomato yield and fruit quality with water deficit at different growth stages under greenhouse condition. *Journal of Agricultural Water Management*, 146, 131–148.
- Chen, J., Kang, S., Du, T., Qiu, R., Guo, P., & Chen, R. (2013). Quantitative response of greenhouse tomato yield and quality to water deficit at different growth stages. *Journal of Agriculture Water Management*, 129, 152–163. <http://doi.org/10.1016/j.agwat.2013.07.011>
- Choi, K., Lee, G., Han, Y. J., & Bunn, J. M. (1995). Tomato maturity evaluation using color image analysis. *Transactions-American Society Of Agricultural Engineers*, 38, 171–176.
- Costa, J. M., & Heuvelink, E. (2005). Introduction: the tomato crop and industry. *Tomatoes*, 1, 1–19.
- Costa, J. M., Ortuño, M. F., & Chaves, M. M. (2007). Deficit irrigation as a strategy to save water: physiology and potential application to horticulture. *Journal of Integrative Plant Biology*, 49(10), 1421–1434.
- De, P., Martino, A., Raimondi, G., & Maggio, A. (2007). Comparative analysis of water and salt stress-induced modifications of quality parameters in cherry tomatoes. *Journal of Horticultural Science and Biotechnology*, 82(2), 283–289.
- Djurović, N., Ćosić, M., Stričević, R., Savić, S., & Domazet, M. (2016). Effect of irrigation regime and application of kaolin on yield, quality and water use efficiency of tomato. *Journal of Scientia Horticulturae*, 201, 271–278.
- Doorenbos, J., & Kassam, A. H. (1979). Yield response to water. *Irrigation and Drainage Paper*, 33, 257.
- Dorji, K., Behboudian, M. H., & Zegbe-Dominguez, J. A. (2005). Water relations, growth, yield, and fruit quality of hot pepper under deficit irrigation and partial rootzone drying. *Journal of Scientia Horticulturae*, 104(2), 137–149.
- English, M., & Raja, S. N. (1996). Perspectives on deficit irrigation. *Journal of Agricultural Water Management*, 32(1), 1–14.
- Escobar, I. C. (2010). An overview of the global water situation. *Sustainability Science and Engineering*, 2, 3–5.
- Ezekiel, O., Jane, O. I., & Sanusi, H. O. (2013). Vegetative growth and yield of tomato as affected by water regime and mulching. *Production Agriculture and Technology*, 9(2), 43–53.

- Falkenmark, M. (2013). Growing water scarcity in agriculture: future challenge to global water security. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 371(2002).
- Farahani, S. M., & Chaichi, M. R. (2009). Deficit (Limited) Irrigation – A Method for Higher Water Profitability. *Irrigation Systems and Practices in Challenging Environments*, pp. 19-32.
- Farahani, S. M., Mazaheri, D., Chaichi, M., Tavakkol Afshari, R., & Savaghebi, G. (2010). Effect of seed vigour on stress tolerance of barley (*Hordeum vulgare*) seed at germination stage. *Journal of Seed Science and Technology*, 38(2), 494–507.
- Farooq, M., Wahid, A., Kobayashi, N., Fujita, D., & Basra, S. M. A. (2009). Plant drought stress: effects, mechanisms and management. *Agronomy for Sustainable Development*, 29(1), 185–212.
- Faures, J. M., Hoogeveen, J., & Bruinsma, J. (2002). The FAO irrigated area forecast for 2030. *FAO, Rome, Italy*.
- Fereres, E., Goldhamer, D. A., & Parsons, L. R. (2003). Irrigation water management of horticultural crops. *HortScience*, 38(5), 1036–1042.
- Fereres, E., & Soriano, M. A. (2007). Deficit irrigation for reducing agricultural water use. *Journal of Experimental Botany*, 58(2), 147–159.
- Fray, R. G., Wallace, A., Grierson, D., & Lycett, G. W. (1994). Nucleotide sequence and expression of a ripening and water stress-related cDNA from tomato with homology to the MIP class of membrane channel proteins. *Plant Molecular Biology*, 24(3), 539–543.
- Garcia, E., & Barrett, D. M. (2006). Evaluation of processing tomatoes from two consecutive growing seasons: quality attributes, peelability and yield. *Journal of Food Processing and Preservation*, 30(1), 20–36.
- Geerts, S., & Raes, D. (2009). Deficit irrigation as an on-farm strategy to maximize crop water productivity in dry areas, *Journal Agriculture Water Management*. 96, 1275–1284. <http://doi.org/10.1016/j.agwat.2009.04.009>
- Giannakoula, A. E. & Ilias, I.F. (2013). The effect of water stress and salinity on growth and physiology of tomato (*Lycopersicon esculentum* Mil.). *Journal of Biological Sciences*, 65 (2), 611-620.
- Guichard, S., Gary, C., Longuenesse, J. J., & Leonardi, C. (1999). Water fluxes and growth of greenhouse tomato fruits under summer conditions. *Acta Horticulturae*, 223–230.

- Hayatu, M., Muhammad, S. Y., & Abdu, H. U. (2014). Effect of water stress on the leaf relative water content and yield of some cowpea (*Vigna Unguiculata* (L) Walp.) genotype. *International Journal of Scientific & Technology Research* 3 (7).
- Helyes, L., Lugasi, A., & Pek, Z. (2012). Effect of irrigation on processing tomato yield and antioxidant components. *Turkish Journal of Agriculture and Forestry*, 36(6), 702–709.
- Helyes, L., & Varga, G. (1993). Irrigation demand of tomato according to the results of three decades. In *V International Symposium on the Processing Tomato*, 376 (pp. 323–328).
- Hockema, B. R., & Etxeberria, E. (2001). Metabolic contributors to drought-enhanced accumulation of sugars and acids in oranges. *Journal of the American Society for Horticultural Science*, 126(5), 599–605.
- Hsiao, T. C. and Xu, L. K. (2000). Sensitivity of growth of roots versus leaves to water stress: biophysical analysis and relation to water transport. *Journal of experimental botany*, 51 (350), pp- 1595-1616.
- Huffman, E. (2014). Water Resources Management: Sector Results Profile. *The World Bank*, <http://www.worldbank.org/en/results/2013/04/15/>.
- Hussain, I., Wijerathna, D. (2004). Irrigation and income-poverty alleviation: a comparative analysis of irrigation systems in developing Asia. *International Water Management Institute, Colombo*.1-42.
- Iglesias, D. J., Tadeo, F. R., Legaz, F., Primo-Millo, E., & Talon, M. (2001). In vivo sucrose stimulation of colour change in citrus fruit epicarps: interactions between nutritional and hormonal signals. *Physiologia Plantarum*, 112(2), 244–250.
- Ilahy, R., Hdider, C., Lenucci, M. S., Tlili, I., & Dalessandro, G. (2011). Phytochemical composition and antioxidant activity of high-lycopene tomato (*Solanum lycopersicum* L.) cultivars grown in Southern Italy. *Scientia Horticulturae*, 127(3), 255–261.
- Jensen, C. R., Battilani, A., Plauborg, F., Psarras, G., Chartzoulakis, K., Janowiak, F., Stikic, F., Jovanovic, Z., Li, G., Qi, X., & Liu, F. (2010). Deficit irrigation based on drought tolerance and root signalling in potatoes and tomatoes. *Agricultural Water Management*, 98(3), 403-413.
- Johnstone, P. R., Hartz, T. K., LeStrange, M., Nunez, J. J., & Miyao, E. M. (2005). Managing fruit soluble solids with late-season deficit irrigation in drip-irrigated processing tomato production. *Journal of HortScience*, 40(6), 1857–1861.
- Jones Jr, J. B. (2007). *Tomato plant culture: in the field, greenhouse, and home garden*. CRC press.

- Kang, S., & Zhang, J. (2004). Controlled alternate partial root-zone irrigation : its physiological consequences and impact on water use efficiency. *Journal of Experimental Botany*, 55(407), 2437–2446.
- Kimura, S., & Sinha, N. (2008). Tomato (*Solanum lycopersicum*): a model fruit-bearing crop. *Journal of Cold Spring Harbor Protocols*, 2008(11), 94–105.
- Kirda, C., Cetin, M., Dasgan, Y., Topcu, S., Kaman, H., Ekici, B., Deria, M., Ozguven, A. I. (2004). Yield response of greenhouse grown tomato to partial root drying and conventional deficit irrigation. *Journal of Agricultural Water Management*, 69(3), 191–201.
- Kole, C. (2007). Genome mapping and molecular breeding in plants (Vol. 1). Springer Heidelberg.
- Kucşu, H., Turhan, A., & Demir, A. O. (2014). The response of processing tomato to deficit irrigation at various phenological stages in a sub-humid environment. *Journal of Agricultural Water Management*, 133, 92–103.
- Kumar, P. S., Singh, Y., Nangare, D. D., Bhagat, K., Kumar, M., Taware, P. B., Kumari, A., Minhas, P. S. (2015). Influence of growth stage specific water stress on the yield, physico-chemical quality and functional characteristics of tomato grown in shallow basaltic soils. *Journal of Scientia Horticulturae*, 197, 261–271.
- Léchaudel, M., & Joas, J. (2007). An overview of preharvest factors influencing mango fruit growth, quality and postharvest behaviour. *Brazilian Journal of Plant Physiology*, 19(4), 287–298.
- Lesage, P., & Destain, M.-F. (1996). Measurement of tomato firmness by using a non-destructive mechanical sensor. *Postharvest Biology and Technology*, 8(1), 45–55.
- Leskovar, D. I., Bang, H., Kolenda, K., Franco, J. A., & Perkins-Veazie, P. (2002). Deficit irrigation influences yield and lycopene content of diploid and triploid watermelon. *Journal of Postharvest Horticulture* 628 (pp. 147–151).
- Liu, F., & Stützel, H. (2004). Biomass partitioning, specific leaf area, and water use efficiency of vegetable amaranth (*Amaranthus* spp.) in response to drought stress. *Scientia Horticulturae*, 102(1), 15–27.
- Liu, F., Jensen, C. R., Shahanzari, A., Andersen, M. N., & Jacobsen, S.-E. (2005). ABA regulated stomatal control and photosynthetic water use efficiency of potato (*Solanum tuberosum* L.) during progressive soil drying. *Journal of Plant Science*, 168(3), 831–836.

- Maestre-valero, J. F., Martin-gorritz, B., Alarcón, J. J., Nicolas, E., & Martinez-alvarez, V. (2016). Economic feasibility of implementing regulated deficit irrigation with reclaimed water in a grapefruit orchard. *Journal of Agricultural Water Management*, 178, 119–125. <http://doi.org/10.1016/j.agwat.2016.09.019>
- Mahajan, S., & Tuteja, N. (2005). Cold, salinity and drought stresses: an overview. *Journal of Biochemistry and Biophysics*, 44(2), 139–158.
- Marouelli, W. A., Silva, W. L. C., & Moretti, C. L. (2004). Production, quality and water use efficiency of processing tomato as affected by the final irrigation timing. *Horticultura Brasileira*.22(2), 226-231.
- Marsal, J., Mata, M., Arbonés, A., Rufat, J., & Girona, J. (2002). Regulated deficit irrigation and rectification of irrigation scheduling in young pear trees: an evaluation based on vegetative and productive response. *European Journal of Agronomy*, 17(2), 111-122.
- Meaza, M., Seyoum, T., & Woldetsadik, K. (2007). Effects of preharvest treatments on yield and chemical composition of tomato. *African Crop Science Journal*, 15(3).
- Mishra, K. B., Iannacone, R., Petrozza, A., Mishra, A., Armentano, N., La Vecchia, G., Trtilek, M., Cellini, F., Nedbal, L. (2012). Engineered drought tolerance in tomato plants is reflected in chlorophyll fluorescence emission. *Journal of Plant Science*, 182, 79–86.
- Mitchell, J. P., Shennan, C., Grattan, S. R., & May, D. M. (1991). Tomato fruit yields and quality under water deficit and salinity. *Journal of the American Society for Horticultural Science*, 116(2), 215–221.
- Mohamed, A. N. (2012). Growth and development of tomato (*Lycopersicon esculentum* Mill.) cultivars in vitro under salinity condition. A thesis submitted in partial fulfilment of the requirements of the degree of Doctor of Horticultural Science at Universiti Putra Malaysia.
- Mpelasoka, B. S., Behboudian, M. H., & Mills, T. M. (2001). Effects of deficit irrigation on fruit maturity and quality of “Braeburn” apple. *Journal of Scientia Horticulturae*, 90(3), 279–290.
- Murshed, R., Lopez-Lauri, F., & Sallanon, H. (2013). Effect of water stress on antioxidant systems and oxidative parameters in fruits of tomato (*Solanum lycopersicon* L, cv. Micro-tom). *Journal of Physiology and Molecular Biology of Plants*, 19(3), 363–378.
- Nahar, K., and Gretzmacher, R. (2002). Effect of water stress on nutrient uptake , yield and quality of tomato (*Lycopersicon esculentum* Mill .) under subtropical conditions. *Journal of Die Bodenkultur*, 53(1), 45-51.

- Nahar, K., & Ullah, S. M. (2011). Effect of water stress on moisture content distribution in soil and morphological characters of two tomato (*Lycopersicon esculentum* Mill) cultivars. *Journal of Scientific Research*, 3(3), 677–682.
- Nangare, D. D., Singh, K. G., & Kumar, S. (2013). Effect of blending fresh-saline water and discharge rate of drip on plant yield, water use efficiency (WUE) and quality of tomato in semi arid environment. *African Journal of Agricultural Research*, 8(27), 3639–3645.
- Nangare, D. D., Singh, Y., Kumar, P. S., & Minhas, P. S. (2016). Growth, fruit yield and quality of tomato (*Lycopersicon esculentum* Mill.) as affected by deficit irrigation regulated on phenological basis. *Journal of Agricultural Water Management*, 171, 73–79.
- Navarro, J. M., Botia, P., & Pérez-Pérez, J. G. (2015). Influence of deficit irrigation timing on the fruit quality of grapefruit (*Citrus paradisi* Mac.). *Journal of Food Chemistry*, 175, 329–336.
- Noctor, G., Arisi, A.-C. M., Jouanin, L., & Foyer, C. H. (1998). Manipulation of glutathione and amino acid biosynthesis in the chloroplast. *Plant Physiology*, 118(2), 471–482.
- Nuruddin, M. M. (2001). Effects of water stress on tomato at different growth stages. McGill University.
- Nuruddin, M. M., Madramootoo, C. A., & Dodds, G. T. (2003). Effects of water stress at different growth stages on greenhouse tomato yield and quality. *Journal of HortScience*, 38 (7), 1389–1393.
- Owusu-Sekyere, J. D., & Andoh, J. (2012). Assessment of Deficit Irrigation on the growth and Yield of some vegetable crops. *Journal of University of Cape Coast, Ghana*, 23–39.
- Pan, X. Y., Wang, G. X., Yang, H. M., & Wei, X. P. (2003). Effect of water deficits on within-plot variability in growth and grain yield of spring wheat in northwest China. *Field Crops Research*, 80(3), 195-205.
- Papadopoulos, A. P. (1991). Growing greenhouse tomatoes in soil and in soilless media. Available from Communications Branch, Agriculture Canada.
- Patanè, C., & Cosentino, S. L. (2010). Effects of soil water deficit on yield and quality of processing tomato under a Mediterranean climate. *Journal of Agricultural Water Management*, 97(1), 131–138.
- Patanè, C., Tringali, S., & Sortino, O. (2011). Effects of deficit irrigation on biomass, yield, water productivity and fruit quality of processing tomato under semi-arid Mediterranean climate conditions. *Journal of Scientia Horticulturae*, 129(4), 590–596.

- Petro-Turza, M. (1986). Flavor of tomato and tomato products. *Journal of Food Reviews International*, 2(3), 309–351.
- Pirzad, A., Shakiba, M. R., Zehtab-Salmasi, S., Mohammadi, S. A., Darvishzadeh, R., & Samadi, A. (2011). Effect of water stress on leaf relative water content, chlorophyll, proline and soluble carbohydrates in *Matricaria chamomilla* L. *Journal of Medicinal Plants Research*, 5(12), 2483–2488.
- Postel, S. L. (2000). Entering an era of water scarcity: the challenges ahead. *Journal of Ecological Applications*, 10(4), 941–948.
- Pulupol, L. U. (1996). Growth, yield, fruit composition and postharvest attributes of glasshouse tomatoes produced under deficit irrigation. *A thesis submitted in partial fulfilment of the requirements of the degree of Master of Horticultural Science at Massey University*.
- Rahman, M. M., Moniruzzaman, M., Ahmad, M. R., Sarker, B. C., & Alam, M. K. (2016). Maturity stages affect the postharvest quality and shelf-life of fruits of strawberry genotypes growing in subtropical regions. *Journal of the Saudi Society of Agricultural Sciences*, 15(1), 28–37.
- Reddy, A. R., Chaitanya, K. V., & Vivekanandan, M. (2004). Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. *Journal of Plant Physiology*, 161(11), 1189–1202.
- Ripoll, J., Urban, L., Staudt, M., Lopez-Lauri, F., Bidel, L. P. R., & Bertin, N. (2014). Water shortage and quality of fleshy fruits—making the most of the unavoidable. *Journal of Experimental Botany*, 65(15), 4097–4117.
- Rockström, J., Karlberg, L., Wani, S. P., Barron, J., Hatibu, N., Oweis, T., Bruggeman, A., Farahani, J., Qiang, Z. (2010). Managing water in rainfed agriculture -The need for a paradigm shift. *Journal of Agricultural Water Management*, 97(4), 543–550.
- Rodriguez, P., Torrecillas, A., Morales, M. A., Ortuno, M. F., & Sánchez-Blanco, M. J. (2005). Effects of NaCl salinity and water stress on growth and leaf water relations of *Asteriscus maritimus* plants. *Journal of Environmental and Experimental Botany*, 53(2), 113–123.
- Saidi, A., Ookawa, T., Motobayashi, T., & Hirasawa, T. (2008). Effects of soil moisture conditions before heading on growth of wheat plants under drought conditions in the ripening stage: insufficient soil moisture conditions before heading render wheat plants more resistant to drought during ripening. *Journal of Plant Production Science*, 11(4), 403–414.
- Savić, S., Liu, F., Stikić, R., Jacobsen, S.-E., Jensen, C. R., & Jovanović, Z. (2009). Comparative effects of partial rootzone drying and deficit irrigation on growth and physiology of tomato plants. *Archives of Biological Sciences*, 61(4), 801–810.

- Seng, K. H. (2014). *The effects of drought, waterlogging and heat stress on tomatoes (Solanum lycopersicon L.)*. Lincoln University.
- Serrano, M., Díaz-Mula, H. M., Zapata, P. J., Castillo, S., Guillén, F., Martínez-Romero, D., Valverde, J. M., Valero, D. (2009). Maturity stage at harvest determines the fruit quality and antioxidant potential after storage of sweet cherry cultivars. *Journal of Agricultural and Food Chemistry*, 57(8), 3240–3246.
- Shahein, M. M., Abuarab, M. E., & Hassan, A. M. (2012). Effects of regulated deficit irrigation and phosphorous fertilizers on water use efficiency, yield and total soluble solids of tomato. *American--Eurasian J. Agric. & Environ. Sci*, 12(10), 1295–1304.
- Shao, H.-B., Chu, L.-Y., Jaleel, C. A., & Zhao, C.-X. (2008). Water-deficit stress-induced anatomical changes in higher plants. *Journal of Comptes Rendus Biologies*, 331(3), 215–225.
- Sibomana, C. I., Opiyo, A. M., & Aguyoh, J. N. (2015). Influence of soil moisture levels and packaging on postharvest qualities of tomato (*Solanum lycopersicum*). *African Journal of Agricultural Research*, 10(12), 1392–1400.
- Sibomana, I. C., Aguyoh, J. N., & Opiyo, M. (2013). Water stress affects growth and yield of container grown tomato (*Lycopersicon esculentum* Mill) plants. *Bangladesh Journal of Agricultural Research*. 2(4), 461-466.
- Silva, E. (2011). Influence of Preharvest Factors on Postharvest Quality. 1–3. <http://www.familyfarmed.org/wholesale-success/> (verified 19 May 2011).
- Smirnoff, N. (1998). Plant resistance to environmental stress. *Current Opinion in Biotechnology*, 9(2), 214–219.
- Steduto, P., Hsiao, T. C., & Fereres, E. (2007). On the conservative behavior of biomass water productivity. *Journal of Irrigation Science*, 25(3), 189–207.
- Taiz, L., & Zeiger, E. (2010). *Plant physiology* 5th Ed. *Sunderland, MA: Sinauer Associates*.
- Taylor, I. B. (1986). Biosystematics of the tomato. In *The tomato crop* (pp. 1–34). Springer.
- Télef, N., Stammitti-Bert, L., Mortain-Bertrand, A., Maucourt, M., Carde, J. P., Rolin, D., & Gallusci, P. (2006). Sucrose deficiency delays lycopene accumulation in tomato fruit pericarp discs. *Journal of Plant Molecular Biology*, 62(3), 453–469.
- Topcu, S., Kirda, C., Dasgan, Y., Kaman, H., Cetin, M., Yazici, A., & Bacon, M. A. (2007). Yield response and N-fertiliser recovery of tomato grown under deficit irrigation. *European Journal of Agronomy*, 26(1), 64–70.

- Torrecillas, A., Guillaume, C., Alarcón, J. J., & Ruiz-Sánchez, M. C. (1995). Water relations of two tomato species under water stress and recovery. *Journal of Plant Science*, 105(2), 169–176.
- Vijitha, R., & Mahendran, S. (2010). Effect of moisture stress at different growth stages of tomato plant (*Lycopersicon esculentum* Mill.) on yield and quality of fruits. *Journal of Sci.Univ.Kelaniya*, 5, 1-11.
- Wahb-Allah, M. A., & Al-Omran, A. M. (2012). Effect of water quality and deficit irrigation on tomato growth, yield and water use efficiency at different developmental stages. *Journal of Agric & Env.Sci. Dam.Univ.,Egypt*. 11(2), 80-109.
- Wang, F., Du, T., Qiu, R., & Dong, P. (2011). Effects of water stress at different growth stage on greenhouse multiple-trusses tomato yield and quality. *New Technology of Agricultural Engineering (ICAE)*, 2011, pp. 282–287.
- Willie, W. K. T. (2016). Effects of deficit irrigation and chicken manure interactions on the growth and yield of okra in a pot experiment. *Asian Journal of Agricultural Research*, 10(1), 47–55. <http://doi.org/10.3923/ajar.2016.47.55>
- Workneh, T. S., Osthoff, G., & Steyn, M. (2012). Effects of preharvest treatment, disinfections, packaging and storage environment on quality of tomato. *Journal of Food Science and Technology*, 49(6), 685–694.
- Wudiri, B. B., & Henderson, D. W. (1985). Effects of water stress on flowering and fruit set in processing-tomatoes. *Journal of Scientia Horticulturae*, 27(3), 189–198.
- Yakushiji, H., Morinaga, K., & Nonami, H. (1998). Sugar accumulation and partitioning in Satsuma mandarin tree tissues and fruit in response to drought stress. *Journal of the American Society for Horticultural Science*, 123(4), 719–726.
- Yang, H., Du, T., Qiu, R., Chen, J., Wang, F., Li, Y., Kang, S. (2017). Improved water use efficiency and fruit quality of greenhouse crops under regulated deficit irrigation in northwest China. *Journal of Agricultural Water Management*, 179, 193–204.
- Yaseen, R., Shafi, J., Ahmad, W., Rana, M. S., Salim, M., & Qaisrani, S. A. (2014). Effect of deficit irrigation and mulch on soil physical properties, growth and yield of maize. *Environment and Ecology Research*, 2(3), 122–137.
- Zegbe, J. A., Behboudian, M. H., & Clothier, B. E. (2006). Responses of “Petopride” processing tomato to partial rootzone drying at different phenological stages. *Journal of Irrigation Science*, 24(3), 203–210.

Zegbe-Domínguez, J. A., Behboudian, M. H., Lang, A., & Clothier, B. E. (2003). Deficit irrigation and partial rootzone drying maintain fruit dry mass and enhance fruit quality in “Petopride” processing tomato (*Lycopersicon esculentum*, Mill.). *Journal of Scientia Horticulturae*, 98(4), 505–510.



© COPYRIGHT UPM



UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : _____

TITLE OF THESIS / PROJECT REPORT :

IMPOSITION OF BENEFICIAL WATER STRESS FOR IMPROVEMENT OF POSTHARVEST QUALITY OF LOWLAND TOMATO FRUITS (*Lycopersicon esculentum* Mill.)

NAME OF STUDENT: MOHAMMED HASSAN NAMA

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (v)

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

OPEN ACCESS

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

PATENT

Embargo from _____ until _____
(date) (date)

Approved by:

(Signature of Student)
New IC No/ Passport No.:

Date :

(Signature of Chairman of Supervisory Committee)
Name:

Date :

[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted.]