



UNIVERSITI PUTRA MALAYSIA

***INFLUENCE OF SOIL pH ON MICROBIAL COMMUNITY AND
INCIDENCE OF BASAL STEM ROT (BSR) DISEASE DEVELOPMENT IN
OIL PALM SEEDLINGS PLANTED ON BSR AFFECTED AND FOREST
SOILS***

KHUNAW ABDULLA RAHMAN

FP 2018 23



**INFLUENCE OF SOIL pH ON MICROBIAL COMMUNITY AND
INCIDENCE OF BASAL STEM ROT (BSR) DISEASE DEVELOPMENT IN
OIL PALM SEEDLINGS PLANTED ON BSR AFFECTED AND FOREST
SOILS**

By

KHUNAW ABDULLA RAHMAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillments of the Requirements for the Degree of Doctor of Philosophy**

September 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

I would like to dedicate this work to those who taught, motivated and helped me throughout my study.

To my memory of late father Abdulla Rahman

This work is also dedicated to my dearest husband, Shihab Ahmed and to my sweet kids Janna, Dana, & Dema.

This work is also dedicated to all my family members



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Degree of Doctor of Philosophy

**INFLUENCE OF SOIL pH ON MICROBIAL COMMUNITY AND
INCIDENCE OF BASAL STEM ROT (BSR) DISEASE DEVELOPMENT IN
OIL PALM SEEDLINGS PLANTED ON BSR AFFECTED AND FOREST
SOILS**

By

KHUNAW ABDULLA RAHMAN

September 2017

Chairman : Associate Professor Radziah Othman, PhD
Faculty : Agriculture

Basal stem rot (BSR) disease caused by *Ganoderma boninense*, remains as the most challenging disease in oil palm industry. In search for a practical approach to halt *Ganoderma* incidence and development in oil palm plantations, experiments were conducted with the following objectives; (i) to evaluate the diversity of microbial communities in BSR affected oil palm and forest soils, (ii) to determine the role of indigenous microbial community in suppressing the development of *G. boninense* in oil palm seedlings, (iii) to determine the effect of soil pH on growth of oil palm seedlings and BSR disease development in *Ganoderma* inoculated seedlings, and (vi) to evaluate the influence of pH and calcium ion in soil amended with liming material on *Ganoderma* development in the roots of oil palm seedlings. Four separate experiments were conducted under laboratory and glasshouse conditions. Soil samples from three different oil palm locations (BSR affected, unaffected and forest) were collected and analyzed for physicochemical properties, the microbial biomass carbon and total microbial enzyme at different depths. Metagenomic determination of soil bacterial diversity in *Ganoderma* affected and unaffected soils was also determined (Experiment 1). Soil sterilization technique was used to eliminate the presence of indigenous microorganisms in the three different soils. Rubber wood blocks fully colonized with *G. boninense* UPM13 mycelium were used to inoculate 3-month old oil palm seedlings. Seedlings with (G+) and without (G-) *Ganoderma* inoculation were then grown in the respective soils for four months. Plant growth (shoot and root dry weights), nutrient uptake, soil microbial properties, disease incidence (DI) and disease severity index (DSI) were determined at harvest (Experiment 2). Soil from BSR affected soil of pH 4.5 was adjusted with liming material, calcium carbonate (CaCO₃) to pH 5, 6, and 7. Oil palm seedlings were then grown with (G+) and without (G-) *Ganoderma* inoculation. Plant growth and disease incidence were

determined as in Experiment 2 (Experiment 3). Acid washed sand culture was adjusted with CaCO_3 to pH 5, pH 6 and pH 7 were each applied with four levels of calcium (0, 0.27, 0.40 and 0.60 g Ca/pot) using CaCl_2 as the calcium source. Oil palm seedlings were grown with and without *Ganoderma* inoculation. Plant growth and disease incidence were determined as in Experiment 2 (Experiment 4). Results of Experiment 1 showed that the populations of bacteria, fungi, and actinomycetes in BSR affected, unaffected and forest soils significantly ($P \leq 0.05$) decreased with increased in soil depth. The observed pattern corresponded with the microbial biomass carbon with the highest value in forest soil. The total microbial enzymes showed decreasing trend with increasing depth, with higher values in forest and BSR unaffected soils. The BSR unaffected soil had the highest relative abundance of Acidobacteria and Actinobacteria compared to other bacterial groups. Both of the bacteria phyla relative abundance exhibited a strong link with soil pH and total carbon. Results of Experiment 2 showed that sterilized soil without presence of indigenous microbial community gave lower plant growth and higher disease incidence. Seedlings grown in non-sterilized and non-*Ganoderma* inoculated forest soil had the highest ($P \leq 0.05$) shoot and root biomass as well as root length, root surface area and root tips number. Mean while, seedlings grown in non-sterilized and non-*Ganoderma* inoculated BSR affected soil had the lowest plant biomass. Higher bacterial populations were observed in the forest soil compared to BSR affected and unaffected soils. Seedlings grown on forest soil showed better growth with lower DI and DSI and having the highest uptake of N (724 mg/plant), P (60 mg/plant) and K (605 mg/plant) compared to others treatments. Experiment 3 showed that the non-*Ganoderma* inoculated seedlings of pH 6 + G- displayed higher top biomass (8.46 %) and highest root growth. Seedlings grown at higher pH of 7 had lower shoot growth compared to that with *Ganoderma* inoculation. *Ganoderma* infected seedlings at pH 5 had the lowest growth. The DI and DSI of seedlings with *Ganoderma* inoculation pH 5 were higher than at other pH. The *Ganoderma* (G+) treated plants displayed lower populations of bacteria, fungi, and actinomycetes than without *Ganoderma* treated groups at all pH levels, though pH 6 + G- soil had the highest bacterial population. Higher nutrient contents, chlorophyll and photosynthesis were observed in plants without *Ganoderma* inoculation as compared to that with *Ganoderma*. The highest shoot N and Ca were observed at pH 6 while the P, K and Mg at pH 5. Experiment 4 showed that addition of Ca to non-*Ganoderma* inoculated seedlings resulted in higher shoot and root biomass with highest yields observed at pH 6 G- (Ca 0.27). Addition of higher Ca (0.6 Ca/pot) increased root length, surface area, and volume (2.99, 3.24 and 2.34 %, respectively) and the bole size compared to at other pH. Higher lignin content of *Ganoderma* infected plants were observed at pH 6 G+ Ca (0.60) compared to at pH 4.5. The pH 6 together with 0.6 Ca g/pot application significantly reduced the disease infection in *Ganoderma* affected seedlings and improved biomass yield and activity of indigenous microorganisms. Under controlled condition, pH 6 with addition of Ca appeared to be effective in controlling BSR disease in oil palm seedlings.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**PENGARUH pH TANAH TERHADAP KOMUNITI MIKROORGANISMA
DAN INSIDEN PEMBENTUKAN PENYAKIT REPUT PANGKAL BATANG
(BSR) DALAM ANAK BENIH KELAPA SAWIT DI ATAS TANAH
TERJEJAS BSR DAN HUTAN**

Oleh

KHUNAW ABDULLA RAHMAN

September 2017

Pengerusi : Profesor Madya Radziah Othman, PhD
Fakulti : Pertanian

Penyakit reput pangkal batang (BSR) yang disebabkan oleh *Ganoderma boninense*, kekal sebagai penyakit yang paling mencabar dalam industri kelapa sawit. Dalam mencari pendekatan praktikal untuk menghentikan kejadian *Ganoderma* dan pembangunan di ladang kelapa sawit, kajian berikut telah dijalankan dengan objektif (i) untuk menilai kepelbagaian komuniti mikrob dalam tanah kelapa sawit yang dijangkiti BSR dan tanah hutan, (ii) untuk menentukan peranan komuniti mikrob asli dalam menghalang pembentukan *G. boninense* dalam anak benih kelapa sawit, (iii) untuk menentukan kesan pH tanah pada pertumbuhan anak benih kelapa sawit dan perkembangan penyakit reput pangkal batang di dalam benih yang diinokulasi *Ganoderma*, dan (vi) untuk menilai pengaruh pH dan kalsium dalam tanah yang dipinda dengan bahan kapur pada pembentukan *Ganoderma* dalam akar anak benih kelapa sawit. Empat eksperimen yang berasingan telah dijalankan di makmal dan rumah kaca. Sampel tanah dari tiga lokasi kelapa sawit berbeza (terjejas BSR, tidak terjejas BSR dan hutan) dikumpulkan dan dianalisa sifat fizikokimia, biojisim karbon, jumlah enzim mikrob di kedalaman yang berbeza. Penentuan metagenomik kepelbagaian bakteria tanah dalam tanah terjejas dan tak terjejas *Ganoderma* ditentukan (Eksperimen 1). Teknik pensterilan tanah telah dilakukan untuk menghapus mikroorganisma asli dalam tiga sampel tanah berbeza. Blok kayu getah yang penuh koloni mesilium *G. boninense* UPM13 digunakan untuk inokulasi pada anak benih kelapa sawit yang berusia 3 bulan. Anak benih kemudiannya dengan inokulasi (G+) dan tanpa (G-) *Ganoderma* ditanam selama 4 bulan. Pertumbuhan pokok (berat kering pucuk dan akar), pengambilan nutrien, sifat mikrob tanah, insiden penyakit (DI) dan indeks keterukan penyakit (DSI) ditentukan pada masa tuaian (Eksperimen 2). Tanah daripada kawasan terjejas BSR pada pH 4.5 diubahsuaikan dengan bahan pengapuran, kalsium karbonat (CaCO₃) kepada pH 5, 6 dan 7. Anak

benih kelapa sawit kemudiannya ditanam dengan inokulasi (G+) dan tanpa (G-) *Ganoderma*. Pertumbuhan pokok dan insiden penyakit ditentukan seperti dalam Eksperimen 2 (Eksperimen 3). Kultur pasir dicuci asid diubahsuai dengan CaCO_3 kepada pH 5, pH 6 dan pH 7, setiap satu ditambah empat tahap kalsium (0, 0.27, 0.40 dan 0.60 g Ca/pasu) menggunakan CaCl_2 sebagai sumber kalsium. Anak benih kelapa sawit ditanam dengan dan tanpa inokulasi *Ganoderma*. Pertumbuhan pokok dan insiden penyakit ditentukan seperti dalam Eksperimen 2 (Eksperimen 4). Keputusan Eksperimen 1 menunjukkan bahawa populasi bakteria, kulat, dan aktinomicetes dalam tanah terjejas BSR, tanah tidak terjejas dan tanah hutan berkurangan dengan ketara ($P \leq 0.05$) dengan peningkatan kedalaman tanah. Pola ini diperhatikan bersesuaian dengan hasil biojisim karbon mikrob dimana tanah hutan menunjukkan nilai tertinggi. Jumlah enzim mikrob menunjukkan trend menurun dengan peningkatan kedalaman, dengan nilai tertinggi dalam tanah hutan dan tanah tidak terjejas BSR. Tanah tidak terjejas BSR mempunyai kelimpahan relatif tertinggi Acidobacteria dan Aktinobacteria berbanding kumpulan bakteria lain. Kedua-dua phyla bakteria ini menunjukkan hubungan yang kuat dengan pH tanah dan jumlah karbon. Keputusan Eksperimen 2 menunjukkan tanah yang disteril tanpa kehadiran komuniti mikrob asli menyebabkan pertumbuhan pokok yang rendah dan tinggi insiden penyakit. Anak benih yang ditanam dalam tanah hutan tidak steril dan tanpa diinokulasi *Ganoderma* mempunyai dengan ketara ($P \leq 0.05$) biojisim pucuk dan akar yang tertinggi serta panjang akar, luas permukaan akar dan bilangan hujung akar. Sebaliknya anak benih ditanam dalam tanah terjejas BSR tidak steril dan tanpa inokulasi *Ganoderma* mempunyai biojisim pokok terendah. Populasi bakteria lebih tinggi diperhatikan di dalam tanah hutan berbanding dengan tanah terjejas dan tidak terjejas BSR. Anak benih ditanam atas tanah hutan menunjukkan pertumbuhan lebih baik dengan DI dan DSI rendah dan mempunyai pengambilan N yang tertinggi (724 mg/pokok), P (60 mg/pokok) dan K (605 mg/pokok) berbanding rawatan lain. Eksperimen 3 menunjukkan bahawa anak benih tanpa inokulasi *Ganoderma* pada pH 6 + G- memaparkan biojisim bahagian atas lebih tinggi (8.46%) dan pertumbuhan akar tertinggi. Anak benih ditanam pada pH lebih tinggi pH 7 menunjukkan pertumbuhan pucuk lebih rendah berbanding yang diinokulasi *Ganoderma*. Anak benih terjangkit *Ganoderma* pada pH 5 mempunyai pertumbuhan terendah. Nilai DI dan DSI anak benih diinokulasi *Ganoderma* pada pH 5 adalah lebih tinggi daripada pH lain. Tanaman dirawat *Ganoderma* (G +) menunjukkan populasi bakteria, kulat, dan aktinomicetes yang lebih rendah daripada kumpulan tanpa *Ganoderma* (G-) yang dirawat di semua tahap pH, walaupun tumbuhan pH 6+ G- mempunyai populasi bakteria tertinggi. Kandungan nutrien, klorofil dan fotosintesis tertinggi diperolehi dalam pokok tanpa inokulasi *Ganoderma* berbanding ada *Ganoderma*. Peratusan N dan Ca pucuk yang paling tinggi dicatat dalam pH 6 dan P, K dan Mg pada pH 5. Eksperimen 4 mendedahkan penambahan Ca pada anak benih tanpa inokulasi *Ganoderma* menghasilkan pertumbuhan pucuk dan akar lebih tinggi dengan hasil tertinggi diperolehi pada pH 6 G- (Ca 0.27). Penambahan Ca (0.6 Ca/pasu) meningkatkan panjang akar, luas permukaan, dan isipadu (2.99, 3.24 dan 2.34%, masing-masing) dan saiz bol berbanding pada pH lain. Kandungan lignin lebih tinggi dalam pokok terjangkit *Ganoderma* didapati pada pH 6 G+ Ca (0.60) berbanding pada pH 4.5. Pada pH 6 dengan 0.6 Ca g/pasu ketara merencat jangkitan *Ganoderma* dalam tanah terjejas BSR dan memperbaiki hasil biojisim dan aktiviti mikroorganisma asli.

Dalam keadaan terkawal, pH 6 dengan penambahan Ca menunjukkan efektif dalam kawalan penyakit BSR pada anak benih kelapa sawit.



ACKNOWLEDGEMENTS

First of all, I am thankful to Allah for giving me the strength, guidance, and perseverance to complete this thesis. May blessing and peace be upon Prophet Muhammad (S. A. W.), who was sent as a mercy to the World. Secondly, this work would have been impossible without the continuous support and supervision of my supervisor, Assoc. Professor, Dr. Radziah Othman Dr. All the taken steps on the way to finishing this thesis were under her direct guidance, she acted not only as a supervisor but also as a mother. I am deeply indebted to her. Alongside her, the other members of the supervisory committee, Professor, Dr. Mohamed Hanafi Musa, Dr. Nusaibah Hj.Syed Ail, who never came short answering my questions and giving helpful comments. I am also thankful to the laboratory assistants in the Department of Land Management, who endured with great patience in all my laboratory tasks. My special thanks to Dr. Dayar who edited my thesis and Dr. Mahdi who did the statistical examination. My sincere thanks are extended to the laboratory assistants, technicians, and officers in the Department of Land Management, Faculty of Agriculture, UPM. Sincere thanks to Universiti Putra Malaysia (UPM), for providing the facilities to conduct the study.

I would like to express my gratitude to my lovely mother, Gulle and my mother in law, Fatima, brothers, Abdulkader & AbulAziez and sisters, Mhabad and Suham for their encouragement and support. My husband, Shihab Ahmed, I am grateful for your awesome love. Thank you for the encouragement, support, and enthusiasm. My beloved kids Janna, Dana, & Dema are my life and my source of inspiration. The authors would like to thank United Malacca Berhad for providing the fund for the project and assistants for helping with the soil sampling.

I certify that a Thesis Examination Committee has met on 19 September 2017 to conduct the final examination of Khunaw Abdulla Rahman on her thesis entitled "Influence of Soil pH on Microbial Community and Incidence of Basal Stem Rot (BSR) Disease Development in Oil Palm Seedlings Planted on BSR Affected and Forest Soils" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Hamdan bin Jol, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Samsuri bin Abd. Wahid, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Ganesan a/l Vadamalai, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Kazuyuki Inubushi, PhD

Professor
Chiba University
Japan
(External Examiner)



NOR AINI AB. SHUKOR, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 December 2017

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Radziah Othman, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairperson)

Hanafi M. M., PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Nusaibah S. A., PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations, and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____

Date: _____

Name and Matric No.: Khunaw Abdulla Rahman (GS35445)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Associate Professor Dr. Radziah Othman

Signature: _____
Name of Member
of Supervisory
Committee: Professor Dr. Hanafi M. M

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Nusaibah S. A

TABLE OF CONTENTS

		Page
	ABSTRACT	i
	ABSTRAK	iii
	ACKNOWLEDGEMENTS	vi
	APPROVAL	vii
	DECLARATION	ix
	LIST OF TABLES	xv
	LIST OF FIGURES	xvii
	LIST OF ABBREVIATIONS	xxi
	CHAPTER	
1	GENERAL INTRODUCTION	1
2	LITERATURE REVIEW	3
	2.1 Oil palm cultivation and its economic significance to Malaysia	3
	2.2 Soil microbial community composition in relation to <i>Ganoderma</i> development	6
	2.3 Microbial diversity and health in relation soil function	7
	2.4 Factors affecting microbial diversity	8
	2.4.1 Competition between plants Nutrient uptake and microorganisms	8
	2.4.2 Effect of soil acidity and alkalinity on microbial structure	9
	2.4.3 Soil enzymes	10
	2.5 Oil palm and <i>Ganoderma</i> disease	11
	2.5.1 <i>Ganoderma</i> species and structure	11
	2.5.2 <i>Ganoderma</i> infection, transmission, incidence, and severity	12
	2.6 Liming, microbial activities, and plant diseases	14
	2.7 Microbial diversity and soil pH	15
	2.8 Factors affecting microbial community	16
	2.9 Microbial classification and diversity	16
	2.10 BSR control and current management practices	17
	2.10.1 Cultural control	18
	2.10.2 Chemical control	18
	2.10.3 Biological control	19
	2.11 Summary of Literature review	19
3	EVALUATION OF THE DIVERSITY OF MICROBIAL COMMUNITIES IN BASAL STEM ROT AFFECTED OIL PALM AND FOREST SOILS	21
	3.1 Introduction	21
	3.2 Materials And Methods	22
	3.2.1 Soil sampling	22
	3.2.2 Soil analysis	25
	3.2.2.1 Soil Texture	25

3.2.2.2	Determination of pH and electrical conductivity	25
3.2.2.3	Total carbon, nitrogen, and sulfur content	25
3.2.2.4	Soil phosphorus	25
3.2.2.5	Determination of Exchangeable Ca, Mg, Na, and K	25
3.2.2.6	Cation exchange capacity	26
3.2.3	3.2.3 Total enzyme of soils (Fluorescein Diacetate Assay, FDA)	26
3.2.4	Microbial biomass carbon measurements (Fumigation- extraction method)	26
3.2.5	Microbiological analysis	27
3.2.6	Metagenomic determination of microbial diversity	27
3.2.6.1	Soil DNA Extraction	27
3.2.6.2	Library preparation	27
3.2.6.3	First stage PCR	27
3.2.6.4	First and second PCR clean-up	28
3.2.7	Library denaturing	28
3.2.7.1	Amplicon library and PhiX control preparation	29
3.2.8	QIIME to analyze 16S DNA gene sequencing	29
3.2.8.1	Protocol 1: Acquiring and demultiplexing DNA sequences	29
3.2.8.2	Protocol 2: Picking OTUs, allocating taxonomy, inferring phylogeny, and generating an OTU table	29
3.2.8.3	Protocol 3: Alpha diversity within samples and rarefaction curves	30
3.2.8.4	Protocol 4: Beta diversity between samples and beta diversity plots	30
3.2.8.5	Statistical analysis	30
3.3	Results	30
3.3.1	Physicochemical properties of soil from different locations	30
3.3.2	Populations of soil microorganisms at different soil depths	35
3.3.3	Microbial Biomass carbon in different soils at different depths	38
3.3.4	Total enzyme (FDA) in different soils at different depths	38
3.3.5	Metagenomic determination	39
3.3.6	3.3.6 Sequence merging	40
3.3.7	Quality filtering	40
3.3.8	Taxonomic classification of bacteria in different soils	41
3.3.9	The dominant bacterial phyla	42
3.4	Discussion	53
3.5	Conclusion	57

4	ROLE OF INDIGENOUS MICROBIAL COMMUNITY IN SUPPRESSING <i>GANODERMA</i> DEVELOPMENT IN OIL PALM SEEDLINGS	58
4.1	Introduction	58
4.2	Materials And Methods	59

4.2.1	Soil sample collection and <i>Ganoderma</i> preparations	59
4.2.2	Microbiological analysis	59
4.2.3	Preparation of rubber wood block and inoculum	59
4.2.4	Planting of oil palm seedling	60
4.2.5	BSR disease incidence	60
4.2.6	Disease Severity Index	60
4.2.7	Photosynthetic leaf determination	61
4.2.8	Seedling harvest	61
4.2.9	Mineral contents	61
4.2.10	Cellulose and lignin determination	61
4.2.11	Statistical analysis	62
4.3	Results	62
4.3.1	Plant Biomass	62
4.3.2	Root development	63
4.3.3	Disease Incidence and Severity	65
4.3.4	Population of microorganisms	67
4.3.5	Nutrient uptake	69
4.3.6	Plant physiological activities and root lignin	70
4.4	Discussion	71
4.5	Conclusion	74
5	EFFECT OF SOIL pH ON GROWTH OF OIL PALM SEEDLINGS AND BASAL STEM ROT DISEASE INCIDENCE IN <i>GANODERMA</i> INOCULATED PLANTS	75
5.1	Introduction	75
5.2	Materials And Methods	76
5.2.1	Preparation of rubber wood block and inoculum	76
5.2.2	Soil microbial population and analysis	76
5.2.3	Fungi and bacteria identification	77
5.2.4	Soil sampling and Soil pH liming preparation	77
5.2.5	Seedling planting	78
5.2.6	Disease incidence	78
5.2.7	Disease severity index (DSI)	78
5.2.8	Mineral contents determination	78
5.2.9	Leaf's photosynthesis determination	78
5.2.10	Plant harvesting	78
5.2.11	Cellulose and Lignin determination	78
5.2.12	The 16S rRNA sequencing	79
5.2.13	Statistical analysis	79
5.3	Results	79
5.3.1	Root biomass yields	79
5.3.2	Disease incidence and severity	83
5.3.3	5.3.3 Microbial population	84
5.3.4	Nutrient uptake	85
5.3.5	Plant photosynthesis and chlorophyll	86
5.3.6	Root lignin and cellulose contents	87
5.3.7	Bacterial and fungal characteristics on culture media	88
5.4	Discussion	93

5.4.1	Oil Palm Biomass yield and nutrient uptake	93
5.4.2	Plant photosynthesis, cellulose and lignin content	93
5.4.3	Disease incidence and severity	94
5.4.4	pH tolerance and BSR disease	95
5.4.5	Microbial populations and performance in culture media	95
5.5	Conclusion	97
6	EVALUATION OF pH AND LIMING MATERIAL ON DEVELOPMENT OF <i>GANODERMA</i> IN OIL PALM SEEDLINGS GROWN ON SAND CULTURE	98
6.1	Introduction	98
6.2	Materials and methods	99
6.2.1	Rubberwood block and inoculum preparation	99
6.2.2	pH, calcium, and seedling planting on washed sand culture	99
6.2.2.1	Sand culture preparation	99
6.2.3	Photosynthesis, chlorophyll content and bole measurement	100
6.2.4	Seedling harvesting	100
6.2.5	Measurement of nutrients	100
6.2.6	Statistical analysis	100
6.3	Results	100
6.3.1	Biomass yield	100
6.3.2	Root development	102
6.3.3	Bole size	104
6.3.4	Plant lignin content	104
6.3.5	Total enzyme	105
6.3.6	Disease Incidences	106
6.3.7	Nutrient uptakes	107
6.3.8	Chlorophyll content and photosynthesis	109
6.4	Discussion	113
6.4.1	Root growth and plant biomass	113
6.4.2	Plant lignin content and total enzyme	114
6.4.3	Disease incidence	114
6.4.4	Photosynthesis, chlorophyll content and bole size	115
6.4.5	Nutrient uptake	115
6.5	Conclusion	116
7	SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	118
7.1	Summary	118
7.2	Conclusion	121
7.3	Recommendations for future research	122
	REFERENCES	123
	APPENDICES	153
	BIODATA OF STUDENT	180
	LIST OF PUBLICATIONS	181

LIST OF TABLES

Table		Page
3.1	Amplification cycles, temperature, and holding time	28
3.2	The nature Library concentrations	28
3.3	Physicochemical properties of soil in unaffected oil palm area	32
3.4	Physicochemical properties of soil in BSR affected oil palm area	33
3.5	Physicochemical properties of forest soil	34
3.6	Adapter sequence trimming using PEAT toolkit	40
3.7	Paired-end sequence merging using the PEAR toolkit	40
3.8	Quality filtering of samples using FastX-Toolkit	41
4.1	Effect of <i>Ganoderma</i> inoculation and soil sterilization on development of oil palm seedlings roots after three months of growth	65
4.2	Populations of bacteria, fungi, and actinomycetes in three soil type before treatments	68
4.3	Populations of bacteria and fungi in three soils as affected by sterilization and <i>Ganoderma</i> inoculation at 4 months of plant growth	68
4.4	Effect of <i>Ganoderma boninense</i> inoculation and soil sterilization on N, P, and K uptake (g/plant) to the upper plant	69
4.5	Effect <i>Ganoderma boninense</i> inoculation and soil sterilization on root cellulose (%) and lignin (%)	70
5.1	Treatment descriptions	77
5.2	Effect of soil pH levels on root biomass yields of oil palm seedlings after infected by <i>Ganoderma</i>	81
5.3a	Microbial population in soil after plant harvesting	85
5.3b	Rhizosphere and entophyte microbial population after harvested plants	85
5.4	Uptake (mg plant ⁻¹) of metals by the shoot of oil palm grown in different soil pH levels with and without <i>Ganoderma</i>	86

5.5	Effect of soil pH levels on root lignin and cellulose contents	88
5.6	Characteristics of bacterial colonies on solid agar media	88
5.7	Characteristics of fungal colonies in soil with different pH and <i>Ganoderma</i> treatments	90
6.1	Effect of pH and calcium on root development in oil palm seedlings of <i>Ganoderma</i> infected and unaffected plants	103
6.2	Effect of pH and calcium ion on lignin content in oil palm seedlings of <i>Ganoderma</i> infected and unaffected plants	105
6.3	Effect of pH and calcium on shoot nutrient uptake in oil palm seedlings of <i>Ganoderma</i> infected and unaffected plants	109
6.4	Effect of pH and calcium ion on chlorophyll content and photosynthesis in oil palm seedlings of <i>Ganoderma</i> infected and unaffected plants	111
6.5	Coefficients of correlation (<i>p</i>) pH level, calcium and <i>Ganoderma</i> on SPAD, bole size, top and root biomass, total enzyme, N, P, K, Ca, and Mg uptake	112

LIST OF FIGURES

Figure		Page
2.1	Malaysia Oil palm Export Trend 2016/2017	3
2.2	World Palm Oil Production by Countries	5
2.3	Top oil palm producers	5
2.4	The basidiocarp of <i>Ganoderma boninense</i> in oil palm	12
3.1.a	Map of soil sampling locations	23
3.1.b	Soil sampling locations	23
3.2	BSR affected sampling area	24
3.3	Unaffected area sampling	24
3.4	Forest area sampling	24
3.5	Total enzyme analysis	26
3.6	Populations of bacteria (a), fungi (b), and actinomycetes (c) in the forest soil at different depths	35
3.7	Population of bacteria (a), fungi (b), and actinomycetes (c) at different depths in the unaffected soil	36
3.8	Population of bacteria (a), fungi (b), and actinomycetes (c) at different depths of the BSR affected soil	37
3.9	Microbial biomass ($\text{mg g}^{-1}\text{soil}$) at different depth ranges	38
3.10	Total enzyme levels ($\mu\text{g g}^{-1}$) in different depth soils	39
3.11	Taxonomic classification of phyla in the soils of the unaffected, BSR affected and forest from oil palm locations	41
3.12	The taxonomic profiles for the relative phylum-level abundance of each sample, classified by representation at $> 0.1\%$ of total sequences	43
3.13	The taxonomic profiles for the relative phylum-level abundance of each sample, classified by representation at $> 0.1\%$ of total sequences	44

3.14	Taxonomic profile for the relative order-level abundance of each sample, classified by representation at > 0.1 % of the total sequences	45
3.15	Taxonomic profile for the relative order-level abundance of each sample, classified by representation at > 0.1 % of the total sequences.	47
3.16	The taxonomic profiles for the relative genera-level abundance of each sample, classified by representation at > 0.1 % of total sequences	48
3.17	Average alpha diversity rarefaction plot for Bacteria/Archaea	49
3.18	Rarefaction curves representing the observed number of operational taxonomic units (OTUs) of bacterial communities	49
3.19	Rarefaction curves calculated for each sample based on the OUT computations	50
3.20	PCoA Plots of weighted UniFrac Distances. PCoA was performed based on the UniFrac distance matrix generated from sequencing cecal 16S rRNA genes in samples from BSR affected, unaffected and forest soils	51
3.21	Jackknifed UPGMA weighted unifracs bootstrap tree the relationship among the 3 samples, and reveals that the 3 samples from the soils	51
3.22	PCoA Plots of Unweighted UniFrac Distances	52
3.23	Jackknifed UPGMA Unweighted unifracs bootstrap tree the relationship among the 4 soil samples and reveals that the 3 samples from the soils cluster together (9KH, 7KH and 8KH), 7KH forest, 8KH BSR affected1, 9KH BSR affected2 and 10KH Unaffected soil	53
4.1	Effect of <i>Ganoderma</i> and soil sterilization on biomass root and shoot weights of oil palm grown on three soils	63
4.2	Roots of inoculated seedlings of BSR affected soil	64
4.3	Roots of inoculated seedlings of unaffected soil	64
4.4	Roots of inoculated seedlings of forest soil	64

4.5	Effect of <i>Ganoderma</i> and soil sterilization on incidences of <i>G. boninense</i> infection in oil palm seedlings grown in three soils, a)BSR- infected and b) unaffected c)forest soils	66
4.6	Disease severity in response to <i>Ganoderma</i> growth in the forest, BSR affected, and unaffected soils after 8-12 weeks	67
4.7	Effect of <i>Ganoderma boninense</i> inoculation and soil sterilization on plant Photosynthesis	71
5.1	Root yields of inoculated (PH 4.5G+, pH 5 G+, pH 6 G+, and pH 7 G+) seedlings	80
5.2	Root yields of non- <i>Ganoderma</i> -inoculated (PH 4.5G-, pH 5G-, pH 6G-, and pH 7G-) seedlings	80
5.3	The root length and root surface area of <i>Ganoderma</i> -inoculated (pH 6 G+) and non- <i>Ganoderma</i> -inoculated (pH 6G-)	80
5.4	Effect of pH levels on top dry weight of oil palm seedlings after infected by <i>Ganoderma</i>	82
5.5	Effect of different soil pH levels on root biomass weights of oil palm seedlings after infection with <i>Ganoderma</i>	82
5.6	Effect of soil pH levels on disease incidence in <i>Ganoderma boninense</i> inoculated oil palm seedlings.	83
5.7	Effect of soil pH levels on disease severity in <i>Ganoderma boninense</i> inoculated oil palm seedlings	84
5.8	Photosynthesis ($\mu\text{mol photon m}^{-2} \text{ S}^{-1}$)in the shoot of oil palm grown in different soil pH.	86
5.9	Chlorophyll content (mg cm^{-2}) in the shoot of oil palm grown in different levels of soil pH	87
5.10	Characteristics of fungi colonies on solid agar media inside petri desh	89
6.1	Effect of pH and calcium ion on biomass yield shoot and root (g/plant) in oil palm seedlings of <i>Ganoderma</i> (A) infected and (B) non-infected plants	101
6.2	Interaction of soil pH, Ca, and <i>Ganoderma</i> on the bole sizes (mm) in oil palm seedlings of <i>Ganoderma</i> -infected (G+) and unaffected (G-) plants	104

- 6.3 Effect of pH and calcium ion on the total enzymes in oil palm seedlings of the *Ganoderma*-infected (G+) and -unaffected (G-) plant 106
- 6.4 Effect of pH and calcium on disease incidence in *Ganoderma boninense* inoculated oil palm seedlings. 107



LIST OF ABBREVIATIONS

ANOVA	Analysis of variance
AAS	Atomic Adsorption Spectrometer
BSR	Basal stem rot
CEC	Cation exchange capacity
CFU	colony forming unit
DI	Disease incidence
DNA	Deoxyribonucleic acid
DS	Disease severity
EC	Electrical conductivity
G-	non-treated with Ganoderma
G+	treated with Ganoderma
H ₂ O ₂	Hydrogen peroxide
MEB	Malt extra agar
NA	Nutrient agar
NH ₄ OAC	Ammonium acetate
PCR	Polymerase chain reaction
PDA	Potato dextrose agar
PDB	Potato dextrose broth
RCBD	Randomized Complete Block Design
SAS	Statistical Analysis System
SE	Standard error
TC	Total carbon

CHAPTER 1

GENERAL INTRODUCTION

One of the major and devastating diseases in oil palm is basal stem rot (BSR) caused by *Ganoderma boninense* (Chong et al., 2011). *Ganoderma boninense* has been identified as the main causative species in Malaysia and Indonesia (Hushiarian et al., 2013; Mercière et al., 2015; Lee and Chang, 2016). Basal stem rot is a disease that appears as root rot with lower stem rot. The disease commonly spreads by direct root contact and basidiospore, via yet unidentified means (Rees et al., 2009; 2012). Likely, the primary route of infection occurs after the oil palm roots come into contact with inoculant from colonized debris left previously in the soil (Idris et al., 2002; Flood et al., 2005) or roots of adjacent infected palms (De Oliveira et al., 2005). *Ganoderma* species are characterized as white rot fungi with the ability to break down plant cell wall (Paterson, 2007; Naidu et al., 2016a). These fungi interrupt the nutrient and water transportation across the cell wall of oil palm, causing the leaf to wilt, yellowness, canopy mottling, unopened leaves, and lower stem appearance of basidiocarps (Turner, 1981; Chung, 2011). This disease is of a major concern in Malaysian oil palm industry. An average of 50 % yield losses of 13 years old oil palms in Malaysian coastal areas was previously reported to be infected by BSR (Parthiban et al., 2016). The infected oil palms are associated with poor yield and finally collapse of their stands (Mazliham et al., 2007). The incidence of BSR disease appears to be influenced by soil pH level (Parthiban et al., 2016). Soil pH is critical for plant nutrient uptake, nutrition, and disease resistance (Sariah and Zakaria, 2000; Huber, 2007; Tajudin et al., 2016). It has been previously demonstrated that low pH may have some influence in suppressing the BSR disease incidence in oil palm (Parthiban et al., 2016).

Several approaches that include cultural (Naher et al., 2013; Assis et al., 2016a), biological (Bivi et al., 2016; Goh et al., 2016; Ramli et al., 2016), chemical (Nadhran et al., 2015), and mechanical (Abdul Razak et al., 2004) methods have been previously employed to control the BSR incidences in oil palms. However, no current control approaches have been proven effective and satisfactory in inhibiting the *Ganoderma* incidences in oil palm seedlings (Susanto et al., 2005; Assis et al., 2016b). As a result of increasing incidences and economic implication of this disease in Malaysia, there is a need for an effective and feasible approach to controlling the disease incidences and development in oil palm seedlings. It is factual to hypothesize that increasing the soil pH level can prevent the oil palm seedlings from BSR pathogens attack.

The main objectives were:

1. To evaluate the diversity of microbial communities in basal stem rot (BSR) infected oil palm and forest soils.
2. To determine the role of indigenous microbial community in suppressing the development of *Ganoderma boninense* in oil palm seedlings.

3. To determine the effect of soil pH on growth of oil palm seedlings and BSR disease development in *Ganoderma* inoculated seedlings.
4. To evaluate the influence of pH and calcium ion in soil amended with liming material on *Ganoderma* development in the roots of oil palm seedlings



REFERENCES

- Abdul Razak, J., Ahmad, H., Ramdhan, K., Idris, A. S., Rahim, S. A., Aminul, R., & Fauzi, I. (2004). Mechanical Trunk Injection for Control of Ganoderma MPOB Information Series. *MPOB TT*, 215.
- Abdullah, F., Ilias, G. N. M., Nelson, M., Izzati, N. A. M. Z., & Yusuf, U. K. (2003). Disease assessment and the efficacy of trichoderma as a biocontrol agent of basal stem rot of oil palms. *Research Bulatin Science Putra*, 1, 31-33.
- Acharya, T., 2010. Bacterial Culture Media: classification, types and uses. Available at www.microbeonline.com/types-of-bacteriological-culture-medium/ (accessed date: 27 December, 2016).
- Aciego Pietri, J. C., & Brookes, P. C. (2008). Nitrogen mineralisation along a pH gradient of a silty loam UK soil. *Soil Biology and Biochemistry*, 40(3), 797-802.
- Adam, G., & Duncan, H. (2001). Development of a sensitive and rapid method for the measurement of total microbial activity using fluorescein diacetate (FDA) in a range of soils. *Soil Biology and Biochemistry*, 33(7), 943-951.
- Adaskaveg, J. E., Blanchette, R. A., & Gilbertson, R. L. (1991). Decay of date palm wood by white-rot and brown-rot fungi. *Canadian Journal of Botany*, 69(3), 615-629.
- Adaskaveg, J. E., Miller, R. W., & Gilbertson, R. L. (1993). Wood decay, lignicolous fungi, and decline of peach trees in South Carolina. *Plant disease*, 77(7), 707-711.
- Aini, I. N., Ezrin, M. H., & Aimrun, W. (2014). Relationship between soil apparent electrical conductivity and pH value of Jawa series in oil palm plantation. *Agriculture and Agricultural Science Procedia*, 2, 199-206.
- Alabouvette, C. (1999). Fusarium wilt suppressive soils: an example of disease-suppressive soils. *Australasian Plant Pathology*, 28(1), 57-64.
- Alabouvette, C., Hoeper, H., Lemanceau, P., & Steinberg, C. (1996). Soil suppressiveness to diseases induced by soilborne plant pathogens. *Soil biochemistry*, 9, 371-413.
- Alam, A. F., Er, A. C., & Begum, H. (2015). Malaysian oil palm industry: prospect and problem. *Journal of Food, Agriculture & Environment*, 13(2), 143-148.
- Alam, S. M. (1999). Nutrient uptake by plants under stress conditions. *Handbook of plant and crop stress*, 2, 285-313.
- Al-Ani, A. N., & Dudas, M. J. (1988). Influence of calcium carbonate on mean weight diameter of soil. *Soil and Tillage Research*, 11(1), 19-26.

- Alef, K., 1995. Estimation of soil respiration. In *Methods in Soil Microbiology and Biochemistry*. Academic Press: New York, NY, USA, pp. 464-470.
- Alemán, F., Nieves-Cordones, M., Martínez, V., & Rubio, F. (2011). Root K⁺ acquisition in plants: the *Arabidopsis thaliana* model. *Plant and Cell Physiology*, 52(9), 1603-1612.
- Ali, S. R. A., Yaacob, N. S., Seman, I. A., & Wahid, M. B. (2004). Oil palm cellulose and lignin degradation of different *Ganoderma* sp. based on ASTM standard rotting experiment. In *International conference on pests and diseases of importance to the oil palm industry, Malaysian Oil Palm Board, Kuala Lumpur, Malaysia* (pp. 2-15).
- Alvarenga, P., Gonçalves, A. P., Fernandes, R. M., De Varennes, A., Vallini, G., Duarte, E., & Cunha-Queda, A. C. (2008). Evaluation of composts and liming materials in the phytostabilization of a mine soil using perennial ryegrass. *Science of the Total Environment*, 406(1), 43-56.
- Amplicon, P. C. R., Clean-Up, P. C. R., & Index, P. C. R. (2013). 16S Metagenomic Sequencing Library Preparation.
- Amtmann, A., & Armengaud, P. (2009). Effects of N, P, K and S on metabolism: new knowledge gained from multi-level analysis. *Current opinion in plant biology*, 12(3), 275-283.
- Amtmann, A., Troufflard, S., & Armengaud, P. (2008). The effect of potassium nutrition on pest and disease resistance in plants. *Physiologia Plantarum*, 133(4), 682-691.
- Anda, M., Shamshuddin, J., Fauziah, C. I., & Omar, S. S. (2008). Mineralogy and factors controlling charge development of three Oxisols developed from different parent materials. *Geoderma*, 143(1), 153-167.
- Andersen, C. P. (2002). Solute Movement in the Rhizosphere. *Journal of Environmental Quality*, 31(2), 699.
- Anderson, J. P. E., & Domsch, K. H. (1978). A physiological method for the quantitative measurement of microbial biomass in soils. *Soil biology and biochemistry*, 10(3), 215-221.
- Andersson, S., Nilsson, S. I., & Saetre, P. (2000). Leaching of dissolved organic carbon (DOC) and dissolved organic nitrogen (DON) in mor humus as affected by temperature and pH. *Soil Biology and Biochemistry*, 32(1), 1-10.
- Andrew K. (2009). http://mushroomobserver.org/name/show_name/14987
- Ariffin, D., Idris, A. S., & Abdul, H. H. (1989). Significance of the black line within oil palm tissue decayed by *Ganoderma boninense*. *Elaeis*, 1(1), 11-16.

- Ariffin, D., Seman, I. A., & Hassan, A. H. (1991). Histopathological studies on colonization of oil palm root by *Ganoderma boninense*. *Elaeis*, 3(1), 289-293.
- Ariffin, D., Singh, G., Lim, T.K., 2000. *Ganoderma* in Malaysia-Current status and research strategy. In 1989 PORIM International Palm Oil Development Conference. Agriculture. 1989. September 5-9, Kuala Lumpur, Malaysia (No. L-0076). PORIM.
- Arifuzzaman, M., Khatun, M. R., & Rahman, H. (2010). Isolation and screening of actinomycetes from Sundarbans soil for antibacterial activity. *African Journal of Biotechnology*, 9(29), 4615-4619.
- Arnold, G., & Van Diest, A. (1991). Nitrogen supply, tree growth and soil acidification. *Nutrient Cycling in Agroecosystems*, 27(1), 29-38.
- Arya, A., & Perelló, A. E. (Eds.). (2010). *Management of fungal plant pathogens*. Cabi.
- Assis, K., Bonaventure, A., Abdul Rahim, A., Affendy, H., Mohammad, A. A., Siti, K. Y., 2016a. Attitude of Oil Palm Smallholders towards *Ganoderma* Disease. *International Journal of Economics and Financial Issues*, 6(7S).
- Assis, K., Chong, K. P., Idris, A. S., & Ho, C. M. (2016). Economic loss due to ganoderma disease in oil palm. *World Academy of Science, Engineering and Technology, International Journal of Social, Behavioral, Educational, Economic, Business and Industrial Engineering*, 10(2), 631-635. scholar.waset.org/1999.10/10003869
- Awang, Y., Shaharom, A. S., Mohamad, R. B., & Selamat, A. (2009). Chemical and physical characteristics of cocopeat-based media mixtures and their effects on the growth and development of *Celosia cristata*. *American journal of agricultural and biological sciences*, 4(1), 63-71.
- Bååth, E., & Anderson, T. H. (2003). Comparison of soil fungal/bacterial ratios in a pH gradient using physiological and PLFA-based techniques. *Soil Biology and Biochemistry*, 35(7), 955-963.
- Badalyan, S. M., Innocenti, G., & Garibyan, N. G. (2004). Interactions between xylotrophic mushrooms and mycoparasitic fungi in dual-culture experiments. *Phytopathologia Mediterranea*, 43(1), 44-48.
- Baker, D. E., Gorsline, G. W., Smith, C. B., Thomas, W. I., Grube, W. E., & Ragland, J. L. (1964). Technique for rapid analyses of corn leaves for eleven elements. *Agronomy Journal*, 56(2), 133-136.
- Balakrishnan, K., Rajendran, C., & Kulandaivelu, G. (2000). Differential responses of iron, magnesium, and zinc deficiency on pigment composition, nutrient content, and photosynthetic activity in tropical fruit crops. *Photosynthetica*, 38(3), 477-479.

- Bardgett, R. D., Jones, A. C., Jones, D. L., Kemmitt, S. J., Cook, R., & Hobbs, P. J. (2001). Soil microbial community patterns related to the history and intensity of grazing in sub-montane ecosystems. *Soil Biology and Biochemistry*, 33(12), 1653-1664.
- Baudoin, E., Benizri, E., & Guckert, A. (2003). Impact of artificial root exudates on the bacterial community structure in bulk soil and maize rhizosphere. *Soil Biology and Biochemistry*, 35(9), 1183-1192.
- Benson, H. J. (2002). *Microbiological Applications 8th Edition*. New York: McGraw Hill.
- Berg, B., & McClaugherty, C. (2003). *Plant litter: decomposition, humus formation, carbon sequestration*. Springer, Berlin.
- Bezemer, T. M., De Deyn, G. B., Bossinga, T. M., Van Dam, N. M., Harvey, J. A., & Van der Putten, W. H. (2005). Soil community composition drives aboveground plant-herbivore-parasitoid interactions. *Ecology Letters*, 8(6), 652-661.
- Bivi, M. R., Farhana, M. S. N., Khairulmazmi, A., & Idris, A. (2010). Control of *Ganoderma boninense*: A causal agent of basal stem rot disease in oil palm with endophyte bacteria in vitro. *International Journal of Agriculture and Biology*, 12(6), 833-839.
- Bivi, M. S. H. R., Paiko, A. S., Khairulmazmi, A., Akhtar, M. S., & Idris, A. S. (2016). Control of Basal Stem Rot Disease in Oil Palm by Supplementation of Calcium, Copper, and Salicylic Acid. *The plant pathology journal*, 32(5), 396.
- Blagodatskaya, E. V., & Anderson, T. H. (1998). Interactive effects of pH and substrate quality on the fungal-to-bacterial ratio and qCO₂ of microbial communities in forest soils. *Soil Biology and Biochemistry*, 30(10), 1269-1274.
- Blagodatskaya, E., Littschwager, J., Lauerer, M., & Kuzyakov, Y. (2010). Growth rates of rhizosphere microorganisms depend on competitive abilities of plants and N supply. *Plant Biosystems*, 144(2), 408-413.
- Blanchette, R. A. (1984). Screening wood decayed by white rot fungi for preferential lignin degradation. *Applied and Environmental Microbiology*, 48(3), 647-653.
- Bloem, J., Hopkins, D. W., & Benedetti, A. (Eds.). (2005). *Microbiological methods for assessing soil quality*. CABI.
- Blume, E., Bischoff, M., Reichert, J. M., Moorman, T., Konopka, A., & Turco, R. F. (2002). Surface and subsurface microbial biomass, community structure and metabolic activity as a function of soil depth and season. *Applied Soil Ecology*, 20(3), 171-181.

- Bradford, M. A., Watts, B. W., & Davies, C. A. (2010). Thermal adaptation of heterotrophic soil respiration in laboratory microcosms. *Global Change Biology*, *16*(5), 1576-1588.
- Brady, C. J., Gibson, T. S., Barlow, E. W. R., Speirs, J., & Wyn Jones, R. G. (1984). Salt-tolerance in plants. I. Ions, compatible organic solutes and the stability of plant ribosomes. *Plant, Cell & Environment*, *7*(8), 571-578.
- Brady, N.C., & Weil, R.R. (2002). *The Nature and Properties of Soil*, 13th ed. Springer Netherlands, p.249.
- Brant, J. B., Myrold, D. D., & Sulzman, E. W. (2006). Root controls on soil microbial community structure in forest soils. *Oecologia*, *148*(4), 650-659.
- Bray, R. H., & Kurtz, L. T. (1945). Determination of total, organic, and available forms of phosphorus in soils. *Soil science*, *59*(1), 39-46.
- Breton, F., Hasan, Y., Lubis, Z., & De Franqueville, H. (2006). Characterization of parameters for the development of an early screening test for basal stem rot tolerance in oil palm progenies. *Journal of Oil Palm Research*, (spec.), 24-36.
- Bruce, A., & Highley, T. L. (1991). Control of growth of wood decay Basidiomycetes by *Trichoderma* spp. and other potentially antagonistic fungi. *Forest products journal*, *41*(2), 63-67.
- Bryant, J. A., Aylward, F. O., Eppley, J. M., Karl, D. M., Church, M. J., & DeLong, E. F. (2016). Wind and sunlight shape microbial diversity in surface waters of the North Pacific Subtropical Gyre. *The ISME journal*, *10*(6), 1308.
- Bulgarelli, D., Rott, M., Schlaeppli, K., van Themaat, E. V. L., Ahmadinejad, N., Assenza, F., ... & Peplies, J. (2013). Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature*, *501*(7468), S25.
- Burns, R. G., DeForest, J. L., Marxsen, J., Sinsabaugh, R. L., Stromberger, M. E., Wallenstein, M. D., ... & Zoppini, A. (2013). Soil enzymes in a changing environment: current knowledge and future directions. *Soil Biology and Biochemistry*, *58*, 216-234.
- Burton, J., Chen, C., Xu, Z., & Ghadiri, H. (2010). Soil microbial biomass, activity and community composition in adjacent native and plantation forests of subtropical Australia. *Journal of Soils and Sediments*, *10*(7), 1267-1277.
- Calderón, F. J., Jackson, L. E., Scow, K. M., & Rolston, D. E. (2000). Microbial responses to simulated tillage in cultivated and uncultivated soils. *Soil Biology and Biochemistry*, *32*(11), 1547-1559.
- Caldwell, B. A. (2005). Enzyme activities as a component of soil biodiversity: a review. *Pedobiologia*, *49*(6), 637-644.

- Campbell, C. L., & Madden, L. V. (1990). *Introduction to plant disease epidemiology*. John Wiley & Sons.
- Carney, K. M., & Matson, P. A. (2006). The influence of tropical plant diversity and composition on soil microbial communities. *Microbial Ecology*, 52(2), 226-238.
- Castro, H. F., Classen, A. T., Austin, E. E., Norby, R. J., & Schadt, C. W. (2010). Soil microbial community responses to multiple experimental climate change drivers. *Applied and environmental microbiology*, 76(4), 999-1007.
- Cebon, A., & Garnier, J. (2005). Nitrobacter and Nitrospira genera as representatives of nitrite-oxidizing bacteria: detection, quantification and growth along the lower Seine River (France). *Water research*, 39(20), 4979-4992.
- Chaanin, A., & Preil, W. (1992). Kalkinduzierte Eisenmangel-Chlorose und Einflüsse der Stickstoff-Form auf das Wachstum von Rhododendron. *Rhododendron und immergrüne Laubgehölze*, 7-22.
- Chao, J. B., Wang, H. J., Zhang, Y. B., Li, Z. Q., Liu, Y. H., Huo, F. J., ... & Wang, J. J. (2017). A single pH fluorescent probe for biosensing and imaging of extreme acidity and extreme alkalinity. *Analytica Chimica Acta*, 975, 52-60.
- Chen, K., Neimark, H., Rumore, P., & Steinman, C. R. (1989). Broad range DNA probes for detecting and amplifying eubacterial nucleic acids. *FEMS microbiology letters*, 57(1), 19-24.
- ChengTuck, & H., Hashim, K. (1997). Usefulness of soil mounding treatments in prolonging productivity of prime-aged Ganoderma infected palms. *Planter*, 73(854), 239-244.
- Chong, K.P., Lum, M.S., Foong, C.P., Wong, C.M.V.L., Atong, M. and Rossall, S., 2011. First identification of Ganoderma boninense isolated from Sabah based on PCR and sequence homology. *African Journal of Biotechnology*, 10(66), pp.14718-14723.
- Chung, G. (2011). Management of Ganoderma diseases in oil palm plantations. *Planter*, 87(1022), 325-339.
- Chung, W. C., Huang, J. W., & Huang, H. C. (2005). Formulation of a soil biofungicide for control of damping-off of Chinese cabbage (*Brassica chinensis*) caused by *Rhizoctonia solani*. *Biological Control*, 32(2), 287-294.
- Conradie, W. J., Van Zyl, J. L., & Myburgh, P. A. (2017). Effect of soil preparation depth on nutrient leaching and nutrient uptake by young *Vitis vinifera* L. cv. Pinot noir. *South African Journal of Enology and Viticulture*, 17(2), 43-52.
- Cooper, R. M., Flood, J., & Rees, R. W. (2011). Ganoderma boninense in oil palm plantations: current thinking on epidemiology, resistance and pathology. *Planter*, 87(1024), 515-526.

- Corley, R. H. V., & Tinker, P. B. (2003a). Vegetative propagation and biotechnology. *The oil palm*, 4, 201-215.
- Corley, R.H.V., & Tinker, P.B. (2003b). *The Oil Palm*. 4th Edition. Blackwell Publishing. Great Britain. pp. 27-28, 407-413.
- Crutchfield, J. (2007). Indonesia: Palm oil production prospects continue to grow. *United States Department of Agriculture, Foreign Agriculture Services, Commodity Intelligence Report*.
- Cullings, K., Raleigh, C., New, M. H., & Henson, J. (2005). Effects of artificial defoliation of pines on the structure and physiology of the soil fungal community of a mixed pine-spruce forest. *Applied and environmental microbiology*, 71(4), 1996-2000.
- Datnoff, L. E., Elmer, W. H., & Huber, D. M. (2007). *Mineral nutrition and plant disease*. American Phytopathological Society (APS Press).
- De Gannes, V., Bekele, I., Dipchansingh, D., Wuddivira, M. N., De Cairies, S., Boman, M., & Hickey, W. J. (2016). Microbial Community Structure and Function of Soil Following Ecosystem Conversion from Native Forests to Teak Plantation Forests. *Frontiers in microbiology*, 7.
- de Oliveira, F. G. R., Candian, M., Lucchette, F. F., Salgon, J. L., & Sales, A. (2005). A technical note on the relationship between ultrasonic velocity and moisture content of Brazilian hardwood (*Goupia glabra*). *Building and Environment*, 40(2), 297-300.
- Dick, R. P. (1994). Soil enzyme activities as indicators of soil quality. *Defining soil quality for a sustainable environment*, (definingsoilqua), 107-124.
- Dimitriu, P. A., & Grayston, S. J. (2010). Relationship between soil properties and patterns of bacterial β -diversity across reclaimed and natural boreal forest soils. *Microbial ecology*, 59(3), 563-573.
- Doran, J. W., Sarrantonio, M., & Liebig, M. A. (1996). Soil health and sustainability. *Advances in agronomy (USA)*.
- Drenovsky, R., Duncan, R., & Scow, K. (2005). Soil sterilization and organic carbon, but not microbial inoculants, change microbial communities in replanted peach orchards. *California agriculture*, 59(3), 176-181.
- Edmeades, D. C. (2003). The long-term effects of manures and fertilisers on soil productivity and quality: a review. *Nutrient cycling in Agroecosystems*, 66(2), 165-180.
- Eichorst, S. A., Breznak, J. A., & Schmidt, T. M. (2007). Isolation and characterization of soil bacteria that define *Terriglobus* gen. nov., in the phylum Acidobacteria. *Applied and Environmental Microbiology*, 73(8), 2708-2717.

- Eichorst, S. A., Kuske, C. R., & Schmidt, T. M. (2011). Influence of plant polymers on the distribution and cultivation of bacteria in the phylum Acidobacteria. *Applied and environmental microbiology*, 77(2), 586-596.
- El-Gali, Z. I. (2014). Comparison of natural soil sterilization methods and their effects on soil inhabitant fungi. *Natural Science*, 12(4), 72-78.
- Ellert, B. H., Clapperton, M. J., & Anderson, D. W. (1997). An ecosystem perspective of soil quality. *Developments in Soil Science*, 25, 115-141.
- Elliott, L. F., Lynch, J. M., Papendick, R. I., Stotzky, G., & Bollag, J. M. (1996). The microbial component of soil quality. *Soil biochemistry*, 9, 1-21.
- Ellison, A. M. (2010). Partitioning diversity. *Ecology*, 91(7), 1962-1963.
- Erhabor, J. O., & Filson, G. C. (1999). Soil fertility changes under an oil palm-based intercropping system. *Journal of Sustainable Agriculture*, 14(2-3), 45-61.
- Esperschütz, J., Gattinger, A., Mäder, P., Schloter, M., & Fließbach, A. (2007). Response of soil microbial biomass and community structures to conventional and organic farming systems under identical crop rotations. *FEMS Microbiology Ecology*, 61(1), 26-37.
- Fageria, N. K., & Baligar, V. C. (2008). Ameliorating soil acidity of tropical Oxisols by liming for sustainable crop production. *Advances in agronomy*, 99, 345-399.
- FAO (2002). Small-Scale Palm oil processing in Africa. FAO Agricultural Services Bulletin 148, Rome, Italy.
- FAO (2009). FAOSTAT online statistical service. Food and Agriculture Organization of the United Nations, Rome, Italy. Available via URL. <http://faostat.fao.org/> (Accessed August 2016)
- Feeney, D. S., Crawford, J. W., Daniell, T., Hallett, P. D., Nunan, N., Ritz, K., ... & Young, I. M. (2006). Three-dimensional microorganization of the soil-root-microbe system. *Microbial Ecology*, 52(1), 151-158.
- Fernández-Calviño, D., & Bååth, E. (2010). Growth response of the bacterial community to pH in soils differing in pH. *FEMS microbiology ecology*, 73(1), 149-156.
- Fernández-Calviño, D., Martín, A., Arias-Estévez, M., Bååth, E., & Díaz-Raviña, M. (2010). Microbial community structure of vineyard soils with different pH and copper content. *Applied Soil Ecology*, 46(2), 276-282.
- Fierer, N., & Jackson, R. B. (2006). The diversity and biogeography of soil bacterial communities. *Proceedings of the National Academy of Sciences of the United States of America*, 103(3), 626-631.

- Fierer, N., Bradford, M. A., & Jackson, R. B. (2007). Toward an ecological classification of soil bacteria. *Ecology*, 88(6), 1354-1364.
- Fierer, N., Leff, J. W., Adams, B. J., Nielsen, U. N., Bates, S. T., Lauber, C. L., ... & Caporaso, J. G. (2012). Cross-biome metagenomic analyses of soil microbial communities and their functional attributes. *Proceedings of the National Academy of Sciences*, 109(52), 21390-21395.
- Fierer, N., Schimel, J. P., & Holden, P. A. (2003). Variations in microbial community composition through two soil depth profiles. *Soil Biology and Biochemistry*, 35(1), 167-176.
- Fitzherbert, E. B., Struebig, M. J., Morel, A., Danielsen, F., Brühl, C. A., Donald, P. F., & Phalan, B. (2008). How will oil palm expansion affect biodiversity?. *Trends in ecology & evolution*, 23(10), 538-545.
- Flood, J., Hasan, Y., & Foster, H. (2002). Ganoderma diseases of oil palm-an interpretation from Bah Lias Research Station. *Planter*, 78(921), 689-710.
- Flood, J., Hasan, Y., Turner, P.D., & O'Grady, E.B. (2000). The spread of *Ganoderma* from its infective sources in the field and its implication of management of the disease in oil palm. In: Flood J, Bridge PD, Holderness M (eds), *Ganoderma Diseases of Perennial Crops*. CABI Publishing, Wallingford, UK, pp.101-112.
- Flood, J., Keenan, L., Wayne, S., & Hasan, Y. (2005). Studies on oil palm trunks as sources of infection in the field. *Mycopathologia*, 159(1), 101-107.
- Flowers, E. (2013). Microbial community structure & function in estuarine sediments (Doctoral dissertation, Cardiff University).
- Forsyth, M. H. (2009). Microbial Diversity in Virginia Old Hardwood Forest Soil Week .1 Bio 203 laboratory module. Virginia State University, USA, 14p.
- Freeman, C., Liska, G., Ostle, N. J., Lock, M. A., Reynolds, B., & Hudson, J. (1996). Microbial activity and enzymic decomposition processes following peatland water table drawdown. *Plant and soil*, 180(1), 121-127.
- Frey, S. D., Knorr, M., Parrent, J. L., & Simpson, R. T. (2004). Chronic nitrogen enrichment affects the structure and function of the soil microbial community in temperate hardwood and pine forests. *Forest Ecology and Management*, 196(1), 159-171.
- Frey-Klett, P., Chavatte, M., Clause, M. L., Courrier, S., Roux, C. L., Raaijmakers, J., ... & Garbaye, J. (2005). Ectomycorrhizal symbiosis affects functional diversity of rhizosphere fluorescent pseudomonads. *New phytologist*, 165(1), 317-328.
- Fritze, H. (1992). Effects of environmental pollution on forest soil microflora-a review.

- Frostegård, Å., Bååth, E., & Tunlio, A. (1993). Shifts in the structure of soil microbial communities in limed forests as revealed by phospholipid fatty acid analysis. *Soil Biology and Biochemistry*, 25(6), 723-730.
- Gabr, S. M. (1999). The influence of nitrate: ammonium ratios and salinity stress on growth, chemical composition and quality of lettuce (*Lactuca sativa* L.) grown in nutrient solutions. *Alexandria Journal of Agricultural Research*, 44(3), 251-262.
- Garbeva, P. V., Van Veen, J. A., & Van Elsas, J. D. (2004). Microbial diversity in soil: selection of microbial populations by plant and soil type and implications for disease suppressiveness. *Annual Review Phytopathology*, 42, 243-270.
- Garbeva, P., Postma, J., Van Veen, J. A., & Van Elsas, J. D. (2006). Effect of above-ground plant species on soil microbial community structure and its impact on suppression of *Rhizoctonia solani* AG3. *Environmental Microbiology*, 8(2), 233-246.
- Garcia, C., Hernandez, T., & Costa, F. (1994). Microbial activity in soils under Mediterranean environmental conditions. *Soil Biology and Biochemistry*, 26(9), 1185-1191.
- George, S. T., Zakaria, K., & Chung, G. F. (1996). *Updated results (1990-1995) on trunk injection of fungicides for the control of Ganoderma basal stem rot*. Palm Oil Research Institute of Malaysia.
- Gharaibeh, M. A., Eltaif, N. I., & Shunnar, O. F. (2009). Leaching and reclamation of calcareous saline-sodic soil by moderately saline and moderate-SAR water using gypsum and calcium chloride. *Journal of Plant Nutrition and Soil Science*, 172(5), 713-719.
- Gilman, E. F., Leone, I. A., & Flower, F. B. (1987). Effect of soil compaction and oxygen content on vertical and horizontal root distribution. *Journal of Environmental Horticulture*, 5(1), 33-36.
- Girvan, M. S., Bullimore, J., Pretty, J. N., Osborn, A. M., & Ball, A. S. (2003). Soil type is the primary determinant of the composition of the total and active bacterial communities in arable soils. *Applied and Environmental Microbiology*, 69(3), 1800-1809.
- Glenn, A. R., & Dilworth, M. J. (1991). Soil acidity and the microbial population: Survival and growth of bacteria in low pH. In *Plant-soil interactions at low pH* (pp. 567-579). Springer Netherlands.
- Goh, K. J. (1995). In Basiron, Jalani and Chan (Eds.), *Advances in oil palm research*, Vol. I. Malaysian Palm Oil Board, Kuala Lumpur. p. 371-410.
- Goh, K. M. (2016). Induction of defence response in lignin biosynthesis of *Elaeis guineensis* during an interaction with *Ganoderma boninense* (Doctoral dissertation, University of Nottingham Malaysia Campus).

- Goh, K. M., Ganeson, M., & Supramaniam, C. V. (2014). Infection potential of vegetative incompatible *Ganoderma boninense* isolates with known ligninolytic enzyme production. *African Journal of Biotechnology*, 13(9).
- Goh, Y. K., Marzuki, N. F., Tan, S. Y., Tan, S. S., Tung, H. J., Goh, Y. K., & Goh, K. J. (2016). Experimental mixture design as a tool to optimize the growth of various *Ganoderma* species cultivated on media with different sugars. *Mycology*, 7(1), 36-44.
- Gomes, N. C. M., Fagbola, O., Costa, R., Rumjanek, N. G., Buchner, A., Mendona-Hagler, L., & Smalla, K. (2003). Dynamics of fungal communities in bulk and maize rhizosphere soil in the tropics. *Applied and environmental microbiology*, 69(7), 3758-3766.
- Grayston, S. J., & Prescott, C. E. (2005). Microbial communities in forest floors under four tree species in coastal British Columbia. *Soil Biology and Biochemistry*, 37(6), 1157-1167.
- Grubinger, V. (2004). *Soil Microbiology: A Primer*. University of Vermont, Burlington, USA.
- Gupta V., Penton, C., Guo, J., Neate, S., Gillings, M., Ophel-Keller, K., Greenfield, P., Tiedje, J. (2016). Functional composition of a disease suppressive community in an agricultural soil. The 9th Australasian Soilborne Diseases Symposium, 14-17 November 2016, Lincoln University, Canterbury, New Zealand.
- Gurmit, S. (1991). *Ganoderma*-the scourge of oil palm in the coastal area. In *Proceedings of Ganoderma workshop, Bangi, Selangor, Malaysia, 11 September 1990*. (pp. 7-35). Palm Oil Research Institute of Malaysia.
- Halimah, M., Zulkifli, H., Vijaya, S., Tan, Y., Wei, P., Let, C., & May, C. (2010). Life cycle assessment of oil palm seedling production (Part 1). *Journal of Oil Palm Research*, 22(December), 878-886.
- Haney, C. H., Samuel, B. S., Bush, J., & Ausubel, F. M. (2015). Associations with rhizosphere bacteria can confer an adaptive advantage to plants. *Nature plants*, 1(6).
- Harpole, W. (2010). Neutral theory of species diversity. *Nature Edu. Know.* 1, 31.
- Harrison, I., Lavery, M. & Sterling, E. (2004). Alpha, beta, and gamma diversity.
- Hasan, Y. O. N. N. E. S., & Flood, J. (2003). Colonisation of rubber wood and oil palm blocks by monokaryons and dikaryons of *Ganoderma boninense*-implications to infection in the field. *Planter*, 79(922), 31-38.
- Hasan, Y., & Turner, P. D. (1998). The comparative importance of different oil palm tissues as infection sources for basal stem rot in replantings. *Planter*, 74(864), 119-135.

- Hasan, Y., Foster, H. L., & Flood, J. (2005). Investigations on the causes of upper stem rot (USR) on standing mature oil palms. *Mycopathologia*, 159(1), 109-112.
- He, X. Y., Su, Y. R., Liang, Y. M., Chen, X. B., Zhu, H. H., & Wang, K. L. (2012). Land reclamation and short-term cultivation change soil microbial communities and bacterial metabolic profiles. *Journal of the Science of Food and Agriculture*, 92(5), 1103-1111.
- Heinemeyer, A., Ineson, P., Ostle, N., & Fitter, A. H. (2006). Respiration of the external mycelium in the arbuscular mycorrhizal symbiosis shows strong dependence on recent photosynthates and acclimation to temperature. *New Phytologist*, 171(1), 159-170.
- Henriques, F. S. (2004). Reduction in chloroplast number accounts for the decrease in the photosynthetic capacity of Mn-deficient pecan leaves. *Plant science*, 166(4), 1051-1055.
- Hepler, P. K. (2005). Calcium: a central regulator of plant growth and development. *The Plant Cell*, 17(8), 2142-2155.
- Hepting, G. H. (1971). Diseases of forest and shade trees of the United States. *Diseases of forest and shade trees of the United States*.
- Ho, Y. W., & Nawawi, A. (1986). Isolation, growth and sporophore development of *Ganoderma boninense* from oil palm in Malaysia. *Pertanika*, 9(1), 69-73.
- Hodson, M. E., Vijver, M. G., & Peijnenberg, W. J. G. M. (2011). Bioavailability in soils. In: Swartjes, F. A. (ed.). *Dealing with contaminated sites: from theory towards practical application*. Springer, pp. 721-746.
- Hopmans, P., & Elms, S. R. (2009). Changes in total carbon and nutrients in soil profiles and accumulation in biomass after a 30-year rotation of *Pinus radiata* on podzolized sands: Impacts of intensive harvesting on soil resources. *Forest ecology and management*, 258(10), 2183-2193.
- Huber, D. M., & Graham, R. D. (1999). The role of nutrition in crop resistance and tolerance to diseases. *Mineral nutrition of crops: fundamental mechanisms and implications*, 169-206.
- Huber, J. A., Welch, D. B. M., Morrison, H. G., Huse, S. M., Neal, P. R., Butterfield, D. A., & Sogin, M. L. (2007). Microbial population structures in the deep marine biosphere. *science*, 318(5847), 97-100.
- Hushiarian, R., Yusof, N.A. and Dutse, S.W., 2013. Detection and control of *Ganoderma boninense*: strategies and perspectives. SpringerPlus, 2(1), p.555.
- Huston, M. A., & Huston, M. A. (1994). *Biological diversity: the coexistence of species*. Cambridge University Press.

- Idris, A. S. (2009). Basal Stem Rot in Malaysia-Biology, economic importance, epidemiology, detection and control. In *International Workshop on Awareness, Detection and Control of Oil Palm Devastating Diseases* (Vol. 6).
- Idris, A. S., Ariffin, H., & D Ismail, S. (2002). *Control of Ganoderma-infected palm development of pressure injection and field applications* (No. A-).
- Idris, A. S., Basri, D., & MW Kushairi, D. (2006). Techniques for Inoculation of Oil Palm Geminated Seeds with *Ganoderma*. MPOB Information Series, No.314, 4.
- Ishak, C. F., & Jusop, S. (2010). *Weathered tropical soils: the ultisols and oxisols*. Universiti Putra Malaysia Press.
- Islam, K. R., & Weil, R. R. (1998). Microwave irradiation of soil for routine measurement of microbial biomass carbon. *Biology and Fertility of Soils*, 27(4), 408-416.
- Janssen, P. H. (2006). Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Applied and environmental microbiology*, 72(3), 1719-1728.
- Janvier, C., Villeneuve, F., Alabouvette, C., Edel-Hermann, V., Mateille, T., & Steinberg, C. (2007). Soil health through soil disease suppression: which strategy from descriptors to indicators?. *Soil biology and Biochemistry*, 39(1), 1-23.
- Jiang, J. F., Li, J. G., & Dong, Y. H. (2013). Effect of calcium nutrition on resistance of tomato against bacterial wilt induced by *Ralstonia solanacearum*. *European journal of plant pathology*, 136(3), 547-555.
- Jiang, W., Struik, P. C., Van Keulen, H., Zhao, M., Jin, L. N., & Stomph, T. J. (2008). Does increased zinc uptake enhance grain zinc mass concentration in rice?. *Annals of Applied Biology*, 153(1), 135-147.
- Jing, X., Sanders, N. J., Shi, Y., Chu, H., Classen, A. T., Zhao, K., ... & He, J. S. (2015). The links between ecosystem multifunctionality and above-and belowground biodiversity are mediated by climate. *Nature communications*, 6.
- Jones, J. B., & Huber, D. M. (2007). Magnesium and plant disease. *Mineral nutrition and plant disease*. APS Press, St. Paul, 95-100.
- Jones, J. P., Engelhard, A. W., & Woltz, S. S. (1989). Management of fusarium wilt of vegetables and ornamentals by macro- and microelement nutrition. pp. 18–32. In: A.W. Engelhard (ed.). *Soilborne Plant Pathogens: Management of Diseases with Macro- and Microelements*. Am. Phytopathol. Soc. pp. 217.
- Jones, R. T., Robeson, M. S., Lauber, C. L., Hamady, M., Knight, R., & Fierer, N. (2009). A comprehensive survey of soil acidobacterial diversity using pyrosequencing and clone library analyses. *The ISME journal*, 3(4), 442.

- Kader, M. A., & Lindberg, S. (2010). Cytosolic calcium and pH signaling in plants under salinity stress. *Plant signaling & behavior*, 5(3), 233-238.
- Kandeler, E., Tscherko, D., & Spiegel, H. (1999). Long-term monitoring of microbial biomass, N mineralisation and enzyme activities of a Chernozem under different tillage management. *Biology and fertility of soils*, 28(4), 343-351.
- Kapoor, P., & Sharma, B. M. (2014). Studies on different growth parameters of *Ganoderma lucidum*. *International Journal Science Environmental Technology*, 3, 1515-1524.
- Kardol, P., Cornips, N. J., van Kempen, M. M., Bakx-Schotman, J. M., & van der Putten, W. H. (2007). Microbe-mediated plant–soil feedback causes historical contingency effects in plant community assembly. *Ecological monographs*, 77(2), 147-162.
- Kemmitt, S. J., Wright, D., & Jones, D. L. (2005). Soil acidification used as a management strategy to reduce nitrate losses from agricultural land. *Soil Biology and Biochemistry*, 37(5), 867-875.
- Kemmitt, S. J., Wright, D., Goulding, K. W., & Jones, D. L. (2006). pH regulation of carbon and nitrogen dynamics in two agricultural soils. *Soil Biology and Biochemistry*, 38(5), 898-911.
- Kennedy, N., Brodie, E., Connolly, J., & Clipson, N. (2004). Impact of lime, nitrogen and plant species on bacterial community structure in grassland microcosms. *Environmental Microbiology*, 6(10), 1070-1080.
- Kenzo, T., Ichie, T., OGAWA, T., Kashimura, S., Hattori, D., Irino, K. O., ... & Ninomiya, I. (2007). Leaf physiological and morphological responses of seven dipterocarp seedlings to degraded forest environments in Sarawak, Malaysia: a case study of forest rehabilitation practice. *Tropics*, 17(1), 1-16.
- Kenzo, T., Yoneda, R., Matsumoto, Y., Azani, A. M., & Majid, M. N. (2011). Growth and photosynthetic response of four Malaysian indigenous tree species under different light conditions. *Journal of Tropical Forest Science*, 271-281.
- Khairudin, H. (1990). Basal stem rot of oil palm: incidence, etiology and control. *Master of Agriculture Science thesis, Universiti Pertanian Malaysia, Selangor, Malaysia, Selangor, Malaysia*.
- Khairudin, H. (1991). Pathogenicity of three *Ganoderma* species on oil palm seedlings. *Journal Perak Planter's Association*, 43-49.
- Kinkel, L. L., & Andrews, J. H. (1988). Disinfestation of living leaves by hydrogen peroxide. *Transactions of the British Mycological Society*, 91(3), 523-528.
- Kiss, S., Dragan-Bularda, M., & Radulescu, D. (1978). Soil polysaccharidases: activity and agricultural importance. *Soil enzymes. Academic, London*, 117-147.

- Kivlin, S. N., & Hawkes, C. V. (2016). Temporal and Spatial Variation of Soil Bacteria Richness, Composition, and Function in a Neotropical Rainforest. *PloS one*, *11*(7), e0159131.
- Klironomos, J. N. (2002). Feedback with soil biota contributes to plant rarity and invasiveness in communities. *Nature*, *417*(6884), 67-70.
- Ko, W. H., & Kao, C. W. (1989). Evidence for the role of calcium in reducing root disease incited by pythium species. pp. 205-217. In: Arthur W. Englehard (ed.). *Soilborne Plant Pathogens: Management of Diseases with Macro and Microelements*. APS Press. St. Paul, MN. Pp.217.
- Koh, L. P., & Wilcove, D. S. (2008). Is oil palm agriculture really destroying tropical biodiversity?. *Conservation letters*, *1*(2), 60-64.
- Krishna, M. P., Varghese, R., & Mohamed, A. A. (2012). Depth wise variation of microbial load in the soils of midland region of Kerala: a function of important soil physicochemical characteristics and nutrients. *Indian Journal of Education and Information Management*, *1*(3), 126-129.
- Kuczynski, J., Stombaugh, J., Walters, W. A., González, A., Caporaso, J. G., & Knight, R. (2012). Using QIIME to analyze 16S rRNA gene sequences from microbial communities. *Current protocols in microbiology*, 1E-5.
- Kuster E. (1968). The actinomycetes. In: *Soil Biology* A. Burges, F. Raw, eds., Academic Press, London.
- Kuzyakov, Y., & Xu, X. (2013). Competition between roots and microorganisms for nitrogen: mechanisms and ecological relevance. *New Phytologist*, *198*(3), 656-669.
- Laila, N., Shafiquzzaman, S., Umi, K. Y., & Mondal, M. M. A. (2015). Issues of *Ganoderma* spp. and basal stem rot disease management in oil palm. *American Journal of Agricultural Science*, *2*(3), 103-107.
- Larkin, R. P. (2003). Characterization of soil microbial communities under different potato cropping systems by microbial population dynamics, substrate utilization, and fatty acid profiles. *Soil Biology and Biochemistry*, *35*(11), 1451-1466.
- Larson, R. A., & Weber, E. J. (1994). *Reaction mechanisms in environmental organic chemistry*. CRC press.
- Latz, E., Eisenhauer, N., Rall, B. C., Scheu, S., & Jousset, A. (2016). Unravelling linkages between plant community composition and the pathogen-suppressive potential of soils. *Scientific reports*, *6*, 23584.

- Lauber, C. L., Hamady, M., Knight, R., & Fierer, N. (2009). Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Applied and environmental microbiology*, 75(15), 5111-5120.
- Lauber, C. L., Strickland, M. S., Bradford, M. A., & Fierer, N. (2008). The influence of soil properties on the structure of bacterial and fungal communities across land-use types. *Soil Biology and Biochemistry*, 40(9), 2407-2415.
- Leckie, S. E., Prescott, C. E., Grayston, S. J., Neufeld, J. D., & Mohn, W. W. (2004). Characterization of humus microbial communities in adjacent forest types that differ in nitrogen availability. *Microbial Ecology*, 48(1), 29-40.
- Lee, S.S. and Chang, Y.S., 2016. Ganoderma–Jekyll and Hyde mushrooms.
- Legg, T. M., Zheng, Y., Simone, B., Radloff, K. A., Mladenov, N., González, A., ... & McKnight, D. M. (2012). Carbon, metals, and grain size correlate with bacterial community structure in sediments of a high arsenic aquifer. *Frontiers in microbiology*, 3.
- Lelong, C. C., Roger, J. M., Brégand, S., Dubertret, F., Lanore, M., Sitorus, N. A., ... & Caliman, J. P. (2010). Evaluation of oil-palm fungal disease infestation with canopy hyperspectral reflectance data. *Sensors*, 10(1), 734-747.
- Leon, M. C. C., Stone, A., & Dick, R. P. (2006). Organic soil amendments: Impacts on snap bean common root rot (*Aphanomyces euteiches*) and soil quality. *Applied Soil Ecology*, 31(3), 199-210.
- Li, Y., Chen, Y.L., Li, M., Lin, X.G., & Liu, R.J., 2012. Effects of arbuscular mycorrhizal fungi communities on soil quality and the growth of cucumber seedlings in a greenhouse soil of continuously planting cucumber. *Pedosphere* 22, 79–87.
- Lim, H. P., & Fong, Y. K. (2005). Research on basal stem rot (BSR) of ornamental palms caused by basidiospores from *Ganoderma boninense*. *Mycopathologia*, 159(1), 171-179.
- Lim, K., Chuah, J., & Ho, C. (1993). Effects of soil heaping on *Ganoderma* infected oil palms. In *Proceedings of the* (pp. 735-738).
- Lim, Y. W., Kim, B. K., Kim, C., Jung, H. S., Kim, B. S., Lee, J. H., & Chun, J. (2010). Assessment of soil fungal communities using pyrosequencing. *The Journal of Microbiology*, 48(3), 284-289.
- Linderman, R. G. (1992). Vesicular-arbuscular mycorrhizae and soil microbial interactions. *Mycorrhizae in sustainable agriculture*, (mycorrhizaeinsu), 45-70.

- Liu, J. J., Ekramoddoullah, A. K., & Zamani, A. (2005). A class IV chitinase is up-regulated by fungal infection and abiotic stresses and associated with slow-canker-growth resistance to *Cronartium ribicola* in western white pine (*Pinus monticola*). *Phytopathology*, 95(3), 284-291.
- Lodge, D. J., Hawksworth, D. L., & Ritchie, B. J. (1996). Microbial diversity and tropical forest functioning. In *Biodiversity and ecosystem processes in tropical forests* (pp. 69-100). Springer, Berlin, Heidelberg.
- Logue, J. B., & Lindström, E. S. (2010). Species sorting affects bacterioplankton community composition as determined by 16S rDNA and 16S rRNA fingerprints. *The ISME journal*, 4(6), 729.
- Lonsdale, D., Pautasso, M., & Holdenrieder, O. (2008). Wood-decaying fungi in the forest: conservation needs and management options. *European Journal of Forest Research*, 127(1), 1-22.
- Lovelock, C. E., Feller, I. C., Ball, M. C., Engelbrecht, B. M., & Ewe, M. L. (2006). Differences in plant function in phosphorus-and nitrogen-limited mangrove ecosystems. *New Phytologist*, 172(3), 514-522.
- Lozupone, C., & Knight, R. (2005). UniFrac: a new phylogenetic method for comparing microbial communities. *Applied and environmental microbiology*, 71(12), 8228-8235.
- Lu, X., Zhen, G., Ni, J., Kubota, K., & Li, Y. Y. (2017). Sulfidogenesis process to strengthen re-granulation for biodegradation of methanolic wastewater and microorganisms evolution in an UASB reactor. *Water research*, 108, 137-150.
- Luo, X., Fu, X., Yang, Y., Cai, P., Peng, S., Chen, W., & Huang, Q. (2016). Microbial communities play important roles in modulating paddy soil fertility. *Scientific reports*, 6.
- MacKenzie, M. D., & Quideau, S. A. (2010). Microbial community structure and nutrient availability in oil sands reclaimed boreal soils. *Applied Soil Ecology*, 44(1), 32-41.
- Maheswari, S. K., Singh, D. V., & Sahu, A. K. (1999). Effect of several nutrient media on the growth and sporulation of *Alternaria alternata*. *Journal of Mycopathological Research*, 37(1), 21-24.
- Malaysian Palm Oil Board (MPOB) (2016). Overview of the Malaysian Oil Palm Industry. http://bepi.mpob.gov.my/images/overview/Overview_of_Industry_2016.pdf (Accessed on 1 April 2017).
- Malaysian Palm Oil Board (MPOB) (2017). Export Trend 2017. Economic and Industry Development Division. <http://bepi.mpob.gov.my/index.php/en/statistics/export/180-export-2017/812-export-trend-2017.html> (Accessed on 29 March 2017)

- Malcolm, G. M., LÓPEZ-GUTIÉRREZ, J. C., & Koide, R. T. (2009). Little evidence for respiratory acclimation by microbial communities to short-term shifts in temperature in red pine (*Pinus resinosa*) litter. *Global change biology*, 15(10), 2485-2492.
- Marshall, R., Hunt, R., & Pilotti, C. A. (2004). Low cost control for basal stem rot-a Poliamba initiative. *Planter*, 80(936), 173-176.
- McLaren, A. D. (1975). Soil as a system of humus and clay immobilized enzymes. *Chemica Scripta*.
- McMahon, P. (2012). Effect of nutrition and soil function on pathogens of tropical tree crops. In *Plant Pathology*. InTech.
- Meister, P. (2013). Two opposing effects of sulfate reduction on carbonate precipitation in normal marine, hypersaline, and alkaline environments. *Geology*, 41(4), 499-502.
- Mellerowicz, E. J., Baucher, M., Sundberg, B., & Boerjan, W. (2001). Unravelling cell wall formation in the woody dicot stem. In *Plant Cell Walls* (pp. 239-274). Springer Netherlands.
- Mercière, M., Laybats, A., Carasco-Lacombe, C., Tan, J.S., Klopp, C., Durand-Gasselín, T., Alwee, S.S.R.S., Camus-Kulandaivelu, L. and Breton, F., 2015. Identification and development of new polymorphic microsatellite markers using genome assembly for *Ganoderma boninense*, causal agent of oil palm basal stem rot disease. *Mycological Progress*, 14(11), p.103.
- Miller, R. N. G. (1995). *The characterization of Ganoderma populations in oil palm cropping systems* (Doctoral dissertation, University of Reading).
- Miller, R. N. G., Holderness, M., Bridge, P. D., Chung, G. F., & Zakaria, M. H. (1999). Genetic diversity of *Ganoderma* in oil palm plantings. *Plant Pathology*, 48(5), 595-603.
- Miller, R. N. G., Holderness, M., Bridge, P. D., Paterson, R. R. M., Sariah, M., Hussin, M. Z., & Hilsley, E. J. (1995). A multidisciplinary approach to the characterization of *Ganoderma* in oil-palm cropping systems. In *Proceedings of the Contributed Symposium* (Vol. 59, pp. 57-66).
- Miransari, M. (2014). Plant growth promoting rhizobacteria. *Journal of Plant Nutrition*, 37(14), 2227-2235.
- Mohammed, C. L., Rimbawanto, A., & Page, D. E. (2014). Management of basidiomycete root-and stem-rot diseases in oil palm, rubber and tropical hardwood plantation crops. *Forest pathology*, 44(6), 428-446.
- Mohd As'wad, A.W., Sariah, M., Paterson, R. R. M., Abidin, M. Z., & Lima, N. (2011). Ergosterol analyses of oil palm seedlings and plants infected with *Ganoderma*. *Crop protection*, 30(11), 1438-1442.

- Moncalvo, J.M. (2000). Systematics of *Ganoderma*. Pp. 23–45. In: Flood J, Bridge PD, Holderness M (eds) *Ganoderma Diseases of Perennial Crops*. CAB International, Wallingford.
- Moore, J., Macalady, J. L., Schulz, M. S., White, A. F., & Brantley, S. L. (2010). Shifting microbial community structure across a marine terrace grassland chronosequence, Santa Cruz, California. *Soil Biology & Biochemistry*, 42(1), 21-31.
- Nadhrah, N.I., Nulit, R., Nurrashyeda, R., Idris, A.S., 2015. Effect of Formulated Bioorganic Containing Burkholderia GanoEB2 in Suppressing Ganoderma Disease in Oil Palm Seedlings. *Plant Protection Science*, 51(2).
- Naher, L., Tan, S. G., Yusuf, U. K., Ho, C. L., & Siddiquee, S. (2012). Activities of chitinase enzymes in the oil palm (*Elaeis guineensis* Jacq.) in interactions with pathogenic and non-pathogenic fungi. *Plant Omics*, 5(4), 333.
- Naher, L., Yusuf, U. K., Ismail, A., Tan, S. G., & Mondal, M. M. A. (2013). Ecological status of *Ganoderma* and basal stem rot disease of oil palms (*Elaeis guineensis* Jacq.). *Australian Journal of Crop Science*, 7(11), 1723.
- Naidu Y., Siddiqui Y., Rafii M.Y., Saud H.M., & Idris A.S. (2016b). Biodegradation of basal stem rot-affected oil palm stumps by white-rot Hymenomycetes: potential disease management by prevention of inoculum spread. The 9th Australasian Soilborne Diseases Symposium, 14-17 November 2016, Lincoln University, Canterbury, New Zealand.
- Naidu, Y., Idris, A. S., Madihah, A. Z., & Kamarudin, N. (2016a). In vitro Antagonistic Interactions Between Endophytic Basidiomycetes of Oil Palm (*Elaeis guineensis*) and *Ganoderma boninense*. *Journal of Phytopathology*, 164(10), 779-790.
- Nannipieri, P., & Badalucco, L. (2003). Biological processes. In: Processes in the Soil–Plant System: Modelling Concepts and Applications (eds D.K. Bembli & R. Nieder). The Haworth Press, Binghamton, NY.
- Nannipieri, P., Ascher-Jenull, J., Ceccherini, M. T., Giagnoni, L., Pietramellara, G., & Renella, G. (2017). Nannipieri, P., Ascher, J., Ceccherini, MT, Landi, L., Pietramellara, G. & Renella, G. 2003. Microbial diversity and soil functions. *European Journal of Soil Science*, 54, 655–670. *European Journal of Soil Science*, 68(1), 2-5.
- Nawawi, A., & Ho, Y. W. (1990). Effect of temperature and pH on growth pattern of *Ganoderma boninense* from oil palm in peninsular Malaysia. *Pertanika*, 13(3), 303-307.

- Nemergut, D. R., Cleveland, C. C., Wieder, W. R., Washenberger, C. L., & Townsend, A. R. (2010). Plot-scale manipulations of organic matter inputs to soils correlate with shifts in microbial community composition in a lowland tropical rain forest. *Soil Biology and Biochemistry*, 42(12), 2153-2160.
- Nguyen, C. (2003). Rhizodeposition of organic C by plants: mechanisms and controls. *Agronomie*, 23(5-6), 375-396.
- Nicol, G. W., Leininger, S., Schleper, C., & Prosser, J. I. (2008). The influence of soil pH on the diversity, abundance and transcriptional activity of ammonia oxidizing archaea and bacteria. *Environmental microbiology*, 10(11), 2966-2978.
- Nielsen, U. N., Ayres, E., Wall, D. H., & Bardgett, R. D. (2011). Soil biodiversity and carbon cycling: a review and synthesis of studies examining diversity–function relationships. *European Journal of Soil Science*, 62(1), 105-116.
- Ntoko, F.A. (2013). Soil microbial diversity of a managed forest ecosystem and the potential for lignocellulose degradation and metal bioaccumulation by white rot fungi. Ph.D. dissertation, Alabama A&M University, Normal, Alabama 35762
- Nusaibah, S. A., Akmar, A. S. N., Idris, A. S., Sariah, M., & Pauzi, Z. M. (2016). Involvement of metabolites in early defense mechanism of oil palm (*Elaeis guineensis* Jacq.) against *Ganoderma* disease. *Plant Physiology and Biochemistry*, 109, 156-165.
- Nusaibah, S. A., Latiffah, Z., & Hassaan, A. R. (2011). ITS-PCR-RFLP analysis of *Ganoderma* sp. infecting industrial crops. *Pertanika J. Trop. Agric. Sci*, 34(1), 83-91.
- Olf, H., Hoorens, B., De Goede, R. G. M., Van der Putten, W. H., & Gleichman, J. M. (2000). Small-scale shifting mosaics of two dominant grassland species: the possible role of soil-borne pathogens. *Oecologia*, 125(1), 45-54.
- Onwuka, M. I., Ozurumba, U. V., & Nkwocha, O. S. (2016). Changes in Soil pH and Exchangeable Acidity of Selected Parent Materials as Influenced by Amendments in South East of Nigeria. *Journal of Geoscience and Environment Protection*, 4(05), 80.
- Orgiazzi, A., Lumini, E., Nilsson, R. H., Girlanda, M., Vizzini, A., Bonfante, P., & Bianciotto, V. (2012). Unravelling soil fungal communities from different Mediterranean land-use backgrounds. *PloS one*, 7(4), e34847.
- Osono, T., & Takeda, H. (1999). A methodological survey on incubation of fungi on leaf litter of *Fagus crenata*. *Applied Forest Science Kansai* 8, 103–108. In Japanese, with English abstract.
- Pal, K. K., & Gardener, B. M. (2006). Biological control of plant pathogens. *The plant health instructor*, 2, 1117-1142.

- Palti, J. (2012). *Cultural practices and infectious crop diseases*(Vol. 9). Springer Science & Business Media.
- Paramanathan, S. (2003). In Fairhurst, T.H., and R. Hardter (Eds.), *Managing oil palm for large and sustainable yields*. PPI/PPIC-IPI, Singapore. p. 27-57.
- Paredes, S. H., & Lebeis, S. L. (2016). Giving back to the community: microbial mechanisms of plant–soil interactions. *Functional Ecology*, 7(30), 1043-1052.
- Parfitt, R. L., Giltrap, D. J., & Whitton, J. S. (1995). Contribution of organic matter and clay minerals to the cation exchange capacity of soils. *Communications in Soil Science & Plant Analysis*, 26(9-10), 1343-1355.
- Parthiban, K., Vanitah, R., Jusoff, K., Nordiana, A.A., Anuar, A.R., Wahid, O., & Hamdan, A.B. (2016). GIS Mapping of Basal Stem Rot Disease in Relation to Soil Series Among Oil Palm Smallholders. *American Journal of Agricultural and Biological Sciences*, 11 (1), 2.12.
- Paterson, R. R. M. (2007). Ganoderma disease of oil palm—A white rot perspective necessary for integrated control. *Crop protection*, 26(9), 1369-1376.
- Paul, V. G., Wronkiewicz, D. J., & Mormile, M. R. (2017). Impact of elevated CO₂ concentrations on carbonate mineral precipitation ability of sulfate-reducing bacteria and implications for CO₂ sequestration. *Applied Geochemistry*, 78, 250-271.
- Pearce, D. A. (2008). Climate change and the microbiology of the Antarctic Peninsula region. *Science progress*, 91(2), 203-217.
- Pennanen, T., Fritze, H., Vanhala, P., Kiikkilä, O., Neuvonen, S., & Bååth, E. (1998). Structure of a microbial community in soil after prolonged addition of low levels of simulated acid rain. *Applied and Environmental Microbiology*, 64(6), 2173-2180.
- Philippot, L., Piutti, S., Martin-Laurent, F., Hallet, S., & Germon, J. C. (2002). Molecular analysis of the nitrate-reducing community from unplanted and maize-planted soils. *Applied and Environmental Microbiology*, 68(12), 6121-6128.
- Pilotti, C. A. (2005). Stem rots of oil palm caused by *Ganoderma boninense*: Pathogen biology and epidemiology. *Mycopathologia*, 159(1), 129-137.
- Pilotti, C. A., Sanderson, F. R., & Aitken, E. A. (2002). Sexuality and interactions of monokaryotic and dikaryotic mycelia of *Ganoderma boninense*. *Mycological Research*, 106(11), 1315-1322.
- Pilotti, C. A., Sanderson, F. R., & Aitken, E. A. B. (2003). Genetic structure of a population of *Ganoderma boninense* on oil palm. *Plant Pathology*, 52(4), 455-463.

- Ploetz, R. C. (2000). Panama disease: a classic and destructive disease of banana. *Plant Health Progress*, *10*, 1-7.
- Poggio, S. L., Chaneton, E. J., & Ghersa, C. M. (2010). Landscape complexity differentially affects alpha, beta, and gamma diversities of plants occurring in fencerows and crop fields. *Biological Conservation*, *143*(11), 2477-2486.
- Ponstein, A. S., Bres-Vloemans, S. A., Sela-Buurlage, M. B., van den Elzen, P. J., Melchers, L. S., & Cornelissen, B. J. (1994). A novel pathogen-and wound-inducible tobacco (*Nicotiana tabacum*) protein with antifungal activity. *Plant physiology*, *104*(1), 109-118.
- Postma, J., Schilder, M. T., Bloem, J., & van Leeuwen-Haagsma, W. K. (2008). Soil suppressiveness and functional diversity of the soil microflora in organic farming systems. *Soil Biology and Biochemistry*, *40*(9), 2394-2406.
- Raaijmakers, J. M. (2016). Going back to the roots: microbiology and chemistry at the plant-soil interface. 14–17. In: Proceedings of the 9th Australasian Soilborne Diseases Symposium, Lincoln University, Christchurch, New Zealand.
- Rai, R. D. (2003). Successful cultivation of the medicinal mushroom Reishi, *Ganoderma lucidum* in India. *Mushroom Research*, *12*(2).
- Ramli, N. R., Mohamed, M. S., Seman, I. A., Zairun, M. A., & Mohamad, N. (2016). The potential of endophytic bacteria as a biological control agent for ganoderma disease in oil palm. *Sains Malays*, *45*, 401-409.
- Rao, A. K. (1990). Basal stem rot (*Ganoderma*) in oil palm smallholdings. In *Proceedings of the Ganoderma Workshop, Palm Oil Research Institute of Malaysia* (pp. 113-131).
- Rashidi, M., & Seilsepour, M. (2008). Modeling of soil cation exchange capacity based on soil organic carbon. *ARPN Journal of Agricultural and Biological Science*, *3*(4), 41-45.
- Rees, R. W., Flood, J., Hasan, Y., & Cooper, R. M. (2007). Effects of inoculum potential, shading and soil temperature on root infection of oil palm seedlings by the basal stem rot pathogen *Ganoderma boninense*. *Plant Pathology*, *56*(5), 862-870.
- Rees, R. W., Flood, J., Hasan, Y., Potter, U., & Cooper, R. M. (2009). Basal stem rot of oil palm (*Elaeis guineensis*); mode of root infection and lower stem invasion by *Ganoderma boninense*. *Plant pathology*, *58*(5), 982-989.
- Rees, R. W., Flood, J., Hasan, Y., Wills, M. A., & Cooper, R. M. (2012). *Ganoderma boninense* basidiospores in oil palm plantations: evaluation of their possible role in stem rots of *Elaeis guineensis*. *Plant pathology*, *61*(3), 567-578.

- Rees, R. W., Flood, J., Hasan, Y., Wills, M. A., & Cooper, R. M. (2012). *Ganoderma boninense* basidiospores in oil palm plantations: evaluation of their possible role in stem rots of *Elaeis guineensis*. *Plant pathology*, 61(3), 567-578.
- Richardson, A. E. (1994). Soil microorganisms and phosphorus availability. *Soil Biota*, 50, 35-39.
- Rolph, H., Wijesekara, R., Lardner, R., Abdullah, F., Kirk, P. M., Holderness, M., & Flood, J. (2000). Molecular variation in *Ganoderma* isolates from oil palm, coconut, and betelnut. *Ganoderma diseases of perennial crops*, 205-221.
- Rosenani, A. B., Rovica, R., Cheah, P. M., & Lim, C. T. (2016). Growth Performance and Nutrient Uptake of Oil Palm Seedling in Prenursery Stage as Influenced by Oil Palm Waste Compost in Growing Media. *International Journal of Agronomy*, 2016.
- Rosenzweig, M. L. (1995). Species diversity in space and time. Cambridge University Press.
- Roslan, A., & Idris, A. S. (2012). Economic impact of *Ganoderma* incidence on Malaysian oil palm plantation—a case study in Johor. *Oil Palm Industry Economic Journal*, 12(1), 24-30.
- Roslan, I., Shamshuddin, J., Fauziah, C. I., & Anuar, A. R. (2010). Occurrence and properties of soils on sandy beach ridges in the Kelantan–Terengganu Plains, Peninsular Malaysia. *Catena*, 83(1), 55-63.
- Rousk, J., Brookes, P. C., & Bååth, E. (2010). Investigating the mechanisms for the opposing pH relationships of fungal and bacterial growth in soil. *Soil Biology and Biochemistry*, 42(6), 926-934.
- Rousk, J., Demoling, L. A., Bahr, A., & Bååth, E. (2008). Examining the fungal and bacterial niche overlap using selective inhibitors in soil. *FEMS Microbiology Ecology*, 63(3), 350-358.
- Rowell, D. L. (2014). Soil science: Methods & applications. Routledge.
- Rowell, D.L. (1994). Soil Science: Methods and applications. Longman Scientific and Technical, UK.
- Rubin, M. A., & Leff, L. G. (2007). Nutrients and other abiotic factors affecting bacterial communities in an Ohio River (USA). *Microbial Ecology*, 54(2), 374-383.
- Ruiz-Lozano, J. M., & Azcón, R. (1993). Specificity and functional compatibility of VA mycorrhizal endophytes in association with *Bradyrhizobium* strains in *Cicer arietinum*. *Symbiosis*, 15(3), 217-226.

- Saha, A., Mandal, P., Dasgupta, S., & Saha, D. (2008). Influence of culture media and environmental factors on mycelial growth and sporulation of *Lasiodiplodia theobromae* (Pat.) Griffon and Maubl. *Journal of Environmental Biology*, 29(3), 407.
- Sanderson, F. R. (2005). An insight into spore dispersal of *Ganoderma boninense* on oil palm. *Mycopathologia*, 159(1), 139-141.
- Sanderson, F. R., Pilotti, C. A., & Bridge, P. D. (2000). Basidiospores: their influence on our thinking regarding a control strategy for basal stem rot of oil palm. *Ganoderma diseases of perennial crops*, 113-119.
- Santos, P., Fortunato, A., Ribeiro, A., & Pawlowski, K. (2008). Chitinases in root nodules. *Plant Biotechnology*, 25(3), 299-307.
- Sapak, Z., Meon, S. A. R. I. A. H., & Ahmad, Z. A. M. (2008). Effect of endophytic bacteria on growth and suppression of *Ganoderma* infection in oil palm. *International Journal of Agriculture and Biology*, 10, 127-132.
- Sariah, M., & Zakaria, H. (2000). The use of soil amendments for the control of basal stem rot of oil-palm seedlings. *Ganoderma diseases of perennial crops*, 89-99.
- Sariah, M., Hussin, M. Z., Miller, R. N. G., & Holderness, M. (1994). Pathogenicity of *Ganoderma boninense* tested by inoculation of oil palm seedlings. *Plant Pathology*, 43(3), 507-510.
- Sariah, M., Joseph, H., & Zakaria, H. (1997). Suppression of basal stem rot (BSR) of oil palm seedlings by calcium nitrate. *Planter*, 73(856), 359-361.
- Saxena, R. K., Sangeetha, L., Vohra, A., Gupta, R., & Gulati, R. (2001). Induction and mass sporulation in lignin degrading fungus *Ceriporiopsis subvermispora* for its potential usage in pulp and paper industry. *Current Science*, 591-594.
- Schmidt, T. M., & Waldron, C. (2015). Microbial diversity in soils of agricultural landscapes and its relation to ecosystem function. *The ecology of agricultural landscapes: long-term research on the path to sustainability*. Oxford University Press, New York, 135-157.
- Seo, G. S., & Kirk, P. M. (2000). Ganodermataceae: nomenclature and classification. *Ganoderma diseases of perennial crops*, 3-22.
- Sewards, J. (2014). Soil pH and its Effect on nutrient availability, soil microbial activity and plant health. UF/IFAS Extension, Volusia County, Florida.
- Shahnavaz, B., & Geremia, R. A. (2012). The effect of pH and temperature on spatial variation of Acidobacteria/Actinobacteria communities from Alpine soil. *Journal of Cell Molecular. Research*, 4, 28-33.

- Shamshuddin, J., Fauziah, C. I., Anda, M., Kapok, J., & Shazana, M. A. R. S. (2011). Using ground basalt and/or organic fertilizer to enhance productivity of acid soils in Malaysia for crop production. *Malaysian Journal of Soil Science*, 15, 127-146.
- Sharma, S., Kumar, V., & Tripathi, R. B. (2017). Isolation of phosphate solubilizing microorganism (PSMs) from soil. *Journal of microbiology and Biotechnology Research*, 1(2), 90-95.
- Sharp, C. E., Urschel, S., Dong, X., Brady, A. L., Slater, G. F., & Strous, M. (2017). Robust, high-productivity phototrophic carbon capture at high pH and alkalinity using natural microbial communities. *Biotechnology for Biofuels*, 10(1), 84.
- Sheng, Y., Bibby, K., Grettenberger, C., Kaley, B., Macalady, J. L., Wang, G., & Burgos, W. D. (2016). Geochemical and temporal influences on the enrichment of acidophilic iron-oxidizing bacterial communities. *Applied and environmental microbiology*, 82(12), 3611-3621.
- Shila B., & Siti R. A. A. (2016). Microbial Biodiversity Of Mineral Soils Planted With Young Oil Palms in Belaga, Sarawak. *Soil Science Conference of Malaysia*.
- Shu'ud, M. M., Loonis, P., & Seman, I. A. (2007). Towards automatic recognition and grading of *Ganoderma* infection pattern using fuzzy systems. *T Eng Comput Technol*, 19.
- Singh, B. K., Dawson, L. A., Macdonald, C. A., & Buckland, S. M. (2009). Impact of biotic and abiotic interaction on soil microbial communities and functions: A field study. *applied soil ecology*, 41(3), 239-248.
- Singh, G. (1990). *Ganoderma*- The scourge of oil palm in the coastal areas. pp. 7–35. In: Ariffin, D. and S. Jalani (eds.) *Proc. Ganoderma Workshop*, Bangi: Palm Oil Research Institute of Malaysia.
- Singh, G. (1991). *Ganoderma* - The scourge of oil palm in the coastal areas. *The Planter*, 67 (786), 421 444.
- Sinsabaugh, R. L., Antibus, R. K., & Linkins, A. E. (1991). An enzymic approach to the analysis of microbial activity during plant litter decomposition. *Agriculture, Ecosystems & Environment*, 34(1-4), 43-54.
- Soil Conservation Service (1984). Procedures for collecting soil samples and methods of analysis for soil survey. USDA-SCS, Soil Survey Investigations Report No. 1. GPO, Washington, DC.
- Song, W. Y., Zhang, Z. B., Shao, H. B., Guo, X. L., Cao, H. X., Zhao, H. B., ... & Hu, X. J. (2008). Relationship between calcium decoding elements and plant abiotic-stress resistance. *International journal of biological sciences*, 4(2), 116.

- Spiers, J. M. (1984). Influence of lime and sulfur soil additions on growth, yield, and leaf nutrient content of rabbiteye blueberry. *Journal of the American Society for Horticultural Science*, 109(4), 559-562.
- Steinauer, K., Tilman, D., Wragg, P. D., Cesarz, S., Cowles, J. M., Pritsch, K., ... & Eisenhauer, N. (2015). Plant diversity effects on soil microbial functions and enzymes are stronger than warming in a grassland experiment. *Ecology*, 96(1), 99-112.
- Stemmer, M., Watzinger, A., Blochberger, K., Haberhauer, G., & Gerzabek, M. H. (2007). Linking dynamics of soil microbial phospholipid fatty acids to carbon mineralization in a ¹³C natural abundance experiment: impact of heavy metals and acid rain. *Soil Biology and Biochemistry*, 39(12), 3177-3186.
- Supramaniam, C. V. (2016). *Molecular interaction between Ganoderma boninense and young oil palm* (Doctoral dissertation, University of Nottingham).
- Suryanto, D., Wibowo, R. H., Siregar, E. B. M., & Munir, E. (2012). A possibility of chitinolytic bacteria utilization to control basal stems disease caused by *Ganoderma boninense* in oil palm seedling. *African Journal of Microbiology Research*, 6(9), 2053-2059.
- Susanto, A. (2009, November). Basal stem rot in Indonesia. Biology, economic importance, epidemiology, detection and control. In *Proceedings of International Workshop on Awareness, Detection and Control of Oil Palm Devastating Diseases November*.
- Susanto, A., Sudharto, P. S., & Purba, R. Y. (2005). Enhancing biological control of basal stem rot disease (*Ganoderma boninense*) in oil palm plantations. *Mycopathologia*, 159(1), 153-157.
- Swier, H., Dkhar, M. S., & Kayang, H. (2011). Fungal population and diversity in organically amended agricultural soils of Meghalaya, India. *Journal of Organic Systems*, 6(2), 3-12.
- Sylvia, D. M., Fuhrmann, J. J., Hartel, P. G., & Zuberer, D. A. (Eds.). (2005). *Principles and applications of soil microbiology* (No. QR111 S674 2005). Upper Saddle River, NJ:: Pearson Prentice Hall.
- Tajudin, N. S., Hanafi, M. M., Idris, A. S., & Balasundram, S. K. (2016). Determination and mapping of calcium and magnesium contents using geostatistical techniques in oil palm plantation related to basal stem rot disease. *Songklanakarin Journal of Science & Technology*, 38(1).
- Takka, F. A. S. (1999). Calcium alginate microparticles for oral administration: II effect of formulation factors on drug release and drug entrapment efficiency. *Journal of microencapsulation*, 16(3), 291-301.
- Tank, J. L., Reisinger, A. J., & Rosi, E. J. (2017). Nutrient limitation and uptake. *Methods in Stream Ecology: Volume 2: Ecosystem Function*, 147.

- Tarafdar, J. C., & Jungk, A. (1987). Phosphatase activity in the rhizosphere and its relation to the depletion of soil organic phosphorus. *Biology and Fertility of Soils*, 3(4), 199-204.
- Thomas, G. W. (1996). Soil pH and soil acidity. *Methods of Soil Analysis Part 3—Chemical Methods*, (methodsofsoilan3), 475-490.
- Torsvik, V., Daae, F. L., Sandaa, R. A., & Øvreås, L. (1998). Novel techniques for analysing microbial diversity in natural and perturbed environments. *Journal of biotechnology*, 64(1), 53-62.
- Tripathi, B. M., Lee-Cruz, L., Kim, M., Singh, D., Go, R., Shukor, N. A., ... & Adams, J. M. (2014). Spatial scaling effects on soil bacterial communities in Malaysian tropical forests. *Microbial ecology*, 68(2), 247-258.
- Trivedi, P., Delgado-Baquerizo, M., Anderson, I. C., & Singh, B. K. (2016). Response of soil properties and microbial communities to agriculture: implications for primary productivity and soil health indicators. *Frontiers in plant science*, 7.
- Trumbore, S. (2000). Age of soil organic matter and soil respiration: radiocarbon constraints on belowground C dynamics. *Ecological Applications*, 10(2), 399-411.
- Tuomisto, H. (2010). A diversity of beta diversities: straightening up a concept gone awry. Part 1. Defining beta diversity as a function of alpha and gamma diversity. *Ecography*, 33(1), 2-22.
- Turner, P. D. (1965). Oil palm and ganoderma III: Treatment and control in established plantings. *The Planter*, 41, 279-282.
- Turner, P.D. (1981). *Planters Oil palm diseases and disorders*. Published for the Incorporated Society of Planters by Oxford University Press, Kuala Lumpur, pp. 280.
- Uchida, R. (2000). Essential nutrients for plant growth: nutrient functions and deficiency symptoms. *Plant nutrient management in Hawaii's soils*, 31-55.
- UKNCC (1998). *Growth and Media Manuals*. Strain databases (www.ukncc.co.uk/).
- USDA (2010). *Indonesia: Rising Global Demand Fuels Palm Oil Expansion*. United States Department of Agriculture.
- USDA (2012). *Malaysia: Stagnating Palm Oil Yields Impede Growth*. Commodity Intelligence Report <http://www.pecad.fas.usda.gov/highlights/2012/12/Malaysia/> (accessed on October, 2016).
- USDA (2014). *Foreign Agricultural Services*. <http://www.schusterinstituteinvestigations.org/palm-fruit-to-product>

- Vale, M., Nguyen, C., Dambrine, E., & Dupouey, J. L. (2005). Microbial activity in the rhizosphere soil of six herbaceous species cultivated in a greenhouse is correlated with shoot biomass and root C concentrations. *Soil Biology and Biochemistry*, 37(12), 2329-2333.
- Van Der Heijden, M. G., Bardgett, R. D., & Van Straalen, N. M. (2008). The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology letters*, 11(3), 296-310.
- Van Overbeek, L., & Van Elsas, J. D. (2008). Effects of plant genotype and growth stage on the structure of bacterial communities associated with potato (*Solanum tuberosum* L.). *FEMS microbiology ecology*, 64(2), 283-296.
- Van Ranst, E., Verloo, M., Demeyer, A., & Pauwels, J.M. (1999). Manual for the soil chemistry and fertility laboratory. Analytical methods for soils and plants equipment and management of consumables. Ghent University, Ghent, Belgium, 243 pp.
- Van Soest, P. V., Robertson, J. B., & Lewis, B. A. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of dairy science*, 74(10), 3583-3597.
- Veena, S. S., & Pandey, M. (2006). Evaluation of the locally available substrates for the cultivation of indigenous *Ganoderma* isolates. *Journal Mycology & Plant Pathology*. 36(3), 434-438.
- Virdiana, I., Hasan, Y., Aditya, R., & Flood, J. (2010). Testing the effects of oil palm replanting practices (windrowing, fallowing and poisoning) on incidence of *Ganoderma*. *Proc IOPC*.
- Vogeler, I., Rogasik, J., Funder, U., Panten, K., & Schnug, E. (2009). Effect of tillage systems and P-fertilization on soil physical and chemical properties, crop yield and nutrient uptake. *Soil and Tillage Research*, 103(1), 137-143.
- Volpin, H., & Elad, Y. (1991). Influence of calcium nutrition on susceptibility of rose flowers to Botrytis blight. *Phytopathology*, 81(11), 1390-1394.
- Von Uexkull H. R., & Fairhurst T.H. (1991). IPI Bulletin No.12, International Potash Institute, Bern, Switzerland. 79p.
- Wahid, M.B., Akmar-Abdullah, S.N., & Henson, I.E. (2005). Oil palm -achievements and potential. *Plant Production Science*, (8), 288-297.
- Wakelin, S.A., Gerard, E., van Koten, C., Banabas, M., O'Callaghan, M., & Nelson, P.N. (2016). Soil physicochemical properties impact more strongly on bacteria and fungi than conversion of grassland to oil palm. *Pedobiologia*, 59 (3),83-91.
- Walker, J. C., 1972. Plant Pathology. New York: McGraw-Hill.

- Wardle, D. A. (1992). A comparative assessment of factors which influence microbial biomass carbon and nitrogen levels in soil. *Biological reviews*, 67(3), 321-358.
- Wardle, D. A., Bardgett, R. D., Klironomos, J. N., Setälä, H., Van Der Putten, W. H., & Wall, D. H. (2004). Ecological linkages between aboveground and belowground biota. *Science*, 304(5677), 1629-1633.
- Wardle, D. A., Bardgett, R. D., Klironomos, J. N., Setälä, H., Van Der Putten, W. H., & Wall, D. H. (2004). Ecological linkages between aboveground and belowground biota. *Science*, 304(5677), 1629-1633.
- Weil, R. R., Brady, N. C., & Weil, R. R. (2016). *The nature and properties of soils*. Pearson.
- Weller, D. M., Raaijmakers, J. M., Gardener, B. B. M., & Thomashow, L. S. (2002). Microbial populations responsible for specific soil suppressiveness to plant pathogens. *Annual review of phytopathology*, 40(1), 309-348.
- Wiedenhoeft, A. C. (2006). *Plant nutrition*. Infobase Publishing.
- Williamson, W. M., Wardle, D. A., & Yeates, G. W. (2005). Changes in soil microbial and nematode communities during ecosystem decline across a long-term chronosequence. *Soil Biology and Biochemistry*, 37(7), 1289-1301.
- Wollum, A. G. (1982). Cultural methods for soil microorganisms. *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties*, (methodsofsoilan2), 781-802.
- Wong, L., Bong, C. F. J., & Idris, A. S. (2012). Ganoderma species associated with basal stem rot disease of oil palm. *American Journal of Applied Sciences*, 9(6), 879-885.
- World Bank (2010). *World Development Report 2010: Development and Climate Change*. Washington DC.
- Wubet, T., Christ, S., Schöning, I., Boch, S., Gawlich, M., Schnabel, B., ... & Buscot, F. (2012). Differences in soil fungal communities between European beech (*Fagus sylvatica* L.) dominated forests are related to soil and understory vegetation. *PloS one*, 7(10), e47500.
- Xu, J., Fan, Y., & Li, Z. (2016). Effect of pH on elemental sulfur conversion and microbial communities by autotrophic simultaneous desulfurization and denitrification. *Environmental technology*, 37(23), 3014-3023.
- Xu, L., Ravnskov, S., Larsen, J., & Nicolaisen, M. (2012). Linking fungal communities in roots, rhizosphere, and soil to the health status of *Pisum sativum*. *FEMS microbiology ecology*, 82(3), 736-745.

- Xue, K., Wu, L., Deng, Y., He, Z., Van Nostrand, J., Robertson, P. G., ... & Zhou, J. (2013). Functional gene differences in soil microbial communities from conventional, low-input, and organic farmlands. *Applied and environmental microbiology*, 79(4), 1284-1292.
- Yannarell, A. C., & Triplett, E. W. (2005). Geographic and environmental sources of variation in lake bacterial community composition. *Applied and environmental microbiology*, 71(1), 227-239.
- Yao, H., Gao, Y., Nicol, G. W., Campbell, C. D., Prosser, J. I., Zhang, L., ... & Singh, B. K. (2011). Links between ammonia oxidizer community structure, abundance, and nitrification potential in acidic soils. *Applied and environmental microbiology*, 77(13), 4618-4625.
- You, Y., Wang, J., Huang, X., Tang, Z., Liu, S., & Sun, O. J. (2014). Relating microbial community structure to functioning in forest soil organic carbon transformation and turnover. *Ecology and evolution*, 4(5), 633-647.
- Youssef, N. H., Couger, M. B., McCully, A. L., Criado, A. E. G., & Elshahed, M. S. (2015). Assessing the global phylum level diversity within the bacterial domain: a review. *Journal of advanced research*, 6(3), 269-282.
- Zaiton, S., Sariah, M., & Zainal, A. (2006). Isolation and characterization of microbial endophytes from oil palm roots: implication as biological control agents against Ganoderma. *Planter*, 82(966), 587-597.
- Zak, D. R., Groffman, P. M., Pregitzer, K. S., Christensen, S., & Tiedje, J. M. (1990). The vernal dam: plant-microbe competition for nitrogen in northern hardwood forests. *Ecology*, 71(2), 651-656.
- Zak, D. R., Pregitzer, K. S., King, J. S., & Holmes, W. E. (2000). Elevated atmospheric CO₂, fine roots and the response of soil microorganisms: a review and hypothesis. *The New Phytologist*, 147(1), 201-222.
- Zhae, S., & Simon, F. S. (2006). Effect of culture media, temperature, pH and bio-herbicidal efficacy of *Phoma exigua*, a potential biological control for salal (*Gaultheria shallon*). *Biocontrol Science and Technology*, 6, 1043-1055.
- Zhang, J., Sun, Q. L., Zeng, Z. G., Chen, S., & Sun, L. (2015). Microbial diversity in the deep-sea sediments of Iheya North and Iheya Ridge, Okinawa Trough. *Microbiological research*, 177, 43-52.
- Zhou, K., Binkley, D., Doxtader, K.G., 1992. A new method for estimating gross phosphorus mineralization and immobilization rates in soils. *Plant Soil* 147, 243-250.

LIST OF PUBLICATIONS

Khunaw, A.R., Radziah, O., Nusaibah .S.A., Mohamed Hanafi. M., Rabara.F.S. (2017). Role of Microbial Community in Suppressing Development of Ganoderma in Oil Palm Seedlings. International Journal of Plant & Soil Science.Applied.[http://www.sciencedomain.org/review-19\(1\):1-14,2017](http://www.sciencedomain.org/review-19(1):1-14,2017); Article no. IJPSS.36192. ISSN: 2320-7035.

Khunaw, A.R., Radziah, O., Nusaibah .S.A., Mohamed Hanafi. M. (under review). Effect of pH on growth of oil palm seedlings and basal stem rot disease incidence in Ganoderma inoculated plants. The Indian Journal of Agricultural Sciences. (Manuscript ID: 71037).





UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : _____

TITLE OF THESIS / PROJECT REPORT :

INFLUENCE OF SOIL pH ON MICROBIAL COMMUNITY AND INCIDENCE OF BASAL STEM ROT (BSR) DISEASE DEVELOPMENT IN OIL PALM SEEDLINGS PLANTED ON BSR AFFECTED, UNAFFECTED AND FOREST SOILS

NAME OF STUDENT: KHUNAW ABDULLA RAHMAN

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (v)

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

OPEN ACCESS

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

PATENT

Embargo from _____ until _____
(date) (date)

Approved by:

(Signature of Student)
New IC No/ Passport No.:

(Signature of Chairman of Supervisory Committee)
Name:

Date :

Date :

[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted.]