



UNIVERSITI PUTRA MALAYSIA

***INDOOR AIR QUALITY AND RESPIRATORY HEALTH IMPLICATION
AMONG MALAY PRE-SCHOOL CHILDREN IN PUCHONG AND HULU-
LANGAT, SELANGOR, MALAYSIA***

ONWUSEREAKA CYNTHIA OLUCHI

FPSK(m) 2018 6



**INDOOR AIR QUALITY AND RESPIRATORY HEALTH IMPLICATION
AMONG MALAY PRE-SCHOOL CHILDREN IN PUCHONG AND
HULU- LANGAT, SELANGOR, MALAYSIA**

By

ONWUSEREAKA CYNTHIA OLUCHI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Master of Science**

December 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

**INDOOR AIR QUALITY AND RESPIRATORY HEALTH IMPLICATION
AMONG MALAY PRE-SCHOOL CHILDREN IN PUCHONG AND
HULU- LANGAT, SELANGOR, MALAYSIA**

By

ONWUSEREAKA CYNTHIA OLUCHI

December 2017

Chairman : Associate Professor Juliana Jalaludin, PhD
Faculty : Medicine and Health Sciences

Background: Indoor air quality (IAQ) refers to the nature of air that circulates throughout the space where we live. Indoor air pollutant in the environment has caused a greater impact on the health of occupants' especially preschool age who are still growing. The increased of respiratory symptoms in children have been linked to the exposure to indoor air pollutants. Children were exposed to indoor air pollutants because they spend most of their time indoors, either in school or at home. **Objective:** This study aimed to determine the association between indoor air pollutants (PM_{2.5}, PM₁₀, VOCs, temperature, mold and relative humidity) and respiratory implications among Malay preschool children in selected areas in Selangor. **Methodology:** A cross sectional comparative study design was used on healthy preschools children in Puchong (n=135), as the study group and Hulu Langat (n=135) as the comparative group. Preschool children in this study were selected based on inclusion criteria such as; those who are between the age of 5 and 6 years old, healthy children, those that have been in the school for at least 6 months and Malays. The respiratory symptoms were assessed using questionnaire adapted from American Thoracic Society. Indoor air quality monitoring was conducted using the following equipment; DustTrak Aerosol monitor for particulates, PbbRAE portable VOC Monitor (pbbRAE 3000) for VOCs, Q-Trak plus Model 8554 Monitor for temperature and relative humidity, TSI VelociCalc Plus Model 8386 for air velocity; Pbi DuoSAS Super 360 and Sabroud Dextrous Agar for sampling and identification of mold. Chestgraph HI-101 spirometer for lung function test. **Results:** Statistical analysis shows a significant higher indoor levels of PM_{2.5}, PM₁₀ and mold in the study area except relative humidity which was higher in comparative area (p = 0.049). Besides, indoor home exposure was also assessed, analysis showed that homes of preschool children in the study area had significantly higher VOCs, Temperature, mold, PM_{2.5}, and PM₁₀, except relative humidity which was significantly higher in the comparative area. There was a

significant difference in FEV₁/FVC% (p= 0.001) and FEV₁% (p=0.008) predicted among the study and comparative group. A significant association was found between the indoor air pollutant levels of PM_{2.5} (χ^2 = 15.008, p=<0.001), PM₁₀ (χ^2 = 8.816, p=0.003) and cough. Exposure to indoor PM₁₀ (χ^2 = 6.842, p=0.009) and VOCs (χ^2 = 12.153, p=<0.001) were significantly associated with FVC%. VOCs was significantly associated with FEV₁% (χ^2 = 14.382, p=<0.001). Mold isolated from preschools in study and comparative area include; *Candida albican*, *Aspergillus niger*, *Microsporium canis*, *Penicillium spp.* Logistics regression was used to determine the main factor that influenced the respiratory symptom and the abnormality of lung function after all confounders have been controlled. Analysis showed that the risk of FEV₁% abnormality increased significantly with increased in the level of PM₁₀ (AOR=2.1, 95% CI= 2.509-8.221), VOC (AOR= 5.3, 95% CI= 1.912-14.835) and RH (AOR=14.3, 95% CI= 1.451-14.306). **Conclusion:** This study found a higher exposure to indoor air pollutants which may increase the respiratory symptoms and reduced lung function among preschool children in the exposed area. It is recommended that the building of the preschools should be located in places that is less expose to outdoor sources of indoor air pollutants. Mold growth was observed more in preschools that had moisture damaged building materials. Therefore, the moisture damaged building materials need to be replaced to avoid mold growth.

Keywords: Preschool children; indoor air quality; respiratory symptoms; mold.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**KUALITI UDARA DALAMAN DAN IMPLIKASI KESIHATAN
RESPIRATORI DALAM KALANGAN KANAK-KANAK PRASEKOLAH
MALAYA DI PUCHONG DAN HULU- LANGAT, SELANGOR, MALAYSIA**

Oleh

ONWUSEREAKA CYNTHIA OLUCHI

Disember 2017

Pengerusi : Profesor Madya Juliana Jalaludin, PhD
Fakulti : Perubatan dan Sains Kesihatan

Latar belakang: Peningkatan simptom respiratori dalam kalangan kanak-kanak telah dikaitkan dengan pendedahan kepada pencemar udara dalaman. Kanak-kanak terdedah kepada pencemar udara dalaman kerana mereka menghabiskan kebanyakan masa mereka di kawasan tertutup, sama ada di sekolah atau di rumah.

Objektif: Kajian ini bertujuan untuk menentukan kaitan antara pencemar udara dalaman ($PM_{2.5}$, PM_{10} , sebatian organik mudah meruap, suhu, kewujudan kulat, humiditi relatif) dan implikasinya terhadap kesihatan respiratori dalam kalangan kanak-kanak prasekolah malaya di Selangor. **Metodologi:** Kajian keratan rentas perbandingan lah dijalankan ke atas kanak-kanak prasekolah yang sihat di Puchong ($n=135$), sebagai kawasan kajian dan di Hulu Langat ($n=135$) sebagai kawasan perbandingan. Kanak-kanak prasekolah dalam kajian ini dipilih berdasarkan beberapa kriteria inklusif seperti; mereka yang berumur antara 5 dan 6 tahun, sihat, mereka yang telah berada di sekolah sekurang-kurangnya 6 bulan dan Melayu. Simptom respiratori telah diperolehi menggunakan borang soal selidik yang diadaptasi daripada American Thoracic Society. Pemantauan kualiti udara dalaman telah dijalankan menggunakan peralatan berikut; DustTrak Aerosol untuk partikel terampai, pbbRAE 3000 untuk sebatian organik mudah meruap, Model 8554 plus Trak Q untuk suhu dan humiditi relatif, Model 8386 Plus TSI VelociCalc untuk velositi udara; Pbi DuoSAS Super 360 dan Agar-agar Dekstrous Sabroud untuk persampelan dan identifikasian kulat, Spirometer Chestgraph HI-101 untuk ujian fungsi paru-paru. **Hasil:** Analisis statistik menunjukkan pencemar udara dalaman secara signifikan lebih tinggi bagi $PM_{2.5}$, PM_{10} dan kulat dalam kajian ini, kecuali humiditi relatif yang didapati lebih tinggi dalam kawasan perbandingan ($p = 0.049$). Selain itu, paras pencemar dalaman di kediaman kanak-kanak prasekolah dalam kajian ini adalah lebih tinggi secara

signifikan bagi sebatian organik meruap, suhu, kulat, PM_{2.5}, dan PM₁₀, kecuali humidity relatif berbanding kumpulan perbandingan. Terdapat perbezaan yang signifikan dalam FEV₁/FVC% ($p = 0.001$) dan FEV₁% ($p = 0.008$) ramalan bagi kumpulan kajian dan perbandingan. Perkaitan yang signifikan didapati antara pencemar udara dalaman PM_{2.5}, ($\chi^2 = 15.008$, $p < 0.001$), PM₁₀ ($\chi^2 = 8.816$, $p = 0.003$) dan batuk. Pendedahan terhadap pencemar PM₁₀ ($\chi^2 = 6.842$, $p = 0.009$) dan sebatian organik meruap ($\chi^2 = 12.153$, $p < 0.001$) adalah berkaitan secara signifikan dengan peratusan FVC. Sebatian organik meruap adalah secara signifikan berkaitan dengan peratus FEV₁ ($\chi^2 = 14.382$, $p < 0.001$). Kulat yang diasingkan dari prasekolah dalam kawasan kajian dan perbandingan meliputi; *Candida albican*, *Aspergillus niger*, *Mikrosporium kanis*, *Penicillium spp*. Regresi logistik telah digunakan untuk menentukan faktor utama yang mempengaruhi simptom respiratori dan abnormaliti fungsi paru-paru selepas pengawalan kesemua faktor pembauran. Hasil analisis mendapati risiko abnormaliti FEV₁% meningkat secara signifikan dengan peningkatan paras PM₁₀ (AOR=2.1, 95% CI= 2.509-8.221), sebatian organik meruap (AOR= 5.3, 95% CI= 1.912-14.835) dan humiditi relatif (AOR=14.3, 95% CI= 1.451-14.306). **Kesimpulan:** Kajian ini mendapati pendedahan pencemar udara dalaman yang tinggi boleh meningkatkan simptom respiratori dan penurunan fungsi paru-paru dalam kalangan kanak-kanak prasekolah di kawasan terdedah. Adalah disarankan bahawa bangunan prasekolah perlulah ditempatkan di kawasan yang kurang terdedah daripada sumber luar pencemar udara dalaman. Pertumbuhan kulat telah banyak didapati di prasekolah yang mempunyai material dalaman bangunan yang lembap. Justeru itu, material bangunan yang rosak akibat kelembapan perlulah diganti bagi mengelakkan pertumbuhan kulat.

Kata kunci: Kanak-kanak prasekolah; kualiti udara tertutup; simptom respiratori; kulat.

ACKNOWLEDGEMENTS

My profound gratitude goes to the one who has the final say in my life. I want to thank God Almighty for his grace and favour all through my study period. His grace was sufficient for me.

My undiluted and sincere gratitude goes to my honourable supervisor, Assoc. Prof. Dr. Juliana Jalaludin. I want to thank you for being patient with me, for your advice and guidance all the way. Thank you so much for letting me share in your knowledge. I was able to complete my masters' degree today because of your guidance and I really appreciate all your efforts.

My special thanks goes to my co supervisor, Dr Titi Rahmawati, external examiner, internal examiner, my lecturers and staff in the department of Environmental Health. I want to also thank my friends who in one way or the other contributed immensely to the success of my data collection. Asrul, Akid, Faseeha, Shikin, Anis, and others I cannot thank you enough for your time and advice during my academic years. May God reward you abundantly.

My love and gratitude goes to my husband who was my pillar of support through my master's programme. Your love, support and encouragement kept me going. My gratitude goes to my lovely children king David and Doreen for your prayers. May God keep you all safe for me in Jesus name.

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Juliana binti Jalaludin, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Titi Rahmawati binti Hamedon, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No: Onwusereaka Cynthia Oluchi, GS39847

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: _____

Signature: _____
Name of Member
of Supervisory
Committee: _____

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF APPENDICES	xvii
LIST OF ABBREVIATIONS	xviii
 CHAPTER	
1 INTRODUCTION	1
1.1 Background	1
1.2 Problem Statement	3
1.3 Conceptual frameworks	4
1.4 Study Justification	6
1.5 Study Variable	7
1.5.1 Dependent variable	7
1.5.2 Independent variable	7
1.6 RESEARCH OBJECTIVES	8
1.6.1 General objective	8
1.6.2 Specific objective	8
1.7 Research Hypothesis	8
1.8 Definition of Variables	9
1.8.1 Conceptual definition	9
1.9 Operational Definition	11
 2 LITERATURE REVIEW	 13
2.1 Indoor Air Quality (IAQ)	13
2.2 Indoor Air Pollutants	13
2.3 Exposure to indoor air pollutants and their impact on health	14
2.4 Particulate matters (PM)	15
2.5 Volatile Organic Compounds (VOCs)	18
2.6 Indoor Mold	19
2.7 Sources of indoor air pollutants in indoor environment	20
2.8 Indoor air pollutants and the respiratory health implication on preschool children	21
2.8.1 Particulate Matter (PM) and its health effects	21
2.8.2 Volatile Organic Compounds (VOCs) and its health effects	23
2.8.3 Biological contaminants and its health effects	23

3	METHODOLOGY	25
3.1	Study location	25
3.2	Study design	25
3.3	Study ethics	25
3.3.1	Consent form.	26
3.4	Study Sampling	26
3.4.1	Study population	26
3.4.2	Sampling frame	26
3.4.3	Sampling criteria	26
3.4.3.1	Inclusive criteria	26
3.4.3.2	Exclusive criteria:	27
3.4.4	Sampling unit	27
3.4.5	Sampling Method	27
3.4.6	Sample size	27
3.4.7	Instrument and Methods	29
3.5	Data Collection	30
3.5.1	Assessment of respiratory symptoms	30
3.6	Kindergarten Indoor Air Quality Assessment	30
3.6.1	Temperature and Relative humidity	30
3.6.2	Particulate Matter (PM2.5, PM10)	31
3.6.3	Volatile Organic Compounds (VOCs)	32
3.6.4	Indoor mold	32
3.6.4.1	Fungi identification	33
3.6.4.2	Staining procedure	33
3.6.5	Lung function test	34
3.6.6	SECA 208 Body meter	36
3.6.7	Tanita®	36
3.7	Data Analysis	36
3.8	Quality control	37
4	RESULTS	38
4.1	Response Rate	38
4.2	Normality Assumption	39
4.3	Background of Respondent	39
4.3.1	Socio-demographic Characteristics	39
4.3.2	Monthly household income of respondents	40
4.3.3	Age and Physical Characteristics of Students	41
4.3.4	Smoking Habit, Breastfeeding and Day Care Assistance	41
4.4	Background of Respondent's Residence	42
4.4.1	Outdoor Residential Background	42
4.4.2	Building Characteristics	43
4.4.3	Inhabitants, Room Sharing and Years Living	44
4.4.4	Indoor Air Pollutant Sources in Homes	45
4.4.5	Poor Indoor Air Quality Indicators in Homes	48
4.5	Comparison of Indoor Air Quality Parameters	49
4.6	Isolation and Identification of Mold in Preschools	50
4.7	Lung Function Test among Respondent in both Location	52
4.8	Respiratory health symptoms	54

4.8.1	Comparison of Respiratory Health Symptoms among Study Respondent	54
4.8.2	Other Health Symptoms	55
4.8.3	Current Health Symptoms	57
4.9	Association between Indoor Air Pollutants and Respiratory Health Symptoms	59
4.9.1	Indoor Air Pollutants and wheezing	59
4.9.2	Indoor Air Pollutants and Phlegm	61
4.9.3	Indoor Air Pollutants and cough	62
4.10	Association between Indoor Air Pollutants and Lung Function	63
4.10.1	Indoor Air Pollutants and FVC Status	63
4.10.2	Indoor Air Pollutants and FEV1 Status	64
4.10.3	Indoor Air Pollutants and FEV1/ FVC% Predicted Status	65
4.11	Factors that Influences the Lung Function and Respiratory Symptom after controlling all confounders	66
4.11.1	Factors that Influence the Lung Function (FEV1%) abnormality among study respondents after controlling all confounders	66
4.11.2	Factors that Influences the Respiratory Symptom (Cough) among Study Respondents after controlling all Confounders	67
5	DISCUSSION	69
5.1	Background of respondents	69
5.1.1	Social-demographic background	69
5.1.2	Socioeconomic Background	69
5.2	Smoking Habit, Day Care Assistance and Breastfeeding	70
5.3	Background of respondents residence	70
5.4	Indoor air pollutant sources in homes	72
5.5	Indoor Air Quality Indicators in Schools and Homes	73
5.5.1	Particulate Matter (PM _{2.5} , PM ₁₀)	74
5.5.2	Volatile Organic Compounds	75
5.5.3	Indoor Air Velocity	76
5.5.4	Indoor mold	77
5.5.5	Indoor Temperature	78
5.5.6	Indoor Relative Humidity (RH)	78
5.5.7	Identification of Mold present in Preschools	80
5.6	Comparison of Lung function test among Respondents in both location	82
5.7	Comparison of Respiratory health symptoms between study and comparative group	84
5.8	Other health symptoms reported by preschool children	86
5.9	Association between Indoor Air Pollutants and Respiratory Health Symptoms among respondents	86
5.9.1	Indoor Air Pollutants and Wheezing	86
5.9.2	Indoor Air Pollutants and Phlegm	87
5.9.3	Indoor Air Pollutants and cough	87
5.10	Association between Indoor Air Pollutants and Lung Function	88

5.10.1	Indoor Air Pollutants and FVC Status	88
5.10.2	Indoor Air Pollutants and FEV1 Status	89
5.10.3	Indoor Air Pollutants and FEV1/ FVC% Predicted Status	89
5.11	Factors that Influence the Lung Function (FEV ₁ %) abnormality among study respondents after controlling all confounders	90
5.12	Factors that Influences the Respiratory Symptom (Cough) among Study Respondents after controlling all Confounders	91
6	CONCLUSION AND RECOMMENDATIONS	93
6.1	Conclusion	93
6.2	Recommendations	94
	REFERENCES	97
	APPENDICES	107
	BIODATA OF STUDENT	129

LIST OF TABLES

Table	Page
3.1 Normal Value of Lung Function Parameters among Children in Malaysia	34
3.2 Evaluation of Lung Function	35
4.1 Socio- demographic characteristics of respondents	40
4.2 Monthly household income of respondents	41
4.3 Age and physical characteristics of respondents	41
4.4 Smoking Habit, Breastfeeding and Day Care Assistance	42
4.5 Comparism of Outdoor Residential Background	43
4.6 Comparism of Residential Building Characteristics	44
4.7 Comparism of Inhabitants, Room sharing and Years living	45
4.8 Comparism of Air Pollutants Sources in Homes	46
4.9 Comparism of Indoor Pollutant Sources from Activities inside Homes	47
4.10 Poor Indoor Air Quality Indicators in Homes	49
4.11 Indoor Air Quality Parameters in Study and Comparative Group Schools (n=12) and Homes (n=60)	50
4.12 Indoor mold isolated from preschools in studied and comparative area.	51
4.13 Lung Function Test among Preschool Children	53
4.14 Lung Function Status and Abnormality among Children	53
4.15 Respiratory Symptoms among Children as Reported by Parents	54
4.16 Other Health Symptoms Reported by Preschool Children and the Association with Study Group n= (270)	56
4.17 Association between IAQ Parameters and the Prevalence of wheezing among study respondents (n=270)	60

4.18	Association between IAQ Parameters and Phlegm among study respondents n = (270)	61
4.19	Association between IAQ Parameters and the Prevalence of cough among study respondents (n=270)	62
4.20	Association between IAQ Parameters and FVC% Predicted Status among study respondents	63
4.21	Association between IAQ Parameters and FEV ₁ % Predicted Status among study respondents	64
4.22	Association between IAQ Parameters and FEV ₁ / FVC% Predicted Status among study respondents	65
4.23	Factors that Influence the Lung Function (FEV ₁ %) abnormality among study respondents after controlling all confounders (n=270)	67
4.24	Factors that Influences the Respiratory Symptom (Cough) after controlling all confounders	68

LIST OF FIGURES

Figure		Page
1.1	Conceptual Frameworks	5
2.1	Particulates matter of different sizes and their source	15
3.1	Data collection workflow	29
3.2	Q-Track Plus IAQ Monitor Model 8554	31
3.3	Dust Trak TM DRX Aerosol Monitor 8534	31
3.4	ppbRAE Portable VOC Monitor	32
3.5	Duo SAS Super 360 Microbiological Air Sampler	34
3.6	MM-SP004 Tabletop Portable Spirometer	35
4.1	Flow chart of Respondent recruitment and the Response Rate	39
4.2	Reported Frequency of Occurrence of Other Health Symptoms	58

LIST OF APPENDICES

Appendix		Page
1	Study Ethics	107
2	Consent Form	110
3	Explanation Letter	111
4	Questionnaire	113
5	Work Plan	125
6	Research Photograph	127

LIST OF ABBREVIATIONS

ACGIH	American Conference of Governmental Industrial Hygienists
ATS	American Thoracic Society
ASHRAE	American Society of Heating, Refrigerating and Air Conditioning Engineers
DOSH	Department of Occupational Safety and Health
EEHS	European Environment and Health Strategy
IAQ	Indoor Air Quality
ICOP	Industrial Code of Practice
ISAAC	International Study of Asthma and Allergies in Children
ppm	Part per million
ppb	Part per billion
PM	Particulate Matter
PM _{0.1}	Particulate Matter less or equal to 0.1
PM _{2.5}	Particulate Matter less or equal to 2.5
PM ₁₀	Particulate Matter less or equal to 10
VOCs	Volatile Organic Compounds
SDA	Sabouraud Dextrose Agar
SPSS	Statistical Package for Social Sciences
USEPA	United State Environmental Protection Agency
WHO	World Health Organisation

CHAPTER 1

INTRODUCTION

1.1 Background

Indoor air quality (IAQ) refers to the nature of air that circulates throughout the space where we live (USEPA 2010). It refers to the air that we breathe during the time we spend indoors. Indoor air is made up of physical, biological, and chemical components. Indoor temperature and humidity can affect the comfort of occupants. Indoor air quality also refers to the characteristic of the indoor air inside a building (USEPA 2010). Indoor environments are considered to be protective, at the same time they can become contaminated with particles which can cause more serious risks than those related to outdoor exposures when the concentration of indoor air pollutants exceeded the recommended maximum limits. Approximately 19% deaths in children under the age of 5 were attributed to respiratory infections (World Health Organization 2010).

Indoor air quality (IAQ) has recently become a concern. There is an increasing concern regarding the effects of indoor air quality (IAQ) to health because people spend more of their time indoors than outdoors. The slow exchange of air indoors has resulted to accumulation of indoor air pollutants to a substantial level. From frequent observations, it has estimated that a quarter of the population in the world has been exposed to pollutants which can be harmful to health (Canha et al., 2010).

Indoor air pollutant level is higher than outdoor air pollutant level (Godoi et al., 2009; Jo and Seo, 2005; Kotzias et al., 2009; Pegas et al., 2010; Yang et al., 2009). Children are more exposed to indoor air pollutants compared to the adult. This is resulted from the immaturity of their immune system and growing process (Mendell et al., 2005). In recent years, there has been an increase in the prevalence of asthma and allergy among children. Currently, researchers focused more in school environments as a second source besides the home environments of children. This is because children spend most of their time either in school or at home (Anna and Elwira 2015; Branco et al., 2014; Santamouris et al., 2008).

Indoor air quality monitoring have being conducted in recent studies. This studies measured the indoor level of particulates (Ayuni et al., 2014; Anis et al 2015; Asrul et al., 2016; Anna and Elwira, 2015), presence of mold (Ayuni et al., 2014; Andrew et al., 2015), relative humidity and thermal comfort (Jing et al., 2012; Asrul et al., 2016; Andrew et al., 2015).

Studies have shown a consistent association between prolonged exposure to indoor air pollutants and increase in respiratory symptoms which reduces the lung function of children (Nazariah et al., 2013; Salleh et al., 2013; Nur Azwani et al., 2015; Nurul Fariza et al., 2010). Barnett et al. (2005) found that the risk of developing respiratory symptoms such as cough, wheeze, asthma, respiratory infections, headache and tiredness is higher for occupants living in moisture damaged buildings. All damp surfaces in buildings including concrete, wood, plastics, bricks, painted surfaces or metal may become colonized by microbes from the air. The colonizing microbes are bacteria, fungi and algae. Human activities like talking, sneezing, coughing, walking, washing and toilet flushing can generate airborne biological particulate matter. Spores from pets, carpets, dust and furniture can be released into the air. (Maeir et al., 2006).

Studies have shown that the presence of indoor air pollutants in the environment has caused a greater impact on the health of occupants' especially preschool age who are still growing. There has been an increase in the rate of respiratory symptoms as a result of exposure to indoor pollutants (Noor Hisyam & Juliana, 2014)

Indoor air comprises of biological components such as mold. Mold includes airborne particles which are of biological origin. Mold can enter into human respiratory tract through inhalation and also by skin contact. Studies have shown consistent associations between indoor mold and respiratory or allergic health effects in infants, children and adults (Institute of medicine, 2005 and World health Organization, 2009).

Many recent studies have reported the association between health symptoms and the physibile growth of mold as a result from accumulation of water in school building and moisture damage in the school structures (Aydogbu et al., 2005; Hussin et al., 2011; Meklin et al., 2010). There has not been a standardized guideline for indoor mold and dampness in indoor environment (Karvala 2012 and WHO 2009). Previous studies related to the measurement of microbial agents have not shown much consistent association between microbiological agents and health symptoms. A water damaged building caused by mold and moisture have reported significant fungi growth in indoor environments (Kavala 2012). There is no generally accepted environmental sampling method which can be used in assessing the exposure to environmental contaminants including mold growth and other indoor environmental contaminants (ACGIH, 2009 & Frankel et al., 2012). Fungi depends on temperature and humidity to grow suitably in indoor environment.

National Institute for Occupational Safety and Health (NIOSH, 2011), has reported that in the last decades microbial contaminants are the main cause of poor air quality. The report also found that, approximately 5% of more than 500 indoor air quality (IAQ) from investigation and the remaining 95% was a result of poor ventilation, contamination in building and the rest are not known (NIOSH, 2011).

Disease of the lungs often manifest by one or more symptoms that can be easily recognized. Thus the presence of particular symptoms can confirm the presence of diseases in the upper and lower airways. Upper respiratory symptoms include (URS), runny and stuffy nose, sinusitis, sore throat, cold, fever, burning or red eyes. Lower respiratory symptoms include wheezing, wet cough and dry cough, phlegm, shortness of breath and chest discomfort or pain. These respiratory diseases are caused by bacteria and mold. These symptoms developed as a result of exposure to indoor air pollutants (Dockery et al. 2010).

1.2 Problem Statement

Children living in the industrial area have different indoor air exposure rate compared to children living in the suburban area. Exposure to industrial emission and automobile exhaust can cause acute respiratory diseases in children. Particulate matter has been associated with respiratory infections in children between the ages of 0-4 years. Exposure to pollutants like gases and particulate matter which are released from combustion of fossil fuel can cause a reduction in the lung function of children. Studies have shown that 1% mortality rate among children has been attributed to exposure to indoor particulate matter (Franklin 2007). Malaysia statistics data by DOE (2014) has reported that 70% to 75% of air pollution are attributed from traffic vehicles emissions (Noor Hisyam & Juliana, 2014; Nur Azwani et al., 2015; Salleh et al., 2013; Lee et al., 2012).

The presence of respiratory symptoms have being linked to the exposure to indoor air pollutants. Local studies have reported a high prevalence of respiratory symptoms like cough and phlegm among preschool children who are exposed compared to indoor air pollutants compared to those who are not exposed (Yahaya and Jalaludin, 2014; Nurul Anis et al., 2013; Nazariah et al., 2013; Ayuni et al., 2014; Nur Fazhilah et al. 2012). A report by World Health Organization (2005) has shown that approximately 2.7% of diseases occurred globally are due to the effects of poor indoor air quality

Preschool children spend most of their time indoors at home and schools. They attend classes in the morning approximately 4-5 hours, will be fetched by their parents or guardian and some of them will spend the remaining hours of the day at home. As a result of these, children are highly exposed to indoor air pollutants. Children are the main subject in this study because they easily respond to exposure to indoor air pollutants. A study has shown that from 1982 to 2013, there has being an increase from 2.5 to 7.1 million in the number of children suffering from asthma. In Malaysia 10% of children had been diagnosed with asthma. Furthermore, nearly 80 to 90% of asthmatic cases in Malaysia were children. There is no conclusive evidence showing that there are more asthmatic children in Malaysia now than before (Junaidah et al., 2010).

Susceptibility to mold related diseases occurs mainly in immune compromised persons. The person may develop chronic illnesses, such as lung diseases (CDC, 2005). Jacob et al. (2008) found that there was a significant association between exposure to *Penicillium* spores and the incidence of respiratory tract infections. Pollutants in the indoor air such as particulate matter, bioaerosol and other chemical components expose children to different respiratory health diseases and symptoms. Exposures to air pollutants are manifested through respiratory symptoms such as coughing, wheezing, sneezing, breathlessness, phlegm and shortness of breath.

Preschool indoor air quality and settings should be assessed in order to reduce indoor air pollutants and spread of respiratory symptoms. Local studies have found that preschool settings and characteristics of the surrounding environment can increase asthmatic symptoms, respiratory symptoms and allergies. (Marzuki et al. 2010; Nurul Anis et al. 2014; Ayuni et al., 2014; Yahaya et al., 2014; Salleh et al., 2013).

The aim of this study was to evaluate the indoor air quality and its' health implication on Malay preschool children living in the selected locations.

1.3 Conceptual frameworks

The aim of this study is to determine the association between indoor air pollutants and respiratory health among Malay preschool children in Puchong and Hulu Langat, Selangor, Malaysia. Locations used in this study were selected based on their level of exposure where Puchong represented the polluted area (Industrial) and Hulu Langat represented the less polluted area (Suburban). Exposure to pollutants was assessed by measuring the concentration of indoor pollutants such as VOCs, PM_{2.5}, PM₁₀, mold, relative humidity and temperature. The main route of exposure to indoor air pollutants in this study is by inhalation. Home exposure to pollutants was also assessed. The presence of respiratory symptoms (cough, phlegm, wheeze, chest tightness) was assessed using a questionnaire. Lung function of respondents was also assessed. Figure 1.1 below shows the study variables in this study.




Figure 1.1 : Conceptual Frameworks

1.4 Study Justification

The purpose of this study is to determine the respiratory health implications due to exposure to indoor pollutants. This study will relate exposure to indoor air pollutants and the respiratory implication among preschool children in the different study locations.

Children are known to be the most susceptible groups (British Lung Foundation, 2014). Approximately 19% deaths in children under the age of 5 were attributed to respiratory infections (World Health Organization 2010). There have not been many studies carried out on preschool aged children compared to adults.

Rate of asthma, allergy and other health implications have been increasing in children which are caused by exposure to environmental factors instead of genetic influence. This study will help to provide information on exposures and prevention of these health implications. Respiratory symptoms can develop into full blown asthma for those who are prone to or those, who have family history of asthma. Unhealthy lungs may also lead to cough or cold which can affect the quality of life and missing school days. Medical treatment can be expensive and therefore it is better to identify health problems earlier by conducting indoor air monitoring study.

Previous studies conducted in Malaysia has shown that the presence of particulate matter (PM_{10} , $PM_{2.5}$) can directly cause an influence on the severity of asthma among school children who reside in the different locations of urban, industrial, and rural areas of Selangor and Kuala Lumpur. Exposure to indoor air pollutants can increase the prevalence of respiratory symptoms, which can affect the lung function of children. Indoor monitoring is important to reduce the level of exposure to children. prolonged exposure can lead to adverse health effects (Marzuki et al., 2010; Salleh et al., 2013; Yahaya et al., 2013; Ayuni et al., 2014; Nurul Anis et al., 2013; Ithnin et al., 2013; Nur Azwani et al., 2015; Tezara et al., 2014).

This study is more reliable since the effects from exposure to indoor air pollutants and respiratory symptoms are conducted among preschool children, this is because smoking is rarely found in children. They are not exposed to occupational pollutants, they have a more stable residence unlike the adults and also because of the nature of their immune system which is still maturing (Branco et al., 2014; Santamouris et al., 2008).

The locations (Puchong and Hulu- Langat) used for this study was selected due to the activities that have been carried out which can increase the possibility of indoor pollutants in preschools located around the areas. Development and construction of amenities, roads and houses has led to increase in the level of atmospheric pollution.

Mold is harmful to human health. It can cause damages on building materials and exposure to mold can also cause health effects. Malaysia is a tropical country with warm and humid climate. There are still limited studies which has related the presence of indoor mold in school environments especially in tropical countries. So far, studies carried out assessed the cause of allergens in settled dust instead of assessing allergen through air sampling (Hussin et al., 2011).

Previous studies (International Study of Asthma and Allergy in Childhood, 2015; European Environment and Health Strategy, 2003) have found an increase in the rate of asthma which is more prominent in the lower middle income countries (Asher et al., 2006). There has been a significant increase in the rate of asthma among children from 3-10% in the year 1990 to 2001 in studies which has been previously conducted in Malaysia. Wheezing symptom has also being found to increase from the rate of 5-8% in children (Quah et al., 2005).

Studies have reported that the exposure to mold in damp buildings is associated with the risk of experiencing health problem, in relation to no clear standard for the concentrations of microorganisms. A study conducted in Singapore on mold exposure has shown an increase in level of allergens in schools compared to homes (Zhang et al., 2006). Therefore, this study will help to provide more information on the existing knowledge on microbial contaminants in the indoor environment.

1.5 Study Variable

1.5.1 Dependent variable

The dependent variables in this study are respiratory symptoms which includes wheezing, cough, phlegm, chest tightness and lung function status.

1.5.2 Independent variable

The independent variable are particulate matter (PM), volatile organic compounds (VOCs) and mold. Physical parameters for Indoor Air Quality status (IAQ) such as temperature, humidity and air velocity were assessed.

1.6 Research Objectives

1.6.1 General objective

To determine the association between exposures to indoor air pollutants and respiratory health implications among preschool children in selected areas.

1.6.2 Specific objective

- 1) To determine the socio demographic status of preschool children in study areas.
- 2) To determine the indoor air quality of the preschools in study and comparative area.
- 3) To identify the different types of mold present in the selected preschools.
- 4) To compare lung functions (FVC, FEV₁, FEV₁/FVC) of respondents in the study and comparative area.
- 5) To compare the presence of respiratory symptoms between the children in the study and comparative area.
- 6) To determine the association between indoor air pollutants and respiratory health symptoms among preschool children in the study locations.
- 7) To determine the association between the indoor air pollutants with lung function (FVC, FEV₁, FEV₁/FVC) among the respondents in the study areas.
- 8) To identify the main factors that influence indoor air pollutants and respiratory symptoms among study respondents after controlling all the confounders.

1.7 Research Hypothesis

- 1) There is a significant difference in the level of exposure to indoor air pollutants between the study and comparative group.
- 2) There is a significant difference between the lung functions of the respondents in study group and the comparative group.
- 3) There is a significant difference between the respiratory symptoms of respondents in the study and the comparative group.
- 4) There is a significant association between exposure to indoor air pollutants and respiratory symptoms in preschools children in the study and comparative group.
- 5) There is a significant association between the indoor air pollutants with lung function (FVC, FEV₁, FEV₁/FVC) among respondents.
- 6) There is a significant association between the indoor concentration of PM_{2.5}, PM₁₀, VOC, temperature, relative humidity with respiratory symptoms among study respondents after controlling all the confounders.
- 7) There is a significant association between the indoor concentration of PM_{2.5}, PM₁₀, VOC, temperature, relative humidity with lung function among study respondents after controlling all the confounders.

1.8 Definition of Variables

1.8.1 Conceptual definition

a) Indoor air quality

Indoor air quality refers to the air quality of buildings and structures as it relates to the health and comfort of building occupants. The component of indoor air determines the quality of indoor environments in relation to the comfort and wellbeing of occupants. Indoor environment comprises of physical components such as humidity, temperature, velocity; biological components which include virus, fungi, bacteria, spores; and chemical components which include volatile organic compounds, carbon dioxide, formaldehyde, smoke from combustion (Environmental Protection Agency 2015).

b) Mold

Mold are group of organisms that belong to the kingdom of fungi and are filamentous in nature. Mold produces airborne spores. These particles are from biological origin, and are usually suspended in the air. They can also be referred to as organic dust. Bioaerosol particles vary in different sizes from below 1 μm to 100 μm in aerodynamic diameter. Mold has being shown to be a major source of indoor air pollution that can cause allergic reactions when they are inhaled by children. Since the presence of mold in the indoor environment is related to various human health symptoms and the indoor environment is a means of exposure, indoor mold has become a concern (Sanchez-Monedero et al., 2008).

c) Volatile Organic Compounds

Volatile organic compounds (VOCs) are emitted as gases from certain solids or liquids. VOCs include a variety of chemicals; some have short- term while some have long- term adverse health effects. Concentrations of VOCs are usually higher indoors than outdoors. These chemical are emitted from household products such as detergents used for cleaning, paints, plastics (Environmental Protection Agency 2013).

d) Respiratory Symptoms.

Respiratory symptoms are symptoms which are experienced after exposure to indoor air pollutants and can last up to 7-10 days. They usually begin 1-3days after the exposure to air pollutants and can last up to 7 to 10 days. These signs include chronic cough, wheezing, shortness of breath and red itchy eyes. (Eccles et al., 2007).

e) Particulate Matter

These are complex mixture of small particles and liquid droplets. Particulates have different aerodynamic diameters. $\text{PM}_{2.5}$ are referred to as inhalable fine particles. They are coarse in nature with the aerodynamic diameter 2.5 μm . They are commonly found in smoke and haze. The can cause impaired vision, it can release harmful substances

into the air. When inhaled these particles can penetrate deep into the lung tissues and cause health problems. (USEPA, 2012).

PM₁₀ are particles with aerodynamic diameter of 10 micrometers or less. This particle causes a greater health risk among other particulate matter because it can cause damage to the lung tissue and respiratory implications when they are inhaled. It can even go as far down the bloodstream of humans. PM₁₀ is known to cause serious health impairment such as decrease in lung function, increase of respiratory symptoms, lung diseases and asthma severity. This particle is produced from activities like grinding and also from roads (USEPA, 2012).

f) Forced Expiratory Volume in 1 second (FEV₁)

This is the maximum volume of air exhaled in the first second which is as a result of forced expiration and from a position of full inspiration. It is expressed in liters. This test is used in the diagnoses of obstructive lung diseases such as asthma and chronic obstructive pulmonary disease (COPD). There is reduction in FEV₁ if there are any forms of obstruction in air flow. Forced Expiratory Volume was performed based on American Thoracic Society standard.

g) Forced Vital Capacity (FVC)

This refers to the volume of air that can be forcefully exhaled from the lungs after a maximum inspiration. This is the total amount of air that is exhaled during FEV test. Measuring how well the children can forcefully exhale can be used to detect the presence of any chronic obstructive pulmonary diseases (COPD). It is expressed in liter. There is reduction in FVC if there is restriction in the lung volume or air flow capacity. Forced Vital Capacity was carried out based on the standard set by American Thoracic Society.

h) FEV₁% Predicted

This is expressed based on the normal value as a percentage of the Forced Expiratory Volume in one second. $FEV_1\% \text{ Predicted} = (FEV_1 \text{ measured} / FEV_1 \text{ Predicted}) \times 100$

i) FVC% Predicted

This is expressed based on the normal value as a percentage of the Forced Vital Capacity. $FVC\% \text{ Predicted} = (FVC \text{ measured} / FVC \text{ Predicted}) \times 100$

h) Wheezing

This is a whistling sound which is made while inhaling and exhaling. This could be as a result of inflammation in the lungs. Wheezing could be associated with breathlessness and occurs days and nights (Bass 2009).

i) Chronic Cough

Cough occurs naturally with the removal mucus when the throat is irritated. Cough becomes chronic when it is consistent and lasts for at least 4days in a week and occurs consecutively for 4 months in a year. Cough occurs when air is released forcefully from the lung. It could be triggered by cold or infection of the sinus (Pubmed article, 2013)

j) Chest Tightness

This is a continuous occurrence of cough or phlegm. It causes a sharp pain and ache in the chest. This occurs consecutively for a period of 3 months in a year. It can signify the presence of some diseases in the body (Chen 2014).

k) Chronic Phlegm

This occurs when there is persistent release of mucus which usually coughs up from the chest and occurs for at least 4days in a week and occurs consecutively for 3 months in a year (Phlegm 2007).

1.9 Operational Definition

a) Particulate Matter (PM_{2.5} and PM₁₀)

Indoor air quality assessment was carried out by measuring the indoor environmental level of PM_{2.5} and PM₁₀. Indoor assessment of particulate matter was measured using TSI DustTrak™ DRX Aerosol Monitor 8534. This instrument detects aerosol concentration between the ranges of 0.001 to 150 mg/m³. The aerosol particles are isolated in the optic chamber using the sheath air system (USEPA, 2012).

b) Volatile Organic Compounds (VOCs)

Volatile Organic Compound is one of the indoor parameter which influences the growth of bioaerosol in indoor environment. In this study, concentration of Volatile Organic Compound was measured using ppbRAE Volatile Organic Compound (VOC) Monitor (Model PGM-7240) which is a sensitive photo-ionization Detector (PID) for real-time monitoring of volatile organic compounds (VOCs) at part per billion (ppb) levels. Data was downloaded using the proRAE Remote software (USEPA, 2013).

c) Indoor mold

Measurement of mold was conducted using DuoSAS Super 360 microbiological air sampler by collecting air samples into petri dishes containing agar medium, 200 volumes of air sample for 2 minutes was used during sampling. The petri dishes was sealed with parafilm for transportation to the laboratory. The samples was incubated and analysis was carried out. The media that was used is Sabaroud Dextrose Agar (SDA) for investigating total fungi. The colonies was identified through their specific colour, turbidity or other characteristics that appear when cultured on selective media (Sanchez-Monedero et al., 2008).

d) Temperature and Relative Humidity

This was measured using TSI Q-Trak plus Indoor Air Quality Monitor 8554. This equipment has a sensor type thermistor and measures the ranges of 0 to 50°C. The sensor is made up of type thin-film capacity which can detect 5 to 95% relative humidity. It uses the principle of detection of a wet bulb and dry bulb to compare the humidity ratio and absolute humidity to measure the level of RH in indoor environment (USEPA, 2013).

e) Respiratory symptoms

Respiratory symptoms of respondents in this study was determined using a questionnaire on respiratory health symptoms adapted from American Thoracic Society (ATS). Examples of respiratory symptoms are coughing, wheezing, phlegm (American Thoracic Society).

f) Forced Vital Capacity (FVC)

Lung function test was assessed based on the American Thoracic Society (ATS) standard to determine the forced vital capacity level of the respondents using a spirometer. This is expressed in liters at body temperature. (American Thoracic Society 2005).

g) Forced expiratory volume (FEV₁)

This was assessed based on the ATS standard 1994 to determine the forced expiratory volume level of the respondents using a spirometer. Forced vital capacity is expressed in liters (American Thoracic Society 2005)

REFERENCES

- Adrian, G., Barnett, Gail, M., Williams, Joel Schwartz (2005). Air pollution and Child Respiratory Health. *American Journal of Respiratory and Critical Care Medicine*. 1272-1278.
- American Conference of Governmental Industrial Hygienists. (1999). *Bioaerosols assessment and control*. American Conference of Governmental Industrial Hygienists, Cincinnati.
- American Lung Association (2013). American Lung Association State of the Air. *Health Effects of Ozone and Particle Pollution*. Retrieved from <http://www.stateoftheair.org/2013/health-risks/>
- American Society of Heating, Refrigerating and Air Conditioning Engineers (ASHRAE), (1999). *Ventilation for Acceptable Indoor Air Quality* (ASHRAE 62-1999) [Standard] Atlanta.
- American Thoracic Society, (2005). Lung Function testing. Selection of Reference values and Interpretive Strategies. *American Review of respiratory Disease*. 85: 762-768.
- American Thoracic Society, (2005). Standardization of Spirometry. *Eur Respiratory Journal*. 26:319 -338.
- America Thoracic Society (1978) Recommended respiratory disease questionnaires for use with adults and children in epidemiological research. Available at: <http://www.thoracic.org/statements/resources/archive/rrdquacer.pdf>
- Andrew, W.D., Jalaludin, J. (2015). Indoor air pollutant exposure and eosinophil cationic protein as an upper airway inflammatory biomarker among preschool children. *Procedia Environmental Sciences* 30, 297-302.
- Anna, M. & Elwira, Z. (2015). Indoor Air Quality in Urban and Rural Preschools in Upper Silesia, Poland: Particulate Matter and Carbon Dioxide. *Int. J. Environ. Res. Public Health*, 7697-7711.
- Anis, S. K. and Juliana, J. (2015). Indoor Air Quality and its Association with Respiratory Health among Malay Preschool Children in Shah Alam and Hulu Langat, Selangor. *Advance in Environment Biology*, (9)9: 17-26.
- Asher, M.I., Montefort, S., Bjorksten B., Lai, C.K., Strachan, D.P (2006). Worldwide time trends in the prevalence of symptoms of asthma, allergy rhino conjunctivitis, and eczema in childhood. *Lancet* 368: 733-743.

- Asrul, S and Juliana, J. (2017). Indoor Air Quality and its Association with Respiratory Health among Preschool Children in Urban and Suburban area Selangor, Malaysia. *Malaysian Journal of Public Health Medicine* (1): 78-88.
- Aydogdu, H., Asan, A., Otkun, M.T., Ture, M. (2005). Monitoring of fungi and bacteria in the indoor air of primary schools in Edirne city, Turkey. *Indoor Built Environment*. 14: 411 - 425.
- Ayuni, N.A., Juliana, J. & Ibrahim, M.H. (2014). Exposure to PM₁₀ and Association with Respiratory Health among Primary School Children Living Near Petrochemical Industry Area at Kertih, Terengganu. *Journal of Medical and Bioengineering*. 3(4): 282-287.
- Azizi, B.H.O. & Henry, R.L. (1990). Effect of Indoor Air Pollution on lung function of primary school children in Kuala Lumpur. *Pediatric Pulmonology*. 9(1): 24 - 29.
- Bass, P. (2009). Wheezing Definition. Retrieved August 25, 2015, from http://asthma.about.com/od/glossary/g/Wheezing_def.htm
- Barnett, A.G., Williams, G.M., & Schwartz, J., (2005). Air pollution and child respiratory health: A case-crossover study in Australia and New Zealand. *American Journal Respir Crit Care Med*. 171: 1272-1278.
- Bornehag, C.G., Blomquist, G., Gyntelberg, F., Jarrholm, B., Malmberg, P., Nordvall, L., Nielsen, A., Pershagen, G., Sundell, J. (2001). Dampness in buidings and health, Nordic interdisciplinary review of the scientific evidence on associations between exposure to dampness in building and health effects. *Indoor Air*. 11: 72 - 86.
- Broms, K., Syardsudd, K., Sundelin, K., Norback, D. (2006). A nationwide study of indoor and outdoor environments in allergen avoidance and conventional daycare centers in Sweden. *Indoor Air*. 16: 227 - 235.
- British Lung Foundation (2014). Children's lung condition. Retrieved October 01, 2015, from <https://www.blf.org.uk/Page/Children-lung-conditions>.
- Canha, N., M. C. Freitas, M.C., S. M. Almeida, S.M. (2010). Indoor school environment: easy and low cost to assess inorganic pollutants. *Journal of Radioanalytical and Nuclear Chemistry*. 286(2): 495-500.
- Centers for Disease Control and Prevention (CDC). National Center for Environmental Health, Air Pollution and Respiratory Health Branch (2005). *Molds in the Environment*. Retrieved from <http://www.cdc.gov/Mould/faqs.htm>. (30th August 2005).

- Chan Yeung, M.B.A. (2006). Primary prevention of childhood asthma and allergic disorders. *Allergy and Clinical Immunology*. 6: 146 - 151.
- Chengi, C.Y. (2002). PDHengineer. *Indoor Air Quality*. Retrieved from <http://www.pdhengineer.com>. [10th February 2011].
- Chen, Q. and L.M. Hildemann, (2009). The Effects of Human Activities on Exposure to Particulate Matter and Bioaerosols in Residential Homes. *Environmental Science & Technology*. 43(13): 4641-4646.
- Chen, M.A. (2014). Chest Pain. Retrieved July 12, 2014, from <http://www.nlm.nih.gov/medlineplus/ency/article/003079.htm>
- Chua, P.C., J. Jalaludin, 2014. An overview of indoor air quality and its impact on respiratory health among Malaysian school-aged children. *Reviews on Environmental Health*. ISSN (Online) 2191-0308, ISSN (Print) 0048-7554, DOI: 10.1515/reveh-2014-0065.
- Chua, P.C., Jalaludin, J., Titi, R.H., Nor Mariah, A. (2015). Indoor Air Quality Assessment and Lung Functions among Children in Preschool at Selangor, Malaysia. *Advances in Environmental Biology*, 9(9): 1-9.
- Cionita T, Jalaludin J, Adam NM, Siregar JP. (2014). Assessment of Children's Health and Indoor Air Contaminants of Day Care Centre in Industrial Area. *Iranian J Publ Health*, 43 (3):81-88.
- Corgnati, S.P., Ansaldi, R., and Filippi, M., (2009). Thermal comfort in Italian classrooms under free running conditions during mid-seasons: assessment through objective and subjective approaches. *Building and Environment*. 44(4): 785 – 792.
- Corgnati, S.P., Filippi, M., and Viazzo, S., (2007). Perception of the thermal environment in high school and university classrooms: subjective preferences and thermal comfort. *Building and Environment*. 42(2): 951 – 959.
- David, E., Stephen, D., Helen, A., Rosemary, H., Robya, B. (2007). *Descriptions of medical fungi 2nd edition*. School of Molecular & Biomedical Sciences University of Adelaide, Australia.
- DOE, (2014). Recommended Malaysian Ambient Air Quality Guidelines (RMAAQs). Department of Environment, Malaysia. Retrieved from <http://apims.doe.gov.my/apims/General%20Info%20of%20Air%20Pollutant%20Index.pdf>
- Department of Occupational Safety and Health (2005). Code of Practice on Indoor Air Quality, Malaysia. Ministry of Human Resources.
- Department of Occupational Safety and Health (2010). Code of Practice on Indoor Air Quality, Malaysia. Ministry of Human Resources.

- Dockery, D.W., Skerrett, P.J., Walters, D., Gilliland, F. (2005). Development of lung function in effects of air pollution on children's health and development: A review of the evidence. World Health Organization Special Programme on Health and Environment.
- Eccles, M.P., Grimshaw, J.M., Johnston, M. (2007). Applying psychological theories to evidence-based clinical practice: Identifying factors predictive of managing upper respiratory tract infections without antibiotic. *Implement Sci*, 2:26.
- Environmental Protection Agency, (2015). *The Inside Story: A Guide to Indoor Air Quality*. http://www.epa.gov/iaq/pubs/inside_story.html.
- EPA. (2013, June 22). *Biological pollutants*. Retrieved from United States Environmental Protection Agency: <http://www.epa.gov/iaq/biologic.html>.
- Fariza, A. R. N. J. Juliana, R. H. Titi, and H. Zailina (2010). Exposure of indoor air pollutants (PM_{2.5}, PM₁₀, ultrafine particle, CO₂, CO, VOCs & NO₂) and lung function of school children in Klang Valley. *Proceedings of the National Conference on Environmental and Health*. 75: 530–539.
- Frankel, M., Timme, M., Hansen, E.W., Madsen, A.M. (2012). Components of sampling methods for the assessment of indoor microbial exposure. *Indoor Air* 22: 405 - 414.
- Franklin, P.J. (2007). Indoor Air Quality and Respiratory Health of Children. *Paediatric Respiratory Review*. 8: 281-286.
- Fritz G. J. and O. Herbarth, (2001). Pulmonary function and urban air pollution in preschool children. *International Journal of Hygiene and Environmental Health*. 203(3): 235–244.
- Fromme, H., Diemer, J., Dietrich, S., Cyrus, J., Heinrich, J., Lang, W., Kiranoglu, M., Twardella, D. (2008). Chemical and morphological properties of particulate matter in school classrooms and outdoor air. *Atmospheric Environment*. 42: 6597 - 6605.
- Gauderman, W.J., Avol, E., Cililand, F. (2004). The Effects of Air Pollution on Lung Development from 10 to 18 years of Age. *The New England Journal of Medicine*. 351 (11): 1057-1067.
- Godoi, R., Avigo, D., J.R., Campos, V., Tavares, T., D.E., Marchi, M., Grieken, R. (2009). Indoor air quality assessment of elementary schools in Curitiba, Brazil. *Water, Air, Soil Pollution*, 9: 171 – 177.
- Godwin, C. & Batterman, S. (2007). Indoor air quality in Michigan schools. *Indoor Air*. 17 (2): 109 – 121

- Gordon, D. (2004). The Environment and children's Health: A Journalist Resource Reporting. *Environmental*. 466-467.
- Goyal, R., G. & Khare, M. (2009). Indoor-outdoor concentrations of RSPM in classroom of a naturally ventilated school building near an urban traffic roadway. *Atmospheric Environment*. 43(38): 6026 - 6038.
- Hernberg, S., Sripaiboonkij, P., Quansah, R., Jaakkola J.J.K., Jaakkola, M.S. (2014). Indoor molds and lung function in healthy adults. *Res Med*. 108: 677-684.
- Ho-Hyun Kim, Myung Hyun Sohn, Kyu- Earn Kim, Chang-soo Kim, Dong Chun Shin. (2011). Prevalence of Allergic Diseases among Korean school- age children. *Journal of Korean Medical Science*. 26(3): 332 - 338.
- Hussin, N.H.M., Sann, L.M., Shams Udin, M.N., Hashim, Z. (2011). Characterization of bacteria and fungi bioaerosol in the indoor air of selected primary schools in Malaysia. *Indoor Built Environ*. 20: 607 - 617.
- Hwang, R.L., Lin, T.P., & Kuo, N.J. (2006). Field experiments on thermal comfort in campus classrooms in Taiwan. *Energy and Buildings* 38 (1): 53 – 62.
- Ismail, M., Nur Zafirah, M.S., Ahmad, M.A. (2010). Indoor Air Quality in selected samples of primary schools in Kuala Terengganu, Malaysia. *Environment Asia*, 3, 103-108.
- Jacob, B., Ritz, B., Gehring, U., Bishop, W., Wichmann, H.E., Heinrich, J. (2008). Indoor exposure to mold and allergic sensitization. *Environmental Health Perspective* 110(7): *The IEQ Review*. Retrieved from <http://www.pureaircontrols.com/> (24th August 2005).
- Jo, W.K. & Y.J. Seo. (2005). Indoor and outdoor bioaerosol levels at recreation facilities, elementary schools and homes. *Chemosphere*. 61(11): 1570 - 1579.
- Junaidah, Z., Lye, M.S and Zailina H. (2012). Asthma Severity and Environmental Health Risk Factor among Asthmatic Primary School Children in the Selected Areas. *American Journal of Applied Sciences*. 9(10), 1553-1560.
- Kalogerakis, N. D., Paschali, V., Le kaditis, A., Pantidou, K., Eleftheriadis, M., Lazaridis. (2005). Bioaerosol measurements in domestic and office premises. *Journal of aerosol science*. 36: 751 - 761.
- Kamaruzzaman, S.N. Emma Zawawi, M.A. Micheal, P. Zuraidah, M.D. (2010). Occupant feedback on indoor environment quality in refurbished historic building. *Int. J. Phys. Sci*. 5(3), 192-199.
- Karvala, K. (2012). Asthma in damp indoor work environments people and work. Research reports 9. Finnish institute of occupational Health, Helsinki: P. 236.

- Kathleen Mortimer (2008). Early lifetime exposure to air pollution and allergic sensitization in children with asthma. *Asian Journal of Atmospheric Env.* 4: 157 - 167.
- Kim, J.L., (2007) Indoor molds, bacteria, microbial volatile organic compounds and plasticizers in schools – associations with asthma and respiratory symptoms in pupils. *Indoor Air*, 17 (2): 153 – 156.
- Kotzias, D., Geiss, O., Tirendi, S., Barrero-Moreno, J., Reina, V., Gotti, A. (2009). Exposure to multiple air contaminants in public buildings, schools and kindergartens. The European indoor air monitoring and exposure assessment (AIRMEX) study. *Fresenius Environmental Bulletin*, 18(5a): 670 – 681.
- Lameshow, S., & Levy, P.S. (1990). Sampling of Populations: Methods and Applications. 3rd edn. John Wiley & Sons inc. New York, US.
- Ling, K.H., Ting, A., Shaharuddin, A., Kadaruddin, M.J., Yaakob. (2010). Urban growth and air quality in Kuala Lumpur city Malaysia. *Environment Asia*. 3: (2) 123–128
- Lippmann, M. & Frampton. (2003). The US environmental Protection Agency particulate matter health effects research centers program: *Environmental Health Perspective*. 111: 1074 - 1092.
- Madureira, J., Paciência, I., Fernandes Ede, O. (2012). Levels of indoor-outdoor relationship of size specific particulate matter in naturally ventilated Portuguese School. *J Toxicol Environ Health*. 75 (22-23): 1423-1436
- Maier, W.C., Arrighi, H.M., Morray, B. (2006). Indoor risk factors for asthma and wheezing among Seattle school children. *Environ Health Perspect*, 105(2): 208 - 214.
- Marzuki, I., Nur Zafirah, M.S and Ahmad, M. A. (2010). Indoor air quality in selected samples of primary schools in Kuala Terengganu, Malaysia. *Environ Asia*, 3:103 - 8.
- Meklin, T., Husman, T., Vepsäläinen, A., Vahleristo, M., Koivisto, J., Halla-Aho, J., Hyvärinen, A., Moschandreas, D., Nevalainen, A. Indoor air microbes and respiratory symptoms of children in moisture damaged and reference schools. *Indoor Air*. 12: 175 - 183.
- Mendell, M.J. & Heath, G. (2005). Do indoor pollutants and thermal conditions in schools influence student performance? A critical review of the literature. *Indoor Air*. 15(1): 27 - 52. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15660567>.

- Menetrez, M.Y., Foarde, K.K., Webber, T.D., Dean, T.R., Betancourt, D.A., Moore, S.A., Svendsen, E.R., Yeatts, K. (2009). The measurement of ambient bioaerosol exposure. *Aerosol Science and Technology*. 41: 884 – 893.
- Meyer, H.W., Suadican, P., Valbjorn, O., Sigsqaard, T., Gyntelberg, F. (2004). Mould in floor dust and building-related symptoms in adolescent school children: A problem for boys only? *Indoor Air*. 14(S10): 65 – 72.
- Mi, Y.H. (2006). Current asthma and respiratory symptoms among pupils in Shanghai, China: influence of building ventilation, nitrogen dioxide, ozone, and formaldehyde in classrooms. *Indoor Air*. 16 (6): 454 – 464.
- MohdNorRawi NA, Jalaludin J, Choo CP. (2015). Indoor Air Quality and Respiratory Health among Malay Preschool Children in Selangor. *BiomedRes Int*, 1-8.
- Mors, J. L., Ter Hensen, M. S., Loomans, M. G., Boerstra, A. C. (2011). Adaptive thermal comfort in primary school classrooms: Creating and validating PMV-based comfort charts. *Build. Environ*. 46 (12): 2454–2461.
- Mumovic, D., Clement, S., Griffiths (2009). Winter indoor air quality, thermal comfort and acoustic performance of newly built secondary schools in England. *Building and Environment*. 44 (7): 1466 – 1477.
- Nazariah, S.S., Juliana, J. and Abdah, M.A. (2013). Interleukin-6 via Sputum Induction as Biomarker of Inflammation for Indoor Particulate Matter among Primary School Children in Klang Valley, Malaysia. *Glob J Health Sci*. 5(4): 93-105.
- Noor Hisyam, N.H. and Juliana, J. (2014). Association between Indoor PM₁₀, PM_{2.5}, NO₂ with Airway Inflammation among Preschool Children at Industrial and Sub-urban Areas. *Adv. Environ. Biol*. 8 (15): 149 - 159.
- Norhayani A.J., Juliana, J. Anis S.K., Mohd H. I. (2015). Exposure to Air Pollutants (PM₁₀, NO₂, SO₂ and VOCs) on the Lung Functions among School Children Living nearby the Petrochemical Industry Area in Kemaman, Terengganu. *Adv.Environ. Biol*, 9(26): 55-63
- Nur Aida Aziz, Juliana Jalaludin and Suhaili Abu Bakar. (2014). Indoor Air Pollutants. Exposure and the Respiratory Inflammation (Feno) among Preschool children in Hulu Langat, Selangor. *Adv. Environ. Biol*. 8 (15): 164-170.
- Nur Azwani Mohd Nor Rawi, Juliana Jalaludin, & Poh Choo Chua. (2014). Indoor Air Quality and Respiratory Health among Malay Preschool Children in Selangor. *Bio Med Research International*, Article ID 248178.
- Nurul, F., Juliana, J., Titi, R.H., & Zailina, H. (2010). Exposure of Indoor Air Pollutants (PM_{2.5}, PM₁₀, Ultrafine Particle, CO, VOCs, CO₂ and NO₂), and Lung Function of School Children in Klang Valley. *National Conference on Environmental and Health*. 75: 530 - 539.

- Nur Fazhilah, A.R., Juliana, J. and Sharifah, N.S.I. (2012). Health among Children Living in Industrial Area Kajang, Selangor. *International Society for Environmental Epidemiology East Asia Chapter (ISEE-EAC) conference Renaissance Kuala Lumpur Hotel 15th–18th October 2012*.
- Nurul Anis Sofiah, F. and Juliana, J. (2013). Indoor Particulate Matter 2.5 (PM_{2.5}) and Lung Function among Children Living Near Busy Road in Cheras, Kuala Lumpur. *Health and the Environment Journal*. 4(2): 1-19.
- Pawanker, R., Baena-Cagnani, C.E., Bousquet, J., Canonica, G.W., Cruz, A.A., Kaliner, M.A., Lanier, B.Q. (2008). Allergy and chronic respiratory disease. *Journal of World Allergy Organization*. S4 - S17.
- Pegas, N. M. G. Evtyugina, C. A. Alves et al. (2010). Outdoor/indoor air quality in primary schools in Lisbon: a preliminary study. *Qu'ímica Nova*. 33(5): 1145–1149.
- Pierse, N. L. Rushton, R. S. Harris, C. E. Kuehni, M. Silverman, and J. Grigg. (2006). locally generated particulate pollution and respiratory symptoms in young children, *Thorax*. 61(3): 216–220.
- Phlegm (2007). In *The American Heritage Medical Dictionary*. Retrieved August 3, 2015, from <http://medical-dictionary.thefreedictionary.com/phlegm>
- Quah, B.S., Wan-Pauzi, I., Ariffin, N., Mazidah, A.R. (2005). Prevalence of asthma, eczema and allergic rhinitis Kota Bahru, Malaysia. *Respirology*. 10: 244-249.
- Rom, W. N. (2007). *Environmental and Occupational Medicine*. Lippincott Williams and Wilkins, Philadelphia, USA. 4: 1904. ISBN 10:0781762995
- Ross, M.A., Curtis, L., Scheff, P.A., Hryhorezuk, D.O., Ramakrishnan, V., Wadden, R.A., Persky, V.W., (2000). Association of asthma symptoms and severity with indoor bioaerosols. *Allergy*. 55: 705 - 711.
- Salleh, N.M., Kamaruzzaman S.N., Mahyuddin, N. (2013). Sick Building Symptoms among Children in private preschools in Malaysia. Association of different ventilation strategies. *Journal of Building Performance*. 4(1): 73- 81.
- Sanchez-Monedero, M.A. (2008). Effect of the aeration system on the levels of airborne microorganisms generated at wastewater treatment plants. *Water Research*. 42(14): 3739 - 3744.
- Sanchez-Monedero, M.A., E.I. Stentiford & C. Mondini. (2003), Biofiltration at composting facilities: Effectiveness for bioaerosol control. *Environmental Science & Technology*. 37(18): 4299 - 4303.

- Savilahti, R., Uitti, J., Roto, P., Laippala, P., Husman, T. (2001). Increased prevalence of atopy among Children exposed to Mold in a school building. *Allergy* 56: 175-179
- Sharifah, N.S. (2013). Indoor Air Pollutants (PM_{2.5}, PM₁₀& NO₂). Exposure and the level of Interleukin-6 and Tumor Necrosis Factor-Alpha among Primary School Children in Klang Valley. Thesis Submitted for the Degree Master of Science, University Putra Malaysia.
- Sundell, J. (2004). On the history of indoor air quality and health. *Indoor Air*. 14(57): 51 - 58.
- Tezara, C., Adam, N.M., Juliana, J., Mariana, M., Siregar, J.P. (2014). Monitoring of indoor air quality parameters in the day care centers in tropical country. *Int. Journal of Mech. Eng.* 3: 47- 52.
- Tippayawong, N., Khuntong, P., Nitatwichit, C., Khunatorn, Y., Tantakitti, C. (2009). Indoor/outdoor relationships of size-resolved particle concentrations in naturally ventilated school environments. *Building and Environment*. 44: 188 - 197.
- Unites States Environmental Protection Agency (EPA). (2010). Managing Asthma in the Utara. *Malaysian Journal of Public Health Medicine*, vol. 3(2): 23–32.
- Verreault, D., S. Moineau, & C. Duchaine. (2008). Methods for sampling of airborne viruses. *Microbiology and Molecular Biology*. 72(3): 413 – 444.
- Wargocki, P. & Wyon, D.P. (2007b). The effects of moderately raised classroom temperatures and classroom ventilation rate on the performance of schoolwork by children. *HVAC&R Research*. 13(2): 193 – 220.
- WHO (2009). WHO Guidelines for Indoor Air Quality: Dampness and mold World Health Organization *Copenhages*. 248.
- Wong, N.H. & Khoo, S.S. (2003). Thermal comfort in classrooms in the tropics. *Energy and Buildings*. 35(4): 337 – 351.
- World Health Organization (2005). The world health report 2000 health systems. Improving performance. World Health Organization.
- World Health Organization (2005). Guidelines for concentration and exposure-response measurements of fine and ultra-fine particulate matter for use in epidemiological studies. Geneva: World Health Organization.
- World Health Organization (WHO) (2010), Strategic approaches to indoor air policy-making: WHO European center for environment and health: Bilthoven.
- World Health Organization (WHO), (2005). Effects of Air pollution on Children's Health and Development. Europe: WHO.

World Health Organization (WHO). (2010). WHO guidelines for indoor air quality: selected pollutants. Copenhagen: WHO Regional Office for Europe.

Yang, D.S., Son, K.C. and Kays, S.J. (2009). Volatile organic compounds emanating from indoor ornamental plants. *Hort. Science* 44: 396 - 400.

Yahaya N.A. and Jalaludin, J. (2014). Exposure to Indoor PM₁₀ and Volatile Organic Compounds and Its Association to Respiratory Health among Preschool Children in Urban and Rural Area. *Springer*. 3(4): 13-6.

Zhang, L., Chew, F.T., Soh, S.Y., Yi, F.C., Ferm, M. (2006). Current asthma and respiratory symptoms among pupils in Shanghai, China influence on building ventilation, nitrogen dioxide, ozone, and formaldehyde in classrooms. *Indoor Air*. 16: 454-464.



UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : _____

TITLE OF THESIS / PROJECT REPORT :

INDOOR AIR QUALITY AND RESPIRATORY HEALTH IMPLICATION AMONG MALAY
PRE-SCHOOL CHILDREN IN PUCHONG AND HULU- LANGAT, SELANGOR, MALAYSIA

NAME OF STUDENT: ONWUSEREAKA CYNTHIA OLUCHI

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (v)

☐

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

☐

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

☐

OPEN ACCESS

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

☐

PATENT

Embargo from _____ until _____
(date) (date)

Approved by:

(Signature of Student)
New IC No/ Passport No.:

Date :

(Signature of Chairman of Supervisory Committee)
Name:

Date :

[Note : If the thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted.]