



UNIVERSITI PUTRA MALAYSIA

***HEALING PROPERTIES OF EPIDERMAL GROWTH FACTOR AND
TOCOTRIENOL-RICH FRACTION FORMULATION IN DEEP PARTIAL-
THICKNESS BURN WOUND MODEL***

GUO HUIFANG

FPSK(p) 2018 2



**HEALING PROPERTIES OF EPIDERMAL GROWTH FACTOR AND
TOCOTRIENOL-RICH FRACTION FORMULATION IN DEEP
PARTIAL-THICKNESS BURN WOUND MODEL**

By

GUO HUIFANG

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfillment of the Requirements for the Degree of
Doctor of Philosophy**

November 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Doctor of Philosophy

**HEALING PROPERTIES OF EPIDERMAL GROWTH FACTOR AND
TOCOTRIENOL-RICH FRACTION FORMULATION IN DEEP
PARTIAL-THICKNESS BURN WOUND MODEL**

By

GUO HUIFANG

November 2017

Chairman : Huzwah Binti Khaza'ai, PhD
Faculty : Medicine and Health Sciences

Burns are one of the most devastating injuries in the world, divided into superficial, partial-thickness (superficial or deep partial-thickness) and full-thickness burns. The current treatment of deep partial-thickness burns is mainly focused on preventing wound infections and less attention to the burn healing process. Therefore, it is imperative to develop medications aimed at promoting the stage of wound healing. It has been shown that epidermal growth factor (EGF) can promote the proliferation of various cells and it is known that tocotrienol-rich fraction (TRF) from palm oil has a strong antioxidant activity. Thus, a combination of EGF and TRF is included in the present formulation (EGF+TRF cream), which is expected to accelerate wound healing and inhibit oxidative stress in the burn. Deep partial-thickness burn wounds were produced on the dorsal part of Sprague-Dawley rats. Animals were then randomly divided into six groups: untreated, treated with Silverdin® cream, base cream, base cream with c% EGF, base cream with 3 % TRF or base cream with c% EGF and 3% TRF, respectively. Creams were applied once daily for 21 consecutive days. Digital images were captured daily for macroscopic evaluation. Six animals from each group were sacrificed on the 3rd, 7th, 11th, 14th and 21st day post-burn, harvesting skin tissues with the wound for histological, cellular, biochemical and gene expression analysis. Prior to the research of the EGF+TRF formulation in wound healing, a new apparatus was designed to create burn wounds. With this device, uniform deep partial-thickness burn was achieved with a contact temperature of 70°C, a fixed pressure of 300 g, and duration of 10 seconds. In addition, a novel histological scoring system was developed to make the evaluation process more systematic and reproducible. The chronological events in burn healing were initiated by the recruitment of neutrophils to the wound site on the 3rd

day post-burn followed by the up-regulation of tumor necrosis factor- α (*TNF- α*) expression and rapid synthesis of collagens to facilitate leukocytes adhesion and crust attachment. The EGF+TRF treatment preceded this process and on the 7th day post-burn, wound contraction accelerated and the dermal healing was sped up by the proliferation of myofibroblasts and augmented expression of *Collagen-1*. By down regulation of interleukin 6 (*IL-6*) and *TNF- α* expression on Day 11 post-burn, the EGF+TRF treatment expedited the wound contraction and entering the re-modelling phase and arriving epithelialization on Day 14. This formulation was capable of promoting the recovery of the dermis and attenuate the oxidative stress by down regulating the expression of *IL-6*, *TNF- α* , inducible nitric oxide synthase (*iNOS*), matrix metalloproteinase-2 (*MMP-2*), tissue inhibitor of metalloproteinase-2 (*TIMP-2*), transforming growth factor- β 1 (*TGF- β 1*), vascular endothelial growth factor-A (*VEGF-A*) and *Collagen-1*. Finally, by Day 21 post-burn, the treatment of EGF+TRF formulation could advance the restoration of the epidermis and dermis and cut down the oxidative stress by down regulating the expression of *IL-6* and *iNOS*. In conclusion, all results from the present study fully supported the beneficial application of EGF+TRF formulation in the treatment of deep-partial thickness burn, adhered to the mechanisms of burn healing process.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**CIRI PENYEMBUHAN FORMULASI FAKTOR PERTUMBUHAN
EPIDERMAL DAN FRAKSI KAYA TOKOTRIENOL DALAM MODEL LUKA
KETEBALAN SEPARA TERBAKAR**

Oleh

GUO HUIFANG

November 2017

Pengerusi : Huzwah Binti Khaza'ai, PhD
Fakulti : Perubatan dan Sains Kesihatan

Luka terbakar adalah satu kecederaan yang dahsyat di dunia terbahagi kepada kebakaran superfisial, ketebalan separa dan ketebalan penuh. Rawatan terkini untuk luka terbakar ketebalan separa lebih mengutamakan pencegahan jangkitan luka berbanding proses penyembuhan. Oleh itu, penemuan rawatan yang berfokus kepada penggalakan proses penyembuhan luka adalah sangat penting. Faktor pertumbuhan epidermis (EGF) telah terbukti dapat membantu proses percambahan sel serta, fraksi kaya tokotrienol (TRF) daripada sawit mengandungi 25% α -tokoferol dan 75 % tokotrienol yang mempunyai aktiviti antioksidan yang tinggi. Oleh itu, kombinasi EGF dan TRF yang digunakan dalam formulasi (krim EGF+TRF) dijangka dapat mempercepatkan penyembuhan luka dan menghalang tekanan oksidatif dalam luka terbakar. Luka terbakar ketebalan separa telah diwujudkan di bahagian dorsal tikus *Sprague-Dawley*. Tikus dibahagikan secara rawak kepada enam kumpulan: tanpa rawatan, rawatan dengan krim *Silverdin*, krim asas, krim asas dengan c% EGF, krim asas dengan 3% TRF atau krim asas dengan c% EGF dan 3% TRF. Krim diaplikasi setiap hari selama 21 hari berturut. Imej digital diambil setiap hari untuk penilaian makroskopik. Enam tikus dari setiap kumpulan dikorbankan pada hari ke-3, ke-7, ke-11, ke-14 dan ke-21 setelah dirangsang luka terbakar, lalu tisu diambil dari luka untuk analisis histologi, selular, biokimia dan ekspresi gen. Satu radas baru untuk mewujudkan luka terbakar juga telah direka. Arus yang stabil dan terkawal, tekanan tetap, suhu yang tepat dan kemudahan operasi yang selamat dan tepat sebagai teknik untuk mendapatkan luka ketebalan separa yang sekata dengan suhu 70°C, tekanan 300g selama 10 saat. Tambahan lagi, sistem penskoran histologi yang unik telah dicipta. Sistem ini memudahkan untuk memadankan skor histologi dengan paras

kesembuhan epidermis dan dermis, maka proses penilaian adalah lebih sistematis dan *boleh direproduksi*. Kronologi penyembuhan luka terbakar bermula dengan pengumpulan neutrofil di tempat luka pada hari ke-3 selepas induksi luka, kemudian *peningkatan* ekspresi *TNF- α* dan penghasilan kolagen yang pantas untuk membantu infiltrasi leukosit dan *lekatan kuping luka*. Rawatan EGF +TRF mendahului proses ini dan pengecutan luka terlihat lebih cepat pada hari ke-7 selepas induksi luka, diikuti oleh penyembuhan dermis yang lebih pantas disebabkan oleh percambahan miofibroblas dan peningkatan ekspresi *kolagen-1*. Dengan penurunan ekspresi *IL-6* dan *TNF- α* pada hari ke-11 selepas induksi luka, didapati rawatan EGF+TRF meningkatkan pengecutan luka dan memasuki proses pembentukan semula lalu mencapai *epitialisasi* pada hari ke-14. Formulasi ini mampu meningkatkan penyembuhan dermis dan mengurangkan tekanan oksidatif dengan menurunkan ekspresi *IL-6*, *TNF- α* , *iNOS*, *MMP-2*, *TIMP-2*, *TGF- β 1*, *VEGF-A* and *Collagen-1*. Akhir sekali, selepas hari ke-21 luka, rawatan formulasi EGF+TRF mampu mempercepatkan proses pemulihan epidermis dan dermis serta menurunkan tekanan oksidatif melalui *penurunan IL-6* dan *iNOS*. Sebagai kesimpulan, kesan formulasi EGF+TRF pada ketebalan separa luka terbakar yang dinilai di peringkat makroskopik, histologi, selular, biokimia dan ekspresi gen, hasil dapatan kajian menyokong manfaat aplikasi dalam penyembuhan luka terbakar seiring dengan mekanisme proses penyembuhan luka.

ACKNOWLEDGEMENTS

First of all, I would like to convey my sincerest gratitude to my supervisor, Dr. Huzwah Binti Khaza'ai. She, who has provided invaluable advice when I am stuck, who has encouraged me to move on when I am depressed and who has illuminated the way ahead when I am lost. Without her guidance, none of this would have been possible.

I am also deeply indebted to the members of my supervisory committee, Dr. Razana Binti Mohd Ali for her constructive suggestions and subtle guidance in histological analysis and Dr. Roslida Binti Abd Hamid who shares knowledge and advices on angiogenesis.

Special thanks are due to the graduate students of biochemistry lab, Nurliyana Najwa Md Razip, Aisha Ali Chiroma, Afifah Abd Jalil and Amirah Salwani Zaulkffali. It is my honor to have the opportunity to work in such a harmonious team. I will never forget the happy moments we experienced together.

Many thanks to Asma Zaini for her previous creative work in the exploration of devices and procedures for inducing burn wound; to Fatimah Kasim for her assistance in inducing burn wounds; to Juita Chupri and Zamzarina Ahmad Bajari for their helps in technical support in hematoxylin and eosin staining as well as Dr. Ling King Hwa and Leong Jia Wen for their knowledge and experience in real time PCR work.

My sincere appreciation and thanks are also extended to my work unit-Chengde Medical University for the opportunity to further my study as well as the financial supports. In addition, I am most grateful to my colleagues from China and friends in Malaysia. Life abroad is full of joy and enrichment with your accompany.

Last but not least, it is more than a word of thanks that I owe to my mother and father for their continuous support and understanding. Specifically, I am immensely thankful for the everlasting love and respect of my husband, Zhang Yu-Fei, who shares every moment of joy and pain in my life. Not to be forgotten my dear sister as well as my lovely nephew.

Life is a journey; thank all of the people who have ever helped me in my life. You are the ones that make this journey wonderful.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Huzwah binti Khaza'ai, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Razana binti Mohd Ali, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Roslida binti Abd Hamid, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No: GUO HUIFANG, GS38505

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xvii
 CHAPTER	
 1 INTRODUCTION	 1
1.1 Background of the study	1
1.2 Problem statement	2
1.3 Research objectives	3
1.3.1 General objective	3
1.3.2 Specific objective	3
1.4 Hypothesis	3
 2 LITERATURE REVIEW	 4
2.1 The skin	4
2.1.1 Anatomy of the human skin	4
2.1.2 Rats as a model to study wound healing	7
2.2 Burns	8
2.2.1 Overview of burns	8
2.2.2 Local changes of burns	8
2.2.3 Classification of burns	9
2.2.4 Underlying mechanism of burn wound healing process	11
2.2.5 Factors regulating burn wound healing process	13
2.2.6 Inflammation and burns	15
2.2.7 Nitric oxide and burns	17
2.2.8 Angiogenesis and burns	18
2.2.9 Tissue re-modelling and burns	21
2.2.10 Management of burns	23
2.3 EGF	24
2.3.1 Introduction of EGF	24
2.3.2 EGF receptor	24
2.3.3 EGF and wound healing	26
2.4 Vitamin E	27
2.4.1 Components of Vitamin E	27
2.4.2 Biological function of TRF	28
2.4.3 Vitamin E and scar healing	29

3	BURN MODEL	30
3.1	Introduction	30
3.2	Materials and methods	31
3.2.1	Materials	31
3.2.2	Experimental design	34
3.2.3	Methods	35
3.3	Results	37
3.3.1	Optimization of inducing deep partial-thickness burn	37
3.3.2	Macroscopic and microscopic evaluation of deep partial-thickness burn wound over time	40
3.4	Discussion	58
3.4.1	Optimization of inducing deep partial-thickness burn	58
3.4.2	Macroscopic evaluation of deep partial-thickness burn wound over time	59
3.4.3	Histological score grade	59
3.5	Conclusion	61
4	MACROSCOPIC AND MICROSCOPIC ASSESSMENT OF EGF+TRF FORMULATION IN RATS WITH DEEP PARTIAL-THICKNESS BURN WOUND	62
4.1	Introduction	62
4.2	Materials and methods	63
4.2.1	Materials	63
4.2.2	Experimental design	63
4.2.3	Methods	65
4.3	Results	69
4.3.1	Macroscopic evaluation	69
4.3.2	Microscopic evaluation	76
4.4	Discussion	85
4.5	Conclusion	88
5	EFFECT OF EGF+TRF FORMULATION ON OXIDATIVE STRESS IN RATS WITH DEEP PARTIAL-THICKNESS BURN	89
5.1	Introduction	89
5.2	Materials and methods	90
5.2.1	Materials	90
5.2.2	Experimental design	90
5.2.3	Methods	90
5.3	Results	91
5.3.1	Measurement of lipid peroxidation	91
5.3.2	Measurement of nitrite	92
5.4	Discussion	94
5.5	Conclusion	95

6	WOUND HEALING RELATED GENE ANALYSIS	96
6.1	Introduction	96
6.2	Materials and methods	97
6.2.1	Materials	97
6.2.2	Experimental design	97
6.2.3	Methods	97
6.3	Results	100
6.3.1	Temporal changes of wound healing related genes in the skin tissue with deep partial-thickness burn	100
6.3.2	Expression of wound healing related genes in all the groups over time post-burn	104
6.4	Discussion	114
6.4.1	Temporal changes of wound healing related genes in the skin tissue with deep partial-thickness burn	114
6.4.2	Expression profile of wound healing related genes in all the groups over time post-burn	116
6.5	Conclusion	119
7	GENERAL DISCUSSION, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	120
7.1	General discussion	120
7.2	Conclusion	126
7.3	Recommendations for future research	126
	REFERENCES	127
	APPENDICES	146
	BIODATA OF STUDENT	162
	LIST OF PUBLICATIONS	163

LIST OF TABLES

Table	Page
2.1 Description of clinical characteristics of burn wounds of various depth	10
2.2 Classification of matrix metalloproteinases	22
3.1 Temperature, duration, and contact pressure for deep partial-thickness burns in the literature	32
3.2 Experimental design for optimizing the temperature to induce deep partial-thickness burn	34
3.3 Experimental design for evaluating macroscopic and microscopic change of rats with deep-partial thickness burn in the healing process	35
3.4 Description of histological score 0 for the epidermis	44
3.5 Description of histological score 1 for the epidermis	45
3.6 Description of histological score 2 for the epidermis	46
3.7 Description of histological score 3 for the epidermis	47
3.8 Description of histological score 4 for the epidermis	48
3.9 Description of histological score 5 for the epidermis	49
3.10 Description of histological score 6 for the epidermis	50
3.11 Description of histological score 7 for the epidermis	51
3.12 Histological score range for the dermis	53
3.13 Images of histological score for the dermis	54
3.14 Temporal changes of histological score of the epidermis and dermis in the skin tissue with deep partial-thickness burn	58
4.1 Design of experimental groups	64
4.2 Experimental design for macroscopic and microscopic evaluations of the effect of the EGF+TRF formulation on deep partial-thickness burns in rats	64

4.3	Clinical evaluations of wounds in all the groups during the healing process	75
4.4	Histological scores of the epidermis in all the groups over time post-burn	78
4.5	Histological scores of the dermis in all the groups over time post-burn	79
4.6	Temporal changes of the number of different cell types in the skin tissue with deep partial-thickness burn	80
4.7	The number of adipose cells in all the groups over time post-burn	81
4.8	The number of neutrophils in all the groups over time post-burn	82
4.9	The number of lymphocytes in all the groups over time post-burn	84
4.10	The number of myofibroblasts in all the groups over time post-burn	85
6.1	Primers used in Q-PCR	99
6.2	Genes involved in Q-PCR analysis	100
7.1	The summary of significant findings in the present study	122

LIST OF FIGURES

Figure	Page
2.1 Structure of the skin	4
2.2 Schematic view of the cross-section of the upper part of the human skin showing the epidermis and part of the dermis	5
2.3 Hair follicle structures	7
2.4 Zones of burn injury	9
2.5 Migration of inflammatory cells correlated with the phase of wound healing	16
2.6 Demonstration of VEGF's potential role in a healing wound	20
2.7 Downstream regulation of EGFR	26
2.8 Natural forms of vitamin E	27
3.1 Burn wound creation apparatus	33
3.2 Microscopic assessment of burn depth at different temperatures on Day 3 post-burn	39
3.3 Macroscopic changes of deep partial-thickness burn wound over time in Sprague-Dawley rats	41
3.4 Wound contraction rates in rats with deep partial-thickness burn over time	42
4.1 General appearance of the wound immediately after inducing burn	66
4.2 Photographs of wound sections showing cells that were identified and counted	68
4.3 Gross appearance of the wound in all the groups on Day 3 post-burn	70
4.4 Gross appearance of the wound in all the groups on Day 7 post-burn	71
4.5 Gross appearance of the wound in all the groups on Day 11 post-burn	72

4.6	Gross appearance of the wound in all the groups on Day 14 post-burn	73
4.7	Gross appearance of the wound in all the groups on Day 21 post-burn	74
4.8	Wound contraction rates in all the groups over time post-burn	76
5.1	MDA content in all the groups over time post-burn	92
5.2	Nitrite content in all the groups over time post-burn	93
6.1	Temporal changes of wound healing related genes in the skin tissue with deep partial-thickness burn	102
6.2	Expression profiles of <i>IL-6</i> in all the groups over time post-burn	105
6.3	Expression profiles of <i>TNF-α</i> in all the groups over time post-burn	106
6.4	Expression profiles of <i>iNOS</i> in all the groups over time post-burn	107
6.5	Expression profiles of <i>VEGF-A</i> in all the groups over time post-burn	108
6.6	Expression profiles <i>TGF-β1</i> in all the groups over time post-burn	109
6.7	Expression profiles <i>MMP-2</i> in all the groups over time post-burn	110
6.8	Expression profiles <i>TIMP-2</i> in all the groups over time post-burn	111
6.9	Expression profiles of <i>Collagen-1</i> in all the groups over time post-burn	112
6.10	Expression profiles of <i>EGFR</i> in all the groups over time post-burn	113
7.1	Burn wound healing signal pathway	124

LIST OF ABBREVIATIONS

bFGF	Basic fibroblast growth factor
CC	Cysteine-cysteine
cDNA	Complementary deoxyribonucleic acid
CVIU	Chronic venous insufficiency ulcer
CXC	Cysteine-X amino acid-cysteine
DAG	Diacylglycerol
ECM	Extra cellular matrix
EGF	Epidermal growth factor
EGFR	Epidermal growth factor receptor
eNOS	Endothelial nitric oxide synthase
GAP	GTPase-activating protein
GAPDH	Glyceraldehyde-3-Phosphate Dehydrogenase
GDP	Guanosine diphosphate
Grb-2	Growth factor receptor-bound protein 2
GTP	Guanosine-5'-triphosphate
H&E	Hematoxylin and eosin
H ₂ O ₂	Hydroxyl peroxide
HER1	Human epidermal growth factor 1
HICs	High-income countries
IGF-1	Insulin-like growth factor-1
IL-10	Interleukin-10
IL-1 α	Interleukin-1 α

IL-1 β	Interleukin-1 β
IL-4	Interleukin-4
IL-6	Interleukin-6
IL-8	Interleukin-8
iNOS	Inducible nitric oxide synthase
IP3	Inositol-1, 4, 5-triphosphate
LDL	Low-density lipoprotein
LMICs	Low-to-middle-income countries
LMWP	Low-molecular-weight protamine
MAP	Mitogen-activated protein
MDA	Malondialdehyde
MMP	Matrix metalloproteinase
MMPis	Matrix metalloproteinase inhibitors
MT-MMP	Membrane-type matrix metalloproteinase
NF-KB	Nuclear factor kappa B
nNOS	Neuronal nitric oxide synthase
NO	Nitric oxide
NO ₂	Nitrite
NO ₃	Nitrate
NOS	Nitric oxide synthase
O ₂ ⁻	Superoxide anion radical
OH $\cdot$	Hydroxyl radical
ONOO ⁻	Peroxynitrite
PBO-TRF	TRF from rice bran oil

PDGF	Platelet-derived growth factor
PEG	Polyethylene glycol
PGK1	Phosphoglycerate kinase 1
PI-3	Phosphatidylinositol-3
PLC- γ	Phospholipase C- γ
PLGF	Placenta growth factor
PO-TRF	TRF from palm oil
Q-PCR	Quantitative real time polymerase chain reaction
RNS	Reactive nitrogen species
ROS	Reactive oxygen species
S.D.	Standard error
SOD	Superoxide dismutase
SOS	Son of sevenless
SSD	Silver sulfadiazine
T-cell	Thymus-derived cell
TGF- β	Transforming growth factor- β
TIMP	Tissue inhibitor of metalloproteinase
TNF- α	Tumor necrosis factor- α
TRF	Tocotrienol-rich fraction
VEGF	Vascular endothelial growth factor

CHAPTER 1

INTRODUCTION

1.1 Background of the study

Burns are defined as skin lesions caused mainly by heat or other acute trauma including scalds, contact burns and flame burns. In the world, approximately 300,000 people die each year from burns (Peck, 2012). The incidence of burns in low- and middle- income countries is higher than in the developed countries (Atiyeh et al., 2009). Depending on the burn depth, burn patients may experience many mental and physical complications such as shock, trauma, infection, electrolyte imbalance and respiratory failure (Evers et al., 2010). In addition to physical complications, the patient may be physically and emotionally suffering from surgery and long-term hospitalization, and some may even have to endure physical scars for a lifetime (Grivna et al., 2014).

Burn injuries lead to different degrees of vascular damage (Shakespeare, 2001). Some blood vessels can be completely coagulated while others may be moderately restricted, leading to different tissue change. The local changes in burn wounds are divided into three zones. They are: i) zone of coagulation—the central point with maximum destruction; ii) zone of hyperemia—the outermost zone without any structural changes; iii) zone of stasis—the area between the zone of coagulation and hyperemia (Evers et al., 2010).

According to the depth of the injured tissue, burns are classified as superficial, partial-thickness and full-thickness burn (Frantz & Byers, 2011). As evidenced by histologic studies, burns are dynamic. In other words, the depth of the burn may further deepen over time. But usually it stabilizes within about 3 days post-burn (Pereira et al., 2012). Necrosis of the stasis area may be responsible for this progress.

Burn wound healing process is similar to the general wound healing, which is the interaction of several cellular and biochemical components of synchronized and macroscopic coordination including hemostasis, inflammation, proliferation and tissue re-modelling (Enoch & Leaper, 2008). In the hemostatic phase, the formation of blood clots prevents further blood loss and helps to attract the inflammatory cells into the wound area (Velnar et al., 2009). In the inflammatory phase, a variety of inflammatory cells accumulate in the wound site and remove bacteria, foreign particles and damaged tissue to protect the tissue from infection (Shakespeare, 2001). Then the wound

moves to the third stage, the proliferative phase, where new blood vessels are growing, fibroblasts are migrating into the wound site and producing matrix proteins and keratinocytes are proliferating, migrating and covering the wound. When the wound is closed, it is usually considered to have achieved maximum healing. At the same time, the re-modelling phase begins and may last from months to years. At this stage, the fibrous tissue laid in the wound site is modified. Any aberration in the process can lead to poor healing processes such as chronic wounds, or excessive healing including hypertrophic scars and keloids (Tuan & Nichter, 1998).

Many growth factors are involved in the wound healing process. Epidermal growth factor (EGF) is a small peptide with 53 amino acids (Değim et al., 2011). Since 1989, topical administration of EGF has been used to increase the wound healing process of various wounds and proved to be safe and beneficial (Berlanga-Acosta et al., 2009). EGF achieves its biological function by binding to specific receptors on the cell membrane (Süzük et al., 2011)

The vitamin E extracted from crude palm oil is mainly composed of a mixture of tocotrienols and some tocopherols and referred as tocotrienol-rich fraction (TRF) (Wu et al., 2008). It has been reported that palm oil TRF has a strong activity in anti-oxidation (Minhajuddin et al., 2005), prevention and treatment of cancer (Srivastava & Gupta, 2006) and reduction in serum cholesterol levels (Qureshi et al., 2002). In addition, it is reported that TRF is more effective than α -tocopherol in reducing oxidative damage in brain mitochondria (Kamat & Devasagayam, 1995) and inhibiting protein oxidation and lipid peroxidation in liver microsomes (Kamat et al., 1997). Although the effect of TRF has been shown in cellular biologic experiments, to the best of our knowledge, TRF studies in burn wound healing is limited, therefore we take the advantage of exploring the combination of EGF and TRF formulation for the treatment of deep partial-thickness burns.

1.2 Problem statement

Many topical medications have been used in the treatment of burns, such as antiseptic agents, antimicrobial agents and enzymatic debriding agents (Leon-Villapalos et al., 2008). Silver sulfadiazine (SSD) is an antibacterial agent and acts on the bacterial wall (Lee et al., 2005). Soon after its discovery, it became a standard local treatment for patients with partial-thickness burns (Jurjus et al., 2007). However, it has been suggested that the local application of the SSD is either no better or worse than the control dressing in preventing wound infection and promoting burn healing (Aziz et al., 2012). Moreover, the benefits of SSD treatment have gone beyond the risk (Miller et al., 2012). Therefore, it is necessary to explore more efficient treatments for burn wounds.

EGF with a small polypeptide of 53 amino acids is of interest in the present study. The use of topical application of growth factor therapy has assisted in minimizing the incidence of burn wound scar and long-term healing. In this study, the combination of EGF and TRF with potent antioxidant activity was used for the treatment of deep partial-thickness burn wounds. The mechanism for understanding these two elements involved in the healing process was elucidated.

1.3 Research objectives

1.3.1 General objective

To elucidate the effect and mechanism of EGF+TRF formulation in the healing process of deep partial-thickness burn in rat model

1.3.2 Specific objective

- To evaluate the effect of EGF+TRF formulation on the macroscopic and microscopic changes involved in the deep partial-thickness burn wound healing process
- To determine the effect of EGF+TRF formulation on oxidative stress involved in the deep partial-thickness burn wound healing process
- To elucidate the effect of EGF+TRF formulation on the wound healing-related genes involved in the deep partial-thickness burn wound healing process

1.4 Hypothesis

The EGF+TRF formulation has synergistic effects in treating the deep partial-thickness burn.

REFERENCES

- Ahn, C. S., & Maitz, P. K. (2012). The true cost of burn. *Burns*, 38(7): 967-974.
- Aktan, F. (2004). iNOS-mediated nitric oxide production and its regulation. *Life Sciences*, 75(6): 639-653.
- Alemdaroğlu, C., Değim, Z., Çelebi, N., Zor, F., Öztürk, S., & Erdoğan, D. (2006). An investigation on burn wound healing in rats with chitosan gel formulation containing epidermal growth factor. *Burns*, 32(3): 319-327.
- Alsarra, I. A. (2009). Chitosan topical gel formulation in the management of burn wounds. *International Journal of Biological Macromolecules*, 45(1): 16-21.
- Ancelin, M., Chollet-Martin, S., Hervé, M. A., Legrand, C., El Benna, J., & Perrot-Appianat, M. (2004). Vascular endothelial growth factor VEGF189 induces human neutrophil chemotaxis in extravascular tissue via an autocrine amplification mechanism. *Laboratory Investigation*, 84(4): 502-512.
- Angel, P., Szabowski, A., & Schorpp-Kistner, M. (2001). Function and regulation of AP-1 subunits in skin physiology and pathology. *Oncogene*, 20(19): 2413-2423.
- Ashcroft, G. S., Jeong, M. J., Ashworth, J. J., Hardman, M., Jin, W., Moutsopoulos, N., Wild, T., McCartney-Francis, N., Sim, D., McGradt, G., Song, X. & Wahl, S. M. (2012). Tumor necrosis factor-alpha (TNF- α) is a therapeutic target for impaired cutaneous wound healing. *Wound Repair and Regeneration*, 20(1): 38-49.
- Atiyeh, B., Masellis, A., & Conte, C. (2009). Optimizing burn treatment in developing low-and middle-income countries with limited health care resources (part 1). *Annals of Burns and Fire Disasters*, 22(3): 121-125.
- Auerbach, R., Lewis, R., Shinnars, B., Kubai, L., & Akhtar, N. (2003). Angiogenesis assays: a critical overview. *Clinical Chemistry*, 49(1): 32-40.
- Aziz, Z. S. F. A., Abu, S. F., & Chong, N. J. (2012). A systematic review of silver-containing dressings and topical silver agents (used with dressings) for burn wounds. *Burns*, 38(3): 307-318.
- Bae, Y. S., Kang, S. W., Seo, M. S., Baines, I. C., Tekle, E., Chock, P. B., & Rhee, S. G. (1997). Epidermal growth factor (EGF)-induced generation of hydrogen peroxide Role in EGF receptor-mediated tyrosine

phosphorylation. *Journal of Biological Chemistry*, 272(1): 217-221.

Baggiolini, M., Dewald, B., & Moser, B. (1997). Human chemokines: an update. *Annual Review of Immunology*, 15(1): 675-705.

Bao, P., Kodra, A., Tomic-Canic, M., Golinko, M. S., Ehrlich, H. P., & Brem, H. (2009). The role of vascular endothelial growth factor in wound healing. *Journal of Surgical Research*, 153(2): 347-358.

Barrientos, S., Stojadinovic, O., Golinko, M. S., Brem, H., & Tomic-Canic, M. (2008). Growth factors and cytokines in wound healing. *Wound Repair and Regeneration*, 16(5): 585-601.

Berlanga-Acosta, J., Gavilondo-Cowley, J., López-Saura, P., González-López, T., Castro-Santana, M. D., López-Mola, E., Guillén-Nieto, G. & Herrera-Martinez, L. (2009). Epidermal growth factor in clinical practice-a review of its biological actions, clinical indications and safety implications. *International Wound Journal*, 6(5): 331-346.

Bertolino, P., Deckers, M., Lebrin, F., & ten Dijke, P. (2005). Transforming growth factor- β signal transduction in angiogenesis and vascular disorders. *CHEST Journal*, 128(6_suppl): 585S-590S.

Bigg, H. F., Shi, Y. E., Liu, Y. E., Steffensen, B., & Overall, C. M. (1997). Specific, high affinity binding of tissue inhibitor of metalloproteinases-4 (TIMP-4) to the COOH-terminal hemopexin-like domain of human gelatinase A. *Journal of Biological Chemistry*, 272(24): 15496-15500.

Birkedal-Hansen, H. (1995). Proteolytic re-modelling of extracellular matrix. *Current Opinion in Cell Biology*, 7(5): 728-735.

Birkenfeld, H. P., McIntyre, B. S., Briski, K. P., & Sylvester, P. W. (1996). Role of Protein Kinase C in Modulating Epidermal Growth Factor- and Phorbol Ester-Induced Mammary Epithelial Cell Growth in Vitro. *Experimental Cell Research*, 223(1): 183-191.

Bloodsworth, A., O'donnell, V. B., & Freeman, B. A. (2000). Nitric oxide regulation of free radical- and enzyme-mediated lipid and lipoprotein oxidation. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 20(7): 1707-1715.

Bogdan, S., & Klämbt, C. (2001). Epidermal growth factor receptor signaling. *Current Biology*, 11(8): R292-R295.

Brenner, M., & Hearing, V. J. (2008). The protective role of melanin against UV damage in human skin. *Photochemistry and Photobiology*, 84(3): 539-549.

Brew, K., & Nagase, H. (2010). The tissue inhibitors of metalloproteinases (TIMPs): an ancient family with structural and functional diversity.

Biochimica et Biophysica Acta (BBA)-Molecular Cell Research, 1803(1): 55-71.

Brew, K., Dinakarbandian, D., & Nagase, H. (2000). Tissue inhibitors of metalloproteinases: evolution, structure and function. *Biochimica et Biophysica Acta (BBA)-Protein Structure and Molecular Enzymology*, 1477(1): 267-283.

Bronaugh, R. L., Stewart, R. F., & Congdon, E. R. (1982). Methods for in vitro percutaneous absorption studies II. Animal models for human skin. *Toxicology and Applied Pharmacology*, 62(3): 481-488.

Butler, G. S., Apte, S. S., Willenbrock, F., & Murphy, G. (1999). Human tissue inhibitor of metalloproteinases 3 interacts with both the N- and C-terminal domains of gelatinases A and B. *Journal of Biological Chemistry*, 274(16): 10846-10851.

Chin, S. F., Ibahim, J., Makpol, S., Hamid, N. A. A., Latiff, A. A., Zakaria, Z., Mazlan, M., Yusof, Y. a. M., Karim, A.A. & Ngah, W. Z. W. (2011). Tocotrienol rich fraction supplementation improved lipid profile and oxidative status in healthy older adults: A randomized controlled study. *Nutrition & Metabolism*, 8: 42.

Chithra, P., Sajithlal, G. B., & Chandrakasan, G. (1998). Influence of Aloe vera on collagen characteristics in healing dermal wounds in rats. *Molecular and Cellular Biochemistry*, 181(1): 71-76.

Cohen, S. (1962). Isolation of a mouse submaxillary gland protein accelerating incisor eruption and eyelid opening in the new-born animal. *Journal of Biological Chemistry*, 237(5): 1555-1562.

Crawford, S. E., Stellmach, V., Murphy-Ullrich, J. E., Ribeiro, S. M., Lawler, J., Hynes, R. O., Boivin, G. P. & Bouck, N. (1998). Thrombospondin-1 is a major activator of TGF- β 1 in vivo. *Cell*, 93(7): 1159-1170.

Cuttle, L., Kempf, M., Phillips, G. E., Mill, J., Hayes, M. T., Fraser, J. F., Wang, X. Q. & Kimble, R. M. (2006). A porcine deep dermal partial thickness burn model with hypertrophic scarring. *Burns*, 32(7): 806-820.

Darby, I. A., Laverdet, B., Bonté, F., & Desmoulière, A. (2014). Fibroblasts and myofibroblasts in wound healing. *Clinical, Cosmetic and Investigational Dermatology*, 7: 301-311.

de Almeida, E. B., Cardoso, J. C., de Lima, A. K., de Oliveira, N. L., de Pontes-Filho, N. T., Lima, S. O., Souza, I. C. L. & de Albuquerque-Júnior, R. L. C. (2013). The incorporation of Brazilian propolis into collagen-based dressing films improves dermal burn healing. *Journal of Ethnopharmacology*, 147(2): 419-425.

- De, S., Fenton, J. E., & Jones, A. S. (2005). Matrix metalloproteinases and their inhibitors in non-neoplastic otorhinolaryngological disease. *The Journal of Laryngology & Otology*, 119(6): 436-442.
- Değim, Z., Çelebi, N., Alemdaroğlu, C., Deveci, M., Öztürk, S., & Özoğul, C. (2011). Evaluation of chitosan gel containing liposome-loaded epidermal growth factor on burn wound healing. *International Wound Journal*, 8(4): 343-354.
- Deryugina, E. I., Ratnikov, B., Monosov, E., Postnova, T. I., DiScipio, R., Smith, J. W., & Strongin, A. Y. (2001). MT1-MMP initiates activation of pro-MMP-2 and integrin $\alpha\beta 3$ promotes maturation of MMP-2 in breast carcinoma cells. *Experimental Cell Research*, 263(2): 209-223.
- Dokumcu, Z., Ergun, O., Celik, H. A., Aydemir, S., Sezak, M., Ozok, G., & Celik, A. (2008). Clostridial collagenase aggravates the systemic inflammatory response in rats with partial-thickness burns. *Burns*, 34(7): 935-941.
- Dovi, J. V., Szpaderska, A. M., & DiPietro, L. A. (2004). Neutrophil function in the healing wound: adding insult to injury? *Thrombosis and Haemostasis- Stuttgart*, 92: 275-280.
- Dröge, W. (2002). Free radicals in the physiological control of cell function. *Physiological Reviews*, 82(1): 47-95.
- Duansak, D., Somboonwong, J., & Patumraj, S. (2003). Effects of Aloe vera on leukocyte adhesion and TNF- α and IL-6 levels in burn wounded rats. *Clinical Hemorheology and Microcirculation*, 29(3, 4): 239-246.
- Dunnill, C., Patton, T., Brennan, J., Barrett, J., Dryden, M., Cooke, J., Leaper, D. & Georgopoulos, N. T. (2017). Reactive oxygen species (ROS) and wound healing: the functional role of ROS and emerging ROS -modulating technologies for augmentation of the healing process. *International Wound Journal*, 14(1): 89-96.
- Durmus, A. S., Han, M. C., & Yaman, I. (2009). Comparative evaluation of collagenase and silver sulfadiazine on burned wound healing in rats. *Firat Universitesi Saglik Bilimleri Veteriner Dergisi*, 23(3): 135-139.
- Eming, S. A., Werner, S., Bugnon, P., Wickenhauser, C., Siewe, L., Utermöhlen, O., Davidson, J. M., Krieg, T. & Roers, A. (2007). Accelerated wound closure in mice deficient for interleukin-10. *The American Journal of Pathology*, 170(1): 188-202.
- Enoch, S., & Leaper, D. J. (2008). Basic science of wound healing. *Surgery (Oxford)*, 26(2): 31-37.

- Evers, L. H., Bhavsar, D., & Mailänder, P. (2010). The biology of burn injury. *Experimental Dermatology*, 19(9): 777-783.
- Fingleton, B. (2008). MMPs as therapeutic targets—still a viable option? *Seminars in Cell & Developmental Biology*, 19(1): 61-68
- Finlay, G. A., Russell, K. J., McMahon, K. J., D'arcy, E. M., Masterson, J. B., FitzGerald, M. X., & O'connor, C. M. (1997). Elevated levels of matrix metalloproteinases in bronchoalveolar lavage fluid of emphysematous patients. *Thorax*, 52(6): 502-506.
- Fishel, R. S., Barbul, A., Beschoner, W. E., Wasserkrug, H. L., & Efron, G. (1987). Lymphocyte participation in wound healing. Morphologic assessment using monoclonal antibodies. *Annals of Surgery*, 206(1): 25-29.
- Fivenson, D. P., Faria, D. T., Nickoloff, B. J., Poverini, P. J., Kunkel, S., Burdick, M., & Strieter, R. M. (1997). Chemokine and inflammatory cytokine changes during chronic wound healing. *Wound Repair and Regeneration*, 5(4): 310-322.
- Forjuoh, S. N. (2006). Burns in low-and middle-income countries: a review of available literature on descriptive epidemiology, risk factors, treatment, and prevention. *Burns*, 32(5): 529-537.
- Frank, S., Hübner, G., Breier, G., Longaker, M. T., Greenhalgh, D. G., & Werner, S. (1995). Regulation of vascular endothelial growth factor expression in cultured keratinocytes. Implications for normal and impaired wound healing. *Journal of Biological Chemistry*, 270(21): 12607-12613.
- Frantz, K., & Byers, C. G. (2011). Thermal injury. *Compendium*, 33(12): E1-E6.
- Fraser, J. F., Cuttle, L., Kempf, M., & Kimble, R. M. (2004). Cytotoxicity of topical antimicrobial agents used in burn wounds in Australasia. *ANZ Journal of Surgery*, 74(3): 139-142.
- Gabbiani, G. (2003). The myofibroblast in wound healing and fibrocontractive diseases. *The Journal of Pathology*, 200(4): 500-503.
- Gainza, G., Villullas, S., Pedraz, J. L., Hernandez, R. M., & Igartua, M. (2015). Advances in drug delivery systems (DDSs) to release growth factors for wound healing and skin regeneration. *Nanomedicine: Nanotechnology, Biology and Medicine*, 11(6): 1551-1573.
- Gallucci, R. M., Simeonova, P. P., Matheson, J. M., Kommineni, C., Guriel, J. L., Sugawara, T., & Luster, M. I. (2000). Impaired cutaneous wound healing in interleukin-6-deficient and immunosuppressed mice. *The FASEB journal*, 14(15): 2525-2531.

- Garcia, X. Santos Heredero, Á. Izquierdo Hernandez, E. Pascual Peña, G. Bilbao Gonzalez de Aledo, C. Hamann, S. S. (2000). Experimental model for local application of growth factors in skin re-epithelialisation. *Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery*, 34(3): 199-206.
- Gillitzer, R., & Goebeler, M. (2001). Chemokines in cutaneous wound healing. *Journal of Leukocyte Biology*, 69(4): 513-521.
- Gokakin, A. K., Deveci, K., Kurt, A., Karakus, B. C., Duger, C., Tuzcu, M., & Topcu, O. (2013). The protective effects of sildenafil in acute lung injury in a rat model of severe scald burn: a biochemical and histopathological study. *Burns*, 39(6): 1193-1199.
- Greene, J., Wang, M., Liu, Y. E., Raymond, L. A., Rosen, C., & Shi, Y. E. (1996). Molecular cloning and characterization of human tissue inhibitor of metalloproteinase 4. *Journal of Biological Chemistry*, 271(48): 30375-30380.
- Grinnell, F. (1994). Fibroblasts, myofibroblasts, and wound Contraction. *The Journal of Cell Biology*, 124(4): 401-404
- Grivna, M., Eid, H. O., & Abu-Zidan, F. M. (2014). Epidemiology of burns in the United Arab Emirates: Lessons for prevention. *Burns*, 40(3): 500-505.
- Gutteridge, J. M. (1995). Lipid peroxidation and antioxidants as biomarkers of tissue damage. *Clinical Chemistry*, 41(12): 1819-1828.
- Guzik, T., Korbout, R., & Adamek-Guzik, T. (2003). Nitric oxide and superoxide in inflammation and immune regulation. *Journal of Physiology and Pharmacology*, 54(4): 469-487.
- Haake, A., Scott, G. A., & Holbrook, K. A. (2001). Structure and function of the skin: overview of the epidermis and dermis. In R. K. Freinkel & D. T. Woodley (Ed.), *The Biology of The Skin* (pp. 19-45). New York: The Parthenon Publishing Group Inc.
- Haase, I., Evans, R., Pofahl, R., & Watt, F. M. (2003). Regulation of keratinocyte shape, migration and wound epithelialization by IGF-1-and EGF-dependent signaling pathways. *Journal of Cell Science*, 116(15): 3227-3238.
- Hardwicke, J., Schmaljohann, D., Boyce, D., & Thomas, D. (2008). Epidermal growth factor therapy and wound healing—past, present and future perspectives. *The Surgeon*, 6(3): 172-177.
- Harper, D., Young, A., & McNaught, C. E. (2014). The physiology of wound healing. *Surgery (Oxford)*, 32(9): 445-450.

- Heldin, C. H., & Westermark, B. (1999). Mechanism of action and in vivo role of platelet-derived growth factor. *Physiological Reviews*, 79(4): 1283-1316.
- Hendriks F. M. (2005). *Mechanical behaviour of human epidermal and dermal layers in vivo*. (Unpublished doctoral dissertation). Technische Universiteit Eindhoven, The Netherlands.
- Henson, E. S., & Gibson, S. B. (2006). Surviving cell death through epidermal growth factor (EGF) signal transduction pathways: implications for cancer therapy. *Cellular Signaling*, 18(12): 2089-2097.
- Hernández-Sotomayor, S. T., & Carpenter, G. (1992). Epidermal growth factor receptor: elements of intracellular communication. *Journal of Membrane Biology*, 128(2): 81-89.
- Hettiaratchy, S., & Dziewulski, P. (2004). Pathophysiology and types of burns. *British Medical Journal*, 328(7453): 1427-1429.
- Horton, J. W. (2003). Free radicals and lipid peroxidation mediated injury in burn trauma: the role of antioxidant therapy. *Toxicology*, 189(1): 75-88.
- Hübner, G., Brauchle, M., Smola, H., Madlener, M., Fässler, R., & Werner, S. (1996). Differential regulation of pro-inflammatory cytokines during wound healing in normal and glucocorticoid-treated mice. *Cytokine*, 8(7): 548-556.
- Huo, Y., Qiu, W. Y., Pan, Q., Yao, Y. F., Xing, K., & Lou, M. F. (2009). Reactive oxygen species (ROS) are essential mediators in epidermal growth factor (EGF)-stimulated corneal epithelial cell proliferation, adhesion, migration, and wound healing. *Experimental Eye Research*, 89(6): 876-886.
- Huskić, J., Zaciragić, A., Babić, N., & Mulabegović, N. (2006). Nitric oxide in serum and renal tissue during compensatory renal hypertrophy in rats. *Bosnian Journal of Basic Medical Sciences*, 6(1): 46-49.
- Hynes, R. O. (2009). The extracellular matrix: not just pretty fibrils. *Science*, 326(5957): 1216-1219.
- Infanger, M., Schmidt, O., Kossmehl, P., Grad, S., Ertel, W., & Grimm, D. (2004). Vascular endothelial growth factor serum level is strongly enhanced post-burn injury and correlated with local and general tissue edema. *Burns*, 30(4): 305-311.
- Jabłońska-Trypuć, A., Matejczyk, M., & Rosochacki, S. (2016). Matrix metalloproteinases (MMPs), the main extracellular matrix (ECM) enzymes in collagen degradation, as a target for anticancer drugs. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 31(sup1): 177-183.

- Jiang, Q. (2014). Natural forms of vitamin E: metabolism, antioxidant, and anti-inflammatory activities and their role in disease prevention and therapy. *Free Radical Biology and Medicine*, 72: 76-90.
- Jorissen, R. N., Walker, F., Pouliot, N., Garrett, T. P., Ward, C. W., & Burgess, A. W. (2003). Epidermal growth factor receptor: mechanisms of activation and signaling. *Experimental Cell Research*, 284(1): 31-53.
- Jurjus, A., Atiyeh, B. S., Abdallah, I. M., Jurjus, R. A., Hayek, S. N., Jaoude, M. A., Gerge, A. & Tohme, R. A. (2007). Pharmacological modulation of wound healing in experimental burns. *Burns*, 33(7): 892-907.
- Kamat, J. P., & Devasagayam, T. P. A. (1995). Tocotrienols from palm oil as potent inhibitors of lipid peroxidation and protein oxidation in rat brain mitochondria. *Neuroscience Letters*, 195(3): 179-182.
- Kamat, J. P., Sarma, H. D., Devasagayam, T. P. A., Nesaretnam, K., & Basiron, Y. (1997). Tocotrienols from palm oil as effective inhibitors of protein oxidation and lipid peroxidation in rat liver microsomes. *Molecular and Cellular Biochemistry*, 170(1): 131-138.
- Khan, M. R., Siddiqui, S., Parveen, K., Javed, S., Diwakar, S., & Siddiqui, W. A. (2010). Nephroprotective action of tocotrienol-rich fraction (TRF) from palm oil against potassium dichromate ($K_2Cr_2O_7$)-induced acute renal injury in rats. *Chemico-Biological Interactions*, 186(2): 228-238.
- Khorasani, G., Hosseinimehr, S. J., Azadbakht, M., Zamani, A., & Mahdavi, M. R. (2009). Aloe versus silver sulfadiazine creams for second-degree burns: a randomized controlled study. *Surgery Today*, 39(7): 587-591.
- Kondo, S., & Kuroyanagi, Y. (2012). Development of a wound dressing composed of hyaluronic acid and collagen sponge with epidermal growth factor. *Journal of Biomaterials Science, Polymer Edition*, 23(5): 629-643.
- Korhonen, R., Lahti, A., Kankaanranta, H., & Moilanen, E. (2005). Nitric oxide production and signaling in inflammation. *Current Drug Targets -Inflammation & Allergy*, 4(4): 471-479.
- Kubo, H., Hayashi, T., Ago, K., Ago, M., Kanekura, T., & Ogata, M. (2014). Temporal expression of wound healing-related genes in skin burn injury. *Legal Medicine*, 16(1): 8-13.
- Kurahashi, T., & Fujii, J. (2015). Roles of antioxidative enzymes in wound healing. *Journal of Developmental Biology*, 3(2): 57-70.
- Lambert, E., Dassé, E., Haye, B., & Petitfrère, E. (2004). TIMPs as multifacial proteins. *Critical Reviews in Oncology/Hematology*, 49(3): 187-198.

- Landén, N. X., Li, D., & Ståhle, M. (2016). Transition from inflammation to proliferation: a critical step during wound healing. *Cellular and Molecular Life Sciences*, 73(20): 3861-3885.
- Latha, B., & Babu, M. (2001). The involvement of free radicals in burn injury: a review. *Burns*, 27(4): 309-317.
- Lawrence, T. (2009). The nuclear factor NF- κ B pathway in inflammation. *Cold Spring Harbor Perspectives in Biology*, 1(6): 1-10
- Lee, A. R. C., Leem, H., Lee, J., & Park, K. C. (2005). Reversal of silver sulfadiazine-impaired wound healing by epidermal growth factor. *Biomaterials*, 26(22): 4670-4676.
- Lee, J. H., Bae, I. H., Choi, J. K., & Park, J. W. (2013). Evaluation of a highly skin permeable low-molecular-weight protamine conjugated epidermal growth factor for novel burn wound healing therapy. *Journal of Pharmaceutical Sciences*, 102(11): 4109-4120.
- Lee, K., Silva, E. A., & Mooney, D. J. (2011). Growth factor delivery-based tissue engineering: general approaches and a review of recent developments. *Journal of the Royal Society Interface*, 8(55): 153-170.
- Lee, S. P., Mar, G. Y., & Ng, L. T. (2009). Effects of tocotrienol-rich fraction on exercise endurance capacity and oxidative stress in forced swimming rats. *European Journal of Applied Physiology*, 107(5): 587-595.
- Leon-Villapalos, J., Jeschke, M. G., & Herndon, D. N. (2008). Topical management of facial burns. *Burns*, 34(7): 903-911.
- Li, J., Chen, J., & Kirsner, R. (2007). Pathophysiology of acute wound healing. *Clinics in Dermatology*, 25(1): 9-18.
- Li, Z., Van Bergen, T., Van de Veire, S., Van de Vel, I., Moreau, H., Dewerchin, M., Maudgal, P. C., Zeyen, T., Spileers, W., Moons, L., & Stalmans, I. (2009). Inhibition of vascular endothelial growth factor reduces scar formation after glaucoma filtration surgery. *Investigative Ophthalmology & Visual Science*, 50(11): 5217-5225.
- Lin, Y. H., Huang, C. C., & Wang, S. H. (2011). Quantitative assessments of burn degree by high-frequency ultrasonic backscattering and statistical model. *Physics in Medicine and Biology*, 56(3): 757.
- Lin, Z. Q., Kondo, T., Ishida, Y., Takayasu, T., & Mukaida, N. (2003). Essential involvement of IL-6 in the skin wound-healing process as evidenced by delayed wound healing in IL-6-deficient mice. *Journal of Leukocyte Biology*, 73(6): 713-721.
- Lloyd, E. C., Rodgers, B. C., Michener, M., & Williams, M. S. (2012). Outpatient burns: prevention and care. *American Family Physician*,

85(1): 25-32.

- Lugrin, J., Rosenblatt-Velin, N., Parapanov, R., & Liaudet, L. (2014). The role of oxidative stress during inflammatory processes. *Biological Chemistry*, 395(2): 203-230.
- Luo, J. D. & Chen, A. F. (2005). Nitric oxide: a newly discovered function on wound healing. *Acta Pharmacologica Sinica*, 26(3): 259-264.
- Luster, A. D. (1998). Chemokines-chemotactic cytokines that mediate inflammation. *New England Journal of Medicine*, 338(7): 436-445.
- Machado, M. J., Watson, M. G., Devlin, A. H., Chaplain, M. A., McDougall, S. R., & Mitchell, C. A. (2011). Dynamics of angiogenesis during wound healing: a coupled in vivo and in silico study. *Microcirculation*, 18(3): 183-197.
- Madlener, M., Parks, W. C., & Werner, S. (1998). Matrix metalloproteinases (MMPs) and their physiological inhibitors (TIMPs) are differentially expressed during excisional skin wound repair. *Experimental Cell Research*, 242(1): 201-210.
- Mahalingam, D., Radhakrishnan, A. K., Amom, Z., Ibrahim, N., & Nesaretnam, K. (2011). Effects of supplementation with tocotrienol-rich fraction on immune response to tetanus toxoid immunization in normal healthy volunteers. *European Journal of Clinical Nutrition*, 65(1): 63-69.
- Manicone, A. M., & McGuire, J. K. (2008). Matrix metalloproteinases as modulators of inflammation. *In Seminars in Cell & Developmental Biology*, 19(1): 34-41.
- Martin, P., & Leibovich, S. J. (2005). Inflammatory cells during wound repair: the good, the bad and the ugly. *Trends in Cell Biology*, 15(11): 599-607.
- Mascia, F., Mariani, V., Girolomoni, G., & Pastore, S. (2003). Blockade of the EGF receptor induces a deranged chemokine expression in keratinocytes leading to enhanced skin inflammation. *The American Journal of Pathology*, 163(1): 303-312.
- McAnulty, R. J. (2007). Fibroblasts and myofibroblasts: their source, function and role in disease. *The International Journal of Biochemistry & Cell Biology*, 39(4): 666-671.
- McGrath, J. A., & Uitto, J. (2010). Anatomy and organization of human skin. In T. Burns, S. Breathnach, N. Cox & C. Griffiths (Ed.), *Rook's Textbook of Dermatology, Eighth Edition* (pp. 1-53). Oxford: Wiley-Blackwell.
- McRobert, J. & Stiles, K. (2014). Ten top tips...The managements of burn wounds. *Wounds International*, 5(1): 9-14.

- Menon, G. K. (2015). Skin Basics; Structure and Function. In A. Pappas (Ed.), *Lipids and Skin Health* (pp. 9-23). New York: Springer-Verlag.
- Midwood, K. S., Williams, L. V., & Schwarzbauer, J. E. (2004). Tissue repair and the dynamics of the extracellular matrix. *The International Journal of Biochemistry & Cell Biology*, 36(6): 1031-1037.
- Miller, A. C., Rashid, R. M., Falzon, L., Elamin, E. M., & Zehtabchi, S. (2012). Silver sulfadiazine for the treatment of partial-thickness burns and venous stasis ulcers. *Journal of the American Academy of Dermatology*, 66(5): e159-e165.
- Minhajuddin, M., Beg, Z. H., & Iqbal, J. (2005). Hypolipidemic and antioxidant properties of tocotrienol rich fraction isolated from rice bran oil in experimentally induced hyperlipidemic rats. *Food and Chemical Toxicology*, 43(5): 747-753.
- Mock, C., Peck, M., Krug, E., & Haberal, M. (2009). Confronting the global burden of burns: A WHO plan and a challenge. *Burns*, 35(5): 615-617.
- Mohajeri, G., Masoudpour, H., Heidarpour, M., Khademi, E. F., Ghafghazi, S., Adibi, S., & Akbari, M. (2010). The effect of dressing with fresh kiwifruit on burn wound healing. *Surgery*, 148(5): 963-968.
- Moore, K. W., de Waal Malefyt, R., Coffman, R. L., & O'Garra, A. (2001). Interleukin-10 and the interleukin-10 receptor. *Annual Review of Immunology*, 19(1): 683-765.
- Murphy, P. S., & Evans, G. R. (2012). Advances in wound healing: a review of current wound healing products. *Plastic Surgery International*, 2012: 1-8.
- Mustapha, N. M., Somchit, M. N. H., & Hasan, A. (2012). Honey hydrogel dressing to treat burn wound in rats-a preliminary report. *Pertanika J. Trop. Agric. Sci*, 35(1): 67-74.
- Mutalib, M. S. A., Khaza'ai, H., & Wahle, K. W. (2003). Palm-tocotrienol rich fraction (TRF) is a more effective inhibitor of LDL oxidation and endothelial cell lipid peroxidation than α -tocopherol in vitro. *Food Research International*, 36(5): 405-413.
- Nakamura, Y., Sotozono, C., & Kinoshita, S. (2001). The epidermal growth factor receptor (EGFR): role in corneal wound healing and homeostasis. *Experimental Eye Research*, 72(5): 511-517.
- Nesaretnam, K., Ambra, R., Selvaduray, K. R., Radhakrishnan, A., Canali, R., & Virgili, F. (2004). Tocotrienol-rich fraction from palm oil and gene expression in human breast cancer cells. *Annals of the New York Academy of Sciences*, 1031(1): 143-157.

- Nessler, M., Puchala, J., Wood, F. M., Wallace, H. J., Fear, M. W., Nessler, K., & Drukala, J. (2013). Changes in the plasma cytokine and growth factor profile are associated with impaired healing in pediatric patients treated with INTEGRA® for reconstructive procedures. *Burns*, 39(4): 667-673.
- Nicol, N. H. (2005). Anatomy and physiology of the skin. *Dermatology Nursing*, 17(1): 62-63.
- Noble, P. W. (2002). Hyaluronan and its catabolic products in tissue injury and repair. *Matrix Biology*, 21(1): 25-29.
- Oberyszyn, T. M. (2007). Inflammation and wound healing. *Front Biosci*, 12(8): 2993-2999.
- Okada, A., Tomasetto, C., Lutz, Y., Bellocq, J. P., Rio, M. C., & Basset, P. (1997). Expression of matrix metalloproteinases during rat skin wound healing: evidence that membrane type-1 matrix metalloproteinase is a stromal activator of pro-gelatinase A. *The Journal of Cell Biology*, 137(1): 67-77.
- Pacher, P., Beckman, J. S., & Liaudet, L. (2007). Nitric oxide and peroxynitrite in health and disease. *Physiological Reviews*, 87(1): 315-424.
- Parihar, A., Parihar, M. S., Milner, S., & Bhat, S. (2008). Oxidative stress and anti-oxidative mobilization in burn injury. *Burns*, 34(1): 6-17.
- Park, J. E., & Barbul, A. (2004). Understanding the role of immune regulation in wound healing. *The American Journal of Surgery*, 187(5): S11-S16.
- Parks, W. C. (1999). Matrix metalloproteinases in repair. *Wound Repair and Regeneration*, 7(6): 423-432.
- Paul, W., & Sharma, C. P. (2004). Chitosan and alginate wound dressings: a short review. *Trends Biomater Artif Organs*, 18(1): 18-23.
- Paus, R., & Cotsarelis, G. (1999). The biology of hair follicles. *New England Journal of Medicine*, 341(7): 491-497.
- Peck, M. D. (2011). Epidemiology of burns throughout the world. Part I: Distribution and risk factors. *Burns*, 37(7): 1087-1100.
- Peck, M. D. (2012). Epidemiology of burns throughout the world. Part II: Intentional burns in adults. *Burns*, 38(5): 630-637.
- Penn, J. W., Grobbelaar, A. O., & Rolfe, K. J. (2012). The role of the TGF- β family in wound healing, burns and scarring: a review. *International Journal of Burns and Trauma*, 2(1): 18-28.

- Pereira, D. D. S. T., Lima-Ribeiro, M. H. M., de Pontes-Filho, N. T., Carneiro-Leão, A. M. D. A., & Correia, M. T. D. S. (2012). Development of animal model for studying deep second-degree thermal burns. *BioMed Research International*, 2012: 460841.
- Proksch, E., Brandner, J. M., & Jensen, J. M. (2008). The skin: an indispensable barrier. *Experimental Dermatology*, 17(12): 1063-1072.
- Quan, T. E., Cowper, S., Wu, S. P., Bockenstedt, L. K., & Bucala, R. (2004). Circulating fibrocytes: collagen-secreting cells of the peripheral blood. *The International Journal of Biochemistry & Cell Biology*, 36(4): 598-606.
- Qureshi, A. A., Sami, S. A., Salser, W. A., & Khan, F. A. (2001). Synergistic effect of tocotrienol-rich fraction (TRF 25) of rice bran and lovastatin on lipid parameters in hypercholesterolemic humans. *The Journal of Nutritional Biochemistry*, 12(6): 318-329.
- Qureshi, A. A., Sami, S. A., Salser, W. A., & Khan, F. A. (2002). Dose-dependent suppression of serum cholesterol by tocotrienol-rich fraction (TRF 25) of rice bran in hypercholesterolemic humans. *Atherosclerosis*, 161(1): 199-207.
- Rawlingson, A. (2003). Nitric oxide, inflammation and acute burn injury. *Burns*, 29(7): 631-640.
- Repertinger, S. K., Campagnaro, E., Fuhrman, J., El-Abaseri, T., Yuspa, S. H., & Hansen, L. A. (2004). EGFR enhances early healing after cutaneous incisional wounding. *Journal of Investigative Dermatology*, 123(5): 982-989.
- Ridnour, L. A., Isenberg, J. S., Espey, M. G., Thomas, D. D., Roberts, D. D., & Wink, D. A. (2005). Nitric oxide regulates angiogenesis through a functional switch involving thrombospondin-1. *Proceedings of the National Academy of Sciences of the United States of America*, 102(37): 13147-13152.
- Roy, S., Khanna, S., Nallu, K., Hunt, T. K., & Sen, C. K. (2006). Dermal wound healing is subject to redox control. *Molecular Therapy*, 13(1): 211-220.
- Ruszczak, Z. (2003). Effect of collagen matrices on dermal wound healing. *Advanced Drug Delivery Reviews*, 55(12): 1595-1611.
- Santoro, M. M., & Gaudino, G. (2005). Cellular and molecular facets of keratinocyte re-epithelialization during wound healing. *Experimental Cell Research*, 304(1): 274-286.
- Sanwal, R., & Chaudhary, A. K. (2011). Wound healing and antimicrobial potential of *Carissa spinarum* Linn. in albino mice. *Journal of Ethnopharmacology*, 135(3): 792-796

- Sarkhail, P., Esmaily, H., Baghaei, A., Shafiee, A., Abdollahi, M., Khademi, Y., Madander, M. & Sarkheil, P. (2011). Burn healing potential of *Nigella sativa* seed oil in rats. *International Journal of Pharmaceutical Sciences and Research*, 2(1): 34-40.
- Sato, Y., Ohshima, T., & Kondo, T. (1999). Regulatory role of endogenous interleukin-10 in cutaneous inflammatory response of murine wound healing. *Biochemical and Biophysical Research Communications*, 265(1): 194-199.
- Schäfer, M., & Werner, S. (2008). Oxidative stress in normal and impaired wound repair. *Pharmacological Research*, 58(2): 165-171.
- Schäffer, M. R., Tantry, U., van Wesep, R. A., & Barbul, A. (1997). Nitric oxide metabolism in wounds. *Journal of Surgical Research*, 71(1): 25-31.
- Schultz, G. S., & Wysocki, A. (2009). Interactions between extracellular matrix and growth factors in wound healing. *Wound Repair and Regeneration*, 17(2): 153-162.
- Schwacha, M. G., Nickel, E., & Daniel, T. (2008). Burn injury–induced alterations in wound inflammation and healing are associated with suppressed hypoxia inducible factor-1 α expression. *Molecular Medicine*, 14 (9-10): 628-633.
- Sen, C. K., & Roy, S. (2008). Redox signals in wound healing. *Biochimica et Biophysica Acta (BBA)-General Subjects*, 1780(11): 1348-1361.
- Shakespeare, P. (2001). Burn wound healing and skin substitutes. *Burns*, 27(5): 517-522.
- Siddiqui, S., Khan, M. R., & Siddiqui, W. A. (2010). Comparative hypoglycemic and nephroprotective effects of tocotrienol rich fraction (TRF) from palm oil and rice bran oil against hyperglycemia induced nephropathy in type 1 diabetic rats. *Chemico-Biological Interactions*, 188(3): 651-658.
- Silva, M. A., Trevisan, G., Klafke, J. Z., Rossato, M. F., Walker, C. I. B., Oliveira, S. M., Silva, C. R., Boligon, A.A., Flores, F. C., Silva, C. D. B., Athayde, M. L. & Ferreira, J. (2013). Antinociceptive and anti-inflammatory effects of *Aloe saponaria* Haw on thermal injury in rats. *Journal of Ethnopharmacology*, 146(1): 393-401.
- Singer, A. J., & Clark, R. A. (1999). Cutaneous wound healing. *New England Journal of Medicine*, 341(10): 738-746.
- Singer, A. J., & Dagum, A. B. (2008). Current management of acute cutaneous wounds. *New England Journal of Medicine*, 359(10): 1037-1046.

- Singer, A. J., Berruti, L., Thode, H. C., & McClain, S. A. (2000). Standardized burn model using a multiparametric histologic analysis of burn depth. *Academic Emergency Medicine*, 7(1): 1-6.
- Siqueira, M. F., Li, J., Chehab, L., Desta, T., Chino, T., Krothpali, N., Behl, Y., Alikhani, M., Yang, J., Barrsch, C. & Graves, D. T. (2010). Impaired wound healing in mouse models of diabetes is mediated by TNF- α dysregulation and associated with enhanced activation of forkhead box O1 (FOXO1). *Diabetologia*, 53(2): 378-388.
- Sobouti, B., Hooman, N., & Movahed, M. (2013). The effect of vitamin E or vitamin A on the prevention of renal scarring in children with acute pyelonephritis. *Pediatric Nephrology*, 28(2): 277-283.
- Soejima, K., Traber, L. D., Schmalstieg, F. C., Hawkins, H., Jodoin, J. M., Szabo, C., Szabo, E., Varig, L., Salzman, A. & Traber, D. L. (2001). Role of nitric oxide in vascular permeability after combined burns and smoke inhalation injury. *American Journal of Respiratory and Critical Care Medicine*, 163(3): 745-752.
- Song, C., & Chua, A. (2005). Epidemiology of burn injuries in Singapore from 1997 to 2003. *Burns*, 31(1): S18-S26.
- Spies, M., Dasu, M. R., Svrakic, N., Nesic, O., Barrow, R. E., Perez-Polo, J. R., & Herndon, D. N. (2002). Gene expression analysis in burn wounds of rats. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 283 (4): R918-R930.
- Srivastava, J. K., & Gupta, S. (2006). Tocotrienol-rich fraction of palm oil induces cell cycle arrest and apoptosis selectively in human prostate cancer cells. *Biochemical and Biophysical Research Communications*, 346(2): 447-453.
- Stamenkovic, I. (2003). Extracellular matrix re-modelling: the role of matrix metalloproteinases. *The Journal of Pathology*, 200(4): 448-464.
- Stashak, T. S., Farstvedt, E., & Othic, A. (2004). Update on wound dressings: indications and best use. *Clinical Techniques in Equine Practice*, 3(2): 148-163.
- Staton, C. A., Stribbling, S. M., Tazzyman, S., Hughes, R., Brown, N. J., & Lewis, C. E. (2004). Current methods for assaying angiogenesis in vitro and in vivo. *International Journal of Experimental Pathology*, 85(5): 233-248.
- Stetler-Stevenson, W. G., Kruttsch, H. C., & Liotta, L. A. (1989). Tissue inhibitor of metalloproteinase (TIMP-2). A new member of the metalloproteinase inhibitor family. *Journal of Biological Chemistry*, 264(29): 17374-17378.

- Strongin, A. Y., Marmer, B. L., Grant, G. A., & Goldberg, G. I. (1993). Plasma membrane-dependent activation of the 72-kDa type IV collagenase is prevented by complex formation with TIMP-2. *Journal of Biological Chemistry*, 268(19): 14033-14039.
- Sullivan, T. P., Eaglstein, W. H., Davis, S. C., & Mertz, P. (2001). The pig as a model for human wound healing. *Wound Repair and Regeneration*, 9(2): 66-76
- Süzük, S., Erdoğan, D., Kaplanoğlu, G. T., Ilgaz, C., & Değim, Z. G. (2011). Kitosan ve Taurin-Kitosan Uygulamasında Yara İyileşmesinde EGF Reseptör ve Fibronektin'in İmmünohistokimyasal Olarak Gösterilmesi. *Gazi Medical Journal*, 22(1): 6-13.
- Suzuki, K., Saito, J., Yanai, R., Yamada, N., Chikama, T. I., Seki, K., & Nishida, T. (2003). Cell-matrix and cell-cell interactions during corneal epithelial wound healing. *Progress in Retinal and Eye Research*, 22(2): 113-133.
- Thannickal, V. J., & Fanburg, B. L. (2000). Reactive oxygen species in cell signaling. *American Journal of Physiology-Lung Cellular and Molecular Physiology*, 279(6): L1005-L1028.
- Tonnesen, M. G., Feng, X., & Clark, R. A. (2000). Angiogenesis in wound healing. *Journal of Investigative Dermatology Symposium Proceedings*, 5(1): 40-46.
- Toussaint, J., & Singer, A. J. (2014). The evaluation and management of thermal injuries: 2014 update. *Clinical and Experimental Emergency Medicine*, 1(1): 8-18.
- Traber, M. G. (2007). Vitamin E regulatory mechanisms. *Annu. Rev. Nutr.* 27: 347-362.
- Traber, M. G., & Atkinson, J. (2007). Vitamin E, antioxidant and nothing more. *Free Radical Biology and Medicine*, 43(1): 4-15.
- Tsirogianni, A. K., Moutsopoulos, N. M., & Moutsopoulos, H. M. (2006). Wound healing: immunological aspects. *Injury*, 37(1): S5-S12.
- Tuan, T. L., & Nichter, L. S. (1998). The molecular basis of keloid and hypertrophic scar formation. *Molecular Medicine Today*, 4(1): 19-24.
- Uría, J. A., Ferrando, A. A., Velasco, G., Freije, J. M., & López-Otín, C. (1994). Structure and expression in breast tumors of human TIMP-3, a new member of the metalloproteinase inhibitor family. *Cancer Research*, 54(8): 2091-2094.
- Varani, J., Dame, M. K., Rittie, L., Fligiel, S. E., Kang, S., Fisher, G. J., & Voorhees, J. J. (2006). Decreased collagen production in

chronologically aged skin: roles of age-dependent alteration in fibroblast function and defective mechanical stimulation. *The American Journal of Pathology*, 168(6): 1861-1868.

Velnar, T., Bailey, T., & Smrkolj, V. (2009). The wound healing process: an overview of the cellular and molecular mechanisms. *Journal of International Medical Research*, 37(5): 1528-1542.

Venter, N. G., Monte-Alto-Costa, A., & Marques, R. G. (2015). A new model for the standardization of experimental burn wounds. *Burns*, 41(3): 542-547.

Verma, R. P., & Hansch, C. (2007). Matrix metalloproteinases (MMPs): chemical-biological functions and (Q) SARs. *Bioorganic & Medicinal Chemistry*, 15(6): 2223-2268.

Wang, R., Ghahary, A., Shen, Q., Scott, P. G., Roy, K., & Tredget, E. E. (2000). Hypertrophic scar tissues and fibroblasts produce more transforming growth factor- β 1 mRNA and protein than normal skin and cells. *Wound Repair and Regeneration*, 8(2): 128-137.

Wang, X. Q., Kravchuk, O., & Kimble, R. M. (2010). A retrospective review of burn dressings on a porcine burn model. *Burns*, 36(5): 680-687.

Ward, R. V., Hembry, R. M., Reynolds, J. J., & Murphy, G. (1991). The purification of tissue inhibitor of metalloproteinases-2 from its 72 kDa progelatinase complex. *Biochemical Journal*, 278(1): 179-187.

Wells, A. (1999). EGF receptor. *The International Journal of Biochemistry & Cell Biology*, 31(6): 637-643.

Werner, S., & Grose, R. (2003). Regulation of wound healing by growth factors and cytokines. *Physiological Reviews*, 83(3): 835-870.

Wilgus, T. A., Roy, S., & McDaniel, J. C. (2013). Neutrophils and wound repair: positive actions and negative reactions. *Advances in Wound Care*, 2(7): 379-388.

Wilhelm, S. M., Collier, I. E., Marmer, B. L., Eisen, A. Z., Grant, G. A., & Goldberg, G. I. (1989). SV40-transformed human lung fibroblasts secrete a 92-kDa type IV collagenase which is identical to that secreted by normal human macrophages. *Journal of Biological Chemistry*, 264(29): 17213-17221.

Witte, M. B., & Barbul, A. (2002). Role of nitric oxide in wound repair. *The American Journal of Surgery*, 183(4): 406-412.

Witte, M. B., Thornton, F. J., Tantry, U., & Barbul, A. (2002). L-Arginine supplementation enhances diabetic wound healing: involvement of the nitric oxide synthase and arginase pathways. *Metabolism*, 51(10):

1269-1273.

- Wu, S. J., Liu, P. L., & Ng, L. T. (2008). Tocotrienol-rich fraction of palm oil exhibits anti-inflammatory property by suppressing the expression of inflammatory mediators in human monocytic cells. *Molecular Nutrition & Food Research*, 52(8): 921-929.
- Wu, X. B., Luo, X. Q., Gu, S. Y., & Xu, J. H. (2012). The effects of *Polygonum cuspidatum* extract on wound healing in rats. *Journal of Ethnopharmacology*, 141(3): 934-937.
- Xie, H., Pallero, M. A., Gupta, K., Chang, P., Ware, M. F., Witke, W., Kwiatkowski, D. J., Lauffenburger, D. A., Murphy-Ullrich, J. E., & Wells, A. (1998). EGF receptor regulation of cell motility: EGF induces disassembly of focal adhesions independently of the motility-associated PLC gamma signaling pathway. *Journal of Cell Science*, 111(5): 615-624.
- Yamasaki, K., Edington, H. D., McClosky, C., Tzeng, E., Lizonova, A., Kovacs, I., Steed, D. L. & Billiar, T. R. (1998). Reversal of impaired wound repair in iNOS-deficient mice by topical adenoviral-mediated iNOS gene transfer. *Journal of Clinical Investigation*, 101(5): 967-971.
- Yan, H., Peng, X., Huang, Y., Zhao, M., Li, F., & Wang, P. (2007). Effects of early enteral arginine supplementation on resuscitation of severe burn patients. *Burns*, 33(2): 179-184.
- Yang, H., Li, X., Ma, J., Lv, X., Zhao, S., Lang, W., & Zhang, Y. (2013). Blockade of the intermediate-conductance Ca^{2+} -activated K^{+} channel inhibits the angiogenesis induced by epidermal growth factor in the treatment of corneal alkali burn. *Experimental Eye Research*, 110: 76-87.
- Yarden, Y., & Shilo, B. Z. (2007). SnapShot: EGFR signaling pathway. *Cell*, 131(5), 1018-e1.
- Zaidi, Z., & Lanigan, S. W. (2010). *Dermatology in clinical practice*. London: Springer-Verlag.
- Zaini, A. A., Khaza'ai, H., Ali, R. M., Abdul Mutalib, M. S., & Baharuddin, A. A. (2016). Topical treatment of tocotrienol-rich fraction (TRF) on deep partial-thickness burn wounds in rats. *Journal of Dermatology and Clinic Research*, 4(1): 1063.
- Zampieri, N., & Castellani, R. (2014). Topical application of lipophilic anhydrous gel rich in vitamin E in the management of scars in patients surgically treated for neck cysts. *Journal of the American Academy of Dermatology*. 70 (5): AB194.
- Zampieri, N., Zuin, V., Burro, R., Ottolenghi, A., & Camoglio, F. S. (2010). A

prospective study in children: pre-and post-surgery use of vitamin E in surgical incisions. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 63(9): 1474-1478.

Zawacki, B. E. (1974). Reversal of capillary stasis and prevention of necrosis in burns. *Annals of Surgery*, 180(1): 98-102.

Zheng, J., Wen, Y., Song, Y., Wang, K., Chen, D. B., & Magness, R. R. (2008). Activation of Multiple Signaling Pathways Is Critical for Fibroblast Growth Factor 2-and Vascular Endothelial Growth Factor-Stimulated Ovine Fetoplacental Endothelial Cell Proliferation 1. *Biology of Reproduction*, 78(1): 143-150.

Zhu, K. Q., Engrav, L. H., Armendariz, R., Muangman, P., Klein, M. B., Carroughe, G. J., Deubner, H., & Gibran, N. S. (2005). Changes in VEGF and nitric oxide after deep dermal injury in the female, red Duroc pig—further similarities between female, Duroc scar and human hypertrophic scar. *Burns*, 31(1): 5-10.

LIST OF PUBLICATIONS

Hui-Fang Guo, Razana, M.L., Roslida, A.H., Asma, A.Z., & Huzwah, B. K. (2017). A new model for studying deep partial-thickness burns in rats. *Internal Journal of Burns and Trauma*, 7(6): 107-114.

Hui-Fang Guo, Razana, M.L., Roslida, A.H., & Huzwah, B. K.(2017). Temporal changes in the cell population and wound healing-related gene expression in deep partial-thickness burn wound model. *Burns*. (Submitted)

