



**UNIVERSITI PUTRA MALAYSIA**

***IN vitro* MODULATORY EFFECTS OF MICROGLIA ON  
NEUROEPITHELIAL CELL NEURODIFFERENTIATION**

**TONG CHIH KONG**

**FPSK(p) 2018 10**



***In Vitro* MODULATORY EFFECTS OF MICROGLIA ON  
NEUROEPITHELIAL CELL NEURODIFFERENTIATION**

By

**TONG CHIH KONG**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra  
Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of  
Philosophy**

**December 2017**

## COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



**Specially dedicated to my beloved parents, family, and friends.**

**For their unconditional love, understanding, patience, support and constant faith.**



© COPYRIGHT UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

***In Vitro* MODULATORY EFFECTS OF MICROGLIA ON  
NEUROEPITHELIAL CELL NEURODIFFERENTIATION**

By

**TONG CHIH KONG**

**December 2017**

**Chairman : Sharmili Vidyadaran, PhD**  
**Faculty : Medicine and Health Sciences**

Neural precursor cells (NPC) give rise to neurons and glial cells during embryonic brain development. Dysregulation of NPC growth and function can cause brain deformities and result in psychiatric and behavioural deficits later in life. Recent discoveries reveal that microglia are involved in modulating neurogenesis during embryonic brain development. This study explores the role of microglia in regulating NPC proliferation and neural differentiation using an in vitro cell culture approach. For this, BV2 microglia-conditioned media from cultures treated acutely and chronically with 1µg/ml lipopolysaccharide were collected and incubated with the NE-4C neuroepithelial cell line. NE-4C cultures were either non-induced or induced into neural differentiation using a retinoid analogue, EC23. The effects of microglia on NPC were elucidated by examining proliferation, cell death, pluripotency and cell adhesion markers, as well as early and terminal neural differentiation. Firstly, it was shown that BV2 soluble factors, regardless of activation status, significantly reduced the proliferation of NE-4C cells at all time points examined (from 24 to 96 hours). For instance, at 24 hours, unstimulated BV2-conditioned media reduced non-induced NE-4C proliferation by  $83.00 \pm 4.43\%$  ( $p < 0.001$ ). DAPI/PI staining and Annexin-V/PI assays showed no increase in apoptotic cells in the presence of BV2 microglia-conditioned media, confirming that the reduced proliferation of NE-4C is unrelated to apoptosis. By immunophenotyping and RT-qPCR analysis, it was shown that BV2 microglia-conditioned media does not alter the expression of pluripotency markers (SSEA-1 and CD133) and neural commitment and maintenance factors (HES1, HES3, HES5, MASH1, SOX1, NGN2, and PAX6) in NE-4C neuroepithelial cells at the early phase of neural differentiation. Immunocytochemical studies showed that BV2 microglia-conditioned media inhibited the production of RC2+ radial glial cell and

TUJ1+ neurons during the early phase of neural differentiation. This was accompanied by the reduced formation of neural aggregates. Next, western blot studies showed that BV2 microglia-conditioned media up-regulate the expression of E-cadherin, beta-catenin, and CDC42 but down-regulate the expression of N-cadherin in NE-4C cells. The alteration of the expression of these molecules in NPC may arrest the cells in an undifferentiated state and therefore fail to commit into neural differentiation. Subsequently, the present study elucidated whether microglia-derived soluble factors can impact the terminal production of neuronal cells. The results demonstrate that unstimulated BV2 microglia-conditioned media reduced TUJ1<sup>+</sup> neuron production in EC23-induced NE-4C cultures by 58.6% ( $p < 0.001$ ). However, there were no significant changes in the percentage of GFAP<sup>+</sup> astrocytes. BV2-conditioned media also did not affect the expression of the mature neuron marker MAP2 and synapse marker SYT1 in NE-4C derived neurons. This study then evaluated the effects of BV2-conditioned media on neurite complexity. BV2-conditioned media significantly reduced the number of neurites from  $9169 \pm 1735$  in untreated cultures to  $1362 \pm 117.6$  ( $p < 0.001$ ). Total neurite length was also significantly decreased from  $5246 \pm 924.7$ px in untreated cultures to  $1026 \pm 203.6$ px ( $p < 0.001$ ). Neurite attachment points were also reduced from  $3115 \pm 451.5$  in untreated cultures to  $1310 \pm 356.8$  ( $p < 0.001$ ). Also, the number of neurite end points were significantly reduced from  $7776 \pm 1845$  in untreated cultures to  $4060 \pm 952.1$  ( $p < 0.001$ ). There was no neural differentiation observed in non-induced NE-4C cultures suggesting that BV2 soluble factors do not contribute to the initiation of neurogenesis. Also, the activation status of microglia had no effect on NE-4C neural differentiation in any of the parameters examined, indicating that ubiquitous microglial soluble factors are responsible for the effects described here. To conclude, the primary finding of this study is that microglia exert inhibitory effects on the neural differentiation of NPC.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**KESAN MODULASI *In Vitro* MIKROGLIA PADA NEURODIFERENSIASI  
SEL NEUROEPITELIAL**

Oleh

**TONG CHIH KONG**

Disember 2017

**Pengerusi : Sharmili Vidyadaran, PhD**  
**Fakulti : Perubatan dan Sain Kesihatan**

Sel induk neural (NPC) berperanan untuk menghasilkan sel-sel neuron dan glial semasa perkembangan otak embrionik. Penyisihan pertumbuhan dan fungsi NPC boleh menyebabkan kecacatan otak dan mengakibatkan defisit psikiatri dan tingkah laku pada tahap kehidupan selanjutnya. Penemuan baru-baru ini mendedahkan bahawa mikroglia terlibat dalam mengubah neurogenesis NPC semasa perkembangan otak embrionik. Kajian ini mengeksplorasi peranan mikroglia dalam mengawal selia proliferasi NPC dan diferensiasi neural dengan menggunakan pendekatan kultur sel *in vitro*. Untuk ini, media BV2 mikroglia dari kultur yang dirawat dengan 1µg/ml lipopolysaccharide secara akut dan kronis dikumpulkan dan digunakan untuk diinkubasi bersama sel neuroepitelial NE-4C. Kultur NE-4C sama ada tidak diinduksi atau diinduksi untuk diferensiasi neural dengan menggunakan analog retinoid, EC23. Kesan mikroglia pada NPC akan dijelaskan dengan memeriksa proliferasi, kematian sel, pluripotency and adhesi sel, serta diferensiasi neural awal dan terminal. Pertama, ditunjukkan bahawa factor larutan BV2, tanpa mengira status pengaktifan, mengurangi proliferasi sel neuroepitelial NE-4C secara mendadak pada semua titik masa yang dikaji (dari 24 hingga 96 jam). Contohnya, media BV2 yang tidak distimulasi mengurangkan proliferasi kultur NE-4C yang tidak diinduksi sebanyak  $83.00 \pm 4.43\%$  ( $p < 0.001$ ) pada titik masa 24 jam. Keputusan dari pewarnaan DAPI/PI dan ujian Annexin-V/PI menunjukkan bahawa tidak terdapat peningkatan sel apoptotik dalam kehadiran media BV2 mikroglia, seterusnya mengesahkan bahawa penurunan kadar proliferasi NE-4C tidak berkaitan dengan apoptosis. Dengan menggunakan analisis flow cytometry dan RT-qPCR, ditunjukkan bahawa media BV2 mikroglia tidak mengubah ekspresi penanda pluripotency (SSEA-1 dan CD133) serta faktor komitmen dan pemeliharaan neural (HES1, HES3, HES5, MASH1, SOX1, NGN2, dan PAX6) dalam sel-

sel neuroepitelial NE-4C pada peringkat awal diferensiasi neural. Kajian immunocytochemical menunjukkan bahawa media BV2 mikroglia menghalang penghasilan produksi RC2<sup>+</sup> sel radial glial dan neuron TUJ1<sup>+</sup> semasa fasa awal diferensiasi neural. Ini disertai dengan berkurangnya pembentukan agregat neural. Seterusnya, kajian western blot menunjukkan bahawa media BV2 mikroglia menaikkan ekspresi E-cadherin, beta-catenin dan CDC42 tetapi menurunkan ekspresi N-cadherin dalam sel NE-4C. Perubahan ungkapan molekul-molekul ini dalam NPC boleh menahan sel-sel dalam keadaan yang tidak berdiferensiasi dan oleh itu gagal melakukan mode diferensiasi neural. Selanjutnya, kajian ini membuktikan sama ada faktor larutan BV2 dapat mempengaruhi produksi sel-sel terminal neuron. Hasilnya menunjukkan bahawa media BV2 yang tidak distimulasi mengurangkan produksi neuron TUJ1<sup>+</sup> di dalam kultur NE-4C yang diinduksi dengan EC23 sebanyak 58.6% ( $p < 0.001$ ). Walau bagaimanapun, tiada perubahan signifikan dalam peratusan GFAP<sup>+</sup> astrocytes dan nestin<sup>+</sup> sel progenitor. Juga diketahui bahawa media BV2 tidak mempengaruhi ekspresi penanda neuron matang MAP2 dan penanda synapse SYT1 dalam neuron keturunan NE-4C. Kajian ini kemudiannya menilai kesan media BV2 pada kompleksiti neurit. Media BV2 mengurangkan bilangan neurit secara medadak dari  $9169 \pm 1735$  dalam kultur tidak dirawat kepada  $1362 \pm 117.6$  ( $p < 0.001$ ). Panjang keseluruhan neurit juga menurun secara signifikan dari  $5246 \pm 924.7\text{px}$  dalam kultur yang tidak dirawat kepada  $1026 \pm 203.6\text{px}$  ( $p < 0.001$ ). Titik pelekatan neurit juga dikurangkan dari  $3115 \pm 451.5$  dalam kultur tidak dirawat kepada  $1310 \pm 356.8$  ( $p < 0.001$ ). Selain itu, bilangan titik akhir neurit juga berkurangan secara signifikan dari  $7776 \pm 1845$  kultur tidak dirawat kepada  $4060 \pm 952.1$  ( $p < 0.001$ ). Tiada diferensiasi neural yang dilihat di NE-4C kultur yang tidak diinduksi dan ini mencadangkan bahawa faktor larutan BV2 tidak menyumbang kepada permulaan neurogenesis. Selain itu, status pengaktifan mikroglia tidak mempunyai kesan yang berbeza terhadap diferensiasi neural NE-4C dalam mana-mana parameter yang diperiksa dan ini mencadangkan bahawa faktor larutan mikroglia setiasa ada yang bertanggungjawab terhadap kesan yang dijelaskan di sini. Kesimpulannya, penemuan utama kajian ini adalah mikroglia melakukan kesan perencatan terhadap diferensiasi neural oleh NPC.



## ACKNOWLEDGEMENTS

In the name of God, the most Merciful and Gracious,

First and foremost, I would like to express my deepest gratitude to Dr Sharmili Vidyadaran for her insightful guidance, encouragement, motivation, and help in my PhD study and research. She contributed to a rewarding postgraduate experience by giving me intellectual freedom in my work, engaging me in new ideas, and demanding a high quality of work in all my endeavours. Additionally, I would like to thank my co-supervisors- Dr Norshariza Nordin and Dr Md.Shariful Hasan Sumon for their interest in my work.

I would like to thank all my friends for giving me moral support. Carmen, Sam, Joey, Daniel, Jervis, Lee Teng and Felicia - Thank you for spending your time running and hiking with me. It soothed my stress during this academic pursuit and gave me the strength to move forward.

I also like take this opportunity to express sincere appreciation to all the members of the Immunology Laboratory. My dear labmates (Shi Wei, Iris, Catherine, Jack and Yip), thank you for sharing the never ending enthusiasm, energy and great friendship. I am indebted to all of you for your constant help and willingness to share the knowledge from which I learnt so much. To the support staffs (Mr Anthonysamy, Aishah, Kak Zura, Afifi and Marsitah), for your help in the technicalities of laboratory maintenance, and the administrative paperwork.

Last, but certainly not the least, I would like to thank my parents and family. Without their unbreakable love and encouragement, never would I have come to this far. My apologies and sincere gratitude go to those whom I failed to mention but contributed to this project in any manner.

I certify that a Thesis Examination Committee has met on 7 December 2017 to conduct the final examination of Tong Chih Kong on his thesis entitled "*In Vitro* Modulatory Effects of Microglia on Neuroepithelial Cell Neurodifferentiation" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

**Rajesh a/l Ramasamy, PhD**

Associate Professor  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Chairman)

**Mohamad Aris bin Mohd Moklas, PhD**

Associate Professor  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**Mohamad Taufik Hidayat bin Baharuldin, PhD**

Associate Professor  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**Sukumal Chongthammakun, PhD**

Professor  
Mahidol University  
Thailand  
(External Examiner)



---

**NOR AINI AB. SHUKOR, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 27 February 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Sharmili Vidyadaran, PhD**

Associate Professor  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Chairman)

**Norshariza Nordin, PhD**

Senior Lecturer  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Member)

**Md. Shariful Hasan Sumon**

Senior Lecturer  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**  
Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Tong Chih Kong, GS32670

## Declaration by Members of the Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: Associate Professor Dr. Sharmili Vidyadaran

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: Dr. Norshariza Nordin

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: Dr. Md. Shariful Hasan Sumon

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xiv
<b>LIST OF FIGURES</b>	xv
<b>LIST OF APPENDICES</b>	xvii
<b>LIST OF ABBREVIATIONS</b>	xviii

### CHAPTER

<b>1</b>	<b>INTRODUCTION</b>	<b>1</b>
<b>2</b>	<b>LITERATURE REVIEW</b>	<b>3</b>
2.1	Development of the brain: a cell biology perspective	3
2.1.1	Heterogeneity of neural precursor cells	3
2.1.2	Neuroepithelial cells are the most primitive NPC	5
2.1.3	Radial glial cells	6
2.1.4	Intermediate progenitor cells	8
2.1.5	Neurogenesis to gliogenesis transition	8
2.2	Microglia biology	9
2.2.1	Microglia phenotype	9
2.2.2	Microglia functions in the CNS	11
2.2.3	Microglia ontogeny	12
2.2.3.1	Microglia are not derived from the mononuclear phagocyte system	12
2.2.3.2	Microglia derive from yolk sac myeloid progenitors	15
2.2.3.3	Microglia are not derived from the neuroectoderm	16
2.2.4	Microglia maintenance and self-sustainability	17
2.2.5	Microglia roles in early neurodevelopment	17
2.2.6	Microglia roles in adult neurodevelopment	21
2.2.6.1	Functions of microglia in synaptic pruning	21
2.2.6.2	Microglia and neurotransmission	23
2.2.6.3	Microglia and adult neurogenesis	23
2.2.7	Microglia and neurodevelopmental diseases	26

<b>3</b>	<b>MATERIALS AND METHODS</b>	<b>29</b>
3.1	Cell culture	29
3.1.1	BV2 microglia cell line culture	29
3.1.2	Acute and chronic LPS stimulation of BV2 microglia	30
3.1.3	NE-4C neuroepithelial cell line culture	33
3.1.4	Induction of neural differentiation	33
3.2	Tritium thymidine ( <sup>3</sup> H-TdR) incorporation assay to quantify cell proliferation	34
3.3	4',6-diamidino-2-phenylindole, dihydrochloride (DAPI) and propidium iodide (PI) double staining assay to analyse cell death	34
3.4	Determination of mRNA expression with quantitative reverse transcription polymerase chain reaction (RT-qPCR) analysis	35
3.4.1	RNA isolation	35
3.4.2	cDNA synthesis	36
3.4.3	Quantitative reverse transcription PCR (RT-qPCR) reaction	37
3.5	Flow cytometry	39
3.5.1	Annexin V-FITC/PI staining to analyse apoptosis	39
3.5.2	Immunophenotyping	39
3.6	Immunocytochemistry	40
3.6.1	Neuron morphology and quantification of neurite outgrowth	43
3.6.2	Total cell count	43
3.6.3	Reliability test	43
3.7	Western blot	44
3.7.1	Total protein extraction	44
3.7.2	Protein quantification	45
3.7.3	Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE)	45
3.7.4	Semi-dry protein transfer	45
3.7.5	Immunoblot detection and visualisation	46
3.7.6	Densitometric analysis	47
3.8	Statistical analysis	47
<b>4</b>	<b>RESULTS AND DISCUSSION</b>	<b>48</b>
4.1	Effect of BV2 microglia-conditioned media on NE-4C neuroepithelial cell number	48
4.1.1	BV2 microglia-conditioned media inhibit NE-4C neuroepithelial cell proliferation	49
4.1.2	BV2 microglia-conditioned media does not induce cell death in NE-4C neuroepithelial cells	52
4.2	Effect of BV2 microglia-conditioned media on early events of NE-4C neuroepithelial cell differentiation	56

4.2.1	BV2 microglia-conditioned media does not affect the expression of pluripotency markers SSEA-1 and CD133 in NE-4C neuroepithelial cells	56
4.2.2	Effect of BV2 microglia-conditioned media on the gene expression of neural commitment and maintenance factors in NE-4C neuroepithelial cells	60
4.2.2.1	BV2 microglia-conditioned media does not alter the gene expression of SOX1 and PAX6 in NE-4C neuroepithelial cells	60
4.2.2.2	BV2 microglia-conditioned media does not alter the gene expression of HES1 and HES5 in NE-4C neuroepithelial cells	62
4.2.2.3	BV2 microglia-conditioned media does not alter the gene expression of MASH1 and NGN2 in NE-4C neuroepithelial cells	63
4.2.3	Effect of BV2 microglia-conditioned media on the early neural differentiation of NE-4C neuroepithelial cells	65
4.2.3.1	BV2 microglia-conditioned media prevent aggregate formation and inhibits expression of RC2 in NE-4C neuroepithelial cells upon EC23 induction	65
4.2.3.2	BV2 microglia-conditioned media inhibits early neuron production in NE-4C neuroepithelial cells upon EC23 neural induction	69
4.2.4	Effects of microglia-conditioned medium on the expression of cell adhesion-related molecules in NE-4C neuroepithelial cells	73
4.2.4.1	BV2 microglia-conditioned media alters expression of E-cadherin in NE-4C neuroepithelial cells during EC23 neural induction	73
4.2.4.2	BV2 microglia-conditioned media alters expression of N-cadherin in NE-4C neuroepithelial cells during EC23 neural induction	76
4.2.4.3	BV2 microglia-conditioned media alters expression of beta catenin in NE-4C neuroepithelial cells during EC23 neural induction	78
4.2.4.4	BV2 microglia-conditioned media alters expression of CDC42 in non-induced NE-4C neuroepithelial cells but not following EC23 neural induction	80
4.3	Effect of BV2 microglia-conditioned media on neuron production of NE-4C neuroepithelial cells	82



4.3.1	BV2 microglia-conditioned media prevents neuron aggregate formation in NE-4C neuroepithelial cells during neural differentiation	82
4.3.2	BV2 microglia-conditioned media reduce production of TUJ1 <sup>+</sup> neurons but do not affect number of GFAP <sup>+</sup> astrocytes and nestin <sup>+</sup> progenitor cells	85
4.3.3	BV2 microglia-conditioned media inhibit neurite outgrowth and decrease the complexity of neurons derived from NE-4C neuroepithelial cells	90
4.3.4	BV2 microglia-conditioned media does not affect expression of MAP-2 in NE-4C derived neurons	92
4.3.5	BV2 microglia-conditioned media does not affect the expression of SYT1 in NE-4C derived neurons	94
4.4	General Discussion	96
<b>5</b>	<b>CONCLUSION</b>	<b>99</b>
	<b>REFERENCES</b>	<b>102</b>
	<b>APPENDICES</b>	<b>133</b>
	<b>BIODATA OF STUDENT</b>	<b>161</b>
	<b>LIST OF PUBLICATIONS</b>	<b>162</b>

## LIST OF TABLES

Table		Page
2.1	Overview of studies on the role of microglia in embryonic neurogenesis	20
3.1	Sequence of primers used for RT-qPCR	38
3.2	Antibodies and dyes used in immunophenotyping	40
3.3	Primary antibodies used in immunocytochemistry experiments	41
3.4	Secondary antibodies used in immunocytochemistry experiments	42
3.5	Reliability test for automated cell quantification	44
3.6	Primary antibodies used in western blot analysis	46
3.7	Secondary antibodies used in western blot analysis	47

## LIST OF FIGURES

Figure		Page
2.1	The development of neural precursor cells (NPC)	4
2.2	Interkinetic nuclear migration of neuroepithelial cells, radial glial cells and basal progenitors	6
3.1	Acute and chronic LPS stimulation of BV2 microglia	32
4.1	BV2 microglia-conditioned media inhibit NE-4C neuroepithelial cell proliferation	51
4.2	BV2 microglia-conditioned media does not induce cell death in NE-4C neuroepithelial cells	54
4.3	BV2 microglia-conditioned media does not induce apoptosis in NE-4C neuroepithelial cells	55
4.4	Gating strategy of NE-4C neuroepithelial cells for immunophenotyping	57
4.5	BV2 microglia-conditioned media does not alter SSEA-1 and CD133 expression in NE-4C neuroepithelial cells	59
4.6	BV2 microglia-conditioned media does not alter the gene expression of SOX1 and PAX6 in NE-4C neuroepithelial cells	61
4.7	BV2 microglia-conditioned media does not alter the gene expression of HES1, and HES5 in NE-4C neuroepithelial cells	63
4.8	BV2 microglia-conditioned media does not alter the gene expression of MASH1 and NGN2 in NE-4C neuroepithelial cells	64
4.9	BV2 microglia-conditioned media inhibit NE-4C neuroepithelial cell differentiation into radial glia and reduces aggregate formation during neural differentiation induction	68
4.10	BV2 microglia-conditioned media inhibit early neuron production in NE-4C neuroepithelial cell	72
4.11	BV2 microglia-conditioned media alters expression of E-cadherin in NE-4C neuroepithelial cells during EC23 neural induction	75

4.12	BV2 microglia-conditioned media alter the expression of N-cadherin in NE-4C neuroepithelial cells during EC23 neural induction	77
4.13	BV2 microglia-conditioned media alter the expression of beta catenin in NE-4C neuroepithelial cells during EC23 neural induction	79
4.14	BV2 microglia-conditioned media alter the expression of CDC42 in NE-4C neuroepithelial cells during EC23 neural induction	81
4.15	BV2 microglia-conditioned media prevent neuron aggregate formation in NE-4C neuroepithelial cells during terminal neural differentiation	84
4.16	BV2 microglia-conditioned media reduces the number of TUJ1 <sup>+</sup> neurons but does not affect the number of GFAP <sup>+</sup> astrocytes and nestin <sup>+</sup> progenitor cells	89
4.17	BV2 microglia-conditioned media inhibit neurite outgrowth and decrease the complexity of neurons derived from NE-4C neuroepithelial cells	91
4.18	BV2 microglia-conditioned media does not affect the expression of MAP2 in NE-4C derived neurons	93
4.19	BV2 microglia-conditioned media does not affect the expression of SYT1 in NE-4C derived neurons	95

## LIST OF APPENDICES

Appendix		Page
A1	Media and reagent composition	133
A2	Western blot reagents	134
A3	NeurphologyJ interactive step by step guide	135
A4	Automated counting of single color images	140
B1	Data set of Figure 4.1.	142
B2	Data set for Figure 4.3.	143
B3	Data set for Figure 4.5. (SSEA-1)	147
B4	Data set for Figure 4.5. (CD133)	153
B5	Data set for Figure 4.16.	159
B6	Data set for Figure 4.17.	160



## LIST OF ABBREVIATIONS


AGM	aorta–gonad–mesonephros
ANOVA	analysis of variance
BDNF	brain derived neurotrophic factor
BLBP	brain lipid-binding protein
BP	basal progenitor
C1q	a subcomponent of complement C1
C3	complement component 3
CA1	C-shaped Cornu ammonis area 1
CA3	C-shaped Cornu ammonis area 3
CD11b	integrin $\alpha$ M
CD133	prominin-1
CD14	cluster of differentiation 14
CD45	cluster of differentiation 45, a receptor type protein tyrosine phosphatase protein
cDNA	complementary DNA
c-kit	mast/stem cell growth factor receptor (SCFR), also known as proto-oncogene c-Kit
CNS	central nervous system
CNTF	Ciliary Neurotrophic Factor
CR3	complement receptor 3
CSF-1R	Colony Stimulating Factor 1 Receptor
CX <sub>3</sub> CR1	fractalkine receptor
DAP12	DNAX activation protein of 12kDa
DAPI	4',6-diamidino-2-phenylindole, ihydrochloride
DLX1	Distal-Less Homeobox 1
DLX2	Distal-Less Homeobox 2
EC23	synthetic retinoid analogue of all-trans retinoic acid
ECM	extracellular matrix
EDTA	Ethylenediaminetetraacetic acid
EGF	Epidermal growth factor

EMX1	empty spiracles homeobox 1
ER81	a member of the ETS family of transcription factors
ESC	Embryonic stem cell
FGF-2	fibroblast growth factor 2
GCV	ganciclovir
GFAP	Glial fibrillary acidic protein
GLAST	astrocyte-specific glutamate transporter
GSH1	GS Homeobox 1
GSH2	GS Homeobox 2
GSK-3 $\beta$	Glycogen synthase kinase 3 beta
Hbms	hydroxymethylbilane synthase
HES1	Hairy and enhancer of split 1 Drosophila
HES3	Hairy and enhancer of split 3 Drosophila
HES5	Hairy and enhancer of split 5 Drosophila
Hoxb8	Homeobox B8
$^3\text{H-TdR}$	tritium thymidine
IGF1	Insulin-like growth factor
IL-1	Interleukin1
IL-1 $\beta$	Interleukin1 beta
IL-6	Interleukin 6
INM	interkinetic nuclear migration
iNOS	Inducible Nitric oxide synthases
IPC	Intermediate progenitor cells
kDa	Kilo dalton
LIF	Leukaemia inhibitory factor
LPS	bacterial endotoxin lipopolysaccharide
MAP2	Neuronal dendritic marker
MASH1	Mammalian Achaete Scute Homolog-1
MHC II	Major histocompatibility class II
MMP	matrix metalloproteases
Myb	Myb proto-oncogene, transcription factor
NFIA	Nuclear factor 1 A-type

NGF	Nerve growth factor
NGN2	neurogenin 2, neurog2
NKX2.1	NK2 Homeobox 1
Noxa	Phorbol-12-myristate-13-acetate-induced protein 1
NPC	Neural precursor cell
OCT-4	octamer-binding transcription factor 4
OLIG2	Oligodendrocyte transcription factor
PAX6	Paired box protein 6
PBS	Phosphate-buffered saline
PET	positron emission tomography
Pgk1	Phosphoglycerate Kinase 1
PI	propidium iodide
Psmb2	Proteasome subunit beta type-2
PU.1	an ETS-family transcription factor
Puma	p53 upregulated modulator of apoptosis)
PVDF	polyvinylidene difluoride
RC2	radial glial cell marker
RG	Radial glial cells
ROS	reactive oxygen species
xg	Gravity force
Runx1	runt-related transcription factor 1
S100 $\beta$	S100 calcium-binding protein B
SD	Standard deviation
SDS-PAGE	sodium dodecyl sulphate polyacrylamide gel electrophoresis
SGZ	subgranular zone
SOX1	SRY-related HMG-box 1
SOX9	SRY-related HMG-box 9
SP8	specificity protein 8, transcription factor
SSEA-1	stage-specific embryonic antigen1
SVET1	subventricular expressed transcript 1
SVZ	subventricular zone



synCAM-1	Synaptic cell adhesion molecule 1
SYT1	Synaptotagmin 1
Tbr2	T-box brain protein 2
TGF- $\beta$	Transforming growth factor beta
TN-C	Tenascin C
TNFR	tumor necrosis factor receptor
TNF- $\alpha$	tumor necrosis factor alpha
TUJ1	Neuron-specific Class III $\beta$ -tubulin
v-myc	viral-myc gene, myelocytomatosis viral oncogene
v-raf	viral raf gene



## CHAPTER 1

### INTRODUCTION

The brain consists of complex neural circuits made up of neurons and glia. This complex organ exerts centralised control over every aspect of an organism, including basic physiological functions, rudimentary reflexes, cognition, and intellect. During embryonic development, the brain derives from neural precursor cells (NPC) that are highly plastic, proliferative, possess the ability to differentiate into different cell types and eventually acquire regional identities (Gotz & Huttner, 2005; Taverna et al., 2014). Development of the brain is a highly coordinated process that requires precise spatiotemporal regulation of NPC self-renewal and pluripotency in order to generate the right amount and right type of neuronal cells at the right time. Dysregulation of NPC function at embryonic developmental stages can result in brain malformations and later lead to neurodevelopmental disorders (Ernst, 2016).

Microglia are specialised tissue macrophages in the brain that play important roles including provide immune defences, tissue repair and maintaining brain homeostasis (Wolf et al., 2017). Emerging data shows that microglia are actively involved in shaping the development of the prenatal and perinatal brain (Paolicelli et al., 2011; Schafer et al., 2012; Cunningham et al., 2013; Arno et al., 2014). During embryonic development, yolk sac-derived microglia progenitors invade the developing brain and strategically take residence at neurogenic areas (Ginhoux et al., 2010; Cunningham et al., 2013; Swinnen et al., 2013). The invasion of microglia into the brain rudiment, their positioning patterns, and maturation timeline coincides with many embryonic brain development milestones including brain vascularisation, appearance of developmental apoptosis death, neuroepithelial to radial glial transformation, neuronal migration, and myelination (reviewed in (Harry & Kraft, 2012)). Given the time and space correlation of microglia development and important neurogenesis milestones, it is tempting to speculate that microglia are capable of modulating the process of embryonic neurogenesis.

The present study aims to elucidate the effects of microglia on NPC neural differentiation. It was hypothesised that microglia affect the neural differentiation outcome of NPC. For this purpose, the study employed a cell culture approach using the NE-4C neuroepithelial cell line as a model of NPC and BV2 microglia cell line as a model of microglia. BV2 microglia-conditioned media from different culture conditions (unstimulated, acute and chronically activated using lipopolysaccharide; LPS) were collected and used to treat NE-4C neuroepithelial cells under non-induced and retinoic acid-induced neural differentiation conditions. Microglia acquire distinct phenotypes upon acute or chronic immune stimulation (Cacci et al., 2008). It

had been demonstrated that these phenotypes have different effects on NPC properties (Battista et al., 2006; Cacci et al., 2008; Guadagno et al., 2013). Here, the effects of distinct microglia phenotypes (non-activated, acute or chronically activated) were examined. As retinoic acid is important for both *in vitro* and *in vivo* neurogenesis, the effects of microglia in the presence and absence of this neural inducer was explored to deduce their role in induced neurogenesis and whether their effects are independent of retinoic acid-induction. With this approach, firstly the potential of microglia to induce neural differentiation and to regulate NPC biology during the early phase of retinoic acid-induced neural differentiation were determined by examining NE-4C proliferation, cell death and radial glia marker. Next, the expression of cell adhesion-related molecules in NPC were examined. Finally, the production of neurons was measured to further characterise the effects of microglia on NPC terminal neuronal differentiation.

### **General Objective**

The general objective of this project is to determine the effects of microglia on neural differentiation of NPC.

### **Specific Objectives**

The specific objectives are:

1. To examine the effect of microglia on the proliferation of neuroepithelial cells.
2. To evaluate the effect of microglia on neuroepithelial cells differentiation to produce radial glial cells, neurons and astrocytes.
3. To determine the effects of microglia on the expression of cell adhesion-related molecules during neuroepithelial cell differentiation.

## REFERENCES

- Aaku-Saraste, E, Hellwig, A, & Huttner, WB. (1996). Loss of occludin and functional tight junctions, but not ZO-1, during neural tube closure--remodeling of the neuroepithelium prior to neurogenesis. *Dev Biol*, 180(2), 664-679.
- abd-el-Basset, E, & Fedoroff, S. (1995). Effect of bacterial wall lipopolysaccharide (LPS) on morphology, motility, and cytoskeletal organization of microglia in cultures. *J Neurosci Res*, 41(2), 222-237.
- Ajami, B, Bennett, JL, Krieger, C, Tetzlaff, W, & Rossi, FM. (2007). Local self-renewal can sustain CNS microglia maintenance and function throughout adult life. *Nat Neurosci*, 10(12), 1538-1543.
- Akimoto, J, Itoh, H, Miwa, T, & Ikeda, K. (1993). Immunohistochemical study of glutamine synthetase expression in early glial development. *Brain Res Dev Brain Res*, 72(1), 9-14.
- Alliot, F, Godin, I, & Pessac, B. (1999). Microglia derive from progenitors, originating from the yolk sac, and which proliferate in the brain. *Brain Res Dev Brain Res*, 117(2), 145-152.
- Almeida, S, Zhang, Z, Coppola, G, Mao, W, Futai, K, Karydas, A, Geschwind, MD, Tartaglia, MC, Gao, F, Gianni, D, Sena-Esteves, M, Geschwind, DH, Miller, BL, Farese, RV, Jr., & Gao, FB. (2012). Induced pluripotent stem cell models of progranulin-deficient frontotemporal dementia uncover specific reversible neuronal defects. *Cell Rep*, 2(4), 789-798.
- Anjos-Afonso, F, & Bonnet, D. (2007). Nonhematopoietic/endothelial SSEA-1+ cells define the most primitive progenitors in the adult murine bone marrow mesenchymal compartment. *Blood*, 109(3), 1298-1306.
- Anthony, TE, Klein, C, Fishell, G, & Heintz, N. (2004). Radial glia serve as neuronal progenitors in all regions of the central nervous system. *Neuron*, 41(6), 881-890.
- Antony, JM, Paquin, A, Nutt, SL, Kaplan, DR, & Miller, FD. (2011). Endogenous microglia regulate development of embryonic cortical precursor cells. *J Neurosci Res*, 89(3), 286-298.
- Arno, B, Grassivaro, F, Rossi, C, Bergamaschi, A, Castiglioni, V, Furlan, R, Greter, M, Favaro, R, Comi, G, Becher, B, Martino, G, & Muzio, L. (2014). Neural progenitor cells orchestrate microglia migration and positioning into the developing cortex. *Nat Commun*, 5, 5611.

- Arnold, SJ, Huang, GJ, Cheung, AF, Era, T, Nishikawa, S, Bikoff, EK, Molnar, Z, Robertson, EJ, & Groszer, M. (2008). The T-box transcription factor Eomes/Tbr2 regulates neurogenesis in the cortical subventricular zone. *Genes Dev*, 22(18), 2479-2484.
- Arnold, T, & Betsholtz, C. (2013). The importance of microglia in the development of the vasculature in the central nervous system. *Vasc Cell*, 5(1), 4.
- Azevedo, EP, Ledo, JH, Barbosa, G, Sobrinho, M, Diniz, L, Fonseca, AC, Gomes, F, Romao, L, Lima, FR, Palhano, FL, Ferreira, ST, & Foguel, D. (2013). Activated microglia mediate synapse loss and short-term memory deficits in a mouse model of transthyretin-related oculoleptomeningeal amyloidosis. *Cell Death Dis*, 4, e789.
- Barnard, JH, Collings, JC, Whiting, A, Przyborski, SA, & Marder, TB. (2009). Synthetic retinoids: structure-activity relationships. *Chemistry*, 15(43), 11430-11442.
- Barry, D, & McDermott, K. (2005). Differentiation of radial glia from radial precursor cells and transformation into astrocytes in the developing rat spinal cord. *Glia*, 50(3), 187-197.
- Battista, D, Ferrari, CC, Gage, FH, & Pitossi, FJ. (2006). Neurogenic niche modulation by activated microglia: transforming growth factor beta increases neurogenesis in the adult dentate gyrus. *Eur J Neurosci*, 23(1), 83-93.
- Beers, DR, Henkel, JS, Xiao, Q, Zhao, W, Wang, J, Yen, AA, Siklos, L, McKercher, SR, & Appel, SH. (2006). Wild-type microglia extend survival in PU.1 knockout mice with familial amyotrophic lateral sclerosis. *Proc Natl Acad Sci U S A*, 103(43), 16021-16026.
- Bennett, ML, Bennett, FC, Liddelov, SA, Ajami, B, Zamanian, JL, Fernhoff, NB, Mulinyawe, SB, Bohlen, CJ, Adil, A, Tucker, A, Weissman, IL, Chang, EF, Li, G, Grant, GA, Hayden Gephart, MG, & Barres, BA. (2016). New tools for studying microglia in the mouse and human CNS. *Proc Natl Acad Sci U S A*, 113(12), E1738-1746.
- Bertrand, N, Castro, DS, & Guillemot, F. (2002). Proneural genes and the specification of neural cell types. *Nat Rev Neurosci*, 3(7), 517-530.
- Beutner, C, Roy, K, Linnartz, B, Napoli, I, & Neumann, H. (2010). Generation of microglial cells from mouse embryonic stem cells. *Nat Protoc*, 5(9), 1481-1494.

- Bilbo, SD, & Schwarz, JM. (2009). Early-life programming of later-life brain and behavior: a critical role for the immune system. *Front Behav Neurosci*, 3, 14.
- Bittman, K, Owens, DF, Kriegstein, AR, & LoTurco, JJ. (1997). Cell coupling and uncoupling in the ventricular zone of developing neocortex. *J Neurosci*, 17(18), 7037-7044.
- Blasi, E, Barluzzi, R, Bocchini, V, Mazzolla, R, & Bistoni, F. (1990). Immortalization of murine microglial cells by a v-raf/v-myc carrying retrovirus. *J Neuroimmunol*, 27(2-3), 229-237.
- Bloomfield, PS, Selvaraj, S, Veronese, M, Rizzo, G, Bertoldo, A, Owen, DR, Bloomfield, MA, Bonoldi, I, Kalk, N, Turkheimer, F, McGuire, P, de Paola, V, & Howes, OD. (2016). Microglial Activity in People at Ultra High Risk of Psychosis and in Schizophrenia: An [(11)C]PBR28 PET Brain Imaging Study. *Am J Psychiatry*, 173(1), 44-52.
- Borsini, A, Zunszain, PA, Thuret, S, & Pariante, CM. (2015). The role of inflammatory cytokines as key modulators of neurogenesis. *Trends Neurosci*, 38(3), 145-157.
- Butovsky, O, Ziv, Y, Schwartz, A, Landa, G, Talpalar, AE, Pluchino, S, Martino, G, & Schwartz, M. (2006). Microglia activated by IL-4 or IFN-gamma differentially induce neurogenesis and oligodendrogenesis from adult stem/progenitor cells. *Mol Cell Neurosci*, 31(1), 149-160.
- Butovsky, O, Jedrychowski, MP, Moore, CS, Cialic, R, Lanser, AJ, Gabriely, G, Koeglsperger, T, Dake, B, Wu, PM, Doykan, CE, Fanek, Z, Liu, L, Chen, Z, Rothstein, JD, Ransohoff, RM, Gygi, SP, Antel, JP, & Weiner, HL. (2014). Identification of a unique TGF- $\beta$ -dependent molecular and functional signature in microglia. *Nat Neurosci*, 17(1), 131-143.
- Bylund, M, Andersson, E, Novitch, BG, & Muhr, J. (2003). Vertebrate neurogenesis is counteracted by Sox1-3 activity. *Nat Neurosci*, 6(11), 1162-1168.
- Cacci, E, Claasen, JH, & Kokaia, Z. (2005). Microglia-derived tumor necrosis factor-alpha exaggerates death of newborn hippocampal progenitor cells in vitro. *J Neurosci Res*, 80(6), 789-797.
- Cacci, E, Ajmone-Cat, MA, Anelli, T, Biagioni, S, & Minghetti, L. (2008). In vitro neuronal and glial differentiation from embryonic or adult neural precursor cells are differently affected by chronic or acute activation of microglia. *Glia*, 56(4), 412-425.

- Capela, A, & Temple, S. (2006). LeX is expressed by principle progenitor cells in the embryonic nervous system, is secreted into their environment and binds Wnt-1. *Dev Biol*, 291(2), 300-313.
- Cappello, S, Attardo, A, Wu, X, Iwasato, T, Itohara, S, Wilsch-Brauninger, M, Eilken, HM, Rieger, MA, Schroeder, TT, Huttner, WB, Brakebusch, C, & Gotz, M. (2006). The Rho-GTPase cdc42 regulates neural progenitor fate at the apical surface. *Nat Neurosci*, 9(9), 1099-1107.
- Casano, AM, Albert, M, & Peri, F. (2016). Developmental Apoptosis Mediates Entry and Positioning of Microglia in the Zebrafish Brain. *Cell Rep*, 16(4), 897-906.
- Castro, DS, Martynoga, B, Parras, C, Ramesh, V, Pacary, E, Johnston, C, Drechsel, D, Lebel-Potter, M, Garcia, LG, Hunt, C, Dolle, D, Bithell, A, Ettwiller, L, Buckley, N, & Guillemot, F. (2011). A novel function of the proneural factor Ascl1 in progenitor proliferation identified by genome-wide characterization of its targets. *Genes Dev*, 25(9), 930-945.
- Chalasan, K, & Brewster, RM. (2011). N-cadherin-mediated cell adhesion restricts cell proliferation in the dorsal neural tube. *Mol Biol Cell*, 22(9), 1505-1515.
- Chen, L, Liao, G, Yang, L, Campbell, K, Nakafuku, M, Kuan, CY, & Zheng, Y. (2006). Cdc42 deficiency causes Sonic hedgehog-independent holoprosencephaly. *Proc Natl Acad Sci U S A*, 103(44), 16520-16525.
- Chen, S, Lewallen, M, & Xie, T. (2013). Adhesion in the stem cell niche: biological roles and regulation. *Development*, 140(2), 255-265.
- Chen, SK, Tvrdik, P, Peden, E, Cho, S, Wu, S, Spangrude, G, & Capecchi, MR. (2010a). Hematopoietic origin of pathological grooming in Hoxb8 mutant mice. *Cell*, 141(5), 775-785.
- Chen, T, Yuan, D, Wei, B, Jiang, J, Kang, J, Ling, K, Gu, Y, Li, J, Xiao, L, & Pei, G. (2010b). E-cadherin-mediated cell-cell contact is critical for induced pluripotent stem cell generation. *Stem Cells*, 28(8), 1315-1325.
- Chen, Z, & Palmer, TD. (2013). Differential roles of TNFR1 and TNFR2 signaling in adult hippocampal neurogenesis. *Brain Behav Immun*, 30, 45-53.
- Chenn, A, & Walsh, CA. (2002). Regulation of cerebral cortical size by control of cell cycle exit in neural precursors. *Science*, 297(5580), 365-369.

- Chiu, IM, Morimoto, ET, Goodarzi, H, Liao, JT, O'Keeffe, S, Phatnani, HP, Muratet, M, Carroll, MC, Levy, S, Tavazoie, S, Myers, RM, & Maniatis, T. (2013). A neurodegeneration-specific gene-expression signature of acutely isolated microglia from an amyotrophic lateral sclerosis mouse model. *Cell Rep*, 4(2), 385-401.
- Choi, BH. (1981). Radial glia of developing human fetal spinal cord: Golgi, immunohistochemical and electron microscopic study. *Brain Res*, 227(2), 249-267.
- Choi, JY, Kim, JY, Park, J, Lee, WT, & Lee, JE. (2017). M2 Phenotype Microglia-derived Cytokine Stimulates Proliferation and Neuronal Differentiation of Endogenous Stem Cells in Ischemic Brain. *Exp Neurol*, 26(1), 33-41.
- Chojnacki, A, Shimazaki, T, Gregg, C, Weinmaster, G, & Weiss, S. (2003). Glycoprotein 130 signaling regulates Notch1 expression and activation in the self-renewal of mammalian forebrain neural stem cells. *J Neurosci*, 23(5), 1730-1741.
- Christie, VB, Barnard, JH, Batsanov, AS, Bridgens, CE, Cartmell, EB, Collings, JC, Maltman, DJ, Redfern, CP, Marder, TB, Przyborski, S, & Whiting, A. (2008). Synthesis and evaluation of synthetic retinoid derivatives as inducers of stem cell differentiation. *Org Biomol Chem*, 6(19), 3497-3507.
- Corbeil, D, Roper, K, Hannah, MJ, Hellwig, A, & Huttner, WB. (1999). Selective localization of the polytopic membrane protein prominin in microvilli of epithelial cells - a combination of apical sorting and retention in plasma membrane protrusions. *J Cell Sci*, 112 ( Pt 7), 1023-1033.
- Corriveau, RA, Huh, GS, & Shatz, CJ. (1998). Regulation of class I MHC gene expression in the developing and mature CNS by neural activity. *Neuron*, 21(3), 505-520.
- Costello, DA, Lyons, A, Denieffe, S, Browne, TC, Cox, FF, & Lynch, MA. (2011). Long term potentiation is impaired in membrane glycoprotein CD200-deficient mice: a role for Toll-like receptor activation. *J Biol Chem*, 286(40), 34722-34732.
- Courchesne, E, Carper, R, & Akshoomoff, N. (2003). Evidence of brain overgrowth in the first year of life in autism. *JAMA*, 290(3), 337-344.
- Courchesne, E, Mouton, PR, Calhoun, ME, Semendeferi, K, Ahrens-Barbeau, C, Hallet, MJ, Barnes, CC, & Pierce, K. (2011). Neuron number and size in prefrontal cortex of children with autism. *JAMA*, 306(18), 2001-2010.



- Cuadros, MA, Martin, C, Coltey, P, Almendros, A, & Navascues, J. (1993). First appearance, distribution, and origin of macrophages in the early development of the avian central nervous system. *J Comp Neurol*, 330(1), 113-129.
- Culican, SM, Baumrind, NL, Yamamoto, M, & Pearlman, AL. (1990). Cortical radial glia: identification in tissue culture and evidence for their transformation to astrocytes. *J Neurosci*, 10(2), 684-692.
- Cunningham, CL, Martinez-Cerdeno, V, & Noctor, SC. (2013). Microglia regulate the number of neural precursor cells in the developing cerebral cortex. *J Neurosci*, 33(10), 4216-4233.
- D'Arcangelo, G, Tancredi, V, Onofri, F, D'Antuono, M, Giovedi, S, & Benfenati, F. (2000). Interleukin-6 inhibits neurotransmitter release and the spread of excitation in the rat cerebral cortex. *Eur J Neurosci*, 12(4), 1241-1252.
- Dai, XM, Ryan, GR, Hapel, AJ, Dominguez, MG, Russell, RG, Kapp, S, Sylvestre, V, & Stanley, ER. (2002). Targeted disruption of the mouse colony-stimulating factor 1 receptor gene results in osteopetrosis, mononuclear phagocyte deficiency, increased primitive progenitor cell frequencies, and reproductive defects. *Blood*, 99(1), 111-120.
- Dalmau, I, Finsen, B, Zimmer, J, Gonzalez, B, & Castellano, B. (1998). Development of microglia in the postnatal rat hippocampus. *Hippocampus*, 8(5), 458-474.
- Dalmau, I, Vela, JM, Gonzalez, B, Finsen, B, & Castellano, B. (2003). Dynamics of microglia in the developing rat brain. *J Comp Neurol*, 458(2), 144-157.
- Das, S, Dutta, K, Kumawat, KL, Ghoshal, A, Adhya, D, & Basu, A. (2011). Abrogated inflammatory response promotes neurogenesis in a murine model of Japanese encephalitis. *PLoS One*, 6(3), e17225.
- Davalos, D, Grutzendler, J, Yang, G, Kim, JV, Zuo, Y, Jung, S, Littman, DR, Dustin, ML, & Gan, WB. (2005). ATP mediates rapid microglial response to local brain injury in vivo. *Nat Neurosci*, 8(6), 752-758.
- De Pietri Tonelli, D, Pulvers, JN, Haffner, C, Murchison, EP, Hannon, GJ, & Huttner, WB. (2008). miRNAs are essential for survival and differentiation of newborn neurons but not for expansion of neural progenitors during early neurogenesis in the mouse embryonic neocortex. *Development*, 135(23), 3911-3921.
- del Zoppo, GJ, Frankowski, H, Gu, YH, Osada, T, Kanazawa, M, Milner, R, Wang, X, Hosomi, N, Mabuchi, T, & Koziol, JA. (2012). Microglial cell

activation is a source of metalloproteinase generation during hemorrhagic transformation. *J Cereb Blood Flow Metab*, 32(5), 919-932.

- Deleyrolle, LP, Rietze, RL, & Reynolds, BA. (2008). The neurosphere assay, a method under scrutiny. *Acta Neuropsychiatr*, 20(1), 2-8.
- Demeter, K, Herberth, B, Duda, E, Domonkos, A, Jaffredo, T, Herman, JP, & Madarasz, E. (2004). Fate of cloned embryonic neuroectodermal cells implanted into the adult, newborn and embryonic forebrain. *Exp Neurol*, 188(2), 254-267.
- Dilger, RN, & Johnson, RW. (2008). Aging, microglial cell priming, and the discordant central inflammatory response to signals from the peripheral immune system. *J Leukoc Biol*, 84(4), 932-939.
- Duan, D, Fu, Y, Paxinos, G, & Watson, C. (2013). Spatiotemporal expression patterns of Pax6 in the brain of embryonic, newborn, and adult mice. *Brain Struct Funct*, 218(2), 353-372.
- Dubreuil, V, Marzesco, AM, Corbeil, D, Huttner, WB, & Wilsch-Brauninger, M. (2007). Midbody and primary cilium of neural progenitors release extracellular membrane particles enriched in the stem cell marker prominin-1. *J Cell Biol*, 176(4), 483-495.
- Dvorak, P, Hampl, A, Jirmanova, L, Pacholikova, J, & Kusakabe, M. (1998). Embryoglycan ectodomains regulate biological activity of FGF-2 to embryonic stem cells. *J Cell Sci*, 111 ( Pt 19), 2945-2952.
- Dvorak, P, Dvorakova, D, Koskova, S, Vodinska, M, Najvirtova, M, Krekac, D, & Hampl, A. (2005). Expression and potential role of fibroblast growth factor 2 and its receptors in human embryonic stem cells. *Stem Cells*, 23(8), 1200-1211.
- Edwards, MA, Yamamoto, M, & Caviness, VS, Jr. (1990). Organization of radial glia and related cells in the developing murine CNS. An analysis based upon a new monoclonal antibody marker. *Neuroscience*, 36(1), 121-144.
- Eggen, BJ, Raj, D, Hanisch, UK, & Boddeke, HW. (2013). Microglial phenotype and adaptation. *J Neuroimmune Pharmacol*, 8(4), 807-823.
- Eglitis, MA, & Mezey, E. (1997). Hematopoietic cells differentiate into both microglia and macroglia in the brains of adult mice. *Proc Natl Acad Sci U S A*, 94(8), 4080-4085.

- Ekdahl, CT, Claasen, JH, Bonde, S, Kokaia, Z, & Lindvall, O. (2003). Inflammation is detrimental for neurogenesis in adult brain. *Proc Natl Acad Sci U S A*, 100(23), 13632-13637.
- Elkabes, S, DiCicco-Bloom, EM, & Black, IB. (1996). Brain microglia/macrophages express neurotrophins that selectively regulate microglial proliferation and function. *J Neurosci*, 16(8), 2508-2521.
- Endo, M, Antonyak, MA, & Cerione, RA. (2009). Cdc42-mTOR signaling pathway controls Hes5 and Pax6 expression in retinoic acid-dependent neural differentiation. *J Biol Chem*, 284(8), 5107-5118.
- Englund, C, Fink, A, Lau, C, Pham, D, Daza, RA, Bulfone, A, Kowalczyk, T, & Hevner, RF. (2005). Pax6, Tbr2, and Tbr1 are expressed sequentially by radial glia, intermediate progenitor cells, and postmitotic neurons in developing neocortex. *J Neurosci*, 25(1), 247-251.
- Erblich, B, Zhu, L, Etgen, AM, Dobrenis, K, & Pollard, JW. (2011). Absence of colony stimulation factor-1 receptor results in loss of microglia, disrupted brain development and olfactory deficits. *PLoS One*, 6(10), e26317.
- Ernst, C. (2016). Proliferation and Differentiation Deficits are a Major Convergence Point for Neurodevelopmental Disorders. *Trends Neurosci*, 39(5), 290-299.
- Esdar, C, Milasta, S, Maelicke, A, & Herget, T. (2001). Differentiation-associated apoptosis of neural stem cells is effected by Bcl-2 overexpression: impact on cell lineage determination. *Eur J Cell Biol*, 80(8), 539-553.
- Evrard, C, Borde, I, Marin, P, Galiana, E, Premont, J, Gros, F, & Rouget, P. (1990). immortalization of bipotential and plastic glio-neuronal precursor cells. *Proc Natl Acad Sci U S A*, 87(8), 3062-3066.
- Eyo, UB, & Dailey, ME. (2013). Microglia: key elements in neural development, plasticity, and pathology. *J Neuroimmune Pharmacol*, 8(3), 494-509.
- Fedoroff, S, Zhai, R, & Novak, JP. (1997). Microglia and astroglia have a common progenitor cell. *J Neurosci Res*, 50(3), 477-486.
- Fernandez, V, Llinares-Benadero, C, & Borrell, V. (2016). Cerebral cortex expansion and folding: what have we learned? *EMBO J*, 35(10), 1021-1044.

- Fillman, SG, Cloonan, N, Catts, VS, Miller, LC, Wong, J, McCrossin, T, Cairns, M, & Weickert, CS. (2013). Increased inflammatory markers identified in the dorsolateral prefrontal cortex of individuals with schizophrenia. *Mol Psychiatry*, 18(2), 206-214.
- Fontainhas, AM, Wang, M, Liang, KJ, Chen, S, Mettu, P, Damani, M, Fariss, RN, Li, W, & Wong, WT. (2011). Microglial morphology and dynamic behavior is regulated by ionotropic glutamatergic and GABAergic neurotransmission. *PLoS One*, 6(1), e15973.
- Frade, JM, & Barde, YA. (1998). Microglia-derived nerve growth factor causes cell death in the developing retina. *Neuron*, 20(1), 35-41.
- Frick, LR, Williams, K, & Pittenger, C. (2013). Microglial dysregulation in psychiatric disease. *Clin Dev Immunol*, 2013, 608654.
- Gabrusiewicz, K, Ellert-Miklaszewska, A, Lipko, M, Sielska, M, Frankowska, M, & Kaminska, B. (2011). Characteristics of the alternative phenotype of microglia/macrophages and its modulation in experimental gliomas. *PLoS One*, 6(8), e23902.
- Gebara, E, Sultan, S, Kocher-Braissant, J, & Toni, N. (2013). Adult hippocampal neurogenesis inversely correlates with microglia in conditions of voluntary running and aging. *Front Neurosci*, 7, 145.
- Gemma, C, Bachstetter, AD, Cole, MJ, Fister, M, Hudson, C, & Bickford, PC. (2007). Blockade of caspase-1 increases neurogenesis in the aged hippocampus. *Eur J Neurosci*, 26(10), 2795-2803.
- Ginhoux, F, Greter, M, Leboeuf, M, Nandi, S, See, P, Gokhan, S, Mehler, MF, Conway, SJ, Ng, LG, Stanley, ER, Samokhvalov, IM, & Merad, M. (2010). Fate mapping analysis reveals that adult microglia derive from primitive macrophages. *Science*, 330(6005), 841-845.
- Ginhoux, F, Lim, S, Hoeffel, G, Low, D, & Huber, T. (2013). Origin and differentiation of microglia. *Front Cell Neurosci*, 7, 45.
- Giovanoli, S, Engler, H, Engler, A, Richetto, J, Voget, M, Willi, R, Winter, C, Riva, MA, Mortensen, PB, Feldon, J, Schedlowski, M, & Meyer, U. (2013). Stress in puberty unmasks latent neuropathological consequences of prenatal immune activation in mice. *Science*, 339(6123), 1095-1099.
- Goings, GE, Kozlowski, DA, & Szele, FG. (2006). Differential activation of microglia in neurogenic versus non-neurogenic regions of the forebrain. *Glia*, 54(4), 329-342.

- Golan, H, Levav, T, Mendelsohn, A, & Huleihel, M. (2004). Involvement of tumor necrosis factor alpha in hippocampal development and function. *Cereb Cortex*, 14(1), 97-105.
- Goldmann, T, & Prinz, M. (2013). Role of microglia in CNS autoimmunity. *Clin Dev Immunol*, 2013, 208093.
- Gotz, M, Bolz, J, Joester, A, & Faissner, A. (1997). Tenascin-C synthesis and influence on axonal growth during rat cortical development. *Eur J Neurosci*, 9(3), 496-506.
- Gotz, M, Stoykova, A, & Gruss, P. (1998). Pax6 controls radial glia differentiation in the cerebral cortex. *Neuron*, 21(5), 1031-1044.
- Gotz, M, & Huttner, WB. (2005). The cell biology of neurogenesis. *Nat Rev Mol Cell Biol*, 6(10), 777-788.
- Govek, EE, Newey, SE, & Van Aelst, L. (2005). The role of the Rho GTPases in neuronal development. *Genes Dev*, 19(1), 1-49.
- Greter, M, Lelios, I, & Croxford, AL. (2015). Microglia Versus Myeloid Cell Nomenclature during Brain Inflammation. *Front Immunol*, 6, 249.
- Guadagno, J, Xu, X, Karajgikar, M, Brown, A, & Cregan, SP. (2013). Microglia-derived TNFalpha induces apoptosis in neural precursor cells via transcriptional activation of the Bcl-2 family member Puma. *Cell Death Dis*, 4, e538.
- Guadagno, J, Swan, P, Shaikh, R, & Cregan, SP. (2015). Microglia-derived IL-1beta triggers p53-mediated cell cycle arrest and apoptosis in neural precursor cells. *Cell Death Dis*, 6, e1779.
- Guillemot, F, & Hassan, BA. (2017). Beyond proneural: emerging functions and regulations of proneural proteins. *Curr Opin Neurobiol*, 42, 93-101.
- Halbleib, JM, & Nelson, WJ. (2006). Cadherins in development: cell adhesion, sorting, and tissue morphogenesis. *Genes Dev*, 20(23), 3199-3214.
- Hanisch, UK, & Kettenmann, H. (2007). Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci*, 10(11), 1387-1394.
- Hao, C, Richardson, A, & Fedoroff, S. (1991). Macrophage-like cells originate from neuroepithelium in culture: characterization and properties of the macrophage-like cells. *Int J Dev Neurosci*, 9(1), 1-14.
- Harry, GJ, & Kraft, AD. (2012). Microglia in the developing brain: a potential target with lifetime effects. *Neurotoxicology*, 33(2), 191-206.

- Hartfuss, E, Galli, R, Heins, N, & Gotz, M. (2001). Characterization of CNS precursor subtypes and radial glia. *Dev Biol*, 229(1), 15-30.
- Hatakeyama, J, Bessho, Y, Katoh, K, Ookawara, S, Fujioka, M, Guillemot, F, & Kageyama, R. (2004). Hes genes regulate size, shape and histogenesis of the nervous system by control of the timing of neural stem cell differentiation. *Development*, 131(22), 5539-5550.
- Hatakeyama, J, Wakamatsu, Y, Nagafuchi, A, Kageyama, R, Shigemoto, R, & Shimamura, K. (2014). Cadherin-based adhesions in the apical endfoot are required for active Notch signaling to control neurogenesis in vertebrates. *Development*, 141(8), 1671-1682.
- He, W, Ingraham, C, Rising, L, Goderie, S, & Temple, S. (2001). Multipotent stem cells from the mouse basal forebrain contribute GABAergic neurons and oligodendrocytes to the cerebral cortex during embryogenesis. *J Neurosci*, 21(22), 8854-8862.
- Hennen, E, Czopka, T, & Faissner, A. (2011). Structurally distinct LewisX glycans distinguish subpopulations of neural stem/progenitor cells. *J Biol Chem*, 286(18), 16321-16331.
- Henry, CJ, Huang, Y, Wynne, AM, & Godbout, JP. (2009). Peripheral lipopolysaccharide (LPS) challenge promotes microglial hyperactivity in aged mice that is associated with exaggerated induction of both pro-inflammatory IL-1beta and anti-inflammatory IL-10 cytokines. *Brain Behav Immun*, 23(3), 309-317.
- Herget, T, Specht, H, Esdar, C, Oehrlein, SA, & Maelicke, A. (1998). Retinoic acid induces apoptosis-associated neural differentiation of a murine teratocarcinoma cell line. *J Neurochem*, 70(1), 47-58.
- Hinze, A, & Stolzing, A. (2011). Differentiation of mouse bone marrow derived stem cells toward microglia-like cells. *BMC Cell Biol*, 12, 35.
- Hinze, A, & Stolzing, A. (2012). Microglia differentiation using a culture system for the expansion of mice non-adherent bone marrow stem cells. *J Inflamm (Lond)*, 9(1), 12.
- Ho, SY, Chao, CY, Huang, HL, Chiu, TW, Charoenkwan, P, & Hwang, E. (2011). NeurphologyJ: an automatic neuronal morphology quantification method and its application in pharmacological discovery. *BMC Bioinformatics*, 12, 230.
- Hockfield, S, & McKay, RD. (1985). Identification of major cell classes in the developing mammalian nervous system. *J Neurosci*, 5(12), 3310-3328.

- Homem, CC, Repic, M, & Knoblich, JA. (2015). Proliferation control in neural stem and progenitor cells. *Nat Rev Neurosci*, 16(11), 647-659.
- Hoshiko, M, Arnoux, I, Avignone, E, Yamamoto, N, & Audinat, E. (2012). Deficiency of the microglial receptor CX3CR1 impairs postnatal functional development of thalamocortical synapses in the barrel cortex. *J Neurosci*, 32(43), 15106-15111.
- Hsieh, CL, Koike, M, Spusta, SC, Niemi, EC, Yenari, M, Nakamura, MC, & Seaman, WE. (2009). A role for TREM2 ligands in the phagocytosis of apoptotic neuronal cells by microglia. *J Neurochem*, 109(4), 1144-1156.
- Huang, T, Cui, J, Li, L, Hitchcock, PF, & Li, Y. (2012). The role of microglia in the neurogenesis of zebrafish retina. *Biochem Biophys Res Commun*, 421(2), 214-220.
- Hughes, MM, Field, RH, Perry, VH, Murray, CL, & Cunningham, C. (2010). Microglia in the degenerating brain are capable of phagocytosis of beads and of apoptotic cells, but do not efficiently remove PrPSc, even upon LPS stimulation. *Glia*, 58(16), 2017-2030.
- Huh, GS, Boulanger, LM, Du, H, Riquelme, PA, Brotz, TM, & Shatz, CJ. (2000). Functional requirement for class I MHC in CNS development and plasticity. *Science*, 290(5499), 2155-2159.
- Hutchins, KD, Dickson, DW, Rashbaum, WK, & Lyman, WD. (1990). Localization of morphologically distinct microglial populations in the developing human fetal brain: implications for ontogeny. *Brain Res Dev Brain Res*, 55(1), 95-102.
- Inestrosa, NC, & Varela-Nallar, L. (2015). Wnt signalling in neuronal differentiation and development. *Cell Tissue Res*, 359(1), 215-223.
- Inuzuka, H, Redies, C, & Takeichi, M. (1991). Differential expression of R- and N-cadherin in neural and mesodermal tissues during early chicken development. *Development*, 113(3), 959-967.
- Iosif, RE, Ekdahl, CT, Ahlenius, H, Pronk, CJ, Bonde, S, Kokaia, Z, Jacobsen, SE, & Lindvall, O. (2006). Tumor necrosis factor receptor 1 is a negative regulator of progenitor proliferation in adult hippocampal neurogenesis. *J Neurosci*, 26(38), 9703-9712.
- Iosif, RE, Ahlenius, H, Ekdahl, CT, Darsalia, V, Thored, P, Jovinge, S, Kokaia, Z, & Lindvall, O. (2008). Suppression of stroke-induced progenitor proliferation in adult subventricular zone by tumor necrosis factor receptor 1. *J Cereb Blood Flow Metab*, 28(9), 1574-1587.

- Izant, JG, & McIntosh, JR. (1980). Microtubule-associated proteins: a monoclonal antibody to MAP2 binds to differentiated neurons. *Proc Natl Acad Sci U S A*, 77(8), 4741-4745.
- Janesick, A, Wu, SC, & Blumberg, B. (2015). Retinoic acid signaling and neuronal differentiation. *Cell Mol Life Sci*, 72(8), 1559-1576.
- Jaszai, J, Graupner, S, Tanaka, EM, Funk, RH, Huttner, WB, Brand, M, & Corbeil, D. (2013). Spatial distribution of prominin-1 (CD133)-positive cells within germinative zones of the vertebrate brain. *PLoS One*, 8(5), e63457.
- Jenrow, KA, Brown, SL, Lapanowski, K, Naei, H, Kolozsvary, A, & Kim, JH. (2013). Selective inhibition of microglia-mediated neuroinflammation mitigates radiation-induced cognitive impairment. *Radiat Res*, 179(5), 549-556.
- Jensen, JB, & Parmar, M. (2006). Strengths and limitations of the neurosphere culture system. *Mol Neurobiol*, 34(3), 153-161.
- Ji, K, Akgul, G, Wollmuth, LP, & Tsirka, SE. (2013a). Microglia actively regulate the number of functional synapses. *PLoS One*, 8(2), e56293.
- Ji, R, Tian, S, Lu, HJ, Lu, Q, Zheng, Y, Wang, X, Ding, J, & Li, Q. (2013b). TAM receptors affect adult brain neurogenesis by negative regulation of microglial cell activation. *J Immunol*, 191(12), 6165-6177.
- Kadowaki, M, Nakamura, S, Machon, O, Krauss, S, Radice, GL, & Takeichi, M. (2007). N-cadherin mediates cortical organization in the mouse brain. *Dev Biol*, 304(1), 22-33.
- Kang, P, Lee, HK, Glasgow, SM, Finley, M, Donti, T, Gaber, ZB, Graham, BH, Foster, AE, Novitsch, BG, Gronostajski, RM, & Deneen, B. (2012). Sox9 and NFIA coordinate a transcriptional regulatory cascade during the initiation of gliogenesis. *Neuron*, 74(1), 79-94.
- Karpowicz, P, Willaime-Morawek, S, Balenci, L, DeVeale, B, Inoue, T, & van der Kooy, D. (2009). E-Cadherin regulates neural stem cell self-renewal. *J Neurosci*, 29(12), 3885-3896.
- Kettenmann, H, Hanisch, UK, Noda, M, & Verkhratsky, A. (2011). Physiology of microglia. *Physiol Rev*, 91(2), 461-553.
- Khan, D, Fernando, P, Cicvaric, A, Berger, A, Pollak, A, Monje, FJ, & Pollak, DD. (2014). Long-term effects of maternal immune activation on depression-like behavior in the mouse. *Transl Psychiatry*, 4, e363.



- Kierdorf, K, Erny, D, Goldmann, T, Sander, V, Schulz, C, Perdiguero, EG, Wieghofer, P, Heinrich, A, Riemke, P, Holscher, C, Muller, DN, Luckow, B, Brocker, T, Debowski, K, Fritz, G, Opdenakker, G, Diefenbach, A, Biber, K, Heikenwalder, M, Geissmann, F, Rosenbauer, F, & Prinz, M. (2013). Microglia emerge from erythromyeloid precursors via Pu.1- and Irf8-dependent pathways. *Nat Neurosci*, 16(3), 273-280.
- Kingwell, K. (2012). Neurodegenerative disease: Microglia in early disease stages. *Nat Rev Neurol*, 8(9), 475.
- Kitamura, T, Miyake, T, & Fujita, S. (1984). Genesis of resting microglia in the gray matter of mouse hippocampus. *J Comp Neurol*, 226(3), 421-433.
- Kornyei, Z, Szlavik, V, Szabo, B, Gocza, E, Czirok, A, & Madarasz, E. (2005). Humoral and contact interactions in astroglia/stem cell co-cultures in the course of glia-induced neurogenesis. *Glia*, 49(3), 430-444.
- Kowalczyk, T, Pontious, A, Englund, C, Daza, RA, Bedogni, F, Hodge, R, Attardo, A, Bell, C, Huttner, WB, & Hevner, RF. (2009). Intermediate neuronal progenitors (basal progenitors) produce pyramidal-projection neurons for all layers of cerebral cortex. *Cereb Cortex*, 19(10), 2439-2450.
- Kriegstein, A, & Alvarez-Buylla, A. (2009). The glial nature of embryonic and adult neural stem cells. *Annu Rev Neurosci*, 32, 149-184.
- Kurusu, M, Katsuki, T, Zinn, K, & Suzuki, E. (2012). Developmental changes in expression, subcellular distribution, and function of Drosophila N-cadherin, guided by a cell-intrinsic program during neuronal differentiation. *Dev Biol*, 366(2), 204-217.
- Kurz, H, & Christ, B. (1998). Embryonic CNS macrophages and microglia do not stem from circulating, but from extravascular precursors. *Glia*, 22(1), 98-102.
- L'Episcopo, F, Tirolo, C, Testa, N, Caniglia, S, Morale, MC, Impagnatiello, F, & Marchetti, B. (2011). Switching the microglial harmful phenotype promotes lifelong restoration of substantia nigra dopaminergic neurons from inflammatory neurodegeneration in aged mice. *Rejuvenation Res*, 14(4), 411-424.
- L'Episcopo, F, Tirolo, C, Testa, N, Caniglia, S, Morale, MC, Deleidi, M, Serapide, MF, Pluchino, S, & Marchetti, B. (2012). Plasticity of subventricular zone neuroprogenitors in MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) mouse model of Parkinson's disease involves cross talk between inflammatory and Wnt/beta-catenin

signaling pathways: functional consequences for neuroprotection and repair. *J Neurosci*, 32(6), 2062-2085.

Laplante, I, Beliveau, R, & Paquin, J. (2004). RhoA/ROCK and Cdc42 regulate cell-cell contact and N-cadherin protein level during neurodetermination of P19 embryonal stem cells. *J Neurobiol*, 60(3), 289-307.

Larue, L, Ohsugi, M, Hirchenhain, J, & Kemler, R. (1994). E-cadherin null mutant embryos fail to form a trophectoderm epithelium. *Proc Natl Acad Sci U S A*, 91(17), 8263-8267.

Lawson, LJ, Perry, VH, Dri, P, & Gordon, S. (1990). Heterogeneity in the distribution and morphology of microglia in the normal adult mouse brain. *Neuroscience*, 39(1), 151-170.

Le Belle, JE, Sperry, J, Ngo, A, Ghochani, Y, Laks, DR, Lopez-Aranda, M, Silva, AJ, & Kornblum, HI. (2014). Maternal inflammation contributes to brain overgrowth and autism-associated behaviors through altered redox signaling in stem and progenitor cells. *Stem Cell Reports*, 3(5), 725-734.

Lee, AS, Azmitia, EC, & Whitaker-Azmitia, PM. (2017). Developmental microglial priming in postmortem autism spectrum disorder temporal cortex. *Brain Behav Immun*, 62, 193-202.

Levitt, P, & Rakic, P. (1980). Immunoperoxidase localization of glial fibrillary acidic protein in radial glial cells and astrocytes of the developing rhesus monkey brain. *J Comp Neurol*, 193(3), 815-840.

Linard, C, Marquette, C, Mathieu, J, Pennequin, A, Clarencon, D, & Mathe, D. (2004). Acute induction of inflammatory cytokine expression after gamma-irradiation in the rat: effect of an NF-kappaB inhibitor. *Int J Radiat Oncol Biol Phys*, 58(2), 427-434.

Ling, EA, Penney, D, & Leblond, CP. (1980). Use of carbon labeling to demonstrate the role of blood monocytes as precursors of the 'ameboid cells' present in the corpus callosum of postnatal rats. *J Comp Neurol*, 193(3), 631-657.

Linnartz, B, Kopatz, J, Tenner, AJ, & Neumann, H. (2012). Sialic acid on the neuronal glycocalyx prevents complement C1 binding and complement receptor-3-mediated removal by microglia. *J Neurosci*, 32(3), 946-952.

Liu, Z, Fan, Y, Won, SJ, Neumann, M, Hu, D, Zhou, L, Weinstein, PR, & Liu, J. (2007). Chronic treatment with minocycline preserves adult new

neurons and reduces functional impairment after focal cerebral ischemia. *Stroke*, 38(1), 146-152.

Livingstone, LR, White, A, Sprouse, J, Livanos, E, Jacks, T, & Tlsty, TD. (1992). Altered cell cycle arrest and gene amplification potential accompany loss of wild-type p53. *Cell*, 70(6), 923-935.

Luo, XG, Ding, JQ, & Chen, SD. (2010). Microglia in the aging brain: relevance to neurodegeneration. *Mol Neurodegener*, 5, 12.

Luo, XG, & Chen, SD. (2012). The changing phenotype of microglia from homeostasis to disease. *Transl Neurodegener*, 1(1), 9.

Lynch, MA. (2009). The multifaceted profile of activated microglia. *Mol Neurobiol*, 40(2), 139-156.

Machon, O, Backman, M, Machonova, O, Kozmik, Z, Vacik, T, Andersen, L, & Krauss, S. (2007). A dynamic gradient of Wnt signaling controls initiation of neurogenesis in the mammalian cortex and cellular specification in the hippocampus. *Dev Biol*, 311(1), 223-237.

Maggi, L, Scianni, M, Branchi, I, D'Andrea, I, Lauro, C, & Limatola, C. (2011). CX(3)CR1 deficiency alters hippocampal-dependent plasticity phenomena blunting the effects of enriched environment. *Front Cell Neurosci*, 5, 22.

Malatesta, P, Hartfuss, E, & Gotz, M. (2000). Isolation of radial glial cells by fluorescent-activated cell sorting reveals a neuronal lineage. *Development*, 127(24), 5253-5263.

Malatesta, P, Hack, MA, Hartfuss, E, Kettenmann, H, Klinkert, W, Kirchhoff, F, & Gotz, M. (2003). Neuronal or glial progeny: regional differences in radial glia fate. *Neuron*, 37(5), 751-764.

Malatesta, P, Appolloni, I, & Calzolari, F. (2008). Radial glia and neural stem cells. *Cell Tissue Res*, 331(1), 165-178.

Maltman, DJ, Christie, VB, Collings, JC, Barnard, JH, Fenyk, S, Marder, TB, Whiting, A, & Przyborski, SA. (2009). Proteomic profiling of the stem cell response to retinoic acid and synthetic retinoid analogues: identification of major retinoid-inducible proteins. *Mol Biosyst*, 5(5), 458-471.

Mander, TH, & Morris, JF. (1996). Development of microglia and macrophages in the postnatal rat pituitary. *Cell Tissue Res*, 286(3), 347-355.

- Marin-Teva, JL, Dusart, I, Colin, C, Gervais, A, van Rooijen, N, & Mallat, M. (2004). Microglia promote the death of developing Purkinje cells. *Neuron*, 41(4), 535-547.
- Marthiens, V, Kazanis, I, Moss, L, Long, K, & Ffrench-Constant, C. (2010). Adhesion molecules in the stem cell niche--more than just staying in shape? *J Cell Sci*, 123(Pt 10), 1613-1622.
- Marzesco, AM, Janich, P, Wilsch-Brauninger, M, Dubreuil, V, Langenfeld, K, Corbeil, D, & Huttner, WB. (2005). Release of extracellular membrane particles carrying the stem cell marker prominin-1 (CD133) from neural progenitors and other epithelial cells. *J Cell Sci*, 118(Pt 13), 2849-2858.
- Mattei, D, Djodari-Irani, A, Hadar, R, Pelz, A, de Cossio, LF, Goetz, T, Matyash, M, Kettenmann, H, Winter, C, & Wolf, SA. (2014a). Corrigendum to "Minocycline rescues decrease in neurogenesis, increase in microglia cytokines and deficits in sensorimotor gating in an animal model of schizophrenia" [Brain Behav. Immun. 38 (2014) 175-184]. *Brain Behav Immun*.
- Mattei, D, Djodari-Irani, A, Hadar, R, Pelz, A, de Cossio, LF, Goetz, T, Matyash, M, Kettenmann, H, Winter, C, & Wolf, SA. (2014b). Minocycline rescues decrease in neurogenesis, increase in microglia cytokines and deficits in sensorimotor gating in an animal model of schizophrenia. *Brain Behav Immun*, 38, 175-184.
- Mattei, D, Ivanov, A, Ferrai, C, Jordan, P, Guneykaya, D, Buonfiglioli, A, Schaafsma, W, Przanowski, P, Deuther-Conrad, W, Brust, P, Hesse, S, Patt, M, Sabri, O, Ross, TL, Eggen, BJL, Boddeke, E, Kaminska, B, Beule, D, Pombo, A, Kettenmann, H, & Wolf, SA. (2017). Maternal immune activation results in complex microglial transcriptome signature in the adult offspring that is reversed by minocycline treatment. *Transl Psychiatry*, 7(5), e1120.
- McKanna, JA. (1993a). Lipocortin 1 immunoreactivity identifies microglia in adult rat brain. *J Neurosci Res*, 36(4), 491-500.
- McKanna, JA. (1993b). Primitive glial compartments in the floor plate of mammalian embryos: distinct progenitors of adult astrocytes and microglia support the notoplate hypothesis. *Perspect Dev Neurobiol*, 1(4), 245-255.
- McKercher, SR, Torbett, BE, Anderson, KL, Henkel, GW, Vestal, DJ, Baribault, H, Klemsz, M, Feeney, AJ, Wu, GE, Paige, CJ, & Maki, RA. (1996). Targeted disruption of the PU.1 gene results in multiple hematopoietic abnormalities. *EMBO J*, 15(20), 5647-5658.

- McPherson, CA, Kraft, AD, & Harry, GJ. (2011). Injury-induced neurogenesis: consideration of resident microglia as supportive of neural progenitor cells. *Neurotox Res*, 19(2), 341-352.
- Mead, EL, Mosley, A, Eaton, S, Dobson, L, Heales, SJ, & Pocock, JM. (2012). Microglial neurotransmitter receptors trigger superoxide production in microglia; consequences for microglial-neuronal interactions. *J Neurochem*, 121(2), 287-301.
- Menezes, JR, & Luskin, MB. (1994). Expression of neuron-specific tubulin defines a novel population in the proliferative layers of the developing telencephalon. *J Neurosci*, 14(9), 5399-5416.
- Merkle, FT, Tramontin, AD, Garcia-Verdugo, JM, & Alvarez-Buylla, A. (2004). Radial glia give rise to adult neural stem cells in the subventricular zone. *Proc Natl Acad Sci U S A*, 101(50), 17528-17532.
- Mildner, A, Schmidt, H, Nitsche, M, Merkler, D, Hanisch, UK, Mack, M, Heikenwalder, M, Bruck, W, Priller, J, & Prinz, M. (2007). Microglia in the adult brain arise from Ly-6ChiCCR2+ monocytes only under defined host conditions. *Nat Neurosci*, 10(12), 1544-1553.
- Mildner, A, Schlevogt, B, Kierdorf, K, Bottcher, C, Erny, D, Kummer, MP, Quinn, M, Bruck, W, Bechmann, I, Heneka, MT, Priller, J, & Prinz, M. (2011). Distinct and non-redundant roles of microglia and myeloid subsets in mouse models of Alzheimer's disease. *J Neurosci*, 31(31), 11159-11171.
- Mission, JP, Takahashi, T, & Caviness, VS, Jr. (1991). Ontogeny of radial and other astroglial cells in murine cerebral cortex. *Glia*, 4(2), 138-148.
- Misson, JP, Austin, CP, Takahashi, T, Cepko, CL, & Caviness, VS, Jr. (1991). The alignment of migrating neural cells in relation to the murine neopallial radial glial fiber system. *Cereb Cortex*, 1(3), 221-229.
- Miyamoto, A, Wake, H, Ishikawa, AW, Eto, K, Shibata, K, Murakoshi, H, Koizumi, S, Moorhouse, AJ, Yoshimura, Y, & Nabekura, J. (2016). Microglia contact induces synapse formation in developing somatosensory cortex. *Nat Commun*, 7, 12540.
- Miyata, T, Kawaguchi, D, Kawaguchi, A, & Gotoh, Y. (2010). Mechanisms that regulate the number of neurons during mouse neocortical development. *Curr Opin Neurobiol*, 20(1), 22-28.
- Mizuguchi, R, Sugimori, M, Takebayashi, H, Kosako, H, Nagao, M, Yoshida, S, Nabeshima, Y, Shimamura, K, & Nakafuku, M. (2001). Combinatorial roles of olig2 and neurogenin2 in the coordinated

induction of pan-neuronal and subtype-specific properties of motoneurons. *Neuron*, 31(5), 757-771.

Monje, ML, Toda, H, & Palmer, TD. (2003). Inflammatory blockade restores adult hippocampal neurogenesis. *Science*, 302(5651), 1760-1765.

Morris, L, Graham, CF, & Gordon, S. (1991). Macrophages in haemopoietic and other tissues of the developing mouse detected by the monoclonal antibody F4/80. *Development*, 112(2), 517-526.

Morrison, SJ, Perez, SE, Qiao, Z, Verdi, JM, Hicks, C, Weinmaster, G, & Anderson, DJ. (2000). Transient Notch activation initiates an irreversible switch from neurogenesis to gliogenesis by neural crest stem cells. *Cell*, 101(5), 499-510.

Mosher, KI, Andres, RH, Fukuhara, T, Bieri, G, Hasegawa-Moriyama, M, He, Y, Guzman, R, & Wyss-Coray, T. (2012). Neural progenitor cells regulate microglia functions and activity. *Nat Neurosci*, 15(11), 1485-1487.

Muzio, L, Di Benedetto, B, Stoykova, A, Boncinelli, E, Gruss, P, & Mallamaci, A. (2002). Conversion of cerebral cortex into basal ganglia in *Emx2(-/-) Pax6(Sey/Sey)* double-mutant mice. *Nat Neurosci*, 5(8), 737-745.

Nakajima, K, Tohyama, Y, Maeda, S, Kohsaka, S, & Kurihara, T. (2007). Neuronal regulation by which microglia enhance the production of neurotrophic factors for GABAergic, catecholaminergic, and cholinergic neurons. *Neurochem Int*, 50(6), 807-820.

Nandi, S, Gokhan, S, Dai, XM, Wei, S, Enikolopov, G, Lin, H, Mehler, MF, & Stanley, ER. (2012). The CSF-1 receptor ligands IL-34 and CSF-1 exhibit distinct developmental brain expression patterns and regulate neural progenitor cell maintenance and maturation. *Dev Biol*, 367(2), 100-113.

Napoli, I, Kierdorf, K, & Neumann, H. (2009). Microglial precursors derived from mouse embryonic stem cells. *Glia*, 57(15), 1660-1671.

Nieto, M, Schuurmans, C, Britz, O, & Guillemot, F. (2001). Neural bHLH genes control the neuronal versus glial fate decision in cortical progenitors. *Neuron*, 29(2), 401-413.

Nimmerjahn, A, Kirchhoff, F, & Helmchen, F. (2005). Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science*, 308(5726), 1314-1318.

Noctor, SC, Flint, AC, Weissman, TA, Wong, WS, Clinton, BK, & Kriegstein, AR. (2002). Dividing precursor cells of the embryonic cortical

ventricular zone have morphological and molecular characteristics of radial glia. *J Neurosci*, 22(8), 3161-3173.

Noctor, SC, Martinez-Cerdeno, V, Ivic, L, & Kriegstein, AR. (2004). Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases. *Nat Neurosci*, 7(2), 136-144.

Noctor, SC, Martinez-Cerdeno, V, & Kriegstein, AR. (2008). Distinct behaviors of neural stem and progenitor cells underlie cortical neurogenesis. *J Comp Neurol*, 508(1), 28-44.

Noles, SR, & Chenn, A. (2007). Cadherin inhibition of beta-catenin signaling regulates the proliferation and differentiation of neural precursor cells. *Mol Cell Neurosci*, 35(4), 549-558.

Norden, DM, & Godbout, JP. (2013). Review: microglia of the aged brain: primed to be activated and resistant to regulation. *Neuropathol Appl Neurobiol*, 39(1), 19-34.

Norden, DM, Muccigrosso, MM, & Godbout, JP. (2015). Microglial priming and enhanced reactivity to secondary insult in aging, and traumatic CNS injury, and neurodegenerative disease. *Neuropharmacology*, 96(Pt A), 29-41.

Ohgidani, M, Kato, TA, Setoyama, D, Sagata, N, Hashimoto, R, Shigenobu, K, Yoshida, T, Hayakawa, K, Shimokawa, N, Miura, D, Utsumi, H, & Kanba, S. (2014). Direct induction of ramified microglia-like cells from human monocytes: dynamic microglial dysfunction in Nasu-Hakola disease. *Sci Rep*, 4, 4957.

Oliveira, AL, Thams, S, Lidman, O, Piehl, F, Hokfelt, T, Karre, K, Linda, H, & Cullheim, S. (2004). A role for MHC class I molecules in synaptic plasticity and regeneration of neurons after axotomy. *Proc Natl Acad Sci U S A*, 101(51), 17843-17848.

Oppenheim, RW. (1991). Cell death during development of the nervous system. *Annu Rev Neurosci*, 14, 453-501.

Panteleev, AV, & Vorob'ev, IA. (2012). [Expression of early hematopoietic markers in cord blood and mobilized blood]. *Tsitologija*, 54(10), 774-782.

Paolicelli, RC, Bolasco, G, Pagani, F, Maggi, L, Scianni, M, Panzanelli, P, Giustetto, M, Ferreira, TA, Guiducci, E, Dumas, L, Ragozzino, D, & Gross, CT. (2011). Synaptic pruning by microglia is necessary for normal brain development. *Science*, 333(6048), 1456-1458.

- Park, D, Xiang, AP, Zhang, L, Mao, FF, Walton, NM, Choi, SS, & Lahn, BT. (2009). The radial glia antibody RC2 recognizes a protein encoded by Nestin. *Biochem Biophys Res Commun*, 382(3), 588-592.
- Park, D, Xiang, AP, Mao, FF, Zhang, L, Di, CG, Liu, XM, Shao, Y, Ma, BF, Lee, JH, Ha, KS, Walton, N, & Lahn, BT. (2010). Nestin is required for the proper self-renewal of neural stem cells. *Stem Cells*, 28(12), 2162-2171.
- Parkhurst, CN, Yang, G, Ninan, I, Savas, JN, Yates, JR, 3rd, Lafaille, JJ, Hempstead, BL, Littman, DR, & Gan, WB. (2013). Microglia promote learning-dependent synapse formation through brain-derived neurotrophic factor. *Cell*, 155(7), 1596-1609.
- Pascual, O, Ben Achour, S, Rostaing, P, Triller, A, & Bessis, A. (2012). Microglia activation triggers astrocyte-mediated modulation of excitatory neurotransmission. *Proc Natl Acad Sci U S A*, 109(4), E197-205.
- Paterson, JA, Privat, A, Ling, EA, & Leblond, CP. (1973). Investigation of glial cells in semithin sections. 3. Transformation of subependymal cells into glial cells, as shown by radioautography after 3 H-thymidine injection into the lateral ventricle of the brain of young rats. *J Comp Neurol*, 149(1), 83-102.
- Perez-Pouchoulen, M, VanRyzin, JW, & McCarthy, MM. (2015). Morphological and Phagocytic Profile of Microglia in the Developing Rat Cerebellum(1,2,3). *eNeuro*, 2(4).
- Perry, VH, Hume, DA, & Gordon, S. (1985). Immunohistochemical localization of macrophages and microglia in the adult and developing mouse brain. *Neuroscience*, 15(2), 313-326.
- Perry, VH, & Gordon, S. (1991). Macrophages and the nervous system. *Int Rev Cytol*, 125, 203-244.
- Perry, VH, & Holmes, C. (2014). Microglial priming in neurodegenerative disease. *Nat Rev Neurol*, 10(4), 217-224.
- Pevny, LH, Sockanathan, S, Placzek, M, & Lovell-Badge, R. (1998). A role for SOX1 in neural determination. *Development*, 125(10), 1967-1978.
- Pfenninger, CV, Roschupkina, T, Hertwig, F, Kottwitz, D, Englund, E, Bengzon, J, Jacobsen, SE, & Nuber, UA. (2007). CD133 is not present on neurogenic astrocytes in the adult subventricular zone, but on embryonic neural stem cells, ependymal cells, and glioblastoma cells. *Cancer Res*, 67(12), 5727-5736.



- Pocock, JM, & Kettenmann, H. (2007). Neurotransmitter receptors on microglia. *Trends Neurosci*, 30(10), 527-535.
- Pollard, SM, & Conti, L. (2007). Investigating radial glia in vitro. *Prog Neurobiol*, 83(1), 53-67.
- Pöschl, J, Grammel, D, Dorostkar, MM, Kretzschmar, HA, & Schüller, U. (2013). Constitutive activation of  $\beta$ -Catenin in neural progenitors results in disrupted proliferation and migration of neurons within the central nervous system. *Dev Biol*, 374(2), 319-332.
- Pribiag, H, & Stellwagen, D. (2013). TNF-alpha downregulates inhibitory neurotransmission through protein phosphatase 1-dependent trafficking of GABA(A) receptors. *J Neurosci*, 33(40), 15879-15893.
- Pribiag, H, & Stellwagen, D. (2014). Neuroimmune regulation of homeostatic synaptic plasticity. *Neuropharmacology*, 78, 13-22.
- Qin, H, Wilson, CA, Lee, SJ, Zhao, X, & Benveniste, EN. (2005). LPS induces CD40 gene expression through the activation of NF-kappaB and STAT-1alpha in macrophages and microglia. *Blood*, 106(9), 3114-3122.
- Qin, L, Wu, X, Block, ML, Liu, Y, Breese, GR, Hong, JS, Knapp, DJ, & Crews, FT. (2007). Systemic LPS causes chronic neuroinflammation and progressive neurodegeneration. *Glia*, 55(5), 453-462.
- Rakic, P. (1995). A small step for the cell, a giant leap for mankind: a hypothesis of neocortical expansion during evolution. *Trends Neurosci*, 18(9), 383-388.
- Ransohoff, RM, & Perry, VH. (2009). Microglial physiology: unique stimuli, specialized responses. *Annu Rev Immunol*, 27, 119-145.
- Rao, JS, Harry, GJ, Rapoport, SI, & Kim, HW. (2010). Increased excitotoxicity and neuroinflammatory markers in postmortem frontal cortex from bipolar disorder patients. *Mol Psychiatry*, 15(4), 384-392.
- Redies, C. (2000). Cadherins in the central nervous system. *Prog Neurobiol*, 61(6), 611-648.
- Redlich, S, Ribes, S, Schutze, S, Eiffert, H, & Nau, R. (2013). Toll-like receptor stimulation increases phagocytosis of Cryptococcus neoformans by microglial cells. *J Neuroinflammation*, 10, 71.
- Redmer, T, Diecke, S, Grigoryan, T, Quiroga-Negreira, A, Birchmeier, W, & Besser, D. (2011). E-cadherin is crucial for embryonic stem cell

pluripotency and can replace OCT4 during somatic cell reprogramming. *EMBO Rep*, 12(7), 720-726.

Rezaie, P, Dean, A, Male, D, & Ulfing, N. (2005). Microglia in the cerebral wall of the human telencephalon at second trimester. *Cereb Cortex*, 15(7), 938-949.

Ribes, S, Ebert, S, Regen, T, Agarwal, A, Tauber, SC, Czesnik, D, Spreer, A, Bunkowski, S, Eiffert, H, Hanisch, UK, Hammerschmidt, S, & Nau, R. (2010). Toll-like receptor stimulation enhances phagocytosis and intracellular killing of nonencapsulated and encapsulated *Streptococcus pneumoniae* by murine microglia. *Infect Immun*, 78(2), 865-871.

Rizo, J, & Xu, J. (2015). The Synaptic Vesicle Release Machinery. *Annu Rev Biophys*, 44, 339-367.

Rogers, JT, Morganti, JM, Bachstetter, AD, Hudson, CE, Peters, MM, Grimmig, BA, Weeber, EJ, Bickford, PC, & Gemma, C. (2011). CX3CR1 deficiency leads to impairment of hippocampal cognitive function and synaptic plasticity. *J Neurosci*, 31(45), 16241-16250.

Ross, SE, Greenberg, ME, & Stiles, CD. (2003). Basic helix-loop-helix factors in cortical development. *Neuron*, 39(1), 13-25.

Roumier, A, Bechade, C, Poncer, JC, Smalla, KH, Tomasello, E, Vivier, E, Gundelfinger, ED, Triller, A, & Bessis, A. (2004). Impaired synaptic function in the microglial KARAP/DAP12-deficient mouse. *J Neurosci*, 24(50), 11421-11428.

Roumier, A, Pascual, O, Bechade, C, Wakselman, S, Poncer, JC, Real, E, Triller, A, & Bessis, A. (2008). Prenatal activation of microglia induces delayed impairment of glutamatergic synaptic function. *PLoS One*, 3(7), e2595.

Rouso, DL, Pearson, CA, Gaber, ZB, Miquelajauregui, A, Li, S, Portera-Cailliau, C, Morrisey, EE, & Novitsch, BG. (2012). Foxp-mediated suppression of N-cadherin regulates neuroepithelial character and progenitor maintenance in the CNS. *Neuron*, 74(2), 314-330.

Saijo, K, & Glass, CK. (2011). Microglial cell origin and phenotypes in health and disease. *Nat Rev Immunol*, 11(11), 775-787.

Sakane, F, & Miyamoto, Y. (2013). N-cadherin regulates the proliferation and differentiation of ventral midbrain dopaminergic progenitors. *Dev Neurobiol*, 73(7), 518-529.

- Samokhvalov, IM, Samokhvalova, NI, & Nishikawa, S. (2007). Cell tracing shows the contribution of the yolk sac to adult haematopoiesis. *Nature*, 446(7139), 1056-1061.
- Sancho-Tello, M, Valles, S, Montoliu, C, Renau-Piqueras, J, & Guerri, C. (1995). Developmental pattern of GFAP and vimentin gene expression in rat brain and in radial glial cultures. *Glia*, 15(2), 157-166.
- Sansom, SN, Griffiths, DS, Faedo, A, Kleinjan, D-J, Ruan, Y, Smith, J, van Heyningen, V, Rubenstein, JL, & Livesey, FJ. (2009). The Level of the Transcription Factor Pax6 Is Essential for Controlling the Balance between Neural Stem Cell Self-Renewal and Neurogenesis. *PLOS Genetics*, 5(6), e1000511.
- Sarnat, HB. (2013). Clinical neuropathology practice guide 5-2013: markers of neuronal maturation. *Clin Neuropathol*, 32(5), 340-369.
- Sato, K. (2015). Effects of Microglia on Neurogenesis. *Glia*, 63(8), 1394-1405.
- Schafer, DP, Lehrman, EK, Kautzman, AG, Koyama, R, Mardinly, AR, Yamasaki, R, Ransohoff, RM, Greenberg, ME, Barres, BA, & Stevens, B. (2012). Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron*, 74(4), 691-705.
- Schlett, K, & Madarasz, E. (1997). Retinoic acid induced neural differentiation in a neuroectodermal cell line immortalized by p53 deficiency. *J Neurosci Res*, 47(4), 405-415.
- Schlett, K, Czirok, A, Tarnok, K, Vicsek, T, & Madarasz, E. (2000). Dynamics of cell aggregation during in vitro neurogenesis by immortalized neuroectodermal progenitors. *J Neurosci Res*, 60(2), 184-194.
- Schnitzer, J, Franke, WW, & Schachner, M. (1981). Immunocytochemical demonstration of vimentin in astrocytes and ependymal cells of developing and adult mouse nervous system. *J Cell Biol*, 90(2), 435-447.
- Schulz, C, Gomez Perdiguero, E, Chorro, L, Szabo-Rogers, H, Cagnard, N, Kierdorf, K, Prinz, M, Wu, B, Jacobsen, SE, Pollard, JW, Frampton, J, Liu, KJ, & Geissmann, F. (2012). A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science*, 336(6077), 86-90.
- Sessa, A, Mao, CA, Hadjantonakis, AK, Klein, WH, & Broccoli, V. (2008). Tbr2 directs conversion of radial glia into basal precursors and guides neuronal amplification by indirect neurogenesis in the developing neocortex. *Neuron*, 60(1), 56-69.

- Shen, MD, Nordahl, CW, Young, GS, Wootton-Gorges, SL, Lee, A, Liston, SE, Harrington, KR, Ozonoff, S, & Amaral, DG. (2013). Early brain enlargement and elevated extra-axial fluid in infants who develop autism spectrum disorder. *Brain*, 136(Pt 9), 2825-2835.
- Shi, L, Fatemi, SH, Sidwell, RW, & Patterson, PH. (2003). Maternal influenza infection causes marked behavioral and pharmacological changes in the offspring. *J Neurosci*, 23(1), 297-302.
- Shibata, T, Yamada, K, Watanabe, M, Ikenaka, K, Wada, K, Tanaka, K, & Inoue, Y. (1997). Glutamate transporter GLAST is expressed in the radial glia-astrocyte lineage of developing mouse spinal cord. *J Neurosci*, 17(23), 9212-9219.
- Shigemoto-Mogami, Y, Hoshikawa, K, Goldman, JE, Sekino, Y, & Sato, K. (2014). Microglia enhance neurogenesis and oligodendrogenesis in the early postnatal subventricular zone. *J Neurosci*, 34(6), 2231-2243.
- Sierra, A, Encinas, JM, Deudero, JJ, Chancey, JH, Enikolopov, G, Overstreet-Wadiche, LS, Tsirka, SE, & Maletic-Savatic, M. (2010). Microglia shape adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell Stem Cell*, 7(4), 483-495.
- Simard, AR, & Rivest, S. (2004). Bone marrow stem cells have the ability to populate the entire central nervous system into fully differentiated parenchymal microglia. *FASEB J*, 18(9), 998-1000.
- Smith, SE, Li, J, Garbett, K, Mirnics, K, & Patterson, PH. (2007). Maternal immune activation alters fetal brain development through interleukin-6. *J Neurosci*, 27(40), 10695-10702.
- Soncin, F, & Ward, CM. (2011). The function of e-cadherin in stem cell pluripotency and self-renewal. *Genes (Basel)*, 2(1), 229-259.
- Sparks, BF, Friedman, SD, Shaw, DW, Aylward, EH, Echelard, D, Artru, AA, Maravilla, KR, Giedd, JN, Munson, J, Dawson, G, & Dager, SR. (2002). Brain structural abnormalities in young children with autism spectrum disorder. *Neurology*, 59(2), 184-192.
- Squarzoni, P, Oller, G, Hoeffel, G, Pont-Lezica, L, Rostaing, P, Low, D, Bessis, A, Ginhoux, F, & Garel, S. (2014). Microglia modulate wiring of the embryonic forebrain. *Cell Rep*, 8(5), 1271-1279.
- Steinmetz, CC, & Turrigiano, GG. (2010). Tumor necrosis factor-alpha signaling maintains the ability of cortical synapses to express synaptic scaling. *J Neurosci*, 30(44), 14685-14690.

- Stellwagen, D, Beattie, EC, Seo, JY, & Malenka, RC. (2005). Differential regulation of AMPA receptor and GABA receptor trafficking by tumor necrosis factor-alpha. *J Neurosci*, 25(12), 3219-3228.
- Stephan, AH, Barres, BA, & Stevens, B. (2012). The complement system: an unexpected role in synaptic pruning during development and disease. *Annu Rev Neurosci*, 35, 369-389.
- Stevens, B, Allen, NJ, Vazquez, LE, Howell, GR, Christopherson, KS, Nouri, N, Micheva, KD, Mehalow, AK, Huberman, AD, Stafford, B, Sher, A, Litke, AM, Lambris, JD, Smith, SJ, John, SW, & Barres, BA. (2007). The classical complement cascade mediates CNS synapse elimination. *Cell*, 131(6), 1164-1178.
- Stocker, AM, & Chenn, A. (2015). The role of adherens junctions in the developing neocortex. *Cell Adh Migr*, 9(3), 167-174.
- Stolt, CC, Lommes, P, Sock, E, Chaboissier, MC, Schedl, A, & Wegner, M. (2003). The Sox9 transcription factor determines glial fate choice in the developing spinal cord. *Genes Dev*, 17(13), 1677-1689.
- Sultan, S, Gebara, E, & Toni, N. (2013). Doxycycline increases neurogenesis and reduces microglia in the adult hippocampus. *Front Neurosci*, 7, 131.
- Sun, T, & Hevner, RF. (2014). Growth and folding of the mammalian cerebral cortex: from molecules to malformations. *Nat Rev Neurosci*, 15(4), 217-232.
- Suter, DM, Tirefort, D, Julien, S, & Krause, KH. (2009). A Sox1 to Pax6 switch drives neuroectoderm to radial glia progression during differentiation of mouse embryonic stem cells. *Stem Cells*, 27(1), 49-58.
- Suzuki, K, Sugihara, G, Ouchi, Y, Nakamura, K, Futatsubashi, M, Takebayashi, K, Yoshihara, Y, Omata, K, Matsumoto, K, Tsuchiya, KJ, Iwata, Y, Tsujii, M, Sugiyama, T, & Mori, N. (2013). Microglial activation in young adults with autism spectrum disorder. *JAMA Psychiatry*, 70(1), 49-58.
- Suzuki, SC, & Takeichi, M. (2008). Cadherins in neuronal morphogenesis and function. *Dev Growth Differ*, 50 Suppl 1, S119-130.
- Swinnen, N, Smolders, S, Avila, A, Notelaers, K, Paesen, R, Ameloot, M, Brone, B, Legendre, P, & Rigo, JM. (2013). Complex invasion pattern of the cerebral cortex by microglial cells during development of the mouse embryo. *Glia*, 61(2), 150-163.

- Takahashi, K, Yamamura, F, & Naito, M. (1989). Differentiation, maturation, and proliferation of macrophages in the mouse yolk sac: a light-microscopic, enzyme-cytochemical, immunohistochemical, and ultrastructural study. *J Leukoc Biol*, 45(2), 87-96.
- Takahashi, T, Nowakowski, RS, & Caviness, VS, Jr. (1995). Early ontogeny of the secondary proliferative population of the embryonic murine cerebral wall. *J Neurosci*, 15(9), 6058-6068.
- Takeichi, M. (1995). Morphogenetic roles of classic cadherins. *Curr Opin Cell Biol*, 7(5), 619-627.
- Takeichi, M. (2007). The cadherin superfamily in neuronal connections and interactions. *Nat Rev Neurosci*, 8(1), 11-20.
- Tarnok, K, Pataki, A, Kovacs, J, Schlett, K, & Madarasz, E. (2002). Stage-dependent effects of cell-to-cell connections on in vitro induced neurogenesis. *Eur J Cell Biol*, 81(7), 403-412.
- Taverna, E, Gotz, M, & Huttner, WB. (2014). The cell biology of neurogenesis: toward an understanding of the development and evolution of the neocortex. *Annu Rev Cell Dev Biol*, 30, 465-502.
- Tay, TL, Mai, D, Dautzenberg, J, Fernandez-Klett, F, Lin, G, Sagar, Datta, M, Drougard, A, Stempfl, T, Ardura-Fabregat, A, Staszewski, O, Margineanu, A, Sporbert, A, Steinmetz, LM, Pospisilik, JA, Jung, S, Priller, J, Grun, D, Ronneberger, O, & Prinz, M. (2017). A new fate mapping system reveals context-dependent random or clonal expansion of microglia. *Nat Neurosci*, 20(6), 793-803.
- Tole, S, Kaprielian, Z, Ou, SK, & Patterson, PH. (1995). FORSE-1: a positionally regulated epitope in the developing rat central nervous system. *J Neurosci*, 15(2), 957-969.
- Tondreau, T, Meuleman, N, Delforge, A, Dejeneffe, M, Leroy, R, Massy, M, Mortier, C, Bron, D, & Lagneaux, L. (2005). Mesenchymal stem cells derived from CD133-positive cells in mobilized peripheral blood and cord blood: proliferation, Oct4 expression, and plasticity. *Stem Cells*, 23(8), 1105-1112.
- Tong, CK, & Vidyadaran, S. (2016). Role of microglia in embryonic neurogenesis. *Exp Biol Med (Maywood)*, 241(15), 1669-1675.
- Torres-Platas, SG, Cruceanu, C, Chen, GG, Turecki, G, & Mechawar, N. (2014). Evidence for increased microglial priming and macrophage recruitment in the dorsal anterior cingulate white matter of depressed suicides. *Brain Behav Immun*, 42, 50-59.

- Town, T, Nikolic, V, & Tan, J. (2005). The microglial "activation" continuum: from innate to adaptive responses. *J Neuroinflammation*, 2, 24.
- Tremblay, ME, Lowery, RL, & Majewska, AK. (2010). Microglial interactions with synapses are modulated by visual experience. *PLoS Biol*, 8(11), e1000527.
- Tremblay, ME, & Majewska, AK. (2011). A role for microglia in synaptic plasticity? *Commun Integr Biol*, 4(2), 220-222.
- Tronnes, AA, Koschnitzky, J, Daza, R, Hitti, J, Ramirez, JM, & Hevner, R. (2015). Effects of Lipopolysaccharide and Progesterone Exposures on Embryonic Cerebral Cortex Development in Mice. *Reprod Sci*.
- Ueno, M, Fujita, Y, Tanaka, T, Nakamura, Y, Kikuta, J, Ishii, M, & Yamashita, T. (2013). Layer V cortical neurons require microglial support for survival during postnatal development. *Nat Neurosci*, 16(5), 543-551.
- Unverzagt, KL, Bender, JG, Loudovaris, M, Martinson, JA, Hazelton, B, & Weaver, C. (1997). Characterization of a culture-derived CD15+CD11b- promyelocytic population from CD34+ peripheral blood cells. *J Leukoc Biol*, 62(4), 480-484.
- Vainchtein, ID, Vinet, J, Brouwer, N, Brendecke, S, Biagini, G, Biber, K, Boddeke, HW, & Eggen, BJ. (2014). In acute experimental autoimmune encephalomyelitis, infiltrating macrophages are immune activated, whereas microglia remain immune suppressed. *Glia*, 62(10), 1724-1735.
- Varga, BV, Hadinger, N, Gocza, E, Dulberg, V, Demeter, K, Madarasz, E, & Herberth, B. (2008). Generation of diverse neuronal subtypes in cloned populations of stem-like cells. *BMC Dev Biol*, 8, 89.
- Varvel, NH, Grathwohl, SA, Baumann, F, Liebig, C, Bosch, A, Brawek, B, Thal, DR, Charo, IF, Heppner, FL, Aguzzi, A, Garaschuk, O, Ransohoff, RM, & Jucker, M. (2012). Microglial repopulation model reveals a robust homeostatic process for replacing CNS myeloid cells. *Proc Natl Acad Sci U S A*, 109(44), 18150-18155.
- Vinet, J, Weering, HR, Heinrich, A, Kalin, RE, Wegner, A, Brouwer, N, Heppner, FL, Rooijen, N, Boddeke, HW, & Biber, K. (2012). Neuroprotective function for ramified microglia in hippocampal excitotoxicity. *J Neuroinflammation*, 9, 27.
- Vives, V, Alonso, G, Solal, AC, Joubert, D, & Legraverend, C. (2003). Visualization of S100B-positive neurons and glia in the central nervous system of EGFP transgenic mice. *J Comp Neurol*, 457(4), 404-419.

- Wake, H, Moorhouse, AJ, Jinno, S, Kohsaka, S, & Nabekura, J. (2009). Resting microglia directly monitor the functional state of synapses in vivo and determine the fate of ischemic terminals. *J Neurosci*, 29(13), 3974-3980.
- Wakselman, S, Bechade, C, Roumier, A, Bernard, D, Triller, A, & Bessis, A. (2008). Developmental neuronal death in hippocampus requires the microglial CD11b integrin and DAP12 immunoreceptor. *J Neurosci*, 28(32), 8138-8143.
- Walton, NM, Sutter, BM, Laywell, ED, Levkoff, LH, Kearns, SM, Marshall, GP, 2nd, Scheffler, B, & Steindler, DA. (2006). Microglia instruct subventricular zone neurogenesis. *Glia*, 54(8), 815-825.
- Wei, J, Gabrusiewicz, K, & Heimberger, A. (2013). The controversial role of microglia in malignant gliomas. *Clin Dev Immunol*, 2013, 285246.
- Weigmann, A, Corbeil, D, Hellwig, A, & Huttner, WB. (1997). Prominin, a novel microvilli-specific polytopic membrane protein of the apical surface of epithelial cells, is targeted to plasmalemmal protrusions of non-epithelial cells. *Proc Natl Acad Sci U S A*, 94(23), 12425-12430.
- Wiese, C, Rolletschek, A, Kania, G, Blyszczuk, P, Tarasov, KV, Tarasova, Y, Wersto, RP, Boheler, KR, & Wobus, AM. (2004). Nestin expression--a property of multi-lineage progenitor cells? *Cell Mol Life Sci*, 61(19-20), 2510-2522.
- Wodarz, A, & Huttner, WB. (2003). Asymmetric cell division during neurogenesis in *Drosophila* and vertebrates. *Mech Dev*, 120(11), 1297-1309.
- Wolf, SA, Boddeke, HW, & Kettenmann, H. (2017). Microglia in Physiology and Disease. *Annu Rev Physiol*, 79, 619-643.
- Wong, GK, Baudet, ML, Norden, C, Leung, L, & Harris, WA. (2012). Slit1b-Robo3 signaling and N-cadherin regulate apical process retraction in developing retinal ganglion cells. *J Neurosci*, 32(1), 223-228.
- Wong, WT. (2013). Microglial aging in the healthy CNS: phenotypes, drivers, and rejuvenation. *Front Cell Neurosci*, 7, 22.
- Woodhead, GJ, Mutch, CA, Olson, EC, & Chenn, A. (2006). Cell-Autonomous  $\beta$ -Catenin Signaling Regulates Cortical Precursor Proliferation. *The Journal of Neuroscience*, 26(48), 12620-12630.
- Wu, S, Wu, Y, & Capecchi, MR. (2006). Motoneurons and oligodendrocytes are sequentially generated from neural stem cells but do not appear to



share common lineage-restricted progenitors in vivo. *Development*, 133(4), 581-590.

- Wu, SX, Goebbels, S, Nakamura, K, Kometani, K, Minato, N, Kaneko, T, Nave, KA, & Tamamaki, N. (2005). Pyramidal neurons of upper cortical layers generated by NEX-positive progenitor cells in the subventricular zone. *Proc Natl Acad Sci U S A*, 102(47), 17172-17177.
- Xavier, AL, Lima, FR, Nedergaard, M, & Menezes, JR. (2015). Ontogeny of CX3CR1-EGFP expressing cells unveil microglia as an integral component of the postnatal subventricular zone. *Front Cell Neurosci*, 9, 37.
- Xi, R. (2009). Anchoring stem cells in the niche by cell adhesion molecules. *Cell Adh Migr*, 3(4), 396-401.
- Yamada, J, Hayashi, Y, Jinno, S, Wu, Z, Inoue, K, Kohsaka, S, & Nakanishi, H. (2008). Reduced synaptic activity precedes synaptic stripping in vagal motoneurons after axotomy. *Glia*, 56(13), 1448-1462.
- Yao, L, Kan, EM, Lu, J, Hao, A, Dheen, ST, Kaur, C, & Ling, EA. (2013). Toll-like receptor 4 mediates microglial activation and production of inflammatory mediators in neonatal rat brain following hypoxia: role of TLR4 in hypoxic microglia. *J Neuroinflammation*, 10, 23.
- Yokota, Y, Eom, TY, Stanco, A, Kim, WY, Rao, S, Snider, WD, & Anton, ES. (2010). Cdc42 and Gsk3 modulate the dynamics of radial glial growth, inter-radial glial interactions and polarity in the developing cerebral cortex. *Development*, 137(23), 4101-4110.
- Yung, SY, Gokhan, S, Jurcsak, J, Molero, AE, Abrajano, JJ, & Mehler, MF. (2002). Differential modulation of BMP signaling promotes the elaboration of cerebral cortical GABAergic neurons or oligodendrocytes from a common sonic hedgehog-responsive ventral forebrain progenitor species. *Proc Natl Acad Sci U S A*, 99(25), 16273-16278.
- Zechner, D, Fujita, Y, Hulsken, J, Muller, T, Walther, I, Taketo, MM, Crenshaw, EB, 3rd, Birchmeier, W, & Birchmeier, C. (2003). beta-Catenin signals regulate cell growth and the balance between progenitor cell expansion and differentiation in the nervous system. *Dev Biol*, 258(2), 406-418.
- Zhan, Y, Paolicelli, RC, Sforzini, F, Weinhard, L, Bolasco, G, Pagani, F, Vyssotski, AL, Bifone, A, Gozzi, A, Ragozzino, D, & Gross, CT. (2014). Deficient neuron-microglia signaling results in impaired functional brain connectivity and social behavior. *Nat Neurosci*, 17(3), 400-406.

- Zhang, GX, Li, J, Ventura, E, & Rostami, A. (2002). Parenchymal microglia of naive adult C57BL/6J mice express high levels of B7.1, B7.2, and MHC class II. *Exp Mol Pathol*, 73(1), 35-45.
- Zhang, J, Woodhead, GJ, Swaminathan, SK, Noles, SR, McQuinn, ER, Pisarek, AJ, Stocker, AM, Mutch, CA, Funatsu, N, & Chenn, A. (2010). Cortical neural precursors inhibit their own differentiation via N-cadherin maintenance of beta-catenin signaling. *Dev Cell*, 18(3), 472-479.
- Zhang, J, Shemezis, JR, McQuinn, ER, Wang, J, Sverdlov, M, & Chenn, A. (2013). AKT activation by N-cadherin regulates beta-catenin signaling and neuronal differentiation during cortical development. *Neural Dev*, 8, 7.
- Zhu, F, Zheng, Y, Liu, Y, Zhang, X, & Zhao, J. (2014). Minocycline alleviates behavioral deficits and inhibits microglial activation in the offspring of pregnant mice after administration of polyriboinosinic-polyribocytidilic acid. *Psychiatry Res*, 219(3), 680-686.
- Zoghbi, HY, & Bear, MF. (2012). Synaptic dysfunction in neurodevelopmental disorders associated with autism and intellectual disabilities. *Cold Spring Harb Perspect Biol*, 4(3).

## LIST OF PUBLICATIONS

**Chih Kong Tong**, Sharmili Vidyadaran: Role of microglia in embryonic neurogenesis. *Experimental Biology and Medicine* 08/2016; 241(15). DOI:10.1177/1535370216664430

**Impact Factor: 2.542**

**Chih Kong Tong**, Md Shariful Hasan Sumon, Sharmili Vidyadaran, Norshariza Nordin: Functions of microglial cells on neurogenesis, synaptic pruning and neurotransmission: a brief review. *Regenerative Research* 08/2015; 4(1).

Shalini Vellasamy, **Chih Kong Tong**, Nur Atiqah Azhar, Radha Kodiappan, Soon Choy Chan, Abhi Veerakumarasivam, Rajesh Ramasamy: Human mesenchymal stromal cells modulate T-cell immune response via transcriptomic regulation. *Cytotherapy* 08/2016; 18(10). DOI:10.1016/j.jcyt.2016.06.017

**Impact factor: 3.625**

Shinsmon Jose, Shi Wei Tan, **Chih Kong Tong**, Sharmili Vidyadaran: Isolation and characterization of primary microglia from post-natal murine brain tissues: A comparison of two methods. *Cell Biology International* 07/2015; 39(12). DOI:10.1002/cbin.10516

**Impact Factor: 1.663**

Randy Tatt Yew Haw, **Chih Kong Tong**, Andrew Yew, Han Chung Lee, James B Phillips, Sharmili Vidyadaran: A three-dimensional collagen construct to model lipopolysaccharide-induced activation of BV2 microglia. *Journal of Neuroinflammation* 07/2014; 11(1). DOI:10.1186/1742-2094-11-134

**Impact Factor: 4.667** (contributed equally as first author)



**UNIVERSITI PUTRA MALAYSIA**

**STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT**

**ACADEMIC SESSION :** \_\_\_\_\_

**TITLE OF THESIS / PROJECT REPORT :**

*In Vitro* MODULATORY EFFECTS OF MICROGLIA ON NEUROEPITHELIAL CELL  
NEURODIFFERENTIATION

**NAME OF STUDENT:** TONG CHIH KONG

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

\*Please tick (✓)

**CONFIDENTIAL**

(Contain confidential information under Official Secret Act 1972).

**RESTRICTED**

(Contains restricted information as specified by the organization/institution where research was done).

**OPEN ACCESS**

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

**PATENT**

Embargo from \_\_\_\_\_ until \_\_\_\_\_  
(date) (date)

**Approved by:**

\_\_\_\_\_  
(Signature of Student)  
New IC No/ Passport No.:

Date :

\_\_\_\_\_  
(Signature of Chairman of Supervisory Committee)  
Name:

Date :

**[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted. ]**