



UNIVERSITI PUTRA MALAYSIA

***MOLECULAR MODELLING OF BERBERINE DERIVATIVES AS
INHIBITORS OF ONCOGENIC SIGNALLING PATHWAYS IN BREAST
CANCER CELL LINES***

PARHAM JABBARZADEH KABOLI

FPSK(p) 2018 13



**MOLECULAR MODELLING OF BERBERINE DERIVATIVES AS
INHIBITORS OF ONCOGENIC SIGNALLING PATHWAYS IN BREAST
CANCER CELL LINES**

By

PARHAM JABBARZADEH KABOLI

**Thesis Submitted to the School of Graduate Studies,
Universiti Putra Malaysia, in Fulfillment of the Requirements for
the Degree of Doctor of Philosophy**

December 2017

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATIONS

This thesis is dedicated to:

My late father, R.I.P., the best teacher I have ever had in my life who never stopped giving of himself in countless ways,

My dearest mother, the best lover in my life who gives me endless love, hopes and supports,

My sisters, Mahnoosh, Hanieh, and Sanaz, who have given me their genuine kindness,

My brothers-in-law, Mohammad and Ali Reza, who empower the family with their hope and energy,

My cute nephews, Aria and Pouya, smart and lovely, and the next generation of gentlemen,

My sweet niece, Parmis, a beautiful princess,

My late grandparents and all people gave me love and happiness,

and

You, as the valuable reader with the very best wishes in your academic journey.

Abstract of thesis presented to the Senate of Universiti Putra Malaysia
in fulfilment of the requirement for the degree of Doctor of Philosophy

**MOLECULAR MODELLING OF BERBERINE DERIVATIVES AS
INHIBITORS OF ONCOGENIC SIGNALLING PATHWAYS IN BREAST
CANCER CELL LINES**

By

PARHAM JABBARZADEH KABOLI

December 2017

Chair: Professor Patimah Binti Ismail, PhD
Faculty: Medicine and Health Sciences

Berberine (BBR) is an alkaloid that is widely distributed in different plant species. Several studies have been carried out on the anti-cancer effects of BBR but direct targets of BBR are unknown. In the development of approximately 20-25% of all cancers, altered hedgehog (Hh) signalling is involved where the smoothened (Smo) transmembrane receptor triggers Hh signalling pathway towards *Gli1* gene expression. Besides Smo receptor, BRAF mutations have been also detected in 7% of all cancers and 66% of melanomas; as such, the FDA has approved a few BRAF inhibitor drugs to date. However, BRAF can activate CRAF leading to resistance to BRAF inhibitors. The aim of this study is to model and compare the effects of BBR against key proteins involved in Hedgehog, MAP kinase, and PI3K pathways using *in silico* and *in vitro* approaches. BBR is found to interact with Lys395 of Smo receptor via hydrogen bonding and cation- π interactions [Score= -8.72 kcal/mol (Smo)]. In addition, π - π interactions between BBR aromatic rings and two aromatic residues in the Smo transmembrane domain, Tyr394 and Phe484, are noted. Target specific binding efficiency indices using an *in-silico* approach are calculated to plot the Smo-specific binding potency of each ligand. In addition, interactions of BBR derivatives (total number= 485 derivatives) against BRAF and CRAF kinases are modelled and predicted using an *in silico*-based approach. The Adenosine Triphosphate (ATP) is modelled in order to analyse and identify the residues important in BRAF docking. Results show Lys483 and Asp594 are the most important residues involved in both ATP and BBR binding [Scores= -11.50 kcal/mol (ATP); -8.50 kcal/mol (BBR)]. In addition to these polar residues, Trp530 and Phe583 are also applicable to the molecular docking of BRAF. The Asp593 is excluded from the enzyme cavity, while Phe594 is included inside the cavity, making the enzyme inactive. Finally, three alternatives

for BBR are identified with dual RAF inhibition effects. Direct effects of BBR derivatives against BRAF and CRAF kinases have not yet been reported previously, and thus, for the first time, three protoberberines are reported as lead compounds against RAF kinases [BBR-7 (Score= -9.75 kcal/mol), BBR-9 (Score= -9.76 kcal/mol), and BBR-10 (Score= -9.27 kcal/mol)]. In the current study, molecular docking and molecular dynamics simulations are also used for EGFR, p38 MAPK, ERK1/2 and AKT. The effects of BBR on MAP kinase and PI3K pathways are evaluated using Immuno-florescence assays and the amounts of phosphorylated kinases are compared with total kinases after they are treated with different concentrations of BBR [IC₅₀s= 7 μ M (EGFR), 45 μ M (ERK1/2), 60 μ M (AKT), 75 μ M (p38)]. BBR interacts accurately with EGFR, AKT, P38, and ERK1/2 active sites *in silico*, and decreases the level of active forms of corresponding enzymes in studied cell lines [Scores= -7.22 kcal/mol (EGFR), -7.51 kcal/mol (ERK), -7.78kcal/mol (AKT), -7.42 kcal/mol (p38)]. In addition, it is observed that BBR has toxicity effects which leads to cyto-necrotic effect at longer treatment time. This study predicts the role of BBR as an inhibitor of Smo receptor, suggesting its effectiveness in hedgehog signalling during cancer treatment. It concludes that BBR through its multi-kinase inhibitory effects may be a useful replacement for lapatinib, an EGFR inhibitor which may sometimes cause drug resistance in patients.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**MODEL MOLEKULAR DERIVATIF BERBERIN SEBAGAI
PENGHAMBAT LALUAN SIGNAL ONKOGENIK DALAM GARISAN SEL
KANSER BUAH DADA**

Oleh

PARHAM JABBARZADEH KABOLI

Disember 2017

Pengerusi: Profesor Patimah Bt. Ismail, PhD
Fakulti: Perubatan dan Kesihatan

Berberin (BBR), merupakan alkaloid yang didapati dalam spesies tumbuhan yang berlainan. Banyak kajian telah dilakukan untuk mengetahui secara mendalam kesan anti-kanser BBR namun sasaran langsung BBR tidak berjaya diketahui. Dalam 20-25% kes kanser, signal Hh adalah terlibat di mana reseptor transmembrane Smo mengaktifkan laluan isyarat Hh ke arah ekspresi gen Gli1. Selain reseptor Smo, mutasi BRAF juga dikesan dalam 7% daripada semua kanser dan 66% kes melanoma. Oleh yang demikian, FDA telah meluluskan beberapa ubat perencat BRAF. Walau bagaimanapun, BRAF boleh mengaktifkan CRAF yang memberi rintangan terhadap perencat BRAF. Kajian ini menggunakan pendekatan *silico* dan *in vitro* untuk memodel dan membandingkan kesan BBR terhadap protein utama yang terlibat dalam Hh, MAP kinase, dan laluan PI3K. BBR didapati berinteraksi dengan Lys395 dari reseptor Smo melalui ikatan hidrogen dan interaksi kation- π [Skor= -8.72 kcal/mol (Smo)]. Di samping itu, interaksi π - π antara BBR aromatic ring dan dua residu aromatik dalam domain transmembran Smo, Tyr394 dan Phe484, telah diperhatikan. Indeks yang berdasarkan pengikat sasaran tertentu efisien menggunakan pendekatan dalam *silico* untuk memplot potensi pengikat khusus Smo bagi setiap ligan. Di samping itu, interaksi derivatif BBR (jumlah = 485 derivatif) terhadap kinas BRAF dan CRAF dimodelkan dan diramalkan menggunakan pendekatan berasaskan siliko. Untuk menganalisis dan mengenal pasti residu penting dalam dok BRAF, ATP dimodelkan dan didapati bahawa Lys483 dan Asp594 adalah residu terpenting yang terlibat dalam ATP dan pengikat BBR [Skor= -11.50 kcal/mol (ATP); -8.50 kcal/mol (BBR)]. Sebagai tambahan kepada residu polar ini, Trp530 dan Phe583 juga boleh digunakan untuk mengikat molekul

BRAF. Asp593 dikecualikan daripada rongga enzim, manakala Phe594 digunakan di dalam rongga, dan kesannya adalah enzim ini menjadi tidak aktif. Tiga alternatif untuk BBR dikenal pasti yang mempunyai kesan penghambatan RAF [BBR-7 (Skor= -9.75 kcal/mol), BBR-9 (Skor= -9.76 kcal/mol), and BBR-10 (Skor= -9.27 kcal/mol)]. Kesan langsung derivatif BBR terhadap kinase BRAF dan CRAF belum dilaporkan sebelum ini, dan oleh itu, untuk pertama kalinya, tiga protoberberin dilaporkan sebagai sebatian utama terhadap kinas RAF. Dalam kajian ini, simulasi dok molekul dan molekul dinamik juga digunakan untuk EGFR, p38 MAPK, ERK1 / 2 dan AKT. Kesan BBR pada laluan kinas MAP dan PI3K dinilai menggunakan ujian Immuno-florescence dan jumlah fosforilasi kinas dibandingkan dengan jumlah kinas selepas rawatan menggunakan kepekatan BBR yang berlainan [IC_{50} = 7 μ M (EGFR), 45 μ M (ERK1/2), 60 μ M (AKT), 75 μ M (p38)]. Interaksi BBR memberi keputusan adalah tepat dalam eksperimen siliko di EGFR, AKT, P38, dan ERK1 / 2, dan berjaya mengurangkan keaktifan enzim yang di lihat dalam garisan sel yang yang dikaji [Skor= -7.22 kcal/mol (EGFR), -7.51 kcal/mol (ERK), -7.78kcal/mol (AKT), -7.42 kcal/mol (p38)]. Di samping itu, kita perhatikan bahawa BBR mempunyai kesan toksik yang menyebabkan kesan sito-nekrotik jika jangka masa rawatan adalah lebih lama. Kajian ini meramalkan peranan BBR sebagai perencat reseptor Smo yang menunjukkan keberkesananannya dalam isyarat Hh dalam rawatan kanser. Kita boleh menyimpulkan bahawa BBR melalui kesan perencatan pelbagai-kinas adalah pengganti yang efisien untuk lapatinib, perencat EGFR yang kadang-kadang boleh menyebabkan rintangan ubat kepada pesakit.

ACKNOWLEDGEMENTS

I would like to dedicate this thesis to my mother, Mrs. Monirbanoo Tamizi, for her positive energy and financial supports during my studies.

I would appreciate Professor Patimah Ismail, Dr. Ling King-Hwa (Michael), and Associate Professor Dr. Abdah Akim who kindly supported this project as supervisory committee. Professor Dr. Asmah Rahmat is also appreciated who was the previous the chairman in supervisory committee between 2013 and 2015.

In vitro experiments of this thesis were performed at Medical Genetics Laboratory (MGL), Faculty of Medicine and Health Sciences (FMHS), UPM. Accordingly, my special appreciations are for Dr Deming Chau, the coordinator of MGL, for his kind collaborations and getting me a place at MGL. He also helped me to have access to the cell culture room, electrophoresis room, and other laboratory facilities at MGL. I strongly appreciate Professor Dr. Johnson Stanlas, Pharmacotherapeutics Laboratory, FMHS, who allowed me to have access to his laboratory instruments. Professor Dr. Johnson Stanlas kindly gifted breast cancer cell lines and some materials as well.

I would like to acknowledge Associate Professor Dr. Rajesh Ramasamy who helped me to have access to FACSCanto II machine at Laboratory of Immunology, FMHS. Professor Dr. Celerino Abad-Zapatero, University of Illinois at Chicago, kindly suggested his book written on Binding Efficiency. I would like to appreciate Professor Dr. Celerino Abad-Zapatero for his valuable comments and suggestions on my manuscripts as well.

This project was supported by Putra IPS grant and Travel Award offered by School of Graduate Studies (SGS), UPM. I also appreciate Dr Ali Etemad, postdoctoral research fellow at Laboratory of Molecular Biology and Bioinformatics, FMHS, for his helpful assistance.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Patimah Binti Ismail, PhD

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Ling King-Hwa, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Abdah Md Akim, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Parham Jabbarzadeh Kaboli, GS36929

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of Chairman of Supervisory Committee:

Patimah Binti Ismail

Signature: _____

Name of Member of Supervisory Committee:

Ling King-Hwa

Signature: _____

Name of Member of Supervisory Committee:

Abdah Md Akim

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xix
 CHAPTER	
1 INTRODUCTION	1
1.1 Brief background	1
1.2 Problem statement	3
1.3 Objectives	4
1.3.1 General objective	4
1.3.2 Specific objectives	4
1.4 Hypothesis	4
 2 LITERATURE REVIEW	5
2.1 Bio-molecular activity of berberine	5
2.1.1 Antioxidant activity of berberine	5
2.1.2 Berberine and apoptosis	6
2.1.3 Berberine and nucleic acid interaction	8
2.2 Berberine and cell signaling pathways	9
2.3 Berberine and cell proliferation	14
2.4 Berberine and gene expression	15
2.5 Berberine metabolism	17
2.6 Standard anti-cancer drugs	18
2.6.1 Smo inhibitors	18
2.6.2 RAF inhibitors	19
2.6.3 EGFR inhibitors	19
2.7 <i>In silico</i> modelling	20
 3 HEDGEHOG SIGNALLING: MODELLING	24
3.1 Introduction	24
3.2 Materials and Methods	25
3.2.1 Virtual High-Throughput Screening	25
3.2.2 Ligand selection	27
3.2.3 Crystals of Smo receptor	27
3.2.4 Grid box determination for docking	27
3.2.5 MD simulations	29
3.2.6 Molecular visualization	29
3.3 Results	31

3.3.1	Docking information	31
3.3.2	Smo-Ligand interactions	31
3.3.3	Binding efficiency metrics	37
3.3.4	MD simulations and validation	39
3.4	Discussion	42
3.5	Conclusion	43
4	ANTI-KINASE ACTIVITIES: MODELLING	44
4.1	Introduction	44
4.2	Materials and Methods	45
4.2.1	Ligand selection and molecular docking	45
4.2.2	Ligand topology and MD simulations	46
4.2.3	Molecular visualization	47
4.3	Results	47
4.3.1	Molecular docking of BBR and BRAF	47
4.3.2	BRAF Molecular interactions	50
4.3.3	Molecular docking of BBR-9 on CRAF	53
4.3.4	MD analysis of BRAF/BBR derivatives	53
4.3.5	BRAF Binding efficiency of BBR derivatives	55
4.3.6	MAPK, EGFR, ERK, and AKT	58
4.4	Discussion	65
4.5	Conclusion	67
5	MAPK/PI3K SIGNALLING: IN VITRO	68
5.1	Introduction	68
5.2	Materials and Methods	71
5.2.1	<i>In vitro</i> Material	71
5.2.2	Cytotoxicity and Apoptosis Assays	71
5.2.3	The enzyme-linked immunosorbent assay	72
5.3	Results	72
5.3.1	Microscopy	72
5.3.2	Berberine cytotoxicity	76
5.3.3	ELISA studies	84
5.4	Discussion	89
5.5	Conclusion	91
6	ADME/TOX MODELLING	92
6.1	Introduction	92
6.2	Methods	92
6.3	Results and Discussion	93
6.3.1	ADME modelling	93
6.3.2	BBR-9 and plasma proteins	94
6.3.3	BBR-9 and ABC transporters	94
6.4	Conclusion	96
7	SUMMARY, CONCLUSION, FUTURE DIRECTIONS AND RECOMMENDATIONS	98
7.1	Summary	98
7.2	Conclusion	99

7.3	Future directions	99
7.4	Recommendations	100
REFERENCES		101
APPENDICES		121
BIODATA OF STUDENT		144
LIST OF PUBLICATIONS		145



LIST OF TABLES

Table	Page
2.1 Molecules directly associated with cancers that have increased expression following berberine exposure	10
2.2 Molecules directly associated with cancers that have decreased levels in response to berberine exposure	11
3.1 Molecular properties of the top berberine (BBR) derivatives plus berberine against Smo receptor	28
3.2 RSCB Protein Data Bank (PDB) files and related information	28
3.3 Human smoothened receptor (Smo) docking information for top ranked BBR derivatives	32
3.4 Energy computations and the average number of hydrogen bonds provided	42
4.1 Physico-chemical properties of selected ligands	45
4.2 Docking information for the optimal berberine derivatives docked to kinase domains of BRAF	49
4.3 Comparative docking information for the optimal berberine derivatives docked to BRAF and CRAF	54
4.4 Energies and the average number of H-bonds computed by Gromacs	55
4.5 Different ligand efficiency indices measured for different ligands docked to BRAF	58
4.6 Docking information for berberine docked to p38 MAPK, ERK1/2, EGFR, and AKT1	60
4.7 Interaction analysis of docked berberine to ERK2, p38, EGFR, and AKT	65
5.1 Antibodies used for detection of phosphorylated and total targets	72

6.1	ADME/tox modelling for BBR and the newly generated berberine derivatives, BBR-9	93
6.2	Docking information for berberine and BBR-7, BBR-9, and BBR-10 docked to HSA and hAGP	95
6.3	Docking information for selected berberine derivatives docked to P-gp	96
B.1	PDB files obtained from RSCB protein data bank and related information	127
C.1	T-test results of cytotoxicity of berberine for MCF-7 cell line	128
C.2	T-test results of cytotoxicity of berberine for MDA-MB231 cell line	129
C.3	T-test results of the effects of berberine on total kinases for MDA-MB231 cell line	133
C.4	T-test results of the effects of berberine on total kinases for MCF-7 cell line	133
C.5	T-test results of the effects of berberine on active kinases for MDA-MB231 cell line	133
C.6	T-test results of the effects of berberine on active kinases for MCF-7 cell line	133

LIST OF FIGURES

Figure	Page
1.1 Prevalence of breast cancer and other kinds of cancer in the United States	2
1.2 Berberine structure	3
2.1 Simplified apoptotic pathway	7
2.2 Her2/neu signaling pathway	12
2.3 BRCA1 and BRCA2 function in DNA repair in breast cancer cells	16
2.4 Berberine metabolism	18
2.5 Work flow of Docking based virtual screening	21
2.6 The flowchart of the whole procedures performed in this research	23
3.1 The Hedgehog signalling pathway	25
3.2 Berberine structure and top anti-Smo ranked compounds	30
3.3 Interactions between top compounds and the neighboring residues in Smo receptor	33
3.4 The most potent inhibitors docked to smoothened receptor	34
3.5 Comparison of four different Smo crystals marked by their PDB IDs	35
3.6 Berberine/Smo interactions	36
3.7 Optimized bivariate analysis of Smo inhibitors according to binding efficiency indices and molecular properties	38
3.8 MD results for the stability of the system after docking of BBR and compound 2	40

3.9	Results computed for RDF during 10 ns MD simulations	41
4.1	Chemical structures of top anti-BRAF berberine derivatives and control	46
4.2	BBR docked to active and inactive BRAFs	51
4.3	Molecular docking of BBR-7, BBR-9, and vemurafenib against BRAF and CRAF	52
4.4	Molecular dynamics simulations of BBR-7, BBR-9, and Vemurafenib	56
4.5	Molecular dynamics simulations of BBR-9 against CRAF	57
4.6	Plots showing the relationship of each ligand to its corresponding size- and polarity-based indices docked to BRAF	58
4.7	The effects of berberine on EGFR	61
4.8	The effects of berberine on AKT	62
4.9	The effects of berberine on ERK	63
4.10	The effects of berberine on p38 MAPK	64
5.1	MAP kinase and PI3K pathways	68
5.2	Pathway interaction network of core signaling targets	70
5.3	Microscopically taken images demonstrating the effects of berberine on MCF-7 cells	74
5.4	Microscopically taken images demonstrating the effects of berberine on MDA-MB231 cells	75
5.5	Concentration-viability curve of berberine for MCF-7 over time	77
5.6	Concentration-viability curve of berberine for MDA-MB231 over time	78
5.7	Concentration-viability curve of lapatinib for MCF-7 over time	79

5.8	Concentration-viability curve of lapatinib for MDA-MB231 over time	80
5.9	Apoptosis analysis of MCF-7 and MDA-MB231	82
5.10	Comparative analysis of apoptosis in MCF-7 and MDA-MB231	83
5.11	Concentration response curve of berberine on total kinases in MCF-7 and MDA-MB231 cell lines (ELISA)	85
5.12	Concentration-response curve of berberine on phosphorylated kinases in MDA-MB231 and MCF-7 cell lines (ELISA)	86
5.13	Concentration-response curve of berberine for MDA-MB231, on normalized AKT, EGFR, p38, and ERK.	88
6.1	Interactions of BBR-9 with P-gp and HSA	97
A.1	Cartesian plots depicted the results of this project for two different targets	126
C.1	Apoptosis analysis of MCF-7 cell line after 24 hours of berberine treatment	129
C.2	Apoptosis analysis of MCF-7 cell line after 48 hours of berberine treatment	130
C.3	Apoptosis analysis of MCF-7 cell line after 72 hours of berberine treatment	130
C.4	Apoptosis analysis of MCF-7 cell line after 48 hours of lapatinib treatment	131
C.5	Apoptosis analysis of MDA-MB231 cell line after 24 hours of berberine treatment	131
C.6	Apoptosis analysis of MDA-MB231 cell line after 48 hours of berberine treatment	132
C.7	Apoptosis analysis of MDA-MB231 cell line after 72 hours of berberine treatment	132
C.8	Concentration-response curve of berberine for MDA-MB231, on total EGFR and total AKT	134

C.9	Concentration-response curve of berberine for MDA-MB231, on total and phosphorylated kinases	135
D.1	Interactions of berberine with AKT	136
D.2	Interactions of berberine with ERK	137
D.3	Root-mean-square deviation and root-mean-square fluctuation of berberine docked to ERK	138
D.4	Interactions of berberine with p38 MAPK	139
D.5	Root-mean-square deviation and root-mean-square fluctuation of berberine docked to p38 MAPK	140
D.6	Comparative pictures from the Interactions of BBR, ATP, and p38 inhibitor with p38 MAPK	141
D.7	Interactions of berberine with EGFR	142
D.8	Interactions of lapatinib with EGFR	143

LIST OF ABBREVIATIONS

Å	Angstrom, 10 ⁻¹⁰ meter
ABC	ATP-binding cassette
ADME	Absorption Distribution Metabolism Excretion
AP	Alkaline phosphatase
ATB	Automated topology builder
BBB	Blood Brain Barrier
BBR	Berberine
BIRC	Baculoviral IAP repeat-containing <i>protein</i>
BRCA	Breast cancer protein
CCND1	Cyclin D1
CCND3	Cyclin D3
CDK6	Cyclin-dependent kinase 6
CDKN2A	Cyclin-dependent kinase inhibitor 2A
cIAP	Cellular inhibitor of apoptosis
DMSO	Dimethyl sulfoxide
DS	Discovery studio
EGF	Epidermal growth factor
EGFR	Epidermal growth factor receptor
ELISA	Enzyme-linked immunosorbent assay
EM	Energy minimization
ERBB	Human epidermal growth factor receptor
ERK	Extracellular signal-regulated kinase
FACS	Fluorescent associated cell sorting
FBS	Fetal bovine serum
FEN1	Flap structure-specific endonuclease 1
FF	Force field
GA	Genetic algorithm
Genrestr	Generate position restraints
Gromacs	GRoningen MACHine for Chemical Simulations
GROMOS	GRoningen MOlecular Simulation computer program package

hAGP	human alpha1-acid glycoprotein
HBA	Hydrogen bond acceptor
HBD	Hydrogen bond donor
HER2	Human epidermal growth factor receptor 2
HRP	Horseradish peroxidase
HAS	Human serum albumin
IC50	Half inhibitory concentration
Ki	Inhibition constant
LEH	Ligand efficiency index
LEI	Ligand efficiency index
LGA	Lamarckian genetic algorithm
LJ	Lennard-Jones potential
LLE	Ligand lipophilicity efficiency index
M	Molar
MAPK	Mitogen activated protein kinase
MAPK1	Mitogen-activated protein kinase 1 (ERK2)
MAPK3	Mitogen-activated protein kinase 1 (ERK1)
MAPK14	Mitogen-activated protein kinase 14 (p38 MAPK)
MD	Molecular dynamics
μM	Micromolar, 10^{-6} molar
μm	Micrometer, 10^{-6} meter
MTOR	Mechanistic target of rapamycin (serine/threonine kinase)
MTT	3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide
MW	Molecular weight
nBEI	Binding efficiency index based on non-hydrogen atoms
NHA	Number of non-hydrogen atoms
NIH	National institute of health
nM	Nanomolar, 10^{-9} molar
nm	Nanometer, 10^{-9} meter
ns	Nanosecond, 10^{-9} second
ps	Picosecond, 10^{-12} second
NPOL	Number of polar atoms

NPT	Constant number of atom, pressure, temperature
NSEI	Binding efficiency index based on polar surface area
nTDOF	Number of torsional degree of freedom
NVT	Constant number of atom, volume, temperature
PBS	Phosphate Buffered Saline
PCNA	Proliferating cell nuclear antigen
PDB	Protein data bank
PI	Propidium iodide
PI3K	Phosphatidylinositide 3-kinase
PK	Pharmacokinetics
pKi	-LogKi
Posres	Position restraints
RAF	Rapidly Accelerated Fibrosarcoma
RDF	Radial distribution function
Rg	Radius of gyration
RFC1	Replication factor C (activator 1) 1
RFC3	Replication factor C (activator 1) 3
RFC4	replication factor C (activator 1) 4
RFC5	Replication factor C (activator 1) 5
RMSD	Root-mean-square deviation
RMSF	Root-mean-square fluctuation
ROS	reactive oxygen species
SHC1	SHC (Src homology 2 domain containing) transforming protein 1
Smo	Smoothened receptor
SPC	Simple Point Charge water model
SPDBV	Swiss protein data bank visualizer
TM	Transmembrane
TORDOF	Torsional degree of freedom
TPSA	Topological polar surface area
vdW	van der Waals
VEGF	Vascular endothelial growth factor



CHAPTER 1

INTRODUCTION

1.1 Brief background

Breast cancer is the most frequently diagnosed cancer in women, and is the second most common cancer worldwide. In 2012, 25% (1.67 million) of all new cancer cases and 15% (522,000) of all cancer deaths in women were due to breast cancer (Tao et al., 2015). Based on the data provided between 2000 and 2014, an estimated 252,710 new cases of invasive breast cancer will be diagnosed among women and 2,470 cases will be diagnosed in men in 2017. In addition, 63,410 cases of in situ breast carcinoma will be diagnosed among women. Approximately 40,610 women and 460 men are expected to die from breast cancer in 2017 (American Cancer Society, 2017) (Figure 1.1). Breast cancer is also the most common cancer in South-East Asian females. A woman in Malaysia had a 1 in 20 chance of developing breast cancer in her lifetime. In Malaysia, 37 individuals per 100000 of population for breast cancer cases, and 14.7 individuals per 100000 of population for death related to breast cancer have been reported (Miao et al., 2014).

Berberine can be isolated from the stems and roots of several plants, such as *Berberis vulgaris* and *Coptis chinensis* (Tomosaka et al., 2008; Zovko Končić et al., 2010; Abd El-Wahab et al., 2013). Berberine is a nitrogenous cyclic compound (Figure 1.2) with a structure that is highly similar to that of intercalating agents (e.g., ethidium). Intercalating agents are often used as nucleic acid dyes to study cell functions, and berberine is a well-known alkaloid drug that is commonly used as a fluorescent dye (Wang et al., 2012).

Berberine (Figure 1.2) induces apoptosis and inhibits cell proliferation in various cell lines derived from breast, lung, colon, and liver cancer. However, berberine has been shown to have synergistic effects on cells treated in combination with more toxic drugs, including vincristine and irinotecan (Kars et al., 2008). Previous studies showed that drug resistance was decreased by combination treatments with berberine (Sun et al., 2009; Zhou et al., 2012; Kukula-Koch et al., 2013).

Despite these findings, berberine efficacy and the molecular regulators that are targeted by berberine remain unclear. This thesis aims to provide an extensive analysis of berberine effects on various molecular mechanisms. In addition, this thesis also covers the metabolism, toxicity and adverse effects of berberine by using of molecular simulation of pharmacodynamics and pharmacokinetics for different berberine derivatives to provide more effective drug candidates than berberine.




Figure 1.1: Prevalence of breast cancer and other kinds of cancer in the United States
(A) Estimated new cancer cases in US in 2017; **(B)** Estimated number of deaths by age for female breast cancer.
 Source: (American Cancer Society, 2017)




Figure 1.2: Berberine structure

Berberine is an alkaloid containing three aromatic rings and five polar atoms. Topological Polar Surface Area (TPSA) is 40.8 Å².

1.2 Problem statement

There are many research gaps associated with anticancer effects of berberine. By damaging and mutating the DNA, oxidative stress, radiation, and mutagens turn normal cells to cancerous forms. As a result, while these mutations change certain genes that control critical biological processes, one or several kinds of cancer occur. The use of anti-oxidant compounds, which is broadly paid attention as an anti-cancer treatment, is a long-term treatment method. Whilst a cancer case observed, however, a significant and short-term treatment way must be found. Drugs which control genes and proteins that are not working properly must be discovered. Consequently, effects of drugs on genes that can cause cancer must be studied.

While drug candidates are studied by different experimental methods to evaluate its specific target, it is generally hard to spend millions of dollars to pass different levels of approval steps: preclinical, and phase 1-3 clinical phases. The challenge of *in silico* studies is the time, often several months, required for molecular modelling and molecular dynamics simulations. Accordingly, the mentioned mixed strategy of *in silico* and *in vitro* investigations not only save the budget and increase the number of drug candidates before starting the laboratorial works, but it can also lead to the more effective drug candidates; meanwhile, the toxicity and distributions of candidates can be also modeled.

There is a lack of such software, at least as a well-known and universal tool, to compute the ligand binding efficiency index and several other parameters from the thousands of input compounds. At the moment, the current tools of drug discovery analyze separate steps of drug discovery and many parameters should be manually calculated. This makes them

difficult to use for thousands of molecules, and the researchers must be involved during throughout of the procedures.

1.3 Objectives

1.3.1 General objective

To determine the effects of berberine and its derivatives on different signalling pathways involved in breast cancer.

1.3.2 Specific objectives

- a. To identify more effective berberine derivatives against smoothened receptor.
- b. To identify more effective berberine derivatives as multi-kinase inhibitors for cancer treatment.
- c. To determine anti-tumour activity of berberine in breast cancer cell lines.

1.4 Hypothesis

- a. Berberine inhibits smoothened receptor.
- b. Berberine derivatives are multi-kinase inhibitors in cancer treatment.
- c. Berberine is anti-tumour in breast cancer cell lines.

REFERENCES

- Abad-zapatero, C. Perspective Special Focus : Horizons in Medicinal Chemistry Alternative Variables in Drug Discovery : Promises and Challenges. *Futur. Med. Chem.* **2014**, 577–593.
- Abad-Zapatero, C. Ligand Efficiency Indices for Effective Drug Discovery. *Expert Opin. Drug Discov.* **2007**, 2 (4), 469–488.
- Abad-Zapatero, C. *Ligand Efficiency Indices for Drug Discovery: Towards an Atlas-Guided Paradigm*; Elsevier Science, 2013.
- Abad-Zapatero, C.; Metz, J. T. Ligand Efficiency Indices as Guideposts for Drug Discovery. *Drug Discov. Today* **2005**, 10 (7), 464–469.
- Abad-Zapatero, C.; Blasi, D. Ligand Efficiency Indices (LEIs): More than a Simple Efficiency Yardstick. *Mol. Inform.* **2011**, 30 (2–3), 122–132.
- Abad-Zapatero, C.; Champness, E. J.; Segall, M. D. Alternative Variables in Drug Discovery: Promises and Challenges. *Future Med. Chem.* **2014**, 6 (5), 577–593.
- Abd El-Wahab, A. E.; Ghareeb, D. A.; Sarhan, E. E. M.; Abu-Serie, M. M.; El Demellawy, M. A. *In Vitro* Biological Assessment of Berberis Vulgaris and Its Active Constituent, Berberine: Antioxidants, Anti-Acetylcholinesterase, Anti-Diabetic and Anticancer Effects. *BMC Complement. Altern. Med.* **2013**, 13 (1), 218.
- Abraham, M.J., van der Spoel, D., Lindahl, E., Hess., B., and the G. development team. Gromacs, User Manual: Version 5.0.5. 2015.
- Aller, S. ; Yu, J.; Ward, A.; Weng, Y.; Chittaboina, S.; Zhuo, R.; Harrell, P. ; Trinh, Y. ; Zhang, Q.; Urbatsch, I. ; et al. Structure of P-Glycoprotein Reveals a Molecular Basis for Poly-Specific Drug Binding. *Science (80-.)*. **2009**, 323, 1718–1722.
- Amakye, D.; Jagani, Z.; Dorsch, M. Unraveling the Therapeutic Potential of the Hedgehog Pathway in Cancer. *Nat. Med.* **2013**, 19 (11), 1410–1422.
- American Cancer Society. *Breast Cancer - Fact S & Figures (2017-2018)*; 2017.
- Andersen, O. A.; Nathubhai, A.; Dixon, M. J.; Eggleston, I. M.; van Aalten, D. M. F. Structure-Based Dissection of the Natural Product Cyclopentapeptide Chitinase Inhibitor Argifin. *Chem. Biol.* **2008**, 15 (3), 295–301.
- Ayati, S. H.; Fazeli, B.; Momtazi-borjani, A. A.; Cicero, A. F. G.; Pirro, M.; Sahebkar, A. Regulatory Effects of Berberine on microRNome in Cancer and Other Conditions. *Crit. Rev. Oncol.*

Hematol. **2017**, 116, 147–158.

Balakrishna, A.; Kumar, M. H. Evaluation of Synergetic Anticancer Activity of Berberine and Curcumin on Different Models of A549, Hep-G2, MCF-7, Jurkat, and K562 Cell Lines. *Biomed Res. Int.* **2015**, 354614, 1–7.

Baselga, J. Targeting Tyrosine Kinases in Cancer: The Second Wave. *Science* **2006**, 312, 1175–1178.

Benight, A.S.; Riccelli, P.V.; Hopfinger, A.J.; Pancoska, P. Modular Computational Models for Predicting the Pharmaceutical Properties of Chemical Compunds. US20020169561, 2002.

Bollag, G.; Hirth, P.; Tsai, J.; Zhang, J.; Ibrahim, P. N.; Cho, H.; Spevak, W.; Zhang, C.; Zhang, Y.; Habets, G.; et al. Clinical Efficacy of a RAF Inhibitor Needs Broad Target Blockade in BRAF-Mutant Melanoma. *Nature* **2010**, 467 (7315), 596–599.

Brastianos, P. K.; Horowitz, P. M.; Santagata, S.; Jones, R. T.; McKenna, A.; Getz, G.; Ligon, K. L.; Palescandolo, E.; Van Hummelen, P.; Ducar, M. D.; et al. Genomic Sequencing of Meningiomas Identifies Oncogenic SMO and AKT1 Mutations. *Nat. Genet.* **2013**, 45 (3), 285–289.

Brinkhuizen, T.; Reinders, M. G.; van Geel, M.; Hendriksen, A. J. L.; Paulussen, A. D. C.; Winnepenninckx, V. J.; Keymeulen, K. B.; Soetekouw, P. M. M. B.; van Steensel, M. A. M.; Mosterd, K. Acquired Resistance to the Hedgehog Pathway Inhibitor Vismodegib due to Smoothened Mutations in Treatment of Locally Advanced Basal Cell Carcinoma. *J. Am. Acad. Dermatol.* **2014**, 71 (5), 1005–1008.

Buonamici, S.; Williams, J.; Morrissey, M.; Wang, A.; Guo, R.; Vattay, A.; Hsiao, K.; Yuan, J.; Green, J.; Ospina, B.; et al. Interfering with Resistance to Smoothened Antagonists by Inhibition of the PI3K Pathway in Medulloblastoma. *Sci. Transl. Med.* **2010**, 2 (51), 51ra70.

Burgeiro, A.; Gajate, C.; Dakir, E. H.; Villa-Pulgarín, J. A.; Oliveira, P. J.; Mollinedo, F. Involvement of Mitochondrial and B-Raf/ERK Signaling Pathways in Berberine-Induced Apoptosis in Human Melanoma Cells. *Anticancer. Drugs* **2011**, 22 (6), 507–518.

Burness, C. B. Sonidegib: First Global Approval. *Drugs* **2015**, 75 (13), 1559–1566.

Chang, S.; Sharan, S. K. BRCA1 and MicroRNAs: Emerging Networks and Potential Therapeutic Targets. *Mol. Cells* **2012**, 34 (5), 425–432.

Chang, Y. P.; Huang, C. C.; Shen, C. C.; Tsai, K. C.; Ueng, Y. F. Differential Inhibition of CYP1-Catalyzed Regioselective Hydroxylation of Estradiol by Berberine and Its Oxidative

- Metabolites. *Drug Metab. Pharmacokinet.* **2015**, 30 (5), 374–383.
- Chen, J.; Wang, T.; Xu, S.; Lin, A.; Yao, H.; Xie, W.; Zhu, Z.; Xu, J. Design, Synthesis and Biological Evaluation of Novel 3-Substituted 4-Anilino-Coumarin Derivatives as Antitumor Agents. *Bioorg. Med. Chem. Lett.* **2017**, 27 (4), 867–874.
- Chen, L.; Bourguignon, L. Y. W. Hyaluronan-CD44 Interaction Promotes c-Jun Signaling and miRNA21 Expression Leading to Bcl-2 Expression and Chemoresistance in Breast Cancer Cells. *Mol. Cancer* **2014**, 13 (1), 52.
- Cheng, T.; Li, Q.; Zhou, Z.; Wang, Y.; Bryant, S. H. Structure-Based Virtual Screening for Drug Discovery: A Problem-Centric Review. *AAPS J.* **2012**, 14 (1), 133–141.
- Chiosea, S. I.; Grandis, J. R.; Lui, V. W. Y.; Diergaarde, B.; Maxwell, J. H.; Ferris, R. L.; Kim, S. W.; Luvison, A.; Miller, M.; Nikiforova, M. N. PIK3CA, HRAS and PTEN in Human Papillomavirus Positive Oropharyngeal Squamous Cell Carcinoma. *BMC Cancer* **2013**, 13, 602.
- Choi, H. E.; Choi, J. H.; Lee, J. Y.; Kim, J. H.; Kim, J. H.; Lee, J. K.; Kim, G. Il; Park, Y.; Chi, Y. H.; Paik, S. H.; et al. Synthesis and Evaluation of Nicotinamide Derivative as Anti-Angiogenic Agents. *Bioorganic Med. Chem. Lett.* **2013**, 23 (7), 2083–2088.
- Chong, H. Z.; Rahmat, A.; Yeap, S. K.; Md Akim, A.; Alitheen, N. B.; Othman, F.; Gwendoline-Ee, C. L. In Vitro Cytotoxicity of Strobilanthes Crispus Ethanol Extract on Hormone Dependent Human Breast Adenocarcinoma MCF-7 Cell. *BMC Complement. Altern. Med.* **2012**, 12 (1), 1033.
- Ciruelos Gil, E. M. Targeting the PI3K/AKT/mTOR Pathway in Estrogen Receptor-Positive Breast Cancer. *Cancer Treat. Rev.* **2014**, 40 (7), 862–871.
- Condon, S. M.; Mitsuuchi, Y.; Deng, Y.; Laporte, M. G.; Rippin, S. R.; Haimowitz, T.; Alexander, M. D.; Kumar, P. T.; Hendi, M. S.; Lee, Y. H.; et al. Birinapant, a Smac-Mimetic with Improved Tolerability for the Treatment of Solid Tumors and Hematological Malignancies. *J. Med. Chem.* **2014**, 57 (9), 3666–3677.
- Cunningham, D. C.; Harrison, L. Y.; Shultz, T. D. Proliferative Responses of Normal Human Mammary and MCF-7 Breast Cancer Cells to Linoleic Acid, Conjugated Linoleic Acid and Eicosanoid Synthesis Inhibitors in Culture. *Anticancer Res.* **1997**, 17 (1A), 197–203.
- DeFeo-Jones, D.; Barnett, S. F.; Fu, S.; Hancock, P. J.; Haskell, K. M.; Leander, K. R.; McAvoy, E.; Robinson, R. G.; Duggan, M. E.; Lindsley, C. W.; et al. Tumor Cell Sensitization to Apoptotic

- Stimuli by Selective Inhibition of Specific Akt/PKB Family Members. *Mol. Cancer Ther.* **2005**, 4 (2), 271–279.
- Deng, W. G.; Jayachandran, G.; Wu, G.; Xu, K.; Roth, J. A.; Ji, L. Tumor-Specific Activation of Human Telomerase Reverses Transcriptase Promoter Activity by Activating Enhancer-Binding Protein-2?? In Human Lung Cancer Cells. *J. Biol. Chem.* **2007**, 282 (36), 26460–26470.
- Dessole, G.; Branca, D.; Ferrigno, F.; Kinzel, O.; Muraglia, E.; Palumbi, M. C.; Rowley, M.; Serafini, S.; Steinkühler, C.; Jones, P. Discovery of N-[(1-Aryl-1H-Indazol-5-Yl)methyl]amides Derivatives as Smoothed Antagonists for Inhibition of the Hedgehog Pathway. *Bioorganic Med. Chem. Lett.* **2009**, 19 (15), 4191–4195.
- Dhillon, A. S.; Hagan, S.; Rath, O.; Kolch, W. MAP Kinase Signalling Pathways in Cancer. *Oncogene* **2007**, 26 (22), 3279–3290.
- Do, M. T.; Kim, H. G.; Tran, T. T. P.; Khanal, T.; Choi, J. H.; Chung, Y. C.; Jeong, T. C.; Jeong, H. G. Metformin Suppresses CYP1A1 and CYP1B1 Expression in Breast Cancer Cells by down-Regulating Aryl Hydrocarbon Receptor Expression. *Toxicol. Appl. Pharmacol.* **2014**, 280 (1), 138–148.
- Eckelman, B. P.; Salvesen, G. S.; Scott, F. L. Human Inhibitor of Apoptosis Proteins: Why XIAP Is the Black Sheep of the Family. *EMBO Rep.* **2006**, 7 (10), 988–994.
- Elgamal, O. A.; Park, J. K.; Gusev, Y.; Azevedo-Pouly, A. C. P.; Jiang, J.; Roopra, A.; Schmittgen, T. D. Tumor Suppressive Function of Mir-205 in Breast Cancer Is Linked to HMGB3 Regulation. *PLoS One* **2013**, 8 (10), 1–12.
- Eom, K.-S.; Hong, J.-M.; Youn, M.-J.; So, H.-S.; Park, R.; Kim, J.-M.; Kim, T.-Y. Berberine Induces G1 Arrest and Apoptosis in Human Glioblastoma T98G Cells through Mitochondrial/caspases Pathway. *Biol. Pharm. Bull.* **2008**, 31 (4), 558–562.
- Fan, S. Sanguinarine Is a Novel VEGF Inhibitor Involved in the Suppression of Angiogenesis and Cell Migration. *Mol. Clin. Oncol.* **2012**, 331–336.
- Fernandes, E.; Ferreira, J. A.; Andreia, P.; Luís, L.; Barroso, S.; Sarmiento, B.; Santos, L. L. New Trends in Guided Nanotherapies for Digestive Cancers: A Systematic Review. *J. Control. Release* **2015**, 209 (May), 288–307.
- Filip, G. A.; Postescu, I. D.; Bolfa, P.; Catoi, C.; Muresan, A.; Clichici, S. Inhibition of UVB-Induced Skin Phototoxicity by a Grape Seed Extract as Modulator of Nitrosative Stress, ERK/NF-kB Signaling Pathway and Apoptosis, in SKH-1 Mice. *Food Chem. Toxicol.* **2013**, 57, 296–306.

- Fitzgerald, T. L.; Lertpiriyapong, K.; Cocco, L.; Martelli, A. M.; Libra, M.; Candido, S.; Montalto, G.; Cervello, M.; Steelman, L.; Abrams, S. L.; et al. Roles of EGFR and KRAS and Their Downstream Signaling Pathways in Pancreatic Cancer and Pancreatic Cancer Stem Cells. *Adv. Biol. Regul.* **2015**, 59, 65–81.
- Fosgerau, K.; Hoffmann, T. Peptide Therapeutics: Current Status and Future Directions. *Drug Discovery Today* **2015**, 20 (1), 122–128.
- Franco-Molina, M. A.; Mendoza-Gamboa, E.; Sierra-Rivera, C. A.; Gómez-Flores, R. A.; Zapata-Benavides, P.; Castillo-Tello, P.; Alcocer-González, J. M.; Miranda-Hernández, D. F.; Tamez-Guerra, R. S.; Rodríguez-Padilla, C. Antitumor Activity of Colloidal Silver on MCF-7 Human Breast Cancer Cells. *J. Exp. Clin. Cancer Res.* **2010**, 29 (1), 148.
- Fu, L.; Chen, W.; Guo, W.; Wang, J.; Tian, Y.; Shi, D.; Zhang, X.; Qiu, H.; Xiao, X.; Kang, T.; et al. Berberine Targets AP-2/hTERT, NF- κ B/COX-2, HIF-1 α /VEGF and Cytochrome-c/Caspase Signaling to Suppress Human Cancer Cell Growth. *PLoS One* **2013**, 8 (7), 1–13.
- Gao, M. Y.; Chen, L.; Yang, L.; Yu, X.; Kou, J. P.; Yu, B. Y. Berberine Inhibits LPS-Induced TF Procoagulant Activity and Expression through NF- κ B/p65, Akt and MAPK Pathway in THP-1 Cells. *Pharmacol. Reports* **2014**, 66 (3), 480–484.
- Ghuman, J.; Zunszain, P. .; Petitpas, I.; Bhattacharya, A.; Otagiri, M.; Curry, S. Structural Basis of the Drug-Binding Specificity of Human Serum Albumin. *J.Mol.Biol.* **2005**, 353, 38.
- Giri, P.; Kumar, G. S. Specific Binding and Self-Structure Induction to poly(A) by the Cytotoxic Plant Alkaloid Sanguinarine. *Biochim. Biophys. Acta - Gen. Subj.* **2007**, 1770 (9), 1419–1426.
- Giussani, P.; Brioschi, L.; Bassi, R.; Riboni, L.; Viani, P. Phosphatidylinositol 3-kinase/AKT Pathway Regulates the Endoplasmic Reticulum to Golgi Traffic of Ceramide in Glioma Cells: A Link between Lipid Signaling Pathways Involved in the Control of Cell Survival. *J. Biol. Chem.* **2009**, 284 (8), 5088–5096.
- Graziani, G.; Artuso, S.; De Luca, A.; Muzi, A.; Rotili, D.; Scimeca, M.; Atzori, M. G.; Ceci, C.; Mai, A.; Leonetti, C.; et al. A New Water Soluble MAPK Activator Exerts Antitumor Activity in Melanoma Cells Resistant to the BRAF Inhibitor Vemurafenib. *Biochem. Pharmacol.* **2014**, 95 (1), 16–27.
- Guo, Q. J.; Mills, J. N.; Bandurraga, S. G.; Nogueira, L. M.; Mason, N. J.; Camp, E. R.; Larue, A. C.; Turner, D. P.; Findlay, V. J. MicroRNA-510 Promotes Cell and Tumor Growth by Targeting peroxiredoxin1 in Breast Cancer. *Breast Cancer Res.* **2013**, 15 (R70), 1–13.

- Guo, Y.; Li, F.; Ma, X.; Cheng, X.; Zhou, H.; Klaassen, C. D. CYP2D Plays a Major Role in Berberine Metabolism in Liver of Mice and Humans. *Xenobiotica* **2011**, 41 (11), 996–1005.
- Gupta, N.; Sitwala, N.; Patel, K. Pharmacophore Modelling, Validation, 3D Virtual Screening, Docking, Design and in Silico ADMET Simulation Study of Histone Deacetylase Class-1 Inhibitors. *Med. Chem. Res.* **2014**, 23 (11), 4853–4864.
- Herberich, B.; Jackson, C.; Wurz, R. .; Pettus, L. .; Sherman, L.; Liu, Q.; Henkle, B.; Saris, C. .; Wong, L. .; Chmait, S.; et al. Identification of Triazolopyridazinones as Potent p38alpha Inhibitors. *Bioorg.Med.Chem.Lett.* **2012**, 22, 1226–1229.
- Hiremath, C. N. Method for Predicting ADMET Properties Using Abbreviated Profile of Drugs (A-POD). US20130117000, 2013.
- Ho, Y. T.; Lu, C. C.; Yang, J. S.; Chiang, J. H.; Li, T. C.; Ip, S. W.; Hsia, T. C.; Liao, C. L.; Lin, J. G.; Wood, W. G.; et al. Berberine Induced Apoptosis via Promoting the Expression of Caspase-8, -9 and -3, Apoptosis-Inducing Factor and Endonuclease G in SCC-4 Human Tongue Squamous Carcinoma Cancer Cells. *Anticancer Res.* **2009**, 29 (10), 4063–4070.
- Hopkins, A. L.; Groom, C. R.; Alex, A. Ligand Efficiency: A Useful Metric for Lead Selection. *Drug Discov. Today* **2004**, 9 (10), 430–431.
- Hou, Q.; Tang, X.; Liu, H.; Tang, J.; Yang, Y.; Jing, X.; Xiao, Q.; Wang, W.; Gou, X.; Wang, Z. Berberine Induces Cell Death in Human Hepatoma Cells in Vitro by Downregulating CD147. *Cancer Sci.* **2011**, 102 (7), 1287–1292.
- Hou, W.; Liu, J.; Chen, P.; Wang, H.; Ye, B. C.; Qiang, F. Mutation Analysis of Key Genes in RAS/RAF and PI3K/PTEN Pathways in Chinese Patients with Hepatocellular Carcinoma. *Oncol. Lett.* **2014**, 8 (3), 1249–1254.
- Houghton, M.; Tuszynski, J. A.; Barakat, K.; Mohamed, A. Systems and Methods of Selecting Compounds with Reduced Risk of Cardiotoxicity. US20150193575A1, 2015.
- Hsu, W. H.; Hsieh, Y. S.; Kuo, H. C.; Teng, C. Y.; Huang, H. I.; Wang, C. J.; Yang, S. F.; Liou, Y. S.; Kuo, W. H. Berberine Induces Apoptosis in SW620 Human Colonic Carcinoma Cells through Generation of Reactive Oxygen Species and Activation of JNK/p38 MAPK and FasL. *Arch. Toxicol.* **2007**, 81 (10), 719–728.
- Hu, H. Y.; Li, K. P.; Wang, X. J.; Liu, Y.; Lu, Z. G.; Dong, R. H.; Guo, H. B.; Zhang, M. X. Set9, NF-κB, and microRNA-21 Mediate Berberine-Induced Apoptosis of Human Multiple Myeloma Cells. *Acta Pharmacol. Sin.* **2013**, 34 (1), 157–166.
- Huang, Z.; Zheng, H.; Wang, W.; Wang, Y.; Zhong, L.; Wu, J.; Li, Q. Berberine Targets Epidermal Growth Factor Receptor

- Signaling to Suppress Prostate Cancer Proliferation *in vitro*. *Mol. Med. Rep.* **2014**, 11, 2125–2128.
- Hwang, J. M.; Kuo, H. C.; Tseng, T. H.; Liu, J. Y.; Chu, C. Y. Berberine Induces Apoptosis through a Mitochondria/caspases Pathway in Human Hepatoma Cells. *Arch. Toxicol.* **2006**, 80 (2), 62–73.
- Hwang, J. T.; Kwon, D. Y.; Yoon, S. H. AMP-Activated Protein Kinase: A Potential Target for the Diseases Prevention by Natural Occurring Polyphenols. *N. Biotechnol.* **2009**, 26 (1–2), 17–22.
- Ishii, T.; Shimizu, Y.; Nakashima, K.; Kondo, S.; Ogawa, K.; Sasaki, S.; Matsui, H. Inhibition Mechanism Exploration of Investigational Drug TAK-441 as Inhibitor against Vismodegib-Resistant Smoothed Mutant. *Eur. J. Pharmacol.* **2014**, 723 (1), 305–313.
- Islam, M. M.; Kumar, G. S. Small Molecule-RNA Interaction: Spectroscopic and Calorimetric Studies on the Binding by the Cytotoxic Protoberberine Alkaloid Coralyne to Single Stranded Polyribonucleotides. *Biochim. Biophys. Acta - Gen. Subj.* **2009**, 1790 (8), 829–839.
- Jang, M. J.; Jwa, M.; Kim, J.-H.; Song, K. Selective Inhibition of MAPKK Wis1 in the Stress-Activated MAPK Cascade of *Schizosaccharomyces Pombe* by Novel Berberine Derivatives. *J. Biol. Chem.* **2002**, 277 (14), 12388–12395.
- Jarzembowski, J. A.; Rajagopalan, L. E.; Shin, H. C.; Malter, J. S. The 5'-untranslated Region of GM-CSF mRNA Suppresses Translational Repression Mediated by the 3' Adenosine-Uridine-Rich Element and the poly(A) Tail. *Nucleic Acids Res.* **1999**, 27 (18), 3660–3666.
- Kars, M. D.; Işeri, O. D.; Gunduz, U.; Molnar, J. Reversal of Multidrug Resistance by Synthetic and Natural Compounds in Drug-Resistant MCF-7 Cell Lines. *Chemotherapy* **2008**, 54 (3), 194–200.
- Kathryn, J. C.; Sireesha V, G.; Stanley, L. Triple Negative Breast Cancer Cell Lines: One Tool in the Search for Better Treatment of Triple Negative Breast Cancer. *Breast Dis* **2012**, 32, 35–48.
- Khan, M. I.; Mohammad, A.; Patil, G.; Naqvi, S. A. H.; Chauhan, L. K. S.; Ahmad, I. Induction of ROS, Mitochondrial Damage and Autophagy in Lung Epithelial Cancer Cells by Iron Oxide Nanoparticles. *Biomaterials* **2012**, 33 (5), 1477–1488.
- Khan, S.; Shukla, S.; Sinha, S.; Meeran, S. M. Role of Adipokines and Cytokines in Obesity-Associated Breast Cancer: Therapeutic Targets. *Cytokine Growth Factor Rev.* **2013**, 24 (6), 503–513.
- Kim, K.; Lee, J.; Park, Y.; Lee, S. ATF3 Mediates Anti-Cancer Activity of Trans -10 , Cis -12-Conjugated Linoleic Acid in Human Colon Cancer Cells. *Biomol. Ther.* **2015**, 23 (2), 134–140.

- Kim, S.; Han, J.; Kim, N. Y.; Lee, S. K.; Cho, D. H.; Choi, M. Y.; Kim, J. S.; Kim, J. H.; Choe, J. H.; Nam, S. J.; et al. Effect of Berberine on p53 Expression by TPA in Breast Cancer Cells. *Oncol. Rep.* **2012**, *27* (1), 210–215.
- Koziara, K. B.; Stroet, M.; Malde, A. K.; Mark, A. E. Testing and Validation of the Automated Topology Builder (ATB) Version 2.0: Prediction of Hydration Free Enthalpies. *J. Comput. Aided. Mol. Des.* **2014**, *28* (3), 221–233.
- Kubo, M.; Nakamura, M.; Tasaki, A.; Yamanaka, N.; Nakashima, H.; Nomura, M.; Kuroki, S.; Katano, M. Hedgehog Signaling Pathway Is a New Therapeutic Target for Patients with Breast Cancer. *Cancer Res.* **2004**, *64* (17), 6071–6074.
- Kukula-Koch, W.; Aligiannis, N.; Halabalaki, M.; Skaltsounis, A. L.; Glowniak, K.; Kalpoutzakis, E. Influence of Extraction Procedures on Phenolic Content and Antioxidant Activity of Cretan Barberry Herb. *Food Chem.* **2013**, *138* (1), 406–413.
- Kuntz, I. D.; Chen, K.; Sharp, K. A.; Kollman, P. A. The Maximal Affinity of Ligands. *Proc. Natl. Acad. Sci.* **1999**, *96* (18), 9997–10002.
- Kuo, H. P.; Chuang, T. C.; Tsai, S. C.; Tseng, H. H.; Hsu, S. C.; Chen, Y. C.; Kuo, C. L.; Kuo, Y. H.; Liu, J. Y.; Kao, M. C. Berberine, an Isoquinoline Alkaloid, Inhibits the Metastatic Potential of Breast Cancer Cells via Akt Pathway Modulation. *J. Agric. Food Chem.* **2012**, *60* (38), 9649–9658.
- Lee, D. U.; Kang, Y. J.; Park, M. K.; Lee, Y. S.; Seo, H. G.; Kim, T. S.; Kim, C. H.; Chang, K. C. Effects of 13-Alkyl-Substituted Berberine Alkaloids on the Expression of COX-II, TNF- α , iNOS, and IL-12 Production in LPS-Stimulated Macrophages. *Life Sci.* **2003**, *73* (11), 1401–1412.
- Lee, E.-J.; Oh, S.-Y.; Sung, M.-K. Luteolin Exerts Anti-Tumor Activity through the Suppression of Epidermal Growth Factor Receptor-Mediated Pathway in MDA-MB-231 ER-Negative Breast Cancer Cells. *Food Chem. Toxicol.* **2012**, *50* (11), 4136–4143.
- Letašiová, S.; Jantová, S.; Čipák, L.; Můčková, M. Berberine-Antiproliferative Activity in Vitro and Induction of Apoptosis/necrosis of the U937 and B16 Cells. *Cancer Lett.* **2006**, *239* (2), 254–262.
- Li, C.; Xi, Y.; Li, S.; Zhao, Q.; Cheng, W.; Wang, Z.; Zhong, J.; Niu, X.; Chen, G. Berberine Ameliorates TNBS Induced Colitis by Inhibiting Inflammatory Responses and Th1/Th17 Differentiation. *Mol. Immunol.* **2015**, *67* (2), 444–454.
- Li, G. X.; Wang, X. M.; Jiang, T.; Gong, J. F.; Niu, L. Y.; Li, N. Berberine Prevents Damage to the Intestinal Mucosal Barrier during Early Phase of Sepsis in Rat through Mechanisms

- Independent of the NOD-like Receptors Signaling Pathway. *Eur. J. Pharmacol.* **2014a**, 730 (1), 1–7.
- Li, H. X.; Zeng, J. F.; Shen, K. PI3K/AKT/mTOR Signaling Pathway as a Therapeutic Target for Ovarian Cancer. *Cancer Epidemiol.* **2016a**, 40, 152–157.
- Li, W.; Li, C.; Zheng, H.; Chen, G.; Hua, B. Therapeutic Targets of Traditional Chinese Medicine for Colorectal Cancer. *J. Tradit. Chinese Med.* **2016b**, 36 (2), 243–249.
- Li, Y.-B.; Zhao, W.-L.; Wang, Y.-X.; Zhang, C.-X.; Jiang, J.-D.; Bi, C.-W.; Tang, S.; Chen, R.-X.; Shao, R.-G.; Song, D.-Q. Discovery, Synthesis and Biological Evaluation of Cycloprotoberberine Derivatives as Potential Antitumor Agents. *Eur. J. Med. Chem.* **2013**, 68, 463–472.
- Li, Z.; Jiang, J.-D.; Kong, W.-J. Berberine Up-Regulates Hepatic Low-Density Lipoprotein Receptor through Ras-Independent but AMP-Activated Protein Kinase-Dependent Raf-1 Activation. *Biol. Pharm. Bull.* **2014b**, 37 (11), 1766–1775.
- Liang, K. W.; Ting, C. T.; Yin, S. C.; Chen, Y. T.; Lin, S. J.; Liao, J. K.; Hsu, S. L. Berberine Suppresses MEK/ERK-Dependent Egr-1 Signaling Pathway and Inhibits Vascular Smooth Muscle Cell Regrowth after in Vitro Mechanical Injury. *Biochem. Pharmacol.* **2006**, 71 (6), 806–817.
- Lipinski, C. A.; Lombardo, F.; Dominy, B. W.; Feeney, P. J. Experimental and Computational Approaches to Estimate Solubility and Permeability in Drug Discovery and Development Settings. *Adv. Drug Deliv. Rev.* **2001**, 46, 3–26.
- Lipinski, C. A.; Lombardo, F.; Dominy, B. W.; Feeney, P. J. Experimental and Computational Approaches to Estimate Solubility and Permeability in Drug Discovery and Development Settings. *Adv. Drug Deliv. Rev.* **2012**, 64 (SUPPL.), 4–17.
- Liu, C.; Hu, M.; Hsu, C.; Huang, C.; Hsu, C.; Chen, M.; Shiau, C.; Tseng, L. Lapatinib Inhibits CIP2A / PP2A / P-Akt Signaling and Induces Apoptosis in Triple Negative Breast Cancer Cells. *Oncotarget* **2016**, 7 (8), 9135–9149.
- Liu, J.; Zhang, X.; Liu, A.; Liu, S.; Zhang, L.; Wu, B.; Hu, Q. Berberine Induces Apoptosis in p53-Null Leukemia Cells by down-Regulating XIAP at the Post-Transcriptional Level. *Cell. Physiol. Biochem.* **2013**, 32 (5), 1213–1224.
- Liu, Q.; Xu, X.; Zhao, M.; Wei, Z.; Li, X.; Zhang, X.; Liu, Z.; Gong, Y.; Shao, C. Berberine Induces Senescence of Human Glioblastoma Cells by Downregulating the EGFR-MEK-ERK Signaling Pathway. *Mol. Cancer Ther.* **2015**, 14 (2), 355–363.
- Lo, S. N.; Chang, Y. P.; Tsai, K. C.; Chang, C. Y.; Wu, T. S.; Ueng, Y. F.

- Inhibition of CYP1 by Berberine, Palmatine, and Jatrorrhizine: Selectivity, Kinetic Characterization, and Molecular Modeling. *Toxicol. Appl. Pharmacol.* **2013**, 272 (3), 671–680.
- Lu, B.; Hu, M.; Liu, K.; Peng, J. Cytotoxicity of Berberine on Human Cervical Carcinoma HeLa Cells through Mitochondria, Death Receptor and MAPK Pathways, and in-Silico Drug-Target Prediction. *Toxicol. Vitro.* **2010**, 24 (6), 1482–1490.
- Lu, W.; Geng, D.; Sun, Z.; Yang, Z.; Ma, H.; Zheng, J.; Zhang, X. Scaffold Hopping Approach to a New Series of Smoothened Antagonists. *Bioorganic Med. Chem. Lett.* **2014**, 24 (10), 2300–2304.
- Lundborg, M.; Lindahl, E. Automatic GROMACS Topology Generation and Comparisons of Force Fields for Solvation Free Energy Calculations. *J. Phys. Chem. B* **2015**, 119, 810–823.
- Luqman, S.; Pezzuto, J. M. NFkB: A Promising Target for Natural Products in Cancer Chemoprevention. *Phyther. Res.* **2010**, 24 (7), 949–963.
- Lv, X. Bin; Jiao, Y.; Qing, Y.; Hu, H.; Cui, X.; Lin, T.; Song, E.; Yu, F. Original Article Genomic DNA Isolation and Materials and Methods Cell Lines and Culture. *Chin. J. Cancer* **2012**, 821–830.
- Lyons, T. G.; O'Kane, G. M.; Kelly, C. M. Efficacy and Safety of Vismodegib: A New Therapeutic Agent in the Treatment of Basal Cell Carcinoma. *Expert Opin. Drug Saf.* **2014**, 13 (8), 1125–1132.
- Ma, H.; Lu, W.; Sun, Z.; Luo, L.; Geng, D.; Yang, Z.; Li, E.; Zheng, J.; Wang, M.; Zhang, H.; et al. Design, Synthesis, and Structure-Activity-Relationship of Tetrahydrothiazolopyridine Derivatives as Potent Smoothened Antagonists. *Eur. J. Med. Chem.* **2015**, 89, 721–732.
- Mahata, S.; Bharti, A. C.; Shukla, S.; Tyagi, A.; Husain, S. A.; Das, B. C. Berberine Modulates AP-1 Activity to Suppress HPV Transcription and Downstream Signaling to Induce Growth Arrest and Apoptosis in Cervical Cancer Cells. *Mol. Cancer* **2011**, 10 (1), 39.
- Majima, H. J.; Indo, H. P.; Suenaga, S.; Kaneko, T.; Matsui, H.; Yen, H. C.; Ozawa, T. Mitochondria as Source of Free Radicals. *Free Radic. Biol. Dig. Dis.* **2010**, 29, 12–22.
- Malancona, S.; Altamura, S.; Filocamo, G.; Kinzel, O.; Hernando, J. I. M.; Rowley, M.; Scarpelli, R.; Steinkühler, C.; Jones, P. Identification of MK-5710 ((8aS)-8a-Methyl-1,3-Dioxo-2-[(1S,2R)-2-Phenylcyclopropyl]-N-(1-Phenyl-1H-Pyrazol-5-Yl)hexahydroimidazo[1,5-A]pyrazine-7(1H)-Carboxamide), a Potent Smoothened Antagonist for Use in Hedgehog Pathway

- Dependent Malignancies, Part . *Bioorganic Med. Chem. Lett.* **2011**, 21 (15), 4422–4428.
- Malde, A. K.; Zuo, L.; Breeze, M.; Stroet, M.; Poger, D.; Nair, P. C.; Oostenbrink, C.; Mark, A. E. An Automated Force Field Topology Builder (ATB) and Repository: Version 1.0. *J. Chem. Theory Comput.* **2011**, 7 (12), 4026–4037.
- Manallack, D. T.; Pranker, R. J.; Yuriev, E.; Oprea, T. I.; Chalmers, D. K.; David, K. The Significance of Acid/base Properties in Drug Discovery. *Chem. Soc. Rev.* **2014**, 42 (2), 485–496.
- Mantena, S. K.; Sharma, S. D.; Katiyar, S. K. Berberine Inhibits Growth, Induces G1 Arrest and Apoptosis in Human Epidermoid Carcinoma A431 Cells by Regulating Cdk1-Cdk-Cyclin Cascade, Disruption of Mitochondrial Membrane Potential and Cleavage of Caspase 3 and PARP. *Carcinogenesis* **2006**, 27 (10), 2018–2027.
- Martonák, R.; Laio, A.; Parrinello, M. Predicting Crystal Structures: The Parrinello-Rahman Method Revisited. *Phys. Rev. Lett.* **2003**, 90 (7), 75503.
- McCubrey, J. A.; Abrams, S. L.; Stadelman, K.; Chappell, W. H.; LaHair, M.; Ferland, R. A.; Steelman, L. S. Targeting Signal Transduction Pathways to Eliminate Chemotherapeutic Drug Resistance and Cancer Stem Cells. *Adv. Enzyme Regul.* **2010**, 50 (1), 285–307.
- McCubrey, J. A.; Steelman, L. S.; Kempf, C. R.; Chappell, W. H.; Abrams, S. L.; Stivala, F.; Malaponte, G.; Nicoletti, F.; Libra, M.; Bäscke, J.; et al. Therapeutic Resistance Resulting from Mutations in Raf/MEK/ERK and PI3K/PTEN/Akt/mTOR Signaling Pathways. *J. Cell. Physiol.* **2011**, 226 (11), 2762–2781.
- Mehrzadi, S.; Fatemi, I.; Esmaeilzadeh, M.; Ghaznavi, H.; Kalantar, H.; Goudarzi, M. Hepatoprotective Effect of Berberine against Methotrexate Induced Liver Toxicity in Rats. *Biomed. Pharmacother.* **2018**, 97, 233–239.
- Miao, H.; Hartman, M.; Bhoo-Pathy, N.; Lee, S.-C.; Taib, N. A.; Tan, E.-Y.; Chan, P.; Moons, K. G. M.; Wong, H.-S.; Goh, J.; et al. Predicting Survival of de Novo Metastatic Breast Cancer in Asian Women: Systematic Review and Validation Study. *PLoS One* **2014**, 9 (4), e93755.
- Mohammadi, A.; Mansoori, B.; Baradaran, B. Regulation of miRNAs by Herbal Medicine: An Emerging Field in Cancer Therapies. *Biomed. Pharmacother.* **2017**, 86, 262–270.
- Moroy, G.; Martiny, V. Y.; Vayer, P.; Villoutreix, B. O.; Miteva, M. A. Toward in Silico Structure-Based ADMET Prediction in Drug Discovery. *Drug Discov. Today* **2012**, 17 (1–2), 44–55.

- Morris, G.M.; Huey, R. AutoDock4 and AutoDockTools4: Automated Docking with Selective Receptor Flexibility. *J. Comput. Chem.* **2009**, *30* (16), 2785–2791.
- Morris, G. M.; Goodsell, D. S.; Halliday, R. S.; Huey, R.; Hart, W. E.; Belew, R. K.; Olson, A. J.; Al, M. E. T. Automated Docking Using a Lamarckian Genetic Algorithm and an Empirical Binding Free Energy Function. *J. Comput. Chem.* **1998**, *19* (14), 1639–1662.
- Morris, G. M.; Huey, R.; Olson, A. J. *Current Protocols in Bioinformatics: UNIT Using AutoDock for Ligand-Receptor Docking*; JohnWiley & Sons, Inc, 2008, Suppl.24, 1-40.
- Murahashi, H.; Azuma, H.; Zamzami, N.; Furuya, K.; Ikebuchi, K. Possible Contribution of Apoptosis-Inducing Factor (AIF) and Reactive Oxygen Species (ROS) to UVB-Induced Caspase-Independent Cell Death in the T Cell Line Jurkat. *J. Leukoc. Biol.* **2003**, *73*, 399–406.
- Muranen, T.; Selfors, L. M.; Worster, D. T.; Iwanicki, M. P.; Song, L.; Morales, F. C.; Gao, S.; Mills, G. B.; Brugge, J. S. Inhibition of PI3K/mTOR Leads to Adaptive Resistance in Matrix-Attached Cancer Cells. *Cancer Cell* **2012**, *21* (2), 227–239.
- Namba, H.; Nakashima, M.; Hayashi, T.; Hayashida, N.; Maeda, S.; Rogounovitch, T. I.; Ohtsuru, A.; Saenko, V. A.; Kanematsu, T.; Yamashita, S. Clinical Implication of Hot Spot BRAF Mutation, V599E, in Papillary Thyroid Cancers. *J. Clin. Endocrinol. Metab.* **2003**, *88* (9), 4393–4397.
- Neel, J. C.; Lebrun, J. J. Activin and TGF β Regulate Expression of the microRNA-181 Family to Promote Cell Migration and Invasion in Breast Cancer Cells. *Cell. Signal.* **2013**, *25* (7), 1556–1566.
- Nikoletopoulou, V.; Markaki, M.; Palikaras, K.; Tavernarakis, N. Crosstalk between Apoptosis, Necrosis and Autophagy. *Biochim. Biophys. Acta - Mol. Cell Res.* **2013**, *1833* (12), 3448–3459.
- Nishi, K.; Ono, T.; Nakamura, T.; Fukunaga, N.; Izumi, M.; Watanabe, H.; Suenaga, A.; Maruyama, T.; Yamagata, Y.; Curry, S.; et al. Structural Insights into Differences in Drug-Binding Selectivity between Two Forms of Human α 1-Acid Glycoprotein Genetic Variants, the A and F1*S Forms. *J. Biol. Chem.* **2011**, *286* (16), 14427–14434.
- Noureini, S. K.; Esmaili, H. Multiple Mechanisms of Cell Death Induced by Chelidonine in MCF-7 Breast Cancer Cell Line. *Chem. Biol. Interact.* **2014**, *223*, 141–149.
- Oberoi-Khanuja, T. K.; Murali, A.; Rajalingam, K. IAPs on the Move: Role of Inhibitors of Apoptosis Proteins in Cell Migration. *Cell Death Dis.* **2013**, *4* (9), e784-9.

- Onishi, H.; Katano, M. Hedgehog Signaling Pathway as a Therapeutic Target in Various Types of Cancer. *Cancer Sci.* **2011**, *102* (10), 1756–1760.
- Ontoria, J. M.; Bufi, L. L.; Torrisi, C.; Bresciani, A.; Giomini, C.; Rowley, M.; Serafini, S.; Bin, H.; Hao, W.; Steinkühler, C.; et al. Identification of a Series of 4-[3-(Quinolin-2-Yl)-1,2,4-Oxadiazol-5-Yl] Piperazinyl Ureas as Potent Smoothed Antagonist Hedgehog Pathway Inhibitors. *Bioorganic Med. Chem. Lett.* **2011**, *21* (18), 5274–5282.
- Pavlopoulou, A.; Oktay, Y.; Vougas, K.; Louka, M.; Vorgias, C. E.; Georgakilas, A. G. Determinants of Resistance to Chemotherapy and Ionizing Radiation in Breast Cancer Stem Cells. *Cancer Lett.* **2016**, *380* (2), 485–493.
- Perola, E. An Analysis of the Binding Efficiencies of Drugs and Their Leads in Successful Drug Discovery Programs. *J. Med. Chem.* **2010**, *53* (7), 2986–2997.
- Peters, E. A. J. F.; Goga, N.; Berendsen, H. J. C. Stochastic Dynamics with Correct Sampling for Constrained Systems. *J. Chem. Theory Comput.* **2014**, *10* (10), 4208–4220.
- Pettersen, E. F.; Goddard, T. D.; Huang, C. C.; Couch, G. S.; Greenblatt, D. M.; Meng, E. C.; Ferrin, T. E. UCSF Chimera--a Visualization System for Exploratory Research and Analysis. *J. Comput. Chem.* **2004**, *25* (13), 1605–1612.
- Peukert, S.; Miller-Moslin, K. Small-Molecule Inhibitors of the Hedgehog Signaling Pathway as Cancer Therapeutics. *ChemMedChem* **2010**, *5* (4), 500–512.
- Pierri, C. L.; Bossis, F.; Punzi, G.; De Grassi, A.; Cetrone, M.; Parisi, G.; Tricarico, D. Molecular Modeling of Antibodies for the Treatment of TNF α -Related Immunological Diseases. *Pharmacol. Res. Perspect.* **2016**, *4* (1), 1–13.
- Piyanuch, R.; Sukthanthar, M.; Wandee, G.; Baek, S. J. Berberine, a Natural Isoquinoline Alkaloid, Induces NAG-1 and ATF3 Expression in Human Colorectal Cancer Cells. *Cancer Lett.* **2007**, *258* (2), 230–240.
- Pujadas, G.; Vaque, M.; Ardevol, A.; Blade, C.; Salvado, M. J.; Blay, M.; Fernandez-Larrea, J.; Arola, L. Protein-Ligand Docking: A Review of Recent Advances and Future Perspectives. *Curr. Pharm. Anal.* **2008**, *4* (1), 1–19.
- Puzanov, I.; Amaravadi, R. K.; McArthur, G. A.; Flaherty, K. T.; Chapman, P. B.; Sosman, J. A.; Ribas, A.; Shackleton, M.; Hwu, P.; Chmielowski, B.; et al. Long-Term Outcome in BRAF(V600E) Melanoma Patients Treated with Vemurafenib: Patterns of Disease Progression and Clinical Management of Limited Progression. *Eur. J. Cancer* **2015**, *51* (11), 1435–1443.

- Qadir, S. A.; Kwon, M. C.; Han, J. G.; Ha, J. H.; Chung, H. S.; Ahn, J.; Lee, H. Y. Effect of Different Extraction Protocols on Anticancer and Antioxidant Activities of Berberis Koreana Bark Extracts. *J. Biosci. Bioeng.* **2009**, *107* (3), 331–338.
- Raval, A.; Piana, S.; Eastwood, M. P.; Dror, R. O.; Shaw, D. E. Refinement of Protein Structure Homology Models via Long, All-Atom Molecular Dynamics Simulations. *Proteins Struct. Funct. Bioinforma.* **2012**, *80* (8), 2071–2079.
- Refaat, A.; Abdelhamed, S.; Yagita, H.; Inoue, H.; Yokoyama, S.; Hayakawa, Y.; Saiki, I. Berberine Enhances Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand-Mediated Apoptosis in Breast Cancer. *Oncol. Lett.* **2013**, *6* (3), 840–844.
- Refaat, A.; Abdelhamed, S.; Saiki, I.; Sakurai, H. Inhibition of p38 Mitogen-Activated Protein Kinase Potentiates the Apoptotic Effect of Berberine/tumor Necrosis Factor-Related Apoptosis-Inducing Ligand Combination Therapy. *Oncol. Lett.* **2015**, 1907–1911.
- Ren, L.; Wenglowisky, S.; Miknis, G.; Rast, B.; Buckmelter, A. J.; Ely, R. J.; Schlachter, S.; Laird, E. R.; Randolph, N.; Callejo, M.; et al. Non-Oxime Inhibitors of B-RafV600E Kinase. *Bioorganic Med. Chem. Lett.* **2011**, *21* (4), 1243–1247.
- Robinson, S. D.; O'Shaughnessy, J. A.; Cowey, C. L.; Konduri, K. BRAF V600E-Mutated Lung Adenocarcinoma with Metastases to the Brain Responding to Treatment with Vemurafenib. *Lung Cancer* **2014**, *85* (2), 326–330.
- Rognan, D.; Mus-Veteau, I. Three-Dimensional Structure of the Smoothed Receptor: Implications for Drug Discovery. *Top. Med. Chem.* **2015**, *16*, 127–146.
- Roskoski, R. RAF Protein-Serine/threonine Kinases: Structure and Regulation. *Biochem. Biophys. Res. Commun.* **2010**, *399* (3), 313–317.
- Rossi, A.; Pasquale, R.; Esposito, C.; Normanno, N. Should Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors Be Considered Ideal Drugs for the Treatment of Selected Advanced Non-Small Cell Lung Cancer Patients? *Cancer Treat. Rev.* **2013**, *39* (5), 489–497.
- Ruat, M.; Hoch, L.; Faure, H.; Rognan, D. Targeting of Smoothed for Therapeutic Gain. *Trends Pharmacol. Sci.* **2014**, *35* (5), 237–246.
- Rudin, C. M.; Hann, C. L.; Laterra, J.; Yauch, R. L.; Callahan, C. A.; Fu, L.; Holcomb, T.; Stinson, J.; Gould, S. E.; Coleman, B.; et al. Treatment of Medulloblastoma with Hedgehog Pathway Inhibitor GDC-0449. *N. Engl. J. Med.* **2009**, *361* (12), 1173–1178.

- Ruiz, A.; Hermosín-Gutiérrez, I.; Mardones, C.; Vergara, C.; Herlitz, E.; Vega, M.; Dorau, C.; Winterhalter, P.; Von Baer, D. Polyphenols and Antioxidant Activity of Calafate (*Berberis Microphylla*) Fruits and Other Native Berries from Southern Chile. *J. Agric. Food Chem.* **2010**, 58 (10), 6081–6089.
- Saeidnia, S.; Abdollahi, M. Antioxidants: Friends or Foe in Prevention or Treatment of Cancer: The Debate of the Century. *Toxicol. Appl. Pharmacol.* **2013**, 271 (1), 49–63.
- Saini, K. S.; Loi, S.; de Azambuja, E.; Metzger-Filho, O.; Saini, M. L.; Ignatiadis, M.; Dancey, J. E.; Piccart-Gebhart, M. J. Targeting the PI3K/AKT/mTOR and Raf/MEK/ERK Pathways in the Treatment of Breast Cancer. *Cancer Treat. Rev.* **2013**, 39 (8), 935–946.
- Sambade, M. J.; Camp, J. T.; Kimple, R. J.; Sartor, C. I.; Shields, J. M. Mechanism of Lapatinib-Mediated Radiosensitization of Breast Cancer Cells Is Primarily by Inhibition of the Raf>MEK>ERK Mitogen-Activated Protein Kinase Cascade and Radiosensitization of Lapatinib-Resistant Cells Restored by Direct Inhibition of MEK. *Radiother. Oncol.* **2009**, 93 (3), 639–644.
- Sarvaiya, H. A.; Yoon, J. H.; Lazar, I. M. Proteome Profile of the MCF7 Cancer Cell Line: A Mass Spectrometric Evaluation. *Rapid Commun. Mass Spectrom.* **2006**, 20 (20), 3039–3055.
- Segall, M.; Champness, E.; Obrezanova, O.; Leeding, C. Beyond Profiling: Using ADMET Models to Guide Decisions. *Chem. Biodivers.* **2009**, 6 (11), 2144–2151.
- Sekar, P.; Bharti, J. N.; Nigam, J. S.; Sharma, A.; Soni, P. B. Evaluation of p53, HoxD10, and E-Cadherin Status in Breast Cancer and Correlation with Histological Grade and Other Prognostic Factors. *J. Oncol.* **2014**, 2014.
- Sekulic, A.; Migden, M. R.; Oro, A. E.; Dirix, L.; Lewis, K. D.; Hainsworth, J. D.; Solomon, J. A.; Yoo, S.; Arron, S. T.; Friedlander, P. A.; et al. Efficacy and Safety of Vismodegib in Advanced Basal-Cell Carcinoma. *N. Engl. J. Med.* **2012**, 366 (23), 2171–2179.
- Selick, H. E.; Korzekwa, K. R.; Mackarechtschian, K. Drug Rescue by Redesign of ADMET/PK Properties. WO2003034065A2, 2003.
- Sengupta, D.; Chowdhury, K. D.; Sarkar, A.; Paul, S.; Sadhukhan, G. C. Berberine and S Allyl Cysteine Mediated Amelioration of DEN + CCl4 Induced Hepatocarcinoma. *Biochim. Biophys. Acta - Gen. Subj.* **2014**, 1840 (1), 219–244.
- Serafim, T. L.; Oliveira, P. J.; Sardao, V. A.; Perkins, E.; Parke, D.; Holy, J. Different Concentrations of Berberine Result in Distinct Cellular Localization Patterns and Cell Cycle Effects in a

- Melanoma Cell Line. *Cancer Chemother. Pharmacol.* **2008**, *61* (6), 1007–1018.
- Shakil, S. A Simple Click by Click Protocol to Perform Docking. *EXCLI J.* **2013**, *12*, 831–857.
- Shi, J.-H.; Chen, J.; Wang, J.; Zhu, Y.-Y.; Wang, Q. Binding Interaction of Sorafenib with Bovine Serum Albumin: Spectroscopic Methodologies and Molecular Docking. *Spectrochim. Acta. A. Mol. Biomol. Spectrosc.* **2015**, *149*, 630–637.
- Shukla, S.; Chen, Z. S.; Ambudkar, S. V. Tyrosine Kinase Inhibitors as Modulators of ABC Transporter-Mediated Drug Resistance. *Drug Resist. Updat.* **2012**, *15* (1–2), 70–80.
- Singh, A.; Duggal, S.; Kaur, N.; Singh, J. Berberine: Alkaloid with Wide Spectrum of Pharmacological Activities. *J. Nat. Prod. Vol.* **2010**, *3*, 64–75.
- Singh, B. N.; Koyano-Nakagawa, N.; Donaldson, A.; Weaver, C. V.; Garry, M. G.; Garry, D. J. Hedgehog Signaling during Appendage Development and Regeneration. *Genes (Basel).* **2015**, *6* (2), 417–435.
- Songtawee, N.; Gleeson, M. P.; Choowongkomon, K. Computational Study of EGFR Inhibition: Molecular Dynamics Studies on the Active and Inactive Protein Conformations. *J. Mol. Model.* **2013**, *19* (2), 497–509.
- Squires, M. S.; Hudson, E. A.; Howells, L.; Sale, S.; Houghton, C. E.; Jones, J. L.; Fox, L. H.; Dickens, M.; Prigent, S. A.; Manson, M. M. Relevance of Mitogen Activated Protein Kinase (MAPK) and Phosphatidylinositol-3-Kinase/protein Kinase B (PI3K/PKB) Pathways to Induction of Apoptosis by Curcumin in Breast Cells. *Biochem. Pharmacol.* **2003**, *65* (3), 361–376.
- Steg, A. D.; Katre, A. A.; Bevis, K. S.; Ziebarth, A.; Dobbin, Z. C.; Shah, M. M.; Alvarez, R. D.; Landen, C. N. Smoothed Antagonists Reverse Taxane Resistance in Ovarian Cancer. *Mol Cancer Ther* **2012**, *11* (7), 1587–1597.
- Sudha, K. N.; Shakira, M.; Prasanthi, P.; Sarika, N.; Kumar, C. N. Bioinformation Virtual Screening for Novel COX-2 Inhibitors Using the ZINC Database Bioinformation. **2008**, 325–329.
- Sun, Y.; Xun, K.; Wang, Y.; Chen, X. A Systematic Review of the Anticancer Properties of Berberine, a Natural Product from Chinese Herbs. *Anticancer. Drugs* **2009**, *20* (9), 757–769.
- Suter, R.; Marcum, J. A. The Molecular Genetics of Breast Cancer and Targeted Therapy. *Biologics* **2007**, *1* (3), 241–258.
- Tao, Z.; Shi, A.; Lu, C.; Song, T.; Zhang, Z.; Zhao, J. Breast Cancer: Epidemiology and Etiology. *Cell Biochem. Biophys.* **2015**, *72* (2), 333–338.

- Tas, S. Use of Cyclopamine, for the Treatment of Basal Cell Carcinoma and Other Tumors. DE60113733, 2006.
- Tian, S.; Wang, J.; Li, Y.; Li, D.; Xu, L.; Hou, T. The Application of in Silico Drug-Likeness Predictions in Pharmaceutical Research. *Adv. Drug Deliv. Rev.* **2015**, 86, 2–10.
- Tomosaka, H.; Chin, Y.-W.; Salim, A. a; Keller, W. J.; Chai, H.; Kinghorn, a D. Antioxidant and Cytoprotective Compounds from *Berberis Vulgaris* (Barberry). *Phyther. Res.* **2008**, 22 (7), 979–981.
- Vigil, D.; Cherfils, J.; Rossman, K. L.; Der, C. J. Ras Superfamily GEFs and GAPs: Validated and Tractable Targets for Cancer Therapy? *Nat. Rev. Cancer* **2010**, 10 (12), 842–857.
- Vogler, M.; Walczak, H.; Stadel, D.; Haas, T. L.; Genze, F.; Jovanovic, M.; Bhanot, U.; Hasel, C.; Möller, P.; Gschwend, J. E.; et al. Small Molecule XIAP Inhibitors Enhance TRAIL-Induced Apoptosis and Antitumor Activity in Preclinical Models of Pancreatic Carcinoma. *Cancer Res.* **2009**, 69 (6), 2425–2434.
- Wan, P. T.; Garnett, M.; Roe, S. .; Lee, S.; Niculescu-Duvaz, D.; Good, V.; Jones, C.; Marshall, C.; Springer, C.; Barford, D.; et al. Mechanism of Activation of the Raf-Erk Signaling Pathway by Oncogenic Mutations of B-Raf. *Cell(Cambridge,Mass.)* **2004**, 116, 855.
- Wang, Yu; Liu, Q.; Liu, Z.; Li, B.; Sun, Z.; Zhou, H.; Zhang, X.; Gong, Y.; Shao, C. Berberine, a Genotoxic Alkaloid, Induces ATM-Chk1 Mediated G2 Arrest in Prostate Cancer Cells. *Mutat. Res. - Fundam. Mol. Mech. Mutagen.* **2012**, 734 (1–2), 20–29.
- Wang, C.; Wu, H.; Evron, T.; Vardy, E.; Han, G. .; Huang, X. .; Hufeisen, S. .; Mangano, T. .; Urban, D. .; Katritch, V.; et al. Structural Basis for Smoothed Receptor Modulation and Chemoresistance to Anticancer Drugs. *Nat Commun* **2014**, 5, 4355.
- Wang, J.; Mook, R. A.; Lu, J.; Gooden, D. M.; Ribeiro, A.; Guo, A.; Barak, L. S.; Kim Lyerly, H.; Chen, W. Identification of a Novel Smoothed Antagonist That Potently Suppresses Hedgehog Signaling. *Bioorganic Med. Chem.* **2012a**, 20 (22), 6751–6757.
- Wang, J.; Peng, Y.; Liu, Y.; Yang, J.; Ding, N.; Tan, W. Berberine, a Natural Compound, Suppresses Hedgehog Signaling Pathway Activity and Cancer Growth. *BMC Cancer* **2015**, 15 (1), 595.
- Wang, J.; Yang, S.; Cai, X.; Dong, J.; Chen, Z.; Zhang, S.; Cao, H.; Lu, D.; Jin, T.; Nie, Y.; et al. Berberine Inhibits EGFR Signaling and Enhances the Antitumor Effects of EGFR Inhibitors in Gastric Cancer. **2016a**, 7 (46), 76076–76086.
- Wang, L.; Berne, B. J.; Friesner, R. A. Correction for Wang et Al., Ligand Binding to Protein-Binding Pockets with Wet and Dry Regions.

Proc. Natl. Acad. Sci. **2012b**, 109 (23), 9220–9220.

- Wang, L.; Cao, H.; Lu, N.; Liu, L.; Wang, B.; Hu, T.; Israel, D. A.; Peek, R. M.; Polk, D. B.; Yan, F. Berberine Inhibits Proliferation and Down-Regulates Epidermal Growth Factor Receptor through Activation of Cbl in Colon Tumor Cells. *PLoS One* **2013**, 8 (2), e56666.
- Wang, W.; Qin Shen; Hui Liang; Changlong Hua; Yuhui Liu; Fengzhi Li; Qingyong Li. Pharmacokinetic Studies of Novel Berberine Derivatives with Ultra-Performance Liquid Chromatography–tandem Mass Spectrometry. *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* **2016b**, 1031, 172–180.
- Wang, Y. X.; Pang, W. Q.; Zeng, Q. X.; Deng, Z. S.; Fan, T. Y.; Jiang, J. D.; Deng, H. Bin; Song, D. Q. Synthesis and Biological Evaluation of New Berberine Derivatives as Cancer Immunotherapy Agents through Targeting IDO1. *Eur. J. Med. Chem.* **2017**, 143, 1858–1868.
- Wen, C. J.; Wu, L. X.; Fu, L. J.; Shen, D. Y.; Zhang, X.; Zhang, Y. W.; Yu, J.; Zhou, H. H. Preferential Induction of CYP1A1 over CYP1B1 in Human Breast Cancer MCF-7 Cells after Exposure to Berberine. *Asian Pacific J. Cancer Prev.* **2014**, 15 (1), 495–499.
- Wood, E. R.; Truesdale, A. T.; McDonald, O. B.; Yuan, D.; Hassell, A.; Dickerson, S. H.; Ellis, B.; Pennisi, C.; Horne, E.; Lackey, K.; et al. A Unique Structure for Epidermal Growth Factor Receptor Bound to GW572016 (Lapatinib): Relationships among Protein Conformation, Inhibitor off-Rate, and Receptor Activity in Tumor Cells. *Cancer Res.* **2004**, 64 (18), 6652–6659.
- Wrighton, S. A.; Ring, B. J. Inhibition of Human CYP3A Catalyzed 1'-hydroxy Midazolam Formation by Ketoconazole, Nifedipine, Erythromycin, Cimetidine, and Nizatidine. *Pharmaceutical Research.* **1994**, 921–924.
- Wu, C.-P.; V. Ambudkar, S. The Pharmacological Impact of ATP-Binding Cassette Drug Transporters on Vemurafenib-Based Therapy. *Acta Pharm. Sin. B* **2014**, 4 (2), 105–111.
- Xiao, N.; Chen, S.; Ma, Y.; Qiu, J.; Tan, J. H.; Ou, T. M.; Gu, L. Q.; Huang, Z. S.; Li, D. Interaction of Berberine Derivative with Protein POT1 Affect Telomere Function in Cancer Cells. *Biochem. Biophys. Res. Commun.* **2012**, 419 (3), 567–572.
- Xiao, Y.; Tian, C.; Huang, T.; Han, B.; Wang, M.; Ma, H.; Li, Z.; Ye, X.; Li, X. 8-Cetylberberine Inhibits Growth of Lung Cancer in Vitro and in Vivo. *Life Sci.* **2017**, 192 (1), 259–269.
- Xie, J. Implications of Hedgehog Signaling Antagonists for Cancer Therapy. *Acta Biochim. Biophys. Sin. (Shanghai).* **2008**, 40 (7),

670–680.

- Xiong, R.; Zhou, W.; Siegel, D.; Kitson, R. R. A.; Freed, C. R.; Moody, C. J.; Ross, D. A Novel Hsp90 Inhibitor Activates Compensatory Heat Shock Protein Responses and Autophagy and Alleviates Mutant A53T α -Synuclein Toxicity. *Mol Pharmacol* **2015**, 88 (6), 1045–1054.
- Yadav, D. K.; Khan, F.; Negi, A. S. Pharmacophore Modeling, Molecular Docking, QSAR, and in Silico ADMET Studies of Gallic Acid Derivatives for Immunomodulatory Activity. *J. Mol. Model.* **2012**, 18 (6), 2513–2525.
- Yadav, I. S.; Nandekar, P. P.; Shrivastava, S.; Sangamwar, A.; Chaudhury, A.; Agarwal, S. M. Ensemble Docking and Molecular Dynamics Identify Knoevenagel Curcumin Derivatives with Potent Anti-EGFR Activity. *Gene* **2014**, 539 (1), 82–90.
- Yamashita, F.; Hashida, M. In Silico Approaches for Predicting ADME Properties of Drugs. *Drug Metab. Pharmacokinet.* **2004**, 19 (5), 327–338.
- Yan, K.; Zhang, C.; Feng, J.; Hou, L.; Yan, L.; Zhou, Z.; Liu, Z.; Liu, C.; Fan, Y.; Zheng, B.; et al. Induction of G1 Cell Cycle Arrest and Apoptosis by Berberine in Bladder Cancer Cells. *Eur. J. Pharmacol.* **2011**, 661 (1–3), 1–7.
- Yang, Y.-S.; Li, Q.-S.; Sun, S.; Zhang, Y.-B.; Wang, X.-L.; Zhang, F.; Tang, J.-F.; Zhu, H.-L. Design, Modification and 3D QSAR Studies of Novel 2,3-dihydrobenzo[b][1,4]dioxin-Containing 4,5-Dihydro-1H-Pyrazole Derivatives as Inhibitors of B-Raf Kinase. *Bioorg. Med. Chem.* **2012**, 20 (20), 6048–6058.
- Yates, L. L.; Górecki, D. C. The Nuclear Factor-kappaB (NF-kappaB): From a Versatile Transcription Factor to a Ubiquitous Therapeutic Target. *Acta Biochim. Pol.* **2006**, 53 (4), 651–662.
- Yu, P.; Li, D.; Ni, J.; Zhao, L.; Ding, G.; Wang, Z.; Xiao, W. Predictive QSAR Modeling Study on Berberine Derivatives with Hypolipidemic Activity. *Chem. Biol. Drug Des.* **2017**, 1, 1–7.
- Zhang, H.; Shan, Y.; Wu, Y.; Xu, C.; Yu, X.; Zhao, J.; Yan, J.; Shang, W. Berberine Suppresses LPS-Induced Inflammation through Modulating Sirt1/NF- κ B Signaling Pathway in RAW264.7 Cells. *Int. Immunopharmacol.* **2017**, 52 (June), 93–100.
- Zhang, J.; Cao, H.; Zhang, B.; Cao, H.; Xu, X.; Ruan, H.; Yi, T.; Tan, L.; Qu, R.; Song, G.; et al. Berberine Potently Attenuates Intestinal Polyps Growth in ApcMin Mice and Familial Adenomatous Polyposis Patients through Inhibition of Wnt Signalling. *J. Cell. Mol. Med.* **2013**, 17 (11), 1484–1493.
- Zhang, R.; Qiao, H.; Chen, S.; Chen, X.; Dou, K.; Wei, L.; Zhang, J. Berberine Reverses Lapatinib Resistance of HER2-Positive

- Breast Cancer Cells by Increasing the Level of ROS. *Cancer Biol. Ther.* **2016**, 17 (9), 925–934.
- Zhang, S.; Kumar, K.; Jiang, X.; Wallqvist, A.; Reifman, J. DOVIS: An Implementation for High-Throughput Virtual Screening Using AutoDock. *BMC Bioinformatics* **2008**, 9, 126.
- Zhang, X.; Harrington, N.; Moraes, R. C.; Wu, M.-F.; Hilsenbeck, S. G.; Lewis, M. T. Cyclopamine Inhibition of Human Breast Cancer Cell Growth Independent of Smoothened (Smo). *Breast Cancer Res. Treat.* **2009**, 115 (3), 505–521.
- Zhou, Y.; He, P.; Liu, A.; Zhang, L.; Liu, Y.; Dai, R. Drug-Drug Interactions between Ketoconazole and Berberine in Rats: Pharmacokinetic Effects Benefit Pharmacodynamic Synergism. *Phyther. Res.* **2012**, 26 (5), 772–777.
- Zhu, L.; Zhang, D.; Zhu, H.; Zhu, J.; Weng, S.; Dong, L.; Liu, T.; Hu, Y.; Shen, X. Berberine Treatment Increases Akkermansia in the Gut and Improves High-Fat Diet-Induced Atherosclerosis in Apoe^{-/-} mice. *Atherosclerosis* **2018**, 268, 117–126.
- Zovko Končić, M.; Kremer, D.; Karlović, K.; Kosalec, I. Evaluation of Antioxidant Activities and Phenolic Content of *Berberis Vulgaris* L. and *Berberis Croatica* Horvat. *Food and Chemical Toxicology.* **2010**, 48, 2176–2180.
- Zubrilov, I.; Sagi-Assif, O.; Izraely, S.; Meshel, T.; Ben-Menahem, S.; Ginat, R.; Pasmanik-Chor, M.; Nahmias, C.; Couraud, P.-O.; Hoon, D. S. B.; et al. Vemurafenib Resistance Selects for Highly Malignant Brain and Lung-Metastasizing Melanoma Cells. *Cancer Lett.* **2015**, 361 (1), 86–96.

LIST OF PUBLICATIONS

Jabbarzadeh Kaboli, P.; Rahmat, A.; Ismail, P.; Ling, K-H. (2014). Targets and mechanisms of berberine, a natural drug with potential to treat cancer with special focus on breast cancer. *European Journal of Pharmacology*, 740, 584–595. Elsevier.

DOI: 10.1016/j.ejphar.2014.06.025

[ISI Impact Factor: 2.90 / Q2]

Jabbarzadeh Kaboli, P.; Rahmat, A.; Ismail, P.; Ling, K-H. (2015). MicroRNA-based therapy and breast cancer: A comprehensive review of novel therapeutic strategies from diagnosis to treatment. *Pharmacological Research*, 97, 104–121. Elsevier.

DOI: 10.1016/j.phrs.2015.04.015

[ISI Impact Factor: 4.49 / Q1]

Jabbarzadeh Kaboli, P.; Bazrafkan, M.; Ismail, P.; Ling, K-H. (2017). Molecular Modelling of Berberine Derivatives as Inhibitors of Human Smoothed Receptor and Hedgehog Signalling Pathway Using a Newly Developed Algorithm on Anti-Cancer Drugs, *Recent Patents on Anti-Cancer Drug Discovery*, 12 (4): 384-400. Bentham Science.

DOI: 10.2174/1574892812666170929131247

[ISI Impact Factor: 2.96 / Q2]

Jabbarzadeh Kaboli, P.; Ismail, P.; Ling, K-H. (2018). Molecular Modelling, Dynamics Simulations, and Binding Efficiency of Berberine Derivatives: A New Group of RAF Inhibitors for Cancer Treatment. *PLOS ONE*.

DOI: 10.1371/journal.pone.0193941

[ISI Impact Factor: 2.81 / Q1]

Jabbarzadeh Kaboli, P.; Leong, P.; Ismail, P.; Ling, K-H. (Submitted). Anti-tumor Activity of berberine as Inhibitor of EGFR, ERK1/2, P38 and AKT in MDA-MB231 and MCF-7 Breast Cancer Cells Using Molecular Modelling and *in vitro* Approaches.

Jabbarzadeh Kaboli, P.; Ismail, P.; Ling, K-H. (2017). Berberine Derivatives as Multi-Kinase Inhibitors of MAP Kinase and PI3K Pathways: Molecular Modelling, Molecular Dynamics Simulations, and *in vitro* Evaluation. *FASEB J.* 31: 671.12-671.12.

[ISI Impact Factor: 5.50 / Q1]

Jabbarzadeh Kaboli, P.; Bazrafkan, M.; Ismail, P.; Ling, K.-H. **(2017).** A Novel One-Click Automated Drug Discovery tool: A New Approach Based on Binding Efficiency Indices. *FASEB J.* 31: 818.7-818.7.

[ISI Impact Factor: 5.50 / Q1]

Jabbarzadeh Kaboli, P.; Ismail, P.; Ling, K.-H. **(2016).** Molecular modelling and molecular dynamics simulations of a new berberine derivative as a dual RAF inhibitor for treatment of drug resistance acquired in melanoma. **(Oral presentation; Presenter)** "Pharmacology 2016, London", British Pharmacological Society, 13-15th December, London, UK.

Jabbarzadeh Kaboli, P.; Ismail, P.; Ling, K.-H. **(2017).** Molecular modelling, molecular dynamics simulations and *in vitro* evaluation of Berberine as an inhibitor of MAP kinase and PI3K pathway. **(Poster presentation; Presenter)** "EACR-AACR-SIC Special Conference 2017: The Challenges of Optimizing Immuno and Targeted Therapies: From Cancer Biology to the Clinic", 25th June 2017, Florence, Italy.

Jabbarzadeh Kaboli, P.; Bazrafkan, M.; Ismail, P.; Ling, K.-H. **(2017).** Molecular modelling of berberine derivatives as inhibitors of smoothened receptor using a new drug discovery script for a large numbers of compounds. **(Poster presentation; Presenter)** "EACR-AACR-SIC Special Conference 2017: The Challenges of Optimizing Immuno and Targeted Therapies: From Cancer Biology to the Clinic", 25th June 2017, Florence, Italy.

Ismail, P.; **Jabbarzadeh Kaboli, P.;** Ling, K.-H. **(2016).** "New Anti-Cancer Drugs against MicroRNA and MAP Kinase Pathway", *Exhibition of Invention, Research, and Innovation (PRPI 2016)*, Universiti Putra Malaysia (UPM), **Silver Medal.**