



UNIVERSITI PUTRA MALAYSIA

***ISOLATION OF TRANSKETOLASE GENE, SUBCELLULAR
LOCALIZATION, AND TRANSKETOLASE PROTEIN STRUCTURE
STUDIES OF SUGARCANE (*Saccharum officinarum*. L)***

NAHID KALHORI

FS 2013 71



**ISOLATION OF TRANSKETOLASE GENE, SUBCELLULAR
LOCALIZATION, AND TRANSKETOLASE PROTEIN STRUCTURE
STUDIES OF SUGARCANE (*Saccharum officinarum*. L)**

By

NAHID KALHORI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirement for the Degree of Master of Science**

May 2013

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within thesis for non-commercial purpose from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright© Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**ISOLATION OF TRANSKETOLASE GENE, SUBCELLULAR
LOCALIZATION, AND TRANSKETOLASE PROTEIN STRUCTURE
STUDIES OF SUGARCANE (*Saccharum officinarum*. L)**

By

NAHID KALHORI

May 2013

Chairman: Rosimah Nulit, PhD

Faculty: Science

This study focused on isolation, identification, subcellular localisation, protein structure prediction of transketolase enzyme from sugarcane, *Saccharum officinarum*. Pentose phosphate pathway is composed of two functionally-connected phases, the oxidative and non-oxidative phase. Both phases are catalysed by a series of enzymes. One of these enzymes is transketolase which play an important role in non-oxidative phase of pentose phosphate pathway (PPP). Synthesis of sugar phosphate intermediate is the main role of this enzyme in non-oxidative phase of pentose phosphate pathway. Transketolase transfer two carbon units from fructose-6-phosphate to erythrose-4-phosphate and convert two carbon fragments from glyceraldehyde-3-phosphate (G3P) to ketose of xylulose-5-phosphate. Erythrose-4-phosphate then enters the shikimate pathway that produced many secondary metabolites such as aromatic amino acids, lignin and flavonoid. Transketolase also play important role in photosynthesis and glycolysis. Although the contribution of transketolase in plant system is important, study of this enzyme is still limited. Until now, *TKT* genes had been isolated only from seven plants so far, thus this leads to

the first objective to isolate *TKT* gene from sugarcane to compare its identity with other organisms. Unlike bacteria, fungi and all animals, PPP is complete in the cytosol of these living system and all enzymes of this pathway localised in the cytosol. However, in plant system, the first phase of pentose phosphate pathway is complete in the cytosol of plant and the sub localisation for non-oxidative pentose phosphate pathway in still remains unclear. Thus, the second objective is to determine its subcellular localization of sugarcane transketolase. The final objective is to predict modelling protein structure of sugarcane transketolase. The isolation of *TKT* by was done by RT- PCR, followed by cloning and sequencing. The present study has isolated 2327 bp length of sugarcane *TKT*. Similarity studies using CLUSTALW2 revealed that sugarcane *TKT* gene showed highest identity with *Zea mays* (99%). Classification analysis revealed that both plants are in the same family, Poaceae. However, the identity of sugarcane *TKT* compared with other plants species are low (<65 %) due to these plants are only in the same phylum, Streptophyte. Moreover, sugarcane *TKT* showed lower similarity (<60%) compared with bacteria, yeast and animals because these organisms from different kingdom, animalia. Analysis by Target P 1.1 followed by Chloro P revealed that sugarcane transketolase is localised in the chloroplast which is 85 amino acids are plant plastid target sequence. The nucleotide sequence of sugarcane *TKT* was converted to amino acid sequence using Expasy.org, Phyre2 and YASARA to predict secondary, tertiary structure and the quality of protein structure of sugarcane transketolase. This present study revealed that sugarcane transketolase protein is 100% similarity with Thiamin diphosphate-binding fold, from *Zea mays* (d1ltz). In conclusion, sugarcane transketolase was successfully isolated and was found plastidic in plant system. This led to conclusion that the OPPP is incomplete in the

cytosol of sugarcane. This present study also similarity sequence of sugarcane TKT closely related with classification or taxonomy level of plants and other organisms. This present study also found that sugarcane transketolase protein is 100% similarity with Thiamin diphosphate-binding fold, from *Zea mays* (d1itz).



Abstrak tesis yang dikemukakan kepada Senate Universiti Putra Malaysia sebagai memenuhi keperluan ijazah Master Sains

**PEMENCILAN, PENENTUAN PENEMPATAN SUBSELLULAR DAN
KAJIAN STRUKTUR PROTEIN TRANSKETOLASE DARIPADA TEBU,
(*Saccharum officinarum*. L)**

Oleh

NAHID KALHORI

Mei 2013

Pengerusi: Rosimah Nuli, PhD

Fakulti: Sains

Kajian ini memfokuskan kepada pemencilan, pengenalpastian, penentuan penempatan subsellular dan kajian struktur protein enzim transketolase daripada tebu, *Saccharum officinarum*. Tapak Jalan Pentosa Fosfat terdiri daripada dua fasa, iaitu, oksidatif dan non-oksidatif. Kedua-dua fasa ini dikawal oleh satu siri enzim. Salah satunya adalah transketolase yang memainkan peranan penting dalam fasa non-oksidatif di mana menghasilkan beberapa pengantara gula fosfat. Transketolase memindahkan dua unit karbon daripada fruktosa-6-fosfat kepada eritrosa-4-fosfat dan menukarkan dua fragmen karbon daripada gliseraldehid-3-fosfat kepada kumpulan ketosa xylulosa-5-fosfat, Eritrosa-4-fosfat akan memasuki tapak jalan shikimate di mana tapak jalan ini merupakan tapak jalan yang penting dalam penghasilan metabolit sekunder dalam sistem tumbuhan seperti, aromatik asid

amino, lignin dan flavonoid. Transketolase juga memainkan peranan penting dalam proses fotosintesis dan glikolisis. Walaupun peranan transketolase dalam tumbuhan adalah penting, kajian mengenai enzim ini dalam tumbuhan adalah terlalu terhad. Sehingga kini, pemencilan gen transketolase telah dilakukan hanya dari tujuh spesies tumbuhan. Maka, objektif pertama kajian ini ialah untuk memencilkan gen TKT daripada tebu, dan untuk membandingkan persamaannya jujukan nukleotida transketolase dengan organisma lain. Berbeza dengan bakteria, kulat dan haiwan, tapak jalan pentosa fosfat adalah lengkap di dalam sitosol. Walaubagaimanapun, dalam sistem tumbuhan, fasa pertama adalah di dalam sitosol, tetapi fasa kedua masih dalam kajian. Jadi, objektif kedua adalah untuk menentukan penempatan subselular transketolase dalam pokok tebu. Manakala, objektif ketiga adalah untuk meramalkan struktur protin transketolase. Pemencilan TKT daripada pokok tebu dilakukan dengan RT-PCR, diikuti dengan pengklonan dan penjujukan nukleotida. Kajian telah berjaya memencilkan jujukan nukleotida TKT sebanyak 2327 bp. Analisa persamaan menggunakan CLUSTALW2 mendapati jujukan nukleotida TKT daripada pokok tebu menunjukkan persamaan yang tertinggi (99%) dengan transketolase dari pokok jagung, *Zea mays*. Ini adalah kerana analisis klasifikasi tumbuhan mendapati kedua-dua tumbuhan adalah daripada famili yang sama, Poaceae. Walaubagaimanapun, persamaan jujukan nukleotida TKT tebu adalah rendah (<65%) berbanding dengan spesie lain memandangkan tumbuhan ini berada dalam filum yang sama, Streptophyte. Ia menunjukkan persamaan jujukan nukleotida terendah (<60%) terhadap bakteria, yis dan haiwan kerana organisma-organisma ini adalah daripada alam yang berbeza. Analisa menggunakan Target P 1.1 dan Chloro P mendapati transketolase daripada pokok tebu adalah kloroplastik dimana sebanyak 85 jujukan asid amino adalah jujukan sasaran plastid tumbuhan.

Jujukan nukleotida TKT pokok tebu telah diterjemahkan kepada jujukan asid amino menggunakan perisian Expasy.org, Phyre 2, dan YASARA untuk meramal struktur sekunder, tertier dan kualiti struktur protin transketolase pokok tebu. Kajian ini menunjukkan struktur protin transketolase pokok tebu adalah 100% sama seperti lipatan ikatan thiamin difosfat daripada *Zea mays*. Sebagai kesimpulan, jujukan nukleotida *TKT* pokok tebu telah berjaya dipencilkan dan gen ini adalah kloroplastik. Oleh itu, kajian ini mendapati bahawa tapak jalan pentose fosfat adalah tidak lengkap dalam sitosol tumbuhan. Kajian ini mendapati persamaan jujukan nukleotida *TKT* adalah berkait rapat dengan klasifikasi dan taksonomi tumbuhan dan organisma lain. Struktur protin transketolase adalah 100% sama dengan lipatan ikatan thiamin difosfat daripada *Zea mays*.

ACKNOWLEDGEMENTS

I would like to express my deepest appreciation to my supervisor, Dr. Rosimah Nulit from Department of Biology of Faculty of Science. Without her, I may not have the chance learning and dealing with the techniques of molecular biology. Despite of the busy moments Dr. Rosimah has, she shows her kindness and care towards the progress of my project and takes time in editing my thesis paper. Apart from that, her hard work in arranging the booking of the apparatus and chemical, ordering primer that I designed. Thank you Dr. Rosimah for giving sincere dedication in teaching the lab work and techniques as well as giving moral support to me, saying that I can do. I also wish to thank Associate Professor Dr. Rusea Go, as co-supervisor, who had given lots of advice throughout my study. This is a great opportunity to express my honest gratitude to University of Putra Malaysia, Faculty of Science and Department of Biology.

Finally, special thanks to my parents and my brother for their love support and understanding while I studied far away from my country. No other people love me more than them. They are the main reason to push me forward and create the level of success that I have.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Rosimah Nulit, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Rusea Go, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at University of Putra Malaysia or at any other institution.

NAHID KALHORI

Date: 30 MAY 2013




TABLE OF CONTENTS

	ABSTRACT	Page
		ii
	ABSTRAK	v
	ACKNOWLEDGMENTS	viii
	APPROVAL	ix
	DECLARATION	xi
	LIST OF TABLES	xiv
	LIST OF FIGURES	xv
	LIST OF ABBREVIATION	xvii
CHAPTER		
1	INTRODUCTION	1
1.1	Problem Statements	2
1.2	Objectives	3
2	LITRETURE REVIEW	4
2.1	The Oxidative Pentose Phosphate Pathway	4
2.2	The Roles and Importance of The Oxidative Pentose Phosphate Pathway	11
2.3	Coordination of Pentose Phosphate Pathway with Glycolysis, Calvin Cycle and Shikimate Pathway	12
2.4	Transketolase	16
2.4.1	The significant of Transketolase in plants	19
2.4.2	Subcellular localization of Transketolase in Plants	20
2.4.3	Isolation of Transketolase from Various Organisms	22
2.5	Protein Structure Prediction	23
2.5.1	Homology Modeling	25
2.5.2	Ramachandran Plot	26
2.6	Verify 3D profile Analysis	29
2.7	Structure Prediction of Transketolase	29
3	MATERIALS AND METHODS	31
3.1	Plant material	31
3.2	Total RNA isolation	31
3.3	The Amplification, cloning and sequencing of Sugarcane <i>TKT</i>	32
3.3.1	Amplification of Sugarcane	32
3.3.2	Purification of PCR products	34
3.3.3	Cloning of the Transketolase into Vector pJET1.2	34
3.3.4	Plasmid Isolation	35
3.3.5	Subcellular Localization of Transketolase	36
3.4	Over expression and Purification Sugarcane Transketolase	37

	Recombinant Plasmid	
3.4.1	Transformation of Recombinant Plasmid in Non-expression Host	37
3.4.2	Transformation of Recombinant Plasmid in Expression Host	38
3.4.3	Induction of Expression Constructs	38
3.4.4	Protein Purification	38
3.4.5	SDS-PAGE	40
3.5	Protein Structure Prediction of Sugarcane Transketolase	41
3.5.1	Identification of the Amino Acid Sequence	41
3.5.2	Protein Structure Comparison	42
4	RESULT AND DISCUSSION	43
4.1	Isolation, Cloning and Sequencing of <i>TKT</i> Sugarcane	43
4.2	Similarity Study of <i>TKT</i> Nucleotide Sequence of Sugarcane with other Organisms	46
4.3	Subcellular Localization of Sugarcane <i>TKT</i>	49
4.4	Construction of Recombinant DNA Sugarcane <i>TKT</i>	50
4.5	Protein Structure Prediction	55
4.5.1	Protein Structure Comparison between Target Sequence and Template Sequence (Homology Modelling)	59
4.5.2	Secondary Structure Prediction	62
4.5.3	Ramachandran Plot Input-atom-only	63
4.6	Comparison of Sugarcane Transketolase Protein Sequences with Transketolase Protein from other Organisms	67
5	CONCLUSION	69
5.1	Future Work	70
	REFERENCES	72
	APPENDICES	80
	BIODATA OF STUDENT	112
	LIST OF PUBLICATIONS	113

LIST OF TABLES

Table		Page
2.1	Subcellular localization of transketolase	20
4.1	Similarities study of nucleotide sequence of sugarcane <i>TKT</i> with other organisms.	47
4.2	Output of SuperPose between the target sequence and top match model sequence.	61
4.3	Comparison of sugarcane transketolase protein with other organism's protein	67

LIST OF FIGURES

Figures	Page
2.1	The Oxidative Pentose Phosphate Pathway 9
2.2	The Inter-connection between Glycolysis, Calvin cycle, Pentose Phosphate Pathway and Shikimate Pathway. 13
2.3	The first reaction of transketolase in non-oxidative phase of pentose phosphate pathway 17
2.4	The second reaction of transketolase in non-oxidative phase of pentose phosphate pathway 17
2.5	Ramachandran Plot 27
2.6	Ball and stick model of a dipeptide 28
4.1a	Sugarcane transketolase was amplified by RT-PCR 43
4.1b	Restriction digestion of recombinant DNA <i>pJet1.2-TKT</i> with double <i>Bgl</i> II. 44
4.2	Nucleotide and deduced amino acid sequence of the functional, full length, <i>Saccharum officinarum</i> TKT cDNA 45
4.3	Phylogenetic analysis of identity of several plants with <i>Escherichia coli</i> as an out group 48
4.4a	Target P server shows sugar cane transketolase takes place in Chloroplast 49
4.4b	ChloroP server shows transketolase with 85 amino acids is in Chloroplast 49
4.5a	Construct of pET32a-TKT containing transketolase cDNA of sugarcane gene with expression plasmid vector 51
4.5b	Recombinant plasmid of <i>TKT</i> and <i>pet32a</i> (+) 52
4.5c	Restriction digestion of the constructed plasmid vectors 53
4.5d	Over expression of the recombinant protein transketolase from sugarcane transketolase protein was over-expressed in the soluble fraction of the expression host 54
4.6	Alignment of <i>S. officinarum</i> transketolase protein sequences 56

shown the top 6 ranking models were chosen from top 20 hits by using the Phyre2 server

4.7	Ribbon presentation for top model named (d1itza) that covered the higher similarity with <i>S. officinarum</i>	57
4.8	Ribbon representation of the three-dimensional structure of target protein transketolase from <i>S. officinarum</i> .	58
4.9	Alignment file between the query target protein sequence and template d1itza protein sequence in ALI format	59
4.10	MolScript superposition of the target PDB transketolase from <i>S. officinarum</i> and top match model of PDB 1itz	60
4.11	Secondary structure of transketolase <i>S. officinarum</i>	62
4.12	Ramachandran Plot from PROCHECK	64
4.13	The result of Ramachandran Plot from PROCHECK	65
4.14	Verify 3D result for target sequence model of transketolase from <i>S. officinarum</i>	66

LIST OF ABBREVIATIONS

ADP	Adenosine Diphosphate
ATP	AdenosineTriphosphate
BLAST	Basic Local Alignment Tool
BNL Buffer	Agarose gel lysis Buffer
bp	Basepair
C	Carbon
cDNA	Complementary DNA
CO ₂	Carbon Dioxide
ddH ₂ O	Distilled water
DNA	Deoxyribonucleic acid
FAD	Flavin Adenine Dinucleotide
G	Gravitational Force
G6PDH	glucose-6-phosphate dehydrogenase
IPTG	Isopropyl β-D-1-thiogalactopyranoside
kDa	Kilo Daltons
LB	Luria Bertani

M	Molar
Mg	Milli gram
MgCl ₂	Magnesium chloride
ml	Milli Litter
μl	Micro Liter
NADP	Nicotinamide Adenine Dinucleotide Phosphate
NADPH	Reduced Nicotinamide Adenine Dinucleotide Phosphate
NMR	Nuclear Magnetic Resonance
OD	Optical Density
OPPP	Oxidative Pentose Phosphate pathway
PCR	Polymerase Chain Reaction
PDB	Protein Data Bank
PDF	Portable Document Format
PEP	Phosphoenolpyruvate
pH	Potentiometric Hydrogen Ion Concentration
Phyre2	Protein Homology

PPP	Pentose Phosphate Pathway
PSI-BLAST	Position-Specific Iterated BLAST
RMSD	Root Mean Square Deviation
RLC	Contains guanidine hydrochloride
RLT Buffer	Contains guanidine thiocyanate
RNase-free water	RNA treatment
RPE	Ribulose-5-phosphate-3-epimerase
RPI	Ribose-5-phosphate isomerase
PRPP	Phosphoribosyl pyrophosphate
RPPP	Reductive Pentose Phosphate Pathway
RT-PCR	Real Time Polymerase Chain Reaction
RW1	Contains ethanol
S	Seconds
S7P	Sedoheptulose-7-Phosphate
SDS-PAGE	Sodium dodecyl sulfate polyacrylamide gel electrophoresis
TBE	Tris hydroxyl methyl amino methane
ThDP	Thiamine diphosphate

TAL	Transaldolase
TKT	Transketolase
TPP	Thiamine Pyro Phosphate
UK	United Kingdom
UPM	Universiti Putra Malaysia
USA	United States of America
UV	Ultraviolet
V	Voltage
X	Time
X5P	Xylulose-5-Phosphate
YASARA	Yet Another Scientific Artificial Reality Application
3D	Three Dimensional
6PG	6-phosphogluconolactonase
6PGDH	6-phosphogluconate dehydrogenase
°C	Degree Celsius
%	Percentage

CHAPTER 1

INTRODUCTION

Sugarcane is type of C₄ plants species in which photosynthetic rate is twice as efficient as the C₃ plants species. Also sugarcane is the most important source of sucrose in world history. The oxidative pentose phosphate pathway (PPP) has crucial role in the metabolism of carbohydrates *via* the oxidation of glucose-6-phosphate. The pathway consists of oxidative and non-oxidative phase. The oxidative phase produces ribulose-5-phosphate by the oxidation of glucose-6-phosphate. The enzymes of the irreversible oxidation phase are successively glucose-6-phosphate dehydrogenase (G6PDH), 6-phosphogluconolactonase (6PG) and 6-phosphogluconate dehydrogenase (6PGDH). The second phase is the reversible non-oxidative phase which is series of inter conversion between 3-, 4-, 5-, 6- and 7-carbons sugars catalysed by the enzymes ribose-5-phosphate isomerase (RPI), ribulose-5-phosphate-3-epimerase (RPE) and transketolase (TKT) and transaldolase (TAL). All enzymes of the OPPP apart from TAL are also part of the reductive pentose phosphate pathway (RPPP) or Calvin cycle. The most important role of the first phase is providing 50%-60% of the required NADPH, a major by reduction of NADP⁺.

The second phase of the OPPP produces ribose-5-phosphate and erythrose-4-phosphate. Ribose-5-phosphate synthesis the nucleotides and nucleic acids – RNA, DNA and nucleotides coenzymes such as ATP, FAD, NADH and NADPH. Erythrose-4-phosphate and phosphoenolpyruvate (PEP) from glycolysis to form the

first substrate of the shikimic pathway which is chorismate. This important substrate provides aromatic amino acids and many aromatic secondary metabolites such as flavonoids, indole acetate, UV light protectants and lignin. These all metabolism have crucial role in the interaction of plant with environment (Nulit 2010).

1.1 Problem Statements

Unlike bacteria, fungi and animal, the number of *TKT* gene had been isolated from plant species is small, only seven *TKT* genes had been isolated from plants which are *Zea mays* (AY148193), *Solanum tuberosum* (Z50099), *Spinacia oleracea* (L76554), *Capsicum annum* (Y15781), *Arabidopsis thaliana* (AY091094), *Persicaria tinctoria* (AB066206), *Craterostigma plantagineum* (Z46646), *Platanus acerifolia* (AJ427413). In addition, the subcellular localization of PPP in bacteria, fungi and animal are in the cytosol. However, in plants, the subcellular localization is not clear. Therefore, another approach to resolve the compartmentation of the PPP genes in plants is molecular approach. According to Garnier *et al.* (2003) as more genome can be sequenced and published, the understanding of the genetic capacity of plants system to catalyse the reactions involved in the PPP could be obtained. In addition, the increasing availability of whole-genome transcript profiles under different physiological and developmental conditions will allow us to determine the relationships between the expressions of the specific PPP isoenzymes, and to establish the contribution of such isoenzymes to particular physiological processes. Therefore, a question regarding a complete PPP in the cytosol and plastid of plant could be answered.

1.2 Objectives of Study

Thus, the objectives of my study are:

- To isolate and clone transketolase gene from sugarcane (*Saccharum officinarum*) and compare DNA sequence of sugarcane transketolase gene with other plants, bacteria, fungi, animal and human being.
- To determine the subcellular localisation of transketolase gene in sugarcane
- To predict the structure of sugarcane transketolase protein

REFERENCES

- Akbay, E., Nuray Ulusu, N., Töröner, F., Ayvaz, G., Taneri, F., Aktürk, M., Arslan, M. & Karasu, Ç. (2004). Effects of rosiglitazone treatment on the pentose phosphate pathway and glutathione-dependent enzymes in liver and kidney of rats fed a high-fat diet. *Current Therapeutic Research*, 65(1): 79-89.
- Baker, L. J., Dorocke, J. A., Harris, R. A. & Timm, D. E. (2001). The Crystal Structure of Yeast Thiamin Pyrophosphokinase. *Structure*, 9(6): 539-546.
- Barrett, M. P. (1997). The pentose phosphate pathway and parasitic protozoa. *Parasitology Today*, 13(1): 11-16.
- Batz, Logemann, E., Reinold, S. & Hahlbrock, K. (1998). Extensive reprogramming of primary and secondary metabolism by fungal elicitor or infection in parsley cells. *Biol. Chem*, 379: 1127–1135.
- Baxevanis, A. D. & Ouellette, B. F. F. (2005). Bioinformatics: A Practical Guide to the Analysis of Genes and Proteins, third edition. Wiley, ISBN 0-471: 47878-47874.
- Beevers, H. (1956). Intermediates of the Pentose Phosphate Pathway as Respiratory Substrates. *Plant Physiol*, 31(5): 339-347.
- Berg, J. M., Deis, F. H., Gerber, N. C., Gumpert, R. & Koeppe, R. E. (2011). *Biochemistry Student Companion*: W. H. Freeman.
- Bernacchia, G., Salamini, F. & Bartels, D. (1996). Molecular Characterization of the Rehydration Process in the Resurrection Plant *Craterostigma plantagineum*. *Plant Physiol*, 111(4): 1043-1050.
- Bernacchia, G., Schwall, G., Lottspeich, F., Salamini, F. & Bartels, D. (1995). The transketolase gene family of the resurrection plant *Craterostigma plantagineum*: differential expression during the rehydration phase. *EMBO J*, 14(3): 610-618.
- Bernardi, R., Fontana, F., Salvini, M., Scala, A. & Durante, M. (2006). Isolation of differentially expressed transcript after cerato-platanin treatments in *Platanus acerifolia*. *EMBL/GENEBANK/DBJ databases*.
- Bhalla, P. L., Singh, M. B. & Malik, C. P. (1981). Characterization of Pentose Phosphate Pathway in Embryo Suspensor of *Tropaeolum majus*. *Biochemie und Physiologie der Pflanzen*, 176(8): 789-792.
- Bolanos, J. P., Delgado-Esteban, M., Herrero-Mendez, A., Fernandez-Fernandez, S. & Almeida, A. (2008). Regulation of glycolysis and pentose-phosphate

pathway by nitric oxide: impact on neuronal survival. *Biochim Biophys Acta*, 1777(7-8): 789-793.

Bouvier., D'harlingue. A., Suire. C., Backaus. R.A.& Camara.B. (1998). Dedicated roles of plastid transketolase during the early onset of isoprenoid biogenesis in pepper fruits. *Plant Physiol*, 117: 1423-1431.

Campbell, N. A. (2009). *Biology: Concepts & Connections*: Pearson/Benjamin Cummings.

Campbell, N. A., Williamson, B.& Heyden, R. J. (2006). *Biology: Exploring Life*: Pearson Prentice Hall.

Champe, P. C.& Harvey, R. A. (1994). *Lippincott's Illustrated Reviews: Biochemistry*: Lippincott.

Christie, P. J., Alfenito., M. R.& Walbot., V. (1994). Impact of lowtemperature stress on general phenylpropanoid and anthocyanin pathways: Enhancement of transcript abundance and anthocyanin pigmentation in maize seedlings. *Planta* 194: 541-549.

Datta, A. G.& Racker, E. (1961). Mechanism of action of transketolase. I. Properties of the crystalline yeast enzyme. *J. Biol. Chem.*, 236: 617-623.

De La Haba, G., Leder, I. G.& Racker, E. (1955). Crystalline transketolase from bakers' yeast: isolation and properties. *J Biol Chem*, 214(1): 409-426.

De Meillon., Small J.G.C& H.A., V. d. V. (1990). The respiratory metabolism of *Strelitzia juncea* Ait. seeds: the effect of dormancy release through oxygen incubation of the seeds on the activity of glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase. *J. Exp. Bot*, 41: 709-714.

Debnam, P. M., Fernie, A. R., Leisse, A., Golding, A., Bowsher, C. G., Grimshaw, C., Knight, J. S.& Emes, M. J. (2004). Altered activity of the P2 isoform of plastidic glucose 6-phosphate dehydrogenase in tobacco (*Nicotiana tabacum* cv. Samsun) causes changes in carbohydrate metabolism and response to oxidative stress in leaves. *Plant J*, 38(1): 49-59.

Dennis, D. T.& Blakeley, S. D. (2000). Carbohydrate metabolism. In B Buchanan, W Gruissem, RL Jones, eds, *Biochemistry and Molecular Biology of Plants*. American Society of Plant Physiologists, Rockville, MD: 630–675.

Dey, P. M.& Harborne, J. J. B. (1997). *Plant Biochemistry*: Acad. Press.

Drenth.J. (1999). Principles of protein X-ray crystallography. *Springer-Verlag*, New York.

- Emanuelsson, O., Nielsen, H. & von Heijne, G. (1999). ChloroP, a neural network-based method for predicting chloroplast transit peptides and their cleavage sites. *Protein Sci*, 8(5): 978-984.
- Emes, M. & Fowler, M. (1979). The intracellular location of the by a set of enzymes of nitrate assimilation in the apices of seedling pea roots. *Planta* 144: 249–253.
- Flechner, A., Dressen, U., Westhoff, P., Henze, K., Schnarrenberger, C. & Martin, W. (1996). Molecular characterization of transketolase (EC 2.2.1.1) active in the Calvin cycle of spinach chloroplasts. *Plant Mol Biol*, 32(3): 475-484.
- Fraenkel, D. G. (1987). Glycolysis, pentose phosphate pathway, and Entner-Doudoroff pathway. *American Society for Microbiology, Washington, D.C.*: 142-150.
- Fullam, E., Pojer, F., Bergfors, T., Jones, T. A. & Cole, S. T. (2012). Structure and function of the transketolase from *Mycobacterium tuberculosis* and comparison with the human enzyme. *Open Biol*, 2(1): 110026.
- Gahan, P. B. (1984). *Plant histochemistry and cytochemistry: an introduction*: Academic Press.
- Garnier, T., Eiglmeier, K., Camus, J. C., Medina, N., Mansoor, H., Pryor, M., Duthoy, S., Grondin, S., Lacroix, C., Monsempe, C., Simon, S., Harris, B., Atkin, R., Doggett, J., Mayes, R., Keating, L., Wheeler, P. R., Parkhill, J., Barrell, B. G., Cole, S. T., Gordon, S. V. & Hewinson, R. G. (2003). The complete genome sequence of *Mycobacterium bovis*. *Proc Natl Acad Sci U S A*, 100(13): 7877-7882.
- Gerhardt, S., Echt, S., Busch, M., Freigang, J., Auerbach, G., Bader, G., Martin, W. F., Bacher, A., Huber, R. & Fischer, M. (2003). Structure and properties of an engineered transketolase from maize. *Plant Physiol*, 132(4): 1941-1949.
- Heinrich, P. C. & Wiss, O. (1971). [Transketolase from human erythrocytes. Purification and properties]. *Helv Chim Acta*, 54(8): 2658-2668.
- Heldt, H. W., Piechulla, B. & Heldt, F. (2010). *Plant Biochemistry*: Academic Press.
- Henkes, S., Sonnewald, U., Badur, R., Flachmann, R. & Stitt, M. (2001). A small decrease of plastid transketolase activity in antisense tobacco transformants has dramatic effects on photosynthesis and phenylpropanoid metabolism. *Plant Cell* 13: 536-551.
- Herrmann. (1995). The shikimate pathway as an entry to aromatic secondary metabolism. *Plant Physiol*(107): 7-12.
- Hollingsworth, S. A. & Karplus, P. A. (2010). A fresh look at the Ramachandran plot and the occurrence of standard structures in proteins. *Biomol Concepts*, 1(3-4): 271-283.

- Horecker, B. L. & Smyrniotis, P. Z. (1955). [53] Transketolase from liver and spinach: Ribulose-5-P + Ribose-5-P $\rightleftharpoons$ Sedoheptulose-7-P + Glyceraldehyde-3-P *Methods in Enzymology* (Vol. Volume 1: pp. 371-375): Academic Press.
- Hrmova, M., Varghese, J. N., De Gori, R., Smith, B. J., Driguez, H. & Fincher, G. B. (2001). Catalytic mechanisms and reaction intermediates along the hydrolytic pathway of a plant beta-D-glucan glucohydrolase. *Structure*, 9(11): 1005-1016.
- Ikeda, M., Kamada, N., Takano, Y. & Nakano, T. (1999). Molecular analysis of the *Corynebacterium glutamicum* transketolase gene. *Biosci Biotechnol Biochem*, 63(10): 1806-1810.
- Joshi, S., Singh, A. R., Kumar, A., Misra, P. C., Siddiqi, M. I. & Saxena, J. K. (2008). Molecular cloning and characterization of *Plasmodium falciparum* transketolase. *Mol Biochem Parasitol*, 160(1): 32-41.
- Kato, N., Higuchi, T., Sakazawa, C., Nishizawa, T., Tani, Y. & Yamada, H. (1982). Purification and properties of a transketolase responsible for formaldehyde fixation in a methanol utilizing yeast, *Candida boidinii*. *Biochim Biophys Acta*, 715: 143-150.
- Kelley, L. A. & Sternberg, M. J. (2009). Protein structure prediction on the Web: a case study using the Phyre server. *Nat Protoc*, 4(3): 363-371.
- Klein, H. & Brand, K. (1977). [Purification and properties of transketolase from *Candida utilis*]. *Hoppe Seylers Z Physiol Chem*, 358(10): 1325-1337.
- Kochetov, G. A. (1982). Transketolase from yeast, rat liver, and pig liver. *Methods Enzymol*, 90 Pt E: 209-223.
- Kruger, N. J. & von Schaewen, A. (2003). The oxidative pentose phosphate pathway: structure and organisation. *Current Opinion in Plant Biology*, 6(3): 236-246.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227(5259): 680-685.
- Lüthy, J. U. & Bowie, D. (1992). *Eisenberg*, *Nature*, 356(83).
- Magel, E. A., Hillinger, C., Wagner, T. & Höll, W. (2001). Oxidative pentose phosphate pathway and pyridine nucleotides in relation to heartwood formation in *Robinia pseudoacacia* L. *Phytochemistry*, 57(7): 1061-1068.
- Markley, J. L., Bahrami, A., Eghbalian, H. R., Peterson, F. C., Ulrich, E. L., Westler, W. M. & Volkman, B. F. (2009). Macromolecular structure determination by NMR spectroscopy. *Structural bioinformatics*, 2: 93-142.

- Martz. (2001). Homology modeling for beginners. Available: <http://www.umass.edu/molvis/workshop/homolmod.htm>. Accessed 12 May 2006.
- Maugeri, D. A., Cazzulo, J. J., Burchmore, R. J. S., Barrett, M. P. & Ogbunude, P. O. J. (2003). Pentose phosphate metabolism in *Leishmania mexicana*. *Molecular and Biochemical Parasitology*, 130(2): 117-125.
- Meek, D. W. & Nimmo, H. G. (1984). Effects of phosphorylation on the kinetic properties of rat liver fructose-1,6-bisphosphatase. *Biochem J*, 222(1): 125-130.
- Meier, N., Dear, T. N. & Boehm, T. (1999). Wn and mHa3 are components of the genetic hierarchy controlling hair follicle differentiation. *Mech Dev*, 89(1-2): 215-221.
- Miles, B. (2003). Pentose Phosphate Pathway Aka the Hexose Monophosphate Shunt.
- Minami, Y., Fujita, K. & Matsubara, H. (2001). Transketolase from *Polygonum tinctorium*. *EMBL/GENEBANK/DDBJ databases*.
- Morris, A. L., MacArthur, M. W., Hutchinson, E. G. & Thornton, J. M. (1992). Stereochemical Quality of Protein Structure Coordinates. *Proteins*, 12: 345-364.
- Murphy, D. J. & Walker, D. A. (1982). The properties of transketolase from photosynthetic tissue. *Planta* 155: 316–320.
- Neuhaus, H. E. & Emes, M. J. (2000). NONPHOTOSYNTHETIC METABOLISM IN PLASTIDS. *Annu Rev Plant Physiol Plant Mol Biol*, 51: 111-140.
- Nguyen, M. N. & Rajapakse, J. C. (2005). Two-stage multi-class support vector machines to protein secondary structure prediction. *Pac Symp Biocomput*: 346-357.
- Nishimura, M. & Beevers, H. (1979). Subcellular distribution of gluconeogenic enzymes in germinating castor bean endosperm. *Plant Physiol*, 64(1): 31-37.
- Nowitzki, U., Wyrich, R., Westhoff, P., Henze, K., Schnarrenberger, C. & Martin, W. (1995). Cloning of the amphibolic Calvin cycle/OPPP enzyme D-ribulose-5-phosphate 3-epimerase (EC 5.1.3.1) from spinach chloroplasts: functional and evolutionary aspects. *Plant Mol Biol*, 29(6): 1279-1291.
- Nulit, R. (2010). Over-expression of *Escherichia coli* Transaldolase in the Cytosol of *Arabidopsis thaliana*. *Pertanika J. Trop. Agric. Sci*, 35: 371 - 379.
- Pal, L. & Basu, G. (1999). Novel protein structural motifs containing two-turn and longer 3(10)-helices. *Protein Eng*, 12(10): 811-814.

- Ramachandran, G. N., Ramakrishnan, C.& Sasisekharan, V. (1963). Stereochemistry of polypeptide chain configurations. *J Mol Biol*, 7: 95-99.
- Riganti, C., Gazzano, E., Polimeni, M., Aldieri, E.& Ghigo, D. (2012). The pentose phosphate pathway: An antioxidant defense and a crossroad in tumor cell fate. *Free Radical Biology and Medicine*, 53(3): 421-436.
- Roscher, A., Kruger, N. J.& Ratcliffe, R. G. (2000). Strategies for metabolic flux analysis in plants using isotope labelling. *J. Biotechnol*, 77: 81–102.
- Rossetti, S., van Unen, L., Sacchi, N.& Hoogeveen, A. T. (2008). Novel RNA-binding properties of the MTG chromatin regulatory proteins. *BMC Mol Biol*, 9: 93.
- Sagisaka, S. (1974). Effect of low temperature on amino Acid metabolism in wintering poplar: arginine-glutamine relationships. *Plant Physiol*, 53(2): 319-322.
- Schenk, G., Layfield, R., Candy, J. M., Duggleby, R. G.& Nixon, P. F. (1997). Molecular evolutionary analysis of the thiamine-diphosphate-dependent enzyme, transketolase. *J Mol Evol*, 44(5): 552-572.
- Schnarrenberger, C., Flechner, A.& Martin, W. (1995). Enzymatic Evidence for a Complete Oxidative Pentose Phosphate Pathway in Chloroplasts and an Incomplete Pathway in the Cytosol of Spinach Leaves. *Plant Physiol*, 108(2): 609-614.
- Schwender, J., Goffman, F., Ohlrogge, J. B.& Shachar-Hill, Y. (2004). Rubisco without the Calvin cycle improves the carbon efficiency of developing green seeds. *Nature*, 432(7018): 779-782.
- Sheng, X., Liu, Y.& Liu, C. (2013). Theoretical studies on the common catalytic mechanism of transketolase by using simplified models. *Journal of Molecular Graphics and Modelling*, 39(0): 23-28.
- Slaski., Zhang.G., Basu.U., Stephens.J.L.& Taylor.g. (1996). Aluminium resistance in wheat (*Triticum aestivum* L.) is associated with rapid Al-induced changes in activities of glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase in root apices. *Physiol. Plant*, 98: 477-484.
- Sprenger, G. A., Schorken, U., Sprenger, G.& Sahm, H. (1995). Transketolase A of *Escherichia coli* K12. Purification and properties of the enzyme from recombinant strains. *Eur J Biochem*, 230(2): 525-532.
- Srere, P., Cooper, J. R., Tabachnick, M.& Racker, E. (1958). The oxidative pentosephosphate cycle. I. Preparation of substrates and enzymes. *Arch. Biochem. Biophys*, 274: 295-274.
- Strable, M. S.& Ntambi, J. M. (2010). Genetic control of de novo lipogenesis: role in diet-induced obesity. *Crit Rev Biochem Mol Biol*, 45(3): 199-214.

- Strafford, J., Payongsri, P., Hibbert, E. G., Morris, P., Batth, S. S., Steadman, D., Smith, M. E. B., Ward, J. M., Hailes, H. C. & Dalby, P. A. (2012). Directed evolution to re-adapt a co-evolved network within an enzyme. *Journal of Biotechnology*, 157(1): 237-245.
- Sundström, M., Lindqvist, Y. & Schneider, G. (1992). Three-dimensional structure of apotransketolase flexible loops at the active site enable cofactor binding. *FEBS Letters*, 313(3): 229-231.
- Sundstrom, M., Lindqvist, Y., Schneider, G., Hellman, U. & Ronne, H. (1993). Yeast TKL1 gene encodes a transketolase that is required for efficient glycolysis and biosynthesis of aromatic amino acids. *J Biol Chem*, 268(32): 24346-24352.
- Taylor, W. R. & Orengo, C. A. (1989). Protein structure alignment. *J Mol Biol*, 208(1): 1-22.
- Teige, M., Kopriva, S., Bauwe, H. & K.H., S. (1996). Primary structure of chloroplast transketolase from potato. *Plant Gene Register PGR*: 96-121.
- Tiessen, A., Nerlich, A., Faix, B., Hummer, C., Fox, S., Trafford, K., Weber, H., Weschke, W. & Geigenberger, P. (2012). Subcellular analysis of starch metabolism in developing barley seeds using a non-aqueous fractionation method. *J Exp Bot*, 63(5): 2071-2087.
- Tramontano, A. (1998). Homology modeling with low sequence identity. *Methods*, 14(3): 293-300.
- Veitch, N. J., Maugeri, D. A., Cazzulo, J. J., Lindqvist, Y. & Barrett, M. P. (2004). Transketolase from *Leishmania mexicana* has a dual subcellular localization. *Biochem J*, 382(Pt 2): 759-767.
- Vlachakis, D. (2007). An Introduction to Molecular Modelling, from theory to application. *University of Cardiff, Wales, UK*, 1(1 isbn:0-615-17606-2).
- Vlachakis, D. (2009). Theoretical study of the Usutu virus helicase 3D structure, by means of computer-aided homology modelling. *Theor Biol Med Model*, 6: 9.
- Voskoboev, A. I. & Gritsenko, E. A. (1981). [Nature of the bond between the coenzyme and protein in pig liver transketolase]. *Biokhimiia*, 46(8): 1383-1388.
- Wamelink, M. M., Struys, E. A. & Jakobs, C. (2008). The biochemistry, metabolism and inherited defects of the pentose phosphate pathway: a review. *J Inherit Metab Dis*, 31(6): 703-717.
- Wikner, C., Nilsson, U., Meshalkina, L., Udekwu, C., Lindqvist, Y. & Schneider, G. (1997). Identification of catalytically important residues in yeast transketolase. *Biochemistry*, 36(50): 15643-15649.

Wood, T. (1974). Catalysis of pentose phosphate pathway reactions by cytoplasmic fractions from muscle, uterus and liver of the rat, and the presence of a reduced nicotinamide-adenine dinucleotide phosphate-triose phosphate oxidoreductase in rat muscle. *Biochem J*, 138(1): 71-76.

Zaki, M.& Bystroff, C. (2010). *Protein Structure Prediction*: Humana Press.

