



UNIVERSITI PUTRA MALAYSIA

***EFFECTS OF COLONIZATION OF AN ENDOPHYTIC FUNGUS,
Hendersonia toruloidea ON THIAMINE BIOSYNTHESIS IN OIL PALM
SEEDLINGS (Elaeis guineensis Jacq.)***

AMIRAH NOR BT. KAMARUDIN

FBSB 2018 6



**EFFECTS OF COLONIZATION OF AN ENDOPHYTIC FUNGUS,
Hendersonia toruloidea ON THIAMINE BIOSYNTHESIS IN OIL PALM
SEEDLINGS (*Elaeis guineensis* Jacq.)**

By

AMIRAH NOR BT. KAMARUDIN

**Thesis Submitted to School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

October 2017

COPYRIGHT

All materials contained within the thesis including without limitation text, logos, icons, photographs and all other artworks are copyright material of Univeristi Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from copyright holder. Commercial use of materials may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright©Universiti Putra Malaysia.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**EFFECTS OF COLONIZATION OF AN ENDOPHYTIC FUNGUS,
Hendersonia toruloidea ON THIAMINE BIOSYNTHESIS IN OIL PALM
SEEDLINGS (*Elaeis guineensis* Jacq.)**

By

AMIRAH NOR BT KAMARUDIN

October 2017

Chairman : Zetty Norhana Binti Balia Yusof, PhD
Faculty : Biotechnology and Biomolecular Sciences

Thiamine, or vitamin B1 plays an indispensable role in many metabolic reactions. Besides that, thiamine is also associated with the induction of systemic acquired resistance (SAR) in plants and having a role in boosting plant's immunity and defense system. In Malaysia, oil palm productivity is hampered by basal stem rot disease caused by a pathogenic fungus, *Ganoderma boninense* and proper disease management have yet to be discovered. Application of endophytes as biocontrol agent is a promising measure to prevent the disease. *Hendersonia toruloidea* is an endophytic fungus originally isolated from oil palm roots which have been shown to have excellent biocontrol activity in oil palm seedlings. Previous studies showed that this endophyte is able to suppress *G. boninense* infection in oil palm seedlings. This work aimed to investigate the responses in oil palm seedlings, specifically on the expressions of thiamine biosynthesis genes upon application of *H. toruloidea*. Seven months old oil palm seedlings were inoculated with *H. toruloidea* and microscopy analyses were carried out to visualize the colonization of the fungus. Total RNA was extracted from oil palm leaves at day 1, 7, 15 and 30 post inoculation. Quantitative real-time PCR (qPCR) was performed to measure the level of expression of four key thiamine biosynthesis genes, namely *THI4*, *THIC*, *THI* and *TPK*. The results showed of up to 12-fold of increase in the expression of all gene transcripts at day 1 post inoculation. At subsequent days of day 7, day 15 and 30 post inoculation, the relative expression of these genes were shown to be downregulated. Thiamine accumulation was observed via HPLC analysis at day 7 post inoculation and subsequently attenuated until day 30. This work provides first evidence of enhancement of thiamine biosynthesis by endophytic colonization in oil palm and suggesting the role of thiamine in stress protection in oil palm seedlings.

Abstrak tesis yang dikemukakan kepada Senat of Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**KESAN KOLONISASI KULAT ENDOFIT *Hendersonia toruloidea*
TERHADAP BIOSINTESIS VITAMIN B1 DALAM KELAPA
SAWIT (*Elaeis guineensis* Jacq.)**

Oleh

AMIRAH NOR BT KAMARUDIN

Oktober 2017

Pengerusi : Zetty Norhana Binti Balia Yusof, PhD
Fakulti : Bioteknologi dan Sains Biomolekular

Vitamin B1 memainkan peranan yang penting dalam metabolisme dalam semua kehidupan. Selain itu, vitamin B1 juga terlibat dalam induksi rintangan sistemik dalam tumbuhan dan memainkan peranan penting dalam immuniti dan pertahanan tumbuhan. Di Malaysia, produktiviti kelapa sawit adalah terjejas disebabkan oleh penyakit pereputan pangkal akar yang disebabkan oleh kulat *Ganoderma boninense*. Namun, pengawalan terbaik dan efektif terhadap kulat *Ganoderma boninense* belum dikenalpasti. Pengawalan penyakit BSR telah dilakukan dengan menggunakan agen kawalan biologi iaitu melalui penggunaan endofit terhadap anak pokok kelapa sawit. Kulat *Hendersonia toruloidea* merupakan kulat endofit yang telah dijumpai dalam akar kelapa sawit dan telah menunjukkan keberkesanan yang memberangsangkan terhadap kawalan penyakit BSR. Oleh itu, kajian ini bertujuan untuk menyelidiki kesan kolonisasi kulat endofit *H. toruloidea* terhadap biosintesis vitamin B1 di dalam anak pokok kelapa sawit. Kulat endofit *H. toruloidea* telah diinokulasi ke atas anak benih kelapa sawit berusia 7 bulan di nurseri. Analisa mikroskopi telah dijalankan untuk mengkaji kolonisasi kulat endofit di dalam akar kelapa sawit. RNA telah diekstrak daripada daun kelapa sawit pada hari ke 1, 7, 15 dan 30 hari usai inokulasi. Tindak balas rangkaian polimerase secara kuantitatif masa sebenar (qPCR) telah dijalankan untuk melihat pengekspressan empat gen biosintesis B1 yang utama iaitu *THI4*, *THIC*, *THI* dan *TPK*. Hasil penemuan menunjukkan pengekspressan gen biosintesis vitamin B1 meningkat sehingga 12 kali ganda. Selepas 15 dan 30 hari usai inokulasi, pengekspressan gen vitamin B1 telah menurun. Analisa HPLC menunjukkan kolonisasi kulat endofit telah menyebabkan pengumpulan metabolit vitamin B1 di dalam daun kelapa sawit. Hasil kajian ini berjaya menunjukkan kolonisasi *H. toruloidea* meningkatkan biosintesis vitamin B1 di dalam pokok dan mencadangkan peranan vitamin B1 dalam perlindungan kelapa sawit terhadap tekanan.

ACKNOWLEDGEMENTS

In the Name of Allah, The Most Gracious and The Most Merciful.

I am totally grateful that Allah has blessed me and guided me in this journey to seek knowledge. In this ayah 'And He found you lost and guided you.'(93:7) resonates in me.

I would like to express my gratitude to my first supervisor, Dr Zetty Norhana for her patience and kindness for guiding me throughout the study. She believed in me and pushed me to do my very best.

I would also like to thank Dr Idris Abu Seman who have always guided and supported me throughout this study.

I also like to thank my co-supervisor, Dr Lai Kok Song, who always provided me with assistance and keeps me motivated. I also would like to thank Dr Dhilia for her helpful support and feedback.

I am thankful to my parents, Kamarudin and Humairah, who have always been my pillar of strength, the source of inspiration that keeps me strong and supported me throughout this journey. I am also indebted to my siblings, Hanim, Ariff, Azam and my twin sister Athirah for their support and motivation.

To Raimathy, Ina, and Angel who have never failed to support me morally and intellectually. I would also like to thank my labmates in Plant-Algae Biochemistry and Biotechnology Lab, especially Atiqah, Mimi, Sakinah and Aisamuddin for their support and contribution in this work.

I would also like to acknowledge MPOB-GSAS for giving me financial support. I would like to thank lab officers and assistants of GanoDROP Unit, AOTD Unit, and Analytical Quality Unit for their involvement and contribution in my work. Special thanks to Dr. Halimah and Puan Rosilawati for assisting me in my HPLC analysis. Lastly, I would like to extend my gratitude to Mr Santhana from Institute of Medical Research for his kind assistance for my transmission electron microscopy work.

Indeed, this beautiful journey will not come into fruition without them.

I certify that a Thesis Examination Committee has met on 3 October 2017 to conduct the final examination of Amirah Nor binti Kamarudin on her thesis entitled "Effects of Colonization of an Endophytic Fungus, *Hendersonia toruloidea* on Thiamine Biosynthesis in Oil Palm Seedlings (*Elaeis guineensis* Jacq.)" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Syahida binti Ahmad, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Noor Azmi Shaharuddin, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Sreeramanan Subramaniam, PhD

Associate Professor
Universiti Sains Malaysia
Malaysia
(External Examiner)



NOR AINI AB. SHUKOR, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 December 2017

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Zetty Norhana Balia Yusof, PhD

Senior Lecturer

Faculty of Biotechnology and Biomolecular Science

Universiti Putra Malaysia

(Chairman)

Dhilia Udie Lamasudin, PhD

Senior Lecturer

Faculty of Biotechnology and Biomolecular Science

Universiti Putra Malaysia

(Member)

Lai Kok Song, PhD

Senior Lecturer

Faculty of Biotechnology and Biomolecular Science

Universiti Putra Malaysia

(Member)

Idris Abu Seman, PhD

Head of Ganoderma & Diseases Reaseach of Oil Palm Unit

Malaysian Palm Oil Board

(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No : Amirah Nor Bt Kamarudin, GS41473

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Zetty Norhana Balia Yusof, PhD

Signature: _____
Name of Member
of Supervisory
Committee: Dhilia Udie Lamasudin, PhD

Signature: _____
Name of Member
of Supervisory
Committee: Lai Kok Song, PhD

Signature: _____
Name of Member
of Supervisory
Committee: Idris Abu Seman, PhD

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	ii
ACKNOWLEDGEMENTS	iii
APPROVAL	v
DECLARATION	vi
LIST OF TABLES	xi
LIST OF FIGURES	xii
KIST OF APPENDICES	xvi
LIST OF ABBREVIATIONS	xvii
CHAPTER	
1 INTRODUCTION	1
1.1 Objectives	2
2 LITERATURE REVIEW	3
2.1 Oil palm industry and its economic importance in Malaysia	3
2.2 Basal stem rot disease (BSR) and disease management	3
2.3 Endophytes and its role in plant protection	3
2.3.1 Mechanism of endophyte-mediated tolerance to biotic stress is through induced systemic resistance	4
2.3.2 Transcriptomic analysis on plant response to fungal endophytes	8
2.3.3 Endophytic microbes in oil palm	8
2.3.4 Endophytic fungus <i>Hendersonia toruloidea</i>	8
2.4 Overview of thiamine	9
2.4.1 The thiamine biosynthesis pathway in plants	9
2.4.2 Role of thiamine in biotic and abiotic stresses in plants	11
2.4.3 Role of thiamine in plant growth	14
2.5 Endophytic <i>H. toruloidea</i> as inducer of thiamine biosynthesis in oil palm	14
3 MATERIALS AND METHODS	15
3.1 Plant Materials	15
3.2 Fungal Materials	15
3.3 Inoculation procedure	17
3.4 Sampling	17
3.5 Colonization of <i>Hendersonia toruloidea</i> in oil palm roots	18
3.5.1 Reisolation of endophytic fungus <i>H. toruloidea</i> in oil palm roots	18
3.5.2 Scanning electron microscopy analysis	18
3.5.3 Transmission electron microscopy	18
3.6 Quantitative real time PCR	19

3.6.1	Total RNA isolation	19
3.6.2	RNA purification	20
3.6.3	Data mining and primer designing	20
3.6.4	Agarose gel electrophoresis	20
3.6.5	cDNA synthesis	21
3.6.6	Amplification Efficiency Test	21
3.6.7	Cloning and plasmid isolation	21
3.6.8	Sequencing of DNA and sequence analysis.	22
3.6.9	Amplification of thiamine biosynthesis genes by quantitative real-time PCR (qPCR)	22
3.7	HPLC analysis for determination of thiamine, thiamine monophosphate and thiamine pyrophosphate	23
3.7.1	Sample preparation of crude plant extract	23
3.7.2	Standard preparation of thiamine, thiamine monophosphate and thiamine pyrophosphate	23
3.7.3	Sample and standard derivatization to thiochromes derivatives of thiamine	23
3.7.4	Chromatography settings	24
3.8	Statistical analyses	24
4	RESULTS AND DISCUSSION	25
4.1	Colonization of <i>Hendersonia toruloidea</i> in oil palm root	25
4.1.1	Scanning electron microscopy analysis	27
4.1.2	Transmission electron microscopy	28
4.2	Quantitative real-time PCR analysis (qPCR)	32
4.2.1	Total RNA isolation and purification	32
4.2.2	Data mining and primer designing	34
4.2.3	Testing for amplification efficiency	39
4.2.4	Cloning and analysis of insert with EcoR1	44
4.2.5	Sequencing analysis	44
4.2.6	Validation of reference genes	45
4.2.7	The effects of colonization of <i>H. toruloidea</i> on expression of thiamine biosynthesis genes in oil palm seedlings	47
4.3	HPLC analysis of total thiamine, thiamine monophosphate and thiamine pyrophosphate	52
4.3.1	Simultaneous determination of thiamine in oil palm leaves sample using HPLC with fluorescent detection (HPLC-FLD)	52
4.3.2	The effect of colonization of <i>H. toruloidea</i> on total content of thiamine and thiamine pyrophosphate in oil palm leaves	55
5	CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCH	57
5.1	Conclusion	57
5.2	Recommendation for future research	57

REFERENCES	59
APPENDICES	68
BIODATA OF STUDENT	81
LIST OF PUBLICATIONS	82



LIST OF TABLES

Table		Page
2.1	Biotic stress tolerance conferred by fungal endophytes in different host plants	6
2.2	Biochemical function of thiamine biosynthesis enzymes	10
3.1	Ratio of acetone and resin mixture and the duration of infiltration with the specimens	18
3.2	Gradient elution settings to manipulate the concentration of mobile phase eluting the column bed	24
4.1	Assessment of nucleic acid quality using NanoDrop. A good quality total RNA was successfully isolated as shown on the A260/A280 and A260/A30 values of (1.8-2.0)	32
4.2	Selected plants and microalgae species obtained from GenBank used for primer designing	35
4.3	List of primers designed using Primer Premier™ software	38
4.4	Amplification efficiency tests. Amplification efficiency tests were evaluated through r^2 value and efficiency (%). All primers met the criteria of $r^2 > 0.98$ and efficiency ranging from 90-110%	42
4.5	BLAST analysis of the sequencing results of the PCR fragments significantly reflected high similarity of the alignment with <i>Elaias guineensis</i> as indicated by the E value score and sequence similarity (%)	45

LIST OF FIGURES

Figure		Page
2.1	The thiamine biosynthesis pathway in plants. HMP-P is hydroxymethylpyrimidine phosphate, HMP-PP is hydroxymethylpyrimidine pyrophosphate, HET-P is hydroxyethylthiazole phosphate. THIC is hydroxymethylpyrimidine-synthase, THI4 is thiazole biosynthetic protein, TH1 is hydroxymethylpyrimidine phosphate kinase, TH2 is thiamine monophosphate phosphatase and TPK is thiamine pyrophosphate kinase. Adapted from Guan <i>et al.</i> (2014)	11
2.2	The transgenic rice that has a knocked-out OsDR8 gene which is homologs of THI1/THI4 showed an apparent disease symptoms caused by <i>Xanthomonas oryzae pv. Oryzae</i> and <i>Magnaporthe grisea</i>	13
3.1	Sequential steps of preparation of <i>H. toruloidea</i> bioformulation. Pure cultures of <i>H. toruloidea</i> was encapsulated in an alginate formulation to maintain a steady release of fungal inoculum	16
4.1	Establishment of <i>H. toruloidea</i> in oil palm seedlings	26
4.2	SEM images of 10 mm root sections of oil palm seedlings inoculated with <i>H. toruloidea</i> revealed fungal ultrastructures after ten days of incubation. A) Invasive growth of hyphae (arrow) B) Arrow shows proliferation of conidiospore C) Appresorium-like structure forming clamp connection (arrow). D) Arrow shows ascospores formation. Bar size 10 μ m	27
4.3	Light micrographs of transverse section of oil palm root stained with toluidine blue. A) Transverse section of uninoculated seedling B) Transverse section of oil palm root colonized by <i>H. toruloidea</i> at day 30 post colonization. Arrow shows direct penetration of hyphae through epidermal and cortical cells. Bar size 0.02 mm	29
4.4	A) and B) Transmission electron microscopy of transverse section of oil palm root colonized by <i>H. toruloidea</i> at day 30 post colonization. Arrow showed appresorium-like structures in cortical cells at day 30 of colonization. Bar size is 5000 nm	30

- 4.5 A) and B) Transmission electron microscopy of oil palm root colonized by *H. toruloidea* at day 30 post colonization. Red arrows showing hyphal coils in cortical cells. Dark cell wall aspositions were apparent as indicated by orange arrows. Bar size is 5000 nm 31
- 4.6 Assessment of RNA integrity by visualizing RNA on 1% agarose gel electrophoresis. Lane M: 1kb ladder (NEB), Lane 1: Total RNA isolated from oil palm spear leaves showing 2 intact bands of 28S and 18S rRNA 33
- 4.7 Sequential effects of RNA integrity after each steps of DNase treatment. Lane M is 100bp Ladder (GeneRuler, ThermoScientific); Lane 1 is RNA at Room temperature, Lane 2 is RNA at heat treatment at 37 °C, Lane 3 is RNA at heat treatment at 37 °C and addition of EDTA, Lane 4 is RNA at heat treatment 37 °C, addition EDTA, and lastly heat treatment at 65 °C. DNase treatment successfully removed genomic DNA and maintained the integrity of RNA 33
- 4.8 Alignment of THIC gene sequences from *Arabidopsis thaliana* (NM_001202705), *Elaeis guineensis* (XM_010924818.1), *Poa secunda* (AF264021.1) and *Zea mays* (EU972242.1) using MSA tool from EMBL Clustal W (<http://www.ebi.ac.uk/Tools/msa/>). Residues which are similar or identical were denoted by asterisks. The consensus sequence was utilized for primer designing using Primer Premier 6.0 37
- 4.9 Phases of PCR amplification curve. qPCR amplification curves of seven 2-fold dilutions of cDNA generated from cDNA of control oil palm seedlings in triplicates. Each amplification curves represent the four different phase: the baseline, the exponential phase, the log linear phase and finally the plateau phases. Y axis represents reference fluorescence units (RFU), which is a fluorescence value of an experimental reaction minus the Rn value of baseline signal generated by Biorad CFX 96. X-axis represents the cycle number. (A) represents the threshold at which Delta Rn reached the linear phase of amplification of all samples 40
- 4.10 Standard curves generated from cDNA of control oil palm seedlings. Seven 2-fold dilutions, starting with 100ng template copies of cDNA and ending with 3.125 ng copies. Y-axis represents the threshold cycle number. X-axis represents the log starting quantity of the cDNA template. A line representing the best fit was calculated for the standard curve using the least squares method of linear regression. A slope of -3.54 (91.3% efficiency), a y-intercept of 33 and r^2 of 0.988 was obtained 41

- 4.11 Amplification of gene fragments from cDNA derived from oil palm seedlings. The gene fragments corresponded to the expected fragment length. Lane 1 is *TUB*, lane 2 is *GAD*, lane 3 is *THI4*, lane 4 is *THIC*, lane 5 is *THI*, and lane 6 is *TPK*. Lane M is 50bp DNA ladder (GeneRuler, ThermoScientific) 42
- 4.12 Melt curve analysis. qPCR amplification was performed on 2-fold serial dilutions of cDNA template. Y-axis represents the rate of change of fluorescence versus temperature. X-axis represents the melting temperature of the PCR product at approximately 82.5 °C. A single peak on the melt curve analysis indicates a single PCR product has melted. No amplification was found in no template control (NTC) 43
- 4.13 Clones with inserts were identified by digestion of plasmid by EcoR1. Lane 1, 3, 5, 7, 9, and 11: Undigested plasmids containing inserts of *THI4*, *THIC*, *THI*, *TPK*, *TUB*, and *GAD*. Lane 2, 4, 6, 8, and 10: Digested plasmids containing inserts of *THI4*, *THIC*, *THI*, *TPK*, *TUB*, and *GAD*. Lane M is 100 bp DNA ladder (Invitrogen) 44
- 4.14 The changes in copy number of tubulin and GADPH in the experimental samples. Bar represents mean Ct values of control and treated samples. Error bars indicated standard deviation (SD) of three independent experiments. No significant difference between all samples (p value<0.05). C1 is untreated control day 1, C7 is untreated control day 7, C15 is untreated control day 15, C30 is untreated control day 30, T1 is Ht-inoculated day 1, T7 is Ht- inoculated day 7, T15 is Ht-inoculated day 15, T30 is Ht-inoculated day 30 46
- 4.15 Changes in transcript abundance of *THI4* and *THIC*, the first enzymes in the thiamine biosynthesis pathway. (A) & (B) qPCR analysis was conducted on *THI4* and *THIC* transcripts of 7 months oil palm seedlings that were inoculated with *H. toruloidea* at day 1, 7, 15 and 30. Error bars denote standard deviation (SD). Asterisks denote significant difference at p<0.05 compared to control seedlings 50
- 4.16 Changes in transcript abundance of *TPK* and *THI1*. (C) & (D) qPCR analysis of *TPK* and *THI1* transcripts of 7 months old oil palm seedlings inoculated with *H. toruloidea* at day 1, 7, 15, and 30. Error bars denote standard deviation (SD). Asterisks denote significant difference at p<0.05 compared to control seedlings 51
- 4.17 Chromatograms of thiochrome derivatives of thiamine standards (A) and oil palm leaves sample (B), where 1 is TPP (Rt=3.04), 2 is TMP (Rt=3.35) and 3 is TF (Rt=7.22) 53

- 4.18 Calibration curves of standard thiamine pyrophosphate (A), thiamine monophosphate (B) and thiamine (C). 10 ppm, 2 ppm, 1 ppm, 0.4 ppm and 0.2 ppm of standards were injected 54
- 4.19 HPLC analysis of thiamine content (TPP, TMP and TF) in 7 months old oil palm seedlings in control and inoculated seedlings T1 is *H. toruloidea* inoculated at day 1, T7 is *H. toruloidea* inoculated at day 7, T15 is *H. toruloidea* inoculated at day 15, and T30 is *H. toruloidea* inoculated at day 30. In each analysis, 5g of fresh oil palm leave was used. Data are means \pm SE of three replicates. Asterisks denotes significant difference between control and inoculated seedlings. Significance level is at p value $<$ 0.05 56



LIST OF APPENDICES

Appendix		Page
1	Preparation of buffers and chemicals	68
2	Sequencing results	70
3	Melt curve analysis	74
4	Calculation for HPLC analysis	77



LIST OF ABBREVIATIONS

°C	Degree celsius
%	Percentage
A _{260nm}	Optical density at wavelength 260 nanometer
A _{280nm}	Optical density at wavelength 280 nanometer
μL	Microliter
μm	Micrometre
μmoles	Micromoles
mg	Milligram
ATP	Adenosine triphosphate
bp	Base pair
cDNA	Complementary deoxyribonucleic acid
CTAB	Cetyl trimethyammonium bromide
DNA	Deoxyribonucleic acid
EDTA	Ethylenediaminetetraacetic acid
g	Gram
HCl	Hydrochloric acid
kb	Kilobase
L	Litres
LB	Luria Bertani
M	Moles
mRNA	Messenger RNA
nm	Nanomoles
NaOH	Sodium hydroxide

PDA	Potato dextrose agar
PCR	Polymerase chain reaction
qPCR	Quantitative real-time PCR
RNA	Ribonucleic acid
RT	Reverse transcriptase



CHAPTER 1

INTRODUCTION

Plants are sessile organisms that are inevitably exposed to unfavourable biotic and abiotic stresses, namely salinity, drought, and also microbial pathogens. Climate changes and attack by pathogenic diseases can severely hamper the productivity of important crop plants, including oil palm. Oil palm is one of Malaysia's important commodity. However, the productivity of oil palm is threatened by basal stem rot disease caused by a fungus, *Ganoderma boninense* which resulted in major economic losses (Paterson, 2007; Rees *et al.*, 2009). Current research in oil palm is accelerating towards finding ways to control the disease and is focusing on the detailed molecular mechanism in the plant-pathogenic interaction (Ho and Tan, 2014). Recently, studies of plant and microorganism interactions are significantly attracting interest because it has been demonstrated that microorganisms such as endophytes play a role in alleviating stresses in their host plants (Boivin *et al.*, 2016).

Endophytes are microorganisms that colonize the insides of plant tissues without causing any disease (Wilson, 1995). It is widely documented that endophytes formed a beneficial mutualistic relationship with plants (Hernández-Montiel *et al.*, 2013; Seerangan and Thangavelu, 2014). It is suggested that plant-endophyte mutualism is formed through direct and indirect mechanisms. Direct mechanism include antibiosis, and indirect is through production of biochemical compounds that are associated to alleviating stresses through induced systemic resistance (ISR), thereby enhancing plant's immune system and preventing pathogenic attack (Alquéres *et al.*, 2010; Gao *et al.*, 2010).

Relatively, no studies have been done on the role of thiamine in stress protection in oil palm. Thiamine or vitamin B1, an enzymatic cofactor in metabolic reactions, is involved in plant adaptation and alleviation of biotic and abiotic stresses in plants (Tunc-Ozdemir *et al.*, 2009; Rapala-Kozik *et al.*, 2008; Goyer, 2010). It was observed that there was an accumulation of thiamine when the plants were subjected to salinity stress, oxidative stress and pathogenic attack (Rapala-Kozik *et al.*, 2008; Tunc-Ozdemir *et al.*, 2009; Zhou *et al.*, 2013). It is now understood that thiamine formed an indirect role in enhancing anti oxidative capacity in the plants, which is important in defense responses (Zhou *et al.*, 2013). Yet, the exact mechanism of biosynthesis of thiamine in response to stresses is still poorly understood.

Thiamine is involved in adaptation to biotic and abiotic stresses and application of endophytes enhance the synthesis of defence metabolites that is associated with ISR/SAR (Zheng *et al.* 2015). This led to a hypothesis that thiamine biosynthesis in oil palm will be upregulated by colonisation by endophytes. In this study, the endophytic fungus *Hendersonia toruloidea* was chosen as the strain of choice due to its excellent

colonization ability and also its ability to suppress *Ganoderma boninense* disease infection in oil palm (Idris *et al.*, 2013).

1.1 Objectives

The objectives of this study are:

1. To determine the localization and colonisation of endophytic fungus in oil palm by scanning electron microscopy and transmission electron microscopy
2. To determine the level of expression of thiamine biosynthesis genes upon colonisation of endophytic fungus by quantitative PCR
3. To quantify the total thiamine accumulation in oil palm upon colonisation by endophytic fungus by HPLC

The first objective was to perform microscopy analysis to study the colonisation and morphological pattern of the endophytic fungus. It was expected that the endophytic *H. toruloidea* will actively colonizing oil palm root tissues. Scanning electron microscopy (SEM) revealed the morphology of the *H. toruloidea* while transmission electron microscopy (TEM) was carried out to examine the structure and localisation of *H. toruloidea* in the oil palm root.

The second objective was to determine the expressions of thiamine biosynthesis genes in oil palm upon successive colonisation of *H. toruloidea*. The successful colonization of endophytic fungus was hypothesized to cause an upregulation of the thiamine biosynthesis genes in oil palm. The gene expression study was performed using quantitative real time-PCR (qRT-PCR), a simple, high throughput technology that enable us to measure gene expression in real time. The expression of thiamine biosynthesis genes was examined over a time course of 1, 7, 15 and 30 days post inoculation and the result will reflect the changes in transcript abundances of thiamine biosynthesis genes upon colonization of *H. toruloidea*.

The third objective of this study was to measure total thiamine content in oil palm upon colonization of the endophytic fungus. The upregulation of thiamine biosynthesis genes was expected to cause the increase in total thiamine and its intermediates accumulation overall. It was performed using High Performance Liquid Chromatography (HPLC). Since gene expressions are not necessarily translated into functional protein, the measurement of total thiamine and its intermediate content will verify that thiamine biosynthesis genes are expressed to synthesise total thiamine or the synthesis of its intermediates might be involved in other mechanisms in the overall metabolic pathways.

REFERENCES

- Abidin, A. A. Z., Wong, S. Y., Abdul Rahman, N. S., Che Idris, and Balia Yusof, Z. N. (2016). Osmotic, oxidative, and salinity stresses upregulate the expressions of thiamine (vitamin B1) biosynthesis genes (*THIC & THII/THI4*) in oil palm (*Elaeis guineensis*). *Journal of Oil Palm Research*, 28(3): 308–319.
- Ahn, I. P., Kim, S., and Lee, Y. H. (2005). Vitamin B1 functions as an activator of plant disease resistance. *Plant Physiology*, 138(3): 1505–15.
- Ahn, I. P., Kim, S., Lee, Y. H., and Suh, S. C. (2007). Vitamin B1-induced priming is dependent on hydrogen peroxide and the NPR1 gene in *Arabidopsis*. *Plant Physiology*, 143(2): 838–48.
- Alquéres, S. M. C., Oliveira, J. H. M., Nogueira, E. M., Guedes, H. V., Oliveira, P. L., Câmara, F., and Martins, O. B. (2010). Antioxidant pathways are up-regulated during biological nitrogen fixation to prevent ROS-induced nitrogenase inhibition in *Gluconacetobacter diazotrophicus*. *Archives of Microbiology*, 192(10): 835–41.
- Ariffin, D., Idris, A. S., and Marzuki, A. (Eds.). (1996). Proceedings of the 1996 PORIM International Palm Oil Congress (Agriculture). PORIM, Malaysia. p. 317-329.
- Asano, Y., Katsumoto, H., Inokuma, C., Kaneko, S., Ito, Y., and Fujiie, A. (1996). Cytokinin and thiamine requirements and stimulative effects of riboflavin and α -ketoglutaric acid on embryogenic callus induction from the seeds of *Zoysia japonica steud.* *Journal of Plant Physiology*, 149(3–4): 413–417.
- Bahuguna, R. N., Joshi, R., Shukla, A., Pandey, M., and Kumar, J. (2012). Thiamine primed defense provides reliable alternative to systemic fungicide carbendazim against sheath blight disease in rice (*Oryza sativa* L.). *Plant Physiology and Biochemistry*, 57: 159–67.
- Balia Yusof, Z. N., Borhan, F. P., Mohamad, F. A., and Rusli, M. H. (2015). The effect of *Ganoderma boninense* infection on the expressions of thiamine (Vitamin B1) biosynthesis genes in oil palm. *Journal of Oil Palm Research*, 27(1): 12–18.
- Barcelos, E., Rios, S. de A., Cunha, R. N. V., Lopes, R., Motoike, S. Y., Babiychuk, E., and Kushnir, S. (2015). Oil palm natural diversity and the potential for yield improvement. *Frontiers in Plant Science*, 6: 190.
- Bustin, S. A., Benes, V., Garson, J. A., Hellems, J., Huggett, J., Kubista, M., et al. (2009). The MIQE guidelines: Minimum Information for publication of quantitative real-time PCR experiments. *Clinical Chemistry*, 55, 611–622.
- Begley, T. P., Downs, D. M., Ealick, S. E., McLafferty, F. W., Van Loon, A. P. G. M., Taylor, S., and Xi, J. (1999). Thiamin biosynthesis in prokaryotes. *Archives of*

- Boivin, S., Fonouni-Farde, C., and Frugier, F. (2016). How Auxin and Cytokinin Phytohormones Modulate Root Microbe Interactions. *Frontiers in Plant Science*, 7: 1240.
- Boubakri, H., Wahab, M. A., Chong, J., Bertsch, C., Mliki, A., and Soustre-Gacougnolle, I. (2012). Thiamine induced resistance to *Plasmopara viticola* in grapevine and elicited host-defense responses, including HR like-cell death. *Plant Physiology and Biochemistry*. 57: 120–33.
- Cabezas, L., Calderon, C., Medina, L. M., Bahamon, I., Cardenas, M., Bernal, A. J., and Restrepo, S. (2012). Characterization of cellulases of fungal endophytes isolated from *Espeletia* spp. *Journal of Microbiology* 50(6): 1009–13.
- Chan, P. L., Rose, R. J., Abdul Murad, A. M., Zainal, Z., Leslie Low, E. T., Ooi, L. C. L., et al. (2014). Evaluation of reference genes for quantitative real-time PCR in oil palm elite planting materials propagated by tissue culture. *PLoS One*, 9(10): e110079
- Choudhary, D. K., Prakash, A., and Johri, B. N. (2007). Induced systemic resistance (ISR) in plants: mechanism of action. *Indian Journal of Microbiology*, 47(4): 289–97.
- Colinas, M., and Fitzpatrick, T. B. (2015). Natures balancing act: examining biosynthesis de novo, recycling and processing damaged vitamin B metabolites. *Current Opinion in Plant Biology*, 25: 98–106.
- Dhillon, R. S., Hooda, M. S., Pundeer, J. S., Ahlawat, K. S., and Chopra, I. (2011). Effects of auxins and thiamine on the efficacy of techniques of clonal propagation in *Jatropha curcas* L. *Biomass and Bioenergy*, 35(4): 1502–1510.
- Djonovic, S., Vargas, W. A, Kolomiets, M. V, Horndeski, M., Wiest, A., and Kenerley, C. M. (2007). A proteinaceous elicitor Sml from the beneficial fungus *Trichoderma virens* is required for induced systemic resistance in maize. *Plant Physiology*, 145: 875–889.
- Dong, W., Stockwell, V. O., and Goyer, A. (2015). Enhancement of thiamin content in *Arabidopsis thaliana* by metabolic engineering. *Plant and Cell Physiology*, 56(12): 2285–2296.
- Dong, W., Thomas, N., Ronald, P. C., and Goyer, A. (2016). Overexpression of thiamin biosynthesis genes in rice Increases leaf and unpolished grain thiamin content but not resistance to *Xanthomonas oryzae* pv. *oryzae*. *Frontiers in Plant Science*, 7: 616.
- Du, Q., Wang, H., and Xie, J. (2011). Thiamin (vitamin B1) biosynthesis and regulation: A rich source of antimicrobial drug targets? *International Journal of Biological Sciences*. 7(1): 41-52.

- Ferdous, A. S., Islam, T., Alam, S. S., and Khan, H. (2015). Identification of stable reference genes for quantitative PCR in jute under different experimental conditions : An essential assessment for gene expression analysis. *Australian Journal of Crop Science*, 9: 646–655.
- Ganley, R. J., Sniezko, R. A., and Newcombe, G. (2008). Endophyte-mediated resistance against white pine blister rust in *Pinus monticola*. *Forest Ecology and Management*, 255(7): 2751–2760.
- Gao, F. K., Dai, C. C., and Liu, X. Z. (2010). Mechanisms of fungal endophytes in plant protection against pathogens. *African Journal of Microbiology Research*, 4(13): 1346–1351.
- Gindro, K., Pezet, R., and Viret, O. (2003). Histological study of the responses of two *Vitis vinifera* cultivars (resistant and susceptible) to *Plasmopara viticola* infections. *Plant Physiology and Biochemistry*, 41(9): 846–853.
- Goyer, A. (2010). Thiamine in plants: Aspects of its metabolism and functions. *Phytochemistry*, 71(14–15): 1615–1624.
- Goyer, A., and Haynes, K. G. (2011). Vitamin B1 Content in Potato: Effect of Genotype, Tuber Enlargement, and Storage, and Estimation of Stability and Broad-Sense Heritability. *American Journal of Potato Research*, 88(4): 374–385.
- Gond, S. K., Verma, V. C., Mishra, A., Kumar, A., and Kharwar, R. N. (2010). Role of Fungal Endophytes in Plant Protection. *Management of Fungal Plant Pathogens*, 183–197.
- Guan, J., Hasnain, G., Garrett, T. J., Chase, C. D., Gregory, J., Hanson, A. D., and Mccarty, D. R. (2014). Divisions of labor in the thiamin biosynthetic pathway among organs of maize. *Frontiers in Plant Science*, 5(370): 1–11.
- Hamada, A. M., and Jonsson, L. M. V. (2013). Thiamine treatments alleviate aphid infestations in barley and pea. *Phytochemistry*, 94: 135–41.
- Hardoim, P. R., van Overbeek, L. S., Berg, G., Pirttilä, A. M., Compant, S., Campisano, A., et al. (2015). The hidden world within plants: Ecological and Evolutionary considerations for defining functioning of microbial endophytes. *Microbiology Molecular Biology Reviews*, 79: 293–320.
- Helliwell, K. E., Scaife, M. A., Sasso, S., Araujo, A. P. U., Purton, S., and Smith, A. G. (2014). Unraveling Vitamin B12-Responsive Gene Regulation in Algae. *Plant Physiology*, 165(1): 388–397.
- Hernández-Montiel, L. G., Rueda-Puente, E. O., Cordoba-Matson, M. V., Holguín-Peña, J. R., and Zulueta-Rodríguez, R. (2013). Mutualistic interaction of rhizobacteria with arbuscular mycorrhizal fungi and its antagonistic effect on *Fusarium oxysporum* in *Carica papaya* seedlings. *Crop Protection*, 47: 61–66.

- Ho, C.L., and Tan, Y.C. (2014). Molecular defense response of oil palm to *Ganoderma* infection. *Phytochemistry*, 114: 168-177.
- Hoppenau, C. E., Tran, V. T., Kusch, H., Abhauer, K. P., Landesfeind, M., Meinicke, P., and Braus, G. H. (2014). *Verticillium dahliae* VdTHI4, involved in thiazole biosynthesis, stress response and DNA repair functions, is required for vascular disease induction in tomato. *Environmental and Experimental Botany*, 108: 14–22.
- Hushiarian, R., Yusof, N. A., and Dutse, S. W. (2013). Detection and control of *Ganoderma boninense*: strategies and perspectives. *SpringerPlus*, 2: 555.
- Idris, A.S., Noor Haida, S and Nur Rashyeda, R. (2010). GanoEF1- A fungal biocontrol agent for *Ganoderma* in oil palm. Bangi: MPOB Information Series: 501.
- Idris, A. S., Ahmad Kushari, D., Nur Rashyeda, R., Madihah, A. Z., Haida, N., and Sebran. (2013). Compositions for controlling *Ganoderma* disease in plants and method thereof by using endophytic fungus, *Hendersonia* GanoEF1. US Patent No. WO2015170961.
- Izzati. N.A. and Abdullah. F. (2008). Disease suppression in *Ganoderma* – infected oil palm seedlings treated with *Trichoderma harzianum*. *Plant Protection Science*, 44: 101-107.
- Jeun, Y., Park, K., Kim, C., Fowler, W., and Kloepper, J. (2004). Cytological observations of cucumber plants during induced resistance elicited by rhizobacteria. *Biological Control*, 29(1): 34–42.
- Johansson, T., Le Quéré, A., Ahren, D., Söderström, B., Erlandsson, R., Lundeberg, J., et al. (2004). Transcriptional responses of *Paxillus involutus* and *Betula pendula* during formation of ectomycorrhizal root tissue. *Molecular Plant and Microbe Interaction*, 17: 202–15.
- Kamarudin, A. N., Song, L. K., Idris A.S., Lamasudin, D. U and Balia Yusof, Z. N. (2017). Enhancement of thiamine biosynthesis in oil palm seedlings by colonization of endophytic fungus *Hendersonia toruloidea*. *Frontiers in Plant Science*, 8: 1799.
- Kloepper, J. W., McInroy, J. A., Liu, K., and Hu, C. H. (2013). Symptoms of fern distortion syndrome resulting from inoculation with opportunistic endophytic fluorescent *Pseudomonas* spp. *Plos One*, 8(3): e58531
- Koskimäki, J. J., Hokkanen, J., Jaakola, L., Suorsa, M., Tolonen, A., Mattila, S., and Hohtola, A. (2009). Flavonoid biosynthesis and degradation play a role in early defence responses of bilberry (*Vaccinium myrtillus*) against biotic stress. *European Journal of Plant Pathology*, 125(4): 629–640.
- Khan, A., Bassett, S., Voisey, C., Gaborit, C., Johnson, L., Christensen, M., et al. (2010). Gene expression profiling of the endophytic fungus *Neotyphodium*

- loli in association with its host plant perennial ryegrass. *Australasian Plant Pathology*, 39: 467.
- Kozyrovska, N. O. (2013). Crosstalk between endophytes and a plant host within information processing networks. *Biopolymers and Cell*, 29(3): 234–243.
- Kumara, P. M., Shweta, S., Vasanthakumari, M. M., Sachin, N., Manjunatha, B. L., Jadhav, S. S., and Ganeshiah, K. N. (2014). Endophytes and Plant Secondary Metabolites Synthesis: Molecular and Evolutionary Perspective. In *Advances in Endophytic Research*, (pp.177–190). Springer India.
- Larriba, E., Jaime, M. D. L. A., Nislow, C., Martín-Nieto, J., and Lopez-Llorca, L. V. (2015). Endophytic colonization of barley (*Hordeum vulgare*) roots by the nematophagous fungus *Pochonia chlamydosporia* reveals plant growth promotion and a general defense and stress transcriptomic response. *Journal of Plant Research*, 128: 665–678.
- Lee, S., Whitaker, V. M., and Hutton, S. F. (2016). Mini Review: Potential Applications of Non-host Resistance for Crop Improvement, *Frontiers in Plant Science*, 7: 997.
- Li, M., Petteys, B. J., McClure, J. M., Valsakumar, V., Bekiranov, S., Frank, E. L., and Smith, J. S. (2010). Thiamine biosynthesis in *Saccharomyces cerevisiae* is regulated by the NAD⁺-dependent histone deacetylase Hst1. *Molecular and Cellular Biology*, 30(13): 3329–41.
- Liddicoat, C., Hucker, B., Liang, H., and Vriesekoop, F. (2015). Thiamin analysis in red wine by fluorescence reverse phase-HPLC. *Food Chemistry*, 177: 325–329.
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and. *Methods*, 25: 402–408.
- Lopez-Llorca, L. V., Bordallo, J. J., Salinas, J., Monfort, E., and López-Serna, M. L. (2002). Use of light and scanning electron microscopy to examine colonisation of barley rhizosphere by the nematophagous fungus *Verticillium chlamydosporium*. *Micron*, 33(1): 61–67.
- Machado, C. R., Praekelt, U. M., de Oliveira, R. C., Barbosa, a C., Byrne, K. L., Meacock, P. A., and Menck, C. F. (1997). Dual role for the yeast THI4 gene in thiamine biosynthesis and DNA damage tolerance. *Journal of Molecular Biology*, 273(1): 114–21.
- Malaysian Palm Oil Board, Economic & Industry Development Division (MPOB). (2015, February). Overview of the Malaysian Oil Palm Industry. Retrieved from http://bepi.mpob.gov.my/images/overview/Overview_of_Industry_2015.pdf.

- Malaysian National News Agency. (2014, May 24). Oil palm industry to contribute RM200bil to Malaysia's export value. *The Star*. Retrieved from <http://www.thestar.com.my/business/business-news/2014/05/21/oil-palm-musa/>
- Mohammed, C. L., Rimbawanto, A., and Page, D. E. (2014). Management of basidiomycete root- and stem-rot diseases in oil palm, rubber and tropical hardwood plantation crops. *Forest Pathology*, 44(6): 428-446
- Mimura, M., Zallot, R., Niehaus, T. D., Hasnain, G., Gidda, S. K., and Nguyen, T. N. (2016). *Arabidopsis* TH2 encodes the orphan enzyme thiamin monophosphate phosphatase'. *The Plant Cell*, 28(10): 2683-2696
- Monaim, M. F. A. (2011). Role of riboflavin and thiamine in induced resistance against charcoal rot disease of soybean. *African Journal of Biotechnology*, 10(53): 10842–10855.
- Nongbri, P. L., Vahabi, K., Mrozinska, A., Seebald, E., Sun, C., Sherameti, I., et al. (2013). Balancing defense and growth—Analyses of the beneficial symbiosis between *Piriformospora indica* and *Arabidopsis thaliana*. *Symbiosis*, 58: 17–28.
- Nurrashyeda, R., Idris, A.S., and Ramle, M. (2011) Viability test of alginate granular formulation of *Hendersonia GanoEF1* against *Ganoderma boninense* *in vitro*. In: Proceedings of the Third International Seminar Integrated Oil Palm Pests and Management. MPOB, Malaysia. p. 111-115.
- Nurrashyeda, R and Idris, A.S. (2013). GanoEF1 Biofertiliser - Colonization of *Hendersonia GanoEF1* in oil palm roots. In: Proceedings of the Fifth International Seminar: Sustainable Management of Pests and Diseases in Oil Palm- The Way Forward. MPOB, Malaysia.
- Paerl, R. W., Bertrand, E. M., Allen, A. E., Palenik, B., and Azam, F. (2015). Vitamin B1 ecophysiology of marine picoeukaryotic algae: Strain-specific differences and a new role for bacteria in vitamin cycling. *Limnology and Oceanography*, 60(1): 215–228.
- Palacios, O. A., Bashan, Y., and de-Bashan, L. E. (2014). Proven and potential involvement of vitamins in interactions of plants with plant growth-promoting bacteria—an overview. *Biology and Fertility of Soils*, 50(3): 415–432.
- Palanisamy, K., Ansari, S. A., Kumar, P., and Gupta, B. N. (1998). Adventitious rooting in shoot cuttings of *Azadirachta indica* and *Pongamia pinnata*. *New Forests*, 16(1): 81–88.
- Paterson, R. (2007). *Ganoderma* disease of oil palm -a white rot perspective necessary for integrated control. *Crop Protection*, 26: 1369–1376.

- Pavlo, A., Leonid, O., Iryna, Z., Natalia, K., and Anna, P. (2011). Endophytic bacteria enhancing growth and disease resistance of potato (*Solanum tuberosum* L.). *Biological Control*, 56(1): 43–49.
- Pinto, E., Pedersén, M., Snoeijs, P., Van Nieuwerburgh, L., and Colepicolo, P. (2002). Simultaneous detection of thiamine and its phosphate esters from microalgae by HPLC. *Biochemical and Biophysical Research Communications*, 291(2): 344–348.
- Podolich, O., Ardanov, P., Zaets, I., Pirttilä, A. M., and Kozyrovska, N. (2014). Reviving of the endophytic bacterial community as a putative mechanism of plant resistance. *Plant and Soil*, 388(1): 367–377
- Pourcel, L., Moulin, M., and Fitzpatrick, T. B. (2013). Examining strategies to facilitate vitamin B1 biofortification of plants by genetic engineering. *Frontiers in Plant Science*, 4: 160.
- Rapala-Kozik, M., Kowalska, E., and Ostrowska, K. (2008). Modulation of thiamine metabolism in *Zea mays* seedlings under conditions of abiotic stress. *Journal of Experimental Botany*, 59(15): 4133–4143
- Rapala-Kozik, M., Wolak, N., Kujda, M., and Banas, A. K. (2012). The upregulation of thiamine (vitamin B1) biosynthesis in *Arabidopsis thaliana* seedlings under salt and osmotic stress conditions is mediated by abscisic acid at the early stages of this stress response. *BMC Plant Biology*, 12: 2.
- Rees, R. W., Flood, J., Hasan, Y., Potter, U., and Cooper, R. M. (2009). Basal stem rot of oil palm (*Elaeis guineensis*); mode of root infection and lower stem invasion by *Ganoderma boninense*. *Plant Pathology*, 58: 982–989.
- Saikkonen, K., Wäli, P., Helander, M., and Faeth, S. H. (2004). Evolution of endophyte-plant symbioses. *Trends in Plant Science*, 9: 275–280.
- Sapak, Z., Meon, S., and Ahmad, Z. A. M. (2008). Effect of endophytic bacteria on growth and suppression of *Ganoderma* infection in oil palm. *International Journal of Agriculture and Biology*, 10(2): 127–132.
- Salter, M. G., and Conlon, H. E. (2007). Extraction of plant RNA. *Methods in Molecular Biology*, 362: 309–314.
- Seerangan, K., and Thangavelu, M. (2014). Arbuscular Mycorrhizal and Dark Septate Endophyte Fungal Associations in South Indian Aquatic and Wetland Macrophytes. *Journal of Botany*, 2014: 1–14.
- Shoresh, M., Yedidia, I., and Chet, I. (2005). Involvement of Jasmonic Acid/Ethylene Signaling Pathway in the Systemic Resistance Induced in Cucumber by *Trichoderma asperellum* T203. *Phytopathology*, 95(1): 76–84.
- Spiering, M. J., Greer, D. H., and Schmid, J. (2006). Effects of the fungal endophyte, *Neotyphodium lolii*, on net photosynthesis and growth rates of perennial

ryegrass (*Lolium perenne*) are independent of in planta endophyte concentration. *Annals of Botany*, 98: 379–387.

- Su, Z.Z., Mao, L.J., Li, N., Feng, X-X., Yuan, Z.L., Wang, L.W., and Zhang, C.L. (2013). Evidence for biotrophic lifestyle and biocontrol potential of dark septate endophyte *Harpophora oryzae* to rice blast disease. *PloS One*, 8(4): e61332
- Sundram, S. (2013). First report: Isolation of endophytic *Trichoderma* from oil palm (*Elaeis guineensis* Jacq.) and their *in vitro* antagonistic assessment on *Ganoderma boninense*. *Journal of Oil Palm Research*, 25(3): 368–372.
- Tarkka, M. T., Lehr, N., Hampp, R., and Schrey, S. D. (2008). Plant behavior upon contact with Streptomyces. *Plant signaling & behavior*, 3(11): 917–919.
- Torres, M. S., White, J. F., Zhang, X., Hinton, D. M., and Bacon, C. W. (2012). Endophyte-mediated adjustments in host morphology and physiology and effects on host fitness traits in grasses. *Fungal Ecology*, 5(3): 322–330.
- Tunc-Ozdemir, M., Miller, G., Song, L., Kim, J., Sodek, A., Koussevitzky, S., and Shintani, D. (2009). Thiamin confers enhanced tolerance to oxidative stress in *Arabidopsis*. *Plant Physiology*, 151(1): 421–32.
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., and Speleman, F. (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology*, 3(7): research0034.1–0034.11
- Waller, F., Achatz, B., Baltruschat, H., Fodor, J., Becker, K., Fischer, M., and Kogel, K.-H. (2005). The endophytic fungus *Piriformospora indica* reprograms barley to salt-stress tolerance, disease resistance, and higher yield. *Proceedings of the National Academy of Sciences of the United States of America*, 102(38): 13386–13391.
- Wang, G., Ding, X., Yuan, M., Qiu, D., Li, X., Xu, C., and Wang, S. (2006). Dual function of rice OsDR8 gene in disease resistance and thiamine accumulation. *Plant Molecular Biology*, 60(3): 437–449.
- Wang, X., Anzhi, R., and Yubao, G. (2014). Effect of endophyte infection on fungal disease resistance of *Leymus chinensis*. *Acta Ecol. Sin.* 34, 6789–6796. doi:10.5846/stxb201303080377.
- Wang, L., Ye, Liu, H., Liu, X., Wei, C., Huang, Y., and Tu, J. (2016). Both overexpression and suppression of an *Oryza sativa* NB-LRR-like gene OsLSR result in autoactivation of immune response and thiamine accumulation. *Scientific Reports*, 6(866): 24079.
- Waqas, M., Khan, A. L., Hamayun, M., Shahzad, R., Kim, Y.-H., Choi, K.-S., and Lee, I.-J. (2015). Endophytic infection alleviates biotic stress in sunflower through

regulation of defence hormones, antioxidants and functional amino acids. *European Journal of Plant Pathology*, 141(4), 803–824.

- Wilson, D. (1995). Endophyte: The Evolution of a Term, and Clarification of Its Use and Definition. *Oikos*, 73(2), 274–276.
- Wong, L., Bong, C.H., and Idris, A. S. (2012). *Ganoderma* Species Associated with Basal Stem Rot Disease of Oil Palm. *American Journal of Applied Sciences*, 9(6), 879–885.
- Wong, S. Y., Abdul Aziz, S. D., and Balia Yusof, Z. N. (2016). Osmotic stress upregulates the transcription of thiamine (vitamin B1) biosynthesis genes (THIC and THI4) in oil palm (*Elaeis guineensis*). *African Journal of Biotechnology*, 15(29): 1566–1574.
- Woodward, J. B., Abeydeera, N. D., Paul, D., Phillips, K., Rapala-Kozik, M., Freeling, M., and Scanlon, M. J. (2010). A maize thiamine auxotroph is defective in shoot meristem maintenance. *The Plant Cell*, 22(10): 3305–3317.
- Zeng, Y., and Yang, T. (2002). RNA isolation from highly viscous samples rich in polyphenols and polysaccharides. *Plant Molecular Biology Reporter*, 20(4): 417.
- Zheng, Y. K., Qiao, X.-G., Miao, C.P., Liu, K., Chen, Y.W., Xu, L. H., and Zhao, L.-X. (2015). Diversity, distribution and biotechnological potential of endophytic fungi. *Annals of Microbiology*, 56: 529–542.
- Zhou, J., Sun, A., and Xing, D. (2013). Modulation of cellular redox status by thiamine-activated NADPH oxidase confers *Arabidopsis* resistance to *Sclerotinia sclerotiorum*. *Journal of Experimental Botany*, 64(11): 3261–3272.