

UNIVERSITI PUTRA MALAYSIA

ELECTROCHEMICAL DNA BIOSENSOR BASED ON IRON OXIDE/NANOCELLULOSE CRYSTALLINE FOR DETECTION OF Mycobacterium tuberculosis

CHE ENGKU NORAMALINA BT CHE ENGKU CHIK

FBSB 2018 7



ELECTROCHEMICAL DNA BIOSENSOR BASED ON IRON OXIDE/NANOCELLULOSE CRYSTALLINE FOR DETECTION OF Mycobacterium tuberculosis



CHE ENGKU NORAMALINA BT CHE ENGKU CHIK


Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfillment of the Requirements for the Degree of Master of Science

December 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This research work is dedicated to my lovely husband Mohd Hazim Abd Aziz, my late father, Che Engku Chik Che Engku Embong, my lovely mother, and my siblings, and my family in law.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

ELECTROCHEMICAL DNA BIOSENSOR BASED ON IRON OXIDE/NANOCELLULOSE CRYSTALLINE FOR DETECTION OF Mycobacterium tuberculosis

By

CHE ENGKU NORAMALINA BT CHE ENGKU CHIK

December 2017

Chairman: Helmi Wasoh @ Mohamad Isa,PhDFaculty: Biotechnology and Biomolecular Science

In this study, electrochemical DNA biosensor based on carboxyl functionalized iron oxide/nanocellulose crystalline surface modified with cetyl trymethylammonium bromide (COOH-Fe₃O₄/NCC/CTAB) for detection of Mycobacterium tuberculosis (MTB) was developed. The modification was made by drop coating a carboxyl COOH-Fe₃O₄ nanocomposite with NCC/CTAB on surface of screen printed carbon electrode (SPCE). The modified SPCE was electrochemically characterized using electrochemical impedance spectroscopy (EIS) and cyclic voltammetry (CV) in the present of $Fe(CN)_6^{3/4-}$ as electrolyte. The electrochemical characterization confirmed the high rate of electron transfer, was due to increase of electroactive surface area and excellent conductivity of modified SPCE. The optimization of SPCE modified in term of COOH-Fe₃O₄ concentration, volume/volume ratio of COOH-Fe₃O₄: NCC/CTAB and scan rate was achieved by 0.5 mg/ml, 1:1 ratio and 100 mV/s; respectively. The present of functional groups were confirmed by physical characterization using fourier transform infrared spectroscopy (FTIR), the surface morphology was confirmed by field emission scanning electron microscopy (FE-SEM), and the elemental composition of modified SPCE was confirmed by energy dispersive X-ray spectroscopy (EDX). The deposition of modifier materials on the surface of SPCE was properly characterized through physical characterization. Their electrochemical activities and sensing capability towards detection of Mycobacterium tuberculosis (MTB) was investigated using differential pulse voltammetry (DPV). Ruthenium complex bipyridyl $Ru(bpy)_3^{2+}$ was used as redox indicator in monitoring hybridization events of the DNA. A 5'-NH₂- end of DNA probe was used to form a covalent bond of amide bond immobilized with carboxyl group via ethyl-3-(3-dimethylaminopropyl) carbodiimide/N-hydroxysuccinimide EDC/NHS as coupling mechanism. The selectivity of the fabricated DNA sensor was confirmed by the different signal produced by $Ru(bpy)_3^{2+}$ in differentiate ssDNA

probe, dsDNA target complementary, non-complementary and the mismatch DNA sequences. The optimum condition were obtained using the following criteria such as Tris EDTA as supporting electrolyte pH 8, deposition potential (0.3V), deposition time (5 sec) probe DNA concentration, immobilization time (60 minutes), EDC/NHSS immersion, ionic strength (20 mM), hybridization time (35 minutes at 42°C) and Ru(bpy)₃²⁺ concentration (25 μ M). Performance of the biosensor was investigated using different concentration of DNA target in the range of 1.0×10^{-12} to 1.0×10^{-6} M and the limit of detection was calculated to be at 7.96 x 10^{-13} M. The calibration curve showed a linear relationship with equation, $y=0.6936 \log (x) +$ 11.47 and correlation coefficient of R²=0.9896. DNA of MTB was extracted from clinical sample and underwent PCR procedure for real sample analysis. The developed biosensor was demonstrated the difference between positive and negative MTB sample, mycobacterium other than tuberculosis (MOTT) and respiratoryrelated bacterial samples, Staphyloccocus aureus and confirmed that this fabricated electrochemical DNA sensor have good selectivity and effectively detected the sequence of DNA from MTB using PCR product analysis. The studied of stability and reproducibility showed a satisfactory result with loss only 5.87% activity after 7 days and below 50% after 5 weeks and finally the reproducibility value of 3.87% (n=8). The repeatability showed quite high with value of 29.30% indicating to a good SPCE to be used for TB detection if compared to previous study.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

DNA BIOSENSOR ELEKTROKIMIA BERASASKAN BESI BEROKSIDA BERFUNGSI KARBOKSIL/NANOCELULOSA KRISTAL UNTUK PENGESANAN Mycobacterium tuberculosis

Oleh

CHE ENGKU NORAMALINA BT CHE ENGKU CHIK

Disember 2017

Pengerusi: Helmi Wasoh @ Mohamad Isa, PhDFakulti: Bioteknologi dan Sains Biomolekul

Dalam kajian ini, elektrokimia biosensor DNA berdasarkan besi oksida difungsikan dengan kumpulan karboksil/nanocelulosa kristal diubahsuai dengan siltrimetilammonium bromida (COOH-Fe₃O₄/NCC/CTAB) untuk mengesan Mycobacterium tuberculosis (MTB) telah dibangunkan. Pengubahsuaian itu dibuat melalui salut titis COOH-Fe₃O₄ dan campuran NCC/CTAB pada permukaan paparan bercetak elektrod karbon (SPCE). SPCE yang telah diubahsuai dicirikan menggunakan kitaran voltametri (CV) dan spektroskopi impedans elektrokimia (EIS) dengan kehadiran $Fe(CN)_6^{3/4-}$ sebagai elektrolit. Pencirian elektrokimia mengesahkan kadar pemindahan elektron yang tinggi, peningkatan kawasan permukaan elektroaktif dan kekonduksian SPCE yang diubahsuai. Pengoptimuman SPCE yang diubah suai dari segi kepekatan COOH-Fe₃O₄, nisbah isipadu/isipadu COOH-Fe₃O₄:NCC/CTAB dan kadar imbasan dicapai oleh 0.5 mg/ml, nisbah 1:1 dan 100 mV/s. Pencirian fizikal telah dikaji menggunakan 'fourier transform infrared spectroscopy' (FTIR) mengesahkan semua kehadiran kumpulan berfungsi, bidang imbasan pancaran elektron mikroskop (FE-SEM) mengesahkan morfologi permukaan, dan serakan tenaga X-ray spektroskopi (EDX) mengesahkan komposisi unsur SPCE yang diubahsuai. Pemendapan bahan pengubah di permukaan SPCE telah disahkan melalui pencirian fizikal. Kegiatan elektrokimia dan keupayaan mengesan MTB diselidiki menggunakan 'differential pulse voltammetry' (DPV). Ruthenium kompleks bipyridyl $Ru(bpy)_3^{2+}$ digunakan sebagai penunjuk redoks dalam memantau peristiwa hibridisasi DNA. DNA yang telah diubahsuai dengan 5'-NH2- telah digunakan untuk membentuk satu ikatan kovalen amida dengan kumpulan karboksil melalui ethyl-3-(3-dimethylaminopropyl) carbodiimide/Nhydroxysuccinimide (EDC/NHS) sebagai mekanisme gandingan. Pemilihan sensor DNA yang dihasilkan telah disahkan oleh isyarat yang berbeza yang dihasilkan oleh Ru(bpy)₃²⁺ dalam membezakan prob ssDNA, sasaran dsDNA pelengkap, DNA tidak



melengkap dan DNA dengan turutan yang salah. Keadaan optimum yang dicapai adalah ; Tris EDTA dengan pH 8 sebagai elektrolit sokongan, pemendapan potensi dan masa di 0.3 V dan 5 saat masing-masing, 10 µM kepekatan prob DNA dengan 60 minit DNA untuk immobilisasi, 10 minit rendaman dalam EDC/NHS, 20 mM kekuatan ionik, masa hibridisasi selama 35 minit pada suhu 42 °C dan kepekatan dan 25 μ M dari Ru(bpy)₃²⁺. Prestasi biosensor disiasat menggunakan kepekatan DNA yang berbeza dalam julat 1.0 x 10⁻¹² hingga 1.0 x 10⁻⁶ M dan batas pengesanan dikira pada 7.96 x 10⁻¹³ M. Lengkok penentukuran menunjukkan hubungan linear dengan y = 0.6936 log(x) + 11.47 dan pekali korelasi R^2 = 0.9896. DNA MTB telah diekstrak daripada sampel klinikal dan menjalani prosedur PCR untuk analisis sampel sebenar. Biosensor yang dihasilkan telah menunjukkan perbezaan antara sampel MTB positif dan negatif, mikobakteria selain tuberkulosis (MOTT) dan sampel bakteria yang berkaitan dengan pernafasan, Staphyloccocus aureus mengesahkan bahawa fabrikasi DNA sensor elektrokimia ini mempunyai pemilihan yang baik dan berkesan mengesan produk PCR MTB. Kajian kestabilan menunjukkan hasil yang memuaskan dengan kehilangan aktiviti cuma sebanyak 5.87% selepas 7 hari dan kehilangan aktiviti di bawah 50% selepas 5 minggu. Manakala kebolehulangan pada elektrod yang berbeza nmempunyai nilai 3.87% (n=8). Kebolehulangannya pada elektrod yang sama menunjukkan nilai yang agak tinggi dengan 29.30% menunjukkan prestasi SPCE yang baik sebagai pengesan MTB jika dibandingkan dengan kajian terdahulu.

ACKNOWLEDGEMENTS

Bismillahirrahmanirrahim,

First of all, I am so grateful to Allah the Almighty for the health, wisdom and strength to finish this project. Secondly, I want to express my profound gratitude for the continous support, guidance, advice, encouragement received from my supervisor, Dr Helmi Wasoh @ Mohamad Isa from the beginning of this research journey and till the end.

Similar appreciation goes to my co supersivor, Prof Dr. Nor Azah Yusof for the financial supports, Dr Jaafar Abdullah; for the valuable comments and ideas, Dr Sarah Othman; for the guidance in molecular works. Their contribution is the reason for this research to the succesful conclusion.

Thirdly, a big thanks goes to Dr Rahizan Isa, Mohd Fairulnizal Md Noh and the staff at Department of Bacteriology, Institute of Medical Research for their expert guidance in handling Tuberculosis (TB) clinical samples. I would also like to acknowledge with much appreciation to my colleagues and friends for making this research journey a memorable one.

Last but not least, special thanks to my entire family for their patience while I am away for this course of study, especially my husband, Mohd Hazim Abd Aziz who stood by my side to see the successful of this research. Not to forget, my mother, my siblings and my family in laws for their kind supports, encouragement and nurture given to me up to this stage, may Allah reward them with Jannah. I certify that a Thesis Examination Committee has met on 13 December 2017 to conduct the final examination of Che Engku Noramalina bt Che Engku Chik on her thesis entitled "Electrochemical DNA Biosensor Based on Iron Oxide/Nanocellulose Crystalline for Detection of *Mycobacterium tuberculosis*" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Siti Aqlima binti Ahmad, PhD

Senior Lecturer Faculty of Biotechnology and Biomolecular Sciences Universiti Putra Malaysia (Chairman)

Mohd. Yunus bin Abd. Shukor, PhD

Associate Professor Faculty of Biotechnology and Biomolecular Sciences Universiti Putra Malaysia (Internal Examiner)

Mohd Khairul Ahmad, PhD

Associate Professor Universiti Tun Hussein Onn Malaysia Malaysia (External Examiner)



NOR AINI AB. SHUKOR, PhD Professor and Deputy Dean School of Graduate Studies Universiti Putra Malaysia

Date: 29 January 2018

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Helmi Wasoh @ Mohamad Isa, PhD

Senior Lecturer Faculty of Biotechnology Universiti Putra Malaysia (Chairperson)

Nor Azah Yusof, PhD

Professor Faculty of Science Universiti Putra Malaysia (Member)

Jaafar Abdullah, PhD

Senior Lecturer Faculty of Science Universiti Putra Malaysia (Member)

Siti Sarah Othman, PhD Senior Lecturer Faculty of Biotechnology

Universiti Putra Malaysia (Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean School of Graduate Studies Universiti Putra Malaysia

Date :

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

| Signature: | Date: |
|------------|-------|
| | |

Name and Matric No: Che Engku Noramalina Bt Che Engku Chik, GS43758

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

| Signature: Name of Chairmar of Supervisory | |
|--|-------------------------------|
| Committee: | Dr. Helmi Wasoh @ Mohamad Isa |
| Signature: Name of Member of Supervisory | |
| Committee: | Professor Dr. Nor Azah Yusof |
| Signature: Name of Member of Supervisory Committee: | Dr. Jaafar Abdullah |
| Signature Name of Member of Supervisory Committee: | Dr. Siti Sarah Othman |
| | |

TABLE OF CONTENTS

Page

| A | BSTRA | CT | i |
|---|----------|---|--------|
| A | BSTRAK | | iii |
| Α | CKNOV | VLEDGEMENTS | v |
| Α | PPROV | AL | vi |
| D | DECLER | ATION | viii |
| L | IST OF | TABLES | xiv |
| L | IST OF | FIGURES | xv |
| L | IST OF A | ABBREVIATIONS | xviii |
| C | CHAPTE | | |
| 1 | INTE | RODUCTION | 1 |
| 1 | 1.1 | Background | 1 |
| | 1.1 | Problem statement | |
| | 1.2 | The objectives of this study: | 2 3 |
| | 1.5 | The objectives of this study. | 5 |
| 2 | LITE | ERATURE REVIEW | 4 |
| - | 2.1 | Tuberculosis | 4 |
| | | 2.1.1 Characteristic of <i>Mycobacterium tuberculosis</i> (MTB) | 4 |
| | | 2.1.2 Symptoms of Tuberculosis (TB) | 5 |
| | | 2.1.3 Cases of Tuberculosis (TB) globally | 5 |
| | | 2.1.4 Cases of Tuberculosis (TB) in Malaysia | 6 |
| | 2.2 | Detection of Tuberculosis (TB) using conventional methods | 8 |
| | | 2.2.1 Gold Standard Method | 8 |
| | | 2.2.1.1 Specimen collection | 8 |
| | | 2.2.1.2 Culture of <i>Mycobacterium tuberculosis</i> (MTB) | 9 |
| | | 2.2.2 Acid Fast Bacilli (AFB) Staining | 9 |
| | | 2.2.2.1 Carbolfuchsin method | 9 |
| | | 2.2.2.2 Fluorochrome staining method | 10 |
| | | 2.2.3 Conventional Polymerase Chain Reaction (PCR) | 11 |
| | 2.3 | Electrochemical DNA biosensor as alternatives for TB detection | 11 |
| | | 2.3.1 Basic principles of biosensor | 11 |
| | | 2.3.2 Electrochemical Biosensor | 12 |
| | | 2.3.3 Electrochemical DNA biosensor configuration | 13 |
| | 2.4 | Voltammetry technique for electrochemical DNA biosensor | 15 |
| | | 2.4.1 Cyclic voltammetry | 15 |
| | | 2.4.2 Differential pulse voltammetry (DPV) | 17 |
| | 2.5 | Deoxyribonucleic acid (DNA) as bioreceptor | 17 |
| | 2.6 | Specific DNA Insertion Sequence IS6110 of MTB | 19 |
| | 2.7 | Technique of DNA Immobilization on transducer | 20 |
| | 2.8 | SPCE as transducer for DNA biosensor | 20 |
| | 2.9 | Nanomaterial for SPCE's modification | 21 |
| | 2.10 | Iron oxide (Fe ₃ O ₄) | 21 |

| | 2.11 | Nanoce | llulose cry | ystalline (NCC) | 22 |
|---|------|---------------------|-------------|---|----------|
| | 2.12 | DNA hy | ybridizatio | on detection methods | 22 |
| | | 2.12.1 | Rutheniu | im complex | 23 |
| 3 | MFT | HODOI | OGV | | 24 |
| 5 | 3.1 | Materia | | | 24 |
| | 5.1 | | Reagents | | 24 |
| | | | Iron Oxi | | 25 |
| | | 3.1.2 | | ion of COOH-Fe ₃ O ₄ | 25 |
| | | 3.1.4 | - | ion of NCC/CTAB | 26 |
| | | 3.1.5 | - | ion of COOH-Fe ₃ O ₄ /NCC/CTAB | 26 |
| | | 3.1.6 | - | ion of other solutions | 26 |
| | | 3.1.7 | - | ion of oligonucleotide solution | 26 |
| | | 3.1.8 | - | ion of sputum specimen from clinical samples | 27 |
| | | 3.1.9 | - | on of reagent for DNA extraction | 27 |
| | | | - | ion of Mastermix for PCR amplification | 28 |
| | | 3.1.11 | - | ion of gel electrophoresis | 28 |
| | 3.2 | Electroc | | nentation and kits | 28 |
| | | 3.2.1 | Screen p | rinted carbon electrode (SPCE) | 29 |
| | | 3.2.2 | Voltamm | netry analyzer | 29 |
| | | 3.2.3 | Electrocl | hemical Impedance Spectroscopy (EIS) | 29 |
| | | 3.2.4 | Fourier t | ransform infrared spectroscopy (FTIR) | 29 |
| | | 3.2 <mark>.5</mark> | Field em | ission scanning electron microscopy (FE-SEM) | 29 |
| | | 3. <mark>2.6</mark> | Energy d | lispersive X-Ray Spectroscopy (EDX) | 30 |
| | | 3.2.7 | U 1 | e Viral DNA Extraction Kit | 30 |
| | | 3.2.8 | | Aultiplex PCR Kit | 30 |
| | | 3.2.9 | | ycler (Mastercycler Gradient, Eppendorf) | 30 |
| | | 3.2.10 | | umentation System (BioRad) | 30 |
| | 3.3 | Procedu | | | 31 |
| | | 3.3.1 | - | ting COOH-Fe ₃ O ₄ / NCC/CTAB on SPCE | 31 |
| | | 3.3.2 | - | ation of COOH-Fe ₃ O ₄ /NCC/CTAB on SPCE | 31 |
| | | | | Effect of different concentration of COOH-Fe ₃ O ₄ Effect of volume/volume ratio | |
| | | | 3.3.2.2 | Effect of different scan rates | 31 31 |
| | | 3.3.3 | | ization of DNA probe on modified SPCE | 31 |
| | | 3.3.4 | | ation with DNA target, DNA non | 52 |
| | | 5.5.4 | • | entary, DNA mismatch | 32 |
| | | 3.3.5 | 1 | ation of Electrochemical DNA Sensor | 32 |
| | | 0.010 | 3.3.5.1 | Effect of supporting electrolyte | 32 |
| | | | 3.3.5.2 | Influence of pH | 32 |
| | | | 3.3.5.3 | Effect of deposition potential and deposition time | 33 |
| | | | 3.3.5.4 | Effect of DNA probe concentration | 33 |
| | | | 3.3.5.5 | Effect of ionic strength | 33 |
| | | | 3.3.5.6 | Effect of time immersion in EDC/NHSS | 33 |
| | | | 3.3.5.7 | Effect of time immobilization of DNA probe | 33 |
| | | | 3.3.5.8 | Effect of DNA hybridization temperature and time | |
| | | | 3.3.5.9 | Effect of Ruthenium complex concentration | 34 |
| | | | | | |

 (\mathbf{C})

| | | 3.3.5.10 | Effect of different complementary DN | IA target | |
|-----|----------------------------------|--------------|--|--------------|-----|
| | | | concentration | - | 34 |
| | 3.3.6 | Validatio | on of developed biosensor sensitivity to | owards | |
| | | MTB DN | VA from clinical sample | | 34 |
| | | | DNA extraction from Clinical Specim | ens | |
| | | | Using High Pure Viral Nucleic Acid F | | 34 |
| | | 3.3.6.2 | Procedure for PCR amplification | | 35 |
| | | 3.3.6.3 | Procedure for Gel Electrophoresis | | 35 |
| | | 3.3.6.4 | Application of developed DNA bioser | nsor towards | |
| | | | clinical sample's PCR product | | 35 |
| | 3.3.7 | Reprodu | cibility, Repeatability and Stability of | developed | |
| | | sensor | , , , , , , , , , , , , , , , , , , , | | 35 |
| | | ~ | | | |
| RES | ULTS A | ND DISC | USSION | | 36 |
| 4.1 | Propose | ed mechan | ism of SPCE modified COOH-Fe ₃ O ₄ / | NCC/CTAB | 36 |
| 4.2 | | | characterization of modified SPCE with | | |
| | Fe ₃ O ₄ / | NCC/CTA | B using cyclic voltammetry (CV) | | 37 |
| 4.3 | | | SPCE modified with COOH-Fe ₃ O ₄ /NC | C/CTAB | |
| | - | yclic volta | | | 39 |
| | | | varying concentration of COOH-Fe ₃ O | 04 | 39 |
| | | | varying volume/volume ratio of COO | | |
| | | NCC/CT | | | 40 |
| | 4.3.3 | | varying scan rates | | 41 |
| 4.4 | | | characterization by Electrochemical | | • • |
| | | | roscopy (EIS) | | 44 |
| 4.5 | - | - | erization of modified SPCE | | 45 |
| | 4. <u>5</u> .1 | | ransforms infrared spectroscopy (FTIR | 0 | 45 |
| | 4.5.2 | | ission scanning electron microscopy (H | , | 47 |
| | 4.5.3 | | Dispersive X-Ray Spectroscopy (EDX) | , | 48 |
| 4.6 | | | characterization of DNA biosensor for | | |
| | detectio | | | | 50 |
| | 4.6.1 | | supporting electrolyte and its pH | | 50 |
| | 4.6.2 | | deposition potential and time | | 53 |
| | 4.6.3 | | ionic strength | | 55 |
| | 4.6.4 | | time immersion in EDC/NHSS | | 56 |
| | 4.6.5 | | ssDNA probe concentration | | 57 |
| | 4.6.6 | | time immobilization of DNA probe | | 58 |
| | 4.6.7 | | temperature on DNA hybridization | | 58 |
| | 4.6.8 | | DNA hybridization time | | 60 |
| | 4.6.9 | | Ruthenium complex concentration on | fabricated | 00 |
| | | DNA bio | - | | 61 |
| | 4.6.10 | | DNA target concentration | | 61 |
| 4.7 | | | ensitivity of the developed electrochem | ical DNA | 01 |
| 1.7 | sensor | , ity and se | montrie of the developed electrochem | | 64 |
| 4.8 | | ion of DN | A biosensor towards real samples by F | PCR | 65 |
| 4.9 | | | Repeatability and Stability | | 67 |
| | reprou | actonity, I | copoundonity and bradinity | | 57 |

4

Ģ

| 5 | CON | CLUSION AND RECOMMENDATIONS | 70 |
|---|---------------|-----------------------------|----------------------|
| | 5.1 | Conclusion | 70 |
| | 5.2 | Recommendations | 70 |
| | NDIC ATA (| | 71 84 87 88 |



LIST OF TABLES

| Table | | Page |
|-------|--|------|
| 2.1 | Percentage of regional incidence TB cases in 2015 | 6 |
| 2.2 | Breakdown of TB cases and death according to sates in Malaysia | 7 |
| 2.3 | TB cases among foreign workers from 2010-2014 in Malaysia | 8 |
| 2.4 | Biosensor classification based on transducer | 12 |
| 2.5 | Previous studied reported based on electrochemical DNA biosensor for TB detection | 14 |
| 3.1 | Reagents and solvents used in this works | 25 |
| 3.2 | Oligonucleotide sequences used in this works | 27 |
| 3.3 | Mastermix preparation | 28 |
| 4.1 | The peak current from SPCEs of different modification | 38 |
| 4.2 | Electroactive surface area of modified SPCEs | 43 |
| 4.3 | Compar <mark>ison of different performance of different fabrication of DNA biosensor reported in literature</mark> | 62 |
| 4.4 | The reproducibility of fabricated biosensor | 68 |
| 4.5 | The repeatability of fabricated biosensor | 68 |
| 4.6 | The stability of fabricated biosensor | 69 |
| A1 | Detection of current peaks at different target DNA concentration | 86 |
| A2 | Detection of current peaks at without DNA (Blank) concentration | 86 |
| | | |

LIST OF FIGURES

| Figu | re | Page |
|------|---|------|
| 2.1 | Basic biosensor mechanism | 12 |
| 2.2 | Electrochemical DNA biosensor configuration | 14 |
| 2.3 | Excitation signals for cyclic voltammetry showing triangle potential waveform | 15 |
| 2.4 | Cyclic voltammogram for a single scan redox process under reversible electron transfer | 16 |
| 2.5 | DNA molecular structure | 18 |
| 2.6 | Watson Crick based pairing | 19 |
| 2.7 | Screen printed carbon electrode (SPCE) | 21 |
| 4.1 | Mechanism of carboxyl functionalization of Fe ₃ O ₄ | 36 |
| 4.2 | Mechanism of surface modification of Nanocellulose crystalline (NCC) by surfactant CTAB | 37 |
| 4.3 | Cyclic voltammetry behavior of (A) bare SPCE, (B) SPCE/Fe ₃ O ₄ , (C) SPCE/COOH-Fe ₃ O ₄ , (D) SPCE/NCC/CTAB, and (E) SPCE/COOH-Fe ₃ O ₄ /NCC/CTAB in 5 mM K ₃ Fe(CN) ₆ /0.1 M KCI with scan rate 100 mV/s | 38 |
| 4.4 | Bar graph for peak current of modified SPCE at different concentration of COOH-Fe ₃ O ₄ | 40 |
| 4.5 | Bar graph for peak current of modified SPCE at different volume/volume ratio of COOH-Fe ₃ O ₄ :NCC/CTAB | 41 |
| 4.6 | Graph for the cyclic voltammograms and current versus square root of scan rates (0.01-0.1 V/s) of (a) bare SPCE (b) SPCE/COOH-Fe ₃ O ₄ (c) SPCE/NCC/CTAB (d) SPCE/COOH-Fe ₃ O ₄ /NCC/CTAB modifications in a 5 mM K ₃ [Fe(CN) ₆]/0.1 M KCI | 43 |
| 4.7 | The Nyquist plot of (a) bare SPCE, (b) SPCE/Fe ₃ O ₄ , (c), SPCE/COOH-Fe ₃ O ₄ , (d) SPCE/NCC/CTAB and (e) SPCE/COOH-Fe ₃ O ₄ /NCC/CTAB in 5.0 mM K_3 Fe(CN) ₆ /0.1 M KCI with frequency 10000 Hz and amplitude 0.01 V | 45 |

| 4.8 | FTIR spectra for (A) Fe ₃ O ₄ , (B) COOH-Fe ₃ O ₄ , (C) NCC, (D) NCC/CTAB, and (E) COOH-Fe ₃ O ₄ /NCC/CTAB | 46 |
|------|--|----|
| 4.9 | FESEM of (A) bare SPCE (B) SPCE/Fe ₃ O ₄ , (C) SPCE/COOH-Fe ₃ O ₄ , (D) SPCE/NCC, (E) SPCE/NCC/CTAB, and (F) COOH-Fe ₃ O ₄ /NCC/CTAB | 47 |
| 4.10 | The EDX spectra of (A) SPCE (B) SPCE/Fe ₃ O ₄ (C) SPCE/COOH-Fe ₃ O ₄ (D) SPCE/NCC/CTAB (E) SPCE/COOH-Fe ₃ O ₄ /NCC/CTAB | 49 |
| 4.11 | CV and DPV of modified SPCE in detection 1μ M of complementary DNA target in (ai-aii) PBS, (bi-bii) Tris NaCl and (ci-cii) TE buffer in Ru(bpy) ₃ ²⁺ | 51 |
| 4.12 | (A) The bar graph and (B) the peak current of pH influence to the supporting electrolyte | 53 |
| 4.13 | Effect of deposition potential on the modified SPCE in detection $1\mu M$ of complementary DNA target | 54 |
| 4.14 | Effect of deposition time on the modified SPCE in detection $1\mu M$ of complementary DNA target | 54 |
| 4.15 | The effect of ionic strength on the modified SPCE in detection $1\mu M$ of complementary DNA target | 55 |
| 4.16 | The effect of time immersion in EDC/NHSS on the modified SPCE in detection 1μ M of complementary DNA target | 56 |
| 4.17 | Effect of ssDNA probe concentration on the modified SPCE in detection 1 μ M of complementary DNA target | 57 |
| 4.18 | Effect of Immobilization time of DNA probe on the modified SPCE in detection $1\mu M$ of complementary DNA target | 58 |
| 4.19 | Effect of of DNA hybridization temperature on the modified SPCE in detection $1\mu M$ of complementary DNA target | 59 |
| 4.20 | Effect of DNA hybridization time on the modified SPCE in detection $1\mu M$ of complementary DNA target | 60 |
| 4.21 | Effect of Ruthenium complex concentration on fabricated DNA biosensor | 61 |
| 4.22 | DPVs of the electrochemical biosensor for target detection at concentrations between a to g (1.0×10^{-6} to 1.0×10^{-12} M) | 63 |

 \bigcirc

| 4.23 | The calibration plot of DPV peak current vs. log DNA concentration (Molar) ranges between 1.0×10^{-12} to 1.0×10^{-6} M | 63 |
|------|---|----|
| 4.24 | DPV curve for oxidation of $Ru(bpy)_3^{2+}$ in TE buffer (pH 8) containing 25 μ M [Ru(bpy) ₃] ²⁺ | 64 |
| 4.25 | Gel electrophoresis photographs of PCR products obtained from D1 to D9 | 65 |
| 4.26 | DPV curves for the detection of PCR product D1 to D9 | 66 |
| 4.27 | DPV peak currents responding to the PCR products obtained from D1 to D9 | 67 |
| | UPM | |

C

LIST OF ABBREVIATIONS

| Fe ₃ O ₄ | Iron Oxide |
|--------------------------------|---|
| СООН | Carboxyl |
| MPA | Mercaptopropanoic acid |
| NCC | Nanocellulose Crystalline |
| CTAB | cetyl trymethylammonium bromide |
| MTB | Mycobacterium Tuberculosis |
| ТВ | Tuberculosis |
| MOTT | Mycobacterium other than tuberculosis |
| SPCE | Screen printed carbon electrode |
| EIS | Electrochemical impedance spectroscopy |
| CV | Cyclic voltammetry |
| DPV | Differential pulse voltammetry |
| FTIR | Fourier transform infrared spectroscopy |
| FE-SEM | Field emission scanning electron microscopy |
| EDX | Energy dispersive X-ray spectroscopy |
| EDC/NHS | Ethyl-3-(3-dimethylaminopropyl) carbodiimide/N-hydroxysuccinimide |
| ssDNA | Single stranded deoxyribonucleic acid |
| dsDNA | Double stranded deoxyribonucleic acid |
| CE | Counter electrode |
| WE | Working electrode |
| RE | Reference electrode |
| g | Gram |

- mg Milligram
- M Molar
- μM Micromolar
- A Ampere
- μA Microampere
- µl Microliter

V

G

°C Degree celcius

Volt

CHAPTER 1

INTRODUCTION

1.1 Background

Mycobacterium tuberculosis (MTB) is an infectious agent, which is responsible for causing Tuberculosis (TB). According to the World Health Organization (WHO), 10.4 million TB cases with 1.4 million death cases reported globally in 2015 (WHO, 2016). An effective diagnosis is considered crucial for TB control, both for the treating of the infected individual as well as for the intervention of the public to decrease further spreading (Ryu, 2015).

Nowadays, biosensor technology for improving global health issues is in great demand especially in the diagnosis of critical diseases. Biosensor has been introduced to be studied extensively for their role in detecting different analytes in clinical analysis and medical diagnosis. The first biosensor was invented by Clark and Lyons in 1962 using the concept of 'soluble' enzyme electrode which is considered as the marker for biosensor technology advancement (Tehrani & Bavarian, 2016).

In the past two decades, the interest towards DNA biosensors is increased due to its ability to detect specific nucleic acid hybridization. Conventionally, the nucleic acid hybridization was applied in various methods, such as southern blotting and polymerase chain reaction (PCR). By taking advantages of different organism's genomic sequence, it leads the ways for numbers of research opportunities especially in medical diagnosis to detect different diseases especially the diseases originated from bacteria.

Electrochemical DNA biosensor method was introduced to overcome the deficiencies of conventional method such as culturing and smear spectroscopy which is laborious, time-consuming and often susceptible to errors. Interestingly, this DNA biosensor works by immobilizing single stranded DNA (ssDNA) onto electrochemical transducer to be recognized by target complementary ssDNA (Rahman et al., 2015). The ability of biosensor system to detect various concentration of analyte shows their level of sensitivity, however, in the diagnosis of critical diseases more attention has to be given as several procedures are lacking and need improvements.

Additional knowledge of different nanomaterial's capability is very crucial in searching the best nanomaterial that can improve performance of biosensor. In this work, to improve the electrical conductivity of the biosensor system, a negatively charged functionalized COOH-Fe₃O₄, and positively charged NCC/CTAB were



introduced as sensor platform. The used of nanocomposites materials based on carboxyl functionalized iron oxide/nanocellulose crystalline/cetyl trimethylammonium bromide (COOH-Fe₃O₄/NCC/CTAB) were considered as one of additional knowledge to contribute to the promising electrochemical DNA biosensor development. In this project the overall hybridization event for the detection of *Mycobacterium tuberculosis* (MTB) was monitored using ruthenium bipyridyl complex as redox indicator.

1.2 Problem statement

Mantoux tuberculin skin test (TST) is the first evaluation method for TB detection. Good result for this test requires 48 to 72 hours incubation period. Besides, the test needs to be done by a trained health worker to reduce error. Chest radiograph also one of the method for TB evaluation to detect chest abnormalities from the lesion that appears in the lungs. Unfortunately, the present of lesion might be mistakenly diagnosed for other lung diseases, and therefore it cannot be used definitely to diagnose TB.

For better MTB confirmation, acid fast bacilli staining and microbial culture technique has been considered as the 'gold standard' method (Cross et al., 2015) for detection (Kritski et al., 2015). Although, acid fast bacilli staining is quick and simple, confirmation using microbial culture technique is still required as it shows low and variable sensitivity. However, a culture technique requires a lot of time as it involves cell culture of a slow growing mycobacterium (Pullan et al., 2016). Currently, many methods have been introduced to detect the TB disease rapidly by developing low time consumption and sensitive diagnosis including a molecular diagnosis method (Tiwari et al., 2017). Nonetheless, despite all advantages, the complex instrumentation and the cost for its reagent are still considered as the main obstacles for these methods.

Recently, as the sensing technology was improved, various studied based on biosensor in clinical diagnosis had been reported includes electrochemical, piezoelectric, and optical biosensor. Electrochemical method has attracted more attention (Arduini et al., 2016; Justino et al., 2016; Yang et al., 2015) due to its characteristics of simple, low-cost, rapid response, and offer ease of operation (Liu et al., 2014). The detection in variety of analytes such as enzymes, proteins and DNA was also observed to be more possible.

The performance of biosensor was improved through application of screen printed carbon electrode (SPCE) which offer capability for surface modification (Kurbanoglu et al., 2017) plus additional advantages such as low cost, disposable, mass production, and portable (Mat Zaid et al., 2016). Another strategy was adopted through modification of SPCE's surface to improve the analytical performance. By introducing nanomaterial on the surface of SPCE, the biocompatibility, additional binding sites and signal intensities are hoping to be subsequently improved as well as

the sensitivity and specificity of MTB detection. Therefore, interest in the electrochemical DNA biosensor has been considered as one of the efforts to improve the detection of TB (Zaffino et al., 2015).

1.3 The objectives of this study

The aim of this study is to develop a simple electrochemical DNA biosensor for the detection of specific DNA sequence related to MTB assisted by ruthenium complex. The sensor incorporates the use of carboxyl functionalized iron oxide nanoparticles (COOH-Fe₃O₄) with nanocellulose crystalline surface which was modified by cetyl trimethylammonium bromide (NCC/CTAB) for performance enhancement. It is very interesting to explore the potential of this nanoparticles as it has not been reported in any previous TB detection.

The objectives set to achieve these goals include:

- i. To develop electrochemical DNA sensor based on carboxyl functionalized iron oxide/nanocellulose crystalline/cetyl trimethylammonium bromide (COOH-Fe₃O₄/NCC/CTAB) modified screen printed carbon electrode (SPCE).
- ii. To evaluate the sensing capability of the specific gene sequence of DNA in the detection of *Mycobacterium tuberculosis* (MTB) through the interaction between ruthenium complexes $Ru(bpy)_3^{2+}$ with DNA molecules.
- iii. To apply and validate the fabricated biosensor on the analysis of TB real sample.

REFERENCES

- Abu-Salah, K. M., Zourob, M. M., Mouffouk, F., Alrokayan, S. A., Alaamery, M. A., & Ansari, A. A. (2015). DNA-based nanobiosensors as an emerging platform for detection of disease. *Sensors (Switzerland)*, 15(6), 14539–14568.
- Alawi, S. M. (2010). Thermodynamics studies of cetyltrimethylammonium bromide (CTAB) in N, N-dimethyl acetamide-water mixtures, *Oriental Journal of Chemistry*, 26(4), 1235–1240.
- Alberts, B., Johnson, A., & Lewis, J. (2002). The Structure and Function of DNA. Molecular Biology of the Cell. (4th edition). New York: Garland Science
- Amir, F. Z., Pham, V. H., & Dickerson, J. H. (2015). RSC Advances Facile synthesis of ultra-small ruthenium oxide nanoparticles anchored on reduced graphene oxide nanosheets for high-performance supercapacitors. *RSC Advances*, 5, 67638–67645.
- Ang, L. F., Por, L. Y., & Yam, M. F. (2015). Development of an Amperometric-Based Glucose Biosensor to Measure the Glucose Content of Fruit, *PLoS ONE*, 10(3): e0111859. doi:10.1371/journal. pone.0111859
- Arduini, F., Micheli, L., Moscone, D., Palleschi, G., Piermarini, S., Ricci, F., & Volpe, G. (2016). Electrochemical biosensors based on nanomodified screenprinted electrodes: Recent applications in clinical analysis. *TrAC - Trends in Analytical Chemistry*, 79, 114–126.
- Asmar, S., & Drancourt, M. (2015). Chlorhexidine decontamination of sputum for culturing Mycobacterium tuberculosis. *BMC Microbiology*, *15*(1), 155.
- Benedict, N. (2014). TB kills more annually than dengue, says ministry. *Malay Mail Online*, pp. 1–7. Retrieved from http://www.themalaymailonline.com/malaysia/article/tb-kills-more-annuallythan-dengue-says-ministry
- Bhattarai, A. (2015). Study of critical micelle concentration of cetyltrimethylammonium bromide (CTAB) in pure wa ter in presence and absence of magnesium sulphate and sodium sulph ate by measuring conductivity meter, (August). http://doi.org/10.3126/bibechana.v11i0.10390
- Carniel, F., Dalla Costa, E. R., Lima-Bello, G., Martins, C., Scherer, L. C., & Rossetti, M. L. (2014). Use of conventional PCR and smear microscopy to diagnose pulmonary tuberculosis in the Amazonian rainforest area. *Brazilian Journal of Medical and Biological Research*, 47(12), 1016–1020.
- Centers for Disease Control and Prevention. (2016). Testing & Diagnosis. Retrieved from http://www.cdc.gov/tb/topic/testing/default.htm

- Central TB Division. (2009). Training Manual for Mycobacterium tuberculosis Culture & Drug susceptibility testing, Central TB Division, Ministry of Health and Family Welfare, Government of India.
- Chen, M., Hou, C., Huo, D., Fa, H., & Zhao, Y. (2017). Sensors and Actuators B : Chemical A sensitive electrochemical DNA biosensor based on threedimensional nitrogen-doped graphene and Fe₃O₄ nanoparticles. Sensors & Actuators: B. Chemical, 239, 421–429.
- Coros, A., DeConno, E., & Derbyshire, K. M. (2008). IS6110, a Mycobacterium tuberculosis Complex-Specific Insertion Sequence, Is Also Present in the Genome of Mycobacterium smegmatis, Suggestive of Lateral Gene Transfer among Mycobacterial Species. Journal of Bacteriology, 190(9), 3408–3410.
- Crick, F. H. C., & Watson, J. D. (1954). The Complementary Structure of Deoxyribonucleic Acid. Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences, 223(1152), 80–96.
- Cross, L. J., Anscombe, C., Mchugh, T. D., Abubakar, I., Shorten, R. J., Thorne, N., & Arnold, C. (2015). A Rapid and Sensitive Diagnostic Screening Assay for Detection of Mycobacteria Including Mycobacterium tuberculosis Directly from Sputum without Extraction, *International Journal of Bacteriology*. 2015. http://doi.org/http://dx.doi.org/10.1155/2015/593745
- de Lyra, J. M. A., Maruza, M., Verza, M., Carneiro, M. M., de Albuquerque, M. de F. M., Rossetti, M. L., Ximenes, R., Braga, M. C., Lucena-Silva, N. (2014). Evaluation of four molecular methods for the diagnosis of tuberculosis in pulmonary and blood samples from immunocompromised patients. *Memórias Do Instituto Oswaldo Cruz*, 109(6), 805–813.
- Dezemon, Z., Muvunyi, C. M., & Jacob, O. (2014). Staining techniques for detection of acid fast bacilli: what hope does fluorescein-diacetate (FDA) vitality staining technique represent for the monitoring of tuberculosis treatment in resource limited settings. *Trends in Bacteriology*, 1(1), 1.
- Dutse, S. W., Yusof, N. A., Ahmad, H., Hussein, M. Z., & Zainal, Z. (2012). An electrochemical DNA biosensor for Ganoderma boninense pathogen of oil palm. *International Journal of Electrochemical Science*, *7*, 8105–8115.
- Dutse, S. W., Yusof, N. A., Ahmad, H., Hussein, M. Z., Zainal, Z., & Hushiarian, R. (2011). DNA-based Biosensor for Detection of Ganoderma boninense, an Oil Palm Pathogen Utilizing Newly Synthesized Ruthenium Complex [Ru(phen)2(qtpy)]²⁺ Based on a PEDOT-PSS/Ag Nanoparticles Modified Electrode. Sensors, 8, 5754–5768.
- Dylan, T. (2014). Extrapulmonary tuberculosis. *MSD Manual*, 316–353. Retrieved from https://www.msdmanuals.com/professional/infectiousdiseases/mycobacteria/extrapulmonary-tuberculosis

- Erdem, A., Kerman, K., Meriç, B., Özkan, D., Kara, P., & Özsöz, M. (2002). DNA biosensor for Microcystis spp. sequence detection by using methylene blue and ruthenium complex as electrochemical hybridization labels. *Turkish Journal of Chemistry*, 26(6), 851–862.
- Ershova, J. V, Volchenkov, G. V, Kaminski, D. A., Somova, T. R., Kuznetsova, T. A., Kaunetis, N. V., Cegielski, J. P., Kurbatova, E. V. (2015). Epidemiology of Primary Multidrug-Resistant Tuberculosis, Vladimir Region, Russia, *Emerging Infectious Diseases*. 21(11), 2048–2051.
- Ertekin, Ö., Öztürk, S., & Öztürk, Z. Z. (2016). Label Free QCM Immunobiosensor for AFB1, *Sensors*. 1–12.
- Esmail, H., Barry, C. E., Young, D. B., & Wilkinson, R. J. (2014). The ongoing challenge of latent tuberculosis. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 369(1645), 20130437.
- Evtugyn, G., & Hianik, T. (2016). Electrochemical DNA sensors and aptasensors based on electropolymerized materials and polyelectrolyte complexes. *TrAC Trends in Analytical Chemistry*, 79, 168–178.
- Fandinho, F. C., Orsi-Souza, A. T., & Salem, J. I. (1990). A comparison of the Ziehl
 Neelsen and Kinyoun methods in staining smears from leprosy patients. *International Journal of Leprosy and Other Mycobacterial Disease*, 58(2), 389–391.
- Fei, J., Dou, W., & Zhao, G. (2016). A sandwich electrochemical immunosensor for Salmonella pullorum and Salmonella gallinarum based on a screen-printed carbon electrode modified with an ionic liquid and electrodeposited gold nanoparticles, Microchim Acta. 182:2267-2275
- Gao, H., Sun, M., Lin, C & Wang, W. (2012). Electrochemical DNA Biosensor Based on Graphene and TiO 2 Nanorods Composite Film for the Detection of Transgenic Soybean Gene Sequence of MON89788, *Electroanalysis*. 24(12), 2283–2290.
- Gholoobi, A., Masoudi-Kazemabad, A., Meshkat, M., & Meshkat, Z. (2014). Comparison of culture and PCR methods for diagnosis of Mycobacterium tuberculosis in different clinical specimens. *Jundishapur Journal of Microbiology*, 7(2), e8939.
- Giamblanco, N., Petralia, S., Conoci, S., Messineo, C., & Marletta, G. (2017). Colloids and Surfaces B: Biointerfaces Ionic strength-controlled hybridization and stability of hybrids of KRAS DNA single-nucleotides: A surface plasmon resonance study. *Colloids and Surfaces B: Biointerfaces*, 158(2017), 41–46.

- Saldaña, N. G., Parra, M. M., Porras, M. H., Castrellón, P. G., Toscano, V. G., & Olguin, H. J. (2014). Pulmonary tuberculous: Symptoms, diagnosis and treatment. 19-year experience in a third level pediatric hospital. BMC Infectious Diseases, 14 (401), 1471-2334
- Hajian, R., Tayebi, Z., Shams, N. (2017). Fabrication of an electrochemical sensor for determination of doxorubicin in human plasma and its interaction with DNA, *Journal of Pharmaceutical Analysis*. 7(2017), 27–33.
- Halid, N. I. A., Hasbullah, S. A., Ahmad, H., Heng, L. Y., Karim, N. H. A., & Harun, S. N. (2014). Electrochemical DNA biosensor for detection of porcine oligonucleotides using ruthenium(II) complex as intercalator label redox, *American Institute of Physics Conferences Proceeding*. 313(2014), 313–316.
- Hammond, J. L., Formisano, N., Estrela, P., Carrara, S., & Tkac, J. (2016). Electrochemical biosensors and nanobiosensors, *Essays in Biochemistry*. 60, 69–80.
- Hayat, A., & Marty, J. L. (2014). Disposable screen printed electrochemical sensors: Tools for environmental monitoring. *Sensors (Switzerland)*, 14(6), 10432–10453.
- Hilmi, Y. (2015). 17,000 TB cases reported among foreign workers. *Bernama*, pp. 1–11.
- Hong, N., Cheng, L., Wei, B., Chen, C., He, L. L., kong, D., Ceng, J., Cui, H., & Fan, H. (2017). An electrochemical DNA sensor without electrode premodification, *Biosensors and Bioelectronics*. 91(2017), 110–114.
- Hosseini, F., Seyedsadjadi, M., & Farhadyar, N. (2014). Fe₃O₄ Nanoparticles Modified with APTES as the Carrier for (+)-(S)-2-(6-methoxynaphthalen-2yl) Propanoic Acid (Naproxen) and (RS) 2-(3-benzoylphenyl)-propionic Acid (Ketoprofen) Drug. *Oriental Journal of Chemistry*, 30(4), 1609–1618.
- Hoyos-arbeláez, J., Vázquez, M., & Contreras-calderón, J. (2017). Electrochemical methods as a tool for determining the antioxidant capacity of food and beverages : A review, *Food chemistry*. 221(2017), 1371–1381.
- Hushiarian, R., Yusof, N. A., Abdullah, A. H., Ahmad, S. A. A., & Dutse, S. W. (2014). A novel DNA nanosensor based on cdse/zns quantum dots and synthesized fe3o4 magnetic nanoparticles. *Molecules*, 19(4), 4355–4368.
- Issa, R., Hamdan, N. A., & Noh, M. F. (2010). Differential Pulse Voltammetric Determination of DNA Hybridization using Methylene Blue on screen Orinted Carbon Electrode for the Detection of Mycrobacterium tuberculosis. *Biotechnology*. ISSN 1682-296X

- Jagadeesha, K., & Vidya, P. (2015). A Comparative Study of Fluorescent Staining and Ziehl Neelsen's Staining for Detection of Acid Fast Bacilli in Sputum in a Tertiary Care Hospital in Mangalore, *International journal o. Microbiology* & *Parasitology*. 1(1), 3–5.
- Jara-palacios, M. J., Escudero-gilete, M. L., Hernández-hierro, J. M. Heredia, J. H., & Hernanz, D. (2017). Cyclic voltammetry to evaluate the antioxidant potential in winemaking by- products. *Talanta*, 165, 211–215.
- Johanna. N. (2015). A retrospective evaluation study of diagnostic accuracy of Xpert ® MTB/RIF assay used for detection of *Mycobacterium tuberculosis* in Greece. Orebro University
- Justino, C. I. L., Duarte, A. C., & Rocha-santos, T. A. P. (2016). Critical overview on the application of sensors and biosensors for clinical analysis. *Trends in Analytical Chemistry*, 85, 36–60.
- Kaushik, M., Kaushik, S., Roy, K., Singh, A., Mahendru, S., Kumar, M., Chaudhary, S., Ahmed, S., & Kukreti, S. (2016). A bouquet of DNA structures: Emerging diversity. *Biochemistry and Biophysics Reports*, 5, 388–395.
- Kavita. (2017). Biomedical Science OMICS International DNA Biosensors-A Review. Journal of Bioengineering & Biomedical Science, 7(2). (222-226)
- Kerman, K., Morita, Y., Takamura, Y., Ozsoz, M., & Tamiya, E. (2004). Modification of Escherichia coli single-stranded DNA binding protein with gold nanoparticles for electrochemical detection of DNA hybridization. *Analytical Chimica Acta*, 510, 169–174.
- Khalil, H. P. S. A., Davoudpour, Y., Aprilia, N. A. S., Mustapha, A., Hossain, M. S., Islam, M. N., & Dungani, R. (2015). Nanocellulose-Based Polymer Nanocomposite: Isolation, Characterization and Applications. *Nanocellulose Polymer Nanocomposites*, 273–310.
- Khalil, I., Julkapli, N. M., Yehye, W. A., Basirun, W. J., & Bhargava, S. K. (2016).
 Graphene–Gold Nanoparticles Hybrid—Synthesis, Functionalization, and
 Application in a Electrochemical and Surface-Enhanced Raman Scattering
 Biosensor. *Materials*, 9,406. http://doi.org/10.3390/ma9060406
- Kivihya-Ndugga, L., van Cleeff, M., Juma, E., Kimwomi, J., Githui, W., Oskam, L., Schuitema, A., van Soolingen, D., Nganga, L., Kibuga, D., Odhiambo, J., & Klatser, P. (2004). Comparison of PCR with the Routine Procedure for Diagnosis of Tuberculosis in a Population with High Prevalences of Tuberculosis and Human Immunodeficiency Virus *Journal of Clinical Microbiology*, 42(3), 1012-1015.
- Kox, L. F. F., Rhienthong, D., Miranda, A. M., Udomsantisuk, N., Ellis, K., van Leeuwen, J., van Heusden, S., Kuijper, S., & Kolk, A. H. J. (1994). A more reliable PCR for detection of Mycobacterium tuberculosis in clinical samples. *Journal of Clinical Microbiologyl*, 32(3), 672–678.

- Kritski, A. L., Huf, G., Oliveira, M. M., Squire, S. B., & Ruffino-netto, A. (2015). Assessing new strategies for TB diagnosis in low- and middle-income countries, *BMC Proceedings*. 8(Suppl 4): O12, http://doi.org/10.1186/1753-6561-8-S4-O12
- Kumar, A., Negi, Y. S., Choudhary, V., & Bhardwaj, N. K. (2014). Characterization of Cellulose Nanocrystals Produced by Acid-Hydrolysis from Sugarcane Bagasse as Agro-Waste, *Journal of Materials Physics and Chemistry*, 2(1), 1–8.
- Kurbanoglu, S., Ozkan, S. A., & Merkoçi, A. (2017). Nanomaterials-based enzyme electrochemical biosensors operating through inhibition for biosensing applications. *Biosensors and Bioelectronic*, 89(2017), 886–898.
- Kwon, H. J., & Akyiano, E. (2011). Simulation of cyclic voltammetry of ferrocyanide / ferricyanide redox reaction in the EQCM Sensor, *Proceedings* of the 2011 COMSOL Conference in Boston, 2–6.
- Laborda, E., González, J., & Molina, Á. (2014). Recent advances on the theory of pulse techniques: A mini review. *Electrochemistry Communications*, 43(2014), 25–30.
- Lee, J. Y. (2015). Diagnosis and treatment of extrapulmonary tuberculosis. *Tuberculosis and Respiratory Diseases*, 78(2), 47–55.
- Lee, P. Y., Costumbrado, J., Hsu, C., & Kim, Y. H. (2012). Agarose Gel Electrophoresis for the Separation of DNA Fragments, *Journal of Visualized Experiments*, (62), e3923. http://doi.org/10.3791/3923
- Lee, Y. T., & Woo, K. (2006). Preparation of water-dispersible and biocompatible iron oxide nanoparticles for MRI agent. 2006 IEEE Nanotechnology Materials and Devices Conference, NMDC, 1(2), 454–455.
- Lee, S., Fang, H., Chen, W., Lin, H & Chang, C. A. (2005). Electrochemical study on screen-printed carbon electrodes with modification by iron nanoparticles in $Fe(CN)_6^{4-/3-}$ redox system, *Analytical Bioanalyst Chemistry*. 532–538.
- Lemos, A. C. M., & Matos, E. D. (2013). Multidrug-resistant tuberculosis. Brazilian Journal of Infectious Diseases, 17(2), 239–246.
- Li, S., Qiu, W., Zhang, X., Ni, J., & Gao, F. (2016). Sensors and Actuators B: Chemical A high-performance DNA biosensor based on the assembly of gold nanoparticles on the terminal of hairpin-structured probe DNA. *Sensors & Actuators: B. Chemical*, 223, 861–867.
- Liu, J., Li, J., & Dong, S. (1996). Interaction of Brilliant Cresyl Blue and Methylene Green with DNA Studied by Spectrophotometric and Volt ammetric Methods. *Electroanalysis*. 8, 8-9

- Liu, C., Jang, D., Xiang, G., Liu, F., Liu, L., & Pu, X. (2014). An electrochemical DNA biosensor for the detection of Mycobacterium tuberculosis, based on signal amplification of graphene and a gold nanoparticles-polyaniline nanocomposite. *The Analyst*, 139, 5460–5465.
- Loaiza, A., Campuzano, S., & Pingarr, M. (2007). DNA sensor based on an Escherichia coli lac Z gene probe immobilization at self-assembled monolayers-modified gold electrodes, *Talanta*, *73*, 838–844.
- Lucarelli, F., Tombelli, S., Minunni, M., Marrazza, G., & Mascini, M. (2008). Electrochemical and piezoelectric DNA biosensors for hybridisation detection. *Analytica Chimica Acta*, 609(2), 139–159.
- Mariño, M., Da Silva, L. L., Durán, N., & Tasic, L. (2015). Enhanced materials from nature: Nanocellulose from citrus waste. *Molecules*, 20(4), 5908–5923.
- Martín-Yerga, D., Rama, E. C., & Costa-García, A. (2016). Electrochemical Study and Determination of Electroactive Species with Screen-Printed Electrodes, *Journal of Chemical Education*, http://doi.org/10.1021/acs.jchemed.5b00755
- Martinez-Romero, A., Ortega-Sanchez, J. L., Urtiz-Estrada, N., Calleros-Rincon, E. Y., Perez-Morales, R., & Alba-Romero, J. de J. (2015). Immunity towards tuberculosis infection: a review. *African Journal of Microbiology Research*, 9(37), 2013–2022.
- Mat Zaid, M. H., Abdullah, J., Yusof, N. A., Sulaiman, Y., Wasoh, H., Md Noh, M. F., & Issa, R. (2016). PNA biosensor based on reduced graphene oxide/water soluble quantum dots for the detection of Mycobacterium tuberculosis. *Sensors and Actuators, B: Chemical*. http://doi.org/10.1016/j.snb.2016.10.045
- Mehrotra, P. (2016). ScienceDirect Biosensors and their applications A review. Journal of Oral Biology and Craniofacial Research, 6(2), 153–159.
- Moghaddam, H. T., Moghadam, Z. E., & Khademi, G. (2016). Tuberculosis : Past, Present and Future, *International Journal of pediatrics*, 4(25), 1243–1254.
- Mohamud, R., Azlan, M., Yero, D., Alvarez, N., Sarmiento, M. E., Acosta, A., & Norazmi, M.-N. (2013). Immunogenicity of recombinant Mycobacterium bovis bacille Calmette-Guèrin clones expressing T and B cell epitopes of Mycobacterium tuberculosis antigens. *BMC Immunology*, 14 Suppl 1(Suppl 1), S5. http://doi.org/10.1186/1471-2172-14-S1-S5
- Moncada, V. (2014). Development of a Modified Carbon Electrode With Ionic Liquid and its Application for Electrocatalytic Oxidation and Voltammetric Determination of Levodopa, *Journal of the Chilean Chemical Society*, 2, 2516–2519.

- Morillo, D., Uheida, A., Pérez, G., Muhammed, M., & Valiente, M. (2015). Arsenate removal with 3-mercaptopropanoic acid-coated superparamagnetic iron oxide nanoparticles. *Journal of Colloid and Interface Science*, 438(2015), 227–234.
- Munshi, S. K., Rahman, F., Mostofa Kamal, S. M., & Noor, R. (2012). Comparisons among the diagnostic methods used for the detection of extra-pulmonary tuberculosis in Bangladesh. *International Journal of Mycobacteriology*, 1(4), 190–195.
- Musa, A., Ahmad, M. B., Hussein, M. Z., Izham, S. M., Shameli, K., & Sani, H. A. (2016). Synthesis of Nanocrystalline Cellulose Stabilized Copper Nanoparticles, Journal of Nanomaterials, http://dx.doi.org/10.1155/2016/2490906
- Ng, H. Z., Ng, M., Eng, C. M., & Gao, Z. (2016). Deoxyribonucleic acid glycosylase assays: Progress and prospects. *TrAC Trends in Analytical Chemistry*, 83, 102–115.
- Ngoensawat, U., Rijiravanich, P., Somasundrum, M., & Surareungchai, W. (2014). Highly sensitive electrochemical detection of DNA hybridisation by coupling the chemical reduction of a redox label to the electrode reaction of a solution phase mediator. *The Analyst*, *139*, 5740–5746.
- Nguyen, D. P., Tran, Q. T., Trinh, X. S., Hoang, T. C., Nguyen, H. N., & Nguyen, H. H. (2012). Crystallization and magnetic properties of amorphous iron-chromium oxide nanoparticles synthesized by sonochemistry. Advances in Natural Sciences: Nanoscience and Nanotechnology, 3(1), 15017.
- Nordin, N., Azah, N., Abdullah, J., & Radu, S. (2016). Sensitive detection of multiple pathogens using a single DNA probe. *Biosensors and Bioelectronic*, 86, 398–405.
- Nurmalasari, R., Gaffar, S., & Hartati, Y. W. (2015). Label-Free Electrochemical DNA Biosensor for the Detection of Mycobacterium tuberculosis using Gold Electrode Modified by Self-Assembled Monolayer of Thiol. *Procedia Chemistry*, 17, 111–117.
- Nyaruhirira, A. U., Toussaint, M., Nemser, B., Vandebriel, G., Gasana, M., & Amor,
 Y. Ben. (2015). Performance of LED fluorescence microscopy for the detection of tuberculosis in rwanda using Zeiss Primo Star. *Pan African Medical Journal*, 21, 1–8.
- Pai, M., & Menzies, D. (2014). Diagnosis of latent tuberculosis infection (tuberculosis screening) in HIV-negative adults., 1–3. Retrieved from www.uptodate.com
- Pang, L., Li, J., Jiang, J., Le, Y., Shen, G. L., & Yu, R. (2007). A novel detection method for DNA point mutation using QCM based on Fe₃O₄/Au core/shell nanoparticle and DNA ligase reaction, *127*, 311–316.

- Panjkovich, A., & Melo, F. (2005). Comparison of different melting temperature calculation methods for short DNA sequences, *Structural bioinformatics*, 21(6), 711–722.
- Parkash, O., Yean, C. Y., & Shueb, R. H. (2014). Screen Printed Carbon Electrode Based Electrochemical Immunosensor for the Detection of Dengue NS1 Antigen, *Diagnostics*, (4), 165–180.
- Periasamy, V. S., Athinarayanan, J., Alhazmi, M., Alatiah, K. A., & Alshatwi, A. A. (2014). Fe₃O₄ Nanoparticle Redox System Modulation via Cell-Cycle Progression and Gene Expression in Human Mesenchymal Stem Cells, *Environmental Toxicology*, 901–912.
- Prabhakar, N., Arora, K., Arya, S. K., Solanki, P. R., Iwamoto, M., Singh, H & Malhotra, B. D. (2008). Nucleic acid sensor for *M. tuberculosis* detection based on surface plasmon resonance. *The Analyst*, 133(11), 1587–1592.
- Prabhakar, N., Solanki, P. R., Kaushik, A., Pandey, M. K., & Malhotra, B. D. (2010). Peptide Nucleic Acid Immobilized Biocompatible Silane Nanocomposite Platform for *Mycobacterium tuberculosis* Detection. *Electroanalysis*, 22(22), 2672–2682.
- Pullan, S. T., Allnutt, J. C., Devine, R., Hatch, K. A., Jeeves, R. E., Hendon-dunn, C. L., Marsh, P. D., & Bacon, J. (2016). The effect of growth rate on pyrazinamide activity in *Mycobacterium tuberculosis*-insights for early bactericidal activity? *BMC Infectious Diseases*, 16:205. http://doi.org/10.1186/s12879-016-1533-z
- Putzbach, W., & Ronkainen, N. J. (2013). Immobilization Techniques in the Fabrication of Nanomaterial-Based Electrochemical Biosensors: A Review, *Sensors, 13,* 4811–4840.
- Rabel, A. M., Jayanthi, V., Raj, N. N., Ramachandran, D., & Brijitta, J. (2013). Synthesis and Characterization of Chitosan-Coated Iron Oxide Nanoparticles, Proceedings of the "International Conference on Advanced Nanomaterials & Emerging Engineering Technologies", 569–571.
- Radhi, M. M., Al-mulla, E. A. J., & Tan, W. T. (2012). Electrochemical characterization of the redox couple of Fe(III)/Fe(II) mediated by grafted polymer electrode, *Research on Chemical Intermediates*. http://doi.org/10.1007/s11164-012-0954-6
- Raghav, R., & Srivastava, S. (2016). Immobilization strategy for enhancing sensitivity of immunosensors: L-Asparagine – AuNPs as a promising alternative of EDC– NHS activated citrate–AuNPs for antibody immobilization. *Biosensors and Bioelectronic*, 78, 396–403.

- Rahman, G., Ahmad, S., Anwar, M., & Shah, A. (2016). Electrocatalytic behavior of glassy carbon electrode modified with ruthenium nanoparticles and ruthenium film. *Journal of Applied Electrochemistry*, 46(4), 459–468. http://doi.org/10.1007/s10800-016-0937-1
- Rahman, M. M., Li, XB., Lopa, N. S., Ahn, S. J., & Lee, JJ. (2015). Electrochemical DNA Hybridization Sensors Based on Conducting Polymers. *Sensors*, 15(2), 3801–3829.
- Torati, S. R., Reddy, V., Yoon, S. S., & Kim, C. (2016). Electrochemical biosensor for Mycobacterium tuberculosis DNA detection based on gold nanotubes array electrode platform. *Biosensors and Bioelectronic*, 78(2016), 483–488.
- Raoof, J. B., Ojani, R., Ebrahimi, M., & Hamidi-Asl, E. (2013). Application of ethyl green as electroactive indicator in genosensors and investigation of its interaction with DNA. *Analytical & Bioanalytical Electrochemistry*, 5(4), 395-415
- Ribovski, L., Zucolotto, V., & Janegitz, B. C. (2017). A label-free electrochemical DNA sensor to identify breast cancer susceptibility, *Microchemical Journal*, *133*(2017), 37–42.
- Ryu, Y. J. (2015). Diagnosis of pulmonary tuberculosis: Recent advances and diagnostic algorithms. *Tuberculosis and Respiratory Diseases*, 78(2), 64–71.
- Salajkova, M., Berglund, L. a., Zhou, Q., Salajková, M., Berglund, L. a., & Zhou, Q. (2012). Hydrophobic cellulose nanocrystals modified with quaternary ammonium salts. *Journal of Materials Chemistry*, 22(37), 19798.
- Sathish, S., & Balakumar, S. (2016). Influence of physicochemical interactions of capping agent on magnetic properties of magnetite nanoparticles. *Materials Chemistry and Physics*, 173, 364–371.
- Sawadogo, T. L., Savadogo, L. G. B., Diande, S., Ouedraogo, F., Mourfou, A.,
 Gueye, A., Sawadogo, I., Nebie, B., Sangare, L., & Ouattara, A. S. (2012).
 Comparison of Kinyoun, auramine O, and Ziehl-Neelsen staining for
 diagnosing tuberculosis at the National Tuberculosis Center in Burkina Faso.
 Health and Tropiacal Medicine, 22, 302–306.
- Saxena, P., Ashish, K. A., & Molly, M. (2014). Comparison of microscopy and PCR in detection of *Mycobacterium tuberculosis*. *Journal of Microbiology and Infectious Diseases*, 4(4), 141–144.
- Schumm, B. (2014). Carbon-Zinc Batteries. Encyclopedia of Applied Electrochemistry, http://doi.org/10.1007/978-1-4419-6996-5
- Shehzad, A., Rehman, G., Ul-Islam, M., Khattak, W. A., & Lee, Y. S. (2015). Challenges in the development of drugs for the treatment of tuberculosis. *The Brazilian Journal of Infectious Diseases : An Official Publication of the Brazilian Society of Infectious Diseases*, 17(1), 74–81.

- Shrivastava, S., Jadon, N., & Jain, R. (2016). Next-generation polymer nanocomposite-based electrochemical sensors and biosensors: A review. *TrAC-Trends in Analytical Chemistry*, 82, 55–67.
- Singh, R., Verma, R., Kaushik, A., Sumana, G., Sood, S., Gupta, R. K., & Malhotra, B. D. (2011). Chitosan-iron oxide nano-composite platform for mismatchdiscriminating DNA hybridization for Neisseria gonorrhoeae detection causing sexually transmitted disease. *Biosensors and Bioelectronics*, 26(6), 2967–2974.
- Singhal, R., & Myneedu, V. P. (2015). Microscopy as a diagnostic tool in pulmonary tuberculosis. *International Journal of Mycobacteriology*, 4(1), 1–6.
- Tankeshwar, A. (2013). Ziehl-Neelsen technique (AFB Staining): Principle, Procedure and Reporting. *Microbe Online*, *3*, 1–6.
- Tankeshwar, A. (2015). Auramine Rhodamine Fluorochrome Staining : Principle, Procedure, Results and Limitations. *Microbe Online*, (January), 1–5. Retrieved from http://microbeonline.com/auramine-rhodaminefluorochrome-staining-principle-procedure-results-limitations/
- Tariq, M. (2016). 40 % of India's population play host to the TB bacillus as a latent TB. One India, 1–5. Retrieved from http://www.oneindia.com/feature/40percent-of-india-s-population-play-host-the-tb-bacillus-as-latenttuberculosis-2049544.html
- Tehrani, F., & Bavarian, B. (2016). Facile and scalable disposable sensor based on laser engraved graphene for electrochemical detection of glucose. *Scientific Reports*, 6(January), 27975. http://doi.org/10.1038/srep27975
- Thiruppathiraja, C., Kamatchiammal, S., & Adaikkappan, P. (2011). Specific detection of Mycobacterium sp. genomic DNA using dual labeled gold nanoparticle based electrochemical biosensor, *Analytical Biochemistry*, 417, 73–79.
- Tichoniuk, M., Gwiazdowska, D., Ligaj, M., & Filipiak, M. (2010). Electrochemical detection of foodborne pathogen Aeromonas hydrophila by DNA hybridization biosensor. *Biosensors and Bioelectronics*, 26(4), 1618–1623.
- Tiffany Abitbol, Heera Marway and, E. D. C. (2014). Surface modification of cellulose nanocrystals. *Nanoscale*, *6*(14), 7764–79.
- Tiwari, D., Haque, S., Tiwari, R. P., Jawed, A., Govender, T., & Kruger, H. G. (2017). Fast and efficient detection of tuberculosis antigens using liposome encapsulated secretory proteins of Mycobacterium tuberculosis. *Journal of Microbiology, Immunology and Infection*, 50(2), 189–198.
- Todar, K. (2012). Mycobacterium tuberculosis and Tuberculosis. In *Todar's online textbook of bacteriology* (pp. 1–4). Wisconsin. Retrieved from http://textbookofbacteriology.net/tuberculosis.html

- Tran, L. D., Nguyen, B. H., Van Hieu, N., Tran, H. V., Nguyen, H. Le, & Nguyen, P. X. (2011). Electrochemical detection of short HIV sequences on chitosan/Fe 3O4 nanoparticle based screen printed electrodes. *Materials Science and Engineering C*, 31(2), 477–485.
- Uchida, Y., Kätelhön, E., & Compton, R. G. (2017). Cyclic voltammetry with nontriangular waveforms: Electrochemically reversible systems. *Journal of Electroanalytical Chemistry*, 801(August), 381–387.
- Uddin, M. K. M., Chowdhury, M. R., Ahmed, S., Rahman, M. T., Khatun, R., van Leth, F., & Banu, S. (2013). Comparison of direct versus concentrated smear microscopy in detection of pulmonary tuberculosis. *BMC Research Notes*, 6, 291. http://doi.org/10.1186/1756-0500-6-291
- Ulianas, A., Heng, L. Y., Hanifah, S. A., & Ling, T. L. (2012). An electrochemical DNA microbiosensor based on succinimide-modified acrylic microspheres. *Sensors (Switzerland)*, 12(5), 5445–5460.
- Van Deun, A., Chuquiyauri, R., Torrea, G., Agapito, J., Verdonck, K., & Gotuzzo, E. (2008). Yield of fluorescence microscopy versus culture for tuberculosis at a middle-income country referral hospital. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 102(6), 564–569.
- Wang, J. (2005). Electrochemical Nucleic Acid Biosensors, Analytica Chimica Acta. 469(1), 63-71.
- WHO. (2014). Tuberculosis (TB) Latent Tuberculosis Infection (LTBI), 1–3. Retrieved from http://www.who.int/tb/challenges/ltbi/en/
- WHO. (2015). Global tuberculosis report 2015. Retrieved from https://reliefweb.int/report/world/global-tuberculosis-report-2015
- WHO. (2016). Drug-Resistant Tb : Retrieved from http://www.cdc.gov/tb/topic/drtb/
- Woolston, C. (2016). Tuberculosis (TB), 3(May 2014), 1–13. Retrieved from https://consumer.healthday.com/encyclopedia/infectious-diseases-26/misc-infections-news-411/tuberculosis-tb-648356.html
- Xu, S., Zhang, Y., Dong, K., Wen, J., Zheng, C., & Zhao, S. (2017). Electrochemical DNA Biosensor Based on Graphene Oxide-Chitosan Hybrid Nanocomposites for Detection of *Escherichia Coli* 0157: H7, *International Journal of Electrochemical Science*, 12, 3443–3458.
- Yang, X., Wang, L. N., Zhou, G. Z., Sui, N., Gu, Y. X., & Wan, J. (2015). Electrochemical detection of H₂O₂ based on Fe₃O₄ nanoparticles with graphene oxide and polyamidoamine dendrimer. *Journal of Cluster Science*, 26(3), 789– 798.

- Yin, H., Yang, Z., Wang, H., Zhou, Y., & Ai, S. (2017). Electrochemical biosensor for hydroxymethylated DNA detection and β-glucosyltransferase activity assay based on enzymatic catalysis triggering signal amplification. *Sensors & Actuators B: Chemical*, 243, 602–608.
- Zaffino, R. L., Galan, T., Pardo, W. A., Mir, M., & Samitier, J. (2015). Nanoprobes for enhanced electrochemical DNA sensors. *Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology*, 7(6), 817–827.
- Zhang, S., Ding, Y., & Wei, H. (2014). Ruthenium polypyridine complexes combined with oligonucleotides for bioanalysis: A review. *Molecules*, 19(8), 11933–11987.
- Zulkifli, H., Salam, F., Saad, S. M., Rahman, R. A., Rani, R. M., Karim, M. S. A., & Ishak, Z. (2016). Preliminary Study of Electrochemical DNA Sensor for Cucumber Mosaic Virus. *Procedia Chemistry*, 20, 98–101.
- Zumla, A., Raviglione, M., Hafner, R., & Fordham von Reyn, C. (2013). Tuberculosis. *New England Journal of Medicine*, 368(8), 745–755.