



**UNIVERSITI PUTRA MALAYSIA**

***USEFULNESS OF CENTOR SCORE FOR DIAGNOSIS OF GROUP A  
STREPTOCOCCAL PHARYNGO-TONSILLITIS AND PREVALENCE OF  
THE DISEASE IN MALAYSIA FROM 2016 TO 2017***

**ABDULRAHMAN MANSOOR MOHAMMED MUTHANNA**

**FPSK(M) 2018 1**



**USEFULNESS OF CENTOR SCORE FOR DIAGNOSIS OF GROUP A  
STREPTOCOCCAL PHARYNGO-TONSILLITIS AND PREVALENCE OF  
THE DISEASE IN MALAYSIA FROM 2016 TO 2017**

**By**

**ABDULRAHMAN MANSOOR MOHAMMED MUTHANNA**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia  
in Fulfilment of the Requirements for the Degree of Master of Science**

**October 2017**

## **COPYRIGHT**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**USEFULNESS OF CENTOR SCORE FOR DIAGNOSIS OF GROUP A STREPTOCOCCAL PHARYNGO-TONSILLITIS AND PREVALENCE OF THE DISEASE IN MALAYSIA FROM 2016 TO 2017**

By

**ABDULRAHMAN MANSOOR MOHAMMED MUTHANNA**

**October 2017**

**Chairman : Siti Zulaikha Zakariah, MB BCh BAO, DrPath**  
**Faculty : Medicine and Health Sciences**

Pharyngo-tonsillitis is very common in general practice and most of the cases are caused by viruses. One of the causes of pharyngo-tonsillitis is group A streptococcus, which has a strong indication for antibiotic treatment. It is difficult to distinguish between streptococcal pharyngo-tonsillitis from non-streptococcal according to the clinical findings. Nevertheless, up to 90% of the patients with sore throat might be treated with antibiotic. Over prescribing of antibiotics has very serious health effects with severe reactions and may promote antibiotic resistance and add significantly to the cost of health care. In Malaysia, Upper Respiratory Tract Infection (URTI) makes up nearly 30% of cases in primary care. Studies have shown trends of inappropriate prescribing of antibiotics for URTI in Malaysian primary care. Centor scoring is a guideline based on a set of criteria that help to identify the likelihood of streptococcal infections in patients with a sore throat. It was conducted to guide physicians to appropriately prescribe antibiotics for adults with pharyngo-tonsillitis. This study aims to describe the epidemiological pattern, etiology and its antibiotic susceptibility, clinical manifestation, antibiotic prescription among adults with sore throat and validity of Centor score in diagnosis group A streptococcal pharyngo-tonsillitis at three primary care clinics in Sepang, Selangor, Malaysia during 2016 to 2017. This cross-sectional study was conducted on 215 patients aged 18 and above with sore throat as one of the complaints at the three primary care clinics during December 2016 until April 2017. Throat swabs were collected from the patients for culture and analysis. Data on clinical manifestations, demographic characteristics, clinical information and throat sample results were analyzed using Chi-square test, and descriptive statistics. From all the participants (42.3% male, 57.7% female), (18.6% smokers, 81.4% non-smokers), (62.8% Malay, 30.2% Indian, 5.1% Chinese, 1.9 % others), 6 isolates (2.4%) were identified as GAS including 50% of those were associated with Centor score of 3 ( $p < 0.001$ ), 50% were associated with Centor score

of 4 ( $p < 0.001$ ), and 0% with Centor scores of 0,1 and 2. Centor criteria were clinical predictors that associated with group A streptococcal pharyngo-tonsillitis ( $p < 0.001$ ). Pharyngo-tonsillitis was diagnosed in 130 (60.5%) of the total adult participants who complained with sore throat. Beta hemolytic streptococci and influenza A and B viruses were isolated from 37.1% and 3.8 % of total participants, respectively. Both Centor scores 3 and 4 had sensitivity of 50%, and specificity of 97.6% and 100%; respectively, positive predictive value 37.5% and 100%; respectively, negative predictive value 98.6% and 98.6%; respectively, positive likelihood ratio 20.5 and 50; respectively, negative likelihood ratio 0.5 and 0.5; respectively. The accuracy of Centor score 3 was 96.3% while the accuracy of Centor score 4 was 98.6%. Antibiotics were prescribed to 48 (22.3%) including 8.3% with group A streptococcus and 91.6% with non-group A streptococcus. A majority of prescribed antibiotics was associated with Centor scores of zero to one (75.0%). Antibiotic susceptibility testing revealed that all beta hemolytic streptococci isolates were susceptible to penicillin G, ampicillin, ofloxacin, cefepime, cefotaxime, ceftriaxone, vancomycin, and linezolid, but 40%, 2.4%, and 9.6% were resistant to tetracycline, clindamycin, and erythromycin, respectively. The current study has suggested that the Centor score is useful for diagnosis and decision making for antibiotic therapy of GAS pharyngo-tonsillitis leading to decrease the unnecessary antibiotic prescription, while achieving better levels of treatment.

Key words: Pharyngo-tonsillitis; Clinical score; Centor score; *Streptococcus pyogenes*; Group A beta-hemolytic streptococcus; antibiotic prescription

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**FAEDAH SKOR CENTOR UNTUK MENDIAGNOSIS FARINGO-TONSILITIS STREPTOKOKUS KUMPULAN A DAN PREVALENS PENYAKIT DI MALAYSIA DARI 2016 HINGGA 2017**

Oleh

**ABDULRAHMAN MANSOOR MOHAMMED MUTHANNA**

**Oktober 2017**

**Pengerusi : Siti Zulaikha Zakariah, MB BCh BAO, DrPath**  
**Fakulti : Perubatan dan Sains Kesihatan**

Faringo-tonsilitis biasa dijumpai di klinik primer dan kebanyakan kes disebabkan oleh virus. Salah satu punca jangkitan faringo-tonsilitis ialah streptokokus kumpulan A. Jangkitan ini adalah satu indikasi yang kukuh untuk rawatan menggunakan antibiotic. Tetapi, jangkitan ini sukar dibezakan antara faringo-tonsilitis streptokokus dan bukan-streptokokus menggunakan tanda-tanda klinikal sahaja. Walau bagaimanapun, sehingga 90% pesakit dengan sakit tekak mungkin dirawat dengan antibiotik. Dengan menetapkan antibiotik yang lebih ianya akan memberi kesan kesihatan yang sangat serius dengan tindak balas yang teruk dan boleh menggalakkan kerintangan antibiotik sekaligus menambah kos kepada penjagaan kesihatan. Di Malaysia, jangkitan saluran pernafasan atas (URTI) membentuk hampir 30% daripada kes di peringkat penjagaan primer. Kajian telah menunjukkan bahawa terdapat trend preskripsi antibiotik yang tidak sesuai untuk URTI di peringkat penjagaan primer Malaysia. Skor Centor adalah garis panduan berdasarkan satu set kriteria yang membantu untuk mengenal pasti kemungkinan jangkitan streptokokus pada pesakit yang mempunyai sakit tekak. Ia digunakan untuk membimbing para doktor untuk menetapkan antibiotik yang sesuai kepada orang dewasa dengan faringo-tonsilitis. Kajian ini bertujuan untuk menggambarkan corak epidemiologi, etiologi dan kerentanan antibiotik, manifestasi klinikal, preskripsi antibiotik di kalangan orang dewasa dengan sakit tekak dan kesahihan skor Centor dalam mendiagnosis faringo-tonsillitis streptokokus kumpulan A di tiga klinik klinik penjagaan primer di Sepang, Selangor, Malaysia pada 2016 hingga 2017. Satu kajian keratan rentas telah dijalankan di kalangan 215 pesakit berusia 18 tahun keatas yang mempunyai sakit tekak sebagai salah satu aduan di tiga klinik utama penjagaan pada Disember tahun 2016 hingga April tahun 2017. Calitan tekak telah dikumpulkan daripada sekumpulan pesakit bagi tujuan pengkulturan dan analisis. Data mengenai manifestasi klinikal, ciri-ciri demografi, maklumat klinikal dan hasil keputusan

calitan tekak dianalisis dengan menggunakan ujian Chi-square, dan statistik deskriptif. Daripada semua peserta (42.3% lelaki, 57.7% perempuan), (18.6% perokok, 81.4% bukan perokok), (62.8% Melayu, 30.2% India, 5.1% Cina, 1.9% lain-lain), 6 isolat (2.4%) telah dikenalpasti sebagai GAS, 50% daripada mereka yang mempunyai skor Centor 3 ( $p < 0.001$ ), 50% dengan skor Centor 4 ( $p < 0.001$ ), dan 0% dengan skor Centor 0,1 dan 2. Kriteria Centor adalah peramal klinikal yang berkaitan dengan faringo- tonsilitis streptokokus Kumpulan A ( $p < 0.001$ ). Faringo-tonsilitis didiagnosis pada 130 (60.5%) dari jumlah peserta dewasa yang mengadu sakit tekak. Streptokokus beta hemolitik dan virus influenza A dan B telah diasingkan dari 37.1% dan 3.8% daripada jumlah peserta. Kedua-dua skor 3 dan 4 bagi permarkahan Centor mempunyai sensitiviti 50%; manakala kekhususan 97.6% dan 100% masing-masing, nilai ramalan positif 37.5% dan 100% masing-masing; nilai ramalan negatif 98.6% dan 98.6%; nisbah kemungkinan positif 20.5 dan 50; dan nisbah kemungkinan negatif 0.5 dan 0.5. Ketepatan skor Centor 3 adalah 96.3% manakala ketepatan skor Centor 4 adalah 98.6%. Antibiotik telah diberikan kepada 48 peserta (22.3%) termasuk 8.3% streptokokus kumpulan A dan 91.6% dengan streptokokus bukan kumpulan A. Majoriti antibiotik yang ditetapkan dikaitkan dengan skor Centor sifar hingga satu (75%). Ujian kerentanan antibiotik mendedahkan bahawa semua isolat streptokokus beta hemolitik adalah mudah terdedah kepada penisilin G, ampicillin, ofloxacin, cefepime, cefotaxime, ceftriaxone, vancomycin dan linezolid, tetapi 40%, 2.4%, dan 9.6% telah menunjukkan kerintangan kepada tetracycline, clindamycin, dan erythromycin. Kajian semasa mencadangkan bahawa skor Centor berguna bagi mendiagnosis dan membuat keputusan kepada terapi antibiotik untuk faringo- tonsilitis GAS yang membawa kepada penurunan preskripsi antibiotik yang tidak diperlukan, bagi mencapai tahap rawatan yang lebih baik.

Kata kunci: Faringo-tonsilitis; Skor klinikal; Skor Centor; *Streptococcus pyogenes*; Streptokokus beta hemolitik Kumpulan A; Preskripsi antibiotic

## ACKNOWLEDGEMENTS

This work has been achieved with the help of Allah Almighty. I would like to express my deep gratitude and appreciation to my supervisor, Dr. Siti Zulaikha Zakariah, for her valuable and constructive suggestions and patient guidance during the planning and development of this research work. Also, I cannot express enough thanks to Associate Professor Dr. Rukman Awang Hamat, Dr. Nurainul Hana Shamsuddin and Dr. Faisal Ali for their professional guidance and valuable support, enthusiastic encouragement and useful critiques of this research work. Not forget as well, to thank all the nurses, physicians and lecturers in the Department of Medical Microbiology and Parasitology, Department of Family Medicine and three primary care clinics especially who have helped me regarding the sample collection.

I would like to thank my parents for their support and patience. They were always there cheering me up and stood by me through the good times and bad. I wish to thank my lovely sisters (DUNIA and DOA'A) for their supporting and encouragement with their best wishes.



I certify that a Thesis Examination Committee has met on 16 October 2017 to conduct the final examination of Abdulrahman Mansoor Mohammed Muthanna on his thesis entitled "Usefulness of Centor Score for Diagnosis of Group A Streptococcal Pharyngo-Tonsillitis and Prevalence of the Disease in Malaysia from 2016 to 2017" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

**Siti Norbaya binti Masri, PhD**

Senior Lecturer  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Chairman)

**Niazlin binti Mohd Taib, PhD**

Senior Lecturer  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**Tong Seng Fah, PhD**

Professor  
Universiti Kebangsaan Malaysia  
Malaysia  
(External Examiner)



---

**NOR AINI AB. SHUKOR, PhD**  
Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 28 December 2017

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Siti Zulaikha Zakariah, MB BCh BAO, DrPath**

Medical Lecturer  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Chairman)

**Rukman Awang Hamat, MPath**

Associate Professor  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Member)

**Nurainul Hana Shamsuddin, MFamMed**

Medical Lecturer  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Abdulrahman Mansoor Mohammed Muthanna, GS45658

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_  
Name of  
Chairman of  
Supervisory  
Committee: Dr. Siti Zulaikha Zakariah

Signature: \_\_\_\_\_  
Name of Member  
of Supervisory  
Committee: Associate Professor Dr. Rukman Awang Hamat

Signature: \_\_\_\_\_  
Name of Member  
of Supervisory  
Committee: Dr. Nurainul Hana Shamsuddin

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xv
<b>LIST OF FIGURES</b>	xvi
<b>LIST OF ABBREVIATIONS</b>	xviii
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
1.1 Background	1
1.2 Problem statement	2
1.3 Significance of study	3
1.4 Objectives	3
1.4.1 General objective	3
1.4.2 Specific objectives	3
1.5 Research hypothesis	4
1.6 Conceptual Framework	4
<b>2 LITERATURE REVIEW</b>	<b>6</b>
2.1 Definition of pharyngo-tonsillitis	6
2.2 Clinical manifestation of pharyngo-tonsillitis	6
2.3 Risk factors of pharyngo-tonsillitis	7
2.4 Epidemiology of pharyngo-tonsillitis	7
2.5 Etiology of pharyngo-tonsillitis	9
2.5.1 Bacterial pharyngo-tonsillitis	9
2.5.2 Viral pharyngo-tonsillitis	9
2.5.3 Candida pharyngitis:	10
2.6 Tools for clinical scoring	11
2.6.1 Centor scoring system	11
2.6.2 McIsaac scoring system	13
2.6.3 Other tools	14
2.7 Transmission of pharyngo-tonsillitis	14
2.8 Pathogenesis of pharyngo-tonsillitis	15
2.8.1 Bacterial pathogenesis	15
2.8.2 Viral pathogenesis	16
2.9 Complications of pharyngo-tonsillitis	17
2.10 Diagnosis of pharyngo-tonsillitis	18
2.10.1 Clinical diagnosis	18
2.10.2 Laboratory diagnosis	18

2.11	Treatment of pharyngo-tonsillitis	20
2.11.1	Bacterial pharyngo-tonsillitis treatment	20
2.11.2	Other pharyngo-tonsillitis treatment	21
2.12	Antibiotics practices in primary care clinics	21
2.13	Antimicrobial sensitivity and resistance patterns	22
2.14	Prevention of pharyngo-tonsillitis	23

### **3 MATERIALS AND METHODS** 24

3.1	Study design: It was a cross sectional study.	24
3.2	Study area	24
3.3	Study duration	25
3.4	Sampling	25
3.4.1	Study population	25
3.4.2	Sampling population	25
3.4.2.1	Inclusion criteria	25
3.4.2.2	Exclusion criteria	25
3.4.3	Sample size	25
3.4.4	Sampling technique: A simple convenience sampling was used	26
3.5	Participant selection	26
3.6	Specimen collection	26
3.7	Data collection	27
3.7.1	The proforma	27
3.7.2	Clinical presentation	27
3.7.3	Medical report	28
3.8	Specimen transport	28
3.9	Bacterial isolation	28
3.10	$\beta$ haemolytic streptococci identification	29
3.11	Antimicrobial susceptibility test	30
3.12	Viral antigen detection test	31
3.13	Clinical scoring	32
3.14	Data analysis	32
3.15	Ethical approval	32
3.16	Variables of study	32
3.16.1	Dependent variables	32
3.16.2	Independent variables	33
3.17	Definition of terms	33
3.17.1	Pharyngo-tonsillitis	33
3.17.2	Acute pharyngo-tonsillitis	33
3.17.3	Carriers	33
3.17.4	Age group	34
3.17.5	Gender	34
3.17.6	Ethnicity	34
3.17.7	Smoking status	34
3.17.8	Chronic diseases	34
3.17.9	Clinical manifestations	34
3.17.10	Sore throat	34

3.17.11	Fever	35
3.17.12	Cough	35
3.17.13	Rhinorrhea	35
3.17.14	Swollen anterior cervical lymph nodes	35
3.17.15	Tonsillar swelling or exudates	35
3.17.16	Prescribed antibiotics	35
3.17.17	Throat sample results	36
3.17.18	Centor score	36
3.18	Flowchart of study	37
<b>4</b>	<b>RESULTS</b>	<b>38</b>
4.1	Response rate	38
4.2	Normality test	38
4.3	Socio-demographic characteristics of participants	39
4.3.1	Age group	39
4.3.2	Gender	39
4.3.3	Ethnicity	40
4.4	Clinical information of participants	40
4.4.1	Smoking status	40
4.4.2	Chronic diseases	41
4.5	The proportion of pharyngo-tonsillitis among participants	42
4.6	The proportion of beta hemolytic streptococci (groups A, B, C, F and G) and influenza A and B viruses among participants	43
4.7	Association between pharyngo-tonsillitis and socio-demographic characteristics	45
4.7.1	Association between pharyngo-tonsillitis and age group	45
4.7.2	Association between pharyngo-tonsillitis and gender	45
4.7.3	Association between pharyngo-tonsillitis and ethnicity	46
4.8	Association between pharyngo-tonsillitis and clinical information	47
4.8.1	Association between pharyngo-tonsillitis and smoking status	47
4.8.2	Association between pharyngo-tonsillitis and chronic diseases	47
4.9	Distribution of clinical manifestations and Centor score with throat sample results	49
4.9.1	Distribution of clinical manifestations with throat sample results	49
4.9.2	Distribution of Centor score with throat sample results	50
4.10	Validity of the Centor score	51
4.11	Distribution of prescribed antibiotics with Centor score and throat sample results	53
4.11.1	Distribution of prescribed antibiotics according to Centor score	53

4.11.2	Distribution of prescribed antibiotics with throat sample results	54
4.12	Distribution of the types of prescribed antibiotics with throat sample results	55
4.13	Antimicrobial susceptibility patterns of beta hemolytic streptococci (groups A, B, C, F and G) that are associated with pharyngo-tonsillitis	56
<b>5</b>	<b>DISCUSSION</b>	<b>60</b>
5.1	The proportion of pharyngo-tonsillitis among participants	60
5.2	The proportion of beta hemolytic streptococci (groups A, B, C, F and G) and influenza A and B viruses that are associated with pharyngo-tonsillitis	62
5.3	Association between pharyngo-tonsillitis and socio-demographic characteristics	64
5.3.1	Association between pharyngo-tonsillitis and age group	64
5.3.2	Association between pharyngo-tonsillitis and gender	64
5.3.3	Association between pharyngo-tonsillitis and ethnicity	65
5.4	Association between pharyngo-tonsillitis and clinical information	65
5.4.1	Association between pharyngo-tonsillitis and smoking status	65
5.4.2	Association between pharyngo-tonsillitis and chronic diseases	66
5.5	Association between throat sample results and clinical manifestations	67
5.6	Association between throat sample results and Centor score	68
5.7	Validity of the Centor score	69
5.8	Association between prescribed antibiotics and Centor score	71
5.9	Association between prescribed antibiotics and throat sample results	72
5.10	Distribution of the types of prescribed antibiotics with throat sample results	73
5.11	Antimicrobial susceptibility patterns of beta hemolytic streptococci (groups A, B, C, F and G) that are associated with pharyngo-tonsillitis	74
<b>6</b>	<b>SUMMARY, CONCLUSION, LIMITATIONS AND RECOMMENDATIONS</b>	<b>77</b>
6.1	Summary	77
6.2	Conclusion	78
6.3	Limitation	79
6.4	Recommendation	79
6.4.1	Health care providers	79
6.4.2	Further studies	79



<b>REFERENCES</b>	80
<b>APPENDICES</b>	101
<b>BIODATA OF STUDENT</b>	125



## LIST OF TABLES

Table	Page
2.1 Etiology of pharyngo-tonsillitis	10
3.1 Centor criteria	32
4.1 Distribution of the participants by age group	39
4.2 The proportion of beta hemolytic streptococci (groups A, B, C, F and G) and influenza A and B viruses among participants with sore throat (N=215)	44
4.3 Association between pharyngo-tonsillitis and age group	45
4.4 Association between pharyngo-tonsillitis and gender	46
4.5 Association between pharyngo-tonsillitis and ethnicity	46
4.6 Association between pharyngo-tonsillitis and smoking status	47
4.7 Association between pharyngo-tonsillitis and chronic diseases	48
4.8 Association between throat sample results and clinical manifestations	50
4.9 Association between throat sample results and Centor score	51
4.10 Diagnostic test of Centor criteria and scores confirmed with throat sample results	52
4.11 Association between prescribed antibiotics and Centor score	54
4.12 Distribution of prescribed antibiotics with throat sample results	55
4.13 Distribution of the types of prescribed antibiotics with throat sample results	56
4.14 Antimicrobial susceptibility patterns of beta hemolytic streptococci (groups A, B, C, F and G) that are associated with pharyngo-tonsillitis	57

## LIST OF FIGURES

Figure	Page
2.1 Inflamed and swollen tonsils	6
2.2 White patches on the tonsils	7
2.3 Centor scoring management	12
2.4 The Chain of infection transmission	15
2.5 Bacterial pathogenesis	16
2.6 Viral pathogenesis	17
3.1 Sepang district map	24
3.2 Throat swab specimen collection	26
3.3 Appearance of group A streptococcus ( <i>Streptococcus pyogenes</i> ) produces a zone of $\beta$ hemolytic around the colonies and it is sensitive to bacitracin disc on the blood agar, following 24 hours of incubation under anaerobic conditions	29
3.4 Agglutination in No. 1 identified positive result of group A streptococcus and it showed blue latex particles by using a latex agglutination test	30
3.5 Antimicrobial susceptibility result of group A $\beta$ haemolytic streptococcus on Mueller-Hinton agar with 5% sheep blood	31
3.6 Positive result of influenza B virus will show distinct test and control lines	31
4.1 Flowchart of recruitment of the participants	38
4.2 Distribution of participants by gender	39
4.3 Distribution of participants by ethnicity	40
4.4 Distribution of participants according to smoking status	41
4.5 Distribution of participants according to chronic diseases	42
4.6 The proportion of pharyngo-tonsillitis among participants (N=215)	42

4.7	The proportion of beta hemolytic streptococci (groups A, B, C, F and G) and influenza A and B viruses that are associated with pharyngo-tonsillitis (n=130)	44
4.8	Distribution of clinical manifestations among participants (N=215)	49
4.9	Distribution of participants according to Centor score (N=215)	51
4.10	Distribution of prescribed antibiotics according to Centor score	53
4.11	Antimicrobial susceptibility patterns of beta hemolytic streptococci that are associated with pharyngo-tonsillitis	57



## LIST OF ABBREVIATIONS

WHO	World Health Organization
CDC	Centers for Disease Control and Prevention
IMR	Institute for Medical Research
AAFP	American Academy of Family Physicians
PCR	Polymerase chain reaction
HIV	Human immunodeficiency virus
STSS	Streptococcal toxic shock syndrome
EBV	Epstein Barr Virus
HSN	Herpes simplex virus
AGN	Acute Glomerulonephritis
SPSS	Statistical package for the social science
GAS	Group A Streptococci
GBS	Group B Streptococci
GCS	Group C Streptococci
GFS	Group F Streptococci
GGs	Group G Streptococci
βHS	Beta haemolytic streptococci
RADT	Rapid Antigen detection test
ASO	Antistreptolysin-O
CRP	C-reactive protein
EES	Erythromycin Ethylsuccinate

# CHAPTER 1

## INTRODUCTION

### 1.1 Background

Pharyngo-tonsillitis is one of the commonest upper respiratory tract infections and it is an inflammation involving both the tonsils and the pharynx wall caused by viral or bacterial infection (Mishra & Agrawal, 2012). Group A streptococcus (also known as *Streptococcus pyogenes*) is the most common and important bacteria causing acute pharyngo-tonsillitis (Al-Moyed, 2011). Rhinovirus, influenza virus, adenovirus are the main viruses causing pharyngo-tonsillitis, present in 70-80% of cases (Anjos et al., 2014). Sore throat, cough, nasal discharge, fever, swollen tonsils, and difficulty swallowing are the main signs and symptoms of pharyngo-tonsillitis. However, cough and nasal discharge are often associated with viral infections than bacterial infections (Anne & Zab, 2017). Serious complications of pharyngo-tonsillitis caused by group A streptococcus are rheumatic fever, scarlet fever, toxic shock syndrome and acute glomerulonephritis (Al-Moyed, 2011). More than 600 million cases annually have been diagnosed as pharyngo-tonsillitis worldwide. Acute pharyngo-tonsillitis is the second most commonly diagnosed pediatric illness. Group A Streptococcal infection frequently causes significant morbidity and is associated with significant mortality rates worldwide (World Health Organization, 2004). There are many ways to identify the organisms that are causing pharyngo-tonsillitis, such as culture of throat swab, rapid antigen detection tests and molecular diagnosis (PCR) (Cheesbrough, 2008). Viral pharyngo-tonsillitis is treated by symptomatic relief, while bacterial pharyngo-tonsillitis may require the prescription of antibiotics such as penicillin, clindamycin or erythromycin. A specific treatment might be required if complications occurred (Choby, 2009).

Antibiotics are prescribed for treating bacterial infections, not for viral illnesses. In general, antibiotics are safe, but should be prescribed by a physician after a careful clinical assessment. However, antibiotics that are taken unnecessarily may contribute to the development of antibiotic resistance (Andersson et al., 2016). The excessive use of antibiotics, an unnecessary antibiotic prescription and the diagnosis of pharyngo-tonsillitis adds an economic burden to the health care system around the world, as well as to patients and their families. For example, in the United Kingdom it is estimated that health costs for diagnosis of sore throat alone amount to 60 million Pound Sterling per year. Therefore, pharyngo-tonsillitis therapy has a great health and economic impact (Andersson et al., 2016). In Malaysia, because of the paucity of studies about sore throat, the economic impact of pharyngo-tonsillitis and its impact on health are not known (Ministry of Health Malaysia, 2003).

## 1.2 Problem statement

According to World Health Organization, many people are probably facing the problem of sore throat each year, and a majority of physicians depend on the clinical findings to diagnose the upper respiratory tract infections (World Health Organization, 2004). However, previous studies have shown that clinical signs and symptoms alone cannot be used to rule out or diagnose pharyngo-tonsillitis with an adequate sensitivity and specificity, as it can mimic other type of diagnosis infections (Cardoso et al., 2013; Windfuhr et al., 2016; Júnior et al., 2014). Many physicians will automatically prescribe antibiotics based on clinical symptoms, but the Centers for Disease Control and Prevention (CDC) and American Academy of Family Physicians (AAFP) have instructed that antibiotics can only be given to patients with streptococcal pharyngo-tonsillitis to avoid any emergence of antimicrobial resistance among bacteria (Kuehn, 2013; Hersh et al., 2016).

Moreover, most of pharyngo-tonsillitis cases are caused by viruses and do not require antibiotics (Kuehn, 2013). Furthermore, diagnosis of the pathogens cause pharyngo-tonsillitis by throat culture has always been delayed. Ideally, the should be confirmed by throat swab culture, which usually takes two to three days for the bacterial growth to be identified (Upton et al., 2017). During this period, the illness might be resolved or patients might suffer several complications such as scarlet fever, toxic shock syndrome, rheumatic fever and acute glomerulonephritis (Wong et al., 2013). Therefore, there is need a quick and effective diagnosis of pharyngo-tonsillitis on the first visit of the patients and determine if the patients need for antibiotic therapy.

Over prescribing of antibiotics has very serious health effects with severe reactions and may promote antibiotic resistance and add significantly to the cost of health care (Llor & Bjerrum, 2014). Furthermore, the resistance of antimicrobial lead to increase morbidity and mortality since resistance increases the risk of inappropriate therapy (Andersson, 2016). The risk is that the patients who do not receive appropriate therapy will have a longer period of disease or fatal effect; therefore, morbidity and transmission of the microorganism will increase due to the patients remain infectious for a long period (Centers for Disease Control and Prevention, 2014). Also, the increasing trend of antimicrobial resistance is a serious challenge in countries at all economic levels (Ventola, 2015). Therefore, there is a need to document the prescription practices of the physicians in the primary care clinics.

In Malaysia, data on the epidemiology of sore throat or pharyngo-tonsillitis with regard to its prevalence, socio- demographic data, aetiology of the pharyngo-tonsillitis and complications is very much lacking (Ministry of Health Malaysia, 2003). Therefore, there is a need to document pharyngo-tonsillitis with respects to its aetiology, risk factors and demographic data in Malaysia.

### **1.3 Significance of study**

Pharyngo-tonsillitis is major public problems globally. Currently, group A streptococcal pharyngo-tonsillitis account for increasing morbidity and mortality. Unnecessarily antibiotic prescriptions add an economic burden not only on the individuals but also the health care system. Also, the risk factors that are associated with pharyngo-tonsillitis increases the risk of pharyngo-tonsillitis and has not been well studied. Centor score is a tool or guideline based on a set of criteria that help to identify the likelihood of streptococcal infections in patients with a sore throat. It was developed as a quick diagnosis of group A streptococcal pharyngo-tonsillitis in adult patients (Borchardt, 2013). Previous studies have shown that the Centor scoring system has an acceptable sensitivity and specificity and it can limit the over prescription of antibiotics, thus will reduce the emergence of antibiotic resistance as well as the cost of health care (Aalbers et al., 2011; Fine et al., 2012). Therefore, this study aims to determine the prevalence and factors associated with pharyngo-tonsillitis and the validity of Centor score to diagnose group A streptococcal pharyngo-tonsillitis among adult patients with sore throat in Sepang, Selangor, Malaysia. The findings in this study will help in understanding the prevalence of pharyngo-tonsillitis in Malaysia, contribute to the advancement of knowledge on this issue and also hope to be beneficial in further research. Furthermore, this study will be a reference to guide physicians to appropriately prescribe antibiotics for adults with pharyngo-tonsillitis in the local setting depending on Centor score.

### **1.4 Objectives**

#### **1.4.1 General objective**

To describe the epidemiological pattern, etiology and its antibiotic susceptibility, clinical manifestation, antibiotic prescription among adults with sore throat and validity of Centor score in diagnosis group A streptococcal pharyngo-tonsillitis at three primary care clinics in Sepang, Selangor, Malaysia during 2016 to 2017.

#### **1.4.2 Specific objectives**

- a. To determine the socio-demographic characteristics (age, gender, and ethnicity), and clinical information (smoking and chronic diseases which are hypertension, diabetes, asthma, chronic obstructive pulmonary disease, allergy, heart diseases and others), proportion of pharyngo-tonsillitis, acute pharyngo-tonsillitis, beta hemolytic streptococci (groups A, B, C, F and G) and influenza A and B viruses that are associated with pharyngo-tonsillitis among the participants.
- b. To determine the association between the socio-demographic characteristics and the clinical information with pharyngo-tonsillitis.
- c. To determine the validity of using Centor score for management of



- sore throat and the association between clinical manifestations and Centor score with throat sample results.
- d. To determine the types of prescribed antibiotics by the attending physician at the clinics with throat sample results and the association between Centor score and throat sample results with prescribed antibiotics
  - e. To determine the antimicrobial susceptibility patterns of beta hemolytic streptococci (groups A, B, C, F and G) that are associated with pharyngo-tonsillitis.

## 1.5 Research hypothesis


Based upon these data, it hypothesized that:

- a. There is a significant association between socio-demographic data (age, gender and ethnicity) and pharyngo-tonsillitis.
- b. There is a significant association between the clinical information (smoking and chronic diseases which include hypertension, diabetes, asthma, chronic obstructive pulmonary disease, allergy, heart diseases and others) and pharyngo-tonsillitis.
- c. There is a significant association between the clinical manifestations and Centor score with group A streptococcal pharyngo-tonsillitis.

## 1.6 Conceptual Framework

The development of pharyngo-tonsillitis is based on interplay of many correlated factors (Klug, 2014). Pharyngo-tonsillitis is an inflammation involving both the pharynx and tonsils most commonly caused by viral or bacterial infection (Wessels, 2011; Anjos et al., 2014). Pharyngo-tonsillitis reported to be influenced by socio-demographic, risk and environmental factors (Klug, 2017). Clinical manifestations including (Temperature  $\geq 38^{\circ}\text{C}$ , absence of cough, swollen anterior cervical lymph nodes and tonsillar swelling or exudates) and Centor score found to be associated with group A streptococcal pharyngo-tonsillitis (Centor et al., 1981).

## Independent variables



Independent Variables



Dependent Variables



## REFERENCES

- Aalbers, J., O'Brien, K., Chan, WS.,Teljeur, C., Dimitrov, B., & Fahey, T. (2011). Predicting streptococcal pharyngitis in adults in primary care: a systematic review of the diagnostic accuracy of symptoms and signs and validation of the Centor score. *BMC Medicine*, 9(3), 6-9.
- Ab Rahman, N., Teng, C. L., & Sivasampu, S. (2016). Antibiotic prescribing in public and private practice: a cross-sectional study in primary care clinics in Malaysia. *BMC Infectious Diseases*, 16(1), 208.
- Ababneh, M. A., Al-Azzam, S. I., Ababneh, R., Rababa'h, A. M., & Demour, S. A. (2017). Antibiotic prescribing for acute respiratory infections in children in Jordan. *International Health*, 9(2), 124-130.
- Abd El-Ghany, S. M. A., Abdelmaksoud, A. A., Saber, S. M., & El Hamidb, D. H. A. (2015). Group A beta-hemolytic streptococcal pharyngitis and carriage rate among Egyptian children: A case-control study. *Annals of Saudi Medicine*, 35(5), 377-382.
- Agrawal, A., Kumar, D., Goyal, A., Gupta, R., & Bhooshan, S. (2014). Bacteriological Evaluation and Their Antibiotic Sensitivity Pattern in Tonsillitis. *IOSR Journal of Dental and Medical Sciences*, 13(3), 51-55.
- Al-Charrakh, A. H., Al-Khafaji, J. K., & Al-Rubaye, R. H. (2011). Prevalence of  $\beta$ -hemolytic groups C and F streptococci in patients with acute pharyngitis. *North American Journal of Medical Sciences*, 3(3), 129-136.
- Allen, K., & Farah, C. S. (2015). Screening and referral of oral mucosal pathology: a check-up of Australian dentists. *Australian dental journal*, 60(1), 52-58.
- Al-Moyed, K. A. (2011). *Short Notes in Medically Important Bacteria (6th edition)*. Sana'a: Hael Press.
- Alper, Z., Uncu, Y., Akalin, H., Ercan, I., Sinirtas, M., & Bilgel, G. (2015). Diagnosis of acute tonsillopharyngitis in primary care: a new approach for low-resource settings. *Journal of Chemotherapy*, 25(13), 148-155.
- Amanda, W., Steer, A., Smeesters, P., & Curtis, N. (2017). Toxic shock syndrome-management and treatment. *Journal of Infection*, 74(1), 147-152.
- Amir, Y., Eilam, I. (2004). Streptococcal pharyngitis in europe children: epidemiology and treatment. *Clinical Microbiology and Infection*, 8(2), 12-17.
- Andersson, I., Hughes, D., & Kubicek-Sutherland, Z. (2016). Mechanisms and consequences of bacterial resistance to antimicrobial peptides. *Journal*

*Homepage: Drug Resistance Updates, 26(1), 43-57.*

- Anjos, L., Marcondes, M., Mariana, F., Lima, M., Mondelli, A. L., & Okoshi, M. (2014). Streptococcal acute pharyngitis. *Revista Da Sociedade Brasileira de Medicina Tropical, 47(4)*, 409-413.
- Anne, M., & Zab, M. (2017). Upper Respiratory Tract Infection. *Journal of Clinical Infection Diseases, 15(2)*, 5-7.
- Ashgar, S., Barhameen, A., Johargy, A., El-Said, M., & Saati, H. (2016). Prevalence of Streptococcus pyogenes Among Pre-School Children Ages 4 to 6 in Makah city , Saudi Arabia Prevalence of Streptococcus pyogenes Among Pre-School Children. *Medicine Sciences and Healthcare Journal, 10(6)*, 554-558.
- Atlas, S. J., McDermott, S. M., Mannone, C., & Barry, M. J. (2005). The role of point of care testing for patients with acute pharyngitis. *Journal of general internal medicine, 20(8)*, 759-761.
- Attia, M. W., Zaoutis, T., Klein, J. D., & Meier, F. A. (2001). Performance of a predictive model for streptococcal pharyngitis in children. *Archives of pediatrics & adolescent medicine, 155(6)*, 687-691.
- Bakare, O., & Schattner, P. (2010). The usefulness of a clinical “scorecard” in managing patients with sore throat in general practice. *Asia Pacific Family Medicine, 9(1)*, 9.
- Ba-Saddik, I. A., Munibari, A. A., Alhilali, A. M., Ismail, S. M., Murshed, F. M., Coulter, J. B. S., ... & Parry, C. M. (2014). Prevalence of Group A beta-haemolytic Streptococcus isolated from children with acute pharyngotonsillitis in Aden, Yemen. *Tropical medicine & international health, 19(4)*, 431-439.
- Bastian, R. (2016). Candida laryngitis and pharyngitis. Retrieved from <http://laryngopedia.com/candida-laryngitis/>.
- Bedair, G. (2006). *Epidemiology and prevention of streptococcal pharyngitis*. (Unpublished doctoral dissertation). Alexandria University, Egypt.
- Bergman, M., Huikko, S., Pihlajamäki, M., Laippala, P., Palva, E., Huovinen, P., & Seppälä, H. (2004). Effect of macrolide consumption on erythromycin resistance in Streptococcus pyogenes in Finland in 1997-2001. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America, 38(9)*, 1251-1256.
- Berthod, D., Genton, B., Hatz, C., Blum, J., & de Vallière, S. (2015). Ability of physicians to diagnose influenza and usefulness of a rapid influenza antigen test in febrile returning travelers: A randomized controlled trial. *Travel*

*Medicine and Infectious Disease*, 13(15), 394-399.

- Bychkova, V., Shephard, A., Hunt, J., Burova, N., & Radkova, E. (2017). (390) Relief over 2 hours from sore throat, difficulty swallowing, and swollen throat with flurbiprofen 8.75 mg delivered as a lozenge or spray. *The Journal of Pain*, 18(4), 72
- Bioseal. (2014). The Chain of Infection Transmission. Retrieved from <http://www.biosealnet.com/2014/05/chain-infection-transmission/>
- Bitnun, A., & McIsaac, W. (2015). Identifying and treating group A streptococcal pharyngitis in children. *Canadian Medical Association Journal*, 187(1), 13-14.
- Blanton, L., Alabi, N., Mustaqim, D., ...& Taylor, C.(2017). Update: Influenza Activity in the United States During the 2016-2017 Season and Composition of the 2017-2018 Influenza Vaccine. *Journal of Morbidity and Mortality Weekly Report*, 66(25), 668-676.
- Borchardt, R. A. (2013). Diagnosis and management of group A beta-hemolytic streptococcal pharyngitis. *Journal of the American Academy of Physician Assistants*, 26(9), 53-54.
- Botica, M., Botica, I., Stamenić, V., Tambić Andrašević, A., Kern, J., & Stojanović Špehar, S. (2013). Antibiotic prescription rate for upper respiratory tract infections and risks for unnecessary prescription in Croatia. *Journal of Coll Antropol*, 37(2), 449-454.
- Boyarchuk, O., Boytsanyuk, S., & Hariyan, T. (2017). Acute rheumatic fever : clinical profile in children in western Ukraine. *Journal of Medicine and Life*, 10(2), 122-126.
- Boyles, H., & Wasserman, S. (2015). Diagnosis of bacterial infection. *South African Medical Journal*, 105(5), 419-421.
- Bramhachari, V., Kaul, Y., McMillan, J., Shaila, S., Karmarkar, G., & Sriprakash, S. (2010). Disease burden due to *Streptococcus dysgalactiae* subsp. *equisimilis* (group G and C streptococcus) is higher than that due to *Streptococcus pyogenes* among Mumbai school children. *Journal of Medical Microbiology*, 59(2), 220-223.
- Breese, B. B. (1977). A simple scorecard for the tentative diagnosis of streptococcal pharyngitis. *American journal of diseases of children*. 131(5), 514-517.
- Calviño, O., Llor, C., Gómez, F., González, E., Sarvisé, C., & Hernández, S. (2014). Association between C-reactive protein rapid test and group A streptococcus infection in acute pharyngitis. *The Journal of the American Board of Family Medicine*, 27(3), 424-426.

- Cardoso, D. M., Gilio, A. E., Hsin, S. H., Machado, B. M., Paulis, M. D., Lotufo, J. P. B., ... & Grisi, S. J. E. (2013). Impact of the rapid antigen detection test in diagnosis and treatment of acute pharyngotonsillitis in a Pediatric emergency room. *Revista Paulista de Pediatria*, 31(1), 4-9.
- Carlos, G. G., Pekka N. P., & Griffin M. R. (2009). Antibiotic Prescription Rates for Acute Respiratory Tract Infections in US Ambulatory Settings. *Jama Journal*, 302(7), 758-766.
- Centers for Disease Control and Prevention. (2001). *Women more likely than men to visit the doctor, more likely to have annual exams*. Maryland: Vital and Health Statistics.
- Centers for Disease Control and Prevention. (2014). *Antibiotic resistance threats in the United States*. Maryland: Division of Healthcare Quality Promotion.
- Centers for Disease Control and Prevention. (2016). *Preface Antibiotic Stewardship Statement for Antibiotic Guidelines – Recommendations of the Healthcare Infection Control Practices Advisory Committee Introduction*. Maryland: Division of Healthcare Quality Promotion.
- Centor, R. (2012). Adolescent and Adult Pharyngitis: More Than “Strep Throat” Comment on “Large-Scale Validation of the Centor and McIsaac Scores to Predict Group A Streptococcal Pharyngitis. *Journal of Internal Medicine*, 172(11), 852-853.
- Centor, R. M., Atkinson, T. P., Ratliff, A. E., Xiao, L., Crabb, D. M., Estrada, C. A., ... & Waites, K. B. (2015). The Clinical Presentation of Fusobacterium-Positive and Streptococcal-Positive Pharyngitis in a University Health Clinic A Cross-sectional Study Presentation of Bacterial Pharyngitis in a University Health Clinic. *Annals of internal medicine*, 162(4), 241-247.
- Centor, R. M., Witherspoon, J. M., Dalton, H. P., Brody, C. E., & Link, K. (1981). The diagnosis of strep throat in adults in the emergency room. *Medical Decision Making*, 1(3), 239-246.
- Chazan, B., Shaabi, M., Bishara, E., Colodner, R., & Raz, R. (2003). Clinical predictors of streptococcal pharyngitis in adults. *IMAJ-RAMAT GAN-*, 5(6), 413-415.
- Cheesbrough, M. (2008). *District laboratory practice in tropical countries, Part 2 (2nd edition)*. Cambridge: Cambridge University Press.
- Chiappini, E., Principi, N., Mansi, N., Serra, A., De Masi, S., Camaioni, A., ... & Speciale, A. M. (2012). Management of Acute Pharyngitis in Children: Summary of the Italian National Institute of Health Guidelines. *Clinical Therapeutics*, 34(6), 1442-1458.

- Choby, B. A. (2009). Diagnosis and treatment of streptococcal pharyngitis. *Journal of American Family Physician, 158*(1), 79.
- Clinical and Laboratory Standards Institute, (2016). *Performance Standards for Antimicrobial Susceptibility Testing (26th edition)*. USA: Wayne Press.
- Cohen, J. F., Cohen, R., & Chalumeau, M. (2013). Rapid antigen detection test for group A streptococcus in children with pharyngitis. *Cochrane Database of Systematic Reviews*. doi: 10.1002/14651858.CD010502.
- Cohen, J. F., Cohen, R., Levy, C., Thollot, F., Benani, M., Bidet, P., & Chalumeau, M. (2015). Selective testing strategies for diagnosing group A streptococcal infection in children with pharyngitis: a systematic review and prospective multicentre external validation study. *Canadian Medical Association Journal, 187*(1), 23-32.
- Cooper, R., Hoffman, R., Bartlett, G., Besser, E., & Gonzales, R. (2001). Position Paper Principles of Appropriate Antibiotic Use for Acute Pharyngitis in Adults : Background. *Annals of Internal Medicine, 134*(6), 509-517.
- Costantinides, F., Luzzati, R., Tognetto, D., Bazzocchi, G., Biasotto, M., & Tirelli, G. C. (2012). Rapidly progressing subperiosteal orbital abscess: an unexpected complication of a group-A streptococcal pharyngitis in a healthy young patient. *Head & face medicine, 8*(1), 28.
- DeMuri, G., Sterkel, A., Kubica, P., Duster, M., Reed, K., & Wald, E. (2017). Macrolide and Clindamycin Resistance in Group A Streptococci Isolated From Children With Pharyngitis. *Journal of Medical Microbiology, 36*(3), 342-344.
- Devi, U., Borah, P. K., & Mahanta, J. (2011). The prevalence and antimicrobial susceptibility patterns of beta-hemolytic streptococci colonizing the throats of schoolchildren in Assam, India. *Journal of Infection in Developing Countries, 5*(11), 804-808.
- Dhanasekharan, S. V., Rajaram, S., Jayan, M., Sargunan, A., & Shankar, R. (2015). Trends of Antimicrobial Prescription in Department of Otorhinolaryngology in a Tertiary Care Teaching Hospital. *Medi. Res. Chron., 2*(1), 44-50.
- Dhanda, V., Chaudhary, P., Toor, D., Kumar, R., & Chakraborti, A. (2013). Antimicrobial susceptibility pattern of beta-hemolytic group A, C and G streptococci isolated from North India. *Journal of Medical Microbiology, 62*(3), 386-393.
- Dia'a, H. (2010). *Upper Respiratory tract Infection (4th edition)*. Sana'a: Mansor Press.

- Dingle, T. C., Abbott, A. N., & Fang, F. C. (2014). Reflexive culture in adolescents and adults with group A streptococcal pharyngitis. *Clinical infectious diseases*, 59(5), 643-650.
- Donnelly, J. P., Baddley, J. W., & Wang, H. E. (2014). Antibiotic utilization for acute respiratory tract infections in U.S. emergency departments. *Antimicrobial Agents and Chemotherapy*, 58(3), 1451-1457.
- El-Hazmi, M. (2015). Case Report Late-onset prosthetic valve *Corynebacterium diphtheriae* endocarditis caused by nontoxigenic. *The Journal of Infections in Developing Countries*, 9(8), 905-909.
- Ellen, W., Morven, E., & Mary, T. (2016). Patient education: Sore throat in children (Beyond the Basics). *Pediatric Emergency Medicine Practice*, 26(1), 1-6.
- Ellis, E., Corinne, S., Camacho, W., & Mercedes, E. (2015). Appraisal of Clinical Practice Guideline for the Diagnosis and Management of Group A Streptococcal Pharyngitis: 2012 Update by the Infectious Diseases Society of America. *Advanced Emergency Nursing Journal*, 37(1), 34-41.
- Engel, E., Muhamed, B., Whitelaw, C., Musvosvi, M., Mayosi, M., & Dale B. (2014). Group A streptococcal emm type prevalence among symptomatic children in Cape Town and potential vaccine coverage. *The Pediatric Infectious Disease Journal*, 33(2), 208-210.
- Eriksen, M., Judith, M., Neil, S., Farhad, I., & Jeffrey, D. (2015). *The Tobacco Atlas (5th edition)*. Georgia: Library of Congress Cataloging.
- Felsensteina, S., Faddoula, D., Spostob, R., Batoonc, K., Polancoc, C., & Bardd, J. (2015). Molecular and Clinical Diagnosis of Group A Streptococcal Pharyngitis in Children. *Journal of Clinical Microbiology*, 55(7), 50-85.
- Fengying, L. C., Azimi, P., Weisman, L., Joseph, P., Regan, J., Clark, P., ... & Brenner, R. A. (2000). Antibiotic Susceptibility Profiles for Group B Streptococci Isolated from Neonates , 1995-1998. *Clinical Infectious Diseases*, 31(1), 76-79.
- Fine, A. M., Nizet, V., & Mandl, K. D. (2012). Large-scale validation of the Centor and McIsaac scores to predict group A streptococcal pharyngitis. *JAMA Internal Medicine*, 172(11), 847-852.
- Fine, A. M., Nizet, V., & Mandl, K. D. (2013). Participatory Medicine: A Home Score for Streptococcal Pharyngitis Enabled by Real-Time Biosurveillance: A Cohort Study. *Annals of Internal Medicine*, 159(9), 577-583.
- Foong, H. B. B., Yassim, M., Chia, Y. C., & Kang, B. H. (1992). Streptococcal pharyngitis in a primary care clinic. *Singapore medical journal*, 33, 597-597.



- Forward, K. R., Haldane, D., Webster, D., Mills, C., Brine, C., & Aylward, D. (2006). A comparison between the Strep A Rapid Test Device and conventional culture for the diagnosis of streptococcal pharyngitis. *Canadian Journal of Infectious Diseases and Medical Microbiology*, 17(4), 221-223.
- Fozi, K. M., & Kamaliah, M. N. (2013). The effect of profiling report on antibiotic prescription for upper respiratory tract infection. *Malaysian Family Physician*, 8(2), 26-31.
- Fujikawa, S., Ito, Y., Ohkuni, M. (1985). A new scoring system for diagnosis of streptopharyngitis. *Japanese circulation journal*. 49(12), 1258-1261.
- Fujita, K., Murono, K., Yoshikawa, M., & Murai, T. (1994). Decline of erythromycin resistance of group A streptococci in Japan. *Pediatr Infect Dis Journal*, 13(12), 1075-1078.
- Gerber, M. (2005). Diagnosis and treatment of pharyngitis. *Journal of Medical Microbiology*, 52(3), 729-747.
- Gillian, B., (2015). Education Gaps Don't Fully Explain Why Black Unemployment Is So High. Retrieved from <https://www.theatlantic.com/business/archive/2015/12/black-white-unemployment-gap/421497/>
- Gonzales, R., Malone, D., Maselli, H., & Sande, A. (2011). Excessive Antibiotic Use for Acute Respiratory Infections in the United States. *Clin Infect Dis*, 33(6), 757-762.
- Gudjonsson, J. E., Thorarinsson, A. M., Sigurgeirsson, B., Kristinsson, K. G., & Valdimarsson, H. (2003). Streptococcal throat infections and exacerbation of chronic plaque psoriasis: a prospective study. *British journal of dermatology*, 149(3), 530-534.
- Gums, John. (2016). To fight antibiotic resistance, we need to fight bad prescribing habits. Retrieved from <http://theconversation.com/to-fight-antibiotic-resistance-we-need-to-fight-bad-prescribing-habits-60300>
- Habermehl, K. O. (2012). Rapid Methods and Automation in Microbiology and Immunology. *Journal of Clinical Microbiology*, 17(3), 3-7.
- Hajar, R. (2017). Rheumatic Fever and Rheumatic Heart Disease a Historical Perspective. *Journal of the Gulf Heart Association*, 18(1), 39-120.
- Hamza, H. (2009). *Group A streptococcal pharyngitis (GRASP) study*. (Unpublished doctoral dissertation). Cairo University, Egypt.
- Hannaford, P. C., Simpson, J. A., Bisset, A. F., Davis, A., McKerrow, W., & Mills, R. (2005). The prevalence of ear, nose and throat problems in the community:

results from a national cross-sectional postal survey in Scotland. *Family practice*, 22(3), 227-233.

- Hawking, M. K., Lecky, D. M., Lundgren, P. T., Aldigs, E., Abdulmajed, H., Ioannidou, E., ... & Mappouras, D. (2017). Attitudes and behaviours of adolescents towards antibiotics and self-care for respiratory tract infections: a qualitative study. *BMJ Open*, 7(5), 4-14.
- Hedin, K., Bieber, L., Lindh, M., & Sundqvist, M. (2015). The aetiology of pharyngotonsillitis in adolescents and adults - *Fusobacterium necrophorum* is commonly found. *Clinical Microbiology and Infection*, 21(3), 263-267.
- Hersh, K., Fleming-Dutra, K. E., Shapiro, D. J., Hyun, D. Y., & Hicks, L. A. (2016). Frequency of First-line Antibiotic Selection Among US Ambulatory Care Visits for Otitis Media, Sinusitis, and Pharyngitis. *JAMA Internal Medicine*, 176(12), 1870-1872.
- Hoffmann, J., Rabezanahary, H., Randriamarotia, M., Najjar, J. & Vernet, G. (2012). Viral and Atypical Bacterial Etiology of Acute Respiratory Infections in Children under 5 Years Old Living in a Rural Tropical Area of Madagascar, *Journal of PLOS ONE*, 7(8), 3-9.
- Hong, C., Lin, R., Tan, E., Chong, P., Lew, Y., & Loo, L. (2004). Acute respiratory symptoms in adults in general practice. *Family Practice an international journal*, 21(3), 317-323.
- Hoza, A. (2009). Mechanisms of Bacterial Pathogenicity. *Journal of Bacteriology*, 8(3), 1-4.
- Hraoui, M., Boubaker, I. B. B., Rachdi, M., Slim, A., & Redjeb, S. B. (2012). Macrolide and tetracycline resistance in clinical strains of *Streptococcus agalactiae* isolated in Tunisia. *Journal of Medical Microbiology*, 61(8), 1109-1113.
- Huovinen, P., Seppälä, H., Kataja, J., & Klaukka, T. (1997). The relationship between erythromycin consumption and resistance in Finland. Finnish Study Group for Antimicrobial Resistance. *Ciba Foundation Symposium*, 207(6), 36-46.
- Ichikawa, M., Minami, M., Ohashi, M., Wakimoto, Y., Matsui, H., & Hasegawa, T. (2011). Clinical and microbiological analysis of beta hemolytic streptococci during 2006-2010 at Nagoya City University Hospital. *Nagoya medical journal*, 51(4), 175-189.
- Irlam, J., Mayosi, M., Engel, M., & Gaziano, A. (2013). Primary Prevention of Acute Rheumatic Fever and Rheumatic Heart Disease With Penicillin in South African. *Journal of Cardiovascular Quality and Outcomes*, 6(3), 343-352.

- Iseri, K., Iyoda, M., Yamamoto, Y., Kobayashi, N., & Oda, T. (2016). Streptococcal Infection-related Nephritis (SIRN) Manifesting Membranoproliferative Glomerulonephritis Type I. *Internal Medicine*, 55(6), 647-650.
- Iskandar, B., Madiyono, B., Sastroasmoro, S., Sukman, T., Mulyadi, M., & Karuniawati, A. (2008). Comparison of minimal inhibitory and bactericidal capacity of oral penicillin V with benzathine penicillin G to Streptococcus beta-hemolyticus group A in children with rheumatic heart disease. *Paediatrica Indonesiana*, 48(3), 158-161.
- Ismail, H., Hodan, M., & Lim C. (2017). A Retrospective Study of Acute Renal Failure in Children: Its Incidence, Etiology, Complications and Prognosis. *Medical Clinic*, 9(5), 12-40.
- Itzhak, B., & Gober, A. (2015). Effect of smoking cessation on the respiratory functions. *Türkiye Aile Hekimliği Dergisi*, 19(3), 130-135.
- Iwata, K., Fukuchi, T., Hirai, M., Yoshimura, K., & Kanatani, Y. (2017). Prevalence of inappropriate antibiotic prescriptions after the great east Japan earthquake, 2011. *Journal of Medicine*, 96(15), 6-25.
- Jachna, S. K., Pietrzak, A., Bogiel, T., & Gospodarek, E. (2010). Antibiotic sensitivity of beta-hemolytic streptococci isolated from throat swabs and purulent material. *Med Dosw Mikrobiol*, 62(4), 309-317.
- Jawan, J. A. (2016). Understanding Political Contestation in Malaysia. *Suwannabhumi*, 8(2), 51-77.
- Jean, H. P., Revaz, S. A., Bovier, P., & Stalder, H. (2006). Management of Acute Pharyngitis in Adults. *Archives of Internal Medicine*, 166(6), 640-644.
- Joachim, L., Campos, D., & Smeesters, P. R. (2010). Pragmatic scoring system for pharyngitis in low-resource settings. *Pediatrics*, 126(3), 608-614.
- John, G., & Shapiro, D. (1990). Using antibiotic stewardship programs to curb resistance. *Clinical Pharmacology*, 18(2), 16-23.
- John, M., Conrad, M., & Perlstein, D. (2016). Eustachian Tube Problems (Ear Tube Dysfunction, Blockage). *Journal of Clinical Microbiology*, 4(1), 2-3.
- Júnior, B., Rocha, A., Oliveira, C. D. L., Fontes, M. J. F., Lasmar, L. M. D. L. B., & Camargos, P. A. M. (2014). Diagnosis of streptococcal pharyngotonsillitis in children and adolescents: clinical picture limitations. *Revista Paulista de Pediatria*, 32(4), 285-291.
- Karaky, N. M., Araj, G. F., & Tokajian, S. T. (2014). Molecular characterization of Streptococcus pyogenes group A isolates from a tertiary hospital in Lebanon. *Journal of medical microbiology*, 63(9), 1197-1204.

- Karevold, G., Kvestad, E., Nafstad, P., & Kvaerner, K. J. (2006). Respiratory infections in schoolchildren: co-morbidity and risk factors. *Archives of Disease in Childhood*, 91(5), 391-395.
- Karunanayake, C. P., Albritton, W., Rennie, D. C., Lawson, J. A., McCallum, L., Gardipy, P. J., ... & Ramsden, V. (2016). Ear infection and its associated risk factors in first nations and rural school-aged Canadian children. *International journal of pediatrics*, 16(5), 10.
- Kayser, H., Bienz, K., Eckert, J., & Zinkernagel, M. (2005). *Medical Microbiology (1st edition)*. New York: Thieme Press.
- Keenswijk, W. & Johan, W. (2017). A 2-year-old boy with circulatory failure owing to streptococcal toxic shock syndrome: case report. *Journal of Paediatrics and International Child Health*, 16(3), 1-4.
- Kim, S., & Lee, N. Y. (2004). Epidemiology and antibiotic resistance of group A streptococci isolated from health schoolchildren in Korea. *Journal of Antimicrobial Chemotherapy*, 54(2), 447-450.
- Klug, T. E. (2014). Incidence and microbiology of peritonsillar abscess: the influence of season, age, and gender. *European Journal of Clinical Microbiology & Infectious Diseases*, 33(7), 1163-1167.
- Klug, T. E. (2017). Peritonsillar abscess : clinical aspects of microbiology , risk factors , and the association with parapharyngeal abscess. *Danish Medical Journal*, 64(3), 1-39.
- Klug, T. E., Rusan, M., Fuursted, K., Ovesen, T., & Jorgensen, A. W. (2016). A systematic review of *Fusobacterium necrophorum*-positive acute tonsillitis: prevalence, methods of detection, patient characteristics, and the usefulness of the Centor score. *European journal of clinical microbiology & infectious diseases: official publication of the European Society of Clinical Microbiology*, 35(12), 1903-1912.
- Koopmans, M., Wilbrink, B., Conyn, M., Natrop, G., van der Nat, H., Vennema, H., ... & Bosman, A. (2004). Transmission of H7N7 avian influenza A virus to human beings during a large outbreak in commercial poultry farms in the Netherlands. *The Lancet Infectious Diseases*, 363(9409), 587-593.
- Kuehn, B. M. (2013). Excessive Antibiotic Prescribing for Sore Throat and Acute Bronchitis Remains Common. *JAMA Medicine*, 310(20), 2135-2136.
- Kulakarni, D., Bardapurkar, S., Kulkarni, V., & Nilekar, S. (2015). ASO Titer in Acute Rheumatic Fever/ Rheumatic Heart Disease in Pediatric Age Group Materials & Methods. *International Journal of Pharma and Bio Sciences*, 6(3), 102-106.

- Kumar, R., Sharma, Y. P., Thakur, J. S., Patro, B. K., Bhatia, A., Singh, I. P., ... & Sharma, M. (2014). Streptococcal pharyngitis, rheumatic fever and rheumatic heart disease: Eight-year prospective surveillance in Rupnagar district of Punjab, India. *The national medical journal of india*, 27(2), 70-75.
- Kunnamo, A., Korppi, M., & Helminen, M. (2016). Tonsillitis in children: unnecessary laboratory studies and antibiotic use. *World Journal of Pediatrics*, 12(1), 114-117.
- Lauri, H. A., Bartoces, M. G., Roberts, R. M., Suda, K. J., Hunkler, R. J., Taylor Jr, T. H., & Schrag, S. J. (2015). US outpatient antibiotic prescribing variation according to geography, patient population, and provider specialty in 2011. *Clinical Infectious Diseases*, 60(9), 1308-1316.
- Lean, W. L., Arnup, S., Danchin, M., & Steer, A. C. (2014). Rapid diagnostic tests for group A streptococcal pharyngitis: a meta-analysis. *American Academy of Pediatrics News and Journal*, 134(4), 771-781.
- Leão, C., Leal, O., Nascimento, M., & Rodrigues, A. (2015). Evaluation of cytokines produced by Beta-hemolytic streptococcus in acute pharyngotonsillitis. *Brazilian Journal of Otorhinolaryngology*, 81(4), 402-407.
- Lemeshow, S., Hosmer, D., Klar, W., & Lwanga, K. (1990). *Adequacy of Sample Size in Health Studies*. Massachusetts: Wiley Press.
- Lin, R. H., Huang, C. C., Tsou, Y. A., Lin, C. D., Tsai, M. H., Chen, J. H., ... & Shiao, Y. T. (2014). Correlation between Imaging Characteristics and Microbiology in Patients with Deep Neck Infections: A Retrospective Review of One Hundred Sixty-One Cases. *Surgical infections*, 15(6), 794-799.
- Lindgren, C., Neuman, I., Monuteaux, C., Mandl, D., & Fine, M. (2016). Patient and Parent-Reported Signs and Symptoms for Group A Streptococcal Pharyngitis. *AAP News & Journal*, 138(1), 33-37.
- Little, P., Stuart, B., Hobbs, F. R., Butler, C. C., Hay, A. D., Campbell, J., ... & Everitt, H. (2013). Predictors of suppurative complications for acute sore throat in primary care: prospective clinical cohort study. *BMJ Open*, 3(10), 39-43.
- Llor, C., & Bjerrum, L. (2014). Antimicrobial resistance: risk associated with antibiotic overuse and initiatives to reduce the problem. *Journal of Medicine*, 5(6), 229-241.
- Llor, C., Madurell, J., Balagué-Corbella, M., Gómez, M., & Cots, J. M. (2011). Impact on antibiotic prescription of rapid antigen detection testing in acute pharyngitis in adults: a randomised clinical trial. *Br J Gen Pract*, 61(586), 244-251.

- Lopardo, G., Calmaggi, A., Clara, L., Hara, G. L., Mykietiuk, A., Pryluka, D., ... & Klein, M. (2012). Consensus guidelines for the management of upper respiratory tract infections. *Medicina (Argentina)*, 72(6),484-494.
- Lu, B., Fang, Y., Fan, Y., Chen, X., Wang, J., Zeng, J., ... & Li, D. (2017). High Prevalence of macrolide-resistance and molecular characterization of *Streptococcus pyogenes* isolates circulating in China from 2009 to 2016. *Frontiers in Microbiology*, 8, 1052.
- Malino, Y., Utama, L., & Soenarto, Y. (2016). McIsaac criteria for diagnosis of acute group-A  $\beta$ -hemolytic streptococcal pharyngitis. *Paediatrica Indonesiana*, 53(5), 258-267.
- Maltezou, H. C., Tsagris, V., Antoniadou, A., Galani, L., Douros, C., Katsarolis, I., ... & Fretzayas, A. (2008). Evaluation of a rapid antigen detection test in the diagnosis of streptococcal pharyngitis in children and its impact on antibiotic prescription. *Journal of antimicrobial Chemotherapy*, 62(6), 1407-1412.
- Marchello, C., & Ebell, M. H. (2016). Prevalence of Group C *Streptococcus* and *Fusobacterium Necrophorum* in Patients With Sore Throat: A Meta-Analysis. *The Annals of Family Medicine*, 14(6), 567-574.
- Margolis, D. J., Fanelli, M., Kupperman, E., Papadopoulos, M., Metlay, J. P., Xie, S. X., ... & Edelstein, P. H. (2012). Association of pharyngitis with oral antibiotic use for the treatment of acne: a cross-sectional and prospective cohort study. *Archives of dermatology*, 148(3), 326-332.
- Marom, T., Cinamon, U., Itskoviz, D., & Roth, Y. (2010). Changing trends of peritonsillar abscess. *American Journal*, 31(3), 162-167.
- Marta, P., & Whyte A. (1998). Immunology of the tonsils. *Journal of Immunology Today*, 19(9), 414-421.
- Mathan, J. J., Ekart, J., Mills, C., Houlding, A., & Payinda, G. (2017). Clinical management and patient persistence with antibiotic course in suspected group A streptococcal pharyngitis for primary prevention of rheumatic fever: the perspective from a New Zealand emergency department. *The New Zealand medical journal*, 130(1457), 58-68.
- Mathur, P., Bhardwaj, N., Mathur, K., Behera, B., Gupta, G., Kapil, A., ... & Misra, M. C. (2014). Clinical and molecular epidemiology of beta-hemolytic streptococcal infections in India. *Journal of Infection in Developing Countries*, 8(3), 297-303.
- Mazor, O., Granek-Catarivas, M., & Giveon, S. (2016). Common Treatment for Acute Tonsillitis in The Community. *Harefuah Journal*, 155(9), 559-562.

- McIsaac, W. J., Kellner, J. D., Aufricht, P., Vanjaka, A., & Low, D. E. (2004). Empirical validation of guidelines for the management of pharyngitis in children and adults. *Jama Journal*, 291(13), 1587-1595.
- McIsaac, W. J., White, D., Tannenbaum, D., & Low, D. E. (1998). A clinical score to reduce unnecessary antibiotic use in patients with sore throat. *Canadian Medical Association Journal*, 158(1), 75-83.
- Meeker, D., Knight, T. K., Friedberg, M. W., Linder, J. A., Goldstein, N. J., Fox, C. R., ... & Doctor, J. N. (2014). Nudging Guideline-Concordant Antibiotic Prescribing: A Randomized Clinical Trial. *Journal of Internal Medicine*, 174(3), 425-431.
- Meeker, D., Linder, J. A., Fox, C. R., Friedberg, M. W., Persell, S. D., Goldstein, N. J., ... & Doctor, J. N. (2016). Effect of behavioral interventions on inappropriate antibiotic prescribing among primary care practices: a randomized clinical trial. *Jama*, 315(6), 562-570.
- Mehta, N., Schilder, A., Fragaszy, E., ER Evans, H., Dukes, O., Manikam, L., ... & Hayward, A. (2017). Antibiotic prescribing in patients with self-reported sore throat. *Journal of Antimicrobial Chemotherapy*, 72(3), 914-922.
- Mencarini, L., Pasqua, S., & Romiti, A. (2014). Children's time use and family structure in Italy. *Journal of Clinical Microbiology*, 35(15), 21-23.
- Merlini, A. B., Stocco, C. S., Schafranski, M. D., Arruda, P., Bail, L., Borges, C. L., & Dornelles, C. F. (2014). Prevalence of group A Beta-hemolytic streptococcus oropharyngeal colonization in children and therapeutic regimen based on antistreptolysin levels: data from a city from southern Brazil. *The Open Rheumatology Journal*, 8(1), 13-17.
- Michael, R. W. (2011). Group C and group G streptococcal infection. *The New England Journal of Medicine*, 364(5), 48-55.
- Miguel, C., Méndez, P., Martínez, M., Álvarez, L., García, G., & Sánchez, S. (2017). Resistance rates and phenotypic characterization of *Streptococcus pyogenes* in a paediatric population in Northern Spain (2005-2015). *The North American Journal of Medical Sciences*, 30(2), 90-92.
- Ministry of Health Malaysia. (1997). *Management of Sore Throat*. Kuala Lumpur: Public Health Institute.
- Ministry of Health Malaysia. (2003). *Clinical Practice Guideline*. Kuala Lumpur: Academy of Medicine Malaysia. Retrieved from <http://www.moh.gov.my/medical/htm>
- Ministry of Health Malaysia. (2013). National Surveillance of Antibiotic Resistance. Kuala Lumpur: Institute for Medical Research. Retrieved from

<http://www.imr.gov.my>

Ministry of Health Malaysia. (2014). National Antibiotic Guideline. Kuala Lumpur: Pharmaceutical Services Division. Retrieved from <http://www.pharmacy.gov.my>

Mishra, S., Agrawal, D. (2012). A Concise Manual of Pathogenic Microbiology. *In Pathogenesis of Bacterial Infections*, John Wiley and Sons, pp. 35-37. Boston: Hoboken Press.

Mistik, S., Gokahmetoglu, S., Balci, E., & Onuk, F. A. (2015). Sore throat in primary care project: a clinical score to diagnose viral sore throat. *Family practice*, 32(3), 263-268.

Mitchell, P. (2013). Swollen Tonsils Caused by Allergies (Ear and Throat Conditions). Retrieved from <http://www.livestrong.com/article/545736-swollen-tonsils-caused-by-allergies/>

Mouy, D., Cavallo, J. D., Leclercq, R., & Fabre, R. (2001). Antibiotic Susceptibility and Mechanisms of Erythromycin Resistance in Clinical Isolates of *Streptococcus agalactiae* : French Multicenter Study Antibiotic Susceptibility and Mechanisms of Erythromycin Resistance in Clinical Isolates of *Streptococcus agala*. *Antimicrobial Agents and Chemotherapy*, 45(8), 2400-2402.

Naik, B., Nadagir, D., & Asmabegaum, B. (2016). Prevalence of Beta-Hemolytic Streptococci Groups A, C, and G in Patients with Acute Pharyngitis, 8(1), 45-49.

Nakhoul, G., & Hickner, J. (2010). Management of Adults with Acute Streptococcal Pharyngitis : Minimal Value for Backup Strep Testing and Overuse of Antibiotics. *Journal of General Internal Medicine*, 28(6), 830-834.

National Center on Homelessness among Veterans. (2013). *The 2013 Annual Homeless Assessment Report (AHAR) to Congress*. Pennsylvania: The U.S. Department of Housing and Urban.

Neves, L., & Nira, F. (2017). In Brazil, women work 7.5 hours more than men. Retrieved from <http://agenciabrasil.ebc.com.br/en/geral/noticia/2017-03/brazil-women-work-75-hours-more-men>

Nibhanipudi, K. V. (2016). A Study to Determine if Addition of Palatal Petechiae to Centor Criteria Adds More Significance to Clinical Diagnosis of Acute Strep Pharyngitis in Children. *Global Pediatric Health*, 3(1), 1-4.

O'Loughlin, R. E., Roberson, A., Cieslak, P. R., Lynfield, R., Gershman, K., Craig, A., ... & Beall, B. (2007). The Epidemiology of Invasive Group A Streptococcal Infection and Potential Vaccine Implications: United States,



2000-2004. *Clinical Infectious Diseases*, 45(7), 853-862.

- O'Meara, W. P., Mott, J., Laktabai, J., Wamburu, K., Fields, B., Armstrong, J., ... & Pan, W. (2015). Etiology of pediatric fever in Western Kenya: A case-control study of falciparum Malaria, Respiratory Viruses, and Streptococcal Pharyngitis. *American Journal of Tropical Medicine and Hygiene*, 92(5), 1030-1037.
- Opavski, N., Gajic, I., Borek, A. L., Obszańska, K., Stanojevic, M., Lazarevic, I., ... & Mijac, V. (2015). Molecular characterization of macrolide resistant *Streptococcus pyogenes* isolates from pharyngitis patients in Serbia. *Infection, Genetics and Evolution*, 33(2), 246-252.
- Orden, B., Martínez, R., López, A., & Franco, A. (1996). Antibiotic resistance to erythromycin, clindamycin and tetracycline of 573 strains of *Streptococcus pyogenes* (1992-1994). *Enferm Infecc Microbiol Clin.*, 14(2), 86-9.
- Orrling, A., Stjernquist-Desatnik, A., Schalen, C., & Kamme, C (1996). Treatment failure in streptococcal pharyngotonsillitis. An attempt to identify penicillin tolerant *Streptococcus pyogenes*. *SAGE Journals*, 28(2), 143-147.
- Palla, A., Khan, R., Gilani, A., & Marra, F. (2012). Over prescription of antibiotics for adult pharyngitis is prevalent in developing countries but can be reduced using McIsaac modification of Centor scores: a cross-sectional study. *Medicine & Public Health journals*, 24(3), 12-18.
- Paul, L., Stuart, B., Hobbs, F. R., Butler, C. C., Hay, A. D., Campbell, J., ... & Everitt, H. (2013). Predictors of suppurative complications for acute sore throat in primary care: prospective clinical cohort study. *BMJ Open Journals*, 3(10),39-43.
- Pelucchi, C., Grigoryan, L., Galeone, C., Esposito, S., Huovinen, P., Little, P., & Verheij, T. (2012). Guideline for the management of acute sore throat. *Clinical Microbiology and Infection*, 18(1), 1-28.
- Pennock, V., Bell, A., Moxon, T., Reed, P., & Maxwell, F. (2014). Retrospective epidemiology of acute rheumatic fever: a 10-year review in the Waikato District Health Board area of New Zealand. *The New Zealand Medical Journal*, 127(13), 26-37.
- Peter, S. D., & Holt, P. G. (2011). Role of Innate Immunity in the Development of Allergy and Asthma. *Current Opinion in Allergy and Clinical Immunology*, 11(2), 127-131.
- Petković, H., Lukežič, T., & Susković, J. (2017). Biosynthesis of Oxytetracycline by *Streptomyces rimosus*: Past, Present and Future Directions in the Development of Tetracycline Antibiotics. *Food Technology and Biotechnology*, 55(1), 3-13.

- Pickup, R. (2016). What will happen when antibiotics stop working? Retrieved from <http://theconversation.com/what-will-happen-when-antibiotics-stop-working-59938>
- Plainvert, C., Duquesne, I., Touak, G., Dmytruk, N., & Poyart C. (2015). In vitro evaluation and comparison of 5 rapid antigen detection tests for the diagnosis of beta-hemolytic group A streptococcal pharyngitis. *Diagnostic Microbiology and Infectious Disease*, 83(2), 105-111.
- Pramila, Y., Vanita, K., Preety, L., & Siddharth, J. (2010). Drug Utilization Trends in Ent Outpatient Department in a Teaching Hospital. *International Journal of Pharma and Bio Sciences*, 1(4), 153-160.
- Purushothama, V., Dasaraju, A., & Lui, C. (1996). *Medical Microbiology (4th edition)*. Texas: Texas University Press.
- Pyeon, J. S., Moon, K. P., Kang, J. H., Ma, S. H., & Bae, S. M. (2016). Etiology of Acute Pharyngotonsillitis in Children: The Presence of Viruses and Bacteria. *Pediatric Infection & Vaccine*, 23(1), 40-45.
- Rantala, A., Jouni J. J., & Jaakkola M. S. (2011). Respiratory infections precede adult-onset asthma. *PLoS ONE*, 6(12), 279-12.
- Rathi, K., & Raefuddin, A. (2014). Pakistan prevalence survey in acute pharyngitis. *Journal of the Pakistan Medical Association*, 64(8), 928-31.
- Ray, D., Saha, S., Sinha, S., Kumar, N., & Bhattacharya, B. (2016). Molecular characterization and evaluation of the emerging antibiotic-resistant *Streptococcus pyogenes* from eastern India. *BMC Infectious Diseases*, 16(1), 753.
- Regoli, M., Chiappini, E., Bonsignori, F., Galli, L., & Martino, M. (2011). Update on the management of acute pharyngitis in children. *Italian Journal of Pediatrics*, 37(1), 10.
- Reitsma, J. B., Glas, A. S., Rutjes, A. W., Scholten, R. J., Bossuyt, P. M., & Zwinderman, A. H. (2005). Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews. *Journal of clinical epidemiology*, 58(10), 982-990.
- Renner, B., Mueller, C. A., & Shephard, A. (2012). Environmental and non-infectious factors in the aetiology of pharyngitis (sore throat). *Inflammation Research*, 61(10), 1041-1052.
- Rennie, R., Crowson, B., (2013). Management of Upper Respiratory Tract Infections. *Journal of Medical Microbiology*, 99(3), 97-105.

- Roberts, L., Connolly, L., Kirse, J., Evans, K., Poehling, A., Peters, R., & Reid, D. (2012). Detection of group A Streptococcus in tonsils from pediatric patients reveals high rate of asymptomatic streptococcal carriage. *BMC Pediatrics*, 12(1), 3.
- Roggen, I., Berlaer, G., Gordts, F., Pierard, D., & Hubloue, I. (2013). Centor criteria in children in a paediatric emergency department. *BMJ Open Journals*, 3(4), 1-5.
- Rossi, R. E., Girelli, C. M., Bernasconi, G., & Conte, D. (2014). Currently neglected, physical examination maintains its clinical relevance. *Internal and emergency medicine*, 9(8), 871-874.
- Said, M., & Wang, S. (1996). National Health Morbidity survey II. *Asia Pacific Journal of Public Health*, 66(5), 10-18.
- Salyers, A., Wilson, B., Whitt, D. & Winkler, M. (2010). *Bacterial Pathogenesis; A Molecular Approach (3rd edition)*. Washington: Library of Congress Cataloging-in-Publication Data.
- Şanlı, A., Bekmez, E., Yıldız, G., Erdoğan, A., Yılmaz, B., & Gökhan, A. (2016). Relationship between smoking and otorhinolaryngological symptoms. *The Turkish Journal of Ear Nose and Throat*, 26(1), 28-33.
- Schick, R. S., Kelsey, T. W., Marston, J., Samson, K., & Humphris, G. W. (2017). MapMySmoke: feasibility of a new quit cigarette smoking mobile phone application using integrated geo-positioning technology, and motivational messaging within a primary care setting. *Pilot and Feasibility Studies*, 4(1), 19.
- Shaikh, N., Leonard, E., & Martin, J. M. (2010). Prevalence of Streptococcal Pharyngitis and Streptococcal Carriage in Children: A Meta-analysis. *American Academy of Pediatrics Journal*, 126(3), 575-564.
- Sharp, M., Coward, J., Moore, B. R., Salman, S., Marshall, L., Davis, T. M., ... & Manning, L. (2017). A penicillin dried blood spot assay for use in patients receiving intramuscular benzathine penicillin G and other penicillin preparations to prevent rheumatic fever. *Antimicrobial Agents and Chemotherapy*, 61(7), 17-25.
- Shih, T., Lin, C., & Lu, C. (2012). Evaluation of a streptococcal pharyngitis score in Southern Taiwan. *Pediatrics and Neonatology*, 53(1), 49-54.
- Shin, S. M., Shin, J. Y., Kim, M. H., Lee, S. H., Choi, S., & Park, B. J. (2015). Prevalence of Antibiotic Use for Pediatric Acute Upper Respiratory Tract Infections in Korea. *Journal of Korean Medical Science*, 30(5), 617-624.

- Shulman, S. T., Bisno, A. L., Clegg, W., Gerber, M., Kaplan, L., Lee, G., Martin J. M., Van-Beneden, C. (2012). Clinical practice guideline for the diagnosis and management of group A streptococcal pharyngitis. *Clinical Infectious Diseases*, 55(10),1279-1282.
- Shulman, S., & Tanz, R. (2014). Group A streptococcal pharyngitis and immune-mediated complications: from diagnosis to management. *Journal of Expert Review of Anti-Infective Therapy*, 8(2), 137-150.
- Singh, K., Mansul, A., Rama, G., Shaban, H., Chai, K., & Sivalingam, S. (2009). Consensus Guideline on Upper Respiratory Tract Infections. *Malaysian Society of Otorhinolaryngology Head Neck Surgery*, 23(4), 10-22.
- Skovbjerg, S., Roos, K., Olofsson, S., Lindh, M., Ljung, A., Hynsjö, L., ... & Wold, A. E. (2015). High Cytokine Levels in Tonsillitis Secretions Regardless of Presence of Beta-Hemolytic Streptococci. *Journal of Interferon & Cytokine Research*, 35(9), 682-689.
- Sokolo, G. J., Longo-Mbenza, B., Matanda, J. N., Lukoki, E. L., Reddy, P., & Buso, D. (2011). Nose and throat complications associated with passive smoking among Congolese school children. *African health sciences*, 11(3) 315-319.
- Stefaniuk, E., Bosacka, K., Wanke-Rytt, M., & Hryniewicz, W. (2017). The use of rapid test QuikRead go® Strep A in bacterial pharyngotonsillitis diagnosing and therapeutic decisions. *European Journal of Clinical Microbiology & Infectious Diseases*. doi: 10.1007/s10096-017-2986-8.
- Stewart, H., Davis, B., Clemans, L., Littenberg, B., Estrada, A., & Centor, R. (2014). Rapid Antigen Group A Streptococcus Test to Diagnose Pharyngitis: A Systematic Review and Meta-Analysis. *PLoS ONE*, 9(11), 1-10.
- Strömberg, A., Schwan, A., & Cars, O. (1988). Throat carrier rates of beta-hemolytic streptococci among healthy adults and children. *Scandinavian Journal of Infectious Diseases*, 20(4), 411-4117.
- Suhaimi, M. E., Desa, M. N., Eskandarian, N., Pillay, S. G., Ismail, Z., Neela, V. K., ... & Nordin, S. A. (2017). Characterization of a Group B Streptococcus infection based on the demographics, serotypes, antimicrobial susceptibility and genotypes of selected isolates from sterile and non-sterile isolation sites in three major hospitals in Malaysia. *Journal of infection and public health*, 10(1), 14-21.
- Sungho, C., Lee, H., Lee, K., Hwang, K., Bae, S., & Lee, Y. (2001). The emergence of erythromycin-resistant Streptococcus pyogenes in Seoul, Korea. *Journal of Infection and Chemotherapy*, 7(2), 81-86.
- Suzumoto, M., Hotomi, M., Billal, S., Fujihara, K., Harabuchi, Y., & Yamanaka, N. (2009). A scoring system for management of acute pharyngo-tonsillitis in

adults. *Auris Nasus Larynx*. 36(3), 14-20.

- Syahroel, R., Alkamar, A., Djamal, A. A., & Machmud, R. (2008). Clinical predictors of childhood streptococcal pharyngitis. *Paediatrica Indonesiana*, 48(2), 114-9.
- Tall, H., Fatoumata, Y., Timi, N., Aly, S., & Bay, D. (2017). Ear, nose and throat disorders in pediatric patients at a rural hospital in Senegal. *International Journal of Pediatric Otorhinolaryngology*, 96(1), 1-3.
- Tan, G. X., & Tunkel, D. E. (2017). Control of Pain After Tonsillectomy in Children: A Review. *JAMA Otolaryngology–Head & Neck Surgery*. doi: 10.1001/jamaoto.2017.084.
- Tanaka, Y., Gotoh, K., Teramachi, M., Ishimoto, K., Tsumura, N., Shindou, S., & Yamashita, Y. (2016). Molecular epidemiology, antimicrobial susceptibility, and characterization of macrolide-resistant *Streptococcus pyogenes* in Japan. *Journal of Infection and Chemotherapy*, 22(11), 727-732.
- Tanis, D. C., Abbott, A. N., & Fang, F. C. (2014). Reflexive culture in adolescents and adults with group A streptococcal pharyngitis. *Clinical infectious diseases*, 59(5), 643-650.
- Tantivitayakul, P., Lapidattanakul, J., Vichayanrat, T., & Muadchiengka, T. (2016). Antibiotic Resistance Patterns and Related Mobile Genetic Elements of Pneumococci and  $\beta$ -Hemolytic Streptococci in Thai Healthy Children. *Indian J Microbiol*, 56(4), 417-425.
- Teng, C. (2014). Antibiotic prescribing for upper respiratory tract infections in the Asia-Pacific region: A brief review. *Malaysian Family Physician: The Official Journal of the Academy of Family Physicians of Malaysia*, 9(2), 18-25.
- Teng, C. L., Leong, K. C., Aljunid, S. M., & Cheah, M. (2004). Antibiotic Prescription In Upper Respiratory Tract Infections. *Asia Pacific Family Medicine*, 3(1) 38-45.
- Teng, C. L., Tong, S. F., Khoo, E. M., Lee, V., Zailinawati, A. H., Mimi, O., ... & Nordin, S (2011). Antibiotics for URTI and UTI Prescribing in Malaysian primary care settings. *Australian Family Physician*, 40(5), 325-329.
- Tesfaw, G., Kibru, G., Mekonnen, D., & Abdissa, A. (2015). Prevalence of group A B-haemolytic *Streptococcus* among children with pharyngitis in Jimma town, Southwest Ethiopia. *Egyptian Journal of Ear, Nose, Throat and Allied Sciences*, 16(1), 35-40.
- Tiemstra, J., & Miranda, F. (2009). Role of Non-Group A Streptococci in Acute Pharyngitis. *The Journal of the American Board of Family Medicine*, 22(6),

663-669.

- Traub, W. H., & Leonhard, B. (1997). Comparative susceptibility of clinical group A, B, C, F, and G  $\beta$ -hemolytic streptococcal isolates to 24 antimicrobial drugs. *Journal of Chemotherapy*, 43(1), 10-20.
- Treebupachatsakul, P., Tiengrim, S., & Thamlikitkul, V. (2006). Upper respiratory tract infection in Thai adults: Prevalence and prediction of bacterial causes, and effectiveness of using clinical practice guidelines. *Journal of the Medical Association of Thailand*, 89(8), 1178-1186.
- Tyrstrup, M., Velden, A., Engstrom, S., Goderis, G., Molstad, S., Verheij, T., ... & Adriaenssens, N. (2017). Antibiotic prescribing in relation to diagnoses and consultation rates in Belgium, the Netherlands and Sweden: use of European quality indicators. *Scandinavian journal of primary health care*, 35(1), 10-18.
- Upton, A., Bissessor, L., Farrelle, E., Shulmand, S., Zhengd X., & Lennon, D. (2017). Comparison of illumigene Group A Streptococcus Assay with Culture of Throat Swabs from Children with Sore Throats in the New Zealand School-Based Rheumatic Fever Prevention Program. *Journal of Clinical Microbiology*, 55(7), 153-156.
- Van Driel, M. L., De Sutter, A. I., Habraken, H., Thorning, S., & Christiaens, T. (2016). Different antibiotic treatments for group A streptococcal pharyngitis. *The Cochrane Library*, 4(1), 9.
- Van Heirstraeten, L., Coenen, S., Lammens, C., Hens, N., Goossens, H., & Malhotra-Kumar, S. (2012). Antimicrobial drug use and macrolide-resistant *Streptococcus pyogenes*, Belgium. *Emerging Infectious Diseases*, 18(9), 1515-1518.
- Ventola, C. L. (2015). The antibiotic resistance crisis: causes and threats. *Journal for Formulary Management*, 40(4), 277-283.
- Warren-Gash, C., Fragaszy, E., & Hayward, A. C. (2013). Hand hygiene to reduce community transmission of influenza and acute respiratory tract infection: a systematic review. *Influenza and other respiratory viruses*, 7(5), 738-749.
- Wessels, M. (2016). Pharyngitis and Scarlet Fever. *Microbiology and Molecular Biology Reviews : MMBR*, 19(2), 1-19.
- Wessels, M. R. (2011). Streptococcal pharyngitis. *New England Journal of Medicine*, 364(7), 648-655.
- Windfuhr, J. P., Toepfner, N., Steffen, G., Waldfahrer, F., & Berner, R. (2016). Clinical practice guideline: tonsillitis I. Diagnostics and nonsurgical management. *European Archives of Oto-Rhino-Laryngology*, 273(4), 973-

987.

- Wong, C. J., & Stevens, D. L. (2013). Serious group A streptococcal infections. *Medical Clinics*, 97(4), 721-736.
- Woranuch, J., Udomcharoensab, P., Kittika, B., & Lerkiatbundit, S. (2016). Appropriateness of diagnosis of streptococcal pharyngitis among Thai community pharmacists according to the Centor criteria. *International Journal of Clinical Pharmacy*, 38(5), 1318–1325.
- World Health Organization. (2004). *The World Medicines Situation*. Geneva: Department of Essential Medicines and Health Products.
- World Health Organization. (2005). *The Current Evidence for the Burden of Group A Streptococcal Diseases*. Geneva: Department of Child and Adolescent Health and Development.
- Yuniar, T., Anggadiredja, K., & Islamiyah, N. (2017). Evaluation of Rational Drug Use for Acute Pharyngitis Associated with the Incidence and Prevalence of the Disease at Two Community Health Centers in Indonesia. *Scientia Pharmaceutica*, 85(2), 22.
- Zafar, A., Hasan, R., Nizamuddin, S., Mahmood, N., Mukhtar, S., Ali, F., ... & Torumkuney, D. (2016). Antibiotic susceptibility in *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Streptococcus pyogenes* in Pakistan: a review of results from the Survey of Antibiotic Resistance (SOAR) 2002-15. *Journal of Antimicrobial Chemotherapy*, 71(1), 103-109.