

## **UNIVERSITI PUTRA MALAYSIA**

CLASSIFICATION OF SURFACE EMG SIGNALS FOR EARLY SIGNS OF PROLONGED FATIGUE

NURUL FAUZANI BINTI JAMALUDDIN

FK 2018 32



## CLASSIFICATION OF SURFACE EMG SIGNALS FOR EARLY SIGNS OF PROLONGED FATIGUE

By

NURUL FAUZANI BINTI JAMALUDDIN

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Doctor of Philosophy

October 2017

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Doctor of Philosophy

### CLASSIFICATION OF SURFACE EMG SIGNALS FOR EARLY SIGNS OF PROLONGED FATIGUE

## By

## NURUL FAUZANI BINTI JAMALUDDIN

October 2017

Chair: Siti Anom Ahmad, PhD Faculty: Engineering

Sports training are very important to athlete in maintaining and improving their performance. During training, adequate rest is essential to allow recuperation and build body strength. Inadequate rest may expose the body to prolonged fatigue (PF). This condition needs to be managed accordingly to avoid chronic fatigue syndrome. Currently, the non-invasive assessment in identifying PF are training log record, questionnaire and Borg scale. Recent findings indicate that there are strong characteristics on surface electromyography (EMG) under PF conditions such as glycogen breakdown, existence of lactate and soreness. This study extends the investigation of PF signs, especially on the inceptions of PF. An experiment has been conducted on twenty participants to investigate the behavior of surface EMG during five days of intensive training that was based on Bruce Protocol treadmill test. The intention was to induce PF on biceps femoris (BF), rectus femoris (RF), vastus lateralis (VL) and vastus medialis (VM). Besides surface EMG signals, physiological measurements were also collected from the participant.

Physiological results demonstrate that the earliest PF signs developed were soreness, lethargy and performance decrement. For the surface EMG, they went through three main processes: de-noising, feature extraction and classification. De-noising technique through stationary wavelet transform (SWT) was employed in enhancing quality of surface EMG signals. During de-noising process, new method in estimating threshold (*Th*) was proposed. The method demonstrated to have higher performance in term on noise removal and accuracy in PF classification, compared to conventional *Th* methods.

Nine features extracted from the de-noised surface EMG signals. There were changes in median frequency ( $\Delta$ Fmed), mean frequency ( $\Delta$ Fmean), mean absolute value ( $\Delta$ MAV), root mean square ( $\Delta$ RMS), and five features from wavelet indices ( $\Delta$ WI). Daily fatigue

mappings indicate that the emergence of PF can be traced based on extracted features. The mappings indicate that  $\Delta$ Fmed and  $\Delta$ Fmean tend to increase under PF condition for all four muscles BF, RF, VL and VM. Additionally, under PF condition, the mapping indicates an increase in  $\Delta$ RMS and  $\Delta$ MAV but decrease in  $\Delta$ WI for RF muscle.

In the classification stage, Naïve Bayes (NB) and Support Vector Machine (SVM) demonstrate accuracy with 98% and 97% respectively, in distinguish PF on RF, 94% and 96% respectively on BF, both 95% on VL, and 98% and 96% respectively on VM. Thus, this study successfully demonstrates that surface EMG can be used in identifying the inception of PF. The findings presented are significant in sports field to prevent higher degree of PF.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

# PENGKELASAN ISYARAT EMG PERMUKAAN UNTUK TANDA AWAL KELETIHAN BERPANJANGAN

Oleh

## NURUL FAUZANI BINTI JAMALUDDIN

Oktober 2017

### Pengerusi: Siti Anom Ahmad, PhD Fakulti: Kejuruteraan

Latihan sukan adalah sangat penting kepada atlit untuk mengekalkan dan meningkatkan prestasi.Semasa latihan, rehat yang cukup adalah perlu untuk pemulihan and meningkatkan kekuatan sistem badan. Rehat yang tidak mencukupi boleh mendedahkan seseorang kepada keletihan yang berpanjangan (PF). Keadaan ini perlu diurus dengan baik untuk mengelakkan sindrom keletohan yang kronik. Ketika ini, kaedah non-invasif yang digunakan untuk mengenalpasti PF adalah log latihan, soal selidik dan juga skala Borg. Penemuan terbaru menunjukan terdapat karakter elektromyografi permukaan (EMG) di bawah keadaan penurunan glycogen, kewujudan laktat dan kesakitan otot. Kajian ini menyiasat beberapa tanda-tanda PF, terutamanya sewaktu bermulanya PF. Eksperimen dijalankan ke atas dua puluh orang peserta berdasarkan ujian treadmill Protokol Bruce untuk mendorong PF ke atas biceps fermoris (BF), rectus femoris (RF), vastus lateralis (VL) and vastus medialis (VM). Selain dari EMG permukaan, pengukuran fisiologi juga diambil daripada peserta.

Keputusan fisiologi menunjukkan bahawa tanda PF yang paling awal berlaku adalah kesakitan otot, keletihan dan juga penurunan prestasi. Untuk EMG permukaan, ianya melalui tiga proses utama: de-noising, pengekstrakan sifat dan pengkelassan. Kaedah de-noising melalui stationary wavelet transform (SWT) digunakan untuk meningkatkan kualiti isyarat EMG permukaan. Semasa proses de-noising, kaedah baru dalam menentukan nilah ambang (*Th*) diusulkan. Kaedah tersebut telah menunjukkan prestasi yang lebih baik di dalam membuang hingar and ketepatan dalam pengkelasan PF, berbanding kaedah konvensional.

Sembilan sifat diekstrak daripada isyarat EMG permukaan yang telah diproses. Ianya adalah frekuensi pertengahan ( $\Delta$ Fmed), purata frekuensi ( $\Delta$ Fmean), purata nilai absolut ( $\Delta$ MAV), purata punca kuasa ( $\Delta$ RMS), dan lima sifat daripada wavelet indice ( $\Delta$ WI).

Plot keletihan harian menunjukkan bahawa permulaan PF boleh dikenalpasti melalui sifat yang telah diekstrak. Plot menunjukkan bahawa  $\Delta$ Fmed dan  $\Delta$ Fmean cenderung untuk meningkay di bawah keadaan PF untuk keempat-empat otot, BF, RF, VL dan VM. Tambahan lagi, plot juga menunjukkan bahawa  $\Delta$ RMS dan  $\Delta$ MAV meningkat semasa PF, dan  $\Delta$ WI menurun untuk otot RF.

Ketika pengkelasan, Naïve Bayes (NB) dan Support Vector Machine (SVM) menunjukkan ketepatan masing-masing 98% dan 97%, dalam membezakan PF terhadap RF, 94% dan 96% masing-masing terhadap BF, kedua-duanya 95% terhadap VL, dan masing-masing 98% dan 96% terhadap VM. Maka, ini menunjukkan kajian ini berjaya menunjukkan bahawa EMG permukaan boleh digunakan untuk mengenalpasti permulaan PF. Penemuan ini sangat penting dalam industry sukan untuk mengelakkan tahap PF yang lebih tinggi.

#### ACKNOWLEDGEMENTS

First and foremost, with deep sense of gratitude I express my sincere thanks to my esteemed and worthy supervisor, Ir. Assoc. Prof Dr. Siti Anom Ahmad, for her valuable guidance in carrying out this work under her effective supervision, encouragement, enlightenment and cooperation. My sincere thanks are also to my co-supervisor, Assoc Prof Dr Samsul Bahari Mohd Noor and Assoc. Prof Dr Wan Zuha Wan Hasan, for continuous supports all this while. High appreciation is also conveyed to Malaysian Government for financial support via MyBrain15.

My great appreciation dedicated to my ever dearest soul mate, Abu Sahri Ismail, my angels, Arif Fahri and Adlyn Kamila, and my family who have been so tolerant and supports me all these years. Thanks for the encouragement, love and emotional supports that they had given to me.

I am also thankful for those involve directly and indirectly with this project.

Above all I render my gratitude to the Almighty who bestowed self-confidence, ability and strength in me to complete this work.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

## Siti Anom Ahmad, PhD

Associate Professor, Ir Faculty of Engineering Universiti Putra Malaysia (Chairman)

## Samsul Bahari Mohd Noor, PhD

Associate Professor Faculty of Engineering Universiti Putra Malaysia (Member)

## Wan Zuha Wan Hasan, PhD

Associate Professor Faculty of Engineering Universiti Putra Malaysia (Member)

## **ROBIAH BINTI YUNUS, PhD** Professor and Dean

School of Graduate Studies Universiti Putra Malaysia

Date:

#### Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature:

Date:

Name and Matric No.: Nurul Fauzani binti Jamaluddin, GS34417

## **Declaration by Members of Supervisory Committee**

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: Name of Chairman of Supervisory Committee:	
Signature: Name of Member of Supervisory Committee:	
Signature: Name of Member of Supervisory Committee:	

## TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	V
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiv
LIST OF EQUATIONS	xvii
LIST OF ABBREVIATIONS	xviii
LIST OF SYMBOLS	xix

## CHAPTER

 $(\mathbf{C})$ 

1	INT	RODUC	TION	1
	1.2	Motiva	tion and Problem Statement	2
	1.3	Aim an	d Objectives	3
	1.4	Scope of	of Work	4
	1.5	Contrib	oution	5
	1.6	Thesis	Structure	5
2	LIT	ERATU	RE REVIEW	6
	2.1	Introdu	ction	6
	2.2	Mecha	nism of Prolonged Fatigue	6
		2.2.1	Fatigue Detection	9
	2.3	Surface	Electromyography	12
		2.3.1	Surface EMG in Fatigue Detection	12
		2.3.2	Gender and Menstruation Effect on Fatigue	17
			Detection	
	2.4	Surface	EMG Signals Acquisition	18
		2.4.1	Surface EMG Acquisition System	18
		2.4.2	Noise in Surface EMG	22
		2.4.3	Electrodes Placement	24
	2.5	Experii	nent in Prolonged Fatigue	24
		2.5.1	Health Screening and Recruitment	24
		2.5.2	Experiment	25
		2.5.3	Running Activity	26
	2.6	Surface	EMG Signals Processing	28
		2.6.1	De-noising Stage	28
		2.6.2	Features Extraction	31
		2.6.3	Fatigue Mapping and Statistical Test	32
		2.6.4	Classification	33
	2.7	Summa	ıry	37

3	METHODOLOGY	38
	3.1 Introduction	38
	3.2 Surface EMG Data Acquisition System	40
	3.3 Data Collection	43
	3.3.1 Screening Test and Recruitment	43
	3.3.2 Experimental Design	44
	3.3.3 Surface EMG Data Collection	51
	3.4 Signal Processing and Analysis	54
	3.4.1 Wavelet De-noising	54
	3.4.2 Feature Extraction and Statistical Test	67
	3.4.3 Fatigue Mapping	70
	3.4.4 Data Verification	71
	3.4.5 Classification	71
	3.5 Summary	75
	And a second property in the second sec	
4	RESULTS AND DISCUSSION	75
	4.1 Introduction	75
	4.2 Surface EMG System Reliability and Evaluation	75
	4.3 Physiological Fatigue Response Analysis	79
	4.3.1 Intensity of Training	79
	4.3.2 Prolonged Fatigue Signs Identification	80
	4.4 Surface EMG Signal Processing and Analysis	82
	4.4.1 Wavelet De-noising	82
	4.4.2 Feature Extraction, Statistical Test and	90
	Fatigue Mapping	
	4.4.3 Data Verification	102
	4.4.4 Classification	102
	4.5 Summary	108
5	CONCLUSION AND RECOMMENDATION	109
	5.1 Conclusion	109
	5.2 Recommendation	110
REFER	ENCES	111
APPEN	DICES	124
BIODA	TA OF STUDENT	183
LIST O	F PUBLICATIONS	184

xi

## LIST OF TABLES

Table		Page	
2.1	Degree of Prolonged Fatigue and Its Signs	8	
2.2	Methods in Diagnosing Prolonged Fatigue Signs	10	
2.3	Previous Findings on Surface EMG under Normal Fatigue Condition	14	
2.4	Previous Findings on Surface EMG under High Intensity	15	
25	Previous Works on Custom Made Surface FMG System	21	
2.5	Bruce Protocol Treadmill Test	21	Þ
2.7	Training Intensity Zone based on Percentage of Maximal Heart Rate	27	
2.8	Performance Metric of Threshold Estimation Method in De-noising Surface Electromyography Signals	30	
2.9	Previous Findings on Prolonged Fatigue Condition based on Extracted Features	32	
2.10	Review on Features Selection and Classification Techniques in Predicting Muscle Fatigue	34	
2.11	Performance Metric of kNN, LDA, Logistic Regression, NB and SVM	35	
2.12	Advantage(s) and Disadvantage(s) of Cross Validation Methods	36	
3.1	Specification of Surface EMG Acquisition Board	41	
3.2	Configuration Setting in Data Logger	42	
3.3	Quadriceps Muscle Movement based on Flexed and Extended Knee	42	
3.4	Demographic Data of Selected Participants	44	
3.5	Diagnostics Tools and Prolonged Fatigue Signs Observed	45	
3.6	The Confusion Matrix for Two Classes Classification	74	
4.1	Intensity of Training based on HR <sub>max</sub>	79	
4.2	Daily Physiological Fatigue Responses of Twenty Participants	80	
4.3	Number of Participant under Normal Fatigue Condition based on Physiological Responses	81	
4.4	Number of Participant Experience Prolonged Fatigue based on Physiological Responses	81	
4.5	Baseline and SNR Value of De-noised Surface EMG with Different $M_T$	85	
4.6	Baseline and SNR Value of De-noised Surface EMG with $M_T$ range between 0.7 and 0.8	85	
4.7	Performance of Threshold Estimation Method in De- noising Surface EMG Signal	87	
4.8	Signal to Noise Ratio of Raw and De-noised Signals	90	
4.9	Statistical Analysis Result on Frequency Features	92	
4.10	Statistical Analysis Result on Time Features	95	
4.11	Statistical Analysis Result on Wavelet Indices Features	99	

4.12	Classification Performance on Prolonged Fatigue Identification based on Different Threshold Estimation	103
	Method	
4.13	Naïve Bayes classification performance based on different	104
	features selection	
4.14	Support Vector Machine Classification Performance based	106
	on Different Features Selection	



## LIST OF FIGURES

Table		Page	
2.1	Chronology of Prolonged Fatigue	9	
2.2	Surface EMG Signals in Time Domain	11	
2.3	Surface EMG Signals in Frequency Domain	12	
2.4	(a) Frequency Spectrum of Surface EMG when Light	13	
	Load Added to Muscle		
	(b) Frequency Spectrum of Surface EMG when Heavy		
	Load Added to Muscle		
2.5	(a) Amplitude of Surface EMG Behaviour under Fatigue	13	
	on Sub-maximal Contraction		
	(b) Amplitude of Surface EMG Contraction		
2.6	Chronologies of Prolonged Fatigue and Surface EMG	16	
	Behaviour		
2.7	Pre-gelled Ag/Ag Cl Electrode	18	
2.8	Basic Diagram for Bio-potential Amplifier Design	19	
2.9	(a) Differential Amplifier	20	
	(b) Instrumentation Amplifier		
2.10	Basic Structure of Surface EMG Acquisition System	22	
2.11	Surface EMG Signal Processing Stage	28	
2.12	Feature extraction method;	31	
	(a) Overlapped Windowed Technique		
	(b) Non-overlapped Windowed Technique		
3.1	General Structure of Research Methodology	38	
3.2	Comprehensive Overview of Research Methodology	39	
3.3	Block Diagram of Surface EMG Acquisition System	40	
3.4	Surface EMG Acquisition Board	41	
3.5	Duration of Experiment	46	
3.6	Flow Process of Phase I: Familiarization	48	
3.7	Symptoms of Fatigue Observed at Before and During	48	
	Running		
3.8	Example of Familiarization with Equipment	49	
	(a) Treadmill		
	(b) Blood Pressure and Heart Rate		
3.9	Flow Chart of Phase II: Intensive Training	50	
3.10	Electrodes Placement of Rectus Femoris, Vastus Medialis	52	
	and Vastus Lateralis Standard based on SENIAM		
3.11	Movement to Activate Quadriceps Muscles Group	53	
3.12	Electrodes Placement of Biceps Femoris Standard based	53	
2.12	on SENIAM		
3.13	Movement to Activate Biceps Femoris Muscle	54	
3.14	Block Diagram of Signal Processing and Analysis	54	
3.15	De-noising Process of Wavelet Transform	55	
3.16	Decomposition Process based on Frequency Sampling	56	
2.17	1400 HZ	57	
3.17	Frequency Spectrum	57	

3.18	Coefficient Approximations of Decomposition Process	58
	and Its Frequency Spectrum	
3.19	Example of De-noising Process based on Threshold Value	59
3.20	Corner Frequency Noise in Collected Surface EMG	60
	Signal	
3.21	Estimation of Threshold Value for Baseline Noises	61
	Removal	
3.22	Method in Determine DC, DB and MB	62
3.23	Coefficient Details with DC-DB Value for Case I	63
3.24	Coefficient Details with DC-DB Value for Case 2	64
3 25	Surface EMG Signal Clearly Visible in Coefficient	64
0.20	Details	Ű.
3 26	Flow Chart of Identifying M <sub>r</sub> Value	65
3 27	Flow Chart of De-noising Process based on Proposed	65
5.21	Estimation Threshold	00
3 28	Non-overlapping Windowing Technique in Features	67
5.20	Extraction	07
3 20	Basic Diagram of Classification Process in Prolonged	72
5.27	Easie Diagram of Classification Process in Prolonged	12
4.1	Validation on Surface EMG Signals Detection by Using	77
7.1	Developed System	//
4.2	Full Contraction and Baseline of Surface EMG Signals	78
7.2	and Its Frequency Spectrum	70
13	Example of Surface EMG Signals Acquired from Bicens	78
т.Ј	Example of Surface Envio Signals Acquired from Diceps	70
	Medialis	
44	Denoised Surface EMG Signals with Different Mr	83
т.т	Applied	05
45	Baseline and SNR Value of Surface EMG with Different	84
т.5	M <sub>m</sub>	0-
4.6	Frequency Domain of Surface FMG Before and After De	86
4.0	noising Process	80
17	Paw and De noised Surface EMC based on Different	86
4.7	Threshold Estimation Method, and its Frequency	80
	Speetrum	
18	Basalina of Surface EMG Bafara and After Da paising	00
4.0	Baseline of Surface EMO Before and After De-hoising	00
4.0	Contraction of Surface EMG Pafere and After De noising	80
4.9	Dragoss	09
4.10	From Distribution of Surface EMC in Spectral at Dra	01
4.10	and Dest of Dunning	91
	(a) Under Normal Fatigue Condition	
	(a) Under Normal Fatigue Condition	
4 1 1	(b) Onder Protonged Faligue Condition	01
4.11	(a) Changes in Finean under Normal and Prolonged	91
	raugue Condition (b) Changes in Emod under Marriel and Prolonged	
	(b) Unanges in Fined under Normal and Prolonged	
4.10	Faugue Collelium Daily Eatimus Manning based on Energy of for Nergy 1	02
4.12	Daily rangue Mapping based on Frequency for Normal	93
	and Froibinged Faugue Condition	

5

4.13	Amplitude of Surface EMG at Pre and Post Running	94	
	Activity;		
	(a) Under Normal Fatigue Condition		
	(b) Under Prolonged Fatigue Condition		
4.14	(a) Changes in RMS under Normal and Prolonged Fatigue	95	
	Condition		
	(b) Changes in MAV under Normal and Prolonged		
	Fatigue Condition		
4.15	Daily Fatigue Mapping based on Amplitude for Normal	98	
	and Prolonged Fatigue Condition		
4.16	(a) Changes in WIRM1551 under Normal and Prolonged	97	
	Fatigue Condition		
	(b) Changes in WIRM1M51 under Normal and Prolonged		
	Fatigue Condition		
	(c) Changes in WIRM1522 under Normal and Prolonged		
	Fatigue Condition		
	(d) Changes in WIRE51 under Normal and Prolonged		
	Fatigue Condition		
	(e) Changes in WIRW51 under Normal and Prolonged		
	Fatigue Condition		
4.17	Daily Fatigue Mapping for Normal and Prolonged Fatigue	100	
	Condition		
	(a) $\Delta$ WIRM1551		
	(b) $\Delta WIRM1M51$		
	(c) $\Delta$ WIRM1522		
4.18	Daily Fatigue Mapping for Normal and Prolonged Fatigue	101	
	Condition for AWIRE51 and AWIRW51		

C

## LIST OF EQUATIONS

## Equations

2.1	HR <sub>max</sub>	27	
3.1	Notch Filter	40	
3.2	Signals to Noise Ratio		
3.3	Percentage of HR <sub>max</sub>	45	
3.4	Endurance Time	49	
3.5	Threshold Estimation based on RigRSURE Method	59	
3.6	Threshold Estimation based on Universal Threshold	59	
	Method		
3.7	Proposed Threshold Estimation Method	61	
3.8	Threshold Estimation Method for Case I	62	
3.9	Threshold Estimation Method for Case II	63	
3.10	Median Frequency	68	
3.11	Mean Frequency	68	
3.12	Mean Absolute Value	68	
3.13	Root Mean Square	68	
3.14	Wavelet Indices Ratio between Moment -1 at Scale 5	69	
	and Moment 5 at Scale 1		
3.15	Wavelet Indices Ratio between Moment -1 at	69	
	Maximum Energy Scale and Moment 5 at Scale 1		
3.16	Wavelet Indices Ratio between Moment -1 at Scale 5	69	
	and Moment 2 at Scale 2		
3.17	Wavelet Indices Ratio of Energies at Scale 5 and 1	69	
3.18	Wavelet Indices Ratio between Square Waveform	70	
	Lengths at Different Scales		
3.19	Changes of Features	70	
3.20	Column Vector y	73	
3.21	Class Conditional Independence of the Features	73	
3.22	Probability of the Features Vector Class	73	
3.23	Discriminant Function	73	
3.24	Margin Estimation of The Two Nearest Points of	74	
	Neighboring Classes		
3.25	Classification Accuracy	75	
3.26	Classification Sensitivity	75	
3.27	Classification Precision	75	

## LIST OF ABBREVIATIONS

ADC	Analog to Digital Conversion
Ag/Ag Cl	Silver silver Chloride
AIS	Anterior Aliac Apine
В	Details Baseline
BF	Biceps Femoris
B <sub>M</sub>	Maximum Baseline
BMI	Body Mass Index
BP	Blood Pressure
С	Details Contract
cA	Approximation Details
cD	Coefficient Details
CV	Cross Validation
DWT	Discrete Wavelet Transform
EMG	Electromyography
FFT	Fast Fourier Transform
FIR	Finite Impulse Response
Fmean	Mean Frequency
Fmed	Median Frequency
HPF	High Pass Filter
HR	Heart Rate
HRmax	Maximum Heart Rate
IA	Instrumentation Amplifier
IEMG	Integral EMG
MAV	Mean Absolute Value
M <sub>T</sub>	Multiplier Threshold
NB	Naïve Bayes
NF	Normal Fatigue
PF	Prolonged Fatigue
PSD	Power Spectral Density
RF	Rectus Femoris
RMS	Root Mean Square
RPE	Rating of Perceived Exertion
SENIAM	Surface Electromyography for the Non-Invasive Assessment of Muscles
SNR	Signals to Noise Ratio
SURE	Stein's Unbiased Risk Estimate
SVM	Support Vector Machine
SWT	Stationary Wavelet Transform
Th	Threshold
VL	Vastus Lateralis
VM	Vastus Medialis
WI	Wavelet Indices

## LIST OF SYMBOLS

- Ι Decomposition Level
- N
- X Fs
- Number of Samples Signal Frequency Sampling Length of Signal Changes Time Standard deviation
- $L \\ \Delta \\ T \\ \Sigma$

Ċ,



#### CHAPTER 1

#### **INTRODUCTION**

#### 1.1 Introduction

Sports training are very important to athlete in maintaining and improving their performance. Nevertheless, sports training cause the athlete to become fatigue. Fatigue is a natural physiological sign showing the limitation of a human body. It is characterized by lower level of performance, reduction of organism forces, and failure to maintain the level of forces exerted (Al-Mulla & Sepulveda, 2010; Boyas & Guével, 2011; Davis, Walsh, & Edin, 2010; C J De Luca, 1984; R. H. Edwards, 1981; Sesboüé & Guincestre, 2006).

In normal fatigue (NF), fatigue usually disappears by itself after a while. Recovering from fatigue indicates that biochemical reactions during sports activity are able to return to a normal level (Pearce, 2002). Occasionally, the load and intensity of sports training are greater than the body can handle and this leads to imbalance. However, adequate recovery process due to imbalance enhances strength and improve performance (Howwit, 2004; Myrick & Aprn, 2015). Unfortunately, inadequate recovery period causes more biochemical reactions accumulation; thus fatigue continues (Kreher & Schwartz, 2012). Inadequate recover (Myrick & Aprn, 2015).

There are several degrees of PF, and it is categorized according to the diagnosed signs. The mildest signs of PF are musculoskeletal disorder (soreness, pain or aches), unexplained lethargy, performance decrement, restlessness, hypertension and sleeping and psychological disturbance (Jeukendrup, 2010; Purvis, Gonsalves, & Deuster, 2010). Playing sports under this condition may lead to injury, continued underperformance, possible damage of body structure and development of more serious state of PF (Jeukendrup, 2010; Myrick & Aprn, 2015). If this situation occurs, the athlete is not able to participate in any competition; hence, risking the athlete's career (Pearce, 2002). The recovery process is time and cost consuming, and affects national sports industry as the athletes fail to participate in national and international competitions.

Prevention from the occurrences of PF condition is the best solution (Pearce, 2002). The identification of PF signs at the earliest of its inception helps in providing proper fatigue and training management. There are no specific tools used for PF identification. Currently, PF is diagnosed invasively through blood test and biopsies. Alternatively, PF can also be diagnosed non-invasively through heart rate, endurance time, the rating of perceived exertion and distress questionnaire (M. Haddad et al., 2013; Hernandez, Estrada, Garcia, Sierra, & Nazeran, 2010; Sakurai et al., 2010).

Nonetheless, there is a significant growth in investigating fatigue via surface electromyography (EMG). In fact, the progression of fatigue process can be determined via surface EMG (Massó et al., 2010).

Surface EMG is a reflection of the behavior, potentials, voltages and electrical field of human skeletal musculature (Joachim H.Nagel, Bronzino, 2000; Medved & Cifrek, 2011; Thakor, 1999). It is acquired by placing electrodes on the surface skin, near to human muscle. The amplitude and frequency of the signals signify the behavior and condition of muscle's motor unit, conduction velocity and ionic alteration of the muscle. Furthermore, there are many advantages of using surface EMG in investigating fatigue including it is a non-invasive process, easy to use, comfortable and inexpensive. Any investigation on fatigue based on muscle evaluation is denoted as muscle fatigue.

Fatigue can be determined by changes in its amplitude and frequency. Most of the opinions agree that the degree of fatigue begins with the increment of amplitude, followed by the unchanged and decreased trends, as well as accompanied by the decrement center of frequency (Dimitrova & Dimitrov, 2003). Recently, there is significant growth in investigating PF using surface EMG. It covers several PF signs such as biochemical reactions, performance decrement and soreness (Gavin, Myers, & Willems, 2015; Sterczala & Nicol, Jx; Fry, 2015; Tenan, Blackburn, & Robert, 2016). The results of these studies prove that useful information can be extracted from surface EMG signals to distinguish PF condition.

Hence, this thesis focuses on investigating surface EMG behavior under the earliest signs of PF. In fact, the characteristics identified can be used as a prevention measure for the serious state of PF.

### 1.2 Motivation and Problem Statement

Reports show that 20% to 60% of athletes, 60% of elite runners, 33% of non-elite runners, and 35% swimmers experienced PF at least once in their career life (Kreher & Schwartz, 2012; Myrick & Aprn, 2015; Purvis et al., 2010). Meanwhile, Nederhof, Lemmink, Visscher, Meeusen, & Mulder, (2006) reported that 20% to 60% of athletes have experienced chronic fatigue due to untreated signs of PF.

To date, invasive methods such as muscle biopsies and blood test are considered as the most reliable indicator in diagnosing PF. However, the methods are time-consuming, cost ineffective and unable to be applied frequently. The existing non-invasive methods mostly require close supervision and time-consuming. This method is ineffective in monitoring a large number of athletes. Nevertheless, identifying early signs of prolonged fatigue is important to avoid injury and disorder. Most importantly, it is able to prevent the occurrence of higher PF state. Therefore, an alternative method such surface EMG, which is cost effective, easier and requires less processing time, procedure and supervision, is essential in diagnosing early sign of PF.

Surface EMG has been used to identify fatigue since 1912. Currently, the information is limited to non-fatigue, the transition to fatigue and fatigue under normal condition. There are recent and growth findings from high-intensity exercise and associated with PF condition. However, the findings are restricted to glycogen breakdown, existing of lactate and soreness only. Investigation on other PF signs, especially during the emergence of the condition is essential for early detection. That is why the investigation of surface EMG under PF condition benefits in determining fatigue classification. At the moment, the classification is successfully performed in distinguishing between non-fatigue to fatigue (Marri & Swaminathan, 2015; Sarillee et al., 2015; Venugopal, Navaneethakrishna, & Ramakrishnan, 2014; Zhang, Jian E.Lockhart, Thurmon Soangra, 2015), and transition from non-fatigue and fatigue (Sepulveda, Colley, & Kattan, 2009) under normal condition. Therefore, the classification to distinguish fatigue under normal condition (NF) and PF has many advantages, particularly in sports.

The success of the classification process highly depends on the quality of surface EMG. Surface EMG is commonly contaminated by corner frequency and baseline noises. Nevertheless, corner frequency noise is easily removed by applying high-pass filter, since it exists in a specific range of frequency. Unfortunately, there is no specific frequency range for baseline noises. Worse, baseline noises merely exist within the surface EMG frequency range. Fret not, wavelet de-noising method can be used in removing baseline noises. In the de-noising process, thresholds need to be estimated. During de-noising process, signals above threshold value are preserved, while signals under threshold value are removed. However, the available conventional threshold estimations are not suitable for surface EMG application. The methods tend to remove important information from surface EMG signals and keep unwanted signals (Bartolomeo, 2012; Phinyomark, Limsakul, & Phukpattaranont, 2009b). Therefore, new estimation of threshold value that is suitable for surface EMG application is required. Essentially, the new threshold estimation is capable to remove noises and preserves a great deal of information. In addition, the de-noised signals based on the proposed technique have better performance in fatigue identification and classification.

#### 1.3 Aim and Objectives

This thesis aims to investigate the behavior of surface EMG signals under the earliest signs of mild PF condition. The classification of PF will help the practitioner to identify and manage fatigue accordingly. This classification certainly assists in preventing a higher state of PF, which eventually is capable of preventing more serious injury.

The specific objectives of the study are as follow:

- 1. To propose a new technique in estimating threshold value for surface EMG wavelet de-noising application.
- 2. To investigate the surface EMG features behavior under the earliest signs of PF developed during five days of intensive training.
- 3. To investigate the significant features and muscle activation types for classification process that can be used to imply the inception of PF.

## 1.4 Scope of Works

The intensive training experiment is conducted based on five consecutive days of Bruce Protocol treadmill test. It is employed in the experiment as it provides highintensity exercise and induces PF signs. As currently practiced, PF signs observed are limited to mild signs such as soreness, performance reduction, restlessness, hypertension, unexplained lethargy, sleeping disturbance and psychological stress. In this study, EMG signals are analyzed to investigate the shown behavior under the earliest signs of PF. PF signs are diagnosed non-invasively by Sports Officer Akademi Sukan Universiti Putra Malaysia (UPM) through 24-hours training distress questionnaire, training log and interview session. The occurrence of PF signs within five days of Bruce Protocol treadmill test (intensive training) indicates the earliest signs developed. This certainly means that not all PF signs are involved. Again, the signs developed are subjected to ethical reasons and potential risk endured by the participants. In addition, five days of intensive training on recruited subject is presumed to be adequate as approved by Jawatankuasa Etika UPM.

During the investigation, firstly, four lower extremities muscles were chosen since these are the activated muscles during running, and suffer the highest muscle injuries during sports particularly during running activities. Secondly, only healthy subjects, who possessed the inclusion and exclusion criteria between the age of 20 to 30 and with the body mass index (BMI) 18 < x < 25, were selected to participate in these experiments. Finally, in total twenty (20) participants, five (5) males and fifteen (15) females, were recruited.

In the study, the pre-processing tool utilized for detecting surface EMG signals was Stationary Wavelet Transform (SWT). The conventional threshold methods applied in de-noising process were RigRSURE, HeurSURE, minimax and fixed from threshold. The proposed new estimation threshold value adopted SWT, with six level decomposition and 'db45' employed as mother wavelet. Apart from comparing the de-noised signals to standard quality of surface EMG signals, removing corner frequency and mitigating baseline noise to acceptable value, the performance of threshold estimation (for conventional and proposed methods) were verified by its ability to classify PF.

In fact, PF was classified in accordance to supervised classification methods namely Naïve Bayes (NB) and support vector machine (SVM). These classifications were made on the individual muscle. Then, the performance of the classifications was evaluated based on its accuracy, specification, precision and cross-validation (CV) error.

#### 1.5 Contribution

General contribution of the study lies in the ability to identify the earliest PF signs based on EMG signal analysis and classification. It indicates that there are other noninvasive tools which are quantifiable, have less procedure and need less supervision in diagnosing PF. Apart from that, the study contributes to the society as the identification can be used as one of on-site monitoring tools in sports training. The outcomes of the study assist particularly in training management to prevent injury and serious state of PF. Other than that, it helps in selecting athletes during competitive matches to ascertain their ability in delivering optimal performance.

### 1.6 Thesis Structure

This thesis consists of five chapters which describe the research activities involved. Chapter 1 presents a general introduction, problem statements, and motivation to do this research. This chapter clarifies the objectives, scopes of research works, contributions of this study, and the structure of thesis.

Chapter 2 discusses the literature review related to the research work. The chapter further elaborates the mechanism of PF, its signs and diagnostics tool commonly used. In addition, this chapter reviews the ability of surface EMG in PF identification. The review includes the characteristics of surface EMF signals itself, methods of acquisition and analysis.

Chapter 3 describes the research methodology employed in this study. It includes the development of surface EMG data acquisition system and data collection procedure. The design and consideration in proposing the new threshold value estimation method are also presented in this chapter. Apart from that, this chapter defines the feature extraction parameters involved, and the classification techniques used in the study.

Meanwhile, the main findings and discussions are presented in chapter 4. The performance of de-noising technique is presented first, then, followed by the results of the experimental design which includes physiological responses during the intensive training. The findings are presented in the form of fatigue mapping for the extracted features. Not only that, this chapter also presents classification performance in PF identification.

#### REFERENCES

- Al-imari, A. A., A.Rashid, K., & Egaidy, N. H. Al. (2003). Design and Implementation of a Surface Electromycogram System for Sport Field Application. In *The 3rd IEEE International Workshops on System-on-Chip for Real-Time Applications* (pp. 357–361).
- Al-Mulla, M. R., & Sepulveda, F. (2010). A Novel Feature Assisting in the Prediction of sEMG muscle fatigue towards a wearable autonomous system. In 2010 IEEE 16th International Mixed-Signals, Sensors and Systems Test Workshop (IMS3TW) (pp. 1–6).
- Arabadzhiev, T. I., Dimitrov, V. G., & Dimitrov, G. V. (2014). The increase in surface EMG could be a misleading measure of neural adaptation during the early gains in strength, pp. 1645–1655.
- Arendt-Nielsen, L., & Sinkjær, T. (1991). Quantification of human dynamic muscle fatigue by electromyography and kinematic profiles. *Journal of Electromyography and Kinesiology : Official Journal of the International Society* of Electrophysiological Kinesiology, 1(1), pp. 1–8.
- Armfield, D. R., Kim, D. H.-M., Towers, J. D., Bradley, J. P., & Robertson, D. D. (2006). Sports-related muscle injury in the lower extremity. *Clinics in Sports Medicine*, 25(4), pp.803–42.
- Bajaj, P., Madeleine, P., Sjøgaard, G., & Arendt-nielsen, L. (2002). Assessment of Postexercise Muscle Soreness by Electromyography and Mechanomyography. *The Journal of Pain*, 3(2), pp.126–136.
- Bartolomeo. (2012). Wavelet thresholding technique for sEMG denoising by baseline estimation. *International Journal of Computer Aided Engineering and Technology*, 4(6), pp. 517–534.
- Basmajian, J. V., & De Luca, C. J. (1985). *Muscles Alive. Their Functions Revealed by Electromyography* (5th ed.). Williams & Wilkins.
- Birch, K., & George, K. (1999). Overtraining the female athlete. *Journal of Bodywork* and Movement Therapies, 3(1), pp. 24–29.
- Blanc, Y., & Dimanico, U. (2010). Electrode Placement in Surface Electromyography ( sEMG) "Minimal Crosstalk Area" (MCA). *The Open Rehabilitation Journal*, 4(3), pp.110–126.
- Bonato, P., Roy, S. H., Knaflitz, M., & Luca, C. J. De. (2001). Time-Frequency Parameters of the Surface Myoelectric Signal for Assessing Muscle Fatigue

during Cyclic Dynamic Contractions. *IEEE Transactions on Bio-Medical Engineering*, 48(7), pp.745–754.

- Bowser, K. L. (1997). *The Effect of Daily, Maximal Resistance Exercise on Muscular Function.* (Unpublished Dissertation), Virginia Politechnic Institute and State University.
- Boyas, S., & Guével, a. (2011). Neuromuscular fatigue in healthy muscle: underlying factors and adaptation mechanisms. *Annals of Physical and Rehabilitation Medicine*, 54(2), pp.88–108.
- Broman, H., Bilotta, G., & Luca, C. J. D. E. (1985). Myoelectric signal conduction velocity and spectral parameters: influence of force and time. *American Physiological Society*, 58(5), pp.1428–1439.
- Budgett, R. (1998). Fatigue and underperformance in athletes: the overtraining syndrome. *British Journal of Sports Medicine*, 32(2),pp. 107–10.
- Canadian Society for Exercise Physiology. (2002). Physical Activity Readiness Questionnaire, Par-Q and You. 2002, (revised).
- Chang, K.-M., Liu, S.-H., & Wu, X.-H. (2012). A wireless sEMG recording system and its application to muscle fatigue detection. *Sensors*, 12(1), pp.489–499.
- Cheney, P. D., Kenton, J. D., Thompson, R. W., McKiernan, B. J., Lininger, R. E., & Trank, J. W. (1998). A low-cost, multi-channel, EMG signal processing amplifier. *Journal of Neuroscience Methods*, 79(1), 123–127.
- Chesler, N. C., & Durfee, W. K. (1997). Surface EMG as a fatigue indicator during FES-induced isometric muscle contractions. *Journal of Electromyography and Kinesiology : Official Journal of the International Society of Electrophysiological Kinesiology*, 7(1), pp.27–37.
- Chowdhury, R., Reaz, M. B. I., & Islam, M. T. (2012). Wavelet transform to recognize muscle fatigue. In 2012 Third Asian Himalayas International Conference on Internet (pp. 1–5).
- Chowdhury, S. K., & Virginia, W. (2012). Discrete Wavelet Transform Analysis of Surface Electromyography for the Objective Assessment of Neck and Shoulder Muscle Fatigue. (Unpublished Dissertation), West Virginia University, West Virginia.
- Chumanov, E., & Heiderscheit, B. (2011). EMG Activity Across Gait and Incline: The Impact of Muscular Activity on Human Morphology. *Journal of Physiology and Anthropology*, 143(4), pp.601–611.

- Cifrek, M., Medved, V., Tonković, S., & Ostojić, S. (2009). Surface EMG based muscle fatigue evaluation in biomechanics. *Clinical Biomechanics*, 24(4),pp. 327–340.
- Cobb, S., & Forbes, A. (1923). Electromyographic Studies of Muscular Fatigue in Man. America Journal of Physiology, 65(2), pp. 234–251.
- Davis, M. P., Walsh, D., & Edin, F. (2010). Mechanisms of Fatigue. *The Journal of Supportive Oncology*, 8(4), pp.164–174.
- Day, S. (2002). Important Factors in Surface EMG Measurement. Bortec Biomedical. Retrieved 1 January 2014 from www.bortec.ca
- De Luca, C. J. (1984). Myoelectrical manifestations of localized muscular fatigue in humans. *Critical Reviews in Biomedical Engineering*, 11(4),pp. 251–279.
- De Luca, C. J., Gilmore, L. D., Kuznetsov, M., & Roy, S. H. (2010). Filtering the surface EMG signal: Movement artifact and baseline noise contamination. *Journal of Biomechanics*, 43(8), pp.1573–1579.
- De Luca, G. (2003). Fundamental Concepts in EMG Signal Acquisition, (March), pp. 1–31.(Unpublished Article)
- Delsys. (2012). Frequently Asked Questions in the field of electromyography., Retrieved 1 January 2014 from http://www.de;sys.com/KnowledgeCenter?FAQ.html
- Devices, A. (2011). Low Cost, Low Power Instrumentation Amplifier AD620,(Datasheet), pp.1–16.
- Dimitrov GV, Arabadzhiev TI, Mileva KN, Bowtell JL, Crichton N, D. N. (2006). Muscle fatigue during dynamic contractions assessed by new spectral indices. *Medicine and Science in Sports and Exercise*, 38(11), pp.1971–1979.
- Dimitrova, N. A., & Dimitrov, G. V. (2003). Interpretation of EMG changes with fatigue: facts, pitfalls, and fallacies. *Journal of Electromyography and Kinesiology*, 13(2003), pp.13–36.
- Dingwell, J. B., Joubert, J. E., Diefenthaeler, F., & Trinity, J. D. (2008). Changes in muscle activity and kinematics of highly trained cyclists during fatigue. *IEEE Transactions on Bio-Medical Engineering*, 55(11), pp.2666–2674.
- E.Huigen, Peper, A., & Grimbergen, C. A. (2002). Investigation into the origin of the noise of surface electrodes. *Medical & Biological Engineering & Computing*, 40(March), pp.332–338.

- E.J.Kupa, S.H.Roy, S.C.Kandarian, Luca, C. J. D. E. (1995). Effects of muscle fiber type and size on EMG median frequency and conduction velocity. *The American Journal of Sports Medicine*, *79*, pp. 23–32.
- Edwards, R. H. (1981). Human muscle function and fatigue. *Ciba Foundation Symposium*, 82, pp. 1–18.
- Edwards, W. B. (2016). Biomechanics and Physiology of Uphill and Downhill Running. *Sports Medicion*, (August), pp.1–16.
- Froelicher, V. F., Brammell, H., Davis, G., Noguera, I., Stewart, A., & Lancaster, M. C. (1974). A Comparison of the Reproducibility and Physiologic Response to Three Maximal Treadmill Exercise Protocols. *Chest Journal*, 65(1), pp.512–517.
- Fu, J., Chen, J., Shi, Y., & Li, Y. (2013). Design of a Low-Cost Wireless Surface EMG Acquisition System. In 6th Annual International IEEE EMBS Conference on Neural Engineering (pp. 6–8).
- Fulco, CS Rock, P. S. M. (1999). Slower Fatigue and Faster Recovery of the Adductor Pollicis Muscle in Women Matched for Strength with Men. *Acta Physiol Scand*, *167*(3), pp.233–9.
- Garcia, M. a C., Catunda, J. M. Y., Lemos, T., Oliveira, L. F., Imbiriba, L. a, & Souza, M. N. (2010). An alternative approach in muscle fatigue evaluation from the surface EMG signal. In *Annual International Conference of the IEEE Engineering in Medicine and Biology Society* (Vol. 2010, pp. 2419–2422).
- Gates, D. H. (2009). *The Role of Muscle Fatigue on Movement Timing and Stability during Repetitive Tasks*. University of Texas.
- Gavin, J. P., Myers, S. D., & Willems, M. E. T. (2015). Neuromuscular responses to mild-muscle damaging eccentric exercise in a low glycogen state. *Journal of Electromyography and Kinesiology*, 25, pp. 53–60.
- Gazendam, M. G. J., & Hof, A. L. (2007a). Averaged EMG profiles in jogging and running at different speeds. *Gait & Posture*, 25(4),pp. 604–614.
- Ghapanchizadeh, H., & Ahmad, S. A. (2014). Developing Multichannel Surface EMG Acquisition System by Using Instrument OpAmp INA2141. In 2014 IEEE Region 10 Symposium (pp. 258–263).
- Gonzalez-Izal, M., Falla, D., Izquierdo, M., & Farina, D. (2010). Predicting force loss during dynamic fatiguing exercises from non-linear mapping of features of the surface electromyogram. *Journal of Neuroscience Methods*, 190(2), pp.271–278.
- González-Izal, M., Malanda, a, Navarro-Amézqueta, I., Gorostiaga, E. M., Mallor, F., Ibañez, J., & Izquierdo, M. (2010). EMG spectral indices and muscle power fatigue during dynamic contractions. *Journal of Electromyography and*

*Kinesiology : Official Journal of the International Society of Electrophysiological Kinesiology, 20*(2), pp.233–240.

- González-Izal, M., Malanda, A., Gorostiaga, E., & Izquierdo, M. (2012). Review Electromyographic models to assess muscle fatigue. *Journal of Electromyography and Kinesiology : Official Journal of the International Society of Electrophysiological Kinesiology*, 22(4), 501–512.
- González-Izal, M., Rodríguez-Carreño, I., Malanda, A., Mallor-Giménez, F., Navarro-Amézqueta, I., Gorostiaga, E. M., & Izquierdo, M. (2010). sEMG wavelet-based indices predicts muscle power loss during dynamic contractions. *Journal of Electromyography and Kinesiology*, 20(6), pp.1097–106.
- Greenwood, M. (2008). Aspects of Overtraining. In *Essentials of Sports Nutrition and Supplements* (pp. 121–142).
- Haddad, M., Chaouachi, A., Wong, D. P., Castagna, C., Hambli, M., Hue, O., & Chamari, K. (2013). Influence of fatigue, stress, muscle soreness and sleep on perceived exertion during submaximal effort. *Physiology & Behavior*, *119*(2013), pp.185–189.
- Haddad, O. (2011). *Development and validation of a fatigue-modified*, *EMG-assisted biomechanical model of the lumbar region*. Iowa State University.
- Halson, S. L., & Jeukendrup, A. E. (2004). Does Overtraining Exist? An Analysis of Overreaching and Overtraining Research. Sports Medicine (Auckland, N.Z.), 34(14), pp.967–981.
- Hassan, M. Z. C., Member, P. I. K.-I., Kamaruddin, N. A., & Ishak, N. A. (2014). Derivation of Simple Muscle Fatigue Index for Biceps Muscle Based on Surface Electromyography Temporal Characteristics. In 2014 IEEE Conference on Biomedical Engineering and Sciences (pp. 8–10).
- Hedayatpour, N., Falla, D., & Farina, D. (2008). Effect of delayed-onset muscle soreness on muscle recovery after a fatiguing isometric contraction. *Scandinavian Journal of Medicine & Science in Sports*, 20(1), pp.1–9.
- Hermie J.Hermens; Bart Freriks, Roberto Merletti; Dick Stegeman; Joleen Blok, G. R.
  C. D.-K. G. H. (1999). European Recommendations for Surface ElectroMyoGraphy. Roessingh Research and Development.
- Hernandez, C., Estrada, E., Garcia, L., Sierra, G., & Nazeran, H. (2010). Traditional SEMG Fatigue Indicators Applied to a Real-world Sport Functional Activity: Roundhouse Kick. In *Electronics, Communications and Computer* (pp. 154– 158).
- Hong-Chun, Y., Du-Ming, W., & Jian, W. (2005). Linear and non-linear features of surface EMG during fatigue and recovery period. *Conference Proceedings : ...*

Annual International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering in Medicine and Biology Society. Conference, 6, pp.5804–5807.

- Howwit, S. (2004). Overtraining Syndrome. *Bulletin de l'Académie Nationale de Médecine*, 188(6), pp.923–30;
- Hug, M., Vogt, M., Ventura, N., & Hoppeler, H. (2003). Training modalities: overreaching and over-training in athletes, including a study of the role of hormones. *Best Practice & Research Clinical Endocrinology and Metabolism*, 17(2), pp.191–209.
- Hunter, I., Seeley, M. K., Hopkins, J. T., Carr, C., & Franson, J. J. (2014). EMG activity during positive-pressure treadmill running. *Journal of Electromyography* and Kinesiology: Official Journal of the International Society of Electrophysiological Kinesiology, 24(3), pp. 348–352.
- Hunter, S. K., Duchateau, J., & Enoka, R. M. (2004). Muscle fatigue and the mechanisms of task failure. *Exercise and Sport Sciences Reviews*, 32(2),pp. 44– 49.
- Ishii, H., & N, Yy. N. (2013). Effect of Lactate Accumulation during Exercise-induced Muscle Fatigue on the Sensorimotor Cortex. Jurnal of Physical Therapy Science, 24, pp.1637–1642.
- Izquierdo, M., González-Izal, M., Navarro-Amezqueta, I., Calbet, J. a L., Ibañez, J., Malanda, A., ... Gorostiaga, E. M. (2011). Effects of strength training on muscle fatigue mapping from surface EMG and blood metabolites. *Medicine and Science in Sports and Exercise*, 43(2),pp.303–11.
- Jamaluddin, N. F. (2012). *Development of Two Electrodes System for Wireless* ECG/FECG Monitoring. (Unpublished Dissertation), Universiti Kuala Lumpur.
- Jeukendrup, A. E. (2010). Overtraining: how to monitor and how to prevent? In *Sport Nutrition Conference* (pp. 12–15).
- Joachim H.Nagel, Bronzino, E. J. D. (2000). "Biopotential Amplifiers." The Biomedical Engineering Handbook: Second Edition. Boca Raton: CRC Press LLC.
- Josef Finsterer. (2012). Biomarkers of peripheral muscle fatigue during exercise. *BMC Musculoskeletal Disorders*, *13*(218), pp.1–13.
- Kasman, G., & Wolf, S. (2002). Surface EMG made easy: A Beginner's Guide for Rehabilitation Clinicians. Noraxon.

- Kattla, S., & Lowery, M. M. (2010). Fatigue related changes in electromyographic coherence between synergistic hand muscles. *Experimental Brain Research*. *Experimentelle Hirnforschung. Expérimentation Cérébrale*, 202(1), pp.89–99.
- Kent-Braun, J. a. (1999). Central and peripheral contributions to muscle fatigue in humans during sustained maximal effort. *European Journal of Applied Physiology and Occupational Physiology*, 80(1), pp.57–63.
- Kirkby, R. (2000). The Collapsed Endurance Athlete: A Practical Approach. *Official Journal of the South African Academy of Family Physicians*, 22(3), pp.40–44.
- Kobayashi, H. (2010). Development of an EMG acquisition system of universal biosignal interface for high-quality life, pp.539–542.
- Konrad, P. (2005). *The ABC of EMG. Noraxon Inc, USA* (Vol. 1.0). Retrieved 1 January 2014 from http://demotu.org/aulas/controle/ABCofEMG.pdf
- Kreher, J. B., & Schwartz, J. B. (2012). Overtraining Syndrome: A Practical Guide. Sports Health, 4(2), pp.128–38.
- Kumar, D.K., V P Singh, I Cosic, S L Giudice, J. H. (2005). Strategies to identify changes in SEMG due to muscle fatigue during cycling. In 2005 IEEE Engineering in Medicine and Biology 27th Annual Conference (pp. 6683–6686).
- Kwatny, E., Thomas, D. H., & Kwatny, H. G. (1970). An application of signal processing techniques to the study of myoelectric signals. *IEEE Transactions on Bio-Medical Engineering*, 17(4), pp.303–313.
- Landreneau, L. L., Watts, K., Heitzman, J. E., & Childers, W. L. (2014). Lower Limb Muscle Activity during Forefoot and Rearfoot Strike Running Technique. *The International Journal of Sports Physical Therapy*, 9(7),pp. 888–897.
- Liu, C., & Lin, K. (2007). Estimation of VO2 Max : A Comparative Analysis of Postexercise Heart Rate and Physical Fitness Index from 3-minute Step Test. *Journal* of Exercise Science and Fitness, 5(2), pp.118–123.
- Liu, S., & Chang, K. (2014). The Progression of Muscle Fatigue During Exercise Estimation With the Aid of High-Frequency. *IEEE Journal of Biomedical and Health Informatics*, *18*(5), pp.1647–1658.
- Loss, J. F., Melo, D. O., Torre, M. La, Pasini, M., Arau, L., Oliveira, L. N. De, & Pinto, L. (2008). Comparing the lactate and EMG thresholds of recreational cyclists during incremental pedaling exercise. *Canadian Journal Physiology Pharmacology*, 86(2008), pp.272–278.
- Lowery, M. M., & O'Malley, M. J. (2003). Analysis and simulation of changes in EMG amplitude during high-level fatiguing contractions. *IEEE Transactions on Bio-Medical Engineering*, 50(9), pp.1052–1062.

- Luca, C. J. De. (2002). Surface Electromyography: Detection and Recording. *DelSys Incorporated*, pp.1–10.
- Maged S, A., Asnor J, I., Siti A, A., & Mohd K, H. (2014). Multichannel EMG Data Acquisition System : Design and Temporal Analysis during Human Ankle joint Movements. In 2014 IEEE Conference on Biomedical Engineering and Sciences (pp. 8–10).
- Malanda, A., & Izquierdo, M. (2009). New Wavelet Indices to Assess Muscle Fatigue during Dynamic Contractions. World Academy of Science, Engineering and Technology, 3(2009-07-21), pp. 456–461.
- Mark A.Osborne, D. A. S. (2006). Muscle glycogen reduction in man: Relationship between surface EMG activity and oxygen uptake kinetics during heavy exercise. *The Physiological Soceity*, *91*(1), pp.179–189.
- Marri, K., & Swaminathan, R. (2015). Classification of Muscle Fatigue using Surface Electromyography Signals and Multifractals. In 2015 12th International Conference on Fuzzy Systems and Knowledge Discovery (FSKD) (pp. 669–674).
- Massó, N., Rey, F., Romero, D., Gual, G., Costa, L., & Germán, A. (2010). Surface electromyography applications in the sport, 45(165),pp. 121–130.
- Maughan, RJ Harmon, M Leiper, JB Sale, D Delma, A. (1986). Endurance Capacity of Untrained Males and Females in Isometric and Dynamic Muscular Contraction. *European Jurnal of Applied Physiology and Occupation Physiology*, 55(4), pp.395–400.
- McKune, A., Semple, S., & Peters-Futre, E. (2012). Acute Exercise-Induced Muscle Injury. *Biology of Sport*, 29(1), pp. 3–10.
- Medved, V., & Cifrek, M. (2011). Kinesiological Electromyography. *Biomechanics in Applications*, pp.349–366.
- Merletti, A. R., & Torino, P. (1999). Standards for Reporting EMG Data. *International Society of Electrophysiology and Kinesiology*.
- Mesa, I., Rubio, A., Tubia, I., De No, J., & Diaz, J. (2014). Channel and feature selection for a surface electromyographic pattern recognition task. *Expert Systems with Applications*, *41*(11), pp.5190–5200.
- Mesin, L., Merletti, R., & Rainoldi, a. (2009). Surface EMG: the issue of electrode location. Journal of Electromyography and Kinesiology : Official Journal of the International Society of Electrophysiological Kinesiology, 19(5), pp.719–26.
- Miller, A. E. J., Macdougall, J. D., Tarnopolsky, M. A., & Sale, D. G. (1993). Gender Differences in Strenght and Muscle Fiber Characteristics. *European Journal of Applied Physiology*, 66, pp. 254–262.

Misiti, M., & Poggi, J. (2014). Wavelet Toolbox <sup>TM</sup> User 's Guide R 2014 b.

- Mizumura, K. (2009). Peripheral mechanism of muscle pain: An update. *Current* Anaesthesia & Critical Care, 20(4),pp. 183–187.
- Mohd Ishak, N. A., Khalid, P. I., Mahmood, N. H., & Harun, M. (2016). Determination of Epoch Length and Regression Model for 15-second Segment of SEMG Signal used in Joiny Analysis of Spectrum and Amplitude. *Jurnal Teknologi*, *5*(78),pp. 7–13.
- Moritani, T., & Muro, M. (1987). Applied Physiology Motor unit activity and surface electromyogram power spectrum during increasing force of contraction. *European Journal of Applied Physiology and Accupational Physiology*, *56*(1), pp.260–265.
- Mortimer, J. T., Magnusson, R., & Petersen, I. (1970). Conduction Velocity in Ischemic Muscle: Effect on EMG Frequency Spectrum. America Journal of Physiology-Legacy Content, 219(5), pp.1324–1329.
- Murley, G. S., Menz, H. B., & Landorf, K. B. (2009). Foot posture influences the electromyographic activity of selected lower limb muscles during gait. *Journal of Foot and Ankle Research*, *2*, 35.
- Myrick, K. M., & Aprn, D. N. P. (2015). Syndrome in Athletes. *TJNP: The Journal for Nurse Practitioners*, *11*(10),pp.1018–1022.
- Nason, G. P., & Silverman, B. W. (1995). The Stationary Wavelet Transform and some Statistical Applications. In A. Antoniadis & G. Oppenheim (Eds.), *Wavelet and Statistics* (pp. 1–19).
- Nederhof, E., Lemmink, K. A. P. M., Visscher, C., Meeusen, R., & Mulder, T. (2006). Possibly a New Marker for Overtraining Syndrome. Sports Medicine, 36(10), pp.817-828.
- Neuman, M. R. (2008). Biopotential amplifiers. In *Medical Instrumentation* Application and Design (4th ed., pp. 241–292). John Wiley & Sons.
- Nie, H., Arendt-nielsen, L., Kawczynski, A., & Madeleine, P. (2007). Gender effects on trapezius surface EMG during delayed onset muscle soreness due to eccentric shoulder exercise. *Journal of Electromyography and Kinesiology*, *17*, pp.401–409.
- Nur, N. M., & Dawal, S. Z. M. (2009). Upper limb and lower back muscle activity during prolonged sitting. 2009 IEEE Toronto International Conference Science and Technology for Humanity (TIC-STH), pp. 234–238.

- Oka, H. (1996). Estimation of muscle fatigue by using EMG and muscle stiffness. Proceedings of 18th Annual International Conference of the IEEE Engineering in Medicine and Biology Society, 4, pp.1449–1450.
- Paluska, S. a. (2005). An overview of hip injuries in running. *Sports Medicine* (Auckland, N.Z.), 35(11), pp. 991–1014.
- Park, S., Lee, S., Kang, H. C., & Kim, S. (2012). EMG Analysis of Lower Limb Muscle Activation Pattern During Pedaling: Experiments and Computer Simulations, 13(4), pp.601–608.
- Pearce, P. Z. (2002). A practical approach to the overtraining syndrome. *Current Sports Medicine Reports*, 1(3),pp. 179–183.
- Petrofsky, J. S. (1979). Applied Physiology During Exercise on the Bicycle Ergometer. *European Journal of Applied Physiology*, 15(41), pp. 1–15.
- Phinyomark, A., Limsakul, C., & Phukpattaranont, P. (2009a). A Comparative Study of Wavelet Denoising for Multifunction Myoelectric Control. In *International Conference on Computer ad Automation Engineering* (pp. 21–25).
- Phinyomark, A., Limsakul, C., & Phukpattaranont, P. (2009b). EMG Denoising Estimation Based on Adaptive Wavelet Thresholding for Multifunction Myoelectric Control. In 2009 Conference on Innovative Technologies in Intelligent Systems and Industrial Applications (CITISIA 2009) (pp. 171–176).
- Pioggia, G., Tartarisco, G., Ricci, G., Volpi, L., Siciliano, G., & Bonfiglio, S. (2010). A wearable pervasive platform for the intelligent monitoring of muscular fatigue. In 2010 10th International Conference on Intelligent Systems Design and Applications (pp. 132–135).
- Polar. (2013). RC3 GPS User Manual.
- Polikar, R. (2006). Pattern recognition. New Jersey: Wiley Encyclopedia of Biomedical Engineering.
- Poo, T. S., & Sundaraj, K. (2010). Design and Development of a Low Cost EMG Signal Acquisition System Using Surface EMG Electrode. In 2010 IEEE Asia Pacific Conference on Circuit and System (APCCAS) (pp. 24–27).
- Potvin, J. R., & Bent, L. R. (1997). A validation of techniques using surface EMG signals from dynamic contractions to quantify muscle fatigue during repetitive tasks. Journal of Electromyography and Kinesiology: Official Journal of the International Society of Electrophysiological Kinesiology, 7(2), pp.131–139.
- Purvis, D., Gonsalves, S., & Deuster, P. a. (2010). Physiological and psychological fatigue in extreme conditions: overtraining and elite athletes. *The Journal of Injury, Function, and Rehabilitation*, 2(5),pp. 442–450.

- Riillo, F., Quitadamo, L. R., Cavrini, F., Gruppioni, E., Pinto, C. a., Pastò, N. C., ... Saggio, G. (2014). Optimization of EMG-based hand gesture recognition: Supervised vs. unsupervised data preprocessing on healthy subjects and transradial amputees. *Biomedical Signal Processing and Control*, 14, pp.117– 125.
- Rumsey, D. (2010). *Statistics Essentials for Dummies* (First Edit.). Indiana: Wiley Publishing.
- S.C.Gandevia. (2001). Spinal and Supraspinal Factors in Human Muscle Fatigue. *The American Physiology Society*, *81*(4), pp.1725–1789.
- Sakurai, T., Toda, M., Sakurazawa, S., Akita, J., Kondo, K., & Nakamura, Y. (2010). Detection of Muscle Fatigue by the Surface Electromyogram and Its Application. In 9th International Conference on Computer and Information Science (pp. 43– 47).
- Sarillee, M., Hariharan, M., Anas, M. N., Omar, M. I., Aishah, M. N., Ck, Y., & Oung, Q. W. (2015). Classification of Muscle Fatigue Condition using. In 2015 IEEE International Conference on Control System, Computing and Engineering (pp. 27–29).
- Senefeld, J., Bement, M. K. H., & Hunter, S. (2013). Fatigue and Recovery from Dynamic Contractions in Men and Women Differ for Arm and Leg Muscles. *Muscle and Nerve*, 48(3), pp.436–439.
- Sepulveda, F., Colley, M., & Kattan, A. (2009). Classification of localized muscle fatigue with Genetic Programming on sEMG during isometric contraction. In 2009 Annual International Conference of the IEEE Engineering in Medicine and Biology Society (pp. 2633–2638).
- Sesboüé, B., & Guincestre, J.-Y. (2006). Muscular fatigue. Annales de Readaptation et de Medecine Physique: Revue Scientifique de La Societe Francaise de Reeducation Fonctionnelle de Readaptation et de Medecine Physique, 49(6), pp.257–264.
- Singh, V. P., Kumar, D. K., Djuwari, D., Polus, B., Fraser, S., Hawley, J., & Giudice, S. L. (2004). Strategies to identify muscle fatigue from SEMG during cycling. *Proceedings of the 2004 Intelligent Sensors, Sensor Networks and Information Processing Conference, 2004.*, pp.547–552.
- Sloniger, M. A., Cureton, K. J., Prior, B. M., Evans, E. M., Carrier, D. R., Anders, C., ... Batterham, M. (1997). Lower extremity muscle activation during horizontal and uphill running. *Journal of Applied Physiology*, 83, pp.2073–2079.
- Soares, F. A., Salomoni, S., Veneziano, W. H., Schwartz, F. P., & Carvalho, J. L. A. De. (2008). A Surface Electromyographic Analysis about The FatigueE on

Women Through The Menstrual Cycle. In 21 Congresso Brasileiro de Engenharia Biomedica (pp. 1675–1678).

- Sobahi, N. M. (2011). Denoising of EMG Signals Based on Wavelet Transform. In *Asian Transactions on Engineering* (Vol. 01, pp. 17–23).
- Stegeman, D., & Hermens, H. (2007). Standards for surface electromyography: The European project Surface EMG for non-invasive assessment of muscles (SENIAM),pp.108–112.
- Sterczala, A., & Nicol, Jx; Fry, A. (2015). The Relationship between Performance Decrements and Peak EMG Ampitude following Resistance Exercise. *International Journal of Exercise Science*, *11*(3), pp.63–68.
- Strakowski, J. a, & Jamil, T. (2006). Management of Common Running Injuries. Physical Medicine and Rehabilitation Clinics of North America, 17(3), pp.537– 552.
- Tenan, M. S., Blackburn, J. T., & Robert, G. (2016). Exercise-Induced Glycogen Reduction Increases Muscle Activity. *International Journal of Exercise Science*, 9(3), pp.336–346.
- Thakor, N. V. (1999). Biopotentials and Electrophysiology Measurement. In *Measurements, Instrumentations and Sensors Handbook* (p. 12). CRC Press LLC.
- Thomas, A. C., McLean, S. G., & Palmieri-Smith, R. M. (2010). Quadriceps and hamstrings fatigue alters hip and knee mechanics. *Journal of Applied Biomechanics*, 26(2), pp.159–70.
- Vahed, A. (2005). 3-Lead Wireless ECG. (Unpublished Report)
- Venugopal, G., Navaneethakrishna, M., & Ramakrishnan, S. (2014). Extraction and analysis of multiple time window features associated with muscle fatigue conditions using sEMG signals. *Expert Systems with Applications*, 41(6), pp.2652–2659.
- Vries, H. A. D. (1967). Quantitative Electromyographic Investigation of The Spasm Theory of Muscle Pain. Jurnal of Physical Medicin, 45, pp.119.
- Wang, L., Huang, Y., & Gong, M. (2011). Sensitivity and Stability Analysis of sEMG Indices in Evaluating Muscle Fatigue of Rectus Femoris Caused by All-out Cycling Exercise. In 4th International Congress on Image and Signal Processing (pp. 2779–2783).
- Yang, Z., Wu, Q., & Fu, S. (2014). Optik Spectral analysis of surface EMG based on empirical mode decomposition, 125, pp.7045–7052.

- Zakynthinos, S., & Roussos, C. (2005). Respiratory muscle fatigue. In *Physiologic Basis of Respiratory Disease* (pp. 289–306).
- Zhang, J., Lockhart, T., & Soangra, R. (2014). Classifying Lower Extremity Muscle Fatigue durig Walking using Machine Learning and Inertial Sensors. *Annual Biomedical Engineering*, 42(3),pp. 600–612.
- Zhang, Jian E.Lockhart, Thurmon Soangra, R. (2015). Classifying Lower Extremity Muscle Fatigue during Walking using Machine Learning and Inertial Sensors. *Annual Biomedical Engineering*, 42(3),pp. 600–612.

