



UNIVERSITI PUTRA MALAYSIA

***METABOLIC DISORDER INDEX AS PREDICTOR OF PERFORMANCE IN
ENDURANCE HORSES***

NUR ZUL IZZATI BT MOHD RAJDI

FPV 2018 1



**METABOLIC DISORDER INDEX AS PREDICTOR OF PERFORMANCE IN
ENDURANCE HORSES**

By

NUR ZUL IZZATI BT MOHD RAJDI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of
Master of Veterinary Science**

November 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATIONS

This thesis is dedicated to my beloved parents, Mohd Rajdi Idris and Nik Hasmah Daud. Thank you for unconditionally love and support.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Veterinary Science

METABOLIC DISORDER INDEX AS PREDICTOR OF PERFORMANCE IN ENDURANCE HORSES

By

NUR ZUL IZZATI BT MOHD RAJDI

November 2017

Chairman : Noraniza Mohd Adzahan, PhD
Faculty : Veterinary Medicine

Endurance is a type of equestrian sport that is recognized by the Fédération Equestre Internationale (FEI). It is a long-distance race ranging from 40 to 120 km and is divided into several loops whereby each loop covers from 20 to 35 km. The performance of endurance horses is usually evaluated on their ability to complete the race in the shortest period. The rate of elimination of horses in the races are still high, ranging from 10% to 60%. Endurance horses are usually eliminated from the race due to lameness, metabolic disorders, and other causes which include failure to reach at the veterinary inspection within the stipulated period, optional withdraws, and concerns over the well-being of the horse. A prediction of horses' performance before the race can help reduce the rate of elimination as well as care for their well-being by preventing the participation of poor performance horses in the competition. Therefore, the purpose of the present study is to determine the current completion and elimination rate of endurance races in Malaysia. A total of 194 endurance horses who participated in 12 endurance races organized in several states in Malaysia from May 2015 until December 2016 are involved. Out of the 194 horses, 188 are from Terengganu while 6 are from Kelantan. From the total of 194 horses, 126 are able to complete the race while 68 of them are eliminated from the race by official veterinarians. In the current study, 35.05% of the horses are eliminated and 64.95% of them are able to complete the race successfully. Among the eliminated horses, 54.41%, 29.41% and 16.18% of them are due to lameness, metabolic disorders and other causes respectively. Most of the eliminated horses competed in the 80 km race while those who successfully completed the race competed in the 40 km race. The average speed for good, moderate, and poor performance horses is 15.72 ± 2.9 , 12.19 ± 2.1 , and 13.91 ± 2.6 km h⁻¹ respectively. In the 40, 80, and 120 km races, the average speed is 12.65 ± 1.8 , 15.99 ± 3.5 , and 12.86 ± 0.2 km h⁻¹

respectively showing that the horses performed fastest in the 80 km races. This data could be of assistance to the veterinarians in monitoring the trend of endurance races in Malaysia and improve the current management practices to reduce the rate of elimination in subsequent races. However, the elimination of the horses is unpredictable because there is no specific method to predict their performance before the race. In 2017, Adamu et al. came out with the Metabolic Disorder Index (MDI) which claims can be used in predicting a horse's performance before the race. The MDI is defined as a product of packed cell volume (PCV), creatine kinase (CK), and interleukin-6 (IL-6) divided by the product of chloride (Cl⁻) and glutathione reductase (GR) multiplied by 100. A prediction is made based on the MDI value. Horses who have MDI value of >5.5 is predicted to be of good performance while those who have MDI value of <5.5 is predicted to be of poor performance. However, this index has not been tested on the field under controlled conditions. Other purpose of this study is to evaluate the MDI's parameters of good and poor performance endurance horses before the race and to validate the MDI as a predictor for horses' performance in endurance races. Out of the 194 horses that are involved in the study, only 54 of them are selected for blood sampling to test the validity of MDI. Blood samples are taken a day prior to the event to determine the packed cell volume (PCV), chloride (Cl⁻), and Interleukin-6 (IL-6) concentrations as well as creatine kinase (CK) and glutathione reductase (GR) activities. These parameters are used to determine the MDI in endurance horses at rest one day before the competition. Poor performance horses have significantly ($p < 0.05$) higher serum CK concentrations at 216.0 ± 65 U L⁻¹ compared to good performance horses at 150.7 ± 29 U L⁻¹. The serum Cl⁻ concentration in poor performance horses is lower at 99.8 ± 5.6 mmol L⁻¹ compared to good performance horses at 103.7 ± 3.3 mmol L⁻¹. The study shows that MDI has a high ability to identify poor performance horses with approximately 90% sensitivity. However, in terms of accuracy, it is more accurate in predicting good performance horses with a negative predictive value of 90.91%.

Keywords: horses, endurance, metabolic, performance.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains Veterinar

INDEKS GANGGUAN METABOLIK (MDI) SEBAGAI PERAMAL PRESTASI PADA KUDA LASAK

Oleh

NUR ZUL IZZATI BT MOHD RAJDI

November 2017

Pengerusi : Noraniza Mohd Adzahan, PhD
Fakulti : Perubatan Veterinar

Sukan kuda lasak merupakan salah satu sukan berkuda yang diiktiraf oleh *Federation Equestre Internationale* (FEI). Ianya merupakan perlumbaan jarak jauh dalam lingkungan 40 hingga 120km dan jarak tersebut terbahagi kepada beberapa bahagian yang meliputi jarak antara 20 ke 35km. Prestasi kuda lasak biasanya dinilai berdasarkan keupayaan untuk menamatkan perlumbaan dalam masa tersingkat. Kadar penyingkiran kuda lasak dalam perlumbaan adalah tinggi iaitu pada kadar 10 hingga 60%. Kuda lasak biasanya disingkir daripada perlumbaan kerana ketempangan, penyakit metabolisme atau sebab lain, termasuk kegagalan untuk sampai di tempat pemeriksaan veterinar dalam tempoh yang ditetapkan, pengunduran pilihan dan kebimbangan atas kebajikan kuda. Ramalan prestasi kuda sebelum perlumbaan boleh membantu untuk mengurangkan kadar penyingkiran dan menjaga kebajikan kuda dengan cara menghalang kuda berprestasi lemah daripada menyertai pertandingan. Oleh itu, tujuan kajian ini adalah untuk menentukan kadar pelengkapan dan penyingkiran kuda lasak di Malaysia. Seratus sembilan puluh empat ekor kuda lasak yang telah menyertai 12 perlumbaan kuda lasak yang dianjurkan di beberapa buah negeri di Malaysia bermula dari Mei 2015 sehingga Disember 2016 telah terlibat dalam kajian ini. Seratus lapan puluh lapan ekor kuda lasak daripada 194 ekor tersebut adalah dari Terengganu manakala 6 ekor adalah dari Kelantan. Berdasarkan daripada jumlah tersebut juga, 126 ekor berjaya menamatkan perlumbaan manakala 68 ekor telah disingkir daripada perlumbaan oleh veterinar yang bertugas. Kajian ini menunjukkan 35.05% kuda telah tersingkir manakala 64.95% telah berjaya menamatkan perlumbaan. Kuda yang tersingkir disebabkan oleh masalah ketempangan, gangguan metabolisme dan sebab-sebab lain masing-masing adalah 54.41%, 29.41% dan 16.18%. Kebanyakan daripada kuda yang tersingkir adalah terdiri daripada kuda yang bertanding

dalam jarak 80km sementara kuda yang berjaya menamatkan perlumbaan adalah bertanding dalam jarak 40km. Purata kelajuan untuk prestasi baik, sederhana dan lemah masing-masing adalah 15.72 ± 2.9 , 12.19 ± 2.1 dan 13.91 ± 2.6 km h⁻¹. Untuk perlumbaan 40, 80 dan 120km, purata kelajuan masing-masing adalah 12.65 ± 1.8 , 15.99 ± 3.5 dan 12.86 ± 0.2 km h⁻¹. Data ini menunjukkan bahawa kuda yang terlibat di dalam perlumbaan 80km mempunyai kelajuan yang terpanas. Data ini mampu membantu para veterinar dalam memerhati corak perlumbaan kuda lasak sekaligus meningkatkan amalan pengurusan semasa dalam mengurangkan kadar penyingkiran dalam setiap perlumbaan. Walau bagaimanapun, penyingkiran dalam kalangan kuda lasak adalah tidak mampu dijangka kerana tiadanya kaedah tertentu bagi meramal prestasi kuda sebelum perlumbaan. Pada tahun 2017, Adamu et al. memperkenalkan indeks gangguan metabolisme (MDI) yang dikatakan dapat digunakan untuk meramal prestasi kuda sebelum perlumbaan. MDI ditakrifkan sebagai hasil isipadu sel padat (PCV), aktiviti kreatin kinase (CK) dan interleukin-6 (IL-6) dibahagi dengan kepekatan klorida (Cl⁻) dan glutathione reductase (GR) yang kemudiannya dikalikan dengan 100. Ramalan tentang prestasi kuda akan dibuat berdasarkan nilai MDI. Kuda yang mempunyai nilai MDI >5.5 diramal sebagai kuda yang berprestasi lemah manakala kuda yang mempunyai nilai MDI <5.5 diramal sebagai kuda yang berprestasi baik. Namun, indeks ini masih belum diuji di bawah keadaan yang terkawal. Jadi, tujuan kajian ini dijalankan adalah untuk menilai parameter-parameter MDI bagi kuda berprestasi baik dan lemah sebelum perlumbaan dan mengkaji kebolehpercayaan indeks gangguan metabolisme (MDI) dalam meramalkan prestasi kuda dalam perlumbaan lasak. Daripada 194 ekor kuda yang terlibat dalam kajian terdahulu, hanya 54 daripada mereka dipilih untuk pensampelan darah untuk menguji kesahihan MDI. Sampel darah diambil sehari sebelum acara perlumbaan untuk menentukan isipadu sel padat (PCV), kepekatan klorida (Cl⁻) dan interleukin-6 (IL-6), dan aktiviti kreatin kinase (CK) dan glutathione reductase (GR). Semua parameter ini diguna untuk penentuan MDI dalam kuda lasak sehari sebelum pertandingan. Kuda berprestasi lemah secara tererti ($p < 0.05$) menunjukkan kepekatan CK lebih tinggi pada 216.0 ± 65 U L⁻¹ berbanding kuda berprestasi baik pada 150.7 ± 29 U L⁻¹. Kepekatan Cl⁻ serum dalam kuda berprestasi lemah pada 99.8 ± 5.6 mmol L⁻¹ lebih rendah daripada kuda berprestasi baik pada 103.7 ± 3.3 mmol L⁻¹. Kajian ini menunjukkan MDI mempunyai keupayaan yang lebih tinggi untuk mengenalpasti kuda yang lemah dengan sensitiviti kira-kira 90%. Walau bagaimanapun, dari segi ketepatan, MDI lebih tepat dalam meramal kuda prestasi yang baik dengan nilai ramalan negatif pada 90.91%.

Katakunci: kuda, lasak, metabolisme, prestasi.

ACKNOWLEDGEMENTS

First and foremost, I would like to express my sincere appreciation and gratitude to my supervisor, Dr. Noraniza Mohd Adzahan for her guidance during my research. Her guidance and ideas have been precious for the development of this thesis content. Additionally, I am indebted to Prof. Dr Rasedee Abdullah, who has been a constant source of encouragement during this thesis project. His ideas and suggestions have improved the content of my thesis. I also would like to thank my committee member, Assoc. Prof. Datuk Dr. Bashir Ahmad for his interest in my work.

I also would like to thank all the crew of STET and RTES who constantly welcome me every time I visited the stable for my sampling. Specials thanks goes to Dr Hamidah Helman and Dr Khairul Anuar Muhammad from the Department of Veterinary Services, Terengganu for their cooperation during the sampling. I will forever remember and thankful for the hospitality and help during the sampling period. Particular thanks goes to En Mohamad Halmi Othman, En Mohd Azri Roslan and all staffs of Hematology and Clinical Biochemistry and Bacteriology Laboratory, Faculty of Veterinary Medicine , and Equine Medicine Unit, University Veterinary Hospital (UVH) , for their help, guidance and assistance during my sampling and sample analysis.

My specials thank goes to Dr. Nor Laili Abdul Rachman from Veterinary Research Institute (VRI), Ipoh for her guidance and tremendous help not only for my lab work, but also for the mental support during my stressful moments.

My deep appreciation goes to my friends who shared this beautiful moment during our postgraduate life. The names that I would like to mention here are Amirah, Syahirah, Shyra, Lukman, Farah and Farhan. Thanks for the moral support and great moments that we shared together.

Finally, my deepest gratitude goes to my family for the love and financial support during my study and my life!

I certify that a Thesis Examination Committee has met on 23 November 2017 to conduct the final examination of Nur Zul Izzati bt Mohd Rajdi on her thesis entitled "Metabolic Disorder Index as Predictor of Performance in Endurance Horses" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Veterinary Science.

Members of the Thesis Examination Committee were as follows:

Mohd Hezme bin Mohd Noor, PhD


Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Hazilawati binti Hamzah, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Mohd Azam Khan bin Goriman Khan, PhD

Senior Lecturer
Universiti Malaysia Kelantan
Malaysia
(External Examiner)



NOR AINI AB. SHUKOR, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 March 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the Master of Veterinary Science. The members of the Supervisory Committee were as follows:

Noraniza Mohd Adzahan, PhD

Senior Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Rasedee Abdullah, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Bashir Ahmad B Fateh Mohamed, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No: Nur Zul Izzati Bt Mohd Rajdi, GS42407

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Dr. Noraniza Mohd Adzahan

Signature: _____
Name of Member
of Supervisory
Committee: Professor Dr. Rasedee Abdullah

Signature: _____
Name of Member
of Supervisory
Committee: Professor Dr. Bashir Ahmad B Fateh Mohamed

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xv
CHAPTER	
1 INTRODUCTION	1
1.1 Equestrian Sports	1
1.2 Endurance Sports	1
1.3 Hypothesis	2
1.4 Problem statements	2
1.5 Objectives	2
2 LITERATURE REVIEW	3
2.1 Endurance Sport	3
2.2 Metabolic Crisis	3
2.3 Prevalence and Risk Factor of Elimination of Endurance Horses	3
2.4 Metabolic Disorder Parameters in Endurance Horses	4
2.4.1 Mucous membrane	4
2.4.2 Capillary Refill Time (CRT)	4
2.4.3 Hydration status	5
2.4.4 Heart rate	5
2.4.5 Cardiac Recovery Index (CRI)	5
2.4.6 Intestinal activity	6
2.4.7 Soreness, laceration and wounds	6
2.4.8 Gait	6
2.5 Factors Affecting Haematological and Biochemical Parameters	6
2.5.1 Sample handling	6
2.5.2 Excitement	7
2.5.3 Feeding	7
2.5.4 Gender	7
2.5.5 Age	7
2.5.6 Breed	8
2.5.7 Exercise	8
2.5.8 Training and over-training	8
2.5.9 Sedatives and tranquilizers	8

2.6	Common Metabolic Disorders in Endurance Horses	9
2.6.1	Exhausted horse syndrome	9
2.6.2	Synchronous diaphragmatic flutter	9
2.6.3	Exertional myopathy	9
2.6.4	Colic	10
2.6.5	Laminitis	10
2.7	Common Musculoskeletal Disorders in Endurance Horses	10
2.7.1	Desmitis	10
2.7.2	Tendonitis	10
2.7.3	Osteoarthritis	10
2.7.4	Fracture	11
2.7.5	Back and wither injury	11
2.8	Haematology and Serum Biochemistry of Endurance Horse	11
2.8.1	Haematological responses of horses in an endurance race	11
2.8.2	Serum biochemical responses of horses in an endurance race	12
2.8.2.1	Total protein	12
2.8.2.2	Lactate	12
2.8.2.3	Creatine Kinase	12
2.8.2.4	Glucose	12
2.8.2.5	Electrolytes	13
2.9	Role of antioxidant in endurance horses	13
2.9.1	Glutathione Reductase (GR)	13
2.10	Roles cytokine in endurance horses	13
2.10.1	Interleukin-6 (IL-6)	13
2.11	Oxidative stress in endurance horses	14
2.12	Metabolic Disorder Index (MDI)	14
3	MATERIALS AND METHODS	15
3.1	Horse	15
3.2	Horse rider	15
3.3	Race track	15
3.4	Endurance event	16
3.5	Physical Examination	19
3.6	Performance Indicators	19
3.7	Data Collection	23
3.8	Blood Sample	23
3.9	Metabolic Disorder Index (MDI)	23
3.10	Sensitivity and Predictive value	24
3.11	Data analysis	24
4	RESULTS	25
4.1	Rates and Causes of Elimination in Malaysian Endurance Horses	25
4.2	Average speed of horses competing in endurance races	31
4.3	PCV, Cl ⁻ , CK, IL-6 & GR Concentrations in Good and Poor Performance Endurance Horses	31

4.4	Metabolic Disorder Index (MDI)	38
4.5	Sensitivity and specificity of MDI	38
5	DISCUSSION	41
5.1	The Rate of Completion and Elimination of Endurance Horses in Malaysia	41
5.2	Pre-Ride Stage Blood Biochemical Parameters for Good and Poor Performance Horses	43
5.3	Metabolic Disorder Index (MDI)	45
6	GENERAL CONCLUSION AND RECOMMENDATION	48
6.1	General conclusion	48
6.2	Recommendation	48
	REFERENCES	50
	APPENDICES	59
	BIODATA OF STUDENT	84
	LIST OF PUBLICATIONS	85

LIST OF TABLES

Table		Page
3.1	The participation of endurance horses in endurance races between 2015 and 2016	18
3.2	Prediction of horse performance based on MDI values (Adamu, 2014)	24
4.1	Rate of elimination of endurance horses from 40, 80 and 120km race distance in 12 races conducted in Malaysia from 2015 until 2016	27
4.2	Rate of elimination of endurance horses from 40km race distance in 12 races conducted in Malaysia from 2015 until 2016	28
4.3	Rate of elimination of endurance horses from 80km race distance in 12 races conducted in Malaysia from 2015 until 2016	30
4.4	Rate of elimination of endurance horses from 120km race distance in 12 races conducted in Malaysia from 2015 until 2016	31
4.5	Average speed of horses competing in endurance races	33
4.6	Blood parameters of good and poor performance endurance horses	35
4.7	Blood parameters of good and poor performance endurance horses in 40km races	36
4.8	Blood parameters of good and poor performance endurance horses in 80km races	37
4.9	Blood parameters of good and poor performance endurance horses in 120km races	38
4.10	The MDI findings on endurance horses	40
4.11	Sensitivity, Specificity, Positive and Negative predictive values of MDI in endurance horses	41

LIST OF FIGURES

Figure		Page
3.1	The tracks used in the endurance race: (A) At the beach (B) At the rubber plantation	17
3.2	Physical examinations are conducted at the veterinary inspection (A) The track used to examine the horse before, during, and after the race (B) The official veterinarians examined the horses before the race	20
3.3	The findings of the physical examination are recorded in the horse's log book. (A) Example of log book for the horse that successfully completed the race (B) Example of log book for the horse that is eliminated from the race due to metabolic problems	21

LIST OF ABBREVIATIONS

%	Percentage
°C	Degree Celsius
ALP	Alkaline phosphatase
AST	Aspartate aminotransferase
Ca	Calcium
CEI	Concours de Raid d'Endurance International
CEIO	Concours de Raid d'Endurance International Official
CK	Creatine kinase
Cl ⁻	Chloride
CRT	Capillary refill time
CRI	Cardiac recovery index
EDTA	Ethylenediaminetetraacetic acid
FEI	<i>Fédération Équestre Internationale</i>
G	Gauge
GGT	Gamma-glutamyl transferase
GLU	Glucose
GR	Glutathione reductase
Hb	Hemoglobin
IL-6	Interkeukin-6
K	Potassium
Kg	Kilogram
km	Kilometer
km h ⁻¹	kilometer per hour
L	Liter
L/L	Liter per liter
Lac	Lactate

M	Meter
MCH	Mean corpuscular hemoglobin
MCHC	Mean corpuscular hemoglobin concentration
MDI	Metabolic disorder index
MCV	Mean corpuscular volume
Mg	Magnesium
Na	Sodium
NPV	Negative predictive value
ORAC	Oxygen radical absorbance capacity
PCV	Packed cell volume
PPV	Positive predictive value
RBC	Red blood cell
ROC	Reactive oxygen species
rpm	Revolution per minute
SDFT	Superficial deep flexor tendon
TAS	Total antioxidant status
TIEP	Terengganu International Endurance Park
TP	Total protein
TPP	Total plasma protein
U/L	Unit per liter
UAE	United Arab Emirates
WEC	World Endurance Championship

CHAPTER 1

INTRODUCTION

1.1 Equestrian Sports

All equestrian sports should get the recognition of the *Fédération Equestre Internationale* (FEI) to be legitimate. The FEI, an organization that has authority in all international equestrian events was established in 1921 in Lausanne, Switzerland. Among the roles of this organization are to regulate rules and regulation and sanction all the equestrian events including the Olympics and Paralympic Games. The main goals of FEI are to promote fair play, foster equality among competing men and women riders, complicity with horses, and respect for horses and the environment. Currently, the FEI recognizes nine equestrian events which are dressage, para-equestrian dressage, jumping, eventing, driving, para-equestrian driving, vaulting, endurance, and reining.

1.2 Endurance Sports

According to the FEI, endurance is the fastest growing discipline and the second most popular discipline among equine sports. The number of endurance events have increased from only 16 in 1994 to 911 in 2016. This shows that endurance is gaining popularity among horse lovers. FEI endurance events are long-distance races conducted over 80 to 160 km which are divided into several phases of 20 to 40 km (Nagy, Dyson, & Murray, 2012).

International endurance rides are divided into two categories, which are *Concours de Raid d'Endurance International* (CEI) and *Concours de Raid d'Endurance International Official* (CEIO). Both must abide to the FEI General Regulation and FEI Endurance Rules. In CEI events, the riders compete individually. The events are conducted according to the level of expertise or 'Stars'. There are four Star levels, which are 4, 3, 2, and 1 Star. In CEIO events, the riders may compete as individual participants or as a member of a team. For each participating nation, an official team must comprise of three teams.

Endurance races are divided into several phases, each comprising a distance between 20 to 40 km. All horses are subjected to a pre-ride examination by veterinary officials. At the end of each phase, the horses must go through a compulsory halt for veterinary inspections and rest for 20 to 50 minutes before they can continue into the next loop. Horses are eliminated from the race if their metabolic or musculoskeletal conditions indicate they are not fit to continue. Horses can also be eliminated if their well-being are compromised

because of conditions such as sore at the back, girth, or mouth or suffering from extensive wounds. The horses can be eliminated either before or during the final examination after completion of the ride.

According to the FEI Rules for Endurance Events, a horse should be eliminated due to lameness at any inspection during the race. The horse shall be removed from competition and deemed failed to qualify for the next phase of the race if “the irregularity of gait is consistently observable when the horse is trotted on a loose lead held in the hand with arm straight out and back, or without flexion or deep palpation observed to be in pain, or the immediate ability of the horse to safely perform athletically is threatened” (FEI Rules of Endurance Events, 2017). The horse may also be eliminated for metabolic reasons if the veterinary inspector decides that its metabolic status is compromised. Elimination of horses can be based on abnormal general appearance, heart rate, cardiac recovery index (CRI), color of mucous membrane, capillary refill time (CRT), hydration status, intestinal activity, or wounds that may affect the horse’s ability to continue.

1.3 Hypothesis

- a) Poor performance horses have higher packed cell volume (PCV), blood creatine kinase (CK), and interleukin-6 (IL-6) as well as lower Cl^- and blood glutathione reductase (GR) concentrations than good performance horses at pre-race period.
- b) Metabolic Disorder Index (MDI) is an accurate predictor of a horse’s performance in endurance races.

1.4 Problem statements

- a) There is a high rate of elimination in endurance horses in Malaysia due to participation of poor performance horses.
- b) There is no specific method to screen the horse’s performance before the race.

1.5 Objectives

- a) To determine the rate of elimination and completion of horses in Malaysian endurance races.
- b) To determine the Metabolic Disorder Index (MDI) of good and poor performance endurance horses prior to races.
- c) To validate MDI as a predictor for potential performance of horses in endurance races.

REFERENCES

- Adamu, L. (2014). Abnormalities in Physical, Hematological and Serum Biochemical Parameters in Endurance Horses with Metabolic Crises, PhD Thesis, *Universiti Putra Malaysia*.
- Adamu, L., Adzahan, N. M., Abdullah, R., & Ahmad, B. (2012). Changes in Biochemical, Hematological and Cytokine in. *Journal of Animal and Veterinary Advances*, 11(18), 3431-3436.
- Akobeng, A. K. (2007). Understanding Diagnostic Tests 1: Sensitivity, Specificity and Predictive Values. *Acta Paediatrica*, 96(3), 338-341.
- Allen, B. V. (1988). Relationships Between the Erythrocyte Sedimentation Rate, Plasma Proteins and Viscosity, and Leucocyte Counts in Thoroughbred Racehorses. *The Veterinary Record*, 122(14), 329-332.
- Al-Qudah, K. M., & Al-Majali, A. M. (2008). Higher Lipid Peroxidation Indices in Horses Eliminated from Endurance Race Because of Synchronous Diaphragmatic Flutter (Thumps). *Journal of Equine Veterinary Science*, 28(10), 573-578.
- Anderson, M. G. (1975). The Influence of Exercise on Serum Enzyme Levels in the Horse. *Equine Veterinary Journal*, 7(3), 160-165.
- Art, T., & Lekeux, P. (2005). Exercise-Induced Physiological Adjustments to Stressful Conditions in Sports Horses. *Livestock Production Science*, 92(2), 101-111.
- Barnes, A., Kingston, J., Beetson, S., & Kuiper, C. (2010). Endurance Veterinarians Detect Physiologically Compromised Horses in a 160 km Ride. *Equine Veterinary Journal*, 42(S38), 6-11.
- Benamou-Smith, A., Robert, C., & Barrey, E. (2006). Time-Induced Changes in Hematological and Biochemical Parameters Over 1 140 Km Endurance Race in A Group of National Level Horses. In: Proceedings of the 7th International Conference on Equine Exercise Physiology, Fontainebleau, France pp13. [PDF file]. Retrieved from <http://www.ivis.org/proceedings/iceep/2006/session1.pdf>.
- Berlin, D., & Aroch, I. (2009). Concentrations of Ionized and Total Magnesium and Calcium in Healthy Horses: Effects of Age, Pregnancy, Lactation, pH and Sample Type. *The Veterinary Journal*, 181(3), 305-311.
- Buzala, M., Krumrych, W., & Janicki, B. (2015). Usefulness of Creatine Kinase Activity Determination for Assessing the Effects of Physical Effort in Horses. *Pakistan Veterinary Journal*, 35(3).

- Capomaccio, S., Cappelli, K., Spinsanti, G., Mencarelli, M., Muscettola, M., Felicetti, M., ... & Bonifazi, M. (2011). Athletic humans and horses: Comparative analysis of interleukin-6 (IL-6) and IL-6 receptor (IL-6R) expression in peripheral blood mononuclear cells in trained and untrained subjects at rest. *BMC physiology*, 11(1), 3.
- de Lacerda-Neto, J. C., de Cassia de Lima Sampaio, R., Ferraz, G. C., Teixeira-Neto, A. R., Pereiral, D. M., Titto, E. A. L., Carvalho, M. B., & de Queiroz-Neto, A. (2003). Effects of Intermittent Water Cooling and Electrolyte Supplements on Serum Osmolality and Electrolytes of Horses during Low-Intensity Exercise. *Revista Portuguesa de Ciências Veterinárias*, 189-196.
- Di Filippo, Alessandra, P., Gomide, L. M. W., Orozco, C. A. G., da Silva, M. A., Martins, C. B., Lacerda, N., & Santana, A. E. (2009). Changes Electrolytes and Blood Gases in Arabian Horses during to 60 km Endurance Race. *Ciência Animal Brasileira*, 10(3), 840-846.
- Donovan, D. C., Jackson, C. A., Colahan, P. T., Norton, N., & Hurley, D. J. (2007). Exercise-induced alterations in pro-inflammatory cytokines and prostaglandin F2 α in horses. *Veterinary immunology and immunopathology*, 118(3-4), 263-269.
- Dorrington, K. (1981). Skin Turgor: Do We Understand the Clinical Sign? *The Lancet*, 317(8214), 264-266.
- Dumont, C. B. S., Okiyama, F. T. M., Barreto-Vianna, A. R. C., & Lima, E. M. M. (2014). The Distance Run is Not the Determining Factor on the Occurrence of Electrolyte Change for Equine Athletes Disqualified in Endurance Competitions. *Pakistan Veterinary Journal*, 34(2), 239-242.
- Evans, D. L., & Rose, R. J. (1988). Cardiovascular and Respiratory Responses to Submaximal Exercise Training In the Thoroughbred Horse. *Pflügers Archiv*, 411(3), 316-321.
- Fédération Equestre Internationale. Retrieved from <http://www.fei.org/>.
- Fédération Equestre Internationale. Rules for Endurance Events. 9th ed. Effective 1st January 2017, Updates effective 1st January 2017.
- Fielding, C. L., Meier, C. A., Balch, O. K., & Kass, P. H. (2011). Risk Factors for the Elimination of Endurance Horses from Competition. *Journal of the American Veterinary Medical Association*, 239(4), 493-498.
- Flaminio, M. J., & Rush, B. R. (1998). Fluid and Electrolyte Balance in Endurance Horses. *Veterinary Clinics: Equine Practice*, 14(1), 147-158.

- Flaminio, M. J., Gaughan, E. M., & Gillespie, J. R. (1996). Exercise Intolerance in Endurance Horses. *Veterinary Clinics: Equine Practice*, 12(3), 565-580.
- Foreman, J. H. (1998). The Exhausted Horse Syndrome. *Veterinary Clinics: Equine Practice*, 14(1), 205-219.
- Gajewski, M., Kamińska, E., Wysocki, Ł., Szczepanik, S., Sygitowicz, G., Wojciechowski, M., & Maśliński, S. (2005). Gospodarka tlenowa w organizmie. Część I. Warunki normy fizjologicznej. *Życie Weterynaryjne*, 80(7), 380-386.
- Gill, J., & Rastawicka, M. (1985). Diurnal Changes in the Haematological Indices in the Blood of Racing Arabian Horses. *Polskie Archiwum Weterynaryjne*, 26(1-2), 169-179.
- Golland, L. C., Evans, D. L., McGowan, C. M., Hodgson, D. R., & Rose, R. J. (2003). The Effects of Overtraining on Blood Volumes in Standardbred Racehorses. *The Veterinary Journal*, 165(3), 228-233.
- Gondim, F. J., Zoppi, C. C., Pereira-da-Silva, L., & de Macedo, D. V. (2007). Determination of the Anaerobic Threshold and Maximal Lactate Steady State Speed in Equines Using the Lactate Minimum Speed Protocol. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 146(3), 375-380.
- Grondin, T. M. & Dewitt, S. F. (2010). Normal Hematology of the Horse and Donkey. *Weiss DK, Wardrop KJ. Schalm's Veterinary Hematology. 6th ed. Iowa: Wiley-Blackwell*, 821-828.
- Halliwell, B., & Gutteridge, J. M. (1990). The Antioxidants of Human Extracellular Fluids. *Archives of Biochemistry and Biophysics*, 280(1), 1-8.
- Harold, C. S. (2010). Challenges of Endurance Exercise: Hydration and Electrolyte Depletion. *Kentucky Equine Research*, 94-111.
- Harvey, R. B., Hambright, M. B., & Rowe, L. D. (1984). Clinical Biochemical and Hematologic Values of the American Miniature Horse: Reference Values. *American Journal of Veterinary Research*, 45(5), 987-990.
- Hernández, A. M., Satué, K., Lorente, C., Garcés, C., & O'connor, J. E. (2008). The Influence of Age and Gender on Haematological Parameters in Spanish Horses. In *Proc Vet Eur Equine Meeting-XIV SIVE Congress, Venice (Italy)*.
- Hinchcliff, K. W., Kaneps, A. J., & Geor, R. J. (2013). *Equine Sports Medicine and Surgery E-Book*. (2nd ed.). Elsevier Health Sciences, Saunders Ltd.
- Hodgson, D. R. & Rose, R.J. (1994). *The Athletic Horse*. W.B. Saunders, Philadelphia, pp: 169-174.

- Jain, N. C. (1993). *Essentials of Veterinary Hematology*. 1st Edition, Wiley-Blackwell.
- Jain, N.C. (1986). The Horse. Normal Haematologic With Comments on Response to Disease. *Schalm's Veterinary Hematology*. Jain N.C. (ed.), pp. 140-177.
- Janicki, B., Kochowicz, A., Buzala, M., & Krumrych, W. (2013). Variability of Selected Biochemical Parameters in Young Stallions during the 100-Day Performance Test. *Journal of Equine Veterinary Science*, 33(12), 1136-1141.
- Jawor, P., Stefaniak, T., Borkowski, J., Wierzbicki, H., & Swagemakers, J. H. (2007). Utility of Selected Blood Parameters in Monitoring the Accuracy of Show Jump-Horse Training. *Medycyna Weterynaryjna*, 63(2), 213-218.
- Jeffcott, L.B. (1977). Clinical Haematology of the Horse. *Comparative Clinical Haematology*. Archer, R.K. & Jeffcott, L.B. (eds.), pp. 161-213.
- Kedzierski, W., & Podolak, M. (2002). Training Arabian Horses and its Effect on the Level of Biochemical Indices Related to the Metabolism of Carbohydrates and Lipids. *Medycyna Weterynaryjna*, 58(10), 788-791.
- Kerr, M. G., & Snow, D. H. (1982). Alterations in Hematocrit, Plasma-Proteins and Electrolytes in Horses Following the Feed of Hay. *Veterinary Record*, 110(23), 538-540.
- Kinnunen, S., Atalay, M., Hyypä, S., Lehmuskero, A., Hänninen, O. & Oksala, N. (2005). Effects of Prolonged Exercise on Oxidative Stress and Antioxidant Defense in Endurance Horse. *Journal of Sports Science and Medicine*, 4(4), 415-21.
- Kinnunen, S., Hyypä, S., Lehmuskero, A., Oksala, N., Mäenpää, P., Hänninen, O., & Atalay, M. (2005). Oxygen Radical Absorbance Capacity (ORAC) and Exercise-Induced Oxidative Stress in Trotters. *European Journal of Applied Physiology*, 95(5-6), 550-556.
- Kramer, J. W. (2000). Normal Hematology of Cattle, Sheep and Goats. *Schalm's Veterinary Hematology*, 5, 1075-1084.
- Kurosawa, M., Nagata, S. I., Takeda, F., Hiraga, A., & Kazuyoshi, T. A. Y. A. (2001). Plasma Catecholamine, Adrenocorticotropin and Cortisol Responses to Exhaustive Incremental Treadmill Exercise of the Thoroughbred Horse. *Journal of Equine Science*, 9(1), 9-18.
- Lamprecht, E. D., Bagnell, C. A., & Williams, C. A. (2008). Inflammatory responses to three modes of intense exercise in Standardbred mares—a pilot study. *Comparative Exercise Physiology*, 5(3-4), 115-125.

- Lawan, A., Noraniza, M. A., Rasedee, A., & Bashir, A. (2012). Prevalence of Lameness and Metabolic Disorders in Endurance Horses. *Malaysian Journal of Veterinary Research*, 3, 33-37.
- Lewis, L. D. (1995). *Equine Clinical Nutrition, Feeding and Care*. Williams and Wilkins, Baltimore, pp: 262-263.
- Malinowski, K., Betros, C. L., Flora, L., Kearns, C. F., & Mckeever, K. H. (2002). Effect of Training on Age-Related Changes in Plasma Insulin and Glucose. *Equine Veterinary Journal*, 34(S34), 147–153.
- Marlin, D. J., McEwen, J., & Sluyter, F. (2008). Completion and Treatment Rates in Modern Endurance Racing. In: Proceedings of 4th International Conference of the International Society for Equitation Science: Dublin, Ireland.
- McArdle, F., Pattwell, D. M., Vasilaki, A., McArdle, A., & Jackson, M. J. (2005). Intracellular Generation of Reactive Oxygen Species by Contracting Skeletal Muscle Cells. *Free Radical Biology and Medicine*, 39(5), 651-657.
- McGowan, C. (2008). Clinical Pathology in the Racing Horse: The Role of Clinical Pathology in Assessing Fitness and Performance in the Racehorse. *Veterinary Clinics: Equine Practice*, 24(2), 405-421.
- McKeever, K. H., (2004). Body Fluid and Electrolytes: Responses to Exercise and Training. In *Equine Sports Medicine and Surgery*, Hinchcliff, K.W., A.J. Kanep and R.J Geor (Eds). WB. Saunders, Philadelphia, pp: 854-871.
- McKeever, K. H., Hinchcliff, K. W., Reed, S. M., & Robertson, J. T. (1993). Role of Decreased Plasma Volume in Hematocrit Alterations during Incremental Treadmill Exercise in Horses. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 265(2), R404-R408.
- Misheff, M. M., Alexander, G. R., & Hirst, G. R. (2010). Management of Fractures in Endurance Horses. *Equine Veterinary Education*, 22(12), 623-630.
- Muñoz, A., Riber, C., Trigo, P., & Castejón, F. M. (2012). Age-and Gender-Related Variations in Hematology, Clinical Biochemistry, and Hormones in Spanish Fillies and Colts. *Research in Veterinary Science*, 93(2), 943-949.
- Muñoz, A., Riber, C., Trigo, P., Castejón-Riber, C., & Castejón, F. M. (2010). Dehydration, Electrolyte Imbalances and Renin-Angiotensin-Aldosterone-Vasopressin Axis in Successful and Unsuccessful Endurance Horses. *Equine Veterinary Journal*, 42(38), 83-90.

- Nagy, A., Dyson, S. J., & Murray, J. K. (2012). A Veterinary Review of Endurance Riding As an International Competitive Sport. *The Veterinary Journal*, 194(3), 288-293.
- Nagy, A., Murray, J. K. & Dyson, S. (2010). Elimination from Elite Endurance Rides in Nine Countries: A Preliminary Study. *Equine Veterinary Journal*, 42(S38), 637-643.
- Nagy, A., Murray, J. K., & Dyson, S. J. (2014). Horse-, Rider-, Venue-and Environment-Related Risk Factors for Elimination from Fédération Equestre Internationale Endurance Rides Due to Lameness and Metabolic Reasons. *Equine Veterinary Journal*, 46(3), 294-299.
- Navarro, A., Gomez, C., López-Cepero, J. M., & Boveris, A. (2004). Beneficial Effects of Moderate Exercise on Mice Aging: Survival, Behavior, Oxidative Stress, and Mitochondrial Electron Transfer. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 286(3), R505-R511.
- Nieto, J. E., Snyder, J. R., Beldomenico, P., Aleman, M., Kerr, J. W., & Spier, S. J. (2004). Prevalence of Gastric Ulcers in Endurance Horses – A Preliminary Report. *The Veterinary Journal*, 167(1), 33-37.
- Nimmo, M. A., & Snow, D. H. (1982). Time Course of Ultrastructural Changes in Skeletal Muscle after Two Types of Exercise. *Journal of Applied Physiology*, 52(4), 910-913.
- Pedersen, B. K. (2005). Natural Immunity–Effect of Exercise. In *Neuroimmune Biology* (Vol. 5, pp. 263-288). Elsevier.
- Penders, J. J. J. N. (2015). Risk Factors for Metabolic Elimination in Endurance Rides in New Zealand. Master Thesis, Massey University, New Zealand.
- Perez, R., Recabarren, S. E., Valdes, P., & Hetz, E. (1992). Biochemical and Physiological Parameters and Estimated Work Output in Draught Horses Pulling Loads for Long Periods. *Veterinary Research Communications*, 16(3), 231-246.
- Persson, S. G. B., & Österberg, I., (1999). Racing Performance in Red Blood Cell Hypervolemic Standardbred Trotters. *Equine Veterinary Journal*, 31(S30), 617-620.
- Petersen, A. M. W., & Pedersen, B. K. (2005). The anti-inflammatory effect of exercise. *Journal of applied physiology*, 98(4), 1154-1162.
- Piccione, G., Casella, S., Giannetto, C., Messina, V., Monteverde, V., Caola, G., & Guttadauro, S. (2010). Haematological and Haematochemical Responses to Training and Competition in Standardbred Horses. *Comparative Clinical Pathology*, 19(1), 95-101.

- Pritchard, J. C., Burn, C. C., Barr, A. R. S., & Whay, H. R. (2008). Validity of Indicators of Dehydration in Working Horses: A Longitudinal Study of Changes in Skin Tent Duration, Mucous Membrane Dryness and Drinking Behaviour. *Equine Veterinary Journal*, 40(6), 558-564.
- Robert, C., Goachet, A. G., Fraipont, A., Votion, D. M., van Erck, E., & Leclerc, J. L. (2010). Hydration and Electrolyte Balance in Horses during an Endurance Season. *Equine Veterinary Journal*, 42(S38), 98-104.
- Robson-Ansley, P., Barwood, M., Canavan, J., Hack, S., Eglin, C., Davey, S., ... & Ansley, L. (2009). The effect of repeated endurance exercise on IL-6 and sIL-6R and their relationship with sensations of fatigue at rest. *Cytokine*, 45(2), 111-116.
- Rose, R. J. (1986). Endurance Exercise in the Horse - A Review. Part I. *British Veterinary Journal*, 142(6), 532-541.
- Rose, R. J. (1986). Endurance Exercise in the Horse - A Review. Part II. *British Veterinary Journal*, 142(6), 542-552.
- Rose, R. J., & Hodgson, D. R. (2000). Fluid and Electrolyte Therapy: Assessment of Fluid and Electrolyte Balance. *Manual of Equine Practice*, 2, 757-758.
- Ross, W. A., & Kaneene, J. B. (1996). An Individual-Animal-Level Prospective Study of Risk Factors Associated with the Occurrence of Lameness in the Michigan (USA) Equine Population. *Preventive Veterinary Medicine*, 29(1), 59-75.
- Satué, K., Blanco, O., & Muñoz, A. (2009). Age-Related Differences in the Hematological Profile of Andalusian Broodmares of Carthusian Strain. *Veterinary Medicine*, Vol. 54, N° 4, (April 2009), pp. 175–182.
- Satué, K., Hernández, A., & Muñoz, A. (2012). Physiological Factors in the Interpretation of Equine Hematological Profile. In *Hematology-Science and Practice*. InTech.
- Schott 2nd, H. C., McGlade, K. S., Molander, H. A., Leroux, A. J., & Hines, M. T. (1997). Body Weight, Fluid, Electrolyte, and Hormonal Changes in Horses Competing in 50- and 100-Mile Endurance Rides. *American Journal of Veterinary Research*, 58(3), 303-309.
- Sen, C. K. (1999). Glutathione Homeostasis in Response to Exercise Training and Nutritional Supplements. *Molecular and Cellular Biochemistry*, 196(1), 31-42.
- Sharif, M. T., Mahabadi, M. A., Moshfeghi, S., Sharifi, H., Hoseini, S. M., & Alavi, S. M. (2012). Artifactual Changes in Hematological Variables in Equine Blood Samples Stored at Different Temperatures and Various Anticoagulants. *Comparative Clinical Pathology*, 21(4), 449-452.

- Sjödin, B., Westing, Y. H., & Apple, F. S. (1990). Biochemical Mechanisms for Oxygen Free Radical Formation during Exercise. *Sports Medicine*, 10(4), 236-254.
- Snow, D. H., Mason, D. K., Ricketts, S. W., & Douglas, T. A. (1983). Post-Race Blood Biochemistry in Thoroughbreds. *Equine Exercise Physiology*, 1, 389-399.
- Szarska, E. (2002). Wykorzystanie badań diagnostycznych krwi do oceny stanu zdrowia i zaawansowania treningowego koni wyczynowych. Wyczynowych. [The Use of Blood Diagnostic Test to Assess the State of Health and Progress of Training Performance of Racing Horses]. *Rozprawy Naukowe i Monografie. Szkoła Główna Gospodarstwa Wiejskiego w Warszawie*, (250), 1-74.
- Szarska, E. (2007). *Konne rajdy długodystansowe*. [Long-Distance Horse Races]. Agencja Reklamowa. CREX, 55-63. [in polish].
- Taylor, F. G., Brazil, T., & Hillyer, M. H. (Eds.). (2010). *Diagnostic Techniques in Equine Medicine E-Book*. Elsevier Health Sciences.
- Thrusfield, M. (2013). *Veterinary Epidemiology*. Elsevier.
- Tomlinson, C. M. (1992). Exhausted Horse Syndrome. In: Proceedings of the annual convention of the American Association of Equine Practitioners (USA). Orlando, Florida, November 29th to December 2nd 1992.
- Trigo, P., Castejon, F., Riber, C., & Muñoz, A. (2010). Use of Biochemical Parameters to Predict Metabolic Elimination in Endurance Rides. *Equine Veterinary Journal*, 42(S38), 142-146.
- Tyler-Mcgowan, C. M., Golland, L. C., Evans, D. L., Hodgson, D. R., & Rose, R. J. (1999). Haematological and Biochemical Responses to Training and Overtraining. *Equine Veterinary Journal*, 31(S30), 621-625.
- Valberg, S. J. (2006). Exertional Rhabdomyolysis. In: Proceedings of the 52nd Annual Convention of the American Association of Equine Practitioners, San Antonio, Texas, USA, 2nd-6th December 2006, 365-372.
- Viu, J., Jose-Cunilleras, E., Armengou, L., Cesarini, C., Tarancón, I., Rios, J., & Monreal, L. (2010). Acid-Base Imbalances during a 120 km Endurance Race Compared by Traditional and Simplified Strong Ion Difference Methods. *Equine Veterinary Journal*, 42(S38), 76-82.
- Whiting, J., (2009). The Exhausted Horse. In: Current Therapy in Equine Medicine, Sixth Ed. Saunders Elsevier, St. Louis, MO, USA, pp. 926–929.

Wickler, S. J., Greene, H. M., Egan, K., Astudillo, A., Dutto, D. J., & Hoyt, D. F. (2006). Stride Parameters and Hindlimb Length in Horses Fatigued on a Treadmill and at an Endurance Ride. *Equine Veterinary Journal*, 38(S36), 60-64.

Zimmel, D. N. (2003). How to Manage Pain and Dehydration in Horses with Colic. In: Proceedings of the 49th Annual Convention of the American Association of Equine Practitioners, New Orleans, Louisiana, USA, 21st-25th November 2003: 127-131.

Zoppi, C. C., Hohl, R., Silva, F. C., Lazarim, F. L., Neto, J. A., Stancanneli, M., & Macedo, D. V. (2006). Vitamin C and E Supplementation Effects In Professional Soccer Players Under Regular Training. *Journal of the International Society of Sports Nutrition*, 3(2), 37.

