



UNIVERSITI PUTRA MALAYSIA

***ASSESSMENT OF SPERM CRYOPRESERVATIVES AND OOCYTE
VITRIFICATION FOR IN VITRO EMBRYO PRODUCTION IN CATTLE***

SALMAN HAMMADI GHAREEB

FPV 2018 8



**ASSESSMENT OF SPERM CRYOPRESERVATIVES AND OOCYTE
VITRIFICATION FOR *IN VITRO* EMBRYO PRODUCTION IN CATTLE**

By

SALMAN HAMMADI GHAREEB

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

December 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

Dedicated to my father (Almarhum) and my mother (Almarhuma), to my beloved wife and my kids Mayar, Acif and Layan for their supplication, patience and understanding



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

ASSESSMENT OF SPERM CRYOPRESERVATIVES AND OOCYTE VITRIFICATION FOR *IN VITRO* EMBRYO PRODUCTION IN CATTLE

By

SALMAN HAMMADI GHAREEB

December 2017

Chairman : Professor Abd Wahid Haron, PhD
Faculty : Veterinary Medicine

Animal breeding programmes and assisted reproductive techniques (ART) such as artificial insemination (AI), embryo transfer (ET) and intra-cytoplasmic sperm injection (ICSI) are extremely dependant on the quality of sperm and oocytes used. Furthermore, fertility ratios still need to be improved particularly those related to the cryopreserved sperms and vitrified oocyte. In this study, a number of experiments were carried out to improve the quality of cryopreserved semen and vitrified immature oocytes and evaluation of *in vitro* embryo production as well. Seven bull semen extenders were prepared and used to extend a total of 102 ejaculates that were collected from twelve mature healthy fertile Brangus-Herford bulls by automatic electro-stimulation at the animal farm of the Universiti Putra Malaysia. On the other hand, 180 ovaries were collected from slaughtered local cattle at Banting Abattoir, Malaysia. Post-thaw semen evaluation were evaluated for the extending capacity of seven different types of extenders implemented in bull semen cryopreservation; these extenders were AndroMed®, BioXcell®, Triladyl®, Tris-egg yolk based extenders 20%, Tris-egg yolk plasma with concentration of 10%, Tris-egg yolk plasma with concentration of 15 % and Tris-egg yolk plasma with concentration of 20% on the sperm's motility, viability, morphology, DNA integrity, plasma membrane integrity and acrosome integrity. Also, the cryopreservation process was conducted in two regimes of fresh semen preparation prior to subjecting to cryopreservation. These were seminal plasma being removed from fresh semen samples and fresh semen samples with their seminal plasma. Semen samples were collected using electrical stimulation by an automatic electro-ejaculator (AEE), and semen quality was analysed using computer assisted semen analyser (CASA). The Eosin-Nigrosine staining technique was applied to evaluate the morphology, viability and acrosome intergrity. The DNA integrity was evaluated using a modified Acridine orange method. Amongst the three commercial extenders, AndroMed®, BioXcell®, Triladyl® and the Tris-egg yolk based extenders, the AndroMed® extender produced the highest extending capacity

to the sperm's motility, morphology, DNA integrity, plasma membrane integrity and acrosome integrity. BioXcell® and Tris-egg yolk based extenders had the best capacity to protect the sperm's viability. These results, therefore, favour AndroMed® over the rest of the other extenders used in post-thaw sperm parameters with seminal plasma. The AndroMed®, Triladyl® and Tris –egg yolk extender were highest in extending capacity than BioXcell® at the level of post-thaw semen evaluation and preserving semen without seminal plasma.

Among the egg yolk extenders, the egg yolk plasma extender 10% produced the highest extending capacity to the semen plasma membrane integrity, viability, and motility, while the Tris-egg yolk based extender 20% and Tris-egg yolk plasma 10% had the best capacity to protect the semen's morphology and acrosome integrity. These results, therefore, favour the concentration of Tris-egg yolk plasma 20% over the rest of the other extender concentration egg yolk used in extending ejaculated semen with seminal plasma. In the study that involved seminal plasma removal, the spermatozoa was also evaluated for its qualities using Tris-egg yolk based extenders 20%, Tris-egg yolk plasma with concentration 10% , Tris-egg yolk plasma with concentration 15% and Tris-egg yolk plasma with concentration 20%. The egg yolk plasma with concentration 15% extender gave a better extending capacity to the semen viability and acrosome integrity, while Tris-egg yolk based extender 20% had the best capacity to protect the semen's morphology and plasma membrane integrity. This result, therefore, favours the concentration of the egg yolk plasma with concentration 15% and Tris-egg yolk based extender 20% over the rest of the other extender concentrations of egg yolk plasma.

In the immature oocyte study, oocytes were harvested from bovine ovaries collected from local abattoir (Banting, Malaysia), then transported in thermos containers at 37°C (phosphate buffer saline) to the Theriogenology and Cytogenetic laboratory within 2-3 hours post-slaughtering. Slicing method of ovaries were applied to harvest the immature oocytes. Pipetting of the considered oocytes was done from the handling medium into the washing medium. This study aimed to study the effect of different vitrification exposure times of 15 sec, 30 sec and 45 sec of immature bovine oocytes and their ability to fertilise and produce an embryos after application of *In vitro* fertilisation technology. Ethylene Glycol (EG) and Dimethylsulfoxide (DMSO) were used as a combination of two cryoprotectants. Two main cryodevices were also used; they were open pulled straws (OPS) and closed French straws (CFS). OPS were modified locally by using hot plate and pulled manually. Vitrified-warmed immature oocytes were evaluated morphologically under epifluorescent microscopy. The viable oocytes were subjected into the maturation media, mainly composed of TCM199+HEPES supplemented with 10% FCS and 10 ng/ml epidermal growth factor (EGF). Polar body extrusion was observed 22 hr post IVM. *In vitro* fertilization (IVF) was applied using direct infusion of frozen-thawed bull semen extended in AndroMed® (as the best extender described in first experiment; Chapter Four). The frozen-thawed bull semen was collected at the bottom of Percoll solution (45% over 90%) then washed in Tyrodes medium. The viable spermatozoa were diluted in the appropriate volume of fertilization medium to achieve a final concentration of 2×10^6

sperm/ml. A 250 μ l aliquot of this suspension was added to each well containing oocytes for a final concentration of sperm cells of 1×10^6 sperm/ml. Spermatozoa and oocytes were co-incubated for 18-22 h at 38.8 °C in a humidified atmosphere of 5% CO₂ in air. For *in vitro* culture (IVC) presumptive zygotes were vortexed and washed three times in HEPES-buffered TCM199 containing 0.4 g/l BSA and 4 μ g/ml gentamicin, to remove cumulus cells and attached spermatozoa. Zygotes were washed two times in SOF medium supplemented with BME amino acids (50 \times), MEM amino acids (100 \times), 3 mM pyruvate, 6 mg/ml fatty acid free BSA and 5% FBS and transferred by groups of 25 into four-well petri dishes containing 50 μ l drops of culture medium covered with 700 μ l of mineral oil. This study showed that the immature bovine oocytes can be vitrified at the GV-stage with our optimized protocol and that the vitrified-warmed GV-oocytes could be matured and *in vitro* fertilized and subsequently *in vitro* embryo development. Also, the vitrification of bovine immature oocytes in the solutions composed of 15% EG + 15% DMSO using OPS method achieved the best viability, the highest maturation, and cleavage in our conditions after IVM and IVF despite a lower recovery rate than the other group. Data were analysed using SPSS software system (Version 12.0, SPSS, Chicago, IL). Study parameters were analysed using a 2-way factorial analysis of variance (ANOVA). Values were expressed as mean \pm standard error of the mean (S.E.M). The level of statistical significance was at $P < 0.05$.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENILIAIAN UNTUK SPERMA KRIOAWETAN DAN OOSIT
PENGKEAKAN *IN VITRO* PRODUKSI EMBRIO LEMBU**

Oleh

SALMAN HAMMADI GHAREEB

Disember 2017

Pengerusi : Profesor Abd Wahid Haron, PhD
Fakulti : Perubatan Veterinar

Program pembiakan haiwan dan teknik pembiakan dibantu (ART) seperti pernianian beradas (AI), pemindahan embrio (ET) dan suntikan sperma intra-sitoplasma (ICSI) amat bergantung kepada kualiti sperma dan oosit yang digunakan. Tambahan pula, nisbah kesuburan masih perlu diperbaiki terutamanya yang berkaitan dengan sperma dikrioawet dan oosit divitfrifikasikan. Dalam kajian ini, beberapa eksperimen telah dijalankan untuk meningkatkan kualiti semen dikrioawet dan oosit divitfrifikasikan yang tidak matang dan penilaian pengeluaran embrio *in vitro* juga. Semasa pembaikan sperma dikrioawet, empat eksperimen utama telah dijalankan. Tujuh penyambung semen lembu jantan telah disediakan dan digunakan untuk menyambungkan sejumlah 102 pancutan air mani yang dikumpul daripada dua belas lembu jantan Brangus-Herford sihat matang subur oleh elektro-rangsangan automatik di ladang haiwan Universiti Putra Malaysia. Sebaliknya, 180 ovari dikumpulkan dari lembu tempatan yang disembelih di Banting Abattoir, Malaysia. Penilaian air mani selepas-dicairkan telah dinilai untuk kapasiti menyambung tujuh jenis penyambung yang dilaksanakan untuk krioawetan semen lembu jantan; penyambung-penyambung ini ialah AndroMed®, BioXcell®, Triladyl®, penyambung berasaskan Tris-kuning telur penyambung 20%, plasma Tris-kuning telur dengan kepekatan 10%, plasma Tris-kuning telur dengan kepekatan 15% dan plasma Tris-kuning telur dengan kepekatan 20% ke atas motiliti, daya maju, morfologi, integriti DNA, integriti membran plasma dan integriti akrosom sperma. Juga, proses krioawetan telah dijalankan mengikut dua rejim penyediaan semen segar sebelum menjalankan krioawetan. Ini adalah pengeluaran plasma semen daripada sampel semen segar dan sampel semen yang segar dengan plasma semennya. Sampel semen dikumpul dengan menggunakan rangsangan elektrik dengan elektro-ejakulator automatik (AEE), dan kualiti semen dianalisis menggunakan penganalisis semen dibantu komputer (CASA). Teknik pewarnaan Eosin-Nigrosin digunakan untuk menilai morfologi, daya maju dan intergrity akrosom. Integriti DNA dinilai menggunakan kaedah oren Acridine

diubahsuai. Di antara tiga penyambung komersial, AndroMed®, BioXcell®, Triladyl® dan penyambung-penyambung berasaskan Tris-kuning telur, penyambung AndroMed® menghasilkan kapasiti penyambungan tertinggi kepada motiliti, morfologi, integriti DNA, integriti membran plasma dan integriti akrosom sperma. Penyambung berdasarkan BioXcell® dan Tris-kuning telur mempunyai keupayaan yang terbaik untuk melindungi daya maju sperma. Oleh itu, keputusan ini menyokong AndroMed® lebih dari penyambung yang lain yang digunakan untuk parameter sperma pasca-cair dengan plasma semen. Penyambung AndroMed®, Triladyl® dan Tris-kuning telur adalah paling baik dalam kapasiti penyambungan berbanding dengan BioXcell® pada tahap penilaian semen pasca-cair dan pemeliharaan semen tanpa plasma semen.

Di kalangan penyambung kuning telur, penyambung plasma kuning telur 10% menghasilkan kapasiti penyambung tertinggi untuk integriti membran plasma semen, daya maju, dan motiliti, manakala penyambung berdasarkan Tris-kuning telur 20% dan Tris-kuning telur plasma 10% mempunyai keupayaan terbaik untuk melindungi morfologi semen dan integriti akrosom. Oleh itu, keputusan ini menyokong kepekatan Tris-kuning telur plasma 20% berbanding kepekatan penyambung kuning telur lain yang digunakan untuk menyambung semen diejakulasi dengan plasma semen. Di dalam kajian yang melibatkan penyingkiran plasma semen, spermatozoa juga dinilai untuk kualitinya dengan menggunakan penyambung Tris-kuning telur 20%, Tris-kuning telur plasma dengan kepekatan 10%, Tris-kuning telur plasma dengan kepekatan 15% dan Tris-kuning telur plasma dengan kepekatan 20%. Penyambung plasma kuning telur dengan kepekatan 15% memberikan kapasiti penyambungan yang lebih baik kepada daya maju semen dan integriti akrosom, manakala penyambung berdasarkan Tris-kuning telur 20% mempunyai kapasiti yang terbaik untuk melindungi morfologi dan integriti membran plasma semen. Oleh itu, keputusan ini memihak kepada kepekatan plasma kuning telur dengan kepekatan 15% dan penyambung berdasarkan Tris-kuning telur 20% lebih berbanding dengan kepekatan pemanjang plasma kuning telur yang lain.

Di dalam kajian oosit yang tidak matang, oosit dituai dari ovari bovin dikumpul daripada rumah penyembelihan tempatan (Banting, Malaysia), kemudian diangkut di dalam bekas termos pada suhu 37°C (salina penampan fosfat) ke makmal Theriogenologi dan Citogenetik dalam masa 2-3 jam selepas penyembelihan. Kaedah penghirisan ovari digunakan untuk menuai oosit yang tidak matang. Penggunaan pipet bagi oosit yang dipertimbangkan telah dilakukan dari medium pengendalian ke dalam medium basuh. Kajian ini bertujuan untuk menyiasat kesan masa pendedahan vitrifikasi yang berbeza selama 15 saat, 30 saat dan 45 saat ke atas oosit bovin tidak matang dan keupayaan mereka untuk bersenyawa dan menghasilkan embrio selepas penggunaan teknologi persenyawaan *In vitro*. Etilena Glikol (EG) dan Dimethylsulfoxide (DMSO) telah digunakan sebagai gabungan dua kriopelindung. Dua krioperanti utama juga telah digunakan; ini ialah straw tertarik terbuka (OPS) dan straw Perancis tertutup (CFS). OPS diubahsuai secara tempatan dengan menggunakan plat panas dan ditarik secara manual. Oosit yang divitrifikasi-hangatkan yang belum matang telah dinilai dari segi morfologi di bawah

kemikroskopan epifluoresen. Oosit yang berdaya maju dimasukkan ke dalam media kematangan, yang terutamanya terdiri daripada TCM199+HEPES ditambah dengan 10% FCS dan 10 ng/ml faktor pertumbuhan epidermis (EGF). Penyemperitan badan kutub diperhatikan 22 jam pasca IVM. Persenyawaan *In vitro* (IVF) telah digunakan secara infusi langsung semen lembu jantan beku-dicairkan disambungkan dalam AndroMed® (sebagai penyambung terbaik digambarkan dalam eksperimen pertama Bab Empat). Semen lembu jantan beku-dicairkan dikumpulkan di bahagian bawah larutan Percoll (45% atas 90%) kemudian dicuci di medium Tyrodes. Spermatozoa berdaya maju telah dicairkan dalam isipadu medium persenyawaan yang sesuai untuk mencapai kepekatan muktamad sebanyak 2×10^6 sperma/ml. Suatu bahagian 50 μ l ampaian ini telah ditambah kepada setiap telaga mengandungi oosit untuk kepekatan muktamad sel-sel sperma sebanyak 1×10^6 sperma/ml. Spermatozoa dan oosit telah dieram bersama selama 18-22 jam pada 38.8°C dalam atmosfera dilembapkan dan sebanyak 5% CO₂ di dalam udara. Kultur *invitro* (IVC) zigot andaian telah divortekskan dan dibasuh tiga kali di dalam TCM199 ditampan-HEPES mengandungi 0.4 g/l BSA dan 4 μ g/ml gentamicin, untuk membuang sel-sel kumulus dan spermatozoa terlekat. Zigot dibasuh dua kali di dalam medium SOF ditambah dengan asid amino BME (50 \times), Asid amino MEM (100 \times), 3 mM pyruvate, 6mg/ml BSA bebas dari asid lemak dan 5% FBS dan dipindahkan secara 250 setiap kumpulan ke piring petri empat-telaga yang mengandungi 25 μ l tetes medium kultur yang ditutupi dengan 700 μ l minyak mineral. Kajian ini memperlihatkan bahawa oosit bovin yang tidak matang boleh divitrifikasi di peringkat-GV dengan protokol kami yang dioptimumkan dan bahawa oosit-GV divitrifikasi-hangatkan boleh dimatangkan dan disenyawakan *in vitro* dan seterusnya perkembangan embrio *in vitro*. Juga, vitrifikasi oosit tidak matang bovin dalam larutan terdiri daripada 15% EG + 15% DMSO menggunakan kaedah OPS mencapai daya maju yang terbaik, kematangan tertinggi, dan belahan dalam keadaan-keadaan kami selepas IVM dan IVF walaupun dengan Kadar pemulihan yang lebih rendah daripada kumpulan lain. Parameter dianalisis menggunakan 2-cara analisis (ANOVA) bagi varians. Nilai itu dinyatakan sebagai min piawai untuk min (S.E.M). Tahap kepentingan statistic adalah pada $P < 0.05$.

ACKNOWLEDGEMENTS

By the name of ALLAH: The Most Merciful and the Most Compassionate. The praise is to ALLAH, The Almighty for providing me this opportunity and granting me the capability to proceed successfully. I would like to express my deep appreciation and sincere gratitude to my respectful supervisor, the Chairman of the Supervisory Committee, Professor, Dr. Abd Wahid Haron for his enthusiasm and passion that he has for science. I am grateful for the opportunity that I had, to work with Professor Dr. Wahid and always consider him an excellent mentor and great friend. I also thank to Associate Professor, Dr. Rosnina Yusoff and Dr. Nurhusein Yimer Degu, members of the Supervisory Committee, for their expert instruction, their thorough reading, guidance, patience and advice, I am forever indebted.

I express many thanks to Mr. Yap Keng Chee and Mr. Ganes Murthi, Mr. Jameel who they were like friends to me, and their assistance in Theriogenology and Cytogenetic laboratory as my own was remarkable.

I am also indebted to the staff of UPM Farm, Cattle unit especially Mr. Rizalana for his invaluable assistance in semen collection. Thanks also extended to the staff of Abattoir Banting, Malaysia for facilitating collecting of cattle ovaries for this study.

Finally, I am thankful to the faculty of Veterinary Medicine, I am extremely thankful to UNIVERSITI PUTRA MALAYSIA for supporting me during my years of study.

Last but not least, I dedicate my thanks to my lovely wife Dr. Anhar Adil Abdullah for her support and patience, and my thanks also dedicated to my team, my lovely kids Mayar, Acif and Layan for their support and inspiration.

I certify that a Thesis Examination Committee has met on 5 December 2017 to conduct the final examination of Salman Hammadi Ghareeb on his thesis entitled "Assessment of Sperm Cryopreservatives and Oocyte Vitrification for *In Vitro* Embryo Production in Cattle" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Noordin bin Mohamed Mustapha, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Md Zuki bin Abu Bakar @ Zakaria, PhD


Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Abdul Aziz bin Saharee, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Arief Boediono, PhD

Professor
Bogor Agricultural University
Indonesia
(External Examiner)



NOR AINI AB. SHUKOR, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 27 February 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Abd Wahid Haron, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Rosnina Yusoff, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Nurhussein Yimer, PhD

Senior Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No: Salman Hammadi Ghareeb, GS39883

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____

Name of
Chairman of
Supervisory
Committee:

Professor Dr. Abd Wahid Haron

Signature: _____

Name of
Member of
Supervisory
Committee:

Associate Professor Dr. Rosnina Yusoff

Signature: _____

Name of
Member of
Supervisory
Committee:

Dr. Nurhusein Yimer

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iv
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xvi
LIST OF FIGURES	xix
LIST OF ABBREVIATIONS	xx

CHAPTER

1	INTRODUCTION	1
	1.1 Background	1
	1.2 Problem statement	2
	1.3 Significance of the study	3
	1.4 Hypothesis:	3
	1.5 Study objectives	3
2	LITERATURE REVIEW	4
	2.1 History of Artificial Insemination	4
	2.2 Principle of semen cryopreservation	5
	2.2.1 Cryopreservation mechanism and methods	5
	2.3 Sperm characteristics and development	7
	2.3.1 Spermatogenesis	7
	2.3.2 Semen evaluation	8
	2.3.2.1 Volume of semen	8
	2.3.2.2 Colour of semen	8
	2.3.2.3 Sperm motility	8
	2.3.2.4 Sperm abnormalities	9
	2.4 Role of the extender	10
	2.4.1 Egg yolk extenders	11
	2.4.1.1 Egg yolk compositions	11
	2.4.1.2 Role of low density lipoproteins in cryopreservation	11
	2.5 Factors affecting quality of cryopreserved sperm	11
	2.5.1 Method of semen collection	11
	2.5.2 Semen Extenders and equilibrium time	12
	2.6 Bull sperm plasma membrane	13
	2.6.1 Sperm plasma membrane	13
	2.6.2 Sperm plasma membrane composition	13

2.6.3	The role of plasma membrane of the sperm	13
2.6.4	Seminal plasma removal	14
2.7	Oocyte collection and <i>in vitro</i> fertilization	14
2.7.1	Oocyte collection and evaluation	14
2.7.2	Oocyte vitrification	15
2.7.3	Oocyte development stages in cattle	15
2.8	Assisted Reproductive Technology (ART)	16
2.9	Role of open system and closed system on quality of vitrified oocytes	16
3	GENERAL MATERIALS AND METHODS	18
3.1	Study Approval	18
3.2	Animal management	18
3.3	Semen collection and preparation	18
3.3.1	Semen collection and storage	18
3.3.2	Semen extension	18
3.3.3	Semen quality evaluation	19
3.3.4	Post-thaw semen evaluation	20
3.3.4.1	Sperm morphology, viability and acrosome integrity	20
3.3.4.2	The hypo-osmotic swelling test (HOST)	20
3.3.4.3	DNA Integrity assessment	20
3.4	Preparation of sterilised egg yolk plasma	21
3.5	Oocyte Harvesting and vitrification	21
3.5.1.1	Oocyte Harvesting	21
3.5.2	Oocytes Vitrification	21
3.5.3	<i>In vitro</i> maturation	22
3.6	Oocytes evaluation	22
3.6.1	Determining of oocyte maturation	22
3.7	Statistical analysis	22
4	A COMPARATIVE STUDY ON CRYOPRESERVED BULLS' SEMEN QUALITY IN ANDROMED[®], BIOXCELL[®], TRILADYL[®] AND TRIS-EGG YOLK EXTENDERS	23
4.1	Introduction	23
4.2	Materials and Methods	24
4.3	Results	24
4.3.1	Sperm motility	24
4.3.2	Sperm viability	25
4.3.3	Sperm morphology	25
4.3.4	Sperm DNA Integrity	26
4.3.5	Sperm Plasma Membrane Integrity	29
4.3.6	Acrosome integrity	29
4.4	Discussion	30
4.5	Conclusions	32

5	EFFECT OF SEMINAL PLASMA REMOVAL ON CRYOPRESERVED BULL SEMEN QUALITY EXTENDED IN FOUR DIFFERENT EXTENDERS	33
5.1	Introduction	33
5.2	Materials and Methods	34
5.3	Results	34
5.3.1	Sperm motility	34
5.3.2	Sperm viability	35
5.3.3	Sperm morphology	35
5.3.4	Acrosome integrity	36
5.3.5	Sperm DNA Integrity	36
5.3.6	Sperm Plasma Membrane Integrity	37
5.4	Discussion	38
5.5	Conclusion	39
6	EFFECT OF EGG YOLK PLASMA IN TRIS-BASED EXTENDER ON POST THAW QUALITY OF EJACULATED BULL SPERMATOZOA	40
6.1	Introduction	40
6.2	Materials and methods	41
6.3	Results	41
6.3.1	Sperm motility	41
6.3.2	Sperm viability	42
6.3.3	Sperm morphology	42
6.3.4	Sperm acrosome Integrity	43
6.3.5	Sperm plasma membrane integrity	43
6.4	Discussion	44
6.5	Conclusion	45
7	EFFECT OF SEMINAL PLASMA REMOVAL AND EGG YOLK PLASMA ON POST THAW QUALITY OF BULL SPERMATOZOA	46
7.1	Introduction	46
7.2	Materials and Methods	47
7.3	Results	47
7.3.1	Sperm motility	47
7.3.2	Sperm Viability	47
7.3.3	Sperm Morphology	48
7.3.4	Acrosome integrity	48
7.3.5	Plasma membrane integrity	49
7.4	Discussion	49
7.5	Conclusion	52

8	EFFECT OF VITRIFICATION EXPOSURE TIME ON IMMATURE VITRIFIED BOVINE OOCYTES AND <i>IN VITRO</i> EMBRYO PRODUCTION	53
8.1	Introduction	53
8.2	Materials and methods	54
8.2.1	Ovaries collection and harvesting of bovine oocytes	54
8.2.2	Morphological evaluation of oocytes pre and post vitrification	54
8.2.3	Vitrification and warming of immature oocytes	54
8.2.4	Oocyte <i>in vitro</i> maturation (IVM)	55
8.2.5	Preparation of viable sperms	55
8.2.6	<i>In vitro</i> fertilization (IVF)	55
8.2.7	<i>In vitro</i> maturation and culture	56
8.2.8	Statistical analysis	56
8.3	Results	56
8.3.1	Harvesting rate of oocytes	56
8.3.2	Recovery rates of oocytes following vitrification-warming	58
8.3.3	Viability rates of oocytes post-vitrification	60
8.3.4	Vitrified-warmed oocyte subjected to <i>in vitro</i> maturation and fertilization	62
8.3.5	<i>In vitro</i> embryo development	63
8.4	Discussion	65
8.5	Conclusion	66
9	GENERAL DISCUSSION	67
10	SUMMARY, GENERAL CONCLUSION AND RECOMMENDATIONS FOR FUTURE WORK	76
	REFERENCES	80
	APPENDICES	100
	BIODATA OF STUDENT	101
	LIST OF PUBLICATIONS	102

LIST OF TABLES

Table		Page
4.1	Post-thaw sperm motility rates \pm SEM pre-extended in four different extenders with two different storage periods in Liquid Nitrogen (-196°C)	24
4.2	Post-thaw sperm viability rates \pm SEM pre-extended in four different extenders with two different storage periods in Liquid Nitrogen (-196°C)	25
4.3	Post-thaw sperm morphology rates \pm SEM pre-extended in four different extenders with two different storage periods in Liquid Nitrogen (-196°C)	26
4.4	Post-thaw sperm DNA integrity rates \pm SEM pre-extended in four different extenders with two different storage periods in Liquid Nitrogen (-196°C)	26
4.5	Post-thaw sperm plasma membrane integrity rates \pm SEM pre-extended in four different extenders with two different storage periods in Liquid Nitrogen (-196°C)	29
4.6	Post-thaw sperm acrosome integrity rates \pm SEM pre-extended in four different extenders with two different storage periods in Liquid Nitrogen (-196°C)	30
5.1	Post-thaw sperm motility rates pre-extended in four different extenders with removed seminal plasma in two different storage periods in Liquid Nitrogen (-196°C)	34
5.2	Post-thaw sperm viability pre-extended in four different extenders with removed seminal plasma in two different storage periods in Liquid Nitrogen (-196°C)	35
5.3	Post-thaw sperm morphology rates \pm SEM pre-extended in four different extenders with removed seminal plasma in two different storage periods in Liquid Nitrogen (-196°C)	36
5.4	Post-thaw sperm acrosome integrity rates \pm SEM pre-extended in four different extenders with removed seminal plasma in two different storage periods in Liquid Nitrogen (-196°C)	36
5.5	Post-thaw sperm DNA integrity rates \pm SEM pre-extended in four different extenders with removed seminal plasma in two different storage periods in Liquid Nitrogen (-196°C)	37

5.6	Post-thaw sperm plasma membrane integrity rates pre-extended in four different extenders with removed seminal plasma in two different storage periods in Liquid Nitrogen (-196°C)	37
6.1	Post-thaw sperm motility ratio pre-extended in four different egg yolk plasma based extender with seminal plasma in two different storage period in Liquid Nitrogen (-196°C)	41
6.2	Post-thaw sperm viability ratio pre-extended in four different egg yolk plasma based extender with seminal plasma in two different storage period in Liquid Nitrogen (-196°C)	42
6.3	Post-thaw sperm morphology ratio pre-extended in four different egg yolk plasma based extender with seminal plasma in two different storage period in Liquid Nitrogen (-196°C)	43
6.4	Post-thaw sperm acrosome integrity ratio pre-extended in four different egg yolk plasma based extender with seminal plasma in two different storage periods in Liquid Nitrogen (-196°C)	43
6.5	Post-thaw sperm plasma membrane integrity ratio pre-extended in four different egg yolk plasma based extender with seminal plasma in two different storage period in Liquid Nitrogen (-196°C)	44
7.1	Post-thaw sperm motility ratio pre-extended in four different egg yolk plasma based extender	47
7.2	Post-thaw sperm viability ratios pre-extended in four different egg yolk plasma based extender with removed seminal plasma	48
7.3	Post-thaw sperm morphology ratio pre-extended in four different egg yolk plasma based extender with removed seminal plasma in two different storage periods in Liquid Nitrogen (-196°C)	48
7.4	Post-thaw sperm acrosome integrity ratio pre-extended in four different egg yolk plasma based extender in two different storage period in Liquid Nitrogen (-196°C)	49
7.5	Post-thaw sperm plasma membrane integrity ratio pre-extended in four different egg yolk plasma based extender in two different storage periods in Liquid Nitrogen (-196°C)	49
8.1	Bovine oocytes quality grading rates \pm SEM harvested from abattoir ovaries	56
8.2	Oocytes recovery rates \pm SEM following vitrification-warming	59

8.3	<i>In vitro</i> oocytes viability rates \pm SEM following vitrification in two different cryodevices and in three different vitrification exposure durations	61
8.4	<i>In vitro</i> maturation rates \pm SEM of vitrified oocytes with two different cryodevices and with three different vitrification exposure durations	62
8.5	<i>In vitro</i> embryo development rates \pm SEM following <i>in vitro</i> maturation of vitrified oocytes in two different cryodevice and in three different vitrification exposure durations	64



LIST OF FIGURES

Figure	Page	
2.1	Illustrating various phases involved in spermatogenesis	7
2.2	Structure of a spermatozoon	9
2.3	The various forms of abnormalities seen in semen	10
2.4	The various oocyte developmental stages	16
4.1	Plane slide of frozen-thawed bull sperms extended in AndroMed, prepared for DNA assessment under epifluorescent microscopy	27
4.2	Second stage of DNA assessment using DAPI filter under epifluorescent microscopy	27
4.3	Third stage of DNA assessment using FITC filter under epifluorescent microscopy	28
4.4	Fourth stage of DNA assessment (same sample in Fig.2) after merging slides in Fig.11 and slide in Fig.12 under epifluorescent microscopy.	28
8.1	Grade 1 oocyte (Right-Up), Grade 4 oocyte (Left-down) under inverted microscope.200x	57
8.2	Oocyte COCs surrounded with more than 5-6 cells layers.200x	57
8.3	Expansion of COCs cells post <i>In vitro</i> maturation	58
8.4	Vitrified-warmed immature bovine oocyte with unruptured clear zona pelucida, increase in previtteline space and granulated cytoplasm, 600x	59
8.5	Completely denuded oocytes post invitro maturation, 200x	60
8.6	Vitrified-warmed immature bovine oocyte with unruptured clear zona pelucida and homogenous cytoplasm. 600x	61
8.7	<i>In vitro</i> mature oocyte observed after 22 h of maturation. (First polar body) many sperm attaching to the zonapellucida .600x	63
8.8	<i>In vitro</i> bovine embryo development (two cells embryo) 2 days post <i>in vitro</i> culture.600x	64

LIST OF ABBREVIATIONS

AO	Acridin Orange
AV	Artificial vagina
AEE	Automatic electro-ejaculator
CFS	Closed French straws
CPA	Cryoprotectants
EE	Electro-ejaculator
EG	Ethelen glycol
ES	Equilibrium solution
EYP	Egg yolk plasma
IVP	Invitro embryo production
LDL	Low density lipoprotein
OPS	Open pulled straws
PKC	Palm kernel cake
PMI	Plasma plasma integrity
SP	Seminal plasma
TCM	Tissue culture medium
TWEY	Tris whole egg yolk
TEYP	Tris egg yolk plasma
VS	Vitrification solution
WS	Warming solution

CHAPTER 1

INTRODUCTION

1.1 Background

Over the decades, history has recorded a tremendous revolution in animal breeding where difficulties in breeding certain desired breeds became inevitably exploited through the revolutionary scientific achievement in the field of Theriogenology where semen transportation became widely accepted (Aires et al., 2003). To achieve this, the application of cryopreservation was applied in breeding desired cattle breeds as farmers all over indulged in the practice of artificial insemination (AI) (Sathe & Shipley, 2015). In Malaysia to be precise, artificial insemination as a technique was practiced as early as 1963 when frozen semen was transported from Australia to Singapore and later railed into Malaysia (Zaman, 2013). Cattle breeding had become widely spread amongst animal farmers as a result of the hike in the per capita consumption of beef, in Malaysia it was reported by Abd Wahid Haron in 2013 to have elevated from about 5kg to 5.7kg in a space of a decade (2001-2010). The viability of semen became rested on the ability to lower its metabolic rates through reduced temperature. Extreme lower temperatures therefore provided a decline in semen metabolic activities thereby making the semen last longer if preserved (Parrish et al., 1986). Cremades in 2005 reported that apart from reduced temperatures, carbon dioxide also poses prolong preservatory property on semen but the success was however considered insignificant (Cremades et al., 2005). Lower temperatures of about 4-7°C have prolonged the survival span of semen and retained semen motility for almost seventy two hours with an appreciable fertility potentials. This technique explains how semen could be preserved; however, semen integrity, reduced semen motility, variation in viability and rates of conception became an outcry.

Cryopreservation was reported in 2000 to be affecting semen viability by almost 53% and eventually posing a detrimental effect on fertilisation to a tune of about seven ties. This necessitated a poll of experimental studies over the years as to the extent of the effects on the anatomy and physiology of semen particularly in cattle breeding. Researchers have pinned down factors like decondensation of chromatin, premature reaction of acrosome and impeded physiology of mitochondria to most of the in vitro experimental studies (Watson, 2000). About two decades ago, reduced extreme temperatures have been indicated to be directly proportional to semen stressing, this phenomenon was suggested to be a membrane disruptive factor which can lead to a condition called 'cold shock' (Kim et al., 2011). Semen cold shock syndrome has direct effect on the liveability of semen depending on the severity of the shock (Drobnis et al., 1993). This reason many trials involving addition some forms of cryoprotective agents like egg yolk, milk, bovine serum albumin, liposomes and glycerol in extenders have conferred some level of longitivity to the semen against the adverse effects of cryopreservation (Shoae & Zamiri, 2008). The advent of assisted reproductive biotechnology is another big achievement in the world of science

that gave a wider window to the improvement of breeding and reproduction (Haron, 2013).

Immature egg cells known as the oocytes have also been cryopreserved over time for the purpose of breeding (Haron, 2013). Oocytes have been reported to be very difficult to preserve because it presents with a very low fertility results which is mostly associated with the physical changes in the composition of oocytes, expansion and contraction of the oocytes and careless handling of oocytes during cryopreservation or vitrification (Kuwayama, 2007). However positive reports have been recorded in the application of cryopreserved oocytes though a lot of augments have proven the report unacceptable (Vieira et al., 2002).

Cryopreserved sperm and vitrified oocytes as a gametes can proceed to produce embryos either *in vivo* or *in vitro* successfully. Embryo quality in cattle breeding is also a very important outlook that determines the success of any breeding industry. Records had it that almost 20% of bovine embryo are widely produced using the *in vitro* technology popularly known as the *In vitro* Embryo Production (IVP) (Wheeler et al., 2006). Embryo transfer is a widely accepted practice extremely dependent on the quality of cryopreserved sperms and vitrified oocytes. Initially surgery was the known mode of embryo transfer but in the recent years non-surgical transfer of embryo have been exploited. Therefore, preserving the semen quality and vitrified oocyte quality, fertility rate post-thaw will be a key factor to improving embryo transfer quality and hence cattle breeding thereby providing animal farmers with a better understanding and options to maximise profit through animal production.

1.2 Problem statement

A decline in cattle breeding had been experienced in the past years in Malaysia with a fluctuating pattern. Semen extenders as a media used to handle semen in cryopreservation have played important role in reproductive efficiencies in farm animals. In recent times, the bull semen quality due to anticipated variation in the protective efficiency of semen extenders have been worrisome coupled with the fact that seminal plasma have been argued to make negative effect on the cryopreserved semen. Egg yolk preserved extender had been of poor quality lately in many reports attributed to its microbial contamination, turbidity and granulation. Immature oocytes vitrification have a great value in oocytes preserving banks. Both sperm cryopreservation and oocyte vitrification still need to be improved to get more superior *in vitro* embryo production. Quality of the oocytes after vitrification-warming is still another challenge may decline the quality of oocyte and embryo as well. Cryodevice and of oocyte exposure time to vitrification solution still a matter of debate to optimise vitrified oocyte quality through reducing cryoprotectant-oocyte toxicity.

1.3 Significance of the study

The quality and fertility capacity of cryopreserved semen and vitrified immature oocytes is very crucial to any successful breeding program as it reflects directly the quality of embryos produced. Therefore, firstly, this study is significantly aims to provide information on using of seven different semen extenders, semen cryopreservation, effect of removal of seminal plasma on the sperm quality, immature oocyte vitrification process, vitrification exposure time, cryodevices used in preserving of vitrified oocytes, *in vitro* fertilization, *in vitro* maturation, *in vitro* embryo culture and *in vitro* embryo production in cattle . Secondly, our study aims to find out the causes resulting in declining sperm quality, oocyte quality and embryo quality as well. Lastly, the present study aims the improvement in cattle breeding knowledge.

1.4 Hypothesis:

- A. H_0 = Assessment of cryopreservatives and oocyte vitrification improve *in vitro* embryo production in cattle.
 H_a = Assessment of cryopreservatives and oocyte vitrification doesnot improve *in vitro* embryo production in cattle

1.5 Study objectives

To evaluate the post-thaw bull spems pre-extended in: AndroMed®, BioXcell®, Triladyl® compared to Tris-egg yolk based extenders 20%.

1. To assess the effect of seminal plasma removal on the post-thaw bull sperms pre-extended in: AndroMed®, BioXcell®, Triladyl® compared to Tris-egg yolk based extenders 20%.
2. To explain the role of different concentration of egg yolk plasma (10%, 15% and 20%) on post-thaw bull sperm quality pre-extended with seminal plasma.
3. To esimate the role of different concentration of egg yolk plasma (10%, 15% and 20%) extenders on post-thaw bull sperm quality pre-extended without seminal plasma.
4. To illustrate the effect of different vitrification exposure time (15 sec, 30 sec and 45 sec) and different cryodevices (OPS/ CFS) on the quality of vitrified immature oocytes (harvested from slaughtered cattle) for *in vitro* embryo production in cattle.

REFERENCES

- Ahmad, M. K., Mahdi, A. A., Shukla, K. K., Islam, N., Rajender, S., Madhukar, D., Shankhwar, S. N. and Ahmad, S. (2010). *Withania somnifera* improves semen quality by regulating reproductive hormone levels and oxidative stress in seminal plasma of infertile males. *Fertility and sterility* 94(3), 989-996.
- Ahmad, M., A. Khan, Z. Shah and K. Ahmad, (1996). Effects of removal of seminal plasma on the survival rate of buffalo bull spermatozoa. *Animal reproduction science* 41(3), 193-199.
- Ahmed, M.-K. A. b. and Ishag, I. (2016). Suggestions for Genetic Improvement of Camels in Sudan.
- Aires, V. A., Hinsch, K.-D., Mueller-Schloesser, F., Bogner, K., Mueller-Schloesser, S. and Hinsch, E. (2003). In vitro and in vivo comparison of egg yolk-based and soybean lecithin-based extenders for cryopreservation of bovine semen. *Theriogenology* 60(2), 269-279.
- Aitken R.J and Nixon, B. (2013). Sperm capacitation: A distant landscape glimpsed but unexplored. *Molecular Human Reproduction* 19:785–93.
- Aitken R.J., De luliis, G.N. and Mclachlan, R.I. (2009). Biological and Clinical Significance of DNA Damage in the Male Germ Line. *International Journal of Andrology* 32:46-56.
- Akhter, S., Ansari, M., Rakha, B., Andrabi, S., Iqbal, S. and Ullah, N. (2010). Cryopreservation of buffalo (*Bubalus bubalis*) semen in Bioxcell® extender. *Theriogenology* 74(6), 951-955.
- Albano, C., Felberbaum, R., Smitz, J., Riethmüller-Winzen, H., Engel, J., Diedrich, K., Devroey, P. and Group, E. C. S. (2000). Ovarian stimulation with HMG: results of a prospective randomized phase III European study comparing the luteinizing hormone-releasing hormone (LHRH)-antagonist cetrorelix and the LHRH-agonist buserelin. *Human Reproduction* 15(3), 526-531.
- Aman, R.R. and Parks, J.E. (1994). Effects of cooling and rewarming on the meiotic spindle and chromosomes of *in vitro* matured bovine oocytes. *Biology of Reproduction* 50:103–10.
- Amann, R. and Pickett, B. (1987). Principles of cryopreservation and a review of cryopreservation of stallion spermatozoa. *Journal of equine veterinary science* 7(3), 145-173.
- Amirat L., Bencharif, D. Vera-Munoz, O. Pineau, S. Thorin, C. Destrumelle, S. Desherces, S. Anton M. , Jouan M. and Shmitt, E. (2010). *In vivo* Fertility of Bull Semen Following Cryopreservation with an LDL (low density lipoprotein)

Extender: Preliminary Results of Artificial Inseminations. *Animal Reproduction Science* 122(3), 282-287.

Amirat, L., Anton, M., Tainturier, D., Chatagnon, G., Battut, I. and Courtens, J. L. (2005). Modifications of bull spermatozoa induced by three extenders: Biociphos, low density lipoprotein and Triladyl, before, during and after freezing and thawing. *Reproduction* 129(4), 535-543.

Andrabi, S. and Maxwell, W. (2007). A review on reproductive biotechnologies for conservation of endangered mammalian species. *Animal reproduction science* 99(3), 223-243.

Anton, M. and Gandemer, G. (1997). Composition, solubility and emulsifying properties of granules and plasma of egg yolk. *Journal of Food Science* 62(3), 484-487.

Arangasamy, A., Kasimanickam, V., DeJarnette, J. and Kasimanickam, R. (2011). Association of CRISP2, CCT8, PEBP1 mRNA abundance in sperm and sire conception rate in Holstein bulls. *Theriogenology* 76(3), 570-577.

Arav, A. (1992). Vitrification of oocytes and embryos. In: Lauria, A., Gandolfi, F. (Eds.), *New Trends in Embryo Transfer*. Portland Press, Cambridge, pp. 255–264.

Arav, A., Zeron, Y., Leslie, S.B., Behboodi, E., Anderson, G.B., Crowe, J.H. (1996). Phase transition temperature and chilling sensitivity of bovine oocytes. *Cryobiology* 33:589–99.

Asadpour, R., Alavi-Shoushtari, S., Rezaii, S. A. and Ansari, M. K. (2007). SDS-polyacrylamide gel electrophoresis of buffalo bull's seminal plasma proteins and their relation with semen freezability. *Animal reproduction science* 102(3), 308-313.

Aydin, R., and Dogan, I. (2010). Fatty Acid Profile and Cholesterol Content of Egg Yolk from Chickens Fed Diets Supplemented with Purslane (*Portulaca oleracea* L.). *Journal Science and Food Agriculture* 90 (10), 1759–1763.

Baas, J. W., Molan, P. and Shannon, P. (1983). Factors in seminal plasma of bulls that affect the viability and motility of spermatozoa. *Journal of reproduction and fertility* 68(2), 275-280.

Bailey J.L. (2010). Factors regulating sperm capacitation. *Systematic Biological Reproductive Medecine* 56:334–48.

Bailey, J., Morrier, A. and Cormier, N. (2003). Semen cryopreservation: Successes and persistent problems in farm species. *Canadian journal of animal science* 83(3), 393-401.

- Barrios, B., Pérez-Pé, R., Gallego, M., Tato, A., Osada, J., Muiño-Blanco, T. and Cebrián-Pérez, J. A. (2000). Seminal plasma proteins revert the cold-shock damage on ram sperm membrane. *Biology of reproduction* 63(5), 1531-1537.
- Bathgate, R. (2008). Functional integrity of sex-sorted, frozen-thawed boar sperm and its potential for artificial insemination. *Theriogenology* 70(8), 1234-1241.
- Bee, L. and Cotterill, O. (1979). Ion- Exchange Chromatography and Electrophoresis of Egg Yolk Proteins. *Journal of Food Science*, 44(3), 656-667.
- Bencharif D., Amirat-Briand, L. Garand, A. Anton, M. Schmitt, E. Desherces, S. Delhomme, G. Langlois, M.L. Barrière, P. and Destrumelle, S. (2010). Freezing Canine Sperm: Comparison of Semen Extenders Containing Equex® and LDL (Low Density Lipoproteins). *Animal Reproduction Science* 119(3): 305-313.
- Beran, J., Stádník, L., Bezdíček, J., Louda, F., Čítek, J. and Ducháček, J. (2012). Effect of sire and extender on sperm motility and share of live or dead sperm in bulls' fresh ejaculate and in AI doses after thawing. *Archives Animal Breeding* 55(3): 207-218.
- Binta, H. (1994). A Study of Some Factors That May Affect Semen Quantity and Quality as Well as Libido in Four Breeds of Dairy Bulls. University of Nairobi.
- Birkhead, T. R., Hosken, D. J. and Pitnick, S. S. (2008). In: Sperm biology: an evolutionary perspective: Academic press.
- Bucak, M. N., Tuncer, P. B., Sariözkan, S., Başpınar, N., Taşpınar, M., Çoyan, K., Bilgili, A., Akalın, P. P., Büyükleblebici, S. and Aydos, S. (2010). Effects of antioxidants on post-thawed bovine sperm and oxidative stress parameters: antioxidants protect DNA integrity against cryodamage. *Cryobiology* 61(3): 248-253.
- Büyükleblebici, S., Tuncer, P.B., Bucak, M.N., Eken, A.S. Sariözkan, Tas, demir U., et al., (2014). Cryopreservation of Bull Sperm: Effects of Extender Supplemented with Different Cryoprotectants and Antioxidants on Sperm motility, antioxidant capacity and fertility results. *Animal Reproduction Science* 150:77–83.
- Byrne, B., Gruber, M. and Ab, G. (1989). The evolution of egg yolk proteins. *Progress in biophysics and molecular biology* 53(1): 33-69.
- Byrne, C.J., Fair, S., English, A.M., Holden, S.A., Dick J.R., Lonergan, P. and Kenny D.A. (2017). Dietary polyunsaturated fatty acid supplementation of young postpubertal dairy bulls alters the fatty acid composition of seminal plasma and spermatozoa but has no effect on semen volume or sperm quality. *Theriogenology* 90: 289-300.

- Caamaño, J.N., Díez, C., Trigal, B., Muñoz, M., Morató, R., Martín, D., Carrocera, S., Mogas, T. and Gomez, E. (2013). Assessment of meiotic spindle configuration and postwarming bovine oocyte viability using polarized light microscopy. *Reproduction in Domestic Animals* 48: 470–476.
- Cabrera, F., Gonzalez, F., Batista, M., Calero, P., Medrano, A., & Gracia, A. (2005). The effect of removal of seminal plasma, egg yolk level and season on sperm freezability of canary buck (*Capra hircus*). *Reproduction in Domestic Animals* 40(3): 191-195.
- Casillas, F., Ducolomb, Y., Lemus, A.E., Cuello, C. and Betancourt, M., (2015). Porcine embryo production following in vitro fertilization and intracytoplasmic sperm injection from vitrified immature oocytes matured with a granulosa cell co-culture system. *Cryobiology* 71: 299–305.
- Celeghini, E. C. C., de Arruda, R. P., de Andrade, A. F. C., Nascimento, J., Raphael, C. F. and Rodrigues, P. H. M. (2008). Effects that bovine sperm cryopreservation using two different extenders has on sperm membranes and chromatin. *Animal Reproduction Science* 104(2): 119-131.
- Chandolia, R., Honaramooz, A., Omeke, B., Pierson, R., Beard, A. and Rawlings, N. (1997). Assessment of development of the testes and accessory glands by ultrasonography in bull calves and associated endocrine changes. *Theriogenology* 48(1): 119-132.
- Chao, N.-H., Chen, H.-P. and Liao, I.-C. (1975). Study on cryogenic preservation of grey mullet sperm. *Aquaculture* 5(4): 389-406.
- Chen, S.U., Lien, Y.R., Chao, K.H., Ho, H.N., Yang, Y.S. and Lee, T.Y. (2003). Effects of cryopreservation on meiotic spindles of oocytes and its dynamics after thawing: clinical implications in oocyte freezing- a review article. *Molecular Cell Endocrinology* Apr 28; 202(1-2):101-7.
- Choi, J. K., Yue, T., Huang, H., Zhao, G., Zhang, M., He, X. (2015). The crucial role of zona pellucida in cryopreservation of oocytes by vitrification. *Cryobiology* 71: 350–5.
- Clemente-Sánchez, F., Cessa-Reyes, V., Cortez-Romero, C. Tarango-Arambula, L.A. and Arenas-Baez, P. (2014). Commercial Extenders and Freezing Curves for the Preservation of Sperm Cells of White-tailed Deer (*Odocoileus virginianus*). *Journal of Application in Animal Research* 1: 1–6.
- Clulow, J., Mansfield, L., Morris, L., Evans, G., & Maxwell, W. (2008). A comparison between freezing methods for the cryopreservation of stallion spermatozoa. *Animal Reproduction Science* 108(3): 298-308.

- Colenbrander, B., Gadella, B. M. and Stout, T. A. (2003). The predictive value of semen analysis in the evaluation of stallion fertility. *Reproduction Domestic Animals* 38(4):305-311.
- Cordova, A., Uzbekova, S., Ponsart, C., Locatelli, Y. and Mermillod, P. (2014). Development rate and gene expression of IVP bovine embryos cocultured with bovine oviduct epithelial cells at early or late stage of preimplantation development. *Theriogenology* 81: 1163–1173.
- Coticchio G, Bonu MA, Borini A, Flamigni C. (2004). Oocyte cryopreservation: a biological perspective. *European Journal of Obstetrics and Gynecology of Reproductive Biology* 115:2–7.
- Courtens, J.L. and Rety, J.M. (2001). Numerical simulation for freezing and thawing mammalian spermatozoa. Evaluation of cell injuries at different depths in bags or straws during all steps of the technique. *Genetics Selection Evolution* 33, S85-S104.
- Cox, G. F., Bürger, J., Lip, V., Mau, U. A., Sperling, K., Wu, B.-L. and Horsthemke, B. (2002). Intracytoplasmic sperm injection may increase the risk of imprinting defects. *The American Journal of Human Genetics* 71(1), 162-164.
- Cremades, T., Roca, J., Rodriguez- Martinez, H., Abaigar, T., Vazquez, J. M. and Martinez, E. A. (2005). Kinematic changes during the cryopreservation of boar spermatozoa. *Journal of andrology* 26(5), 610-618.
- Dellenbach, P., Nisand, I., Moreau, L., Feger, B., Plumere, C. and Gerlinger, P. (1985). Transvaginal sonographically controlled follicle puncture for oocyte retrieval. *Fertility and sterility*, 44(5): 656-662.
- Desai, A., Shirode, A. R., Mittal, B. and Kadam, V. J. (2011). Assisted Reproductive Technology (ART): Combating Infertility. *Asian Journal of Pharmaceutical and Clinical Research* 4(1).
- Desnoyers, L. and Manjunath, P. (1992). Major proteins of bovine seminal plasma exhibit novel interactions with phospholipid. *Journal of Biological Chemistry* 267(14):10149-10155.
- Devireddy, R., Swanlund, D., Olin, T., Vincente, W., Troedsson, M., Bischof, J. and Roberts, K. (2002). Cryopreservation of equine sperm: optimal cooling rates in the presence and absence of cryoprotective agents determined using differential scanning calorimetry. *Biology of Reproduction* 66(1): 222-231.
- Diez, C., Muñoz, M., Caamaño, J.N. and Gomez, E. (2012). Cryopreservation of the bovine oocyte: current status and perspectives. *Reproduction in Domestic Animals* 47: 76–83.

- Dowglish, F. C., Emilie C., Carmen A., Yann L., Joanna (2017). Vitrification of immature and *in vitro* matured bovine cumulus-oocyte complexes: Effects on oocyte structure and embryo development. *Livestock Science* 199: 50–56.
- Dresser, B. L. (1988). Cryobiology, embryo transfer, and artificial insemination in ex situ animal conservation programs. *Biodiversity* 296.
- Drobnis, E. Z., Crowe, L. M., Berger, T., Anchordoguy, T. J., Overstreet, J. W. and Crowe, J. H. (1993). Cold shock damage is due to lipid phase transitions in cell membranes: a demonstration using sperm as a model. *Journal of Experimental Zoology* 265(4): 432-437.
- Druart, X., Rickard, J.P., Mactier, S., Kohnke, P.L., Kershaw-Young, C.M., Bathgate, R., Gibb, Z., Crossett, B., Tsikis, G., Labas, V., Harichaux, G., Grupen, C.G. and de Graaf, S.P. (2013). Proteomic characterization and crossspecies comparison of mammalian seminal plasma. *Journal Proteomics* 91C: 13-22.
- Duru, N.K., M.S. Morshedi, A. Schuffner and S. Oehninger, (2001). Cryopreservation- Thawing of fractionated human spermatozoa is associated with membrane phosphatidylserine externalization and not DNA fragmentation. *Journal of Andrology* 22: 646-651.
- Dziuk, P., Graham, E., & Petersen, W. (1954). The technique of electroejaculation and its use in dairy bulls. *Journal of Dairy Science*, 37(9): 1035-1041.
- El-Moemen, S. A. (2012). International Workshop on" Recent Strategies in Animal. Alexandria University.
- Ezoe, K., Yabuuchi, A., Tani, T., Mori, C., Miki, T., Takayama, Y., Beyhan, Z., Kato, Y., Okuno, T., Kobayashi, T. and Kato, K., (2015). Developmental competence of vitrified warmed bovine oocytes at the germinal vesicle stage is improved by cyclic adenosine monophosphate modulators during *in vitro* maturation. PLoS One 10, 0126801.
- Ferreira, E.M., Vireque, A.A., Adona, P.R., Meirelles, F.V., Ferriani, R.A. and Navarro, P.A., (2009). Cytoplasmic maturation of bovine oocytes: structural and biochemical modifications and acquisition of developmental competence. *Theriogenology* 71, 836–848.
- Fleisch A., Malama, E., Witschi, U., Leiding, C., Siuda, M., Janett, F. and Bollwein, H. (2017). Effect of an Axtension of the Equilibration Period up to 96 hours on the Characteristics of Cryopreserved Bull Semen. *Theriogenology* 89: 255-262.
- Flesch, F. M. and Gadella, B. M. (2000). Dynamics of the mammalian sperm plasma membrane in the process of fertilization. *Biochimica et Biophysica Acta (BBA)-Reviews on Biomembranes* 1469(3): 197-235.

- Foote, R. H. (2002). The history of artificial insemination: Selected notes and notables. *Journal of Animal Science*, 80: 1-10.
- Ford, W. (2006). Glycolysis and sperm motility: does a spoonful of sugar help the flagellum go round? *Human Reproduction Update*, 12(3) 269-274.
- Fraser, L. R., Abeydeera, L. and Niwa, K. (1995). Ca²⁺- Regulating mechanisms that modulate bull sperm capacitation and acrosomal exocytosis as determined by chlortetracycline analysis. *Molecular Reproduction and Development* 40(2): 233-241.
- Fuller, B., Paynter, S. (2004). Fundamentals of cryobiology in reproductive medicine. *Reproductive Biomedicine*, Online 9, 680–691.
- Gadea, J. (2003). Review: semen extenders used in the artificial insemination of swine. *Spanish Journal of Agricultural Research* 1(2): 17-27.
- Galli, C., Crotti, G., Notari, C., Turini, P., Duchi, R. and Lazzari, G. (2001). Embryo production by ovum pick up from live donors. *Theriogenology* 55(6): 1341-1357.
- Galli, C., Duchi, R., Crotti, G., Turini, P., Ponderato, N., Colleoni, S., Lagutina, I., and Lazzari, G. (2003). Bovine Embryo Technologies. *Theriogenology* 59(2): 599-616.
- Gangwar, D.K. and Atreja, S.K. (2015). Signalling events and associated pathways related to the mammalian sperm capacitation. *Reproduction in Domestic Animals* 50:705–11.
- Gao, D., Liu, J., Liu, C., McGann, L., Watson, P., Kleinhans, F., Mazur, P., Critser, E. and Critser, J. (1995). Andrology: Prevention of osmotic injury to human spermatozoa during addition and removal of glycerol. *Human Reproduction* 10(5): 1109-1122.
- Garcia, M. and Graham, E. (1987). Effects of low-molecular-weight fractions (LMWF) from milk, egg yolk, and seminal plasma on freezability of bovine spermatozoa. *Cryobiology* 24(5): 429-436.
- Ghetler, Y., Yavin, S., Shalgi, R. and Arav, A. (2005). The effect of chilling on membrane lipid phase transition in human oocytes and zygotes. *Human Reproduction*, 20,3385–3389.
- Gil, J., Lundeheim, N., Söderquist, L. and Rodríguez-Martínez, H. (2003). Influence of extender, temperature, and addition of glycerol on post-thaw sperm parameters in ram semen. *Theriogenology* 59(5): 1241-1255.
- Gook, D.A., Osborn, S.M. and Johnston, W.I. (1993). Cryopreservation of mouse and human oocytes using 1, 2-propanediol and the configuration of the meiotic spindle. *Human Reproduction* 8:1101–9.

- Graham, J. K. (1996). Analysis of stallion semen and its relation to fertility. *Veterinary Clinics of North America-Equine Practice* 12(1):119-130.
- Graham, J.K. and Mocé, E. (2005). Fertility evaluation of frozen/thawed semen. *Theriogenology* 64:492–504.
- Gurler, H., Melama E., Heppelmann, M., Calisici, O., Leiding, C., Kastelic, J.P., and Bollwein, H. (2016). Effect of Cryopreservation on Sperm Viability, Synthesis of Reactive Oxygen Species and DNA Damage of bovine sperm. *Theriogenology* 86: 562-571.
- Hafez, E. S. E. (1993). In: Artificial insemination. Philadelphia: Lea and Febiger.
- Hafez, E. S. E. and Hafez, B. (2000). In: Reproduction in Farm Animals: Wiley.
- Halasz, N. A. and Collins, G. M. (1984). Studies in cryoprotection II: propylene glycol and glycerol. *Cryobiology* 21(2): 144-147.
- Hamamah, S. and Gatti, J.-L. (1998). Role of the ionic environment and internal pH on sperm activity. *Human Reproduction* 13(suppl 4): 20-30.
- Hammer, C. E. (1970). In: The Semen. Reproduction and Breeding Techniques for Laboratory Animals (E. S. E. Hafez Ed.). Philadelphia: Lea and Febiger.
- Hammerstedt, R., Graham, J. K. and Nolan, J. P. (1990). Cryopreservation of mammalian sperm: what we ask them to survive. *J Androl*, 11(1), 73-88.
- Haron, A. W. (2013). In: Livestock breeding: the past, the present and the future.
- Herold F., Aurich J., and Gerber D. (2004). Epididymal Sperm from the African Buffalo (*Syncerus caffer*) can be Frozen Successfully with AndroMed® and with Triladyl™ but the Addition of Bovine Seminal Plasma is Detrimental. *Theriogenology* 61(4), 715-724.
- Herold, F.C., Hass, K., Colenbrander, B. and Gerber, D. (2006). Comparasion of Equilibration Times when Freezing Epididymal Sperm from African (*Syncerus caffer*) using Triladyl™ or Andromed®. *Theriogenology* 66:1123-1130.
- Hoflack, G., Van Soom, A., Maes, D., de Kruif, A., Opsomer, G. and Duchateau, L. (2006). Breeding soundness and libido examination of Belgian Blue and Holstein Friesian artificial insemination bulls in Belgium and The Netherlands. *Theriogenology* 66(2): 207-216.
- Holden, S.A., Fernandez-Fuertes B., Murphy C., Whelan H., O’Gorman A., Brennan L., Butler S.T. , Lonergan P. and Fair S. . (2017). Relationship between in vitro sperm functional assessments, seminal plasma composition, and field fertility after AI with either non-sorted or sex-sorted bull semen. *Theriogenology* 87: 221–228.

- Hoque, M., Talukder, A. K., Kamal, M., Jha, A. K., Bari, F. Y. and Shamsuddin, M. (2011). Ovulation Synchronization in Water Buffaloes Guided by Milk Progesterone ELISA. *Journal of Embryo Transfer* 26(2): 105-109.
- Hu J.H., L.S. Zan, X.L. Zhao, Q.W. Li, Z.L. Jiang, Y.K. Li and et al. (2010). Effects of trehalose supplementation on semen quality and oxidative stress variables in frozen-thawed bovine semen. *Journal of Animal Science* 88: 1657–62.
- Hurttt, A.E., Landim-Alvarenga, F., Seidel, Jr G.E. and Squires, E.L. (2000). Vitrification of immature and mature equine and bovine oocytes in an ethylene glycol, ficoll and sucrose solution using open pulled straws. *Theriogenology* 54:119–28.
- Izadyar, F., Matthijs-Rijsenbilt, J., Ouden, K., Creemers, L. B., Woelders, H. and de Rooij, D. G. (2002). Development of a cryopreservation protocol for type A spermatogonia. *Journal of Andrology* 23(4): 537-545.
- Jain, Y. and Anand, S. (1976). The lipids of buffalo spermatozoa and seminal plasma. *Journal of reproduction and fertility*, 47(2), 255-260.
- Jainudeen, M. R., Wahid, H. and Hafez, E. S. E. (2000). In: *Sheep and goats Reproduction*. 3rd Ed.
- Jiménez-Trigos, E., Naturil-Alfonso, C., Vicente, J. and Marco-Jiménez, F. (2012). Effects of cryopreservation on the meiotic spindle, cortical granule distribution and development of rabbit oocytes. *Reproduction in Domestic Animals* 47: 472–478.
- Johnson, W. H. (1997). The significance to bull fertility of morphologically abnormal sperm. *Veterinary Clinic in North America for Food and Animal Practice* 13(2): 255-270.
- Juyena, N. S. and Stelletta, C. (2012). Seminal plasma: an essential attribute to spermatozoa. *Journal of Andrology* 33(4): 536-551.
- Kaka A., Wahid, H., Rosnina, Y., Yimer, N., Khumran, A.M., Sarsaifi, K. Atique, A.B. Ubedullah, K. and Ebrahimi, M. (2015). α -Linolenic Acid Supplementation in BioXcell® Extender can Improve the Quality of Post-cooling and Frozen-thawed Bovine Sperm. *Animal Reproduction Science* 153: 1–7.
- Kasai, M. and Mukaida, T. (2004). Cryopreservation of animal and human embryos by vitrification. *Reproductive Biomedicine*. Online 9, 164–170.
- Katila, T. (2001). Sperm-uterine interactions: a review. *Animal Reproduction Science* 68(3-4): 267-272.

- Kaya, A., Aksoy, M. and Tekeli, T. (2002). Influence of ejaculation frequency on sperm characteristics, ionic composition and enzymatic activity of seminal plasma in rams. *Small Ruminant Research* 44(2): 153-158.
- Khan, M. I. R. and Ijaz, A. (2008). Effects of osmotic pressure on motility, plasma membrane integrity and viability in fresh and frozen-thawed buffalo spermatozoa. *Animal Reproduction Sciences* 2: 548-553.
- Khan, R., Rahman, Z.-u., Javed, I. and Muhammad, F. (2012). Effect of vitamins, probiotics and protein on semen traits in post-molt male broiler breeders. *Animal reproduction science* 135(1), 85-90.
- Khatun, M., Bhuiyan, M.M., Ahmed, J.U., Haque, A., Rahman, M.B., Shamsuddin, M. (2011). *In vitro* maturation and fertilization of prepubertal and pubertal black Bengal goat oocytes. *Journal of Veterinary Science* 12: 75–82.
- Khumran, A., Yimer, N., Rosnina, Y., Ariff, O., Wahid, H., Ebrahimi, M., Kaka, A. and Mahre, M. (2015). Evaluation of Physical and Ultra-structural attributes of Bulls' Semen with Variable Freezing Potential. *Malaysian Journal of Animal Science* 18(2): 55-65.
- Kim, D., Park, H., Kim, S., Hwang, I., Yang, B., IM, G., Chung, H., Seong, H., Moon, S. and Yang, B., (2007). Vitrification of immature bovine oocytes by the microdrop method. *Journal of Reproduction Devevelopment*. 53, 843–851.
- Kim, S., Lee, Y.J. and Kim, Y.-J. (2011). Changes in sperm membrane and ROS following cryopreservation of liquid boar semen stored at 15°C. *Animal Reproduction Science* 124(1): 118-124.
- Kovac, J.R., Pastuszak, A.W., Lamb, D.J. (2013). The use of genomics, proteomics, and metabolomics in identifying biomarkers of male infertility. *Fertility and Sterility* 99:998–1007.
- Kozdrowski, R., A., Dubiel, W. Bielas and Dzięcioł M., (2007). Two protocols of cryopreservation of goat semen with the use of computer-assisted semen analysis system. *Acta Veterinaria Brno* 76(4), 601-604.
- Kraus, M., M. Ticha, B. Zelezna, J. Peknicova and Jonakova V. (2005). Characterization of human seminal plasma proteins homologous to boarAQN spermadhesins. *Journal of Reproductive Immunology* 65:33–46.
- Kukov, A., Daskalova, D., Ivanova-Kicheva, M., Stefanov, R. and Neicheva, A. (2009). Immunocytochemical localization of separated seminal plasma proteins on the ram's sperm plasma membrane during in vitro preservation at 4°C. *Biotechnology in Animal Husbandry*, 25(5-6-2), 925-934.

- Kumar, P., Saini, M. Kumar D., Balhara A.K., Yadav S.P., Singh P. and Yadav P.S. (2015). Liposome -Based Semen Extender is Suitable Alternative to Egg Yolk-Based Extender for Cryopreservation of Buffalo (*Babalu babalus*) Semen. *Animal Reproduction Science* 159: 38-45.
- Kuwayama, M. (2007). Highly efficient vitrification for cryopreservation of human oocytes and embryos: the Cryotop method. *Theriogenology* 67(1), 73-80.
- Kuwayama, M., Vajta, G., Kato, O. and Leibo, S. P. (2005). Highly efficient vitrification method for cryopreservation of human oocytes. *Reproductive Biomedicine Online* 11(3): 300-308.
- Langdon, W. C. (2007). In: Effect of semen parameters of bovine spermatozoa after using a contemporary collecting receptacle. Texas Tech University.
- Leahy, T. and de Graaf, S.P. (2012). Seminal plasma and its effect on ruminant spermatozoa during processing. *Reproduction in Domestic Animals*. *Zuchthygiene* 47:207–13.
- Ledda, S., Bogliolo, L., Succu, S., Ariu, F., Bebbere, D., Leoni, G.G. and Naitana, S. (2007). Oocyte cryopreservation: oocyte assessment and strategies for improving survival. *Reproductive Fertility and Development* 19:13–23.
- Lefievre L, Bedu-Addo K, Conner SJ, Machado-Oliveira GS, Chen Y, Kirkman-Brown JC, et al. (2007). Counting sperm does not add up any more: time for a new equation? *Reproduction* 133:675–84.
- Leibfried, L. and First, N. (1979). Characterization of bovine follicular oocytes and their ability to mature. *Journal of Animal Science*, 48(1), 76-86.
- Leibo, S. (1989). Equilibrium and nonequilibrium cryopreservation of embryos. *Theriogenology* 31(1), 85-93.
- Leisinger C.A., Pinto, R.F., Cramer, E., Love, C. C. and Paccamonti, D. L. (2017). Effects of Repeated Partial Thaw and Refreeze on Post-Thaw Parameters of Stallion Semen Cryopreserved in Cryovials. *Journal of Equine Veterinary Science* 49:19–24.
- Leite, T. G., do Vale Filho, V. R., de Arruda, R. P., de Andrade, A. F. C., Emerick, L. L., Zaffalon, F. G., Martins, J. A. M. and de Andrade, V. J. (2010). Effects of extender and equilibration time on post-thaw motility and membrane integrity of cryopreserved Gyr bull semen evaluated by CASA and flow cytometry. *Animal reproduction science*, 120(1), 31-38.
- Lemma, A. (2013). Effect of cryopreservation on sperm quality and fertility: INTECH Open Access Publisher.

- Len J.A., Jenkins Eilts, B.E., Paccamonti, D.L., Lyle, S.K. and Hosgood, G., (2010). Immediate and delayed (after cooling) effects of centrifugation on equine sperm. *Theriogenology* 73:225–31.
- León H., Porras, A.A., Galina, C.S. and Navarro-Fierro, R., (1991). Effect of the collection method on semen characteristics of Zebu and European type cattle in the tropics. *Theriogenology* 36:349–55.
- Li, M.W., Meyers S., Tollner T.L and Overstreet J.W. (2007). Damage to chromosomes and DNA of rhesus monkey sperm following cryopreservation. *Journal of Andrology* 28:493-501.
- Lipar, J.L., Ketterson, E.D., Nolan Jr.V. and Casto, J.M., (1999). Egg yolk layers vary in the concentration of steroid hormones in two avian species. *Endocrinology* 115:220–227.
- Loomis, P.R. (2006). Advanced methods for handling and preparation of stallion semen. *Veterinary Clinic of North America for Equine Practice* 22:663–76.
- Love, C. C. (2005). The sperm chromatin structure assay: a review of clinical applications. *Animal Reproduction Science* 89(1): 39-45.
- Maccarrone, M., Lorenzon, T., Bari, M., Melino, G. and Finazzi-Agro, A. (2000). Anandamide induces apoptosis in human cells via vanilloid receptors. Evidence for a protective role of cannabinoid receptors. *Journal of Biology and Chemistry* 275:31938–45.
- Macías García, B., González Fernández, L., Ortega Ferrusola, C., Salazar- Sandoval, C., Morillo Rodríguez, A., Rodríguez Martínez, H., et al. (2011). Membrane lipids of the stallion spermatozoon in relation to sperm quality and susceptibility to lipid peroxidation. *Reproduction in Domestic Animals* 46:141–8.
- Manjunath, P., Nauc, V., Bergeron, A., Ménard, M. (2002). Major proteins of bovine seminal plasma bind to the low-density lipoprotein fraction of the hen's egg yolk. *Biology of Reproduction* 67:1250–8.
- Marco-Jiménez, F., Puchades, S., Gadea, J., Vicente, J. and Viudes-de-Castro, M. (2005). Effect of semen collection method on pre-and post-thaw Guirra ram spermatozoa. *Theriogenology* 64(8):1756-1765.
- Marco- Jiménez, F., Puchades, S., Moce, E., Viudes- de- Cartro, M., Vicente, J. and Rodriguez, M. (2004). Use of powdered egg yolk vs fresh egg yolk for the cryopreservation of ovine semen. *Reproduction in Domestic Animals* 39(6): 438-441.

- Martínez-Pastor, F., Martínez, F., Alvarez, M., Maroto-Morales, A., García-Alvarez, O., Soler, A., Garde, J., De Paz, P. and Anel, L. (2009). Cryopreservation of Iberian red deer (*Cervus elaphus hispanicus*) spermatozoa obtained by electroejaculation. *Theriogenology* 71(4): 628-638.
- Martini, F. H. (2006). In: *Fundamentals of Anatomy and Physiology* (7th Ed.). Pearson: San Francisco.
- Martinus, R.D., Molan, P.C. and Shannon, P. (1991). Deleterious effect of seminal plasma in the cryo-preservation of bovine spermatozoa. *New Zealand Journal of Agricultural Research* 34:281-5.
- Maxwell W.M., Evans, G. Mortimer, S.T. Gillan, L. Gellatly E.S. and McPhie, C.A. (1999). Normal fertility in ewes after cervical insemination with frozen-thawed spermatozoa supplemented with seminal plasma. *Reproduction and Fertility Development* 11:123-6.
- Maxwell, W., De Graaf, S., Ghaoui, R. and Evans, G. (2007). Seminal plasma effects on sperm handling and female fertility. *Society of Reproduction and Fertility* 64: 13.
- Maxwell, W., Parrilla, I., Caballero, I., Garcia, E., Roca, J., Martinez, E., Vazquez, J. and Rath, D. (2007). Retained functional integrity of bull spermatozoa after double freezing and thawing using PureSperm® density gradient centrifugation. *Reproduction in Domestic Animals*, 42(5), 489-494.
- Mazur, P., Seki, S., Pinn, I.L., Kleinhans, F.W. and Edashige, K. (2005). Extra- and intracellular ice formation in mouse oocytes. *Cryobiology* 51: 29-53.
- Medeiros, C., Forell, F., Oliveira, A. and Rodrigues, J. (2002). Current status of sperm cryopreservation: why isn't it better? *Theriogenology* 57(1): 327-344.
- Morató, R., Izquierdo, D., Albarraci, J.L., Anguita, B., Palomo, M.J., Jiménez-Macedo, A.R. and Paramio, M.T. (2008a). Effects of pre-treating in vitro-matured bovine oocytes with the cytoskeleton stabilizing agent taxol prior to vitrification. *Molecular Reproductive and Development* 75:191-201.
- Morató, R., Izquierdo, D., Paramio, M.T. and Mogas, T. (2008b). Cryotops versus open-pulled straws (OPS) as carriers for the cryopreservation of bovine oocytes: effects on spindle and chromosome configuration and embryo development. *Cryobiology* 57, 137-141.
- Morató, R., Izquierdo, D., Paramio, M.T. and Mogas, T. (2008c). Embryo development and structural analysis of *in vitro* matured bovine oocytes vitrified in flexipet denuding pipettes. *Theriogenology* 70, 1536-1543.

- Morató, R., Mogas, T., Hyttel, P.M., (2008d). Ultrastructure of bovine oocytes exposed to taxol prior to OPS vitrification. *Molecular Reproduction and Development* 75:1318–1326.
- Morrell, J. (2006). Update on Semen Technologies for Animal Breeding. *Reproduction in Domestic Animals* 41(1): 63-67.
- Morrell, J. and Wallgren, M. (2011). Removal of bacteria from boar ejaculates by Single Layer Centrifugation can reduce the use of antibiotics in semen extenders. *Animal Reproduction Science* 123(1): 64-69.
- Mostek A., Aleksandra M., Mariola S., and Andrzej C. (2017). Cryopreservation of Bull Semen is Associated with Carbonylation of Sperm Proteins. *Theriogenology* 92 95-102.
- Moura, A. A., Chapman, D. A., Koc, H. and Killian, G. J. (2007). A comprehensive proteomic analysis of the accessory sex gland fluid from mature Holstein bulls. *Animal Reproduction Science* 98(3): 169-188.
- Moura, A. and Memili, E. (2016). Functional aspects of seminal plasma and sperm proteins and their potential as molecular markers of fertility. *Animal Reproduction* 13(3): 191-199.
- Nagy, S., Johannisson, A., Wahlsten, T., Ijäs, R., Andersson, M. and Rodriguez-Martinez, H. (2013). Sperm chromatin structure and sperm morphology: Their association with fertility in AI-dairy Ayrshire sires. *Theriogenology* 79(8): 1153-1161.
- Naing, S. W., Haron, A. W., Goriman, M. A. K., Yusoff, R., Bakar, M. Z. A., Sarsaifi, K., Bukar, M. M., Thein, M., Kyaw T. and San M. M. (2011). Effect of seminal plasma removal, washing solutions, and centrifugation regimes on Boer goat semen cryopreservation. *Pertanika Journal of Tropical Agricultural Sciences* 34(2): 271-279.
- Nielsen, U. S., Aamand, G. P. and Mark, T. (2000). National genetic evaluation of udder health and other health traits in Denmark. *Interbull bulletin* (25), 143.
- Nohalez, A., Martinez, C.A., Gil, M.A., Almiñana, C., Roca, J., Martinez, E.A. and Cuello, C. (2015). Effects of two combinations of cryoprotectants on the in vitro developmental capacity of vitrified immature porcine oocytes. *Theriogenology* 84, 545–552.
- O'Brien, J., Steinman, K., Schmitt, T. and Robeck, T. (2008). Semen collection, characterisation and artificial insemination in the beluga (*Delphinapterus leucas*) using liquid-stored spermatozoa. *Reproduction, Fertility and Development* 20(7): 770-783.

- Orzolek, A., Wysocki, P., Strzeżek, J. and Kordan, W. (2013). Superoxide dismutase (SOD) in boar spermatozoa: Purification, biochemical properties and changes in activity during semen storage (16°C) in different extenders. *Reproductive Biology*, 13(1), 34-40.
- Pace, M.M. and Graham, E.F. (1974). Components in egg yolk which protect bovine spermatozoa during freezing. *Journal of Animal Science* 39: 1144–1149.
- Paradis, F., Vigneault, C., Robert, C. and Sirard, M. A. (2005). RNA interference as a tool to study gene function in bovine oocytes. *Molecular Reproduction and Development* 70(2), 111-121.
- Peddinti, D., Nanduri, B., Kaya, A., Feugang, J.M., Burgess, S.C. and Memili, E. (2008). Comprehensive proteomic analysis of bovine spermatozoa of varying fertility rates and identification of biomarkers associated with fertility. *BMC Systematic Biology* 2:19.
- Pesch, S. and Bergmann, M. (2006). Structure of mammalian spermatozoa in respect to viability, fertility and cryopreservation. *Micron*, 37(7), 597-612.
- Phillips, P. H. and Lardy, H. A. (1940). A yolk-buffer pabulum for the preservation of bull semen. *Journal of Dairy Science*, 23(5), 399-404.
- Pieterse, M., Vos, P., Kruip, T. A., Wurth, Y., van Beneden, T. H., Willemse, A. and Taverne, M. (1992). Repeated transvaginal ultrasound-guided ovum pick-up in egg-treated cows. *Theriogenology*, 37(1), 273.
- Pillet, E., C. Labbe, F. Batellier, G. Duchamp, V. Beaumal, M. Anton, S. Desherces, E. Schmitt and M. Magistrini, (2012). Liposomes as an Alternative to Egg Yolk in Stallion Freezing Extender. *Theriogenology* 77 (2): 268–279.
- Prentice, J.R., Singh, J., Dochi, O. and Anzar, M., (2011). Factors affecting nuclear maturation, cleavage and embryo development of vitrified bovine cumulus-oocyte complexes. *Theriogenology* 75, 602 –609.
- Prentice-Biensch, J.R., Singh, J., Mapletoft, R.J. and Anzar, M. (2012). Vitrification of immature bovine cumulus-oocyte complexes: effects of cryoprotectants, the vitrification procedure and warming time on cleavage and embryo development. *Reproductive Biology and Endocrinology*. 73, 1–7.
- Rall, W.F., Wood, M.J., Kirby, C. and Whittingham, D.G. (1987). Development of mouse embryos cryopreserved by vitrification. *Journal of Reproduction and Fertility*. 80, 499–504.
- Rasul, Z., Ahmad, N. and Anzar, M. (2001). Changes in motion characteristics, plasma membrane integrity, and acrosome morphology during cryopreservation of buffalo spermatozoa. *Journal of Andrology* 22(2): 278-283.

- Ravandi, A., Kuksis, A. and Shaikh, N. A. (2000). Glucosylated Glycerophosphoethanolamines are the Major LDL Glycation Products and Increase LDL Susceptibility to Oxidation Evidence of Their Presence in Atherosclerotic Lesions. *Arteriosclerosis, thrombosis, and vascular biology* 20(2): 467-477.
- Rego, J.P.A., Moura, A.A., Nouwens, A.S., McGowan, M.R. and Boe-hansen, G.B. (2015). Seminal plasma protein profiles of ejaculates obtained by internal artificial vagina and electroejaculation in Brahman bulls. *Animal Reproduction Science* 160:126–37.
- Rekha, A., Zoharab B. F. and Baric, F.Y. (2016). Comparison of commercial Triladyl extender with a tris-fructose-egg-yolk extender on the quality of frozen semen and pregnancy rate after trans-cervical AI in Bangladeshi indigenous sheep (*Ovis aries*). *Small Ruminant Research*, 134: 39–43.
- Rho, G.J., Kim, S., Yoo, J.G., Balasubramanian, S., Lee, H.J. and Choe, S.Y. (2002). Microtubulin configuration and mitochondrial distribution after ultrarapid cooling of bovine oocytes. *Molecular Reproduction and Development* 63:464–70.
- Rodger, J. C. (1975). Seminal plasma, an unnecessary evil? *Theriogenology*, 3(6), 237-247.
- Rojas, C., Palomo, M.J., Albarracín, J.L. and Mogas, T. (2004). Vitrification of immature and in vitro matured pig oocytes: study of distribution of chromosomes, microtubules, and actin microfilament. *Cryobiology* 49:211-20.
- Romek, M., Gajda, B., Rolka, M. and Smorag, Z., (2011). Mitochondrial activity and morphology in developing porcine oocytes and pre-implantation non-cultured and cultured embryos. *Reproduction in Domestic Animals* 46: 471–480.
- Röpke, T., Oldenhof, H., Leiding, C., Sieme, H., Bollwein, H and Wolkers, W.F. (2011). Liposomes for Cryopreservation of Bovine Sperm. *Theriogenology* 76 (8): 1465–1472.
- Rota, A., Furzi, C., Panzani, D. and Camillo, F. (2004). Studies on motility and fertility of cooled stallion spermatozoa. *Reproduction in Domestic Animals* 39(2): 103-109.
- Rota, A., Ström, B. and Linde-Forsberg, C. (1995). Effects of seminal plasma and three extenders on canine semen stored at 4 C. *Theriogenology* 44(6), 885-900.
- Roth, Z., Arav, A., Bor, A., Zeron, Y., Braw-Tal, R. and Wolfenson, D. (2001). Improvement of quality of oocytes collected in the autumn by enhanced removal of impaired follicles from previously heat-stressed cows. *Reproduction*, 122(5), 737-744.

- Saacke, R., Dalton, J., Nadir, S., Nebel, R. and Bame, J. (2000). Relationship of seminal traits and insemination time to fertilization rate and embryo quality. *Animal Reproduction Science* 60:663–77.
- Salmani, H., Towhidi A., Zhandi M., Bahreini M., Sharafi, M. (2014). *In Vitro* Assessment of Soybean Lecithin and Egg Yolk Based Diluents for Cryopreservation of Goat Semen. *Cryobiology* 68 (2), 276–280.
- Sansone, G., Nastri, M. and Fabbrocini, A. (2000). Storage of buffalo (*Bubalus bubalis*) semen. *Animal Reproduction Science*, 62(1), 55-76.
- Sarsaifi K., Rosnina Y., Ariff M.O., Wahid H., Hani H. and Yimer N. (2015). Effect of semen collection methods on the quality of pre and post-thawed Bali cattle (*Bos javanicus*) spermatozoa. *Reproduction in Domestic Animals* 44:1006–12.
- Sathe, S. and Shipley, C. F. (2015). Cryopreservation of Semen. *Bovine Reproduction* 662-670.
- Schiller, J., Müller, K., Süß, R., Arnhold, J., Gey, C., Herrmann, A., Leßig, J., Arnold, K. and Müller, P. (2003). Analysis of the lipid composition of bull spermatozoa by MALDI-TOF mass spectrometry-a cautionary note. *Chemistry and physics of lipids*, 126(1), 85-94.
- Şen, Ç. Ç., Tekin K. and Akcay E. (2015). Effect of Egg Yolk and Removal of Seminal Fluid on Semen Cryopreservation in Norduz Goat.
- Senger, P., Becker, W., Davidge, S., Hillers, J. and Reeves, J. (1988). Influence of cornual insemination on conception in dairy cattle. *Journal of Animal Science* 66(11): 3010-3016.
- Setchell, B. P. (1977). In: *Male Reproductive Organ and Semen* (3 Ed.). New York: Academic Press.
- Shaliutina, A., Hulak, M., Gazo, I., Linhartova, P. and Linhart, O. (2013). Effect of short-term storage on quality parameters, DNA integrity, and oxidative stress in Russian (*Acipenser gueldenstaedtii*) and Siberian (*Acipenser baerii*) sturgeon sperm. *Animal Reproduction Science*, 139(1), 127-135.
- Shoae, A. and Zamiri, M. (2008). Effect of butylated hydroxytoluene on bull spermatozoa frozen in egg yolk-citrate extender. *Animal Reproduction Science* 104(2), 414-418.
- Sirard, M., Parrish, J., Ware, C., Leibfried-Rutledge, M. and First, N. (1988). The culture of bovine oocytes to obtain developmentally competent embryos. *Biology of Reproduction*, 39(3): 546-552.
- Skalet, L. H. (1986). Effects of age and season on the spermogram of Nubian male goats. (MSc), Tuskegee University.

- Sohnrey, B. and Holtz, W. (2005). Technical Note: Transcervical deep cornual insemination of goats. *Journal of Animal Science* 83(7):1543-1548.
- Souza-Fabjan, J.M., Locatelli, Y., Duffard, N., Corbin, E., Touze, J.L., Perreau, C., Beckers, J.F., Freitas, V.J. and Mermillod, P., (2014). In vitro embryo production in goats: slaughterhouse and laparoscopic ovum pick up-derived oocytes have different kinetics and requirements regarding maturation media. *Theriogenology* 81: 1021-1031.
- Stradaoli, G., Noro, T., Sylla, L. and Monaci, M., (2007). Decrease in glutathione (GSH) content in bovine sperm after cryopreservation: comparison between two extenders. *Theriogenology* 67: 1249–1255.
- Sugie, T. (1965). Successful transfer of a fertilized bovine egg by non-surgical techniques. *Journal of Reproduction and Fertility*, 10(2), 197-201.
- Sun, Q.Y., Wu, G.M., Lai, L., Park, K.W., Cabot, R., Cheong, H.T., Day, B.N., Prather, R.S. and Schatten, H. (2001). Translocation of active mitochondria during pig oocyte maturation, fertilization and early development *in vitro*. *Reproduction* 122: 155–163.
- Suzane, P.C., Fabio, M.M., Erika, A.R.D., Maria, E.Z.M., Claudia de C. P. P., José, A.D.J., Frederico, O.P., Camila de, P.F.D., Roberta, V. and Joaquim, M.G. (2017). Effect of seminal plasma removal before cryopreservation of bovine semen obtained by electroejaculation on semen quality and *in vitro* fertility. *Theriogenology* 89: 114–121.
- Tachikawa, S., Otoi, T., Kondo, S., Machida, T. and Kasai, M. (1993). Successful vitrification of bovine blastocysts, derived by *in vitro* maturation and fertilization. *Molecular Reproduction and development*, 34(3), 266-271.
- Taşdemir, U., Büyükleblebici, S., Tuncer, P. B., Coşkun, E., Özgürtaş, T., Aydın, F. N., Büyükleblebici, O. and Gürcan, İ. S. (2013). Effects of various cryoprotectants on bull sperm quality, DNA integrity and oxidative stress parameters. *Cryobiology* 66(1): 38-42.
- Tatham, A. S. and Shewry, P. R. (2000). Elastomeric proteins: biological roles, structures and mechanisms. *Trends in Biochemical Sciences* 25(11): 567-571.
- Tejada, R. I., Mitchell, J. C. Norman, A. Marik, J. J. and Friedman, S. (1984). A Test for the Practical Evaluation of Male Fertility by Acridine Orange (AO) Fluorescence. *Fertility and Sterility* 42(1), 87-91.
- Thérien, I., Bousquet, D. and Manjunath, P. (2001). Effect of seminal phospholipid-binding proteins and follicular fluid on bovine sperm capacitation. *Biology of reproduction*, 65(1), 41-51.

- Vajta, G., Holm, P., Kuwayama, M., Booth, P. J., Jacobsen, H., Greve, T. and Callesen, H. (1998). Open pulled straw (OPS) vitrification: a new way to reduce cryoinjuries of bovine ova and embryos. *Molecular Reproduction and Development* 51(1): 53-58.
- Vajta, G. and Nagy, Z.P. (2006). Are programmable freezers still needed in the embryo laboratory? Review on vitrification. *Reproductive Biomedicine*, Online 12, 779–796.
- Verstegen, J., Iguer-Ouada, M. and Onclin, K., (2002). Computer assisted semen analyzers in andrology research and veterinary practice. *Theriogenology* 57:149–79.
- Vieira, O. V., Botelho, R. J. and Grinstein, S. (2002). Phagosome maturation: aging gracefully. *Biochemical Journal* 366(3): 689-704.
- Villemure, M., Lazure, C. and Manjunath, P. (2003). Isolation and characterization of gelatin-binding proteins from goat seminal plasma. *Reproductive Biology and Endocrinology* 1(1): 1
- Vishwanath, R. and Shannon, P. (2000). Storage of bovine semen in liquid and frozen state. *Animal Reproduction Science*, 62(1), 23-53.
- Wall, R. and Foote, R. (1999). Fertility of bull sperm frozen and stored in clarified egg yolk-Tris-glycerol extender. *Journal of Dairy Science* 82(4): 817-821.
- Waterhouse K.E., Gjeldnes, A., Tverdal, A., Angelis, P.M. De, Farstad, W., Haard M. and Kommisrud, E. (2010). Alteration of Sperm DNA Integrity during Cryopreservation Procedure and Invitro Incubation of Bull Semen. *Animal Reproduction Science* 117: 34-42.
- Watson, A.J., (2007). Oocyte cytoplasmic maturation: a key mediator of oocyte and embryo developmental competence. *Journal of Animal Science* 85 (13 Suppl): E1–E3.
- Watson, P. (2000). The causes of reduced fertility with cryopreserved semen. *Animal Reproduction science*, 60, 481-492.
- Watson, P. and Duncan, A. E. (1988). Effect of salt concentration and unfrozen water fraction on the viability of slowly frozen ram spermatozoa. *Cryobiology* 25(2): 131-142.
- Way A., Grieljr L.C., Killian, G.J. (2000). Effects of accessory sex gland fluid on viability, capacitation, and the acrosome reaction of cauda epididymal bull spermatozoa. *Journal of Andrology* 21:213.
- Weidel, L. and Prins, G. S. (1987). Cryosurvival of human spermatozoa frozen in eight different buffer systems. *Journal of Andrology*, 8(1), 41-47.

- Wheeler, M. B., Rutledge, J. J., Fischer-Brown, A., Vanetten, T., Malusky, S. and Beebe, D. J. (2006). Application of sexed semen technology to *in vitro* embryo production in cattle. *Theriogenology*, 65(1), 219-227.
- White, I. (1993). Lipids and calcium uptake of sperm in relation to cold shock and preservation: a review. *Reproduction, Fertility and Development*, 5(6), 639-658.
- Woelders, H. (1997). Fundamentals and recent development in cryopreservation of bull and boar semen. *Veterinary quarterly* 19(3): 135-138.
- Wright, V. C., Chang, J., Jeng, G., Macaluso, M., Control, C. f. D. and Prevention. (2008). Assisted reproductive technology surveillance—United States, 2005. *MMWR Surveill Summary* 57(5): 1-23.
- Yamada, C., Caetano, H.V., Simões, R., Nicacio, A.C., Feitosa, W.B., Assumpção, M.E., et al. (2007). Immature bovine oocyte cryopreservation: comparison of different associations with ethyleneglycol, glycerol and dimethylsulfoxide. *Animal Reproduction Science*; 99:384–8.
- Yanagimachi R., (1994). Fertility of mammalian spermatozoa: its development and relativity. *Zygote* (Cambridge, Engl.) 2: 371–372.
- Yániz, J., Mateos, J. and Santolaria, P. (2011). Zwitterionic buffers preserve ram semen quality more efficiently than TRIS during storage at 15° C. *Small Ruminant Research*, 95(1), 54-60.
- Zaman, A. (2013). Livestock improvement using biotechnology: world and Bangladesh perspective. *Journal of Livestock Science* 4: 78-96.
- Zeron, Y., Ocheretny, A., Kedar, O., Borochoy, A., Sklan, D. and Arav, A. (2001). Seasonal changes in bovine fertility: relation to developmental competence of oocytes, membrane properties and fatty acid composition of follicles. *Reproduction* 121(3): 447-454.
- Zheng, C., Khoo, C., Furtado, J. and Sacks, F. M. (2010). Apolipoprotein C-III and the metabolic basis for hypertriglyceridemia and the dense low-density lipoprotein phenotype. *Circulation* 121(15): 1722-1734.