



UNIVERSITI PUTRA MALAYSIA

***SERO-EPIDEMIOLOGY OF JAPANESE ENCEPHALITIS VIRUS AMONG
LIVESTOCK, AVIAN AND COMPANION ANIMALS IN
SELECTED STATES OF MALAYSIA***

KIVEN KUMAR A/L KARUNANITHI

FPV 2018 13



**SERO-EPIDEMIOLOGY OF JAPANESE ENCEPHALITIS VIRUS AMONG
LIVESTOCK, AVIAN AND COMPANION ANIMALS IN
SELECTED STATES OF MALAYSIA**

By

KIVEN KUMAR A/L KARUNANITHI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Master of Science**

November 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**SERO-EPIDEMIOLOGY OF JAPANESE ENCEPHALITIS VIRUS AMONG
LIVESTOCK, AVIAN AND COMPANION ANIMALS IN
SELECTED STATES OF MALAYSIA**

By

KIVEN KUMAR A/L KARUNANITHI

November 2017

Chairman : Professor Siti Suri Arshad, PhD
Faculty : Veterinary Medicine

Japanese encephalitis (JE) is vector-borne disease causes encephalitis in human and horse as well as reproductive failure in sows. Pigs, bats, wild boar, and *ardeidae* birds play an important role as a main vertebrate amplifier of JEV. In Malaysia the disease is endemic in east Malaysia, Sarawak and epidemic in Peninsular Malaysia. The encephalitis cases were observed among human and animals reported in earlier 1950s. However, they are no data regarding seroprevalence pattern among livestock (buffaloes, pigs, cattle), avian (captive water birds, poultry and migratory birds) and companion animals (cats, dogs). At the same time, risk factors associated with JEV seroprevalence in high risk areas in Malaysia are not documented. In this study, the objectives are to determine JEV seroprevalence rate, risk factors associated with JEV seroprevalence and determination of JEV genotype circulating among livestock (buffaloes, pigs, cattle), avian (water captive birds, poultry and migratory birds) and companion animals (cats, dogs) were documented. Total of 461 serum samples were collected from dogs, cats, captive water birds, village chickens, domestic pigs, buffaloes and cattle in selective states of Perak, Selangor and Sabah and screened for evidence of JEV antibodies by using a double sandwich IgG ELISA kits (DAS-ELISA). The risk factors associated with JEV seroprevalence of the animals were analysed by using IBM SPSS 22 which included *Chi-square* (χ^2 test is not appropriate if one of expected cell value is less than 5, Fisher's exact test was used at $\alpha = 0.05$) and multiple logistic regression. A total of 13 survey criteria which include gender (male vs female), health status (healthy vs sick), age (young vs adult), breeding types (local, vs import), ownership of animal (own, shelter or none) and source of animals (same area, same district, same state vs epidemic areas) were collected. At the same time, environmental factors of sampling areas including presence of stagnant water (yes or no), paddy cultivation (yes or no), presence of ardeid birds (yes or no), locality (urban vs rural) and presence of mosquitos (yes or no). In the molecular assay, total of

791 samples of plasma, serum, and buffy coat from livestock (buffaloes, pigs, cattle), companion animals (cats, dogs) and avian (water captivity birds, poultry and migratory birds) were subjected to one step RT-PCR to detect the JEV by targeting the conserved region of NS3 of the JEV. In seroprevalence study among the animals studied, dogs show the highest seropositive rate of 80% (36/45; 95%CI: ± 11.69) followed by pigs with 44.4% (40/90; 95% CI: ± 1.715), buffaloes with 33.3% (15/45; 95% CI: ± 6.661), cattle with 32.2% (29/90; 95% CI: ± 1.058), avian 28.9% (13/45; 95% CI: ± 5.757), and cats with 14.4% (13/90; 95% CI: ± 7.38). This study reveals that the risk factors associated with JEV seroprevalence are varied among different species of animals. The risk factors involved in different species of animals can be either singly or more than one. In addition to that, geographical factors may influence on the seropositivity of animals to JEV. In dogs, significant risk factor associated with JEV seroprevalence is the source of dog. In cats, significant risk factors associated with JEV seroprevalence are the age, locality, breeding types, and source of cats. In cattle, significant risk factors associated with JEV seroprevalence are gender, healthy status, breeding types and source of cattle. In pigs, significant risk factors associated with JEV seroprevalence are age and ownership of pigs. In buffaloes, significant risk factors associated with JEV seroprevalence are the source of buffalo and breeding types. In avian, significant risk factors associated with JEV seroprevalence are age and the breeding types. In this study, the high JEV seroprevalence in dogs observed is believed due to the preference and selective feeding pattern of *Culex* spp. in dogs compared to cats or human, thus making the dogs more prone to infection. JEV seroprevalence in pigs in this study are lower than reported previously probably due to the location of the subject matter in a non-paddy cultivation areas. This study also observed that JEV seroprevalence rate in buffaloes and cattle are low and support previous finding of this species play minimum role in JEV transmission cycles. The low viremia following JEV infection is too low of viral load to infect vector. Low JEV seroprevalence observed in chickens in this study is rather unexpected as avian was sampled in high risks areas with paddy plantation, water bodies and mosquitoes breeding sites. This finding may be further support by the lack of pig farming in this area, as the residents are engaged in fishing activities besides paddy cultivation. In this study, JEV antigen was unable to be detected in all the samples collected from various species of animals. Preliminary findings show that, all the animals were negative for JEV antigen. This could be due to the presence of antibodies of JEV naturalised the antigen. Apart from that, the sero-complex characteristics of flaviviruses make the antigen-antibodies cross-react with each other members. This makes the antigen being neutralised and unable to be detected in blood of the host. Samples collected during acute stage of JEV infection in animals have high chances of virus being isolated. In addition to that, the oronasal fluid is preeminent for isolation and detection of JEV antigen compared to serum, cerebrum spinal fluid, plasma and buffy coat.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

SERO-EPIDEMIOLOGI VIRUS ENSEFALITIS JEPUN DALAM HAIWAN TERNAKAN, BURUNG DAN HAIWAN KESAYANGAN DI NEGERI-NEGERI TERPILIH DI MALAYSIA

Oleh

KIVEN KUMAR A/L KARUNANITHI

November 2017

Pengerusi : Profesor Siti Suri Arshad, PhD
Fakulti : Perubatan Veterinar

Ensefalitis Jepun (JE) adalah penyakit bawaan vektor yang menyebabkan ensefalitis pada manusia dan kuda serta kegagalan pembiakan dalam khinzir. Khinzir domestik, kelawar, khinzir liar, dan burung ardeidae memainkan peranan penting sebagai penguat vertebrata utama virus ensefalitis Jepun (JEV). Di Malaysia, penyakit ini adalah endemik di timur Malaysia iaitu Sarawak dan epidemik di Semenanjung Malaysia. Kes ensefalitis dilaporkan dalam kalangan manusia dan haiwan sebelum tahun 1950-an. Walaubagaimanapun, data mengenai corak seroprevalensi dalam kalangan haiwan ternakan (kerbau, khinzir, lembu), haiwan kesayangan (kucing, anjing) dan burung (burung air, unggas dan burung berhijrah) tidak didokumentasikan. Pada masa yang sama, faktor-faktor risiko yang dikaitkan dengan seroprevalensi JEV di kawasan berisiko tinggi di Malaysia juga tidak didokumentasikan. Dalam kajian ini, objektifnya adalah untuk menentukan kadar seroprevalensi JEV, faktor-faktor risiko yang dikaitkan dengan seroprevalensi JEV dan penentuan kitaran genotip JEV di kalangan haiwan ternakan (kerbau, khinzir, lembu), haiwan kesayangan (kucing, anjing) dan burung (burung air, unggas dan burung berhijrah) telah didokumentasikan. Empat ratus enam puluh satu sampel serum telah dikumpulkan daripada anjing, kucing, burung penangkapan air, ayam kampung, khinzir domestik, kerbau dan lembu daripada negeri-negeri terpilih iaitu Perak, Selangor dan Sabah dan diperiksa untuk megesan kehadiran antibodi JEV dengan menggunakan kit komersil berkonsep 'sandwich berganda' IgG ELISA (DAS -ELISA). Faktor-faktor risiko yang dikaitkan dengan seroprevalensi JEV haiwan dianalisis dengan menggunakan IBM SPSS 22 yang termasuk *Chi-square* (ujian χ^2 tidak sesuai apabila jumlah sample kurang daripada 5, ujian 'exact Fisher' telah digunakan pada peringkat $\alpha = 0.05$) dan multivariasi regresi logistik. Sebanyak 13 kriteria telah dikaji termasuk jantina (jantan dengan betina), status kesihatan (sihat dengan sakit), umur (muda dengan dewasa), jenis baka (tempatan, dengan import), pemilikan haiwan (sendiri, tempat perlindungan

atau tiada) sumber haiwan (daripada kawasan yang sama; daerah yang sama; negeri yang sama dengan kawasan epidemik) telah dikumpulkan. Pada masa yang sama, faktor persekitaran persampelan termasuk kewujudan air bertakung (ya atau tidak), penanaman padi (ya atau tidak), kewujudan burung ardeid (ya atau tidak), persekitaran tempatan (bandar dengan luar bandar) dan kewujudan nyamuk (ya atau tidak). Dalam ujian molekul, sebanyak 791 sampel termasuk plasma, serum, dan 'buffy coat' daripada haiwan ternakan (kerbau, khinzir, lembu), haiwan kesayangan (kucing, anjing) dan burung (burung air, unggas dan burung berhijrah) telah dianalisis melalui 'reverse transcription polymerase chain reaction' (RT-PCR) untuk mengesan JEV antigen dengan berasaskan bahagian NS3 JEV. Dalam kajian ini, anjing menunjukkan kadar seropositif tertinggi sebanyak 80% (36/45; 95% CI: ± 11.69) diikuti oleh khinzir sebanyak 44.4% (40/90; 95% CI: ± 1.715), kerbau sebanyak 33.3% (15/45; 95% CI: ± 6.661), lembu sebanyak 32.2% (29/90; 95% CI: ± 1.058), burung sebanyak 28.9% (13/45; 95% CI: ± 5.757), dan kucing sebanyak 15.6% (14/90; 95% CI: ± 7.38). Kajian ini mendedahkan bahawa faktor-faktor risiko yang dikaitkan dengan seroprevalensi JEV adalah berbeza di antara spesies haiwan yang berlainan. Faktor risiko yang terlibat dalam spesies haiwan yang berlainan boleh sama ada secara tunggal atau lebih daripada satu. Tambahan pula, faktor geografi boleh mempengaruhi seropositif JEV dalam haiwan. Faktor risiko yang signifikan dalam anjing yang dikaitkan dengan seroprevalensi JEV adalah sumber anjing. Dalam kucing pula, faktor risiko yang signifikan yang berkaitan dengan seroprevalensi JEV adalah umur, lokasi, baka, dan sumber kucing. Dalam lembu, faktor risiko yang signifikan yang dikaitkan dengan seroprevalensi JEV adalah jantina, status kesihatan, baka dan sumber lembu. Dalam khinzir, faktor risiko yang signifikan yang berkaitan dengan seroprevalensi JEV adalah umur dan pemilik khinzir. Dalam kerbau, faktor risiko yang signifikan berkaitan dengan seroprevalensi JEV adalah sumber kerbau dan baka. Dalam burung, faktor risiko yang signifikan yang berkaitan dengan seroprevalensi JEV adalah umur dan baka. Dalam kajian ini, seroprevalensi JEV yang tinggi dalam anjing dipercayai kerana pilihan dan pola pemakanan oleh *Culex* spp. pada anjing berbanding dengan kucing atau manusia, sehingga menjadikan anjing lebih terdedah terhadap jangkitan. Seroprevalensi JEV khinzir dalam kajian ini adalah lebih rendah daripada yang dilaporkan sebelum ini mungkin disebabkan tidak mempunyai lokasi penanaman padi di kawasan sampel yang diperolehi. Kajian ini juga mendapati bahawa kadar seroprevalensi dalam kerbau dan lembu adalah rendah dan menyokong penemuan sebelumnya kerana spesies ini memainkan peranan minimum dalam kitaran transmisi JEV. Viremia yang rendah berikutan jangkitan JEV dengan bebanan virus yang rendah didalam perumah tidak dapat menjangkiti vektor. Seroprevalensi JEV dalam ayam dalam kajian ini tidak dijangka kerana sampel ayam telah dikumpulkan daripada kawasan-kawasan yang berisiko tinggi dengan penanaman padi, air bertakung dan tapak pembiakan nyamuk. Penemuan ini disokong selanjutnya dengan kekurangan peternakan khinzir di kawasan ini, kerana penduduk terlibat dalam kegiatan penangkapan ikan selain daripada penanaman padi. Dalam kajian ini, antigen JEV tidak dapat dikesan dalam semua sampel yang dikumpulkan dari pelbagai spesies haiwan. Penemuan awal menunjukkan bahawa semua haiwan adalah negatif untuk antigen JEV. Ini disebabkan kewujudan antibodi JEV dalam haiwan mampu meneutralkan antigen JEV. Selain itu, ciri-ciri sero-kompleks flavivirus membuatkan antibodi antigen bertindak balas dengan virus yang lain. Ini menjadikan antigen itu dineutralkan dan tidak dapat mengesan di dalam darah perumah. Sampel

yang dikumpulkan semasa peringkat akut jangkitan JEV pada dipercayai mampu untuk diasingkan virus. Selain itu, cecair oronasal adalah yang paling penting untuk pengasingan dan pengesanan antigen JEV berbanding dengan serum, cecair cerebrum tulang belakang, plasma dan 'buffy coat'.



ACKNOWLEDGEMENTS

The completion of this dissertation would not be achieved without the constructive advices, continuous support and constant guidance from my supervisory committee. First and foremost, I would like to thank my supervisor, Professor Dr Siti Suri Arshad the chairman of my supervisory committee for her assistance in completing my research and thesis. Her expertise in virology was a great asset in designing the experiments and analyzing the results. My deepest appreciation also goes to my co-supervisor, Associate Professor Dr Gayathri Thevi Selvarajah, Associate Professor Dr Jalila Abu and Associate Professor Dr Ooi Peck Toung. Thank you very much for the guidance. Completing my studies on time would be remain as dream without the guidance and motivation from my mentors Dr Faruku Bande, Dr Yasmin Abdul Rahman and Dr Yusuf Abba where their continuous support and kindness were my inspiration to complete this study.

I am grateful to all my friends and virology laboratory staff, Krishnan Nair, Ashwaq Ahmed Abdullah, Jamilu Abubakar Bala, Rusdam Awang, Mohammad Sabri Abdul Rahman, Nur Diyana Mohamad Tahir, Ruviniya Kalaiselvam, Mohd Kamaruddin Awang Isa, and Amira Peli. I would like to express our gratitude to Prof. Dr. Sazaly Bin Abu Bakar from Universiti Malaya and Department of Wildlife and National Parks (Perhilitan) Malaysia for JEV positive control and technical support during migratory and wild birds sampling respectively. My thanks and love goes to my family, especially my mom whom provides me the best childhood as well as continue supporting me till date and my brothers, sister and loved ones to encourage me throughout many years in my master's journey. Finally, I would like to acknowledge the financial support from Graduate Research Fellowship (GRF) and FRGS from Universiti Putra Malaysia and Kementerian Pengajian Tinggi.

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Siti Suri Arshad, DVM, MS, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Gayathri Thevi Selvarajah, DVM, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Jalila Abu, DVM, MS, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Ooi Peck Toung DVM, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date :

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No.: Kiven Kumar A/L Karunanithi GS43997

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Professor Dr. Siti Suri Arshad

Signature: _____
Name of Member
of Supervisory
Committee: Associate Professor Dr. Gayathri Thevi Selvarajah

Signature: _____
Name of Member
of Supervisory
Committee: Associate Professor Dr. Jalila Abu

Signature: _____
Name of Member
of Supervisory
Committee: Associate Professor Dr. Ooi Peck Toung

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	vi
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xv
LIST OF FIGURES	xvii
LIST OF ABBREVIATIONS	xx
 CHAPTER	
1 INTRODUCTION	1
1.1 Hypothesis	2
1.2 The objectives of this study are:	2
 2 LITERATURE REVIEW	 3
2.1 Taxonomy of <i>Flavivirus</i>	3
2.1.1 Physiochemical Properties of FVs	3
2.1.2 Genomic Properties of JEV	3
2.1.3 The JE serological group of viruses	7
2.1.4 Envelope (E) Protein	7
2.2 Replication of JEV	8
2.3 Geographical Range of JEV	8
2.3.1 Epidemiological Areas of JEV	8
2.3.1.1 Malaysia	10
2.3.1.2 Thailand	11
2.3.1.3 China	11
2.3.1.4 Japan	11
2.3.1.5 Indonesia	12
2.4 The Transmission Cycles of JEV	12
2.5 Japanese encephalitis virus in Animals	13
2.5.1 Pigs and Piglets	13
2.5.2 Wild Boars	14
2.5.3 Dogs	15
2.5.4 Birds	15
2.5.5 Bats	16
2.5.6 Cow and Calf	17
2.5.7 Reptiles	17
2.6 Mosquito Ecology and Japanese Encephalitis virus	17
2.6.1 JEV Vectors	17
2.6.2 Overwintering of JEV	18
2.7 Factors Contributing the Transmission of JEV	18
2.8 Diagnostic Methods for JEV	21

2.8.1	Serology Assay	21
2.8.1.1	2.8.1.1 Hemagglutination Inhibition (HI)	21
2.8.1.2	IgM and IgG Determination	21
2.8.1.3	Plaque Reduction Neutralisation Titre Assay	22
2.8.1.4	Western Blot	22
2.8.2	Virus Isolation	22
2.8.3	Molecular Detection of JEV Nucleic Acid	22
3	MATERIALS AND METHODS	24
3.1	Study Areas	24
3.2	Study Design	24
3.3	Ethical Approval	27
3.4	Blood Sampling	27
3.5	Methods of Trapping Wild and Migratory Birds	28
3.5.1	Mist-net Trapping	28
3.5.2	Hand-net Trapping	29
3.6	Serology Test	30
3.6.1	Schematic Diagrams of DAS-ELISA	30
3.6.2	Blood Samples of Animals used for DAS-ELISA Assay.	31
3.6.2.1	Detection of JEV Antibodies IgG in Avian	32
3.6.2.2	Detection of JEV Antibodies IgG in Buffaloes	33
3.6.2.3	Detection of JEV Antibodies IgG in Cats	33
3.6.2.4	Detection of JEV Antibodies IgG in Cattle	33
3.6.2.5	Detection of JEV Antibodies IgG in Dogs	34
3.6.2.6	Detection of JEV Antibodies IgG in Pigs	34
3.6.3	Determination of ELISA Cut off Value	34
3.7	Survey on Risk Factors for JEV Prevalence in Animals	35
3.7.1	Data Collection	35
3.7.2	Data Management and Statistical Analysis	35
3.8	Molecular Detection of Japanese encephalitis virus by One Step RT-PCR	36
3.8.1	Amplification and Detection of JEV Gene	36
3.8.1.1	Oligonucleotides Primers	36
3.8.2	Reference JEV Nakayama Strain	37
3.8.2.1	Determination of RNA Concentration and Purity of JEV Reference Strain	37
3.8.3	Optimisation of One-Step Reverse Transcriptase Polymerase Chain Reaction (RT-PCR) to Amplify JEV	37
3.8.3.1	RT-PCR Assays	37
3.8.3.2	Cycling Conditions for the One-Step RT-PCR for detecting Reference JEV	37
3.8.3.3	Gel Electrophoresis of RT-PCR Products	38
3.8.3.4	Visualisation of Electrophoresed Gel	38
3.8.4	Gene Cloning of JEV Reference Strain	38
3.8.4.1	Amplification of NS-3 and E Genes by RT-Polymerase Chain Reaction	38
3.8.4.2	Activation of Competent Cell of <i>E.coli</i>	38
3.8.4.3	Plasmid Ligation and Transformation	39

3.8.4.4	Selection of the Cells with Gene of Interest	40
3.8.4.5	Plasmid Extraction	40
3.8.4.6	Screening and Verification of Putative Recombinant Clones by PCR	40
3.8.4.7	Cycling Conditions of PCR for Verifying Plasmid Product	41
3.9	Extraction of RNA from Blood Samples	41
3.9.1	One-Step RT-PCR of Extracted RNA from Blood Samples of Animals	42
4	RESULTS	43
4.1	Prevalence of JEV Antibodies in Livestock, Avian and Companion Animals	43
4.2	Risk Factors	44
4.2.1	Dogs	45
4.2.2	Cats	47
4.2.3	Cattle	49
4.2.4	Pigs	51
4.2.5	Buffaloes	53
4.2.6	Avian	55
4.3	Molecular Detection of Japanese encephalitis virus	57
4.3.1	Amplification of NS3 and E of JEV reference strain by One-step RT-PCR	57
4.3.2	Cloning of NS3 and E Genes of JEV Reference Strain using Molecular Cloning Steps	58
4.3.2.1	Molecular Cloning of Envelope Gene of Reference JEV Strain	58
4.3.2.2	Molecular Cloning of Non-Structures 3 (NS3) Gene of Reference JEV Strain	59
4.4	Detection of Japanese encephalitis virus in Blood samples of Companion, Livestock and Avian by One-Step RT-PCR	60
4.4.1	Detection of JEV in Dogs	60
4.4.2	Detection of JEV in Cats	61
4.4.3	Detection of JEV in Cattle	62
4.4.4	Detection of JEV in Pigs	63
4.4.5	Detection of JEV in Buffaloes	64
4.4.6	Detection of JEV in Avian	65
5	DISCUSSION	67
5.1	Seroprevalence of JEV in Livestock, Companion and Avian	67
5.1.1	Dogs	67
5.1.2	Cats	67
5.1.3	Cattle	68
5.1.4	Pigs	69
5.1.5	Buffaloes	70
5.1.6	Avian	71
5.2	Risk Factors and Multivariate Analysis	72
5.2.1	Overall Risk Factors	72

5.2.1.1 Dogs	72
5.2.1.2 Cats	73
5.2.1.3 Cattle	73
5.2.1.4 Pigs	74
5.2.1.5 Buffaloes	75
5.2.1.6 Avian	75
5.3 Molecular Studies on JEV	75
6 CONCLUSION AND RECOMMENDATIONS	78
6.1 Conclusion	78
6.2 Recommendations for future research	78
REFERENCES	80
APPENDICES	96
BIODATA OF STUDENT	128
LIST OF PUBLICATIONS	129

LIST OF TABLES

Table	Page
2.1 Structure genes and their functions of Japanese encephalitis virus	6
2.2 Non-structure genes and their function of Japanese encephalitis virus	7
3.1 Total numbers of blood samples from various animals and birds for Japanese encephalitis virus studies	28
3.2 Distribution of animal samples and their percentage of blood samples from three states of Malaysia	28
3.3 Total number of animals' blood samples used in Japanese encephalitis virus detection by DAS-ELISA assay	32
3.4 Primers used in this study to amplify Japanese encephalitis virus genes	36
3.5 Total number of blood samples from animals used for molecular detection study	42
4.1 Univariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among dogs in Malaysia	46
4.2 Multivariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among dogs in Malaysia	47
4.3 Univariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among cats in Malaysia	48
4.4 Multivariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among cats in Malaysia	49
4.5 Univariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among cattle in Malaysia	50
4.6 Multivariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among cattle in Malaysia	51
4.7 Univariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among pigs in Malaysia	52
4.8 Multivariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among pigs in Malaysia	53

4.9	Univariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among buffaloes in Malaysia	54
4.10	Multivariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among buffaloes in Malaysia	55
4.11	Univariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among avian in Malaysia	56
4.12	Multivariate logistic regression analysis for the risk of Japanese encephalitis virus seropositivity among avian in Malaysia	57



LIST OF FIGURES

Figure	Page
2.1 Three-dimensional image of immature (left) and mature (right) of Flavivirus	4
2.2 The structure of Flavivirus (A) Schematic diagram of a cross-section of a Flavivirus; (B) whole mature Flavivirus; (C) mature Japanese encephalitis virus	4
2.3 The linear genomic structures of Japanese encephalitis virus and their genes functions	6
2.4 Global distribution of Japanese encephalitis virus disease in South-East Asia. Affected countries in South East Asia are Pakistan, India, Bangladesh China, Burma, Vietnam, Nepal, Thailand, Malaysia, Philippines, Sri Lanka, Korea, Japan, Taiwan, Russia and Australia as well as the European countries including Italy	9
2.5 Overall transmission of Japanese encephalitis virus. The amplifying animals are ardeidae birds (Heron, Egret), pigs and bats. Human and horses are dead-end hosts, whereas buffalo, cattle, dog and cats are the infected hosts without showing any symptoms	13
2.6 Examples of wild birds as the amplifying agents for Japanese encephalitis virus, Great Egret (A), Black Crowned Night Heron (B) and Cattle Egret (C)	16
3.1a Areas of sampling for JEV studies in the states of Perak and Selangor. Blood samples were collected from animals and birds in different states. Pig samples: Tapah, Perak; Sepang, Selangor. Cattle samples: Sabak Bernam, Selangor; Serdang, Selangor. Cat Samples: Gopeng, Perak; UVH-UPM, Selangor; Putrajaya, Selangor. Dog Samples: Gopeng, Perak and Kepong, Selangor. Poultry samples: Tanjong Piandang, Perak and Jenderam Hulu, Selangor. Wild bird samples: Parit Buntar, Perak. Migratory birds samples: Kapar, Selangor. Captivity water birds samples: Putrajaya, Selangor	25
3.1b Areas of sampling for JEV studies in the state of Sabah. Buffalo samples: Telupid, Sabah	26
3.2 Trapping of migratory birds for JEV study using Mist-net method. The bird (Greater Sand Plover) was trapped in the baggy pocket of the net. Picture was taken at the sea side of Kapar, Selangor	29

3.3	Trapping of wild birds for Japanese encephalitis virus study using Hand-net method. The bird (Cattle Egret) was trapped in the aluminium hoop. Picture was taken at a pond in Parit Buntar, Perak	30
3.4	The mechanism of double-antigen sandwich ELISA (DAS-ELISA)	31
3.5	Diagrams of Japanese encephalitis virus linear genome	36
4.1	Distribution of Japanese encephalitis virus antibodies in different species of animals detected by DAS-ELISA. For livestock, the highest seroprevalence were detected in pigs (44.4%), followed by buffaloes (33.3%) and cattle (32.2%). For companion animals, the highest seroprevalence were detected in dogs (80%), followed by cats (14.4%). For avian, the highest seroprevalence were detected in poultry (24.4%) followed by captivity water birds (4.4%)	44
4.2	RT-PCR assay of JEV reference strain using two primers sets namely primer NS3 F/R and primer E F/R. Electrophoresis was carried out on 1.5% (w/v) agarose gel. Lane M1: 1 kb DNA ladder (Promega, USA), Lane M2: 100 bp (Promega, USA), Lane 1-3 are positive amplification of E gene with 1.5 kb product, Lane 4-6 are positive amplification of NS3 gene with 602 kb product	58
4.3	Agarose gel electrophoresis of PCR products of pCAMBIA1301+ E gene clone of Japanese encephalitis virus amplified by primer E F/R which produce 1.5 kb of E gene and 11 kb of plasmid pCAMBIA1301. Electrophoresis was carried out on 1.5% (w/v) agarose gel. Lane M: 1 Kb DNA ladder (Promega, USA), Lane 1-5 are positive amplification of E gene	59
4.4	Agarose gel electrophoresis of PCR products of pCAMBIA1301+ NS3 gene clone of Japanese encephalitis virus amplified by primer NS3 F/R which produce 602 bp of NS3 gene and 11 kb of plasmid pCAMBIA1301. Electrophoresis was carried out on 1.5% (w/v) agarose gel. Lane M1: 100 bp DNA ladder (Vivantis, USA), Lane M2: 1 kb DNA ladder (Promega, USA), Lane 1-5 are positive amplification of NS3 gene	60
4.5	One-step RT-PCR assay for the detection of Japanese encephalitis virus in blood samples of dogs by using primer pairs NS3 F/ R to produce 602 bp. Electrophoresis was carried out on 1.5% (w/v) agarose gel. All blood samples show no amplification (Lane 1-20). PC = Positive control with 602 bp, NC = no template control, Lane M = 100 bp DNA marker (Vivantis, USA)	61

4.6	One-step RT-PCR assay for the detection of Japanese encephalitis virus in blood samples of cats by using primer pairs NS3 F/ R to produce 602 bp. Electrophoresis was carried out on 1.5% (w/v) agarose gel. All blood samples show no amplification (Lane 1-23). PC = Positive control with 602 bp, NC = No template control, Lane M: 100 bp DNA marker (Vivantis, USA)	62
4.7	One-step RT-PCR assay for the detection of Japanese encephalitis virus in blood samples of cattle by using primer pairs NS3 F/ R to produce 602 bp. Electrophoresis was carried out on 1.5% (w/v) agarose gel. All blood samples show no amplification (Lane 1-23). PC = Positive control with 602 bp, NC = No template control, Lane M: 100 bp DNA marker (Vivantis, USA)	63
4.8	One-step RT-PCR assay for the detection of Japanese encephalitis virus in blood samples of pigs by using primer pairs NS3 F/ R to produce 602 bp. Electrophoresis was carried out on 1.5% (w/v) agarose gel. All blood samples show no amplification (Lane 1-23). PC = Positive control with 602 bp, NC = No template control, Lane M: 100 bp DNA marker (Vivantis, USA)	64
4.9	One-step RT-PCR assay for the detection of Japanese encephalitis virus in blood samples of buffaloes by using primer pairs NS3 F/ R to produce 602 bp. Electrophoresis was carried out on 1.5% (w/v) agarose gel. All blood samples show no amplification (Lane 1-27). PC = Positive control with 602 bp, NC = No template control, Lane M: 100 bp DNA marker (Vivantis, USA)	65
4.10	One-step RT-PCR assay for the detection of Japanese encephalitis virus in blood samples of avian by using primer pairs NS3 F/ R to produce 602 bp. Electrophoresis was carried out on 1.5% (w/v) agarose gel. All blood samples show no amplification (Lane 1-27). PC = Positive control with 602 bp, NC = No template control, Lane M: 100 bp DNA marker (Vivantis, USA)	66
5.1	Parameters used in JEV studies to examine risk factors associated with JEV infection among animals. The source of animals (25%), breeding types (25%) and age (25%) show highest correlation to JEV seropositivity.	72

LIST OF ABBREVIATIONS

α	Alpha
Ab	Antibodies
amps	Ampere
BBB	Blood-Brain Barrier
bp	Base pair
CD	Cluster of Differentiation
CI	Confidence Intervals
C-Medium	Culture Medium
CNS	Central Nervous System
CSF	Cerebrospinal fluid
Cx.	<i>Culex</i>
°C	Degree Celsius
DENV	Dengue virus
DI	Deionized water
DNA	Deoxyribonucleic acid
dNTP	Dinucleotide triphosphate
ddPCR	Droplet Digital Polymerase Chain Reaction
<i>E.coli</i>	<i>Escherichia coli</i>
EDTA	Ethylenediaminetetraacetic acid
ELISA	Enzyme-Linked Immunosorbent Assay
ER	Endoplasmic reticulum
FVs	Flaviviruses
γ	Gamma
G	Gauge
<i>g</i>	Gravity

HRP	Horse Radish Peroxidase
IBM	International Business Machines Corporation
ID	Identity document
IgG	Immunoglobulin G
IgM	Immunoglobulin M
in	Inch
JE	Japanese encephalitis
JE-IgG	Japanese encephalitis Immunoglobulin G
JEV	Japanese encephalitis virus
kb	Kilo bases
kDa	Kilodalton
km	kilometer
LB agar	Luria broth agar
LIV	Louping ill virus
m	Meter
M	Marker
MHC	Major Histocompatibility Complex
μL	Microliter
mL	Milliliter
MVEV	Murray Valley encephalitis virus
<i>n</i>	Sub-total Population
<i>N</i>	Total number of Population
NA	Not Applicable
NAT	Nucleic acid test
NCBI	National Center for Biotechnology Information
NCR	Noncoding Region
NiV	Nipah virus

nm	Nanometer
NS	Non-structure
OD	Optical density
OD _{NC}	Optical density for Negative control
OD _{PC}	Optical density for Positive Control
OR	Odds Ratio
ORF	Open Reading Frame
%	Percentage
PBMC	Peripheral Blood Mononuclear Cell
PCR	Polymerase Chain Reaction
pH	Power of Hydrogen
prM	Premembrane
qRT-PCR	Quantitative Polymerase Chain Reaction
Ref	Reference
RNA	Ribonucleic acid
rpm	Revolutions per minute
RT-PCR	Reverse Transcription Polymerase Chain Reaction
S E	Standard Error
SLEV	St. Louis encephalitis virus
Spp.	Species
SPSS	Statistical Package for the Social Sciences
β	Beta
TAE	Tris-acetate Ethylenediaminetetraacetic acid
TCID	Tissue Culture Infective Dose
TNF	Tumor Necrosis Factor
U ml ⁻¹	Unit Per Milliliter
UK	United Kingdom

UPM	Universiti Putra Malaysia
USA	United State of America
UV	Ultraviolet
UVH	University Veterinary Hospital
V	Volt
V	Vision
VN	Virus- neutralizing
VS	Versus
WNV	West Nile virus
WSLV	Wesselsbron virus
X	Times
X-gal	Indoxyl β -galactosidase
ZIKV	Zika virus

CHAPTER 1

INTRODUCTION

Japanese encephalitis (JE) is a vector-borne zoonotic disease spread by the bite of *Culex* spp. JE disease responsible for 30,000 to 70,000 cases every year and 10,000 deaths cases in the eastern Asia (Schuh *et al.*, 2010; Campbell *et al.*, 2011). Japanese encephalitis virus (JEV) causes central nervous system (CNS) disease in humans and animals. The virus induces abortion and weak piglets and causes fatal outcomes in humans and horses (Mackenzie *et al.*, 2007). Wading birds, pigs and bats are considered as the amplifying agents, whereas humans and horses are the dead end host. Humans and other animals such as horses, cats and dogs are susceptible to JEV infection; however, they are not capable of developing enough viremia to turn mosquitoes into infected vectors. Some studies indicated that the high temperature during summer can lead to the high number of JE cases. The spread of JEV is correlated with rapid globalisation, population boom and global climate changes due to industrialisation and deforestation (Ghosh and Basu, 2009).

JEV is a positive-sense single-stranded RNA virus with 50 nm in diameter. JEV RNA consists of 10 functional proteins, which made up three structural proteins (Capsid, Premembrance, and Envelope) and seven non-structural proteins (NS1, NS2a, NS2b, NS3, NS4a, NS4b and NS5). JEV strains can be classified into five genotypes based on gene E namely the JEV (GI), JEV (GII), JEV (GIII), JEV (GIV) and JEV (GV) (Schuh *et al.*, 2013). JEV belongs to the *Flavivirus* genus under the family of *Flaviviridae*. Other members of Flaviviruses include West Nile virus (WNV), Zika virus (ZIKV), Tick-borne encephalitis virus (TBEV), Spondweni virus (SPOV), Yellow fever virus (YF), Murray Valley encephalitis virus (MVEV), St. Louis encephalitis virus (SLEV), Dengue virus (DENV) and Usutu virus (USUV), which are the few examples of common viruses.

JE cases in human were first reported in Japan in 1924. Following that, more cases were reported in many parts of Asia especially in the South Asia countries namely Thailand, Japan, China, South Korea, Singapore, Taiwan, Vietnam, Malaysia, India, Burma, Cambodia, Laos, Philippines and Sri Lanka (Schuh *et al.*, 2013). A prototype virus of Nakayama strain was isolated in Japan from the brain of encephalitis patient in 1935 and later in 1940, another strain known as Beijing JEV strain was isolated in China (Yin *et al.*, 1997).

JEV has been shown to circulate in its enzootic stage, presenting a risk to unvaccinated or nonimmune humans and animals. JE cases usually occur in the absence of intensive pig farming and where pig density is relatively low to other livestock in countries including Bangladesh and India (Lord *et al.*, 2015). This indicates that other hosts can play a vital role in the transmission of JEV. Several factors contributing to JEV transmission include paddy cultivation, rubber plantation, climate change, bat population, semi-intensive pig farming and migratory bird. Most of the time,

mosquitoes become inactive during winter season; however, JE cases have been continuously detected during this season in countries like Japan, China, Korea and Russia (Mackenzie *et al.*, 2007). The actual transmission cycle of JEV is yet to be revealed due to the variation of animals that can be infected by JEV.

In Malaysia, there are a higher number of high-risk zones for JEV especially in pig farmers, wetlands and paddy cultivation areas. JE cases in human are reported annually among children below 15 years old even though JE is not a serious infection disease in Malaysia. The seroprevalence rate of JEV was not reported since 20 years back among livestock and companion animals in high-risk areas in Malaysia. Besides, the risk factors contributing to seroprevalence rate were not fully documented in Malaysia. The molecular characteristics of JEV from birds, buffalo, cats, cattle, dogs and pigs are still unclear. Moreover, the roles of these animals in disease transmission and as a sentinel to human JE in these high zones within Malaysia have not been fully discovered.

1.1 Hypothesis

In this study, the following alternative hypothesis was proposed:

Ho: JEV antibodies IgG and antigen are not detected in livestock (buffaloes, cattle and pigs), avian and companions animals (cats and dogs) at the high-risk areas in Malaysia.

Hi: JEV antibodies IgG and antigen are detected in livestock (buffaloes, cattle and pigs), avian and companions animals (cats and dogs) at the high-risk areas in Malaysia.

1.2 The objectives of this study are:

- I. To determine the seroprevalence rate of JEV in livestock (buffaloes, cattle and pigs), avian and companions animals (cats and dogs) using JEV-IgG ELISA.
- II. To identify the risk factors associated with seroprevalence of JEV in livestock (buffaloes, cattle and pigs), avian and companions animals (cats and dogs) using univariate and multivariate logistic regression analysis.
- III. To detect JEV antigen in blood samples of livestock (buffaloes, cattle and pigs), avian and companion animals (cats and dogs) using one step RT-PCR.

REFERENCES

- Allen, R., Sims, R. A., & Sulkin, S. E. (1964). Studies with brown adipose tissue. I. Persistence of rabies virus in bat brown fat. *American Journal of Hygiene*, 80, 11–24.
- Arias, C.F., Preugschat, F. & Strauss, J.H. (1993). Dengue 2 virus NS2B and NS3 form a stable complex that can cleave NS3 within the helicase domain. *Virology*, 193, 888–899.
- Arshad, S.S., Amira, P., Karunanithi, K.K., Faruku, B., Selvarajah, G.T., Jalila, A., & et al. (2016, 18th – 21st April). *Detection of Japanese encephalitis virus in Free Roamer Cats in Malaysia*. Poster session presented at the meeting of Federation of Asian Small Veterinary Associations Congress, Kuala Lumpur, Malaysia.
- Bai, Y., Xu, Z., Zhang, J., Mao, D., Luo, C., He, Y., & et al. (2014). Regional impact of climate on Japanese encephalitis in areas located near the three gorges dam. *PLoS ONE*, 9(1), 1-9.
- Banerjee, K., Mahadev, P.V.M., Ilkal, M.A., Mishra, A.C., Dhanda, V., Modi, G.B., & et al. (1979). Isolation of Japanese encephalitis virus from mosquitoes collected in Bankura district, West Bengal, during October 1974 to December 1975. *Indian Journal of Medical Research*, 69, 201-205.
- Beck-Johnson, L.M., Nelson, W.A., Paaijmans, K.P., Read, A.F., Thomas, M.B., & Bjørnstad, O. N. (2013). The Effect of Temperature on Anopheles Mosquito Population Dynamics and the Potential for Malaria Transmission. *PLoS ONE*, 8(11): e79276.
- Bhattacharya, S., & Basu, P. (2014). Japanese Encephalitis Virus (JEV) infection in different vertebrates and its epidemiological significance: a Review. *International Journal of Fauna and Biological Studies*, 1(6), 32–37.
- Boyle, D.B., Dickerman, R.W., & Marshall, I.D. (1983). Primary viraemia responses of herons to experimental infection with Murray Valley encephalitis, Kunjin and Japanese encephalitis viruses. *The Australian Journal of Experimental Biology and Medical Science*, 61, 655-664.
- Breed, A.C., Smith, C.S. & Epstein, J.H. (2006). Winged wanderers: Long distance movements of flying foxes. In D.W. Macdonald (Ed.), *The Encyclopedia of Mammals*, Oxford: Oxford University Press.
- Brinton, M.A. (2002). The molecular biology of West Nile virus: A new invader of the western hemisphere. *Annual Review of Microbiology*, 56, 371–402.
- Bunnag, T., Singharaj, P., Sindhusen, S., & Punyagupta, S. (1976). Japanese encephalitis in Nakorn Rajsima. *Journal of the Medical Association of Thailand*, 50(9), 590-598.

- Burke, D.S., & Leake, C.J. (1988). Japanese encephalitis. In T.P. Monath. (Ed.), *The Arboviruses: Epidemiology and Ecology*. Boca Raton: CRC Press.
- Burke, D.S., Nisalak, A., Ussery, M.A., Laorakpongse, T., & Chantavibul, S. (1985). Kinetics of IgM and IgG response to Japanese encephalitis virus in human serum and cerebrospinal fluid. *The Journal of Infectious Diseases*, 151, 1093–1099.
- Campbell, G.L., Hills, S.L., Fischer, M., Jacobson, J.A., Hoke, C.H., Hombach, J.M., & et al. (2011). Estimated global incidence of Japanese encephalitis: a systematic review. *Bulletin World Health Organization*, 89, 766–774.
- Cappelle, J., Duong, V., Pring, L., Kong, L., Yakovlev, M., Prasetyo, D. B., & et al. (2016). Intensive Circulation of Japanese Encephalitis Virus in Peri-urban Sentinel Pigs near Phnom Penh, Cambodia. *PLOS Neglected Tropical Diseases*, 10(12), e0005149.
- Cha, G.W., Lee, E.J., Lim, E.J., Sin, K.S., Park, W.W., Jeon, D.Y., & et al. (2015). A novel immunochromatographic test applied to a serological survey of Japanese encephalitis virus on pig farms in Korea. *PLoS ONE*, 10(5), 1–12.
- Chan, M., & Johansson, M.A. (2012). The Incubation Periods of Dengue Viruses. *PLoS ONE*, 7(11), e50972.
- Cheit, T.W. (2012). *Detection of JEV in Wild Boars and Comparison of Blood Profiles between Wild Boars and Domestic Pigs in Selangor* (Unpublished degree dissertation). Universiti Putra Malaysia, Serdang, Malaysia.
- Chernesky, (1996). Traditional Serological Tests. In M. Brian & H. Kangro (Eds.), *Virology Methods Manual* (pp. 107-122). New York: Elsevier Ltd.
- Christopher, S., & Reuben, R. (1971). Studies on the mosquitoes of North Arcot district, Madras state, India Part 4 Host preference as shown by precipitin tests. *Journal of Medical Entomology*, 8(2), 314-318.
- Chu, C.M. (1982). Viral diseases in China. In J.S. MacKenzie, (Ed.), *Viral Diseases in South-East Asia and the Western Pacific*. New York: Academic.
- Chu, F.T., Wu, J.P., & Teng, C.H. (1940). Acute encephalitis in children; clinical and serologic study of ten epidemic cases. *Chinese Medical Journal*, 58, 68-78.
- Clarke, D. H., & Casals, J. (1958). Techniques for hemagglutination and hemagglutination-inhibition with arthropod-borne viruses. *The American Journal of Tropical Medicine and Hygiene*, 7, 561-573.
- Cleton, N. B., Bosco-Lauth, A., Page, M. J., & Bowen, R. A. (2014). Age-related susceptibility to Japanese encephalitis virus in domestic ducklings and chicks. *American Journal of Tropical Medicine and Hygiene*, 90(2), 242–246.
- Cruickshank, E. K. (1951). Acute encephalitis in Malaya. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 45(1), 113.-118.

- Cui, J., Counor, D., Shen, D., Sun, G., He, H., Deubel, V., & et al. (2008). Detection of Japanese encephalitis virus antibodies in bats in Southern China. *The American Journal of Tropical Medicine and Hygiene*, 78, 1007–1011.
- Daniels, P.W., Williams, D.T., & Mackenzie, J.S. (2002). Japanese encephalitis virus. In A. Morrilla, K.J. Yoon, J.J. Zimmermen (Eds.), *Trends in Emerging Viral Infections of Swine* (pp. 49-60). Ames: Iowa State University Press.
- Dhanda, V., Banerjee, K., Deshmukh, P.K., & Ilkal, M.A. (1977). Experimental viremia and transmission of Japanese encephalitis virus by mosquitoes in domestic ducks. *Indian Journal of Medical Research*, 66, 881–888.
- Diercks, F. H., & Hammon, W. M. (1958). Hamster kidney cell tissue cultures for propagation of Japanese B encephalitis virus. *Proceedings of the Society for Experimental Biology and Medicine*, 97, 627-632.
- Emidi, B., Kisinza, W. N., Mmbando, B. P., Malima, R., & Mosha, F. W. (2017). Effect of physicochemical parameters on *Anopheles* and *Culex* mosquito larvae abundance in different breeding sites in a rural setting of Muheza , Tanzania. *Parasites & Vectors*, 10(304), 1–12.
- Endy, T.P., & Nisalak, A. (2002). Japanese encephalitis virus: ecology and epidemiology. *Current Topics in Microbiology and Immunology*, 267, 11-48.
- Flavivirus-ViralZone. (n.d.). Retrieved January 10, 2017 from *ViralZone*: <https://owl.english.purdue.edu/owl/resource/560/10/>
- Fu, D.W., & Zhang, P.F. (1996). Establishment and characterization of Japanese B encephalitis virus persistent infection in the Sf9 insect cell line. *Journal of Biologicals*, 24, 225-233.
- George, S., Prasad, S.R., Rao, J.A., Yergolkar, P.N., & Setty, C.V. (1987). Isolation of Japanese encephalitis & West Nile viruses from fatal cases of encephalitis in Kolar district of Karnataka. *Indian Journal of Medical Research*, 86, 131-134.
- Ghosh, D., & Basu, A. (2009). Japanese encephalitis a pathological and clinical perspective. *PLoS Neglected Tropical Diseases*, 3(9), 1-7.
- Go, Y.Y., Balasuriya, U. B. R., & Lee, C.K. (2014). Zoonotic encephalitis caused by arboviruses: transmission and epidemiology of alphaviruses and *Flaviviruses*. *Clinical and Experimental Vaccine Research*, 3(1), 58–77.
- Grossman, R.A., Gould, D., Smith, T., Johnsen, D.O., & Pantuwatana, S. (1973). Study of Japanese encephalitis virus in Chiangmai Valley, Thailand. I. *Introduction and study design. American Journal of Epidemiology*, 98(2), 111-120.
- Gubler, D.J. (2002). The Global Emergence Resurgence of Arboviral Diseases as Public Health Problems. *Archives of Medical Research*, 33 (4), 330–342.

- Gulati, B.R., Singha, H., Singh, B.K., Virmani, N., Kumar, S., & Singh, R.K. (2012). Isolation and genetic characterization of Japanese encephalitis virus from equines in India. *Journal of Veterinary Science*, 13(2), 111–118.
- Halstead, S.B., & Jacobson, J. (2008). Japanese encephalitis vaccines. In S.A. Plotkin, W.A. Orenstein, & P.A. Offit (Eds.), *Vaccine* (pp. 311-352). Philadelphia: Elsevier.
- Hammon, W.M., Reeves, W.C., & Sather, G.E. (1951). Japanese B encephalitis virus in the blood of experimentally inoculated birds; epidemiological implications. *American Journal of Epidemiology*, 53(3), 249–261.
- Hanna, J.N., Ritchie, S.A., Phillips, D.A., Shield, J., Bailey, M.C., Mackenzie, J.S., & et al. (1996). An outbreak of Japanese encephalitis in the Torres Strait, Australia. *Medical Journal of Australia*, 165, 256-260.
- Hasegawa T., Takehara Y., & Takahashi K. (1975). Natural and experimental infections of Japanese tree sparrows with Japanese encephalitis virus. *Archives Virology*, 49 (4), 373-376.
- Henderson, J.R., & Taylor, R.M. (1959). Arthropod-borne virus plaques in agar overlaid tube cultures. *Proceedings of the Society for Experimental Biology and Medicine*, 101, 257–259
- Hotta, S., Aoki, H., Samoto, S., Yasui, T., & Kawabe, M. (1970). Virologic-epidemiological studies on Indonesia. II. Measurement of anti-arbovirus antibodies in sera from residents of Lombok, South Sumatra, and West Java, in comparison with results concerning sera from residents of Japanese main islands. *Kobe Journal of Medical Sciences*, 16(4), 215-234.
- Hotta, S., Aoki, H., Yasui, T., & Samoto, S. (1967). Virologic-epidemiological studies on Indonesia. Survey of anti-arboviral antibodies in sera from residents of Lombok, Sumatera and Djawa. *Kobe Journal of Medical Sciences*, 13(3), 221-234.
- Hotta, S., Kuromaru, S., Funasaka, K., & Mizoguti, H. (1964). Experimental infection of dogs with Japanese B encephalitis virus. *Acta Neuropathologica*, 3(5), 494–510.
- Hsu, S.M., Yen, A.M.F. & Chen, T.H.H. (2008). The impact of climate on Japanese encephalitis. *Epidemiology and Infection* 136(7), 980-987.
- Huang, C.H. (1982). Studies of Japanese encephalitis in China. *Advances in Virus Research*, 27, 71–101.
- Igarashi A. (1992). Epidemiology and control of Japanese encephalitis. *World Health Stat Q*, 45, 229.
- Igarashi, A. (2002). Control of Japanese encephalitis in Japan: immunization of humans and animals, and vector control. *Current Topics in Microbiology and Immunology*, 267, 139-152.

- Igarashi, A., Buei, K., Ueba, N., Yoshida, M., Ito, S., Nakamura, H., & et al. (1981). Isolation of viruses from female *Culex tritaeniorhynchus* in *Aedes albopictus* cell cultures. *The American Journal of Tropical Medicine and Hygiene*, 30, 449-460.
- Igarashi, A., Sasao, F., Wungkobkiat, S., & Fukai, K. (1973). Growth of Japanese encephalitis virus in established lines of mosquito cells. *Biken Journal*, 16, 17-23.
- Ilkal, M.A., Dhanda, V., Rao, B.U., George, S., Mishra, A. C., Prasanna, Y., & et al. (1988). Absence of viraemia in cattle after experimental infection with Japanese encephalitis virus. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 82(4), 628–631.
- Impoinvil, D.E., Solomon, T., Schluter, W.W., Rayamajhi, A., Bichha, R. P., Shakya, G., & et al. (2011). The spatial heterogeneity between Japanese encephalitis incidence distribution and environmental variables in Nepal. *PLoS One*, 6, e22192.
- Japanese Encephalitis- Diagnostic Testing CDC. (2015, August 25). Retrieved from *Centers for Disease Control and Prevention* from June 21, 2017, from <http://www.easybib.com/reference/guide/apa/website>
- Jeffries, C.L., & Walker, T. (2015). The Potential Use of Wolbachia-Based Mosquito Biocontrol Strategies for Japanese Encephalitis. *PLoS Neglected Tropical Diseases*, 9(6):e0003576.
- Johnsen, D.O., Edilman, R., Grossman, R. A., Muang- man, D., Pomsdhit, J. & Could, D. J. (1974). Studies of Japanese encephalitis virus in Chiangmai valley, Thailand. V. Animal infections. *American Journal of Epidemiology*, 100, 57-68.
- Joo, H.S., & Chu, R.M. (1999). Japanese B encephalitis. In B.E. Straw, S. D’Allaire, W.L. Mengeling, & D.T. Taylor. (Eds.), *Diseases of Swine*. Malvern: Blackwell Science Ltd.
- Kako, N., Suzuki, S., Sugie, N., Kato, T., Yanase, T., Yamakawa, M., & Shirafuji, H. (2014). Japanese encephalitis in a 114-month-old cow: pathological investigation of the affected cow and genetic characterization of Japanese encephalitis virus isolate. *BMC Veterinary Research*, 10(1), 63.
- Kanamitsu, M., Taniguchi, K., Urasawa, S., Ogata, T., Wada, Y., & Saroso J.S. (1979). Geographic distribution of arbovirus antibodies in indigenous human populations of the Indo- Australian archipelago. *The American Journal of Tropical Medicine and Hygiene*, 28(2), 351-363.
- Kant Upadhyay, R. (2013a). Japanese Encephalitis Virus Generated Neurovirulence, Antigenicity, and Host Immune Responses. *International Scholarly Research Notices: Virology*, 1–24.

- Kant Upadhyay, R. (2013b). Biomarkers in Japanese encephalitis: A review. *BioMed Research International*, 591290, 1-24.
- Katayama, K., Ikeda, A., Saika, K., Ishizaka, M., Onodera, M., & Ito, T. (2000). A case of Japanese encephalitis in a black Japanese Heifer. *Journal of the Japan Veterinary Medical Association*, 53, 293–296.
- Katayama, T., Saito, S., Horiuchi, S., Maruta, T., Kato, T., Yanase, T., & et al. (2013). Nonsuppurative encephalomyelitis in a calf in Japan and isolation of Japanese encephalitis virus genotype 1 from the affected calf. *Journal of Clinical Microbiology*, 51, 3448–3453.
- Kimoto, T., Yamada, T., Ueba, N., Kunita, N., & Kawai, A. (1968). Laboratory diagnosis of Japanese encephalitis: comparison of the fluorescent antibody technique with virus isolation and serologic tests. *Biken Journal*, 11, 157-168.
- King, A.M.Q., Lefkowitz, E., Adams, M. J., & Carstens, E. B. (Eds.). (2005). *Virus Taxonomy: Classification and Nomenclature of Viruses*. United State of America: Elsevier Press.
- Konishi, E., Shoda, M., & Kondo, T. (2006). Analysis of yearly changes in levels of antibodies to Japanese encephalitis virus nonstructural 1 protein in racehorses in central Japan shows high levels of natural virus activity still exist. *Vaccine*, 24(4), 516-524.
- Kuno, G., & Chang, G. J. (2005). Biological Transmission of Arboviruses : Reexamination of and New Insights into Components , Mechanisms , and Unique Traits as Well as Their Evolutionary Trends. *Clinical Microbiology Reviews*, 18(4), 608–637.
- Kuwata, R., Sugiyama, H., Yonemitsu, K., Van Dung, N., Terada, Y., Taniguchi, M., & et al. (2015). Isolation of Japanese encephalitis virus and a novel insect-specific flavivirus from mosquitoes collected in a cowshed in Japan. *Archives of Virology*, 160(9), 2151–2159.
- Lam, K.H.K., Ellis, T.M., Williams, D.T., Lunt, R.A., Daniels, P.W., Watkins, K.L., & et al. (2005). Japanese encephalitis in a racing thoroughbred gelding in Hong Kong. *Veterinary Record*, 157(6), 168-174.
- Leake, C.J., Burke, D.S., Nisalak, A., & Hoke, C.H. (1986). Isolation of Japanese encephalitis virus from clinical specimens using a continuous mosquito cell line. *The American Journal of Tropical Medicine and Hygiene*, 35, 45-50.
- Ledermann, J. P., Lorono-Pino, M. A., Ellis, C., Saxton-Shaw, K. D., Blitvich, B. J., Beaty, B. J., & et al. (2011). Evaluation of Widely Used Diagnostic Tests to Detect West Nile Virus Infections in Horses Previously Infected with St. Louis encephalitis Virus or Dengue Virus Type 2. *Clinical and Vaccine Immunology*, 18(4), 580–587.

- Lee, E. J., Cha, G.W., Ju, Y.R., Han, M. G., Lee, W. J., & Jeong, Y. E. (2016). Prevalence of Neutralizing Antibodies to Japanese Encephalitis Virus among High-Risk Age Groups in South Korea, 2010. *PLoS ONE* 11(1), e0147841.
- Lee, H., Min, B., & Lim, Y. (1972). Isolation and serologic studies of Japanese encephalitis virus from snakes in Korea. *Journal of the Korean Medical Association*, 15, 69-74.
- Lee, H.W. (1968). Multiplication and antibody formation of Japanese encephalitis virus in Snakes. II. Proliferation of the virus. *The Seoul Journal of Medicine*, 9, 157-161.
- Legeay, O., Caudrelier, Y., Cordevant, C., Rigottier-Gois, L., & Lange, M. (2001). Waiter, There's an Enteric Pathogen in My Shellfish Soup. *Promega Note*, 78, 13-16.
- Leung, J.Y., Pijlman, G.P., Kondratieva, N., Hyde, J., Mackenzie, J.M., & Khromykh, A. (2008). Role of nonstructural protein NS2A in flavivirus assembly. *Journal of Virology*, 82(10), 4731–4741.
- Li, J., Gao, N., Fan, D., Chen, H., Sheng, Z., & Fu, S. (2016). Cross-protection induced by Japanese encephalitis vaccines against different genotypes of Dengue viruses in mice. *Nature Publishing Group, October 2015*, 1–9.
- Li, L., Lok, S.M., Yu, I.M., Zhang, Y., Kuhn, R.J., Chen, J., & et al. (2008). The flavivirus precursor membrane-envelope protein complex: structure and maturation. *Science*, 319, 1830–1834.
- Lindahl, J. F., Ståhl, K., Chirico, J., Boqvist, S., Thu, H. T. V., & Magnusson, U. (2013). Circulation of Japanese Encephalitis Virus in Pigs and Mosquito Vectors within Can Tho City, Vietnam. *PLoS Neglected Tropical Diseases*, 7(4), 1-8.
- Lindahl, J., Boqvist, S., Ståhl, K., Thu, H. T. V., & Magnusson, U. (2012). Reproductive performance in sows in relation to Japanese Encephalitis Virus seropositivity in an endemic area. *Tropical Animal Health and Production*, 44(2), 239–245.
- Lindenbach, B.D., Thiel, H.J., & Rice, C.M. (2007). *Flaviviridae: The Viruses and Their Replication*. In D.M. Knipe, & P.M. Howley, (Ed.), *Fields Virology* (4th ed.). Philadelphia: Lippincott Williams and Wilkins.
- Liu, Z., Qiu, J., Cheng, F., Chu, Y., Yoto, Y., O'Sullivan, M.G., & et al. (2004). Comparison of the transcription profile of simian parvovirus with that of the human erythrovirus B19 reveals a number of unique features. *Journal of Virology*, 78, 12929–12939.
- Lopez, A. L., Aldaba, J. G., Roque, V. G., Tandoc, A. O., Sy, A. K., Espino, F. E., & et al (2015). Epidemiology of Japanese Encephalitis in the Philippines: A Systematic Review. *PLoS Neglected Tropical Diseases*, 9(3), 1–17.

- Lord, J.S., Gurley, E.S., & Pulliam, J.R.C. (2015). Rethinking Japanese Encephalitis Virus Transmission: A Framework for Implicating Host and Vector Species. *PLoS Neglected Tropical Diseases*, 9(12): e0004074.
- Lorenz, I. C., Kartenbeck, J., Mezzacasa, A., Allison, S.L., Heinz, F.X., & Helenius, A. (2003). Intracellular assembly and secretion of recombinant subviral particles from tick-borne encephalitis virus. *Journal Virology*, 77, 4370–4382.
- Mackenzie, J. S., Childs, J. E., Field, H. E., Wang, L., & Breed, A. C. (2008). The role of bats as reservoir hosts of emerging neurological viruses. *Neurotropic Viral Infections*, 382–406.
- Mackenzie, J.M., & Westaway, E.G. (2001). Assembly and maturation of the Flavivirus Kunjin virus appear to occur in the rough endoplasmic reticulum and along the secretory pathway, respectively. *Journal Virology*, 75, 10787–10799.
- Mackenzie, J.S. (2000). The family *Flaviviridae*: a brief overview of the family with particular reference to members of the Asian and Australasian regions. In S.D Blacksell, (Ed.), *Classical Swine Fever and Emerging Viral Diseases in South-East Asia*. Canberra: Australian Centre for International Agricultural Research.
- Mackenzie, J.S., Johansen, C.A., Ritchie, S.A., van den Hurk, A.F., & Hall, R.A. (2002). Japanese encephalitis as an emerging virus: the emergence and spread of Japanese encephalitis virus in Australasia. *Current Topics in Microbiology and Immunology*, 267, 49–73.
- Mackenzie, J.S., Williams, D.T., & Smith, D.W. (2007). Japanese Encephalitis Virus: The Geographic Distribution, Incidence, and Spread of a Virus with a Propensity to Emerge in New Areas. *Perspectives in Medical Virology*, 16(06), 201–268.
- MacLachlan, N.J. & Dubovi, E.J. (Eds.) (2011). *Fenner's Veterinary Virology*. San Diego: Elsevier Academic Press.
- Maes, D., Nauwynck, H., Rijsselaere, T., Mateusen, B., Vyt, P., De Kruif, A., & et al. (2008). Diseases in swine transmitted by artificial insemination: An overview. *Theriogenology* 70, 1337–1345.
- Mani, T.R., Rao, C.V., Rajendran, R., Devaputra, M., Prasanna Y, Hanumaiah, & et al. (1991). Surveillance for Japanese encephalitis in villages near Madurai, Tamil Nadu, India. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 85(2), 287-291.
- Masuoka, P., Klein, T.A., Kim, H.C., Claborn, D.M., Achee, N., Andre, R., & et al. (2010). Modeling the distribution of *Culex tritaeniorhynchus* to predict Japanese encephalitis distribution in the Republic of Korea. *Geospatial Health*, 5, 45–57.

- Meyer, D., Fritsche, S., Luo Y., Engemann, C., Blome, S., Beyerbach, M., & et al., (2017). The double-antigen ELISA concept for early detection of E^{ms}-specific classical swine fever virus antibodies and application as an accompanying test for differentiation of infected from marker vaccinated animals. *Transboundary and Emerging Diseases*, September 2016, 1–10.
- Mitchell, C.J., Chen, P.S. & Boreham, P.F.L. (1973). Host-feeding patterns and behavior species in an endemic area of Japanese encephalitis, *Bulletin of the World Health Organization*, 49(3), 293-299.
- Miura, T., & Takahashi, R.M. (1976). Effect of a synthetic pyrethroid SD 43775, on non-target organisms, when utilised as a mosquito larvicide. *Mosquito News*, 30, 322-326.
- Mogi, M. (1983). Relationship between number of human Japanese encephalitis cases and summer meteorological conditions in Nagasaki, Japan. *The American Journal of Tropical Medicine and Hygiene* 32, 170–174.
- Mukhopadhyay, S., Kuhn, R.J. & Rossmann, M.G. (2005). A structural perspective of the flavivirus life cycle. *Nature Reviews Microbiology*, 3, 13–22.
- Mwandawiro, C., Boots, M., Tuna, N., & Suwonkerd, W. (2000). Heterogeneity in the host preference of Japanese encephalitis vectors in Ching Mai, northern Thailand. *Transactions of the royal society of tropical medicine and hygiene*, 94, 238–242.
- Nemeth, N., Bosco-Lauth, A., Oesterle, P., Kohler, D., & Bowen, R. (2012). North American birds as potential amplifying hosts of Japanese encephalitis virus. *American Journal of Tropical Medicine and Hygiene*, 87(4), 760–767.
- Nidaira, M., Taira, K., Itokazu, K., Okano, S., Kudaka, J., Nakamura, M., & et al. (2008). Detection of Japanese encephalitis virus genome in Ryukyu wild boars (*Sus scrofa riukiuanus*) in Okinawa, Japan. *Japanese Journal of Infection Diseases*, 61, 164–165.
- Nitatpattana, N., Le Flohic, G., Thongchai, P., Nakgoi, K., Palaboodeewat, S., Khin, M., & et al. (2011). Elevated Japanese Encephalitis Virus Activity Monitored by Domestic Sentinel Piglets in Thailand. *Vector-Borne and Zoonotic Diseases*, 11(4), 391–394.
- Nowak, T. & Wengler, G. (1987). Analysis of disulphides present in the membrane proteins of West Nile flavivirus. *Virology* 156, 127–137.
- Oda, K., Igarashi, A., Kheong, C. T., Hong, C. C., Vijayamalar, B., Sinniah, M., & et al. (1996). Cross- sectional serosurvey for Japanese encephalitis specific antibody from animal sera in Malaysia 1993. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 27, 463-470.
- Ogasa, A., Yokoki, Y., Fujisaki, Y., & Habu, A. (1977). Reproductive disorders in boars infected experimentally with Japanese encephalitis virus. *The Japanese journal of animal reproduction*, 27, 21–26.

- Ogata, M., Nagao, Y., Jitsunari, F., Kitamura, N., & Okazaki, T. (1970). Infection of herons and domestic fowls with Japanese encephalitis virus with specific reference to maternal antibody of hen (epidemiological study on Japanese encephalitis 26). *Acta Medica Okayama*, 24, 175-184.
- Ohno, Y., Sato, H., Suzuki, K., Yokoyama, M., Uni, S., Shibasaki, T., & et al. (2009). Detection of antibodies against Japanese encephalitis virus in raccoons, raccoon dogs and wild boars in Japan. *The Journal of Veterinary Medical Science*, 71, 1035-1039.
- Olson, J.G., Ksiazek, T.G., Lee, V.H., Tan, R., & Shope, R.E. (1985). Isolation of Japanese encephalitis virus from *Anopheles annularis* and *Anopheles vagus* in Lombok, Indonesia. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 79, 845-847.
- Ooi, M.H., Lewthwaite, P., Lai, B.F., Anand Mohan, Clear, D., Lim, L., & et al. (2008). The epidemiology, clinical features, and long-term prognosis of Japanese encephalitis in central Sarawak, Malaysia, 1997-2005. *Clinical Infectious Diseases* 47, 458-468.
- Oya, A., Doi, R., Shirasaka, A., Yabe, S., & Sasa, M. (1983). Studies of Japanese encephalitis virus infection of reptiles. I. Experimental infection of snakes and lizards. *The Japanese journal of experimental medicine*, 53, 117-123.
- Paily, K.P., Chandhiran, K., Vanamail, P., Kumar, N.P., & Jambulingam, P. (2013). Efficacy of a mermithid nematode *Romanomermis iyengari* (Welch) (Nematoda: *Mermithidae*) in controlling tree hole-breeding mosquito *Aedes albopictus* (Skuse) (Diptera: *Culicidae*) in a rubber plantation area of Kerala, India. *Parasitology Research*, 112, 1299-1304.
- Patel, P., Landt, O., Kaiser, M., Faye O., Koppe, T., Lass, U., & et al. (2013). Development of one-step quantitative reverse transcription PCR for the rapid detection of flaviviruses. *Virology Journal*, 10(58), 1-11.
- Paterson, P.Y., Ley, H.L., Wisserman, J.R., Pond, W.L., Smadel, J.E., Diereks, F.H., & et al. (1952). Japanese encephalitis in Malaya. I Isolation of virus and serologic evidence of human and equine infections. *The American Journal of Tropical Medicine and Hygiene*, 56, 320-330.
- Peiris, J.S., Dittus, W.P., & Ratnayake, C.B. (1993). Seroepidemiology of dengue and other arboviruses in a natural population of toque macaques (*Macaca sinica*) at Polonnaruwa, Sri Lanka. *Journal of Medical Primatology*, 22, 240-245.
- Pfeffer, M., & Dobler, G. (2010). Emergence of zoonotic arboviruses by animal trade and migration. *Parasites & Vectors*, 3(1), 35.
- Phanthumachinda, B. (1995). Ecology and biology of Japanese encephalitis vectors. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 26(3), 11-16.

- Phukan, A.C., Borah, P.K., & Mahanta, J. (2004). Japanese encephalitis in Assam, northeast India. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 35(3), 618-622.
- Porterfield, J.S. (1959). Plaque production with yellow fever and related arthropod-borne viruses. *Nature*, 183, 1069-1070.
- Prakash, S. (2017). Role of real time PCR in diagnosis of Japanese encephalitis virus in acute encephalitis cases. *Virology and Mycology*, 6(2), 27.
- Prasad, C.B. (2013). *Veterinary Medical Virology*. Delhi: Sathish Serial Publishing House.
- Ravanini, P., Huhtamo, E., Ilaria, V., Crobu, M.G., Nicosia, A.M., Servino, L., & et al. (2012). Japanese encephalitis virus RNA detected in *Culex pipiens* mosquitoes in Italy. *Eurosurveillance*, 17(28), 1-4.
- Ren, X., Fu, S., Dai, P., Wang, H., Li, Y., Li, X., & et al. (2017). Pigsties near dwellings as a potential risk factor for the prevalence of Japanese encephalitis virus in adult in Shanxi, China. *Infectious Diseases of Poverty*, 6(100), 1-11.
- Reuben, R., & Gajanana, A. (1997). Japanese encephalitis in India. *The Indian Journal of Pediatrics*, 64, 43.
- Reuben, R., Thenmozhi, V., Samuel, P.P., Gajanana, A. & Mani, T.R. (1992). Mosquito Blood feeding pattern as a factor in the epidemiology of Japanese encephalitis in southern India. *The American Journal of Tropical Medicine and Hygiene*, 46(6), 654-663.
- Rey, F. A., Heinz, F. X., Mandl, C., Kunz, C., Harrison, S. C., Kuhn, R. J. & et al. (1995). The envelope glycoprotein from tick-borne encephalitis virus at 2 ÅG resolutions. *Nature* 375, 291-298.
- Richards, E.E., Masuoka, P., Brett-Major, D., Smith, M., Klein, T.A., Kim, H.C., & et al. (2010). The relationship between mosquito abundance and rice field density in the Republic of Korea. *International Journal of Health Geographics*, 9, 32.
- Ricklin, M. E., García-Nicolás, O., Brechbühl, D., Python, S., Zumkehr, B., Nougaiède, A., & et al. (2016). Vector-free transmission and persistence of Japanese encephalitis virus in pigs. *Nature Communications*, 7, 10832.
- Rosen, L. (1986). The natural history of Japanese encephalitis virus. *Annual Review of Microbiology*, 40, 395-414.
- Sam, J. I. (2015). The burden of human influenza in Malaysia. *Medical Journal of Malaysia* 70(3), 127-130.
- Samuel, P., & Gajanana, A. (2000). Japanese Encephalitis virus infection in mosquitoes and its epidemiological implications. *ICMR Bulletin*, 30(4), 1-9.

- Sangkawibha, N., Nakornsri, S., & Rojanasuphot, S. (1982). Japanese encephalitis in Thailand. *International Journal of Medical Sciences*, 24, 1.
- Saxena, S.K., Agrawal, P.T., & Nair, M.P.N. (2014). Japanese Encephalitis: A Neglected Viral Disease and Its Impact on Global Health. In S.K. Saxena (Ed.), *Trends in Infectious Diseases*, (pp. 3-16) Rijeka: InTech Publisher.
- Scherer, W.F., Moyer, J.T., & Izumi, T. (1959). Immunologic Studies of Japanese Encephalitis Virus in Japan V. Maternal Antibodies, Antibody Responses and Viremia Following Infection of Swine. *The Journal of Immunology*, 83 (6), 620–626.
- Schuh, A. J., Li, L., Tesh, R. B., Innis, B. L., & Barrett, A. D. T. (2010). Genetic characterization of early isolates of Japanese encephalitis virus: Genotype II has been circulating since at least 1951. *Journal of General Virology*, 91(1), 95–102.
- Schuh, A.J., Guzman, H., Tesh, R. B., & Barrett, A.D. (2013). Genetic diversity of Japanese encephalitis virus isolates obtained from the Indonesian archipelago between 1974 and 1987. *Journal of Vector Borne Diseases*, 13(7), 479–488.
- Scott, T.W. (1988). Vertebrate host ecology. In T.P. Monath (Ed.), *The Arboviruses: Epidemiology and Ecology*. Boca Raton: CRC Press.
- Shimizu, T., Mochizuki, H., Sugawa, Y., Okazaki, K., & Matumoto, M. (1951). Studies on Japanese encephalitis of cattle - 1. Bovine encephalitis caused by natural infection with Japanese encephalitis virus. *National Institute of Animal Health*, 23, 111–118.
- Shimoda, H., Ohno, Y., Mochizuki, M., Iwata, H., Okuda, M., & Maeda, K. (2010). Dogs as Sentinels for Human Infection with Japanese Encephalitis Virus. *Emerging Infectious Diseases*, 16 (7), 1137–1139.
- Shimoda, H., Tamaru, S., Morimoto, M., Hayashi, T., Shimojima, M., & Maeda, K. (2011). Experimental infection of Japanese encephalitis virus in dogs. *The Journal of veterinary medical science-the Japanese Society of Veterinary Science*, 73(9), 1241–1242.
- Shiryaev, S.A., Chernov, A.V., Aleshin, A.E., Shiryaeva, T.N., & Strongin, A.Y. (2009). NS4A regulates the ATPase activity of the NS3 helicase: a novel cofactor role of the non-structural protein NS4A from West Nile virus. *Journal of General Virology*, 90(9), 2081–2085.
- Shortridge, K.F., Ng, M.H., Oya, A., Koboyashi, M., Munro, R., Wong, F., & et al (1974). Arbovirus infections in reptiles, Immunological evidences for a high incidence of Japanese encephalitis virus in the *Cobra Naja naja*. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 68, 454-460.

- Shortridge, K.F., Oya, A., Koboyashi, M., & Yip, D.Y. (1975). Arbovirus infections in reptiles, Studies on the presence of Japanese encephalitis virus antibody in the plasma of the Turtle, *Trionyx sinensis*. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 6, 161-169.
- Simasathien, P., Rohitayodhin, S., Nisalak, A., Singharaj, P., Halstead, S.B., & Russell, P.K. (1972). Recovery of Japanese encephalitis virus from wild caught mosquitoes in Thailand. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 3, 52.
- Simpson, D. I. H., Bowen, E. T. W., Platt, G. S., Way, H., Smith, C. E. G., Peto, S., & et al. (1970). Japanese encephalitis in Sarawak: Virus isolation and serology in a land Dyak village. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 64(4), 503–510.
- Solomon, T., Ooi, M.H., & Mallewa, M. (2007). Chapter 10 Viral Infection of lower motor neurons. *Handbook Clinical Neurology*, 82, 179-206.
- Soman, R.S., Rodrigues, F.M., Guttikar, S.N., & Guru, P.Y. (1977). Experimental viraemia and transmission of Japanese encephalitis virus by mosquitoes in ardeid birds. *Indian Journal of Medical Research*, 66(5), 709-718.
- Stadler, K., Allison, S.L., Schlich, J., & Heinz, F.X. (1997). Proteolytic activation of tick-borne encephalitis virus by furin. *Journal of Virology*, 71, 8475–8481.
- Su, C. L., Yang, C. F., Teng, H. J., Lu, L. C., Lin, C., Tsai, K. H., & et al. (2014). Molecular Epidemiology of Japanese Encephalitis Virus in Mosquitoes in Taiwan during 2005- 2012. *PLoS Neglected Tropical Diseases*, 8(10), e3122.
- Sutherst, R. W. (2004). Global Change and Human Vulnerability to Vector-Borne Diseases. *Clinical Microbiology Reviews*, 17(1), 136–173.
- Swami, R., Radha Kanta Ratho, Baijayantimala Mishra & Mini P. Singh. (2008). Usefulness of RT-PCR for the diagnosis of Japanese encephalitis in clinical samples. *Scandinavian Journal of Infectious Diseases*, 40, 815-820.
- Tan, D.S.K. (1981). Some Zoonotic Diseases Prevalent in Malaysia. *Malaysia Journal of Pathology*, 4, 19-27.
- Tan, R., Nalim, S., Suwasono, H., & Jennings, J.B. (1993). Japanese encephalitis virus isolated from seven species of mosquitoes collected at Semarang Regency, Central Java. *Bulletin penelitian kesehatan: Health studies in Indonesia*, 21, 1–5.
- Tangena, J.A.A., Thamavong, P., Wilson, A.L., Brey, P.T., & Lindsay, S. W. (2016). Risk and Control of Mosquito-Borne Diseases in Southeast Asian Rubber Plantations. *Trends in Parasitology*, 32(5), 402–415.
- Thakur, K. K., Pant, G. R., Wang, L., Hill, C. A., Pogranichniy, R. M., Manandhar, S., & et al. (2012). Seroprevalence of Japanese Encephalitis Virus and Risk Factors Associated with Seropositivity in Pigs in Four Mountain Districts in Nepal. *Zoonoses and Public Health*, 59(6), 393–400.

- Thammapalo, S., Wonghiranrat, W., Moonmek, S., & Sriplong, W. (2009). Biting time of *Aedes albopictus* in the rubber plantations and the orchards, the southern-most of Thailand. *Journal of Vector Borne Diseases*, 6(2), 1–6.
- Thongcharoen P. (1985). Japanese encephalitis in Thailand. *Journal of the Medical Association of Thailand*, 68(10), 534-545.
- Tiawsirisup, S., Punpee, A., & Nuchprayoon, S. (2012). Mosquito distribution and Japanese encephalitis virus infection in a bat cave and its surrounding area in Lopburi province, central, Thailand. *The Thai Journal of Veterinary Medicine*, 42(1): 43-49.
- Ting, S. H., Tan, H. C., Wong, W. K., Ng, M. L., Chan, S. H., & Ooi, E. E. (2004). Seroepidemiology of neutralizing antibodies to Japanese encephalitis virus in Singapore: continued transmission despite abolishment of pig farming?. *Acta Tropica*, 92(3), 187–191.
- Towbin, H., Staehelin, T., & Gordon, J. (1979). Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proceedings of the National Academy of Sciences of the United States of America*, 76, 4350-4354.
- Truong, Q. L., Seo, T. W., Yoon, B.I., Kim, H.C., Han, J. H., & Hahn, T.W. (2013). Prevalence of swine viral and bacterial pathogens in rodents and stray cats captured around pig farms in Korea. *The Journal of Veterinary Medical Science the Japanese Society of Veterinary Science*, 75(12), 1647–1650.
- Tsuchie, H., Oda, K., Vythilingam, I., Thayan, R., Vijayamalar, B., Sinniah, M., & et al. (1997). Genotypes of Japanese encephalitis virus isolated in three states in Malaysia. *American Journal of Tropical Medicine and Hygiene* 56, 153–158.
- Umenai, T., Krzysko, R., Bektimirov, T.A., & Assaad, F.A. (1985). Japanese encephalitis: current worldwide status. *Bulletin of the World Health Organization*, 63, 625–631.
- Van den Hurk, A.F., Ritchie, S.A., & Mackenzie, J.S. (2009). Ecology and geographical expansion of Japanese encephalitis virus. *Annual Review of Entomology*, 54, 17–35.
- Van Peenen, P.F.D., Irsiana, R., Saroso, J.S., Joseph, S.W., Shope, R.E., & Joseph, P.L. (1974). First isolation of Japanese encephalitis virus from Java. *Military Medicine-International Journal of AMSUS* 139, 821–823.
- Vasudevan, S. (2013). Climate change and Infectious Diseases. *Kerala Medical Journal*, VI (2), 28-31.
- Vaughn, D.W., & Hoke, C. H. (1992). The Epidemiology of Japanese Encephalitis: Prospects for Prevention. *Epidemiologic Reviews*, 14, 197-221.

- Vijayarani, H., & Gajanana, A. (2000). Low rate of Japanese encephalitis infection in rural children in Thanjavur district (Tamil Nadu), an area with extensive paddy cultivation. *Indian Journal of Medical Research*, 111, 212-214.
- Vythilingam, I., Tan, S.B. & Krishnasamy, M. (2002) Susceptibility of *Culex sitiens* to Japanese encephalitis virus in peninsular Malaysia. *Tropical Medicine & International Health*, 7, 539-540.
- Wacharapluesadee, S., Samseeneam, P., Phernpool, M., Kaewpom, T., Rodpan, A., Maneeorn, P., & et al. (2016). Molecular characterization of Nipah virus from *Pteropus hypomelanus* in Southern Thailand. *Virology Journal*, 13(1), 53.
- Wada, Y., & Watanabe, M. (1971) Ecological studies of *Culex tritaeniorhynchus summorosus* (Diptera: Culicidae): IV. Comparative studies on the behavior of *C. tritaeniorhynchus summorosus* and *C. pseudovishnui*. *Medical Entomology and Zoology*, 22, 24-30.
- Wang, H., & Liang, G. (2015). Epidemiology of Japanese encephalitis: Past, present, and future prospects. *Therapeutics and Clinical Risk Management*, 11, 435-448.
- Wang, J., Pan, X., Zhang, H., Fu, S., & Wang, H. (2009). Japanese Encephalitis viruses from bats in Yunnan. China. *Emerging Infectious Diseases journal*, 15, 939-942.
- Wang, L.H., Fu, S.H., Wang, H.Y., Liang, X.F., Cheng, J.X., & Jing, H.M. (2007). Japanese encephalitis outbreak, Yuncheng, China, 2006. *Emerging Infectious Diseases journal*, 13, 1123-1125.
- WHO- Japanese encephalitis. (2015). Retrieved January 15, 2016 from World Health Organization Media Centre- Japanese encephalitis <http://www.who.int/mediacentre/factsheets/fs386/en/>
- Winkler, G., Heinz, F. X., & Kunz, C. (1987). Studies on the glycosylation of flavivirus E proteins and the role of carbohydrate in antigenic structure. *Virology* 159(2), 237-243.
- Wisseman, C.L., Kitaoka, M., & Tamiya, T. (1966). Immunological studies with group B arthropod-borne viruses. V. Evaluation of cross-immunity against type 1 dengue fever in human subjects convalescent from subclinical natural Japanese encephalitis virus infection and vaccinated with 17D strain yellow fever vaccine. *The American Journal of Tropical Medicine and Hygiene*, 15, 588-600.
- Wu, X., Lin, H., Chen, S., Xiao, L., Yang, M., An, W., & et al. (2017). Development and application of a reverse transcriptase droplet digital PCR (RT-ddPCR) for sensitive and rapid detection of Japanese encephalitis virus. *Journal of Virological Methods*, 248, 166-171.

- Yamada T, Rojanasuphot S, Takagi M, Wungkobkiat S, Hirota T, Yamashita T, & et al. (1971). Studies on an epidemic of Japanese encephalitis in the northern region of Thailand in 1969 and 1970. *Biken Journal*, 14(3), 267-297.
- Yamamoto, S., Tsubahara, H., Yoshida, T., & Harada, K. (1949). First observation on case of Japanese B encephalitis of cattle. *National Institute of Animal Health*, 22, 197–203.
- Yang, D. K., Kim, B. H., Kweon, C. H., Kwon, J. H., Lim, S. I., & Han, H. R. (2004). Biophysical characterization of Japanese encephalitis virus (KV1899) isolated from pigs in Korea. *Journal of Veterinary Science (Suwon-Si, Korea)*, 5(2), 125–130.
- Yin, Z., & Liu, J.H. (1997). *Animal Virology. 2nd. Marrickville*. Australia: Science Press.
- Yu, I.M., Zhang, W., Holdaway, H.A., Li, L., Kostyuchenko, V.A., Chipman, P.R., & et al. (2008). Structure of the immature dengue virus at low pH primes proteolytic maturation. *Science*, 319, 1834–1837.
- Zhang, Y., Jing, J., Li, X., Wang, J., Feng, X., Cao, R., & et al. (2015). Integration analysis of miRNA and mRNA expression profiles in swine testis cells infected with Japanese encephalitis virus. *Infection, Genetics and Evolution*, 32, 342–347.