



UNIVERSITI PUTRA MALAYSIA

***BIOLOGICAL CHARACTERIZATION OF WILD COBIA,
Rachycentron canadum (LINNAEUS, 1766) OFF DUNGUN COAST,
TERENGGANU, MALAYSIA FOR STOCK MANAGEMENT AND
AQUACULTURE***

BABATUNDE TAOFIK ADEMOLA

FP 2016 29



**BIOLOGICAL CHARACTERIZATION OF WILD COBIA,
Rachycentron canadum (LINNAEUS, 1766) OFF DUNGUN
COAST, TERENGGANU, MALAYSIA FOR STOCK
MANAGEMENT AND AQUACULTURE**

By

BABATUNDE TAOFIK ADEMOLA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

August 2016

All material contained in this thesis, including without limitation text, logos, icons, photographs, and all other artworks, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copywrite holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

I dedicate this work to my parents who have set a pace for me that take me to reach this extent in life, and to my family and friends



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree Doctor of Philosophy

**BIOLOGICAL CHARACTERIZATION OF WILD COBIA,
Rachycentron canadum (LINNAEUS, 1766) OFF DUNGUN
COAST, TERENGGANU, MALAYSIA FOR STOCK
MANAGEMENT AND AQUACULTURE**

By

BABATUNDE TAOFIK ADEMOLA

August 2016

Chairperson : Assoc. Prof. S. M. Nurul Amin, PhD
Faculty : Agriculture

Investigation of some aspects of biology of a recreationally and commercially important fish, cobia, *Rachycentron canadum* in Dungun coast, Malaysia was conducted between April 2014 and May 2015. A total of 249 specimens were opportunistically collected from the catches of trawl net, gill net and by hooks and line. The total length ranged from 39.20 to 143.00 cm (88.29 ± 1.28 cm), fork length ranged from 37.50 to 124.00 cm (78.69 ± 1.09 cm) while the wet weight ranged from 0.50 to 20.40 kg (5.51 ± 0.26 kg). There was no significant difference in all the morphometric characters between sexes except for head width (HW) ($p < 0.05$). Ten (10) out of the seventeen (17) truss distances in the body landmarks of cobia showed significant differences between males and females. The equation of length-weight (LW) relationship for combined sex was $W = 0.000002FL^{3.3204}$ ($R^2 = 0.92$). The growth of cobia in Dungun water followed positive allometric as the growth coefficient (b) was higher than 3. Observed length frequency of both sex followed a normal distribution pattern and there was a predominance of cobia in the medium size range from 65 - 85 cm fork length for the females and 55 - 90 cm for the males. The sex ratio of females to males was 1:1.18 and it was not significantly different from the expected 1:1 ($X^2 = 2.12$ df = 1; $p < 0.05$). Growth parameters of von Bertalanffy growth formula for cobia estimated were $L_{\infty} = 142.00$ cm, $K = 1.20$ yr⁻¹, while natural mortality rates (M), fishing mortality rate (F), total mortality (Z) and Exploitation rate (E) estimated were 1.34, 0.95, 2.29 and 0.41 respectively. The maximum allowable limit of exploitation (E_{max}) was estimated to be 0.56. The sizes attained by cobia at the end of age 1 (year) was 99.28 cm. The estimation of fork length at first maturity (L_{mat}) from the plot of gonadosomatic index (GSI) against fork length (FL) indicated early maturation of males compared to females with respective value of 70 cm and 72 cm. Males with mature gonads were encountered throughout the year, while the proportion of mature females was higher in March and November, and this may be taken as the peak of the spawning period, even though

females with hydrated oocyte were also obtained in April, May, June, July and September. The mean batch fecundity (BF) for all matured females with hydrated oocytes was ranged from 54886 to 4316648, ($11,37,317 \pm 183712$) while average eggs per gram of ovary were ranged from 2108 to 5413 (3427.032 ± 128.73). Weak positive correlation ($r^2 = 0.48$) was found between the BF and female fork length with equation $BF = 38732.03FL - 2200000$. Similarly, BF was significantly correlated with ovary-free body weight ($r^2 = 0.56$). Histological data support continuous spawning characteristics of cobia as oocyte size frequency distribution showed progressive batch development. Increase in GSI was observed from September to November and it reached the peak in March which corresponds to the period when the highest proportion of matured females was caught. Study on the stomach content of the cobia, *R. canadum* in Dungun coast, showed that the bony fish was the dominant group in the diet with crustacean and mollusc contributing less. The percentage index of relative important (% IRI) values of fish, crustacean and mollusc were 72.27%, 2.98% and 24.75% respectively. Out of the 231 cobia specimens examined in this study, 111 were found to contain food in their stomach, out of which 98 had at least one identifiable prey items. Gut content analysis revealed that fish was consumed by cobia all the year round with *Hilsa* sp., *Alepes* sp. and *Carangides* sp. being the frequently encountered species. Higher stomach fullness index (SFI) was recorded in the month of April, June and November with values of 2.88, 2.06 and 2.42 respectively while lower SFI was in the month of May and August with values of 0.67 and 0.86 respectively. The percentage of empty coefficient (EC) was higher in October, August and July with values of 80%, 70% and 69% respectively, intermediate in the remaining month as values were about 50%. This stomach fullness index SFI and EC suggest moderate feeding intensity of cobia in the waters of Dungun. The proximate and chemical composition of cobia species from Dungun and its seasonal changes was examined. The result showed that lipids content was significantly ($p < 0.05$) higher in females (6.38 ± 0.19) than males (5.44 ± 0.11). No significant ($p < 0.05$) difference was found in the moisture, ash, total cholesterol and malondialdehyde (MDA) based on season, sex and feeding regime. In the fatty acid (FA) composition of the muscle tissue, the unsaturated class was dominant followed by saturated FA followed by the monoenes. Similar trend was found in the liver. Total n-3 poly-unsaturated fatty acid (PUFA) was higher compared to the total PUFA (n-6) in the muscle, and the total PUFA (n-6) was significantly ($p < 0.05$) higher in the muscle during the monsoon. Other FA classes that showed significant ($p < 0.05$) seasonal differences in the muscle were pentadecanoic acid, palmitoleic acid and linoleic acid. However, in the liver, the total saturated, total monones, total PUFA (n-6) and total PUFA (n-3) showed significant ($p < 0.05$) seasonal variation. The FA composition of males and females cobia did not show significant ($p < 0.05$) differences. During starvation in cobia, a significant reduction in total PUFA n-6, pentadecanoic acid, and palmitoleic acid was observed in the muscle. The result of this work suggests adult cobia in Malaysia preferred bony fishes as food while its reproductive period was continuous and cobia lipid profiles contained appreciable quantity of PUFA such as docosahexaenoic, eicosapentaenoic and arachidonic acids and it can be promoted for human consumption.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

PENCIRIAN BIOLOGI IKAN HARUAN TASEK LIAR, *Rachycentron canadum* (LINNAEUS, 1766) DI PANTAI DUNGUN, TERENGGANU, MALAYSIA UNTUK PENGURUSAN STOK DAN AKUAKULTUR

Oleh

BABATUNDE TAOFIK ADEMOLA

Ogos 2016

Pengerusi : Prof. Madya S. M. Nurul Amin, PhD

Fakulti : Pertanian

Kajian aspek-aspek biologi bagi ikan yang penting secara komersial dan rekreasi, *Rachycentron canadum* di Dungun, Malaysia telah dijalankan di antara April 2014 dan Mei 2015. Sejumlah 249 spesimen telah dikumpul secara rawak dari tangkapan pukut tunda, pukut hanyut dan tangkapan kail dan pancing. Jumlah panjang adalah dari 39.20 hingga 143.00 cm (88.29 ± 1.28 cm, min $\pm$ SE), panjang badan adalah dari 37.50 hingga 124.00 cm (78.69 ± 1.09 cm, min $\pm$ SE) manakala berat basah adalah dari 0.50 hingga 20.40 kg (5.51 ± 0.26 kg, min $\pm$ SE). Tiada perbezaan ketara pada semua karekter-karekter morfometrik di antara jantina kecuali pada lebar kepala (HW) ($p < 0.05$). Sepuluh (10) daripada tujuh belas (17) jarak truss pada bahagian badan haruan tasik telah menunjukkan perbezaan ketara di antara jantan dan betina. Persamaan panjang-berat (LW) bagi kedua-dua jantina adalah $W = 0.000002FL^{3.3204}$, $R^2 = 0.92$ manakala “b” pada eksponen ini menunjukkan pertumbuhan haruan tasik di perairan Dungun bersifat alometrik positif. Kekerapan panjang bagi kedua-dua jantina telah diperhatikan diikuti dengan corak taburan normal dan terdapat banyak haruan tasik pada saiz sederhana di antara 65 - 85 cm panjang badan bagi betina dan 55 - 90 cm bagi jantan. Nisbah jantina bagi betina dengan jantan adalah 1:1.18 dan ini tidak menunjukkan perbezaan ketara dengan nilai jangkaan 1:1 ($X^2 = 2.12$ df = 1; $p = 0.145$). Parameter pertumbuhan formula pertumbuhan von Bertalanffy untuk Ikan Aruan Tasek dianggarkan sebagai $L_{\infty} = 142.00$ cm, $K = 1.20 \text{ thn}^{-1}$, manakala kadar semula jadi kematian (M), kadar memancing kematian (F), jumlah kematian (Z) dan kadar eksploitasi (E) dianggarkan adalah 1.34, 0.95, 2.29 dan 0.41 masing-masing. Had maksimum yang dibenarkan eksploitasi ($E_{\max}$) dianggarkan 0.56. Saiz yang dicapai oleh Ikan Aruan Tasek di akhir zaman 1 (tahun) adalah 99,28 cm. Anggaran panjang badan pada kematangan pertama (L_{mat}) dari plot indeks gonadosomatik (GSI) berlawanan panjang badan (FL) menganggarkan kematangan awal pada jantan berbanding betina dengan nilai sebanyak 70 cm dan 72 cm. Ikan jantan dengan gonad yang matang telah ditemui sepanjang tahun, manakala nisbah ikan betina yang matang adalah lebih tinggi pada Mac dan November dan ini boleh

diambil kira kerana puncak tempoh pembiakan, walaupun ikan betina dengan oosit terhidrat juga telah didapati pada April, Mei, Jun, Julai dan September. Min kumpulan fekunditi (BF) untuk semua ikan betina yang telah matang dengan oosit yang terhidrat adalah dari 54886 hingga 4316648, ($11,37,317 \pm 183712$, min $\pm$ SE) manakala purata telur per gram dari ovary adalah dari 2108 hingga 5413 (3427.032 ± 128.73 , min $\pm$ SE). Korelasi positif yang lemah ($r^2 = 0.48$) telah ditemui di antara kumpulan fekunditi (BF) dan panjang badan betina dengan persamaan $BF = 38732.03FL - 2200000$. Secara kebetulan, BF adalah berkorelasi ketara dengan ovary-berat badan bebas ($r^2 = 0.56$). Data histologi menyokong pembiakan berterusan haruan tasik kerana kekerapan taburan saiz oosit telah menunjukkan perkembangan kumpulan yang menggalakkan. Pertambahan GSI telah diperhatikan dari September hingga November dan ia mencapai puncak pada Mac yang berkaitan dengan tempoh semasa nisbah tertinggi ikan betina yang telah matang ditangkap. Kajian berkenaan kandungan perut haruan tasik, *Rachycentron canadum* di pantai Dungun, Malaysia telah menunjukkan bahawa ikan bertulang adalah kumpulan dominan di dalam diet dengan krustasia dan molusk kurang menyumbang dan peratus indeks kepentingan relatif mereka (% IRI) bernilai 72.27, 2.98 dan 24.75 secara berturutan. Daripada 231 spesimen haruan tasik yang telah diperiksa di dalam kajian ini, 111 telah didapati mempunyai makanan di dalam perut mereka, dengan 98 daripada mereka sekurang-kurangnya mempunyai satu barangan mangsa yang boleh dikenal pasti. Analisis kandungan perut telah mendedahkan bahawa ikan yang telah dimakan oleh haruan tasik pada sepanjang tahun dengan *Hilsa* spp, *Alepes* spp dan *Carangids* spp paling banyak dijumpai. Indeks kepenuhan perut (SFI) yang lebih tinggi telah direkodkan pada bulan April, Jun dan November dengan nilai-nilai 2.88, 2.06 dan 2.42 secara berturutan manakala SFI yang lebih rendah adalah pada bulan Mei dan Ogos dengan nilai-nilai 0.67 dan 0.86 secara berturutan. Peratusan koefisien kosong (EC) adalah lebih tinggi pada Oktober, Ogos dan Julai dengan nilai-nilai 80%, 70% dan 69% secara berturutan, berkedudukan tengah pada bulan yang selebihnya dengan nilai kira-kira 50%. Indeks kepenuhan perut SFI dan EC mencadangkan kekerapan makan yang sederhana bagi haruan tasik di perairan Dungun. Komposisi proksimat and kimia bagi spesies haruan tasik dari Dungun, Malaysia dan perubahan-perubahan musimnya telah diperiksa. Keputusan telah menunjukkan kandungan lipid adalah ketara ($p < 0.05$) lebih tinggi pada ikan betina (6.38 ± 0.19) daripada ikan jantan (5.44 ± 0.11). Tiada perbezaan ketara ($p < 0.05$) telah ditemui pada kelembapan, abu, jumlah kolestrol dan malondialdehyde (MDA) berdasarkan musim, jantina dan regim pemakanan. Dalam komposisi asid lemak (FA) bagi tisu otot, kelas tidak tepu adalah dominan diikuti dengan FA tepu dan kemudian monoenes. Perkara sama juga telah ditemui pada hati. Jumlah n-3 asid lemak poli – tak tepu (PUFA) adalah lebih tinggi berbanding dengan jumlah PUFA (n-6) di dalam otot, dan jumlah PUFA (n-6) adalah lebih tinggi ($p < 0.05$) di dalam otot pada musim hujan. Kelas-kelas FA lain telah menunjukkan perbezaan musim yang ketara ($p < 0.05$) pada otot adalah asid pentadecanoic, asid palmitoleic dan asid linoleic. Walau bagaimanapun, di dalam hati, jumlah tepu, jumlah monones, jumlah PUFA (n-6) dan jumlah PUFA (n-3) telah menunjukkan variasi musim yang ketara ($p < 0.05$). Walau Bagaimana pun, komposisi FA bagi ikan jantan dan ikan betina tidak menunjukkan perbezaan yang ketara ($p < 0.05$). Ketika haruan tasik berada dalam keadaan kelaparan, pengurangan ketara dalam jumlah PUFA n-6, asid pentadecanoic dan asid palmitoleic telah dijumpai pada otot. Keputusan kajian ini mencadangkan haruan tasik dewasa di Malaysia memilih ikan bertulang sebagai makanan manakala tempoh pembiakan adalah berterusan dan lipid haruan tasik mengandungi kuantiti PUFA yang penting

seperti asid docosahexaenoic, eicosapentaenoic dan arachidonic dan ia boleh digunakan untuk makanan manusia.



ACKNOWLEDGEMENTS

I would begin by thanking the chairman of my supervisory committee, Associate Prof. Dr. S. M. Nurul Amin, who offer me an opportunity to work with him at a crucial time, and offered me sufficient guidance to accomplish this research. I also thank the members of supervisory committee; Associate Prof. Dr. Yuzine Esa, Prof. Aziz Arshad and Prof. Fatimah Md. Yusoff for their dedication, support, encouragement and understanding. I also acknowledge the support of Dr. Nicholas Paul Romano and Dr. Mahdi Ebraheemi. I wish to express my gratitude to Mr Azim Md Katib and Mr Hazmadi Zakaria for their assistance during fish sampling.

This research wouldn't have been possible without the support of the management of Umaru Musa Yaradua University, Nigeria and funding provided by the Research Management Centre of Universiti Putra Malaysia, (grant number GP-IBT/2013/9410700).

Finally, may Allah be praised, through His mercy all good is accomplished.

This thesis was submitted to the senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirements for the degree of Doctor of Philosophy. The members of Supervisory committee were as follow:

S. M. Nurul Amin, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Yuzine Esa, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Aziz Arshad, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Fatimah Md. Yusoff

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by Members of Supervisory Committee

This is to confirm that:

- The research conducted and the writing of the thesis was under our supervision;
- Supervision responsibilities as stated in the UPM (Graduate Studies) Rule 2003 (Revised 2012-2013) were adhere to.

Signature: _____

Name of Chairman of Supervisory
committee

Associate Prof. Dr. S. M. Nurul Amin

Signature: _____

Name of member of Supervisory
committee

Associate Prof. Dr. Yuzine Esa

Signature: _____

Name of member of Supervisory
committee

Prof. Dr. Aziz Arshad

Signature: _____

Name of member of Supervisory
committee

Prof. Dr. Fatimah Md. Yusoff

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	vi
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xx
 CHAPTER	
 1 INTRODUCTION	
1.1 Background of the study	1
1.2 Statement of problem	2
1.3 Research hypothesis	3
1.4 Objectives of the research	3
1.5 Organization of the study	3
 2 LITERATURE REVIEW	
2.1 Taxonomy of cobia	5
2.2 Morphological characteristics and species description	6
2.3 Distribution of <i>Rachycentrum canadum</i>	7
2.4 Biological characteristics of cobia	9
2.4.1 Sex ratio	9
2.4.2 Size at maturity	9
2.4.3 Fecundity and spawning frequency	10
2.4.4 Spawning period	10
2.4.5 Length-weight relationship	12
2.5 Ecological aspects of cobia	13
2.5.1 Habitat	13
2.5.2 Migration	13
2.5.3 Food and feeding habits	13
2.6 Genetics and stock assessment	15
2.6.1 Cobia population genetics	15
2.6.2 Stock assessment	16
2.7 Past and present research on cobia	18
2.7.1 Seed production	18
2.7.2 Rearing	20
2.8 Prospects for cobia aquaculture	21
 3 GENERAL METHODOLOGY	
3.1 Study area	22
3.1.1 Description of study area	22
3.1.2 Fisheries in Dungun coast	23

3.2	Status of cobia in Terengganu	24
3.3	Sample collection	24
4	MORPHOMETRIC CHARACTERISTICS OF WILD COBIA, <i>Rachycentrum canadum</i> POPULATION OFF DUNGUN COAST, TERENGGANU MALAYSIA USING TRADITIONAL AND TRUSS MORPHOMETRICS	
4.1	Introduction	25
4.2	Materials and methods	26
4.2.1	Sample collection and morphometric measurement	26
4.2.2	Truss morphometric measurement	28
4.2.3	Data analysis	29
4.3	Results	29
4.3.1	Traditional morphometric characteristics	29
4.3.2	Truss morphometric characteristics	32
4.4	Discussion	33
4.5	Conclusions	34
5	POPULATION DYNAMICS AND STOCK ASSESSMENT OF <i>R. canadum</i> IN THE COASTAL WATER OF DUNGUN	
5.1	Introduction	35
5.2	Materials and methods	36
5.3	Results	37
5.3.1	Size frequency distribution	37
5.3.2	Length-weight relationship	42
5.3.3	Growth parameters	43
5.3.4	Mortality and exploitation	45
5.3.5	Recruitment pattern	45
5.3.6	Relative yield per recruit and biomass per recruit	46
5.3.7	Age and growth	47
5.3.8	Stock size and maximum sustainable yield (MSY)	47
5.4	Discussion	48
5.5	Conclusions	50
6	REPRODUCTIVE BIOLOGY OF <i>R. canadum</i> COLLECTED FROM DUNGUN COAST	
6.1	Introduction	51
6.2	Materials and methods	51
6.2.1	Sampling	51
6.2.2	Estimation of condition indices	52
6.2.3	Sex ratio and length at maturity	52
6.2.4	Histological procedure	53
6.2.5	Estimation of batch fecundity and spawning frequency	53
6.3	Results	54
6.3.1	Spawning cycle	54
6.3.2	Length at maturity	56
6.3.3	Sex ratio	57

6.3.4	Batch fecundity and spawning frequency	59
6.3.5	Condition indices	60
6.3.6	Gonad histology of <i>R. Canadum</i>	63
6.4	Discussion	70
6.5	Conclusions	72
7	FOOD AND FEEDING HABIT OF <i>R. canadum</i> IN DUNGUN COAST, MALAYSIA	
7.1	Introduction	73
7.2	Materials and methods	74
7.2.1	Fish sampling	74
7.2.2	Stomach content analysis	74
7.3	Results	76
7.3.1	Feeding intensity of <i>R. Canadum</i>	76
7.3.2	Stomach content analysis	78
7.3.3	Important of relative index (IRI) of prey items	83
7.3.4	Seasonal variation in food items	84
7.4	Discussion	86
7.5	Conclusions	87
8	NUTRITIONAL COMPOSITION OF WILD COBIA, <i>R. canadum</i>, OFF DUNGUN COAST, MALAYSIA	
8.1	Introduction	88
8.2	Materials and methods	89
8.2.1	Sampling	89
8.2.2	Analysis of chemical composition	90
8.2.3	Analysis of fatty acid	90
8.2.4	Analysis of total cholesterol	91
8.2.5	Analysis of malondialdehyde	91
8.2.6	Statistical analysis	92
8.3	Results	92
8.3.1	Chemical composition	92
8.3.2	Fatty acid composition of liver and muscle	93
8.4	Discussion	101
8.5	Conclusions	103
9	SUMMARY, CONCLUSIONS AND RECOMMENDATION FOR FUTURE RESEARCH	
9.1	Summary	104
9.2	Conclusions	107
9.3	Recommendations for future research	108
	REFERENCES	110
	BIODATA OF STUDENT	125
	LIST OF PUBLICATIONS	126

LIST O TABLES

Table	Page
2.1 Spawning period of <i>Rachycentron canadum</i> in some regions in the world	11
2.2 Length weight relationship parameters of <i>R. canadum</i> from different countries	12
2.3 Stock assessment of cobia and some population's parameters	17
2.4 Hatching rates of <i>R. canadum</i> under natural and hormonal induced spawning	19
3.1 Landings of cobia, <i>R. canadum</i> in Malaysia (Tonnes)	24
4.1 Morphomertic characteristics of <i>R. canadum</i> used in the study and their descriptions	27
4.2 Value of morphometric characteristics (mean $\pm$ SE) of <i>R. canadum</i> in Dungun coast, Malaysia	30
4.3 Percentage of each morphomertic characteristics to total length (TL) of <i>R. canadum</i>	30
4.4 Percentage of each morphometric characteristics to head length (HL) of <i>R. canadum</i>	31
4.5 Regression equations of total length (TL) against each morphometric characteristics of <i>R. canadum</i>	31
4.6 Summary of truss morphometric characters of <i>R. canadum</i>	32
5.1 Grouped length-frequency data for cobia collected from Dungun coast	36
5.2 Length-weight relationship of <i>R. canadum</i> in Dungun coast	43
5.3 Comparison of length-weight relationship of <i>R. canadum</i>	48
6.1 Reproductive stages of <i>R. canadum</i> by month	54
6.2 Number of <i>R. canadum</i> collected by months and variations in their sex ratio	58
6.3 Sex ratio of cobia in different size classes	58

6.4	Condition indices of <i>R. canadum</i> by months	60
7.1	Description of stomach fullness classes used in this study	74
7.2	Stomach fullness of <i>R. canadum</i> in different months	77
7.3	Monthly variations in numerical abundance of prey items in the gut content of <i>R. canadum</i>	79
7.4	Monthly variation in frequency of occurrence of prey items in the gut content of <i>R. canadum</i>	81
7.5	Monthly variation in volume (ml) of prey items consumed by <i>R. canadum</i>	82
7.6	Monthly variation in weight (g) of prey items consumed by <i>R. canadum</i>	82
7.7	Estimated index of relative important of different food items in the stomach of <i>R. canadum</i>	83
8.1	Chemical composition of <i>R. canadum</i>	93
8.2	Fatty acid profile (% of identified fatty acids) of the muscle and liver of <i>R. canadum</i> by season	94
8.3	Fatty acid profile (% of identified fatty acids) of muscle and liver of <i>R. canadum</i> with respect to feeding status	96
8.4	Fatty acid profile (% of identified fatty acids) of muscle and liver of <i>R. canadum</i> with respect to sex	98
8.5	Correlation between fatty acids classes and % lipid of <i>R. canadum</i>	99
8.6	Overall fatty acid content (% of identified fatty acids) in the muscle and liver of <i>R. canadum</i>	100

LIST OF FIGURES

Figure	Page
2.1 External morphology of juvenile (A) and adult cobia (B)	6
2.2 Areas within EEZs favorable for offshore grow-out of cobia (FAO, 2013a)	8
2.3 Capture statistics of cobia in the important fish nations	8
2.4 Early metamorphosis in <i>R. canadum</i> : A; fertilized egg 0 hours B; Hatching 21hours C; newly hatched larva 22hours (Sakthivel <i>et al.</i> , 2012)	20
3.1 Map of Peninsular Malaysia showing the sampling location	23
4.1 Morphometric characters of <i>Rachycentron canadum</i>	28
4.2 Annotated diagram of anatomical land marks for truss morphometric study of <i>R. canadum</i>	29
5.1 Size frequency distribution of female (A) and male (B) <i>R. canadum</i>	38
5.2 Size frequency distribution <i>R. canadum</i> (sex combined)	38
5.3 Monthly size frequency distribution of cobia	41
5.4 Relationship between wet weight and fork length of <i>R. canadum</i>	42
5.5 Logarithmic relationship between wet weight and fork length of <i>R. canadum</i>	42
5.6 Predicted maximum length for <i>R. canadum</i> based on the extreme value theory (Formacion <i>et. al.</i> , 1991). The predicted maximum length value and the 95% confidence intervals were obtained from the intersection of the overall maximum length from the y and x, z lines, respectively	43
5.7 K-scan routine for best value of von Bertalanffy growth function (VBGF), asymptotic length (L_{∞}) and growth coefficient (K) of <i>R. canadum</i> using ELEFAN-1	44

5.8	von Bertalanffy growth curves of <i>R. canadum</i> superimposed on the restructured length-frequency histograms. The black and white bars are positive and negative deviations from the “weighted” moving average of three length classes and they represent pseudo-cohorts	44
5.9	Length-converted catch curved of <i>R. kanagurta</i> , the darkened full dots represent the points used in calculating through least square linear regression and the open dots represent the point either not fully recruited or close to L_{∞}	45
5.10	Recruitment pattern of <i>R. canadum</i> in Dungun coast, Malaysia	46
5.11	Relative Y/R and B/R of <i>R. canadum</i> in Dungun coast, Malaysia using a knife-edge procedure	46
5.12	Plot of age and growth of cobia based on computed growth parameters	47
6.1	Proportion of mature female in total female <i>R. canadum</i>	55
6.2	Relationship between fork length and gonad weight of <i>R. canadum</i>	55
6.3	Relationship between gonadosomatic index (GSI) and fork length of <i>R. canadum</i>	56
6.4	Logistic curve fitted to the proportion of spawning capable female (A) and male (B) cobia, <i>Rachycentron canadum</i> plotted against fork length (dash lines represent the estimated length at 50% maturity (L_{50}))	57
6.5	Relationship between fork length and fecundity of <i>R. canadum</i>	59
6.6	Relationship between batch fecundity and ovary-free body weight of <i>R. canadum</i>	60
6.7	Trends in gonadosomatic index (GSI) and stomach fullness index (SFI) of <i>R. canadum</i>	61
6.8	Relationship between GSI and stage of gonad development <i>R. canadum</i>	62
6.9	Trends in VSI of <i>R. canadum</i> (sex combined)	62
6.10	Trends in condition factor of female (A) and male (B) <i>R. canadum</i> (Error bars: 95% CI)	63

6.11	(A) Immature ovary in stage I (unfoliated stage) containing perinucleolar oocytes (a) and a large intra ovarian space (b) under $\times 10$ (B). Absence of large oocytes with cortical alveoli and degenerating oocyte suggest no previous spawning. (58.50cm FL, Bar = 0.1mm)	64
	(B) Immature ovary in stage II with developing, early yolked oocyte (a) dominating the lamellae under $\times 10$ objectives. Some large oocytes shows the presence of cortical alveoli (b) (101.00cm FL, Bar = 0.1mm)	64
	(C) Maturing ovary in stage III, containing vitellogenic oocytes (a) undergoing synchronous maturation, and cortical alveolar (b) (95.5cm FL, Bar = 0.1mm)	65
	(D) Maturing ovary (stage IV) containing hydrated oocytes (a) and large lipid globule (96.00cm FL, Bar = 0.1mm)	65
	(E) Matured ovary (stage V) containing fully hydrated oocytes (a), a large lipid globule (b), and visible nucleus (c) (102.4cm FL, Bar = 0.1mm)	66
	(F) Spent ovary (stage VI) containing few matured oocytes (a), maturing oocytes (b) and post ovulatory follicle (c) (109.00FL, Bar = 0.1mm)	66
6.12	Oocyte size distribution in <i>R. canadum</i> matured ovary	67
6.13	(A) Testes in the early maturation stage of development (stage I) with a number of intact and ruptured spermatogonia (SG) few spermatocytes (SC), spermatid (ST) (66.50cm FL, Bar 0.1mm)	68
	(B) Testes in the mid maturation stage of development (stage II) with a numerous spermatocysts (SC) and spermatid (ST) (Bar 0.1mm) (64.50cm FL, Bar 0.1mm)	68
	(C) Testes in the late maturation stage of development (stage III) with a few spermatocysts (SC) and abundant of spermatozoa (SZ) in the lumen (L) (65.00 cm FL, Bar 0.1mm)	69
	(D) Testes in the regressed stage of development (stage IV) with a few spermatocysts (SC), spermatid (ST) and reappearance of spermatogonia (SG) in the lumen (L) (82.50cm FL, Bar 0.1mm)	69
	(E) Testes in the regression stage of development (stage V) with a few spermatocysts (SC) and spermatid (ST) in the lumen (L). (84.00cm FL, Bar 0.1mm)	70

7.1	Cobia stomach (a) full stomach (b) empty stomach before dissection (c) empty stomach after dissection	75
7.2	Number of prey taxa in food items of <i>R. canadum</i> by months	80
7.3	Relative important index of the groups of food in the stomach content of consumed by <i>R. canadum</i>	84
7.4	Relationship between the fork length and the weight of prey consumed by <i>R. canadum</i>	84
7.5	Prey items recorded in the stomach of <i>R. canadum</i>	85
8.1	Standard curve for the estimation of total cholesterol in this study	91
8.2	Standard curve for the estimation of total malondialdehyde in this study	92
8.3	Percentage weight (g) of food classes in the stomach content of <i>R. canadum</i> by season	97

LIST OF ABBREVIATIONS

BF	Batch Fecundity
DoF	Department of Fisheries
dph	Days Post Hatch
E	Exploitation rate
EC	Empty Coefficient
EEZs	Exclusive Economic Zones
ELEFAN	Electronic Length Frequency Analysis
$E_{\max}$	Maximum allowable limit of exploitation
FAO	Food and Agricultural Organization
FCR	Feed Conversion Ratio
FiSAT	FAO ICLARM Stock Assessment Tools
GSI	Gonadosomatic Index
IRI	Index of Relative Importance
K	Fulton Condition Factor (K)
K	Growth co-efficient of VBGF
L_{∞}	Asymptotic length
L_{mat}	Length at Maturity
L_{max}	Predicted extreme length
M	Natural mortality
MSY	Maximum Sustainable Yield
SFI	Stomach Fullness Index
VSI	Viscerosomatic Index

CHAPTER 1

INTRODUCTION

1.1 Background of the study

Fish population is a renewable natural resource if exploited in a sustainable way. Hence, fishery represents an important source of revenue to the economy of many countries and an important food sector for humans (Dwivedi *et al.*, 2009). It is an important food resource for humans, accounting for about 15.7 % of the global animal protein consumption (FAO, 2010). Increasing trends of human activities, overfishing, fish habitat destruction, habitat fragmentation and pollution have been identified as threats to aquatic biodiversity. Fish population are been subjected to natural control processes that constantly modify the structure and abundance of their population in various life cycle stages in response to these factors (Milner *et al.*, 2003).

Cobia is the only species of the family Rachycentridae. They are migratory warm water species widely distributed in subtropical, tropical and temperate marine waters except in the eastern Pacific (Briggs, 1960). They migrate to deeper water in fall and winter (Shaffer and Nakamura, 1989). Cobia is a non-target species in capture fisheries as most cobia landings were from recreational fishing. This may be due to the fact that adult cobia is solitary. Cobia constitutes 0.1% of landing between 2007 and 2010 in Karnataka coast, India (Rohit and Bhat, 2012). Similarly, in Malaysia, cobia accounts for 0.1% the total landings in 2014 (DoF, 2014). Countries where cobia fishing is high exceeding 1000 metric tons per annum include Malaysia, Brazil, Iran, Philippines and Pakistan (FAO, 2009). Cobia is found associated with a wide range of habitat including rock, mud, sand and gravel bottom as well as buoying objects. This benthopelagic nature may have enhanced nonselective carnivorous feeding behavior reported for cobia, exploring prey from water column, benthic and open water (Meyer and Franks, 1996; Smith, 1995; Rohit and Bhat, 2012). They also establish commensal relationship with rays (Smith and Merriner, 1982). However, various feeding studies reveal geographic difference in the diet of cobia and their spawning grounds are areas of abundant prey (Richards, 1967; Lefebvre and Denson, 2012). Currently, there is no published information on the diet composition of cobia in Malaysian waters. Cobia exhibit extended batch spawning and peak spawning seasons vary with geographical locations. Their spawning behaviour in Malaysia has not been reported.

The first attempt to culture cobia using naturally spawned eggs collected from the wild was carried out by Hassler and Rainville (1975) and held in captivity for 131 days. The performance of cobia observed in the rearing trial in terms of hatching rate, growth rate, readiness to accept supplemented feed and hardiness indicated that cobia has potential for culture. Collection of wild larvae could not be utilized for commercial production due to limited and inconsistency in the availability; hence there is a need to adapt cobia to captive spawning. Biological characteristics of cobia

has made it reputed for mariculture, like high growth rate ($K = 2.6 \text{ (kg/yr}^{-1})$) in the wild (Ganga *et al.*, 2012), 6kg in 1 year in captivity (Chou *et al.*, 2001), and high fecundity up to 2.88×10^6 (van der Velde *et al.*, 2010). It also possesses a good flesh quality, diseases resistant and value of feed conversion ratio (FCR) which is a measure of the ratio between feed input and output of 1.05 has been reported for cobia in juvenile stage (Resley *et al.*, 2006) and FCR of 2.0 in adult (Benetti *et al.*, 2008a). Adaptability for captivity breeding and high market demand especially in Europe has been attributed to cobia (Gopakumar *et al.*, 2011; Benetti *et al.*, 2010). In addition to this, cobia has nutritional biochemical compounds in a balanced composition, essential amino acids, polyunsaturated fatty acids, fat content and microelements in a good proportion (Liu *et al.*, 2009). Therefore, consumption of cobia fish may be a good complementary in human diet. Production of cobia increased 7000 folds from 1995 to 2005, estimated global production in 2007 was 30,000 tonnes while in 2012 it was estimated to be 41000 tonnes valued US\$71 million (FAO, 2012). In Malaysia, aquaculture of cobia remains undeveloped.

As cobia gaining more acceptances for aquaculture, its potential remains unexploited in Malaysia. At present, there is lack of information on stock assessment, morphometric features, feeding ecology and reproductive biology of wild cobia of Malaysia waters. This information is necessary for the development of indigenous broodstocks and for management purposes.

1.2 Statement of problem

Although cobia is an important component in the landing statistic in Malaysia, there is lack of published information on basic biological aspects like morphometry, reproductive cycle, sex ratio, fecundity, food habits and nutritional values. Cobia culture is still underdeveloped in Malaysia. Its market value as a food fish has been identified in Europe, China, and Taiwan, where its aquaculture production is on the increase. The understanding of biology, ecology and nutritional quality of the Malaysian stock will be a prerequisite for the broodstock development and its aquaculture development. Till date, assessment of food habit, reproductive biology, population structure and nutritional chemical composition of cobia population in Malaysia has not been documented. Even though the capture of cobia was high in Malaysia, study of the population dynamic and stock assessment has not been carried out. This is necessary to avoid overfishing. Current research therefore focuses on some biological aspects and chemical composition of cobia in Peninsular Malaysia. This information is essential for sustainable management and proper utilization of wild cobia, and possibly recommending it for aquaculture in Malaysia.

1.3 Research hypothesis

The hypotheses of this research are:

- i) Morphology and condition indices of cobia is influenced by habitat water quality parameters and food preference, hence examination of morphometric parameter and condition factors would reflect habitat differences.
- ii) As the capture of cobia is high in Malaysia, there is possibility of overfishing and overexploitation.
- iii) Gonad development, spawning, and fecundity of cobia in Dungun coast, Malaysia are attuned with geographical location and environment. It is assumed that spawning season would be different in from other locations.
- iv) The food preference of cobia in Malaysia would be different from other locations due to differences in the population assemblage and species coexistence.
- v) *R. canadum* is a top carnivore by nature and it will have numbers of essential fatty acids that will make it a good choice of food fish.

1.4 Objectives of the research

The specific objectives of this research are:

- i) To document the morphometric characteristics, of Wild cobia population off Dungun coastal waters of Terengganu, Malaysia using traditional and truss morphometric methods.
- ii) To study the population dynamics and stock assessment of *R. canadum* in coaster water of Dungun.
- iii) To determine the reproductive cycle, spawning season, sex ratio and fecundity of *R. canadum* in Dungun coast and its correlation with its feeding pattern.
- iv) To investigate the food and feeding habits of cobia and seasonal changes in diet composition.
- v) To determine the nutritional properties of wild cobia in Dungun coast, Malaysia.

1.5 Organization of the study

This thesis is subdivided into seven chapters:

- i) Chapter one: This chapter gives an overview of cobia capture and its status in Malaysia, the problem statement, significance of the study and objectives.
- ii) Chapter two: This contains a review of literature on cobia fisheries and aquaculture, population biology, feeding habits, population parameter and

genetics. This chapter provides a background for formulation of hypothesis, understanding research problems and adopting appropriate methodologies to accomplish the study objectives of this research.

- iii) Chapter three: This chapter present a brief overview of general research methodology used to accomplish the objectives of this research including the statistical analysis.
- iv) Chapter four: In this chapter, a detail of the first objective of this research was presented. Information on the morphometric characteristics determined by traditional and truss analysis of *Rachycentron canadum* in Dungun, Malaysia were provided therein.
- v) Chapter five: Herein, information on the population dynamics and key stock parameters was given. Population structure and length–weight relationship of cobia was also reported.
- vi) Chapter six: This chapter was based on the third objective. Findings on the reproductive cycle, spawning season, sex ratio and fecundity of *Rachycentron canadum* in Peninsular Malaysia were presented.
- vii) Chapter seven: Investigation on the food habits of cobia, *Rachycentron canadum* and seasonal changes in feeding pattern as contained in the third objective.
- viii) Chapter eight: Nutrition composition of cobia and its suitability as human food supplement was reported.
- ix) Chapter nine: This was the final chapter and it represents the summary, conclusions and recommendations for future research.

REFERENCES

- Akyol, O. and Unal, V. 2013. Second record of the cobia, *Rachycentron canadum* (actinopterygii: Perciformes: rachycentridae), from the mediterranean sea. *Acta ichthyologica et piscatorial* 43(4): 315-317.
- Al-Barwani, S.M., Arshad, A., Amin, S.M.N., Japar, S.B., Siraj, S.S. and Yap, C.K. 2007. Population dynamics of the green mussel *Perna viridis* from the high spat-fall coastal water of Malacca, Peninsular Malaysia. *Fisheries Research* 84(2): 147-152.
- Aliabadi, S.M.A., Gilkolaei, S.R., Savari, A., Zolgharnein H., and Nabavi, S.M.B. 2008. Microsatellite Polymorphism in Iranian Populations of Cobia (*Rachycentron canadum*) *Biotechnology* 7: 775-780.
- Ambak, M.A. 2010. Fishes of Malaysia. UMT Press, Terengganu.
- Amin, S.M.N., Arshad, A., Ismail, N.H., Idris, M.H., Bujang, J.S., Sirai, S.S. 2010. Morphometric variation among the three species of genus *Acetes* (Decapoda: Sergestidae) in the coastal waters of Malacca, Peninsular Malaysia. *Pertanika Journal of Tropical Agricultural Science* 33: 341-347.
- Amin, S.M.N., Arshad, A., Haldar, G.C., Shohaimi, S. and Ara, R. 2005. Estimation of Size Frequency Distribution, Sex Ratio and Length-Weight Relationship of Hilsa (*Tenualosa ilisha*) in the Bangladesh Water. *Research Journal of Agriculture and Biological Sciences* 1(1): 61-66.
- Amin, S.M.N., Rahman, A., Haldar, G.C., Mazid, M.A. 2001. Studies on age and growth and exploitation level of *Tenualosa ilisha* in the coastal region of Chittagong. *Journal of Inland Fisheries Society of India* 33: 1-5.
- AOAC, 1990. Official Methods of Analysis of the Association of Official Analytical Chemists, In Herlick, K. 15th (eds.) Arlington, Va, USA.
- AOAC, 1995. Official Methods of Analysis of the Association of Official Analytical Chemists. In Patricia Cunniff 16th (eds.) Arlington, VA, USA.
- Ara, R., Amin, S.M.N., Mazlan, A.G. and Arshad, A. 2013. Morphometric variation among six families of larval fishes in the seagrass-mangrove ecosystem of Gelang Patah, Johor, Malaysia. *Asian Journal of Animal and Veterinary Advances* 8(2): 247-256.
- Arnold C.R., Kaiser J.B. and Holt G.J. 2002. Spawning of cobia (*R. canadum*) in captivity. *Journal of the World Aquaculture Society* 33: 205-208.
- Arendt, M.D., Olney, J.E. and Lucy, J.A. 2001. Stomach content analysis of cobia, *R. canadum* from lower Chesapeake Bay. *Fisheries Bulletin* 99(4): 665-670.

- Atwood, H.L., Young, S.P., Tomasso, J.R. and Smith, T.I.J. 2004. Resistance of cobia, *Rachycentron canadum*, juveniles to low salinity, low temperature, and high environmental nitrite concentrations. *Journal of Applied Aquaculture* 15:191–195.
- Ayala, A., Muñoz, M.F. and Argüelles, S. 2014. Lipid Peroxidation: Production, Metabolism, and Signaling Mechanisms of Malondialdehyde and 4-Hydroxy-2-Nonenal. *Oxidative Medicine and Cellular Longevity* doi: 10.1155/2014/360438.
- Bakhoun, S.A. and Faltas, S.N. 2003. Food and feeding habits of Bayad fish *Bagrus Bajad* (Forsskal, 1775) in El-Nozha Hydrodrome, Alexandria, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries* 7(3): 197-211.
- Bayir, A., Halilog˘ lu, H.I., Sirkeciog˘ lu, A.N. and Aras, N.M. 2006. Fatty acid composition in some selected marine fish species living in Turkish waters. *Journal of Science of Food and Agriculture* 86: 163–168.
- Benetti, D.D., Hanlon, B.O., Rivera, J.A., Welch, A.W., Maxe, Y.C. and Orhun, M.R. 2010. Growth rates of cobia (*Rachycentron canadum*) cultured in open ocean submerged cages in the Caribbean. *Aquaculture* 302: 195-201.
- Benetti, D.D., Orhun, M.R., Sardenberg, B., Hanlon, B.O., Welch, A., Hoenig, R., Zink, I., Rivera, J.A., Denlinger, B., Bacoat, D., Palmer, K. and Cavalin, F. 2008a. Advances in hatchery and grow-out technology of cobia *R. canadum* (L). *Aquaculture Recourses* 39: 701-711.
- Benetti, D.D., Sardenberg, B., Welch, A., Hoenig, R., Orhun, M.R. and Zink, I. 2008b. Intensive larval husbandry and fingerling production of cobia *R. canadum*. *Aquaculture* 281: 22-27.
- Blaber S.J.M., Brewer D.T., Milton D.A, Merta G.S., Efizon D, Fry G, van der Velde T.D 1999. The life history of the protandrous tropical shad *Tenualosa macrura* (Alosinae: Clupeidae): fishery implications. *Estuarine, Coastal and Shelf Science* 49: 689–701.
- Blueweiss, L., Fox, H., Kudzman, V., Nakashima, D., Peters, R. and Sams, S. 1978. Relationships between body size and some life history parameters. *Oecologia* 37: 257-272.
- Booke, H.E. 1981. The conundrum of the stock concept – Are nature and nurture definable in fishery science? *Canadian Journal of Fisheries and Aquatic Sciences* 38: 1479-1480.
- Braga, R.R., Bornatowski, H., Vitule J.R.S. 2012. Feeding ecology of fishes: an overview of worldwide publications. *Reviews in Fish Biology and Fisheries* 22: 915–929.

- Briggs, J.C. 1960. Fishes of worldwide (circumtropical) distribution. *Copeia* 3: 171-180.
- Brown-Peterson, N.J., Overstreet, R.M., Lotz, J.M., Franks, J.S. and Burns, K.M. 2001. Reproductive biology of cobia, *Rachycentron canadum*, from coastal waters of the southern United States. *Fisheries Bulletin* 99: 15-28.
- Brzeski, V.J. and Doyle, R.W. 1988. A morphometric criterion for sex discrimination in Tilapia. In *Proceeding of the Second International Symposium on Tilapia in Aquaculture* 15: 439-444.
- Budge, S.M., Iverson, S.J., Koopman, H.N. 2006. Studying trophic ecology in marine ecosystems using fatty acids: a primer on analysis and interpretation. *Marine Mammal Science* 22: 759-801.
- Cadrin, S.X. and Silva, V.M. 2005. Morphometric variation of Yellowtail Flounder. *ICES Journal of Marine Science* 62: 683-694.
- Cadrin, S.X. and K.D. Friedland. 1999. The utility of image processing techniques for morphometric analysis and stock identification. *Fisheries Research* 43: 129-139.
- Caylor, R.E., Biesiot, P.M. and Franks, J.S. 1994. Culture of cobia (*Rachycentron canadum*): cryopreservation of sperm and induced spawning. *Aquaculture* 125: 81-92.
- Chou, R.L., Her, B.Y., Su, M.S., Hwang, G., Wu, Y.H. and Chen, H.Y. 2004. Substituting fishmeal with soybean meal in diets of juvenile cobia *R. canadum*. *Aquaculture* 229: 325-333.
- Chou, R.L., Su, M.S. and Chen, H.Y. 2001. Optimal dietary protein and lipid levels for juvenile cobia *R. canadum*. *Aquaculture* 193: 81-89.
- Chu, K. B, Abdullah, A., Abdullah, S. Z. and Abu Bakar, R. 2013. A case study on the mortality of cobia (*Rachycentron canadum*) cultured in traditional cages. *Tropical Life Sciences Research* 24(2): 77-84.
- Colle, D.E. 1976. The food habits of three centrarchids, *Lepomis macrochirus* (Rafinesque), *Micropterus salmoides* (Lacepede), and *Lepomis gulosus* (Cuvier) in a central Texas farm pond. M.S. thesis, Texas A&M University, College Station.
- Cortes, E., Manire, C.A. and Hueter, R.E. 1996. Diet, feeding habits, and diel feeding chronology of the bonnethead shark, *Sphyrna tiburo*, in southwest Florida. *Bulletin of Marine Science* 58: 353-367.
- Craig, S.R. 2012. Soy use optimization in feeds for cobia. *American Soybean Association-International Marketing*. AQ50.

- Cren L.E.D. 1951. The length–weight relationship and seasonal cycle in gonad weight and conditions in the perch *Perca fluviatilis*. *Journal of Animal Ecology* 20: 201–19.
- Darden, T.L., Walker, J.M., Brenkert, K., Justin R. Yost, J.R., and Denson, M.R. 2014. Population genetics of Cobia (*Rachycentron canadum*): implications for fishery management along the coast of the southeastern United States. *Fisheries Bulletin* 112: 24–35.
- Darracott, A.1977. Availability, morphometrics, feeding, and breeding activity of a multi-species, demersal fish stock of the western Indian Ocean. *Journal of Fisheries Biology* 10:1:1–16.
- Denson, M.R, Stuart K.R, Smith T.I.J., Weirich, C.R. and Segars, A. 2003. Effects of salinity on growth, survival, and selected hematological parameters on juvenile cobia *Rachycentron canadum*. *Journal of World Aquaculture Society* 34(3): 496–504.
- Ditty, J.G. and Shaw, R.F. 1992. Larval development, distribution, and ecology of cobia *R. canadum* (Family: Rachycentridae) in the northern Gulf of Mexico. *Fisheries Bulletin* 90: 668-677.
- DoF, 2012. Department of Fisheries Malaysia, Annual Fisheries Statistics of Malaysia 2002-2012.
- DoF, 2014. Department of Fisheries Malaysia, Annual Fisheries Statistics of Malaysia. 2014.
- Dwivedi, A.C., Mayank, P., Masud, S. and Khan, S. 2009. An investigation of the population status and age pyramid of *Cyprinus carpio* var. communis from the Yamuna river at Allahabad. *The Asian Journal of Animal Science* 4: 98-101.
- Ebrahimi, M., Rajion, M. A. and Goh Y. M. 2014. Effects of Oils Rich in Linoleic and α -Linolenic Acids on Fatty Acid Profile and Gene Expression in Goat Meat. *Nutrients* 6: 3913-3928.
- Elliot, J. M. 1972. Rates of gastric evacuation in brown trout, *Salmo trutta*. *Freshwater Biology* 2: 1–18.
- Erkan, N. and Özden, O. 2007. Proximate composition and mineral contents in aqua cultured sea bass (*Dicentrarchus labrax*), sea bream (*Sparus aurata*) analyzed by ICP-MS. *Food Chemistry* 102(3): 721–725.
- Espinosa-Lemus V., Arredondo-Figueroa, J.L., Barriga-Sosa, I.D.L.A. 2009. Morphometric and genetic characterization of Tilapia (Cichlidae: Tilapiini) stocks for effective fisheries management in two Mexican reservoirs. *Hidrobiologica*. 19(2): 95–107.

- FAO, 2013a. A global assessment of offshore mariculture potential from a spatial Perspective. *Fisheries and Aquaculture Technical Paper* 549 Rome, Italy.
- FAO, 2013b. Global aquaculture production 2003-2013. Food and Agriculture Organization of the United Nations, Rome. Downloaded 25 Aug. 2014.
- FAO, 2012. Fishstat plus Vers. 2.3.2000: Universal software for fishery statistical time series: Capture production 2003-2013.
- FAO, 2010. Proceeding of the global conference for aquaculture. The state of world fisheries and aquaculture, Rome Italy.
- FAO, 2009. The state of the world fisheries and aquaculture, Fishery Information, Data and Statistics Unit.
- FAO, 2007. The world's mangroves 1980-2005. *FAO Forestry Paper* 2007; 153: 75.
- Fanelli, E. and Cartes, J.E. 2010. Temporal variations in the feeding habits and trophic levels of three deep-sea demersal fishes from the western Mediterranean Sea, based on stomach contents and stable isotope analyses. *Marine Ecology Progress Series* 402: 213–232.
- Faulk, C.K. and Holt G.J. 2008. Biochemical composition and quality of captive-spawned cobia *Rachycentron canadum* eggs. *Aquaculture* 279: 70–76.
- Faulk, C.K., Benninghoff, A.D. and Holt, G.J. 2007. Ontogeny of the gastrointestinal tract and selected digestive enzymes in cobia *Rachycentron canadum*. *Journal of Fish Biology* 70: 1-17.
- Faulk, C.K. and Holt, G.J. 2005. Advances in rearing cobia *Rachycentron canadum* larvae in recirculating aquaculture systems: live prey enrichment and green water culture. *Aquaculture* 249: 231–243.
- Faulk, C.K. and Holt, G.J. 2003. Lipid nutrition and feeding of cobia *Rachycentron canadum* larvae. *Journal of the World Aquaculture Society* 34: 368-378.
- Folch, J., Lees, M., Stanley, G.H.S. 1957. A simple method for the isolation and purification of total lipides from animal tissues. *The Journal of Biological Chemistry* 226: 497–509.
- Franks, J.S. and Brown-Peterson, N.J. 2002. A review of age, growth, and reproduction of cobia *Rachycentron canadum* from U.S. waters of the Gulf of Mexico and Atlantic Ocean. In *Proceedings of the 53rd annual Gulf and Caribbean Fisheries Institute*, ed. Creswell, R.L. 553 – 569.
- Franks, J.S., Ogle, J.T., Lotz, J.M., Nicholson, L.C., Barnes, D.N. and Larsen, K.M., 2001. Spontaneous spawning of cobia, *Rachycentron canadum*, induced by human chorionic gonadotropin (HCG), with comments on fertilization,

- hatching, and larval development. In *Proceeding of Gulf and Caribbean Fisheries Institute*, ed. Creswell, R.L. 52: 598–609.
- Franks, J.S. Warren, J.R. and Buchanan, M.V. 1999. Age and growth of cobia, *Rachycentron canadum*, from the northeastern Gulf of Mexico. *Fishery Bulletin* 97: 459-471.
- Franks, J.S., Garber, N.K. and Warren, J.R. 1996. Stomach contents of juvenile cobia, *R. canadum*, from the northern Gulf of Mexico. *Fisheries Bulletin* 94(2): 374-380.
- Fry, G.C. and Griffiths, S.P. 2010. Population dynamics and stock status of cobia, *Rachycentron canadum*, caught in Australian recreational and commercial coastal fisheries. *Fisheries Management Ecology* 17: 231 – 239.
- Fulton, T. 1902. Rate of growth of seas fishes. *Scientific Investigation and Fisheries Division of Scottish Report*. 20.
- Ganga, U., Pillai, N.G.K., Akhilesh, K.V., Rajoolshanis, C.P., Beni, N., Hashim, M. and Prakasan, D. 2012. Population dynamics of Cobia *R. canadum* (Linnaeus, 1766) off Cochin coast, south-eastern Arabian Sea. *Indian Journal of Fisheries* 59(3): 15-20.
- Ghanawi, J., Roy, L., Davis, D.A. Saoud, I.P. 2011. Effects of dietary lipid levels on growth performance of marbled spinefoot rabbitfish *Siganus rivulatus*. *Aquaculture* 310: 395–400.
- Gilliland, E.R., Kleinholz, C.W. and Clady, M.D. 1981. The efficiency of removing food items from fish with glass tubes. In *Proceeding of Texas - American Fisheries Society* 4: 95–100.
- Golani, D., and Ben-Tuvia, A. 1986. New records of fishes from the Mediterranean coast of Israel including Red Sea immigrants. *Cybiu* 10(3): 285-291.
- Golani D., Orsi-Relini, L., Massuti E. and Quignard J. P. 2002. CIESM atlas of exotic species in the Meditenanean. *Fishes Commission Internationale pour l'Exploration cientifique de la Mediter.raoe*, Monaco.
- Gold, J.R., Giresi, M.M., Renshaw, M.A. and Gwo, J.C. 2013. Population genetic comparisons among cobia from the Northern Gulf of Mexico, U.S. Western Atlantic, and Southeast Asia, North American. *Journal of Aquaculture* 75:(1)57-63.
- Gopakumar, G., Abdul Nazar, A.K., Tamilmani, G., Sakthivel, M., Kalidas, C, Ramamoorthy, N, Palanichamy, S., Ashok, V. M., Rao, K.S. and Rao, G.S. 2011. Broodstock development and controlled breeding of cobia *Rachycentron canadum* Linnaeus 1766) from Indian Seas. *Indian Journal of Fisheries* 58(4): 27-32.

- Grier, H.J. and Taylor, R.G. 1998. Testicular maturation and regression in the common snook. *Journal of Fish Biology* 53: 521–542.
- Gulland, J.A. 1982. Why do fish numbers vary? *Journal of Theoretical Biology* 97:69-75.
- Gulland, J.A. 1971. The fish resources of the Ocean. *Fishing News* 255.
- Hammond, D.L. 2001. Status of the South Carolina fisheries for cobia. *South Carolina Marine Resources Center Technical Report* 83.
- Harris, W.S., Assaad, B. and Poston, W.C. 2006. Tissue omega-6/omega-3 fatty acid ratio and risk for coronary artery disease. *American Journal of Cardiology* 21: 19-26.
- Hassler, W.W. and R.P. Rainville, 1975. Techniques for hatching and rearing cobia, *Rachycentron canadum*, through larval and juvenile stages. University of North Carolina Sea Grant College Program, Raleigh, North Carolina 26.
- Hitzfelder, G.M., Holt, G.J., Fox, J.M. and McKee, D.A. 2007. The effect of rearing density on growth and survival of cobia, *Rachycentron canadum*, larvae in a closed recirculating aquaculture system. *Journal of the World Aquaculture Society* 37: 204–209.
- Holt, G.J., Faulk, C.K. and Schwarz, M.H. 2007. A review of the larviculture of cobia *Rachycentron canadum*, a warm water marine fish. *Aquaculture* 268: 181-187.
- Hossain, M.Y., Ohtomi, J. and Ahmed, Z.F. 2009. Morphometric, meristic characteristics and conservation of the threatened fish, *Puntius sarana* (Hamilton, 1822) (Cyprinidae) in the Ganges River, northwestern Bangladesh. *Turkish Journal of Fisheries and Aquatic Science* 9: 223–225.
- Humphries, J.M., Bookstein, F.L., Chernoff, B., Smith, G.R., Elder, R.L. and Poss, S.G. 1981. Multivariate discrimination by shape in relation to size. *Systematic Biology* 30: 291-308.
- Hunter, J.R., Macewicz, B.J. and Sibert, H.R. 1986. The spawning frequency of skipjack tuna, *Katsuwonus pelamis*, from the south Pacific. *Fisheries Bulletin* 84:895–903.
- Hureau, J.C. 1969. Biologie comparée de quelques poissons anarctiques (Nototheniidae). *Bulletin of Institute for Oceanography Monaco* 68: 1–44.
- Hyslop E.J. 1980. Stomach contents analysis: a review of methods and their application. *Journal of Fish Biology* 17: 411–429
- Ibrahim A., Juahir H, Toriman M.E, Mustapha A., Azid A, Isiyaka, H.A. 2015. Assessment of surface water quality using multivariate statistical techniques

in the Terengganu river basin. *Malaysian Journal of Analytical Sciences* 19: 338 - 348.

Innis, S.M. 2000. Essential fatty acids in infant nutrition: lessons and limitations from animal studies in relation to studies on infant fatty acid requirements. *American Journal Clinical Nutrition* 71: 238-244.

Izquierdo, M.S., Ferna'ndez-Palacios, H. and Tacon, A.G.H. 2001. Effect of broodstock on reproductive performance in fish. *Aquaculture* 197: 25-42.

Jabeen, F. and Chaudhry, A.S. 2011. Chemical compositions and fatty acid profiles of three freshwater fish species. *Food Chemistry* 125: 991-996.

Jons, G.D. and Miranda, L.E. 1997. Ovarian weight as an index of fecundity, maturity and spawning periodicity. *Journal of Fish Biology* 50: 150-156.

Joseph, E.B., Norcross, J.J. and Massmann, W. H. 1964. Spawning of the cobia, *Rachycentron canadum*, in the Chesapeake Bay area, with observations of juvenile specimens. *Chesapeake Scientist* 5: 67-71.

Kaiser, J.B. and Holt, G.J. 2005. Species profile: cobia. *SRAC Publication*. 7202.

Kaiser, J.B. and G.J. Holt. 2004. Cobia: A new species for aquaculture in the US. *World Aquaculture* 35: 12-14

Kitts, D.D., Huynhl, M.D., Hu, C. and Trites, A.W. 2004. Seasonal variation in nutrient composition of Alaskan walleye pollock. *Canadian Journal of Zoology* 82: 1408-1415.

Kinsey, S.T., Orsoy, T., Bert, T.M. and Mahmoudi, B., 1994. Population structure of the Spanish sardine *Sardinella aurita*: natural morphological variation in a geneti-cally homogenous population. *Marine Biology* 118: 309-317.

Knapp, F.T. 1951. Food habits of the sergeant fish, *Rachycentron canadus*. *Copeia* 1951: 101-102.

Kolanowski, W. and Laufenberg, G. 2006. Enrichment of food products with polyunsaturated fatty acids by fish oil addition. *European Food Research Technology* 222: 472-477.

Kris-Etherton, P.M., Harris, W.S. and Appel, L.J. 2003. Nutrition committee. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Arteriosclerosis Thrombosis and Vascular Biology* 23: 20-30.

Le Xan, 2005. Advances in the seed production of Cobia *Rachycentron canadum* in Vietnam. *Marine Finfish Aquaculture Network* 1: 5-8.

Lefebvre, L.S. and Denson, M.R. 2012. Inshore spawning of *Rachycentron canadum* in South Carolina. *National Marine Fishery Bulletin* 4: 110.

- Lewis, G. 1957. The book of Florida fishing. Great Outdoors Publishing Co., St. Petersburg, Florida, 80.
- Li, L.D., Chen, B.S., Feng, J., Ke, H., Cai, J.X. and Fan, K. 2002. Analysis and evaluation in nutritive value of *Rachycentron canadum*. *Journal of Tropical Oceanography* 21: 76–82.
- Liao, I.C., Su, H.M. and Chang, E.Y. 2001. Techniques in finfish larviculture in Taiwan. *Aquaculture* 200: 1- 31.
- Liao, I.C., Huang, T., Tsai, W., Hsueh, C., Chang, S., Leano, E.M. 2004. Cobia culture in Taiwan: current status and problems. *Aquaculture* 237: 155-165.
- Liu, S.C., Li D.T., Hong P.Z., C.H., W.H., Gao, J.L. and Zhang, L. 2009. Cholesterol, lipid content, and fatty acid composition of different tissues of farmed cobia (*Rachycentron canadum*), from China. *Journal of American Oil Chemical Society* 86: 1155–1161.
- Lotz, J.M., Overstreet, R.M. and Franks, J.S. 1996. Gonadal maturation in the cobia, *Rachycentron canadum*, from the north central Gulf of Mexico. *Gulf Research Reports* 9: 147-159.
- Martin, W.R. 1949. The mechanics of environmental control of body form in fishes. *University of Toronto Studies Biology* 56: 1-91.
- Maceina, M.J., and Pereira, D. L. 2007. Recruitment. In *Analysis and interpretation of freshwater fisheries data* ed. C.S. Guy and M.L. Brown. American Fisheries Society, Bethesda, Maryland 121–185
- Mazorra, C., Bruce, M., Bell, J. G., Davie, A., Alorend, E., Jordan, N., Rees, J., Papanikos, N., Porter, M. and Bromage, N. 2003. Dietary lipid enhancement of broodstock reproductive performance and egg and larval quality in Atlantic halibut (*Hippoglossus hippoglossus*). *Aquaculture* 227: 21–33.
- Mazuki, R., Man, N., Omar, S.Z., Bolong, J., D'Silva, J.L., Shaffril, H.A.M. 2012. Technology adoption among fishermen in Malaysia. *Journal of American Science* 8(12): 1-4.
- Meyer, G.H. and Franks, J.S. 1996. Food of cobia, *Rachycentron canadum*, from the northcentral Gulf of Mexico. *Gulf Research Repot* 9(3): 161-167.
- Milner, N.J., Elliott, J.M., Armstrong, J.D., Gardiner, R., Welton, J.S. and Ladle, M. 2003. The natural control of salmon and trout populations in streams. *Fisheries Research* 62: 111–125.
- Moss, J.H., Farley, E.V., Feldman, A.M. and Ianelli, J.N. 2009. Spatial distribution, energetic status, and food habits of eastern Bering Sea age-0 walleye pollock. *Transaction of American Fisheries Society* 138: 497–505.

- Naguib, Y.M.A. 2000. Antioxidant activities of astaxanthin and related carotenoids. *Journal of Agriculture and Food Chemistry* 48: 1150-1154.
- Nagelkerken, I., Blaber, S.J.M., Bouillon, S. Green, P., Haywood, M., Kirton, L.G. Meynecke, J.O., Pawlik, J., Penrose, H.M., Sasekumar A. and Somerfield, P. J. 2008. The habitat function of mangroves for terrestrial and marine fauna: A review. *Aquatic Botany* 89: 155–185.
- Newman, S.J., 2002. Growth, age estimation and preliminary estimates of longevity and mortality in the moses perch, *Lutjanus russelli* (Indian ocean form), from continental shelfwaters off north-western Australia. *Asian Fisheries Science* 15: 283–294.
- Nhu, V.C., 2005. Present Status of Hatchery Technology for Cobia *Rachycentron canadum* in Vietnam. *Aquaculture Asia Magazine* Oct.–Dec.: 32–35.
- Nikolsky, G.V. 1963. The Ecology of fishes. Academy press, London and New York 352.
- Nguyen, M.V., Rønnestad I., Buttle, L., Lai, H.V. and Espe, M. 2014. Evaluation of a high plant protein test diet for juvenile cobia *Rachycentron canadum* in comparison to commercial diets. *Journal of Agricultural and Crop Research* 2(6): 117-125.
- Nguyen, H.Q., Tran, T.M., Reinertsen, H. and Kjørsvik, E. 2010. Effects of dietary essential fatty acid levels on broodstock spawning performance and egg fatty acid composition of cobia, *Rachycentron canadum*. *Journal of the world aquaculture society* 41:687-699
- Pauly, D. Palomares, M.L. Froese, R. Sa-a, P. Vakily, M. Preikshot, D. and Wallace, S. 2001. Fishing down Canadian aquatic food webs. *Canadian Journal of Fisheries and Aquatic Science* 58: 51-62.
- Pauly, D. 1995. Anecdotes and the shifting baseline syndrome of fisheries. *Trends in Ecology and Evolution* 10(10): 430.
- Pauly, D., 1980. On the interrelationships between natural mortality, growth parameters and mean environmental temperature in 175 fish stocks. *Journal of Conservation and Exploring Meridian* 39: 175–192
- Pauly, D., 1987. A review of the ELEFAN system for analysis of length-frequency data in fish and aquatic invertebrates. In *Proceeding of ICLARM* 13: 7-34.
- Pauly, D. 1984. Fish population dynamics in tropical waters: a manual for use with programmable calculators. *ICLARM Contributor* 143: 325.

- Pauly, D. and David, N., 1981. ELEFAN I, a basic program for the objective extraction of growth parameters from length-frequency data. *Meeresforsch* 28(4): 205-211.
- Phinchongsakuldit, J., Chaipakdee, P., James J., Collins F., Jaroensutasinee M. and Brookfield J.F.Y. 2013. Population genetics of cobia (*Rachycentron canadum*) in the Gulf of Thailand and Andaman Sea: fisheries management implications *Aquaculture International* 21:197–217.
- Pillay, T.V.R. 1952. A critique of the methods of study of food of fishes. *Journal of Zoological Society of India* 4: 1885-200.
- Pinkas, L., Oliphant, M.S. and Iverson, I.L.K. 1971. Food habits of albacore, bluefin tuna, and bonito in California waters. *California Department of Fisheries, Game Fish Bulletin* 152: 105-110.
- Pruett, C.L, Saillant, E. and Renshaw M. 2005. Microsatellite DNA markers for population genetic studies and parentage assignment in cobia, *Rachycentron canadum*. *Molecular Ecology* 5:8 4–86.
- Rajan, P.R., Lopez, C., Lin, J.H. and Yang, H. 2001. *Vibrio alginolyticus* infection in cobia (*Rachycentron canadum*) cultured in Taiwan. *Bulletin of the European Association of Fish Pathologists* 21(6): 228–234.
- Ramiah S.K., Meng G.Y., Ebrahimi, M. 2014. Dietary Conjugated Linoleic Acid Alters Oxidative Stability and Alleviates Plasma Cholesterol Content in Meat of Broiler Chickens. *The Scientific World Journal* doi:10.1155/2014/949324.
- Resley, M.J., Webb, K.A. Jr. and Holt, G.J. 2006. Growth and survival of juvenile cobia, *Rachycentron canadum*, at different salinities in a recirculating aquaculture system. *Aquaculture* 253: 398-407.
- Richards, C.E. 1967. Age, growth, and fecundity of the cobia, *Rachycentron canadum*, from Chesapeake Bay and adjacent mid-Atlantic waters. *Transaction of American Fisheries Society* 96(3): 343-350.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. *Bulletin Fisheries Research Board of Canada* 191: 382.
- Rohit, P. and Bhat, U.S. 2012. Fishery and Diet composition of the cobia *R. canadum* (Linnaeus, 1766) exploited along Karnataka coast. *Indian Journal of Fisheries* 59(4): 61-65.
- Rudel, L.L. and Morris, M.D. 1973. Determination of cholesterol using o-phthalaldehyde. *Journal of Lipid Research* 14: 1973.
- Sagarese, S.R., Cerrato, R.M. and Frisk, M.G. 2011. Diet Composition and Feeding Habits of Common Fishes in Long Island Bays, New York. *Northeastern Naturalist* 18(3): 291-314.

- Sahoo, S.K, Giri, S.S.S. Mohapatra C.B.C. 2008. Evaluation of breeding performance of Asian catfish *Clarias batrachus* at different dose of HCG and latency period combinations. *Turkish Journal of Fisheries and Aquatic Sciences* 8: 249-251.
- Sajeevan, M.K. 2014. Osteological features of Cobia, *Rachycentron canadum* (Linnaeus 1766). *Journal of the ocean science foundation* 11: 40-49.
- Sajeevan, M.K. and Kurup, B.M. 2014. Evaluation of feeding indices of cobia *Rachycentron canadum* (Linnaeus 1766) from northwest coast of India. *Journal of Marine Biology Association India* 55 (2): 16-21.
- Sajina, A.M., Chakraborty, S.K., Jaiswar, A.K., Pazhayamadam, D.G. and Sudheesan, D. 2011. Stock structure analysis of Indian mackerel, *Rastrelliger kanagurta* (Cuvier, 1816) along the Indian coast. *Asian Fisheries Science* 24: 331-342.
- Sakthivel, M., Abdul Nazar, A.K., Tamilmani, G., Kalidas, C., Ramamoorthy, N. , Ashok Maharshi, V., Rao K.S.S. and Gopakumar G. 2012. Embryonic development of cobia, *Rachycentron canadum* (Linnaeus,1766) in controlled conditions. *Journal of Marine Biology Association, India*, 54: 2-12.
- Salze, G., McLean, E., Battle, P.R., Schwarz, M.H. and Craig, S.R. 2010. Use of soy protein concentrate and novel ingredients in the total elimination of fish meal and fish oil in diets for juvenile cobia, *Rachycentron canadum*. *Aquaculture* 298:294–299.
- Sandlund, O.T., Gunnarsson, K., Jónasson, P.M., Jonsson, B., Lindem, T., Magnússon, K.P., Malmquist, H.J., Sigurjónsdóttir, H., Skúlason, S. and Snorrason, S.S. 1992. The arctic charr *Salvelinus alpinus* in Thingvallavatn. *Oikos* 64: 305–351.
- Sargent, J., Bell, G., McEvoy, L., Tocher, D. and Estevez, A. 1999. Recent developments in the essential fatty acid nutrition of fish. *Aquaculture* 177: 191–199.
- Shaffer, R.V. and Nakamura. E.L. 1989. Synopsis of biological data on the cobia *Rachycentron canadum* (Pisces: Rachycentridae). *FAO Fisheries Synopsis* 153.
- Scherrer, B., 1984. Biostatistique. Morin, Montreal, Paris.
- Schmidt, E.B., Arnesen, H., de Caterina, R., Rasmussen, L.H. and Kristensen, S.D. 2005. Marine omega-3 polyunsaturated fatty acids and coronary heart disease: Part I. Background, epidemiology, animal data, effects on risk factors and safety. *Thrombosis Research* 115(3): 163–170.
- Simopoulos, A.P. 2004. Omega-6/Omega-3 Essential Fatty Acid Ratio and Chronic Diseases. *Food Reviews International* 20(1) 77–90.

- Smith, J.W. and Merriner, J.V. 1982. Association of cobia, *Rachycentron canadum*, with cownose ray, *Rhinoptera bonasus*. *Estuaries* 5(3): 240-242.
- Smith, J. W. 1995. Life history of cobia, *Rachycentron canadum* (Osteichthyes: Rachycentridae), in North Carolina waters. *Brimleyana*, 23: 1-23.
- Somvanshi, V.S., Vargese, S., Gulati, D.K. and Bhargava, A.K. 2000. Some Biological aspects of kingfish *Rachycentron canadum* (Linnaeus, 1766) from the north-west Indian EEZ. *Occasional Paper, Fishery Survey of India, Mumbai*.
- Sparre, P., Ursin, E. and Venema, S.C., 1989. Introduction to tropical fish stock assessment. *FAO Fisheries Technical Paper* 306: 192–218.
- Sparre, P. and Venema, S.C., 1998. Introduction to tropical fish stock assessment, Part I -Manual. *FAO Fisheries Technical Paper* 306: 376.
- Sparre, P., Venema, S.C., 1992. Introduction to Tropical Fish Stock Assessment, Part 1—Manual. *FAO Fisheries Technical Paper* 306/1: 376.
- Stieglitz, J.D., Benetti, D. D., Hoenig, R. H., Sardenberg, B., Welch, A.W. and Miralao, S. 2012. Environmentally conditioned, year-round volitional spawning of cobia (*Rachycentron canadum*) in broodstock maturation systems. *Aquaculture Research* 43: 1557-1566.
- Strauss, R.F. and Bookstein, F.L. 1982. The Truss: Body form reconstructions in morphometrics. *Systematic Zoology* 31: 113-135.
- Su, Y., Sun, X., Feng J., Guo, Z., Xu, L. and Wang, J. 2008. Morphological and histological observations of digestive system of cobia *Rachycentron canadum*. *South China Fisheries Science* 6: 88-94.
- Suriah R.A, Huah, T.S., Hassan, O. and Daud, N.M. 1995. Fatty acid composition of some Malaysian freshwater fish. *Journal of Food Chemistry* 54: 45–49.
- Sushchik, N.N., Gladyshev, M.J. and Kalachova, G.S. 2007. Seasonal dynamics of fatty acid content of common food fish from the Yenisei River, Siberian grayling, *Thymallus arctius*. *Food Chemistry* 104: 1353–1358.
- Szlinder-Richert, J., Usydus, Z., Wyszyn'ski, M., Adamczyk, M. 2010. Variation in fat content and fatty acid composition of the Baltic herring (*Clupea harengus membras* L). *Journal of Fish Biology* 77: 585–599.
- Tandler, A., Harel, M., Koven, W.M. and Kolkovsky, S. 1995. Broodstock and larvae nutrition in gilthead sea bream *Sparus aurata* – new findings on its involvement in improving growth, survival and swim bladder inflation. *Israel Journal of Aquaculture Bamidgeh* 47: 95–111.

- Tesoriere, L., D'Arpa, D., Butera, D., Pintaudi, A.M., Allegra, M. and Livrea, M.A. 2002. Exposure to malondialdehyde induces an early redox unbalance preceding membrane toxicity in human erythrocytes. *Free Radical Research* 36: 89-97.
- Thompson, B.A., Wilson, C.A., Render, J.H. and Beasley, M. 1992. Age, growth and reproductive biology of greater amberjack and cobia from Louisiana waters. *Final report to Marine Fisheries Research Initiative (MARFIN) Program, NMFS, St. Petersburg, FL. NA90AA-H-MF722.*
- Thurman, H.V. 1997. Introduction to Oceanography. Prentice Hall, Upper Saddle River Press.
- Tocher, D.R., Bell, J.G., Dick, J.R. and Crampton, V.O. 2003. Effect of dietary vegetable oil on Atlantic salmon hepatocyte fatty acid desaturation and liver fatty acid compositions. *Lipids* 38: 723-732.
- Torres, M.V., Giri, F. and Collins, P.A., 2014. Geometric morphometric analysis of The freshwater prawn *Macrobrachium borellii* (Decapoda: Palaemonidae) at a micro-geographical scale in a floodplain system. *Ecological Resources* 29: 959-968. Troadec, J.P. 1980. Utilization of resource survey results in stock assessment. *Canada Funds-in-Trust, FAO/TF/INT* 180: 139-52.
- Trushenski, J.T., Laporte, J., Lewis, H., Schwarz, M., Delbos, B., Takeuchi, R. and Sampaio, L.A., 2011. Fish meal replacement with soy-derived protein in feeds for juvenile cobia: influence of replacement level and attractant supplementation. *Journal of the World Aquaculture Society* 42: 435-443.
- Turan, C., Oral, M., Ozturk, B. and Duzgunes, E. 2006. Morphometric and meristic variation between stocks of bluefish (*Pomatomus saltatrix*) in the Black, Marmara, Aegean and northeastern Mediterranean Seas. *Fisheries Resource* 79: 139-147.
- Usydus, Z., Szlifder-Richert, J. and Adamczyk, M. 2012. Variations in proximate composition and fatty acid profiles of Baltic sprat (*Sprattus sprattus balticus*) *Food Chemistry* 130: 97-103.
- Van Den Avyle, M.J. and Rousse., J.E. 1980. Evaluation of a simple method for removing food items from live black bass. *Progressive Fish-Culturist* 42: 222-223.
- Van der Velde, T.D., Griffiths, S.P. and Fry, G.C. 2010. Reproductive biology of the commercially and recreationally important cobia *Rachycentron canadum* in northeastern Australia. *Fry Fish Science* 76: 33-43.
- Watson, A.M., Barrows, F.T. and Place, A.R. 2013. Taurine supplementation of plant derived protein and n-3 fatty acids are critical for optimal growth and development of cobia, *Rachycentron canadum*. *Lipids* 48: 899-913.

- Weirich, C.R., Stokes, A.D., Smith, T.J.I., Jenkins, W.E. and Denson, M.R. 2006. Outdoor tank and pond spawning of cobia, *Rachycentron canadum* Coastal South Carolina. *Journal of Applied Aquaculture* 18(3):1-16
- WHO, 2003. Report of the Joint WHO/FAO Expert consultation on diet, nutrition and the prevention of chronic. *WHO Technical Report Series* 916.
- Williams, E. H. 2001. Assessment of cobia, *Rachycentron canadum*, in the waters of the U.S. Gulf of Mexico. *NOAA Technical Memorandum NMFS-SEFSC*. 469.
- Winston, G.W. and Di Giulio, R.T. 1991. Prooxidant and antioxidant mechanism in aquatic organism. *Aquatic Toxicology* 19: 137-161.
- Woods, A.E. and Ellis, R.C. 1994. Laboratory histopathology: A complete reference. Churchill Livingstone (Eds.), New York.
- Zink I, Cavalin, F., Bacoat, D., Denlinger, B., Palmer, K., Sardenberg, B., Kirkpatrick, R., Orhun, R. and Benetti, D. 2006. U. S. trials compare commercial diets fed to juvenile cobia. *Global Aquaculture Advocate* 53-54.
- Zhou, Q.C., Wu, Z.H., Chi, S.Y. and Yang, Q.H. 2007. Dietary lysine requirement of juvenile cobia (*Rachycentron canadum*). *Aquaculture* 273: 634 – 640.