



UNIVERSITI PUTRA MALAYSIA

***SPECTRAL APPROACHES TO NUTRIENT STRESS DETECTION IN
PINEAPPLE CULTIVATION ON A TROPICAL PEAT SOIL***

CHONG YEN MEE

FP 2018 2



**SPECTRAL APPROACHES TO NUTRIENT STRESS DETECTION IN
PINEAPPLE CULTIVATION ON A TROPICAL PEAT SOIL**

By

CHONG YEN MEE

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Master of
Science**

November 2017

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Science

SPECTRAL APPROACHES TO NUTRIENT STRESS DETECTION IN PINEAPPLE CULTIVATION ON A TROPICAL PEAT SOIL

By

CHONG YEN MEE

November 2017

Chairman: Associate professor Siva Kumar Balasundram, PhD
Faculty: Agriculture

Pineapple in Malaysia is largely cultivated on peat which has resulted in problems including poor drainage, low fertility, subsidence and irreversible drying which rarely occurred on mineral soil. Hence, replenishment of nutrients and organic matter to soil through rational fertiliser application is part of the precision agriculture practices to enhance soil quality and crop productivity. To further avoid excessive application of fertilisers, a study was conducted to measure the reflectance changes of nutrient stress induced by nitrogen (N) and potassium (K) deficiencies, using a spectroradiometer. While other parameters such as growth measurement including height, leaf width and leaf number were made at 45, 135 and 225 days after planting, alongside with leaf nitrogen content measurement with hand-held Minolta SPAD-502 chlorophyll meter. Four different levels of nitrogen and potassium were applied to the pot as Urea (46 % N) and Muriate of Potash (60 % K₂O) respectively. All suckers of MD2 variety were pre-treated with fungicide before potted on raised beds in double row, spaced 0.6 m x 0.6 m on the bed and 0.9 m apart within row in a randomised complete block design. The fertilisers were applied at three split periods of 3-, 5- and 8-month after planting [N0 (control with no fertiliser added); N1 (369 kg/ha); N2 (528 kg/ha); N3 (686 kg/ha); K0 (control with no fertiliser added); K1 (256 kg/ha); K2 (366 kg/ha); K3 (476 kg/ha). Other fertilisers were applied equally for all pots either *via* broadcast or micronutrient fertiliser spray. In this study, several published indices were used to evaluate their capabilities to quantify leaf N concentration of pineapple grown under four contrasting levels of N & K. Results showed that different vegetation indices were found most useful depending on crop phenology. The three indices, the Narrow Band Normalised Difference Vegetation Index (NBNDVI), Greenness Index (GI) and Nitrogen Reflectance Index (NRI) performed better at differentiating N rates from mid to advanced growth stages. While the Normalised Difference Red Edge Index (NDRE) performed equally well but at more advanced growth stage and the Normalised Difference Vegetation Index (NDVI) was the least performing index with greater sensitivity to N stress in young plant rendering it

ineffective in present study focusing on mid to advance growth stages of pineapple. On the other hand, K deficiency tested reported three out of six indices evaluated in this study to be effective at differentiating K rates across varying growth stages. The Normalised Pigment Chlorophyll Index (NPCl) and Plant Senescence Reflectance Index (PSRI) were found to be useful throughout the entire study period, while the Red Edge Vegetation Stress Index (RVS_I) only showed minimal differences between K rates. The other three indices including NDRE, NDVI and NBNDVI were found to be ineffective in K study. Additionally, canopy reflectance of K-deficient pineapple also exhibited a shift of red edge towards shorter wavelengths of 500-700 nm as often observed in the absence of N. The results ascertained that canopy and leaf reflectance changes are important indicators of fertiliser application. The changes of canopy and leaf reflectance may show whether the plant has received sufficient nutrients, so that fertiliser can be applied at appropriate amount without being wasted.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Sarjana Sains

**PENDEKATAN SPEKTRA DALAM MENGESAN TEKANAN NUTRIEN DI
PENANAMAN NANAS ATAS TANAH GAMBUT**

Oleh

CHONG YEN MEE

November 2017

Pengerusi: Profesor Madya Siva Kumar Balasundram, PhD
Fakulti: Pertanian

Sebahagian besar nanas di Malaysia ditanam di tanah gambut dan bukannya tanah mineral seperti mana di negara-negara pengeluar yang lain di rantau Asia Tenggara. Penanaman nanas di tanah gambut telah mendatangkan masalah termasuklah saliran miskin, kesuburan tanah yang rendah, penenggelaman serta pengeringan tidak berbalik yang jarang berlaku di atas tanah mineral. Oleh itu, penggantian semula nutrien dan bahan organik ke tanah melalui penggunaan baja yang rasional, adalah sebahagian daripada amalan ketepatan pertanian (*precision agriculture*) demi meningkatkan kualiti tanah dan produktiviti tanaman. Bagi mengelakkan penggunaan baja yang berlebihan, satu kajian mengaplikasi spektoradiometer telah dijalankan untuk mengukur perubahan pantulan tekanan nutrien akibat kekurangan nitrogen (N) dan kalium (K). Walaupun parameter lain seperti ukuran pertumbuhan, termasuklah ketinggian, kelebaran daun dan bilangan daun akan diambil pada 45, 135 dan 225 hari selepas penanaman, berserta ukuran kandungan nitrogen daun menggunakan SPAD-502 Minolta meter. Empat tahap nitrogen dan kalium telah dibubuh pada pot sebagai urea (46 % N) dan muriate of potash (60 % K). Kesemua tunas nanas varieti MD2 dirawat dengan racun kulat sebelum ditanam atas tapak bertimbun dalam dua barisan, berjarak 0.6 m x 0.6 m atas tapak, dan 0.9 m dalam barisan berbentuk blok rawak lengkap. Baja-baja dibubuh mengikut tiga tempoh berasingan pada 3-, 5- dan 8-bulan selepas penanaman [N0 (kawalan tanpa baja); N1 (369 kg/ha); N2 (528 kg/ha); N3 (686 kg/ha); K0 (kawalan tanpa baja); K1 (256 kg/ha); K2 (366 kg/ha); K3 (476 kg/ha)]. Baja lain turut dibubuh sama rata melalui semburan mikronutrien. Dalam kajian ini, beberapa indeks yang telah diterbitkan dinilai keupayaan mereka dalam mengukur kepekatan nitrogen daun nanas bawah empat rawatan N dan K. Keputusan mendapati indeks tumbuhan yang berlainan berguna bergantung pada fenologi tumbuhan. Ketiga-tiga indeks, Indeks Perbezaan Band Terpilih Vegetasi (NBNDVI), Indeks Kehijauan (GI) dan Indeks Refleksi Nitrogen (NRI) didapati berkesan dalam membezakan kadar N dari peringkat pertengahan hingga peringkat matang pertumbuhan. Walaupun Indeks Perbezaan Red Edge Vegetasi (NDRE)

berprestasi baik, namun hanya terhad kepada peringkat matang pertumbuhan. Manakala Indeks Perbezaan Vegetasi (NDVI) adalah indeks yang berprestasi kurang memuaskan dengan kepekaan yang lebih tinggi terhadap tekanan nutrien nitrogen pada peringkat pertumbuhan muda. Ini menyebabkan ia kurang berguna dalam kajian ini yang bertumpu pada peringkat pertengahan ke matang pertumbuhan nanas. Sebaliknya, kekurangan K yang diuji mendapati tiga daripada enam indeks berguna dalam membezakan kadar K sepanjang peringkat pertumbuhan. Indeks Nisbah Pewarnaan Klorofil (NPCI) dan Indeks Refleksi Penderiaan Tanaman (PSRI) didapati berguna sepanjang tempoh kajian. Manakala Indeks Tekanan Vegetasi (RVSI) hanya menunjukkan perbezaan yang sedikit antara kadar K. Tiga indeks lain termasuklah NDRE, NDVI dan NBNDVI, didapati tidak berkesan dalam kajian atas K. Selain itu, pantulan kanopi nanas bawah tekanan nutrient K juga memaparkan pergeseran red edge ke arah gelombang lebih pendek dalam lingkungan 500-700 nm, sebagaimana yang dilihat semasa tanpa N. Hasil kajian membuktikan bahawa perubahan pantulan kanopi dan daun adalah indikator penting dalam aplikasi baja. Perubahan pantulan kanopi dan daun akan menentukan sama ada tumbuhan menerima nutrien yang mencukupi, supaya baja dapat dibubuh pada kadar yang sesuai dengan tidak mensia-siakan.

ACKNOWLEDGEMENTS

First of all, I owe my greatest gratitude to my supervisor and co-supervisor, Associate Professor Dr. Siva Kumar Balasundram and Associate Professor Dr. Ahmad Husni Mohd Haniff, who spent his precious time to provide me the guidance, the advices and the encouragement I needed most to accomplish my thesis. I really appreciated his efforts to find time to meet and discuss with me, despite them being so busy.

Besides that, I had been fortunate to have the assistance from the plantation manager of Peninsula Plantation Sdn Bhd in Simpang Renggam, Johor who greeted us with warm welcoming and providing free suckers for my study.

I would like to take this special chance to thank my mother, for the countless sacrifices she made to the family, and my elder sister, for both of them showing enthusiastic support, patience and love. Meanwhile, my father remained as an important person in my life though he had passed away, I am sure he would be proud to know that I have not disappointed him.

I am also grateful to my other team mates, including Mr. Kamlesh and Ms. Husna. It had been great to have known all of you, and the team work was a pleasure experience.

Lastly, I would like to thank those who have contributed directly or indirectly to the completion of my thesis, the examiners for the precious suggestions they gave to correct my thesis, and not to forget the lab assistants of faculty of soil science for all the help at the laboratory.

TABLE OF CONTENTS

		Page
	ABSTRACT	i
	ABSTRAK	iii
	ACKNOWLEDGEMENTS	v
	APPROVAL	vi
	DECLARATION	viii
	LIST OF TABLES	xii
	LIST OF FIGURES	xiii
CHAPTER		
1	INTRODUCTION	1
2	LITERATURE REVIEW	4
	2.1 Environmental conditions for plant growth and development	4
	2.1.1 Temperature	4
	2.1.2 Relative humidity	5
	2.1.3 Soil properties	6
	2.2 Remote sensing in nutrient stress detection	8
	2.2.1 Multispectral/hyperspectral approach	8
	2.2.2 Chlorophyll fluorescence	8
	2.2.3 Thermography	9
	2.2.4 Development of vegetative indices	10
	2.3 Types of nutrient required for pineapple growth and regulation	11
	2.3.1 Nitrogen	11
	2.3.2 Potassium	11
	2.3.3 Calcium	11
	2.3.4 Phosphorus	12
	2.3.5 Magnesium	12
	2.3.6 Zinc, Sulphur and Iron	12
3	PINEAPPLE CANOPY AND LEAF N AND K LEVELS	13
	3.1 Introduction	13
	3.2 Materials and methods	14
	3.2.1 Plant materials and growth conditions	14
	3.2.2 Treatments	14
	3.2.3 Measurements	15
	3.2.4 Plant maintenance	16

3.2.5	Pineapple leaf samplings	17
3.2.6	Analysis of plant samples	17
3.2.7	Soil samplings and preparation	17
3.2.8	Soil analysis	18
	3.2.8.1 Chemical properties	18
	3.2.8.2 Physical properties	19
3.3	Results and discussion	19
3.3.1	Data analysis	19
	3.3.1.1 Effect of N and K fertilisers on vegetative growth of pineapple	19
	3.3.1.2 Leaf reflectance measurement using SPAD chlorophyll meter	21
	3.3.1.3 Comparison of SPAD readings and laboratory determination of leaf N and K	23
	3.3.1.4 Canopy reflectance measurement using spectroradiometer	24
	3.3.1.5 Soil factors influencing plant nutritional status	25
3.4	Conclusion	27
4	RESULTS AND DISCUSSION	28
4.1	Introduction	28
4.2	Materials and methods	29
	4.2.1 Determination of spectral indices corresponding to N and K sufficiency or deficiency in pineapple	27
	4.2.1.1 Study site and experimental design	29
	4.2.1.2 Spectral data collection and processing	29
	4.2.1.3 Statistical analysis	29
4.3	Results and discussion	30
4.4	Conclusion	37
5	SUMMARY AND CONCLUSION	38
	REFERENCES	40
	BIODATA OF STUDENT	47

LIST OF TABLES

Table	Page
1 N treatment used in pot trial	15
2 K treatment used in pot trial	15
3 Effects of N treatments on pineapple growth	20
4 Effects of K treatments on pineapple growth	20
5 First fertilisation	21
6 Second fertilisation	21
7 Third fertilisation	21
8 Leaf total N	22
9 Leaf total K	22
10 Correlation coefficients for total N versus SPAD readings	22
11 Soil properties at 0-10cm before the experiment	23
12 Initial status of total P, K, Cu, Fe, Mn, Zn, Ca, Mg	25
13 Variation of spectral indices responses to N treatments	29
14 Correlation coefficients for indices derived from spectroradiometer versus leaf chlorophyll readings	31
15 Summary of descriptive statistics for three best spectral indices at different growth stages	32

LIST OF FIGURES

Table		Page
1	Plastic shields for fruit protection	5
2	Curvilaria Blight on pineapple in (A) pineapple estate (B) pot trial	6
3	Canopy reflectance of pineapple under four N rates at 254 days after planting	23
4	Canopy reflectance of pineapple under four K rates at 254 days after planting	24



CHAPTER 1

INTRODUCTION

In recent years, there has been an increasing interest in understanding fertiliser management in relation to crop nutrient needs. Established crops are confined to a particular compound and therefore are dependent upon the characteristics of the environment they occupy. These environmental characteristics such as temperature, moisture and soil quality are known to play important roles in crop growth and reproduction. However, unsustainable agricultural practices and intensive cropping have resulted in rapid nutrient washout which render the soil infertile for crop (Mafongoya et al., 2006). This has eventually affected crop performance with problems like stunted growth, reduced shoots and roots growth depending on the severity of deficiency (Hawkesford et al., 2012).

In view of all that have been mentioned so far, it seemed that replenishment of nutrients and organic matter to soil through rational fertiliser application is a good precision agriculture practice in enhancing soil quality and crop productivity. Environmental pollution resulted from excessive fertiliser inputs is fast becoming a great concern where nutrient applied above the crop requirement often leached through soil and contaminate ground water (Zhao et al., 2006). Recently it has been found that excessive inorganic fertiliser especially that of nitrogen (N) will lead to higher emission of greenhouse gases such as nitrous oxide (N₂O) and carbon dioxide (CO₂) across different agricultural systems (Wang et al., 2011; Linquist et al., 2012).

Considering the fact that crop is generally sensitive to changes of the surrounding environment especially stress imposed upon them due to unfavourable condition, it is paramount to monitor crop nutritional status on real time basis so that fertilisation regime can be adjusted accordingly. Traditionally, time-consuming and expensive laboratory analyses as well as detailed sampling have dominated the market. Recent development of science and technology especially in the field of remote sensing has brought upon the improved and more up-to-date remote sensing tools which are found to be both economically and environmentally feasible at large scale. These remote sensing tools are proven effective in assessing crop biochemical properties changes triggered by stressors like pest and disease, water deficit and nutrient scarcity. In this respect, spectral parameters including leaf pigment content, water status, leaf area index (LAI) are often gauged as useful indicators of crop stress physiology.

Studies of nutritional status of crop revealed how the interactions of different environmental factors contributing to the alteration of the plant features. It is recalled that crop responds to environmental changes through structural modifications especially that of architecture and morphology for continued

survival. These long-term adaptation responses usually manifest in the alterations of leaf size, number and thickness, as well as development of special features like extended root system and modified architecture for better access to limited nutrients and water in soil (Hiradate et al., 2007). However, these internal metabolic and growth responses are unable to provide ultimate protection and adverse effects caused by the constraints especially concerning vegetative growth and yield are common in plant.

From these perspectives, it is noted that while there have been many studies done on crop nutritional status and effects on crops, few have explored the crop nutrient demand at different growth stage. This information is crucial as it can be related to nutrient application in field and also its associated crop yield. In addition to common spectral tools such as SPAD meter, extensive research using more advanced remote sensing techniques such as hyperspectral sensor Spectroradiometer also needs to be conducted in gathering more details of the plant functional properties alterations to better address the nutrient deficiency problem in plant.

Fertiliser management remains one of the greatest challenges in maintaining crop production in agriculture. The common perception is that increased fertiliser input always results in higher crop yield, and that explains why farmers are willing to spend more on fertilisation and even ignore the risk of polluting the environment. However, studies showed that crop nutrient uptake relied very much on crop nutrient demand which vary across crop types and soil medium. Under this context, nutrient use efficiency of potassium (K) in pineapple cultivated on peat soil is low due to the nutrient rapid leaching in low clay and mineral content condition (Ahmed et al., 2005). On the other hand, nitrogen (N) content in peat is abundant but its uptake in pineapple depends on a range of factors including temperature, moisture and C/N ratio. This concurred with some findings that suggested recovery of N for most crops ranged from 40-60 % and a lot more are lost through leaching and volatilisation (Aulakh and Malhi, 2005).

Similar study on Oil Palm found a leaching loss of 1 to 1.6 % of N fertiliser applied to Oil Palm and a higher rate of K loss between 2.4 and 5.3 % at variable fertiliser application rate and may associate with factors including crop age, soil type and rainfall intensity. The uncertain amount of available N in peat has resulted in additional input of N applied to pineapple through fertilisation (Comte et al., 2012). Further to that findings, it is suggested that N Use Efficiency also correlates with crop nutritional status as well as the amount of N applied; where nitrogen uptake was slow in well-nourished crop and was more rapid in nitrogen-deficient crop. Conversely, K-deficient Olive plant was less ready in taking up K fertiliser applied (Fernández-Escobar et al., 2014). Besides, some findings suggested that crop nutrient needs change at different growth stages. In this case, nutrient use efficiency is closely linked to its growth capacity which deals directly with crop management practice, site-specific spatial variability and its genetic potential (Lemaire and Gastal, 2009). Therefore, it is now time to utilise remote sensing technique for continuous monitoring of the pineapple nutrient status across different growth periods to determine the crucial timing for nutrient

application over different nutrient requirement in crop. Such move will minimise fertiliser input while maintaining pineapple yield at economic cost in the long run.

Pineapple (*Ananas comosus* L.) is largely cultivated on peat soil in Malaysia. This has resulted in restrained production of pineapple in the country as we faced several problems associated with peat soil characteristics such as poor drainage, low fertility and irreversibly drying (Ahmed et al., 2005). Other producer countries of the Southeast Asian regions can easily produce pineapple in bulk without these obstacles as the cultivation is all done on mineral soil. As reported by online statistics of Malaysian Pineapple Industry Board (MPIB), Malaysia ranked four in pineapple production right after Thailand, the Philippines and Indonesia (Ahmed et al., 2013). However, we are still able to improve our pineapple production through better crop management especially fertilisation in the plantation. This is in line with Malaysia's target of expanding the current 15,000 hectare of areas planted with pineapples to 24,000 hectare by year 2020. The move is expected to export Malaysian pineapple on a large scale to meet the demands in China as stated in an agreement signed by both countries in year 2013 (Zazali, 2014).

It is hoped that with better understanding of pineapple nutrient demand especially of nitrogen and potassium, we are then able to adjust fertilisation input in a manner that maximizes our pineapple growth potential as well as its yield and quality. Under this context, this study aims to determine spectral reflectance changes in the visible (380-720 nm) and infrared range (720-1500 nm) at different leaf N and K levels, and then develop spectral indices that indicate pineapple nutritional status along the growth cycle and ensure efficiency recovery of fertiliser applied in the pineapple field.

The current study undertaken with the objectives of studying leaf reflectance in response to four different leaf nitrogen (N) and potassium (K) levels ranged from 0 to 250 kg/ha across three planting ages of 3-, 5- and 8-month for pineapple (*Ananas comosus*) in pots. In doing so, spectral reflectance of healthy and nutrient-deficient plant are compared for pot trial following fertilisation scheme of local plantation estate, and subsequently validated in field. Furthermore, spectral indices corresponding to leaf N and K sufficiency as well as deficiency are also developed. The remote sensing tools utilised in developing the spectral indices for the assessment are Spectroradiometer and Greenseeker.

REFERENCES

- Ahmed, O. H., Ahmad, H. M., Musa, H. M., Rahim, A. A., & Rastan, S. O. (2005). Applied K fertilizer use efficiency in pineapples grown on a tropical peat soil under residues removal. *Scientific World Journal*, 5, 42-49.
- Ahmed, O. H., Ahmad, H. M., Rahim, A. A., & Musa, M. H. (2013). Sustainable production of pineapples on tropical peat soils. Serdang: Universiti Putra Malaysia Press.
- Ahmed, O. H., Husni, M. H. A., Hanafi, M. M., Anuar, A. R., & Syed Omar, S. R. (2007). Evaluation of soil applied phosphorus fertilizer use in pineapple cultivation with in situ burning of the crop's residues on tropical peat soils. *Communications in Soil Science and Plant Analysis*, 38, 1243-1254.
- Aulakh, M. S., & Malhi, S. S. (2005). Interactions of Nitrogen with Other Nutrients and Water: Effect on Crop Yield and Quality, Nutrient Use Efficiency, Carbon Sequestration, and Environmental Pollution. In L. S. Donald (Ed.), *Advances in Agronomy*, 86, 341-409: Academic Press.
- Benediktyova, Z., & Nedbal, L. (2009). Imaging of multi-color fluorescence emission from leaf tissues. *Photosynthesis Research*, 102, 169-175.
- Blackburn, G. A. (2007). Hyperspectral remote sensing of plant pigments. *Journal of Experimental Botany*, 58(4), 855-867.
- Bray, R. H., & Kurtz, L. T. (1945). Determination of Total, Organic and Available Forms of Phosphorus in Soils. *Soil Science*, 59, 39-45.
- Briat, J.F., Dubos, C., & Gaymard, F. (2015). Iron nutrition, biomass production, and plant product quality. *Trends in Plant Science*, 20(1), 33-40.
- Bürling, K., Hunsche, M., & Noga, G. (2011). Use of blue-green and chlorophyll fluorescence measurements for differentiation between nitrogen deficiency and pathogen infection in winter wheat. *Journal of Plant Physiology*, 168(14), 1641-1648.
- Buschmannm, C., Langsdorf, G., & Lichtenthaler, H. K. (2000). Imaging of the blue, green and red fluorescence emission of plants: An overview. *Photosynthetica*, 38, 483-491.
- Cadet, É., & Samson, G. (2011). Detection and discrimination of nutrient deficiencies in Sunflower by blue-green and chlorophyll-a fluorescence imaging. *Journal of Plant Nutrition*, 34(14), 2114-2126.
- Cao, Q., Miao, Y., Wang, H., Huang, S., Cheng, S., R.Khosla, & Jiang, R. (2013). Non-destructive estimation of rice plant nitrogen status with crop circle multispectral active canopy sensor. *Field Crops Research*, 154, 133-144.

- Chaerle, L., Leinonen, I., Jones, H. G., & Dominique Van Der Straeten. (2007). Monitoring and screening plant populations with combined thermal and chlorophyll fluorescence imaging. *Journal of Experimental Botany*, 58(4), 773-784.
- Chaerle, L., & Van Der Straeten, D. (2000). Imaging techniques and the early detection of plant stress. *Trends in Plant Science*, 5(11), 495-501.
- Comte, I., Colin, F., Whalen, J. K., Grünberger, O., & Caliman, J.-P. (2012). Chapter three - Agricultural Practices in Oil Palm Plantations and Their Impact on Hydrological Changes, Nutrient Fluxes and Water Quality in Indonesia: A Review. In L. S. Donald (Ed.), *Advances in Agronomy*, 116, 71-124: Academic Press.
- Djanaguiraman, M., Vara Prasad, P. V., Murugan, M., Perumal, R., & Reddy, U. K. (2014). Physiological differences among sorghum (*Sorghum bicolor* L. Moench) genotypes under high temperature stress. *Environmental and Experimental Botany*, 100(0), 43-54.
- Dong, H., Kong, X., Li, W., Tang, W., & Zhang, D. (2010). Effects of plant density and nitrogen and potassium fertilization on cotton yield and uptake of major nutrients in two fields with varying fertility. *Field Crops Research*, 119(1), 106-113.
- Fageria, N. K., Baligar, V. C., & Clark, R. B. (2002). Micronutrients in Crop Production. In L. S. Donald (Ed.), *Advances in Agronomy*, 77, 185-268: Academic Press.
- Feng, W., He, L., Zhang, H.-Y., Guo, B.-B., Zhu, Y.-J., Wang, C.-Y., & Guo, T.-C. (2015). Assessment of plant nitrogen status using chlorophyll fluorescence parameters of the upper leaves in winter wheat. *European Journal of Agronomy*, 64, 78-87.
- Feng, W., Yao, X., Zhu, Y., Tian, Y. C., & Cao, W. X. (2008). Monitoring leaf nitrogen status with hyperspectral reflectance in wheat. *European Journal of Agronomy*, 28(3), 394-404.
- Fernández-Escobar, R., Sánchez-Zamora, M. F., & Molina-Soria, C. (2014). The amount of nitrogen applied and nutritional status of olive plants affect nitrogen uptake efficiency. *Scientia Horticulturae*, 167, 1-4.
- Franck, F., Juneau, P., & Popovic, R. (2002). Resolution of the Photosystem I and Photosystem II contributions to chlorophyll fluorescence of intact leaves at room temperature. *Biochimica et Biophysica Acta*, 1556, 239-246.
- Gislerød, H. R., & Mortensen, L. M. (1990). Relative Humidity and Nutrient Concentration Affect Nutrient Uptake and Growth of *Begonia x hiemalis*. *Horticultural Sciences*, 25(5), 524-526.

- Gislerød, H. R., Selmer-Olsen, A. R., & Mortensen, L. M. (1987). The effect of air humidity on nutrient uptake of some greenhouse plants. *Plant and Soil*, 102, 193-196.
- Hanafi, M. M., Selamat, M. M., Husni, M. H. A., & Adzemi, M. A. (2009). Dry matter and nutrient partitioning of selected pineapple cultivars grown on mineral and tropical peat soils. *Communications in Soil Science and Plant Analysis*, 40(21-22), 3263-3280.
- Hawkesford, M., Horst, W., Kichey, T., Lambers, H., Schjoerring, J., Møller, I. S., & White, P. (2012). Chapter 6-Functions of Macronutrients. In P. Marschner (Ed.), *Marschner's Mineral Nutrition of Higher Plants*. Third ed., 135-189. San Diego: Academic Press.
- Heisel, F., Sowinska, M., Miehe, J. A., Lang, M., & Lichtenthaler, H. K. (1996). Detection of Nutrient Deficiencies of Maize by Laser Induced Fluorescence Imaging. *Journal of Plant Physiology*, 148(5), 622-631.
- Hideg, E., Juhasz, M., Bornman, J., & Asada, K. (2002). The distribution and possible origin of blue-green fluorescence in control and stressed barley leaves. *Photochemical and Photobiological Sciences*, 1, 934-941.
- Hiradate, S., Ma, J. F., & Matsumoto, H. (2007). Strategies of Plants to Adapt to Mineral Stresses in Problem Soils. In L. S. Donald (Ed.), *Advances in Agronomy* 96, 65-132: Academic Press.
- Hochmal, A. K., Schulze, S., Trompelt, K., & Hippler, M. (2015). Calcium-dependent regulation of photosynthesis. *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, 1847(9), 993-1003.
- Kim, Y. S., Glenn, D. M., Park, J., Ngugi, H. K., & Lehman, B. L. (2011). Hyperspectral image analysis for water stress detection of apple trees. *Computers and Electronics in Agriculture*, 77, 155-160.
- Kitchen, N. R., Sudduth, K. A., Drummond, S. T., Scharf, P. C., Palm, H. L., Roberts, D. F., & Vories, E. D. (2010). Ground-based canopy reflectance sensing for variable-rate nitrogen corn fertilization. *Journal of Agronomy*, 102, 71-84.
- Krumov, A., Nikolova, A., Vassilev, V., & Vassilev, N. (2008). Assessment of plant vitality detection through fluorescence and reflectance imagery. *Advances in Space Research*, 41, 1870-1875.
- Lee, W., Alchanatis, V., Yang, C., Hirafuji, H., Moshou, D., & Li, C. (2010). Sensing technologies for precision specialty crop production. *Computers and Electronics in Agriculture*, 74, 2-33.
- Lemaire, G., & Gastal, F. (2009). Chapter 8 - Quantifying Crop Responses to Nitrogen Deficiency and Avenues to Improve Nitrogen Use Efficiency. In V. Sadras & D. Calderini (Eds.), *Crop Physiology*. 171-211. San Diego: Academic Press.

- Li, S., Wang, Z., Malhi, S. S., Li, S., Gao, Y., & Tian, X. (2009). Chapter 7 nutrient and water management effects on crop production, and nutrient and water use efficiency in dryland areas of China. *Advances in Agronomy*, 223-265.
- Li, F., Miao, Y. X., Feng, G. H., Yuan, F., Yue, S. C., Gao, X. W., Liu, Y. Q., Liu, B., Ustin, S. L. & Chen, X. P. (2014). Improving estimation of summer maize nitrogen status with red edge-based spectral vegetation indices. *Field Crops Research*, 157, 111-123.
- Linguist, B. A., Adviento-Borbe, M. A., Pittelkow, C. M., Kessel, C. V., & Groenigen, K. J. V. (2012). Fertilizer management practices and greenhouse gas emissions from rice systems: A quantitative review and analysis *Field Crops Research*, 135, 10-21.
- Mafongoya, P. L., Bationo, A., Kihara, J., & Waswa, B. S. (2006). Appropriate technologies to replenish soil fertility in Southern Africa. *Nutrient Cycling in Agroecosystem*, 76, 137-151.
- Margesin, R., & Schinner, F. (2005). *Manual for Soil Analysis-Monitoring and Assessing Soil Bioremediation*. Berlin Heidelberg, Germany: Springer-Verlag.
- Marschner, P., & Rengel, Z. (2012). Chapter 12 - Nutrient Availability in Soils. In P. Marschner (Ed.), *Marschner's Mineral Nutrition of Higher Plants (Third Edition)* (pp. 315-330). San Diego: Academic Press.
- Mathur, S., Agrawal, D., & Jajoo, A. (2014). Photosynthesis: Response to high temperature stress. *Journal of Photochemistry and Photobiology B: Biology*, 137(0), 116-126.
- Maxwell, K., & Johnson, G. N. (2000). Chlorophyll fluorescence- α practical guide. *Journal of Experimental Botany*, 51, 659-668.
- Mehlich, A. (1953). *Determination of P, K, Na, Ca, Mg and NH₄ NC*, USA: Soil Test Division Mimeo, North Carolina Department.
- Moshou, D., Bravo, C., Oberti, R., West, J. S., Ramon, H., Vougioukas, S., & Bochtis, D. (2011). Intelligent multi-sensor system for the detection and treatment of fungal diseases in arable crops. *Biosystems Engineering*, 108, 311-321.
- Mujib, M., & Sapii, A. T. (2010). *Manual Teknologi Penanaman Nanas*. Serdang: Institut Penyelidikan dan Kemajuan Pertanian Malaysia (MARDI).
- Mulla, D. J. (2013). Twenty five years of remote sensing in precision agriculture: Key advances and remaining knowledge gaps. *Biosystems Engineering*, 114, 358-371.
- Muñoz-Huerta, R. F., Guevara-Gonzalez, R. G., Contreras-Medina, L. M., Torres-Pacheco, I., Prado-Olivarez, J., & Ocampo-Velazquez, R. V.

- (2013). A review of methods for sensing the Nitrogen status in plants: Advantages, disadvantages and recent advances *Sensors*, 13, 10823-10843.
- Nagai, T., & Makino, A. (2009). Differences between rice and wheat in temperature responses of photosynthesis and plant growth. *Plant And Cell Physiology*, 50, 744-755.
- Navarro-Cerillo, R. M., Trujillo, J., Sánchez de la Orden, M., & Hernández-Clemente, R. (2014). Hyperspectral and multispectral satellite sensors for mapping chlorophyll content in a Mediterranean *Pinus sylvestris* L. Plantation. *International Journal of Applied Earth Observation And Geoinformation*, 26, 88-96.
- Okimoto, M.C. 1948). Anatomy and histology of the pineapple inflorescence and fruit. *Botanical Gazette (Chicago)* 110:217-231.
- Ondimu, S. & Murase, H. (2008). Water stress detection in Sunagoke moss (*Rhacomitrium canescens*) using combined thermal infrared and visible light imaging techniques. *Biosystems Engineering*, 100, 4-13.
- Osório, J., Osório, M. L., Correia, P. J., de Varennes, A., & Pestana, M. (2014). Chlorophyll fluorescence imaging as a tool to understand the impact of iron deficiency and resupply on photosynthetic performance of strawberry plants. *Scientia Horticulturae*, 165, 148-155.
- Pacumbaba, J. R. O., & Beyl, C. A. (2011). Changes in hyperspectral reflectance signatures of lettuce leaves in response to macronutrient deficiencies *Advances in Space Research*, 48, 32-42.
- Pandey, R., Zinta, G., AbdElgawad, H., Ahmad, A., Jain, V., & Janssens, I. A. (2015). Physiological and molecular alterations in plants exposed to high [CO₂] under phosphorus stress. *Biotechnology Advances*, 33(3–4), 303-316.
- Pimstein, A., Karnieli, A., Bansal, S. K., & Bonfil, D. J. (2011). Exploring remotely sensed technologies for monitoring wheat potassium & phosphorus using field spectroscopy. *Field Crops Research*, 121(1), 125-135.
- Preston MD, Smemo KA, McLaughlin JW, Basiliko N. (2012). Peatland Microbial Communities and Decomposition Processes in the James Bay Lowlands, Canada. *Frontiers in Microbiology*, 70(3).
- Ranjan, R., Chopra, U. K., Sahoo, R. N., Kumar, A., & Pradhan, S. S. (2012). Assessment of plant nitrogen stress in wheat (*Triticum aestivum* L.) through hyperspectral indices. *International Journal of Remote Sensing*, 33(20), 6342-6360.
- Rose, T. J., Rengel, Z., Ma, Q., & Bowden, J. W. (2008). Hydraulic lift by canola plants aids P and K uptake from dry topsoil. *Australian Journal of Agricultural Research*, 59(1), 38-45.

- Ryan, J., Rashid, A., Torrent, J., Yau, S. K., Ibrikci, H., Sommer, R., & Erenoglu, E. B. (2013). Chapter One - Micronutrient Constraints to Crop Production in the Middle East–West Asia Region: Significance, Research, and Management. In L. S. Donald (Ed.), *Advances in Agronomy* 122, 1-84: Academic Press.
- SAS, Inc. (1985). *SAS Introductory Guide for Personal Computers, Version 6 ed.* Cary, NC: SAS Institute Inc.
- Schmidt, J. P., Dellinger, A. E., & Beegle, D. B. (2009). . Nitrogen recommendations for corn: an on-the-go sensor compared with current recommendation methods. *Journal of Agronomy*, 101, 916-924.
- Scotford, I. M., & Miller, P. C. H. (2004). Combination of spectral reflectance & ultrasonic sensing to monitor the growth of winter wheat. *Biosystems Engineering*, 87(1), 27-38.
- Shiratsuchi, L., Ferguson, R., Shanahan, J., Adamchuk, V., Rundquist, D., Marx, D., & Slater, G. (2011). Water and nitrogen effects on active canopy sensor vegetation indices. *Journal of Agronomy*, 103, 1815-1826.
- Thomason, W. E., Phillips, S. B., & Raymond, F. D. (2007). Defining useful limits for spectral reflectance measures in corn. *Journal of Plant Nutrition*, 30(8), 1263-1277.
- Tilling, S. K., O'Leary, G. J., Ferwerda, J. G., Jones, S. D., Fitzgerald, G. J., & Rodriguez, D. (2007). Remote sensing of nitrogen and water stress in wheat. *Field Crops Research*, 104, 77-85.
- Vadivambal, R., & Jayas, D. S. (2011). Application of thermal imaging in agriculture and food industry-A review. *Food Bioprocess Technology*, 4, 186-199. doi: 10.1007/s11947-010-0333-5
- Wahid, A., Gelani, S., Ashraf, M., & Foolad, M. R. (2007). Heat Tolerance in Plants: An Overview. *Environmental and Experimental Botany*, 61, 199-223.
- Walsh, L. M., Dinaeur, R. C., & Wilson, B. B. (1971). *Instrumental Methods for Analysis of Soils and Plant Tissue*. Madison, Wisconsin USA: Soil Science Society of America.
- Wang, J., Xiong, Z., & Yan, X. (2011). Fertilizer-induced emission factors and background emissions of N₂O from vegetable fields in China. *Atmospheric Environment*, 45, 6923-6929.
- Xin, C., Qing-wei, Y., Jia-lin, S., Shuang, X., Fu-chun, X., & Ya-jun, C. (2014). Research Progress on Nitrogen Use and Plant Growth. *Journal of Northeast Agricultural University (English Edition)*, 21(2), 68-74.
- Yang, F., Li, J., Gan, X., Qian, Y., Wu, X., & Yang, Q. (2010). Assessing nutritional status of *Festuca arundinacea* by monitoring photosynthetic

- pigments from hyperspectral data. *Computers and Electronics in Agriculture*, 70(1), 52-59.
- Zazali, M. (2014, August 12). Pineapple board plans big exports to China, *The Star*. Retrieved from www.thestar.com.my.
- Zhang, J.H., Wang, K., Bailey, J. S., & Wang, R.C. (2006). Predicting Nitrogen Status of Rice Using Multispectral Data at Canopy Scale. *Pedosphere*, 16(1), 108-117.
- Zhang, M., Qin, Z., Liu, X., & Ustin, S. L. (2003). Detection of stress in tomatoes induced by late blight disease in California, USA, using hyperspectral remote sensing *International Journal of Applied Earth Observation and Geoinformation*, 4, 295-310.
- Zhang, X., Liu, F., He, Y., & Gong, X. (2013). Detecting macronutrients content and distribution in oilseed rape leaves based on hyperspectral imaging. *Biosystems Engineering*, 115(1), 56-65.
- Zhao, R. F., Chen, X. P., Zhang, F. S., Zhang, H. I., Schroder, J., & Römheld, V. (2006). Fertilization and Nitrogen Balance in a Wheat–Maize Rotation System in North China. *Journal of Agronomy*, 98(4), 938-945.
- Zörb, C., Senbayram, M., & Peiter, E. (2014). Potassium in agriculture-Status and perspectives. *Journal of Plant Physiology*, 171(9), 656-669.