



UNIVERSITI PUTRA MALAYSIA

***ESTABLISHMENT OF SHOOT MULTIPLICATION SYSTEM AND
CRYOPRESERVATION OF JACKFRUIT (*Artocarpus heterophyllus*
LAMK) SHOOT TIPS***

OLUWAGADE ADENIKE OKUNADE

FP 2018 32



**ESTABLISHMENT OF SHOOT MULTIPLICATION SYSTEM AND
CRYOPRESERVATION OF JACKFRUIT (*Artocarpus heterophyllus* LAMK)
SHOOT TIPS**

By

OLUWAGADE ADENIKE OKUNADE

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

June 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Doctor of Philosophy

**ESTABLISHMENT OF SHOOT MULTIPLICATION SYSTEM AND
CRYOPRESERVATION OF JACKFRUIT (*Artocarpus heterophyllus* LAMK)
SHOOT TIPS**

By

OLUWAGADE ADENIKE OKUNADE

June 2017

Chairman : Professor Uma Rani a/p Sinniah, PhD
Faculty : Agriculture

Jackfruit produces recalcitrant seeds that are dessication-sensitive and cannot be stored by conventional seed storage. This poses a major problem for conservation of valuable genetic diversity of this plant species. Thus, sustainable conservation of its germplasm is imperative. Cryopreservation is the only viable option for long-term storage of recalcitrant plant species. The aim of this study, therefore, is to develop *in vitro* multiplication system using shoots obtained from seedlings and cryopreservation protocol for *in vitro* shoot tips. The results presented in this thesis ascertain that the physiological and developmental state of explant and suitable *in vitro* culture system coupled with optimization of the various steps involved in cryopreservation protocols are critical factors that influence successful cryopreservation of jackfruit shoot tips. To establish an efficient *in vitro* culture, suitable seeds were selected for seedling establishment. Seeds from top, middle and basal section of the fruit were extracted and germinated. The seeds from the middle and basal sections germinated earlier than those from the top and had 100% germination. *In vitro* culture was established using shoot tips from the germinated seedlings at 4, 6, 8 and 12 weeks with variation in terminal shoot developmental stage viz shoot with unexpanded leaf, shoot with partially expanded leaf and shoot with expanded leaf. The results showed that terminal shoot obtained from 6-week-old seedlings in MS medium supplemented with 3.0 mg/L benzylaminopurine performed best compared with 4, 8 and 12-week-old seedlings. Shoots with partially expanded leaf and with expanded leaf obtained from 6 weeks old seedlings induced 100% new shoots and produced up to seven multiple shoots than shoots with unexpanded leaf. The shoot tips from the multiple shoot produced *in vitro* were used to establish the cryo-protocol. Successful cryopreservation was achieved by using 6-week *in vitro* cultured shoot tips of sizes 1.5 mm - 2.0 mm as the explant. Two cryopreservation protocol viz vitrification and encapsulation-vitrification techniques using plant vitrification solutions: PVS2, PVS3 and PVS4 were carried out. The findings of the study on encapsulation shows that stepwise exposure to PVS2 [30%

(w/v) glycerol 15% (w/v) ethylene glycol, 15% (w/v) DMSO, 0.4 M sucrose] resulted in survival of 82.1% and 23.5% in non-cryopreserved and cryopreserved encapsulated shoot tips. In contrast, survival of 88.9% and 35.2% was obtained with direct exposure to PVS3 [50% (w/v) glycerol, 50% sucrose] while PVS4 [35% (w/v) glycerol, 20% (w/v) ethylene glycol, 0.6 M sucrose] resulted in 77.1% and 37.0% survival respectively. Moreover, 28.7% survival was obtained with PVS2 at 0°C while a higher percentage survival of 35.8% and 37.9% was obtained at 25°C with PVS3 and PVS4 in cryopreserved shoot tips respectively. In non-encapsulated shoot tips, sucrose concentration at 0.5 M exposed for 48 h induced cryotolerance to dehydration. Elevated sucrose concentration (0.7 - 0.9 M) and extended period of exposure (72 - 240 h) was detrimental. The different plant vitrification solution used had varied responses on the shoot tips at different times of exposure. The critical hydration window range 49.7% - 52.4% water content was attained in all the vitrification solutions with survival range of 26.8% - 37.4% at different time of exposure. Histological observation revealed that structural changes occurred in the cell during incubation with plant vitrification solution and freezing. Sucrose preculture and PVS4 treatment for 50 minutes (52.4% WC) was able to preserve some living cells against damage during freezing. This study was successful in establishing an efficient system for *in vitro* shoot tip multiplication and cryopreservation protocol by vitrification using PVS4. Thus, long-term storage of explants of *Artocarpus* species is possible by focusing on the critical factors influencing successful cryopreservation.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PEMBANGUNAN SISTEM PENGGANDAAN PUCUK DAN KRIOAWETAN
HUJUNG PUCUK NANGKA (*Artocarpus heterophyllus* LAMK)**

Oleh

OLUWAGADE ADENIKE OKUNADE

Jun 2017

Pengerusi : Professor Uma Rani a/p Sinniah, PhD

Fakulti : Pertanian

Buah nangka mempunyai biji benih jenis rekalsitran yang sensitif kepada pengeringan dan tidak boleh disimpan menggunakan kaedah penyimpanan biji benih secara konvensional. Ini menyebabkan masalah besar dalam pemuliharaan kepelbagaian genetik yang bernilai bagi spesies tumbuhan ini. Oleh itu, kaedah konservasi janaplasma yang mampan adalah penting. Krioawetan merupakan satu-satunya pilihan berdaya maju bagi penyimpanan jangka panjang bagi spesies tumbuhan jenis rekalsitran. Oleh itu, tujuan kajian ini adalah untuk menghasilkan satu sistem bagi penggandaan pucuk bagi explan yang diperolehi dari anak benih dan mendapatkan protokol bagi krioawetan hujung pucuk in vitro nangka. Keputusan yang dibentangkan dalam tesis ini menunjukkan bahawa keadaan fisiologi dan tahap perkembangan explan dan sistem in vitro yang sesuai digabungkan dengan pengoptimuman pelbagai langkah yang terlibat dalam protokol krioawetan adalah faktor kritikal yang mempengaruhi kejayaan krioawetan nangka. Untuk mewujudkan sistem in vitro yang efisien, biji benih yang sesuai dipilih untuk mendapatkan anak benih. Biji benih dari bahagian atas, tengah dan basal buah telah diekstrak dan dicambahkan. Biji benih dari bahagian tengah dan basal bercambah lebih awal daripada yang dari atas dan mempunyai percambahan 100% Kultur in vitro dihasilkan dengan menggunakan pucuk dari anak benih yang berumur 4, 6, 8 dan 12 minggu selepas percambahan dengan variasi dalam tahap perkembangan pucuk iaitu dengan daun yang belum kembang, daun separa kembang dan daun kembang sepenuhnya. Keputusan menunjukkan bahawa pucuk yang diperolehi daripada anak benih berusia 6 minggu dalam medium MS ditambah dengan benzylaminopurine 3.0 mg / L memberikan respon terbaik dibandingkan dengan anak benih 4, 8 dan 12-minggu. Pucuk dengan daun yang kembang sepenuhnya yang diperolehi daripada anak benih berusia 6 minggu memberikan 100% pucuk baru dengan tujuh pucuk berbanding dengan pucuk dari daun yang tidak berkembang. Krioawetan telah berjaya dicapai dengan menggunakan hujung pucuk in vitro berusia 6-minggu yang bersaiz 1.5 mm - 2.0 mm yang digunakan

sebagai explan. Dua protokol krioawetan iaitu teknik vitrifikasi dan teknik pengkapsulan-vitrifikasi menggunakan larutan vitrifikasi tanaman: PVS2, PVS3 dan PVS4 telah dijalankan. Keputusan kajian pengkapsulan menunjukkan bahawa pendedahan langkah demi langkah terhadap PVS2 [30% (w/v) gliserol 15% (w/v) ethylene glycol, 15% (w/v) DMSO, 0.4 M sukrosa] memberikan kemandirian hujung pucuk yang telah dikapsulkan sebanyak 82.1% dan 23.5% tanpa dan dengan krioawetan. Sebaliknya, kemandirian sebanyak 88.9% dan 35.2% diperoleh dengan pendedahan langsung kepada PVS3 [50% (w/v) gliserol, 50% sukrosa] manakala PVS4 [35% (w/v) gliserol, 20% (w/v) ethylene glycol, 0.6 M sukrosa] masing-masing menghasilkan 77.1% dan 37.0% kemandirian. Selain itu, kemandirian sebanyak 28.7% diperoleh dengan PVS2 pada 0°C manakala peratus kemandirian yang lebih tinggi sebanyak 35.8% dan 37.9% diperolehi pada 25°C dengan PVS3 dan PVS4 masing-masing selepas krioawetan. Kepekatan sukrosa optimum iaitu 0.5 M dengan pendedahan selama 48 jam memberikan toleransi kepada pengeringan dan menyebabkan teknik vitrifikasi berjaya. Peningkatan kepekatan sukrosa kepada 0.7 - 0.9 M dan tempoh pendedahan yang lanjut (72 - 240j) adalah memudaratkan. Jenis larutan vitrifikasi yang berbeza memberikan respon yang berbeza terhadap hujung pucuk mengikut jangkamasa pendedahan. Julat penghidratan kritikal dengan kandungan kelembapan sebanyak 49.7% - 52.4% dapat diperolehi dalam semua jenis larutan vitrifikasi dengan julat kemandirian sebanyak 26.8% - 37.4% pada masa pendedahan yang berbeza. Pemerhatian histologi mendedahkan bahawa terdapat perubahan pada struktur sel sewaktu pendedahan kepada larutan vitrifikasi dan penyejukan. Pendedahan kepada sukrosa dan larutan PVS4 selama 50 min (52.4% WC) membolehkan sel hidup mengelakkan dari kerosakan semasa penyejukan. Kajian ini berjaya menghasilkan sistem yang cekap untuk penggandaan hujung pucuk in vitro dan protokol cryopreservation melalui keadaan vitrification menggunakan PVS4. Oleh itu, penyimpanan jangka panjang spesies *Artocarpus* boleh dilakukan dengan memberi tumpuan kepada faktor kritikal yang mempengaruhi kejayaan krioawetan.

ACKNOWLEDGEMENTS

I want to express my greatest gratitude to the Almighty God, the ever faithful one, who gave me this opportunity and also provide the strength to complete my study. My sincere appreciation goes to my supervisors Associate Professor Uma Rani Sinniah, Associate Professor Parameswari Namasivayam and Professor Paul Lynch for their excellent guidance, contribution, motivation and constant support throughout the entire study. I felt opportune to have work with dedicated and committed researchers as you.

My deepest appreciation goes to my lovely husband, Peter, who stood by me all through this period. His love, calmness, encouragement and contributions make my stay in Malaysia a fulfilling one. I must say he is simply the best. Special thanks to my beloved family members: the Okunades', Onipedes', Ajayis' Faloyes' and Yekeens' for their unconditional support. To my sister, Yemi, for her unflagging love and support throughout and being my haven. You are the best of siblings. A million thanks to my brother and sister in law: Favour and Paul, for their unwavering support for me and my husband throughout this journey.

Also, I would like to thank all the friends, staffs, and technicians of Cryopreservation Laboratory and the Department of Crop Science who made my stay in UPM a home away from home. Many thanks to friends in Serdang and Johor for sharing their knowledge, and ideas during my study. Truly, the future belongs to those who believe in the beauty of their dreams. Thanks a lot to you all.

This thesis was submitted to the senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Uma Rani a/p Sinniah, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Parameswari a/p Namasivayam, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences,
Universiti Putra Malaysia
(Member)

Saleh b. Kadzimin, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Paul Thomas Lynch, PhD

Professor
College of Life and Natural Sciences
University of Derby
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No: _____

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF APPENDICES	xviii
LIST OF ABBREVIATIONS	xix
 CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	3
2.1 Origin and distribution of Jackfruit	3
2.2 Botanical characteristics	3
2.2.1 The Plant and fruit Characteristics	3
2.2.2 The Seeds and Seedling Characteristics	4
2.3 Economic Importance	4
2.4 Conservation of Plant Genetic Resources	5
2.4.1 <i>In Situ</i> Conservation	5
2.4.2 <i>Ex Situ</i> Conservation	6
2.4.2.1 Seed Banks	6
2.4.2.2 Field Banks	6
2.4.2.3 <i>In Vitro</i> Banks	6
2.4.2.4 Cryobank	7
2.5 Cryopreservation	7
2.5.1 Factors Responsible for Successful Cryopreservation	8
2.5.1.1 <i>In Vitro</i> Regeneration System	8
2.5.1.2 Source of Explant	9
2.5.1.3 Type of Explant	10
2.5.1.4 Size of Explant	10
2.5.1.5 Physiological and Developmental State of Explant	11
2.5.1.6 Medium	12
2.5.1.7 Plant Growth Regulator	12
2.5.2 Cryopreservation Methods	13
2.5.2.1 Controlled Cooling Rate	13
2.5.2.2 Vitrification-based Methods	14
2.5.2.3 Vitrification	14
2.5.2.4 Encapsulation Vitrification	15
2.5.3 Conditioning of Tissue for Cryopreservation	15

2.5.3.1	Preculture	16
2.5.3.2	Loading	16
2.5.3.3	Dehydration	17
2.5.3.4	Plant Vitrification Solution	18
2.5.3.5	Freezing and Thawing	19
2.5.3.6	Recovery	20
3	INFLUENCE OF SEED POSITION WITHIN THE FRUIT ON SEEDLING QUALITY AND ASSOCIATED <i>IN VITRO</i> PERFORMANCE OF JACKFRUIT (<i>Artocarpus heterophyllus</i> Lamk) SHOOT TIPS	21
3.1	Introduction	21
3.2	Materials and Methods	22
3.2.1	Plant Material	22
3.2.2	Preparation of Sand for Germination	22
3.2.3	Preparation of Stock Solution for Basal Medium	22
3.2.4	Preparation of Culture Media	22
3.2.5	Preparation of Plant Growth Regulators	23
3.2.6	Morphological Assessment of Seed and Embryo Extracted from the Different Position within the Fruit	23
3.2.7	Seed Selection and Seedling Establishment	23
3.2.8	Determination of Suitable Physiological Age of Shoot Tips For <i>In Vitro</i> Culture	23
3.2.9	Sterilisation, Culture and Culture Conditions	24
3.2.10	Effect of BAP Concentrations on Shoot Multiplication	24
3.2.11	Effect of TDZ Concentrations on Shoot Multiplication	24
3.2.12	Morphological Characteristics of Shoot Developmental Stages of Growth of Jackfruit	24
3.2.12.1	Effect of Different Culture Medium on Stage of Growth of Shoot Tips	26
3.2.12.2	Performance of Shoot tip and Nodal Parts during <i>In Vitro</i> Culture	26
3.2.13	Data Collection and Measurement	27
3.2.14	Experimental Design and Statistical Analysis	28
3.3	Results and Discussion	28
3.3.1	Morphological Characteristics of Seeds and Embryos Extracted From Different Positions within the Fruit	28
3.3.2	Effect of Seed Position within the Fruit on the Germination Potential of Jackfruit	32
3.3.3	Effect of Seed Position within the Fruit, and Age of Seedlings on <i>In Vitro</i> Performance of Shoot Tips.	33
3.3.4	Effect of BAP Concentrations on Shoot tip Induction and Number of Multiple Shoot tip	36
3.3.5	Effect of TDZ Concentrations on Shoot tip Induction and Number of Multiple Shoot	36
3.3.6	Effects of Culture Media on Stage of Growth of Shoot Tips during <i>In Vitro</i> Culture	37
3.3.7	Performance of the Different Explants Cultured <i>In Vitro</i>	39

4	EFFECT OF SUCROSE PRECULTURE AND PLANT VITRIFICATION SOLUTION EXPOSURE ON CRYOPRESERVATION OF JACKFRUIT (<i>Artocarpus heterophyllus</i>) SHOOT TIPS	44
4.1	Introduction	44
4.2	Materials and Methods	45
4.2.1	Plant Material	45
4.2.2	Preparation of Sucrose for Preculture Treatment	45
4.2.3	Preparation of Loading Solution	45
4.2.4	Plant Vitrification Solution Treatment	45
4.2.5	Preparation of Unloading Solution	46
4.2.6	Selection of Suitable Shoot Tips Size for Cryopreservation Study	46
4.2.7	Selection of Suitable Age of Shoot Tips <i>In Vitro</i> Culture Donor Plant for Cryopreservation Study	47
4.2.8	Effect of Sucrose Preculture Treatment	47
4.2.9	Effect of Plant Vitrification Solution Treatment	48
4.2.10	Post-Treatment and Recovery	48
4.2.11	Sample Preparation for Histological Studies	49
4.2.12	Data Collection and Statistical Analysis	49
4.3	Results and Discussions	50
4.3.1	Effect of Size of Shoot Tips on Survival of Cryopreserved Shoot Tips	50
4.3.2	Effect of Age of <i>In vitro</i> Donor Plant on Survival of non-Cryopreserved and Cryopreserved Shoot Tips	52
4.3.3	Effect of Sucrose Concentration And Duration of Exposure on Survival of Cryopreserved and Non-Cryopreserved Shoot Tips	53
4.3.4	Effect of Extended Duration of Preculture in 0.5 M Sucrose on Survival and Regrowth of Shoot Tips	55
4.3.5	Effect of PVS on Water Content of Non-Cryopreserved Shoot Tips	57
4.3.6	Effect of Time of Exposure to Different Plant Vitrification Solution (PVS) on Survival of Non-Cryopreserved Shoot Tips of Jackfruit	58
4.3.7	Effect of Different PVS on Regrowth of Non-Cryopreserved Shoot Tips	59
4.3.8	Effect of Different PVS on Survival and Regrowth of Cryopreserved Shoot Tips	60
4.3.9	Comparison of Time of Shoot Regrowth of Non-Cryopreserved and Cryopreserved Shoot Tips	67
4.4	Conclusion	68
5	INFLUENCE OF APPLICATION METHOD AND TEMPERATURE CONTROL ON JACKFRUIT SHOOT TIPS CRYOPRESERVATION USING ENCAPSULATION VITRIFICATION	69
5.1	Introduction	69

5.2	Materials and Method	70
5.2.1	Plant Material	70
5.2.2	Preparation of Encapsulation Gel	70
5.2.3	Preparation of Sucrose Precultured and Plant Vitrification Solution	70
5.2.4	Encapsulation of Shoot Tips for Cryopreservation Study	70
5.2.5	Effect of Direct and Stepwise Method of Application of Plant Vitrification Solutions on Encapsulated Shoot Tips	71
5.2.6	Determination of Suitable Exposure Temperature of Plant Vitrification Solutions on Encapsulated Shoot Tips	72
5.2.7	Post-Treatment and Recovery	72
5.2.8	Sample Preparation for Histological Studies	72
5.2.9	Statistical Analysis	73
5.3	Results and Discussion	73
5.3.1	Effect of Time of Exposure of Beads to CaCl_2 for Complete Polymerisation on Survival and Regrowth of Non-cryopreserved Encapsulated Shoot Tips	73
5.3.2	Effect of Method of Application of PVS2 on Survival of Encapsulated Cryopreserved and Non-cryopreserved Shoot Tips	75
5.3.3	Effect of Temperature Treatment of PVS2 on Survival of Encapsulated Cryopreserved and Non-Cryopreserved Shoot Tips	76
5.3.4	Effect of Application Method of PVS3 on Survival of Encapsulated Cryopreserved and Non-cryopreserved Shoot Tips	77
5.3.5	Effect of Temperature during Treatment with PVS3 on Survival of Encapsulated Cryopreserved and Non-Cryopreserved Shoot Tips	79
5.3.6	Effect of Application Method of PVS4 on Survival of Encapsulated Cryopreserved and Non-Cryopreserved Shoot Tips	80
5.3.7	Effect of Temperature during Treatment with PVS4 on Survival of Encapsulated Cryopreserved and Non-Cryopreserved Shoot Tips	81
5.4	Conclusions	84
6	SUMMARY, GENERAL CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCH	85
	REFERENCES	88
	APPENDICES	104
	BIODATA OF STUDENT	130
	LIST OF PUBLICATIONS	131

LIST OF TABLES

Table	Page
3.1 Data collection and measurement for the experiment	27
3.2 Morphological characteristics of seeds and embryo within the seed obtained from different position within the fruit	29
3.3 Effect of seed position in the fruit on germination potential of shoot tips	32
3.4 Effect of seeds position within the fruit, and age of seedlings on performance of <i>in vitro</i> culture shoots tips in MS media supplemented with 3 mg/L BAP	34
3.5 Effect of concentrations of BAP on shoot tip induction and number of multiple shoots	36
3.6 Effect of concentrations of TDZ on percentage shoot induction and number of multiple shoots	37
3.7 Effect of media composition supplemented with 3.0 mg L ⁻¹ BAP on <i>in vitro</i> performance of the different stage of growth of jackfruit shoot tips	37
3.8 Performance of the different explants cultured <i>in vitro</i>	40
4.1 Effect of size of shoot tips on survival of non-cryopreserved shoot tips	50
4.2 Effect of age of <i>in vitro</i> donor on survival of cryopreserved shoot tips	52
4.3 Effect of sucrose preculture on survival of cryopreserved and non-cryopreserved shoot tips of jackfruit	54
4.4 Effect of extended preculture duration 24-240 h (1-10 days) in 0.5 M sucrose on water content (WC) and survival of cryopreserved and non-cryopreserved shoot tips of jackfruit	56
4.5 Effect of time of exposure to different plant vitrification solution (PVS) on survival of non-cryopreserved shoot tips of jackfruit	59
4.6 Effect of time of exposure to different plant vitrification solution (PVS) on regrowth of non-cryopreserved shoot tips of jackfruit	60
4.7 Effect of time of exposure to different plant vitrification solution (PVS) on survival and regrowth of cryopreserved shoot tips of jackfruit	61
4.8 Effect of time of exposure to different plant vitrification solution (PVS) on the time of shoot regrowth of non-cryopreserved and cryopreserved shoot tips of jackfruit	67
5.1 Method of application of the different PVSs	71
5.2 Effect of time of exposure of beads to CaCl ₂ for complete polymerisation on encapsulated non-cryopreserved shoot tips	73

5.3	Effect of method of application of PVS2 on survival of encapsulated cryopreserved and non-cryopreserved shoot tips	75
5.4	Effect of temperature during treatment with PVS2 on survival of encapsulated cryopreserved and non-cryopreserved shoot tips	77
5.5	Effect of application method of PVS3 on survival of encapsulated cryopreserved and non-cryopreserved shoot tips	78
5.6	Effect of temperature during treatment with PVS3 on survival of encapsulated cryopreserved and non-cryopreserved shoot tips	79
5.7	Effect of application method of PVS4 on survival of encapsulated cryopreserved and non-cryopreserved shoot tips	81
5.8	Effect of temperature during treatment with PVS4 on survival of encapsulated cryopreserved and non-cryopreserved shoot tips	82
5.9	Comparison of Time of Regrowth of Non-Cryopreserved and Cryopreserved Shoot Tips	83

LIST OF FIGURES

Figure	Page
3.1 Shoot tip developmental stages of jackfruit showing (a) shoot tip with unexpanded leaf (b) shoot tip with partially expanded leaf (c) shoot tip with expanded leaf of 6-week old seedlings. <i>Bar</i> represents 0.5 cm	25
3.2 Excised shoot tips from 6-week old seedling at different developmental stage of growth (a) shoot tip with unexpanded leaf (b) shoot tip with partially expanded leaf (c) shoot tip with expanded leaf. <i>Bar</i> rep. 0.5 cm	25
3.3 Jackfruit seedling showing (a) Shoot tip (b) upper node and (c) lower node. <i>Bar</i> represents 2 cm	27
3.4 Seed extracted from (a) Middle (b) Basal and (c) Top positions of fruit. <i>Bar</i> represent 1 cm	30
3.5 Embryos within the seed from (a) Top (b) Middle and (c) Basal part. <i>Bar</i> represent 5 mm	31
3.6 Embryos extracted from seed from (a) Top (b) Middle and (c) Basal part	31
3.7 Multiple shoot production from shoot tips obtained from (a) age 6 weeks old (b) 6 weeks old (side view) (c) 8 weeks old (d) 12 weeks old and (e) 4 weeks old seedling of jackfruit. <i>Bar</i> represent 0.5 cm	35
3.8 New shoot tip induction from (a) node (b) apical shoot of 6 weeks old seedling. Arrow shows the new shoot tip induction. <i>Bar</i> represents 0.5 cm	41
3.9 Effect of subculture cycle of explants on multiple shoot proliferation. <i>Bar</i> represent SE	42
4.1 Histological section of shoot tip size 1.5 mm. Arrow showing living cells in the apical dome (AD) and in the leaf primordia (LP)	51
4.2 Histological section showing non-cryopreserved shoot tips precultured in 0.5 M sucrose for 48 h. Arrow showing intact living cells at the AD (apical dome) and LPs (leaf primordia)	54
4.3 Effect of time of exposure of different PVSs on water content of non-cryopreserved jackfruit shoot tips. Percentage water content was measured on fresh weight basis. <i>Bar</i> represent standard deviation	58
4.4 Regrowth of non-cryopreserved shoot tips exposed to PVS4 for 50 minutes at 4 week post treatment. <i>Bar</i> represents 0.5 mm	60
4.5 Recovery of cryopreserved shoot tips in MS medium supplemented with 3 mg/L BAP (a) Shoot tips treated with PVS survival after frozen (7 days post-treatment) and (b) Shoot tips regrowth 6 weeks post treatment. <i>Bars</i> represent 0.5 mm	62

4.6	Comparative histology of non-treated freshly excised jackfruit shoot tips (a) Living cells (LC) in the apical dome (AD) and leaf primordia (LP) of freshly excised shoot tips without exposure to LN. (b) Pattern of dead cells (DC) in freshly excised shoot tips exposed to LN and recovered after 24 h post-treatment. Dead cells (DC) showed that heavily condensed nucleus and weakly stained nucleus both at the apical dome (AD) and leaf primordia (LP)	64
4.7	Histology of shoot tips treated with PVS4 for 50 minutes without exposure to LN (a) Living cells (LC) in the apical dome and young leaf primordia (YLP) shoot tips treated with PVS4 for 50 minutes. (b) Dead cells (DC) in the older leaf primordia (OLP) after exposure to PVS4 for 50 minutes	65
4.8	Histology of shoot tips treated with PVS4 for 50 minutes and exposed to LN (a) frozen shoot tips recovery after 24 h post-treatment. (b) Longitudinal section of surviving and dead cells at the apical dome (AD) of frozen jackfruit shoot tips (detail of photo) after 24 h post-treatment	66
5.1	Shoot tips encapsulated with alginate beads. <i>Bar</i> represent 0.5 mm	74
5.2	Survival of non-cryopreserved of encapsulated shoot tips at 7 days post culture. <i>Bar</i> represents 0.5 mm	76
5.3	Regrowth in non-cryopreserved encapsulated shoot tips after direct exposure to PVS3 for 55 minutes. <i>Bar</i> represents 0.5 mm	79
5.4	Encapsulated Shoot tip regrowth at 6 weeks post-treatment	80
5.5	Encapsulated shoot tips in Alginate beads showing (a) preserved cell in the AD and LP (b) Plasmolysed cells at the cortex after PVS treatment	83

LIST OF APPENDICES

Appendix		Page
A	Table of chemical composition	104
B	Table of results	107
C	Table of results	110
D	Table of results	115
E	Table of results	120
F	Protocol diagrams	127

LIST OF ABBREVIATIONS

μM	Micromolar
ANOVA	Analysis of Variance
BAP	N ⁶ -benzylaminopurine
CaCl ₂	Calcium chloride
DMRT	Duncan's Multiple Range Test
DMSO	Dimethylsulfoxide
g.g-1	Gram per gram
g/L	Gram per litre
LN	Liquid Nitrogen
LP	Leaf Primordia
M	Molar
ml	Milliliter
mM	Milli molar
NaOCl	Sodium Hypochlorite
PGR	Plant Growth Regulator
PVS2	Plant Vitrification Solution 2
PVS3	Plant Vitrification Solution 3
PVS4	Plant Vitrification Solution 4
TDZ	N-phenyl-N-1, 2, 3-thiadazol-5-yl urea
WC	Water Content
w/v	Weight per volume

CHAPTER 1

INTRODUCTION

Jackfruit (*Artocarpus heterophyllus* Lamk) is a monoecious tree that produces the largest tree-borne fruits in the world; yielding an average of 200 fruits per tree per year. The fruit can grow up to 50 kg in weight and 90 cm in length with an average of 250 seeds in a fruit. Jackfruit is widely cultivated in many countries in Asia and also found in some parts of Caribbean, Africa, America, and Australia (Sidhu, 2012). In South East Asia, jackfruit is cultivated due to its use in the food industry particularly as dessert owing to its rich source of phenolic compounds, flavonoid and various antioxidants (Swami et al., 2012).

Conservation of jackfruit valuable genetic diversity is vital not only to ensure food security but also to support breeding programmes (Haq, 2006). Jackfruits are mainly conserved in the field gene banks because the seeds are highly hydrated and sensitive to even slight drying, thus, cannot be stored under conventional conditions for orthodox seeds. The cost of maintenance of field bank is high and field accessions are vulnerable to adverse weather conditions, pests, and diseases. Currently, due to rapid land development and deforestation, genetic resources of jackfruit are being lost. Therefore, complementary conservation strategies via cryopreservation for long-term storage is of utmost importance (Mal et al., 2011).

For effective cryopreservation, it is important to establish a suitable system for the regeneration of the explant to be cryopreserved (Reed et al., 2013). Shoot tip is a suitable explant for cryopreservation of recalcitrant plant species (Ibrahim & Normah, 2013). *In vitro* culture is valuable in providing initial explants or stock cultures and the recovery of cryopreserved samples in appropriate growing media (Engelmann & Dussert, 2013; Engelmann, 2011). The success or failure of cryopreserved material is partly dependent on the appropriate *in vitro* culture method. A major step to successful cryopreservation is removal of free water from the tissue which is difficult to achieve in recalcitrant plant species like jackfruit. The sensitivity and intolerance to desiccation and freezing can be controlled with plant vitrification solutions. Vitrification-based method such as vitrification and encapsulation-vitrification is, thus, effective for recalcitrant plant species (Engelmann & Dussert, 2013).

Presently, there is limited knowledge on differential responses of tissues to the different plant vitrification solutions. Fundamental knowledge is required of the composition of the vitrification solution, time of exposure, method of application and prevailing temperature during vitrification process in order to establish a successful cryopreservation protocols. Furthermore, low levels of post-storage survival and regrowth still remains a challenge in recalcitrant plant species regardless of the protocol employed (Normah et al., 2012). Therefore, basic understanding relating to

the structural changes induced in cells during cryopreservation steps are required to improve responses of recalcitrant explants to cryopreservation.

Therefore the objectives of this study are:

1. To investigate the variation among seeds within the fruit in order to establish vigorous seedlings as source of explant for *in vitro* culture of jackfruit shoot tips.
2. To optimize an effective *in vitro* multiplication system for successful cryopreservation.
3. To establish a suitable protocol for cryopreservation of jackfruit shoot tips using vitrification and encapsulation-vitrification techniques.

REFERENCES

- Acedo, A. L. (1992). *Jackfruit biology, production, use and Philippine research*. USA: Multipurpose Trees Species Network Series.
- Ahmed, M. R., Anis, M., & Al-Etta, H. A. (2015). Encapsulation technology for short-term storage and germplasm exchange of *Vitex trifolia* L. *Rendiconti Lincei*, 26(2): 133-139.
- Albarelo, N., Simões, C., GonCALves Rosas, P. F., De Castro, T. C., Gianfaldoni, M. G., Callado, C. H., & Flores, H. (2006). In vitro propagation of *Cleome spinose* (Capparaceae) using explants from nursery-grown seedlings and axenic plants. *In Vitro Cellular and Developmental Biology-Plant*, 42(6): 601-606.
- Amin, M. N., & Jaiswal, V. S. (1987). Rapid clonal propagation of guava through in vitro shoot proliferation on nodal explants of mature trees. *Plant Cell, Tissue and Organ Culture*, 9(3): 235-243.
- Amin, M. N., & Jaiswal, V. S. (1993). In vitro response of apical bud explants from mature trees of jackfruit (*Artocarpus heterophyllus*). *Plant Cell, Tissue and Organ Culture*, 33(1): 59-65.
- Anderson, W. C. (1980). Tissue culture propagation of red and black raspberries, *Rubus idaeus* and *R. occidentalis*. *Acta Horticulturae*, 112: 124-132.
- Ashrafuzzaman, M., Kar, S., Khanam, D., & Prodhan, S. H. (2012). In vitro regeneration and multiplication of jackfruit (*Artocarpus heterophyllus* L.). *Research Journal of Biology*, 2(2): 59-65.
- Azad, A., Jones, J., & Haq, N. (2007). Assessing morphological and isozyme variation of jackfruit (*Artocarpus heterophyllus* Lam.) in Bangladesh. *Agroforestry Systems*, 71(2): 109-125.
- Bachiri, Y., Song, G., Plessis, P., Shoar-Ghaffari, A., Rekab, T., & Morisset, C. (2000). Routine cryopreservation of kiwifruit (*Actinidia* spp) germplasm by encapsulation-dehydration: importance of plant growth regulators. *CryoLetters*, 22(1): 61-74.
- Bairu, M. W., Aremu, A. O., & Van Staden, J. (2011). Somaclonal variation in plants: causes and detection methods. *Plant Growth Regulation*, 63(2): 147-173.
- Baliga, M. S., Shivashankara, A. R., Haniadka, R., Dsouza, J., & Bhat, H. P. (2011). Phytochemistry, nutritional and pharmacological properties of *Artocarpus heterophyllus* Lam (jackfruit): A review. *Food Research International*, 44(7): 1800-1811.

- Barraco, G., Sylvestre, I., Collin, M., Escoute, J., Lartaud, M., Verdeil, J. L., & Engelmann, F. (2014). Histocytological analysis of yam (*Dioscorea alata*) shoot tips cryopreserved by encapsulation–dehydration. *Protoplasma*, 251(1): 177-189.
- Bekheet, S. A. (2011). In vitro conservation of date palm germplasm. In M. J. Shri, M. A. Jameel & V. J. Dennis (Eds.), *Date Palm Biotechnology* (pp. 337-360). New York: Springer.
- Benelli, C., De Carlo, A., & Engelmann, F. (2013). Recent advances in the cryopreservation of shoot-derived germplasm of economically important fruit trees of *Actinidia*, *Diospyros*, *Malus*, *Olea*, *Prunus*, *Pyrus* and *Vitis*. *Biotechnology advances*, 31(2): 175-185.
- Benson, E. E. (2008). Cryopreservation theory. In B.M. Reed (Ed.), *Plant Cryopreservation: A Practical Guide* (pp. 15-32). New York: Springer.
- Berjak, P., Farrant, J. M., & Pammenter, N. (1989). The basis of recalcitrant seed behaviour In: *Recent advances in the development and germination of seeds* (pp. 89-108). New York: Springer.
- Berjak, P., & Pammenter, N. (2008). From Avicennia to Zizania: seed recalcitrance in perspective. *Annals of Botany*, 101(2): 213-228.
- Berjak, P., Pammenter, N., & Wesley-Smith, J. (2012). The effects of various parameters during processing for cryopreservation on the ultrastructure and viability of recalcitrant zygotic embryos of *Amaryllis belladonna*. *Protoplasma*, 249(1): 155-169.
- Bhojwani, S. S., & Dantu, P. K. (2013). *Plant Tissue Culture: An Introductory Text*. New York: Springer.
- Bonfil, C. (1998). The effects of seed size, cotyledon reserves, and herbivory on seedling survival and growth in *Quercus rugosa* and *Q. laurina* (Fagaceae). *American Journal of Botany*, 85: 79-87.
- Bramhanapalli, M., Thogatabalija, L., & Gudipalli, P. (2016). Efficient in vitro plant regeneration from seedling-derived explants and genetic stability analysis of regenerated plants of *Simarouba glauca* DC. by RAPD and ISSR markers. *In Vitro Cellular & Developmental Biology-Plant*, 53(1): 50-63.
- Bunn, E., Turner, S., Panaia, M., & Dixon, K. W. (2007). The contribution of in vitro technology and cryogenic storage to conservation of indigenous plants. *Australian Journal of Botany*, 55(3): 345-355.
- CBD (2007). Convention on Biological Diversity (CBD) international biodiversity day. Retrieved from <https://www.cbdint/doc/bioday/ibd-booklet-en.pdf>.

- Chandel, K., Chaudhury, R., Radhamani, J., & Malik, S. (1995). Desiccation and freezing sensitivity in recalcitrant seeds of tea, cocoa and jackfruit. *Annals of Botany*, 76(5): 443-450.
- Chee, R., & Pool, R. (1987). Improved inorganic media constituents for in vitro shoot multiplication of *Vitis*. *Scientia Horticulturae*, 32(2): 85-95.
- Chen, Y., & Wang, J.-H. (2003). Cryopreservation of carrot (*Daucus carota* L.) cell suspensions and protoplasts by vitrification. *CryoLetters*, 24(1): 57-64.
- Chin, H. F., Quek, P., & Sinniah, U. R. (2013). Seed Banks for Future Generation. In M. N. Normah & H. F. Chin (Eds.), *Conservation of Tropical Plant Species* (pp 43-63). New York: Springer.
- Chu, C. C. (1975). Establishment of an efficient medium for anther culture of rice through comparative experiments on the nitrogen sources. *Scientia Sinica*, 18: 659-668.
- Chua, S.P., & Normah, M. N. (2011). Effect of preculture, PVS2 and vitamin C on survival of recalcitrant *Nephelium ramboutan*-ake shoot tips after cryopreservation by vitrification. *CryoLetters*, 32: 506–515.
- Coelho, N., González-Benito, M. E., & Romano, A. (2014). Cryopreservation of shoot tips from the endangered endemic species *Tuberaria major*. *Acta Physiologiae Plantarum*, 36(12): 3333-3336.
- Compton, M. E. (1994). Statistical methods suitable for the analysis of plant tissue culture data. *Plant Cell, Tissue and Organ Culture*, 37(3): 217-242.
- Corredoira, E., San-José, M. C., Ballester, A., & Vieitez, A. M. (2004). Cryopreservation of zygotic embryo axes and somatic embryos of European chestnut. *CryoLetters*, 25(1): 33-42.
- Ćosić, T., Motyka, V., Raspor, M., Savić, J., Cingel, A., Vinterhalter, B., & Bohanec, B. (2015). In vitro shoot organogenesis and comparative analysis of endogenous phytohormones in kohlrabi (*Brassica oleracea* var. gongylodes): effects of genotype, explant type and applied cytokinins. *Plant Cell, Tissue and Organ Culture*, 3: 741-760.
- Crowe, J. H., Crowe, L. M., Carpenter, J. F., & Wistrom, C. A. (1987). Stabilization of dry phospholipid bilayers and proteins by sugars. *Biochemical Journal*, 242(1): 1.
- Cruz-Cruz, C., González-Arno, M., & Engelmann, F. (2013). Biotechnology and conservation of plant biodiversity. *Resources* 2: 73–95.
- Day, J. G., Fleck, R. A., & Benson, E. E. (2000). Cryopreservation-recalcitrance in microalgae: novel approaches to identify and avoid cryo-injury. *Journal of*

- Diab, A. A., Khalil, S., & Ismail, R. M. (2011). Regeneration and micropropagation of grapevine (*Vitis vinifera* L.) through shoot tips and axillary buds. *International Journal of Advanced Biotechnology and Research*, 2(4): 484-491.
- Diggle, P., Abrahamson, N., Baker, R., Barnes, M., Koontz, T., Lay, C., & Simpson, H. (2010). Dynamics of maternal and paternal effects on embryo and seed development in wild radish (*Raphanus sativus*). *Annals of Botany*, 106(2): 309-319.
- Driver, J. A., & Kuniyuki, A. H. (1984). In vitro propagation of paradox walnut rootstock. *Horticultural Science*, 19:507-509.
- Duchefa, B. B. V. (2012). Plant cell and tissue culture phytopathology biochemicals. In F. T. M. Kors (Ed.), *Duchefa Catalogue* (pp. 34-52). The Netherlands: Duchefa Catalogue.
- Dulloo, M. E. (2013). Global challenges for agricultural plant biodiversity and international collaboration. In M. N. Normah & H. F. Chin (Eds.), *Conservation of Tropical Plant Species* (pp. 491-509). New York: Springer.
- Dumet, D., Benson, E. E. (2000). The use of physical and biochemical studies to elucidate and reduce cryopreservation-induced damage in hydrated/desiccated plant germplasm. In F. Engelmann, & H. Takagi (Eds.), *Cryopreservation of Tropical Plant Germplasm. Current Research Progress and Applications* (pp. 43-56). Italy: IPGRI.
- Dussert, S., Chabrilange, N., Rocquelin, G., Engelmann, F., Lopez, M., & Hamon, S. (2001). Tolerance of coffee (*Coffea* spp.) seeds to ultra- low temperature exposure in relation to calorimetric properties of tissue water, lipid composition, and cooling procedure. *Acta Physiologia Plantarum*, 112(4): 495-504.
- Dussert, S., & Engelmann, F. (2006). New determinants for tolerance of coffee (*Coffea arabica* L.) seeds to liquid nitrogen exposure. *CryoLetters*, 27(3): 169-178.
- El-Danasouri, I., Selman, H., Orief, Y., Schultze-Mosgau, A., Al-Hasani, S., Elnomrosy, K., & ElSadek, M. (2005). Vitrification versus conventional cryopreservation technique. *Middle East Fertility Society Journal*, 10(3): 205.
- Elevitch, C. (2006). *Species profiles for Pacific Island agroforestry*. Hawaii: Permanent Agriculture Resources.
- El-Hawaz, R., Park, D., Bridges, W. C., & Adelberg, J. (2016). Optimizing in vitro mineral nutrition and plant density increases greenhouse growth of *Curcuma*

- longa* L. during acclimatization. *Plant Cell, Tissue and Organ Culture*, 126(1): 33-42.
- Engelmann, F. (1991). In vitro conservation of tropical plant germplasm—a review. *Euphytica*, 57(3): 227-243.
- Engelmann, F. (1997). In vitro conservation methods. In J. A. Callow, B. V. Ford-Lloyd & H. J. Newbury (Eds.), *Biotechnology and Plant Genetic Resources, Conservation and Use* (pp. 119-160). Wallingford: CAB International.
- Engelmann, F. (2000). Importance of cryopreservation for the conservation of plant genetic resources. In F. Engelmann, & H. Takagi (Eds.), *Cryopreservation of Tropical Plant Germplasm. Current Research Progress and Applications* (pp. 8–20). Italy: IPGRI.
- Engelmann, F. (2004). Plant cryopreservation: progress and prospects. *In Vitro Cellular and Developmental Biology-Plant*, 40(5): 427-433.
- Engelmann, F. (2011). Use of biotechnologies for the conservation of plant biodiversity. *In Vitro Cellular & Developmental Biology-Plant*, 47(1): 5-16.
- Engelmann, F., & Dussert, S. (2013). Cryopreservation. In M. N. Normah & H. F. Chin (Eds.), *Conservation of Tropical Plant Species* (pp. 107-120). New York: Springer.
- ETP (2013). Economic transformation programme in national key economic areas. Retrieved from http://etp.pemandu.gov.my/upload/NKEA_Factsheet_Agriculture.pdf
- Fabre, J., & Dereuddre, J. (1990). Encapsulation-dehydration: a new approach to cryopreservation of *Solanum* shoot-tips. *CryoLetters*, 11: 413-426.
- FAO. (1996). State of the world's plant genetic resources. *Report on the State of the World's Plant Genetic Resources for Food and Agriculture*, Food & Agriculture Organization.
- FAO. (2010). Commission on genetic resources for food. *The Second Report on the State of the World's Plant Genetic Resources for Food and Agriculture*, Food & Agriculture Organization.
- Feng, C. H., Cui, Z. H., Li, B. Q., Chen, L., Ma, Y. L., Zhao, Y. H., & Wang, Q. C. (2013). Duration of sucrose preculture is critical for shoot regrowth of in vitro-grown apple shoot-tips cryopreserved by encapsulation-dehydration. *Plant Cell, Tissue and Organ Culture*, 112(3): 369-378.
- Fuller, B. J. (2004). Cryoprotectants: the essential antifreezes to protect life in the frozen state. *CryoLetters*, 25(6): 375-388.

- Ganeshan, S., Punathil, E., & Sunitha, B. (2010). In vitro conservation of *Artocarpus heterophyllus* Lam. *Tree and Forestry Science and Biotechnology*, 6: 126-129.
- Ganino, T., Silvanini, A., Beghé, D., Benelli, C., Lambardi, M., & Fabbri, A. (2012). Anatomy and osmotic potential of the *Vitis* rootstock shoot tips recalcitrant to cryopreservation. *Biologia Plantarum*, 56(1): 78-82.
- Gantait, S., Sinniah, U. R., Suranthran, P., Palanyandy, S. R., & Subramaniam, S. (2015). Improved cryopreservation of oil palm (*Elaeis guineensis* Jacq.) polyembryoids using droplet vitrification approach and assessment of genetic fidelity. *Protoplasma*, 252(1): 89-101.
- Ge, X., Chu, Z., Lin, Y., & Wang, S. (2006). A tissue culture system for different germplasms of indica rice. *Plant Cell Reports*, 25(5): 392-402.
- George, E. F., & De Klerk, G. J. (2008). The components of plant tissue culture media I: macro-and micro-nutrients. In E. F. George, M. A. Hall & G. J. De Klerk (Eds.), *Plant Propagation by Tissue Culture* (pp. 65-113). The Netherlands: Springer.
- Gilissen, L. J., van Staveren, M. J., Hakkert, J. C., & Smulders, M. J. (1996). Competence for regeneration during tobacco internodal development (Involvement of plant age, cell elongation stage, and degree of polysomaty). *Plant Physiology*, 111(4): 1243-1250.
- Grospietsch, M., Stodulkova, E., & Zamecnik, J. (1999). Effect of osmotic stress on the dehydration tolerance and cryopreservation of *Solanum tuberosum* shoot tip. *CryoLetters*, 20(6): 339-346.
- Hackett, W. P. (1985). Juvenility, maturation, and rejuvenation in woody plants. *Horticultural Reviews*, 7: 109-155.
- Halwani, D. O., Brockbank, K. G., Duman, J. G., & Campbell, L. H. (2014). Recombinant *Dendroides canadensis* antifreeze proteins as potential ingredients in cryopreservation solutions. *Cryobiology*, 68(3): 411-418.
- Hamed, A. M., Ali, E. A. M., & Boshra, E. S. (2007). In vitro propagation of jackfruit (*Artocarpus heterophyllus* L.) *Journal of Applied Science Research*, 3(3): 218-226.
- Hao, Y.-J., You, C.-X., & Deng, X.-X. (2002). Effects of cryopreservation on developmental competency, cytological and molecular stability of citrus callus. *CryoLetters*, 23(1): 27-35.
- Haq, N. (2006). *Jackfruit: Artocarpus heterophyllus crops for the future*. UK: University of Southampton press.

- Harb, E. M., Alhady, M. R., Abd, A., & Elsalam, N. A. A. (2015). In vitro rapid propagation of jackfruit (*Artocarpus heterophyllus* lam.). *American-Eurasian Journal of Agriculture and Environmental Science*, 15(2): 147-153.
- Harding, K. (2004). Genetic integrity of cryopreserved plant cells: a review. *CryoLetters*, 25(1): 3-22.
- Hari, A., Revikumar, K., & Divya, D. (2014). Artocarpus: A review of its phytochemistry and pharmacology. *Journal of Pharmacolgy Research*, 9(1):7.
- Hensleigh, T. E., & Holaway, B. K. (1988). *Agroforestry Species for the Philippines*. Manila: JA Printers.
- Hirai, D., Shirai, K., Shirai, S., & Sakai, A. (1998). Cryopreservation of in vitro-grown meristems of strawberry (*Fragaria* × *ananassa* Duch.) by encapsulation-vitrification. *Euphytica*, 101(1): 109-115.
- Huijser, P., & Schmid, M. (2011). The control of developmental phase transitions in plants. *Development*, 138(19): 4117-4129.
- Ibrahim, S., & Normah, M. (2013). The survival of in vitro shoot tips of *Garcinia mangostana* L. after cryopreservation by vitrification. *Plant Growth Regulation*, 70(3): 237-246.
- Ide, H., Price, W. S., Arata, Y., & Ishikawa, M. (1998). Freezing behaviors in leaf buds of cold-hardy conifers visualized by NMR microscopy. *Tree physiology*, 18(7): 451-458.
- Ismail, N., & Kaur, B. (2013). Consumer Preference for Jackfruit Varieties in Malaysia. *Journal of Agribusiness Marketing*, 6: 37-51.
- Jagtap, U. B., & Bapat, V. A. (2013). Green synthesis of silver nanoparticles using *Artocarpus heterophyllus* Lam. seed extract and its antibacterial activity. *Industrial Crops and Products*, 46: 132-137.
- Jarvis, D. I., Hodgkin, T., Sthapit, B. R., Fadda, C., & Lopez-Noriega, I. (2011). An heuristic framework for identifying multiple ways of supporting the conservation and use of traditional crop varieties within the agricultural production system. *Critical Reviews in Plant Sciences*, 30(2): 125-176.
- Jitsuyama, Y., Suzuki, T., Harada, T., & Fujikawa, S. (2002). Sucrose incubation increases freezing tolerance of asparagus (*Asparagus officinalis* L.) embryogenic cell suspensions. *CryoLetters*, 23(2): 103-112.
- Kaczmarczyk, A., Menon, A., Al-Hanbali, A., Funnekotter, B., Bunn, E., Phang, P. Y., & Mancera, R. L. (2012). Current issues in plant cryopreservation. In I. KatKov (Ed.), *Current Frontiers in Cryobiology* (pp. 418-438). Croatia: Intech.

- Kartha, K., Fowke, L., Leung, N., Caswell, K., & Hakman, I. (1988). Induction of somatic embryos and plantlets from cryopreserved cell cultures of white spruce (*Picea glauca*). *Journal of Plant Physiology*, 132(5): 529-539.
- Kaviani, B. (2011). Conservation of plant genetic resources by cryopreservation. *Australian Journal of Crop Science*, 5(6): 778-800.
- Keller, E. (2005). Improvement of cryopreservation results in garlic using low temperature preculture and high-quality in vitro plantlets. *CryoLetters*, 26(6): 357-366.
- Khan, M. L., Uma S. (2001). Effect of seed weight, light regime, and substratum microsite on germination and seedling growth of *Quercus semiserrata* Roxb. *Tropical Ecology*, 42: 117-125.
- Khan, M. L., Bhuyan, P., Singh, N. D., Todaria, N. P. (2002). Fruit set, seed germination and seedling growth of *Mesua ferrea* Linn. (Clusiaceae) in relation to light intensity. *Journal of Tropical Forest Science*, 14: 35-48.
- Khan, M. L. (2004). Effects of seed mass on seedling success in *Artocarpus heterophyllus* L., a tropical tree species of North-East India. *Acta Oecologica*, 25: 103-110.
- Khan, M., Xu, H., & Hepworth, S. R. (2014). Blade-on-petiole genes: setting boundaries in development and defense. *Plant science*, 215: 157-171.
- Khoddamzadeh, A. A., Sinniah, U. R., Lynch, P., Kadir, M. A., Kadzimin, S. B., & Mahmood, M. (2011). Cryopreservation of protocorm-like bodies (PLBs) of *Phalaenopsis bellina* (Rchb. f.) Christenson by encapsulation-dehydration. *Plant Cell, Tissue and Organ Culture*, 107(3): 471-481.
- Khoury, C., Laliberté, B., & Guarino, L. (2010). Trends in ex situ conservation of plant genetic resources: a review of global crop and regional conservation strategies. *Genetic Resources and Crop Evolution*, 57(4): 625-639.
- Kim, H. H., Cho, E. G., Baek, H. J., Kim, C. Y., Keller, E., & Engelmann, F. (2004). Cryopreservation of garlic shoot tips by vitrification: effects of dehydration, rewarming, unloading and regrowth conditions. *CryoLetters*, 25(1): 59-70.
- Korban, S., & Sul, I. W. (2007). Micropropagation of coast redwood (*Sequoia sempervirens*). In S. M. Jain & H. Haggman (Eds.), *Protocols for Micropropagation of Woody Trees and Fruits* (pp. 23-32). The Netherlands: Springer.
- Kulus, D., (2015). Application of cryopreservation for chrysanthemum genetic resources conservation. *Acta Horticulturae*, 1087: 225-232.
- Kutas, E., Veyevnik, A., Titok, V., & Ogorodnyk, L. (2016). Morphogenesis of rhododendron yellow, introduced varieties of high bush blueberry, red bilberry

ordinary, depending on the composition of the nutrient media. *International Journal of Advanced Research Biological Science*, 3(3): 108-112.

Lawson, A., Ahmad, H., & Sambanis, A. (2011). Cytotoxicity effects of cryoprotectants as single-component and cocktail vitrification solutions. *Cryobiology*, 62(2): 115-122.

Leunufna, S., & Keller, E. R. J. (2003). Investigating a new cryopreservation protocol for yams (*Dioscorea* spp.). *Plant Cell Reports*, 21(12): 1159-1166.

Li, B. Q., Feng, C. H., Hu, L. Y., Wang, M. R., Chen, L., & Wang, Q. C. (2014). Shoot regeneration and cryopreservation of shoot tips of apple (*Malus*) by encapsulation- dehydration. *In Vitro Cellular Developmental Biology- Plant*, 50: 357–368.

Long, T. J., Jones, R. H. (1996). Seedling growth strategies and seed size effects in fourteen oak species native to different soil moisture habitats. *Trends in Ecology and Evolution*, 11:1-8.

Lynch, P. T., Siddika, A., Johnston, J. W., Trigwell, S. M., Mehra, A., Benelli, C., Lambardi, M., & Benson, E. E. (2011). Effects of osmotic pretreatments on oxidative stress, antioxidant profiles and cryopreservation of olive somatic embryos. *Plant Science*, 181:47–56.

Lynch, P. T., Souch, G. R., & Harding, K. (2012). Effects of post-harvest storage of *Allium sativum* bulbs on the cryopreservation of stem-discs by encapsulation/dehydration. *The Journal of Horticultural Science and Biotechnology*, 87(6): 588-592.

Magyar-Tábori, K., Dobránszki, J., da Silva, J. A. T., Bulley, S. M., & Hudák, I. (2010). The role of cytokinins in shoot organogenesis in apple. *Plant Cell, Tissue and Organ Culture*, 101(3), 251-267.

Mal, B., Rao, R., Sthapit, B., & Sajise, P. (2011). Conservation and sustainable use of tropical fruit species diversity: Bioversity's efforts in Asia, the Pacific and Oceania. *Indian Journal of Plant Genetic Resource* 24(1):1-22.

Manjunath B. (1948). *The wealth of India*. India: Council of Scientific and Industrial Research.

Marín, J., García, E., Lorente, P., Andreu, P., & Arbeloa, A. (2016). A novel approach for propagation of recalcitrant pistachio cultivars that sidesteps rooting by ex vitro grafting of tissue cultured shoot tips. *Plant Cell, Tissue and Organ Culture*, 124(1): 191-200.

Marković, Z., Chatelet, P., Sylvestre, I., Kontić, J., & Engelmann, F. (2013). Cryopreservation of grapevine (*Vitis vinifera* L.) in vitro shoot tips. *Open Life Sciences*, 8(10): 993-1000.

- Martinez-Montero, M. E., & Harding, K. (2015). Cryobionomics: Evaluating the Concept in Plant Cryopreservation. In D. Bahr, M. S. Khan & E. Davies (Eds.), *Plant Omics: The Omics of Plant Science* (pp. 655-682). New York: Springer.
- Masondo, N. A., Aremu, A. O., Finnie, J. F., & Van Staden, J. (2014). Plant growth regulator induced phytochemical and antioxidant variations in micropropagated and acclimatized *Eucomis autumnalis* subspecies *autumnalis* (Asparagaceae). *Acta Physiologiae Plantarum*, 36(9): 2467-2479.
- Matsumoto, T., Sakai, A., & Yamada, K. (1995). Cryopreservation of in vitro-grown apical meristems of lily by vitrification. *Plant Cell, Tissue and Organ Culture*, 41(3): 237-241.
- Mazur, P. (2004). Principles of cryobiology. In: B. J. Fuller, N. Lane, E. E. Benson (Eds) *Life in the frozen state*. London: CRC press.
- McGowran, E., Douglas, G., & Parkinson, M. (1998). Morphological and physiological markers of juvenility and maturity in shoot cultures of oak (*Quercus robur* and *Q. petraea*). *Tree physiology*, 18: 251-258.
- Mendonça, D., Luna, S., Bettencourt, S., Lopes, M. S., Monteiro, L., Neves, J. D., & da Câmara Machado, A. (2015). In vitro propagation of *Picconia azorica* (Tutin) Knobl (Oleaceae) an Azorean endangered endemic plant species. *Acta Physiologiae Plantarum*, 37(3): 1-10.
- Menon, A., Funnekotter, B., Kaczmarczyk, A., Bunn, E., Turner, S., & Mancera, R. L. (2014). Cold-induced changes affect survival after exposure to vitrification solution during cryopreservation in the South-West Australian Mediterranean climate species *Lomandra sonderi* (Asparagaceae). *Plant Cell, Tissue and Organ Culture*, 119(2): 347-358.
- Meryman, H. T., & Williams, R. J. (1985). Basic principles of freezing injury to plant cells; natural tolerance and approaches to cryopreservation. In K. K. Kartha (Ed), *Cryopreservation of plant cells and organs* (pp 13-47) Boca Raton: CRC Press.
- Mubbarakh, S. A., Rahmah, S., Rahman, Z. A., Sah, N. N. M., & Subramaniam, S., (2014). Cryopreservation of Brassidium shooting star orchid using the PVS3 method supported with preliminary histological analysis. *Applied Biochemical Biotechnology*, 172: 1131-1145.
- Murashige, T., & Skoog, F. (1962). A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Acta Physiologia Plantarum*, 15(3): 473-497.
- Nadarajan, J., Staines, H., Benson, E., Marzalina, M., Krishnapillay, B., & Harding, K. (2007). Optimization of cryopreservation for *Sterculia cordata* zygotic

embryos using vitrification techniques. *Journal of Tropical Forest Science*, 19(2):96-69.

- Nagesh, K. S., Shanthamma, C., & Bhagyalakshmi, N. (2009). Role of polarity in de novo shoot bud initiation from stem disc explants of *Curculigo orchoides* Gaertn. and its encapsulation and storability. *Acta Physiologiae Plantarum*, 31(4): 699-704.
- Nas, M. N., Bolek, Y., & Sevgin, N. (2010). The effects of explant and cytokinin type on regeneration of *Prunus microcarpa*. *Scientia Horticulturae*, 126(2): 88-94.
- Navarro-García, N., Morte, A., & Pérez-Tornero, O. (2016). In vitro adventitious organogenesis and histological characterization from mature nodal explants of *Citrus limon*. *In Vitro Cellular & Developmental Biology-Plant*, 52(2): 161-173.
- Nishizawa, S., Sakai, A., Amano, Y., & Matsuzawa, T. (1993). Cryopreservation of asparagus (*Asparagus officinalis* L.) embryogenic suspension cells and subsequent plant regeneration by vitrification. *Plant Science*, 91(1): 67-73.
- Noor, N. M., Kean, C. W., Vun, Y. L., & Mohamed-Hussein, Z. A. (2011). In vitro conservation of Malaysian biodiversity—achievements, challenges and future directions. *In Vitro Cellular & Developmental Biology-Plant*, 47(1): 26-36.
- Normah, M., Chin, H. F., & Reed, B. M. (2012). *Conservation of tropical plant species*. New York: Springer.
- Normah, M., Malik, S., Chaudhury, R., Salma, I., & Makeen, M. (2013). Conservation of tropical fruit genetic resources. In M. N. Normah & H. F. Chin (Eds.), *Conservation of Tropical Plant Species* (pp. 137-170). New York: Springer.
- Oliver, M. J. (1996). Desiccation tolerance in vegetative plant cells. *Acta Physiologia Plantarum*, 97(4): 779-787
- Ozudogru, E. A., & Kaya, E. (2012). Cryopreservation of *Thymus cariensis* and *T. vulgaris* shoot tips: comparison of three vitrification-based methods. *CryoLetters*, 33(5): 363-375.
- Padayachee, K., Watt, M., Edwards, N., & Mycock, D. (2008). Physiological responses of Eucalyptus in vitro axillary buds to cryopreparative desiccation and osmotic preculture: Effects of abscisic acid. *South African Journal of Botany*, 74(4): 639-646.
- Pammenter, N., Berjak, P., & Wesley-Smith, J. (2007). Cryopreservation of embryonic axes of selected *amaryllid* species. *CryoLetters*, 28(5): 387-399.

- Parveen, S., & Shahzad, A. (2011). A micropropagation protocol for *Cassia angustifolia* Vahl. from root explants. *Acta physiologiae plantarum*, 33(3): 789-796.
- Prickett, R. C., Marquez-Curtis, L. A., Elliott, J. A. W., McGann, L. E., 2015. Effect of supercooling and cell volume on intracellular ice formation. *Cryobiology*, 70: 156–163.
- Phulwaria, M., Rai, M. K., Gupta, A. K., Ram, K., & Shekhawat, N. (2012). An improved micropropagation of *Terminalia bellirica* from nodal explants of mature tree. *Acta Physiologiae Plantarum*, 34(1): 299-305.
- Popenoe, W. (1974). *Manual of Tropical and Sub-tropical Fruits*. New York: Halfner.
- Rabba'a, M. M., Shibli, R. A., & Shatnawi, M. A. (2012). Cryopreservation of *Teucrium polium* L. shoot-tips by vitrification and encapsulation-dehydration. *Plant Cell, Tissue and Organ Culture*, 110(3), 371-382.
- Rahaman, M. A., & Blake, J. (1988). Factors affecting in vitro proliferation and rooting of shoots of jackfruit (*Artocarpus heterophyllum* Lam). *Plant Cell, Tissue and Organ Culture*, 13: 179-187.
- Rai, M. K., Jaiswal, V. S., & Jaiswal, U. (2009). Shoot multiplication and plant regeneration of guava (*Psidium guajava* L.) from nodal explants of in vitro raised plantlets. *Journal of Fruit and Ornamental Plant Research*, 17(1): 29-38.
- Rajasekharan, P. (2006). In vitro conservation of horticultural crops. In P. Rajasekharan (Ed.), *In vitro Conservation and Cryopreservation-New Options to Conserve Horticultural Genetic Resources* (pp 66-72). Bangalore: Indian Institute of Horticultural Research.
- Rakshit, S., Rashid, Z., Sekhar, J., Fatma, T., & Dass, S. (2010). Callus induction and whole plant regeneration in elite Indian maize (*Zea mays* L.) inbreds. *Plant Cell, Tissue and Organ Culture*, 100(1): 31-37.
- Rao, R., & Sthapit, B. (2013). Conservation of tropical plant genetic resources: in situ approach. In M. N. Normah & H. F. Chin (Eds.), *Conservation of Tropical Plant Species* (pp. 3-26). New York: Springer.
- Reddy, C. V. K., Sreeramulu, D., & Raghunath, M. (2010). Antioxidant activity of fresh and dry fruits commonly consumed in India. *Food Research International*, 43(1): 285-288.
- Reed, B. M., Schumacher, L., Wang, N., D'Achino, J., & Barker, R. E. (2006). Cryopreservation of bermudagrass germplasm by encapsulation dehydration. *Crop Science*, 46(1): 6-11.

- Reed, B. M., Sarasan, V., Kane, M., Bunn, E., & Pence, V. C. (2011). Biodiversity conservation and conservation biotechnology tools. *In Vitro Cellular & Developmental Biology-Plant*, 47(1): 1-4.
- Reed, B. M., Gupta, S., & Uchendu, E. E. (2013). In vitro genebanks for preserving tropical biodiversity. In M. N. Normah & H. F. Chin (Eds.), *Conservation of Tropical Plant Species* (pp. 77-106). New York: Springer.
- Ruan, Y. L., Xu, S. M., White, R., & Furbank, R. T. (2004). Genotypic and developmental evidence for the role of plasmodesmatal regulation in cotton fiber elongation mediated by callose turnover. *Plant Physiology*, 136(4): 4104-4113.
- Sakai, A., Kobayashi, S., & Oiyama, I. (1990). Cryopreservation of nucellar cells of navel orange (*Citrus sinensis* Osb. var. *brasiliensis* Tanaka) by vitrification. *Plant Cell Reports*, 9(1): 30-33.
- Sakai, A., Engelmann, F., Takagi, H. (2000). Development of cryopreservation techniques. In F. Engelmann & H. Takagi (Eds.), *Cryopreservation of Tropical Plant Germplasm. Current Research Progress and Applications* (pp. 1-7). Italy: IPGRI.
- Sakai, A., & Engelmann, F. (2007). Vitrification, encapsulation-vitrification and droplet-vitrification: a review. *CryoLetters*, 28(3): 151-172.
- Sakai, A., Hirai, D., Niino, T. (2008). Development of PVS-based vitrification and encapsulation-vitrification protocols. In B. M. Reed (Ed.), *Plant Cryopreservation: A Practical Guide* (pp. 33-57). New York: Springer.
- Salma, M., Fki, L., Engelmann-Sylvestre, I., Niino, T., & Engelmann, F. (2014). Comparison of droplet-vitrification and D-cryoplate for cryopreservation of date palm (*Phoenix dactylifera* L.) polyembryonic masses. *Scientia horticultrae*, 179: 91-97.
- San José, M. C., Valladares, S., Janeiro, L. V., & Corredoir, E. (2014). Cryopreservation of in vitro-grown shoot tips of *Alnus glutinosa* (L.) Gaertn. *Acta Physiologiae Plantarum*, 36(1): 109-116.
- Sarasan, V. (2010). Importance of in vitro technology to future conservation programmes worldwide. *Kew Bulletin*, 65(4): 549.
- Sarasan, V., Cripps, R., Ramsay, M. M., Atherton, C., McMichen, M., Prendergast, G., & Rowntree, J. K. (2006). Conservation in vitro of threatened plants—progress in the past decade. *In Vitro Cellular and Developmental Biology-Plant*, 42(3): 206-214.

- Seki, S., & Mazur, P. (2009). The dominance of warming rate over cooling rate in the survival of mouse oocytes subjected to a vitrification procedure. *Cryobiology*, 59(1): 75-82.
- Sharaf, S. A., Shibli, R. A., Kasrawi, M. A., & Baghdadi, S. H. (2012). Cryopreservation of wild Shih (*Artemisia herba-alba* Asso.) shoot-tips by encapsulation-dehydration and encapsulation-vitrification. *Plant Cell, Tissue and Organ Culture*, 108(3): 437-444.
- Shirly, R., & Iyer, R. (2007). Micropropagation of cashew (*Anacardium occidentale* L.). In S. M. Jain & H. Haggman (Eds.), *Protocols for Micropropagation of Woody Trees and Fruits* (pp. 313-322) The Netherlands: Springer.
- Sidhu, A. S., (2012). *Jackfruit improvement in the Asia-Pacific Region-A status report. Asia-Pacific*. Thailand: Association of Agricultural Research Institutions.
- Sinniah, U.R., & Gantait, S., (2013). Cryopreservation of immature *Parkia speciosa* Hassk. zygotic embryonic axes following desiccation or exposure to vitrification solution. *Acta Physiologiae Plantarum*, 35: 2629–2634.
- Suzuki, M., Akihama, T., & Ishikawa, M. (2005). Cryopreservation of encapsulated gentian axillary buds following 2 step-preculture with sucrose and desiccation. *Plant Cell, Tissue and Organ Culture*, 83(1): 115-121.
- Swami, S. B., Thakor, N., Haldankar, P., & Kalse, S. (2012). Jackfruit and its many functional components as related to human health: a review. *Comprehensive Reviews in Food Science and Food Safety*, 11(6): 565-576.
- Takagi, H. (2000). Recent developments in cryopreservation of shoot apices of tropical species. In F. Engelmann & H. Takagi (Eds.), *Cryopreservation of Tropical Plant Germplasm. Current Research Progress and Applications* (pp. 178–193). Italy: IPGRI.
- Thach, B. D., Le Nguyen Tu Linh, D. T., Cang, T. T. B., Giang, T. T. L., Uyen, N. P. A., & Suong, N. K. (2016). Protocol Establishment For Multiplication And Regeneration Of *Polyscias Fruticosa* L. Harms. An Important Medicinal Plant In Vietnam. *European Journal of Biotechnology and Genetic Engineering*, 3:1.
- Thammasiri, K. (1999). Cryopreservation of embryonic axes of jackfruit. *CryoLetters*, 20(1): 21-28.
- Thinh, N. T., Takagi, H., & Sakai, A. (2000). Cryopreservation of in vitro-grown apical meristems of some vegetatively propagated tropical monocots by vitrification. In F. Engelmann & H. Takagi (Eds.), *Cryopreservation of Tropical Plant Germplasm. Current Research Progress and Applications* (pp. 227–232). Italy: IPGRI.

- Uchendu, E. E., & Reed, B. M. (2008). A comparative study of three cryopreservation protocols for effective storage of in vitro-grown mint (*Mentha spp.*). *CryoLetters*, 29(3): 181-188.
- Vidal, N., Sánchez, C., Jorquera, L., Ballester, A., & Vieitez, A. M. (2005). Cryopreservation of chestnut by vitrification of in vitro-grown shoot tips. *In Vitro Cellular & Developmental Biology-Plant*, 41(1): 63-68.
- Vieitez, A., Sánchez, M., García-Nimo, M., & Ballester, A. (2007). Protocol for micropropagation of *Castanea sativa*. In S. M. Jain & H. Haggman (Eds.), *Protocols for Micropropagation of Woody Trees and Fruits* (pp. 229-312). The Netherlands: Springer.
- Volk, G. M., & Walters, C. (2006). Plant vitrification solution 2 lowers water content and alters freezing behavior in shoot tips during cryoprotection. *Cryobiology*, 52(1): 48-61.
- Volk, G. M., & Caspersen, A. M. (2007). Plasmolysis and recovery of different cell types in cryoprotected shoot tips of *Mentha* × *Piperita*. *Protoplasma*, 231(3-4): 215-226.
- Vujović, T., Sylvestre, I., Ružić, D., & Engelmann, F. (2011). Droplet-vitrification of apical shoot tips of *Rubus fruticosus* L. and *Prunus cerasifera* Ehrh. *Scientia Horticulturae*, 130(1): 222-228.
- Walters, C., Wesley-Smith, J., Crane, J., Hill, L. M., Chmielarz, P., Pammenter, N. W., & Berjak, P. (2008). Cryopreservation of recalcitrant (ie desiccation-sensitive) seeds. In B. M. Reed (Ed.), *Plant Cryopreservation: A Practical Guide* (pp. 465-484). New York: Springer.
- Wang, B., Li, J. W., Zhang, Z. B., Wang, R. R., Ma, Y. L., Blystad, D. R., & Wang, Q. C. (2014). Three vitrification-based cryopreservation procedures cause different cryo-injuries to potato shoot tips while all maintain genetic integrity in regenerants. *Journal of Biotechnology*, 184: 47-55.
- Wen, B., Cai, C., Wang, R., Tan, Y., & Lan, O. (2010). Critical moisture content windows differ for the cryopreservation of pomelo (*Citrus grandis*) seeds and embryonic axes. *CryoLetters*, 31: 29-39.
- Wesley-Smith, J., Vertucci, C. W., Berjak, P., Pammenter, N. W., & Crane, J. (1992). Cryopreservation of desiccation-sensitive axes of *Camellia sinensis* in relation to dehydration, freezing rate and the thermal properties of tissue water. *Journal of Plant Physiology*, 140(5): 596-604.
- Wesley-Smith, J., Pammenter, N., Berjak, P., & Walters, C. (2001). The effects of two drying rates on the desiccation tolerance of embryonic axes of recalcitrant jackfruit (*Artocarpus heterophyllus* Lamk.) seeds. *Annals of Botany*, 88(4): 653-664.

- Xue, S. H., Luo, X. J., Wu, Z. H., Zhang, H. L., & Wang, X. Y. (2008). Cold storage and cryopreservation of hairy root cultures of medicinal plant *Eruca sativa* Mill., *Astragalus membranaceus* and *Gentiana macrophylla* Pall. *Plant Cell, Tissue and Organ Culture*, 92(3): 251-260.
- Yadav, K., & Singh, N. (2011). In vitro propagation and biochemical analysis of field established wood apple (*Aegle marmelos* L.). *Analele Universității din Oradea–Fascicula Biologie*, 18(1): 23-28.
- Yıldırım, A. N., Şan, B., Yıldırım, F., Ecevit, F. M., & Ercişli, S. (2015). Micropropagation of Promising Jujube (*Ziziphus jujuba* Mill.) Genotypes. *Erwerbs-Obstbau*, 57(3): 135-140.
- Yin, M., & Hong, S. (2009). Cryopreservation of *Dendrobium candidum* Wall. ex Lindl. protocorm-like bodies by encapsulation-vitrification. *Plant Cell, Tissue and Organ Culture*, 98(2), 179-185.
- Zámečník, J., Faltus, M., Bilavčík, A., & Kotková, R. (2012). Comparison of cryopreservation methods of vegetatively propagated crops based on thermal analysis: Current issues in plant cryopreservation. In I. KatKov (Ed.), *Current Frontiers in Cryobiology* (pp. 333-358). Croatia: Intech.
- Zhang, J. M., Zhang, X. N., Lu, X. X., Xin, X., Yin, G. K., He, J. J., Xu, Y. M., & Chen, X. L. (2014). Optimization of droplet-vitrification protocol for carnation genotypes and ultrastructural studies on shoot tips during cryopreservation. *Acta Physiologiae Plantarum*, 36: 3189–3198.
- Zhumagulova, Z. B., Kovalchuk, I., Reed, M. B., Kampitova, G., & Turdiev, T. (2014). Effect of pretreatment methods of dormant pear buds on viability after cryopreservation. *World Applied Science Journal*, 30: 330-334.