



UNIVERSITI PUTRA MALAYSIA

***PHYSIOLOGICAL AND CHROMOSOMAL CHANGES OF DELAYED
HARVEST AND STORED SOYBEAN (*Glycine max* L. Merr.) SEEDS
FOLLOWING PRIMING***

PHYU SIN THANT

FP 2018 34



**PHYSIOLOGICAL AND CHROMOSOMAL CHANGES OF DELAYED
HARVEST AND STORED SOYBEAN (*Glycine max* L. Merr.) SEEDS
FOLLOWING PRIMING**

By

PHYU SIN THANT

**Thesis submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

February 2018

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of University Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of University Putra Malaysia.

Copyright © Universiti Putra Malaysia




DEDICATION

Dedicated to:

(My Beloved Parents)

U Kyi Myint and Daw Kyin



© COPYRIGHT UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

PHYSIOLOGICAL AND CHROMOSOMAL CHANGES OF DELAYED HARVEST AND STORED SOYBEAN (*Glycine max* L. Merr.) SEEDS FOLLOWING PRIMING

By

PHYU SIN THANT

February 2018

Chairman: Professor Adam bin Puteh, PhD
Faculty: Agriculture

Seed deteriorations begin on the mother plant in the field and continue during storage. Basic factors influencing seed ageing are temperature, relative humidity, seed moisture content and duration of storage. Seed priming known as a pre-sowing seed treatment using natural or synthetic compounds is commonly practiced to improve seed germination and seedling emergence in a wide range of crop species. This study was undertaken to evaluate the effectiveness of seed priming treatments and post-storage priming treatments on changes in seed quality and mechanisms involved in deterioration process during field weathering and under controlled storage. For field weathering, seeds of AGS-190 and Cikurai were harvested at harvest maturity (HM), one week after HM (H1) and three weeks after HM (H3). At the time of harvest, the seeds from main stem and branches were differentiated to evaluate seed quality from different positions under field weather conditions. For storage study, the HM seeds of AGS-190 and Cikurai were stored in cold room ($8\pm 2^{\circ}\text{C}$) or room temperature ($25\pm 2^{\circ}\text{C}$) for 3, 6, 9, 12 and 15 months. Aged seeds of delayed harvest or stored seeds were primed with -0.8MPa PEG (Polyethylene glycol), 0.5% chitosan and water.

The delayed harvested seeds from main stem and branches showed no difference in seed quality and seedling performance. Soybean seeds harvested past the HM showed adverse effects on seed physiological assessments and consequently affected seed quality. Cultivar AGS-190 was more sensitive to adverse weather conditions as shown by deterioration of seed quality at one week after HM while cultivar Cikurai with black seed coat features could maintain better seed quality up to one week after HM. Soybean seed viability and vigor considerably declined in H3 seeds with decreased activities of catalase (CAT) and superoxide dismutase (SOD) and increased accumulation of malondialdehyde (MDA) content and chromosomal aberrations. Loss of soybean seed quality in delayed harvest seeds was influenced by seed moisture content during harvest and *Phomopsis* sp. infection. The reactive oxygen species (ROS) production in moist seeds are much higher than dry seeds. Higher accumulation of ROS with concomitant increase in MDA content not only injured cell membrane but also caused oxidative damage to DNA of delayed harvest seeds and chromosomal aberrations.

Seed priming with 0.5% chitosan and -0.8MPa PEG enhanced viability of H1 seeds in both cultivars with better germination percentage, germination index, better seedling vigour index and faster speed of germination time. Priming treatments decreased the contents of MDA and the accumulation of chromosomal aberrations in delayed harvest seeds. Recovery of germinability in field weathered seeds during priming is strongly associated with increased synthesis in CAT or SOD activities, inhibiting accumulation of lipid peroxidation and genetic damage.

Significant increases in MDA content and electrical conductivity (EC) of seed leachate with increasing storage periods indicated that ageing of the seeds in room temperature ($25\pm 2^{\circ}\text{C}$) caused oxidative damage to cell membrane integrity. The activities of CAT and SOD in the seeds stored at room temperature decreased with longer storage time which was favorable for ROS accumulation. Oxidative damage caused by ROS accumulation during storage at room temperature not only oxidized lipid but also damaged the nucleic acid which led to chromosomal aberrations. Deteriorative effects of seed ageing inhibited to some extent of the metabolic processes for root and shoot growth resulting longer mean germination time (MGT) and slower speed of germination, lower performance of seedling growth in aged seeds. Storing soybean seeds at $25\pm 2^{\circ}\text{C}$ could maintain seed viability and vigour until 3 months. Soybean seeds stored at $8\pm 2^{\circ}\text{C}$ could maintain seed viability up to 15 months and seed vigour up to 12 months.

Post-storage priming with -0.8MPa PEG in 6 months stored seeds of AGS-190 resulted in higher germination percentage, germination index and better seedling performance. Moreover, post-storage priming with 0.5% chitosan and -0.8MPa PEG of seeds stored for 6-9 months improved germination percentage, germination index, seedling vigour index of cultivar Cikurai. Osmopriming with -0.8MPa PEG improved the mechanisms involved in seed germination involving CAT and SOD activities, MDA accumulation and chromosomal changes of stored seeds of AGS-190 and Cikurai.

There were six different types of chromosomal aberrations observed in deteriorated seeds during field weathering and controlled storage. Under both conditions, the most abundant types of chromosomal aberrations are single bridge and sticky chromosomes. Priming with water, 0.5% chitosan and -0.8MPa PEG generally reduced single bridge and sticky types of chromosomal aberrations in both field deteriorated seeds and stored seeds. Priming not only repairs the chromosomal damage, but appears to slow down the ageing process. This study indicates that deterioration process of soybean seeds involves the production of reactive oxygen species (ROS) through depletion of antioxidant enzymes, and lipid peroxidation which ultimately interferes with cell mitotic activity. Priming improves seed quality through increase activities of antioxidant enzymes by repairing membrane damage and minimizing chromosomal damage.

Abstrak thesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

PERUBAHAN FISILOGI DAN KROMOSOM BAGI BIJI BENIH KACANG SOYA (*Glycine max* L. Merr.) TERTANGGUH PENUAIAN DAN DISIMPAN SETELAH PRIMING

Oleh

PHYU SIN THANT

Februari 2018

Pengerusi: Professor Adam bin Puteh, PhD
Fakulti: Pertanian

Kemerosotan biji benih bermula dari pokok induk di ladang dan berterusan sehingga penyimpanan. Faktor asas yang mempengaruhi kemerosotan biji benih adalah suhu, kandungan kelembapan benih dan tempoh penyimpanan. Priming biji benih yang dikenali sebagai rawatan benih pra-menanam menggunakan bahan semulajadi atau sintetik biasanya diamalkan untuk meningkatkan percambahan biji benih dan kemunculan anak benih untuk pelbagai spesies tanaman. Kajian ini dijalankan untuk menilai keberkesanan rawatan pembenihan benih dan rawatan penyembuhan pasca penyimpanan mengenai perubahan dalam kualiti benih dan mekanisma yang terlibat dalam proses kemerosotan semasa diladang dan semasa simpanan terkawal. Untuk kajian kemerosotan di ladang, benih AGS-190 dan Cikurai dituai pada peningkat matang penuaian (HM), satu minggu selepas HM (H1) dan tiga minggu selepas HM (H3). Pada masa penuaian, benih dari batang utama dan dahan dibezakan untuk menilai kualiti benih dari kedudukan yang berbeza di bawah keadaan cuaca lapangan. Untuk kajian penyimpanan, benih HM AGS-190 dan Cikurai disimpan di bilik sejuk ($8 \pm 2^{\circ}\text{C}$) atau suhu bilik ($25 \pm 2^{\circ}\text{C}$) selama 3, 6, 9, 12 dan 15 bulan. Biji benih yang mengalami kemerosotan di ladang dan biji benih yang disimpan dirawat dengan -0.8MPa PEG, 0.5% kitosan dan air.

Biji yang ditangguhkan penuaian dari batang utama dan dahan tidak menunjukkan perbezaan dalam kualiti benih dan prestasi anak benih. Biji benih kacang soya yang dituai melepasi HM menunjukkan kesan buruk terhadap penilaian fisiologi benih dan akibatnya kualiti benih terjejas. Kultivar kacang soya bersaiz besar lebih sensitif terhadap keadaan cuaca yang buruk seperti yang ditunjukkan oleh kemerosotan kualiti benih pada satu minggu selepas HM manakala kultivar bersaiz kecil dengan ciri-ciri benih hitam boleh mengekalkan kualiti biji benih yang lebih baik sehingga satu minggu selepas HM. Kebernasan dan kecergasan biji benih kacang soya agak menurun bagi benih H3 dengan penurunan aktiviti catalase (CAT) dan superoxide dismutase (SOD) dan peningkatan pengumpulan kandungan malondialdehid (MDA) dan perubahan kromosom. Kehilangan kualiti biji benih kacang soya yang tertangguh penuaian dipengaruhi oleh kandungan kelembapan benih semasa tuaian dan jangkitan *Phomopsis*

sp.. Kandungan ROS dalam benih lembab jauh lebih tinggi daripada benih kering. Pengumpulan lebih tinggi ROS dengan peningkatan bersamaan dengan kandungan MDA bukan sahaja mencederakan membran sel tetapi turut menyebabkan kerosakan oksidatif kepada DNA biji benih penuaian tertangguh dan kerosakan kromosom.

Priming biji benih dengan 0.5% chitosan dan -0.8MPa PEG meningkatkan keberhasilan benih HI untuk kedua-dua kultivar dengan meningkatkan peratusan percambahan, indeks percambahan, indeks perkembangan biji benih yang lebih baik serta percambahan yang lebih cepat. Rawatan priming menurunkan kandungan MDA dan pengumpulan perubahan kromosom dalam biji benih kacang soya yang ditangguhkan penuaian diladang. Pemulihan kebolehan bercambah bagi biji benih yang mengalami kemerosotan di ladang semasa priming dikaitkan dengan peningkatan sintesis dalam aktiviti CAT atau SOD, yang menghalang pengumpulan peroxidation lipid dan kerosakan genetik.

Peningkatan ketara dalam kandungan MDA dan EC leachate biji benih dengan peningkatan tempoh penyimpanan menunjukkan bahawa kemerosotan biji benih dalam suhu bilik ($25 \pm 2^\circ\text{C}$) menyebabkan kerosakan oksidatif kepada integriti membran sel. Aktiviti CAT dan SOD dalam biji benih yang disimpan pada suhu bilik berkurangan dengan masa penyimpanan lebih lama yang menggalakkan pengumpulan ROS. Kerosakan oksidatif yang disebabkan oleh pengumpulan ROS semasa penyimpanan pada suhu bilik bukan sahaja menyebabkan lipid teroksida tetapi juga merosakkan asid nukleik yang menyebabkan kerosakan kepada kromosom. Kesan kemerosotan biji benih menghindar proses metabolik untuk pertumbuhan akar dan pucuk menyebabkan MGT yang lebih lama dan masa percambahan lebih perlahan, di samping prestasi benih yang lebih rendah dalam benih tua. Menyimpan biji benih kacang soya pada $25 \pm 2^\circ\text{C}$ dapat mengekalkan keberhasilan biji benih selama 3 bulan tetapi kecergasan biji benih menurun dengan peningkatan tempoh penyimpanan. Biji kacang soya yang disimpan pada $8 \pm 2^\circ\text{C}$ dapat mengekalkan kecergasan benih hingga 15 bulan dan keberhasilan benih hingga 12 bulan.

Priming pasca penyimpanan dengan -0.8Mpa PEG dalam biji benih AGS-190 disimpan 6 bulan menghasilkan peratusan percambahan yang lebih tinggi, indeks percambahan dan prestasi biji benih yang lebih baik. Selain itu, priming pasca penyimpanan dengan 0.5% kitosan dan -0.8MPa PEG bagi biji benih yang disimpan untuk 6-9 bulan meningkatkan peratusan percambahan, indeks percambahan, indeks kesegahan anak benih bagi kultivar Cikurai. Osmopriming dengan -0.8MPa PEG telah meningkatkan mekanisme yang terlibat dalam percambahan benih yang melibatkan aktiviti CAT dan SOD, pengumpulan MDA dan perubahan kromosom bagi biji benih AGS-190 dan Cikurai yang disimpan.

Terdapat enam jenis kerosakan kromosom dalam benih yang terdedah pada persekitaran diladang dan semasa simpanan terkawal. Pada kedua-dua keadaan, jenis kerosakan kromosom yang paling banyak adalah jambatan tunggal dan kromosom melekat. Priming dengan air, 0.5% chitosan dan -0.8MPa PEG secara amnya mengurangkan kerosakan kromosom jambatan tunggal dan kromosom melekit. Priming bukan sahaja membaiki kerosakan kromosom, juga memperlakankan proses

kemososotan. Kajian ini menunjukkan bahawa proses kemerosotan biji benih kacang soya melibatkan penghasilan spesies oksigen reaktif (ROS) melalui pengurangan enzim antioksidan, dan peroxidation lipid yang mengganggu aktiviti mitosis sel. Priming meningkatkan kualiti biji benih melalui peningkatan aktiviti enzim antioksidan dengan membaiki kerosakan membran dan mengurangkan perubahan genetik.



ACKNOWLEDGEMENTS

The completion of this thesis was a long process and there are many people for whose support I am grateful. I would like to thank first and foremost my supervisor, Prof. Dr. Adam Puteh for giving me the opportunity to study in the area of seed technology. He provided me enormous support and guidance during this period of study. My special appreciation is extended to my supervisory committee member, Prof Dr. Uma Rani Sinniah for her intellectual guidance and professional expertise throughout my research and the thesis preparation process. I would like to express my deep sincere gratitude to Dr. Mohd Firdaus Bin Ismail for his valuable comments and help throughout my study.

I am truly grateful to Organization for Women in Science for the Developing World (OWSDW) for financial support by awarding me the scholarships from MSc leading to PhD. My thanks also go to OWSDW secretariats for their helpful communications and suggestion during my study.

I very much appreciate Mr. Harris Ahmad from Seed Technology laboratory and all staff members from Physiology and Postharvest laboratories for their kind helps and also providing facilities and material for my research assistance in my experiments. Regarding to Mycology Lab 2, Department of Plant Protection, thank you to Mr. Mohamed Nazri Abdul Rahman for his friendly assistance in microscopic work. I would like to express my honest thanks to staff members of field 2, UPM for their kind assistance. I would also like to thank my friends working in the seed technology laboratory for your unconditional and genuine friendship and for the great times that we have shared together. My deep gratitude is dedicated to my family, who always support and encourage me under whatever circumstances.

I certify that a Thesis Examination Committee has met on 5 February 2018 to conduct the final examination of Phyu Sin Thant on her thesis entitled "Physiological and Chromosomal Changes of Delayed Harvest and Stored Soybean (*Glycine max* L. Merr.) Seeds Following Priming" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Yahya bin Awang, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Ghizan bin Saleh, PhD


Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Nor Aini binti Ab Shukor, PhD

Professor
Faculty of Forestry
Universiti Putra Malaysia
(Internal Examiner)

Satriyas Ilyas, PhD

Professor
Bogor Agricultural University
Indonesia
(External Examiner)



NOR AINI AB. SHUKOR, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 March 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Adam bin Puteh, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Uma Rani Sinniah, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Mohd Firdaus bin Ismail, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____

Date: _____

Name and Matric No.: Phyu Sin Thant (GS43955)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: Adam bin Puteh

Signature: _____
Name of
Member of
Supervisory
Committee: Uma Rani Sinniah

Signature: _____
Name of
Member of
Supervisory
Committee: Mohd Firdaus bin Ismail

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	vi
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xv
LIST OF FIGURES	xviii
LIST OF APPENDICES	xx
LIST OF ABBREVIATIONS	xxi
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 Importance of Soybean	4
2.2 Soybean Seed Quality at Different Crop Growth Stages	5
2.3 Field Weathering	6
2.4 Storage	7
2.5 Responses of Cultivars to Seed Deterioration	8
2.6 Seed Borne Diseases and Seed Quality	10
2.7 Symptoms and Mechanisms of Seed Deterioration	10
2.7.1 Lipid Peroxidation	11
2.7.2 Chromosomal Aberrations	12
2.7.3 Types of Chromosomal Aberrations	13
2.7.4 Antioxidant Enzyme Systems	14
2.7.5 Seedling Performance of Deteriorated Seeds	16
2.8 Improvement of Seed Quality	17
2.8.1 Seed Priming	17
2.8.2 Effects of Priming on Mechanism Changes of Seed Germination	18
3 PHYSIOLOGICAL AND CHROMOSOMAL CHANGES OF DELAYED HARVEST SOYBEAN SEEDS	21
3.1 Introduction	21
3.2 Materials and Methods	22
3.2.1 Seed Materials and Field Weathering Treatments	22
3.2.2 Measurement of Seed Quality and Seedling Performance	24
3.2.2.1 Determination of A Hundred-seed Weight	24
3.2.2.2 Pathogen Bioassay	24
3.2.2.3 Determination of Seed Moisture Content	24
3.2.2.4 Electrical Conductivity (EC) Test	25
3.2.2.5 Standard Germination Test	25
3.2.2.6 Tetrazolium Viability Test	25

	3.2.2.7	Tetrazolium Vigor Test	25
	3.2.2.8	Evaluation of Germination Index	26
	3.2.2.9	Evaluation of Speed of Germination	26
	3.2.2.10	Evaluation of Mean Germination Time	26
	3.2.2.11	Evaluation of Seedling Vigor Index	26
	3.2.2.12	Measurement of Seedling Length and Dry Weight	26
	3.2.3	Antioxidant Enzyme Analysis	26
	3.2.3.1	Catalase (EC1.11.1.6) Assay	27
	3.2.3.2	Superoxide Dismutase (EC1.15.1.1) Assay	27
	3.2.4	Malondialdehyde (MDA) Assay	28
	3.2.5	Observation of Chromosomal Aberrations	28
	3.2.6	Data Analysis	30
3.3	Results		30
	3.3.1	Seed Viability and Vigour	30
	3.3.2	Germination Performance and Seedling Performance	33
	3.3.3	Seed Quality and Seedling Performance (Main Stem)	36
	3.3.4	Changes in Antioxidant Enzyme Activities	43
	3.3.5	Changes in Lipid Peroxidation	43
	3.3.6	Changes in Chromosomal Aberrations	45
	3.3.7	Types of Chromosomal Aberrations	45
3.4	Discussion		47
3.5	Conclusion		51
4	EFFECT OF PRIMING ON PHYSIOLOGICAL AND CHROMOSOMAL CHANGES OF DELAYED HARVEST SOYBEAN SEEDS		52
	4.1	Introduction	52
	4.2	Materials and Methods	53
	4.2.1	Seed Materials and Priming Treatments	53
	4.2.2	Measurement of Seed Quality and Seedling Performance	54
	4.2.3	Antioxidant Enzyme Analysis	54
	4.2.4	Malondialdehyde (MDA) Assay	54
	4.2.5	Observation of Chromosomal Aberrations	54
	4.2.6	Data Analysis	55
4.3	Results		55
	4.3.1	Effects of Priming on Seed Quality	55
	4.3.2	Effects of Priming on Seedling Performance	62
	4.3.3	Effect of Priming on Antioxidant Enzyme Activities	67
	4.3.3.1	Catalase (CAT) Activity	67
	4.3.3.2	Superoxide Dismutase (SOD) Activity	68
	4.3.4	Effect of Priming on Lipid Peroxidation	69

4.3.5	Effect of Priming on Chromosomal Aberrations	70
4.3.6	Types of Chromosomal Aberrations following Priming	71
4.4	Discussion	75
4.5	Conclusion	77
5	PHYSIOLOGICAL AND CHROMOSOMAL CHANGES OF STORED SOYBEAN SEEDS	79
5.1	Introduction	79
5.2	Materials and Methods	80
5.2.1	Seed Materials and Seed Aging Treatments	80
5.2.2	Measurement of Seed Quality and Seedling Performance	80
5.2.3	Antioxidant Enzyme Analysis	81
5.2.4	Malondialdehyde (MDA) Assay	81
5.2.5	Observation of Chromosomal Aberrations	81
5.2.6	Data Analysis	81
5.3	Results	81
5.3.1	Seed Viability and Vigour	81
5.3.2	Germination Performance	85
5.3.3	Seedling Performance	85
5.3.4	Changes in Antioxidant Enzyme Activities	90
5.3.4.1	Catalase (CAT) Activity	90
5.3.4.2	Superoxide Dismutase (SOD) Activity	90
5.3.5	Changes in Lipid Peroxidation	91
5.3.6	Changes in Chromosomal Aberrations	92
5.3.7	Types of Chromosomal Aberrations	93
5.4	Discussion	95
5.5	Conclusion	98
6	EFFECT OF PRIMING ON PHYSIOLOGICAL AND CHROMOSOMAL CHANGES OF STORED SOYBEAN SEEDS	100
6.1	Introduction	100
6.2	Materials and Methods	101
6.2.1	Seed Materials and Priming Treatments	101
6.2.2	Measurement of Seed Quality and Seedling Performance	102
6.2.3	Antioxidant Enzyme Analysis	102
6.2.4	Malondialdehyde (MDA) Assay	102
6.2.5	Observation of Chromosomal Aberrations	102
6.2.6	Data Analysis	102
6.3	Results	103
6.3.1	Effect of Priming on Seed Quality	103
6.3.2	Effect of Priming on Seedling Performance	108
6.3.3	Effect of Priming on Antioxidant Enzyme Activities	116
6.3.3.1	Catalase (CAT) Activity	116

	6.3.3.2	Superoxide Dismutase (SOD) Activity	117
	6.3.4	Effect of Priming on Lipid Peroxidation	118
	6.3.5	Effect of Priming on Chromosomal Aberrations	119
	6.3.6	Types of Chromosomal Aberrations following Priming	120
	6.4	Discussion	124
	6.5	Conclusion	126
7		GENERAL DISCUSSION, CONCLUSION AND RECOMMENDATIONS	127
		REFERENCES	131
		APPENDICES	159
		BIODATA OF STUDENT	169
		LIST OF PUBLICATIONS	170

LIST OF TABLES

Table		Page
3.1	Reproductive Stages of Soybean (Fehr and Caviness, 1977)	23
3.2	Mean square values of analysis of variance results for the effect of cultivars, harvest dates, seed positions on the plant and their interactions on seed viability and vigour	31
3.3	Main effects of cultivars, harvest dates and seed positions on the plant on seed viability and vigour	32
3.4	Interaction effect of soybean cultivar and harvest date on seed viability and vigour	34
3.5	Mean square values of analysis of variance results for the effect of cultivars, harvest dates, seed positions on the plant and their interaction on germination performance and seedling performance	35
3.6	Main effects of cultivars, harvest dates and seed positions on the plant on seed germination performance and seedling performance	37
3.7	Interaction effect of soybean cultivar and harvest date on germination performance	38
3.8	Interaction effect of harvest dates and seed positions on germination performance	39
3.9	The effects of harvest dates on seed viability and vigor of main stem seeds of AGS-190 and Cikurai	40
3.10	The effects of harvest dates on germination performance of main stem seeds of AGS-190 and Cikurai	42
3.11	Main effects of cultivars and harvest dates on seedling performance	43
4.1	Mean square values of analysis of variance results for the effect of cultivars, harvest dates, priming treatments and their interactions on seed quality	56
4.2	Interaction effect of cultivars and harvest dates on seed quality	57
4.3	Interaction effect of cultivars and priming treatments on seed quality	58
4.4	The effects of priming on seed quality of delayed harvest seeds of AGS-190	59
4.5	The effects of priming on seed quality of delayed harvest seeds of Cikurai	61

4.6	Mean square values of analysis of variance results for the effect of cultivars, harvest dates, priming treatments and their interactions on seedling performance	62
4.7	Interaction effect of cultivars and harvest dates on seedling performance	62
4.8	Interaction effect of cultivars and priming treatments on seedling dry weight	63
4.9	The effects of priming on seedling performance of delayed harvest seeds of AGS-190	65
4.10	The effects of priming on seedling root performance of delayed harvest seeds of Cikurai	66
4.11	Main effects of harvest dates and priming treatments on seedling shoot performance of Cikurai	67
4.12	Types of chromosomal aberrations found in delayed harvest seeds of AGS-190 after priming	73
4.13	Types of chromosomal aberrations found in delayed harvest seeds of Cikurai after priming.	74
5.1	Mean square values of analysis of variance results for the effects of storage conditions, cultivars, storage periods and their	83
5.2	Seed viability and vigour of AGS-190 and Cikurai as affected by storage periods and storage conditions	84
5.3	Mean square values of analysis of variance results for the effects of storage conditions, cultivars, storage periods and	86
5.4	Germination performance of AGS-190 and Cikurai as affected by storage periods and storage conditions	87
5.5	Mean square values of analysis of variance results for the effects of storage conditions, cultivars, storage periods and their interactions on seedling performance	88
5.6	Seedling shoot performance of AGS-190 and Cikurai as affected by storage periods and storage conditions	89
5.7	Main effects of storage conditions, cultivars and storage periods on seedling root performance	89
6.1	Mean square values of analysis of variance results for the effects of cultivars, storage periods, priming treatments and their	104

6.2	Interaction effect of soybean cultivars and storage periods on seed quality	105
6.3	Interaction effect of cultivars and priming treatments on seed quality	106
6.4	Seed quality of AGS-190 stored at room temperature as affected by storage periods and priming treatments	107
6.5	Main effects of storage periods and priming treatments on mean germination time of AGS-190 seeds stored at	109
6.6	Seed quality of Cikurai stored at room temperature as affected by storage periods and priming treatments	110
6.7	Main effects of storage periods and priming treatments on mean germination time of Cikurai seeds stored at room temperature	111
6.8	Mean square values of analysis of variance results for the effects of cultivars, storage periods, priming treatments and their interactions on seedling performance of seeds stored at room temperature	111
6.9	Interaction effects of soybean cultivars and priming treatments on seedling performance	112
6.10	Seedling performance of AGS-190 stored at room temperature as affected by storage periods and priming treatments	113
6.11	Main effect of storage periods and priming treatments on root dry weight of AGS-190 seeds stored at room temperature	114
6.12	Shoot dry weight of Cikurai stored at room temperature as affected by storage periods and priming treatments	115
6.13	Main effects of storage periods and priming treatments on seedling performance of Cikurai seeds stored at room temperature	116
6.14	Types of chromosomal aberrations found in AGS-190 seeds stored at room temperature after priming.	121
6.15	Types of chromosomal aberrations found in Cikurai seeds stored at room temperature as after priming	123

LIST OF FIGURES

Figure		Page
2.1	Accumulation of reactive oxygen species on cell membrane and DNA	13
2.2	Protection of reactive oxygen species by antioxidant enzyme systems	15
3.1	Anaphase cell division in the radicle-tips of soybean during a 12-hour period of the day	29
3.2	Changes in CAT activities of delayed harvest soybean seeds of AGS-190 and Cikurai.	44
3.3	Changes in SOD activities of delayed harvest soybean seeds of AGS-190 and Cikurai.	44
3.4	Changes in MDA content of delayed harvest soybean seeds of AGS-190 and Cikurai.	45
3.5	Changes in chromosomal damage in delayed harvest soybean seeds of AGS-190 and Cikurai.	46
3.6	Types of chromosomal aberrations found in different harvest dates and cultivars.	46
3.7	Relationship between seed viability and different types of chromosomal aberration in delayed harvest soybean seeds of AGS-190 (A) and Cikurai (B).	47
4.1	The effect of priming on catalase (CAT) activities of delayed harvest seeds of AGS-190 (A) and Cikurai (B).	68
4.2	The effect of priming on superoxide dismutase (SOD) activities of delayed harvest seeds of AGS-190 (A) and Cikurai (B).	69
4.3	The effect of priming on malondialdehyde (MDA) of delayed harvest seeds of AGS-190 (A) and Cikurai (B).	69
4.4	The effect of priming on chromosomal aberration of delayed harvest seeds of AGS-190 (A) and Cikurai (B).	70
4.5	Types of chromosomal aberrations found in delayed harvest seeds following priming.	72
5.1	Changes in catalase (CAT) activities of AGS-190 (A) and Cikurai (B) as affected by different storage conditions and storage periods.	90
5.2	Changes in SOD activities of AGS-190 (A) and Cikurai (B) as affected by different storage conditions and storage periods.	91

5.3	Changes in MDA contents of AGS-190 (A) and Cikurai (B) as affected by different storage conditions and storage periods.	92
5.4	Changes in chromosomal aberration of AGS-190 (A) and Cikurai (B) as affected by different storage conditions and storage periods.	92
5.5	Types of chromosomal aberrations found in stored seeds as affected by different storage conditions and storage periods.	94
5.6	Types of chromosomal aberration found in AGS-190 as affected by different storage conditions and storage periods.	94
5.7	Types of chromosomal aberrations found in Cikurai as affected by different storage conditions and storage periods.	95
6.1	Changes in catalase (CAT) of AGS-190 (A) and Cikurai (B) as affected by storage periods and priming.	117
6.2	Changes in superoxide dismutase (SOD) of AGS-190 (A) and Cikurai (B) as affected by storage periods and priming.	118
6.3	Changes in malondialdehyde (MDA) of AGS-190 (A) and Cikurai (B) as affected by storage periods and priming.	118
6.4	Changes in chromosomal aberration of AGS-190 (A) and Cikurai (B) as affected by storage periods and priming.	119
6.5	Types of chromosomal aberrations found in stored seeds as affected by priming.	122

LIST OF APPENDICES

Appendix		Page
A. 1	Origin and Characters of Soybean Cultivars Used	160
A. 2	Soybean Cultivars and Harvest Dates	160
A. 3	Meteorological data of 2016 planting season	160
A. 4	Tetrazolium staining patterns and their interpretation for soybean seeds (viable) as procedure describe in AOSA (1970)	161
A. 5	Tetrazolium staining patterns and their interpretation for soybean seeds (non-viable) as procedure describe in AOSA (1970)	162
B. 1	Mean square value of analysis of variance results for the effects of cultivars and harvest dates on catalase (CAT), superoxide dismutase (SOD), malondialdehyde (MDA) and chromosomal aberration	165
C. 1	Mean square value of analysis of variance results for the effects of harvest dates and priming treatments on catalase (CAT), superoxide dismutase (SOD), malondialdehyde (MDA) and chromosomal aberration of AGS-190	166
C. 2	Mean square value of analysis of variance results for the effects of harvest dates and priming treatments on catalase (CAT), superoxide dismutase (SOD), malondialdehyde (MDA) and chromosomal aberration of Cikurai	166
D. 1	Mean square value of analysis of variance results for the effects of storage conditions and storage period on catalase (CAT), superoxide dismutase (SOD), malondialdehyde (MDA) and chromosomal aberration of AGS-190	167
D. 2	Mean square value of analysis of variance results for the effects of storage conditions and storage period on catalase (CAT), superoxide dismutase (SOD), malondialdehyde (MDA) and chromosomal aberration of Cikurai	167
E. 1	Mean square value of analysis of variance results for the effects of storage period and priming on catalase (CAT), superoxide dismutase (SOD), malondialdehyde (MDA) and chromosomal aberration of AGS-190	168
E. 2	Mean square value of analysis of variance results for the effects of storage period and priming on catalase (CAT), superoxide dismutase (SOD), malondialdehyde (MDA) and chromosomal aberration of Cikurai	168

LIST OF ABBREVIATIONS

μM	Micromolar
μmol	Micromole
AOSA	Association of Official Seed Analysts
AVRDC	Asian Vegetable Research and Development Center
CAT	Catalase
DNA	Deoxyribose Nucleic Acid
EC	Electrical conductivity of seed leachates
GI	Germination index
H1	One week after HM
H3	Three weeks after HM
HM	Harvest maturity
ISTA	International Seed Testing Association
LSD	Least significant difference
M	Molarity
MDA	Malondialdehyde
MGT	Mean germination time
mM	Millimolar
mmol	Millimole
mol	Mole
nmol	Nanomole
OWSD	Organization of Women Scientists in Developing World
PEG	polyethylene glycol
PM	Physiological maturity
RCBD	Randomized complete block design
RH	Relative humidity
RNA	Ribose Nucleic Acid
ROS	Reactive oxygen species
Rpm	Revolution per minute
SOD	Superoxide Dismutase
SVI	Seedling vigor Index
TZ	Tetrazolium Test
UPM	University Putra Malaysia
UV	Ultra Violet
w/v	Weight per volume
w/w	Weight per weight

CHAPTER 1

INTRODUCTION

Soybean [*Glycine max* (L.) Merr.] is an economically important leguminous crop in the world. Soybean seeds consist of about 40% protein, 21 % oil and 34 % carbohydrates and 5% ash (Burton, 1997). Since soybean seed is a rich source of protein and oil, the production of soybean has been rapidly increasing as the world demand for protein and oil increase. Thus, soybean stands at the top in the world production of oil seed crops (Singh et al., 2011). Nowadays, soybean account for 35% of total harvested area devoted to annual and perennial oil crops (Prakash, 2008). Although soybean originated from China, United States of America is the leading producer in the world today producing about 117.21 million metric tonnes in 2016. Brazil and Argentina rank second and third in terms of production of 96.30 and 58.80 million metric tonnes, respectively. In Asia, the major soybean producing countries are China (11.97 million metric tonnes) and India (14.00 million metric tonnes) in 2016 (FAOSTAT, 2016).

High quality seeds which produce rapid and uniform field emergence of seedlings are essentially important for efficient production of soybean crop (Soltani et al., 2002). However, seed ageing which is described by reduction of seed viability with time is one of the major problems facing successful field establishment in agricultural production system. Deterioration of seed viability is an inevitable and irreversible process (Kapoor et al., 2011) and it is usually described by a decrease in germination percentage, reduced viability and vigour producing weak seedlings, and ultimately seed death (Tilebeni and Golpayegani, 2011). Seed ageing are influenced by a number of factors during field weathering, harvesting and storage.

Seed deterioration begins after harvest maturity. Unfavorable weather conditions which contribute to seed deterioration especially during post-maturation and pre-harvest period (Bhatia et al., 2010) are a challenge for production of high quality soybean seeds in the tropics and sub-tropics where the climate is characterized by high temperature and erratic rainfall (Pádua et al., 2009; Sanadhya and Dubeya, 2014). Delaying harvest past the harvest maturity stage causes longer exposure of seeds to adverse field conditions and consequently decreases the quality of the seeds (Eskandari, 2012). Losses of soybean seed viability and vigor due to delaying harvest have been reported by many researchers (Philbrook and Oplinger, 1989; Marcos-Filho et al., 1994; Dias et al. 2005; Diniz et al., 2013). However, the mechanisms involved in seed deterioration due to field weathering are unclear.

Seed aging after harvest and during storage is a major problem for maintaining high quality of planting materials. Major factors influencing seed longevity during storage are initial quality (Khatun et al., 2009), the storage conditions (Surki et al., 2012; Akter et al., 2014) and storage duration (Mbofung et al., 2013). Shelar et al. (2008) reported that decline in seed germination is much more serious under tropical conditions since environmental circumstances in these regions are challenging to maintain its viability during storage.

Seed deterioration is related to physiological and biochemical changes which include the decrease in enzyme activities, membrane integrity, protein synthesis and DNA degradation (Sun et al., 2007). Free radical reactions from reactive oxygen species (ROS) including superoxide radicals (O₂⁻), hydrogen peroxide (H₂O₂) and hydroxyl radicals (·OH) are regarded as primary causes of seed deterioration during controlled storage (Pukacka and Ratajczak, 2007; Kaewnaree et al., 2011). A number of studies proved that lipid peroxidation increased with seed ageing during storage in sunflower (Kibinza et al., 2006), wheat (Lehner et al., 2008), Sal tree (Parkhey et al., 2012), soybean (Xin et al., 2014) and oat (Xia et al., 2015). Whether similar physiological mechanisms occurred in seed deterioration in the field is not known.

The damage to DNA is a fundamental reason for nonfunctional cells or an organ. Accumulation of chromosomal aberrations in the seeds through time in controlled storage may cause the embryos unable to germinate and leads to loss of seed viability and vigor (Villiers, 1974; Rao et al., 1987). The occurrence and accumulation of chromosomal aberrations are a function of environmental temperature and moisture content and time during seed storage (Villiers 1974; Khan et al., 2003). Association between seed quality loss and increased frequencies of chromosomal aberrations were reported in lettuce seeds (Rao, et al., 1987), barley, peas and wheat under artificial aging conditions (Menezes et al., 2014). However, the study related to accumulation of chromosomal aberrations during field weathering has escaped attention.

Although reactive oxygen species (ROS) buildup is a major contributor to seed deterioration, these ROS are scavenged by various antioxidant defense systems during ageing (Yao et al., 2012). The main function of antioxidant enzymes is to protect cells from oxidative damage during growth, development and desiccation. Balancing production of ROS and protection of antioxidants is important for the cell to counter oxidative challenges (Fuchs et al., 1997). Degradation and inactivation of enzymes during aging contribute to seed deterioration due to changes in their macromolecular structure (McDonald, 2004). Several studies have revealed a close relationship between seed deterioration and a reduction in the activity of various antioxidant enzyme systems in cotton (Goel et al., 2003), sunflower (Kibinza et al., 2006), wheat (Lehner et al., 2008), soybean (Tian et al., 2008) and pea (Yao et al., 2012) seeds.

Deteriorated seeds critically reduce seed germination, seedling strength and field emergence. It is needed to develop technologies to improve seed quality of aged seeds to ensure crop yield. Seed priming has been widely used to enhance seed quality and seedling establishment in a wide range of crops. This technique is a pre-sowing technique to control imbibition process by treating seeds with natural or synthetic compounds (Jisha et al. 2012). Using this technique, seeds are partly imbibed to a point to allow pre-germination metabolic processes without radical protrusion (Giri and Schilinger, 2003). Seed priming has been approved to be a beneficial technique to improve seed germination and seedling establishments in delayed harvest soybean seeds (Thant, 2015) and storability of soybean seeds (Assefa, 2008). Various studies reported that increase in seed quality of primed seeds is related to increase in activation of antioxidant enzymes in order to protect cell membrane damage from reactive oxygen species (Kibinza et al., 2011; De-Oliveria et al., 2012; Siri et al., 2013), repair in chromosomal changes (Rao et al., 1987; Sivritepe an Durado, 1995) in aged seeds.

Soybean is sensitive to environmental conditions and the qualities of soybean seeds are influenced by field weather conditions during harvesting time especially under adverse environmental conditions. Moreover, these conditions make very difficult to maintain soybean seeds longevity during storage. It is important to know which mechanisms are involved in seed deterioration during field weathering and under controlled storage. On the other hand, it is critically important to produce rapid and uniform seedling emergence for good crop establishment mainly under adverse environmental conditions. Thus, the study needs to examine how seed priming treatments improve seed quality and reverse mechanisms involved in soybean seeds deterioration. Therefore, the present study was undertaken with the following objectives.

Objectives

1. To detect physiological mechanisms such as antioxidant enzyme activities, age-induced membrane deterioration and chromosomal aberrations in seed deterioration of delayed harvest and stored seeds.
2. To determine the beneficial effects of priming on changes in mechanisms involved in seed deterioration of delayed harvest and stored seeds.

REFERENCES

- Abdalla, F. H. and Roberts, E. H. (1968). Effects of temperature, moisture, and oxygen on the induction of chromosome damage in seeds of barley, broad beans, and peas during storage. *Annals of Botany*, 32(1): 119-136.
- Abdellaoui, R., Souid, A., Zayoud, D. and Neffati, M. (2013). Effects of natural long storage duration on seed germination characteristics of *Periploca angustifolia* Labill. *African Journal of Biotechnology*, 12(15): 1760-1768.
- Abdul-Baki, A. A. and Anderson, J. D. (1973). Vigor determination in soybean seed by multiple criteria. *Crop Science*, 13(6): 630-633.
- Abdullah, W. D., Powell, A. A. and Matthews, S. (1991). Association of differences in seed vigour in long bean (*Vigna sesquipedalis*) with testa colour and imbibition damage. *The Journal of Agricultural Science*, 116(02): 259-264.
- Adam, N. M., McDonald, M. B. and Henderlong, P. R. (1989). The influence of seed position, planting and harvesting dates on soybean seed quality. *Seed Science and Technology*, 17: 143-152.
- Adisarwanto, T. and Wudianto, R. (1999). Improving Crop Soybean production in lowland dry-downs. *Sower Swadaya*, Jakarta.
- Aebi, H. (1984). Catalase in vitro. *Method of Enzymology*, 105:121-126.
- Ajouri, A., Asgedom, H. and Becker, M. (2004). Seed priming enhances germination and seedling growth of barley under conditions of P and Zn deficiency. *Journal of Plant Nutrition and Soil Science*, 167(5): 630-636.
- Akhter, F. N., Kabir, G., Mannan, M. A. and Shaheen, N. N. (1992). Aging effect of wheat and barley seeds upon germination mitotic index and chromosomal damage. *Journal of Islamic Academy Science*, 5: 44-48.
- Akter, N., Haque, M. M., Islam, M. R. and Alam, K. M. (2014). Seed Quality of Stored Soybean (*Glycine max* L.) as Influenced by Storage Containers and Storage Periods. *The Agriculturists*, 12(1): 85-95.
- Amable, R. A. and Obendorf, R. L. (1986). Soybean seed respiration during simulated preharvest deterioration. *Journal of Experimental Botany*, 37(9): 1364-1375.
- Amooaghaie, R. (2011). The effect of hydro and osmopriming on alfalfa seed germination and antioxidant defenses under salt stress. *African Journal of Biotechnology*, 10(33): 6269-6275.
- Andrews, C. H. (1966). *Some aspects of pod and seed development in Lee soybeans*. PhD Thesis, Mississippi State University, USA.
- Anjum, S. A., Tanveer, M., Hussain, S., Bao, M., Wang, L., Khan, I. Ullah, E., Tung, S. A., Samad, R. A. and Shahzad, B. (2015). Cadmium toxicity in Maize (*Zea*

- mays* L.): consequences on antioxidative systems, reactive oxygen species and cadmium accumulation. *Environmental Science and Pollution Research*, 22(21):17022-17030.
- Ansari, O. and Sharif-Zadeh, F. (2012). Osmo and hydro priming improvement germination characteristics and enzyme activity of Mountain Rye (*Secale montanum*) seeds under drought stress. *Journal of Stress Physiology and Biochemistry*, 8(4):253-261.
- AOSA (1970). Rules for testing seed. *Proceeding of Association of Official Seed Analysts*, 60 (2): 1-116
- AOSA (2000). *Tetrazolium Testing Handbook*. Association of Official Seed Analysts. Lincoln, NE.
- AOSA. (2002). *Seed vigor testing handbook*. Association of Official Seed Analysts. Las Cruces, NM.
- Arif, M., Jan, M. T., Marawat, K. B. and Khan, M. A. (2008). Seed priming improves emergence and yield of soybean. *Pakistan Journal of Botany*, 40 (3): 1169-1177.
- Ashraf, M. and Foolad, M. R. (2005). Pre-Sowing seed treatment-A shotgun approach to improve germination, plant growth, and crop yield under saline and non-saline conditions. *Advances in Agronomy*, 88: 223-271.
- Asiedu, E. A. and Powell, A. A. (1998). Comparisons of the storage potential of cultivars of cowpea (*Vigna unguiculata*) differing in seed coat pigmentation. *Seed Science and Technology*, 26(1): 211-221.
- Assefa, M.K. (2008). *Effect of seed priming on storability, seed yield and quality of soybean (Glycine max (L.) Merrill)*. MSc Thesis. University of Agricultural Sciences, Dharwad.
- Azadi, M.S., Tabatabaei, S. A., Younesi, E., Rostami, M. R. and Mombeni, M. (2013). Hormone priming improves germination characteristics and enzyme activity of sorghum seeds (*Sorghum bicolor* L.) under accelerated aging. *Cercetari agronomice in Moldova*, 46(3):155.
- Bailly, C. (2004). Active oxygen species and antioxidants in seed biology. *Seed Science Research*, 14(02): 93-107.
- Bailly, C. and Kranner, I. (2011). Analyses of reactive oxygen species and antioxidants in relation to seed longevity and germination. *Seed Dormancy: Methods and Protocols*, 343-367.
- Bailly, C., Benamar, A., Corbineau, F. and Come, D. (1998). Free radical scavenging as affected by accelerated ageing and subsequent priming in sunflower seeds. *Physiologia Plantarum*, 104(4): 646- 652.

- Bailly, C., Benamar, A., Corbineau, F. and Côme, D. (2000). Antioxidant systems in sunflower (*Helianthus annuus* L.) seeds as affected by priming. *Seed Science Research*, 10 (01):35-42.
- Bailly, C., Bogatek-Leszczynska, R., Come, D. and Corbineau, F. (2002). Changes in activities of antioxidant enzymes and lipoxygenase during growth of sunflower seedlings from seeds of different vigour. *Seed Science Research*. 12: 47-55.
- Bailly, C., El-Maarouf-Bouteau, H. and Corbineau, F. (2008). From intracellular signaling networks to cell death: the dual role of reactive oxygen species in seed physiology. *Biological Complications*, 331(10): 806-814.
- Balasubramanian, B., Pogożelski, W. K. and Tullius, T. D. (1998). DNA strand breaking by the hydroxyl radical is governed by the accessible surface areas of the hydrogen atoms of the DNA backbone. *Proceedings of the National Academy of Sciences*, 95(17): 9738-9743.
- Balešević- Tubić, S., Malencic, D., Tatic, M. and Miladinovic, J. (2005). Influence of ageing processes on biochemical changes in sunflower seed. *Helia*, 28(42):107-114.
- Balešević-Tubić, S., Milošević, M., Vujaković, M., Zlokolica, M. and Nikolić, Z. (2000). Seed ageing. *Field Vegetable Crop Research*, 33: 207-212.
- Balešević-Tubić, S., Tatić, M., Đorđević, V., Nikolić, Z. and Đukić, V. (2010). Seed viability of oil crops depending on storage conditions. *Helia*, 33(52): 153-160.
- Barton, L. V. (1943). Effect of moisture fluctuations on the viability of seeds in storage. *Boyce Thompson Institute Plant Research*, 13: 35-45.
- Basra, A. S., Singh, B. and Malik, C.P. (1994) Amelioration of the effects of ageing in onion seeds. *Biologia Plantarum*, 36 (3): 365-371.
- Basra, S. M. A., Pannu, I. A. and Afzal, I. (2003). Evaluation of seedling vigor of hydro and matrimprimed wheat (*Triticum aestivum* L.) seeds. *International Journal of Agriculture and Biology*, 5:121-123.
- Basra, S. M. A., Farooq, M., Tabassam, R. and Ahmad, N. (2005). Physiological and biochemical aspects of pre-sowing seed treatments in fine rice (*Oryza sativa* L.). *Seed Science and Technology*, 33(3): 623-628.
- Basu, R. N. and Choudhary, P. (2005). Partitioning of assimilates in soybean seedlings. *Annals of Agricultural Research*, 11: 285-288.
- Bautista-Banjós, S., Hernandez-Lopez, M., Bosquez-Molina, E. and Wilson, C. L. (2003). Effects of chitosan and plant extracts on growth of *Colletotrichum gloeosporioides*, anthracnose levels and quality of papaya fruit. *Crop Protection*, 22:1087-1092.

- Beevers, H. (1979). Microbodies in higher plants. *Annual Review of Plant Physiology*, 30(1):159-193.
- Berti, M. T. and Johnson, B. L. (2008). Physiological changes during seed development of cuphea. *Field Crops Research*, 106(2): 163-170.
- Bessa, J. F., Donadon, J. R., Resende, O., Alves, R., Sales, J. D. F. and Costa, L. M. (2015). Storage of crambe seeds in different containers and environments: Part I-Physiological quality. *Brazilian Journal of Agricultural and Environmental Engineering*, 19(3): 224-230.
- Bewley, J. D., Bradford, K. J., Hilhorst, H. W. M. and Nonogaki, H. (2013). Seeds: *Physiology of development, germination and dormancy*, 3rdEd. *Seed Science Research*, 23: 289.
- Bewley, J.D. and Black, M. (1994). Seeds: *physiology of development and germination*, 2nd Ed, (pp. 167-171). New York, USA: Plenum Press.
- Bhatia, V. S., Yadav, S., Jumrani, K., and Guruprasad, K. N. (2010). Field deterioration of Soybean seed: Role of oxidative stress and antioxidant defense mechanism. *Journal of Plant Biology*, 32(2): 179-190.
- Biabani, A., Boggs, L.C., Katozi, M., Sabouri, H., (2011). Effects of seed deterioration and inoculation with Mesorhizobium cicerion yield and plant performance of chickpea. *Australian Journal of Crop Science*, 5(1): 66-70.
- Blokhina, O., Virolainen, E. and Fagerstedt, K. V. (2003). Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Annals of Botany*, 91(2): 179-194.
- Borowski, E. and Michalek, S. (2014). The effect of chilling temperature on germination and early growth of domestic and Canadian soybean (*Glycine max* L.) Merr.) cultivars. *Acta Scientiarum Polonorum Hortorum Cultus*, 13: 31-43.
- Braccini, A.L., Albrecht, L.P, Ávila, M.R, Scapim, C.A, Bio, F.E.I and Schuab, S.R.P (2003). Physiological and sanitary quality of the seeds of fifteen soybean cultivars (*Glycine max* (L.) Merrill) harvested in the normal season and after the crop delay. *Acta Scientiarum Agronomy*, 25 (2): 449-457.
- Braga, S.L.O., Luiz, N.J. and de Souza, M.M. (2000). Seed vigor and growth analysis of black oats. *Scientia Agricola*, 57(2): 305-312.
- Bray, C.M. (1995). Biochemical processes during the osmopriming of seeds. In J. Kigel, and G. Galili (Eds.), *Seed development and germination*, (pp. 767-789). New York, USA.
- Bray, C.M., Davison, P.A., Ashraf M. and Taylor, R.M. (1989). Biochemical changes during osmopriming of leek seeds. *Annals of Botany*, 63: 185-193.

- Brits, G. J., Brown, N. A. C., Calitz, F. J. and Van Staden, J. (2015). Effects of storage under low temperature, room temperature and in the soil on viability and vigour of *Leucospermum cordifolium* (Proteaceae) seeds. *South African Journal of Botany*, 97: 1-8.
- Brocklehurst, P. A., Dearman, J. and Drew, R. L. K. (1984). Effects of osmotic priming on seed germination and seedling growth in leek. *Scientia Horticulturae*, 24(3): 201-210.
- Burgass, R.W. and Powell, A. A. (1984). Evidence for repair processes in the invigoration of seed by hydration. *Annals of Botany*, 53: 753-757.
- Burris, J. S. (1973). Effect of seed maturation and plant population on soybean seed quality. *Agronomy Journal*, 65(3): 440-441.
- Burton, J. W. (1997). Soyabean (*Glycine max* (L.) Merr.). *Field Crops Research*, 53(1): 171-186.
- Butler, L. H., Hay, F. R., Ellis, R. H., Smith, R. D. and Murray, T. B. (2009). Priming and re-drying improve the survival of mature seeds of *Digitalis purpurea* during storage. *Annals of Botany*, 103(8): 1261-1270.
- Byrd, H.W. and Delouche, J.C. (1971). Deterioration of soybean seed in storage. *Proceeding of Association of Official Seed Analysts*, 61:41-57.
- Cadet, J., Loft, S., Olinski, R., Evans, M. D., Bialkowski, K., Richard Wagner, J., Dedon, P. C., Møller, P., Greenberg, M. M. and Cooke, M. S. (2012). Biologically relevant oxidants and terminology, classification and nomenclature of oxidatively generated damage to nucleobases and 2-deoxyribose in nucleic acids. *Free Radical Research*, 46(4): 367-381.
- Cantliffe, D.J. (1983). Sowing primed seed. *American Vegetable Grower*, 31: 42-43.
- Capron, I., Corbineau, F., Dacher, F., Job, C., Côme, D. and Job, D. (2000). Sugarbeet seed priming: effects of priming conditions on germination, solubilization of 11-S globulin and accumulation of LEA proteins. *Seed Science Research*, 10(3): 243-254.
- Catão, H. C. R. M., Gomes, L. A. A., Guimarães, R. M., Fonseca, P. H. F., Caixeta, F. and Marodin, J. C. (2016). Physiological and isozyme alterations in lettuce seeds under different conditions and storage periods. *Journal of Seed Science*, 38(4): 305-313.
- Caviness, C. E. (1966). Estimates of natural cross-pollination in Jackson soybeans in Arkansas. *Crop Science*. 6: 211.
- Chachalis, D., and Smith, M. L. (2000). Imbibition behavior of soybean (*Glycine max* (L.) Merrill) accessions with different testa characteristics. *Seed Science and Technology*, 28(2): 321-331.

- Chan, K. L. and Hickson, I. D. (2011). New insights into the formation and resolution of ultra-fine anaphase bridges. In *Seminars in cell & developmental biology* (pp.906-912). Academic Press.
- Chen, H., Osuna, D., Colville, L., Lorenzo, O., Graeber, K., Küster, H., Leubner-Metzger, G. and Kranner, I. (2014). Correction: Transcriptome-Wide Mapping of Pea Seed Ageing Reveals a Pivotal Role for Genes Related to Oxidative Stress and Programmed Cell Death. *PLoS One*, 9(1): 10-1371.
- Chen, K. and Arora, R. (2011). Dynamics of the antioxidant system during seed osmopriming, post-priming germination, and seedling establishment in Spinach (*Spinacia oleracea*). *Plant Science*, 180: 212-220.
- Chen, K., Arora, R. and Arora, U. (2010). Osmopriming of spinach (*Spinacia oleracea* L. cv. Bloomsdale) seeds and germination performance under temperature and water stress. *Seed Science and Technology*, 38(1): 36-48.
- Chen, W., Guo, C., Hussain, S., Zhu, B., Deng, F., Xue, Y., Geng, M. and Wu, L. (2016). Role of xylo-oligosaccharides in protection against salinity-induced adversities in Chinese cabbage. *Environmental Science and Pollution Research*, 23(2): 1254-1264.
- Chibu, H. and Shibayama, H. (2001). Effects of chitosan applications on the growth of several crops. In T. Uragami, K. Kurita, T. Fukamizo (Eds.), *Chitin and chitosan in life science* (pp. 235-239). Yamaguchi, Japan.
- Chiu, K. Y., Chuang, S. J. and Sung, J. M. (2006). Both anti-oxidation and lipid-carbohydrate conversion enhancements are involved in priming-improved emergence of *Echinacea purpurea* seeds that differ in size. *Scientia Horticulturae*, 108(2): 220-226.
- Chiu, K. Y., Wang, C. S. and Sung, J. M. (1995). Lipid peroxidation and peroxide scavenging enzymes associated with accelerated aging and hydration of watermelon seeds differing in ploidy. *Physiologia Plantarum*, 94(3): 441-446.
- Coolbear, P. (1995). Mechanisms of seed deterioration. *Seed quality: Basic Mechanisms and Agricultural Implications* (pp.223-277). Food Product Press New York.
- Copeland, L. O. and McDonald, M. B. (1995). Seed vigour and vigour tests. *Principles of Seed science and Technology* (pp. 153- 180). Chapman and Hill. New York.
- Copeland, L. O. and McDonald, M. B. (2001). Seed vigor and vigor testing. *Principles of Seed Science and Technology* (pp. 165-191). Springer USA.
- Czarnecki, E. and Evans, L. E. (1986). Effect of weathering during delayed harvest on test weight, seed size, and grain hardness of wheat. *Canadian Journal of Plant Science*, 66(3): 473-482.

- Dell'Aquila, A. and Taranto, G. (1986). Cell division and DNA-synthesis during osmopriming treatment and following germination in aged wheat embryos. *Seed Science and Technology*, 14(2): 333-341.
- Dell'Aquila, A., Savino, G. and De Leo, P. (1978). Metabolic changes induced by hydration-dehydration treatment in wheat embryos. *Plant and Cell Physiology*, 19(2): 349-354.
- Delouche, J. C. (1968). Physiology of seed storage. *Proceedings of Corn and Sorghum Research Conference American Trade Association*, 23, Mississippi, USA.
- Delouche, J. C. (1980). Environmental effects on seed development and seed quality. *HortScience*, 15(6):775-780.
- De-Oliveira, A., Gomes-Filho, E., Enéas-Filho, J., Prisco, J. T. and Alencar, N. L. M. (2012). Seed priming effects on growth, lipid peroxidation, and activity of ROS scavenging enzymes in NaCl-stressed sorghum seedlings from aged seeds. *Journal of Plant Interactions*, 7(2): 151-159.
- Dhingra, O.D.; Acuña, R.S. (1997). Pathology of soybean seeds. UFV, Viçosa, MG, Brazil.
- Dias, A.C.P., Reis, M.S., Sediya, C.S., Moreira, M.A. and Rocha, V.S. (2005). Physiological quality of soybean (*Glycine max* (L.) Merrill) seeds without lipoxygenases. *Revista Ceres*, 52: 453-466.
- Diniz, F. O., Reis, M. S., Dias, L. A. D. S., Araújo, E. F., Sediya, T. and Sediya, C. A. (2013). Physiological quality of soybean seeds of cultivars submitted to harvesting delay and its association with seedling emergence in the field. *Journal of Seed Science*, 35(2): 147-152.
- Donà, M., Balestrazzi, A., Mondoni, A., Rossi, G., Ventura, L., Buttafava, A. and Carbonera, D. (2013). DNA profiling, telomere analysis and antioxidant properties as tools for monitoring ex situ seed longevity. *Annals of Botany*, 111(5): 987-998.
- Dourado, A. M. and Roberts, E. H. (1984). Chromosome aberrations induced during storage in barley and pea seeds. *Annals of Botany*, 54(6): 767-779.
- Dowd, M. A., Gaulden, M. E., Proctor, B. L. and Seibert, G. B. (1986). Formaldehyde-induced acentric chromosome fragments and chromosome stickiness in *Chortophaga neuroblasts*. *Environmental and Molecular Mutagenesis*, 8(3): 401-411.
- Edwards, C. J. and Hartwig, E. E. (1971). Effect of seed size upon rate of germination in soybeans. *Agronomy Journal*, 63(3): 429-450.
- Elder, R. H. and Osborne, D. J. (1993). Function of DNA synthesis and DNA repair in the survival of embryos during early germination and in dormancy. *Seed Science Research*, 3(1): 43-53.

- Eskandari, H. (2012). Seed quality variation of crop plants during seed development and maturation. *International journal of Agronomy and Plant Production*, 3 (11): 557-560.
- FAOSTAT (2016). Food and Agriculture Organization of the United Nations, FAOSTAT database.
- Farooq, M., Wahid, A., Kobayashi, N., Fujita, D. and Basra, S.M.A. (2009). Plant drought stress: effects, mechanisms and management. *Agronomy for Sustainable Development*, 29: 185-212.
- Fehr, W. R. and Caviness, C. E. (1977). *Stages of soybean development*. Ames: Iowa State University.
- Fessel, S. A., Vieira, R. D., Cruz, M. C. P. D., Paula, R. C. D. and Panobianco, M. (2006). Electrical conductivity testing of corn seeds as influenced by temperature and period of storage. *Brazilian Agricultural Research*, 41(10): 1551-1559.
- Floris, C. (1970). Ageing in *Triticum durum* seeds: behaviour of embryos and endosperms from aged seeds as revealed by the embryo-transplantation technique. *Journal of Experimental Botany*, 21(2):462-468.
- Franca-Necto, J.B and Henning A.A. (1984). Physiological and sanitary quality of soybean seeds. Londrina, *Embrapa-CNPSo, Technical Circular*, 9.
- França-Neto, J. B., Krzyzanowski, F. C., and Costa, N. D. (1998). The tetrazolium test for soybean seeds. Londrina: *Embrapa-CNPSo, Documentos*, 115: 71.
- Franca-Neto, J.B, Henning, A.A and Krzyzanowski, H.R. (1994). Seed production and technology for the tropics. In Embrapa-Cnpso (Ed.), *Tropical soybean: improvement and production* (pp.217-240). Rome: FAO.
- Fu, J. R., Lu, X. H., Chen, R. Z., Zhang, B. Z., Liu, Z. S., Li, Z. S. and Cai, D. Y. (1988). Osmoconditioning of peanut (*Arachis hypogea* L.) seeds with PEG to improve vigour and some biochemical activities. *Seed Science and Technology*, 16(1): 197-212.
- Fuchs, D., Bitterlich, G.B., Wede, I., Wachter, H., (1997). Reactive oxygen and apoptosis. In J.G. Scandalios (Ed.), *Oxidative stress and the molecular biology of antioxidant defenses* (pp.139-167). Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Gallardo, K., Job, C., Groot, S. P., Puype, M., Demol, H., Vandekerckhove, J. and Job, D. (2001). Proteomic analysis of *Arabidopsis* seed germination and priming. *Plant Physiology*, 126(2): 835-848.
- Gallego, S. M., Benavides, M. P. and Tomaro, M. L. (1996). Effect of heavy metal ion excess on sunflower leaves: evidence for involvement of oxidative stress. *Plant Science*, 121(2): 151-159.

- Gaulden, M.E. (1987). Hypothesis: some mutagens directly alter specific chromosomal proteins (DNA topoisomerase II and peripheral proteins) to produce chromosome stickiness, which causes chromosome aberrations. *Mutagenesis*, 2: 357-365.
- Ghasemnezhad, A. and Honermeier, B. (2007). Influence of storage conditions on quality and viability of high and low oleic sunflower seeds. *International Journal of Plant Production*, 3(4): 41-50.
- Ghassemi-Golezani, K., Khomari, S., Valizadeh, M. and Alyari, H. (2008). Effect of seed vigor and the duration of cold acclimation on freezing tolerance of winter oilseed rape. *Seed Science and Technology*, 36:767-775.
- Ghassemi-Golezani, K., Lotfi, R., and Norouzi, M. (2012). Seed quality of soybean cultivars affected by pod position and water stress at reproductive stages. *International Journal of Plant, Animal and Environmental Sciences*, 3: 119-125.
- Ghassemi-Golezani, K., Mousabeygi, T., Yagoob, R. A. E. I., and Aharizad, S. (2010). Effects of water stress and pod position on the seed quality of chickpea (*Cicer arietinum* L.) cultivars. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 38(1): 114.
- Gill S. S. and Tuteja N. (2010). Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology and Biochemistry*. 48: 909-930.
- Giri, G.S. and Schillinger, W. F. (2003). Seed priming winter wheat for germination, emergence and yield. *Crop Science*, 43: 2135-2141.
- Goel, A. and Sheoran, L.S. (2003). Lipid peroxidation and peroxide scavenging enzymes in cotton seeds under natural ageing. *Biologia Plantarum*, 46: 429-434.
- Grant, W. F. (1982). Cytogenetic studies of agricultural chemicals in plants. In R. A. Fleck et al. (Eds), *Genetic Toxicology*. (pp. 353-378). Plenum Press, New York.
- Green, D. E., Pinnell, E. L. and Cavanaugh, L. E. (1966). Effect of seed moisture content, field weathering, and combine cylinder speed on soybean quality. *Crop Science*, 6: 7-10.
- Groot, S. P. C., Surki, A. A., De Vos, R. C. H. and Kodde, J. (2012). Seed storage at elevated partial pressure of oxygen, a fast method for analysing seed ageing under dry conditions. *Annals of Botany*, 110(6): 1149-1159.
- Guan, Y. J., Hu, J., Wang, X. J. and Shao, C. X. (2009). Seed priming with chitosan improves maize germination and seedling growth in relation to physiological changes under low temperature stress. *Journal of Zhejiang University Science B*, 10(6): 427-433.

- Guo, Z., Xing, R., Liu, S., Yu, H., Wang, P., Li, C., and Li, P. (2005). The synthesis and antioxidant activity of the Schiff bases of chitosan and carboxymethyl chitosan. *Bioorganic and Medicinal Chemistry Letters*, 15(20): 4600-4603.
- Gupta, A. S., Webb, R. P., Holaday, A. S. and Allen, R. D. (1993). Overexpression of superoxide dismutase protects plants from oxidative stress (induction of ascorbate peroxidase in superoxide dismutase-overexpressing plants). *Plant Physiology*, 103(4): 1067-1073.
- Gupta, A., Dadlani, M., Kumar, M. B. A., Roy, M., Naseem, M., Choudhary, V. K. and Maiti, R. K. (2008). Seed priming: the aftermath. *International Journal of Agriculture Environment and Biotechnology*, 1(4): 199-209.
- Gurusinghe, S. H., Cheng, Z. and Bradford, K. J. (1999). Cell cycle activity during seed priming is not essential for germination advancement in tomato. *Journal of Experimental Botany*, 101-106.
- Gutema, E. (2006). *The effect of canopy architecture and seasonal variation on several qualities attributes in soybean (Glycine max L. Merr.)*. PhD Thesis, University Putra Malaysia.
- Hadwiger, L.A., Klosterman S.J. and Choi, J.J. (2002). The mode of action of chitosan and its oligomers in inducing plant promoters and developing disease resistance in plants, *Advances in Chitin Science*, 5: 452-457.
- Hamidi, R. and Pirasteh-Anosheh, H. (2013). Comparison effect of different seed priming methods on sunflower germination and seedling growth. *International Journal of Agronomy and Plant Production*, 4: 1247-1250.
- Hanson, A. D. (1973). The effects of imbibition drying treatments on wheat seeds. *New Phytologist*, 72(5): 1063-1073.
- Harrington, J.F. (1959). The value of moisture resistant containers in vegetable seed packaging. *California Agricultural Experiment Station*, 792: 1-23.
- Harrington, J.F. (1972). Seed storage and longevity. In T.T. Kozloswki (Ed.), *Seed Biology* (pp.145-245). New York: New York Academic Press.
- Harrington, J.F. (1973). Biochemical basis of seed longevity. *Seed Science and Technology*. 1: 453-461.
- Hasanuzzaman, M., Nahar, K. and Fujita, M. (2013). Extreme temperatures, oxidative stress and antioxidant defense in plants, In K.Vahdati and C. Leslie (Eds.), *Abiotic Stress-Plant Response and Applications in Agriculture*, (pp.169-205). Rijeka : INTECH Open Access Publisher.
- Hendry, G. A. (1993). Oxygen, free radical processes and seed longevity. *Seed Science Research*, 3(03): 141-153.

- Henning, F. A., Mertz, L. M., Jacob Junior, E. A., Machado, R. D., Fiss, G. and Zimmer, P. D. (2010). Chemical composition and mobilization of reserves in high and low vigor soybean seeds. *Bragantia*, 69(3): 727-734.
- Hosamani, J., Kumar, M. B., Talukdar, A., Lal, S. K. and Dadlani, M. (2013). Molecular characterization and identification of candidate markers for seed longevity in soybean [*Glycine max* (L.) Merrill]. *Indian Journal of Genetics and Plant Breeding*, 73(1): 64-71.
- Howell, R. W., Collins, F. I. and Sedgwick, V. E. (1959). Respiration of soybean seeds as related to weathering losses during ripening. *Agronomy Journal*, 51(11):677-679.
- Hu, D., Ma, G., Wang, Q., Yao, J., Wang, Y. U., Pritchard, H. W. and Wang, X. (2012). Spatial and temporal nature of reactive oxygen species production and programmed cell death in elm (*Ulmus pumila* L.) seeds during controlled deterioration. *Plant, Cell and Environment*, 35(11): 2045-2059.
- Huang, R., Mendis, E., and Kim, S. K. (2005). Factors affecting the free radical scavenging behavior of chitosan sulfate. *International Journal of Biological Macromolecules*, 36(1): 120-127.
- Huang, Z., Boubriak, I., Osborne, D. J., Dong, M. and Gutterman, Y. (2008). Possible role of pectin-containing mucilage and dew in repairing embryo DNA of seeds adapted to desert conditions. *Annals of Botany*, 101(2): 277-283.
- Hussain, S., Zheng, M., Khan, F., Khaliq, A., Fahad, S., Peng, S., ... and Nie, L. (2015). Benefits of rice seed priming are offset permanently by prolonged storage and the storage conditions. *Scientific Reports*, 5: 8101.
- Illipronti, R. A., Lommen, W. J. M., Langerak, C. J. and Struik, P. C. (2000). Time of pod set and seed position on the plant contribute to variation in quality of seeds within soybean seed lots. *NJAS-Wageningen Journal of Life Sciences*, 48(2): 165-180.
- Isely, D. (1957). Vigor tests. *Proceeding of Association of Official Seed Analyses*, 47:176-182.
- ISTA. (1999). International rules for seed testing. *Seed Science and Technology*, 27: 340.
- ISTA. (2006). *International rules for seed testing*, International Seed Testing Association, Bassersdorf, Switzerland.
- Jabeen., N. and Ahmad, R. (2013). The activity of antioxidant enzymes in response to salt stress in safflower (*Carthamus tinctorius* L.) and sunflower (*Helianthus annuus* L.) seedlings raised from seed treated with chitosan. *Journal of the Science of Food and Agriculture*, 93: 1699-1705.

- Jacoby, R. P., Li, L., Huang, S., Pong Lee, C., Millar, A. H. and Taylor, N. L. (2012). Mitochondrial Composition, Function and Stress Response in Plants . *Journal of Integrative Plant Biology*, 54(11): 887-906.
- Jafari, H., Najafi, R., Soltani, A., Dastan, S., 2013. Seed deterioration and seedling germination of soybean (*Glycine max* L.) cv. Sahar. *International Journal of Agronomy and Plant Production*, 4: 3762-3766.
- Jisha, K. C., Vijayakumari, K. and Puthur, J. T. (2012). Seed priming for abiotic stress tolerance: an overview. *Acta Physiologiae Plantarum*, 35(5): 1381-1396.
- Johnson, R. R. and Wax, L. M. (1978). Relationship of soybean germination and vigor tests to field performance. *Agronomy Journal*, 70(2): 273-278.
- Johnson, W. H. (1959). Efficiency in combining wheat. *Agricultural Engineering*, 40: 16-29.
- Jones, S.I., Gonzalez, D.O. and Vodkin, L.O. (2010). Flux of transcript patterns during soybean seed development. *BMC Genomics*, 11:136.
- Justice, O. L. and Bass, L. N. (1978). *Principles and practices of seed storage* (No. 506). US Department of Agriculture.
- Jyoti and Malik C.P . (2013). Seed deterioration: A review. *International Journal of Life science and Pharma Reviews*, 2 (3): 374-385.
- Kaewnaee, P., Vichitphan, S., Klanrit, P., Siri, B. and Vichitphan, K. (2011). Effect of accelerated aging process on seed quality and biochemical changes in sweet pepper (*Capsicum annuum* Linn.) seeds. *Biotechnology*, 10(2): 175-182.
- Kandil, A. A., Sharief, A. E. and Sheteiwy, M. S. (2013). Effect of seed storage periods, conditions and materials on germination of some soybean seed cultivars. *American Journal of Experimental Agriculture*, 3(4): 1020.
- Kang, H.M. and Saltveit, E. (2002). Effect of chilling on antioxidant enzymes and DPPH-radical scavenging activity of high- and low-vigour cucumber seedling radicles. *Plant Cell Environment*, 25: 1233-1238.
- Kannababu, N. and Karivaratharaju T. V. (2001). Glutamate dehydrogenase activity in seedling organs of sunflower. *Seed Research*, 29(2): 167-170.
- Kanwar, R., Mehta, D.K and Lal, M. (2014). Effect of seed priming on physiological parameters of aged and non-aged seeds of bitter gourd, *Momordica charantia* L. *International Journal of Farm Sciences*, 4(3): 24-32.
- Kapilan, R. (2015). Accelerated aging declines the germination characteristics of the maize seeds. *Scholars Academic Journal of Biosciences*, 3(8):708-711.
- Kapoor, N., Arya, A., Siddiqui, M. A., Amir, A. and Kumar, H. (2010). Seed deterioration in chickpea (*Cicer arietinum* L.) under accelerated ageing. *Asian Journal of Plant Sciences*, 9(3): 158-162.

- Kapoor, N., Arya, A., Siddiqui, M.A., Kumar, H. and Amir, A. (2011). Physiology and biochemical changes during seed deterioration in aged seeds of rice (*Oryza sativa* L.) *American Journal of Plant Physiology*, 6 (1): 28-35.
- Kaur, S., Gupta, A. K. and Kaur, N. (2005). Seed priming increases crop yield possibly by modulating enzymes of sucrose metabolism in chickpea. *Journal of Agronomy and Crop Science*, 191(2): 81-87.
- Kausar, M., Mahmood, T., Basra, S. M. A. and Arshad, M. (2009). Invigoration of low vigor sunflower hybrids by seed priming. *International Journal of Agricultural Biology*, 11(5): 521-528.
- Kaushal, N., Gupta, K., Bhandhari, K., Kumar, S., Thakur, P. and Nayyar, H. (2011). Proline induces heat tolerance in chickpea (*Cicer arietinum* L.) plants by protecting vital enzymes of carbon and antioxidative metabolism. *Physiology and Molecular Biology of Plants*, 17: 203-213.
- Kaya, M. D., Okçu, G., Atak, M., Çıkılı, Y. and Kolsarıcı, Ö. (2006). Seed treatments to overcome salt and drought stress during germination in sunflower (*Helianthus annuus* L.). *European Journal of Agronomy*, 24(4): 291-295.
- Kaya, M., Kaya, G., Kaya, M. D., Atak, M., Sağlam, S., Khawar, K. M. and Ciftci, C.Y. (2008). Interaction between seed size and NaCl on germination and early seedling growth of some Turkish cultivars of chickpea (*Cicer arietinum* L.). *Journal of Zhejiang University Agricultural Science B*, 9 (5):371-377.
- Keigley, P. J. and Mullen, R. E. (1986). Changes in soybean seed quality from high temperature during seed fill and maturation. *Crop Science*, 26(6): 1212-1216.
- Khajeh-Hosseini, M., Powell, A. A. and Bingham, I. J. (2003). The interaction between salinity stress and seed vigour during germination of soyabean seeds. *Seed Science and Technology*, 31(3): 715-725.
- Khan, A. A., Peck, N. H. and Samimy, C. (1980). Seed osmoconditioning: physiological and biochemical changes. *Israel Journal of Botany*, 29(1-4): 133-144.
- Khan, M. M., Iqbal, M. J., Abbas, M. and Usman, M. (2003). Effect of Ageing on Viability, Vigour and Chromosomal Damage in Pea (*Pisum sativum* l.) seeds. *Pakistan Journal of Agricultural Science*, 40: 50-54.
- Khatun, A., Kabir, G. and Bhuiyan, M. A. H. (2009). Effect of harvesting stages on the seed quality of lentil (*Lens culinaris* L.) during storage. *Bangladesh Journal of Agricultural Research*, 34(4): 565-576.
- Kibinza, S., Bazin, J., Bailly, C., Farrant, J. M., Corbineau, F. and El-Maarouf-Bouteau, H., (2011). Catalase is a key enzyme in seed recovery from ageing during priming. *Plant Science*, 181(3): 309-315.
- Kibinza, S., Vinel, D., Côme, D., Bailly, C. and Corbineau, F. (2006). Sunflower seed deterioration as related to moisture content during ageing, energy metabolism

- and active oxygen species scavenging. *Physiologia Plantarum*, 128(3): 496-506.
- Kim, K. W. and Thomas, R. L. (2007). Antioxidative activity of chitosans with varying molecular weights. *Food Chemistry*, 101(1): 308-313.
- Kmetz, K. T., Schmitthenner, A. F. and Ellett, C. W. (1978). Soybean seed decay: Prevalence of infection and symptom expression caused by *Phomopsis* sp., *Diaporthe phaseolorum* var. *sojae*, and *D. phaseolorum* var. *caulivora*. *Phytopathology*, 68(6): 836-840.
- Kong, F., Chang, S. K. C., Liu, Z. and Wilson, L. A. (2008). Changes of soybean quality during storage as related to soymilk and tofu making. *Journal of Food Science*, 73(3):134-144.
- Koning, G., TeKrony, D. M., Pfeiffer, T. W. and Ghabrial, S. A. (2001). Infection of Soybean with Increases Susceptibility to spp. Seed Infection. *Crop Science*, 41(6): 1850-1856.
- Krainart, C., Siri, B. and Vichitphan, K. (2015). Effects of accelerated ageing and subsequent priming on seed quality and biochemical change of hybrid cucumber (*Cucumis sativa* Linn.) seeds. *International Journal of Agricultural Technology*, 11(1): 165-179.
- Kranner, I., Chen, H., Pritchard, H. W., Pearce, S. R. and Birtić, S. (2011). Inter-nucleosomal DNA fragmentation and loss of RNA integrity during seed ageing. *Plant Growth Regulation*, 63(1): 63-72.
- Krittigamas, N., Vearasilp, S., Thanapornpoonpong, S., Suriyong, S., Pa-oblek, S., Pawelzik, E. and Becker, M. (2001). Investigation of Post-harvest Soybean Seed Storability after Passing the Different Steps of Processing. Conference on International Agricultural Research for Development. Deutscher Tropentag-Bonn, 9-11.
- Krober, O. A. and Collins, F. I. (1948). Effect of weather damage on the chemical composition of soybeans. *Journal of the American Oil Chemists' Society*, 25(8): 296-298.
- Krzyzanowski, F.C., Franca-Neto, J.B., Henning, A.A. and Costa, N.P. (2008). Soybean seed and technology as the basis for high productivity: series seeds. *Embrapa Soja, Technical Circular*, 55.
- Kuchlan, M.K. (2006). *Identification of physical, physiological and biochemical factors and molecular analysis for longevity of soybean seeds*. PhD Thesis, Indian Agricultural Research Institute, New Delhi.
- Kueneman, E. A. (1983). Genetic control of seed longevity in soybeans. *Crop Science*, 23(1): 5-8.
- Kumar Vasant T., (1993). *Deterioration of chilli seeds under accelerated ageing conditions*, PhD Thesis, Genetics, IARI, New Delhi;

- Kuo, W. H. (1989). Delayed-permeability of soybean seeds: characteristics and screening methodology. *Seed Science and Technology*, 17: 134-142.
- Lanteri, S., Nada, E., Belletti, P., Quagliotti, L. and Bino, R. J. (1996). Effects of controlled deterioration and osmoconditioning on germination and nuclear replication in seeds of pepper (*Capsicum annuum* L.). *Annals of Botany*, 77(6): 591-597.
- Lanteri, S., Saracco, F., Kraak, H. L. and Bino, R. J. (1994). The effects of priming on nuclear replication activity and germination of pepper (*Capsicum annuum*) and tomato (*Lycopersicon esculentum*) seeds. *Seed Science Research*, 4(02): 81-87.
- Lehner, A., Mamadou, N. Poels, P. Côme, D. Bailly, C. and Corbineau, F. (2008). Changes in soluble carbohydrates, lipid peroxidation and antioxidant enzyme activities in the embryo during ageing in wheat grains. *Journal of Cereal Science*, 47: 555-565.
- Leubner-Metzger, G. (2005). β -1, 3-Glucanase gene expression in low-hydrated seeds as a mechanism for dormancy release during tobacco after-ripening. *The Plant Journal*, 41(1): 133-145.
- Li, Y., Liu, Y. and Zhang, J. (2010). Advances in the research on the AsA-GSH cycle in horticultural crops. *Frontiers of Agriculture in China*, 4(1): 84-90.
- Lima, W. A. A., Borem, A., Dias, D. C. F. S., Moreira, M. A., Dias, L. A. S. and Piovesan, N. D. (2007). Harvest delay as a method of differentiation of soybean genotypes for seed quality. *Brazilian Journal of Seeds*, 29(1):186-192.
- Lindhahl, T. (1993). Instability and decay of the primary structure of DNA. *Nature*, 362(6422): 709-715.
- Lins, S. R. D. O., Moreira de Carvalho, M. L., das Gracas Cardoso, M., Miranda, D. H. and de Andrade, J. (2014). Physiological, enzymatic, and microstructural analyses of sunflower seeds during storage. *Australian Journal of Crop Science*, 8(7): 1038.
- Lü, J. M., Lin, P. H., Yao, Q. and Chen, C. (2010). Chemical and molecular mechanisms of antioxidants: experimental approaches and model systems. *Journal of Cellular and Molecular Medicine*, 14(4): 840-860.
- Mahdavi, B., Modarres-Sanavy, S. A. M., Aghaalikhani, M., Sharifi, M. and Dolatabadian, A. (2011). Chitosan improves osmotic potential tolerance in safflower (*Carthamus tinctorius* L.) seedlings. *Journal of Crop Improvement*, 25(6): 728-741.
- Malaysia Weather-AccuWeather.com. Retrieved from <http://www.accuweather.com/en/my/serdang/228381/may-weather/228381?monyr=5/1/2016&view=table>.

- Mansouri-Far, C., Goodarzian-Ghahfarokhi, M., Saeidi, M. and Abdoli, M. (2015). Antioxidant enzyme activity and germination characteristics of different maize hybrid seeds during ageing. *Environmental and Experimental Biology*. 13: 177-182.
- Marcano, L., Carruyo, I., Del Campo, A. and Montiel, X. (2004). Cytotoxicity and mode of action of maleic hydrazide in root tips of *Allium cepa* L. *Environmental Research*, 94(2): 221-226.
- Marcos-Filho, J., Chamma, H. M. C. P., Casagrande, J. R. R., Marcos, E.A. and Regitano-d'arce, A. B. (1994). Effect of harvesting time on seed physiological quality, chemical composition and storability of soybean. *Scientia Agricola. Piracicaba*, 5(2): 298-304.
- Maroufi, K., Farahani, H. A. and Moradi, O. (2011). Increasing of seedling vigour by hydro priming method in cowpea (*Vigna sinensis* L.). *Advances in Environmental Biology*, 5(11): 3668-3672.
- Marwanto, M. (2004). Soybean seed coat characteristics and its quality losses during incubator aging and storage. *Journal of the Association of Physicians of India*, 6 (2): 57-65.
- Mayne, R. Y., Harper, G. A., Franz, A. O., Lee, L. S. and Goldblatt, L. A. (1969). Retardation of the elaboration of aflatoxin in cottonseed by impermeability of the seedcoats. *Crop Science*, 9(2): 147-150.
- Mbofung, G. C., Goggi, A. S., Leandro, L. F. and Mullen, R. E. (2013). Effects of storage temperature and relative humidity on viability and vigor of treated soybean seeds. *Crop Science*, 53(3): 1086-1095.
- McDonald M.B. (2004). Orthodox seed deterioration and its repair. In R. L. Benech-Arnold, and R.A. Sanchez (Eds.), *Handbook of Seed Physiology: Applications to Agriculture* (pp. 273-304), Food Products Press, New York.
- McDonald, M. B. (1999). Seed deterioration: physiology, repair and assessment. *Seed Science and Technology*, 27:177-237.
- McDonald, M.B. (2000). Seed priming. In M. Black, J. D. Bewley (Eds), *Seed technology and its biological basis* (pp. 287-325). Sheffield: Sheffield Academic Press, UK.
- McGee, D. C. (1986). Prediction of *Phomopsis* seed decay by measuring soybean pod infection. *Plant Disease*, 70:329-333.
- Menezes, V. O., Lopes, S. J., Tedesco, S. B., Henning, F. A., Zen, H. D. and Mertz, L. M. (2014). Cytogenetic analysis of wheat seeds submitted to artificial aging stress. *Journal of Seed Science*, 36(1): 71-78.
- Mengistu, A. and Heatherly, L. G. (2006). Planting date, irrigation, maturity group, year, and environment effects on *Phomopsis longicolla*, seed germination, and

seed health rating of soybean in the early soybean production system of the midsouthern USA. *Crop Protection*, 25(4): 310-317.

- Mengistu, A., Castlebury, L. A., Rossman, A. Y., Smith, J. R. and Reddy, K. N. (2007). Isolates of *Diaporthe-Phomopsis* from weeds and their effect on soybean. *Canadian Journal of Plant Pathology*, 29(3): 283-289.
- Mengistu, A., Castlebury, L., Smith, R., Ray, J. and Bellaloui, N. (2009). Seasonal progress of *Phomopsis longicolla* infection on soybean plant parts and its relationship to seed quality. *Plant Disease*, 93(10): 1009-1018.
- Metz, G. L., Green, D. E. and Shibles, R. M. (1985). Reproductive duration and date of maturity in populations of three wide soybean crosses. *Crop Science*, 25 (1): 171-176.
- Michel, B. E. (1983). Evaluation of the water potentials of solutions of polyethylene glycol 8000 both in the absence and presence of other solutes. *Plant Physiology*, 72(1): 66-70.
- Miranda, G., Souza, P., Moreira, C. and Spehar, C.R (1996). Harvest time effects and mechanical threshing on the physical and physiological quality of soybean seeds. *Ceres Journal*, 43 (249): 663-73.
- Moghanibashi, M., Karimmojeni, H., Nikneshan, P. and Behrozi, D. (2012). Effect of hydropriming on seed germination indices of sunflower (*Helianthus annuus* L.) under salt and drought conditions. *Plant Knowledge Journal*, 1(1): 10-15.
- Mohamed-Yasseen, Y., Barringer, S. A., Splittstoesser, W. E. and Costanza, S. (1994). The role of seed coats in seed viability. *The Botanical Review*, 60(4): 426-439.
- Mohammadi, H., Soltani, A., Sadeghipour, H. R. and Zeinali, E. (2012). Effects of seed aging on subsequent seed reserve utilization and seedling growth in soybean. *International Journal of Plant Production*, 5(1): 65-70.
- Møller, I. M. (2001). Plant mitochondria and oxidative stress: electron transport, NADPH turnover, and metabolism of reactive oxygen species. *Annual Review of Plant Biology*, 52(1): 561-591.
- Møller, I.M., Jensen, P.E. and Hansson, A. (2007). Oxidative modifications to cellular components in plants. *Annual Review of Plant Biology*, 58: 459-481.
- Moore, R. P. E. (1985). *Handbook on tetrazolium testing*. Zurich, Switzerland: International Seed Testing Association.
- Moore, S. H., Halloin, J. M. and Yaklich, R. W. (2000). Genetic resistance to seed deterioration. *Genetic improvement of seed quality. Proceedings of a Symposium sponsored by Divisions C-4, C-1, and C-8 of the Crop Science Society of America* (pp. 21-37). Anaheim, California, USA.

- Moosavi, A., Tavakkol Afshari, R., Sharif-Zadeh, F. and Aynehband, A. (2009). Seed priming to increase salt and drought stress tolerance during germination in cultivated species of Amaranth. *Seed Science and Technology*, 37(3): 781-785.
- Mourad, A. L., Neto, A. R., de Oliveira Miguel, A. M. R., Henriques, E. A. and Alves, R. M. V. (2016). Storage stability of three genotypes of sunflower seeds. *Industrial Crops and Products*, 80: 6-16.
- Murata, M. (1979). *Genetic changes induced by artificial ageing in barley*. PhD Thesis. Colorado State University.
- Murata, M., Roos, E.E. and Tsuchiya, T. (1979). Relationship between loss of germinability and the occurrence of chromosomal aberrations in artificially aged seeds of barley. *Barley Genetic Newsletter*, 9: 65-67.
- Murata, M., Tsuchiya, T. and Roos, E. E. (1982). Chromosome damage induced by artificial seed aging in barley. II. Types of chromosomal aberrations at first mitosis. *Botanical Gazette*, 143(1): 111-116.
- Murthy, U. N., Kumar, P. P. and Sun, W. Q. (2003). Mechanisms of seed ageing under different storage conditions for *Vigna radiata* (L.) Wilczek: lipid peroxidation, sugar hydrolysis, Maillard reactions and their relationship to glass state transition. *Journal of Experimental Botany*, 54(384): 1057-1067.
- Muthiah, S., Longer, D. E. and Harris, W. M. (1994). Staging soybean seedling growth from germination to emergence. *Crop Science*, 34(1): 289-291.
- Nagel, M. and Börner, A. (2010). The longevity of crop seeds stored under ambient conditions. *Seed Science Research*, 20(1): 1.
- Nakagawa, J. (1999). Vigor tests based on the performance of seedlings. In F.C Krzyzanowski, R. D. Veira and J. B. Francca-Neto (Eds.), *Seed Vigor: Concepts and Testing* (pp.1-24). Londrin, Abrates.
- Nangju, D. (1977). Effect of date of harvest on seed quality and viability of soya beans. *The Journal of Agricultural Science*, 89(1): 107-112.
- Nawaz, J., Hussain, M., Jabbar, A., Nadeem, G. A., Sajid, M., Subtain, M. U. and Shabbir, I. (2013). Seed Priming a technique. *International Journal of Agriculture and Crop Sciences*, 6(20): 1373.
- Nedeva, D. and Nikolova, A. (1999). Fresh and dry weight changes and germination capacity of natural or premature desiccated developing wheat seeds. *Bulgarian Journal Plant Physiology*, 25(1-2): 3-15.
- Nichols, C.J. (1941). Spontaneous chromosome aberrations in *Allium*. *Genetics*, 26: 89-100.
- Noctor, G., De Paepe, R. and Foyer, C. H. (2007). Mitochondrial redox biology and homeostasis in plants. *Trends in Plant Science*, 12(3): 125-134.

- Obendorf, R. L., Ashworth, E. N. and Rytko, G. T. (1980). Influence of seed maturation on germinability in soybean. *Crop Science*, 20(4): 483-486.
- Oerke, E-C., Dehne, H-W., Schonbeck, F. and abd Weber, A. (1994). Crop Production and Crop Protection, *Estimated loss in major food and cash crops*. Amsterdam, The Netherlands Elsevier Science.
- Osborne, D. J. (1983). Biochemical control systems operating in the early hours of germination. *Canadian Journal of Botany*, 61(12): 3568-3577.
- Owolade, O. F., Olasoji, J. O. and Afolabi, C. G. (2011). Effect of storage temperature and packaging materials on seed germination and seed-borne fungi of sorghum (*Sorghum bicolor* (L.) Moench.) in South West Nigeria. *African Journal of Plant Science*, 5(15): 873-877.
- Pádua, G. P. D., França-Neto, J. D. B., Carvalho, M. L. M. D., Krzyzanowski, F. C. and Guimarães, R. M. (2009). Incidence of green soybean seeds as a function of environmental stresses during seed maturation. *Brazilian Journal of Seeds*, 31(3):150-159.
- Parera, C. A. and Cantliffe, D. J. (1994). Presowing seed priming. *Horticultural Reviews*, 16(16): 109-141.
- Park, E., Choi, Y. S., Jeong, J. Y. and Lee, S. S. (1999). Effect of priming on germination of aged soybean seeds. *Korean Journal of Crop Science*, 44(1): 74-77.
- Park, S. J., Huang, Y. and Ayoubi, P. (2006). Identification of expression profiles of sorghum genes in response to greenbug phloem-feeding using cDNA subtraction and microarray analysis. *Planta*, 223(5): 932-947.
- Prakash, A. (2008). Competitive Commercial Agriculture in Sub-Saharan Africa (CCAA) Study.
- Parkhey, S., Naithani, S. C and Keshavkant, S. (2012). ROS production and lipid catabolism in desiccating *Shorea robusta* seeds during aging. *Plant Physiology and Biochemistry*, 57: 261-267.
- Parrish, D. J. and Leopold, A. C. (1978). On the mechanism of aging in soybean seeds. *Plant Physiology*, 61(3):365-368.
- Paschal, E. H., and Ellis, M. A. (1978). Variation in seed quality characteristics of tropically grown soybeans. *Crop Science*, 18(5): 837-840.
- Pasquini, S., Mizzau, M., Petrusa, E., Braidot, E., Patui, S., Gorian, F., ... and Vianello, A. (2012). Seed storage in polyethylene bags of a recalcitrant species (*Quercus ilex*): analysis of some bio-energetic and oxidative parameters. *Acta Physiologiae Plantarum*, 34(5): 1963-1974.

- Patil B.C., Bhat G.I., A. (1992). Comparative study of MH and EMS in induction of chromosomal aberrations on lateral root meristem in *Clitoria ternate* L. *Cytologia*, 57: 259-264.
- Patil, V. N. and Dadlani, M. (2009). Tetrazolium test for seed viability and vigour. *Handbook of Seed Testing*, 209-241.
- Pellinen, R. I., Korhonen, M. S., Tauriainen, A. A., Palva, E. T. and Kangasjärvi, J. (2002). Hydrogen peroxide activates cell death and defense gene expression in birch. *Plant Physiology*, 130(2):549-560.
- Pereira, W. A., Pereira, S. M. A. and Dias, D. C. F. D. S. (2013). Influence of seed size and water restriction on germination of soybean seeds and on early development of seedlings. *Journal of Seed Science*, 35(3): 316-322.
- Pereira, W. A., Pereira, S. M. A. and Dias, D. C. F. D. S. (2015). Dynamics of reserves of soybean seeds during the development of seedlings of different commercial cultivars. *Journal of Seed Science*, 37(1): 63-69.
- Philbrook, B.D. and Oplinger, E.S. (1989). Soybean field losses as influenced by harvest delays. *Agronomy Journal*, 81(2): 251-258.
- Pool, M., Patterson, F. L. and Bode, C. E. (1958). Effect of Delayed Harvest on Quality of Soft Red Winter Wheat 1. *Agronomy Journal*, 50(5): 271-275.
- Popovic, B. (2006). *The influence of γ -radiation to the antioxidant system and the genotypes of soybean oxidative stress*. PhD Thesis, University of Novi Sad.
- Powell, A. A. and Matthews, S. (1980). The significance of damage during imbibition to the field emergence of pea (*Pisum sativum* L.) seeds. *The Journal of Agricultural Science*, 95(01): 35-38.
- Prashanth, K. V. H., Dharmesh, S. M., Rao, K. S. J. and Tharanathan, R. N. (2007). Free radical-induced chitosan depolymerized products protect calf thymus DNA from oxidative damage. *Carbohydrate Research*, 342(2): 190-195.
- Priestley, D. A. (1986). *Seed aging: implications for seed storage and persistence in the soil*. Comstock Associates. Ithaca: Cornell University Press.
- Probst, A.H. and Judd, R.W. (1973). Origin, US history and development, and world distribution. In B.E. Caldwell (Ed.), *Soybeans: improvement, production, and uses* (pp. 1-15). American Society of Agronomy, Madison, WI.
- Pukacka, S. (1991). Changes in membrane lipid components and antioxidant levels during natural ageing of seeds of *Acer platanoides*. *Physiologia Plantarum*, 82(2): 306-310.
- Pukacka, S. and Ratajczak, E. (2006). Antioxidative response of ascorbate–glutathione pathway enzymes and metabolites to desiccation of recalcitrant *Acer saccharinum* seeds. *Journal of Plant Physiology*, 163(12): 1259-1266.

- Pukacka, S. and Ratajczak, E. (2007). Age-related biochemical changes during storage of beech (*Fagus sylvatica* L.) seeds. *Seed Science Research*, 17(1): 45-53.
- Pushman, F. M. (1975). The effects of alteration of grain moisture content by wetting or drying on the test weight of four winter wheats. *The Journal of Agricultural Science*, 84(01): 187-190.
- Puteh, A. B., Tekrony, D. M. and Egli, D. B. (1997). Expression of cotyledon necrosis in deteriorating soybean (*Glycine max* (L.) Merrill] seed. *Seed Science and Technology*, 25(1): 133-145.
- Quiles, M. J. and López, N. I. (2004). Photoinhibition of photosystems I and II induced by exposure to high light intensity during oat plant grown effects on the chloroplastic NADH dehydrogenase complex, *Plant Science*, 166: 815-823.
- Raiesi, S., Puteh, A. B., Sijam, K. B., and Abdullah, N. A. P. (2011). Seed quality of soybean in relation to *Phomopsis* seed decay in Malaysia. *Asian Journal of Plant Pathology*, 5(1): 28-36.
- Rajjou, L., Lovigny, Y., Groot, S. P., Belghazi, M., Job, C. and Job, D. (2008). Proteome-wide characterization of seed aging in Arabidopsis: a comparison between artificial and natural aging protocols. *Plant Physiology*, 148(1): 620-641.
- Ramseur, E. L., Wallace, S. U. and Quisenberry, V. L. (1984). Distribution pattern of yield components in 'Braxton' soybeans. *Agronomy Journal* 76: 493-497.
- Rao, N. K., Roberts, E. H. and Ellis, R. H. (1987). Loss of viability in lettuce seeds and the accumulation of chromosome damage under different storage conditions. *Annals of Botany*, 60(1): 85-96.
- Rao, N.K. (1986). *Chromosomal aberration and gene mutations induced in lettuce (Lactuca sativa L.) seeds during storage*. PhD Thesis, Reading University, UK.
- Rao, R.G.S., Singh, P.M. and Mathura Rai. (2006). Storability of onion seeds and effects of packaging and storage conditions on viability and vigour. *Scientia Horticulture*, 110: 1-6.
- Ray, P. D., Huang, B. W. and Tsuji, Y. (2012). Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cellular Signalling*, 24(5): 981-990.
- Roberts, E. H. and Ellis, R. H. (1989). Water and seed survival. *Annals of Botany*, 63(1): 39-39.
- Samarah, N.H., Mullen, R.E., Goggi, S. and Gaul, A. (2009). Effect of drying treatment and temperature on soybean seed quality during maturation. *Seed Science and Technology*, 37: 469-473.

- Sanadhya, A. and Dubeya, W. (2014). Impacts of global climate changes on food security in India, A Review. *Octa Journal of Environmental Research*, 2(2): 121-126.
- Saracco, F., Bino, R. J., Bergervoet, J. H. W. and Lanteri, S. (1995). Influence of priming-induced nuclear replication activity on storability of pepper (*Capsicum annuum* L.) seed. *Seed Science Research*, 5(1): 25-29.
- Sathish, S. and Bhaskaran, M. (2013). Seed dry dressing with botanicals to improve physiological performance of fresh and aged seeds of blackgram (*Vigna mungo* L.). *African Journal of Agricultural Research*, 8(29): 4049-4057.
- Schmidt, D. H. and Tracy, W. F. (1988). Endosperm type, inbred background, and leakage of seed electrolytes during imbibition in sweet corn. *Journal of the American Society for Horticultural Science*, 113(2): 269-272.
- Schmidt, L. (2000). *Guide to handling of tropical and subtropical forest seed* (p. 511). Danida Forest Seed Centre, Denmark.
- Scott, G. E. (1981). Improvement for accelerated aging response of seed in maize populations. *Crop Science*, 21(1): 41-43.
- Sharma, S., Gambhir, S. and Munshi, S. K. (2007). Changes in Lipid and Carbohydrate Composition of Germinating. *Asian Journal of Plant Sciences*, 6(3): 502-507.
- Shelar, V. R. (2008). Role of mechanical damage in deterioration of soybean seed quality during storage-a review. *Agricultural Review*, 29(3): 177-184.
- Shelar, V.R. (2002). Strategies to improve the seed quality and storability of soybean- a review. *Agricultural Review*, 28: 188-96.
- Siadat, S. A., Moosavi, A. and Zadeh, M. S. (2012). Effects of seed priming on antioxidant activity and germination characteristics of maize seeds under different ageing treatment. *Research Journal of Seed Science*, 5(2): 51-62.
- Sibandé, G. A. K., Kabambe, V. H., Maliro, M. F. A. and Karoshi, V. (2015). Effect of priming techniques and seed storage period on soybean (*Glycine max* L) germination. *Journal of Dynamics in Agricultural Research*, 2(5): 46-53.
- Simic, B., Popovic, R., Sudaric, A., Rozman V, Kalinovic I, Cosic J., (2007). Influence of Storage Condition on Seed Oil Content of Maize, Soybean and Sunflower. *Agriculturae Conspectus Scientificus*. 72(3): 211-213.
- Singh, R. J. (2017). Botany and Cytogenetics of Soybean. In H.T. Nguyen and M.K. Bhattacharyya (Eds.), *The Soybean Genome* (pp. 11-40). Springer, Cham.,
- Singh, J. N. and Setia, R. K. (1974). germination of different qualities of soybean seeds under varying storage conditions. *Bulletin of Grain Technology*. 12: 3-10.
- Singh, R. J. (2011). *Genetic Resources, Chromosome Engineering, and Crop Improvement: Medicinal Plants, Volume 6*. CRC press.

- Siri, B., Vichitphan, K., Kaewnaree, P., Vichitphan, S. and Klanrit, P. (2013). Improvement of quality, membrane integrity and antioxidant systems in sweet pepper (*Capsicum annuum* Linn.) seeds affected by osmopriming. *Australian Journal of Crop Science*, 7 (13): 2068-2073.
- Sisman, C. B. (2005). Quality losses in temporary sunflower seed stores and influences of storage conditions on quality losses during storage. *Journal of Central European Agriculture*, 6(2):143-150.
- Sivritepe, H. O. and Dourado, A. M. (1995). The effect of priming treatments on the viability and accumulation of chromosomal damage in aged pea seeds. *Annals of Botany*, 75(2): 165-171.
- Sivritepe, H. Ö. and Dourado, A. M. (1998). The Effect of Storage Environment on Seed Survival and The Accumulation of Chromosomal Aberrations in Pea Landraces and Cultivars (*Pisum sativum* L.). *Turkish Journal of Botany*, 22(4): 223-232.
- Sivritepe, H.O. (1992). *Genetic Deterioration and Repair in Pea (Pisum sativum L.) Seeds During Storage*, PhD Thesis, University of Bath.
- Smiciklas, K. D., Mullen, R. E., Carlson, R. E. and Knapp, A. D. (1992). Soybean seed quality response to drought stress and pod position. *Agronomy Journal*, 84(2):166-170.
- Smirnoff, N. (1993). The role of active oxygen in the response of plants to water deficit and desiccation. *New Phytologist*, 125: 27-58.
- Soeda, Y., Konings, M.C., Vorst, O., van Houwelingen, A.M., Stoopen, G.M., Maliepaard, C.A., Kodde, J., Bino, R.J., Groot, S.P. and van der Geest, A.H.,(2005). Gene expression programs during *Brassica oleracea* seed maturation, osmopriming, and germination are indicators of progression of the germination process and the stress tolerance level. *Plant Physiology*, 137(1): 354-368.
- Soltani, A., Galeshi, S., Zeinali, E. and Latifi, N. (2002). Germination, seed reserve utilization and seedling growth of chickpea as affected by salinity and seed size. *Seed Science and Technology*, 30(1): 51-60.
- Soltani, E., Galeshi, S., Kamkar, B. and Akramghaderi, F. (2009). The Effect of Seed Aging on the Seedling Growth as Affected by. *Research Journal of Environmental Sciences*, 3(2): 184-192.
- Spilker, D. A., Schmitthenner, A. F. and Ellett, C. W. (1981). Effects of Humidity, Temperature, Fertility, and Cultivar on the Reduction of Soybean Seed Quality by *Phomopsis* sp. *Phytopathology*, 71(10): 1027-1029.
- Srivastava, A. K. and Gill, M. K. (1975). Physiology & biochemistry of seed deterioration in soyabean. III. seedling growth & seed leachate analysis. *Indian Journal of Experimental Biology*, 13: 481-485.

- Stewart, R. R. and Bewley, J. D. (1980). Lipid peroxidation associated with accelerated aging of soybean axes. *Plant Physiology*, 65(2): 245-248.
- Subedi, K. D. and Ma, B. L. (2005). Seed priming does not improve corn yield in a humid temperate environment. *Agronomy Journal*, 97(1): 211-218.
- Sun, H., Li, L., Wang, X., Wu, S. and Wang, X. (2011). Ascorbate–glutathione cycle of mitochondria in osmoprimed soybean cotyledons in response to imbibitional chilling injury. *Journal of Plant Physiology*, 168(3): 226-232.
- Sun, Q., Wang, J. H. and Sun, B. Q. (2007). Advances on seed vigor physiological and genetic mechanisms. *Agricultural Sciences in China*, 6(9): 1060-1066.
- Sung J. M. and Chiu, C.C. (1995). Lipid peroxidation and peroxide-scavenging enzymes of naturally aged soybean seed. *Plant Science*, 110: 45-52.
- Sung, J. M. and Jeng, T. L. (1994). Lipid peroxidation and peroxide-scavenging enzymes associated with accelerated aging of peanut seed. *Physiologia Plantarum*, 91(1): 51-55.
- Sung, J.M. (1996). Lipid peroxidation and peroxide-scavenging in soybean seeds during aging. *Physiologia Plantarum*, 97: 85-89.
- Sung, J.M. and Chang, Y.H. (1993). Biochemical activities associated with priming of sweet corn seeds to improve vigor. *Seed Science and Technology*, 21: 97-105.
- Surki, A. A., Sharifzadeh, F. and Afshari, R. T. (2012). Effect of drying conditions and harvest time on soybean seed viability and deterioration under different storage temperature. *African Journal of Agricultural Research*, 7(36): 5118-5127.
- Suzuki, N., and Mittler, R. (2006). Reactive oxygen species and temperature stresses: a delicate balance between signaling and destruction. *Physiologia Plantarum*, 126: 45-51.
- Suzuki, N., Koussevitzky, S., Mittler, R., and Miller, G. (2011). ROS and redox signalling in the response of plants to abiotic stress. *Plant Cell Environment*. 14: 691-699.
- Tabatabaei, S. A. (2013). The effect of priming on germination and enzyme activity of sesame (*Sesamum indicum* L.) seeds after accelerated aging. *Journal of Stress Physiology & Biochemistry*, 9(4).
- Tabatabaei, S. A. (2015). The Changes of Germination Characteristics and Enzyme Activity of Barley Seeds under Accelerated Aging. *Cercetari Agronomice in Moldova*, 48(2): 61-67.
- Tada, M. and Kawamura, S. (1963). Changes of soybean carbohydrates during growth and germination I. University of Kagawa, Japan, *Kagawa Daigaku Nogakubu Gakujutsu Hokoku*, 14: 148-155.

- Tatic, M., Balesevic- Tubic, S., Dordevic, V., Miklic, V., Vujakovic, M., Dukic, V. (2012). Vigor of sunflower and soybean ageing seed. *Helia*, 35(56): 119-126.
- Taylor, A.G., Allen, P.S., Bennet, M.A., Bradford, K.J., Burris, J.S. and Misra, M.K. (1998) Seed enhancements. *Seed Science and Research*, 8: 245-256.
- TeKrony, D.M. (2003). Precision is an essential component in seed vigor testing. *Seed Science and Technology*, 31: 435-447.
- TeKrony, D.M., Egli, D.B. and Phillips, A.D. (1980). Effect of field weathering on the viability and vigor of soybean seed. *Agronomy Journal*, 72: 749-753.
- TeKrony, D.M., Egli, D.B., Balles, J., Pfeiffer, T. and Fellows. R.J. (1979). Physiological maturity in soybeans. *Agronomy Journal*, 71:771-775.
- TeKrony, D.M., Egli, D.B., Balles, J., Tomes, L. and Stukey. R.E. (1984). Effect of date of harvest of harvest maturity on soybean seed quality and *Phomopsis* seed infection. *Crop Science*, 24: 189-193.
- TeKrony. D. M., and Egli, D. B. (1977). Relationship between laboratory indices of soybean seed vigor and field emergence. *Crop Science*. 17:573-577.
- Thant, P. S. (2015). *Priming of delayed harvest soybean (Glycine max L. Merr) seed for quality improvement*. MSc Thesis. Universiti Putra Malaysia.
- Thomas, U.C., Varughese, K., Thomas, A. and Sadanandan, S. (2000). Seed priming for increased vigour, viability and productivity of upland rice. *Leisa India*, 4:14.
- Tian, X., Song, S. and Lei, Y. (2008). Cell death and reactive oxygen species metabolism during accelerated ageing of soybean axes. *Russian Journal of Plant Physiology*, 55(1): 33-40.
- Tiecker Junior, A., Guimarães, L. E., Ferrari Filho, E., Castro, B. D., Del Ponte, E. M. and Dionello, R. G. (2014). Physicochemical quality of maize grain stored with different moisture contents in airtight and non-airtight environments. *Brazilian Journal of Corn and Sorghum*, 13(2): 174-186.
- Tilden R. L. and West, S.H. (1985) Reversal of the effects of ageing in soybean seeds. *Plant Physiology*, 77(3): 584-586.
- Tilebeni, G. H. and Golpayegani, A. (2011). Effect of seed ageing on physiological and biochemical changes in rice seed (*Oryza sativa* L.). *International Journal of AgriScience*, 1(3):138-143.
- Toole, E. H., Toole, V. K., Hendricks, S. B. and Borthwick, H. A. (1957). Effect of temperature on germination of light-sensitive seeds. *Proceeding of International Seed Testing Association*, 22: 1-9.

- Trawatha, S. E., TeKrony, D. M. and Hildebrand, D. F. (1995). Relationship of soybean seed quality to fatty acid and C6-aldehyde levels during storage. *Crop Science*, 35(5): 1415-1422.
- Tripathi, R. S. and Kumar, G. (2011). Comparative effect of ageing and gamma irradiation on the somatic cells of *Lathyrus sativus* L. *Journal of Central European Agriculture*, 11(4): 437-442.
- Ventura, L., Donà, M., Macovei, A., Carbonera, D., Buttafava, A., Mondoni, A., and Balestrazzi, A. (2012). Understanding the molecular pathways associated with seed vigor. *Plant Physiology and Biochemistry*, 60: 196-206.
- Verma, R. S. and Gupta, P. C. (1975). Storage behavior of soybean varieties vastly differing in seed size. *Seed Research*, 3(1): 39-44.
- Verma, S. S., Verma, U. and Tomer, R. P. S. (2003). Studies on seed quality parameters in deteriorating seeds in Brassica (*Brassica campestris*). *Seed Science and Technology*, 31(2): 389-396.
- Vertucci, C. W. and Farrant, J. M. (1995). Acquisition and loss of desiccation tolerance. *Seed Development and Germination*, 237-271.
- Vertucci, C. W. and Roos, E. E. (1993). Theoretical basis of protocols for seed storage II. The influence of temperature on optimal moisture levels. *Seed Science Research*, 3(03): 201-213.
- Vieira, R. D., Sedyama, T. U. N. E. O., da Silva, R. F., Sedyama, C. S. and Thiebaut, J. T. L. (1982). Effect of delaying the harvest of the seed quality sojz CV'UFV-2. *Brazilian Seed Journal*. 4(2): 9-22.
- Vijay, D., Dadlani, M., Kumar, P. A. and Panguluri, S. K. (2009). Molecular marker analysis of differentially aged seeds of soybean and safflower. *Plant molecular Biology Reporter*, 27(3): 282-291.
- Villiers, T. A. (1974). Seed aging: chromosome stability and extended viability of seeds stored fully imbibed. *Plant Physiology*, 53(6): 875-878.
- Villiers, T. A. (1975). Genetic maintenance of seeds in imbibed storage. *Crop Genetic Resources for Today and Tomorrow*, IBP, 2: 297-316.
- Vishwanath, K., Pallavi, H. M., Devraju, P. J. and Prashanth, Y. (2011). Prediction of Storability of Different Seed Size Grades of French bean Varieties through Accelerated Ageing Response. *Research Journal of Agricultural Sciences*, 2(2): 213-216.
- Walters, C., Ballesteros, D. and Vertucci, V. A. (2010). Structural mechanics of seed deterioration: standing the test of time. *Plant Science*, 179(6): 565-573.

- Walton, D. A. and Wallace, H. M. (2009). Delayed harvest reduces quality of raw and roasted macadamia kernels. *Journal of the Science of Food and Agriculture*, 89(2): 221-226.
- Wanichpongpan, P., Suriyachan, K. and Chandkrachang, S. (2001). Effects of chitosan on the growth of Gerbera flower plant (*Gerbera jamesonii*). *Chitin and chitosan: Chitin and Chitosan in Life Science, Yamaguchi, Japan*, 198-201.
- Waterworth, W. M., Bray, C. M. and West, C. E. (2015). The importance of safeguarding genome integrity in germination and seed longevity. *Journal of Experimental Botany*, 66 (12): 3549-3558.
- Whitcomb, W. O. and Johnson, A. H. (1928). Effect of severe weathering on certain properties of wheat. *Cereal Chemistry Journal*. 5: 117-128.
- Wien, H. C. and Kueneman, E. A. (1981). Soybean seed deterioration in the tropics. II. Varietal differences and techniques for screening. *Field Crops Research*, 4: 123-132.
- Wilcox, J. R., Laviolette, F. A. and Athrow, K. L. (1974). Deterioration of soybean seed quality associated with delayed harvest. *Plant Disease Reporter*, 58(2): 130-133.
- Xia, F., Chen, L., Sun, Y. and Mao, P. (2015). Relationships between ultrastructure of embryo cells and biochemical variations during ageing of oat (*Avena sativa* L.) seeds with different moisture content. *Acta Physiologiae Plantarum*, 37(4): 1-11.
- Xin, X., Tian, Q., Yin, G., Chen, X., Zhang, J., Ng, S. and Lu, X. (2014). Reduced mitochondrial and ascorbate–glutathione activity after artificial ageing in soybean seed. *Journal of Plant Physiology*, 171(2): 140-147.
- Yadav, S., Bhatia, V. S. and Guruprasad, K. N. (2003). Role of peroxidase and catalase enzymes in deterioration of soybean seeds due to field weathering. *Indian Journal of Plant Physiology*, 8(1): 195.
- Yadav, S., Bhatia, V. S., and Guruprasad, K. N. (2006). Oxyradical accumulation and rapid deterioration of soybean seeds due to field weathering. *Indian Journal of Biochemistry and Biophysics*, 43(1):41-7.
- Yadav., S. (2005) *Soybean seed deterioration due to field weathering. Role of Antioxidants and Antioxidant Enzymes*. PhD. Thesis, Devi Ahilya, Vishwavidyalaya, Idore, India.
- Yaklich, R.W. (1985). Effect of aging on soluble oligosaccharide content in soybean seeds. *Crop Science*. 25:701-704.
- Yao, Z., Liu, L., Gao, F., Rampitsch, C., Reinecke, D. M., Ozga, J. A. and Ayele, B. T. (2012). Developmental and seed aging mediated regulation of antioxidative

genes and differential expression of proteins during pre-and post-germinative phases in pea. *Journal of Plant Physiology*, 169(15): 1477-1488.

Yen, M.T., Yang, J.H. and Mau, J.L. (2008). Antioxidant properties of chitosan from crab shells. *Carbohydrate Polymers*. 74: 840-844.

Yin, G., Xin, X., Song, C., Chen, X., Zhang, J., Wu, S., Lu, X. (2014). Activity levels and expression of antioxidant enzymes in the ascorbate–glutathione cycle in artificially aged rice seed. *Plant Physiology and Biochemistry*. 80: 1-9.

Zamani, A., Nouri, S. A. S., Afshari, R. T., Nezhad, H. I., Akbari, G. A. and Tavakoli, A. (2010). Lipid peroxidation and antioxidant enzymes activity under natural and accelerated aging in safflower (*Carthamus tinctorius* L.) seed. *Iranian Journal of Field Crop Science*, 41(3):545-554.

Zeng, D., Luo, X. and Tu, R. (2012). Application of bioactive coatings based on chitosan for soybean seed protection. *International Journal of Carbohydrate Chemistry*, 1-5.

Zhan, J., Li, W., He, H. Y., Li, C. Z. and He, L. F. (2014). Mitochondrial alterations during Al-induced PCD in peanut root tips. *Plant physiology and biochemistry*, 75: 105-113.

Zhang F., Yu, J., Christopher, R.J., Wang, Y., Zhu, K., et al. (2015) Seed Priming with Polyethylene Glycol Induces Physiological Changes in Sorghum (*Sorghum bicolor* L. Moench) Seedlings under Suboptimal Soil Moisture Environments. *Plos One*, 10(10): e0140620. DOI: 10.1371/journal.pone.0140620.

Zhang, M., Wang, Z., Yuan, L., Yin, C., Cheng, J., Wang, L., Huang, J. and Zhang, H. (2012). Osmopriming improves tomato seed vigor under aging and salinity stress. *African Journal of Biotechnology*, 11(23): 6305-6311.

Zhang, R. G., Westbrook, M. L., Westbrook, E. M., Scott, D. L., Otwinowski, Z., Maulik, P. R., Reed, R.A. and Shipley, G. G. (1995). The 2.4 Å crystal structure of cholera toxin B subunit pentamer: cholera toxin B subunit pentamer: cholera toxin B subunit pentamer: cholera toxin B subunit pentamer. *Journal of Molecular Biology*, 251(4): 550-562.

Zhang, S., Hu, J., Zhang, Y., Xie, X.J. and Allen, K. (2007). Seed priming with brassinolide improves lucerne (*Medicago sativa* L.) seed germination and seedling growth in relation to physiological changes under salinity stress. *Australian Journal of Agricultural Research*, 58 (8): 811-815.

Zhang, Y., Liu, H., Shen, S. and Zhang, X. (2011). Improvement of eggplant seed germination and seedling emergence at low temperature by seed priming with incorporation SA into KNO₃ solution. *Frontiers of Agriculture in China*, 5(4): 534-537.

Zhu, S., Hong, D., Yao, J., Zhang, X. and Luo, T. (2010). Improving germination, seedling establishment and biochemical characters of aged hybrid rice seed by priming with KNO₃ + PVA. *African Journal of Agricultural Research*, 5(1): 078-083.