



**UNIVERSITI PUTRA MALAYSIA**

***DEVELOPMENT OF WEED COMPETITIVE RICE VARIETY  
UNDER WATER DEFICIT CONDITIONS THROUGH  
MARKER-ASSISTED BACKCROSS BREEDING***

**MASITAH AB JALIL**

**FP 2018 36**



**DEVELOPMENT OF WEED COMPETITIVE RICE VARIETY  
UNDER WATER DEFICIT CONDITIONS THROUGH  
MARKER-ASSISTED BACKCROSS BREEDING**

By

**MASITAH AB JALIL**

**Thesis submitted to the School of Graduate Studies,  
Universiti Putra Malaysia, in Fulfillment of the  
Requirements for the Degree of Doctor of Philosophy**

**January 2018**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in the fulfillment of the requirement for the degree of Doctor of Philosophy

**DEVELOPMENT OF WEED COMPETITIVE RICE VARIETY  
UNDER WATER DEFICIT CONDITIONS THROUGH  
MARKER-ASSISTED BACKCROSS BREEDING**

By

**MASITAH AB JALIL**

**January 2018**

**Chairman : Professor Abdul Shukor Juraimi, PhD**  
**Faculty : Agriculture**

Weed is one of the most troublesome pests in rice cultivation especially in water deficit areas, where farmers are commonly used herbicides to control it. Hence, it is important to develop rice variety that has capability to withstand water deficit and higher weed interference conditions, using an environmentally friendly method, such as marker-assisted backcross breeding. Eight rice varieties viz. AERON1, MR185, MR211, MR219, MR232, MR253, MR263 and MRQ74 were evaluated for their performances in water and weed stress conditions. From this study, two varieties (AERON1 and MRQ74) with the best performances were selected as donor and recurrent parental plants, respectively, for molecular breeding study. In molecular study, two foreground and 57 background markers were found as suitable to be used along the breeding study. The foreground markers were RM242 and RM263, which are linked markers to root length genes for drought tolerance. MRQ74 and AERON1 were crossed and backcrossed to generate F<sub>1</sub>, BC<sub>1</sub>F<sub>1</sub> and BC<sub>2</sub>F<sub>1</sub> populations, respectively. Rice lines of BC<sub>2</sub>F<sub>1</sub> population were self-pollinated to produce BC<sub>2</sub>F<sub>2</sub> population. Chi-square ( $\chi^2$ ) analysis for BC<sub>1</sub>F<sub>1</sub> and BC<sub>2</sub>F<sub>1</sub> populations showed good fit to the expected segregation ratio (1:1) for a single dominant gene model (d.f.=1, P>0.05) in Mendelian law for both foreground markers. The Chi-square ( $\chi^2$ ) values for RM242 and RM263 were  $\chi^2=3.1$  and  $\chi^2=0.03$ , and  $\chi^2=1.01$  and  $\chi^2=0.22$ , in BC<sub>1</sub>F<sub>1</sub> and BC<sub>2</sub>F<sub>1</sub> populations, respectively. In each BC<sub>1</sub>F<sub>1</sub> and BC<sub>2</sub>F<sub>1</sub> populations respectively, 18 and 25 rice lines were revealed as to carry both foreground markers. The highest RPG percentage for BC<sub>1</sub>F<sub>1</sub> and BC<sub>2</sub>F<sub>1</sub> populations were found in line (L)

50-39 with 71.4% and L45-6-24 with 90.8%, respectively. In BC<sub>2</sub>F<sub>2</sub> population, the Chi-square ( $\chi^2$ ) values for RM242 and RM263 were  $\chi^2=2.81$  and  $\chi^2=2.97$ , respectively, and showed good fit to the expected marker segregation ratio (1:2:1) for a single dominant gene model (d.f.=2, P>0.05) in Mendelian law. The highest RPG percentage was discovered in L45-6-24-23 with 93.0%. BC<sub>2</sub>F<sub>2</sub> population had also been evaluated for its agro-morphological traits and tolerance to water deficit conditions by using several drought indices, rank and standard deviation of rank method. Through BC<sub>2</sub>F<sub>2</sub> population, 12 rice lines viz. L45-27-3-7, L50-39-8-5, L50-39-8-8, L45-27-10-18, L45-27-10-28, L45-27-8-13, L45-27-8-28, L45-27-8-34, L50-39-7-15, L45-6-24-10, L45-6-24-20 and L45-6-24-23 were found to carry both foreground markers, whereas L2 (L50-39-8) was discovered as the most tolerant line to water deficit conditions in this study. The usage of molecular markers in this study is responsible to accelerate the development of new rice lines that can withstand water deficit conditions and possessed good weed competitive ability.

Abstrak tesis yang dikemukakan kepada senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PEMBANGUNAN VARIETI PADI YANG BERDAYA SAING  
DENGAN RUMPAI PADA KEADAAN KURANG AIR MELALUI  
PEMBIAKBAKAAN BERSILANG DENGAN PENANDA BANTUAN**

Oleh

**MASITAH AB JALIL**

**Januari 2018**

**Pengerusi: Profesor Abdul Shukor Juraimi, PhD  
Fakulti: Pertanian**

Rumpai adalah salah satu perosak penanaman padi terutamanya di kawasan yang kurang air, dimana petani menggunakan herbisid untuk mengawal rumpai. Oleh itu, adalah penting untuk membangunkan varieti yang berkebolehan untuk hidup pada keadaan kurang air dan tinggi kehadiran rumpai, menggunakan kaedah mesra alam seperti pembiakbakaan silang dengan penanda bantuan. Lapan varieti padi iaitu AERON1, MR185, MR211, MR219, MR232, MR253, MR263 dan MRQ74 dinilai keupayaannya pada keadaan stres air dan rumpai. Dalam kajian ini, dua varieti (AERON1 dan MRQ74) dengan keupayaan terbaik dipilih sebagai pokok induk penderma dan penerima, masing-masing, untuk kajian pembiakbakaan molekular. Dua penanda utama dan 57 penanda latarbelakang ditemui sesuai untuk digunakan sepanjang kajian pembiakbakaan ini. Dua penanda yang dipilih adalah RM242 dan RM263, yang merupakan penanda yang berkait dengan gen kepanjangan akar untuk kerintangan terhadap kemarau. MRQ74 dan AERON1 masing-masing dikacukkan dan dikacukbalikkan untuk menghasilkan populasi-populasi  $F_1$ ,  $BC_1F_1$  dan  $BC_2F_1$ . Titisan-titisan padi dari populasi  $BC_2F_1$  dikacuksendirikan untuk menghasilkan populasi  $BC_2F_2$ . Analisis Chi-kuasa dua ( $\chi^2$ ) menunjukkan padanan yang baik kepada nisbah pemisahan dijangka (1:1) bagi model gen tunggal dominan (d.f.=1,  $P>0.05$ ) dalam peraturan Mendel untuk dua penanda utama. Nilai-nilai Chi-kuasa dua ( $\chi^2$ ) untuk RM242 dan RM263 adalah  $\chi^2=3.1$  dan  $\chi^2=0.03$ , dan  $\chi^2=1.01$  dan  $\chi^2=0.22$ , masing-masing bagi populasi  $BC_1F_1$  dan  $BC_2F_1$ . Populasi  $BC_1F_1$  dan  $BC_2F_1$  masing-

masing menunjukkan bahawa 18 dan 25 titisan padi membawa kedua-dua penanda utama. Peratusan RPG tertinggi bagi populasi BC<sub>1</sub>F<sub>1</sub> dan BC<sub>2</sub>F<sub>1</sub> masing-masing adalah dari L50-39 dengan 71.4% dan L45-6-24 dengan 90.8%. Dalam populasi BC<sub>2</sub>F<sub>2</sub>, nilai Chi-kuasa dua ( $\chi^2$ ) untuk RM242 dan RM263 masing-masing adalah  $\chi^2=2.81$  dan  $\chi^2=2.97$ , dan menunjukkan padanan yang baik kepada nisbah pemisahan dijangka (1:2:1) bagi model gen tunggal dominan (d.f.=2, P>0.05) dalam peraturan Mendel. Peratusan RPG tertinggi adalah dari L45-6-24-23 dengan 93.0%. Titisan padi populasi BC<sub>2</sub>F<sub>2</sub> juga dinilai dari segi ciri-ciri agro-morfologikal dan kerintangan terhadap keadaan stres air menggunakan kaedah indeks kemarau, kedudukan dan pembahagian standard kedudukan. Melalui populasi BC<sub>2</sub>F<sub>2</sub>, 12 titisan padi iaitu L45-27-3-7, L50-39-8-5, L50-39-8-8, L45-27-10-18, L45-27-10-28, L45-27-8-13, L45-27-8-28, L45-27-8-34, L50-39-7-15, L45-6-24-10, L45-6-24-20 dan L45-6-24-23 ditemui membawa kedua-dua penanda utama, manakala L2 (L50-39-8) ditemui sebagai titisan paling rintang terhadap keadaan kurang air dalam kajian ini. Kesimpulannya, penggunaan penanda molekular dapat mempercepatkan pembangunantitisan padi baru yang dapat bertahan pada keadaan kurang air dan berdaya saing dengan rumpai.

## ACKNOWLEDGEMENTS

Foremost, all praises and endless thanks to Allah the Almighty for giving me the opportunity, strength and knowledge to pursue my study and complete it satisfactorily.

I would like to express my sincere gratitude to my supervisor Prof. Dr. Abdul Shukor Juraimi, for his patience, generous help and continuous support during my Ph.D study. Beside that, I would like to expand my grateful appreciation to the supervisory committee members: Prof. Dr. Mohd Rafii Yusop and Dr. Md. Kamal Uddin, for their critical reviews, insightful comments and encouragement.

My sincere thanks also go to all staffs of Department of Crop Science, Faculty of Agriculture, and Institute of Tropical Agriculture, who have provided me with the research facilities and valuable helps in faculty laboratories and net house of Ladang 10.

I share the credit of this work to all my fellow friends and labmates for the priceless discussions and suggestions, legion moral support and motivations, encouragement and their warm friendship.

Lastly, I would like to acknowledge with gratitude, the support and love of my family- my parents, En. Ab. Jalil Ab. Hamid and Pn. Kalsom Kadir, and my sister, Cik Musliha Ab Jalil. Thank you for the selfless love, care, pain, sacrifice, understanding and unconditional support along this journey. I would never be able to pay back the love and affection lavished by them.

I also place on record, my sense of gratitude to everyone, who directly or indirectly, has lent his or her hand in this venture.

Thank you,

**Masitah Ab Jalil**  
UPM Serdang, Malaysia  
November 2017



I certify that a Thesis Examination Committee has met on 25 January 2018 to conduct the final examination of Masitah binti Ab Jalil on her thesis entitled "Development of Weed Competitive Rice Variety Under Water Deficit Conditions through Marker-Assisted Backcross Breeding" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

**Mahmud bin Tengku Muda Mohamed, PhD**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Mohamad bin Osman, PhD**


Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Mohamed Hanafi bin Musa, PhD**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Iftikhar Hussain Khalil, PhD**

Professor  
University of Agriculture  
Pakistan  
(External Examiner)



**NOR AINI AB. SHUKOR, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 28 March 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Abdul Shukor Juraimi, PhD**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Mohd Rafii Yusop, PhD**

Professor  
Institute of Tropical Agriculture  
Universiti Putra Malaysia  
(Member)

**Md Kamal Uddin, PhD**

Senior Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Anuar bin Abdul Rahim, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia(Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals,modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Masitah Ab Jalil (GS32702)

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to

Signature: \_\_\_\_\_

Name of  
Chairman of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_

Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_

Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_

Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

## TABLE OF CONTENTS

<b>ABSTRACT</b>	<b>Page</b> i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xiv
<b>LIST OF FIGURES</b>	xvii
<b>LIST OF ABBREVIATIONS</b>	xxiii
 <b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	<b>3</b>
2.1 Weed problem in rice	3
2.1.1 Weed species and their classifications	3
2.1.2 Yield loss due to rice weeds	4
2.1.3 Weed effects on rice plants	5
2.1.4 Methods in weed management	5
2.2 Importance of water	8
2.2.1 Water scarcity	8
2.2.2 Drought adaptation	8
2.2.3 Water-saving irrigations in rice cultivation	9
2.2.4 Roles of rice's root	10
2.3 Rice culture	10
2.4 Aromatic and aerobic rice in Malaysia	11
2.5 Relationship between rice, water and weed	12
2.6 Rice breeding	13
2.6.1 Rice genome	13
2.6.2 History and challenges in rice breeding	14
2.6.3 Conventional backcross breeding and marker-assisted backcross breeding	16
2.6.4 Types of markers	17
2.6.5 Breeding for drought resistance rice varieties	21
2.6.6 Breeding for weed competitive rice varieties	21
2.6.7 Herbicide-tolerant rice varieties	22

2.7	Summary of literature review	23
-----	------------------------------	----

<b>3</b>	<b>IDENTIFICATION OF WEED COMPETITIVE ABILITY IN WATER DEFICIT CONDITIONS AMONG SELECTED RICE VARIETIES</b>	24
3.1	Introduction	24
3.2	Materials and methods	24
3.2.1	Experimental preparation	25
3.2.2	Soil sampling and preparation	25
3.2.3	Experimental design	25
3.2.4	Plant growth measurements	26
3.2.5	Weed evaluation	27
3.2.6	Statistical analysis	28
3.3	Results	28
3.3.1	Plant height	28
3.3.2	Tillers number	32
3.3.3	Relative chlorophyll content	36
3.3.4	Plant erectness	40
3.3.5	Early vigor	40
3.3.6	Days to flowering and days to maturity	41
3.3.7	Total root length and root dry weight	43
3.3.8	Above-ground biomass, grain yield and yield components	45
3.3.9	Weed relative density, relative dry weight and sum dominance ratio	51
3.3.10	Weed dry weight	52
3.3.11	Relative yield loss	53
3.3.12	Correlation of rice and weed traits	56
3.4	Discussion	58
3.5	Conclusion	61
<b>4</b>	<b>SELECTION OF SSR MARKERS AND DEVELOPMENT OF F<sub>1</sub> POPULATION OF MRQ74 × AERON1 FOR WEED COMPETITIVE AND WATER DEFICIT TOLERANT RICE VARIETY</b>	62
4.1	Introduction	62
4.2	Materials and methods	63
4.2.1	Crossing methods and development of crossed seeds	63
4.2.2	Laboratory works protocols	66
4.3	Results	67

4.4	Discussion	72
4.5	Conclusion	73
<b>5</b>	<b>RECURRENT PARENT GENOME RECOVERY AND GENETIC ANALYSIS IN BC<sub>1</sub>F<sub>1</sub> AND BC<sub>2</sub>F<sub>1</sub> POPULATIONS OF MRQ74 × AERON1</b>	<b>74</b>
5.1	Introduction	74
5.2	Materials and methods	75
5.2.1	Experimental site and plant materials	75
5.2.2	DNA extraction, PCR amplification and electrophoresis	75
5.2.3	Genotyping for marker segregation	75
5.2.4	Statistical analysis	76
5.3	Results	76
5.3.1	Marker-assisted foreground selection in BC <sub>1</sub> F <sub>1</sub> and BC <sub>2</sub> F <sub>1</sub> populations	76
5.3.2	BC <sub>1</sub> F <sub>1</sub> performance	78
5.3.3	BC <sub>2</sub> F <sub>1</sub> performance	83
5.4	Discussion	87
5.5	Conclusion	88
<b>6</b>	<b>GENETIC AND AGRO-MORPHOLOGICAL PERFORMANCE OF DROUGHT TOLERANCE AND WEED COMPETITIVE RICE LINES IN WATER AND WEED STRESS CONDITIONS</b>	<b>89</b>
6.1	Introduction	89
6.2	Materials and methods	90
6.2.1	Experimental preparation	90
6.2.2	Soil sampling and preparation	90
6.2.3	Plant materials	90
6.2.4	DNA extraction, PCR amplification and gel electrophoresis	91
6.2.5	Plant growth measurements	91
6.2.6	Weed evaluation	92
6.2.7	Drought tolerance evaluation	92
6.2.8	Molecular marker and statistical analysis	97
6.2.9	Rice line/variety codes and names for BC <sub>2</sub> F <sub>2</sub> population	97
6.3	Results	97
6.3.1	Marker-assisted foreground and background selection of BC <sub>2</sub> F <sub>2</sub> population	97

6.3.2	Agro-morphological performance of BC <sub>2</sub> F <sub>2</sub> population	103
6.3.3	Drought tolerance evaluation of BC <sub>2</sub> F <sub>2</sub> population	127
6.4	Discussion	134
6.5	Conclusion	139
<b>7</b>	<b>SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	<b>140</b>
	<b>REFERENCES</b>	<b>144</b>
	<b>APPENDICES</b>	<b>170</b>
	<b>BIODATA OF STUDENT</b>	<b>182</b>
	<b>LIST OF PUBLICATIONS</b>	<b>183</b>



## LIST OF TABLES

Table	Page
2.1 Traits differences between different rice types	14
2.2 Achievement of using marker assisted backcross breeding in developing aerobic rice with respect to root traits	15
2.3 Other achievements in rice breeding programs	16
2.4 Comparison of common markers used in rice breeding	20
3.1 Plant height of selected rice varieties affected by different level of water and weeding treatments	29
3.2 Analysis of variance of plant height at 20, 40, 60 DAS and harvest	29
3.3 Tillers number of selected rice varieties affected by different level of water and weeding treatments	33
3.4 Analysis of variance of tillers number at 20, 40, 60 DAS and harvest	33
3.5 RCC values at 30 DAS and 60 DAS, plant erectness and visual vigor of selected rice varieties affected by different level of water and weeding treatments	37
3.6 Analysis of variance of RCC at 30 and 60 DAS, plant erectness and early vigor	37
3.7 Days to flowering and days to maturity of selected rice varieties affected by different level of water and weeding treatments	42
3.8 Analysis of variance of days to flowering, days to maturity, total root length and root dry weight	42
3.9 Above-ground biomass, grains/panicle, thousand grains weight and grain filling and grain yield of selected rice varieties affected by different level of water and weeding treatments	46
3.10 Analysis of variance of above-ground biomass, grain/panicle, thousand grains weight, grain filling and grain yield	46
3.11 Family name, type, relative density, relative dry weight and sum dominance ratio of different weed species	52
3.12 Weed dry weight and relative yield loss of selected rice varieties affected by different level of water and weeding treatments	54

3.13	Analysis of variance of weed dry weight and relative yield loss	54
3.14	Pearson's correlation coefficient among different traits of rice and weed	57
4.1	List of polymorphic SSR markers used in this study	68
5.1	Chi-square analysis for foreground marker in BC <sub>1</sub> F <sub>1</sub> and BC <sub>2</sub> F <sub>1</sub> populations	77
5.2	Genetic background analysis in selected lines of BC <sub>1</sub> F <sub>1</sub> population	79
5.3	Genetic background analysis in selected lines of BC <sub>2</sub> F <sub>1</sub> population	83
6.1	Ranking scale of yield in non-stress condition, yield in stress condition and drought indices	96
6.2	Rice line/variety codes and names for BC <sub>2</sub> F <sub>2</sub> population plants	97
6.3	Chi-square analysis for foreground markers in BC <sub>2</sub> F <sub>2</sub> population derived from self-pollinating of BC <sub>2</sub> F <sub>1</sub> plants	98
6.4	Genetic background analysis in selected lines of BC <sub>2</sub> F <sub>2</sub> population	100
6.5	Plant height of selected rice lines affected by different level of water and weeding treatments	104
6.6	Analysis of variance of plant height at 20, 40, 60 DAS and harvest	104
6.7	Tillers number of selected rice lines affected by different level of water and weeding treatments	108
6.8	Analysis of variance of tillers number at 20, 40, 60 DAS and harvest	109
6.9	Days to flowering, days to maturity, root dry weight and total root length of selected rice lines affected by different level of water and weeding treatments	113
6.10	Analysis of variance of days to flowering, days to maturity, root dry weight and total root length	114
6.11	Thousand grains weight, grain yield and above-ground biomass of selected rice lines affected by different level of water and weeding treatments	118
6.12	Analysis of variance of thousand grains weight, grain yield and above-ground biomass	118

6.13	Weed dry weight and relative yield loss of selected rice lines affected by different level of water and weeding treatments	122
6.14	Analysis of variance of weed dry weight and relative yield loss	123
6.15	Family name, type, relative density, relative dry weight and sum dominance ratio of different weed species	124
6.16	Pearson's correlation coefficient among different traits of rice and weed	126
6.17	Tolerance indices of BC <sub>2</sub> F <sub>2</sub> plant population in water and weeding of non-stress and stress conditions	128
6.18	Correlation coefficients of drought tolerance indices between yield in non-water stress and water stress conditions for saturated condition (below diagonal) and field capacity condition (upper diagonal) under weeded condition	130
6.19	Correlation coefficients of drought tolerance indices between yield in non-water stress and water stress conditions for saturated condition (below diagonal) and field capacity condition (upper diagonal) under unweeded condition	131
6.20	Rank, mean rank and standard deviation of rank of drought tolerance indices	132
A	List of SSR markers used throughout this study	170

## LIST OF FIGURES

Figure		Page
2.1	Several types of weeds according to their classes; from left: <i>E. crus-galli</i> (grass), <i>C. iria</i> (sedge) and <i>M. vaginalis</i> (broadleaf)	4
2.2	Three different types of panicles (left) and grains (right) of Japonica, Javanica and Indica rice types	13
2.3	Conventional backcross breeding and molecular backcross breeding	17
2.4	Molecular backcross breeding steps	18
3.1	Interactions effect of variety $\times$ water $\times$ weeding for plant height at 20 DAS	30
3.2	Interactions effect of variety $\times$ water $\times$ weeding for plant height at 40 DAS	30
3.3	Interactions effect of variety $\times$ water $\times$ weeding for plant height at 60 DAS	31
3.4	Interactions effect of variety $\times$ water $\times$ weeding for plant height at harvest	31
3.5	Interactions effect of variety $\times$ water $\times$ weeding for tillers number at 20 DAS	34
3.6	Interactions effect of variety $\times$ water $\times$ weeding for tillers number at 40 DAS	34
3.7	Interactions effect of variety $\times$ water $\times$ weeding for tillers number at 60 DAS	35
3.8	Interaction effects of variety $\times$ water $\times$ weeding for tillers number at harvest	35
3.9	Interaction effects of variety $\times$ water for RCC values at 30 DAS	38
3.10	Interaction effects of variety $\times$ water for RCC values at 60 DAS	38
3.11	Interaction effects of variety $\times$ weeding for RCC values at 30 DAS	39
3.12	Interaction effects of variety $\times$ weeding for RCC values at 60 DAS	39
3.13	Interaction effects of variety $\times$ water $\times$ weeding for early vigor	41
3.14	Interaction effects of variety $\times$ water for days of flowering	43

3.15	Interaction effects of variety × water × weeding for total root length	44
3.16	Interaction effects of variety × water × weeding for root dry weight	45
3.17	Interaction effects of variety × water for above-ground biomass	47
3.18	Interaction effects of variety × water for grain/panicle	48
3.19	Interaction effects of variety × water for thousand grains weight	49
3.20	Interaction effects of variety × water for grain filling	48
3.21	Interaction effects of variety × water × weeding for grain yield	52
3.22	Relative density based on weed classes (left) and relative dry weight (right) based on weeds classes	52
3.23	Interaction effect of variety × water × weeding for weed dry weight	55
3.24	Interaction effect of variety × water × weeding for relative yield loss	55
4.1	Crossing and selection scheme of F <sub>1</sub> , BC <sub>1</sub> F <sub>1</sub> , BC <sub>2</sub> F <sub>1</sub> and BC <sub>2</sub> F <sub>2</sub> populations	64
4.2	Emasculation process by using pollen emasculator vacuum	65
4.3	Pollination process until grain development	65
4.4	Polymorphism between parent plants for (a) RM242 and (b) RM263 as foreground markers, and (c) RM248, (d) RM5 and (e) RM280 as background markers, ran on 3% metaphor gel stained with FloroSafe DNA stain (L=50 bp ladder, P1=parent AERON1, P2=parent MRQ74)	67
5.1	Screening of BC <sub>1</sub> F <sub>1</sub> population using (a) RM263 and (b) RM242 markers, ran on 3% metaphor gel stained with FloroSafe DNA stain (L=50 bp ladder, bp=base pair, P1=parent AERON1 and P2=parent MRQ74). Double banding patterns showed rice lines with both donor and recurrent parent alleles, while single banding patterns showed rice line with either donor or recurrent parent allele	77
5.2	Screening of BC <sub>2</sub> F <sub>1</sub> population using (a) RM263 and (b) RM242 marker, ran on 3% metaphor gel stained with FloroSafe DNA stain (L=50 bp ladder, bp=base pair, P1=parent AERON1 and P2=parent MRQ74). Double banding patterns showed rice lines with both	78

	donor and recurrent parent alleles, while single banding patterns showed rice line with either donor or recurrent parent allele	
5.3	Frequency distribution of the recurrent parent genome percentage of BC <sub>1</sub> F <sub>1</sub> population	79
5.4	Selected background markers banding pattern of (a) RM11 and (b) RM72 in BC <sub>1</sub> F <sub>1</sub> population, ran on 3% metaphor gel stained with FloroSafe DNA stain (L=50 bp ladder; A=donor parent banding pattern; B=recurrent parent banding pattern; H=heterozygous banding pattern)	80
5.5	Recurrent parent genome recovery of selected BC <sub>1</sub> F <sub>1</sub> plants in regards to all 12 rice chromosomes. Blue color indicates homozygous regions of MRQ74 alleles, red color indicates homozygous regions of AERON1 alleles, and yellow color indicates heterozygous regions	81
5.6	Chromosome-wise of the highest recurrent parent genome recovery of L50-39 in BC <sub>1</sub> F <sub>1</sub> population. Blue color indicates homozygous regions for MRQ74 alleles and yellow color indicates heterozygous regions.	82
5.7	Frequency distribution of the recurrent parent genome percentage of BC <sub>2</sub> F <sub>1</sub> populations of selected rice lines	84
5.8	Selected background markers banding pattern of (a) RM271 and (b) RM274 in selected plants of BC <sub>2</sub> F <sub>1</sub> population, ran on 3% metaphor gel stained with FloroSafe DNA stain (L=50 bp ladder; A=donor parent banding pattern; B=recurrent parent banding pattern; H=heterozygous banding pattern)	84
5.9	Recurrent parent genome recovery of selected BC <sub>2</sub> F <sub>1</sub> plants in regards to all 12 rice chromosomes. Blue color indicates homozygous regions for MRQ74 alleles, red color indicates homozygous regions for AERON1 alleles, and yellow color indicates heterozygous regions	85
5.10	Chromosome-wise of the highest recurrent parent genome recovery	85

of L45-6-24 in BC<sub>2</sub>F<sub>1</sub> generation. Blue color indicates homozygous regions for MRQ74 alleles and yellow color indicates heterozygous regions.

6.1	Frequency distribution of the recurrent parent genome percentage in BC <sub>2</sub> F <sub>2</sub> population of selected rice lines	99
6.2	Homozygous banding pattern in accordance to donor parent (AERON1) alleles of BC <sub>2</sub> F <sub>2</sub> population for (a) RM242 and (b) RM263 and homozygous banding pattern in accordance to recurrent parent (MRQ74) alleles of BC <sub>2</sub> F <sub>2</sub> population (c) RM1132 and (d) RM7193 (L=50 bp ladder; A=donor parent; B=recurrent parent)	100
6.3	Recurrent parent genome recovery of selected BC <sub>2</sub> F <sub>2</sub> plants in regards to all 12 rice chromosomes. Blue color indicates homozygous regions for MRQ74 alleles, red color indicates homozygous regions for AERON1 alleles, and yellow color indicates heterozygous regions	101
6.4	Chromosome-wise of the highest recurrent parent genome recovery of L45-6-24-23 in BC <sub>2</sub> F <sub>2</sub> population. Blue color indicates homozygous regions for MRQ74 alleles, red color indicates homozygous regions for AERON1 alleles, and yellow color indicates heterozygous regions	102
6.5	Interaction effects of line/variety × water for plant height at 20 DAS	105
6.6	Interaction effects of line/variety × weeding for plant height at 20 DAS	105
6.7	Interaction effects of lines/variety × water × weeding for plant height at 40 DAS	106
6.8	Interaction effects of lines/variety × water × weeding for plant height at 60 DAS	106
6.9	Interaction effects of lines/variety × water × weeding for plant height at harvest	107
6.10	Interaction effects of line/variety × water for tillers number at 40 DAS	109



6.11	Interaction effects of line/variety $\times$ water for tillers number at 60 DAS	110
6.12	Interaction effects of line/variety $\times$ weeding for tillers number at 40 DAS	110
6.13	Interaction effects of line/variety $\times$ weeding for tillers number at 60 DAS	111
6.14	Interaction effects of line/variety $\times$ water $\times$ weeding for tillers number at harvest	111
6.15	Interaction effects of line/variety $\times$ water $\times$ weeding for days to flowering	114
6.16	Interaction effects of line/variety $\times$ water $\times$ weeding for days to maturity	115
6.17	Interaction effects of line/variety $\times$ water $\times$ weeding for root dry weight	116
6.18	Interaction effects of line/variety $\times$ water $\times$ weeding for total root length	117
6.19	Interaction effects of line/variety $\times$ water $\times$ weeding for thousand grains weight	119
6.20	Interaction effects of line/variety $\times$ water $\times$ weeding for grain yield	120
6.21	Interaction effects of line/variety $\times$ water $\times$ weeding for above-ground biomass	121
6.22	Interaction effects of line/variety $\times$ water $\times$ weeding for weed dry weight	123
6.23	Interaction effects of line/variety $\times$ water $\times$ weeding for relative yield loss	124
6.24	Relative density based on weed classes (left) and relative dry weight (right) based on weeds classes	125
B1	MRQ74 plants (female/recurrent parent)	181
B2	AERON1 plants (male/donor parent)	181
B3	F <sub>1</sub> population plants	181
B4	BC <sub>1</sub> F <sub>1</sub> population	181



B5	BC <sub>2</sub> F <sub>1</sub> population	181
B6	BC <sub>2</sub> F <sub>2</sub> population	181



## LIST OF ABBREVIATIONS

AFLP	Amplified Fragment Length Polymorphism
ANOVA	Analysis of Variance
AWD	Alternate Wetting and Drying
bp	base pair
BIL	Backcross Inbred Lines
CEC	Cation Exchange Capacity
cM	centi Morgan
CTAB	Centyltrimethylammonium bromide
$\chi^2$	Chi-square
DAS	Day after sowing
d.f.	Degree of freedom
DI	Drought Resistance Index
DMRT	Duncan Multiple Range Test
DNA	Deoxyribonucleic acid
dSm	decisemens per meter
EDTA	Ethylenediaminetetraacetic acid
gmp	Geometric Mean Productivity
IMW	Integrated Weed Managment
kPa	kiloPascal
M	molar
MABC	Marker-assisted backcross breeding
MAS	Marker-assisted selection
Mb	megabytes
meq	millequivalents
mM	micromolar
MOP	Muriate of potash
MPI	Mean Productivity Index
MRP	Mean Relative Performance

NPK	Natrium/Potassium/Kalium
PCR	Polymerase Chain Reaction
ppm	parts per million
QTL	Quantitative Trait Loci
RCBD	Randomized Complete Block Design
RPG	Recurrent Parent Genome
SDR	Standard deviation of rank
SRI	System of Rice Intensification
SSC	Saturated Soil Culture
SSR	Simple Sequence Repeat
SSI	Stress Susceptibility Index
SSSI	Schneider's Stress Susceptibility Index
STI	Stress Tolerance Index
RAPD	Randomly Amplified Polymorphic DNA
RCC	Relative Chlorophyll Content
RDI	Relative Drought Index
REI	Relative Efficiency Index
RFLP	Restriction Fragment Length Polymorphism
rpm	rounds per minute
SNP	Single Nucleotide Polymorphism
SPAD	Silicon Photon Activated Diode
TBE	Tris/Boric acid/EDTA
TOL	Stress Tolerance
TE	Tris/EDTA
TSP	Triple Super Phosphate
WAS	Week after sowing
YSI	Yield Stability Index
μl	microliter

## CHAPTER 1

### INTRODUCTION

Weeds are the major constraint in rice cultivation followed by nitrogen deficiency and pests and diseases (Anwar *et al.*, 2010). Generally, weeds are competing with rice for sunlight, water, soil and nutrients (Davies *et al.*, 2008). In water deficit conditions, more C<sub>4</sub> weeds are found, as they are able in adapting hot and dryland conditions due to having higher water use efficiency compared to C<sub>3</sub> plants (Abouziena *et al.*, 2015). Therefore, weed control in water deficit condition is more important and challenging as rice is most susceptible and severely affected to dry condition during booting and flowering stages that tend to cause yield loss (Valencia, 2015). Meanwhile, in typical rice cultivation with easiness of water access, weed had also become a major problem and lowered rice yield production. This is due to shifting of cultivation method from transplanting to direct-seeded method as a way to reduce labor cost and production risks (Karim *et al.*, 2004). In this situation, weeds emerge at the same time with rice seedlings. To overcome weed problem, farmers tend to use hazardous herbicides that can cause health problem, pollute the environment, damage the plant itself, as well as encourage the emergence of herbicides resistant weeds (Maclean *et al.*, 2002). Hence, a method known as Integrated Weed Management (IWM) was developed to overcome this problem. This method implements preventive, cultural, mechanical and chemical methods throughout rice cultivation period (Owombo *et al.*, 2014), as single method seldom meet the demanded outcomes (Webster and Levy, 2009).

Rice is one of the important cereals in the world as one-third of the world population consumes it. Rice use approximately one-fourth to one-third of world's freshwater annual supply and cause a waste of this limited and costly resource (Africare, 2010). It demands water as much as three times than wheat and maize and typically grown in puddle-irrigated soil condition (Swamy and Kumar, 2012). However, water is scarce nowadays. China and India, the world's leader of rice production are severely affected by freshwater resource scarcity (Priyanka *et al.*, 2012). Uncertainty in climate and increase of human population all over the world are among contributing factors to water scarcity condition (Sariam and Anuar, 2010). Hence, to overcome the problem, scientists had come up with aerobic rice varieties that were believed to withstand water deficit condition efficiently. Aerobic rice is a combination of traits of upland and lowland rice varieties with low water usage and is responsive to input (Priyanka *et al.*, 2012). Despite of its ability, these varieties are facing serious weed problem since weeds are abundantly grow in inadequate water availability.

To manage water and weed stress problems, marker-assisted backcross breeding had been utilized to develop new rice varieties that can tolerate those stress conditions. Marker-assisted backcross breeding has given a huge positive impact in rice breeding

program (Yi *et al.*, 2009). Markers are used in backcross breeding to control the target gene through foreground selection and genetic background through background selection, which aims to reduce donor parent genome through recurrent backcrosses (Hospital, 2001). Markers-assisted selection helps to choose the best plants during cultivation, time, space and cost saving, and raise no ethical issues as in transgenic plants (Bernier *et al.*, 2008; Francia *et al.*, 2005; Hasan *et al.*, 2015a). Meanwhile, breeding for drought tolerance varieties is quite a challenging effort due to complexity of genetics and uncertainty in time and intensity of drought period (Tuberosa and Salvi, 2005). Thus, with the discovery of many QTLs related to drought, new routes to study the contributor traits of drought tolerance are now enlarge (Saadalla, 2008). For water stress, root traits play important role in resisting drought condition in rainfed lowland irrigation (Manickavelu *et al.*, 2006). Deep-rooting variety is more resistant to drought as it has abilities to extract water from deeper soil and produce higher yield than shallow-rooting variety (Saadalla, 2008). Meanwhile, rice plant's traits, such as plant height, tillers number, early growth vigor and allelopathy ability (Anwar, 2012) should be wisely manipulated to overcome weed occurrence in rice field.

Perhaps, resistance towards both water and weed stress conditions is an ideal type of rice plant in competing those stress simultaneously in rice field. Hence, to develop a new rice variety that can withstand water and weed stress conditions, these objectives need to be achieved:

- i. To identify weed competitive traits in water deficit condition among selected rice varieties,
- ii. To select suitable parental plants of rice for molecular breeding study,
- iii. To improve selected high-yielding rice variety by enhancing the ability to compete with weeds and water deficit environments through molecular breeding,
- iv. To evaluate the new improved rice lines towards different water and weed stress conditions.

## REFERENCES

- Abeysekera and Anuruddhika, S. K. (2001). *Management of Echinochloa spp. in rice in Sri Lanka*. Paper presented at the FAO workshop on *Echinochloa spp.* control Beijing.
- Abidin, M. I. Z, Lassim, M. and Hashim, N. (1989). *Tanaman bijian*. Kuala Lumpur: Dewan Bahasa dan Pustaka Kementerian Pendidikan Malaysia.
- Abouziena, H. F., El-saeid, H. M. and Amin, A. A. E. (2015). Water loss by weeds: a review. *International Journal of ChemTech Research*, 7: 323–336.
- ACIAR. (2006). Growing more rice with less water. Retrieved from <http://aciar.gov.au/content/growing-more-rice-less-water>. Accessed January 22, 2013.
- Africare, Oxfam America, WWF-ICRISAT Project (2010). *More rice for people, more water for the planet*. WWF-ICRISAT Project: Hyderabad, India.
- Ahmadi, N., Audebert, A., Bennett, M. J., Bishopp, A., de Oliveira, A., Courtois, B., Diedhiou, A., Diévar, A., Gantet, P., Ghesquière, A., Guiderdoni, E., Henry, A., Inukai, Y., Kochian, L., Laplaze, L., Lucas, M., Luu, D., Manneh, B., Mo, X., Muthurajan, R., Périn, C., Price, A., Robin, S., Sentenac, H., Sine, B., Uga, Y., Véry, A., Wissuwa, M., Wu, P. and Xu, J. (2014). The roots of future rice harvests. *Rice*, 7: 1-9.
- Ahmed, F. (2015). *Development of submergence tolerant rice variety through marker-assisted backcross breeding between MR219 and Swarna-Sub1*, PhD Thesis, Universiti Putra Malaysia.
- Ahmed, F., Rafii, M. Y., Ismail, M. R., Juraimi, A. S., Rahim, H. A., Tanweer, F. A., and Latif, M. A. (2016). Recurrent parent genome recovery in different populations with the introgression of *Sub1* gene from a cross between MR219 and Swarna-Sub1. *Euphytica*, 207: 605–618.
- Alam, M. S., Salim, M., Moniruzzaman, M., Rashid, J. A. and Islam, M. M. (2012). Marker-assisted foreground selection for identification of salt tolerant rice genotypes. *The Agriculturists*, 10: 1–8.
- Ali, M. B. and El-sadek, A. N. (2016). Evaluation of drought tolerance indices for wheat (*Triticum aestivum* L.) under irrigated and rainfed conditions. *Communications in Biometry and Crop Science*, 11: 77–89.

- Al-Kaisi, M., Hanna, M. and Tidman, M. (2002). Post-planting tillage: What are the options? Retrieved from <http://www.ipm.iastate.edu/ipm/icm/2002/5-6-2002/postplanttillage.html>. Accessed August 31, 201.
- Akçura, M., Partigoç, F. and Kaya, Y. (2011). Evaluating of drought stress tolerance based on selection indices in Turkish bread wheat landraces. *Journal of Animal and Plant Sciences*, 21: 700–709.
- Akhtar, S., Bhat, M. A., Bhat, K. A., Chalkoo, S., Mir, M. R. and Wani, S. A. (2010). Marker-assisted selection in rice. *Journal of Phytology*, 2: 66–81.
- Ampong-Nyarko, K. and De Datta, S. K. (1991). *A Handbook for Weed Control in Rice*. Manila: International Rice Research Institute.
- Ananna, R. A. (2014). *Marker-assisted selection and screening of salt tolerant gene(s) in BC<sub>2</sub>F<sub>2</sub> population of Binadhan-7 × FL478*. Bangladesh Agricultural University, Mymensingh.
- Andrew, I. K. S., Storkey, J. and Sparkes, D. L. (2015). A review of the potential for competitive cereal cultivars as a tool in integrated weed management. *Weed Research*, 55: 239–248.
- Anwar, M. P., Juraimi, A. S., Man, A., Puteh, A., Selamat, A. and Begum, M. (2010). Weed suppressive ability of rice (*Oryza sativa* L.) germplasm under aerobic soil conditions. *Australian Journal of Crop Science*, 4: 706-717.
- Anwar, M. P. (2012). *Strategies for weed suppression in aerobic rice cultivation*, PhD Thesis, Universiti Putra Malaysia, Malaysia.
- Asch, F., Dingkuhn, M., Sow, A. and Audebert, A. (2005). Drought-induced changes in rooting patterns and assimilate partitioning between root and shoot in upland rice. *Field Crops Research*, 93: 223–236.
- Ashraf, M. (2010). Inducing drought tolerance in plants: recent advances. *Biotechnology Advances*, 28: 169–183.
- Awasthi, S. and Lal, J. P. (2014). Marker assisted selection for the improvement of Sarjoo-52 for drought tolerance by introgression of MQTL1.1 from the source Nagina-22. *Journal of Plant Molecular Breeding*, 2: 43–55.
- Azmi, M. (1992). Competitive ability of barnyard grass in direct seeded rice. *Teknologi Padi*, 8: 19-25.
- Azmi, M., Baki, B. B. and Mashhor, M. (1993). *Weed communities in principal rice-growing areas in Peninsular Malaysia*. MARDI Report No. 165.



- Azmi, M., Shukor, J. A. and Najib, M. Y. M. (2007). Critical period for weedy rice control in direct-seeded rice. *Journal of Tropical Agriculture and Food Science*, 35: 333–339.
- Babu, R. C. (2010). Breeding for drought resistance in rice : an integrated view from physiology to genomics. *Electronic Journal of Plant Breeding*, 1: 1133–1141.
- Balch, E. P. M., Gidekel, M., Segura-Nieto, M., Estrella, L. H. and Alejo, N. O. (1996). Effect of water stress on plant growth and roots proteins in three cultivars of rice (*Oryza sativa*) with different level of drought tolerance. *Physiologia Planetarium*, 96: 284-290.
- Bàrberi, P. (2003). Preventive and cultural methods for weed management. In R. Labrada (Ed.), *Weed management for developing countries* (p. 179-194). Rome: Food and Agriculture Organization of the United Nations.
- Bardakci, F. (2001). Random amplified polymorphic DNA (RAPD) markers. *Turkey Journal Biology*, 25: 185-196.
- Becker, M. and Johnson D. E. (1999). The role of legume fallows in intensified upland rice-based systems in West Africa. *Nutrient Cycling in Agroecosystems*, 53: 71-81
- Begum, M., Juraimi, A. S., Amartalingum, R., Omar, S. Y. S. and Man, A. (2009). Effect of *Fimbristylis miliacea* competition with MR220 rice in relation to different nitrogen levels and weed density. *International Journal of Agriculture and Biology*, 11: 183–187.
- Bernier, J., Atlin, G. N., Serraj, R., Kumar, A. and Spaner, D. (2008). Breeding upland rice for drought resistance. *Journal of the Science of Food and Agriculture*, 88: 927–939.
- Bernama. (2012, June 18). 10,000 petani KADA alami kerugian RM150 juta kerana kemarau. *Berita Wilayah*.
- Bhatt, M. D. and Tewari, A. (2006). Losses in growth and yield attributes due to weed composition in transplanted paddy in Terai region. *Scientific World*, 4: 99–101.
- Boopathi, N. M. (2013). *Genetic mapping and marker assisted selection: basics, practice and benefits*. Tamil Nadu: Springer India.
- Bouman, B. A. M. and Tuong, T. P. (2000). Field water management to save water and increase its productivity in irrigated lowland rice. *Agricultural Water Management*, 1615: 1-20.



- Bouman, B. A. M., Humphreys, E., Tuong, T. P. and Barker, R. (2006). Rice and water. *Advances in Agronomy*, 92: 187-237.
- Bouman, B. A. M., Lampayan, R. M. and Tuong, T. P. (2007). *Water management in irrigated rice: coping with water scarcity*. Los Baños: International Rice Research Institute.
- Bousslama, M. and Schapaugh, W. T. (1984). Stress tolerance in soybean. Part 1: evaluation of three screening techniques for heat and drought tolerance. *Crop Science*, 24: 93-937.
- Chagné, D., Gasic, K., Crowhurst, R. N., Han, Y., Bassett, H. C., Bowatte, D. R., Lawrence, T., Rikkerink, E. H. A., Gardiner, S. E. and Korban, S. S. (2008). Development of a set of SNP markers present in expressed genes of the apple. *Genomics*, 92: 353-358.
- Chakrabhorthy, S. (2008). *Rice breeding and genetics*. New Delhi: Concept Publishing Company.
- Chan, C. S., Zainudin, H., Saad, A. and Azmi, M. (2012). Productive water use in aerobic rice cultivation. *Journal of Tropical Agriculture and Food Science*, 49: 117-126.
- Chang, T. T. (2003). Origin, domestication and diversification. In W. C. Smith and R. H. Dilday (Eds.), *Rice: origin, history, technology and production* (pp. 3-25). Hoboken: John Wiley and Sons Inc.
- Chauhan, B. S. (2012). *Weed management in direct-seeded rice systems*. Los Baños: International Rice Research Institute.
- Chauhan, B. S. and Abugho, S. B. (2013). Effect of water stress on the growth and development of *Amaranthus spinosus*, *Leptochloa chinensis* and rice. *American Journal of Plant Sciences*, 4: 989-998.
- Codrington, S. (2005). *Planet geography*. Hong Kong: Solid Star Press.
- Comas, L. H., Becker, S. R., Cruz, V. M. V, Byrne, P. F. and Dierig, D. A. (2013). Root traits contributing to plant productivity under drought. *Frontiers in Plant Science*, 4: 1-16.
- Collard, Iftekharuddaula, Thomson, Pamplona and Mackill. (2008). Recovery of the recurrent parent genome. Retrieved from [http://mcclintock.generationcp.org/index.php?option=com\\_content&task=view&id=109&Itemid=115](http://mcclintock.generationcp.org/index.php?option=com_content&task=view&id=109&Itemid=115). Accessed October. 7, 2015.

- Collard, B. C. Y. and Mackill, D. J. (2008). Marker-assisted selection: an approach for precision plant breeding in the twenty-first century. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 363: 557–572.
- Courtois, B., Ahmadi, N., Khowaja, F., Price, A. H., Rami, J. F., Frouin, J., Hamelin, C. and Ruiz, M. (2009). Rice root genetic architecture: meta-analysis from a drought QTL database. *Rice*, 2: 115–128.
- Croughan, T. P., Utomo, H. S., Sanders, D. H. and Braveman, M. B. (1996). Herbicide-resistant rice offers potential solution to red rice problem. *Louisiana Agriculture*, 39, 10-12.
- Curran, W. (2004). *Weed Management in Organic Cropping Systems*. The Pennsylvania State University: Pennsylvania.
- Davatgar, N., Neishabouri, M. R., Sepaskhah, A. R. and Soltani, A. (2009). Physiological and morphological responses of rice (*Oryza sativa* L.) to varying water stress management strategies. *International Journal of Plant Production*, 3: 19–32.
- Davies, G., Turner, B., Bond B., Doubleday, H. (2008). *Weed management for organic farmers, growers and smallholders: a complete guide*. Wiltshire: The Crowood Press Ltd.
- De Datta, S. K. (1975). *Major research in upland rice*. Los Banos: International Rice Research Institute.
- De Datta, S. K. and Feuer, R. (1975). *Soils on which upland rice is grown*. Los Banos: International Rice Research Institute.
- De Datta, S. K. and Baltazar, A. M. (1996). Weed control technology as a component of rice production systems. In B. A. Auld and K.-U. Kim, (Eds.), *Weed management in rice* (pp. 27-52). Rome: Food and Agriculture Organization of the United Nations.
- Dingkuhn, M., Johnson, D. E., Sow, A. and Audebert A. Y. (1999). Relationships between upland rice canopy characteristics and weed competitiveness. *Field Crops Research*, 61: 79-95.
- Dingkuhn, M., Tivet, F., Siband, P., Asch, F., Audebert, A. and Sow, A. (2001). Varietal differences in specific leaf area: a common physiological determinant of tillering ability and early growth vigor? In S. Peng and B. Hardy, (Eds.), *Rice research for food security and poverty alleviation. Proceedings of the International Rice Research Conference* (pp. 95–108). Los Banos: International Rice Research Institute.

- Dorman, W. K. (1976). Selection and breeding of multiple traits. In *The Genetic and Breeding of Southern Pines* (pp. 339–351). USA: Reforestration, Nurseries and Genetics Resources.
- Eastin, E. F. (1983). Weed, disease, insect interaction in rice. *Proceedings of the Conference on Weed Control*. Manila: International Rice Research Institute.
- El-Rawy, M. A. and Hassan, M. I. (2014). Effectiveness of drought tolerance indices to identify tolerant genotypes in bread wheat (*Triticum aestivum* L.). *Journal of Crop Science and Biotechnology*, 17: 255–266.
- Engku, A. K., Norida, M., Hsa, N., Juraimi, A. S., Rafii, M. Y., and Anwar, M. P. (2014). Variation of herbicide resistant Clearfield® rice and weedy rice variants in Malaysia. *International Journal of Biology, Pharmacy and Allied Sciences*, 3, 2825–2835.
- Estorninos, L. E., Gealy, J. D. R. and Talbert, R. E. (2002). Growth response of rice (*Oryza sativa*) and red rice (*O. sativa*) in a replacement series study. *Weed Technology*, 16: 401–406.
- Fard, A. K. and Sedaghat, S. (2013). Evaluation of drought tolerance indices in bread wheat recombinant inbred lines. *European Journal of Experimental Biology*, 3: 201–204.
- Farhad, M., Hakim, M. A., Alam, M. A. and Barma, N. C. D. (2014). Screening wheat genotypes for coleoptile length: a trait for drought tolerance. *American Journal of Agriculture and Forestry*, 2: 237–245.
- Farooq, M., Wahid, A., Kobayashi, N., Fujita, D. and Basra, S. M. A. (2009). Plant drought stress: effects, mechanisms and management. *Agronomy for Sustainable Development*, 29: 185–212.
- Farshadfar, E., Moradi, Z., Elyasi, P., Jamshidi, B. and Chaghakabodi, R. (2012). Effective selection criteria for screening drought tolerant landraces of bread wheat (*Triticum aestivum* L.). *Scholars Research Library Annals of Biological Research*, 3: 2507–2516.
- Fatimah, Prasetiyono, J., Dadang, A. and Tasliah. (2014). Improvement of early maturity in rice variety by marker assisted backcross breeding of Hd2 gene. *Indonesia Journal Agriculture Science*, 15: 55–64.
- Fernandez, G. (1992). Effective selection criteria for assessing plant stress tolerance. *Proceedings of the International Symposium on Adaptation of Vegetables and other Food Crops in Temperature and Water Stress*. Taiwan: Asian Vegetable Research and Development Centre.

- Ferrero, A. (2003). Weedy rice, biological features and control. In R. Labrada, (Ed.), *Weed management for developing countries*. (pp. 89–107). Italy: FAO Plant Production and Protection Paper.
- Fischer, R. A. and Maurer, R. (1978). Drought resistance in spring wheat cultivars. I. Grain yield responses. *Australian Journal of Agricultural Research*, 29: 897–912.
- Florez, J. A., Fischer, A. J., Ramirez, H. and Duque, M. C. (1999). Predicting rice yield losses caused by multispecies weed competition. *Agronomy Journal*, 91: 87–92.
- Francia, E., Tacconi, G., Crosatti, C., Barabaschi, D., Bulgarelli, D., Dall’Aglio, E. and Valè, G. (2005). Marker assisted selection in crop plants. *Plant Cell, Tissue and Organ Culture*, 82: 317–342.
- Frisch, M., Bohn, M. and Melchinger, A. E. (1999). Comparison of selection strategies for marker-assisted backcrossing of a gene. *Crop Science*, 39: 1295–1301.
- Fofana, B. and Rauber, R. (1999). Weed suppression ability of upland rice under low-input conditions in West Africa. *Weed Research*, 40: 271–280.
- Ganzel, B. (2007). *Miracle rice*. Retrieved from [http://www.livinghistoryfarm.org/farminginthe50s/crops\\_17.html](http://www.livinghistoryfarm.org/farminginthe50s/crops_17.html). Accessed November 3, 2013.
- Gomez, S. M., Boopathi, N. M., Kumar, S. S., Ramasubramanian, T., Chengsong, Z., Jeyaprakash, P., Senthil, A. and Babu, R. C. (2010). Molecular mapping and location of QTLs for drought-resistance traits in indica rice (*Oryza sativa* L.) lines adapted to target environments. *Acta Physiologiae Plantarum*, 32: 355–364.
- Gowda, V. R. P., Henry, A., Yamauchi, A., Shashidhar, H. E. and Serraj, R. (2011). Root biology and genetics improvement for drought avoidance in rice. *Field Crops Research*, 122: 1–13.
- Gramene (2002). Retrieved from <http://www.gramene.org/>. Accessed July 29, 2014.
- Greenland, D. (1997). *The sustainability of rice farming*. New York and Wallingford: CAB International.
- Guimarães, E. P. (2009). Rice breeding. In M. J. Carena (Ed.), *Cereals, the Banks and the Italian Economy* (pp. 99–126). New York: Springer US.

- Hakim, M. A., Juraimi, A. S., Ismail, M. R., Hanafi, M. M., Selamat, A. (2010). Distribution of weed population in the costal rice growing area of Kedah in Peninsular Malaysia. *Journal of Agronomy*, 9: 9-16.
- Hakim, M. A., Juraimi, A. S., Hanafi, M. M., Selamat, A., Ismail, M. R. and Anwar, M. P. (2011). Weed flora in rice field of Tanjong Karang coastal area in Malaysia. *Journal of Food, Agriculture and Environment*, 9: 694–699.
- Hakim, M. A., Juraimi, A. S., Musa, M. H., Ismail, M. R., Rahman, M. M. and Selamat, A. (2013). Impacts of weed competition on plant characters and the critical period of weed control in rice under saline environment. *Australian Journal of Crop Science*, 7: 1141–1151.
- Hasan, M. M. (2015). *Development of blast resistant rice variety through marker-assisted backcross between varieties MR263 and Pongsu Seribu 1*, PhD Thesis, Universiti Putra Malaysia.
- Hasan, M. M., Rafii, M. Y., Ismail, M. R., Mahmood, M., Rahim, H. A., Alam, M. A., Ashkani, S., Malek, M. A. and Latif, M. A. (2015a). Marker-assisted backcrossing: a useful method for rice improvement. *Biotechnology and Biotechnological Equipment*, 29: 237-254.
- Hasan, N. A., Rafii, M. Y., Rahim, H. A., Ali, N. S., Mazlan, N. and Abdullah, S. (2015b). Introgression of Pi-kh resistance gene into a Malaysian cultivar, MR264 using marker-assisted backcrossing (MABC). *International Journal of Agriculture and Biology*, 17: 1172–1178.
- Hausmann, B. I. G., Parzies, H. K., Presterl, T., Sušić, Z. and Miedaner, T. (2004). Plant genetic resources in crop improvement. *Plant Genetic Resources*, 2: 3–21.
- Hemamalini, G. S., Shashidhar, H. E. and Hittalmani, S. (2000). Molecular marker assisted tagging of morphological and physiological traits under two contrasting moisture regimes at peak vegetative stage in rice (*Oryza sativa* L.). *Euphytica*, 112: 69–78.
- Hirabayashi, H., Sato, H., Nonoue, Y., Kuno-Takemoto, Y., Takeuchi, Y., Kato, H., Nemoto H., Ogawa, T., Yano, M., Imbe, T. and Ando, I. (2010). Development of introgression lines derived from *Oryza rufipogon* and *O. glumaepatula* in the genetic background of *japonica* cultivated rice (*O. sativa* L.) and evaluation of resistance to rice blast. *Breeding Science*, 60: 604–612.
- Hittalmani, S. (2010). "More yield per field" with less and less water. Retrieved from Aerobic rice: <http://aerobicrice.org/>. Accessed November 4, 2013.



- Hospital, F. (2001). Marker-assisted backcross breeding : a case-study in genotype building theory. In M. S. Kang (Ed.), *Quantitative genetics, genomics and plant breeding* (pp. 135–141). Wallingford: CABI Publishing.
- Hospital, F. (2005). Selection in backcross programmes. *Philosophical Transactions of the Royal Society of London, Series B*, 360: 1503–1511.
- Hossain, M. and Abedin, M. Z. (2004). Rice research and development in the flood-prone ecosystem: an overview. In S. I. Bhuiyan, M. Z. Abedin, V. P. Singh, and B. Hardy (Eds.), *Proceedings of the International Workshop on Flood-Prone Rice Systems* (pp. 1-10). Los Baños: International Rice Research Institute.
- Hossain, M., Bose, M. L. and Chowdhury, A. (2004). Changes in agriculture and the economy in the flood-prone environment in Bangladesh, 1988 to 2000: insights from a repeat survey of 16 villages. In S. I. Bhuiyan, M. Z. Abedin, V. P. Singh and B. Hardy, (Eds.), *Proceedings of the International Workshop on Flood-Prone Rice Systems* (pp. 13-31). Los Baños: International Rice Research.
- Huyen, L. T. N., Cuc, L. M., Ismail, A. M. and Ham, L. H. (2012). Introgression the salinity tolerance QTLs Saltol into AS996, the elite rice variety of Vietnam. *American Journal of Plant Sciences*, 3: 981–987.
- Huke, R. E. and Huke, E. H. (1997). *Rice area by types of culture: South and South-east Asia*. Los Banos: International Rice Research Institute.
- IRRI. (1984). *Terminology for rice growing*. Los Baños: International Rice Research Institute.
- IRRI. (2003). Main weeds of rice in Asia. Retrieved from <http://www.knowledgebank.irri.org/training/fact-sheets/pest-management/weeds/main-weeds-of-rice-in-asia>. Accessed May 16, 2012.
- IRRI. (2006). Breeding program management. Retrieved from [http://www.knowledgebank.irri.org/ricebreedingcourse/Lesson\\_3\\_Pedigree\\_program\\_management.htm](http://www.knowledgebank.irri.org/ricebreedingcourse/Lesson_3_Pedigree_program_management.htm). Accessed September 26, 2015.
- IRRI. (2007). Rice Races. Retrieved from [http://www.knowledgebank.irri.org/ericeproduction/bodydefault.htm#0.5\\_Rice\\_races.htm](http://www.knowledgebank.irri.org/ericeproduction/bodydefault.htm#0.5_Rice_races.htm). Accessed April 7, 2015.
- IRRI. (2009a). The Dirty Dozen-the most troublesome weed of rice in Asia. Retrieved from [www.knowledgebank.irri.org/ipm/index.php/The-dirty-dozen](http://www.knowledgebank.irri.org/ipm/index.php/The-dirty-dozen). Accessed May 16, 2012.
- IRRI. (2009b). Water management in irrigated lowland rice. Retrieved from: <http://www.knowledgebank.irri.org/watermanagement/>. Accessed May 21, 2012.

- IRRI. (2011). Mozambique gets new designer rice. Retrieved from [http://www.irri.org/index.php?option=com\\_k2&view=item&id=10700:mozambique-gets-new-designer-rice&lang=en](http://www.irri.org/index.php?option=com_k2&view=item&id=10700:mozambique-gets-new-designer-rice&lang=en). Accessed November 4, 2013.
- Ismaila, U., Wada, A. C., Daniya, E. and Gbanguba, A. U. (2013). Meeting the local rice needs in nigeria through effective weed management. *Sustainable Agriculture Research*, 2: 37–48.
- Jalilvandy, A. and Rozrokh, M. (2013). Assessment of drought tolerance indices in bread wheat genotypes under different. *International Journal of Agriculture and Crop Sciences*, 6: 370–374.
- Jennings, P. R., Coffman, W. R. and Kauffman, H. E. (1979). *Rice improvement*. Los Banos: International Rice Research Institute.
- Johnson, D., Mortimer, M., Orr, A. and Riches, C. (2003). *Weeds, rice and poor people in South Asia*. Natural Resources Institute: Chatham.
- Johnson, D. E., Wopereis, M. C. S., Mbodj, D., Diallo, S., Powers, S. and Haefele, S. M. (2004). Timing of weed management and yield losses due to weeds in irrigated rice in the Sahel. *Field Crops Research*, 85: 31–42.
- Juraimi, A. S., Najib, M. Y. M., Begum, M., Anuar, A. R., Azmi, M. and Puteh, A. (2009). Critical period of weed competition in direct seeded rice under saturated and flooded conditions. *Pertanika Journal of Tropical Agriculture Science*, 32: 305–316.
- Juraimi, A. S., Hamdani, M. S. A., Anuar, A. R., Azmi, M., Anwar, M. P. and Uddin, M. K. (2012). Effect of water regimes on germination of weed seeds in a Malaysia rice field. *Australian Journal of Crop Science*, 6: 598-605.
- Kanbar, A. and Shashidhar, H. E. (2010). Participatory selection assisted by DNA markers for enhanced drought resistance and productivity in rice (*Oryza sativa* L.). *Euphytica*, 178: 137–150.
- Karim, S. M. R., Man, A. and Sahid, I. (2004). Weed problems and their management in rice fields of Malaysia: an overview. *Weed Biology and Management*, 4: 177-186.
- Katalani, K., Nematzadeh, G., Kiani, G. and Hashemi, S. H. (2013). Marker-assisted selection for quality improvement of Ghaem rice variety in multiple crosses at segregating population. *International Journal of Agronomy and Plant Production*, 4: 2420–2428.

- Kawano, K., Gonzalez, H. and Lucena, M. (1974). Intraspecific competition, competition with weeds and spacing response in rice. *Crop Science*, 14: 841–845.
- Khalik, A. and Matloob, A. (2011). Weed crop competition period in three fine rice cultivars under direct-seeded rice culture. *Pakistan Journal of Weed Science Research*, 17: 229-243.
- Khan, I. M. and Dhurve, O. P. (2016). Drought response indices for identification of drought tolerant genotypes in rainfed upland rice (*Oryza sativa* L.). *International Journal of Science, Environment and Technology*, 5: 73–83.
- Khanh, T. D., Linh, L. H., Linh, T. H., Ismail, A. M., Ham, L. H. and Xuan, T. D. (2012). Molecular Breeding to Improve Salt Tolerance of Rice (*Oryza sativa* L.) in the Red River Delta of Vietnam. *International Journal of Plant Genomics*, 2012: 1-9.
- Khush, G. S. (1987). *List of gene markers maintained in the Rice Genetic Stock Centre, IRRI*. Rice Genetics Newsletter 4.
- Kim, K. U. (1996). Ecological forces influencing weed competition and herbicide resistance. In R. Naylor (Ed.), *Herbicides in Asian rice: transitions in weed management* (pp. 129–144). Palo Alto and Manila: Institute for International Studies, Stanford University and International Rice Research Institute.
- Kreuzer, H. and Massey, A. (2005). *Biology and biotechnology: science and applications and issues*. Washington: ASM Press.
- Kruepl, C., Hoad, S., Davies, K., Bertholdsson, N. -O. and Paolini, R. (2007). Weed Competitiveness. In D. Donner and A. Osman (Eds.), *Handbook cereal variety testing for organic and low input agriculture* (pp. 1-16). Denmark: COST860-SUSVAR
- Kumar, U. (2008). *DNA fingerprinting*. Jodhpur: Agrobios.
- Kumar, B., Gomez, S. M., Boopathi, N. M., Kumar, S. S., Kumaresan, D., Biji, K. R., Babu, N. S. R., Shanmugasundaram, P. and Babu, R. C. (2005). Identification of microsatellite markers associated with drought tolerance in rice (*Oryza sativa* L.) using bulked line analysis. *Tropical Agricultural Research*, 17: 39–47.
- Kumar, S., Dwivedi, S. K., Singh, S. S., Jha, S. K., Lekshmy, S., Singh, O. N. and Bhatt, B. P. (2014). Identification of drought tolerant rice genotypes by analysing drought tolerance indices and morpho-physiological traits. *SABRAO Journal of Breeding and Genetics*, 46: 217–230.



- Labrada, R. (2003). The need for improved weed management in rice. *20th Session of the International Rice Commission*. Bangkok, Thailand: Agriculture and Consumer Protection.
- Lacey, S. (2012). *Climate progress: drought, flooding and 'multiple, combined outbreaks' of pests threaten to reduce Asian agricultural output 50%*. Retrieved from <http://thinkprogress.org/climate/2012/04/16/463231/drought-flooding-and-outbreaks-of-pests-threaten-to-reduce-asian-agricultural-output-50/?mobile=nc>. Accessed February 26, 2013.
- Lafitte, H. R., Price, A. H. and Courtois, B. (2004). Yield response to water deficit in an upland rice mapping population: associations among traits and genetic markers. *Theoretical and Applied Genetics*, 109: 1237–1246.
- Lafitte, H. R., Li, Z. K., Vijayakumar, C. H. M., Gao, Y. M., Shi, Y., Xu, J. L. and Mackill, D. (2006). Improvement of rice drought tolerance through backcross breeding: Evaluation of donors and selection in drought nurseries. *Field Crops Research*, 97: 77–86.
- Lakshman L. R. (1995). *Enhancement of the weed competitive ability of lowland rice (Oryza sativa L.)*. Oregon State University, United States of America.
- Lal, B., Priyanka, G. and Ekta, J. (2013). Different rice establishment methods for producing more rice per drop of water. *International Journal of Research in BioSciences*, 2: 1–12.
- Lang, N. T., Tao, N. V. and Buu, B. C. (2011). Marker-assisted backcrossing (MAB) for rice submergence tolerance in Mekong Delta. *Omonrice*, 18: 11–21.
- Langridge, P. and Chalmers, K. (2008). The principle: identification and application of molecular markers. In H. Lörz and G. Wenzel (Eds.), *Molecular marker systems in plant breeding and crop improvement* (pp. 3-21). Germany: Springer-Verlag Berlin Heidelberg.
- Lawn, R. J. and Byth, D. E. (1989). Saturated soil culture: a technology to expand the adaptation of soybean. In A. J. Pascale (Ed.), *Proceedings IV World Soybean Research Conference* (pp. 576-581). Buenos Aires: Association of Argentina de la soja.
- Lemerle, D., Veebeek, B., Cousens, R. D. and Coombes, N. E. (1996). The potential for selecting wheat varieties strongly competitive against weeds. *Weed Science*, 36: 505-513.
- Linares, O. F. (2012). African rice (*Oryza glaberrima*): History and future potential. *Proceedings of the National Academy of Science*. United States of America.

- Linh, L. H., Linh, T. H., Xuan, T. D., Ham, L. H., Ismail, A. M. and Khanh, T. D. (2012). Molecular breeding to improve salt tolerance of rice (*Oryza sativa* L.) in the Red River Delta of Vietnam. *International Journal of Plant Genomics*, 2012: 1-9.
- Li-yun, C., Ying-hui, X., Wen-bang, T. and Dong-yang, L. (2007). Practices and prospects of super hybrid rice breeding. *Rice Science*, 14: 71-77.
- Lo, N. P., Azmi, M. and Xavier, A. (1990). Sulfonylurea herbicides for transplanted rice. In K. F. Kon and S. A. Lee (Eds.), *Proceeding Third Tropical Weed Science Conference* (pp. 211-219). Kuala Lumpur: Malaysian Plant Protection Society.
- Lobo, I. (2008). Environmental influences on gene expression. *Nature Education*, 1: 39.
- Long-biao, G. and Guo-you, Y. (2014). Use of major quantitative trait loci to improve grain yield of rice. *Rice Science*, 21: 65–82.
- Lörz, H. and Wenzel, G. (2005) Molecular marker systems in plant breeding and crop improvement. New York: Springer
- Lu, J., Ookawa, T. and Hirasawa, T. (2000). The effects of irrigation regimes on the water use, dry matter production and physiological responses of paddy rice. *Plant and Soil*, 223: 207–216.
- Lutz, D. (2013). *How rice twice became a crop and twice became a weed and what it means for the future*. Retrieved from <http://news.wustl.edu/news/Pages/24751.aspx>. Accessed March 12, 2014.
- Mackill, D. J., Coffman, W. R. and Garrity, D.P. (1996). *Rainfed lowland rice improvement*. Manila: International Rice Research Institute.
- Mackill, D. J., Septiningsih, E., Pamplona, A. M., Sanchez, D. L., Labios, R. V., Heuer, S. and Ismail, A. M. (2008). Developing rice variety tolerant to abiotic stresses through marker assisted backcrossing. *Seminar Nasional Padi 2008*, (pp. 95-102). Sukamandi, Jawa Barat.
- Mackill, D. J., Ismail, A. M., Pamplona, A. M., Sanchez, D. L., Carandang, J. J. and Septiningsih, E. M. (2010). Stress tolerant rice varieties for adaptation to a changing climate. *Crop, Environment and Bioinformatics*, 7: 250-259.
- Maclean, J. L., Dawe, D. C., Hardy, B. and Hettel, G. P. (2002). *Rice almanac*. Oxon: CABI Publishing.
- Mahajan, G., Ramesha, M. S. and Chauhan, B. S. (2014). Response of rice genotypes to weed competition in dry direct-seeded rice in India. *The Scientific World Journal*, 2014: 1–8.

- Mai, C. D., Phung, N. T. P., To, H. T. M., Gonin, M., Hoang, G. T., Nguyen, K. L., Do, V. N., Courtois, B. and Gantet, P. (2014). Genes controlling root development in rice. *Rice*, 7: 1–11.
- Mallory-Smith, C. A. and Junior, E. J. R. (2003). Revised classification of herbicides by site of action for weed resistance management strategies. *Weed Technology*, 17, 605–617.
- Mammadov, J., Aggarwal, R., Buyyarapu, R. and Kumpatla, S. (2012). SNP markers and their impact on plant breeding. *International Journal of Plant Genomics*, 2012: 1–11.
- Manandhar, S., Shrestha, B. B. and Lekhak, H. D. (2007). Weeds of paddy field at Kirtipur, Kathmandu. *Scientific World*, 5: 100–106.
- Mangrauthia, S. K., Revathi, P., Agarwal, S., Singh, A. K. and Bhadana, V. P. (2014). Breeding and transgenic approaches for development of abiotic stress tolerance in rice. In P. Ahmad, M. R. Wani, M. M. Azooz and L.-S. P. Tran (Eds.), *Improvement of crops in the era of climatic changes* (pp. 153–190). New York: Springer New York.
- Manickavelu, A., Nadarajan, N., Ganesh, S. K., Gnanamalar, R. P. and Chandra Babu, R. (2006). Drought tolerance in rice: morphological and molecular genetic consideration. *Plant Growth Regulation*, 50: 121–138.
- Mardeh, A. Sio-se., Ahmadi, A., Poustini, K. and Mohammadi, V. (2006). Evaluation of drought resistance indices under various environmental conditions. *Field Crops Research*, 98: 222–229.
- Meyer, H. and Reiter, K. (1997). *Genetic engineering of rice*. Retrieved from <http://twinside.org.sg/title/rice-cn.htm>. Accessed January 6, 2013.
- Miah, G. (2015). *Development of blast resistant rice variety derived from crossing between MR219 and Pongsu Seribu 1 through marker-assisted selection*, PhD Thesis, Universiti Putra Malaysia.
- Miah, G., Rafii, M. Y., Ismail, M. R., Puteh, A. B., Rahim, H. A., Islam, N. K. and Latif, M. A. (2013). A review of microsatellite markers and their applications in rice breeding programs to improve blast disease resistance. *International Journal of Molecular Sciences*, 14: 22499–22528.
- Mian, M. A. K., Matin, M. A. and Hossain, M. A. (2007). Occurrence of weed species in transplanted Aman rice field as affected by cultivar. *Bangladesh Journal of Botany*, 36: 89–92.

- Mina-Roguel, S. S. and Duldulao, V. A. (2007). *Philippine rice production training manual*. Nueva Ecija: Philippine Rice Research Institute.
- Mohammadi, R., Armion, M., Kahrizi, D. and Amri, A. (2010). Efficiency of screening techniques for evaluating durum wheat genotypes under mild drought conditions. *International Journal of Plant Production*, 4: 11–24.
- Mohler, V. and Schwarz, G. (2005). Genotyping tools in plant breeding: from restriction fragment length polymorphisms to single nucleotide polymorphisms. In T. Nagat, H. Lörz and J. M. Widholm (Eds.), *Molecular marker systems in plant breeding and crop improvement* (pp. 23-38). Germany: Springer-Verlag Berlin Heidelberg.
- Molden, D., Oweis, T. Y., Steduto, P., Kijne, J. W., Hanjra, M. A., Bindraban, P. S., Bouman, B. A. M., Cook, S., Erenstein, O., Farahani, H., Hachum, A., Hoogeveen, J., Mahoo, H., Nangia, V., Peden, D., Sikka, A., Silva, P., Turrall, H., Upadhyaya, A. and Zwart, S. (2007). Pathways for increasing agricultural water productivity. In D. Molden (Ed.), *Comprehensive assessment of water management in agriculture, water for food, water for life: a comprehensive assessment of water management in agriculture*. London: Earthscan, Colombo.
- Mondal, B., Sarkar, M. A. R., Paul, S. K. and Zaman, F. (2013). Effect of variety and weeding regime on the yield components and yield of Aus rice. *Journal of Bangladesh Agricultural University*, 11: 17–21.
- Moody, K. (1981). *Major weeds of rice in South and Southeast Asia*. Laguna: International Rice Research Institute.
- Moradi, H., Akbari, G. A., Khorasani, S. K. and Ramshini, H. A. (2012). Evaluation of drought tolerance in corn (*Zea mays* L.) new hybrids with using stress tolerance indices. *European Journal of Sustainable Development*, 1: 543–560.
- Mueller, U. G. and Wolfenbarger, L. L. (1999). AFLP genotyping and fingerprinting. *Tree*, 14: 389–394.
- Mueen, A. K. (2012). Current status of genomic based approaches to enhance drought tolerance in rice (*Oryza sativa* L.), an overview. *Molecular Plant Breeding*, 3: 1–10.
- Munene, J. T., Kinyamario J. I., Holst N. and Mworio J. K. (2008). Competition between cultivated rice (*Oryza sativa*) and wild rice (*Oryza punctata*) in Kenya. *African Journal of Agricultural Research*, 3: 605–611.
- Naghavi, M. R., Aboughadareh, A. P. and Khalili, M. (2013). Evaluation of drought tolerance indices for screening some of corn (*Zea mays* L.) cultivars under environmental conditions. *Notulae Scientia Biologicae*, 5: 388–393.

- Naipictuasdharwad. (2009). Weed management in paddy. Retrieved from <http://agropedia.iitk.ac.in/content/weed-management-paddy>. Accessed April 13, 2015.
- Naylor, R. (1994). Herbicide use in Asian rice production. *World Development*, 22: 55–70.
- Nguyen, D. T., Nguyen, T. K. L., Pham, Q. C., Nguyen, T. H., Tran, Q. T., Dao, X. H. and Nguyen, H. T. (2004). Mapping QTLs associated with root traits related to drought resistance in Vietnamese upland rice. In D. Poland, M. Sawkins, J. M. Ribaut and D. Hoisington (Eds.), *Resilient crops for water limited environments*. Cuernavaca.
- Nogoy, F. M., Song, J. -Y., Ouk, S., Rahimi, S., Kwon, S. W., Kang, K. -K. and Cho, Y. -G. (2016). Current applicable DNA markers for marker-assisted breeding in abiotic and biotic stress tolerance in rice (*Oryza sativa* L.). *Plant Breeding and Biotechnology*, 4: 271–284.
- Norman, M. J. T., Pearson, C. J. and Searle, P. G. E. (1995). *The ecology of tropical food crops*. Cambridge: Cambridge University Press.
- Nouraein, M., Mohammadi, S. A., Aharizad, S., Mohammad, M. and Sadeghzadeh, B. (2013). Evaluation of drought tolerance indices in wheat recombinant inbred line population. *Annals of Biological Research*, 4: 113–122.
- Odero, D. C. and Rainbolt, C. (2000). *Weed management in rice*. Retrieved from <https://edis.ifas.ufl.edu/wg001>. Accessed August 9, 2015.
- Oerke, E. -C. (1999). Estimated crop losses due to pathogens, animal pests and weeds. In E. -C. Oerke, H. -W. Dehne, F. Schönbeck and A. Weber (Eds.), *Crop Production and Crop Protection: Estimated Losses in Major Food and Cash Crops* (pp. 72–741). Amsterdam: Elsevier Science.
- Owombo, P., Aregbesola, O. Z. and Adeloye, K. A. (2014). Eco-friendliness of weed management methods in organic farming: the need for extension education. *Journal of Agriculture and Sustainability*, 6: 179–199.
- Pantuwan, G., Fukai, S., Cooper, M., O'Toole, J. C. and Sarkarung, S. (1997). Germplasm development for rainfed lowland ecosystems: breeding strategies for rice in drought-prone environments. In S. Fukai, M. Cooper, and J. Salisbury (Eds.), *Breeding Strategies for Rainfed Lowland Rice in Drought-prone Environments: Proceedings of an International Workshop* (pp. 4–49). Canberra: ACIAR Proceedings.
- Papademetriou, M. K. (2000). Rice production in the Asia-Pacific region: issues and perspectives. In M. K. Papademetriou, F. J. Dent and E. M. Herath, (Eds.),



*Bridging the rice yield gap in the Asia-Pacific* (pp. 4-25). Bangkok: Food and Agriculture Organization of the United Nations Regional Office for Asia and the Pacific.

Parthasarathi, T., Vanitha, K., Lakshamanakumar, P. and Kalaiyarasi, D. (2012). Aerobic rice-mitigating water stress for the future climate change. *International Journal of Agronomy and Plant Production*, 3: 241–254.

Pathak, M. D. (1991). *Rice production in uttar pradesh: progress and suggestions for improvement*. India: Wiley Eastern Limited.

Perera, K. K., Ayres, P. G. and Gunasena, H. P. M. (1992). Root growth and the relative importance of root and shoot competition in interactions between rice (*Oryza sativa*) and *Echinochloa crus-galli*. *Weed Research*, 32: 67–76.

Pioneer. (2001). Using molecular markers in plant genetics research. Retrieved from [https://www.pioneer.com/CMRoot/Pioneer/About\\_Global/news\\_media/media\\_library/articles/MolecularMarkers.pdf](https://www.pioneer.com/CMRoot/Pioneer/About_Global/news_media/media_library/articles/MolecularMarkers.pdf). Accessed November 13, 2015.

Price, A. H., Cairns, J. E., Horton, P., Jones, H. G. and Griffiths, H. (2002a). Linking drought-resistance mechanisms to drought avoidance in upland rice using a QTL approach: progress and new opportunities to integrate stomatal and mesophyll responses. *Journal of Experimental Botany*, 53: 989–1004.

Price, A. H., Steele, K. A., Moore, B. J. and Jones, R. G. W. (2002b). Upland rice grown in soil-filled chambers and exposed to contrasting water-deficit regimes: Mapping QTLs for root morphology and distribution. *Field Crops Research*, 76: 25–43.

Priyanka, S., Jitesh, B. and Babu, S. (2012). Aerobic rice, a new approach of rice cultivation. *International Journal of Research in BioSciences*, 1: 1–6.

Postel, S. (1992). *The last oasis: facing water scarcity*. Pentonville Road: Worldwatch Institute.

Potrykus, I. (2001). Golden rice and beyond. *Plant Physiology*, 125: 1157-1161.

Qu, Y., Mu, P., Zhang, H., Chen, C. Y., Gao, Y., Tian, Y., Wen, F. and Li, Z. (2008). Mapping QTLs of root morphological traits at different growth stages in rice. *Genetica*, 133: 187–200.

Radosevich, S., Holt, J. and Ghera, C. (1997). *Weed ecology: implications for management*. USA: John Wiley & Sons Inc.

- Rajasekaran, B. (1993). *Indigenous technical practices in a rice-based farming system*. Retrieved from <http://www.ciesin.org/docs/004-195/004-195.html>. Accessed August 24, 2015.
- Rahman, M. R. and Bulbul, S. H. (2014). Effect of alternate wetting and drying (AWD) irrigation for Boro rice cultivation in Bangladesh. *Agriculture, Forestry and Fisheries*, 3: 86–92.
- Raman, A., Verulkar, S. B., Mandal, P. N., Variar, M., Shukla, V. D., Dwivedi, J. L., Singh, B. N., Singh, N. O., Swain, P., Mall, A. K., Robin, S., Chandrababu, R., Jain, A., Ram, T., Hittalmani, S., Haefele, S., Piepho, Hans-Peter and Kumar, A. (2012). Drought yield index to select high yielding rice lines under different drought stress severities. *Rice*, 5: 1–12.
- Rangel, P. H. N., Neto, F. P. M., Fagundes, P. R. R., Junior, A. M. de M., de Moraes, O. P., Schmidt, A. B., Mendonca, J. A., Santiago, C. M., Rangel, P. N., Cutrim, V. de A. and Ferreira, M. E. (2010). Development of herbicide-tolerant irrigated rice cultivars. *Brasilia*, 45, 701–708.
- Rao, A. N., Johnson, D. E., Sivaprasad, B., Ladha, J. K. and Mortimer, A. M. (2007). Weed management in direct-seeded rice. *Advances in Agronomy*, 93: 155-255.
- Rao, Y., Li, Y. and Qian, Q. (2014). Recent progress on molecular breeding of rice in China. *Plant Cell Reports*, 33, 551–564.
- Reddy, E. S. S., Verma, S. K., Xalxo, S. M., Saxena, R. R. and Verulkar, S. B. (2013). Identification of molecular marker(s) for root length in rice (*Oryza sativa* L.). *The Bioscan*, 8: 1511–1514.
- Reddy, S. and Patil, J. V. (2015). *Genetic enhancement of rabi sorghum: adapting the Indian Durras*. USA: Elsevier Inc.
- Rodenburg, J., Group, C., Agricult, I., Lucas, R. and Kaka, G. (2009). Weed competitiveness of the lowland rice varieties of NERICA in the Southern Guinea Savanna. *Field Crops Research*, 114: 411–418.
- Rosielle, A. A. and Hamblin, J. (1981). Theoretical aspects of selection for yield in stress and non-stress environment. *American Society of Agronomy*, 21: 943–946.
- Saadalla, M. M. (2008). Breeding for drought tolerance in cereals: an overview. *Journal of Agriculture and Environmental Science*, 7: 96–126.
- Sahid, I. (1989). *Sains rumpai*. Kuala Lumpur: Dewan Bahasa dan Pustaka

- Sahid, I. (1995). *Rumpai tropika: impak biologi dan pengurusan*. Bangi: Penerbit Universiti Kebangsaan Malaysia.
- Sandhu, N., Jain, S., Battan, K. R. and Jain, R. K. (2012). Aerobic rice genotypes displayed greater adaptation to water-limited cultivation and tolerance to polyethyleneglycol-6000 induced stress. *Physiology and Molecular Biology of Plants*, 18: 33–43.
- Saito, K., Azoma, K. and Rodenburg, J. (2010). Plant characteristics associated with weed competitiveness of rice under upland and lowland conditions in West Africa. *Field Crops Research*, 116: 308–317.
- Sariam, O. (2004). *Growth of non-flooded rice and its response to nitrogen fertilization*, PhD Thesis, Universiti Putra Malaysia.
- Sariam, O. and Anuar, A. R. (2010). Effects of irrigation regime on irrigated rice. *Journal of Tropical Agriculture and Food Science*, 38: 1-9.
- Scheible, W.R., Törjek, O. and Altmann, T. (2005). From markers to cloned genes: map-based cloning. In T. Nagat, H. Lörz and J. M. Widholm (Eds.), *Molecular marker systems in plant breeding and crop improvement* (pp. 55-87). Germany: Springer-Verlag Berlin Heidelberg.
- Schneider, K. A., Rosales-Serna, R., Ibarra-Perez, F., Cazares-Enriquez, B., Acosta-Gallegos, J. A., Ramirez-Vallejo, P., Wassini, N. and Kelly, J. D. (1997). Improving common bean performance under drought stress. *Crop Science*, 37: 43–50.
- Seckler, D., Barker, R. and Amarasinghe, U. (1999). Water scarcity in the twenty-first century. *International Journal of Water Resources Development*, 15: 29-42.
- Shashidhar, H. E., Kanbar, A., Toorchi, M., Raveendra, G. M., Kundur, P., Vimarsha, H. S., Soman, R., Kumar, N.G., Bekele, B.D. and Bhavani, P. (2013). Breeding for drought resistance using whole plant architecture conventional and molecular approach. In S. B. Andersen (Ed.), *Plant breeding from laboratories to fields* (pp. 151–166). InTech.
- Shen, L., Courtois, B., McNally, K., McCouch, S. and Li, Z. (1999). Developing near-isogenic lines of IR64 introgressed with QTLs for deeper and thicker roots through marker-aided selection. In O. Ito, J. O'Toole and B. Hardy (Eds.), *Proceedings of the Workshop on Genetic Improvement of Rice for Water-limited Environments* (pp. 275-292). Los Baños: International Rice Research Institute.
- Shen, L., Courtois, B., McNally, K. L., Robin, S. and Li, Z. (2001). Evaluation of near-isogenic lines of rice introgressed with QTLs for root depth through marker-aided selection. *Theoretical and Applied Genetics*, 103: 75–83.



- Shibayama, H. (1992). *Integrated management of paddy weeds in Japan*. Proceeding of International Symposium on Biological Control and Integrated Management of Paddy and Aquatic Weeds in Asia, Tsukuba.
- Shibayama, H. (2001). Weeds and weed management in rice. *Weed Biology and Management*, 1: 53-60.
- Siangliw, J. L., Jongdee, B., Pantuwan, G. and Toojinda, T. (2007). Developing KDML105 backcross introgression lines using marker-assisted selection for QTLs associated with drought tolerance in rice. *Science Asia*, 33: 207–214.
- Sikuku, P. A., Netondo, G. W., Musyimi, D. M. and Onyango, J. C. (2010). Effects of water deficit on days to maturity and yield of three NERICA rainfed rice varieties. *ARPJ Journal of Agricultural and Biological Science*, 5: 1–9.
- Singh, B. U., Rao, K. V. and Sharma, H. C. (2011). Comparison of selection indices to identify sorghum genotypes resistant to the spotted stem borer, *Chilo partellus* (Swinhoe) (Lepidoptera: Noctuidae). *International Journal of Tropical Insect Science*, 31: 38–51.
- Singh, V. K., Singh, A., Singh, S. P., Ellur, R. K., Singh, D., Krishnan, S. G., Bhowmick, P. K., Nagarajan, M., Vinod, K. K., Singh, U. D., Mohapatra, T., Prabhu, K. V. and Singh, A. K. (2013). Marker-assisted simultaneous but stepwise backcross breeding for pyramiding blast resistance genes Piz5 and Pi54 into an elite Basmati rice restorer line “PRR78.” *Plant Breeding*, 132: 486–495.
- Siti Mardina, I. (2005). *Kajian paras air berbeza ke atas populasi rumpai dan hasil padi*, Final Year Project Thesis, Universiti Putra Malaysia.
- Sleper, D. A. and Poehlman J. M. (2006). *Breeding field crop*. Iowa: Blackwell Publishing.
- Smith, R. J. (1968). Weed competition in rice. *Weed Science*, 16: 252–255.
- Smith, R. J. and Hill, J. E. (1990). Weed control technology in U.S. rice. *Pest management in rice* (pp. 314-327). Barking: Elsevier Applied Science Publishers Ltd.
- Somers, D. J. (2004). Molecular marker systems and their evaluation for cereal genetics. In P. K. Gupta and R. K. Varshney (Eds.), *Cereal genomics* (pp. 19–34). USA: Kluwer Academic Publishers.
- Steele, K. (2009). *Novel upland rice variety bred using marker-assisted selection and client-oriented breeding released in Jharkhand, India*. Associate Parliamentary Food & Health Forum GM Crops and Food Security 2010-2050: Is GM

technology simply an extension of traditional breeding methods for the production of new varieties of crops and, if not, does it matter?, London.

Steele, K. A., Price, A. H., Shashidhar, H. E. and Witcombe, J. R. (2006). Marker-assisted selection to introgress rice QTLs controlling root traits into an Indian upland rice variety. *Theoretical and Applied Genetics*, 112: 208–221.

Sukhla, B. (2012). What is the importance of water for plants? Retrieved from <http://www.preservearticles.com/201101012194/importance-of-water-for-plants.html>. Accessed February 26, 2015.

Sunyob, N., Juraimi, A., Hakim, M. A., Man, A., Selamat, A. and Alam, A. (2015). Competitive ability of some selected rice varieties against weed under aerobic condition. *International Journal of Agriculture and Biology*, 17: 61–70.

Suria, A. S. M. J., Juraimi, A. S., Rahman, M. M., Man, A. B. and Selamat, A. (2011). Efficacy and economics of different herbicides in aerobic rice system. *African Journal of Biotechnology*, 10: 8007–8022.

Swamy, B. P. M. and Kumar, A. (2012). Sustainable rice yield in water-short drought-prone environments: conventional and molecular approaches. In T. S. Lee (Ed.), *Irrigation systems and practices in challenging environments* (pp. 149-168). InTech.

Taharim, N. (2011). System of Rice Intensification (SRI) in Malaysia: giving credit where credit is due. Retrieved from <https://pertanianselangor.wordpress.com/2011/10/29/system-of-rice-intensification-in-malaysia-sri/>. Accessed March 24, 2015.

Takai, T., Kondo, M., Yano, M. and Yamamoto, T. (2010). A quantitative trait locus for chlorophyll content and its association with leaf photosynthesis in rice. *Rice*, 3: 172–180.

Tan, K. Z., Radziah, O., Halimi, M. S., Khairuddin, A. R. and Shamsuddin, Z. H. (2015). Assessment of plant growth-promoting rhizobacteria (PGPR) and rhizobia as multi-strain biofertilizer on growth and N<sub>2</sub> fixation of rice plant. *Australian Journal of Crop Science*, 9, 1257–1264.

Tanweer, F. A. (2015). *Development of blast resistance rice variety derived from crossing MR219 and Pongsu Seribu 2 through marker-assisted backcross breeding*, PhD Thesis, Universiti Putra Malaysia.

Tanweer, F. A., Rafii, M. Y., Sijam, K., Rahim, H. A., Ahmed, F., Ashkani, S. and Latif, M. A. (2015). Introgression of blast resistance genes (Putative Pi-b and Pi-kh) into elite rice cultivar MR219 through marker-assisted selection. *Frontiers in Plant Science*, 6: 1–11.

- Terano, R., Mohamed, Z., and Din, N. S. Z. (2016). Determinants of farmers' adoption of Clearfield production system in Malaysia. *Agriculture and Agricultural Science Procedia*, 9, 103–107.
- Thanh, N. D., Thi, N., Lien, K. and Chung, P. Q. (2006). Mapping QTLs associated with root traits related to drought resistance. *ASEAN Journal for Science and Technology Development*, 23: 323–332.
- Tian, F., De, J. L., Fu, Q., Zuo, F. Z., Yong, C. F., Xiang, K. W. and Chuan, Q. S. (2006). Construction of introgression lines carrying wild rice (*Oryza rufipogon* Griff.) segments in cultivated rice (*Oryza sativa* L.) background and characterization of introgressed segments associated with yield-related traits. *Theoretical and Applied Genetics*, 112: 570–580.
- Tripathi, K. K., Govila, O. P., Warriar, R. and Ahuja, V. (2011). *Biology of Oryza sativa L. (rice)*. India: Ministry of Environment and Forests Government of India.
- Troedson, R. J., Garside, A. L., Lawn, R. J., Byth, D. E. and Wilson, G. L. (1985). Saturated soil culture-an innovative water management option for soybean in the tropics and subtropics. In S. Shanmugasundaram and E. W. Sulzberger (Eds.), *Soybean in tropical and subtropical cropping systems* (pp. 171-180). Shanhua: Asian Vegetable Research and Development Centre.
- Tuberosa, R. and Salvi, S. (2005). Molecular marker systems and their evaluation for cereal genetics. In P. K. Gupta and R. K. Varshney (Eds.), *Cereal genomics* (pp. 19–34). USA: Kluwer Academic Publishers.
- Tuong, T. P. and Bouman, B. A. M. (2003). Rice production in water scarce environments. In J. W. Kijne, R. Barker and D. Molden (Eds.), *Water productivity in agriculture: limits and opportunities for improvement* (pp. 53–67). Wallingford: CABI Publishing.
- Tuong, T. P. *Alternate wetting and drying irrigation (AWD): a technology for water saving in rice production*. Paper presented at the Crop Cutting Ceremony, Tangail. May 2007.
- Uga, Y., Okuno, K. and Yano, M. (2011). Dro1, a major QTL involved in deep rooting of rice under upland field conditions. *Journal of Experimental Botany*, 62, 2485–2494.
- Uga, Y., Sugimoto, K., Ogawa, S., Rane, J., Ishitani, M., Hara, N., Kitomi, Y., Inukai, Y., Ono, K., Kanno, N., Inoue, H., Takehisa, H., Motoyama, R., Nagamura, Y., Wu, J., Matsumoto, T., Takai, T., Okuno, K. and Yano, M. (2013). Control of root system architecture by *DEEPER ROOTING 1* increases rice yield under drought conditions. *Nature Genetics*, 45: 1097–1102.

- Uphoff, N. T. (2007). *The System of rice intensification (SRI): an efficient, economical and ecologically friendly way to increase productivity*. Pesticide Action Network Asia and the Pacific Rice Sheets: Malaysia.
- Utusan Online. (2005). Kualiti Maswangi setanding beras wangi Siam. Retrieved from [http://www1.utusan.com.my/utusan/info.asp?y=2005&dt=0206&sec=Muka\\_Hadapan&pg=mh\\_02.htm#ixzz3iQdEl1IY](http://www1.utusan.com.my/utusan/info.asp?y=2005&dt=0206&sec=Muka_Hadapan&pg=mh_02.htm#ixzz3iQdEl1IY). Accessed August 11, 2015.
- van Berloo, R. (2008). GGT 2.0: Versatile software for visualization and analysis of genetic data. *Journal of Heredity*, 99: 223-236.
- Valencia, C. (2015). Philrice breeding drought-tolerant rice varieties. Philippines. Retrieved from <http://www.philstar.com/agriculture/2015/05/31/1460476/philrice-breeding-drought-tolerant-rice-varieties>. Accessed August 26, 2015.
- Vasant, D. V. (2012). *Genome wide association mapping of drought resistance traits in rice (Oryza sativa L.)*, Master Thesis, Tamil Nadu Agricultural University.
- Venuprasad, R., Lafitte, H. R. and Atlin, G. N. (2007). Response to direct selection for grain yield under drought stress in rice. *Crop Science*, 47: 285-293.
- Venuprasad, R., Bool, M. E., Quiatchon, L. and Atlin, G. N. (2012). A QTL for rice grain yield in aerobic environments with large effects in three genetic backgrounds. *Theoretical and Applied Genetics*, 124: 323-332.
- Vial, L. (2007). *Grains Research Update (GRDC) for irrigation croppers: aerobic and alternate wet-and-dry (AWD) rice systems can we increase water use efficiency?* Griffith: Irrigation Research & Extension Committee.
- Vogel, K. E. (2009). Backcross breeding. In M. P. Scott, (Ed.), *Transgenic maize: methods and protocols* (pp. 161-169). Iowa: Humana Press.
- Wade, L. J. (1999). Critical characteristics of rainfed rice environments and implications for rice improvement. In O. Ito, J. O'Toole and B. Hardy (Eds.), *Proceedings of the Workshop on Genetic Improvement of Rice for Water-limited Environments* (pp.1-10). Los Baños: International Rice Research Institute.
- WARDA. (2000). *WARDA anual report 1999*. West Africa Rice Development Association: Bouake, Cote d'ivoire.
- Webster, E. and Levy, R. (2009). Weed management. In J. Saichuk (Ed.), *Louisiana rice production handbook* (pp. 46-71). Louisiana State University Agricultural: United States of America.

- Weerakoon, W. M. W., Priyadarshani, T. N. N., Priyadarshani, C. H. and Silva, L. S. (2010). Impact of water saving irrigation systems on water use, growth and yield of irrigated lowland rice. In P. Weligamage, G. G. Godaliyadda and K. Jinapala (Eds.), *National Conference on Water, Food Security and Climate Change in Sri Lanka: Irrigation for Food Security* (pp. 57–63). Colombo: International Water Management Institute.
- Wibawa, W., Mohayidin, M. G., Mohamad. R. B., Juraimi, A. S. and Omar, D. (2010). Efficacy and cost-effectiveness of three broad-spectrum herbicides to control weeds in immature oil palm plantation. *Pertanika Journal of Tropical Agricultural Science*, 33: 233-241.
- Wijayanto, A. (2013). *Aplikasi marka molekuler dalam seleksi populasi silang balik IR64/Hawara Bunar untuk mendapatkan galur padi toleran aluminium*, Master Thesis, Institut Pertanian Bogor.
- Williams, J. F., Roberts, S. R., Hill, J. E., Scardaci, S. C. and Tibbits, G. (1990). Managing water for weed control in rice. *California Agriculture*, 44: 7-10.
- Wilson, S. (2014). BASF announces new herbicide tolerant rice system in development. *BASF News Release*. United States. Retrieved from <https://www.basf.com/us/en/company/news-and-media/news-releases/2014/06/p-13-597.html>. Accessed February 18, 2018.
- Wilson, R. G., Kerr, E. D. and Nelson, L. A. (1985). Potential for using weed seed content in the soil to predict future weed problems. *Weed Science*, 33: 171-175.
- Witcombe, J. R., Hollington, P. A., Howarth, C. J., Reader, S. and Steele K. A. (2008). Breeding for abiotic stresses for sustainable agriculture. *Philosophical Transaction of the Royal Society B*, 363: 703-716.
- Wu, W. and Cheng, S. (2014). Root genetic research, an opportunity and challenge to rice improvement. *Field Crops Research*, 165: 111–124.
- Xu, Y. (2010). *Molecular plant breeding*. Mexico: CAB International.
- Yaduraju, J. S. and Mishra, N. T. (2008). Sedges in rice culture and their management. In Y. Singh, V. P. Singh, B. Chauhan, A. Orr, A. M. Mortimer, D. E. Johnson and B. Hardy (Eds.), *Direct seeding of rice and weed management in the irrigated rice-wheat cropping system of the Indo-Gangetic Plains* (p. 272). Los Baños and Pantnagar: International Rice Research Institute and Directorate of Experiment Station, G.B. Pant University of Agriculture and Technology.
- Yamauchi, A., Inukai, Y., Kano, M., Suralta, R., Ogawa, A., Kobayashi, N. and Serraj, R. (2008). Role of roots for stabilizing rice production under water stress



conditions. In *5th International Crop Science Congress*. Jeju: International Crop Science Society.

- Ye, G., Ogonnaya, F. and Ginkev, M. V. (2009). Marker-assisted recurrent backcrossing in cultivar development. In R. K. Singh, R. Singh, G. Ye, A. Selvi and G. P. Rao (Eds.), *Molecular plant breeding: principle, method and application* (pp. 295–319). USA: Studium Press LLC.
- Yi, M., Nwe, K. T., Vanavichit, A., Chai-arree, W. and Toojinda, T. (2009). Marker assisted backcross breeding to improve cooking quality traits in Myanmar rice cultivar Manawthukha. *Field Crops Research*, 113: 178–186.
- Yoshida, S. (1972). Physiological aspects of grain yield. *Annual Review of Plant Physiology*, 23: 437–464.
- Yoshida, S. (1975). Factors that limit the growth and yields of upland rice. In *Major research in upland rice* (pp. 46-71). Los Baños: International Rice Research Institute.
- Yoshida, S. and Hasegawa, S. (1982). The rice root system: its development and function. In M. R. Vega (Ed.), *Drought resistance in crops with emphasis on rice* (pp. 97–114). Los Banos: International Rice Research Institute.
- Zainudin, H., Sariam, O., Chan, C. S., Azmi, M., Saad, A., Alias, I. and Marzukhi, H. (2014). Performance of selected aerobic rice varieties cultivated under local condition. *Journal of Tropical Agriculture and Food Science*, 42: 175–182.
- Zhang, Z. P. (2001). *Weed management in China*. Paper presented at FAO Workshop on *Echinochloa spp.* control. Beijing.
- Zhang, J. E., Xu, R., Chen, X. and Quan, G. (2009). Effects of duck activities on a weed community under a transplanted rice-duck farming system in southern China. *Weed Biology and Management*, 9: 250-257.
- Zhao, D. L. (2006). *Weed competitiveness and yielding ability of aerobic rice genotypes*, PhD Thesis, Wageningen University.
- Zhao, D. L., Atlin, G. N., Amante, M., Cruz, M. T. S. and Kumar, A. (2010). Developing aerobic rice cultivars for water-short irrigated and drought-prone rainfed areas in the tropics. *Crop Science*, 50: 2268–2276.
- Ziska, L. H. and Dukes, J. S. (2011). *Weed biology and climate change*. Iowa: Blackwell Publishing Ltd.

Zubaer, M. A., Chowdhury, A. K. M. M. B., Islam, M. Z., Hasan, M. A. and Ahmed, T. (2007). Effects of water stress on growth and yield attributes of Aman rice genotypes. *International Journal of Sustainable Crop Production*, 2: 25–30.

Zulkarnain, W. M., Ismail, M. R., Ashrafuzzaman, M., Saud, H. M. and Haroun, I. C. (2009). Rice growth and yield under rain shelter house as influenced by different water regimes. *International Journal of Agriculture and Biology*, 11: 566–570.

