



UNIVERSITI PUTRA MALAYSIA

***DISTRIBUTION AND SOURCE OF HYDROCARBONS IN THE
SEDIMENTS OF THE SELANGOR RIVER, MALAYSIA***

NAJAT ABDULLAH EBRAHIM MASOOD

FPAS 2018 10



**DISTRIBUTION AND SOURCE OF HYDROCARBONS IN THE
SEDIMENTS OF THE SELANGOR RIVER, MALAYSIA**

By

NAJAT ABDULLAH EBRAHIM MASOOD

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for Degree of Doctor of Philosophy**

July 2017


All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

In the name of Allah, the Beneficent, the Merciful



To my beloved husband and my daughters

To the memory of my late father

To my beloved mother and my dear siblings

In dedicate this dissertation

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement of the degree of Doctor of Philosophy

DISTRIBUTION AND SOURCE OF HYDROCARBONS IN THE SEDIMENTS OF THE SELANGOR RIVER, MALAYSIA

By

NAJAT ABDULLAH EBRAHIM MASOOD

July 2017

Chairman : Normala bt. Halimoon, PhD

Faculty : Environmental Studies

Environmental pollutants such as anthropogenic hydrocarbons (AHs) are known to result from increased urbanization and act as indicators of anthropogenic activity. Hydrocarbons contamination is a great global concern due to their negative effects on human health and environmental consequences, such as toxic, mutagenic, carcinogenic, teratogenic and hepatotoxic effects. Consequently, a clear understanding and monitoring of distribution, sources, pathways and fate in the environment and ecosystems of AHs is of high importance in sediment samples collected from Kuala Selangor River, which is located on the west coast of Peninsular Malaysia, in order to determine the distributions and sources of hopanes, alkanes, polycyclic aromatic hydrocarbons (PAHs), and linear alkylbenzenes (LABs) in four periods. The sediment samples were collected from the river and its estuary during the rainy inter-monsoonal period, the dry inter-monsoonal period, the North-East Monsoon (NEM) period, and the South-West Monsoon (SWM) period because during these periods the samples are expected to contain varying amounts of pollutants. The collected samples were homogenized, freeze-dried, extracted, cleanup, fractionated and analyzed using gas chromatography mass spectrometry. The results of this study revealed that the concentrations of PAHs for the 13 sampling sites pooled over the four studied periods ranged from 341.33 ng g⁻¹ dw (in the SWM period) to 651.64 ng g⁻¹ dw (in the wet inter-monsoon period) with the mean value of 471.05 ng g⁻¹ dw. In other respects, the outcomes of the one-way ANOVA indicated that the differences between the PAHs among the sampling stations were significant ($p < 0.05$). The diagnostic ratios of individual PAHs in sediments indicated both petrogenic and pyrogenic origin PAHs with significant dominance of pyrogenic source. The comparison of PAHs with Sediment Quality Guidelines (SQGs) indicates that the levels of the PAHs in the surface sediments of Selangor River are unlikely to cause any adverse biological effects to the aquatic organisms, including those that are closely associated with sediments. No significant correlation was found between AHs and organic carbon, which means that the distribution of AHs was not affected by the organic carbon content. In the four studied periods, the concentrations of the LABs in Selangor River ranged from 23.70 to 113.30 ng g⁻¹ dw with an arithmetical mean of 54.04 ± 22.34 ng

g⁻¹ dw. The average concentration of the total LABs was found to be higher in the wet season (65.20 ± 25.04 ng g⁻¹ dw) than in the other seasons. The average values of the ratio of long-chain to short-chain LAB (L/S), C_{13}/C_{12} , and the ratio of Internal to External isomers (I/E) in the sediments of Selangor River during the four studied periods were 2.03 ± 1.06 , 1.74 ± 0.28 , and 0.6 ± 0.17 , respectively, indicating inputs of LABs from direct wastewater discharges. The I/E ratio decreased from upstream to downstream of Selangor River. This indicates that the sedimentary LABs downstream of the river (estuary) were more degraded than those in the upstream. The concentrations of total alkanes (nC₁₀-nC₃₆) ranged from 967 to 3,711 µg g⁻¹ dw in sediment samples. In addition, the results of this study indicated that the alkanes in the sediment samples originated from diverse sources. Fresh oil, terrestrial plants (riverine area), and heavy/degraded oil (estuarine area) were the predominant source of alkanes in sediments. The alkanes originating from higher plants were mostly predominant in sediments from the stations located in upper parts of the river. The levels of hopanes ranged from 235 to 1,044 ng g⁻¹ dw in sediments. The C₂₉/C₃₀ hopanes ratios were similar with those of used crankcase oil and Middle East Crude Oil (MECO), suggesting MECO as a major source of petroleum hydrocarbons (PHCs) in the sediments of Selangor River. In sum, presence of selected hydrocarbons (PAHs, LABs, alkanes, and hopanes) in the sediments of Selangor River, Malaysia, was studied during four periods. The results show that the agricultural, urban, and industrial activities in the state of Selangor, coupled with a high population growth, have caused deterioration in the quality of its river water. The concentrations of the investigated hydrocarbons varied from sampling site to another and from one period to another. Further studies should be conducted to address this problem.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

TABURAN DAN SUMBER HYDROCARBON DALAM PERMUKAAN SEDIMEN SUNGAI SELANGOR, MALAYSIA

Oleh

NAJAT ABDULLAH EBRAHIM MASOOD

Julai 2017

Pengerusi : Normala bt. Halimoon, PhD

Fakulti : Pengajian Alam Sekitar

Pencemar alam sekitar seperti antropogenik hidrokarbon (AHs), diketahui berpunca akibat daripada peningkatan perbandaran dan bertindak sebagai petunjuk aktiviti antropogenik. Pencemaran hidrokarbon menjadi kebimbangan global yang besar disebabkan oleh kesan negatifnya terhadap kesihatan manusia dan alam sekitar, seperti kesan toksik, mutagenik, karsinogenik, teratogenik dan hepatoksik. Oleh itu, pemahaman dan pemantauan yang jelas mengenai taburan, sumber, laluan dan nasib dalam alam sekitar dan ekosistem AHs adalah sangat penting di dalam sampel sedimen yang diambil dari Sungai Kuala Selangor, yang terletak di pantai barat Semenanjung Malaysia, dalam menentukan taburan dan sumber hopan, alkana, hidrokarbon aromatik polisiklik (PAHs), dan alkilbenzena linear (LABs) dalam empat tempoh. Sampel sedimen diambil dari sungai dan muara sungai tersebut ketika tempoh peralihan monsun hujan, tempoh peralihan monsun kering, tempoh monsun timur laut (NEM), dan tempoh monsun barat daya (SWM) kerana pada waktu tersebut, sampel dijangkakan mengandungi pelbagai bahan pencemar. Sampel yang diambil telah dihomogenkan, dibekukeringkan, diekstrak, dibersihkan, difraksinasi dan dianalisa menggunakan kromatografi gas spektrometer jisim. Keputusan kajian ini menunjukkan bahawa kepekatan PAHs yang diperolehi dari 13 kawasan persampelan untuk empat tempoh kajian berjulat antara 341.33 ng g⁻¹ dw (dalam tempoh SWM) hingga 651.64 ng g⁻¹ dw (dalam tempoh basah intermonsun) dengan nilai purata 471.05 ng g⁻¹ dw. Dalam aspek lain pula, hasil daripada ANOVA sebelah menunjukkan bahawa perbezaan PAHs antara setiap stesen persampelan adalah signifikan ($p < 0.05$). Nisbah diagnostik PAHs secara individu dalam sedimen menunjukkan kedua-dua PAHs adalah petrogenik dan pirogenik dengan sumber pirogenik yang paling ketara. Perbandingan PAHs dengan panduan kualiti sedimen (SQGs) menunjukkan paras PAHs di permukaan sedimen Sungai Selangor bermungkinan rendah untuk memberikan kesan biologi ketara pada organisma akuatik, termasuklah yang berkaitan dengan sedimen. Tiada hubungan signifikan antara pengagihan AHs dan karbon organik di mana, taburan AHs tidak terkesan dengan kandungan karbon organik. Dalam keempat-empat tempoh kajian, kepekatan LABs di Sungai Selangor berjulat antara 23.70 ke 113.30 ng g⁻¹ dw dengan purata aritmetik 54.04 ± 22.34 ng g⁻¹ dw. Kepekatan purata LABs didapati lebih tinggi pada musim hujan (65.20 ± 25.04 ng g⁻¹ dw) berbanding musim-musim lain.

Nilai purata nisbah rantai panjang terhadap rantai pendek LAB (L/S), C_{13}/C_{12} , dan nisbah dalaman untuk isomer dalaman terhadap isomer luaran (I/E) dalam sedimen Sungai Selangor untuk keempat-empat tempoh kajian adalah 2.03 ± 1.06 , 1.74 ± 0.28 dan 0.6 ± 0.17 menunjukkan kehadiran LABs adalah berasal secara terus daripada pelepasan air kumbahan. Nisbah I/E menunjukkan penurunan dari hulu ke hilir Sungai Selangor. Ini menunjukkan bahawa sedimen LABs di kawasan hilir sungai (muara) lebih terurai daripada hulu sungai. Kepekatan jumlah alkana (nC_{10} - nC_{36}) adalah berjalut antara 967 ke 3,711 $\mu g\ g^{-1}$ dw dalam sampel-sampel sedimen. Tambahan pula, keputusan kajian ini menunjukkan bahawa alkana dalam sampel sedimen berasal dari pelbagai sumber. Minyak segar, tumbuhan daratan (kawasan sungai) dan minyak berat/terurai (kawasan muara) merupakan punca utama alkana di dalam sedimen. Alkana yang berasal daripada tumbuh-tumbuhan kompleks adalah paling ketara di dalam sedimen stesen yang terletak di hulu sungai. Paras hopana dalam sedimen pula adalah antara 235 hingga 1,044 $ng\ g^{-1}$ dw dalam sedimen. Nisbah C_{29}/C_{30} hopana adalah sama dengan nisbah bagi minyak crankcase dan Minyak Mentah Timur Tengah (MECO), mencadangkan bahawa MECO menjadi sumber utama hidrokarbon petroleum (PHCs) dalam sedimen Sungai Selangor. Keseluruhannya, kehadiran hidrokarbon terpilih (PAHs, LABs, alkana, dan hopana) dalam sedimen Sungai Selangor, Malaysia, telah dikaji sepanjang empat tempoh. Hasil kajian menunjukkan bahawa aktiviti-aktiviti pertanian, perbandaran dan perindustrian di negeri Selangor, ditambah pula dengan pertumbuhan penduduk yang tinggi, telah menyebabkan kemerosotan kualiti air sungai tersebut. Kepekatan hidrokarbon adalah berbeza di antara setiap lokasi dan tempoh persampelan kajian selanjutnya perlu dijalankan untuk menangani masalah ini.

ACKNOWLEDGEMENT

I would like to thank God for making this dream come true. Many people have contributed to the successful completion of this work. Special thanks are due to my Supervisor, Dr. Normala Halimoon, for her guidance, support, motivation, stimulating feedback, and critical questions. Her insight and support of all aspects of this project have been crucial and invaluable. I am most indebted and grateful to my supervisory committee members, Professor Ahmad Zaharin Aris, Professor Shuhaimi Mustafa, and Associate Prof. Masni Mohd Ali, Prof. Mohamad Pauzi B Zakaria for volunteering their time to read my thesis and giving me unlimited support, valuable suggestions, and encouragement throughout working on this project.

I would like to express my deepest gratitude to my lovely father, my lovely mother, and my dear siblings and relatives who have been the nucleus of constant inspiration, love, encouragement, and moral support during the study period.

I express my heartiest thanks and gratitude to my husband and my daughters for their boundless patience, profound love, and continuous encouragement and care. I am proud of you all, now and forever.

Last but not least, I would like to extend my gratitude to my dear friends especially Mr. Muhamad Shirwan, Dr. Vahab Vaezzadeh, Dr. Mehrzad Keshavarzifard, Dr. Nabeel M. Gazzaz, Dr. Najat Al-Odaini, Sadeq Alkhadher, Mudher Hussein, and Muhammad Raza.

This thesis was submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of supervisory committee were as follow:

Normala bt. Halimoon, PhD

Senior Lecturer
Faculty of Environmental Studies
Universiti Putra Malaysia
(Chairman)

Ahmad Zaharin Aris, PhD

Professor
Faculty of Environmental Studies
Universiti Putra Malaysia
(Member)

Shuhaimi Mustafa, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Masni Mohd Ali, PhD

Associate Professor
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(Member)

Mohamad Pauzi B Zakaria, PhD

Professor
Faculty of Environmental Studies
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Najat Abdullah Ebrahim Masood, GS35263

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of Chairman of
Supervisory
Committee: Normala bt. Halimoon

Signature: _____
Name of Member of
Supervisory
Committee: Ahmad Zaharin Aris

Signature: _____
Name of Member of
Supervisory
Committee: Shuhaimi Mustafa

Signature: _____
Name of Member of
Supervisory
Committee: Masni Mohd Ali

TABLE OF CONTENTS

ABSTRACT	Page
ABSTRAK	i
ACKNOWLEDGEMENT	iii
APPROVAL	v
DECLARATION	vi
LIST OF TABLES	viii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xvi
	xviii

CHAPTER

1	INTRODUCTION	1
	1.1 Background of the study	1
	1.2 Problem statement	2
	1.3 Significance of the study	2
	1.4 Research questions	3
	1.5 Scope of the study	3
	1.6 Objectives of the study	4
	1.7 Organization of the thesis	4
2	LITERATURE REVIEW	6
	2.1 Introduction	6
	2.1.1 Pollutants in the environment	6
	2.1.2 Pollutant movement	7
	2.1.3 Studies on hydrocarbon pollution	8
	2.1.4 Hydrocarbon pollution in Southeast Asia and Malaysia	8
	2.1.5 Sources of hydrocarbon pollution in Malaysia	9
	2.1.6 Use of molecular markers in the pollution studies	10
	2.1.7 Hydrocarbons in polluted environments	11
	2.1.8 Anthropogenic molecular markers	11
	2.2 Introduction to the hydrocarbon compounds under study	12
	2.2.1 Normal alkanes (alkanes)	12
	2.2.2 Hopanes	19
	2.2.3 Linear alkylbenzenes (LABs)	23
	2.2.3.1 Sources, transport, and distribution of LABs	26
	2.2.3.2 Internal to external isomers ratio (I/E Ratio)	26
	2.2.3.3 Biodegradation of LABs	27
	2.2.4 Polycyclic aromatic hydrocarbons (PAHs)	28
	2.2.4.1 Presence of PAHs in the environment	30
	2.2.4.2 Molecular structure and chemical properties of PAHs	32

2.2.4.3	Sources, distribution, and fate of PAHs in aquatic compartments	35
2.2.4.4	Molecular weight profile of PAHs	40
2.2.4.5	Diagnostic mass ratio of PAHs isomers	41
2.2.4.6	Distribution of the PAHs in the environment (air, water, sediments, and biota)	44
2.2.4.7	Surface sediment and grain size fractions	45
2.2.4.8	The PAHs and organic carbon in sediments	46
2.2.4.9	The PAHs in soil	47
2.2.4.10	The PAH pollution in the mangrove and estuary ecosystems	48
2.2.4.11	Routes of exposure to PAHs	51
2.2.5	Evaluation of the ecotoxicological risk of PAHs	52
2.2.5.1	Sediment quality guidelines (SQGs)	52
2.2.5.2	The toxic equivalency factor approach	54
2.3	Studies of PAHs in Malaysia and Southeast Asia	57
3	METHODOLOGY	59
3.1	Sampling	59
3.1.1	Study area	59
3.1.2	Sampling locations	61
3.2	Sample collection	62
3.3	Chemicals and materials	64
3.3.1	Organic solvents	64
3.3.2	Preparation of sodium sulfate (Na_2SO_4) and silica gel	64
3.4	Preparation of sample	64
3.5	Soxhlet extraction and fractionation of the hydrocarbons	64
3.6	Instrumental Analysis	68
3.6.1	Analysis of alkanes	68
3.6.2	Analysis of hopane	68
3.6.3	Analysis of LABs	69
3.6.4	Analysis of PAHs	70
3.7	Quality assurance (QA) and quality control (QC)	73
3.8	Total Organic Carbon (TOC) Analysis	74
3.9	Calibration and chromatograms	75
3.10	Analyses of grain size	77
3.11	Statistical analysis	77
3.12	Risk assessment of PAHs	77

4	RESULTS AND DISCUSSION	79
4.1	Grain-size distribution	79
4.2	Polycyclic aromatic hydrocarbons (PAHs) in the sediments of Selangor River	79
4.2.1	Levels of PAHs in the sediments of Selangor River	79
4.2.2	Composition of PAHs in the sediments of Selangor River	82
4.2.3	Sources of PAHs in the sediments of Selangor River	87
4.2.3.1	Source identification based on the composition of the PAHs	87
4.2.3.2	Evaluation of the diagnostic ratio	90
4.2.3.3	Identification of the sources of PAHs in the sediments of Selangor River using clustering analysis	96
4.2.3.4	Identification of the sources of PAHs in the sediments of Selangor River using Principal Components Analysis (PCA)	97
4.2.4	Spatial distributions of PAHs in the sediments of Selangor River	100
4.2.5	Assessment of toxicity of the sediments of Selangor River in terms of PAHs	101
4.2.6	Carcinogenic potencies of the PAHs (The toxicity equivalency quotient approach)	103
4.2.7	Correlations between water quality parameters and the PAHs in the sediments of Selangor River	106
4.2.8	Percentages of TOC in the sediments	108
4.2.9	Comparison of the levels of PAHs in sediments of Selangor River with those in other areas	110
4.3	Linear alkylbenzenes (LABs) in the sediments of Selangor River	113
4.3.1	Levels of LABs in the sediments of Selangor River	113
4.3.2	Variability in the LABs compositional profiles	107
4.3.3	Assessment of degradation of the sediments of Selangor River in terms of LABs	119
4.3.4	Inter-correlations among LABs, PAHs, and TOC	124
4.4	Alkanes in the sediments of Selangor River	125
4.4.1	Levels of alkanes in the sediments of Selangor River	125
4.4.2	Sources of alkanes in the Sediment of Selangor River	132
4.4.2.1	Unresolved complex mixture (UCM)	132
4.4.2.2	The carbon preference index (CPI) for alkanes	132
4.4.2.3	Average chain length (ACL)	135
4.4.2.4	Terrigenous aquatic ratio (TAR)	136
4.4.2.5	Correlation between total alkanes and TOC	137

4.5	Hopanes in the sediments of Selangor River	137
4.5.1	Levels and homogeneous hopanes composition in Selangor River sediments	137
4.5.2	Identification and apportionment of the sources of the PAHs in Selangor River	141
5	CONCLUSIONS AND RECOMMENDATIONS FOR FUTURE RESEARCH	143
5.1	Summary and general conclusion	143
5.2	Recommendations for future research	145
	REFERENCES	147
	APPENDICES	187
	BIODATA OF STUDENT	205
	LIST OF PUBLICATIONS	206

LIST OF TABLES

Table		Page
2.1	Classification of the anthropogenic markers	12
2.2	Indices related to n-alkane and their application in evaluation of the types of sources of petroleum hydrocarbon	17
2.3	The concentrations (ng g ⁻¹) and indices of alkanes in the sediments	18
2.4	Concentrations (ng g ⁻¹) and indices of hopanes in different types of samples in Peninsular Malaysia	21
2.5	Diagnostic ratios of hopanes in the crude oil, fresh and used crankcase oil, street dust, and asphalt	23
2.6	Percentages of low molecular weight PAHs in the global emission of PAHs	31
2.7	Physical and chemical properties of the 16 PAHs designated as 'Priority Pollutants' by the USEPA	34
2.8	Major sources of PAHs in the environment	37
2.9	Source markers	38
2.10	Diagnostic mass ratios for identification of petrogenic and pyrogenic sources of PAHs	42
2.11	Empirical SQGs for PAHs in the Sediment	53
2.12	Classification of potentially carcinogenic PAHs	54
2.13	Maximum levels of benzo(a)pyrene (ng g ⁻¹) in foodstuffs	55
2.14	Sediment potential toxicity classification	56
2.15	The total landuse and landcover areas in Kuala Selangor	58
3.1	Sampling locations	62
3.2	Molecular structures of PAH surrogate internal standard (SIS) and internal injection standard (IIS)	65
3.3	The Surrogate Internal Standard (SIS) used to quantify each LAB compound	70
3.4	Condition of GCMS for PAHs analysis	71
3.5	Selected ions monitored in this study	72
3.6	Estimated Detection and Quantification Limits for the target PAHs	74
3.7	Toxicity Equivalency Factors (TEF) for individual PAHs	78
4.1	Grain size distributions in the sediments of the study area	79
4.2	Results of One-Way ANOVA in the total concentrations of PAHs in the river sediments in different seasons	81
4.3	Average concentrations of individual PAHs in the sediments of Selangor River	84
4.4	Correlations between the levels of LMW PAH, HMW PAH, TPAH, 16USEPA PAHs, cPAHs, and TOC	86
4.5	Average values of different diagnostic ratios of PAHs in the sediments of Selangor River	88
4.6	Rotated component matrix of PAHs in Selangor River sediments (High loadings (> 0.65) shown in bold)	98
4.7	Proposed sediment quality guidelines (SQGs) in terms of PAH content	102
4.8	Concentrations of the total and the carcinogenic PAHs and their corresponding toxicity equivalency quotients (TEQs) in the sediments of Selangor River	104
4.9	Correlations between water quality parameters and the PAHs in the sediments of Selangor River (<i>n</i> = 62)	107

4.10	Levels of PAHs in sediments in different locations around the world	112
4.11	Average concentrations of LABs in the sediments of Selangor River in the four study periods ($n = 52$)	115
4.12	Correlations between n-C10, n-C11, n-C12, n-C13, and n-C14 LABs, TLABs, and TOC	119
4.13	Average concentrations of LABs (in ng g^{-1} dw) and selected ratios in the surface sediments of Selangor River during the four study periods	120
4.14	Concentrations of alkanes ($\mu\text{g g}^{-1}$ dw) in the surface sediments of Selangor River	127
4.15	Correlation between the total concentration of alkanes and TOC	137
4.16	Hopane concentrations (ng g^{-1}) and relative ratios in the sediments of Selangor River	140



LIST OF FIGURES

Figure	Page
2.1 Model of pollutant movement and partitioning in the marine environment	7
2.2 Molecular structures of some alkanes	13
2.3 The basic hopane molecular structure	19
2.4 The molecular structures of some hopanes	20
2.5 The Friedel Craft Alkylation Process: An illustration of the release of LABs into the aquatic environment	25
2.6 Logarithmic values of the K_{ow} for the LABs	25
2.7 Internal Isomers	26
2.8 External Isomers	26
2.9 The I/E ratio in a sewage water treatment plant	27
2.10 The eight most regulated carcinogenic PAHs as listed by the US EPA and by the European community	29
2.11 The eight most regulated non-carcinogenic PAHs as listed by the US EPA and by the European community	29
2.12 Molecular structures of the 16 priority pollutant PAHs	33
2.13 Pathways of PAHs in the environment	36
2.14 The LMW/HMW ratio applied in the current study	41
2.15 The MP/P ratio	43
2.16 Main possible pathways of PAHs in the soil-plant system	48
2.17 The transport pathway of PAHs in mangrove ecosystems	50
2.18 The sediments quality guidelines applied in this study	53
3.1 Locations of the 13 stations for surface sediment sampling in Selangor River, Malaysia	60
3.2 Sediment sampling using Ekman grabber (A) and sediment (B)	63
3.3 Analytical procedure for analysis of the hydrocarbons in the sediment samples	67
3.4 Chromatogram showing concentrations of various PAHs in a sample	73
3.5 The calibration curve for naphthalene	75
3.6 The calibration curve for 5C10-LAB	76
3.7 Chromatogram for the 3 mg L ⁻¹ reference standard	76
4.1 The concentration of PAHs in the sediments of Selangor River in the four study periods	80
4.2 Percentages of the 2-ring PAHs to the 6-ring PAHs in the sediments: (a) according to sampling station; and (b) according to sampling period ($n = 52$)	83
4.3 Boxplot showing concentration of individual PAHs in sediment	85
4.4 Comparison between the concentrations of the LMW and the HMW PAHs in the sediments of Selangor River in the four study periods	89
4.5 The Phe/Ant vs. Fluo/Pyr ratios in the surface sediments of Selangor River and their associations with PAH sources	91
4.6 Relations between PAH ratios and their associations with PAH sources: (a) BaA/(BaA+Chr) vs. Fluo/(Fluo+Pyr), and (b) InP/(InP+BgP) vs. Fluo/(Fluo+Pyr), and (c) InP/(InP+BgP) vs. MP/P.	92
4.7 Hierarchical dendrogram for PAHs in Selangor River sediments	97
4.8 Contributions of three PAH sources to the PAH inputs into the	100

	sediments of Selangor River	
4.9	The TEQs of the PAHs at the 13 sampling sites	105
4.10	Relationships between TOC and the total concentration of the PAHs (Σ [PAHs]) in the sediments of Selangor River	109
4.11	Concentrations of LABs in the sediment of Selangor River in the four periods of the study	116
4.12	Relative abundance of LAB homologues in the sediments of Selangor River	117
4.13	Percent composition of the LABs in the sediments of Selangor River during the four study periods	118
4.14	Box plots of I/E, L/S, and C_{13}/C_{12} for the sediments of Selangor River in (a) the rainy inter-monsoon period; (b) the dry inter-monsoon period; (c) the SWM period; and (d) the NEM period	121
4.15	Relationships between the total LAB concentration and (a) the total PAH concentration, and (b) the TOC in the sediments of Selangor River	125
4.16	Total concentrations of the alkanes in the sediment samples	126
4.17	Concentraions of alkanes in the sediments of Selangor River during the four studied periods	129
4.18	Values of the carbon preference index (CPI) in the surface sediments of Selangor River	133
4.19	Spatial variations of the n-C17/pristane and n-C18/phytane ratios in Selangor River	134
4.20	Spatial patterns in the values of the pristane/phytane (Pri/Phy) ratio in the surface sediments of Selangor River	135
4.21	Spatial pattern of the terrigenous aquatic ratio (TAR) in the surface sediments of Selangor River	136
4.22	Plot of C_{29}/C_{30} vs. $\Sigma C_{31}-C_{35}/C_{30}$ for the sediments of Selangor River	142

LIST OF ABBREVIATIONS

ACL	Average chain length
Ant/(Ant+Phe)	Ratio of anthracene to phenanthrene plus anthracene
BaA/(BaA+Chr)	Ratio of Benzo(a)anthracene to benzo(a)anthracene plus chrysene
C ₁₃ /C ₁₂	The ratio of Σ C ₁₃ - LAB/ Σ C ₁₂ - LAB
C ₁₇ /pristane	Ratio of C ₁₇ n-alkane to pristane
C ₁₈ /phytane	Ratio of C ₁₈ n-alkane to pristane
C ₂₉ /C ₃₀	Ratio of 17 α , 21 β (H)-30-norhopane to 17 α , 21 β (H)-hopane
Σ C ₃₁ -C ₃₅ /C ₃₀	Ratio of sum of 17 α , 21 β (H)-C ₃₁ homohopane to 17 α , 21 β (H)-C ₃₅ homohopane relative to 17 α , 21 β (H)-hopane
Σ cPAHs	Sum of carcinogenic PAHs
CPI _s	Carbon preference indices
DCM	Dichloromethane
D(%)	The average degradation
dw	Dry weight
ERL	Effects range low
ERM	Effect range median
eV	Electron Volt
Fluo/(Fluo +Pyr)	Ratio of fluoranthene to fluoranthene plus pyrene
Fluo/Pyr	Ratio of fluoranthene to pyrene
GC-MS	Gas Chromatography-Mass Spectrometry
HMW	High Molecular Weight
I/E	The ratio of Internal to External isomers
IIS	Internal Injection Standard
K _{ow}	Octanol-water partitioning coefficient
LABs	Linear alkylbenzenes
LMW	Low Molecular Weight
LMW/HMW	Ratio of Low Molecular Weight to High Molecular Weight isomers
InP/(InP+BgP)	Ratio of indeno(1,2,3-cd)pyrene to indeno(1,2,3-cd)pyrene plus benzo(g,h,i)perylene
L/S	The ratio of long-chain to short-chain LAB
MECO	Middle East Crude Oil
MH	Major hydrocarbon
MP/P	Ratio of methylphenanthrenes to phenanthrene
m/z	Mass to charge ratio
alkanes	Normal alkanes
Na ₂ SO ₄	Sodium sulfate
nd	Not detected
PAHs	Polycyclic aromatic hydrocarbons

PC	Principal component
PCA	Principal Component Analysis
PEL	Probable effects level
PHC	Petroleum hydrocarbon
Phe/Ant	Ratio of phenanthrene to anthracene
Phy	Phytane
Pri	Pristane
Pr/Ph	Ratio of pristane to phytane
S	Station
SEACO	South East Asia Crude Oil
SIM	Selected Ion Monitoring
SIS	Surrogate Internal Standard
SLC	Screening level contamination
SQGs	Sediment quality guidelines
TARs	Terrigenous/aquatic ratios
TEF	Toxicity equivalency factor
TEL	Threshold effect level
TEQ	Toxicity equivalency quotient
TOC	Total organic carbon
T _s	18 α (H)-22,29,30-trisnorneohopane
T _m	17 α (H)-22,29,30-trisnorhopane
T _m /T _s	Ratio of 17 α -22,29,30-trisnorhopane to 18 α -22,29,30-trisnorhopane
UCM	Unresolved complex mixture
USEPA	United States Environmental Protection Agency

CHAPTER 1

INTRODUCTION

1.1 Background of the study

The eclectic integration of the economic, social, and cultural aspects has led to a rapid physical growth in the urban areas of Malaysia. Due to these rapid development activities, several environmental problems, including pollution, have been created. As a matter of fact, human activities such as lodging and waste generation and improper discharge can harmfully contribute to the environmental degradation problems. The various human activities have serious environmental complications and may overwhelm the surroundings if there are no instantaneous precautionary steps taken by all stakeholders.

Selangor is one of the 13 states of Malaysia and one of the states lying on the west coast of Peninsular Malaysia. Its geographical position, in the core of Peninsular Malaysia, has contributed to rapid development of the state. Hence, the State of Selangor is considered as the transportation and industrial hub of Malaysia which, in turn, attracts both interior migrants such as those coming from the other Malaysian States and exterior tourists coming from different countries (Alsalahi et al., 2014). The basin of Selangor River faces serious environmental degradation from urbanization, industrialization, and population growth. Currently, the major land-based pollution sources in the basin are (Ong, 2005), poorly-treated or untreated domestic sewage, siltation from land clearing, construction and sand mining activities, run-off from agricultural land, and untreated effluent from industry. In the past few decades, the anthropogenic activities associated with rapid industrialization and urbanization growth like fossil fuel combustion, discharge of domestic sewage, and incineration of industrial waste contributed greatly to pollution, especially in the marine and coastal environments in Malaysia. Among the pollutants, hydrocarbons are among the most threatening pollutants which have been widely studied. The Strait of Malacca is one of the world's busiest supertanker routes (Zakaria et al., 2001). It is surrounded by an area which is rich with natural resources that include oil and natural gas. Therefore, there is a variety of potential sources of petroleum in these coastal areas. Moreover, the coasts of Malaysia are subjected to various potential sources of petroleum pollution, including routine and accidental oil spills from tankers, spillage of crude oils from inland and offshore oil fields, and run-off from urban lands.

Because of their seriousness and being among the most widespread organic pollutants, the polycyclic aromatic hydrocarbons (PAHs) are given more attention than other pollutants. Pollution with PAHs is one of the primary environmental problems facing humanity at present, mainly owing to the fact that PAHs are toxic, mutagenic, and carcinogenic to humans and other organisms. Additionally, they are subjected to bioaccumulation and concentration in the aquatic food web. Furthermore, they are

relatively persistent in the environment. Thus, the focus of the current study is investigation of the concentrations and the origins of four groups of hydrocarbons, namely, alkanes, hopanes, linear alkylbenzenes (LABs), and PAHs in the sediments of Selangor River, which faces the Strait of Malacca. The results of this study give an indication of the magnitude of pollution of Selangor River with these pollutants and the distribution of these pollutants along the river. Studying the levels and distribution of the PAHs and LABs in the sediments is highly important as both groups of compounds are highly hydrophobic and have large values of the octanol-water partition co-efficient (K_{ow}). As such, they are expected to partition and accumulate into the sediments and the lipid parts of biota. Besides, some PAHs are carcinogenic and mutagenic in nature. Accordingly, the identification and quantification of PAHs in sediments are significant in order to mark clean-up areas.

1.2 Problem statement

The aquatic and terrestrial ecosystems are exposed to the threat of hydrocarbon accumulation as a result of the increase in human activities along the coastline facing the Strait of Malacca. This has led to pollution of the marine and coastal environments, including the mangrove forests, rivers such as Selangor River. More than 80% of the hydrocarbons (HCs) found in the environment are produced by anthropogenic activities while the other 20% are released naturally through oil seeps and biogenic sources. The major HCs of environmental prevalence and, hence, concern are the straight chain alkanes, cycloalkanes, and aromatics. Relatively, the HCs are of the greatest concern, which is an issue that is related to the ubiquity of the PAHs in the environment. The PAHs are considered as one of the most important classes of micro-organic pollutants, produced mostly via incomplete combustion of organic materials containing carbon, hydrogen, and other minor elements (Bakhtiari et al., 2010). In this study, the PAHs were selected because they are persistent, toxic, mutagenic, and carcinogenic and because they have teratogenic and hepatotoxic effects (Cao et al., 2009; Karami et al., 2012a; Masiol et al., 2012). These pollutants have worrying human health implications and are also of concern for potential adverse effects on a range of organisms. They have not only affected biological changes in, and sometimes death of, aquatic and terrestrial ecosystems, but also the tourism industry, which is one of the main economic engines of Malaysia that has experienced damages and loss. Therefore, a study on hydrocarbons contamination in sediments is required in order to determine the current level of hydrocarbons in Peninsular Malaysia especially the sediment regions located along the Straits of Malacca such as the chosen site, the Selangor River.

1.3 Significance of the study

Study on hydrocarbons in sediments will significantly provide the scientific community and local environmental managers with useful data on the various characteristics of hydrocarbons in the aquatic environment of Selangor River that can be utilized by decision makers in the future planning for the purpose of achievement of sustainable development. The current study is important for scientists, environmentalists, educators, and the tourism managers since it will contribute to further understanding of the effects of the human activities and the human by products on the ecosystems of

Selangor River. It is expected that the study results can help understand the current condition of the hydrocarbons pollution, project the future outcomes of its anticipated effects, and change certain human activities or impose certain regulations to save the aquatic and terrestrial environment. Furthermore, none of the previous studies employed source apportionment techniques for the identification and quantitative evaluation of hydrocarbons sources in Selangor River. Therefore, in view of the pollution condition and the importance of the investigated river as source of food and drinking water, the characterization of hydrocarbons and their sources by apportionment techniques is considered as a novel approach that can contribute substantially to the understanding of the primary source(s) of hydrocarbons in sediments. The results of this study will be compared with the proposed PAH levels of sediment quality guidelines in order to evaluate the biological effects of sedimentary PAHs to the aquatic organisms, especially the ones that are closely associated with sediment. Finally, a number of new findings on PAHs fate in Malaysian sediment will be scientifically shown through this study.

1.4 Research questions

Whatever the origin of contamination, benthic habitats are preferential environmental sink for many contaminants, since many pollutants originally introduced into the water column have affinities for sediment particles. Because sediments are also an important biological habitat, incorporation of toxicants into the food web is influenced by toxicant concentrations in sediments, which represent the main sink for PAHs and constitute PAH reservoirs. Therefore, this study was initiated to address the following four research questions:

1. What the compositional patterns and sources of alkanes, hopanes, LABs and PAHs in the sediments of Selangor River in the four studied periods?
2. Are there significant differences in hydrocarbon concentrations among the four periods under study?
3. Are there significant relationships between total organic carbon (TOC) and the concentrations of alkanes, hopanes, LABs and PAHs in the surface sediments in the study area?
4. Can LABs serve as molecular markers for identification of the sources of PAHs in sediments? and is there a toxic effect of PAHs on human health and ecological environment.

1.5 Scope of the study

This research has been conducted on the sediments of Selangor River to investigate the level of hydrocarbon pollution in these sediments. Besides, the study aimed at differentiating hydrocarbon contamination, linking the molecular markers to their sources. In other words, the purpose of this study was to apply environmental fingerprinting techniques in order to determine the levels, origins, and distributions of alkanes, hopanes, LABs and PAHs in the sediments of Selangor River in four periods. Therefore, this study adopts the principles of environmental forensics, including source

apportionment of compounds in environmental samples to Anthropogenic Waste Indicators (AWI).

The AWIs are indicators of man-made wastes that consist of chemical compounds originating from pharmaceuticals, detergent disinfectants, fragrances, antimicrobials, PHC, pesticides, and other waste-borne materials present in wastewater and agricultural and urban run-off, which are commonly referred to as AWI compounds. The AWIs include endocrine-disrupting compounds (EDCs) that have detrimental reproductive effects on fish (Brian et al., 2007) and humans (Guillette, 1995). The AWI can also be utilized to indicate other anthropogenic pollutants like PAHs and LABs. Selangor River was chosen because the river flows through the commercial, industrial, and residential areas facing the Strait of Malacca. Furthermore, urban and other development areas surround the river.

1.6 Objectives of the study

The main and the leading objective of this study was to analyze the degree of anthropogenic pollution and used of hydrocarbon markers to investigate the distribution of anthropogenic inputs to the sediments of Selangor River, Malaysia. However, under the main objective of the study, the following four sub-objectives are listed:

1. To assess the compositional patterns and sources of alkanes, hopanes, LABs and PAHs in the sediments of Selangor River in four periods: the dry inter-monsoonal period, the wet inter-monsoonal period, the south-west (SW) monsoon, and the north-east (NE) monsoon periods.
2. To compare the concentrations of the alkanes, hopanes, LABs and PAHs in different sampling stations during the four study periods through the use of multivariate data analysis.
3. To determine the relationships between total organic carbon (TOC) and the concentrations of alkanes, hopanes, LABs and PAHs in the surface sediments in the study area.
4. To investigate the utility of LABs as molecular marker for identification of the sources of PAHs in sediments and to evaluate the PAHs potent toxicity on human health and ecological environment.

1.7 Organization of the thesis

The outline presented next describes the different chapters of this thesis and assists the reader in understanding the flow of its chapters and main sections.

The thesis consists of five chapters. Chapter 1 defines the background and specifies the purpose of this investigation.

Next, Chapter 2 reviews and discusses the literature relevant to the study. It highlights the characteristics of alkanes, hopanes, LABs and PAHs in the light of their sources

and uses. The chapter also spotlights the relevant chemical and environmental properties of alkanes, hopanes, LABs and PAHs, as well as the main risks of human exposure to PAHs. It also provides international and national standards with regard to PAHs. Finally, this Chapter discusses trends in alkanes, hopanes, LABs and PAHs, and sheds some light on the shortcomings of the related research.

Then, Chapter 3 describes the methodology and experimental procedures followed in the study. It describes the sampling and the data collection procedures.

Afterwards, Chapter 4 presents a detailed analysis of the data collected and provides illustrations along with the discussion of the results. It also compares the results of this study with those of previous studies.

Finally, Chapter 5 concludes the thesis. It presents a summary of the main research findings, the major conclusions, and some recommendations for future research.

REFERENCES

- Abas, M.R.b., Simoneit, B.R.T., Elias, V., Cabral, J.A. and Cardoso, J.N., 1995. Composition of higher molecular weight organic matter in smoke aerosol from biomass combustion in Amazonia. *Chemosphere*, 30, 995–1015.
- Abas, M., Rahman, N.A., Omar, N.Y.M., Maah, M.J., Abu Samah, A., Oros, D.R., Otto, A. and Simoneit, B.R., 2004. Organic composition of aerosol particulate matter during a haze episode in Kuala Lumpur, Malaysia. *Atmospheric Environment*, 38, 4223-4241.
- Abdullah P. and Nainggolan, H., 1991. Phenolic water pollutants in a Malaysian River basin. *Environmental Monitoring and Assessment*, 19 (1), 423-431.
- Abdullah, A.R., Tahir, N.M., Loong, T.S., Hoque, T.M. and Sulaiman, A.H., 1999. The GEF/UNDP/IMO Malacca Straits Demonstration Project: Sources of Pollution. *Marine Pollution Bulletin*, 39 (1-12), 229-233.
- Abha, S. and Singh, C.S., 2012. Hydrocarbon Pollution: Effects on Living Organisms, Remediation of Contaminated Environments and Effects of Heavy Metals Co-contamination on Bioremediation. In: Introduction to Enhanced Oil on Recovery (EOR) Processes and Bioremediation of Oil Contaminated Sites, Romero-Zeron, L. (Ed.). *InTech Publisher, China, ISBN: 978-953-51-0629-6*, pp, 186-206.
- Abu-Allaban, M., Gillies, J.A., Gertler, A.W., Clayton, R. and Proffitt, D., 2003. Tailpipe, resuspended road dust and brake-wear emission factors from on-road vehicles. *Atmospheric Environment*, 37, 5283-5293.
- Ahad, J.M.E., Ganeshram, R.S., Bryant, C.L., Cisneros-Dozal, L.M., Ascough, P.L., Fallick, A.E. and Slater, G.F., 2011. Sources of n-alkanes in an urbanized estuary: insights from molecular distributions and compound-specific stable and radiocarbon isotopes. *Marine Chemistry*, 126, 239-249.
- Ahrens, M.J. and Depree, C.V., 2004. Inhomogenous distribution of polycyclic aromatic hydrocarbons in different size and density fractions of contaminated sediment from Auckland Harbour, New Zealand: an opportunity for mitigation. *Marine Pollution Bulletin*, 48, 341-350.
- Akpor, O.B., Okolomike, U.F. Olaolu, T.D. and Aderiye, B.I., 2014. Remediation of Polluted Wastewater Effluents: Hydrocarbon Removal. *Trends in Applied Sciences Research*, 9, 160-173. doi: 10.3923/tasr.2014.160.173.
- Alcock, R., Bashkin, V., Bisson, M., Brecher, R., van Bree, L., Chrast, R., Colosio, C., Desqueyroux, H., Evstafjeva, E. and Feeley, M., 2003. Health Risks of Persistent Organic Pollutants from Long-Range Transboundary Air Pollution, Masaryk University, Czech Republic. *World Health Organization, Europe*.

- Ali, M.M., Humrawali, N. and Latif, M.T., 2009. Phytosterols composition in surface sediment of Kuala Selangor, Selangor, Malaysia. *European Journal of Scientific Research*, 33, 187-194.
- Alkhadher, S.A.A., Zakaria, M.P., Yusoff, F.M., Kannan, N., Suratman, S., Keshavarzifard, M., Magam, S.M., Masood, N. and Vaezzadeh, V., 2016. Distribution and sources of linear alkylbenzenes (LABs) in surface sediments from Johor Bahru Coast and the Kim Kim River, Malaysia. *Environmental Forensics*, 17(1), 36-47.
- Al-Saad, H.T. Aziz, S., Majeed, A., Khazim, H., Abd, I.M. and Aleal, A.H., 2008. Oil Spill in Khor Al-Zubair: A case study. *Marina Mesopotamica*, 23(2), 321-332.
- Alsalahi, M.A., Latif, M.T., Ali, M.M., Magam, S.M., Wahid, N.B.A., Khan, M.F. and Suratman, S., 2014. Distribution of surfactants along the estuarine area of Selangor River, Malaysia. *Marine Pollution Bulletin*, 80, 344-350.
- Aronstein, B.N. and Alexander, M., 1992. Surfactants at low concentrations stimulate biodegradation of sorbed hydrocarbons in samples of aquifer sands and slurries. *Environmental Toxicology and Chemistry*, 11, 1227-1233.
- Arora, A., 2006. Hydrocarbons (Alkanes, Alkenes And Alkynes). *Discovery Publishing House*. ISBN: 818356142X, 9788183561426, 1-252.
- Ashley, J.T. and Baker, J.E., 1999. Hydrophobic organic contaminants in surficial sediments of Baltimore Harbor: Inventories and sources. *Environmental Toxicology and Chemistry*, 18, 838-849.
- ATSDR. 1995. Agency for Toxic Substances and Disease Registry, Toxicology profile for polycyclic aromatic hydrocarbons. U.S. *Department of Health and Human Services*.
- ATSDR. 2005. Agency for Toxic Substances and Disease Registry, Toxicology Profile for Polyaromatic Hydrocarbons. ATSDR's Toxicological Profiles on CD-ROM, CRC Press, *Boca Raton, FL*.
- Baek, S.O., Goldstone, M.E., Kirk, P.W.W., Lester, J.N. and Pery, R., 1991. Concentrations of particulate and gaseous polycyclic aromatic hydrocarbons in London air following a reduction in the Lead content of petrol in the United Kingdom. *Science of the Total Environment*, 11, 169-199.
- Bahry, P.S., Zakaria, M.P., Bin Abdullah, A.M., Abdullah, D.K., Sakari, M., Chandru, K. and Shahbazi, A., 2009. Forensic Characterization of Polycyclic Aromatic Hydrocarbons and Hopanes in Aerosols from Peninsular Malaysia. *Environmental Forensics*, 10(3), 240-252.
- Bai, Y.J., Li, X.Q., Liu, W.X., Tao, S., Wang, L.G. and Wang, J.F., 2008. Polycyclic aromatic hydrocarbon (PAH) concentrations in the dissolved, particulate, and sediments phases in the Luan River watershed, China. *Journal of*

- Baker, J.E. and Eisenreich, S.J., 1990. Concentrations and fluxes of polycyclic aromatic hydrocarbons and polychlorinated biphenyls across the air-water interface of Lake Superior. *Environmental Science and Technology*, 24, 342-352.
- Baker, J.E., Eisenreich, S.J. and Eadie, B.J., 1991. Sediment trap fluxes and benthic recycling of organic carbon, polycyclic aromatic hydrocarbons, and polychlorobiphenyl congeners in Lake Superior. *Environmental Science and Technology*, 25, 500-509.
- Bakhtiari, A.R., Zakaria, M.P., Yaziz, M.I., Lajis, M.N.H. and Bi, X., 2009. Polycyclic aromatic hydrocarbons and n-alkanes in suspended particulate matter and sediments from the langat river, peninsular malaysia. *Environment Asia*, 2(2), 1-10.
- Bakhtiari, A.R., Zakaria, M.P., Yaziz, M.I., Lajis, H., Nordin, M., Bi, X., Shafiee, M., Reza, M. and Sakari, M., 2010. Distribution of PAHs and n-alkanes in Klang River surface Sediments, Malaysia. *Pertanika Journal of Science & Technology*, 18.
- Baumard, P., Budzinski, H. and Garrigues, P., 1998. Polycyclic aromatic hydrocarbons in sediments and mussels of the western Mediterranean sea. *Environmental Toxicology and Chemistry*, 17, 765-776.
- Baumard, P., Budzinski, H., Garrigues, P., Dizer, H. and Hansen, P.D., 1999. Polycyclic aromatic hydrocarbons in recent sediments and mussels (*Mytilus edulis*) from the Western Baltic Sea: occurrence, bioavailability and seasonal variations. *Marine Environmental Research*, 47, 17-47.
- Belih, L.I., Malih, U.M., Penzina, A.A. and Smagunova, A.N., 2003. Benzo[a]pyrene in atmospheric exhausts of industrial Pribaikalye. Luminescence and laser physics. Irkutsk: *Publishing house of Irkutsk State University*, 45-50 (in russian).
- Bell, R., Green, M., Hume, T. and Gorman, R., 2000. What regulates sediments in estuaries? *Water and Atmosphere*, 8 (4), 13-16.
- Bence, A.E., Kvenvolden, K.A. and Kennicutt, M.C., 1996. Organic geochemistry, applied, to environmental assessments of Prince William Sound, Alaska, after the Exxon Valdez oil spill--a review. *Organic Geochemistry*, 24 (1), 7-42.
- Blumer, M., Mullin, M.M. and Thomas, D.W., 1963. Pristane in zooplankton. *Science*, 140, 974.
- Bernard, D., Pascaline, H. and Jeremie, J.J., 1996. Distribution and origin of hydrocarbons in sediments from lagoons with fringing mangrove communities. *Marine Pollution Bulletin*, 32 (10), 734-739.

- Bin Abas, M.R., Rahman, N.A., Omar, N.Y.M.J., Maah, M.J., Samah, A.A., Oros, D.R., Otto, A., and Simoneit, B.R.T., 2004. Organic composition of aerosol particulate matter during a haze episode in Kuala Lumpur, Malaysia. *Atmospheric Environment*, 38, 4223–4241.
- Bixian, M., Jiamo, F., Gan, Z., Zheng, L. and Yushun, M. et al., 2001. Polycyclic aromatic hydrocarbons in sediments from the pearl river and estuary, China: Spatial and temporal distribution and sources. *Applied Geochemistry*, 16, 1429-1445. DOI: 10.1016/S0883-2927(01)00050-6.
- Boehm, P.D., Page, D.S., Gilfillan, E.S., Bence, A.E., Burns, W.A. and Mankiewicz, P.J., 1998. Study of the fates and effects of the Exxon Valdez oil spill on benthic sediments in two bays in Prince William Sound, Alaska. 1. Study design, chemistry, and source fingerprinting. *Environmental Science and Technology*, 32, 567-576.
- Bojes, H.K. and Pope, P.G., 2009. Characterization of EPA's 16 priority pollutant polycyclic aromatic hydrocarbons (PAHs) in tank bottom solids and associated contaminated soils at oil exploration and production sites in Texas. *Regul. Toxicology and Pharmacology*, 47, 288-295.
- Boonchan, S., Britz, M.L. and Stanley, G.A., 2000. Surfactant-enhanced biodegradation of high molecular weight polycyclic aromatic hydrocarbons by *Stenotrophomonas maltophilia*. *Journal of Biotechnology Bioengineering*, 59, 482-494.
- Boonyatumanond, R., Wattayakorn, G., Togo, A. and Takada, H., 2006. Distribution and origins of polycyclic aromatic hydrocarbons (PAHs) in riverine, estuarine, and marine sediments in Thailand. *Marine Pollution Bulletin*, 52(8), 942-956.
- Boonyatumanond, R., Murakami, M., Wattayakorn, G., Togo, A. and Takada, H., 2007. Sources of polycyclic aromatic hydrocarbons (PAHs) in street dust in a tropical Asian mega-city, Bangkok, Thailand. *Science of the Total Environment*, 384, 420-432.
- Boot, C.S., Ettwein, V.J., Maslin, M.A., Weyhenmeyer, C.E. and Pancost, R.D., 2006. A 35,000 year record of terrigenous and marine lipids in Amazon Fan sediments. *Organic Geochemistry*, 37, 208-219.
- Bouloubassi, I. and Saliot, A., 1993. Dissolved, Particulate and Sedimentary Naturally Derived Polycyclic Aromatic-hydrocarbons in a Coastal Environment - Geochemical Significance. *Marine Chemistry*, 42, 127-143.
- Bouloubassi, I., Roussiez, V., Azzoug, M. and Lorre, A., 2012. Sources, dispersal pathways and mass budget of sedimentary polycyclic aromatic hydrocarbons (PAH) in the NW Mediterranean margin, Gulf of Lions. *Marine Chemistry*, 142, 18-28.
- Boström, C.E., Gerde, P., Hanberg, A., Jernström, B., Johansson, C., Kyrklund, T., Rannug, A., Törnqvist, M., Victorin, K. and Westerholm, R., 2002. Cancer

risk assessment, indicators, and guidelines for polycyclic aromatic hydrocarbons in the ambient air. *Environmental health perspectives*, 110(Suppl 3), 451.

Bourbonniere, R.A. and Meyers, P.A., 1996. Sedimentary geolipid records of historical changes in the watersheds and productivities of Lakes Ontario and Erie. *Limnology and oceanography*, 41, 352-359.

Brändli, R.C., Bucheli, T.D., Ammann, S., Desaulles, A., Keller, A., Blum, F. and A. Stahel, W., 2008. Critical evaluation of PAH source apportionment tools using data from the Swiss soil monitoring network. *Journal of Environmental Monitoring*, 10(11), 1278-1286.

Brian, J.V., Harris, C.A., Scholze, M., Kortenkamp, A., Booy, P., Lamoree, M., Pojana, G., Jonkers, N., Marcomini, A. and Sumpter, J.P., 2007. Evidence of estrogenic mixture effects on the reproductive performance of fish. *Environmental Science & Technology*, 41(1), 337-344.

Budzinski, H., Jones, I., Bellocq, J., Pierard, C. and Garriques, P., 1997. Evaluation of sediment contaminant by polycyclic aromatic hydrocarbons in the Gironde estuary. *Marine Chemistry*, 58, 85-97.

Burns, K.A. and Yelle-Simmons, L., 1994. The Galeta Oil Spill. IV. Relationship Between Sediments and Organism Hydrocarbon Loads. *Estuarine, Coastal and Shelf Science*, 38 (4), 397-412.

Bush, R.T. and McInerney, F.A., 2015. Influence of temperature and C4 abundance on n-alkane chain length distributions across the central USA. *Organic Geochemistry*, 79, 65-73.

Burton Jr, G.A., 2002. Sediment quality criteria in use around the world. *Limnology*, 3, 65-76.

Cadle, S.H. and Mulawa, P.A., 1999. Composition of lightduty motor vehicle exhaust particulate matter in the Denver, Colorado area. *Environmental Science and Technology*, 33 (14), 2328-2339.

Cailleaud, K., Forget-Leray, J., Peluhet, L., LeMenach, K., Souissi, S. and Budzinski, H., 2009. Tidal influence on the distribution of hydrophobic organic contaminants in the Seine Estuary and biomarker responses on the copepod *Eurytemora affinis*. *Environmental Pollution*, 157(1), 64-71.

Cao, B., Nagarajan, K. and Loh, K.C., 2009. Biodegradation of aromatic compounds: current status and opportunities for biomolecular approaches. *Applied Microbiology and Biotechnology*, 85, 207-228.

Caravaggio, G.A., Charland, J.P., MacDonald, P. and Graham, L., 2007. n-Alkane profiles of engine lubricating oil and particulate matter by molecular sieve extraction. *Environmental Science Technology*, 41, 3697-3701.

- Cardellicchio, N., Buccolieri, A., Giandomenico, S., Lopez, L., Pizzulli, F. and Spada, L., 2007. Organic pollutants (PAHs, PCBs) in sediments from the Mar Piccolo in Taranto (Ionian Sea, Southern Italy). *Marine Pollution Bulletin*, 55, 451-458.
- Castellano, A.V., Cancio, J.L., Aleman, P.S. and Rodriguez, J.S., 2003. Polycyclic aromatic hydrocarbons in ambient air particles in the city of Las Palmas de Gran Canaria. *Atmospheric Environment*, 29, 475-480.
- Cavalcante, R.M. and Sousa, F.W., Nascimento, R.F., Silveira, E.R., Freire, G.S.S., 2009. The impact of urbanization on tropical mangroves (Fortaleza, Brazil): Evidence from PAH distribution in sediments. *Journal of Environmental Management*, 91 (2), 328-335.
- Cerniglia, C.E., 1992. Biodegradation of polycyclic aromatic hydrocarbons. *Biodegradation*, 3 (2), 351-368.
- Chandru, K., Zakaria, M.P., Anita, S., Shahbazi, A., Sakari, M., Bahry, P.S. and Mohamed, C.A.R., 2008. Characterization of alkanes, hopanes, and polycyclic aromatic hydrocarbons (PAHs) in tar-balls collected from the East Coast of Peninsular Malaysia. *Marine Pollution Bulletin*, 56(5), 950-962.
- Chen, B., Xuan, X., Zhu, L., Wang, J., Gao, Y., Yang, K., Shen, X. and Lou, B. 2004. Distributions of polycyclic aromatic hydrocarbons in surface waters, sediments and soils of Hangzhou City, China. *Water Research*, 38, 3558-3568.
- Chen, S.C., Liao and C.M., 2006. Health risk assessment on human exposed to environmental polycyclic aromatic hydrocarbons pollution sources. *Science of The Total Environment*, 366(1), 112-123.
- Chen, J.L., Wong, Y.S. and Tam, N.F.Y., 2009. Static and dynamic sorption of phenanthrene in mangrove sediment slurry. *Journal of Hazardous Materials*, 168(23), 1422-1429. doi:10.1016/j.jhazmat.2009.03.043.
- Chen, C.W. and Chen, C.F., 2011. Distribution, origin, and potential toxicological significance of polycyclic aromatic hydrocarbons (PAHs) in sediments of Kaohsiung Harbor, Taiwan. *Marine Pollution Bulletin*, 63 (5-12), 417-423.
- Chen, C.W., Chen, C.F., Dong, C.D., and Tu, Y.T., 2012. Composition and source apportionment of PAHs in sediments at river mouths and channel in Kaohsiung Harbor, Taiwan. *Journal of Environmental Monitoring*, 14(1), 105-115.
- Cherng, S.H., Lin, S.T. and Lee, H., 1996. Modulatory effects of polycyclic aromatic hydrocarbons on the mutagenicity of I-nitropyrene: A structure-activity-relationship study. *Mutation Research*, 367 (4), 177-185.
- Chouksey, M.K., Kadam, A.N and Zingde, M.D., 2004. Petroleum hydrocarbon residues in the marine environment of Bassein-Mumbai. *Marine Pollution Bulletin*, 49, 637-647.

- Clendening, L.D., Jury, W.A. and Ernst, F.F., 1990. A field mass balance study of pesticide volatilization, leaching and persistence. In Long range transport of pesticides. Ed. D. A. Kurtz. Lewis Publishers Inc, Michigan, USA, 47-60.
- Collins, J.F., Brown, J.P., Alexeeff, G.V. and Salmon, A.G., 1998. Potency equivalency factors for some polycyclic aromatic hydrocarbons and polycyclic aromatic hydrocarbon derivatives. *Regulatory Toxicology and Pharmacology*, 28(1), 45-54.
- Colombo, J.C., Pelletier, E., Brochu, C., Khalil, M. and Catoggio, J.A., 1989. Determination of hydrocarbon sources using *n*-alkane and polyaromatic hydrocarbon distribution indexes. Case study: Rio de la Plata estuary, Argentina. *Environmental Science & Technology*, 23, 888-894.
- Commendatore, M.G., Esteves, J.L. and Colombo, J.C., 2000. Hydrocarbons in coastal sediments of Patagonia, Argentina: levels and probable sources. *Marine Pollution Bulletin*, 40(11), 989-998.
- Conde, F.J., Ayala, J.H., Afonso, A.M. and Ganzalez, V., 2005. Emissions of polycyclic aromatic hydrocarbons from combustion of agriculture and silvicultural debris. *Atmospheric Environment*, 39, 6654 – 6663.
- Connell, D.W., Wu, R.S.S., Richardson, B.J., Leung, K., Lam, P.S.K. and Connell, P.A., 1998. Occurrence of persistent organic contaminants and related substances in Hong Kong marine areas: An overview. *Marine Pollution Bulletin*, 36 (5), 376-384.
- CPCB. 2003. Central Pollution Control Board, Polycyclic Aromatic Hydrocarbons (PAHs) in air and their effects on human Health. *Ministry of Environment and Forests*, 42.
- Currie, T.J, Alexander, R. and Kagi, R.T., 1992. Coastal bitumens from Western Australia-long distance transport by ocean currents. *Organic Geochemistry*, 18, 595-601.
- Daka, E.R. and Ugbomeh, A.P., 2013. Polycyclic Aromatic Hydrocarbons in Sediment and Tissues of the Crab *Callinectes pallidus* from the Azuabie Creek of the Upper Bonny Estuary in the Niger Delta. *Research Journal of Applied Sciences, Engineering and Technology*, 6, 2594-2600.
- Dauner, A.L.I., Hernández, E.A., MacCormack, W.P. and Martins, C.C., 2015. Molecular characterisation of anthropogenic sources of sedimentary organic matter from Potter Cove, King George Island, Antarctica. *Science of the Total Environment*, 502, 408-416.
- Delgado, T., 2000. The occurrence and regulation of PAHs in groundwater and soil. *Smart News*, 2(5), 1-31.

- Delistraty, D., 1997. Toxic equivalency factor approach for risk assessment of polycyclic aromatic hydrocarbons. *Environmental Toxicology and Chemistry*, 16, 81-108.
- De Luca, G.D., Furesi, A., Micera, G., Panzanelli, A., Piu, P.C., Pilo, M.I., Spano, N. and Sanna, G., 2005. Nature, distribution and origin of polycyclic aromatic hydrocarbons (PAHs) in the sediments of Olbia harbor (Northern Sardinia, Italy). *Marine pollution bulletin*, 50, 1223-1232.
- Dickhut, R., Canuel, E., Gustafson, K., Liu, K., Arzayus, K., Walker, S., Edgecombe, G., Gaylor, M. and MacDonald, E., 2000. Automotive sources of carcinogenic polycyclic aromatic hydrocarbons associated with particulate matter in the Chesapeake Bay region. *Environmental science and technology*, 34 (21), 4635-4640.
- Djozan, D.J. and Assadi, Y., 1999. Monitoring of polycyclic aromatic hydrocarbons in water using headspace solid-phase microextraction and capillary gas chromatography. *Microchemical Journal*, 63, 276-284.
- Dobbins, R.A., Fletcher, R.A., Benner, B.A., Hoeft, S., 2006. Polycyclic aromatic hydrocarbons in flames, in diesel fuels, and in diesel emissions. *Combustion and flame*, 144(4), 773-781.
- DOE. 1994. Department of Environment: A study for the assessment of the sources of marine pollution. Ministry of Science, *Technology and the Environment*.
- Doong, R.A. and Lin, Y.T., 2004. Characterization and distribution of polycyclic aromatic hydrocarbon contaminations in surface sediment and water from Gao-ping River, Taiwan. *Water Research*, 38(7), 1733-1744. doi:10.1016/j.watres.2003.12.042.
- Dsikowitzky, L., Nordhaus, I., Jennerjahn, T.C., Khrycheva, P., Sivatharshan, Y., Yuwono, E. and Schwarzbauer, J., 2011. Anthropogenic organic contaminants in water, sediments and benthic organisms of the mangrove-fringed Segara Anakan Lagoon, Java, Indonesia. *Marine pollution bulletin*, 62 (4), 851-862.
- Dutta, T.K. and Harayama, S., 2000. Analysis of long-side-chain alkylaromatics in crude oil for evaluation of their fate in the environment. *Environmental Science and Technology*, 35, 102-107.
- Edlund, S., 2001. PAH as a POP. Possibilities, implications and appropriateness of regulating global emissions of Polycyclic Aromatic Hydrocarbons through the Stockholm Convention on Persistent Organic Pollutants. *IIIEE Reports* 2001, 9.
- Eganhouse, R.P. and Gossett, R.W., 1991. Historical deposition and biogeochemical fate of polycyclic aromatic hydrocarbons in sediments near a major submarine wastewater outfall in southern California., Baker, R. A. (Ed.) *Organic Substances in Sediments and Water*. Lewis Publishers: Chelsea. Michigan, 2, 191-220.

- Eganhouse, R.P. and Venkatesan, M.I., 1993. Chemical Oceanography and Geochemistry. In: M.D. Dailey, D.J. Reish, and J.W. Anderson (eds.). *Ecology of the Southern California Bight: A Synthesis and Interpretation*. University of California Press, Berkeley, CA., 71-189.
- Eglinton, G., and Hamilton, R. J., 1963. The distribution of alkanes. *Chemical Plant Taxonomy*, 187-217.
- El Nemr, A., El-Sadaawy, M.M., Khaled, A. and El-Sikaily, A., 2013. Distribution patterns and risks posed of polycyclic aromatic hydrocarbons contaminated in the surface sediment of the Red Sea coast (Egypt). *Desalination and Water Treatment*, 1-19.
- El Nemr, A., Moneer, A.A., Safaa Ragab, Ragab, S. and El-Sikaily, A., 2016. Distribution and sources of n-alkanes and polycyclic aromatic hydrocarbons in shellfish of the Egyptian Red Sea coast. *Egyptian Journal of Aquatic Research*, 42, 121-131.
- EQR. 2012. Environmental Quality Report, Department of Environment, *Ministry of Natural Resources and Environment*, Malaysia.
- EU. European Union. 2001. Ambient air pollution by Polycyclic Aromatic Hydrocarbons (PAH): Position Paper. ISBN 92- 894-2057-X. *Office for Official Publications of the European Communities*, Luxembourg.
- Eurachem, 1998. The fitness for purpose of analytical methods: A laboratory guide to method validation and related topics, *Laboratory of the Government Chemist*.
- Fang, M., Zheng, M., Wang, F., To, K.L., Jaafar, A.B. and Tong, S.L., 1999. The solventextractable organic compounds in the Indonesia biomass burning aerosols – characterization studies. *Atmospheric Environment*, 33, 783-795.
- Farias, C.O., Hamacher, C., Wagener, A.d.L. and Scofield, A.d.L., 2008. Origin and degradation of hydrocarbons in mangrove sediments (Rio de Janeiro, Brazil) contaminated by an oil spill. *Organic Geochemistry*, 39, 289-307.
- Feng, C., Xia, X., Shen, Z. and Zhou, Z., 2007. Distribution and sources of polycyclic aromatic hydrocarbons in Wuhan section of the Yangtze River, China. *Environmental Monitoring and Assessment*, 133, 447-58.
- Fernandez, P., Vilanova, R.M. and Grimalt, J.O., 1999. Sediment fluxes of polycyclic aromatic hydrocarbons in European high altitude mountain lakes, *Environmental Science and Technology*, 33, 3716-3722.
- Fowler, S.W., Readman, J.W., Oregioni, B., Villeneuve, J.P. and McKay, K., 1993. Petroleum hydrocarbons and trace metals in near shore Gulf sediment and biota before and after the 1991 war: an assessment of temporal and spatial trends. *Marine Pollution Bulletin*, 27, 171-182.

- Fraser, M.P., Cass, G.R. and Simoneit, B.R.T., 1999. Particulate organic compounds emitted from motor vehicle exhaust and in the urban atmosphere. *Atmospheric Environment*, 33, 2715-2724.
- Froehner, S., Machado, K.S., Falcão, F., Monnich, C. and Bessa, M., 2011. Inputs of domestic and industrial sewage in upper Iguassu, Brazil, identified by emerging compounds. *Water, Air, and Soil Pollution*, 215, 251-259.
- Gaga, E.O., 2004. Investigation of Polycyclic Aromatic Hydrocarbon (PAH) Deposition in Ankara, Ph.D Thesis pp. 233.
- Gallon, C., Tessier, A., Gobeil, C. and Beaudin, L., 2005. Sources and chronology of atmospheric lead deposition to a Canadian Shield lake: inferences from Pb isotopes and PAH profiles. *Geochimica Cosmochimica Acta*, 69 (13), 3199–3210. <http://dx.doi.org/10.1016/j.gca.2005.02.028>.
- Garrigues, P., Budzinski, H., Manitz, M. and Wise, S., 1995. Pyrolytic and petrogenic inputs in recent sediments: a definitive signature through phenanthrene and chrysene compound distribution. *Polycyclic Aromatic Compounds*, 7, 275-284.
- Georgiadis, P. and Kyrtopoulos, S.A., 1999. Molecular epidemiological approaches to the study of the genotoxic effects of urban air pollution. *Mutation Research*, 428 (1-2), 91-98.
- Gledhill, W., Saeger, V.W. and Trehry, M.L., 1991. An aquatic environmental safety assessment of linear alkylbenzene. *Environmental Toxicology and Chemistry*, 10, 169-78.
- Gomes, A.D.O. and Azevedo, D.D.A., 2003. Aliphatic and aromatic hydrocarbons in tropical recent sediments of Campos dos Goytacazes, RJ, Brazil. *Journal of the Brazilian Chemical Society*, 14, 358-368.
- González, J., Viñas, L., Franco, M., Fumega, J., Soriano, J., Grueiro, G. Muniategui, S., López-Mahía, P., Prada, D. and Bayona, J., 2006. Spatial and temporal distribution of dissolved/dispersed aromatic hydrocarbons in seawater in the area affected by the Prestige oil spill. *Marine pollution bulletin*, 53(5), 250-259.
- Gough, M. and Rowland, S., 1990. Characterization of unresolved complex mixtures of hydrocarbons in petroleum. *Nature*, 344, 648-650.
- Gough, M.A., Rhead, M.M. and Rowland, S.G., 1992. Biodegradation studies of unresolved complex mixtures of hydrocarbons: model UCM hydrocarbons and the aliphatic UCM. *Organic Geochemistry*, 18, 17-22.
- Grosenheider, K.E., Bloom, P.R. and Halbach, T.R., 2005. A Review of the Current Literature Regarding Polycyclic Aromatic Hydrocarbons in Asphalt Pavement. Mn/DOT Contract No. 81655. *University of Minnesota, Minneapolis, MN, USA*.

- Guillette, L.H., 1995. Endocrine-disrupting environmental contaminants and developmental abnormalities in embryos. *Human and Ecological Risk Assessment*, 1(2), 25-36.
- Guo, Z., Lin, T., Zhang, G., Yang, Z. and Fang, M., 2006. High-resolution depositional records of polycyclic aromatic hydrocarbons in the central continental shelf mud of the East China Sea. *Environmental Science & Technology*, 40, 5304-5311.
- Guo, W., He, M.C., Yang, Z.F., Lin, C.Y., Quan, X.C. and Wang, H.Z., 2007. Distribution of polycyclic aromatic hydrocarbons in the water suspended particulate matter and sediment from Daliao River watershed, China. *Chemosphere*, 68(1), 93-104. doi:10.1016/j.chemosphere.2006.12.072.
- Guo, W., Pei, Y., Yang, Z. and Chen, H., 2011a. Historical changes in polycyclic aromatic hydrocarbons (PAHs) input in Lake Baiyangdian related to regional socio-economic development. *Journal of Hazardous Materials*, 187, 441-449.
- Guo, J., Liang, Z., Liao, H., Tang, Z., Zhao, X., Wu, F., 2011b. Sedimentary record of polycyclic aromatic hydrocarbons in Lake Erhai, Southwest China. *Journal of Environmental Sciences*, 23 1308-1315.
- Gustafsson, Ö., Haghsseta, F., Chan, C., Macfarlane, J. and Gschwend, P.M. 1997. Quantification of the dilute sedimentary soot phase: Implications for PAH speciation and bioavailability. *Environmental Science & Technology*, 31, 203-209.
- Gustafsson, Ö., Long, C.M., MacFarlane, J. and Gschwend, P.M., 2001. Fate of linear alkylbenzenes released to the coastal environment near Boston Harbor. *Environmental Science & Technology*, 35, 2040-2048.
- Han, Y.M., Bandowe, B.A.M., Wei, C., Cao, J.J., Wilcke, W., Wang, G.H., Ni, H.Y., Jin, Z.D., An, Z.S. and Yan, B.Z., 2015. Stronger association of polycyclic aromatic hydrocarbons with soot than with char in soils and sediments. *Chemosphere*, 119, 1335-1345.
- Harji, R.R., Yvenat, A. and Bhosle, N.B., 2008. Sources of hydrocarbons in sediments of the Mandovi estuary and the Marmugoa harbour, west coast of India. *Environment International*, 34, 959-965.
- Haritash, a. K. and Kaushik, C.P., 2009. Biodegradation aspects of Polycyclic Aromatic Hydrocarbons (PAHs): A review. *Journal of Hazardous Materials*, 169 (1-3), 1-15. <https://doi.org/10.1016/j.jhazmat.2009.03.137>.
- Harrison, R.M., Smith, D. and Luhana, L., 1996. Source apportionment of atmospheric polycyclic aromatic hydrocarbons collected from an urban location in Birmingham, UK. *Environmental Science & Technology*, 30, 825-832.
- Harrison, E.Z., Oakes, S.R., Hysell, M. and Hay, A., 2006. Organic chemicals in sewage sludges. *Science of the Total Environment*, 367, 481-497.

- Hartmann, P.C., Quinn, J., King, J.W., Tsutsumi, S. and Takada, H., 2000. Intercalibration of LABs in marine sediment SRM1941a and their application as a molecular marker in Narragansett Bay sediments. *Environmental Science and Technology*, 34, 900-906.
- Hasanati, M., Savari, A., Nikpour, Y. and Ghanemi, K., 2011. Assessment of the Sources of Polycyclic Aromatic Hydrocarbons in Mousa Inlet by Molecular Ratios. *Journal of Environmental Studies*, 37, 1-6.
- Haynes, D., Brodie, J., Christie, C., Devlin, M., Michalek-Wagner, K., Morris, S., Ramsay, M., Storrie, J., Waterhouse, J. and Yorkston, H., 2001. Great Barrier Reef Water Quality: Current Issues. Great Barrier Reef Marine Park Authority, Townsville.
- He, J., Zielinska, B. and Balasubramanian, R., 2010. Composition of semi-volatile organic compounds in the urban atmosphere of Singapore: influence of biomass burning. *Atmospheric Chemistry and Physics*, 10, 11401-11413.
- He, X., Pang, Y., Song, X., Chen, B., Feng, Z. and Ma, Y., 2014. Distribution, sources and ecological risk assessment of PAHs in surface sediments from Guan River Estuary, China. *Marine Pollution Bulletin*, 80, 52-58.
- Hedges, J.I. and Prahl, F.G., 1993. Early diagenesis: consequences for applications of molecular biomarkers. In: Organic Geochemistry. Springer, 237-253.
- Hegazi, A.H. and Andersson, J.T., 2007. Characterization of polycyclic aromatic sulfur heterocycles for source identification. In: Wang Z, Stout SA (eds) Oil spill environmental forensics. Fingerprinting and source identification. Elsevier jobs in Boston, MA, 147-168.
- Hellmann, H., 1999. PAH background levels in contaminated soils. *Wasser Boden*, 51(10), 26-29.
- Hellou, J., Feraut, M., Leonard, J. and Li, W., 2006. Defining the bioaccumulation of PAC in mussels: considering time, distance and effects. *Polycyclic Aromatic Compounds*, 26, 1-16.
- Henner, P., Schiavon, M., Morel, J.L. and Lichtfouse, E., 1997. Polycyclic Aromatic Hydrocarbon (PAH) occurrence and remediation methods. *Analysis Magazine*, 25 (9-10), 56-59.
- Hoffman, E.J., Mills, G.L., Latimer, J.S. and Quinn, J.G., 1984. Urban runoff as a source of polycyclic aromatic hydrocarbons to coastal waters. *Environmental Science & Technology*, 18, 580-587.
- Hong, Y., Yu, S., Yu, G., Liu, Y., Li, G. and Wang, M., 2012. Impacts of urbanization on surface sediment quality: evidence from polycyclic aromatic hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs) contaminations in the Grand Canal of China. *Environmental Science and Pollution Research*, 19, 1352-1363.

- Howsam, M. and Jones, K.C., 1998. Sources of PAHs in the Environment. In Neilson, A. H. (ed.): PAHs and related compounds. *Springer-Verlag, Berlin, Heidelberg*, 137-174.
- Hu, L., Guo, Z., Feng, J., Yang, Z. and Fang, M., 2009. Distributions and sources of bulk OM and aliphatic hydrocarbons in the surface sediments of the Bohai Sea, China. *Marine Chemistry*, 113, 197-211.
- Huang, W., Wang, Z. and Yan, W., 2012. Distribution and sources of polycyclic aromatic hydrocarbons (PAHs) in sediments from Zhanjiang Bay and Leizhou Bay, South China. *Marine Pollution Bulletin*, 64, 1962-1969.
- IDPH. 2005. (Illinois Department of Public Health). Environmental health fact sheet: polycyclic aromatic hydrocarbons. Available at <http://www.idph.state.il.us/envhealth/factsheets/polycyclicaromatichydrocarbons.htm>.
- IARC 1973. International Agency for Research on Cancer. Certain polycyclic aromatic hydrocarbons and heterocyclic compounds. Monograph on the evaluation of carcinogenic risks of the chemical to man. Vol. 3. Lyon, France: *World Health Organization*.
- ICH. 2005. Harmonized Tripartite Guideline. Validation of analytical procedures: Text and methodology Q2 (R1), *ICH Working Group*.
- IPCS. 1998. Environmental Health Criteria 198; Diazinon, Geneva, *World Health Organization*, 140 pp.
- Iqbal, J., Overton and E.B. Gisclair, D., 2008. Polycyclic aromatic hydrocarbons in Louisiana Rivers and coastal environments: source fingerprinting and forensic analysis. *Environmental Forensics*, 9, 63-74.
- Irwin, R., 1997. National Park Service. *Environmental Contaminants Encyclopedia PAHs Entry, National Park Service, Colorado*.
- Isobe, K.O., Zakaria, M.P., Chiem, N.H., Minh, L.Y., Prudente, M., Boonyatumanond, R., Saha, M., Sarkar, S. and Takada, H., 2004. Distribution of linear alkylbenzenes (LABs) in riverine and coastal environments in South and Southeast Asia. *Water Research*, 38, 2449-2459.
- Iwata, H., Tanabe, S., Sakai, N., Nishimura, A. and Tatsukawa, R., 1994. Distribution of persistent organochlorines in the oceanic air and surface sea water and the role of ocean on their global transport and fate. *Environmental Pollution*, 85, 15-33.
- Jaffe', R., Mead, R., Hernandez, M.E., Peralba, M.C. and DiGuida, O.A., 2001. Origin and transport of sedimentary organic matter in two subtropical estuaries: a comparative, biomarker-based study. *Org. Geochemical Journal*, 32, 507-526.

- Jeng, W.-L., 2006. Higher plant n-alkane average chain length as an indicator of petrogenic hydrocarbon contamination in marine sediments. *Marine Chemistry*, 102(3), 242-251.
- Jennings, A.A., 2012. Worldwide regulatory guidance values for surface soil exposure to carcinogenic or mutagenic polycyclic aromatic hydrocarbons, *Journal of Environmental Management*, 110, 82-102.
- Jiang, J.-J., Lee, C.-L., Fang, M.-D. and Liu, J.-T., 2009. Polycyclic Aromatic Hydrocarbons in Coastal Sediments of Southwest Taiwan: An Appraisal of Diagnostic Ratios in Source Recognition. *Marine Pollution Bulletin*, 58, 752-760.
- Jinot, J. and Bayad, S., 1996. Respiratory health effects of exposure to environmental tobacco smoke. *Revolution Environmental Health*, 11 (3), 89-100.
- Kakareka, S.V. and Kukharchyk, T.I., 2003. PAH emissions from open burning of agricultural debris. *Science of the Total Environment*, 308, 257-261. DOI: 10.1016/S0048-9697(02)00650-2.
- Kannan, K., Johnson-Restrepo, B., Yohn, S.S., Giesy, J.P. and Long, D.T., 2005. Spatial and Temporal Distribution of Polycyclic Aromatic Hydrocarbons in Sediments from Michigan Inland Lakes. *Environmental Science & Technology*, 39 (13), 4700-4706.
- Kanzari, F., Syakti, A. D., Asia, L., Malleret, L., Mille, G., Jamoussi, B., Abderrabba, M. and Doumenq, P., 2012. Aliphatic hydrocarbons, polycyclic aromatic hydrocarbons, polychlorinated biphenyls, organochlorine, and organophosphorous pesticides in surface sediments from the Arc river and the Berre lagoon, France. *Environmental Science and Pollution Research*, 19, 559-576.
- Kanzari, F., Syakti, A., Asia, L., Malleret, L., Piram, A., Mille, G. and Doumenq, P., 2014. Distributions and sources of persistent organic pollutants (aliphatic hydrocarbons, PAHs, PCBs and pesticides) in surface sediments of an industrialized urban river (Huveaune), France. *Science of the Total Environment*, 478, 141-151.
- Karacik, B., Okay, O., Henkelmann, B., Bernhöft, S. and Schramm, K.-W., 2009. Polycyclic aromatic hydrocarbons and effects on marine organisms in the Istanbul Strait. *Environment International*, 35, 599-606.
- Karami, A., Christianus, A., Ishak, Z., Shamsuddin, Z.H., Masoumian, M. and Courtenay, S.C., 2012a. Use of intestinal *Pseudomonas aeruginosa* in fish to detect the environmental pollutant benzo [a] pyrene. *Journal of Hazardous Materials*, 215-216, 108-114. doi: 10.1016/j.jhazmat.2012.02.038.
- Karyab, H., Nasser, S., Ahmadkhaniha, R., Rastkari, N., Mahvi, A. H., Nabizadeh, R. and Yunesian, M., 2014. Determination and Source Identification of

- Polycyclic Aromatic Hydrocarbons in Karaj River, Iran. *Bulletin of Environmental Contamination and Toxicology*, 92, 50-56.
- Katsoyiannis, A., Terzi, E., Cai and Q.-Y., 2007. On the use of PAH molecular diagnostic ratios in sewage sludge for the understanding of the PAH sources. Is this use appropriate?. *Chemosphere*, 69, 1337-1339.
- Ke, L., Wong, T.W.Y., Wong, Y.S. and Tam, N.F.Y., 2002. Fate of polycyclic aromatic hydrocarbon (PAH) contamination in a mangrove swamp in Hong Kong following an oil spill. *Marine Pollution Bulletin*, 45 (1-12), 339-347.
- Ke, L., Wang, W.Q., Wong, T.W.Y., Wong, Y.S. and Tam, N.F.Y., 2003a. Removal of pyrene from contaminated sediments by mangrove microcosms. *Chemosphere*, 51 (1), 25-34.
- Ke, L., Yu, K.S.H., Wong, Y.S. and Tam, N.F.Y., 2005. Spatial and vertical distribution of polycyclic aromatic hydrocarbons in mangrove sediments. *Science of The Total Environment*, 340 (1-3), 177-187.
- Kennish, M.J., 1997. Practical handbook of estuarine and marine pollution. *CRC Press, London*, 524 pp.
- Keshavarzifard, M., Zakaria, M.P., Hwai, T.S., Mustafa, S., Vaezzadeh, V., Magam, S.M., Masood, N., Alkhadher, S.A. and Abootalebi-Jahromi, F., 2014. Baseline distributions and sources of Polycyclic Aromatic Hydrocarbons (PAHs) in the surface sediments from the Prai and Malacca Rivers, Peninsular Malaysia. *Marine Pollution Bulletin*, 88, 366-372.
- Keshavarzifard, M., Zakaria M.P., Hwai, T.S., Yusuff, F.M. and Mustafa, S., 2015. Distributions and source apportionment of sediment-associated polycyclic aromatic hydrocarbons (PAHs) and hopanes in rivers and estuaries of Peninsular Malaysia. *Environmental Science and Pollution Research*, DOI 10.1007/s11356-015-4093-7.
- Khalili, N.R., Scheff, P.A. and Holsen, T.M., 1995. PAH source fingerprints for coke ovens, diesel and, gasoline engines, highway tunnels, and wood combustion emissions. *Atmospheric Environment*, 29, 533-542.
- Kim, G.B., Maruya, K.A., Lee, R.F., Lee, J.-H., Koh, C.-H. and Tanabe, S., 1999. Distribution and sources of polycyclic aromatic hydrocarbons in sediments from Kyeonggi Bay, Korea. *Marine Pollution Bulletin*, 38, 7-15.
- Klekowski Jr, E.J., Corredor, J.E., Morell, J.M. and Del Castillo, C.A., 1994. Petroleum pollution and mutation in mangroves. *Marine Pollution Bulletin*, 28 (3), 166-169.
- Koe, L.C.C., Arellano, A.F., and McGregor, J.L., 2001. Investigating the haze transport from 1997 biomass burning in Southeast Asia: Its impact upon Singapore, *Atmospheric Environment*, 35, 2723– 2734.

- Krauss, M., Wilcke, W., Martius, C., Bandeira, A.G., Garcia, M.V B. and Amelung, W., 2005. Atmospheric versus biological sources of polycyclic aromatic hydrocarbons (PAHs) in a tropical rain forest environment. *Environmental Pollution*, 135, 143-154.
- Kumata, H., Uchida, M., Sakuma, E., Uchida, T., Fujiwara, K., Tsuzuki, M., Yoneda, M. and Shibata, Y., 2006. Compound class specific ¹⁴C analysis of polycyclic aromatic hydrocarbons associated with PM₁₀ and PM₁. 1 aerosols from residential areas of suburban Tokyo. *Environmental Science and Technology*, 40, 3474-3480.
- Kwon, S.H., Kim, Y.P. and Lee, J.Y., 2013. Impact of Dust Storms on the Organic Composition in the Ambient Aerosol. *Aerosol and Air Quality Research*, 13, 97-106. doi: 10.4209/aaqr.2012.04.0105.
- Larsen, III R.K. and Baker, J.E., 2003. Source apportionment of polycyclic aromatic hydrocarbons in the urban atmosphere: a comparison of three methods. *Environmental Science and Technology*, 37, 1873-81.
- Latimer, J.S., Hoffman, E.J., Hoffman, G., Fasching, J.L. and Quinn, J.G., 1990. Sources of petroleum hydrocarbons in urban runoff. *Water, Air, and Soil Pollution*, 52, 1-21.
- LBNL. 2004. Lawrence Berkeley National Laboratory. Constraining Uncertainties about the Sources and Magnitude of Ambient Air Exposures to Polycyclic Aromatic Hydrocarbons (PAHs): the State of Minnesota as a Case Study. Paper LBNL-54473. *University of California, Berkeley, CA, USA*.
- Leal-Granadillo, I.A., Alonso, J.I. and Sanz-Medel, A., 2000. Determination of n-alkanes and polycyclic aromatic hydrocarbons in atmospheric particulate and vapour phases in Oviedo, Spain, by GC-MS. *Journal of environmental monitoring*, 2(3), 218-222.
- Le Dréau, Y., Jacquot, F., Doumenq, P., Guiliano, M., Bertrand, J. C. and Mille, G., 1997. Hydrocarbon balance of a site which had been highly and chronically contaminated by petroleum wastes of a refinery (from 1956 to 1992). *Marine Pollution Bulletin*, 34(6), 456-468.
- Leong, K.H., Benjamin Tan, L.L. and Mustafa, A.M., 2007. Contamination levels of selected organochlorine and organophosphate pesticides in the Selangor River, Malaysia between 2002 and 2003. *Chemosphere*, 66, 1153-1159. doi:10.1016/j.chemosphere.2006.06.009.
- Levendis, Y.A., Atal, A. and Carlson, J.B., 1998. On the correlation of CO and PAH emissions from the combustion of pulverized coal and waste tires. *Environmental Science and Technology*, 32(23), 3767-3777.
- Li, M., 2003. Ambient concentrations and measurement precision of molecular markers in fine particles from Philadelphia, PA. *Drexel Thesis and Dissertations*.

- Li, C.-H., Zhou, H.-W. and Wong, Y.-S. and Tam, N.F.-Y., 2009. Vertical distribution and anaerobic biodegradation of polycyclic aromatic hydrocarbons in mangrove sediments in Hong Kong, South China. *Science of The Total Environment* 407, (21), 5772-5779.
- Li, W., Peng, Y. and Bai, Z., 2010. Distributions and Sources of n-alkanes in PM_{2.5} at Urban, Industrial and Coastal Sites in Tianjin, China. *Journal of Environmental Sciences*, 22, 1551-1557.
- Liang, Y., Tse, M., Young, L. and Wong, M., 2007. Distribution patterns of polycyclic aromatic hydrocarbons (PAHs) in the sediments and fish at Mai Po Marshes Nature Reserve, Hong Kong. *Water Research*, 41, 1303-1311.
- Lipiatou, E. and Saliot, A., 1991. Fluxes and transport of anthropogenic and natural polycyclic aromatic hydrocarbons in the western Mediterranean Sea. *Marine Chemistry*, 32, 51-71.
- Liu, S., Tao, S., Liu, W., Liu, Y., Dou, H., Zhao, J., Wang, L., Wang, J., Tian, Z. and Gao, Y., 2007. Atmospheric polycyclic aromatic hydrocarbons in North China: a winter-time study. *Environmental Science & Technology*, 41, 8256-8261.
- Liu, S., Tao, S., Liu, W., Dou, H., Liu, Y., Zhao, J. and Gao, Y., 2008a. Seasonal and spatial occurrence and distribution of atmospheric polycyclic aromatic hydrocarbons (PAHs) in rural and urban areas of the North Chinese Plain. *Environmental Pollution*, 156(3), 651-656.
- Liu, Y., Chen, L., Jianfu, Z., Qinghui, H., Zhiliang, Z. and Hongwen, G., 2008b. Distribution and Sources of Polycyclic Aromatic Hydrocarbons in Surface Sediments of Rivers and an Estuary in Shanghai, China. *Environmental Pollution*, 154, 298-305.
- Liu, A., Lang, Y., Xue, L. and Liu, J., 2009. Ecological risk analysis of polycyclic aromatic hydrocarbons (PAHs) in surface sediments from Laizhou Bay. *Environmental Monitoring and Assessment*, 159, 429-436.
- Liu, L.Y., Wang, J.Z., Wei, G.L., Guan, Y.F. and Zeng, E.Y., 2012. Polycyclic aromatic hydrocarbons (PAHs) in continental shelf sediment of China: implications for anthropogenic influences on the coastal marine environment. *Environmental Pollution*, 167, 155-162.
- Liu, L.Y., Wang, J.Z., Wong, C.S., Qiu, J.W. and Zeng, E.Y., 2013. Application of multiple geochemical markers to investigate organic pollution in a dynamic coastal zone. *Environmental Toxicology and Chemistry*, 32, 312-319.
- Llamas, A., Al-Lal, A.-M., García-Martínez, M.-J., Ortega, M.F., Llamas, J.F., Lapuerta, M. and Canoira L., 2017. Polycyclic Aromatic Hydrocarbons (PAHs) produced in the combustion of fatty acid alkyl esters from different feedstocks: Quantification, statistical analysis and mechanisms of formation. *Science of the Total Environment*, 586, 446-456.

- Long, E., Macdonald, D., Smith, S. and Calder, F., 1995. Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments. *Environmental Management*, 19 (1), 81-97.
- Long, E.R., Field, L.J. and MacDonald, D.D., 1998. Predicting toxicity in marine sediments with numerical Sediment quality guidelines. *Environmental Toxicology and Chemistry*, 17 (4), 714-727.
- Long, E.R. and MacDonald, D.D., 1998. Recommended Uses of Empirically Derived, Sediment quality guidelines for Marine and Estuarine Ecosystems. Humanband Ecological Risk Assessment: *An International Journal*, 4 (5), 1019-1039.
- Long, E.R., MacDonald, D.D., Severn, C.G. and Hong, C.B., 2000. Classifying probabilities of acute toxicity in marine sediments with empirically derived Sediment quality guidelines. *Environmental Toxicology and Chemistry*, 19 (10), 2598-2601.
- Lourenço, R.A., Araujo Júnior, M.A.G., Meireles Júnior, R.O., Macena, L.F., de A. Lima, E.F. and Carneiro, M.E.R., 2013. Aliphatic and polycyclic aromatic hydrocarbons and trace elements as indicators of contamination status near oil and gas platforms in the Sergipe– Alagoas Basin (Southwest Atlantic Ocean). *Continental Shelf Research*, 71, 37-44.
- Luellen, D.R. and Shea, D., 2003. Semipermeable membrane devices accumulate conserved ratios of sterane and hopane petroleum biomarkers. *Chemosphere*, 53, 705-713.
- Lukman, M., Balzer, W. and Jose, C., 2006. Polycyclic Aromatic Hydrocarbon (PAHs) pollution in coastal sediment from Riau Province, Indonesia. *Extended Abstract. Proceeding*.
- Lundstedt, S., Haglund, P. and Orberg, L., 2003. Degradation and formation of polycyclic aromatic compounds during bioslurry treatment of an aged gasworks soil. *Journal of Environmental Toxicological Chemistry*, 22, 1413-1420.
- Luo, X.-J., Chen, S.-J., Ni, H.-G., Yu, M. and Mai, B.-X., 2008. Tracing sewage pollution in the Pearl River Delta and its adjacent coastal area of South China Sea using linear alkylbenzenes (LABs). *Marine Pollution Bulletin*, 56, 158-62.
- Ma, X., Ran, Y., Gong, J. and Zou, M., 2007. Concentrations and inventories of polycyclic aromatic hydrocarbons and organochlorine pesticides in watershed soil in the Pearl River Delta, China. *Environmental Monitoring and Assessment*, 145, 453-64.
- Macdonald, D.D., Carr, R.S., Calder, F.D., Long, E.R. and Ingersoll, C.G., 1996. Development and evaluation of Sediment quality guidelines for Florida coastal waters. *Ecotoxicology*, 5 (4), 253-278.

- MacDonald, D.D., Ingersoll, C.G. and Berger, T.A., 2000. Development and Evaluation of Consensus-Based Sediment quality guidelines for Freshwater Ecosystems. *Archives of Environmental Contamination and Toxicology*, 39 (1), 20-31.
- Mackay, D., Shiu, W.Y. and Ma, K.C., 1992. Illustrated Handbook of Physical-Chemical Properties and Environmental Fate for Organic Chemicals. *Chelsea, Michigan: Lewis Publishers*.
- Magam, S.M., Zakaria, M.P., Halimoon, N., Masood, N. and Alsalahi, M.A., 2012. Distribution of linear alkylbenzenes (LABs) in sediments of Sarawak and Sembulan Rivers Malaysia. *Environment Asia*, 5(1), 48-55.
- Magam, S.M., Zakaria, M.P., Halimoon, N., Aris, A.Z., Kannan, N., Masood, N., Mustafa, S., Alkhadher, S.A., Keshavarzifard, M., Vaezzadeh, V., Sani, M. and Latif, M.T., 2015. Evaluation of distribution and sources of sewage molecular marker (LABs) in selected rivers and estuaries of Peninsular Malaysia. *Environmental Science and Pollution Research*, 23(6), 5693-5704. doi 10.1007/s11356-015-5804-9.
- Magi, E., Bianco, R., Ianni, C. and Di Carro, M., 2002. Distribution of polycyclic aromatic hydrocarbons in the sediments of the Adriatic Sea. *Environmental Pollution*, 119, 91-98.
- Mahler, B.J., Van Metre, P.C., Bashara, T.J., Wilson, J.T. and Johns, D.A., 2005. Parking lot sealcoat: An Unrecognized Source of Urban Polycyclic Aromatic Hydrocarbons. *Environmental Science and Technology*, 39 (15), 5560-66.
- Mahro, B., Schaefer, G. and Kästner, M., 1994. Pathways of microbial degradation of polycyclic aromatic hydrocarbons in soil. in: Hinchey RE, Leeson A, Semprini L, Ong SK (Eds.), *Bioremediation of Chlorinated and Polycyclic Aromatic Hydrocarbon Compounds CRC Press, Florida*, 203-217.
- Mandalakis, M., Gustafsson, O., Reddy, C.M. and Xu, L., 2004. Radiocarbon apportionment of fossil versus biofuel combustion sources of polycyclic aromatic hydrocarbons in the Stockholm metropolitan area. *Environmental Science and Technology*, 38, 5344-5349.
- Manoli, E. and Samara, C., 1999. Polycyclic Aromatic Hydrocarbons in Natural Waters: Sources, Occurrence and Analysis. *Trends in Analytical Chemistry*, 18, 417-428.
- Manoli, E., Samara, C., Konstantinou, I. and Albanis, T., 2000. Polycyclic aromatic hydrocarbons in the bulk precipitation and surface waters of Northern Greece. *Chemosphere*, 41, 1845-1855.
- Marr, L.C., Kirchstetter, T.W., Harley, R.A., Miguel, A.H., Hering, S.V and Hammond, S.K., 1999. Characterization of polycyclic aromatic hydrocarbons in motor vehicle fuels and exhaust emissions. *Environmental Science and Technology*, 33, 3091-3099.

- Martins, C.C., Ferreira, J.A., Taniguchi, S., Mahiques, M.M., Bicego, M.C. and Montone, R.C., 2008. Spatial distribution of sedimentary linear alkylbenzenes and faecal steroids of Santos Bay and adjoining continental shelf, SW Atlantic, Brazil: origin and fate of sewage contamination in the shallow coastal environment. *Marine Pollution Bulletin*, 56, 1353-1376.
- Martins, C.C., Bicego, M.C., Mahiques, M.M., Figueira, R.C.L., Tessler, M.G. and Montone, R.C., 2010. Depositional history of sedimentary linear alkylbenzenes (LABs) in a large South American industrial coastal area (Santos Estuary, Southeastern Brazil). *Environmental Pollution*, 158, 3355-3364.
- Martins, C.C., Aguiar, S.N., Bicego, M.C. and Montone, R.C., 2012. Sewage organic markers in surface sediments around the Brazilian Antarctic station: results from the 2009/10 austral summer and historical tendencies. *Marine Pollution Bulletin*, 64, 2867-2870.
- Maruya, K.A., Risebrough, R.W. and Horne, A.J., 1996. Partitioning of polynuclear aromatic hydrocarbons between sediments from San Francisco Bay and their porewater. *Environmental Science and Technology*, 30, 2942-2947.
- Masiol, M., Hofer, A., Squizzato, S., Piazza, R., Rampazzo, G. and Pavoni, B., 2012. Carcinogenic and mutagenic risk associated to airborne particle-phase polycyclic aromatic hydrocarbons: A source apportionment. *Atmospheric Environment*, 60, 375-382.
- Masood, N., Zakaria, M.P., Ali, M.M., Magam, S.M., Alkhadher, S.A., Keshavarzifard, M., Vaezzadeh, V. and Hussein, M.A., 2014. Distribution of Petroleum Hydrocarbons in Surface Sediments from Selected Locations in Kuala Selangor River, Malaysia. A.Z. Aris et al. (eds.), *From Sources to Solution. Springer*, 351-356. doi:10.1007/978-981-4560-70-2_64, 351-356.
- Masood, N., Zakaria, M.P., Halimoon, N., Aris, A.Z., Magam, S.M., Kannan, N., Mustafa, S., Ali, M.M., Keshavarzifard, M., Vaezzadeh, V., Alkhadher, S.A. and Al-Odaini, N.A., 2016. Anthropogenic waste indicators (AWI) particularly PAHs and LABs in Malaysian sediments: application of aquatic environment for identifying anthropogenic pollution. *Marine Pollution Bulletin*, 102, 160-175.
- Masoud, M.S., Mahmoud, Th.H. and Abdel-halim, A.M., 2001. Chemical studies of El-Mex Bay, Alexandria. *Proceeding of the Second Conference and Exhibition for Life and Environment*, 3-5 April, Alexandria, 339-60.
- Massara Paletto, V., Commendatore, M.G. and Esteves, J.L., 2008. Hydrocarbon levels in sediments and bivalve mollusks from Bahía Nueva (Patagonia, Argentina): An assessment of probable origin and bioaccumulation factors. *Marine Pollution Bulletin*, 56(12), 2100-2105.

- Mastral, A.M. and Callen, M.S., 2000. A review on polycyclic aromatic hydrocarbon (PAH) emissions from energy generation. *Environmental Science and Technology*, 34, 3051-3057.
- Mazurek, M.A. and Simoneit, B.R.T., 1984. Characterization of biogenic and petroleum-driven organic matter in aerosols over remote, rural, and urban area. In: Identification and analysis of organic pollutants in air. Keith, L.H. (Ed.). *Ann Arbor Science Woburn, MA*, 353-370.
- McCarthy, B., 2003. Polycyclic aromatic hydrocarbons (PAHs) in urban runoff. *The GEI MGP Reporter*, 1-3.
- McCready, S., Slee, D.J., Birch, G.F. and Taylor, S.E., 2000. The Distribution of Polycyclic Aromatic Hydrocarbons in Surficial Sediments of Sydney Harbour, Australia. *Marine Pollution Bulletin*, 40 (11), 999-1006.
- Medeiros, P.M. and Bícigo, M.C., 2004. Investigation of natural and anthropogenic hydrocarbon inputs in sediments using geochemical markers. II. São Sebastião, SP e Brazil. *Marine Pollution Bulletin*, 49, 761-769.
- Medeiros, P.M., Bícigo, M.C., Castelao, R.M., Del Rosso, C., Fillmann, G. and Zamboni, A.J., 2005. Natural and anthropogenic hydrocarbon inputs to sediments of Patos Lagoon Estuary, Brazil. *Environment International*, 31, 77-87.
- Mehta, K.V., 2010. Physicochemical characteristics and statistical study of groundwater of some places of vadgam taluka in banaskantha district of Gujarat State (India). *Journal of Chemical and Pharmaceutical Research*, 2(4), 663-670.
- Menzie, C.A., Potocki, B.B. and Santodonato, J., 1992. Exposure to carcinogenic PAHs in the environment. *Environmental Science and Technology*, 26, 1278-83.
- Mersch-Sundermann, V., Mochayedi, S. and Kevekordes, S., 1992. Genotoxicity of polycyclic aromatic hydrocarbons in *Escherichia coli* PQ37. *Mutation Research*, 278, 1-9.
- Meyers, P.A., and Ishiwatari, R., 1993. Lacustrine organic geochemistry—an overview of indicators of organic matter sources and diagenesis in lake sediments. *Organic Geochemistry*, 20(7), 867-900.
- Meyers, P.A. and Ishiwatari, R., 1995. Organic matter accumulation records in lake sediments. In: Physics and chemistry of lakes. Lerman, A., Imboden, D.M., Gat, J.R. (Eds.). *Springer, Berlin*, 279-328.
- Meyers, P.A., 2003. Applications of organic geochemistry to paleolimnological reconstructions: a summary of examples from the Laurentian Great Lakes. *Organic Geochemistry*, 34(2), 261-289.

- Mai, B.X., Fu, J.M., Sheng, G.Y., Kang, Y.H., Lin, Z., Zhang, G., Min, Y.S. and Zeng, E.Y., 2002. Chlorinated and polycyclic aromatic hydrocarbons in riverine and estuarine sediments from Pearl River Delta China. *Environmental Pollution*, 117, 457-474.
- Minai-Tehrani, D., Minoui, S. and Herfatmanesh, A., 2009. Effect of salinity on biodegradation of polycyclic aromatic hydrocarbons (PAHs) of heavy crude oil in soil. *Bulletin of Environmental Contamination and Toxicology*, 82(2), 179-184. doi:10.1007/s00128-008-9548-9.
- Mirsadeghi, S.A., 2010. Bioaccumulation and Hazard Assessment of Polycyclic Aromatic Hydrocarbons in Cockle (*Anadara Granosa* L.) From Selected Intertidal Mudflats of Peninsular Malaysia. *PhD Thesis, Universiti Putra Malaysia*.
- Mirsadeghi, S.A., Zakaria, M.P., Yap, C.K. and Shahbazi, A., 2011. Risk assessment for the daily intake of polycyclic aromatic hydrocarbons from the ingestion of cockle (*Anadara granosa*) and exposure to contaminated water and sediments along the west coast of Peninsular Malaysia. *Journal of Environmental Sciences*, 23, 336-345.
- Mirsadeghi, S.A., Zakaria, M.P., Yap, C.K. and Gobas, F., 2013. Evaluation of the potential bioaccumulation ability of the blood cockle (*Anadara granosa*) for assessment of environmental matrices of mudflats. *Science of The Total Environment*, 454, 584-597.
- Mitchell, P.K.K., Mills, G., Fisher-Niwa, G. and Eason-Landcare, C., 1998. *Technical paper No. 37 Toxic*.
- Mohamed, I.B. and Mohamed, A.E., 2010. Distribution of polycyclic aromatic hydrocarbons in drinking water in Egypt. *Desalination*, 251, 34-40. doi:10.1016/j.desal.2009.09.148.
- Mohammad, N.A., Mastura, S.S.A. and Akhir, J.M., 2007. Land use evaluation for Kuala Selangor, Malaysia using remote sensing and GIS technologies. *GEOGRAFIA Online™ Malaysia Journal of Society and Space*, 3 (1-18) © 2007, ISSN 2180-2491.
- Monza, L.B., Loewy, R.M., Savini, M.C. and Pechen de d'Angelo, A.M., 2013. Sources and distribution of aliphatic and polyaromatic hydrocarbons in sediments from the Neuquen River, Argentine Patagonia. *Journal of Environmental Science and Health, Part A*, 48(4), 370-379.
- MOOPAM, 1999. Manual of Oceanographic Observation and Pollutant Analyses Methods. 3rd edition. *Regional Organization for Protection of the Marine Environment (ROPME)*, Kuwait.
- Morillo, E., Romero, A., Maqueda, C., Madrid, L., Ajmone-Marsan, F., Grçman, H., Davidson, C.M., Hursthouse, A.S. and Villaverde, J., 2007. Soil pollution by

- PAHs in urban soils: a comparison of three European cities. *Environmental Monitoring and Assessment*, 9(9),1001-1008.
- Morrison, R.T. and Boyd, R.N., 1992. Organic Chemistry (6th ed.). New Jersey: *Prentice Hall*, ISBN 0-13-643669-2.
- Mostafa, A.R., Wade, T.L., Sweet, S.T., Al-Alimi, A.K.A. and Barakat, A.O., 2009. Distribution and characteristics of polycyclic aromatic hydrocarbons (PAHs) in sediments of Hadhramout coastal area, Gulf of Aden, Yemen. *Journal of Marine Systems*, 78, 1-8.
- Motelay-Massei, A., Garban, B., Tiphagne-Larcher, K., Chevreuil, M. and Ollivon, D., 2006. Mass balance for polycyclic aromatic hydrocarbons in the urban watershed of Le Havre (France): transport and fate of PAHs from the atmosphere to the outlet. *Water Research*, 40, 1995-2006.
- Mukherjee, P. and Viswanathan, S., 2001. Contributions to CO concentrations from biomass burning and traffic during haze episodes in Singapore, *Atmospheric Environment*, 35, 715-725.
- Nadal, M., Schuhmacher, M. and Domingo, J.L., 2004. Levels of PAHs in soil and vegetation samples from Tarragona County, Spain. *Environmental Pollution*, 132(1), 1-11.
- Nash, R.G. and Hill, B.D., 1990. Modelling pesticide volatilization and soil decline under controlled conditions. In Long range transport of pesticides. ed. D. A. Kurtz. *Lewis Publishers, Inc., Michigan, USA*. 17-28.
- Nasher, E., Yook Heng, L. Zakaria, Z. and Surif, S., 2013. Concentrations and sources of polycyclic aromatic hydrocarbons in the seawater around Langkawi Island, Malaysia, *Journal of Chemistry*, Article ID 975781, 10 pages.
- Neff, J.M., 1985. Polycyclic Aromatic Hydrocarbons. In Fundamentals of Aquatic Toxicology, G. M. Rand and S. R. Petrocelli (Eds.). *New York: Hemisphere Publishing Corporation*, 416-454.
- Nelson, E., 2001. The miseries of passive smoking. *Human Exposure Toxicology*, 20, 61-83.
- Neser, G., Kontas, A., Unsalan, D., Altay, O., Darilmaz, E., Uluturhan, E., Kucuksezgin, F., Tekogul, N. and Yercan, F., 2012. Polycyclic aromatic and aliphatic hydrocarbons pollution at the coast of Aliaga (Turkey) ship recycling zone. *Marine Pollution Bulletin*, 64, 1055-1059.
- Nicholls, P. and Ellis, J., 2002. Fringing habitats in estuaries: the sediments-mangrove connection. *Water and Atmosphere*, 10(4), 24-25.
- Nozar, S.L.M., Ismail, W.R. and Zakaria, M.P., 2014. Distribution, Sources Identification, and Ecological Risk of PAHs and PCBs in Coastal Surface

Sediments from the Northern Persian Gulf. *Human and Ecological Risk Assessment: An International Journal*, 20, 1507-1520.

NRC. 2003. National Research Council, Oil in the Sea III: Inputs, Fates, and Effects. *National Academies Press, Washington, D.C.*, p. 280.

NTP. 2005. National Toxicology Program. Report on Carcinogens, eleventh ed., Public Health Service, US. *Department of Health and Human Services, Washington, D.C.*

Ni, H.G., Lu, F.H., Wang, J.Z., Guan, Y.F. and Zeng, E.Y., 2008. Linear alkylbenzenes in riverine runoff of the Pearl River Delta (China) and their application as anthropogenic molecular markers in coastal environments. *Environmental Pollution*, 154, 348-355.

Ni, H.G., Shen, R.L., Zeng, H. and Zeng, E.Y., 2009. Fate of linear alkylbenzenes and benzothiazoles of anthropogenic origin and their potential as environmental molecular markers in the Pearl River Delta, South China. *Environmental Pollution*, 157, 3502-3507.

Nielsen, T., Jorgensen, H.E., Larsen, J.C. and Poulsen, M., 1996. City air pollution of polycyclic aromatic hydrocarbons and other mutagens: occurrence, sources and health effects. *Science of the Total Environment*, 189(190), 41-49.

Nisbet, I.C.T. and LaGoy, P.K., 1992. Toxic equivalency factors (TEFs) for polycyclic aromatic hydrocarbons (PAHs). *Regulatory Toxicology and Pharmacology*, 16(3), 290-300.

Notar, M., Leskovšek, H. and Faganeli, J., 2001. Composition, distribution and sources of polycyclic aromatic hydrocarbons in sediments of the Gulf of Trieste, Northern Adriatic Sea. *Marine Pollution Bulletin*, 42, 36-44.

NPI. 2004. National Pollution Inventory, Polycyclic aromatic hydrocarbons. Available at <http://www.npi.gov.au/database/substance-info/profiles/74.html>.

Nylund, L., Heikkilä, P., Hameila, M., Pyy, L., Linnainmaa, K. and Sorsa, M., 1992. Genotoxic effects and chemical composition of four creosotes. *Mutation Research*, 265, 223-236.

Oen, A.M.P., Cornelissen, G. and Breedveld, G.D., 2006. Relation between PAH and black carbon contents in size fractions of Norwegian harbor sediments. *Environmental Pollution*, 141, 370-380.

Ogogo, N.K., 2013. Characterisation and source apportionment of polycyclic aromatic hydrocarbons in the sediments of Brisbane River. *Master thesis. Queensland University of Technology*.

Okuda, T., Kumata, H., Zakaria, M. P., Naraoka, H., Ishiwatari, R. and Takada, H., 2002. Source identification of Malaysian atmospheric polycyclic aromatic

- hydrocarbons nearby forest fires using molecular and isotopic compositions. *Atmospheric Environment*, 36, 611-618.
- Omar, N.-Y.-M.-J., Abas, M.-R.-B., Ketuly, K.-A. and Tahir, N.-Md., 2002. Concentrations of PAHs in atmospheric particles (PM-10) and road side soil particles collected in Kuala Lumpur, Malaysia. *Atmospheric Environment*, 36, 247-254.
- Omar, N.-Y.-M.-J., Mon, T.-C., Rahman, N.-A. and Abas, M.-R.-B., 2006. Distributions and health risks of polycyclic aromatic hydrocarbons (PAHs) in atmospheric aerosols of Kuala Lumpur, Malaysia. *Science of the Total Environment*, 369, 76-81.
- O'Neill, P., 1997. *Chemia srodowiska (Chemistry of the Environment)*. Warsaw, PL: Wydawnictwo Naukowe PWN.
- Ong, D.J., 2005. *Land Use in the Selangor River: A Study on the Application and Strategy for Integrated River Basin Management*, WWF Malaysia, Petaling Jaya.
- Onuska, F.I., 1989. Analysis of Polycyclic Aromatic Hydrocarbons in Environmental Samples: In *Analysis of Trace Organics in the Aquatic Environment*, B. K. Afghan and A. S. Y. Chau (Eds.). Boca Raton: CRC Press, 205-241,
- Oros, D.R., Ross, J.R., Spies, R.B. and Mumley, T., 2007. Polycyclic aromatic hydrocarbon (PAH) contamination in San Francisco Bay: a 10-year retrospective of monitoring in an urbanized estuary. *Environmental Research*, 105, 101-118.
- OSPAR Commission. 2001. *Draft OSPAR background document on Polycyclic Aromatic Hydrocarbons (PAH)*.
- Ouyang, Y., Zhang, J.E. and Ou, L.-T., 2006. Temporal and spatial distributions of sediment total organic carbon in an estuary river. *Journal of Environmental Quality*, 35, 93-100.
- Oyo-Ita, O.E., Ekpo, B.O. and Orosa, D.R., 2010. Distributions and sources of aliphatic hydrocarbons and ketones in surface sediments from the Cross River estuary, S.E. Niger Delta, Nigeria. *Journal of Applied Sciences in Environmental Sanitation*, 5 (1), 1-11.
- Page, D.S., Boehm, .PD., Douglas, G.S., Bence, A.E., Burns, W.A. and Mankiewicz, P.J., 1996. The natural petroleum hydrocarbon background in subtidal sediments of Prince William Sound, Alaska, USA. *Environmental Toxicology and Chemistry*, 15, 1266-1281.
- Page, D.S., P.D. Boehm, G.S. Douglas and A.E. Bence., 1995. Identification of Hydrocarbon Sources in the Benthic Sediments of Prince William Sound and the Gulf of Alaska Following the Exxon Valdez Oil Spill. In *Exxon Valdez Oil Spill: Fate and Effects in Alaskan Waters ASTM Special Technical*

Publication # 1219 (P. G. Wells, J. N. Butler and J. S. Hughes, ed.), *American Society for Testing and Materials, Philadelphia, PA*, 41-83.

- Peng, C., Chen, W., Liao, X., Wang, M., Ouyang, Z., Jiao, W. and Bai, Y., 2011. Polycyclic aromatic hydrocarbons in urban soils of Beijing: Status, sources, distribution and potential risk. *Environmental Pollution*, 159, 802-808.
- Patel, V. and Shah, K., 2013. Petroleum Hydrocarbon Pollution and its Biodegradation. *International Journal of Chemtech Applications*, 2 (3), 63-80.
- Pavlova, A., and Papazova, D., 2003. Oil-spill identification by gas chromatography-mass spectrometry. *Journal of chromatographic science*, 41, 271-273.
- Pendoley, K., 1992. Hydrocarbons in Rowley Shelf (Western Australia) oysters and sediments. *Marine Pollution Bulletin*, 24, 210-215.
- Pereira, W.E., Hostettler, F.D., Luoma, S.N., Van Geen, A., Fuller, C.C. and Anima, R.J., 1999. Sedimentary record of anthropogenic and biogenic polycyclic aromatic hydrocarbons in San Francisco Bay, California. *Marine Chemistry*, 64, 99-113.
- Perez, S., Marinel, La Farré, M., Garcia, M.J. and Barceló, D., 2001. Occurrence of polycyclic aromatic hydrocarbons in sewage sludge and their contribution to its toxicity in the ToxAlert® 100 bioassay, *Chemosphere*, 45(6-7), 705-712.
- Pérez-Fernández, B., Viñas, L., Franco, M.A. and Bargiela, J., 2015. PAHs in the Ría de Arousa (NW Spain): A consideration of PAHs sources and abundance. *Marine Pollution Bulletin*, 95, 155-165.
- Peters, K.E. and Moldowan, J.M., 1993. The Biomarker Guide: Interpreting molecular fossils in petroleum and ancient sediments; *Prentice Hall, Englewood Cliffs, New Jersey*, 363p.
- Peters, C.A., Knightes, C.D., Brown, D.G., 1999. Long-term composition dynamics of PAH-containing NAPLs and implications for risk assessment. *Environmental Science & Technology*, 33, 4499-4507.
- Peters, K.E., Walters, C.C. and Moldowan, J.M., 2005. The biomarker guide: biomarkers and isotopes in the environment and human history, *Cambridge University Press*.
- Phelps, H.L., 2000. DC AES Contaminated Anacostia Estuary Sediments: A Biomonitoring Approach, *Final Technical Report District of Columbia Water Resources Research Center, Washington, DC*.
- Philips, D.H., 1996. DNA adducts in human tissues: biomarkers of exposure to carcinogens in tobacco smoke. *Environmental Health Perspectives*, 104 (3), 453-458.

- Phillips, D.H., 1999. Polycyclic aromatic hydrocarbons in the diet. *Mutation Research*, 443(1-2), 139-47.
- Phuleria, H.C., Sheesley, R.J., Schauer, J.J., Fine, P.M. and Sioutas, C., 2007. Roadside measurements of size-segregated particulate organic compounds near gasoline and diesel-dominated freeways in Los Angeles, CA. *Atmospheric Environment*, 41 (22), 4653-4671.
- Piccardo, M., Coradeghini, R. and Valerio, F., 2001. Polycyclic aromatic hydrocarbon pollution in native and caged mussels. *Marine Pollution Bulletin*, 42, 951-956.
- Poynter, J.G. and Eglinton, G., 1990. Molecular composition of three sediments from hole 717C: the Bengal Fan. In: Cochran, J.R., Stow, D.A.V., et al. (Eds.), . In: *Proceedings of the Ocean Drilling Program Scientific Results*, 116, 155-161.
- Prabhukumar, G. and Pagilla, K., 2010. Polycyclic aromatic hydrocarbons in urban runoff– sources, sinks and treatment: A review. *Department of Civil, Architectural and Environment Engineering, Institute of technology, Chicago, IL* 2010.
- Prahl, F.G., Ertel, J.R., Goni, M.A., Sparrow, M.A. and Eversmeyer, B., 1994, Terrestrial organic carbon contributions to sediments on the Washington margin, *Geochimica et Cosmochimica Acta*, 58, 3035-3048.
- Prince, R.C., Elmendorf, D.L., Lute, J.R., Hsu, C.S., Halth, C.E., Senlus, J.D., Dechert, G.J., Douglas, G.S. and Butler, E.L., 1994. 17a(H),21a(H)-Hopane as a conservative internal marker for estimating the biodegradation of cruce oil. *Environmental Science and Technology*, 142-145.
- Qian, Y., Wade, T.L. and Sericano, J.L., 2001. Sources and bioavailability of polynuclear aromatic hydrocarbons in Galveston Bay, Texas. *Estuaries*, 24, 817-827.
- Qiao, M., Wang, C., Huang, S., Wang, D. and Wang, Z., 2006. Composition, sources, and potential toxicological significance of PAHs in the surface sediments of the Meiliang Bay, Taihu Lake, China. *Environment International*, 32, 28-33.
- Raymundo, C.C. and Preston, M.R., 1992. The distribution of linear alkylbenzenes in coastal and estuarine sediments of western North Sea. *Marine Pollution Bulletin*, 24, 138-146.
- Raza, M., Zakaria, M.P. and Hashim, N.R., 2011. Hydrocarbon Contamination in Mangrove Sediment of Linggi River: A Pilot Study. in: Zakaria MP, Mohamed MI, Kasmin S (Eds.), *Contemporary Environmental Quality Management in Malaysia and Selected Countries. Universiti Putra Malaysia Press, Serdang*, 25-33.
- Raza, M., Zakaria, M.P., Hashim, N.R., Yim, U.H., Kannan, N. and Ha, S.Y. (2013). Composition and source identification of polycyclic aromatic hydrocarbons in

mangrove sediments of Peninsular Malaysia: indication of anthropogenic input. *Environmental Earth Sciences*, 70(6), 2425-2436.

- Ravindra, K., Sokhi, R. and Grieken, R.V., 2008. Atmospheric polycyclic aromatic hydrocarbons: Source attribution, emission factors and regulation. *Atmospheric Environment*, 42, 2895-2921.
- Readman, J.W., Fillmann, G., Tolosa, I., Bartocci, J., Villeneuve, J.-P., Catinni, C. and Mee, L.D., 2002. Petroleum and PAH contamination of the Black Sea. *Marine Pollution Bulletin*, 44(1), 48-62.
- Reeves, W.R., Barhoumi, R., Burghardt, R.C., Lemke, S.L., Mayura, K., McDonald, T.J., Phillips, T.D. and Donnelly, K.C., 2001. Evaluation of methods for predicting the toxicity of polycyclic aromatic hydrocarbon mixtures. *Environmental Science and Technology*, 35, 1630-1636.
- Revill, A.T., Carr, M.R. and Rowland, S.J., 1992. Use of oxidative degradation followed by capillary gas chromatographymass spectrometry and multi dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons. *Journal of Chromatography*, 589, 281-86.
- Retnam, A., Zakaria, M.P., Juahir, H., Aris, A.Z., Zali, M.A. and Kasim, M.F., 2013. Chemometric techniques in distribution, characterisation and source apportionment of polycyclic aromatic hydrocarbons (PAHS) in aquaculture sediments in Malaysia. *Marine Pollution Bulletin*, 69(1), 55-66.
- Rieley, G., Collier, R.J., Jones, D.M. and Eglinton, G., 1991. The biogeochemistry of Ellesmere Lake, UK I: source correlation of leaf wax inputs to the sedimentary record. *Organic Geochemistry*, 17, 901-912.
- Rinawati, K.T., Koike, H., Kurumisawa, R., Ito, M., Sakurai, S., Togo, A., Saha, M., Arifin, Z. and Takada, H., 2012. Distribution, source identification, and historical trends of organic micro pollutants in coastal sediment in Jakarta Bay Indonesia. *Journal of Hazardous Materials*, 217-218, 208-216.
- Rockne, K.J., Shor, L.M., Young, L.Y., Taghon, G.L. and Kosson, D.S., 2002. Distribution sequestration and release of PAHs in weathered sediment: the role of sediment structure and organic carbon properties. *Environmental Science and Technology*, 36, 2636-2644.
- Rogge, W.F., Hildemann, L.M., Mazurek, M.A., Cass, G.R. and Simoneit, B.R.T., 1993. Sources of fine organic aerosol. 2. Noncatalyst and catalyst-equipped automobiles and heavy-duty diesel trucks. *Environmental Science and Technology*, 27, 636-651.
- Safari, M., Zakaria, M.P., Mohamed, C.A.R., Lajis, N.H., Chandru, K., Bahry, P.S. and Anita, S., 2010. The history of petroleum pollution in Malaysia; urgent need for integrated prevention approach. *Environment Asia*, 3, 131-142.

- Saha, M., Togo, A., Mizukawa, K., Murakami, M., Takada, H., Zakaria, M.P., Chiem, N.H., Tuyen, B.C., Prudente, M. and Boonyatumanond, R., 2009. Sources of sedimentary PAHs in tropical Asian waters: Differentiation between pyrogenic and petrogenic sources by alkyl homolog abundance. *Marine Pollution Bulletin*, 58, 189-200.
- Sakari, M., Zakaria, M.P., Junos, M.B.M., Annuar, N.A., Yun, H.Y., Heng, Y.S., Syed Zainuddin, S.M.H. and Chai, K.L., 2008a. Spatial distribution of petroleum hydrocarbon in sediments of major rivers from east coast of peninsular Malaysia. *Estuarine and Coastal Marine Science*, 31(1), 9-18.
- Sakari, M., Zakaria, M.P., Lajis, N.H., Mohamed, C.A.R., Bahry, P.S. and Anita, S., 2008b. Characterization, distribution, sources and origins of aliphatic hydrocarbons from surface sediment of Prai Strait, Penang, Malaysia: A widespread anthropogenic input. *Environment Asia*, 2, 1-14.
- Sakari, M., Zakaria, M.P., Mohamed, C.A.R., Lajis, N.H., Chandru, K., Bahry, P.S., Mokhtar, M.B. and Shahbazi, A., 2010. Urban vs. Marine based oil pollution in the strait of Johor, Malaysia: a century record. *Soil Sediment Contam*, 19, 644-666.
- Sakari, M. and Zakaria, M.P., Mohamed, C.A.R., Lajis, N.H., Abdullah, M.H., Shahbazi, A., 2011. Polycyclic Aromatic Hydrocarbons and Hopane in Malacca Coastal Water: 130 Years of Evidence for Their Land-Based Sources. *Environmental Forensics*, 12, 63-78.
- Sakari, M. and Zakaria, M.P., 2013. Distribution, Characterization and Origins of Polycyclic Aromatic Hydrocarbons (PAHs) in Surficial Sediment of Penang, Malaysia: The Presence of Fresh and Toxic Substances. *World Applied Sciences Journal*, 23.
- Sanger, D.M., Holland, A.F. and Scott, G.I., 1999. Tidal creek and salt marsh sediments in South Carolina coastal estuaries: II. Distribution of organic contaminants. *Archives of Environmental Contamination and Toxicology*, 37, 458-71.
- Savinov, V.M., Savinova, T.N., Matishov, G.G., Dahle, S. and Næs, K., 2003. Polycyclic aromatic hydrocarbons (PAHs) and organochlorines (OCs) in bottom sediments of the Guba Pechenga, Barents Sea, Russia. *Science of The Total Environment*, 306 (1-3), 39-56. doi:10.1016/S0048-9697(02)00483-7.
- Sawidis, T. and Bellos, D., 2005. Chemical pollution monitoring of the River Pinios (Thessalia-Greece). *Journal of Environmental Management*, 76, 282-292.
- Schoellhamer, D.H., William, H.M. and Ashish, J.M., 2000. Influence of salinity, bottom topography, and tides on locations of estuarine turbidity maxima in northern San Francisco Bay. *Proceedings in Marine Science*, Elsevier, 343-357.

- Schumacher, B.A., 2002. Methods for the determination of total organic carbon (TOC) in soils and sediments. *Ecological Risk Assessment Support Center*, 1-23.
- Schwarzenbach, R.P., Gschwend, P.M. and Imboden, D.M., 2003. Environmental Organic Chemistry. (2nd edition). *New York: John Wiley*.
- Shahbazi, A., 2009. *Green-Lipped Mussels (Perna Viridis) as Biomarkers of Petroleum Hydrocarbon Contamination in Selected Coastal Waters of Peninsular Malaysia*. PhD Thesis, Universiti Putra Malaysia.
- Shahbazi, A., Zakaria, M.P., Yap, C.K., Surif, S., Bakhtiari, A.R., Chandru, K., Bahry, P.S. and Sakari, M., 2010a. Spatial distribution and sources of polycyclic aromatic hydrocarbons (PAHs) in green mussels (*Perna viridis*) from coastal areas of Peninsular Malaysia: implications for source identification of perylene. *Int. Journal Environmental Toxicology and Chemistry*, 90, 14-30.
- Shahbazi, A., Zakaria, M.P., Yap, C.K., Tan, S.G., Surif, S., Mohamed, C.A.R., Sakari, M., Bakhtiari, A.R., Bahry, P.S. and Chandru, K., 2010b. Use of different tissues of *Perna viridis* as biomonitors of polycyclic aromatic hydrocarbons (PAHs) in the coastal waters of Peninsular Malaysia. *Environmental Forensics*, 11, 248-263.
- Shaw, M., Tibbetts, I.R. and Mueller, J.F., 2004. Monitoring PAHs in the Brisbane River and Moreton Bay, Australia, using semipermeable membrane devices and EROD activity in yellowfin bream, *Acanthopagrus australis*. *Chemosphere*, 56, 237-246.
- Sherblom, P.M., Gschwend, P.M. and Eganhouse, R.P., 1992. Aqueous solubilities, vapor pressures, and 1-octanol-water partition coefficients for C9-C14 linear alkylbenzenes. *Journal of Chemical & Engineering Data*, 37, 394-399.
- Shi, Z., Tao, S., Pan, B., Fan, W., He, X. C. and Zuo, Q., 2005. Contamination of rivers in Tianjin, China by polycyclic aromatic hydrocarbons. *Environmental Pollution*, 134(1), 97-111. doi:10.1016/j.envpol.2004.07.014.
- Short, J.W., Irvine, G.V., Mann, D.H., Maselko, J.M., Pella, J.J., Lindeberg, M.R., Payne, J.R., Driskell, W.B. and Rice, S.D., 2007. Slightly weathered Exxon Valdez oil persists in gulf of Alaska beach sediments after 16 years. *Environmental Science and Technology*, 41, 1245-1250.
- Silva, T.R., Lopes, S.R.P., Spörl, G., Knoppers, B.A. and Azevedo, D.A., 2012. Source characterization using molecular distribution and stable carbon isotopic composition of n-alkanes in sediment cores from the tropical Mundaú-Manguaba estuarine-lagoon system, Brazil. *Organic Geochemistry*, 53, 25-33.
- Silliman, J.E., Meyers, P.A., Eadie, B.J. and Val Klump, J., 2001. A hypothesis for the origin of perylene based on its low abundance in sediments of Green Bay, Wisconsin. *Chemical Geology*, 177, 309-322.

- Simcik, M.F., Eisenreich, S.J., Golden, K.A., Liu, S.P., Lipiatou, E., Swackhamer, D.L. and Long, D.T., 1996. Atmospheric loading of polycyclic aromatic hydrocarbons to Lake Michigan as recorded in the sediments. *Environmental Science and Technology*, 30, 3039-3046.
- Simoneit, B.R.T., 1982. Some applications of computerized GC-MS to the determination of biogenic and anthropogenic organic matter in the environment. Intern. *Environmental Toxicology and Chemistry*, 12, 177-193.
- Simoneit, B.R.T., Sheng, G., Chen, X., Fu, J., Zhang, H. and Xu, Y., 1991. Molecular marker study of extractable organic matter in aerosols from the urban areas of China. *Atmospheric Environment*, 25A, 2111-2129.
- Simoneit, B.R.T., 2002. Biomass burning—a review of organic tracers for smoke from incomplete combustion. *Applied Geochemistry*, 17(3), 129-162.
- Simpson, C.D., Harrington, C.F., Cullen, W.R., Bright, D.A. and Reimer, K.J., 1998. Polycyclic aromatic hydrocarbon contamination in marine sediments near Kitimat, British Columbia. *Environmental Science & Technology*, 32, 3266-3272.
- Singh, S.N., Kumari, B. and Mishra, S., 2012. Microbial degradation of alkanes. In *Microbial Degradation of Xenobiotics*, Springer, 439-469.
- Skupinska, K., Misiewicz, I. and Kasprzycka-Guttman, T., 2004. Polycyclic aromatic hydrocarbons: physiochemical properties, environmental appearance and impact on living organisms. *Acta Poloniae Pharmaceutica*, 61(3), 233-40.
- Soclo, H.H., Garrigues, P. and Ewald, M., 2000. Origin of Polycyclic Aromatic Hydrocarbons (PAHs) in Coastal Marine Sediments: Case Studies in Cotonou (Benin) and Aquitaine (France) Areas. *Marine Pollution Bulletin*, 40 (5), 387-396.
- Solomons, Graham T.W., 1996. Organic chemistry. 6th ed. New York : John Wiley and Sons, xxvii, 121. ISBN 0-471- 01342-0.
- Sosrowidjojo, I.B., Alexander, R. and Kagi, P., 1994. The biomarker composition of some crude oils from Sumatra. *Organic Geochemistry*, 21, 303-312.
- Sprovieri, M., Feo, M.L., Prevedello, L., Manta, D.S., Sammartino, S. and Tamburrino, S., 2007. Heavy metals, polycyclic aromatic hydrocarbons and polychlorinated biphenyls in surface sediments of the Naples harbour (southern Italy). *Chemosphere*, 67, 998-1009.
- Srogi, K., 2007. Human exposure to polycyclic aromatic hydrocarbons (PAHs) from coke plant. *Air Protection Waste Problem*, 2, 45-51.
- Suhaila, J., Deni, S.M., Zin, WZ.W. and Jemain, A.A., 2010. Trends in Peninsular Malaysia rainfall data during the southwest monsoon and northeast monsoon seasons: 1975-2004. *Sains Malaysia*, 39, 533-542.

- Sullivan, J.B., Krieger and G.R., eds., 2001. Clinical environmental health and toxic exposures. 2nd ed. Philadelphia, PA: *Lippincott Williams and Wilkins*, p. 1241.
- Sun, J. H., Wang, G. L., Chai, Y., Zhang, G., Li, J. and Feng, J., 2009. Distribution of polycyclic aromatic hydrocarbons (PAHs) in Henan Reach of the Yellow River, Middle China. *Ecotoxicology and Environmental Safety*, 72(5), 1614-1624. <https://doi.org/10.1016/j.ecoenv.2008.05.010>.
- Stout, S.A. and Uhler, A.D., 2003. Distinguishing “background” hydrocarbons from contamination using chemical fingerprinting. *Environmental Claims Journal*, 15(2), 241-259.
- Stout, S.A., Uhler, A.D. and Emsbo-Mattingly, S.D., 2004. Comparative evaluation of background anthropogenic hydrocarbons in surficial sediments from nine urban waterway. *Environmental Science and Technology*, 38 (11), 2987-2994.
- Tahir, N.M., Fadzil, M.F., Ariffin, J., Maarop, H. and Wood, A.K.H.J., 2011. Sources of polycyclic aromatic hydrocarbons in mangrove sediments of pulau cik wan dagang, kemaman. *Journal of Sustainability and Science Management*, 6(1), 98–106.
- Takada, H. and Ishiwatari, R., 1990. Biodegradation experiments of linear alkylbenzenes (LABs): isomeric composition of C12 LABs as an indicator of the degree of LAB degradation in the aquatic environment. *Environmental Science and Technology*, 24(1), 86-91.
- Takada, H., Onda, T. and Ogura, N., 1990. Determination of polycyclic aromatic hydrocarbons in urban street dusts and their source materials by capillary gas chromatography. *Environmental Science and Technology*, 24, 1179-1186.
- Takada, H., Onda, T., Harada, M. and Ogura, N., 1991. Distribution and sources of polycyclic aromatic hydrocarbons (PAHs) in street dust from the Tokyo Metropolitan area. *Science of the Total Environment*, 107, 45-69.
- Takada, H., Ishiwatari, R. and Ogura, N., 1992. Distribution of linear alkylbenzenes (LABs) and linear alkylbenzenesulphonates (LAS) in Tokyo Bay sediments. *Estuarine Coastal and Shelf Science*, 35 (2), 141-156.
- Takada, H., Farrington, J.W., Bothner, M.H., Johnson, C.G. and Tripp, B.W., 1994. Transport of sludge-derived organic pollutants to deep-sea sediments at Deep Water Dump Site 106. *Environmental Science and Technology*, 28, 1062-1072.
- Takada, H., Satoh, F., Bothner, M.H., Tripp, B.W., Johnson, C.G. and Farrington, J.W., 1997. Anthropogenic Molecular Markers: Tools To Identify the Sources and Transport Pathways of Pollutants. *American Chemical Society*, 671, 178-195.

- Takada, H. and Eganhouse, R.P., 1998. Molecular markers of anthropogenic waste. In: Meyers, R.A. (Ed.), *Encyclopedia of Environmental Analysis and Remediation*. Wiley, New York, 2883-2940.
- Tam, N.F.Y., Ke, L., Wang, X.H. and Wong, Y.S., 2001. Contamination of polycyclic aromatic hydrocarbons in surface sediments of mangrove swamps. *Environmental Pollution*, 114 (2), 255-263.
- Tanner, B.R., Uhle, M.E., Mora, C.I., Kelley, J.T., Schuneman, P.J., Lane, C.S. and Allen, E.S., 2010. Comparison of bulk and compound-specific $\delta^{13}\text{C}$ analyses and determination of carbon sources to salt marsh sediments using n-alkane distributions (Maine, USA). *Estuarine Coastal and Shelf Science*, 86, 283-291.
- Thia-Eng, C., Gorre, I.R.L., Adrian Ross, S., Bernad, S.R., Gervacio, B. and Corazon Ebarvia, M., 2000. The Malacca Straits. *Marine Pollution Bulletin*, 41 (1-6), 160-178.
- Tsymbalyuk, K.K., Den'ga, Y.M., Berlinsky, N.A. and Antonovich, V.P., 2011. Determination of 16 priority polycyclic aromatic hydrocarbons in bottom sediments of the Danube estuarine coast by GC/MS, *Geo-Eco-Marina* 17/2011, 67-72.
- Tiwari, M., Sahu, S., Bhangare, R., Ajmal, P. and Pandit, G., 2012. Estimation of polycyclic aromatic hydrocarbons associated with size segregated combustion aerosols generated from household fuels. *Microchemical Journal*.
- Tolosa, I., De Mora, S., Sheikholeslami, M.R., Villeneuve, J.P., Bartocci, J. and Cattini, C., 2004. Aliphatic and aromatic hydrocarbons in coastal Caspian Sea sediments. *Marine Pollution Bulletin*, 48(1), 44-60.
- Tran, K., Yu, C.C. and Zeng, E.Y., 1997. Organic pollutants in the coastal environment off San Diego, California. 2. Petrogenic and biogenic sources of aliphatic hydrocarbons. *Environmental Toxicology and Chemistry*, 16, 189-195.
- Tsai, P., Hoenicke, R., Yee, D., Bamford, H. A. and Baker, J. E., 2002. Atmospheric concentrations and fluxes of organic compounds in the northern San Francisco Estuary. *Environmental Science and Technology*, 36, 4741-4747.
- Tsutsumi, S., Yamaguchi, Y., Nishida, I., Akiyama, K.-I., Zakaria, M.P. and Takada, H., 2002. Alkylbenzenes in mussels from South and South East Asian coasts as a molecular tool to assess sewage impact. *Marine Pollution Bulletin*, 45, 325-331.
- USEPA. 1993. Provisional Guidance for Quantitative Risk Assessment of Polycyclic Aromatic Hydrocarbons, EPA/600/R-93/089. Washington, DC: Office of Research and Development, *US Environment Protection Agency*.
- USEPA. 2003. Integrated Risk Information System (IRIS). US Environmental Protection Agency. <<http://www.epa.gov/iris/>> (10.07.12).

- USEPA. 2007. Method 8270D. Semivolatile Organic Compound by Gas Chromatography/Mass Spectrometry (GCMS) Revision 4, February 2007.
- Vaezzadeh, V., Zakaria, M.P., Mustafa, S., Ibrahim, Z.Z., Shau-Hwai, A.T., Keshavarzifard, M., Magam, S.M. and Masood, N., 2014. Distribution of Polycyclic Aromatic Hydrocarbons (PAHs) in Sediment from Muar River and Pulau Merambong, Peninsular Malaysia. A.Z. Aris et al. (eds.), *From Sources to Solution*, 451-455.
- Vaezzadeh, V., Zakaria, M.P., Mustafa, S., Ibrahim, Z.Z., Shau-Hwai, A.T., Keshavarzifard, M., Magam, S.M. and Alkhadher, S.A.A., 2015a. Source type evaluation of polycyclic aromatic hydrocarbons (PAHs) in surface sediments from the Muar River and Pulau Merambong, Peninsular Malaysia. *Environmental Forensic*, 16, 135-142.
- Vaezzadeh, V., Zakaria, M.P., Shau-Hwai, A.T., Ibrahim, Z.Z., Mustafa, S., Jahromi, F. A., Masood, N., Magam, S.M. and Alkhadher, S.A.A., 2015b. Forensic investigation of aliphatic hydrocarbons in the sediments from selected mangrove ecosystems in the west coast of Peninsular Malaysia. *Marine Pollution Bulletin*, 100, 311-320.
- Van Brummelen, T.C., Van Gestel, C.A.M. and Verweij, R.A., 1996. Long-Term toxicity of five polycyclic aromatic hydrocarbons for the terrestrial isopods *Oniscus Asellus* and *Porcellio Scaber*. *Environmental Toxicology and Chemistry*, 15, 1199-1210.
- Van Metre, P.C., Mahler, B.J. and Furlong, E.T., 2000. Urban sprawl leaves its PAH signature. *Environmental Science and Technology*, 34, 4064-4070.
- Van Schooten, F.-J., 1996. Coal tar therapy: is it carcinogenic? *Drug Safety*, 6, 374-7.
- Vane, C.H., Harrison, I. and Kim, A.W., 2007. Polycyclic aromatic hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs) in sediments from the Mersey Estuary, U.K. *Science of The Total Environment*, 374, 112-126.
- Veyrand, B., Sirot, V., Durand, S., Pollono, C., Marchand, P. and Dervilly-Pinel, G., 2013. Human dietary exposure to polycyclic aromatic hydrocarbons: results of the second French Total Diet Study. *Environment International*, 54, 11-17.
- Viguri, J., Verde, J. and Irabien, A., 2002. Environmental assessment of polycyclic aromatic hydrocarbons (PAHs) in surface sediments of the Santander Bay, Northern Spain. *Chemosphere*, 48, 157-165. doi:10.1016/S0045-6535(02)00105-4.
- Viñas, L., Angeles Franco, M., Antonio Soriano, J., José González, J., Pon, J. and Albaigés, J., 2010. "Sources and distribution of polycyclic aromatic hydrocarbons in sediments from the Spanish northern continental shelf. Assessment of spatial and temporal trends. *Environmental Pollution*, 158 (5), 1551-1560.

- Volkman, J.K., Holdsworth, D.G., Neill, G.P. and Bavor Jr, H.J., 1992. Identification of natural, anthropogenic and petroleum hydrocarbons in aquatic sediments. *Science of the Total Environment*, 112(2), 203-219.
- Volkman, J.K., Revill, A.T., Murray, A.P., 1997. Applications of biomarkers for identifying sources of natural and pollutant hydrocarbons in aquatic environments. In: ACS Symposium Series, 1997. Washington, DC: American Chemical Society, 110-132.
- Wagrowski, D.M. and Hites, R.A., 1997. Polycyclic aromatic hydrocarbon accumulation in urban, suburban, and rural vegetation. *Environmental Science and Technology*, 3, 279-282.
- Wakeham, S.G., 1996. Aliphatic and polycyclic aromatic hydrocarbons in Black Sea sediments. *Marine Chemistry*, 53, 187-205.
- Wang, Z., Fingas, M. and Sergy, G., 1994. Study of 22-year-old Arrow oil samples using biomarker compounds by GC/MS. *Environmental Science and Technology*, 28, 1733-1746.
- Wang, Z. and Fingas, M., 1999. Identification of the Source(s) of Unknown Spilled Oils. *Environment Canada Arctic and Marine Oil Spill Program Technical Seminar (AMOP) Proceedings*, Volume: 22. DOI: 10.7901/2169-3358-1999-1-211.
- Wang, X.-C., Zhang, Y.-X. and Chen, R.F., 2001. Distribution and partitioning of polycyclic aromatic hydrocarbons (PAHs) in different size fractions in sediments from Boston Harbor, United States. *Marine Pollution Bulletin*, 42, 1139-1149.
- Wang, Y., Field, R.D. and Roswintarti, O., 2004. Trends in atmospheric haze induced by peat fires in Sumatra Island, Indonesia and El Niño phenomenon from 1973 to 2003. *Geophysical Research Letters*, 31 (4), L04103.
- Wang, Z., Stout, S.A. and Fingas, M., 2006. Forensic fingerprinting of biomarkers for oil spill characterization and source identification. *Environmental Forensics*, 7(2), 105-146.
- Wang, Z. and Stout, S.A., 2007. Oil Spill Environmental Forensics. Fingerprint and Source Identification. *Academic Press*, Waltham. ISBN: 9780123695239.
- Wang, Z., Yang, C., Fingas, M., Hollebone, B., Yim, U.H. and Oh, J.R., 2007. Petroleum biomarker fingerprinting for oil spill characterization and source identification. In *Oil Spill Environmental Forensics*, ed. Wang, Z. and Stout, S.A. London: Academic Press, 73-14.
- Wang, Z. and Brown, C., 2009. Chemical fingerprinting of petroleum hydrocarbons. In: Mudge SM (ed) *Methods in environmental forensics*. CRC Press, Boca Raton, FL, 43-112.

- Wang, H.-S., Liang, P., Kang, Y., Shao, D.-D., Zheng, G.J., Wu, S.-C., Wong, C.K.C. and Wong, M.H., 2010. Enrichment of polycyclic aromatic hydrocarbons (PAHs) in mariculture sediments of Hong Kong. *Environmental Pollution*, 158, 3298-3308.
- Wang, C., Sun, H., Chang, Y., Song, Z. and Qin, X., 2011. Polycyclic aromatic hydrocarbons distribution in sediments associated with gas hydrate and oil seepage from the Gulf of Mexico. *Marine Pollution Bulletin*, 62, 2714-2723.
- Wang, Y., Li, X., Li, B.H., Shen, Z.Y., Feng, C.H. and Chen, Y.X., 2012. Characterization, sources, and potential risk assessment of PAHs in surface sediments from nearshore and farther shore zones of the Yangtze estuary, China. *Environmental Science Pollution Research*, 19, 4148-4158.
- Wang, J.-Z., Zhang, K. and Liang, B., 2012c. Tracing urban sewage pollution in Chaohu Lake (China) using linear alkylbenzenes (LABs) as a molecular marker. *Science of Total Environment*, 414, 356-363.
- Wang, C., Li, F., Shi, H., Jin, Z., Sun, X., Zhang, F., Wu, F. and Kan, S., 2013. The significant role of inorganic matters in preservation and stability of soil organic carbon in the Baoji and Luochuan loess/paleosol profiles, Central China. *CATENA*, 109, 186-194.
- Wang, J.-Z., Chen, T.-H., Zhu, C.-Z. and Peng, S.-C., 2014. Trace organic pollutants in sediments from Huaihe River, China: Evaluation of sources and ecological risk. *Journal of Hydrology*, 512, 463-469.
- Waples, D.W. and Machihara, T., 1991. Biomarkers for geologists. Tulsa, Oklahoma: *American Association of Petroleum Geologists*, 91-99.
- Wei, G.L., Liu, L.Y., Bao, L.J. and Zeng, E.Y., 2014a. Tracking anthropogenic influences on the continental shelf of China with sedimentary linear alkylbenzenes (LABs). *Marine Pollution Bulletin*, 80: 80-87.
- Wei, G.L., Bao, L.J., Guo, L.C., He, Z.C., Wu, F.C. and Zeng, E.Y., 2014b. Utility of soil linear alkylbenzenes to assess regional anthropogenic influences with special reference to atmospheric transport. *Science of Total Environment*, 487, 528-536. doi:10.1016/j.scitotenv.2014.04.067.
- Weinstein, J.E., Crawford, K.D., Garner, T.R. and Flemming, A.J., 2010. Screening-level ecological and human health risk assessment of polycyclic aromatic hydrocarbons in stormwater detention pond sediments of Coastal South Carolina, USA. *Journal of Hazardous Materials*, 178(1-3), 906-916. doi:10.1016/j.jhazmat.2010.02.024.
- Westerholm, R., Almén, J., Li, H., Rannug, U. and Rosén, Å., 1992. Exhaust emissions from gasoline-fuelled light duty vehicles operated in different driving conditions: a chemical and biological characterization. *Atmospheric Environment*, 26B(1), 79-90.

- Wilcke, W., Amelung, W., Martius, C., Garcia, M.V.B. and Zech, W., 2000. Biological sources of polycyclic aromatic hydrocarbons (PAHs) in the Amazonian rain forest. *Journal of Plant Nutrition and Soil Science*, 163(1), 27-30.
- Wild, S.R., and Jones, K.C., 1995. Polynuclear aromatic hydrocarbons in the United Kingdom environment: a preliminary source inventory and budget. *Environmental Pollution*, 88(1), 91-108.
- Witt, G., 1995. Polycyclic aromatic hydrocarbons in water and sediment of the Baltic Sea. *Marine Pollution Bulletin*, 31, 237-248.
- Witt, G. and Trost, E., 1999. Polycyclic aromatic hydrocarbons (PAHs) in sediments of the Baltic Sea and of the German coastal waters. *Chemosphere*, 38(7), 1603-1614.
- Woodhead, R.J., Law, R.J. and Matthiessen, P., 1999. Polycyclic aromatic hydrocarbons in surface sediments around England and Wales, and their possible biological significance. *Marine Pollution Bulletin*, 38(9): 773-790.
- Xu, J., Yu, Y., Wang, P., Guo, W., Dai, S. and Sun, H., 2007. Polycyclic aromatic hydrocarbons in the surface sediments from Yellow River, China. *Chemosphere*, 67: 1408-1414.
- Yang, H.-H. and Chen, C.-M., 2004. Emission inventory and sources of polycyclic aromatic hydrocarbons in the atmosphere at a suburban area in Taiwan. *Chemosphere*, 56, 879-887.
- Yang, H.-H., Jung, R.-C., Wang, Y.-F. and Hsieh, L.-T., 2005. Polycyclic aromatic hydrocarbons emission from joss paper furnaces. *Atmospheric Environment*, 39, 3305-3312.
- Yang, Y. and T. Hofmann., 2009. Aqueous accelerated solvent extraction of native polycyclic aromatic hydrocarbons (PAHs) from carbonaceous river floodplain soils. *Environmental Pollution*, 157(10), 2604-2609.
- Yang, H.-H., Chien, S.-M., Lo, M.-Y., Lan, J.C.-W., Lu, W.-C. and Ku, Y.-Y., 2007a. Effects of biodiesel on emissions of regulated air pollutants and polycyclic aromatic hydrocarbons under engine durability testing. *Atmospheric Environment*, 41, 7232-7240.
- Yang, C.R., Lin, T.C. and Chang, F.H., 2007b. Particle size distribution and PAH concentrations of incense smoke in a combustion chamber. *Environmental Pollution*, 145(2), 606-615.
- Ye, B., Zhang, Z. and Mao, T., 2006. Pollution sources identification of polycyclic aromatic hydrocarbons of soils in Tianjin area, China. *Chemosphere*, 64:525-34.

- Yim, U.H., Hong, S.H., Shim, W.J., Oh, J.R. and Chang, M., 2005. Spatio-temporal distribution and characteristics of PAHs in sediments from Masan Bay, Korea. *Marine Pollution Bulletin*, 50 (3), 319-326.
- Youngblood, W.W. and Blumer, M., 1975. Polycyclic aromatic hydrocarbons in the environment: homologous series in soils and recent marine sediments. *Geochimica et Cosmochimica Acta*, 39, 1303-1314.
- Yunker, M.B., Macdonald, R.W., Vingarzan, R., Mitchell, R.H., Goyette, D. and Sylvestre, S., 2002. PAHs in the Fraser River basin: a critical appraisal of PAH ratios as indicators of PAH source and composition. *Organic Geochemistry*, 33 (4), 489-515.
- Zaghden, H., Kallel, M., Louati, A., Elleuch, B., Oudot, J. and Saliot, A., 2005. Hydrocarbons in surface sediments from the Sfax coastal zone (Tunisia), Mediterranean Sea. *Marine Pollution Bulletin*, 50 1287-1294.
- Zakaria, M. P., Horinouchi, A. I., Tsutsumi, S., Takada, H., Tanabe, S. and Ismail, A. 2000. Oil pollution in the Straits of Malacca, Malaysia: Application of molecular markers for source identification. *Environmental Science and Technology*, 34(7), 1189-1196.
- Zakaria, M.P., Okuda, T. and Takada, H., 2001. Polycyclic aromatic hydrocarbon (PAHs) and hopanes in stranded tar-balls on the coasts of Peninsular Malaysia: applications of biomarkers for identifying sources of oil pollution. *Marine Pollution Bulletin*, 42, 1357-1366.
- Zakaria, M. P., Takada, H., Tsutsumi, S., Ohno, K., Yamada, J., Kouno, E. and Kumata, H., 2002. Distribution of polycyclic aromatic hydrocarbons (PAHs) in rivers and estuaries in Malaysia: a widespread input of petrogenic PAHs. *Environmental Science & Technology*, 36(9), 1907-1918.
- Zakaria, M.P., Geik, K.H., Lee, W.Y. and Hayet, R., 2005. Landfill leachate as a source of polycyclic aromatic hydrocarbons (PAHs) to Malaysian waters. *Estuarine and Coastal Marine Science*, 29, 116-123.
- Zakaria, M.P. and Mahat, A.A., 2006. Distribution of polycyclic aromatic hydrocarbon (PAHs) in sediments in the Langat Estuary. *Coastal Marine Science*, 30, 387-395.
- Zemo, D.A., 2009. Use of parent polycyclic aromatic hydrocarbon (PAH) proportions to attribute PAH sources in sediments: a case study from the Pacific Northwest. *Environment Forensics*, 10(3), 229-239.
- Zeng, E.Y. and Vista, C.L., 1997. Organic pollutants in the coastal environment off San Diego, California. 1. Source identification and assessment by compositional indices of polycyclic aromatic hydrocarbons. *Environmental Toxicology and Chemistry*, 16, 179-188.

- Zhang, J., Chai, L., Yuan, D. and Chen, M., 2004. Distribution and sources of polynuclear aromatic hydrocarbons in mangrove superficial sediments of Deep bay, China. *Marine Pollution Bulletin*, 49, 479–486.
- Zhang, H., Y. Luo, M. Wong, Q. Zhao, and G. Zhang. 2006. Distributions and concentrations of PAHs in Hong Kong soils. *Environmental Pollution*, 141(1), 107-114.
- Zhang, Y. and Tao, S., 2009. Global atmospheric emission inventory of polycyclic aromatic hydrocarbons (PAHs) for 2004. *Atmospheric Environment*, 43, 812-819. doi:10.1016/j.atmosenv.2008.10.050.
- Zhang, K., Wang, J.Z., Liang, B., Shen, R.L. and Zeng, E.Y., 2012. Assessment of aquatic wastewater pollution in a highly industrialized zone with sediment linear alkylbenzenes. *Environmental Toxicology and Chemistry*, 31, 724-730.
- Zhao, H., Yin, C., Chen, M., Wang, W., Chris, J. and Shan, B., 2009. Size distribution and diffuse pollution impacts of PAHs in street dust in urban streams in the Yangtze River Delta. *Journal of Environmental Sciences*, 21(2), 162–167.
- Zhao, X., Ding, J. and You H., 2014. Spatial distribution and temporal trends of polycyclic aromatic hydrocarbons (PAHs) in water and sediment from Songhua River, China. *Environmental Geochemistry and Health*, 36, 131-143. doi:10.1007/s10653-013-9524-0.
- Zheng, G.J., Lam, M.H.W., Lam, P.K.S., Richardson, B.J., Man, B.K.W., Li, A.M.Y., 2000. Concentrations of Persistent Organic Pollutants in Surface Sediments of the Mudflat and Mangroves at Mai Po Marshes Nature Reserve, Hong Kong. *Marine Pollution Bulletin*, 40 (12), 1210-1214.
- Zheng, G.J., Man, B.K.W., Lam, J.C.W., Lam, M.H.W. and Lam, P.K.S., 2002. Distribution and sources of polycyclic aromatic hydrocarbons in the sediments of a sub-tropical coastal wetland. *Water Research*, 36 (6), 1457-1468.
- Zhou, J.L. and Maskaoui, K., 2003. Distribution of polycyclic aromatic hydrocarbons in water and surface sediments from Daya Bay, China. *Environmental Pollution*, 121, 269-281.
- Zhu Y., Liu H., Cheng H., Xi Z., Liu X. and Xu X., 2005. The distribution and source apportionment of aliphatic hydrocarbons in soils from the outskirts of Beijing. *Organic Geochemistry*, 36, 475-483.
- Zielinska, B., Sagebiel, J., Arnott, W.P., Rogers, C.F., Kelly, K.E., Wagner, D.A., Lighty, J.S., Sarofim, A.F. and Palmer, G., 2004. Phase and size distribution of polycyclic aromatic hydrocarbons in diesel and gasoline vehicle emissions. *Environmental Science and Technology*, 38, 2557-2567.
- Zmirou, D., Masdet, P., Boudet, C., Dor, F. and Dechenaux, J., 2000. Personal exposure to atmospheric polycyclic aromatic hydrocarbons in a general adult

population and lung cancer risk assessment. *Journal of Occupational Environmental Medicine*, 42, 121-126.

Zuydam, C.S.V., 2007. Determination of polycyclic aromatic hydrocarbons (PAHs) resulting from wood storage and wood treatment facilities for electricity transmission in Swaziland, M.Sc. *Thesis, University of South Africa*, 112P.

