



UNIVERSITI PUTRA MALAYSIA

***MICROBIAL QUALITY AND ANTIOXIDANT ACTIVITY OF AN
IMPROVED BITTER GOURD (*Momordica charantia* L.) DRINK***

CHERN PEI ERN

FSTM 2018 15



**MICROBIAL QUALITY AND ANTIOXIDANT ACTIVITY OF AN
IMPROVED BITTER GOURD (*Momordica charantia* L.) DRINK**

By

CHERN PEI ERN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

September 2017

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**MICROBIAL QUALITY AND ANTIOXIDANT ACTIVITY OF AN
IMPROVED BITTER GOURD (*Momordica charantia* L.) DRINK**

By

CHERN PEI ERN

September 2017

Chair: Nor Ainy Mahyudin, Associate Professor, PhD
Faculty: Food Science and Technology

The bitter gourd (*Momordica charantia* L.) (MC) is being consumed in various countries due to its nutritional value and is high in antioxidants. In this study, the (MC) is used in the preparation of a ready-to-serve (RTS) drink with antioxidant properties. The objectives of this research are to determine the physicochemical and antioxidant properties of a RTS MC drink with different concentrations of the MC, to conduct a sensory evaluation on the MC drink for acceptability and finally to determine the effect of water quality of the MC drink stored at different temperatures. The MC drink was prepared by water extraction of its juice, added with dissolved brown sugar, artificial honey, low fat yogurt, lime juice and mint extract. A proximate analysis was carried out followed by DPPH radical scavenging assay. A sensory evaluation was then carried out among 40 untrained panelists on a Hedonic scale of 1 to 7. The microbial quality was determined using the aerobic plate count, enumeration of moulds and yeasts, presumptive test for coliforms and detection of *Escherichia coli*. Five concentrations (0.58, 0.70, 0.80, 0.90 and 1.00 g/mL) of the MC drink was developed. The proximate analyses carried out showed that the RTS drink consist of high moisture content from 92.94 – 94.65%. The pH and °Brix of the MC drink increased as the concentration increased, whereby the pH was 4.30 – 4.54, and the °Brix was 5.0 – 5.6. The increase in concentration of MC has led to an increase in the radical scavenging activity. At the original concentration of 0.58 g/mL, the inhibition was at 71.51% which increased to 87.22% at the final concentration of 1.00 g/mL. The most accepted concentration of the MC drink according to the sensory evaluation was 0.80 g/mL with a mean score of 4.09 (like slightly), while the least accepted concentration was 1.00 g/mL and obtained the lowest mean score of 3.80 (dislike slightly). At the selected concentration (0.80 g/mL), the amount of bacterial growth exceeded 6 log CFU/mL after 24 hours for the MC drink stored at room temperature (28±2°C). When stored at refrigerated

temperature ($4\pm 2^{\circ}\text{C}$) it took 72 hours to reach 6 log CFU/mL. The total coliform count for all juice samples were between 93-210 MPN/g. *E. coli* was detected in the MC drink but not in the water used for its preparation. This study demonstrated MC has the potential to be used as an ingredient in preparing RTS drinks containing antioxidant properties.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**KUALITI MIKROB DAN KEGIATAN ANTIOKSIDAN
MINUMAN PERIA (*Momordica charantia* L.) SEDIA DIHIDANG YANG
DITAMBAH BAIK**

Oleh

CHERN PEI ERN

September 2017

**Pengerusi: Nor Ainy Mahyudin, Profesor Madya, PhD
Fakulti: Sains dan Teknologi Makanan**

Buah peria (*Momordica charantia* L.) (MC) sering dihidang dalam pemakanan di banyak negara disebabkan nilai nutrien dan kandungan antioksidan yang tinggi. Dalam kajian ini, MC telah digunakan dalam penyediaan minuman sedia dihidang (RTS) dengan sifat-sifat antioksidan. Objektif penyelidikan ini adalah untuk menilai sifat fizikokimia dan antioksidan RTS MC yang berkepekatan berbeza, untuk membuat penilaian deria pada minuman MC untuk penerimaan pengguna, dan akhirnya untuk menentukan kesan kualiti air terhadap minuman MC yang disimpan pada suhu yang berbeza. Minuman MC telah disediakan melalui pengekstrakan air jus, ditambah dengan gula perang, madu tiruan, yogurt lemak rendah, jus limau dan ekstrak pudina. Analisis proksimat telah dijalankan dan seterusnya pemeriksaan aktiviti antioksidan. Satu penilaian deria telah dijalankan oleh 40 ahli panel tidak terlatih menggunakan skala Hedonik 1 hingga 7. Kiraan plat aerobik, penghitungan kulat dan yis, ujian penandaian koliform dan pengesanan *Escherichia coli* dijalankan untuk penentuan kualiti mikrob. Lima kepekatan (0.58, 0.70, 0.80, 0.90 dan 1.00 g/mL) minuman MC telah diwujudkan. Analisis proksimat yang dijalankan menunjukkan bahawa minuman ini terdiri daripada kandungan air yang tinggi iaitu 92.94 – 94.65%. pH dan Brix minuman semakin meningkat mengikut kepekatan, di mana pH adalah 4.30 – 4.54, dan Brix adalah 5.0 – 5.6. Peningkatan kepekatan MC telah meningkatkan kegiatan antioksidan. Pada kepekatan asal 0.58 g/mL perencatannya adalah 71.51% yang meningkat kepada 87.22% pada kepekatan akhir 1.00 g / mL. Mengikut keputusan penilaian deria, kepekatan paling diterima adalah 0.80 g/mL dengan skor purata 4.09 (menyukai sedikit), manakala kepekatan yang paling kurang diterima adalah 1.00 g/mL dan mendapat purata mata terendah sebanyak 3.80 (tidak menyukai sedikit). Pada kepekatan yang dipilih (0.80 g/mL), jumlah pertumbuhan mikrob telah melebihi 6 log CFU/mL selepas 24 jam untuk minuman MC yang disimpan pada suhu bilik (28±2°C). Jumlah pertumbuhan mikrob dalam minuman MC yang disimpan pada suhu sejuk (4±2°C) melebihi 6 log CFU/mL selepas 72 jam. Jumlah kiraan koliform bagi semua sampel jus adalah 93-210 MPN/g. *E. coli* boleh didapati di minuman RTS tetapi tidak

wujud dalam air yang digunakan untuk penyediaan minuman. Kajian ini menunjukkan bahawa MC mempunyai potensi untuk digunakan sebagai ramuan dalam menyediakan minuman RTS yang mengandungi antioksidan.



ACKNOWLEDGEMENTS

First and foremost, I would like to offer my utmost gratitude to the chairman of my supervisory committee, Dr. Nor Ainy Mahyudin, who offered advice and guidance throughout the process of completing this study. Also not forgetting my co-supervisors Dr. Hazrina Ghazali and Dr Nor Khaizura Mahmud @ Ab Rashid for their advices and encouragement.

I would also like to thank all the laboratory assistants from the Biochemical Laboratory and the Microbiology Laboratory for their professional support. Thanks also to all supporting staffs from the department of Postgraduates, Research and Innovation.

Besides, I would like to sincerely thank my colleagues Faez Nur Syafiqah, Nur Amira, Selvaneswary, Zuraidah, Salahaldin and Shu'aibu for their help and support. It was a great time working with all of you and it had made our time in the laboratory more enjoyable. Not forgetting all other UPM friends that offered encouragement.

Special thanks to Putra Grant - Putra Graduate Initiative (IPS) and Graduate Research Fellowship (GRF) for providing financial support.

Last but not least, I would want to thank my family for their unending support throughout my studies in university.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Nor Ainy Mahyudin, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Hazrina Ghazali, PhD

Senior Lecturer
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Nor Khaizura Mahmud @ Ab Rashid, PhD

Senior Lecturer
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Chern Pei Ern (GS42180)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of Chairman of
Supervisory
Committee:



Nor Ainy Mahyudin

Signature: _____

Name of Member of
Supervisory
Committee:

Hazrina Ghazali

Signature: _____

Name of Member of
Supervisory
Committee:

Nor Khaizura Mahmud @ Ab Rashid

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
 CHAPTER	
1 INTRODUCTION	1
1.1 Background of Study	1
1.2 Problem Statement	2
1.3 Objectives of Study	3
1.4 Hypotheses	3
1.5 Significance of Study	3
 2 LITERATURE REVIEW	5
2.1 Ready-to-Serve Drink	5
2.2 Biology of the Bitter Gourd (<i>Momordica charantia</i> L.)	5
2.2.1 Origin and Distribution of the Bitter Gourd	6
2.2.2 Medicinal and Culinary Uses of Bitter Gourd	6
2.2.3 Antimicrobial Activity of Plant Extracts and Bitter Gourd	7
2.2.3.1 Antimicrobial Activity of Plant Extracts	7
2.2.3.2 Antimicrobial Activity of Bitter Gourd	8
2.2.4 Antioxidant Activity of Plant Extracts and Bitter Gourd	9
2.2.4.1 Antioxidant Activity of Plant Extracts	9
2.2.4.2 Antioxidant Activity of Bitter Gourd	9
2.2.5 Other Bioactivities of Bitter Gourd	10
2.2.5.1 Antidiabetic Activity of Bitter Gourd	10
2.2.5.2 Anticancer Activity of Bitter Gourd	11
2.3 Ready-to-Serve Drink using Bitter Gourd	11
2.4 Increasing Palatability and Acceptability of the Bitter Gourd	12
2.4.1 Debittering of the Bitter Gourd	12
2.4.2 Sensory Evaluation of Bitter-Tasting Foods	13
2.5 Microbiological Food Spoilage	13
2.5.1 Yeasts and Moulds Spoilage	14
2.5.2 Microbial Quality of Water	14
2.5.3 Quality of Water for Unpasteurized Juice	15
2.5.4 Control of Microbial Growth Using Low Temperature	16
2.5.5 Storage of Unpasteurized Juices by Vendors	16

3	METHODOLOGY	17
3.1	Determining the Physicochemical and Antioxidant Properties of Ready-to-Serve Bitter Gourd Drink	17
3.1.1	Preparation of the Ready-to-Serve Bitter Gourd Drink	17
3.1.2	Proximate Analysis of Ready-to-Serve Bitter Gourd Drink	19
3.1.2.1	Moisture content	19
3.1.2.2	Ash content	20
3.1.2.3	Crude fat content	20
3.1.2.4	Crude protein content	20
3.1.2.5	Crude fibre content	21
3.1.2.6	Carbohydrate content	21
3.1.2.7	Other analyses	21
3.1.3	DPPH Radical-Scavenging Activity of Ready-to-Serve Bitter Gourd Drink	22
3.2	Sensory Evaluation of Ready-to-Serve Bitter Gourd Drink	22
3.3	Determining the Microbiological Quality of Ready-to-Serve Bitter Gourd Drink with Different Water Quality and Stored at Different Temperatures	22
3.3.1	Aerobic Plate Count	23
3.3.2	Enumeration of Yeasts and Moulds	23
3.3.3	Presumptive Test of Coliforms	23
3.3.4	Detection of <i>Escherichia coli</i>	23
3.4	Statistical Analysis	24
4	RESULTS AND DISCUSSION	25
4.1	To determine the physicochemical and antioxidant properties of ready-to-serve drink using the bitter gourd	25
4.1.1	Proximate Analysis of the Ready-to-Serve Bitter Gourd Drink	25
4.1.2	DPPH Radical-Scavenging Activity of Ready-to-Serve Bitter Gourd Drink	28
4.2	Sensory evaluation of the ready-to-serve bitter gourd (<i>Momordica charantia</i> L.) drink	29
4.3	Determination of the microbial quality of the ready-to-serve bitter gourd (<i>Momordica charantia</i> L.) drink	31
4.3.1	Aerobic Plate Count	31
4.3.2	Enumeration of Moulds and Yeasts	32
4.3.3	Presumptive Test of Coliforms	34
4.3.4	Detection of <i>Escherichia coli</i>	36
4.3.5	Changes in pH	38
5	CONCLUSION	41
5.1	Summary	41
5.2	Limitations	41
5.3	Recommendations	42

REFERENCES	43
APPENDICES	53
BIODATA OF STUDENT	70
PUBLICATIONS	71



LIST OF TABLES

Table	Page
2.1 The scientific classification of the bitter gourd (<i>Momordica charantia</i> L.)	6
3.1 Original bitter gourd (<i>Momordica charantia</i> L.) drink recipe	18
3.2 Amount of each ingredient in percentage for five different concentrations (0.58, 0.70, 0.80, 0.90 and 1.00 g/mL) of MC in the juice recipe	19
4.1 Proximate analysis for moisture, ash, fat, protein, fibre and carbohydrate content of five concentrations (0.58, 0.70, 0.80, 0.90 and 1.00 g/mL) of MC in the ready-to-serve bitter gourd drink	26
4.2 pH and °Brix of MC drink with five concentrations (0.58, 0.70, 0.80, 0.90 and 1.00 g/mL) of MC in the ready-to-serve bitter gourd drink	27
4.3 MPN per gram for ready-to-serve bitter gourd drink prepared using untreated and boiled water stored at 4±2°C and 28±2°C	35
4.4 MPN per gram for plain untreated and boiled water used for the preparation of the ready-to-serve bitter gourd drink	36
4.5 <i>Escherichia coli</i> formation in ready-to-serve bitter gourd drink showing positive (+) and negative (-) formation.	37

LIST OF FIGURES

Figure		Page
4.1	DPPH radical-scavenging activity against the concentration of MC in the ready-to-serve drink	28
4.2	Mean score of sensory evaluation for the ready-to-serve bitter gourd drink with five different concentrations (0.58, 0.70, 0.80, 0.90 and 1.00 g/mL) of MC in the ready-to-serve bitter gourd drink	29
4.3	Mean score of sensory evaluation for the ready-to-serve bitter gourd drink with five concentrations (0.58, 0.70, 0.80, 0.90 and 1.00 g/mL) of MC in the ready-to-serve bitter gourd drink showing standard deviation bars	30
4.4	log CFU/mL for the aerobic plate count for ready-to-serve bitter gourd drink prepared using untreated and boiled water and stored in refrigerated ($4\pm 2^{\circ}\text{C}$) and room temperature ($28\pm 2^{\circ}\text{C}$)	31
4.5	log CFU/mL for the enumeration of moulds and yeasts for ready-to-serve bitter gourd drink prepared using untreated and boiled water and stored in refrigerated ($4\pm 2^{\circ}\text{C}$) and room temperature ($28\pm 2^{\circ}\text{C}$)	33
4.6	Observable formation of mould and yeast growth on the surface of the ready-to-serve bitter gourd drink after a period of storage	34
4.7	Formation of green metallic sheen on EMB agar for the ready-to-serve bitter gourd drink	37
4.8	pH of the ready-to-serve bitter gourd drink prepared using untreated and boiled water and stored in refrigerated ($4\pm 2^{\circ}\text{C}$) and room temperature ($28\pm 2^{\circ}\text{C}$)	39

LIST OF ABBREVIATIONS

°C	Degree in Celcius
μL	Microliter
Abs	Absorbance
AOAC	Association of Official Agricultural Chemists
A _w	Water activity
BAM	Bacteriological Analytical Manual
CFU	Colony-forming units
<i>E. coli</i>	<i>Escherichia coli</i>
g	grams
IC ₅₀	Half maximal inhibitory concentration
L	Liter
LAB	Lactic acid bacteria
LST	Lauryl tryptose/lauryl sulphate broth
L-EMB	Eosin methylene blue, Levine's formulation
MC	<i>Momordica charantia</i> L.
mg	milligrams
mL	milliliter
N	Amount of substance in moles, mol.
RTS	Ready-to-serve
TFTC	Too Few to Count
TNTC	Too Numerous to Count
TSS	Total soluble solids
USFDA	US Food and Drug Administration



CHAPTER 1

INTRODUCTION

1.1 Background of Study

The consumption of natural fruit and vegetable juices has increased in recent years as opposed to caffeine containing beverages like coffee, tea and carbonated soft drinks (Waghray et al., 2012). Juices are the liquid products, purees or concentrates extracted from one or a mixture of fruits or vegetables (Fraternale et al., 2011). The addition of sugars or acids can be permitted but has to be in accordance with standards and regulations (Aneja et al., 2014). As fresh juices can be too acidic or too strongly flavoured, diluting or blending is carried out for a more pleasant consumption (Bagde and Tumane, 2011). Besides providing refreshing satisfaction, functional juices are able to provide necessary nutrients. It is known to have low sodium, cholesterol and fat content, high concentration of vitamin C, polyphenols and antioxidants that are able to prevent nutrient-related diseases such as heart diseases, cancer and diabetes (Aneja et al., 2014).

Momordica charantia L. (MC), which is also known as bitter gourd or bitter melon grows at tropical and subtropical areas of Asia, Africa and the Caribbean (Efird et al., 2014). It is often grown for its consumption as a vegetable, and also used for its medicinal properties in developing countries such as Brazil, China, Colombia, Cuba, Ghana, Haiti, India, Mexico, Malaya, New Zealand, Panama and Peru (Grover and Yadav, 2004). More than 200 different compounds that are potentially possessing medicinal properties are isolated from different parts of the bitter gourd including its fruit, seeds, leaves and stems. Some of the most studied constituents that showed evidence in improving glycemic control are charantin, polypeptide-p, vicine, momordin and other derivatives (Efird et al., 2014). It is also high in phenolic compounds such as gallic acid, gentisic acid (2, 5-dihydroxyl benzoic acid), catechins, chlorogenic acid and epicatechin (Din et al, 2011). Studies for the past few decades showed that MC possess various beneficial properties such as antidiabetic, antiviral, antitumor, antileukemic, antibacterial, anthelmintic, antimutagenic, antimycobacterial, antioxidant, antiulcer, anti-inflammatory, hypocholesterolemic, hypotriglyceridemic, hypotensive, immunostimulant, and insecticidal properties (Grover and Yadav, 2004).

Antioxidants in natural foods are gaining attention due to their function as chemopreventive agents against oxidative damage (Wu and Ng, 2007). The MC is shown to possess potent antioxidant and free radical scavenging activities, which explained the therapeutic benefits suggested by traditional claims (Wu and Ng, 2007). Various parts of MC including the leaf, stem, the green fruit and the ripe fruit are known to have antioxidant (Kubola and Siriamornpun, 2008). The seeds of the MC also shown potent antioxidant activity, and is regarded as directly or indirectly responsible to its hypoglycemic activity (Sathishsekar, and Subramanian, 2005).

There is a growing concern in the microbial safety in raw juices as microbial contamination has become a major health problem. Raw juices are among one of the street foods and are sold at public places and at the roadside (Reddi et al., 2015). The sanitary condition of water used to prepare juices are also a concern. Water sources for washing fruits are mostly obtained from wells and taps, at the same time there are also vendors not washing fruits before juicing and water used for dilution of juices are not boiled (Nonga, Simforian, and Ndabikunze, 2014). These water sources including environmental exposure, improper washing, prolonged preservation without refrigeration and use of unhygienic water may lead to the increase in foodborne illnesses due to pathogens (Tambekar et al., 2009). Juices can become a transmission path for pathogens such as *Escherichia coli* O157, *Salmonella* and *Cryptosporidium*. Juices are also more susceptible to spoilage due to high water content, raw ingredients in contact with air and microorganisms in the environment during handling processes (Aneja et al., 2014).

In this study, the bitter gourd is used to prepare a ready-to-serve drink with minimal amount of processing techniques. The drink was tested for its antioxidant properties as well as the overall proximate analysis to identify its physicochemical properties. Due to the strong taste of the bitter gourd, it is a challenge to improve on the original recipe by increasing the concentration of MC, but is easily accepted by most consumers. Therefore, a sensory evaluation was carried out to determine the acceptability of the ready-to-serve drink. As the ready-to-serve drink only undergo minimal amount of processing, it is succumb to high perishability. Microbiological quality tests were carried out to observe the shelf life of the drink.

1.2 Problem Statement

The bitter gourd (*Momordica charantia* L.) (MC) is consumed in many ways due to its nutritional contents including vitamin A and C, phosphorous and iron (Raman & Lau, 1996). It is developed into drinks and beverages due to its convenience. It is necessary to conduct proximate analyses in order to determine if a certain RTS drink adhere to local laws and regulations before it can be sold in the market. The level of antioxidant activity in the drink has to be determined and in relation to study the influence of MC concentration against the antioxidant activity level.

The MC is a good option for preparing healthy ready-to-serve drinks as it contain antioxidant properties (Kubola and Siriamornpun, 2008) but the taste is not preferred by most people and it needs to be processed to increase palatability (Satkar, Kulthe and Chalke, 2013). The bitter flavour of MC is due to presence of the alkaloid momordicin produced in the fruit and leaves (Thakur and Sharma, 2016; Din et al., 2011). Therefore, there is a need for developing a palatable recipe for the bitter gourd drink. A sensory evaluation of the ready-to-serve drink is necessary in order to determine the most palatable formulation among consumers.

Juices are very susceptible to spoilage and a transmitting vehicle of foodborne pathogens (Aneja et al., 2014) and is also not handled properly by food handlers (Nonga, Simforian, and Ndabikunze, 2014; Tambekar et al., 2009). A study on the microbial shelf life of this drink formulation is needed so that food handlers have the knowledge on how long the drink can be kept after preparation before serving in order to reduce wastage due to spoilage and for it to be safe for consumption.

1.3 Objectives of Study

Based on the stated problems, the following objectives of the study are formed:

1. To determine the physicochemical and antioxidant properties of a ready-to-serve bitter gourd (*Momordica charantia* L.) drink with different concentrations of the bitter gourd (*Momordica charantia* L.)
2. To conduct a sensory evaluation on the ready-to-serve bitter gourd (*Momordica charantia* L.) drink for acceptability
3. To determine the effect of water quality of the ready-to-serve bitter gourd (*Momordica charantia* L.) drink stored at different temperatures against the microbial shelf life.

1.4 Hypotheses

The following hypotheses were developed and will be answered in the study:

1. A higher concentration of the bitter gourd (*Momordica charantia* L.) in the ready-to-serve drink will enhance the level of antioxidant radical scavenging activity
2. The increase in concentration of bitter gourd (*Momordica charantia* L.) in the ready-to-serve drink will reduce its palatability and overall acceptability
3. The microbial quality of the ready-to-serve bitter gourd (*Momordica charantia* L.) drink can be enhanced using boiled water and stored at low storage temperatures

1.5 Significance of Study

An improved ready-to-serve drink using the bitter gourd (MC) with antioxidant properties will be developed from the original recipe (Ghazali et al., 201612270829). This study is able to provide the developed recipe with basic physicochemical information such as moisture, protein, fat, fibre, ash and carbohydrate content. The antioxidant activity inhibition level will be determined using the DPPH radical scavenging assay and the relationship between the concentrations of MC in the RTS drink against the antioxidant activity can be studied. By conducting a sensory evaluation, the most accepted concentration of MC in the drink can be determined and it can provide with a general level of acceptability of the MC taste among consumers. The study on microbial quality can evaluate the current sanitary conditions of water used in the preparation of fresh RTS drinks. The effects of water quality and storage

temperatures of fresh RTS drinks on its microbial safety will also be demonstrated through this study.



REFERENCES

- A.O.A.C. (1990). Official Methods of Analysis. Association of Official Agricultural Chemists, Washington, D. C. 15th Edition. Vol-1, 40-64.
- Ahmad, Z., Zamhuri, K. F., Yaacob, A., Siong, C. H., Selvarajah, M., Ismail, A., & Hakim, M. N. (2012). In vitro anti-diabetic activities and chemical analysis of polypeptide-k and oil isolated from seeds of *Momordica charantia* (bitter gourd). *Molecules*, 17(8), 9631-9640.
- Al-Jedah, J. H., & Robinson, R. K. (2002). Nutritional value and microbiological safety of fresh fruit juices sold through retail outlets in Qatar. *Pakistan Journal of Nutrition*, 1(2), 79-81.
- Allgeyer, L. C., Miller, M. J., & Lee, S. Y. (2010). Sensory and microbiological quality of yogurt drinks with prebiotics and probiotics. *Journal of Dairy Science*, 93(10), 4471-4479.
- Amira, K., Aminah, A., & Zuhair, A. (2013). Evaluation of bitter melon (*Momordica charantia*) extract administration in the antioxidant and free radical scavenging activities of plasma and liver in male rat. *International Food Research Journal*, 20(1).
- Amirthaveni, M., Premakumari, S., Gomathi, K., & Yang, R. Y. (2013). Acceptability of Bitter Gourd (*Momordica charantia* L.) Recipes with Different Cooking Methods. *The Indian Journal of Nutrition and Dietetics*, 50(7), 262-268.
- Aneja, K. R., Dhiman, R., Aggarwal, N. K., & Aneja, A. (2014). Emerging preservation techniques for controlling spoilage and pathogenic microorganisms in fruit juices. *International Journal of Microbiology*, 2014.
- Anilakumar, K. R., Kumar, G. P., & Ilaiyaraja, N. (2015). Nutritional, pharmacological and medicinal properties of *Momordica charantia*. *International Journal of Nutrition and Food Sciences*, 4(1), 73-83.
- Andarwulan, N., Batari, R., Sandrasari, D. A., Bolling, B., & Wijaya, H. (2010). Flavonoid content and antioxidant activity of vegetables from Indonesia. *Food Chemistry*, 121(4), 1231-1235.
- Aziz, M. G., Roy, J., Sarker, M. S. H., & Yusof, Y. A. (2011). Isolation and use of bitter gourd polysaccharide in formulating dietetic soft drinks. *African Journal of Agricultural Research*, 6(23), 5314-5319.
- Azlim Almey, A. A., Ahmed Jalal Khan, C., Syed Zahir, I., Mustapha Suleiman, K., Aisyah, M. R., & Kamarul Rahim, K. (2010). Total phenolic content and

primary antioxidant activity of methanolic and ethanolic extracts of aromatic plants' leaves. *International Food Research Journal*, 17(4).

Bagde, N. I., & Tumane, P. M. (2011). Studies on microbial flora of fruit juices and cold drinks. *Asiatic Journal of Biotechnology Resources*, 2(4), 454-460.

Battelli, M.G., Polito, L., Bolognesi, A., Lafleur, L., Fradet, Y., Stirpe, F. (1996). Toxicity of ribosome-inactivating proteins-containing immunotoxins to a human bladder carcinoma cell line. *International Journal of Cancer* 68, 485–490.

Belitz, H. -D., Grosch, W., Schieberle, P. (2009). Food Chemistry, 4th revised and extended edition. Germany, Springer.

Bharathi, L. K., & John, K. J. (2013). Origin, Domestication and Distribution. In *Momordica genus in Asia-An Overview* (pp. 37-44). Springer India.

Boghani, A. H., Raheem, A., & Hashmi, S. I. (2013). Development and storage studies of blended papaya-aloe vera ready to serve (RTS) beverage. *Journal of Food Processing & Technology*, 2012.

Boshtam, M., Moshtaghian, J., Naderi, G., Asgary, S., & Nayeri, H. (2011). Antioxidant effects of Citrus aurantifolia (Christm) juice and peel extract on LDL oxidation. *Journal of research in medical sciences: the official journal of Isfahan University of Medical Sciences*, 16(7), 951.

Braca, A., Siciliano, T., D'Arrigo, M., & Germanò, M. P. (2008). Chemical composition and antimicrobial activity of *Momordica charantia* seed essential oil. *Fitoterapia*, 79(2), 123-125.

Buck, J. W., Walcott, R. R., & Beuchat, L. R. (2003). Recent trends in microbiological safety of fruits and vegetables. *Plant Health Progress*, 10(1), 1094.

Bull, M. K., Zerdin, K., Howe, E., Goicoechea, D., Paramanandhan, P., Stockman, R.,.... & Stewart, C. M. (2004). The effect of high pressure processing on the microbial, physical and chemical properties of Valencia and Navel orange juice. *Innovative Food Science & Emerging Technologies*, 5(2), 135-149.

Chen, Z., Bertin, R., & Frolidi, G. (2013). EC 50 estimation of antioxidant activity in DPPH assay using several statistical programs. *Food chemistry*, 138(1), 414-420.

Cushnie, T. T., & Lamb, A. J. (2005). Antimicrobial activity of flavonoids. *International journal of antimicrobial agents*, 26(5), 343-356.

Dagher, S., & Ali, A. (1985). Effect of pasteurization, centrifugation and additives on quality of concentrated yogurt (Labneh). *Journal of Food Protection*, 48, 300–302.

- de Boer, E., & Beumer, R. R. (1999). Methodology for detection and typing of foodborne microorganisms. *International Journal of Food Microbiology*, 50(1), 119-130.
- Din, A., Bukhari, S. A. H., Salam, A., & Ishfaq, B. (2011). Development of functional and dietetic beverage from bitter gourd. *Internet Journal of Food Safety*, 13, 355-360.
- Dorman, H. J. D., & Deans, S. G. (2000). Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *Journal of applied microbiology*, 88(2), 308-316.
- Drinking Water Quality Standard. (2010). Drinking Water Quality Surveillance Programme, Engineering Services Division, Ministry of Health Malaysia. Retrieved from: <http://kmam.moh.gov.my/public-user/drinking-water-quality-standard.html>, on November 4, 2016 at 11.15am.
- Edwards, C.G., Haag, K.M., Collins, M.D., Hutson, R.A., Huang, Y.C. (1998). *Lactobacillus kunkeei* sp. nov.: a spoilage organism associated with grape juice fermentations. *Journal of Applied Microbiology*. 84, 698 – 702.
- Food Act 1983 (Act 281) & Regulations (As at 20th April 2012) International Law Book Series.
- Frame, A.D., Rios-Olivares, E., De Jesus, L., Ortiz, D., Pagan, J., Mendez, S. (1998). Plants from Puerto Rico with anti-*Mycobacterium tuberculosis* properties. *Puerto Rico Health Science Journal* 17, 243–252.
- Franz, E., & van Bruggen, A. H. (2008). Ecology of *E. coli* O157: H7 and *Salmonella enterica* in the primary vegetable production chain. *Critical Reviews in Microbiology*, 34(3-4), 143-161.
- Fraternal, D., Ricci, D., Flamini, G., & Giomaro, G. (2011). Volatiles profile of red apple from marche region (Italy). *Records of Natural Products*, 5(3), 202.
- Gabriel, A. A. (2008). Estimation of water activity from pH and Brix values of some food products. *Food Chemistry* 108. 1106–1113.
- Ghazali, H., Abdul Kadir, S. A., Mahpof, S. N., Mohd Shahrin, F. S., Sariman, N. U., Chern, P. E., Zainal Abidin, N. (2016) Patent No.: 201612270829. Selangor, Malaysia: Universiti Putra Malaysia.
- Gil, M. I., Tomás-Barberán, F. A., Hess-Pierce, B., Holcroft, D. M., & Kader, A. A. (2000). Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *Journal of Agricultural and Food chemistry*, 48(10), 4581-4589.

- Gram, L., Ravn, L., Rasch, M., Bruhn, J. B., Christensen, A. B., & Givskov, M. (2002). Food spoilage—interactions between food spoilage bacteria. *International Journal of Food Microbiology*, 78(1), 79-97.
- Greenfield, H., & Southgate, D. A. (2003). Food composition data: production, management, and use. Food & Agriculture Org.
- Grover, J. K., Yadav, S. P. (2004). Pharmacological actions and potential uses of *Momordica charantia*: a review. *Journal of Ethnopharmacology* 93. 123-132.
- Guimarães, R., Barros, L., Barreira, J. C., Sousa, M. J., Carvalho, A. M., & Ferreira, I. C. (2010). Targeting excessive free radicals with peels and juices of citrus fruits: grapefruit, lemon, lime and orange. *Food and Chemical Toxicology*, 48(1), 99-106.
- Hamissou, M., Smith, A. C., Carter Jr, R. E., & Triplett II, J. K. (2013). Antioxidative properties of bitter melon (*Momordica charantia*) and zucchini (*Cucurbita pepo*). *Emirates Journal of Food and Agriculture*, 25(9), 641.
- Hammer, K. A., Carson, C. F., & Riley, T. V. (1999). Antimicrobial activity of essential oils and other plant extracts. *Journal of applied microbiology*, 86(6), 985-990.
- Hernández-Anguiano, A. M., Landa-Salgado, P., Eslava-Campos, C. A., Vargas-Hernández, M., & Patel, J. (2016). Microbiological Quality of Fresh Nopal Juice. *Microorganisms*, 4(4), 46.
- Hilliam, M. (2000). Functional food – How big is the market? *The World of Food Ingredients*, 12. 50-2.
- Horax, R., Hettiarachchy, N., & Islam, S. (2005). Total phenolic contents and phenolic acid constituents in 4 varieties of bitter melons (*Momordica charantia*) and antioxidant activities of their extracts. *Journal of Food Science*, 70(4).
- Institute for Public Health. The Third National Health and Morbidity Survey (NHMS III) 2006: Nutritional status. Kuala Lumpur, Malaysia: Malaysia Institute for Public Health, Ministry of Health; 2008.
- Ismail, A., Marjan, Z. M., & Foong, C. W. (2004). Total antioxidant activity and phenolic content in selected vegetables. *Food Chemistry*, 87(4), 581-586.
- Käferstein, F., & Abdussalam, M. (1999). Food safety in the 21st century. *Bulletin of the World Health Organization*, 77(4), 347.
- Kenny, O., Smyth, T. J., Hewage, C. M., & Brunton, N. P. (2013). Antioxidant properties and quantitative UPLC-MS analysis of phenolic compounds from extracts of fenugreek (*Trigonella foenum-graecum*) seeds and bitter melon (*Momordica charantia*) fruit. *Food chemistry*, 141(4), 4295-4302.

- Khan, M. R., & Omoloso, A. D. (1998). *Momordica charantia* and *Allium sativum*; Broad Spectrum Antibacterial Activity. *Korean Journal of Pharmacognosy*, 29(3), 155-158.
- Krawinkel, M. B., Keding, G. B., (2006). Bitter Gourd (*Momordica charantia*): A dietary approach to hyperglycemia. *Nutrition Reviews*, Vol. 64, No. 7. 331-337
- Kubola, J., & Siriamornpun, S. (2008). Phenolic contents and antioxidant activities of bitter gourd (*Momordica charantia* L.) leaf, stem and fruit fraction extracts in vitro. *Food Chemistry*, 110(4), 881-890.
- Lawless, H. T. (2013). Acceptance and preference testing. In *Laboratory Exercises for Sensory Evaluation* (pp. 101-107). Springer US.
- Lawless, H.T., & Heymann, H. (1998). *Sensory Evaluation of Food: Principles and Practices*. New York: Chapman & Hall.
- Lee, D.K., Kim, B., Lee, S.G., Gwon, H.J., Moon, E.Y., Hwang, H.S., Seong, S.K., Lee, M., Lim, M.J., Sung, H.J., Shin, D.H., Yoon, S.J., Yang, C.H. (1998). Momordins inhibit both AP-1 function and cell proliferation. *Anticancer Research* 18, 119–124.
- Lee, S. E., Hwang, H. J., Ha, J. S., Jeong, H. S., & Kim, J. H. (2003). Screening of medicinal plant extracts for antioxidant activity. *Life sciences*, 73(2), 167-179.
- Lesser, D. (1983). Marketing and sensory quality. In: *Food and Beverages: Definition, Measurement and Control*. Eds. A.A. Williams & R.K. Atkins, Ellis Horwood, Chichester, pp. 448-66.
- Lewis, J. E., Thompson, P., Rao, B. V. V. B. N., Kalavati, C., & Rajanna, B. (2006). Human bacteria in street vended fruit juices: A case study of Visakhapatnam city, India. *Internet Journal of Food Safety*, 8(1), 35-38.
- Lima Tribst, A. A., de Souza Sant'Ana, A., & de Massaguer, P. R. (2009). Review: microbiological quality and safety of fruit juices—past, present and future perspectives. *Critical Reviews in Microbiology*, 35(4), 310-339.
- Maehashi, K., Aritome, Y., Matano, M., & Yamamoto, Y. (2008). Reduction of bitterness of the bitter gourd by Katsuobushi [dried skipjack tuna]. *Journal of the Japanese Society for Food Science and Technology (Japan)*. 186-190.
- Mahesh, B., & Satish, S. (2008). Antimicrobial activity of some important medicinal plant against plant and human pathogens. *World journal of agricultural sciences*, 4(5), 839-843.
- Mahmood, A., Raja, G. K., Mahmood, T., Gulfraz, M., & Khanum, A. (2012). Isolation and characterization of antimicrobial activity conferring component (s) from

seeds of bitter gourd (*Momordica charantia*). *Journal of Medicinal Plants Research*, 6(4), 566-573.

Malaysian Meteorological Department. (2017). Monthly Weather Bulletin. Mean Monthly Temperature. Ministry of Science, Technology and Innovation (MOSTI). Retrieved from: http://www.met.gov.my/en/web/metmalaysia/publications/bulletinpreview/monthlyweather?p_p_id=122_INSTANCE_phshFS8dJslQ&p_p_lifecycle=0&p_p_state=normal&p_p_mode=view&p_p_col_id=_118_INSTANCE_F96odbUpU62g__column-2&p_p_col_count=1&p_r_p_564233524_resetCur=true&p_r_p_564233524_categoryId=431249 on June 15, 2017 at 11.00 am.

Mataragas, M., Dimitriou, V., Skandamis, P. N., & Drosinos, E. H. (2011). Quantifying the spoilage and shelf-life of yoghurt with fruits. *Food Microbiology*, 28(3), 611-616.

Mudgil, S., Aggarwal, D., & Ganguli, A. (2004). Microbiological Analysis of street vended fresh squeezed carrot and kinnow-mandarin juices in Patiala city, India. *Internet Journal of Food Safety*, 3, 1-3.

Nair, S. S., Kavrekar, V., & Mishra, A. (2013). In vitro studies on alpha amylase and alpha glucosidase inhibitory activities of selected plant extracts. *European Journal of Experimental Biology*, 3(1), 128-132.

National Primary Drinking Water Regulations (NPDWR) (2009). United States Environmental Protection Agency. Retrieved from: <https://www.epa.gov/ground-water-and-drinking-water/table-regulated-drinking-water-contaminants#one>, on November 4, 2016 at 11.15am.

Nedwell, D. B. (1999). Effect of low temperature on microbial growth: lowered affinity for substrates limits growth at low temperature. *FEMS Microbiology Ecology*, 30(2), 101-111.

Ng, T.B., Liu, W.K., Sze, S.F., Yeung, H.W. (1994). Action of alphas-momorcharin, a ribosome inactivating protein, on cultured tumor cell lines. *General Pharmacology* 25, 75-77.

Nonga, H. E., Simforian, E. A., & Ndabikunze, B. K. (2014). Assessment of physicochemical characteristics and hygienic practices along the value chain of raw fruit juice vended in Dar es Salaam City, Tanzania. *Tanzania Journal of Health Research*, 16(4).

Oishi, Y., Sakamoto, T., Udagawa, H., Taniguchi, H., Kobayashi-Hattori, K., Ozawa, Y., & Takita, T. (2007). Inhibition of increases in blood glucose and serum

neutral fat by *Momordica charantia* saponin fraction. *Bioscience, biotechnology, and biochemistry*, 71(3), 735-740.

Oki, T., Matsui, T., & Matsumoto, K. (2000). Evaluation of ALPHA.-Glucosidase Inhibition by Using an Immobilized Assay System. *Biological and Pharmaceutical Bulletin*, 23(9), 1084-1087.

Omemu, A. M., & Aderoju, S. T. (2008). Food safety knowledge and practices of street food vendors in the city of Abeokuta, Nigeria. *Food Control*, 19(4), 396-402.

Omeregbe, R.E., Ikuebe, O.M., Ihimire, I.G., 1996. Antimicrobial activity of some medicinal plants extracts on *Escherichia coli*, *Salmonella paratyphi* and *Shigella dysenteriae*. *African Journal of Medical Science* 25, 373–375.

Porat, R., Daus, A., Weiss, B., Cohen, L., Fallik, E., Droby, S. (2000). Reduction of postharvest decay in organic citrus fruit by a short hot water brushing treatment. *Postharvest Biology Technology*. 18, 151-157.

Preethi. (2013, Aug 12). How to Remove Bitter Gourd Bitterness. Retrieved on 11 July 2017 from <http://preethis-vegetableplate.blogspot.my/2013/08/how-to-remove-bitter-gourd-bitterness.html>.

Radke-Mitchell, L. C., & Sandine, W. E. (1986). Influence of temperature on associative growth of *Streptococcus thermophilus* and *Lactobacillus bulgaricus*. *Journal of Dairy Science*, 69(10), 2558-2568.

Raghubeer, E. V., & Matches, J. R. (1990). Temperature range for growth of *Escherichia coli* serotype O157: H7 and selected coliforms in *E. coli* medium. *Journal of Clinical Microbiology*, 28(4), 803-805.

Raman, A., Lau, C. (1996). Anti-diabetic properties and phytochemistry of *Momordica charantia* L. (Cucurbitaceae). *Phytomedicine* 2, 349–362.

Ray, B., & Bhunia, A. (2007). Fundamental food microbiology. CRC press.

Reddi, S. L., Kumar, R. N., Balakrishna, N., & Rao, V. S. (2015). Microbiological quality of street vended fruit juices in Hyderabad, India and their association between food safety knowledge and practices of fruit juice vendors. *International Journal of Current Microbiology and Applied Sciences*, 4, 970-82.

Rizvi, S. I., & Mishra, N. (2013). Traditional Indian medicines used for the management of diabetes mellitus. *Journal of diabetes research*, 2013.

Roller, S. (1999). Physiology of food spoilage organisms. *International Journal of Food Microbiology*, 50, 151–153.

- Sarkar, S., Pranava, M., Marita, R. (1996). Demonstration of the hypoglycemic action of *Momordica charantia* in a validated animal model of diabetes. *Pharmacology Research* 33, 1–4.
- Sathishsekar, D., & Subramanian, S. (2005). Antioxidant properties of *Momordica charantia* (bitter gourd) seeds on Streptozotocin induced diabetic rats. *Asia Pacific Journal of Clinical Nutrition*, 14(2), 153.
- Satkar, K. P., Kulthe, A. A., & Chalke, P. R. (2013). Preparation of bitter gourd ready-to-serve beverage and effect of storage temperature on its keeping quality. *The bioscan*, 8(1), 115-117.
- Schaefer, H., Renner, S. S. (2010). A three-genome phylogeny of *Momordica* (Cucurbitaceae) suggests seven returns from dioecy to monoecy and recent long-distance dispersal to Asia., *Molecular Phylogenetics and Evolution*, 54(2):553-560.
- Semiz, A., & Sen, A. (2007). Antioxidant and chemoprotective properties of *Momordica charantia* L.(bitter melon) fruit extract. *African journal of Biotechnology*, 6(3).
- Shui, G., & Leong, L. P. (2002). Separation and determination of organic acids and phenolic compounds in fruit juices and drinks by high-performance liquid chromatography. *Journal of Chromatography A*, 977(1), 89-96.
- Sidel, J.L., & Stone, H. (1993). The role of sensory evaluation in the food industry. *Food Quality & Preference*, 4, 65-73. San Diego: Academic Press.
- Singh-Ackbarali, D., & Maharaj, R. (2014). Sensory evaluation as a tool in determining acceptability of innovative products developed by undergraduate students in food science and technology at the University of Trinidad and Tobago. *Journal of Curriculum and Teaching*, 3(1), p10.
- Sitasawad, S. L., Shewade, Y., & Bhonde, R. (2000). Role of bittergourd fruit juice in stz-induced diabetic state in vivo and in vitro. *Journal of Ethnopharmacology*, 73(1), 71-79.
- Snee, L. S., Nerurkar, V. R., Dooley, D. A., Efird, J. T., Shovic, A. C., & Nerurkar, P. V. (2011). Strategies to improve palatability and increase consumption intentions for *Momordica charantia* (bitter melon): A vegetable commonly used for diabetes management. *Nutrition Journal*, 10(1), 78.
- Stein, L. J., Nagai, H., Nakagawa, M., & Beauchamp, G. K. (2003). Effects of repeated exposure and health-related information on hedonic evaluation and acceptance of a bitter beverage. *Appetite*, 40(2), 119-129.

- Stone, H., Bleibaum, R., & Thomas, H. A. (2012). *Sensory evaluation practices*. Academic press.
- Stratford, M. (2006). Food and beverage spoilage yeasts. In *Yeasts in food and beverages* (pp. 335-379). Springer Berlin Heidelberg.
- Such A Simple Technique To Remove Bitterness From Bitter Gourd (Karela) (2016, Oct 31). Retrieved on 11 July 2017 from <http://squapl.com/blog/such-a-simple-technique-to-remove-bitterness-from-bitter-gourd-karela--180>.
- Tagne, R. S., Telefo, B. P., Nyemb, J. N., Yemele, D. M., Njina, S. N., Goka, S. M. C., ... & Farooq, A. D. (2014). Anticancer and antioxidant activities of methanol extracts and fractions of some Cameroonian medicinal plants. *Asian Pacific journal of tropical medicine*, 7, S442-S447.
- Takemoto, D.J., Dunford, C., Vaughn, D., Kramer, K.J., Smith, A., Powell, R.G. (1982b). Guanylate cyclase activity in human leukemic and normal lymphocytes. Enzyme inhibition and cytotoxicity of plant extracts. *Enzyme* 27, 179–188.
- Tambekar, D. H., Jaiswal, V. J., Dhanorkar, D. V., Gulhane, P. B., & Dudhane, M. N. (2009). Microbial quality and safety of street vended fruit juices: a case study of Amravati city. *Internet Journal of Food Safety*, 10(7).
- Tazzari, P.L., de-Totero, D., Bolognesi, A., Testoni, N., Pileri, S., Roncella, S., Reato, G., Stein, H., Gobbi, M., Stirpe, F. (1999). An Epstein–Barr virus-infected lymphoblastoid cell line (D430B) that grows in SCID mice with the morphologic features of a CD30+ anaplastic large cell lymphoma, and is sensitive to anti-CD30 immunotoxins. *Haematologica* 84, 988–995.
- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L., & Byrne, D. H. (2006). Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of food composition and analysis*, 19(6), 669-675.
- Thakur, M., & Sharma, R. K. (2016). Bitter Gourd: Health Properties and Value Addition At Farm Scale. *Marumegh*, Volume 1(2), 17-21.
- Tortora, G. J., Funke, B. R., Case, C. L. (2010). *Microbiology, an Introduction*. Tenth Edition. Pearson Education, Inc. 158-159.
- Tournas, V. H., Heeres, J., & Burgess, L. (2006). Moulds and yeasts in fruit salads and fruit juices. *Food Microbiology*, 23(7), 684-688.
- Tsao, S.W., Ng, T.B., Yeung, H.W. (1990). Toxicities of trichosanthin and alpha-momorcharin, abortifacient proteins from Chinese medicinal plants, on cultured tumor cell lines. *Toxicon* 28, 1183–1192.

- Ukwo, S. P., Ndaeyo, N. U., & Udoh, E. J. (2011). Microbiological quality and safety evaluation of fresh juices and edible ice sold in Uyo Metropolis, South-South, Nigeria. *Internet Journal of Food Safety*, 13, 374-8.
- US Food and Drug Administration. (2016). Bacteriological analytical manual (BAM). Retrieved on 4th January 2017 from <http://www.fda.gov/Food/FoodScienceResearch/LaboratoryMethods/ucm2006949.htm#intro>.
- Vats, P. (2013). Thermally Processed Fruit Beverages / Fruit Drink/ Ready to Serve Fruit Beverages. Retrieved on 29 November 2016 from <http://foodsafetyhelpline.com/2013/03/thermally-processed-fruit-beverages-fruit-drink-ready-to-serve-fruit-beverages/>
- Waghray, K., Gulla, S., Kumar, C. S., Kumar, M. P., & Kumar, A. A. (2012). Sensory Quality and Acceptability of Fresh Juices. *Studies on Home and Community Science*, 6(3), 179-181.
- World Health Organization Institute for Public Health. National Health and Morbidity Survey (2011): Volume II Non-Communicable Disease. Kuala Lumpur, Malaysia: Malaysia Institute for Public Health, Ministry of Health; 2011.
- Wu, S. J., & Ng, L. T. (2008). Antioxidant and free radical scavenging activities of wild bitter melon (*Momordica charantia* Linn. var. *abbreviata* Ser.) in Taiwan. *LWT-Food Science and Technology*, 41(2), 323-330.
- Yesilada, E., Gurbuz, I., Shibata, H. (1999). Screening of Turkish antiulcerogenic folk remedies for anti-*Helicobacter pylori* activity. *Journal of Ethnopharmacology* 66, 289–293.