



UNIVERSITI PUTRA MALAYSIA

***BIODEGRADATION OF CYANIDE BY CYANIDE DIHYDRATASE FROM
LOCALLY ISOLATED *Serratia marcescens* ISOLATE AQ07***

KARAMBA KABIRU IBRAHIM

FBSB 2016 21



**BIODEGRADATION OF CYANIDE BY CYANIDE DIHYDRATASE FROM
LOCALLY ISOLATED *Serratia marcescens* ISOLATE AQ07**

By

KARAMBA KABIRU IBRAHIM

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

June 2016

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

I dedicated this thesis to Bauchi State University Gadau (BASUG) for giving me the opportunity to serve and obtain a Philosophy Doctorate Degree (PhD) under its umbrella. I hope the University will grow and be the No. 1 University in the Federal Republic of Nigeria.



Abstract of thesis presented to Senate of Universiti Putra Malaysia in fulfillment of the requirements for the Degree of Doctor of Philosophy

**BIODEGRADATION OF CYANIDE BY CYANIDE DIHYDRATASE FROM
LOCALLY ISOLATED *Serratia marcescens* ISOLATE AQ07**

By

KARAMBA KABIRU IBRAHIM

June 2016

Chairman : Siti Aqlima Ahmad, PhD
Faculty : Biotechnology and Biomolecular Sciences

Cyanide is a very toxic chemical and is one of the environmental pollutants found in sewage. *Serratia marcescens* isolated from soil sample around Universiti Putra Malaysia (3°00'23.91"N, 101°42'31.45"E) was found to have cyanide degrading capability. Spectrophotometric method was used to examine the biodegradation ability of the bacteria in free and immobilised forms using cyanide incorporated buffer medium. Factors affecting cyanide biodegradation such as carbon and nitrogen sources, pH of medium, inoculums size, cyanide concentration and temperature were optimised using one factor at time and response surface methods. Cyanide tolerance and effect of heavy metals (silver, arsenic, cadmium, cobalt, chromium, copper, mercury, nickel, lead and zinc) were investigated. The results illustrates that glucose at 5.5 g/L, yeast extract at 0.55 g/L, pH 6, 20% inoculums size, 200 mg/L cyanide concentration and 32.5°C are the optimum biodegradation conditions required by the bacteria. Immobilised form of the bacteria showed better biodegradation in terms of duration as it degrades the cyanide in 24 hours compared to free cells that require 72 hours degradation process. The bacteria can tolerate 700 mg/L cyanide concentration in free cells and 900 mg/L in immobilised forms. Heavy metals tested at 1 ppm illustrates that the bacteria could stand their effect with the exception of mercury, which degraded only 24.7% in free cells and 61.6% in immobilised forms. Enzyme activity assay illustrates that the bacteria follow the hydrolytic pathway catalysed by cyanide dihydratase to degrade the cyanide. The purified enzyme was able to detoxify 82% of 2 mM potassium cyanide in 10 min of incubation and the rate of cyanide depletion improved linearly as the enzyme concentration is increased. Hydrolysis of cyanide by the purified enzyme fits Michaelis-Menten saturation kinetics when examined over cyanide concentration of 5 mM potassium cyanide. Lineweaver-Burk plot revealed a linear response at 5 mM KCN and less. Michaelis-Menten constant (K_m) for best-fit values of 26.52 and V_{max} value of 1.13 and R^2 value of 0.9 were determined. Total enzyme activity for crude extract stands at 79.9 and 49, 880 mg/L total protein. After final purification process, the total enzyme activity stands at 0.165 with a total protein of 52 mg/L demonstrating yield of 0.207% and purification fold of 65.78. Effect of pH and temperature revealed that enzyme activity was most active at pH of 8 and

temperature of 27°C. The temperature stability test carried out on the enzyme illustrated that it was stable for 70 days at – 20°C and when stored at 4°C, the stability starts reducing after 4 days of incubation. Furthermore, SDS-PAGE electrophoresis post purification revealed the molecular weight of the enzyme to be ~38 kDa, which is a further affirmation. *Serratia marcescens* isolate AQ07 was observed to have the ability to degrade cyanide. Suitable growth and biodegradation conditions were obtained using the optimisation methods. It demonstrates that immobilised cells of the bacteria have a greater ability for cyanide biodegradation compared to free cells, which can be applied for cyanide treatment in sewage. It has been registered in the gene bank as isolate AQ07 with assigned accession number KP213291



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**BIOPENURUNAN SIANIDA OLEH SIANIDA DIHIDRATASE DARIPADA
PEMENCILAN TEMPATAN *Serratia marcescens* PENCILAN AQ07**

Oleh

KARAMBA KABIRU IBRAHIM

Jun 2016

Pengerusi : Siti Aqlima Ahmad, PhD

Fakulti : Bioteknologi dan Sains Biomolekul

Sianida adalah bahan kimia yang sangat toksik dan adalah salah satu daripada pencemaran alam sekitar yang dijumpai di dalam kumbahan. *Serratia marcescens* dipencilkan daripada sampel tanah di Universiti Putra Malaysia (3°00'23.91"N, 101°42'31.45"E) didapati mempunyai keupayaan menurunkan sianida. Kaedah spektrofotometri telah digunakan untuk mengkaji keupayaan biopenurunan oleh bakteria dalam bentuk yang bebas dan sekat gerak menggunakan sianida yang dicampar dengan media penimbal. Faktor yang mempengaruhi biopenurunan sianida seperti sumber karbon dan nitrogen, medium pH, saiz inokulum, kepekatan sianida dan suhu telah dioptimumkan menggunakan satu faktor pada masa dan kaedah gerak balas permukaan. Toleransi sianida dan kesan logam berat (perak, arsenik, kadmium, kobalt, kromium, tembaga, merkuri, nikel, plumbum dan zink) telah dikaji. Keputusan menunjukkan bahawa glukosa pada 5.5 g/L, ekstrak yis pada 0.55 g/L, pH 6, 20% saiz inokulum, 200 mg/L kepekatan sianida dan 32.5°C adalah diperlukan bagi biopenurunan optima oleh bakteria. Bentuk sekat gerak bakteria menunjukkan biopenurunan lebih baik dari segi jangka masa kerana ia menurunkan sianida dalam tempoh 24 jam berbanding dengan sel-sel bebas yang memerlukan 72 jam proses penurunan. Bacteria boleh bertoleransi dengan 700 mg/L kepekatan sianida dalam sel bebas dan 900 mg/L dalam bentuk sekat gerak. Logam berat diuji pada 1 ppm menggambarkan bahawa bakteria boleh tahan kesannya kecuali merkuri, yang menurunkan hanya 24.7% dalam sel bebas dan 61.6% dalam bentuk sekat gerak. Aktiviti enzim asai menunjukkan bahawa bakteria mengikuti laluan hidrolitik menjadi pemangkin oleh sianida hidratase untuk menurunkan sianida. Enzim tulen dapat menyahtoksik 82% daripada 2 mM kalium sianida dalam 10 min masa pengeraman dan kadar pengurangan sianida meningkat secara linear sebagai kepekatan enzim bertambah. Hidrolisis sianida oleh enzim tulen sepadan kinetik tepu Michaelis-Menten apabila diperiksa atas kepekatan sianida pada 5 mM kalium sianida. Plot Lineweaver-Burk mendedahkan tindak balas linear pada 5 mM KCN dan kurang. Pemalar Michaelis-Menten (K_m) untuk nilai-nilai terbaik patut 26.52 dan nilai V_{max} 1.13 dan nilai R^2 0.9 telah ditentukan. Jumlah aktiviti enzim untuk ekstrak mentah berada pada 79.9 dan 49, 880 jumlah protein. Selepas proses penulenan berakhir, aktiviti enzim jumlah mencecah 0.165 dengan jumlah protein 52

mg/L menunjukkan hasil 0.207% dan pembersihan kali ganda 65.78. Kesan pH dan suhu menunjukkan bahawa aktiviti enzim adalah yang paling aktif pada pH 8 dan suhu 27°C. Ujian kestabilan suhu dijalankan ke atas enzim menunjukkan bahawa ia adalah stabil selama 70 hari di -20°C dan apabila disimpan pada 4°C, kestabilan mula mengurangkan selepas 4 hari pengeraman. Tambahan pula, SDS-PAGE elektroforesis penulinan menunjukkan berat molekul enzim menjadi ~38 kDa, yang merupakan ikrar selanjutnya. *Serratia marcescens* pencilan AQ07 telah diperhatikan mempunyai keupayaan untuk menurunkan sianida. Keadaan pertumbuhan dan biopuraian yang sesuai telah diperolehi dengan menggunakan kaedah pengoptimuman. Ia menunjukkan bahawa sel sekat gerak daripada bakteria mempunyai kemampuan yang lebih besar untuk biopenurunan sianida berbanding sel bebas, yang boleh digunakan untuk rawatan sianida dalam kumbahan. Ia telah didaftarkan di GenBank sebagai pencilan AQ07 dengan nombor kesertaan diberikan KP213291.

ACKNOWLEDGEMENTS

First of all, I have to thank the Almighty Allah (SWT), the omnipotent, omnipresent and omniscience to whom glory, honour, majesty and praises are due, who makes mankind in the wombs of their mothers in stages, one after the other, in their veils of darkness, and may peace be upon the most chosen servant of God, Muhammad (SAW) and all prophets sent to mankind.

Next, I would like to thank my parents Alh. Karamba Ibrahim and Hajiya Hauwa Yakubu for their love and support. Oh! Allah have mercy on them as they did care for me when I was young until my present stage.

My special thanks goes to Prof. Ezzeldin Abdurrahman, the Vice Chancellor, Bauchi State University, Gadau (BASUG) for giving me the rare opportunity to serve in the school and to pursue my Ph.D in the Universiti Putra Malaysia (UPM)

I have no words to express my special appreciation to my main supervisor, Dr. Siti Aqlima Ahmad for her kindness, hands-on support, easy to approach attitude and proper supervision during my period of study. No student is luckier than us under her supervision. I pray for Allah to grant her future endeavours.

My special thanks also goes to my initial main supervisor Associate Prof. Dr. Mohd Yunus Abd Shukor, my current my co-supervisor who gave me all the needed support, guide, hints and prompt attention that keep me on track throughout my research. This is enormously essential in the beginning of every student's research project. I have no words to complement his good gesture. I wish him long life and prosperity.

I must also acknowledge the good gesture of my entire supervisory committee; Dr. Siti Aqlima Ahmad, Assoc. Prof. Dr. Mohd Yunus Abd Shukor, Dr. Nur Adeela Binti Yasid and Dr. Azham Bin Zulkarnain for their tremendous support and guidance during the course of my research supervision.

Not to be forgotten, my wife Hanifa Rufa'i Makama and my children Mahmoud and Salim for standing by my side and giving me the moral and emotional support during the course of this programme.

I am also thankful to the good gesture, help and support of my lab partner Dr. Khalizan Bin Sabullah for his tremendous support during my research. I have no words to pass my appreciation to him, may the almighty Allah guide him in his future life endeavours.

Last but not least, I warmly appreciate the good gesture of Assoc. Prof. Dr. Abdulkarim Mohammed, our patron who has helped me with good advices during the course of my programme.

Ma shaa Allah, Alhamdulillah

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Siti Aqlima Ahmad, PhD

Senior Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Mohd Yunus Shukor, PhD

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

Nur Adeela Yasid, PhD

Senior Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

Azham Zulkharnain, PhD

Senior Lecturer

Faculty of Resource Science and Technology

Universiti Malaysia Sarawak

(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work
- quotations, illustrations and citations have been duly referenced
- the thesis has not been submitted previously or concurrently for any other degree at any institutions
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be owned from supervisor and deputy vice –chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No: Karamba Kabiru Ibrahim, GS37491

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____

Name of Chairman
of Supervisory
Committee:

Dr. Siti Aqlima Ahmad

Signature: _____

Name of Member
of Supervisory
Committee:

Associate Professor Dr. Mohd Yunus Shukor

Signature: _____

Name of Member
of Supervisory
Committee:

Dr. Nur Adeela Yasid

Signature: _____

Name of Member
of Supervisory
Committee:

Dr. Azham Zulkharnain

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xv
LIST OF FIGURES	xviii
LIST OF ABBREVIATIONS	xxii
 CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 Cyanide	4
2.2 Uses of cyanide	5
2.3 General sources of cyanide	5
2.4 Cyanide Toxicity	7
2.5 Cyanide pollutions	8
2.6 Environmental occurrence of cyanide	9
2.7 Economic importance of cyanide	10
2.8 Disposal of industrial waste/effluents	11
2.9 Conventional treatment of cyanide	12
2.9.1 Alkaline chlorination	12
2.9.2 Copper catalysed hydrogen peroxide process	13
2.10 Biological cyanide removal	20
2.11 Common pathways for cyanide degradation	20
2.12 Factors affecting the cyanide biodegradation in the environment	23
2.13 Microbial degradation of free/complex cyanide and thiocyanate	24
2.14 Cell immobilisation	25
3 MATERIALS AND METHODS	29
3.1 Chemicals, reagents and equipment	29
3.2 Analytical method	29
3.3 Preparation of analytical reagents	29
3.3.1 Preparation of Standard cyanide solution	29
3.3.2 Buffer solution (pH – 5.2)	29
3.3.3 α - picoline and barbituric reagent	29
3.4 Procedure for cyanide assay	30
3.5 Phenate method for ammonia assay	30
3.5.1 Preparation of hypochlorous acid reagent	30
3.5.2 Preparation of manganese sulphate solution	30
3.5.3 Preparation of phenate reagent	30

3.6	Procedure for phenate assay	30
3.7	Isolation of cyanide degrading bacteria	31
3.7.1	Sampling	31
3.7.2	Handling and growth of bacterial isolates	31
3.8	Screening of cyanide degrading bacteria	32
3.8.1	Primary Screening	32
3.8.2	Secondary screening	32
3.8.3	Tertiary screening	32
3.8.4	Identification of cyanide-degrading bacteria by 16s rRNA	32
3.8.5	Genomic extraction	32
3.8.6	Measurement of DNA concentration	33
3.8.7	Polymerase chain reaction	33
3.8.8	Purification of the amplified PCR products	34
3.8.9	Sequence analysis	34
3.9	Phylogenetic tree analysis and evolutionary relationships of taxa	35
3.10	Characterisation of Isolate AQ07	35
3.11	Preparation of Resting cells	35
3.12	Optimisation of degradation and growth conditions using one factor at a time approach	35
3.12.1	Culture condition	36
3.12.2	Effect of pH	36
3.12.3	Effect of temperature	36
3.12.4	Effect of nitrogen source	36
3.12.5	Effect of carbon source	36
3.12.6	Effect of inoculums size	37
3.12.7	Effect of cyanide concentration	37
3.13	Statistical analysis	37
3.14	Optimisation of degradation and growth conditions using response surface methodology	37
3.14.1	Plackett-Burman design	37
3.14.2	Central composite design	39
3.14.3	Comparison of cyanide degrading activities between free cell and immobilised cells	42
3.15	Effects of cyanide concentrations and evaluation of kinetic models for growth and degradation by free cells	42
3.16	Cell immobilisation	43
3.17	Characterisation of immobilised cells protocols	43
3.17.1	Effect of gellan gum concentration	43
3.17.2	Effect of beads size	44
3.17.3	Effect of number of beads	44
3.17.4	Reusability of beads	44
3.18	Effects of cyanide concentrations and evaluation of kinetic models for cyanide degradation by immobilised cells of <i>Serratia marcescens</i> isolate AQ07	44
3.19	Effect of heavy metals on cyanide-degrading activities among resting cells and immobilised cells	45
3.19.1	Effect of different mercury concentration on cyanide degradation by free and immobilised cells	45

3.20	Determination of cyanide degradation pathway	46
3.21	Preparation of crude enzyme extract	46
3.22	Protein determination	46
3.23	Enzyme determination	46
3.23.1	Oxidative reaction	47
3.23.1.1	Cyanide dioxygenase	47
3.23.1.2	Cyanide monooxygenase	47
3.23.1.3	Cyanase	47
3.23.2	Hydrolytic reaction	47
3.23.2.1	Cyanide dihydratase (Cyanidase) assay	47
3.24	Purification of cyanide degrading enzyme (cyanide dihydratase)	47
3.24.1	Ammonium sulphate precipitation	48
3.24.2	Purification using diethylaminoethyl – cellulose (DEAE – Cellulose) anion exchanger	48
3.24.3	Enzymatic activity of cyanide dihydratase	49
3.24.4	Determination of V _{max} and K _m	49
3.24.5	Determination of K _m and V _{max} using cyanide as substrate	49
3.24.6	Effects of temperature on cyanide dihydratase activity	49
3.24.7	Effects of pH on cyanide dihydratase activity	50
3.24.8	Temperature stability of the cyanide dihydratase	50
3.24.9	SDS- polyacrylamide gel electrophoresis	50
4	RESULTS AND DISCUSSION	52
4.1	Isolation of cyanide-degrading bacteria	52
4.2	Biodegradation of cyanide	52
4.3	Screening of cyanide-degrading bacteria	53
4.4	Identification of cyanide degrading bacteria	56
4.4.1	Gram staining	56
4.4.2	16S rRNA analysis	57
4.4.2.1	Genomic extraction	57
4.4.2.2	Polymerase chain reaction (PCR)	57
4.4.2.3	16S rRNA gene sequencing	58
4.4.3	Phylogenetic analysis	62
4.5	Optimisation via one-factor-at-a-time (OFAT)	64
4.5.1	Effects of carbon source	64
4.5.2	Effects of nitrogen source	65
4.5.3	Effects of pH	67
4.5.4	Effect of temperature	69
4.5.5	Effect of inoculums size	71
4.5.6	Effect of cyanide concentration	73
4.6	Optimisation via response surface method (RSM)	74
4.6.1	Plackett-Burmann	74
4.6.2	Central composite design (CCD)	76
4.6.3	Verification of RSM experiment	87
4.7	Effects of cyanide concentrations on cyanide degradation by free cells of <i>Serratia marcescens</i> isolate AQ07	88
4.8	Evaluation of kinetic models for biodegradation and growth	89

	of cyanide-degrading <i>Serratia marcescens</i> isolate AQ07.	
4.8.1	Effect of various concentrations of cyanide on biodegradation	89
4.8.2	Modelling the degradation and growth kinetics of bacteria	91
4.8.3	Effect of initial cyanide concentration on the growth of bacteria	91
4.8.4	Modelling kinetics of the bacteria growth on cyanide degradation	93
4.9	Effects of heavy metals on cyanide degrading <i>Serratia marcescens</i> isolate AQ07	96
4.9.1	Effects of mercury on cyanide degrading <i>Serratia marcescens</i> isolate AQ07	97
4.10	Immobilisation of bacterial cells	98
4.10.1	Effect of gelling components composition	99
4.10.2	Effect of beads size	101
4.10.3	Effect of cell loading (Number of beads)	103
4.10.4	Reusability of immobilised beads	105
4.11	Response surface methodology (RSM)	107
4.11.1	Verification of RSM experiment	112
4.12	Effect of high cyanide concentration on cyanide biodegradation by immobilised cells of <i>Serratia marcescens</i> isolate AQ07	114
4.13	Review of kinetic models for biodegradation on immobilised cells of cyanide-degrading <i>Serratia marcescens</i> isolate AQ07	114
4.14	Influence of various cyanide concentrations on biodegradation by immobilised cells of <i>Serratia marcescens</i> isolate AQ07	114
4.15	Modelling the detoxification of cyanide by immobilised cells of <i>Serratia marcescens</i> isolate AQ07	115
4.16	Effects of heavy metals on cyanide biodegradation by immobilised cells of <i>Serratia marcescens</i> isolate AQ07	117
4.17	Determination of protein	120
4.18	Determination of cyanide degrading pathway	121
4.18.1	Cyanide dioxygenase, cyanide monooxygenase and cyanase	121
4.18.2	Cyanide dihydratase	122
4.19	Purification of cyanide dihydratase (cyanidase)	128
4.19.1	Ammonium sulphate precipitation	128
4.19.2	Ion exchange chromatography	130
4.19.3	SDS-page polyacrylamide gel electrophoresis	130
4.19.4	Characterisation of cyanide dihydratase(cyanidase)	131
4.19.4.1	Kinetics of cyanide detoxification by partially purified enzyme of <i>Serratia marcescens</i> Isolate AQ07	134
4.19.4.2	Effect of various temperatures on cyanide dihydratase activity	135
4.19.4.3	Effect of pH on cyanide dihydratase activity	136

4.19.4.4	Determination of cyanide dihydratase temperature stability	137
5	CONCLUSION AND RECOMMENDATION	139
	REFERENCES	141
	APPENDICES	153
	BIODATA OF STUDENT	182
	LIST OF PUBLICATIONS	183



LIST OF TABLES

Table	Page
2.1 General sources of cyanide.	7
2.2 Types of cyanide removal technologies and the effectiveness.	14
2.3 Some microorganisms and their biodegradation potential of free/complex cyanide and thiocyanate	16
3.1 Range and level of variables that affects cyanide degradation using Plackett-Burmann design.	38
3.2 Plackett-Burman experimental design for cyanide degradation.	38
3.3 Central composite experimental design for cyanide degradation.	39
3.4 Variability of kinetic models and there mathematical equations.	45
4.1 Degradation capacities of the screened cyanide-degrading bacteria.	53
4.2 Foremost ten sequences producing important alignment with <i>Serratia marcescens</i> Isolate AQ07 from NCBI blast.	62
4.3 Mean squares from analysis of variance and mean values for carbon source effect in biodegradation by free cells of <i>S. marcescens</i> isolate AQ07.	65
4.4 Mean squares from analysis of variance and mean values for nitrogen source effect in biodegradation by free cells of <i>S. marcescens</i> isolate AQ07.	67
4.5 Mean squares from analysis of variance and mean values for pH effect in biodegradation by free cells of <i>S. marcescens</i> isolate AQ07.	69
4.6 Mean squares from analysis of variance and mean values for temperature effect in biodegradation by free cells of <i>S. marcescens</i> isolate AQ07.	71
4.7 Mean squares from analysis of variance and mean values for inoculums size effect in biodegradation by free cells of <i>S. marcescens</i> isolate AQ07.	72
4.8 Mean squares from analysis of variance and mean values for cyanide concentration effect in biodegradation by free cells of <i>S. marcescens</i> isolate AQ07.	74

4.9	ANOVA for central composite design.	77
4.10	Evaluation of optimal conditions and response obtained among OFAT and RSM.	79
4.11	Specific bioremoval rate kinetic parameters acquired for various models through biodegradation of cyanide by <i>Serratia marcescens</i> isolate AQ07 in batch screw cap shake flask.	94
4.12	Specific growth rate kinetic parameters acquired for various models through biodegradation of cyanide by <i>S. marcescens</i> isolate AQ07 in batch screw cap shake flask.	94
4.13	Mean squares from analysis of variance and mean values for effect of gellan gum concentration in biodegradation by immobilised cells of <i>S. marcescens</i> isolate AQ07.	101
4.14	Mean squares from analysis of variance and mean values for effect of beads size in biodegradation by immobilised cells of <i>S. marcescens</i> isolate AQ07.	102
4.15	Mean squares from analysis of variance and mean values for effect of number of beads. (a) in biodegradation by immobilised cells of <i>S. marcescens</i> isolate AQ07.	104
4.16	Mean squares from analysis of variance and mean values for effect of number of beads (a) in biodegradation by immobilised cells of <i>S. marcescens</i> isolate AQ07.	105
4.17	Mean squares from analysis of variance and mean values for effect of reusability of beads in biodegradation by immobilised cells of <i>S. marcescens</i> isolate AQ07.	106
4.18	Central composite experimental design for cyanide biodegradation.	108
4.19	Analysis of Variance (ANOVA) for central composite design.	109
4.20	Comparison of optimum conditions and results obtained between OFAT and RSM.	112
4.21	Mean squares from analysis of variance and mean values for comparison between free and immobilised cells in biodegradation by immobilised cells of <i>S. marcescens</i> isolate AQ07.	113
4.22	Specific biodegradation rate kinetic parameters achieved using different models through biodegradation of cyanide by immobilised cells of <i>Serratia marcescens</i> isolate AQ07 in batch screw cap shake flask.	117

4.23	Mean squares from analysis of variance and mean values for heavy metals effect in biodegradation by immobilised cells of <i>S. marcescens</i> isolate AQ07.	119
4.24	Mean squares from analysis of variance and mean values for mercury concentration in biodegradation by immobilised cells of <i>S. marcescens</i> isolate AQ07.	120
4.25	Various enzyme activities obtained in the course of different reactions.	123
4.26	Cyanide dihydratase purification table	124
4.27	General results of cyanide detoxification bacteria (<i>Serratia marcescens</i> Isolate AQ07). The results of experiment achieved on cyanide detoxification bacteria consist of isolation, screening, characterisation, identification and optimisation of bacteria cyanide detoxification studies	125
4.28	General results of cyanide detoxification bacteria (<i>Serratia marcescens</i> Isolate AQ07) by resting cells and immobilised cells. The results of experiment achieved on cyanide detoxification bacteria consist of characterisation of immobilised protocols, rates of cyanide detoxification, effects of heavy metals and reusability of immobilised beads	126
4.29	General results of cyanide detoxification enzyme (cyanide dihydratase). The results of experiment achieved on cyanide detoxification enzyme consist of characterisation of identification of cyanide degrading pathway, purification of cyanide dihydratase, molecular weight of cyanide dihydratase and enzyme studies	127

LIST OF FIGURES

Figure		Page
2.1	Microscopic view of cyanide (Logsdon <i>et al.</i> , 2015).	5
2.2	Universal pathway accountable for the bioremoval of cyanide and thiocyanate.	21
2.3	The structure of different support materials for immobilisation.	27
4.1	Primary phase screening utilising 25 ppm KCN by six bacteria isolates for 96 hours.	54
4.2	Secondary phase of screening utilising 100 ppm KCN by three bacterial isolates for 120 hours.	55
4.3	Tertiary phase of screening utilising 100 ppm KCN and resting cells of three bacterial isolates for 72 hours.	56
4.4	Gram stain smear of Isolate AQ07.	57
4.5	Agarose gel electrophoresis of genomic DNA extraction and PCR product of 16S rRNA gene of isolate AQ07.	58
4.6	The region of homology between the forward and the reverse complement of Isolate AQ07.	60
4.7	The accession number of 16s rRNA sequence of isolate AQ07 deposited in GeneBank.	61
4.8	This cladogram was carried out based on neighbour joining technique illustrating the phylogenetic relationship among isolate AQ07 and other interconnected analogous bacteria based on 16s rDNA gene sequence examination.	63
4.9	Effects of carbon source on biodegradation of 25 ppm KCN by resting cells of <i>Serratia marcescens</i> isolate AQ07.	65
4.10	Effects of nitrogen sources effect on bioremoval of 25 ppm KCN by resting cells of <i>Serratia marcescens</i> isolate AQ07.	66
4.11	Effects of pH on the bioremoval of 25 ppm KCN by resting cells of <i>Serratia marcescens</i> isolate AQ07.	68
4.12	Effects of temperature on the biodegradation of 25 ppm KCN by resting cells of <i>Serratia marcescens</i> isolate AQ07.	70

4.13	Bacterial inoculums size effect on the biodegradation 100 ppm KCN by <i>Serratia marcescens</i> isolate AQ07.	72
4.14	Cyanide concentration effect on biodegradation by resting cells <i>Serratia marcescens</i> isolates AQ07.	73
4.15	Comparison plot illustrating the dissemination of predicted against actual values in Placket-Burmann.	75
4.16	Comparison plot illustrating the distribution of predicted against actual values of in central composite design.	78
4.17	3D contour curves surface plots indicating interactions among factors.	87
4.18	Validation of experiment (RSM) conducted to ascertain the initial result obtained through the conduct of the experiment.	88
4.19	Result of cyanide concentration on the bioremoval of 200 to 700 mg/L KCN concentration by resting cells of <i>Serratia marcescens</i> isolate AQ07.	89
4.20	Different cyanide concentration on bioremoval by <i>Serratia marcescens</i> isolates AQ07.	90
4.21	various cyanide concentration on <i>Serratia marcescens</i> isolate AQ07 on growth	90
4.22	Growth and degradation rate of <i>Serratia marcescens</i> isolate AQ07 with respect to various cyanide concentration in 72 hours.	91
4.23	Contrast among specific degradation and growth rates at different initial cyanide concentrations.	93
4.24	Experimental and foreseen specific degradation rate (q) of the bacteria at various cyanide concentrations.	95
4.25	Experimental and foreseen specific growth rate (μ) of the bacteria at various cyanide concentrations.	95
4.26	Growth and biodegradation effect of different heavy metals on the removal of 200 ppm KCN by <i>Serratia marcescens</i> isolate AQ07 in 72 hours.	97
4.27	Growth and biodegradation effect of mercury on the removal of 200 ppm KCN by <i>Serratia marcescens</i> isolate AQ07 in 72 hours.	98
4.28	Effects of gellangum concentration in the biodegradation of	100

	200 ppm potassium cyanide by immobilised cells <i>Serratia marcescens</i> isolate AQ07.	
4.29	Effect of size of beads in the bioremoval of 200 ppm cyanide by immobilised cells <i>Serratia marcescens</i> isolate AQ07.	102
4.30	Effect of beads numbers in the bioremoval of 200 ppm cyanide by immobilised cells <i>Serratia marcescens</i> isolate AQ07.	103
4.31	Effect of higher number of beads in the biodegradation of 200 ppm cyanide by immobilised cells <i>Serratia marcescens</i> isolate AQ07.	104
4.32	Effects of reusability of beads in the biodegradation of 200 ppm cyanide by immobilised cells <i>Serratia marcescens</i> isolate AQ07.	106
4.33	3D contour response surface plots illustrating interactions among factors.	111
4.34	Similarity plot between predicted and actual value.	111
4.35	Verification of experiment (RSM) conducted to validate the result obtained.	113
4.36	Effect of concentration of cyanide on the biodegradation of 200 to 900 ppm cyanide by immobilised cells of <i>Serratia marcescens</i> isolate AQ07.	114
4.37	Influence of various cyanide concentration on immobilised cells <i>Serratia marcescens</i> isolate AQ07 on cyanide biodegradation.	116
4.38	Experimental and visualized specific biodegradation rate of immobilised cells <i>Serratia marcescens</i> isolate AQ07 at various cyanide concentration.	116
4.39	Effect of various heavy metals on the biodegradation of 200 ppm cyanide by immobilised cells of <i>Serratia marcescens</i> isolate AQ07 after 24 hrs incubation.	118
4.40	Effect of varying mercury concentration on biodegradation of 200 ppm cyanide by immobilised cells of <i>Serratia marcescens</i> isolate AQ07 after 24 hrs incubation.	119
4.41	Determinations of protein using crude extract of <i>S. marcescens</i> isolate AQ07.	121
4.42	Enzyme activity of cyanide dihydratase via hydrolytic	123

	reaction.	
4.43	Cyanide degrading activity, bacteria growth and ammonia production by <i>S. marcescens</i> isolate AQ07.	128
4.44	Enzyme activity post ammonium sulphate precipitation.	129
4.45	SDS Page of purified cyanide dihydratase.	131
4.46	Phase progression detoxification of 1 to 5 mM KCN by partially purified enzyme of <i>Serratia marcescens</i> Isolate AQ07 conducted as described in materials and methods.	132
4.47	Enzyme activity as a result of cyanide detoxification of 1 to 6 mM KCN by partially purified enzyme of <i>Serratia marcescens</i> Isolate AQ07.	132
4.48	Rate of cyanide disappearance measured as a function of enzyme concentration.	133
4.49	Michaelis –Menten Plot with cyanide substrate.	134
4.50	Lineweaver-Burk Plot with cyanide as substrate.	135
4.51	Effect of temperature on cyanide dihydratase activity.	136
4.52	Effect of pH on cyanide dihydratase activity.	137
4.53	Effect of prolong pre-incubation temperatures on cyanide dihydratase.	138

LIST OF ABBREVIATIONS

(NH ₄) ₂ SO ₄	Ammonium sulphate
>	Greater than
%	Percent
<	Less than
μL	Microliter
°C	Degrees Celsius
μM	Micro molar
As	Arsenic
Ag	Argentum
ATP	Adenosine triphosphate
CFU	Colony Forming Unit
Cd	Cadmium
cm	Centimetre
Cr	Chromium
Co	Cobalt
Cu	Copper
dH ₂ O	Distilled water
DEAE	Diethylaminoethylamine
DNA	Deoxyribonucleic acid
EDTA	Ethylene diamine tetra acetic acid
Fe	Iron
<i>et al</i>	and friends
G	gram
Hg	Mercury

HCL	Hydrogen chloride
HPLC	High Performance Liquid Chromatography
hrs	Hours
Kb	Kilobase
kDa	Kilodaltons
KCN	Potassium Cyanide
Kg	Kilogram
L	Litre
Km	Michaelis-Menten constant
M	Meter
mA	Milliampere
M	Molar
mg	Milligram
mA _u	Milli Absorbance Unit
MgCl ₂	Magnesium Chloride
Min	Minutes
MgSO ₄	Magnesium Sulphate
mM	Millimolar
MSM	Mineral Salt Medium
MW	Molecular Weight
K ₂ HPO ₄	di-Potassium Hydrogen Phosphate
NA	Nutrient Agar
KH ₂ PO ₄	Potassium dihydrogen Phosphate
NaCl	Sodium Chloride
NAD ⁺	Nicotinamide Adenine-dinucleotide oxidized form

NADH	Nicotinamide Adenine-dinucleotide reduced form
Ni	Nickel
OD	Optical Density
Nm	Nanometer
PAGE	Polyacrylamide Gel Electrophoresis
PCR	Polymerase Chain Reaction
Pb	Lead
pH	Log concentration of H ⁺ ion (<i>Puissance hydrogen</i>)
PO ₄ ³⁻	Phosphate
ppm	Parts Per Million
RNA	Ribonucleic Acid
rRNA	Ribosomal ribonucleic Acid
SDS	Sodium Dodecyl Sulphate
TBE	Tri-borate EDTA
<i>Taq</i>	<i>Thermus Aquaticus</i>
TEMED	N,N,N',N'-tetramethyl-ethylenediamine
v/v	Volume/Volume
UV	Ultraviolet
V_{max}	Maximum Velocity
Zn	Zinc
w/v	Weight/Volume
γ	Gamma
β	Beta
α	Alpha

CHAPTER 1

INTRODUCTION

Free cyanide like Hydrogen Cyanide and CN^- are reflected as the most toxic forms of cyanide because of their great metabolic inhibition capacity (Gurbuz *et al.*, 2009). They are universally found in the environment. Amygdalin found in apricots, fruits, vegetables, seeds, cashew nuts, cherries, bean sprouts, is a normal source of HCN (DOH, 2004). Cyanide ensues as normal metabolite in a wide-range of animals, plants and fungi inspite of its noxiousness. It is reflected as one of the poisonous chemicals worldwide. It virtually affects all living organisms as it disables the respiratory functions of a living cell by resolutely binding to the terminal oxidase (Chen *et al.*, 2008). Contact with deadly dosage of cyanide via ingestion, inhalation or skin adsorption devoid of quick first aid action can be tremendously lethal to human being in just a few minutes (Badugu *et al.*, 2005).

Additionally, cyanide also arises anthropogenically. Utilisation of huge amounts of cyanide in several manufacturing practices like metal plating, polymer synthesis, electroplating, steel tempering, and mining signify the significant sources of possible cyanide effluence in the environment (Ebbs *et al.*, 2010). Moreover, cyanide is utilised in gold extraction and jewellery manufacturing that produce wastes highly contaminated with cyanide. The cyanidation method utilised for gold mining has generated over 20% of worldwide cyanide production (Luque-Almagro *et al.*, 2011). Through the cyanidation course, cyanide leaching produces various complexes of metal cyanide.

Due to industrial actions, cyanide compounds and complexes are released as industrial waste into the environment, whereby a predicted 14 million kg/year of entire cyanide discharge from these industries have been reported (Dash *et al.*, 2009). The wastes from these manufacturing industries commonly contain between 0.01 and 10 ppm of total cyanide. Conversely, these figures can increase up to 10,000 to 30,000 ppm as certain cyanide effluents from discrete processes at metal plating and electroplating finishing plants can be stowed for years. Actually, greater levels of cyanide amounting to 100,000 ppm can be obtained in some industrial wastes in which it surpasses the acceptable standards for release to the environment.

Ozonation, alkaline chlorination, sulphur-based technologies and wet air oxidation are some of the presently obtainable chemical approaches for cyanide containing effluent treatments. The utilisation of expensive and unsafe compounds as depolluting agents attested to be unfavourable (Luque-Almagro *et al.*, 2011). Moreover, total removal of cyanide complexes is not attainable by these methods. The employment of commercially prevailing physical and chemical approaches of cyanide removal inclines to produce noxious spinoffs, which also need appropriate treatment and consequently increase the total cost of wastes treatment. Furthermore, exceptional equipment and upkeep are also needed to employ these techniques. Owing to these

demerits, cyanide bearing sewages from several manufacturing plants are incompletely detoxified to the cyanate stage or in vilest scenarios straight release devoid of treatment so as to reduce operational cost (Dash *et al.*, 2009). Hence, the exploration of a substitute treatment method capable of attaining high detoxification efficacy at subsidised cost is very significant.

On the other hand, biological detoxification can be used to treat cyanide bearing effluents. Biological methods of cyanide detoxification that are less expensive and free of risk have consequently been accepted, predominantly in mining industries where cyanide can be converted to inoffensive end products (Ebbs, 2004). Nevertheless, the detoxification of cyanide harbouring effluents by biological methods has not been broadly investigated, which could be attributable to the noxiousness of cyanide and deficiency of the essential knowledge on bacteriological processes (Khor, 2009). Though most bacteria are very subtle to cyanide, others not only endure it, but even utilise it in selected instances as a source of sustenance. Cyanide forbearance can be instructed in two ways, which are by the production of cyanide resilient cytochrome oxidases and by the metabolism of cyanide to harmless end-products (Kunz, 2004).

With fast proliferations of many manufacturing plants that regularly use cyanide, cost-effective and effective degradation methodologies are necessary. Constant researches for assessing the degrading capacities of the new microbes for different cyanide compounds from effluents have to be mutually conducted, in the laboratory and in real scale. As a result, the assortment of microbes has to be based on their capacity to degrade cyanide and also to endure the supplementary pressures, like the thrilling environmental conditions such as higher or lower pH and toxicity effect of other contaminants. The microbes must be capable of competing with native microbial inhabitants efficiently in the environment wherein they will be operative (Dash *et al.*, 2009). Primarily, biological detoxification method has to be established and improvised so as to surpass present technologies with added value benefits as well as filling particular essentials related to the treatment of industrial waste waters.

A broad variety of microbes has been identified to break down the extremely noxious cyanide and therefore established the cyanide metabolic degradation pathways, which have been applied in manufacturing plants for the previous 40 years (Hong, 2006). Several previous efforts in designing a biological method for the degradation of cyanide have focused on cyanide-degrading moulds such as *Trichoderma* and *Fusarium* and quite a few reports have been also reported on the utilisation of bacterial strains like *Klebsiella*, *Acinetobacter*, *Pseudomonas*, *Burkholderia*, *Bacillus* and *Alcaligenes*.

The utilisation of bacteria inter alia appeared to be the best acknowledged and best viable method of biological cyanide treatment. Strains of bacteria are capable of adapting and growing in the cyanide bearing medium either by inducing degradation enzymes or by the inducing of cyanide resisting enzymes (Naveen *et al.*, 2011). Meanwhile, the use of bacterial strains demonstrates to be achievable for practical application in degrading cyanide containing effluent, thus the necessity to explore

extra cyanide detoxification microbes is of great importance. Nevertheless, no additional study on the bacteria was conducted in relation to the removal capacity of its immobilised form or other environmental conditions that could affect the degradation or the degradation pathways that bacteria utilise in the removal process. The dearth of thorough valuation on these bacteria encouraged the present study to deliver a better all-inclusive account on the series of actions that could result to the cyanide degradation. Furthermore, both one factor at a time and response method methodology were employed as the optimisation methods to bring out more detailed options required by the bacteria in the removal of cyanide.

This study aims to bring out a biological methodology for degradation of cyanide by means of bacterial isolate. Bacteria were anticipated to induce cyanide detoxification enzymes or to develop cyanide-resistant enzymes as well as to allow them to adapt and proliferate in the cyanide bearing medium.

The specific objectives of this study are:

1. To isolate, screen and identify cyanide degrading bacteria from soil.
2. To optimise cyanide degradation condition using one factor at a time (OFAT) approach and response surface methodology (RSM) by free and immobilised cells.
3. To determine the effect of cyanide concentration and heavy metals on biodegradation of cyanide by free and immobilised cells.
4. To purify the enzyme responsible for the degradation of cyanide
5. To examine the capacity and kinetic properties of cyanide degradation enzymes

REFERENCES

- Abdel-Salam, A.M., Al-Dekheil, A., Babkr, A., Farahna, M., Mousa, H.M., 2010. High fiber probiotic fermented mare's milk reduces the toxic effects of mercury in rats. *North American Journal of Medical Sciences*. 2, 569.
- Acinas, S.G., Rodríguez-Valera, F., Pedrós-Alió, C., 1997. Spatial and temporal variation in marine bacterioplankton diversity as shown by RFLP fingerprinting of PCR amplified 16S rDNA. *FEMS Microbiology Ecology*. 24, 27–40.
- Adjei, M.D., Ohta, Y., 1999. Isolation and characterization of a cyanide-utilizing *Burkholderia cepacia* strain. *World Journal of Microbiology and Biotechnology*. 15, 699–704.
- Ahmad, S.A., 2011. Biodegradation of phenol by locally isolated *Acinetobacter* sp. strain AQ05NOL 1 and purification of phenol hydrolase. *Universiti Putra Malaysia*.
- Ahmad, S., Ibrahim, S., Shukor, M., Johari, W.L., Rahman, N.A., Syed, M., 2015. Biodegradation kinetics of caffeine by *Leifsonia* sp. strain siu. *Journal of Chemical and Pharmaceutical Sciences*. 8, 312 – 316.
- Ahmed, S.A., 2008. Invertase production by *Bacillus macerans* immobilized on calcium alginate beads. *Journal of Applied Sciences Research*. 4, 1777–1781.
- Aitimbetov, T., White, D.M., Seth, I., 2005. Biological gold recovery from gold – cyanide solutions. *International Journal of Mineral Processing* 76, 33–42.
- Akcil, A., 2003. Destruction of cyanide in gold mill effluents: biological versus chemical treatments. *Biotechnology Advances*. 21, 501–511.
- Aksu, Z., Bülbül, G., 1999. Determination of the effective diffusion coefficient of phenol in Ca-alginate-immobilized *P. putida* beads. *Enzyme and Microbial Technology*. 25, 344–348.
- Alkarkhi, A.F., Ahmad, A., Ismail, N., Easa, A.M., 2008. Multivariate analysis of heavy metals concentrations in river estuary. *Environmental Monitoring and Assessment*. 143, 179–186.
- Alzubaidy, S.K., 2012. The resistance of locally isolated *Serratia marcescens* to heavy metals chlorides and optimization of some environmental factors. *Journal of Environmental and Occupational Sciences*. 1, 37–42.
- Anderson, P.M., 1980. Purification and properties of the inducible enzyme cyanase. *Biochemistry (Mosc.)* 19, 2882–2888.
- ASTDR, 2008. Reader Question: What fruits and vegetables contain Cyanide? <https://www.healthaliciousness.com> (accessed 1.2.16).

- Atkinson, A., 1975. Bacterial cyanide detoxification. *Biotechnology and Bioengineering* 17, 457–460.
- Awasthi, M., Rai, L.C., 2005. Toxicity of nickel, zinc, and cadmium to nitrate uptake in free and immobilized cells of *Scenedesmus quadricauda*. *Ecotoxicology and Environmental Safety*. 61, 268–272.
- Badugu, R., Lakowicz, J.R., Geddes, C.D., 2005. Enhanced fluorescence cyanide detection at physiologically lethal levels: reduced ICT-based signal transduction. *Journal of the American and Chemical Society*. 127, 3635–3641.
- Bai, J., Wen, J.-P., Li, H.-M., Jiang, Y., 2007. Kinetic modeling of growth and biodegradation of phenol and m-cresol using *Alcaligenes faecalis*. *Process Biochemistry*. 42, 510–517.
- Banerjee, A., Ghoshal, A.K., 2010. Phenol degradation by *Bacillus cereus*: Pathway and kinetic modeling. *Bioresource Technology*. 101, 5501–5507.
- Barclay, M., Day, J.C., Thompson, I.P., Knowles, C.J., Bailey, M.J., 2002. Substrate-regulated cyanide hydratase (chy) gene expression in *Fusarium solani*: the potential of a transcription-based assay for monitoring the biotransformation of cyanide complexes. *Environmental Microbiology*. 4, 183–189.
- Barclay, M., Hart, A., Knowles, C.J., Meeussen, J.C., Tett, V.A., 1998. Biodegradation of metal cyanides by mixed and pure cultures of fungi. *Enzyme and Microbial Technology*. 22, 223–231.
- Barnum, D.A., Thackeray, E.L., Fish, N.A., 1958. An outbreak of mastitis caused by *Serratia marcescens*. *Canadian Journal of Comparative Medicine and Veterinary Science* 22, 392 – 395.
- Baş, D., Boyacı, İ.H., 2007. Modeling and optimization I: Usability of response surface methodology. *Journal of Food Engineering*. 78, 836–845.
- Baxter, J., Cummings, S.P., 2006. The current and future applications of microorganism in the bioremediation of cyanide contamination. *Antonie Van Leeuwenhoek* 90, 1–17.
- Beshay, U., 2003. Production of alkaline protease by *Teredinobacter turnirae* cells immobilized in Ca-alginate beads. *African Journal of Biotechnology*. 2, 60–65.
- Biotech, U., 2015. BCH 3102 Enzymology, Universiti Putra Malaysia. 1 - 37
- Botz, M.M., 2001. Overview of cyanide treatment methods. *Mining Environmental Management*. Mining Journal. Ltd London UK. 28–30.
- Button, D.K., 1985. Kinetics of nutrient-limited transport and microbial growth. *Microbiology Reviews*. 49, 270.

- Cappuccino, J.G., Sherman, N., 1999. Basic laboratory techniques for isolation, cultivation and cultural characterization of microorganisms, 5th ed. Benjamin/Cummings Science Publishing.
- Castric, P.A., Strobel, G.A., 1969. Cyanide metabolism by *Bacillus megaterium*. Journal of Biological Chemistry. 244, 4089–4094.
- Chapatwala, K.D., Babu, G.R.V., Vijaya, O.K., Kumar, K.P., Wolfram, J.H., 1998. Biodegradation of cyanides, cyanates and thiocyanates to ammonia and carbon dioxide by immobilized cells of *Pseudomonas putida*. Journal of Industrial Microbiology and Biotechnology. 20, 28–33.
- Chatterjee, S., Sau, G.B., Mukherjee, S.K., 2011. Bioremediation of Cr (VI) from chromium-contaminated wastewater by free and immobilized cells of *Cellulosimicrobium cellulans* KUCr3. Bioremediation Journal. 15, 173–180.
- Chen, C.Y., Kao, C.M., Chen, S.C., 2008. Application of *Klebsiella oxytoca* immobilized cells on the treatment of cyanide wastewater. Chemosphere 71, 133–139.
- Cherruault, Y., 1994. Global optimization in biology and medicine. Mathematical and Computer Modeling. 20, 119–132.
- Chibata, I., Tosa, T., Sato, T., Takata, I., 1986. Immobilization of cells in carrageenan. Methods in Enzymology. 135, 189–198.
- Choi, S.H., Moon, S.-H., Gu, M.B., 2002. Biodegradation of chlorophenols using the cell-free culture broth of *Phanerochaete chrysosporium* immobilized in polyurethane foam. Journal of Chemical Technology and Biotechnology. 77, 999–1004.
- Chung, T.-P., Tseng, H.-Y., Juang, R.-S., 2003. Mass transfer effect and intermediate detection for phenol degradation in immobilized *Pseudomonas putida* systems. Process Biochemistry. 38, 1497–1507.
- Cipollone, R., Ascenzi, P., Tomao, P., Imperi, F., Visca, P., 2008. Enzymatic detoxification of cyanide: clues from *Pseudomonas aeruginosa* Rhodanese. Journal of Molecular Microbiology and Biotechnology. 15, 199–211.
- Cunningham, C.J., Ivshina, I.B., Lozinsky, V.I., Kuyukina, M.S., Philp, J.C., 2004. Bioremediation of diesel-contaminated soil by microorganisms immobilised in polyvinyl alcohol. International Biodeterioration and Biodegradation. 54, 167–174.
- Dash, R.R., Gaur, A., Balomajumder, C., 2009. Cyanide in industrial wastewaters and its removal: a review on biotreatment. Journal of Hazardous Material. 163, 1–11.

- Davey, K.R., 1994. Modelling the combined effect of temperature and pH on the rate coefficient for bacterial growth. *International Journal of Food Microbiology*. 23, 295–303.
- Demoling, L.A., Baath, E., 2008. The use of leucine incorporation to determine the toxicity of phenols to bacterial communities extracted from soil. *Applied Soil Ecology*. 38, 34–41.
- De Souza, M.P., Pickering, I.J., Walla, M., Terry, N., 2002. Selenium assimilation and volatilization from selenocyanate-treated Indian mustard and muskgrass. *Plant Physiology*. 128, 625–633.
- Dhillon, J.K., Shivaraman, N., 1999. Biodegradation of cyanide compounds by a *Pseudomonas species* (S1). *Canadian Journal of Microbiology*. 45, 201–208.
- Dias, J., Rezende, R., Linardi, V., 2001. Bioconversion of nitriles by *Candida guilliermondii* CCT 7207 cells immobilized in barium alginate. *Applied Microbiology and Biotechnology*. 56, 757–761.
- DOE, M., 2003. Malaysia Environmental Quality Report 2002. Ministry of Science, Technology and the Environment Malaysia. (Accessed 18/10/2015)
- DOH, N., 2004. The facts about cyanide. New York State Department of Health. (<http://www.health.state.ny.us>) (Accessed 27/1/2016)
- Drancourt, M., Bollet, C., Carlouz, A., Martelin, R., Gayral, J.-P., Raoult, D., 2000. 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *Journal of Clinical Microbiology*. 38, 3623–3630.
- Dubey, S.K., Holmes, D.S., 1995. Biological cyanide destruction mediated by microorganisms. *World Journal of Microbiology and Biotechnology*. 11, 257–265.
- Dumestre, A., Chone, T., Portal, J., Gerard, M., Berthelin, J., 1997. Cyanide Degradation under Alkaline Conditions by a Strain of *Fusarium solani* Isolated from Contaminated Soils. *Applied and Environmental Microbiology*. 63, 2729–2734.
- du Plessis, C. a., Barnard, P., Muhlbauer, R.M., Naldrett, K., 2001. Empirical model for the autotrophic biodegradation of thiocyanate in an activated sludge reactor. *Letters in Applied Microbiology*. 32, 103–107.
- Dursun, A.Y., Aksu, Z., 2000. Biodegradation kinetics of ferrous (II) cyanide complexes by immobilized *Pseudomonas fluorescens* in a packed bed column reactor. *Process Biochemistry*. 35, 615–622.
- Dursun, A.Y., Tepe, O., 2005. Internal mass transfer effect on biodegradation of phenol by Ca-alginate immobilized *Ralstonia eutropha*. *Journal of Hazardous Material*. 126, 105–111.

- Ebbs, S., 2004. Biological degradation of cyanide compounds. *Current Opinion in Biotechnology*. 15, 231–236.
- Ebbs, S.D., Kosma, D.K., Nielson, E.H., Machingura, M., Baker, A.J.M., Woodrow, I.E., 2010. Nitrogen supply and cyanide concentration influence the enrichment of nitrogen from cyanide in wheat (*Triticum aestivum* L.) and sorghum (*Sorghum bicolor* L.). *Plant Cell and Environment*. 33, 1152–1160.
- Ezzi, M.I., Lynch, J.M., 2005. Biodegradation of cyanide by *Trichoderma* spp. and *Fusarium* spp. *Enzyme and Microbial Technology*. 36, 849–854.
- Ezzi, M.I., Lynch, J.M., 2002. Cyanide catabolizing enzymes in *Trichoderma* spp. *Enzyme and Microbial Technology*. 31, 1042–1047.
- Frost, R.C., 2009. EU practice in setting wastewater emission limit values. <http://www.wgw.org.ua> (2015-09-22 11:30:34)
- Fry, W.E., Millar, R.L., 1972. Cyanide degradation by an enzyme from *Stemphylium loti*. *Archives of Biochemistry and Biophysics*. 151, 468–474.
- Ganczarczyk, J.J., Takoaka, P.T., Ohashi, D.A., 1985. Application of polysulfide for pretreatment of spent cyanide liquors. *Journal of the Water Pollution control Federation*. 1089–1093.
- Gonzalez, G., Herrera, G., Garc , M.T., Pena, M., 2001. Biodegradation of phenolic industrial wastewater in a fluidized bed bioreactor with immobilized cells of *Pseudomonas putida*. *Bioresource Technology* 80, 137–142.
- Guilloton, M., Espie, G.S., Anderson, P.M., 2002. What is the role of cyanase in plants. *Rev Plant Biochemistry and Biotechnology* 1, 57–79.
- Gurbuz, F., Ciftci, H., Akcil, A., 2009. Biodegradation of cyanide containing effluents by *Scenedesmus obliquus*. *Journal of Hazardous Material*. 162, 74–79.
- Hao, O.J., Kim, M.H., Seagren, E.A., Kim, H., 2002. Kinetics of phenol and chlorophenol utilization by *Acinetobacter* species. *Chemosphere* 46, 797–807.
- Haroun, M., Idris, A., Omar, S.S., 2007. A study of heavy metals and their fate in the composting of tannery sludge. *Waste Management*. 27, 1541–1550.
- Heide, B., 2012. Natural Source of Cyanide in Plants. <http://www.ehow.com> (accessed 10.9.15).
- Hong, L.-Y., 2006. Dissipation of cyanide contaminants in the rhizosphere environment. PURDUE UNIVERSITY.
- House, J.M.Y.I., Marsden, J., 1992. The chemistry of gold extraction. Ellis Horwood, London, United Kingdom.

- Ingvorsen, K., Højer-Pedersen, B., Godtfredsen, S.E., 1991. Novel cyanide-hydrolyzing enzyme from *Alcaligenes xylosoxidans* subsp. *denitrificans*. *Applied and Environmental Microbiology*. 57, 1783–1789.
- Jandhyala, D., Berman, M., Meyers, P.R., Sewell, B.T., Willson, R.C., Benedik, M.J., 2003. CynD, the Cyanide Dihydratase from *Bacillus pumilus*: Gene Cloning and Structural Studies. *Applied and Environmental Microbiology* 69, 4794–4805.
- Jones, R., 2012. Bukit Koman: 10 Reasons Why We Protest Against the Use of Cyanide In the Gold Mine Operation <http://www.loyarburok.com> (accessed 11.5.15).
- Kaewkannetra, P., Imai, T., Garcia-garcia, F.J., Chiu, T.Y., 2009. Cyanide removal from cassava mill wastewater using *Azotobacter vinelandii* TISTR 1094 with mixed microorganisms in activated sludge treatment system 172, 224–228.
- Kao, C.M., Chen, K.F., Liu, J.K., Chou, S.M., Chen, S.C., 2006. Enzymatic degradation of nitriles by *Klebsiella oxytoca*. *Applied Microbiology and Biotechnology*. 71, 228–233.
- Kao, C.M., Liu, J.K., Lou, H.R., Lin, C.S., Chen, S.C., 2003. Biotransformation of cyanide to methane and ammonia by *Klebsiella oxytoca*. *Chemosphere* 50, 1055–1061.
- Karamba, K.I., Syed, M.A., Shukor, M.Y., Ahmad, S.A., 2015. Biological Remediation of Cyanide: A Review. *Biotropia* 22, 151 - 163.
- Kenfield, C.F., Qin, R., Semmens, M.J., Cussler, E.L., 1988. Cyanide recovery across hollow fiber gas membranes. *Environmental Science and Technology*. 22, 1151–1155.
- Khor, B.H., 2009. Treatment of Wastewater Containing Cyanide Using Locally Isolated Bacteria Immobilized Onto Bioparticles. University Technology Malaysia. B. Sc. Thesis.
- Knorre, H., Griffiths, A., 1984. Cyanide detoxification with hydrogen peroxide using the Degussa process, in: *Proceedings of the Cyanide and the Environment Conference*, Tucson, Arizona.
- Kumar, V., Kumar, V., Bhalla, T.C., 2013. In vitro cyanide degradation by *Serratia marcescens* RL2b. *International Journal of Environmental Sciences*. 3, 1985–1995.
- Kunz, D.A., 2004. Bacterial Cyanide Assimilation: Pterin Cofactor and Enzymatic Requirements for Substrate Oxidation. University of North Texas.
- Kunz, D.A., Chen, J.-L., Pan, G., 1998. Accumulation of α -keto acids as essential components in cyanide assimilation by *Pseudomonas fluorescens* NCIMB 11764. *Applied and Environmental Microbiology*. 64, 4452–4459.

- Kunz, D.A., Fernandez, R.F., Parab, P., 2001. Evidence that bacterial cyanide oxygenase is a pterin-dependent hydroxylase. *Biochemical and Biophysical Research and Communication*. 287, 514–518.
- Kunz, D.A., Nagappan, O., Silva-Avalos, J., Delong, G.T., 1992. Utilization of cyanide as nitrogenous substrate by *Pseudomonas fluorescens* NCIMB 11764: evidence for multiple pathways of metabolic conversion. *Applied Environmental Microbiology*. 58, 2022–2029.
- Kwon, H.K., Woo, S.H., Park, J.M., 2002. Thiocyanate degradation by *Acremonium strictum* and inhibition by secondary toxicants. *Biotechnology Letters*. 24, 1347–1351.
- Li, P., Feng, X.B., Qiu, G.L., Shang, L.H., Li, Z.G., 2009. Mercury pollution in Asia: a review of the contaminated sites. *Journal of Hazardous Material*. 168, 591–601.
- Liu, Y., He, L., Mustapha, A., Li, H., Hu, Z.Q., Lin, M., 2009. Antibacterial activities of zinc oxide nanoparticles against *Escherichia coli* O157: H7. *Journal of Applied Microbiology*. 107, 1193–1201.
- Logsdon, M.J., Hagelstein, K., Mudder, T., 2015. The management of cyanide in gold extraction. *International Council on Metals and the Environment Ottawa*.
- Lowry, C.O., Rosebrough, N., Farr, A., Randall, R., 1951. Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*. 193, 265–275.
- Luque-Almagro, V.M., Blasco, R., Martínez-Luque, M., Moreno-Vivián, C., Castillo, F., Roldán, M.D., 2011. Bacterial cyanide degradation is under review: *Pseudomonas pseudoalcaligenes* CECT5344, a case of an alkaliphilic cyanotroph. *Biochemical Society Transaction*. 39, 269–274.
- Luque-Almagro, V.M., Huertas, M.-J., Martínez-Luque, M., Moreno-Vivián, C., Roldán, M.D., García-Gil, L.J., Castillo, F., Blasco, R., 2005. Bacterial degradation of cyanide and its metal complexes under alkaline conditions. *Applied and Environmental Microbiology*. 71, 940–947.
- Maniyam, M.N., 2013. Biodegradation of cyanide by Malaysian isolate *Rhodococcus* UKMP-5M. *Universiti Selangor Malaysia*.
- Maniyam, M.N., Sjahrir, F., Ibrahim, A.L., 2011. Bioremediation of Cyanide by Optimized Resting Cells of *Rhodococcus* Strains Isolated from Peninsular Malaysia, *International Journal of Bioscience, Biochemistry and Bioinformatics*. 1, 98–101.
- Maniyam, M.N., Sjahrir, F., Ibrahim, A.L., Cass, A.E.G., 2013. Biodegradation of cyanide by acetonitrile-induced cells of *Rhodococcus* sp. UKMP-5M. *Journal of General and Applied Microbiology*. 404, 393–404.
- Nallapan Maniyam, M., Sjahrir, F., Latif Ibrahim, A., & Cass, A. E. (2015).

- Enzymatic cyanide degradation by cell-free extract of *Rhodococcus* UKMP-5M. *Journal of Environmental Science and Health, Part A*, 50(4), 357-364.
- Megharaj, M., Avudainayagam, S., Naidu, R., 2003. Toxicity of hexavalent chromium and its reduction by bacteria isolated from soil contaminated with tannery waste. *Current Microbiology*. 47, 0051–0054.
- Meyers, P.R., Gokool, P., Rawlings, D.E., Woods, D.R., 1991. An efficient cyanide-degrading *Bacillus pumilus* strain. *Journal of general microbiology* 137, 1397–1400.
- Meyers, P.R., Rawlings, D.E., Woods, D.R., Lindsey, G.G., 1993. Isolation and Characterization of a Cyanide Dihydratase from *Bacillus pumilus* Cl. *Journal of Bacteriology* 175, 6105–6112.
- Meyer, W., 2010. Isolation and Genetic Characterization of a Microbial Consortium Capable of Cyanide Degradation. University of the free state, Bloemfontein, South Africa.
- Mirzadeh, S., Yaghmaei, S., Ghobadi Nejad, Z., 2014. Biodegradation of cyanide by a new isolated strain under alkaline conditions and optimization by response surface methodology (RSM). *Journal of Environmental and Health Science Engineering*. 12, 1–9.
- Mohamed, N.I., 2011. Effect of heavy metals on phenol degradation by consortium bacteria. Universiti Putra Malaysia.
- Mohanty, S.S., Jena, H.M., Satpathy, G.R., 2011. Comparative Study of the Phenol Biodegradation Potential of Free Cells and Immobilized *Pseudomonas Resinovorans*.
- Monod, J., 2012. The growth of bacterial cultures. *Selected Papers in Molecular Biology*. Jacques Monod 371 - 395.
- Mordocco, A., Kuek, C., Jenkins, R., 1999. Continuous degradation of phenol at low concentration using immobilized *Pseudomonas putida*. *Enzyme and Microbial Technology* 25, 530–536.
- Mudder, T.I., Botz, M.M., Smith, A., 2001. Chemistry and treatment of cyanidation wastes. Mining Journal Books, London, UK.
- Muntari, B., Amid, A., Mel, M., Jami, M.S., Salleh, H.M., 2012. Recombinant bromelain production in *Escherichia coli*: process optimization in shake flask culture by response surface methodology. *AMB Express* 2, 1–9.
- Nagashima, S., 1977. Spectrophotometric determination of cyanide with α - picoline - barbituric acid. *Analytica Chimica Acta*. 91, 303–306.
- Naveen, D., Majumder, C.B., Mondal, P., Shubha, D., 2011. Biological treatment of cyanide containing wastewater. *Research Journal of Chemical Sciences* 1, 15–21.

- Nies, D.H., 1999. Microbial heavy-metal resistance. *Applied Microbiology and Biotechnology*. 51, 730–750.
- Nolan, L.M., Harnedy, P.A., Turner, P., Hearne, A.B., O'Reilly, C., 2003. The cyanide hydratase enzyme of *Fusarium lateritium* also has nitrilase activity. *FEMS Microbiology Letters*. 221, 161–165.
- O'Neil, M.J., 2013. The Merck index: an encyclopedia of chemicals, drugs, and biologicals. RSC Publishing. 15th Edition. 1 - 1020.
- O'Reilly, C., Turner, P.D., 2003. The nitrilase family of CN hydrolysing enzymes—a comparative study. *Journal of Applied Microbiology*. 95, 1161–1174.
- Osman, R.M., Ang, L.H., 2000. The occurrence of some potentially toxic elements in ex-mining land located in Bidor. International Atomic Energy Agency, INIS research.
- Özel, Y.K., Gedikli, S., Aytar, P., Ünal, A., Yamaç M., Çabuk, A., Kolankaya, N., 2010. New fungal biomasses for cyanide biodegradation. *Journal of Bioscience and Bioengineering*. 110, 431–435.
- Pandiyan, S., Mahendradas, D., 2011. Application of bacteria to remove Ni (II) Ions from aqueous solution. *European Journal of Scientific Research*. 52, 345–358.
- Parga, J.R., Shukla, S.S., Carrillo-Pedroza, F.R., 2003. Destruction of cyanide waste solutions using chlorine dioxide, ozone and titania sol. *Waste Management*. 23, 183–191.
- Parmar, P., Soni, A., Desai, P., 2013. Enzymatic study of cyanide utilizing *Pseudomonas* species isolated from contaminated soil. *Journal of Science and Innovative Research*. 2, 1058–1066.
- Parmar, P., Soni, A., Vyas, A., Desai, P.V., 2012. Isolation and characterization of cyanide degrading bacterial strains from contaminated soil. *International Journal of Environmental Sciences*. 2, 2006 – 2014.
- Patil, Y.B., Paknikar, K.M., 2000. Development of a process for bi detoxification of metal cyanides from waste waters. *Process Biochemistry* 35, 1139–1151.
- Pong, T.K., Adrien, R.J., Besida, J., O'donnell, T.A., Wood, D.G., 2000. Spent potlining—a hazardous waste made safe. *Process Safety and Environmental Protection*. 78, 204–208.
- Potivichayanon, S., Kitleartpornpaioat, R., 2010. Biodegradation of Cyanide by a Novel Cyanide- degrading Bacterium. *World Academy of Science, Engineering and Technology*. 42, 1362–1365.
- Pritchard, J.D., 2007. Hydrogen Cyanide Toxicological Overview. Health Protection. Agency CHAPD HQ. 1 -11

- Rahman, R., Ghazali, F.M., Salleh, A.B., Basri, M., 2006. Biodegradation of hydrocarbon contamination by immobilized bacterial cells. *Journal of Microbiology.-SEOUL-* 44, 354.
- Raja, C.E., Anbazhagan, K., Selvam, G.S., 2006. Isolation and characterization of a metal-resistant *Pseudomonas aeruginosa* strain. *World Journal of Microbiology and Biotechnology*. 22, 577–585.
- Rand, M.C., Greenberg, A.E., Taras, M.J., others, 1976. Standard methods for the examination of water and wastewater. Prepared and published jointly by American Public Health Association, American Water Works Association, and Water Pollution Control Federation. <http://www.cabdirect.org>. (Accessed: 2015-09-28 15:38:43)
- Raybuck, S.A., 1992. Microbes and microbial enzymes for cyanide degradation. *Biodegradation* 3, 3–18.
- Reyes, G.F., Corbett, D., Benz, F.W., Doyle, R.J., 2000. Acrylonitrile induces autolysis *Bacillus subtilis*. *FEMS Microbiology Letters*. 182, 255–258.
- Rezende, R.P., Teixeira Dias, J.C., Ferraz, V., Linardi, V.R., 2000. Metabolism of benzonitrile by *Cryptococcus* sp. UFMG-Y28. *Journal of Basic Microbiology*. 40, 389–392.
- Saitou, N., Nei, M., 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*. 4, 406–425.
- Sambrook, J., Fritsch, E.F., Maniatis, T., 1989. *Molecular cloning*. Cold spring harbor laboratory press New York. 2, 1 -34
- Saravanan, P., Pakshirajan, K., Saha, P., 2009. Batch growth kinetics of an indigenous mixed microbial culture utilizing m-cresol as the sole carbon source. *Journal of Hazardous Material*. 162, 476–481.
- Saravanan, P., Pakshirajan, K., Saha, P., 2008. Growth kinetics of an indigenous mixed microbial consortium during phenol degradation in a batch reactor. *Bioresource Technology*. 99, 205–209.
- Scopes, R.K., 2013. *Protein purification: principles and practice*. Springer Science & Business Media. 1 - 345
- Silver, S., 1996. Bacterial resistances to toxic metal ions-a review. *Gene* 179, 9–19.
- Skowronski, B., Strobel, G.A., 1969. Cyanide resistance and cyanide utilization by a strain of *Bacillus pumilus*. *Canadian Journal of Microbiology*. 15, 93–98.
- Sorokin, D.Y., Tourova, T.P., Lysenko, a M., Kuenen, J.G., 2001. Microbial thiocyanate utilization under highly alkaline conditions. *Applied and Environmental Microbiology*. 67, 528–38.

- Suh, Y., Park, J.M., Yang, J., 1994. Biodegradation of cyanide compounds by *Pseudomonas fluorescens* immobilized on zeolite. *Enzyme and Microbial Technology* 16, 529–533.
- Tamura, K., Nei, M., Kumar, S., 2004. Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proceedings of the National Academy of Sciences. U. S. A.* 101, 11030–11035.
- Tamura, K., Stecher, G., Peterson, D., Filipinski, A., Kumar, S., 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution*. 30, 2725–2729.
- Tatusova, T.A., Madden, T.L., 1999. BLAST 2 Sequences, a new tool for comparing protein and nucleotide sequences. *FEMS Microbiology Letters*. 174, 247–250.
- Tepe, O., Dursun, A.Y., 2008. Combined effects of external mass transfer and biodegradation rates on removal of phenol by immobilized *Ralstonia eutropha* in a packed bed reactor. *Journal of Hazardous Materials*. 151, 9–16.
- Ubalua, A.O., 2010. Cyanogenic glycosides and the fate of cyanide in soil. *Australian Journal of Crop Sciences*. 4, 223.
- USEPA, 2009. Water Treatability Database. <http://iaspub.epa.gov> (accessed 10.12.15).
- Vidali, M., 2001. Bioremediation. an overview. *Pure and Applied Chemistry*. 73, 1163–1172.
- Wagner-Döbler, I., Lünsdorf, H., Lübbehüsen, T., Von Canstein, H.F., Li, Y., 2000. Structure and species composition of mercury-reducing biofilms. *Applied and Environmental Microbiology*. 66, 4559–4563.
- Wang, X., Gai, Z., Yu, B., Feng, J., Xu, C., Yuan, Y., Lin, Z., Xu, P., 2007. Degradation of carbazole by microbial cells immobilized in magnetic gellan gum gel beads. *Applied and Environmental Microbiology* 73, 6421–6428.
- Watanabe, A., Yano, K., Kazuyoshi, Y., Kazunori, I., Isao, K., 1998. Cyanide hydrolysis in a cyanide-degrading bacterium, *Pseudomonas stutzeri* AK61, by cyanidase. *Microbiology*. 144, 1677–1682.
- White, D.M., Schnabel, W., 1998. Treatment of cyanide waste in a sequencing batch biofilm reactor. *Water Research*. 32, 254–257.
- White, J.M., Jones, D.D., Huang, D., Gauthier, J.J., 1988. Conversion of cyanide to formate and ammonia by a *Pseudomonas* obtained from industrial wastewater. *Journal of Industrial Microbiology*. 3, 263–272.
- Wijffels, R.H., 2000. Immobilized cells, R.H. Wijffels. ed. Springer. 1 - 259

- Wuertz, S., Mergeay, M., van Elsas, J.D., Trevors, J.T., Wellington, E.M.H., others, 1997. The impact of heavy metals on soil microbial communities and their activities. *Modern Soil Microbiology*. 607–642.
- Yadzir, M., Hani, Z., 2007. Characterization, Identification and Application of *Acinetobacter baumannii* Serdang 1 for Phenol Biodegradation. Universiti Teknologi PETRONAS.
- Yamasaki, M., Matsushita, Y., Namura, M., Nyunoya, H., Katayama, Y., 2002. Genetic and immunochemical characterization of thiocyanate-degrading bacteria in lake water. *Applied and Environmental Microbiology*. 68, 942–946.
- Yan, J., Jianping, W., Jing, B., Daoquan, W., Zongding, H., 2006. Phenol biodegradation by the yeast *Candida tropicalis* in the presence of m-cresol. *Biochemical Engineering Journal*. 29, 227–234.
- Yap, C.K., Ismail, A., Edward, F.B., Tan, S.G., Siraj, S.S., 2006. Use of different soft tissues of *Perna viridis* as biomonitors of bioavailability and contamination by heavy metals (Cd, Cu, Fe, Pb, Ni, and Zn) in a semi-enclosed intertidal water, the Johore Straits. *Toxicological and Environmental Chemistry*. 88, 683–695.
- Yap, C.K., Razeff, S.M.R., Edward, F.B., Tan, S.G., 2009. Heavy metal concentrations (Cu, Fe, Ni and Zn) in the clam, *Glaucanome virens*, collected from the northern intertidal areas of peninsular Malaysia. *Malays Applied Biology*. 38, 29–35.
- Ying, W., Ye, T., Bin, H., ZHAO, H., BI, J., CAI, B., 2007. Biodegradation of phenol by free and immobilized *Acinetobacter* sp. strain PD12. *Journal of Environmental Sciences*. 19, 222–225.
- Zeng, L., Huang, J., Zhang, Y., Qiu, G., Tong, J., Chen, D., Zhou, J., Luo, X., 2008. An effective method of DNA extraction for bioleaching bacteria from acid mine drainage. *Applied Microbiology and Biotechnology*. 79, 881–888.