



UNIVERSITI PUTRA MALAYSIA

***MOLECULAR DYNAMICS SIMULATION OF PALM KERNEL OIL
ESTERS-BASED NANO-EMULSION WITH IBUPROFEN AND
DIPALMITOYLPHOSPHATIDYL-CHOLINE LIPID BILAYER***

NUR HANA BINTI FAUJAN

FS 2016 51



**MOLECULAR DYNAMICS SIMULATION OF PALM KERNEL OIL
ESTERS-BASED NANO-EMULSION WITH IBUPROFEN AND
DIPALMITOYLPHOSPHATIDYL-CHOLINE LIPID BILAYER**

By

NUR HANA BINTI FAUJAN

**Thesis Submitted to the School of Graduate Studies,
Universiti Putra Malaysia, in Fulfilment of the Requirement for the
Degree of Doctor of Philosophy**

June 2016

All material contained within the thesis, which including the without limitations texts, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATED

This thesis is lovingly dedicated to

*My great parents,
Prof. Dr. Faujan B. H. Ahmad @ Amat and Mrs. Samilah binti Kutim.*

*My beloved husband,
Khairul Syahmi bin Kamso.*

*My dearest kids,
Muhammad Ahnaf and Khairunnajah.*

*My kindness siblings,
Nur Huda, Nur Hadi, Nur Hani, Nur Hafizah, Nur Hidayah, Nur Hakim, Nur Hariz
and Nur Haziqah.*

*Who lead me with the light of their endless love, support and encourage me
throughout my life.*

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Doctor of Philosophy

**MOLECULAR DYNAMICS SIMULATION OF
PALM KERNEL OIL ESTERS-BASED NANO-EMULSIONS WITH
IBUPROFEN AND DIPALMITOYLPHOSPHATIDYL-CHOLINE
LIPID BILAYER**

By

NUR HANA BINTI FAUJAN

June 2016

Chairman : Roghayeh Abedi Karjiban, PhD
Faculty : Science

Transdermal drug delivery shows a great potential to enhance the permeation process of drugs with poor solubility and low degree of bioavailability. Nevertheless, the penetration of drug through the skin is a big challenge to overcome. Nano-emulsion system can offer the solution to this problem by acting as chemical penetration enhancers (CPEs). Therefore, palm kernel oil esters (PKOEs)-based and oleyl laurate (OLA)-based nano-emulsion systems were used as drug carrier model. PKOEs-based nano-emulsions with ibuprofen (PKOEs:IBU/T80) and without ibuprofen (PKOEs/T80) were simulated followed by the simulation with dipalmitoylphosphatidylcholine (DPPC) in water (PKOEs/DPPC).

The PKOEs/T80, PKOEs:IBU/T80 and PKOEs/DPPC were simulated to determine the effect of surfactant and drug in the model systems. All simulations were performed using all-atom level molecular dynamics (MD) technique for 50 ns. The aggregation process was observed rapidly in the PKOEs-based nano-emulsion systems. These simulations provided better understanding and insight onto the properties of esters, surfactants, drug and water as well as the diffusion of IBU in PKOEs-based nano-emulsion system. A prolate ellipsoidal shape was obtained in both PKOEs/T80 and PKOEs:IBU/T80 models whereas a doughnut-like toroidal shape was gained in PKOEs/DPPC system. The average radius of gyration (R_g) values of 4.43 (± 0.01), 4.50 (± 0.00) and 4.09 (± 0.01) nm were reported for the PKOEs/T80, PKOEs:IBU/T80 and PKOEs/DPPC aggregates, respectively. The radial distribution function (RDF) analysis detected higher interaction between the PKOEs molecules compared to surfactant molecules in all models which could

be due to the hydrophobic interaction in the aggregated structures. In addition, oleyl oleate (OLE) produced the strongest interaction between IBU molecules with the RDF value of 1.26 (± 0.41) in the PKOEs:IBU/T80 aggregate.

Oleyl laurate was used as the main composition of PKOEs for coarse-grained molecular dynamics (CG-MD) simulation study. CG-MD simulation was applied to investigate the aggregation process of OLA-based nano-emulsion with IBU (OLA:IBU/T80) and without IBU (OLA/T80) for 500 ns. The structure of the OLA/T80 and OLA:IBU/T80 aggregates were not completely spherical. The R_g values obtained were 4.36 (± 0.04) and 4.34 (± 0.04) nm, respectively. The distribution of IBU molecules between the OLA was higher compared to T80 molecules in OLA:IBU/T80 model with the RDF values of 1.77 (± 1.16) and 1.12 (± 0.40), respectively. The OLA:IBU/T80 was then simulated with DPPC as a lipid bilayer model. The new model created provided a detailed understanding of the diffusion process of drug through the skin. The OLA:IBU/T80 aggregate was able to move freely inside DPPC molecules. The diffusion of OLA:IBU/T80 also affected the DPPC lipid bilayer structure by disturbing the structure of DPPC and losing the bilayer compactness during 500 ns. The RDF value of DPPC as a lipid bilayer model was decreased from $g(r)=2.92$ to $g(r)=1.22$ in the presence of OLA:IBU/T80 aggregate.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan ijazah Doktor Falsafah

**SIMULASI DINAMIK MOLEKUL BAGI NANO-EMULSI BERASASKAN
ESTER MINYAK ISIRONG KELAPA SAWIT DENGAN IBUPROFEN DAN
LIPID DWILAPISAN DIPALMITOILFOSFATIDIL-KOLIN**

Oleh

NUR HANA BINTI FAUJAN

Jun 2016

Pengerusi : Roghayeh Abedi Karjiban, PhD
Fakulti : Sains

Penghantaran transdermal ubatan mempunyai potensi yang sangat hebat bagi meningkatkan proses penelapan ubat yang kurang larut dan rendah tahap bioketersediaan. Namun begitu, penembusan ubat melalui kulit merupakan satu cabaran yang besar. Sistem nano-emulsi dapat menyelesaikan masalah ini dengan bertindak sebagai bahan kimia peningkat penembusan. Oleh yang demikian, nano-emulsi berasaskan ester minyak isirong kelapa sawit (PKOEs) dan oleil laurat (OLA) telah digunakan sebagai model pembawa ubatan. Nano-emulsi berasaskan PKOEs dengan ibuprofen (PKOEs:IBU/T80) dan tanpa ibuprofen (PKOEs/T80) telah disimulasi dan diikuti oleh simulasi dengan dipalmitoilfosfatidilkolin (DPPC) dalam air (PKOEs/DPPC).

PKOEs/T80, PKOEs:IBU/T80 dan PKOEs/DPPC telah disimulasi untuk menentukan kesan surfaktan dan ubat pada model sistem. Kesemua simulasi telah dijalankan menggunakan teknik dinamik molekul (MD) seluruh-atom selama 50 ns. Proses penggumpalan telah dilihat berlaku sangat pantas pada sistem nano-emulsi berasaskan PKOEs. Simulasi ini memberikan pemahaman yang baik dan pengertian yang mendalam tentang sifat ester, surfaktan, ubat dan air serta penyebaran IBU pada sistem nano-emulsi berasaskan PKOEs. Bentuk elipsoidal lonjong telah terbentuk pada kedua-dua model PKOEs/T80 and PKOEs:IBU/T80 sementara bentuk toroid seakan donat telah terbentuk pada sistem PKOEs/DPPC. Nilai jejari putaran (R_g) yang telah dilaporkan bagi agregat PKOEs/T80, PKOEs:IBU/T80 dan PKOEs/DPPC, masing-masing adalah sebanyak $4.43 (\pm 0.01)$, $4.50 (\pm 0.00)$ dan $4.09 (\pm 0.01)$ nm. Analisis fungsi pengagihan radial (RDF) telah mengesan interaksi yang lebih tinggi di antara molekul PKOEs berbanding molekul surfaktan pada semua model yang

mungkin disebabkan oleh interaksi hidrofobik pada struktur agregat yang diperoleh. Di samping itu, oleil oleat (OLE) telah menghasilkan interaksi yang kuat terhadap molekul IBU dengan nilai RDF sebanyak 1.26 (± 0.41) nm pada agregat PKOEs:IBU/T80.

Oleil laurat telah digunakan untuk pengajian simulasi berbutir kasar-dinamik molekul (CG-MD) kerana ia merupakan komposisi yang utama bagi PKOEs. CG-MD telah digunakan untuk menyiasat proses pengagregatan bagi nano-emulsi berasaskan OLA dengan IBU (OLA:IBU/T80) dan tanpa IBU (OLA/T80) selama 500 ns. Struktur bagi OLA/T80 dan OLA:IBU/T80 agregat merupakan sfera yang tidak sempurna. Nilai R_g yang diperolehi masing-masing adalah 4.36 (± 0.04) dan 4.34 (± 0.04) nm. Pengagihan di antara molekul IBU terhadap OLA adalah lebih tinggi berbanding dengan molekul T80 pada model OLA:IBU/T80 dengan nilai RDF masing-masing 1.77 (± 1.16) dan 1.12 (± 0.40). OLA:IBU/T80 kemudiannya disimulasi terhadap DPPC sebagai model lipid dwilapisan. Model yang baharu dibina dapat memberikan pemahaman yang lebih mendalam bagi penyebaran ubat melalui kulit. Agregat OLA:IBU/T80 mempunyai kebolehan untuk bergerak dengan bebas di dalam molekul DPPC. Penyerapan agregat OLA:IBU/T80 juga mempengaruhi struktur lipid dwilapisan DPPC dengan mengganggu struktur DPPC and kehilangan kepadatan dwilapisan sepanjang 500 ns. Nilai RDF bagi membran DPPC sebagai model lipid dwilapisan telah berkurangan daripada $g(r)=2.92$ ke $g(r)=1.22$ dengan kehadiran struktur agregat OLA:IBU/T80.

ACKNOWLEDGEMENTS

ALHAMDULILLAH, most of thank to ALLAH through these blessings to complete this thesis. First and foremost, I would like to sincerely thank my supervisor Dr Roghayeh Abedi Karjiban. Performing the projects described herein and writing this dissertation would not have been possible without her continuing support and guidance. I am also very grateful for the rest of my committee members, Professor Dr. Mohd Basyaruddin Abdul Rahman and Professor Dr. Mahiran Basri for their closer guidance and suggestions. Through meaningful discussions, they helped me to refine and expand upon my projects, resulting in better work, better publications, and greater scientific impact. I thank each of them for their patience and their belief in my abilities. Their knowledge in a broad range of disciplines helped me gain a new perspective and direction on this research project.

I am deeply grateful to my colleagues for providing me the opportunity to pursue graduate research and creating an environment in which I was able to grow and excel as a researcher. Many thanks to Nurul Syahidah Shaari, Rozana Yahya, Fong Pik Mun, Lim Wu Zhuan and Muhammad Alif Muhammad Latif have been incredibly helpful, especially in doing any computational works. Shafiqah Abdul Ghani, Zalikha Ibrahim, and several other fellow lab mates have been fantastic friends and have made my research work both interesting and valuable. I also thank Dr Rozana Othman for encouraging me an interest in molecular modeling. They have each been exceptionally constructive towards the improvement of my work through lab meetings, informal discussions, e-mail exchanges and general advice in the list goes on.

Many other members of the Universiti Putra Malaysia have been instrumental to my growth and success as a graduate student. Universiti Putra Malaysia and Ministry of Education have each supported part of my work financially and through the intellectual contributions during the attachment at Lancaster University, United Kingdom under Professor Dr Jamshed Anwar. The rest of the staff of Faculty of Science, Department of Chemistry and Centre of Foundation Studies for Agricultural Science, which have created a friendly and inviting atmosphere in which it was a pleasure to work. For each of these things, I am exceedingly grateful.

A special deeply thank to my parents and my family members have been a great support to me through their mutual encouragement. My husband has been incredibly patient and supportive throughout my graduate career. These works are dedicated to him because he is both my motivation and my reason to work hard and to try my best in everything I do. Thank you so much.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Roghayeh Abedi, PhD

Senior Lecturer

Faculty of Science

Universiti Putra Malaysia

(Chairman)

Mahiran Basri, PhD

Professor

Faculty of Science

Universiti Putra Malaysia

(Member)

Mohd Basyaruddin Abdul Rahman, PhD

Professor

Faculty of Science

Universiti Putra Malaysia

(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No: Nur Hana Binti Faujan (GS32443)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of Chairman of Supervisory Committee: _____

Signature: _____

Name of Member of Supervisory Committee: _____

Signature: _____

Name of Member of Supervisory Committee: _____

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENT	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF APPENDICES	xxi
LIST OF ABBREVIATIONS	xxii
 CHAPTER	
1 INTRODUCTION	1
1.1 Background of Research	1
1.2 Problem Statements	2
1.3 Scope of Research	2
1.4 Objectives	3
 2 LITERATURE REVIEW	4
2.1 Transdermal Drug Delivery	4
2.1.1 The Skin	5
2.1.2 Skin Permeation	5
2.1.3 Stratum Corneum and Its Compositions	7
2.1.4 Penetration Enhancers	8
2.1.5 Nano-emulsion and Its Components	8
2.1.5.1 Palm Kernel Oil	9
2.1.5.2 Surfactants	11
2.1.6 Non-steroidal Anti-Inflammatory Drugs	13
2.1.6.1 Ibuprofen	14
2.2 Computational Studies of Bilayer Transport	15
2.2.1 Dipalmitoylphosphatidyl-choline as the Skin Membrane Model	15
2.2.2 Computational Methods Used in the Bilayer Transport Studies	16
2.2.2.1 Molecular Dynamics Simulation	16
2.2.2.2 Coarse-grained Molecular Dynamics Simulation	18
 3 METHODOLOGY	20
3.1 Materials	20
3.1.1 Computer Software	21
3.1.2 Computer Hardware	22
3.2 The Model Systems	23
3.2.1 Palm Kernel Oil Esters-based	23

Nano-emulsions with and without Ibuprofen	
3.2.2 Palm Kernel Oil Esters-based Nano-emulsion with Dipalmitoylphosphatidyl-choline	25
3.2.3 Oleyl Laurate-based Nano-emulsions with and without Ibuprofen	26
3.2.4 Oleyl Laurate-based Nano-emulsions with Ibuprofen and Dipalmitoylphosphatidyl-choline	26
3.3 Simulation Technique	26
3.3.1 Force Field	26
3.3.1.1 The Optimized Potentials for Liquid Force Field	27
3.3.1.2 MARTINI Force Field	28
3.3.2 Preparing the Initial Structure	31
3.3.3 Creating the Topology	33
3.3.4 Mapping Coarse-Grained Structure	33
3.3.4.1 Tween 80	35
3.3.4.2 Oleyl Laurate	38
3.3.4.3 Ibuprofen	40
3.3.4.4 Dipalmitoylphosphatidyl-choline	42
3.3.5 Periodic Boundary Condition	44
3.3.6 Thermodynamic Ensembles	45
3.3.6.1 Constant Temperature Dynamics	46
3.3.6.2 Constant Pressure Dynamics	47
3.3.7 Energy Minimization	48
3.3.8 Production of Model Simulations	49
3.3.8.1 Molecular Dynamics Simulations of Palm Kernel Oil Ester-based Nano-emulsion	49
3.3.8.2 Coarse-Grained Molecular Dynamics Simulations of Oleyl Laurate-based Nano-emulsions	50
3.3.8.3 Coarse-Grained Molecular Dynamics Simulation of Oleyl Laurate-based Nano-emulsion with Ibuprofen and Dipalmitoylphosphatidylcholine	51
3.4 Analysis of the System	53
3.4.1 Physical Properties	53
3.4.1.1 Shape of the Aggregate	53
3.4.1.2 Size of the Aggregate	54
3.4.2 Hydration Properties	55
3.4.2.1 Solvent Accessible Surface Area	55
3.4.2.2 Radial Distribution Function	55
3.4.2.3 Diffusion Analysis	56
3.4.2.4 Area per Lipid	57
4 RESULTS AND DISCUSSION	59
4.1 Effect of Ibuprofen in Palm Kernel Oil Esters-based Nano-emulsions with Tween 80	59

4.1.1 Self-assembly Process	59
4.1.2 Physical Properties	62
4.1.2.1 The Shape of the Aggregates	63
4.1.2.2 The Size of the Aggregates	66
4.1.3 Hydration Properties	67
4.1.3.1 Solvent Accessible Surface Area	67
4.1.3.2 Radial Distribution Function	70
4.1.3.3 Diffusion Analysis	75
4.2 Palm Kernel Oil Ester-based Nano-emulsion with Dipalmitoylphosphatidyl-choline	78
4.2.1 Self-assembly Process	78
4.2.2 Physical Properties	81
4.2.2.1 The Shape of the Aggregate	82
4.2.2.2 The Size of the Aggregate	84
4.2.3 Hydration Properties	85
4.2.3.1 Solvent Accessible Surface Area	85
4.2.3.2 Radial Distribution Function	86
4.2.3.3 Self-diffusion Coefficient	88
4.3 Effect of Ibuprofen in Oleyl Laurate-based Nano-emulsion Model	90
4.3.1 Self-assembly Process	90
4.3.2 Physical Properties	93
4.3.2.1 The Shape of the Aggregates	94
4.3.2.2 The Size of the Aggregates	99
4.3.3 Hydration Properties	100
4.3.3.1 Solvent Accessible Surface Area	100
4.3.3.2 Radial Distribution Function	102
4.3.3.3 Diffusion Analysis	106
4.4 Oleyl Laurate-based Nano-emulsion with Ibuprofen and Dipalmitoylphosphatidyl-choline	108
4.4.1 Physical Properties	108
4.4.1.1 DPPC Bilayer Structure	109
4.4.1.2 Bilayer Thickness	111
4.4.1.3 Area per Lipid	113
4.4.2 The transport of Oleyl Laurate-based Nano-emulsion with Ibuprofen into Dipalmitoylphosphatidyl-choline Bilayer	115
4.4.2.1 Hydration Properties	116
4.4.2.2 Distribution of Oleyl Laurate-based Nano-emulsion with Ibuprofen within the Dipalmitoylphosphatidyl-choline Bilayer	117
4.4.3 Diffusion Analysis	121
5 CONCLUSIONS AND RECOMMENDATION	124
5.1 Conclusions	124
5.2 Recommendations for Future Studies	125

REFERENCES	127
APPENDICES	146
BIODATA OF STUDENT	181
LIST OF PUBLICATIONS	182



LIST OF TABLES

Table		Page
1	The composition of fatty acid in palm kernel oil	10
2	List of Computer Hardware and Software	22
3	The number of molecules used in the PKOEs-based nano-emulsions with and without ibuprofen	24
4	The number of molecules used in the PKOEs-based nano-emulsion with DPPC	25
5	The number of molecules used in the OLA-based nano-emulsions with and without ibuprofen	26
6	The level of interactions between the different CG sites	30
7	Masses assigned to CG particles of T80	35
8	LJ interaction parameters for coarse-grained T80	37
9	Bond parameters for coarse-grained T80	37
10	Angle bending parameters used for coarse-grained T80	38
11	Masses assigned to CG particles of OLA	39
12	LJ interaction parameters for coarse-grained OLA	39
13	Bond parameters for coarse-grained OLA	40
14	Angle bending parameters used for coarse-grained OLA	40
15	Masses assigned to CG particles of IBU	41
16	LJ interaction parameters for coarse-grained IBU	41
17	Bond parameters for coarse-grained IBU	42
18	Angle bending parameters used for coarse-grained IBU	42
19	Masses assigned to CG particles of DPPC	43
20	LJ interaction parameters for coarse-grained DPPC	44
21	Bond parameters for coarse-grained DPPC	44
22	Angle bending parameters used for coarse-grained DPPC	44
23	The radius of gyration (R_g) of PKOEs-based nano-emulsion with and without ibuprofen during the last 2 ns for both simulations	67
24	Total solvent-accessible surface areas (SASA) of PKOEs-based nano-emulsion with and without ibuprofen during the last 2 ns	68
25	Total solvent-accessible surface areas (SASA) of PKOEs, T80 and IBU molecules in both simulations of PKOEs-based nano-emulsions during the last 2 ns	70
26	A summary of the radial distribution function (RDF) values between each oleyl ester, T80 and IBU molecules in the PKOEs-based nano-emulsion model with and without drug	76
27	A summary of the diffusion rate of the PKOEs, T80 and IBU molecules in the PKOEs-based nano-emulsions models	77

28	The principal moments of inertia and the eccentricity (e) values of the OLA/T80 and OLA:IBU/T80 aggregate at 500 ns	98
29	The radial distribution function (RDF) average values between OLA, T80 and IBU in oleyl laurate nano-emulsion system during the last 100 ns of both simulation	106
30	The diffusion rate of OLA, T80 and IBU molecules in OLA/T80 and OLA:IBU/T80 models for 500 ns	108



LIST OF FIGURES

Figure		Page
1	The physiology of human skin	5
2	Transepidermal permeation route in transdermal drug delivery system	6
3	Transappendegeal pathway involves the sweat ducts (1) and the hair follicles (3) whereas the transcellular route through the continuous SC (2) in transdermal drug delivery system	7
4	A Flowchart of the summary of all simulations carried out in this study	21
5	Phase diagram of PKOEs/T80/H ₂ O system at 25±1.0 °C. The isotropic, multiphase, and liquid crystal regions represented in blue, green and yellow, respectively	23
6	Phase diagram of PKOEs with ibuprofen/T80/H ₂ O system at 25±1.0 °C. The isotropic, multiphase, and liquid crystal regions represented in blue, green and yellow, respectively	24
7	Phase diagram of PKOEs/DPPC/H ₂ O system at 25±1.0 °C. The homogenous region represented in white and the multiphase region in green	25
8	Molecular structure of oleyl esters, surfactants and IBU	32
9	CG model for T80 molecule	36
10	CG model of OLA molecule	39
11	CG model of IBU molecule	41
12	CG model of DPPC molecule	43
13	Periodic boundary conditions in two dimensions represented by atom <i>i</i> and <i>j</i> with their images of <i>i'</i> and <i>j'</i>	45
14	Snapshot pictures of the aggregation process for PKOEs/T80 model show a spontaneous self-assembly of PKOEs together with T80 molecules at 4 ns. A prolate ellipsoidal shape was obtained from PKOEs-based nano-emulsion system at 50 ns. The PKOEs molecules are shown in magenta and T80 molecules are lime. Water molecules have been removed for clarity	60
15	Snapshot pictures of the aggregation process for PKOEs:IBU/T80 model show a spontaneous self-assembly of PKOEs together with T80 molecules at 2 ns. A prolate ellipsoidal shape was obtained from PKOEs-based nano-emulsion system at 50 ns. The PKOEs molecules are shown in magenta, T80 molecules are lime and IBU molecules are orange. Water molecules have been removed for clarity	61
16	Root mean square deviations (RMSD) during the self-assembly process of the PKOEs/T80 and PKOEs:IBU/T80 aggregates over 50 ns	62
17	The principal moments of inertia fluctuation of PKOEs/T80 for 50 ns	63

18	The principal moments of inertia fluctuation of PKOEs:IBU/T80 for 50 ns simulation	63
19	The average ratios of the principal moments of inertia (I_1/I_2 , I_1/I_3 and I_2/I_3) of PKOEs/T80 aggregate as a function of time	64
20	The average ratios of the principal moments of inertia (I_1/I_2 , I_1/I_3 and I_2/I_3) of PKOEs:IBU/T80 aggregate as a function of time	65
21	The time evolution profile of the eccentricity (e) fluctuation for PKOEs/T80 and PKOEs:IBU/T80 models	65
22	Radius of gyration (R_g) fluctuation versus time for PKOEs/T80 and PKOEs:IBU/T80 nano-emulsion during 50 ns	66
23	Total solvent-accessible surface areas (SASA) is shown in purple, hydrophobic (green) and hydrophilic (black) contacted with the PKOEs/T80 aggregate in PKOEs-based nano-emulsion system as a function of time	69
24	Total solvent-accessible surface areas (SASA) is shown in purple, hydrophobic (green) and hydrophilic (black) contacted with the PKOEs:IBU/T80 aggregate in PKOEs-based nano-emulsion system as a function of time	69
25	Radial distribution function (RDF) changes in both PKOEs/T80 and PKOEs:IBU/T80 aggregates	71
26	Radial distribution functions (RDF) changes between the PKOEs and the T80 molecules in the PKOEs/T80 aggregated structure	72
27	Radial distribution functions (RDF) changes between the PKOEs, T80 and IBU molecules in the PKOEs:IBU/T80 aggregated structure	72
28	Radial distribution function (RDF) of the water molecules that were near to the T80 molecules and PKOEs molecules in the PKOEs/T80 nano-emulsion system during the last 2 ns of the simulation times	73
29	Radial distribution function (RDF) of the water molecules that were near to the T80 molecules and PKOEs molecules in the PKOEs:IBU/T80 nano-emulsion system during the last 2 ns of the simulation times	74
30	Mean square displacements (MSD) changes of the PKOEs/T80 and PKOEs:IBU/T80 aggregates for 50 ns	77
31	Snapshot pictures of PKOEs/DPPC system during molecular dynamics (MD) simulations. These snapshots show the spontaneous self-assembly of PKOEs together with the DPPC into the doughnut-like toroidal shape. The PKOEs are shown in orange and DPPC molecules are in blue. Water molecules have been removed for clarity	80

32	The number of cluster changes during the self-assembly process	81
33	Root mean square deviations (RMSD) of the PKOEs/DPPC model system as a function of simulation time	82
34	The principal moments of inertia fluctuations of PKOEs/DPPC model system for 50 ns	83
35	The changes of the ratio of the principal moments of inertia of PKOEs/DPPC aggregate during 50 ns	83
36	Eccentricity fluctuations of the formation PKOEs/DPPC aggregate during 50 ns	84
37	The radius of gyration (R_g) fluctuations of PKOEs/DPPC model system for 50 ns	85
38	Solvent accessible surface areas (SASA) of PKOEs/DPPC model system during 50 ns simulation	86
39	Radial distribution functions (RDF) of PKOEs and DPPC molecules in the PKOEs/DPPC model during 50 ns	87
40	Radial distribution functions (RDF) analysis between the PKOEs molecules and the DPPC molecules in the aggregated structure formed during the self-assembly of PKOEs/DPPC system	88
41	Mean Square Displacement (MSD) of PKOEs/DPPC self-assembled structure as a function of time	89
42	Mean Square Displacement (MSD) changes of PKOEs and DPPC molecules in the PKOEs/DPPC model during 50 ns	89
43	Potential energy of OLA/T80 and OLA:IBU/T80 models for 500 ns	90
44	The snapshots of OLA/T80 aggregate for 500 ns; oleyl laurate (orange) and Tween 80 (purple). Water was omitted for clarity	91
45	The snapshots of OLA:IBU/T80 aggregate for 500 ns; oleyl laurate (orange), Tween 80 (purple) and ibuprofen (blue). Water was omitted for clarity	92
46	The Root Mean Square Deviations (RMSD) of OLA/T80 and OLA:IBU/T80 aggregates for 500 ns	94
47	The principal moments of inertia fluctuations of OLA/T80 aggregate for 500 ns	95
48	The principal moments of inertia fluctuations of OLA:IBU/T80 aggregate for 500 ns	96
49	The principle moment of inertia ratios plot for OLA/T80 model as a function of time	97
50	The principle moment of inertia ratios changes for OLA:IBU/T80 model as a function of time	97
51	The eccentricity (e) fluctuation for OLA/T80 and OLA:IBU/T80 aggregates through 500 ns simulation times	99
52	Radius of gyration fluctuations of OLA/T80 and OLA:IBU/T80 aggregates during 500 ns simulation	100

53	The hydrophobic SASA of OLA/T80 and OLA:IBU/T80 models during the simulation	101
54	Radial Distribution Function (RDF) change of the hydrophobic regions for OLA/T80 and OLA:IBU/T80 models during the last 100 ns of both simulation	103
55	Radial Distribution Function (RDF) between OLA/T80 and OLA:IBU/T80 aggregates with water during the last 100 ns	104
56	Radial Distribution Function (RDF) between OLA and T80 molecules with water in the OLA/T80 system during the last 100 ns of simulation	104
57	Radial Distribution Function (RDF) between OLA, T80 and IBU molecules with water in OLA:IBU/T80 system during the last 100 ns of simulation	105
58	The Mean Square Displacement (MSD) changes of OLA/T80 and OLA:IBU/T80 aggregates during 500 ns simulation	107
59	The snapshot pictures of DPPC aggregation for 500 ns simulation time; the polar head groups of choline (blue); phosphate (grey); glycerol (pink); the hydrocarbon tail (cyan) and water (yellow)	110
60	Density distribution profile of DPPC in water during the last 100 ns	112
61	Density distribution profile of DPPC lipid bilayer with OLA:IBU/T80 system during 500 ns	113
62	The area per lipid changes of DPPC lipid bilayer in the presence of OLA:IBU/T80 aggregate for 500 ns simulation time	114
63	The snapshots picture of the starting simulation. The OLA:IBU/T80 aggregate was placed outside of the DPPC bilayer.	115
64	The snapshot pictures of OLA:IBU/T80 and DPPC bilayer during 500 ns simulation time	116
65	Radial distribution function changes of DPPC in the bilayer system at the last 100 ns	117
66	Radial distribution function of DPPC in the presence of OLA:IBU/T80 at the last 100 ns	118
67	Radial distribution function of <i>sn</i> -1 tail relative to the other <i>sn</i> -1 tails	118
68	Radial distribution function of the center of mass of <i>sn</i> -1 tail relative to the other <i>sn</i> -1 tails in the presence of OLA:IBU/T80	119
69	The radial distribution function (RDF) of DPPC lipid bilayer at interface	120
70	The distribution of OLA:IBU/T80 calculated according to the distance between the OLA:IBU/T80 center of mass and the membrane center of mass during 500 ns simulation time	121
71	The Mean Square Displacement (MSD) change of DPPC lipid bilayer system	122

72	The Mean Square Displacement (MSD) changes of DPPC in the presence of OLA:IBU/T80 aggregate	123
----	---	-----



LIST OF APPENDICES

Appendix		Page
A	Supplementary data for the Model System	146
A1	Calculation for number of molecule used in the model	146
B	Topology of the molecules used for MD Simulation	147
B1	Oleyl caproate	147
B2	Oleyl laurate	148
B3	Oleyl stearate	150
B4	Oleyl caprate	152
B5	Oleyl myristate	153
B6	Oleyl oleate	155
B7	Oleyl caprylate	157
B8	Oleyl palmitate	159
B9	Oleyl linoleate	160
B10	Tween 80	162
B11	Ibuprofen	166
C	Supplementary data for Mapping Coarse-Grained Structure	168
C1	Mapping of Tween 80 Structure	168
D	Simulation Parameters for MD Simulation	170
D1	Parameter file for Energy Minimization	170
D2	Parameter file for Position Restrained Heating	170
D3	Parameter file for Equilibration	171
D4	Parameter file for Production runs	173
D5	Simulation Parameters for CG-MD Simulation	174
	Parameter file for Energy Minimization	174
D6	Parameter file for Position Restrained Heating	175
D7	Parameter file for Equilibration	176
D8	Parameter file for Production runs	178
E1	Radius of gyration (R_g) of the PKOEs and T80 molecules in the PKOEs/T80 model	180
E2	Radius of gyration (R_g) of the PKOEs, T80 and IBU molecules in the PKOEs:IBU/T80 model	180
E3	A mean square displacements (MSD) analysis of the PKOEs and the T80 molecules in PKOEs/T80 aggregate during 50 ns	180
E4	A mean square displacements (MSD) analysis of the PKOEs, T80 and IBU molecules in PKOEs:IBU/T80 during 50 ns	180

LIST OF ABBREVIATIONS

AA	all-atom
ACAT	advanced compartment absorption and transit
AO	Atomic Orbitals
AOT	aerosol OT
BBB	blood brain barrier
CERs	ceramides
CG	coarse-grained
CG-MD	coarse-grained molecular dynamics
CHOL	cholesterol
CMC	critical micelle concentration
CML	Chemical Markup Language
COX	cyclo-oxygenase
COX-1	cyclo-oxygenase-1
COX-2	cyclo-oxygenase-2
CPEs	chemical penetration enhancers
DeTAB	<i>n</i> -decyltrimethylammonium bromide
DMPC	dimyristoylphosphatidylcholine
DMSO	dimethyl sulfoxide
DOPC	dioleoylphosphatidylcholine
DPD	dissipative particle dynamics
DPPC	Dipalmitoylphosphatidylcholine
ER	enhancement ratio
FFAs	free fatty acids
FFT	Fast Fourier Transform
GAMESS	General Atomic and Molecular Electronic Structure System
GI	Gastrointestinal
GROMACS	Groningen Machine for Chemistry Simulation
GTO	Gaussian Type Orbitals
HF	Hartree-Fock
HLB	hydrophilic-lipophilic balance
IBU	Ibuprofen
IV	intravenous
L_{α}	lamellar phase
LCAO	linearly combining atomic orbitals
LJ	Lennard-Jones
MC	Monte Carlo
MD	Molecular dynamics
MM	molecular mechanical
MM/QM	molecular mechanics/quantum mechanics
NPT	number of atoms, pressure, temperature
NSAIDs	Non-steroidal anti-inflammatory drugs
NVE	number of atoms, volume and energy
NVT	number of atoms, volume, temperature
O/W	oil-in-water
OCL	Oleoyl caprylate

OCP	Oleyl caprate
OCR	Oleyl caproate
OLA	Oleyl laurate
OLE	Oleyl oleate
OLI	Oleyl linoleate
OMY	Oleyl myristate
OPA	Oleyl palmitate
OPLS	optimized potentials for liquid
OPLS	Optimized Potential for Liquid
OST	Oleyl stearate
PBC	Periodic boundary condition
PBC	Periodic Boundary Condition
PC	posphatidylcholine
PE	phosphatidylethanolamine
PEG400	polyethylene glycol 400
PFOB	perfluorooctylbromide
PFOB-based	perfluorooctylbromide-based
PFOB-NEP	PFOB nano-emulsion interface
PGs	prostaglandins
PGTOs	Primitive Gaussian Type Orbital
PKO	Palm kernel oil
PKOEs	palm kernel oil esters
PME	Particle-mesh Ewald
PMFs	potentials of mean force
POPC	1-palmitoyl-2-oleyl-phosphatidylcholine
PPEs	physical penetration enhancers
PVA	polyvinyl alcohol
QM	quantum mechanical
RESP	restrained electrostatic potential
RMs	reverse micelles
SANS	small-angle neutron scattering
SC	<i>Stratum corneum</i>
SDS	sodium dodecyl sulphate
SLM	solid lipid micro-particles
STO	Slater Type Orbitals
T80	Tween 80
UA	united atom
VMD	Visual Molecular Dynamics
W/O	water-in-oil

CHAPTER 1

INTRODUCTION

1.1 Background of Research

Ibuprofen (IBU) is one of the non-steroidal anti-inflammatory drugs (NSAIDs) used for pain relief, anti-stiffness and anti-inflammatory effect in rheumatoid arthritis, osteoarthritis, fever and gout treatments. It has a short plasma half-life, low degree of bioavailability and low solubility in water (Winstanley and Walley, 2002). Drug based nano-emulsions with nano droplet size have the ability to enhance the absorption of drugs with poor solubility and low bioavailability like IBU. Drugs can diffuse effectively by using the nano-emulsion formulation. Nano-emulsion systems can solubilize the hydrophobic substances within water-based phase (Delmas *et al.*, 2010) and improve the permeation of many drugs for transdermal delivery application (Shakeel *et al.*, 2009).

Nano-emulsion systems can act as chemical penetration enhancers (CPE) and drug carrier (Shakeel *et al.*, 2007; Kong *et al.*, 2011). Palm kernel oil esters (PKOEs)-based nano-emulsions have been produced to be applied for drug delivery system (Salim *et al.*, 2011; Musa *et al.*, 2013; De Costa *et al.*, 2014; Razaee *et al.*, 2014). PKOEs with relatively short chain length hydrocarbon can be considered as a good carrier to deliver drugs into the body (Keng *et al.*, 2009).

Transdermal drug delivery system has been developed to control the release of drugs. Transdermal delivery is the administration of drug molecules directly to the targeting area through the *stratum corneum* (SC) of skin. The mobility of drugs, ions and water molecule is controlled by the SC skin barrier arranged in multiple bilayers called lamellae structure (Marrow *et al.*, 2007; Subedi *et al.*, 2010; Iwai *et al.*, 2012).

The skin barrier consists of lipid structure in the extracellular space between the SC cells. Lipid consists of free fatty acids (FFAs), long-chain ceramides (CERs) and cholesterol (CHOL) (Wertz and Noelen, 2003). The SC is a dead keratinocytes layer of epidermal cells embedded in a lipid matrix (Morrow *et al.*, 2007). The lipid matrix is formed by a parallel orientation of the lipid head groups in a bilayer structure.

Molecular dynamics (MD) simulation is one of the computational approaches which have been used to model the nano-emulsion (Abdul Rahman *et al.*, 2008; Abdul Rahman *et al.*, 2009; Lee *et al.*, 2010) and lipid systems (Eriksson and

Eriksson, 2011; Mihailescu *et al.*, 2011) for several years. Many interesting properties of the systems are possible to be understood and estimated in both atomic and molecular levels. MD technique also has the abilities to simulate the interactions of nano-materials with biological membranes (Shi *et al.*, 2008; Wallace and Sansom, 2008).

1.2 Problem Statements

Oral intakes of IBU may cause gastrointestinal (GI) ulcers or perforations, stomach bleeding and kidney toxicity in long term treatment (Beetge *et al.*, 2000). Transdermal drug delivery system could be the alternative to transfer IBU drug through the skin. Transdermal delivery has many advantages as compared to other routes by avoiding first-pass hepatic metabolism and providing patient compliance (Prausnitz and Langer, 2008). Nevertheless, there is a big challenge in transdermal application, considering the physicochemical properties of IBU. The transportation of IBU molecules has to overcome the skin barrier. Nano-emulsion systems could improve the properties of IBU in order to enhance the permeability of drug passing through the skin.

Many experimental works tried to produce nano-emulsion systems to achieve the optimal level of drug permeation through the SC (Shakeel *et al.*, 2009). They focused mostly on the formulation and preparation process. However, the self-assembly or self-aggregation of nano-emulsions and the formation of droplet are difficult to be observed using laboratory tools. The laboratory experiments hardly detect the distribution of drug molecules in nano-emulsion droplet and the fundamental mechanism on how the system can penetrate the SC.

Computer simulations could be applied to describe the behaviour of the simulated nano-emulsion model by determining their structural and dynamical properties (Abedi Karjiban *et al.*, 2015). An all-atomic MD technique is still limited due to a practical upper simulation time limit of ~100 ns for complex systems if computer clusters are not used. The main problem for all-atom (AA) MD techniques is to reach the real equilibration state. This problem can be solved by using coarse-grained molecular dynamics (CG-MD). CG-MD represents the system by reducing the numbers of atoms as compared with an all-atom description. Coarse-graining approach can be very helpful to extend the simulation time and bridge the gap between the simulation and experimental techniques.

1.3 Scope of Research

By using the experimental findings reported, detailed understanding and insight into the interaction of nano-emulsion system with lipid bilayer can be

further explained by applying computer simulation approaches. In this project, MD simulation was applied to obtain the molecular structure, the stability and the dynamical information of PKOEs-based nano-emulsion system. The simulations were utilized to investigate the mechanism of self-assembly process of PKOEs-based nano-emulsion model with and without IBU. Later on, the CG-MD study was performed to simulate the self-assembly process of oleyl laurate (OLA)-based nano-emulsion system with and without IBU. Oleyl laurate (OLA) is the main component of PKOEs. This CG model was then used to study the distribution of OLA-based nano-emulsion containing IBU through dipalmitoylphosphatidyl-choline (DPPC) lipid bilayer. DPPC lipid bilayer was simulated as the uppermost SC layer of skin. Overall, this study could explain the application of PKOEs-based nano-emulsion system as a carrier of IBU drug for transdermal delivery.

1.4 Objectives

The main objective of this research was to apply MD simulation techniques to model the self-assembly process of PKOEs-based and OLA-based nano-emulsions with and without IBU followed by exploring the distribution of OLA-based nano-emulsion with IBU through DPPC lipid bilayer. Therefore, the following specific objectives were pursued:

1. To examine the mechanism of self-assembly process in both PKOEs-based and OLA-based nano-emulsion systems.
2. To determine the physicochemical and dynamical properties of PKOEs-based and OLA-based nano-emulsions models.
3. To investigate the diffusion process of IBU in both PKOEs-based and OLA-based nano-emulsions model systems.
4. To identify the distribution OLA:IBU/T80 aggregate in DPPC.

REFERENCES

- Abdul Rahman, M. B., Mohd Latif, M. A., Basri, M., Salleh, A. B., and Abd Rahman, R. N. Z., (2008). Molecular dynamics simulation of palm-based nano-emulsion system. *Proceeding MCBC'08 Proceedings of the 9th WSEAS International Conference on Mathematics & Computers In Biology & Chemistry*, 112-117.
- Abdul Rahman, M.B., Huan, Q.-Y., Tejo, B.A., Basri, M., Salleh, A.B., and Abdul Rahman, R.N.Z., (2009). Self-assembly formation of palm-based esters nano-emulsion: A molecular dynamics study. *Chemical Physics Letters*, 480(4-6): 220-224.
- Abedi Karjiban, R., Huan, Q.-Y., Abdul Rahman, M. B., Basri, M., and Bimo, A. T., (2015). Self-assembly of palm kernel oil wax esters in aqueous media: A molecular dynamics study. *International Journal of Chemistry*, 7(1): 133-139.
- Abedi Karjiban, R., Shaari, N.S., Gunasakaran, U.V., and Basri, M., (2013). A coarse-grained molecular dynamics study of DLPC, DMPC, DPPC, and DSPC mixtures in aqueous solution. *Journal of Chemistry*, v 2013.
- Adhangale, P. S., and Gaver, D. P. III, (2006). Equation of state for a coarse-grained DPPC monolayer at the air/water interface. *Molecular Physics*, 104(19): 1-19.
- Akiladevi, D., and Basak, S., (2010). Ethosomes – a noninvasive approach for transdermal drug delivery. *International Journal of Current Pharmaceutical Research*, 2(4): 1-4.
- Allen, M.P., (2004). Introduction to Molecular Dynamics Simulation. Published in *Computational soft Matter: From Synthetic Polymers to Proteins*, John von Neumann Institute for Computing, Julich, NIC Series, Vol. 23, ISBN 3-00-012641-4, pp. 1-28.
- Allen, M.P., and Tildesley, D.J. (1989). Computer simulation of liquids, Oxford University Press.
- Anton, N., and Vandamme, T. F., (2009). The universality of low-energy nano-emulsification. *International Journal of Pharmaceutics*, 377(1-2): 142-147.
- Anuchapreeda, S., Fukumori, Y., Okonogi, S., and Ichikawa, H., (2012). Preparation of lipid nanoemulsions incorporating curcumin for cancer therapy. *Journal of Nanotechnology*, 270383.

- Asbill, C. S., and Michniak, B. B., (2000). Percutaneous penetration enhancers: local versus transdermal activity. *Pharmaceutical Science and Technology Today*, 3(1): 36-41.
- Azeem, A., Ahmad, F. J., Khar, R., and Talegaonkar, S., (2009). Nanocarrier for the Transdermal Delivery of an Antiparkinsonian Drug. *American Association of Pharmaceutical Scientists PharmciTech*, 10(4):1093-1103.
- Baheti, S. R., Wadher, K. J., and Umekar, M. J., (2011). A recent approach towards transdermal drug delivery by physical and chemical techniques. *Internationale Pharmaceutica Scientia*, 1(1): 44-53.
- Baker, M. T., and Naquib, M., (2005). Propofol: The challenges of formulation, *Anesthesiology*, 103: 860-876.
- Bayly, C.I., Cieplak, P., Cornell, W., and Kollman, P.A. (1993). A well-behaved electrostatic potential based method using charge restraints for deriving atomic charges: the RESP model. *The Journal of Physical Chemistry*, 97(40): 10269-10280.
- Bedrov, D., Smith, G. D., Davande, H., and Li, L., (2008). Passive transport of C₆₀ fullerenes through a lipid membrane: Molecular dynamics simulation study. *Journal of Physical Chemistry B*, 112(7): 2078-2084.
- Beetge, E., du Plessis, J., Muller, D. G., Goosen, C., and van Rensburg, F. J., (2000). The influence of the physicochemical characteristics and pharmacokinetic properties of selected NSAID's on their transdermal absorption. *International Journal of Pharmaceutics*, 193(2): 261-264.
- Bemporad, D., Essex, J., and Luttmann, C., (2004). Permeation of Small Molecules through a Lipid Bilayer: A Computer Simulation Study. *Journal of Physical Chemistry B*, 108, 4875-4884.
- Benjapornkulaphong, S., Ngamcharussrivichai, C., Bunyakiat, K., (2009). Al₂O₃-supported alkali and alkali earth metal oxides for transesterification of palm kernel oil and coconut oil. *Chemical Engineering Journal*, 145(3): 468-474.
- Berendsen, H.J.C., van der Spoel, D., and van Drunen, R. (1995). GROMACS: A message-passing parallel molecular dynamics implementation. *Computer Physics Communications*, 91: 43-56.
- Berendsen, H. J. C., Postma, J. P. M., van Gunsteren, W. F., and Hermans, J., (1981). Interaction models for water in relation to protein hydration, In: *Intermolecular Forces*, B. Pullman ed., Reidel, Dordrecht, 331-342.

- Berendsen, H. J. C., Postma, J. P. M., van Gunsteren, W. F., DiNola, A., and Haak, J.R., (1984). Molecular Dynamics with coupling to an external bath. *The Journal of Chemical Physics*, 81(8): 3684-3690.
- Biltonen, R., Lichtenberg, D., (1993). The use of differential scanning calorimetry as a tool to characterize liposome preparations. *Chemistry and Physics of Lipids*, 64,129-142.
- Bode, B.M., and Gordon, M.S., (1998). MacMolplt: a graphical user interface for GAMESS. *Journal of Molecular Graphics and Modelling*, 16(3): 133-138.
- Bombardier, C., Laine, L., Reicin, A., Shapiro, D., Burgos-Vargas, R., Davis, B., Day, Richard., Ferraz, M. B., Hawkey, C. J., Hochberg, M. C., Kvein, T.g K., and Schnitzer, T. J., (2000). Comparison of upper gastrointestinal toxicity of rofecoxib and naproxen in patients with rheumatoid arthiritis. *The New England Journal of Medicine*, 343: 1520-1528.
- Bond, P. J., and Sansom, M. S., (2006). Insertion and assembly of membrane proteins via simulation. *Journal of the American Chemical Society*, 128(8): 2697-2704.
- Bond, P. J., Holyoake, J., Ivetec, A., Khalid, S., Sansom, M. S. P., (2007). Coarse-grained molecular dynamics simulations of membrane proteins and peptides. *Journal of Structural Biology*, 157(3): 593-605.
- Brooks, B.R., Bruccoleri, R.E., Olafson, B.D., States, D.J., Swaminathan, S., and Karplus, M. (1983). CHARMM: A program for macromolecular energy, minimization, and dynamics calculations. *Journal of Computational Chemistry*, 4(2): 187-217.
- Bruce, C. D., Berkowitz, M. L., Perera, L., and Forbes, M. D. E., (2002) Molecular dynamics simulation of sodium dodecyl sulfate micelle in water: micellar structural characteristics and counterion distribution. *Journal of Physical Chemistry B*, 106: 3788-3793.
- Bussi, G., Donadio, D., and Parrinello, M., (2007). Canonical sampling through velocity rescaling. *Journal of Chemical Physics*, 126: 014101.
- Capek, I., (2004). Degradation of kinetically-stable o/w emulsions. *Advances in colloid and Interface Science*, 107(2-3): 125-155.
- Chandrashekar, N.S. and Shobha Rani, R.H., (2008). Physicochemical and Pharmacokinetic Parameters in Drug Selection and Loading for Transdermal Drug delivery. *Indian Journal Pharmaceutical Sciences*, 70(1); 94-96.

- Chantasart, D., Sa-Nguandeeikul, P., Prakongpan, S., Li, S. K., and Higuchi, W. I., (2007). Comparison of the effects of chemical permeation enhancers on the lipoidal pathways of human epidermal membrane and hairless mouse skin and the mechanism of enhancer action. *Journal of Pharmaceutical Sciences*, 96(9): 2310-2326.
- Cheon, M., Chang, I., and Hall, C. K., (2010). Extending the PRIME model for protein aggregation to all 20 amino acids. *Proteins*, 78: 2950-2960.
- Cornell, W. D., Cieplak, P., Bayly, C. I., Gould, I. R., Merz, K. M., Ferguson, D. M., Spellmeyer, D. C., Fox, T., Caldwell, J. W., and Kollman, P.A., (1995). A second generation force field for the simulation of proteins, nucleic acids, and organic molecules, *Journal of the American Chemical Society*, 117(19): 5179-5197.
- Daful, A.G., Avalos, J.B., and Mackie, A.D., (2012). Model shape transitions of micelles: spheres to cylinders and disks. *Langmuir*, 28: 3730-3743.
- Damm, W., Frontera, A., Tirado-Rives, J., and Jorgensen, W.L., (1997). OPLS all-atom force field for carbohydrates, *Journal of Computational Chemistry*, 18(16): 1955-1970.
- Darden, T., Perera, L., Li, L., and Pedersen, L., (1999). New tricks for modelers from the crystallography toolkit the particle mesh Ewald algorithm and its use in nucleic acid simulations. *Structure*, 15;7(3):R55-60.
- Darden, T., York, D., and Pedersen, L., (1993). Particle mesh Ewald: An $N \log(N)$ method for Ewald sums in large systems. *Journal of Chemical Physics*, 98:10089-10092.
- De Costa, S., Basri, M., Shamsudin, N., and Basri, H., (2014). Formulation of stable palm kernel oil esters nanoemulsion system containing hydrocortisone. *Asian Journal of Chemistry*, 26(10): 2883-2888.
- de Joannis, J., Jiang F. Y., and Kindt J.T., (2006). Coarse-grained model simulations of mixed-lipid systems: composition and line tension of a stabilized bilayer edge. *Langmuir*, 22: 998-1005.
- de Maura, A.F., and Freitas, L.C.G., (2005). Molecular dynamics simulation of the sodium octanoate micelle in aqueous solution. *Chemical Physical Letters*, 411:474-478.
- Delmas, T., Piraux, H., Couffin, A. C., Texier, I., Vinet, F., Poulin, P., Cates, M. E., and Bibette, J., (2011). How to prepare and stabilize very small nanoemulsions. *Langmuir*, 27(5): 1683-1692.

- Devarajan, V., and Ravichandran, V., (2011). Nanoemulsions: As modified drug delivery tool. *International Journal of Comprehensive Pharmacy*, 2(4): 1-6.
- Dickey, A. N., and Faller, R., (2006). Investigating interactions of biomembranes and alcohols: A multiscale approach. *Journal of Polymer Science part B: Polymer Physics*, 43(8): 1025-1032.
- Dokka, S., Cooper, S. R., Kelly, S., Hardee, G. E., Karras, J. G., (2005). Dermal delivery of topically applied oligonucleotides via follicular transport in mouse skin. *Journal of Investigative Dermatology*, 124: 971-975.
- Dooley, J. M., Gordon, K. E., Wood, E. P., Brna, P. M., MacSween, J., and Fraser, A., (2007). Caffeine as an adjuvant to ibuprofen in treating childhood headaches. *Pediatric Neurology*, 37(1): 42-46.
- Eriksson, E.S.E., and Eriksson, L.A., (2011). The Influence of cholesterol on the properties and permeability of hypericin derivatives in lipid membranes. *Journal of Chemical theory and Computation*, 7(3): 560-574.
- Essman, U., Perera, L., Berkowitz, M.L., Darden, T., Lee, H., Pedersen, L.G., (1995). A smooth particle mesh ewald potential. *Journal of Chemical Physics*, 103: 8577-8592.
- Faller, R., and Marrink, S. J., (2004). Simulation of Domain Formation in DLPC-DSPC mixed bilayers. *Langmuir*, 20(18): 7686-7693.
- Feller, S.E., (2000). Molecular Dynamics Simulations of Lipid Bilayers. *Current Opinion Colloid Interface Science*, 5:217-223.
- Ferguson, D.M., (1995). Parameterization and evaluation of a flexible water model. *Journal of Computational Chemistry*. 16, 501-511.
- Flynn, G. L., (2002). Cutaneous and transdermal delivery process and system of delivery. In Banker, G. S., and Rhodes, C. T., *Modern Pharmaceutics*, Fourth Edition, New York, CRC Press, 292-319.
- Gao, X., and Wong, T. C., (2001). NMR studies of adrenocorticotropin hormone peptides in sodium dodecylsulfate and dodecylphosphocholine micelles: proline isomerism and interactions of the peptides with micelles. *Biopolymers*, 58(1): 20-32.
- Goyal, P. S., and Aswal, V. K., (2001). Micellar structure and inter-micelle interactions in micellar solutions: Results of small angle neutron scattering studies. *Soft Condensed Matter*, 8(8): 972-979.

- Gunawan, E.R., Basri, M., Abdul Rahman, M.B., Salleh, A. B., and Abdul Rahman, R.N.Z. (2005). Study on response surface methodology (RSM) of lipase-catalyzed synthesis of palm-based wax ester. *Enzyme and Microbial Technology*, 37(7), 739-744.
- Halgren, T.A., (1999). MMFF VII. Characterization of MMFF94, MMFF94s, and other widely available force fields for conformational energies and for intermolecular-interaction energies and geometries, *Journal of Computational Chemistry*, 20(7): 730-748.
- Hanwell, M.D., Curtis, D.E., Lonie, D., Vandermeersch, T., Zurek, E., and Hutchison, H.R. (2012). Avogadro: an advanced sematic chemical editor, visualization, and analysis platform. *Journal of Cheminformatics*, 4: 17.
- Hess B., Kutzner C., van der Spoel D. and Lindahl E. (2008). GROMACS 4: Algorithms for highly efficient, load-balanced, and scalable molecular simulation. *Journal of Chemical Theory and Computation*, 4: 435-447.
- Hess, B., Bekker, H., Berendsen, H.J.C., Fraaije, J.G.E.M., (1997). LINCS: A linear constraint solver for molecular simulation. *Journal of Computational Chemistry*. 18:1463-1472.
- Hockney, R. W., (1970). The potential calculation and some applications. *Methods in Computational Physics* vol. 9 pp. 135-211 Academic Press, New York.
- Hoover, W. G., (1985). Canonical dynamics: Equilibrium phase-space distribution. *Physical Review A*, 31(3): 1695-1697.
- Huang, H., Chung, B., Jung, J., Park, H.-W., and Chang, T., (2009). Toroidal micelles of uniform size from Diblock copolymers, *Angewandte Chemie International Edition*, 48: 4594-4597.
- Humphrey, W., Dalke, A., and Schulten, K., (1996). VMD: Visual Molecular Dynamics. *Journal of Molecular Graphics*, 14:33-38.
- Hünenberger, P. H., (2005) Thermostat Algorithms for Molecular Dynamics simulations. *Advance Polymer Sciences*, 173: 105-149.
- Huzil, J.T., Sivaloganathan, S., Kohandel, M., and Foldvari, M., (2011). Drug delivery through the skin: molecular simulations of barrier lipids to design more effective noninvasive dermal and transdermal delivery systems for small molecules, biologics, and cosmetics. *Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology*, 3(5): 449-462.

- Israelachvili, J., and Ladyzhinski, I., (2005). The physico-chemical basis of self-assembling structures, Skjeltorp, A. T., Belushkin, A.V., (Eds.), *Forces, Growth and Form in Soft Condensed Matter: At the Interface between Physics and Biology*, vol. 160 Springer, Netherlands, 1–28.
- Iwai, I., Han, H.M., den Hollander, L., Svensson, S., Ofverstedt, G., Anwar, J., Brewer, J., Bloksgaard, M., Laloef, A., Nosek, D., Masich, S., Bagatolli, L. A., Skoglund, U., and Norlen, L., (2012). The human skin barrier is organized as stacked bilayers of fully extended ceramides with cholesterol molecules associated with the ceramide sphingoid moiety. *Journal of Investigative Dermatology*, 132: 2215-2225.
- Jahning, F., (1996). What is the surface tension of a lipid bilayer membrane?. *Biophysical Journal*, 71(3):1348-1349.
- Jiang, Y., Zhu, J., Jiang, W., and Liang, H., (2005). Cornucopian cylindrical aggregate morphologies from self-assembly of amphiphilic triblock copolymer in selective media. *Journal of Physical Chemistry B*, 109(46): 21549–21555.
- Jitputti, J., Kitiyanan, B., Rangsunvigit, P., Bunyakiat, K., Attanatho, L., Jenvanitpanjakul, P., (2006). Transesterification of crude palm kernel oil and crude coconut oil by different solid catalysts. *Chemical Engineering Journal*, 116(1), 61-66.
- Jorgensen, W. L., and Swenson, C. J., (1985). Optimized intermolecular potential functions for amides and peptides. Hydration of amides, *Journal of American Chemical Society*, 107(6): 1489-1496.
- Jorgensen, W. L., and Tirado-Rives, J., (1988). The OPLS (optimized potentials for liquid simulations) potential functions for proteins, energy minimizations for crystals of cyclic peptides and crambin, *Journal of American Chemical Society*, 110(6): 1657-1666.
- Jorgensen, W. L., Maxwell, D. S., and Tirado-Rives, J., (1996). Development and Testing of the OPLS all-atom force field on conformational energetic and properties of organic liquids, *Journal of American Chemical Society*, 118(45): 11225-11236.
- Kaminski, G., and Jorgensen, W.L., (1996). Performance of the AMBER94, MMFF94, and OPLS-AA force fields for modeling organic liquids. *Journal of Physical Chemistry B*, 100: 18010-18013.
- Kaminski, G., Duffy, E.M., Matsui, T., and Jorgensen, W.L., (1994). Free energies of hydration and pure liquid properties of hydrocarbons from the OPLS all-atom model. *Journal of Physical Chemistry* 98(49): 13077-13082.

- Kaminski, G.A., Friesner, R.A., Tirado-Rives, J., and Jorgensen, W.L., (2001). Evaluation and Reparametrization of the OPLS-AA Force for Protein via Comparison with Accurate Quantum Chemical Calculations on Peptides. *Journal of Physical Chemistry B*, 105(28): 6474-6487.
- Karande, P., Jain, A., Ergun, K., Kispersky, V., and Motragotri, S., (2005). Design principles of chemical penetration enhancers for transdermal drug delivery. *Proceedings of the National Academy of Sciences*, 102(13): 4688-4693.
- Kasson, P.M., Kelly, N. W., Singhal, N., Vrljic, M., Brunger A. T., and Pande V. S., (2006). *Proceeding of the National Academy of Sciences USA*, 103(32): 11916-11921.
- Keng, P.S., Basri, M., Zakaria, M.R.S., Abdul Rahman, M. B., Ariff, A.B., Abdul Rahman, R. N. Z., and Salleh, A. B., (2009). Newly synthesized palm esters for cosmetics industry. *Industrial Crops and Products*, 29(1): 37-44.
- Khalid, S., Bond, P. J., Holyoake, J., Hawtin, R. W., and Sansom, S. P., (2008). DNA and lipid bilayers: self-assembly and insertion. *Journal of Royal Society Interface*, 241-250.
- Kim, H.-J., Kang, S.-K., Lee, Y.-K., Seok, C., Lee, J.-K., Zin, W.-C., and Lee, M., (2010). Self-dissociating tubules from helical staking of noncovalent macrocycles, *Angewandte Chemie International Edition*, 49(45): 8471-8475.
- Kim, J.K., Lee, E., Kim, M.-C., Sim, E., and Lee, M., (2009). Reversible transformation of helical coils and straight rods in cylindrical assembly of elliptical macrocycles. *Journal of American Chemical Society*, 123, 8159-8160.
- Klang, V., and Valenta, C., (2011). Lecithin-based nanoemulsions, *Journal of Drug Delivery Science and Technology*, 21: 55-76.
- Kong, M., Chen, X. G., Kweon, D. K., and Park, H. J., (2011). Investigations on skin permeation of hyaluronic acid based nanoemulsion as transdermal carrier. *Carbohydrate Polymers*, 86: 837-843.
- Kravs, D. M., and Pharm, J. T., (2005). Neonatal therapy. In: Koda-Kimble, M. A., Young, L. V., Kradjan, W. A., Guglielmo, B. J., Alldredge, B. K., and Corelli, R. L., editors. *Applied therapeutics: the clinical use of drugs*, 8th ed., Lipponcott William and Wilkins A Wolters Kluwer company Philadelphia New York, 94-23.

- Kucerka, N., Liu, Y., Chu, N., Petrache, H.I., Tristram-Nagle, S., Nagle, J.F., (2005). Structure of fully hydrated fluid phase DMPC and DLPC lipid bilayer using X-ray scattering from oriented multilamellar arrays and from unilamellar vesicles. *Biophysical Journal*, 88: 2626-2637.
- Kucerka, N., Nieh, M-P, and Katsaras, J., (2011). Fluid phase lipid areas and bilayer thickness of commonly used phosphatidylcholine as a function of temperature. *Biochimica et Biophysica Acta (BBA)- Biomembranes*, 1808(11), 2761-2771.
- Kuhn, R. J., Zhang, W., Rossmann, M. G., Pletnev, S. V., Corver, J., Lenches, E., Jones, C. T., Mukhopadhyay, S., Chipman, P. R., Strauss, E. G., Baker, T. S., and Strauss, J. H., (2002). Structure of dengue virus: Implications for flavivirus organization, maturation, and fusion, *Cell*, 108(5): 717-725.
- Kumar, S., Tyagi, L. K., and Chandra, A., (2010). Chemical penetration enhancers: An approach for better transdermal drug delivery. *International Journal of Pharmaceutical Research and Development*, 3(7): 87-95.
- Kwatra, S., Taneja, G., and Nimisha, N., (2012). Alternative Routes of Drug Administration-Transdermal, Pulmonary & Parenteral. *Indo global Journal of Pharmaceutical sciences*, 2(4): 409-426.
- Kweon, J.-H., Chi, S.-C., and Park, E.-S., (2004). Transdermal delivery of diclofenac using microemulsions. *Archives of Pharmacal Research*, 27(3): 351-356.
- Langer, R., and Peppas, N. A., (2003). Advances in biomaterials, drug delivery, and bionanotechnology. *American Institute of Chemical Engineers Journal*, 49(12): 2990-3006.
- Leach, A.R. (2001). *Molecular modelling: principles and applications*. Pearson Education.
- Lee H., and Larson, R.G., (2006). Molecular dynamics simulations of PAMAM dendrimer-induced pore formation in DPPC bilayers with a coarse-grained model. *Journal of Physical Chemistry B*, 110:18204-18211.
- Lee, E., Kim, J.-K., and Lee, M., (2009). Tubular stacking of water-soluble toroids triggered by guest encapsulation. *Journal of American Chemical Society*, 131: 18242-18243.
- Lee, E., Kim, J.-K., and Lee, M., (2009b). Reversible scrolling of two-dimensional sheets from self-assembly of laterally-grafted amphiphilic rods. *Angewandte Chemie International Edition*, 48: 3657-3660.

- Lee, S.-J., Olsen, B., Schlesinger, P. H., and Baker, N. A., (2010). Characterization of perfluorooctylbromide-based nanoemulsion particles using atomistic molecular dynamics simulations. *Journal of Physical Chemistry B*, 114(31): 10086-10096.
- Lichtenberger, L. M., Romero, J. J., de Ruijter, W. M. J., Behbod, F., Darling, R., Ashraf, A. Q., and Sanduja, S. K., (2001). Phosphatidylcholine association increases the anti-inflammatory and analgesic activity of ibuprofen in acute and chronic rodent models of joint inflammation: Relationship to alterations in bioavailability and cyclooxygenase-inhibitory potency. *The Journal of Pharmacology and Experimental Therapeutics*, 298(1): 279-287.
- Lindahl, E., and Edholm, O., (2000). Mesoscopic undulations and thickness fluctuations in lipid bilayers from molecular dynamics simulations. *Biophysical Journal*, 79: 426-433.
- Lindahl, E., Hess, B., van der Spoel, D., (2001). GROMACS 3.0: a package for molecular simulation and trajectory analysis, *Journal of Molecular Modelling*, 7: 306-317.
- Lopez, C. A., Rzepiela, A. J., de Vries, A. H., Dijkhuizen, L., Hunenberger, P. H., and Marrink, S. J., (2009). Martini coarse-grained force field: Extension to carbohydrates, *Journal of Chemical Theory and Computation*, 5(12): 3195-3210.
- Lopez, C. A., Sovova, Z., van Eerden, F. J., de Vries, A. H., Marrink, S. J., (2013). Martini force field parameters for glycolipids. *Journal of Chemical Theory and Computation*, 9(3): 1694-1708.
- Lopez, C. F., Nielsen, S. O., Moore, P. B., and Klein, M. L. (2004). Understanding nature's design for a nanosyringe, *Proceedings of the National Academy of Sciences of the United States of America*, 101(3): 4431-4434.
- Macierzanka, A., and Szelag, H., (2006). Microstructural behaviour of water-in-oil emulsions stabilized by fatty acid esters of propylene glycol and zinc fatty acid salts. *Colloids and Surfaces A Physicochemical Engineering Aspects*, 281: 125-137.
- Mahdi, E.S., Noor, A.M., Sakeena, M.H., Abdullah, G.Z., Abdulkarim, M.F., and Sattar, M.A., (2011). Formulation and in vitro release evaluation of newly synthesized palm kernel oil esters-based nanoemulsion delivery system for 30% ethanolic dried extract derived from local *Phyllanthus urinaria* for skin antiaging. *International Journal Nanomedicine*, 6: 2499-2512.

- Marrink, S. J. and Tieleman, D. P., (2013). Perspective on the MARTINI model, *Chemical Society Reviews*, 42(16): 6801-6822.
- Marrink, S. J., and Mark, A. E., (2004). Molecular view of hexagonal phase formation in phospholipid membranes, *Biophysical Journal*, 87(6): 3894-3900.
- Marrink, S. J., Lindahl, E., Edholm, O., and Mark, A. E., (2001). Simulation of the spontaneous aggregation of phospholipids into bilayers. *Journal of American Chemical Society*, 123: 8638-8639.
- Marrink, S. J., Risselada, H. J., Yefimov, S., Tieleman, D. P., and de Vries A. H., (2007). The MARTINI Force Field: Coarse Grained Model for Biomolecular Simulations. *Journal of Physical Chemistry B*, 111(27): 7812-7824.
- Marrink, S.J., and Mark A. E., (2003). Molecular Dynamics simulation of the formation, structure and dynamics of small phospholipid vesicles. *Journal of the American Chemical Society*, 125(49): 15233-15242.
- Marrink, S.J., and Mark, A. E., (2005). Coarse grained simulation of phase transitions of lipid membranes. *Biophysical Journal*, 88(1); 384A-384A.
- Martinez, J. M., and Martinez, L., (2003). Packing optimization for automated generation of complex system's initial configurations for molecular dynamics and docking. *Journal of Computational Chemistry*, 24(7):819-825.
- Mason, T.G., Wilking, J., Meleson, K., Chang, C., and Graves, S. (2006). Nanoemulsions: formation, structure, and physical properties. *Journal of Physics: Condensed Matter*, 18(41): R635.
- Maxwell, D.S., Tirado-Rives, J., and Jorgensen, W.L., (1995). A comprehensive study of the rotational energy profiles of organic systems by ab initio MO theory, forming a basis for peptide torsional parameters, *Journal of Computational Chemistry*, 16(8): 984-1010.
- Mihailescu, M., Vaswani, R.G., Jardon-Valadez, E., Castro-Roman, F., Freitas, J.-A., Worcester, D.L., Chamberlin, A.R., Tobias, D.J., and White, S.H., (2011). Acyl-chain methyl distributions of liquid-ordered and -disordered membranes. *Biophysical Journal*, 100, 1455-1462.
- Miyamoto, S., and Kollman, P.A., (1992). SETTLE: An Analytical Version of the SHAKE and RATTLE Algorithm for Rigid Water Models. *Journal of Computational Chemistry* 13(8), 952-962.

- Morrow, D. I. J., McCarron, P. A., Woolfson, A. D., and Donnelly, R. F., (2007). Innovative strategies for enhancing topical and transdermal drug delivery. *The Open Drug Delivery Journal*, 1: 36-59.
- Musa, S. H., Basri, M., Fard Masoumi, H. R., Abedi Karjiban, R., Abd Malek, E., Basri, H., and Shamsuddin, A. F., (2013). Formulation optimization of palm kernel oil esters nanoemulsion-loaded with chloramphenicol suitable for meningitis treatment. *Colloids and Surfaces B: Biointerfaces*, 112: 113-119.
- Nagle, J. F., and Tristram-Nagle, S., (2000). Structure of lipid bilayers. *Biochim Biophys Acta* 1469:159-195.
- Ngamcharussrivichai, C.; Totarat, P. and Bunyakiat, K., (2008). Ca and Zn mixed oxide as a heterogeneous base catalyst for transesterification of palm kernel oil. *Applied Catalysis A: General*, 341(1-2): 77-85.
- Nielsen, S. O., Lopez, C.F., Srivivas, G., and Klein, M. L., (2004). Coarse grain models and the computer simulation of soft materials, *Journal of Physics: Condensed Matter*, 16: R481-R512.
- Nokhodchi A., Shokri J., Dashbolaghi A., Hassan-zadeh, D., Ghafourian T., Barzegar-jalali, M., (2003). The enhancement effect of surfactants on the penetration of lorazepam through rat skin. *International Journal of Pharmaceutics*, 250; 359-69.
- Nose, S., (1984). A unified formulation of the constant temperature molecular dynamics methods. *The Journal of Chemical Physics*, 81(1): 511-519.
- Notman, R., den Otter, W. K., Noro, M. G., Briels, W. J., and Anwar J., (2007). The permeability enhancing mechanism of DMSO in ceramide bilayers simulated by Molecular Dynamics. *Biophysical Journal*, 93(6): 2056-2068.
- Notman, R., Noro, M., O'Malley, B., and Anwar, J., (2006). Molecular basis for dimethylsulfoxide (DMSO) action on lipid membranes, *Journal of American Chemical Society*, 128(43): 13982-13983.
- Olive, G., (2006). Analgesic/Antipyretic treatment: ibuprofen or acetaminophen? An update. *Therapie*, 61(2): 151-160.
- Oliver, R. C., Lipfert, J., Fox, D. A., Lo, R. H., Doniach, S., and Columbus, L., (2013). Dependence of Micelle Size and Shape on Detergent Alkyl Chain Length and Head Group. *PLoS ONE*, 8(5): e62488.
- Ordog, R., (2008). PyDeT, a PyMOL plug-in for visualizing geometric concepts around proteins, *Bioinformation*, 2(8):346-347.

- Orsi, M., Sanderson, W., and Essex, W., (2006). Coarse-Grain Modelling of Lipid Bilayers: A Literature Review. Bozen, 15th -19th May, Bozen, Italy.
- Park., E.-S., Cui, Y., Yun B.-J., Ko I.-J., Chi, S.-C., (2005). Transdermal delivery of piroxicam using microemulsions. *Archives of Pharmacal Research*, 28(2): 243-248.
- Parrinello, M., and Rahman, A., (1980). Crystal structure and pair potentials: A Molecular Dynamics study. *Physical Review Letters*, 45: 1196-1199.
- Parrinello, M., and Rahman, A., (1981). Polymorphic transitions in single crystals: A new molecular dynamics method. *Journal of Applied Physics*, 52:7182.
- Parrinello, M., and Rahman, A., (1982). Strain fluctuations and elastic constants. *Journal of Chemical Physics*, 76:2662.
- Peltola, S., Saarinen-Savolainen, P., Kiesvaara, J., Suhonen, T.M., and Urtti, A. (2003). Microemulsions for topical delivery of estradiol. *International Journal of Pharmaceutics*, 254(2): 99-107.
- Prausnitz, M. R., and Langer, R., (2008). Transdermal drug delivery. *Nature Biotechnology*, 26(11): 1261-1268.
- Price, M. L. P., Ostrovsky, D., and Jorgensen, W. L., (2001). Gas-phase and liquid-state properties of esters, nitriles, and nitro compounds with the OPLS-AA force field. *Journal of Computational Chemistry*, 22(13): 1340-1352.
- Qiao, R., Roberts, A. P., Mount, A. S., Klaine, S. J., and Ke P. C., (2007). Translocation of C₆₀ and its derivatives across a lipid bilayer. *Nano Letters*, 7(3): 614-619.
- Rachke, T.M., Tsai, J., and Levitt, M., (2001). Quantification of the hydrophobic interaction by simulations of the aggregation of small hydrophobic solutes in water. *Proceeding of National Academic Science USA*, May 22;98(11):5965-5969.
- Ranade, V. V., and Cannon, J. B., (2011). Drug delivery systems, Third Edition, CRC Press.
- Ratke, L., and Voorhees, P. W., (2002). Growth and coarsening: Ostwald ripening in material processing. Berlin: Springer.

- Raut, S., Bhadoriya, S. S., Uplanchiwar, V., Mishra, V., Gahane, A., and Jain, S. K., (2012). Lecithin organel: a unique micellar system for the delivery of bioactive agents in the treatment of skin aging. *Acta Pharmaceutica Sinica B*, 2(1): 8-15.
- Razaei, M., Basri, M., Raja Abd Rahman, R. N. Z., Salleh, A. B., Chaibakhsh, N., and Abedi Karjiban, R., (2014). Formulation development and optimization of palm kernel oil esters-based nanoemulsions containing sodium diclofenac. *International Journal of Nanomedicine*, 9: 539-548.
- Rhee, Y.-S., Choi, -S., and Park, S.-C., (2001). Transdermal delivery of Ketoprofen using Microemulsions. *International Journal of Pharmaceutics*. 228: 161-170.
- Rosen, M.J. (2004). Micelle formation by surfactants. *Surfactants and Interfacial Phenomena* (pp. 105-177). John Wiley & Sons, Inc.
- Rowe, R. C., Sheskey, P. J., and Quinn, M. E.; (2009). (Eds.), *Handbook of Pharmaceutical Excipients* 6th edition Pharmaceutical Press, London, England.
- Sadurni, N., Solans, C., Azemar, N., and Garcia-Celma, M. J., (2005). Studies on the formation of O/W nano-emulsions, by low-energy emulsification methods, suitable for pharmaceutical applications. *European Journal of Pharmaceutical Sciences*, 26(5): 438-445.
- Saify, Z.S., Ozra, A., and Abdullah, D., (2002). Cineole as Skin Penetration enhancer. *Pakistan Journal Pharmaceutical Sciences*, 13: 29-32.
- Saiz, L., and Klein, M. L., (2002). Computer simulation studies of Model Biological Membranes. *Accounts of Chemical Research*, 35:482-489.
- Salim, N., Basri, M., Abdul Rahman, M. B., Abdullah, D. K., and Basri, H., (2011). Phase behaviour, formation and characterization of Palm-based esters nanoemulsion formula tion containing Ibuprofen. *Journal of Nanomedicine and Nanotechnology*, 2: 113.
- Salim, N., Basri, M., Abdul Rahman, M. B., Abdullah, D. K., and Basri, H., (2012). Modification of palm kernel oil esters nanoemulsions with hydrocolloid gum for enhanced topical delivery of ibuprofen. *International Journal of Nanomedicine*, 7: 4739-4747.
- Schlick, T., (2013). The 2013 nobel prize in chemistry celebrates computations in chemistry and biology, *SIAM News*, 46(10).

- Schmidt, M. W., Baldridge, K. K., Boatz, J. A., Elbert, S. T., Gordon, M.S., Jensen, J.H., Koseki, S., Matsunaga, N., Nguyen, K.A., Su, S.W., Theresa L., Dupuis, M. and Montgomery, J.A. (1993). General atomic and molecular electronic structure system. *Journal of Computational Chemistry*, 14(11): 1347-1363.
- Scott, H.L., (2002). Modeling the lipid component of membranes. *Current Opinion in Structural Biology*, 12:495-502.
- Shaari, N.S., Abedi karjiban, R., and Basri, M., (2014). The Effect of Temperature and Pressure on the Self-assembly of Dipalmitoylphosphatidylcholine Using coarse-Grained Molecular Dynamics. *Journal of Medical and bioengineering*, 3(2), June 2014.
- Shah, P., Bhalodia, D., and Shelat P., (2010). Nanoemulsion: A pharmaceutical review, *Systematic Reviews in Pharmacy*, 1(1): 24-32.
- Shakeel, F., Baboota, S., Ahuja, A., Ali, J., and Shafiq, S., (2009). Celecoxib Nanoemulsion for Transdermal Drug Delivery: Characterization and In Vitro Evaluation. *Journal of Dispersion Science and Technology*, 30: 834-842.
- Shakeel, F., Baboota, S., Ahuja, A., Ali, J., Aqil, M., and Shafiq, S., (2007). Nanoemulsions as vehicles for transdermal delivery of aceclofenac. *American Association of Pharmaceutical Scientists PharmSciTech*, 8(2): E104.
- Sharma, N., Bansai, M., Visht, S., Sharma, P. K., and Kulkarni, G. T., (2010). Nanoemulsion: A new concept of delivery system. *Chronicles of Young Scientists*, 1(2): 2-6.
- Shi, X., Kong, Y., and Goa H., (2008). Coarse grained molecular dynamics and theoretical studies of carbon nanotubes entering cell membrane. *Acta Mechanica Sinica*, 24(2): 161-169.
- Shih, A. Y., Arkhipov, A., Freddolino, P. L., and Schulten, K., (2006). Coarse grained protein-lipid model with application to lipoprotein particles. *Journal of Physical Chemistry B*, 110(8): 3674-3684.
- Shrestha, L. K., Saito, E., Shrestha, R. G., Kato, H., Takase, Y., and Aramaki, K., (2007). Foam stabilized by dispersed surfactant solid and lamellar liquid crystal in aqueous systems of diglycerol fatty acid esters. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 293(1-3): 262-271.
- Singla, V., Saini, S., Joshi, B., and Rana, A.C., (2011). Emulgel: A new platform for topical drug delivery. *International Journal of Pharma and Bio Sciences*, 3(1): 486-498.

- Smith E.W., and Maibach, H.I., (2005). Percutaneous penetration enhancer. 2nd ed. Boca Raton: Taylor and Francis.
- Smith, G. D., and Bedrov, D., (2007). The Relationship between the α - and β -Relaxation Processes in Amorphous Polymers: Insight from Atomistic Molecular Dynamics Simulations of 1,4-Polybutadiene Melts and Blends. *Journal of Polymer Science B: Polymer Physics* 45: 627-643.
- Solans, C., Izquierdo, P., Nolla, J., Azemar, N., and Garcia-Celma, M.J. (2005). Nano-emulsions. *Current Opinion in Colloid & Interface Science*, 10(3-4): 102-110.
- Som, I., Bhatia, K., and Yasir, M., (2012). Status of surfactant as penetration enhancers in transdermal drug delivery. *Journal of Pharmacy & Bioallied Sciences*, 2012 Jan-Mar, 4(1); 2-9.
- Stephenson, B. C., Rangel-Yagui, C. O., Pessoa Junior, A., Tavares, L. C., Beers, K., and Blankschtein, D., (2006). Experimental and theoretical investigation of the micellar-assisted solubilization of ibuprofen in aqueous media. *Langmuir*, 22(4): 1514-1525.
- Subedi, R. K., Oh, S. Y., Chun, M. K., Choi, H. K., (2010). Recent advances in transdermal drug delivery. *Archives of Pharmacal Research*, 33(3): 339-351.
- Sutradhar, K. B., and Amin, M. L., (2013). Nanoemulsions: increasing possibilities in drug delivery. *European Journal of Nanomedicine*, 5(2): 97-110.
- Sznitowska, M., Dbrowska, E. A., and Janicki, S., (2012). Solubilizing potential of submicron emulsions and aqueous dispersions of lecithin. *International Journal of Pharmaceutics*, 246(1-2): 203-206.
- Tadros, T.F., Vandamme, A., Leveck, B., Booten, K., and Stevens, C.V. (2004). Stabilization of emulsions using polymeric surfactants based on inulin. *Advances in Colloid and Interface Science*, 108-109(0): 207-226.
- Takahashi, K., Oda, T., and Naruse, K., (2014). Coarse-grained molecular dynamics simulations of biomolecules. *AIMS Biophysics*, 1(1): 1-15.
- Tcholakova, S., Denkov, N., D., and Danner, T., (2004). Role of surfactant type and concentration for the mean drop size during emulsification in turbulent flow. *Langmuir*, 20: 7444-7458.
- Thomas, L.L., Christakis, T.J., and Jorgensen, W.L., (2006). Conformation of alkanes in the gas phase and pure liquids. *Journal of Physical Chemistry B*, 110(42): 21198-21204.

- Tieleman, D. P., van der Spoel, D., and Berendsen, H. J. C., (2000). Molecular dynamics simulations of dodecylphosphocholine micelles at three different aggregate sizes: micellar structure and chain relaxation. *Journal of Physical Chemistry B*, 104: 6380–6388.
- Tozzini, V., (2005). Coarse-grained models for proteins. *Current Opinion in Structural Biology*, 15: 144-150.
- Tripathi, K. D., (2003). Non steroidal anti inflammatory drugs and anti pyretic analgesics. In: *Essentials of medical pharmacology*. 5th edn., Jaypee Brothers, New Delhi, 176.
- Tsume, Y., Langguth, P., Garcia-Arieta, A., and Amidon, G. L., (2013). In silico prediction of drug dissolution and absorption with variation in intestinal pH for BCS Class II Weak Acid Drugs: Ibuprofen and Ketoprofen, *Biopharmaceutics and Drug disposition*, 33(7): 366-377.
- Ulander, J., and Haymet, A.D.J., (2003). Permeation across Hydrated DPPC Lipid Bilayers: Simulation of the Titrable Amphiphilic Drug Valproic Acid. *Biophysical Journal*, 85, 3475.
- van der Spoel, D., Lindahl, E., Hess, B., Groenhof, G., Mark, A. E., and Berendsen, H.J.C., (2005). GROMACS: Fast, Flexible, and Free. *Journal of Computational Chemistry*, 26: 1701-1718.
- van Gunsteren, W.F., Biller, S.R., Eising, A.A., Hunenberger, P.H., Kruger, P., Mark, A. E., Scott, W. R. P., Tironi, I.G., (1996). Biomolecular Simulation: The GROMOS 96 Manual and User Guide. Vdf Hochschulverlag AG an der ETH Zürich, Zürich, 1-1042.
- van Nieuwenhuyzen, W., and Thomas, M. C., (2008). Update of vegetable lecithin and phospholipid technologies. *European Journal of Lipid Sciences and Technology*, 5: 472–486.
- Venable, R.M., Brooks, .R., Pastor, R.W., (2000). Molecular dynamics simulations of gel (L_{BI}) phase lipid bilayers in constant pressure and constant surface area ensembles. *Journal of Chemical Physics*, 112; 4822-4831.
- Venturoli, M., Smit, B., and Sperotto M., (2005). Simulation studies of protein-induced bilayer deformations, and lipid-induced pprotein tilting, on a mesoscopic model for lipid bilayers with embedded proteins. *Biophysical Journal*, 88: 1778-1798.
- Verlet L (1967) Computer “Experiments” on Classical Fluids. I. Thermodynamical Properties of Lennard-Jones Molecules. *Physical Reviews* 159:98-103.

- Verma, S. K., Shaban, A., Purohit, R., Chimata, M. L., Rai, G., and Verma, O. P., (2012). Immunomodulatory activity of *Withania somnifera* (L.), *Journal of Chemical and Pharmaceutical Research*, 4(1): 559-561.
- Vikas, S., Seema, S., Gurpreet, S., Rana, A. C., and Baibhav, J., (2011). Penetration enhancers: A novel strategy for enhancing transdermal drug delivery. *International Research Journal of Pharmacy*, 2(12): 32-36.
- Vlugt, T. J. H., van der Eerden, J. P. J. M., Dijkstra, M., Smit, B and Frenkel, D., (2008). Introduction to Molecular Simulation and Statistical Thermodynamics, Delft, The Netherland.
- Wallace, E. J., and Sansom, M. S., (2008). Blocking of carbon nanotube based nanoinjectors by lipids: a simulation study. *Nano Letters*, 8(9): 2751-2756.
- Wang, J., Wolf, R. M., Caldwell, J. W., Kollman, P. A., and Case, D. A., (2004). Development and testing of a general amber force field, *Journal of Computational Chemistry*, 25: 1157-1174.
- Wang, L., Dong, J., Chen, J., Eastoe J., and Li X., (2009). Design and optimization of a new self-nanoemulsifying drug delivery system. *Journal Colloid Interface Sciences*, 330: 443-448.
- Watkins, E. K., and Jorgensen, W. L., (2001). Perfluoroalkanes: Conformational Analysis and liquid-state properties from ab Initio and Monte Carlo simulations. *Journal of Physical Chemistry A*, 105(16): 4118-4125.
- Wertz, P., and Noelen, L., (2003). "Confidence Intervals" for the "true" lipid compositions of the human skin barrier?. In: Forslind B, Lindberg M (eds) *Skin, Hair, and Nails Structure and Function*. Marcel Dekker, New York, 85-106.
- Williams, A. C., (2003). Theoretical aspects of transdermal drug delivery. In *Transdermal and Topical Drug Delivery*. Ghosh, T. K., Pfister, W., and Yum, S. I., editors. Pharmaceutical Press: From Theory to Clinical Practice, London. 27-49.
- Winstanley, P., and Walley, T., (2002) Drug for arthritis. In: *Medical pharmacology: a clinical core text for integrated curriculum with self assessment*. Churchill Livingstone, Adinburgh. 105-107.
- Xiang, Y., Zhang, D. W., and Zhang, J. Z., (2004). Fully quantum mechanical energy optimization for protein-ligand structure, *Journal of Computational Chemistry*, 25(12): 1431-1437.

Yuan, Y., Li, S.-M., Mo, F.-K., and Zhong, D.-F., (2006). Investigation of microemulsion system for transdermal delivery of meloxicam. *International Journal of Pharmaceutics*, 321(1-2): 117-123.

Zainol, S., Basri, M., Basri, H. B., Shamsuddin, A. F., Abdul-Gani, S. S., Karjiban, R. A., and Abdul-Malek, E., (2012). Formulation Optimization of a Palm-Based Nanoemulsion System Containing Levodopa. *International Journal of Molecular Sciences*. 13, 13049-13064.

