



UNIVERSITI PUTRA MALAYSIA

***GROUP ACTIONS AND THEIR APPLICATIONS IN ASSOCIATIVE
ALGEBRAS AND ALGEBRAIC STATISTICS***

NADIA FAIQ MOHAMMED

FS 2017 61



**GROUP ACTIONS AND THEIR APPLICATIONS IN ASSOCIATIVE
ALGEBRAS AND ALGEBRAIC STATISTICS**

By

NADIA FAIQ MOHAMMED

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

September 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright ©Universiti Putra Malaysia



DEDICATIONS

*To
my lovely country*

Iraq



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

GROUP ACTIONS AND THEIR APPLICATIONS IN ASSOCIATIVE ALGEBRAS AND ALGEBRAIC STATISTICS

By

NADIA FAIQ MOHAMMED

September 2017

Chair: Professor Isamiddin S. Rakhimov, PhD

Faculty: Science

The study of group action is an exciting and very intensive research that takes place in Mathematics and Physics. There can be different ways for a group to act on different kinds of objects.

This dissertation is mostly concerned with group actions on vector spaces and affine algebraic varieties. It is mainly comprised of three parts. In the first part, we consider an action of associative algebra A on a vector space to study low-dimensional cohomology groups of associative algebras. The origin of cohomology groups is found in algebraic topology. The dimension of the cohomology groups is considered one of the important invariant to study properties of algebras. Particularly, this invariant plays a rigorous role in geometric classification of associative algebras. We focus on the applications of low dimensional cohomology groups $H^i(A, A)$. We start with the zero order cohomology groups of two and three dimensional complex associative algebras. Then, we consider an important special case of derivations so-called inner derivation mapping which can be roughly interpreted as 1-coboundaries on vector space where the algebra A acting. We study some of their properties and give an algorithm to obtain the inner derivations of associative algebras. Another main result of this part is precisely formulated two algorithms for describing the 2-cocycles and 2-coboundaries of A . Using an existing classification result of low dimensional associative algebras, we apply all these algorithms to complex associative algebras up to dimension three.

In the second part, general linear group acts on an affine algebraic variety over an algebraically closed field. This part is devoted to the study of the rigidity of associative algebras. We give necessary invariance arguments for the existence of degenerations

which is helpful in finding out for associative algebras to be rigid. Subsequently, applications of the invariance arguments to the varieties of low-dimensional complex associative algebras are described.

In the last part of the thesis, we consider an action of dihedral group D_p on the rational vector space $\mathbb{Q}^p$ to study an invariance group of Markov basis for some contingency tables. We define a Markov basis B for a $\frac{p(v-1)(p-v)}{2v} \times v \times \frac{p}{v}$ - contingency table with fixed two-dimensional marginals, where p is a multiple of v and greater than or equal to $2v$. Using this action, we find the invariance subgroup H of Markov basis B with respect to the dihedral group D_p . Moreover, an algorithm to obtain the set of all independence models of two-way contingency tables with the same row sums and column sums is proposed. Finally, we introduce a new class of algebras which is closely related to the Markov basis in algebraic statistics.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

TINDAKAN KUMPULAN DAN APLIKASINYA DALAM ALJABAR BERSEKUTU DAN STATISTIK BERALJABAR

Oleh

NADIA FAIQ MOHAMMED

September 2017

Pengerusi: Profesor Isamiddin S. Rakhimov, PhD
Fakulti: Sains

Kajian tindakan kumpulan adalah satu penyelidikan yang menarik dan sangat intensif yang berlaku dalam Matematik dan Fizik. Terdapat cara yang berbeza bagi kumpulan untuk bertindak pada pelbagai jenis objek.

Disertasi ini kebanyakannya berkaitan dengan tindakan kumpulan pada ruang vektor dan pelbagai afin aljabar. Ianya terdiri daripada tiga bahagian utama. Dalam bahagian pertama, kami mempertimbangkan tindakan terhadap aljabar bersekutu A pada suatu ruang vektor untuk mengkaji dimensi rendah kumpulan kohomologi bagi aljabar-aljabar bersekutu. Asal-usul kumpulan kohomologi terdapat dalam topologi aljabar. Dimensi bagi kumpulan-kumpulan kohomologi dianggap sebagai salah satu tak varian yang penting untuk mengkaji sifat-sifat aljabar. Terutamanya, tak varian ini memainkan peranan yang ketat dalam pengelasan bergeometri bagi aljabar-aljabar bersekutu. Kami menumpukan pengaplikasian bagi dimensi rendah kumpulan-kumpulan kohomologi $H^i(A, A)$. Kami mulakan dengan kumpulan kohomologi berperingkat sifar bagi dimensi dua and tiga aljabar-aljabar bersekutu kompleks. Kemudian, kami mempertimbangkan satu kes penting bagi pembezaan yang berasal dari pemetaan pembezaan dalaman yang boleh ditakrifkan secara kasar sebagai $1 -$ kosempadan dalam ruang vektor di mana aljabar tersebut A bertindak. Kami mengkaji beberapa sifat mereka dan memberikan satu algoritma untuk mendapatkan pembezaan-pembezaan dalaman bagi aljabar-aljabar bersekutu. Hasil utama lain dari bahagian ini adalah merumuskan dua algoritma dengan tepat untuk menerangkan $2 -$ kokitar dan $2 -$ kosempadan bagi A . Dengan menggunakan pengelasan sedia ada bagi aljabar-aljabar bersekutu berdimensi rendah, kami menggunakan semua algoritma-algoritma ini ke atas aljabar-aljabar bersekutu kompleks sehingga dimensi empat.

Dalam bahagian kedua, kumpulan linear terturun am bertindak ke atas pelbagai afin aljabar terhadap medan aljabar tertutup. Bahagian ini dikhaskan untuk mengkaji ketegaran aljabar-aljabar bersekutu. Kami memberikan hujah-hujah tak varian yang diperlukan untuk kewujudan kemerosotan yang membantu dalam mencari aljabar-aljabar bersekutu untuk menjadi tegar. Selanjutnya, aplikasi hujah tak varian kepada pelbagai aljabar-aljabar bersekutu kompleks berdimensi rendah diterangkan.

Dalam bahagian akhir tesis ini, kami mempertimbangkan tindakan terhadap kumpulan dwihedron D_p pada ruang vektor nisbah $\mathbb{Q}^p$ untuk mengkaji suatu kumpulan ketak varianan bagi asas Markov untuk beberapa jadual kontingensi. Kami mentakrifkan satu asas Markov B untuk $\frac{p(v-1)(pv)}{2v} \times v \times \frac{p}{v}$ - jadual kontingensi dengan kekerapan dua dimensi yang tetap, di mana p adalah suatu gandaan daripada v dan lebih besar daripada atau sama dengan $2v$. Dengan menggunakan tindakan ini, kami mendapati satu ketak varianan subkumpulan H dari asas Markov B terhadap kumpulan dwihedron D_p . Selain itu, satu algoritma untuk mendapatkan satu set bagi semua model-model kebebasan untuk dua-cara jadual kontingensi dengan jumlah baris dan jumlah lajur yang sama dicadangkan. Akhir sekali, kami memperkenalkan satu kelas baru bagi aljabar-aljabar yang sangat berkait rapat dengan asas Markov dalam statistik beraljabar.

ACKNOWLEDGEMENTS

In the name of Allah, Most Gracious, Most Merciful.

First of all, I would like to thank Allah for giving me the strength, guidance, and patience that enable me to learn and complete this thesis. To him I owe everything. Peace be upon Prophet Mohammed and his honorable followers.

I would like to express my sincere appreciation and deepest gratitude to my supervisor, Professor, Dr. I.S. Rakhimov, chairman of the supervisory committee, for his full support, expert, guidance, valuable time, helpful discussions and his encouragement throughout my research. I will be forever grateful to him for his continued patience. I don't have enough words to express them just to say "Thanks for being my supervisor".

I also want to thank my co-supervisor Associate Professor Dr. Mahendran S.Shitan, for all his guidance, support and helpful discussions. Also, I am grateful to my supervisory committee members Professor, Dr. Adem Kilicman, Senior Lecturers Dr. Athirah Binti Nawawi and Dr. Rabha Waell Ibrahim to all of their help during this process.

This acknowledgement will not be complete if I do not include Dr. Sharifah Kartini Binti Said Husain, senior lecturer of Mathematics Department, UPM for her help and support in this search. Also many thanks to, Dr. Nasir Ganikhodjaev, Professor of Computational and Theoretical Sciences Department, IIUM for his great suggestions during this work.

I would like to take this opportunity to thank the Ministry of Higher Education and Scientific Research, Baghdad University-Iraq for the financial support during my PhD study.

Last but not least, I cannot end without thanking my parents. I owe so much to my dear mother and late father (Allah bless him) who always encouraged me to learn and supported me throughout my life. I am especially grateful to my beloved husband, Ali Naji, for his patience, love, support and for giving me all the time to do my research. I would like to extend my gratitude to our two kids, Mustafa and Sara; they are a great blessing to us. I wish to thank also my brothers, Nabeel and Waseem, and my sister, Sroor, for their encouragement and spiritually support throughout my life.

Finally, I would like to thank everybody who was important to the successful realization of this thesis, as well as expressing my apology that I could not mention personally one by one.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy.

The members of the Supervisory Committee were as follows:

Isamiddin S. Rakhimov, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairperson)

Adem Kılıçman, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Mahendran A/L S.Shitan, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Athirah Binti Nawawi, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Member)

Rabha Wael Ibrahim, PhD

Senior Lecturer
Faculty of Science
University Malaya
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No: Nadia Faiq Mohammed, GS 38684

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xiv
 CHAPTER	
1 INTRODUCTION	1
1.1 Introduction	1
1.2 Basic concepts	1
1.3 Motivation	7
1.4 Objectives of the research	8
1.5 Methodology	8
1.6 Organization of thesis	10
 2 LITERATURE REVIEW	12
2.1 Introduction	12
2.2 Associative algebras	12
2.3 The classification of rigid associative algebras	13
2.4 Algebraic statistics	15
2.5 Conclusion	25
 3 COHOMOLOGY OF LOW-DIMENSIONAL COMPLEX ASSOCIATIVE ALGEBRAS	26
3.1 Introduction	26
3.2 Cohomology of associative algebras	26
3.3 The zero-cohomology spaces	28
3.3.1 Zero-cohomology spaces of two-dimensional associative algebras	28
3.3.2 Zero-cohomology spaces of three-dimensional associative algebras	30
3.4 The first-cohomology spaces	32
3.5 An algorithm for finding inner derivations	34
3.5.1 Inner derivations of two-dimensional associative algebras	35
3.5.2 Inner derivations of three-dimensional associative algebras	38
3.6 The second-cohomology spaces	41
3.7 An algorithm for finding 2-cocycles	41

3.7.1	Cocycles of two-dimensional associative algebras	43
3.7.2	Cocycles of three-dimensional associative algebras	47
3.8	An algorithm for finding 2-coboundaries	60
3.8.1	Coboundaries of two-dimensional associative algebras	61
3.8.2	Coboundaries of three-dimensional associative algebras	65
3.9	Conclusion	70
4	GEOMETRIC CLASSIFICATIONS OF LOW-DIMENSIONAL ASSOCIATIVE ALGEBRAS	72
4.1	Introduction	72
4.2	On rigidity of associative algebras	72
4.3	Invariance argument	74
4.4	Applications of the invariance argument to the varieties of low-dimensional associative algebras	76
4.4.1	The variety of two-dimensional associative algebras	76
4.4.2	The variety of three-dimensional associative algebras	79
4.5	Conclusion	93
5	INVARIANCE GROUP FOR SOME CONTINGENCY TABLES	94
5.1	Introduction	94
5.2	Markov bases and toric ideals for three-way contingency tables with fixed two-dimensional marginals	94
5.3	Invariance group for three-way contingency tables with fixed two-dimensional marginals	99
5.4	Algebraic method for independence model of two-way contingency tables	107
5.4.1	The proposed method	107
5.4.2	Application	109
5.5	Markov Algebras	112
5.6	Conclusion	115
6	CONCLUSION AND FUTURE RESEARCH	116
6.1	Conclusion	116
6.2	Future Research	117
	BIBLIOGRAPHY	118
	APPENDICES	120
	BIODATA OF STUDENT	129
	LIST OF PUBLICATIONS	130

LIST OF TABLES

Table	Page
3.1 The inner derivations of two-dimensional associative algebras	36
3.2 The inner derivations of three-dimensional associative algebras	39
3.3 2–cocycles for two-dimensional associative algebras	44
3.4 2–cocycles for three-dimensional associative algebras	56
3.5 2–coboundaries for two-dimensional associative algebras	62
3.6 2–coboundaries for three-dimensional associative algebras	66
4.1 Contraction invariants of two-dimensional associative algebras	77
4.2 The proposed contraction invariants of two-dimensional associative algebras	77
4.3 The (α, β, γ) -derivations invariants of two-dimensional associative algebras	78
4.4 All possibilities of degenerations of two-dimensional associative algebras	78
4.5 Contraction invariants of three-dimensional associative algebras	80
4.6 The proposed contraction invariants of three-dimensional associative algebras	81
4.7 The (α, β, γ) -derivations of three-dimensional associative algebras	82
4.8 All possibilities of degenerations of three-dimensional associative algebras	88
5.1 $v \times \frac{p}{v}$ - contingency table	95
5.2 $I \times J$ - contingency table to show the leading variables and free variables	109
5.3 3×2 - contingency table to show the relationship between agriculture's professional qualifications and their perception of secondary school agriculture	110
5.4 3×2 - contingency table to show the leading variables and free variables	111
5.5 Multiplication table of basis elements of 2-dimensional Markov algebra	114
5.6 Multiplication table of basis elements of 3-dimensional Markov algebra	115

LIST OF FIGURES

Figure	Page
2.1 Undirected graph for $t = (3, 2, 2, 2, 1)$	23
4.1 Degenerations of 2-dimensional associative algebras	79
4.2 Degenerations of 3-dimensional associative algebras	91
5.1 Bipartite graph	98
5.2 The regular polygon with p vertices	99
5.3 The lattice of subgroups of dihedral group D_6	104
5.4 Basic Markov basis elements for 2×3 -contingency table	104
5.5 Basic Markov basis elements for 3×2 -contingency table	106
A.1 The fiber of all 3×2 -contingency tables with the same row sums and column sums	127
A.2 A connected graph $G_{(t,B)}$ for given t	128

LIST OF ABBREVIATIONS

As_p^q	q^{th} isomorphism class of associative algebras in dimension p
$Der(A)$	The algebra of all derivations of the algebra A
$In(A)$	The algebra of all inner derivations of the algebra A
IC	Isomorphism classes of algebras
$\mathbb{N}$	The set of natural numbers
$\mathbb{Z}$	The set of integer numbers
$\mathbb{Q}$	The set of rational numbers
$\mathbb{R}$	The set of real numbers
$\mathbb{C}$	The set of complex numbers
$\mathbb{K}$	Field
$Z^1(A, D)$	1-cocycles
$Z^2(A, D)$	2-cocycles
$B^1(A, D)$	1-coboundaries
$B^2(A, D)$	2-coboundaries
□	The end of a proof

CHAPTER 1

INTRODUCTION

1.1 Introduction

Group action is very exciting and intensive research that takes place in Mathematics and Physics. Symmetries coming from a group action appearing in such a construction are a powerful tool for understanding the structure of geometric objects. There can be different ways for a group to act on different kinds of objects, such as vector spaces and affine algebraic varieties. It will be an interesting topic to study the applications of group actions on these special cases which will lead to resolve some of the issues in relation to cohomology of associative algebras, geometric classification of associative algebras and invariance groups of generating sets for a toric ideal.

In this thesis, an associative algebra A acts on a vector space to study low-dimensional cohomology groups of associative algebras $H^i(A, A)$, where $i = 0, 1, 2$. The dimension of the cohomology groups is considered one of the important invariant to study properties of algebras. Particularly, this invariant plays a rigorous role in geometric classification of associative algebras. Then, a general linear group acts on an affine algebraic variety over an algebraically closed field to investigate the affine algebraic varieties of low dimensional complex associative algebras in relation to the of geometric classification. Of course, it is interesting aspect but difficult problem to determine the number of irreducible components of an algebraic variety. In order to identify solutions to such a problem, a degeneration approach can be considered as helpful. Finally, a new area which has been given the name "algebraic statistics" is concerned with independence models for contingency tables that can be described, in some way, via polynomials. Algebraic statistics is concerned with the development of techniques in algebraic geometry, commutative algebra, and combinatorics, to address problems in statistics and its applications. There is a wide array of problems for which this approach has been highly successful such as optimization, computational biology, the maximum likelihood estimation and parametric inference. One of the main objects which is considered in algebraic statistics is toric ideal. In this regard, a dihedral group acts on the rational vector space $\mathbb{Q}^P$ to study an invariance group of generating set for a toric ideal of the so-called Markov basis for some contingency tables.

1.2 Basic concepts

In this section, we introduce some well-known definitions along with associated concepts, which have implications for the present study. Readers who are familiar with related materials in the area may attest to the wealth of information in relation to Lie algebras, which can be found even in standard books such as Jacobson (1962). In the case of associative algebras, most of the notations and along with related concepts are

referred to Pierce (1982). Finally, the main notions of algebraic geometry are provided in Hartshorne (2013).

Definition 1.1 Let $\mathbb{K}$ be a field, V be a vector space over $\mathbb{K}$ with a binary operation $\lambda : V \times V \longrightarrow V$. If the binary operation is bilinear, i.e.,

$$\lambda(\alpha_1 x + \alpha_2 y, z) = \alpha_1 \lambda(x, z) + \alpha_2 \lambda(y, z),$$

$$\lambda(z, \alpha_1 x + \alpha_2 y) = \alpha_1 \lambda(z, x) + \alpha_2 \lambda(z, y),$$

where $x, y, z \in V$ and $\alpha_1, \alpha_2 \in \mathbb{K}$ then $A = (V, \lambda)$ is said to be an algebra over $\mathbb{K}$.

For ease of reference, we first introduce some classes of algebras, which are closely related to the ones reported in the present study. The definitions and examples reported in the section are mostly well-known. In this regard we attempt to help the readers to have a better understanding of these definitions and examples in relation to the present study.

Definition 1.2 An associative algebra A is a vector space over a field $\mathbb{K}$ equipped with bilinear map

$$\lambda : A \times A \rightarrow A$$

satisfying the associative law:

$$\lambda(\lambda(x, y), z) = \lambda(x, \lambda(y, z)), \quad \forall x, y, z \in A.$$

Further the notations $x \cdot y$ (even just xy) will be used for $\lambda(x, y)$. Note that fields, polynomial algebras, set of linear transformations on a vector space (quadratic matrices over a fixed field) are simple examples of associative algebras.

Let A be an n -dimensional algebra. By taking a basis $\{e_1, e_2, \dots, e_n\}$ in A , elements x and y of A can be represented as follows:

$$x = \sum_{i=1}^n x^i e_i \quad \text{and} \quad y = \sum_{j=1}^n y^j e_j.$$

Notice that the product $x \cdot y$ of any two elements x and y in A , is completely determined by the product $e_i \cdot e_j$ of pairs of basis elements, that is:

$$x \cdot y = \sum_{i,j=1}^n x^i y^j e_i \cdot e_j.$$

Since $e_i \cdot e_j \in A$, the product $e_i \cdot e_j$ is expanded with respect to the basis $\{e_1, e_2, \dots, e_n\}$:

$$e_i \cdot e_j = \sum_{k=1}^n \gamma_{ij}^k e_k, \quad i, j = 1, 2, \dots, n.$$

The numbers γ_{ij}^k are called the structure constants of A with respect to the basis $\{e_1, e_2, \dots, e_n\}$. It is worth highlighting that any algebra is completely determined by its structure constants.

Definition 1.3 A Lie algebra L is a vector space over a field $\mathbb{K}$ equipped with bilinear map

$$[\cdot, \cdot] : L \times L \rightarrow L$$

satisfying the following conditions:

- (1) $[x, x] = 0, \forall x \in L$
- (2) $[x, y], z] + [y, z], x] + [z, x], y] = 0, \forall x, y, z \in L$

Condition (2) is known as the Jacobi identity. As the Lie bracket $[\cdot, \cdot]$ is bilinear, we have

$$0 = [x + y, x + y] = [x, x] + [x, y] + [y, x] + [y, y] = [x, y] + [y, x].$$

The condition (1) implies

$$(1') [x, y] = -[y, x] \quad \forall x, y \in L \quad (\text{anti-symmetry}).$$

Example 1.1 Any vector space V has a Lie bracket defined by $[x, y] = 0$ for all $x, y \in V$. In particular, the field $\mathbb{K}$ may be regarded as a one-dimensional abelian Lie algebra.

Most of the readers are aware of the following fact which plays a fundamental role in the classification.

Let A be a associative algebra with a bilinear map λ . Then, the law μ defined by

$$\mu(x, y) = \lambda(x, y) - \lambda(y, x), \quad \forall x, y \in A$$

is a Lie law.

Let $\mathbb{K}[X] = \mathbb{K}[x_1, \dots, x_n]$ be the polynomial ring with n indeterminants over algebraically closed field $\mathbb{K}$. Let $\mathbb{A}^n = \{P = (a_1, \dots, a_n) : a_i \in \mathbb{K}\}$ be the affine n -space over $\mathbb{K}$. We will interpret the elements of $\mathbb{K}[X]$ as functions from the affine n -space to $\mathbb{K}$, by defining $f(P) = f(a_1, \dots, a_n)$, where $f \in \mathbb{K}[X]$ and $P \in \mathbb{A}^n$.

Definition 1.4 Let S be any subset of $\mathbb{K}[X]$, we define the zero set of S to be the common zeros of all elements of S :

$$V(S) = \{P \in \mathbb{A}^n : f(P) = 0, \forall f \in S\}.$$

A subset Y is called an algebraic set if there exists a subset $S \subset \mathbb{K}[X]$ such that $Y = V(S)$.

Upon the discussion, it is necessary to introduce as earlier as possible a topology on the algebraic set in the present study, which is called Zariski topology.

Proposition 1.1

- (i) The affine n -space $\mathbb{A}^n$ and the empty set are algebraic sets.
- (ii) The union of two algebraic sets is algebraic.
- (iii) The intersection of any family of algebraic sets is algebraic.

The meaning of the proposition is that if we take the algebraic sets as closed sets, we get a topology on $\mathbb{A}^n$, which is referred to as Zariski topology. If X is any algebraic set, the Zariski topology on X is the topology induced on it from $\mathbb{A}^n$. Closed/open sets in X are intersections of X with closed/open sets in $\mathbb{A}^n$.

Definition 1.5 A nonempty subset Y of a topological space X is irreducible if it cannot be expressed as a union $Y = Y_1 \cup Y_2$ of two proper subsets, each of them is closed in Y .

Definition 1.6 An affine algebraic variety or simply affine variety is an irreducible closed subset of $\mathbb{A}^n$ (with the induced topology).

The next step is to explore the correspondence identified by subsets of $\mathbb{A}^n$ and ideals in $\mathbb{K}[X]$.

Definition 1.7 Let Y be any subset of $\mathbb{A}^n$. Let

$$I(Y) = \{f \in \mathbb{K}[X] : f(a_1, \dots, a_n) = 0, \forall (a_1, \dots, a_n) \in Y\}.$$

It is easy to see that $I(Y)$ is an ideal of $\mathbb{K}[X]$. If Y is affine variety, then this ideal is called defining ideal of the affine variety Y .

Definition 1.8 Let Y be an affine variety in $\mathbb{A}^n$. A function $f : Y \rightarrow \mathbb{K}$ is said to be regular if it coincides with the restriction on Y of some polynomial function, i.e., a function $\mathbb{A}^n \rightarrow \mathbb{K}$ mapping a point to $F(a)$, where $F \in \mathbb{K}[X]$.

Definition 1.9 Let X and Y be affine varieties. The function $\phi : X \rightarrow Y$ is called morphism (or regular map) if $f \circ \phi : X \rightarrow \mathbb{K}$ is regular for each regular function $f : Y \rightarrow \mathbb{K}$.

Example 1.2 Let ϕ_i , where $1 \leq i \leq m$, be polynomials in $\mathbb{K}[X]$ then a function $\phi : A^n \rightarrow A^m$ defined by $\phi(P) = (\phi_1(P), \phi_2(P), \dots, \phi_m(P))$ is a morphism.

Definition 1.10 An algebraic group is an algebraic variety G equipped with the structure of a group, such that the multiplication map and the inverse map are morphisms of varieties.

Example 1.3 The general linear group $GL_n(\mathbb{C})$ consisting of all invertible $n \times n$ matrices with complex coefficients, is the open subset of the space M_n of $n \times n$ complex matrices (an affine space of dimension n^2) whose determinant is nonzero. Then $GL_n(\mathbb{C})$ is an algebraic group.

Definition 1.11 An action of algebraic group G on a variety Z is a morphism $\sigma : G \times Z \rightarrow Z$ with

- (i) $\sigma(e, z) = z$, where e is the unit element of G and $z \in Z$.
- (ii) $\sigma(g, \sigma(h, z)) = \sigma(gh, z)$, for any $g, h \in G$ and $z \in Z$.

We shortly write gz for $\sigma(g, z)$, and all call Z a G -variety.

Definition 1.12 Let $A = (V, \lambda)$ and $A_1 = (V, \lambda_1)$ be two algebras over a field $\mathbb{K}$. A linear mapping $\psi : A \rightarrow A_1$ is a homomorphism if

$$\psi(\lambda(x, y)) = \lambda_1(\psi(x), \psi(y)), \quad \forall x, y \in A.$$

Remark 1.1

1. The mapping ψ is an isomorphism if it is a bijective homomorphism (one to one and onto).
2. The mapping ψ is an endomorphism if ψ is a homomorphism and $A = A_1$.
3. The mapping ψ is an automorphism if ψ is an isomorphism and $A = A_1$.

Let V be a vector space of dimension n over an algebraically closed field $\mathbb{K}$ ($\text{char } \mathbb{K} = 0$). Bilinear maps $V \times V \rightarrow V$ form a vector space $\text{Hom}(V \otimes V, V)$ of dimension n^3 , which can be considered together with its natural structure of an affine algebraic variety over $\mathbb{K}$ and denoted by $\text{Alg}_n(\mathbb{K}) \cong \mathbb{K}^{n^3}$. An n -dimensional algebra A over $\mathbb{K}$ can be an element $\lambda(A)$ of $\text{Alg}_n(\mathbb{K})$ via the bilinear mapping $\lambda : A \otimes A \rightarrow A$. Let $\{e_1, e_2, \dots, e_n\}$ be a basis of the algebra A . Then the table of multiplication of A is represented by point (γ_{ij}^k) of affine space as follows:

$$\lambda(e_i, e_j) = \sum_{k=1}^n \gamma_{ij}^k e_k.$$

The general linear group $GL_n(\mathbb{K})$ acts on $Alg_n(\mathbb{K})$ by:

$$(g * \lambda)(x, y) = g(\lambda(g^{-1}(x), g^{-1}(y))).$$

Let $O(\lambda)$ be the set of laws isomorphic to λ . It is called an orbit of λ with respect to the actions of $GL_n(\mathbb{K})$. Two algebras λ_1 and λ_2 are isomorphic if and only if they belong to the same orbit under this action.

Let $A_n(\mathbb{K})$ be a subvariety of $Alg_n(\mathbb{K})$ consisting of all n -dimensional associative algebras over a field $\mathbb{K}$. It is stable under the above mentioned action of $GL_n(\mathbb{K})$. As a subset of $Alg_n(\mathbb{K})$ the set $A_n(\mathbb{K})$ is specified by the system of polynomial equations with respect to the structure constants $\{\gamma_{ij}^k\}$:

$$\sum_{l=1}^n (\gamma_{ij}^l \gamma_{lk}^s - \gamma_{il}^s \gamma_{jk}^l) = 0, \text{ where } i, j, k, s = 1, \dots, n.$$

Definition 1.13 A derivation of associative algebra A over $\mathbb{K}$ is a linear transformation $d : A \rightarrow A$ where

$$d(x \cdot y) = d(x) \cdot y + x \cdot d(y), \quad \forall x, y \in A.$$

We denote the set of all derivations of an associative algebra A by $Der(A)$. The $Der(A)$ is considered as a Lie algebra with respect to the bracket $[d_1, d_2] = d_1 \circ d_2 - d_2 \circ d_1$.

Moving on, we now define an important special case of the derivation mapping, which is widely referred to the inner derivations.

Definition 1.14 Let A be an associative algebra over $\mathbb{K}$ and z be a vector in A . Let $ad_z : A \rightarrow A$ be a function defined as follows:

$$ad_z(x) = xz - zx, \quad \forall x \in A.$$

The function ad_z is called an inner derivation of A .

The set of all inner derivations of an associative algebra A is denoted by $Inn(A)$. The set $Inn(A)$ is an ideal of $Der(A)$.

Definition 1.15 Let A be an associative algebra over $\mathbb{K}$. We define

$$A^1 = A, A^k = \lambda(A, A^{k-1}) \quad (k > 1).$$

The series

$$A^1 \supseteq A^2 \supseteq A^3 \supseteq \dots$$

is called the descending central series of A . If there exists an integer $s \in \mathbb{N}$, such that

$$A^1 \supseteq A^2 \supseteq \cdots \supseteq A^s = \{0\},$$

then the associative algebra A is said to be nilpotent. The smallest integer s for which $A^s = \{0\}$ is called nilpotency rank of A and denoted by $r_n(A)$.

Example 1.4

- (1) Every abelian algebra is nilpotent with nilpotency rank equals to 2.
- (2) The space of n -square matrices $M = [a_{ij}]$ on $\mathbb{K}$ such that $a_{ij} = 0$ for $i \geq j$, with the usual matrix multiplication is a nilpotent associative algebra over $\mathbb{K}$ with nilpotency rank equals to n .

1.3 Motivation

The motivation to carry out the present study was sparked by the applications of group actions on vector spaces, affine algebraic varieties and generating set of toric ideal to resolve some of the issues in relation to cohomology of associative algebras, geometric classification of associative algebras and invariance groups of generating sets for a toric ideal, respectively. These problems can be summarized as follows:

On cohomology of associative algebras:

In algebra, the dimension of the cohomology groups is considered as one of the important invariants in studying the geometric classification. It is for such a reason that the researcher in the present study, consider an action of associative algebra A on a vector space to study low-dimensional cohomology groups $H^i(A, A)$ of associative algebras, where $i = 0, 1, 2$.

On geometric classification of associative algebras:

It is interesting aspect but difficult problem to determine the number of irreducible components of an algebraic variety. However, if one interested in finding the dimension of the algebraic variety then degeneration approach can be considered as helpful. In order to identify solutions to such a problem, a general linear group reportedly acts on an affine algebraic variety over an algebraically closed field.

On invariance groups of generating set for a toric ideal:

As it has been indicated in the literature, one of the main objects, which is considered in algebraic statistics is toric ideal. In this regard, we of the present study look into the invariance groups of generating set for a toric ideal of the so-called Markov basis for some contingency tables. In doing so, we consider an action of dihedral group D_p on the rational vector space $\mathbb{Q}^p$ to describe these invariance groups.

1.4 Objectives of the research

The main objectives of this research are:

1. To propose an algorithm to describe the inner derivations of associative algebras, which can be roughly interpreted as 1-coboundaries of the first cohomology groups, and give another two algorithms for describing the 2-cocycles and 2-coboundaries of the second cohomology groups. The implementation of these algorithms is then provided in low dimensional complex associative algebras.
2. To investigate the affine algebraic varieties of low dimensional complex associative algebras in relation to the geometric classification.
3. To describe the invariance groups of generating set for a toric ideal so-called Markov basis for some contingency tables.
4. To propose an algebraic method to obtain the set of all independent models of two-way contingency tables with the same row sums and column sums, which is referred to as fiber in algebraic statistics.
5. To define and study a new class of algebras which is closely related to the Markov basis in algebraic statistics.

1.5 Methodology

- First Objective: The algorithm to describe the inner derivations of associative algebras, which can be roughly interpreted as 1-coboundaries of the first cohomology groups, involves on solving a system of linear algebraic equations:

$$d_{ji} = \sum_{t=1}^n a_t \gamma_{it}^j - \sum_{t=1}^n a_t \gamma_{ti}^j \quad \text{for } i, j, t = 1, 2, \dots, n. \quad (1.1)$$

To derive this system, fix a basis $\{e_1, e_2, \dots, e_n\}$ of A and a vector $z = a_1 e_1 + a_2 e_2 + \dots + a_n e_n$ in A , where A is an n -dimensional associative algebra. Suppose $ad_z : A \rightarrow A$ be a linear transformation defined as follows:

$$ad_z(x) = xz - zx, \quad \forall x \in A. \quad (1.2)$$

On other hand, ad_z can be considered as a linear transformation of A that is represented in a matrix form

$$ad_z = (d_{ij}), \quad i, j = 1, 2, \dots, n. \quad (1.3)$$

Combining and simplify the equations (1.2) and (1.3), then we get the system of linear algebraic equations (1.1).

The remaining algorithms of the first objective can be done in a similar manner as shown above.

- Second Objective: We take a complete list of nonisomorphic associative algebras of a fixed dimension. For each member of this list we calculate invariance argument that concern necessary criteria of degeneration. Then, for each pair of algebras from the list we test possible existence of degeneration with the necessary criteria of degeneration via comparing the calculated invariance argument.
- Third Objectives: We consider a dihedral group D_p of degree p , where p is a positive integer greater than or equal to 3, acting on the p - dimensional vector space $\mathbb{Q}^p$ over the field of rational numbers $\mathbb{Q}$. Suppose $B \subset \mathbb{Q}^p$, where B is a Markov basis for $\frac{p(v-1)(p-v)}{2v} \times v \times \frac{p}{v}$ - contingency tables with fixed two dimensional marginal. Let H be the largest invariance subgroup of the dihedral group D_p for the Markov basis B , i.e., $gB = B$ for all $g \in H$. Then, for each subgroup of D_p we check possible existence of a largest invariance subgroup H of the dihedral group D_p for the Markov basis B . As a result, it is worthy of note that the largest invariance subgroup H of the dihedral group D_p is generated by $\langle sr, r^d \rangle$ where d equals to the number of columns of two-way contingency tables. Additionally, the order of H is $2 \times$ (the number of rows of two-way contingency tables).
- Fourth Objectives: We intend to solve the following non-homogeneous linear system of equations:

$$A\mathbf{x} = \mathbf{t} \quad (1.4)$$

where A is a configuration matrix and has size $v \times p$, $\mathbf{x}$ is a $I \times J$ two-way contingency table that can be represented as p -dimensional column frequency vector and $\mathbf{t}$ is a v - dimensional column vector. This system is regarded underdetermined system because it consists of a set of $(I+J-1)$ linear equations in $(I \times J)$ variables. Let x_{ij} , $i = 1, 2, \dots, I$, $j = 1, 2, \dots, J$, be the observed cell. Furthermore, let the leading variables be the frequency of cells in the first row and in the first column denoted as x_{11}, x_{k1}, x_{1l} and the free variables be the other frequency of cells in the contingency table $\mathbf{x}$ and denoted by

$$x_{kl} = d_{(k-1)(l-1)}, \quad k = 2, \dots, I, \quad l = 2, \dots, J. \quad (1.5)$$

To obtain the set of all independent models of two-way contingency tables with the same row sums and column sums, we propose an algebraic method that is consisted of four steps. The summarized steps for this method can be found in section 5.4.

- **Fifth Objectives:** To define a new class of algebras which is closely related to the Markov basis in algebraic statistics, let a fiber F_t be the set of all $I \times J$ two-way contingency tables with the same row sums and column sums such that $|F_t|$ represents the number of tables in a fiber F_t . Using the Markov basis B , we can construct a connected, aperiodic and symmetric Markov chain with the transition probability matrix $R = (r_{ij})$ of size $|F_t| \times |F_t|$. The new algebra A is called Markov algebra if their structure constants determine the multiplication table in A via the following rule:

$$\lambda(e_i, e_j) = \sum_{k=1}^n p_{ij,k} e_k, \quad p_{ij,k} = \frac{(r_{ik} + r_{jk})}{2}, \quad \forall i, j, k = 1, \dots, |F_t|.$$

1.6 Organization of thesis

The thesis contains six chapters. Now, we briefly describe the layout of the thesis.

In Chapter 1, as it was highlighted earlier, we discuss some basic notations, affine variety and Zarisk topology which is used throughout the thesis.

In Chapter 2, we provide a brief review on algebraic and geometric classification of associative algebras and some basic introduction on algebraic statistics are given.

In Chapter 3, we focus on applications of low dimensional cohomology groups $H^i(A, A)$. It begins with the zero order cohomology groups of low dimensional complex associative algebras. Moving on, we consider an important special case of derivations, so-called inner derivation mapping, which can be interpreted as 1-coboundaries on vector space where the algebra A acting. We look into some of their properties and give an algorithm to obtain the inner derivations of associative algebras. Another significant aspect here is the precisely formulated two algorithms for describing the 2-cocycles and 2-coboundaries of A . We apply all these algorithms on low dimensional complex associative algebras.

In Chapter 4, we investigate the affine algebraic varieties of low-dimensional complex associative algebras in the sense of geometric classification. For this reason, we consider an action of general linear group on an affine algebraic variety over an algebraically closed field. We provide necessary invariance arguments for the existence of degenerations, which is helpful to discover that the associative algebras are rigid.

Moreover, applications of the invariance arguments to the varieties of low-dimensional complex associative algebras are described.

In Chapter 5, we deal with a generating sets for a toric ideals so-called Markov basis. Therefore, an action of dihedral group on the rational vector space to describe the invariance group of some contingency tables is considered. Additionally, we propose an algebraic method, which serves to obtain the set of all independence models of two-way contingency tables with the same row sums and column sums, which is referred as fiber in algebraic statistics. Finally, we introduce a new class of algebras which is closely related to the Markov basis in algebraic statistics.

In Chapter 6, we summarize and conclude our research. We also give future research problems in these areas.

BIBLIOGRAPHY

- Abbass, H. H. and Hussein, H. M. (2014). On toric ideals for $\frac{n^2-3n}{3} \times 3 \times \frac{n}{3}$ -contingency tables with fixed two dimensional marginals, n is a multiple of 3. *European Journal of Scientific Research*, 123:83–98.
- Agresti, A. (2002). *Categorical data analysis*. 2nd edn. Wiley & Sons, Inc., Hoboken, New Yourk.
- Aoki, S. (2016). An introduction to computational algebraic statistics. *arXiv preprint arXiv:1607.07600*.
- Aoki, S., Hara, H., and Takemura, A. (2012). *Markov bases in algebraic statistics*, volume 199. Springer Science & Business Media.
- Aoki, S. and Takemura, A. (2003). Minimal basis for a connected markov chain over $3 \times 3 \times k$ contingency tables with fixed two-dimensional marginals. *Australian & New Zealand Journal of Statistics*, 45(2):229–249.
- Aoki, S., Takemura, A., and Yoshida, R. (2008). Indispensable monomials of toric ideals and markov bases. *Journal of Symbolic Computation*, 43(6-7):490–507.
- Basri, W. (2014). *Classification and derivations of low-dimensional complex dialgebras*. PhD thesis, Universiti Putra Malaysia Malaysia.
- Burde, D. (2005). Degenerations of 7-dimensional nilpotent lie algebras. *Communications in Algebra*, 33(4):1259–1277.
- Diaconis, P. and Sturmfels, B. (1998). Algebraic algorithms for sampling from conditional distributions. *The Annals of statistics*, 26(1):363–397.
- Drton, M., Sturmfels, B., and Sullivant, S. (2008). *Lectures on algebraic statistics*, volume 39. Springer Science & Business Media.
- Dummit, D. S. and Foote, R. M. (2004). *Abstract algebra*, volume 3. Wiley Hoboken.
- Fiidow, M., Rakhimov, I., and Husain, S. S. (2015). Centroids and derivations of associative algebras. In *Research and Education in Mathematics (ICREM7), International Conference on*, pages 227–232. IEEE.
- Gabriel, P. (1974). Finite representation type is open. In *Proceedings of ICRAI, Ottawa, Lect. Notes in Math.*, volume 488, pages 132–155. Springer.
- Gerstenhaber, M. (1964). On the deformation of rings and algebras. *Annals of Mathematics*, pages 59–103.
- Gilbert, W. J. and Nicholson, W. K. (2004). *Modern algebra with applications*, volume 66. John Wiley & Sons.
- Grunewald, F. and O’Halloran, J. (1988). Varieties of nilpotent lie algebras of dimension less than six. *Journal of Algebra*, 112(2):315–325.
- Guichard, D. (2014). An introduction to combinatorics and graph theory. *Creative Commons*.
- Hartshorne, R. (2013). *Algebraic geometry*, volume 52. Springer Science & Business Media.

- Hazlett, O. C. (1916). On the classification and invariantive characterization of nilpotent algebras. *American Journal of Mathematics*, 38(2):109–138.
- Hrivnák, J. (2007). *Invariants of Lie algebras*. PhD thesis, Faculty of Nuclear Science and Physical Engineering, Czech Technical University, Prague.
- Inonu, E. and Wigner, E. P. (1953). On the contraction of groups and their representations. *Proceedings of the National Academy of Sciences*, 39(6):510–524.
- Jacobson, N. (1962). *Lie algebras*. Dover Publications, Inc. New York.
- Makhlouf, A. (1993). The irreducible components of the nilpotent associative algebras. *Revista Mat. de la Univ. Compl. de Madrid*, 6(1):27–40.
- Makhlouf, A. (2007). A comparison of deformations and geometric study of varieties of associative algebras. *International journal of mathematics and mathematical sciences*, 2007.
- Makhlouf, A. and Goze, M. (1990). Classification of rigid associative algebras in low dimensions. *Hermann, Paris, France*, 50:5–22.
- Mazzola, G. (1979). The algebraic and geometric classification of associative algebras of dimension five. *manuscripta mathematica*, 27(1):81–101.
- Mazzola, G. (1980). Generic finite schemes and hochschild cocycles. *Commentarii Mathematici Helvetici*, 55(1):267–293.
- Muchiri, J. M., Odilla, G. A., Kathuri, N., and Kiriungi, L. (2013). Agriculture teachers perception of secondary school agriculture: A case of meru central district, kenya. *Middle-East Journal of Scientific Research*, 17(4):534–538.
- Nesterenko, M. and Popovych, R. (2006). Contractions of low-dimensional lie algebras. *Journal of mathematical physics*, 47(12):123515.
- Nijenhuis, A., Richardson, R. W., et al. (1966). *Cohomology and deformations in graded Lie algebras*, volume 72. Bulletin of the American Mathematical Society.
- Pierce, B. (1881). Linear associative algebra. *American Journal of Mathematics*, 4(1):97–229.
- Pierce, R. S. (1982). Associative algebra. In *Associative Algebras*. Springer- Verlag New Yourk Inc.
- Pistone, G., Riccomagno, E., and Wynn, H. P. (2000). *Algebraic statistics: computational commutative algebra in statistics*. CRC Press.
- Poonen, B. (2008). Isomorphism types of commutative algebras of finite rank. *Computational arithmetic geometry*, 463:111–120.
- Rakhimov, I., Husain, S. K. S., and Abdulkadir, A. (2015). On generalized derivations of finite dimensional associative algebras. *FEIIC International Journal of Engineering and Technology*, 13(2):121–126.
- Rakhimov, I., Rikhsiboev, I., and Basri, W. (2009). Complete lists of low dimensional complex associative algebras. *ArXiv: math.0910.0932v2, [math.RA]*.
- Segal, I. E. et al. (1951). A class of operator algebras which are determined by groups. *Duke Mathematical Journal*, 18(1):221–265.

Sullivant, S. (2006). Statistical models are algebraic varieties. *Posted at <http://www.math.harvard.edu/~seths/lecture1.pdf>*.

Takemura, A. and Aoki, S. (2004). Some characterizations of minimal markov basis for sampling from discrete conditional distributions. *Annals of the Institute of Statistical Mathematics*, 56(1):1–17.

Wedderburn, J. (1908). On hypercomplex numbers. *Proceedings of the London Mathematical Society*, 2(1):77–118.



LIST OF PUBLICATIONS

The following are the list of publications that arise from this study.

Journal Articles:

- N. F. Mohammed**, I.S. Rakhimov and M. Shitan, 2017, Algebraic Method for Independence Model of Two-Way Contingency Tables, *Malaysian Journal of Mathematical Sciences*, 11(S), February: 5372.
- N. F. Mohammed**, I.S. Rakhimov and Sh.K. Said Husain, 2017, On Cohomology Groups of Three-Dimensional Complex Associative Algebras, *Far East Journal of Mathematical Sciences*, 102(4), 669-686.
- N. F. Mohammed**, I.S. Rakhimov and Sh.K. Said Husain, 2017, On Contractions of Three-Dimensional Complex Associative Algebras, *Journal of Generalized Lie Theory and Applications*, 11(3).
- M. A. Fiidow, **N. F. Mohammed**, I.S. Rakhimov and Sh.K. Said Husain, 2017, On Inner Derivations of Finite Dimensional Associative Algebras, *Far East Journal of Mathematical Sciences*, ((In Press)).

Conferences:

- N. F. Mohammed**, I.S. Rakhimov and M. Shitan, *Markov Bases and Toric Ideals for Some Contingency Tables*, The 2nd International Conference of Mathematical Sciences and Statistics (ICMSS 2016). Sunway Putra Hotel, Kuala Lumpur, Malaysia. January 26-28, 2016.
- N. F. Mohammed**, I.S. Rakhimov and M. Shitan, *Toric Ideals and Markov Bases for an Agriculture Data Set*, Conference on Agriculture Statistics 2015 (CAS2015), Faculty of Agriculture and Food Sciences, UPM Bintulu Sarawak Campus, Malaysia. October 6-8, 2015.
- N. F. Mohammed**, I.S. Rakhimov and Sh.K. Said Husain, *Contractions of Low Dimensional Complex Associative Algebras*, The 2nd International Conference and Workshop of Mathematical Analysis (ICOWMA 2016), Century Langkawi Beach Resort, Langkawi, Malaysia. August 2-4, 2016.
- N. F. Mohammed**, I.S. Rakhimov and Sh.K. Said Husain, *Cohomology Spaces of Low Dimensional Complex Associative Algebras*, The 4th International Conference of Mathematical Sciences (ICMS4), Palm Garden Hotel, Putrajaya, Malaysia. November 15-17, 2016.
- N. F. Mohammed**, I.S. Rakhimov and M. Shitan, *Toric Ideals and Markov Bases for an Agriculture Data Set*, The Fundamental Science Congress (FSC2015), Faculty of Science Complex, Universiti Putra Malaysia. Malaysia. November 12-13, 2015.

N. F. Mohammed, I.S. Rakhimov and Sh.K. Said Husain, *On Degenerations of Complex Associative Algebras*, The Fundamental Science Congress (FSC2016), Faculty of Science Complex, Universiti Putra Malaysia. Malaysia. August 9-10, 2016.

N. F. Mohammed, I.S. Rakhimov and Sh.K. Said Husain, *Invariance Groups for Some Contingency Tables*, The 37th International Conference on Quantum Probability and Related Topics (37th QP), International Islamic University Malaysia, Kuantan, Malaysia. August 22-26, 2016.(Poster).

