



UNIVERSITI PUTRA MALAYSIA

***NMR-BASED METABOLOMICS FOR QUALITY CONTROL OF DUKUNG
ANAK (*Phyllanthus niruri* Linn.) AND IDENTIFICATION OF
BIOMARKERS
FOR ITS DIABETES-RELATED ACTIVITIES***

AHMED MEDIANI

FSTM 2016 1



NMR-BASED METABOLOMICS FOR QUALITY CONTROL OF DUKUNG ANAK (*Phyllanthus niruri* Linn.) AND IDENTIFICATION OF BIOMARKERS FOR ITS DIABETES-RELATED ACTIVITIES

By
AHMED MEDIANI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

August 2016



© COPYRIGHT UPM

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright© Universiti Putra Malaysia



DEDICATION

This thesis is dedicated to my parents, family, and friends.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

NMR-BASED METABOLOMICS FOR QUALITY CONTROL OF DUKUNG ANAK (*Phyllanthus niruri* Linn.) AND IDENTIFICATION OF BIOMARKERS FOR ITS DIABETES-RELATED ACTIVITIES

By

AHMED MEDIANI

August 2016

Chairman : Associate Professor Faridah Abas, PhD
Faculty : Food Science and Technology

Phyllanthus is a plant genus that has been used in traditional medicine due to its phytomedicinal metabolites content. However, there is a lack of consistency and efficacy of plant of this genus as well as not enough scientific reports to support its traditional uses specifically in treating diabetes. The main aim of this study was to investigate the antidiabetic properties of *Phyllanthus* *in vitro* and *in vivo* using metabolomics approach. The first part of the present study determined the variation between two *Phyllanthus* species (*P. niruri* and *P. urinaria*) at different growth stages (8, 10 and 12 weeks) using proton nuclear magnetic resonance (¹H NMR) combined with multivariate data analysis (MVDA). The results revealed that the Principal Component Analysis (PCA) and Partial Least Square (PLS) showed ideal differentiation between these species, suggesting the contributed metabolites, which were responsible for the discrimination. *Phyllanthus niruri* extracted with 80% ethanol possessed high antioxidant and α -glucosidase inhibitory activities compared to *P. urinaria* and 10 weeks was the valuable growth age that provides *P. niruri* extract with the highest content or/and number of bioactive metabolites. Thus, *P. niruri* harvested at 10 weeks was selected in the second part of the study to optimize the appropriate drying method and solvent for extracting these bioactive components. Three extracting solvents (methanol, ethanol and water) with different ratios were assessed for their effects on biological activities of *P. niruri* subjected to three drying treatments (air, oven and freeze). The freeze-dried *P. niruri* samples that were extracted with 80% ethanol exhibited higher biological activity values compared to the other extracts. The 80% ethanol extract had the quantities of epicatechin (172.0 $\mu\text{g/g}$), ellagic acid (46.67 $\mu\text{g/g}$), chlorogenic acid (44.33 $\mu\text{g/g}$), quercetin (1.73 $\mu\text{g/g}$), catechin (1.63 $\mu\text{g/g}$), phyllanthin (46.0 $\mu\text{g/g}$), and hypophyllanthin (23.17 $\mu\text{g/g}$) analyze using high performance liquid chromatography (HPLC). In the third part of this study, the correlation was also determined among the phytochemical constituents and the bioactivities using PLS regression. The identified phenolics and hypophyllanthin were strongly correlated with the antioxidant and α -glucosidase inhibitory activities, suggesting their great contributions to these activities. The optimized *P. niruri* extract was further

evaluated for *in vivo* antidiabetic properties at two concentrations (250 and 500 mg/kg body weight) and compared with metformin. The effect of *P. niruri* extract on the biochemical parameters of obese–diabetic rats was also investigated. The development of the obese diabetic model was done by feeding the high fat diet to Sprague-Dawley rats and inducing diabetic condition with a low dose of streptozotocin (STZ). The *in vivo* results indicated that *P. niruri* extracts of 500 mg/kg bw displayed the management of metabolites disorders of obese diabetic rats toward the normal state. The phytochemicals of *P. niruri* extracts might improve the metabolic disorders caused by diabetes. The extract at a concentration of 500 mg/kg bw also exhibited a noticeable effect in declining the plasma glucose level and improving lipid profile in obese diabetic rats as compared to 250 mg extract/kg bw. Both *P. niruri* doses showed reduction in the plasma cholesterol and low-density lipoprotein (LDL) levels and an increase in high-density lipoprotein (HDL). The effect of various metabolites in controlling the diabetes syndromes was discussed. The present study suggested that *P. niruri* can be a prominent and constructive medicinal plant. In conclusion, ¹H NMR-based metabolomics approach was successfully used to optimize the postharvest parameters, provide insights into the efficacy of *P. niruri* as a remedy for diabetes, and set the preliminary step towards developing this herb into high claim products.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

PENDEKATAN METABOLOMIK DALAM MENYIASAT SEBATIAN ANTIDIABETIK BAGI EKSTRAK DUKUNG ANAK (*Phyllanthus niruri* Linn.) DAN PENGENALPASTIAN PENANDA BIO DI DALAM TIKUS SPRAGUE-DAWLEY TERARUH DIABETES

Oleh

AHMED MEDIANI

Ogos 2016

Pengerusi : Profesor Madya Faridah Abas, PhD
Fakulti : Sains dan Teknologi Makanan

Phyllanthus merupakan genus tumbuhan yang telah digunakan dalam perubatan tradisional disebabkan oleh kandungan metabolit fitoperubatannya. Walau bagaimanapun, terdapat kekurangan konsistensi dan keberkesanan tumbuhan daripada genus ini dan juga laporan saintifik tidak cukup untuk menyokong kegunaan tradisional khusus dalam merawat diabetes. Tujuan kajian ini adalah untuk menyiasat sifat-sifat antidiabetik *Phyllanthus* secara *in vitro* dan *in vivo* menggunakan pendekatan metabolomik. Bahagian pertama kajian ini menentukan variasi antara dua spesies *Phyllanthus* (*P. niruri* dan *P. urinaria*) pada peringkat pertumbuhan yang berbeza (8, 10 dan 12 minggu) dengan menggunakan proton resonans magnetik nuklear (^1H NMR) digabungkan dengan analisis data multivariat (MVDA). Keputusan daripada Analisis Komponen Utama (PCA) dan Separa Dua Terkecil (PLS) menunjukkan perbezaan yang ideal antara spesies ini, mencadangkan metabolit yang menyumbang terhadap diskriminasi. Ekstrak *P. niruri* memiliki bioaktiviti yang lebih tinggi berbanding *P. urinaria* dan sepuluh minggu merupakan umur bermanfaat yang menyediakan ekstrak *P. niruri* dengan kandungan tertinggi atau/ dan beberapa metabolit bioaktif. Oleh itu, *P. niruri* yang dituai pada 10 minggu telah dipilih untuk mengenal pasti kaedah pengeringan dan pelarut yang sesuai untuk mengekstrak komponen bioaktif. Tiga pelarut pengekstrakan (metanol, etanol dan air) dengan nisbah yang berbeza telah dinilai untuk kesannya terhadap aktiviti-aktiviti biologi *P. niruri* tertakluk kepada tiga rawatan pengeringan (udara, ketuhar dan sejuk beku). Sampel pengeringan sejuk beku *P. niruri* yang diekstrak dengan 80% etanol mempamerkan nilai aktiviti biologi yang lebih tinggi berbanding dengan ekstrak lain. Ekstrak bagi 80% etanol mempunyai kuantiti epikatekin (172.0 $\mu\text{g/g}$), asid elargik (46.67 $\mu\text{g/g}$), asid klorogenik (44.33 $\mu\text{g/g}$), kuersetin (1.73 $\mu\text{g/g}$), katekin (1.63 $\mu\text{g/g}$), filantin (46.0 $\mu\text{g/g}$), dan hipofilantin (23.17 $\mu\text{g/g}$) dianalisis menggunakan kromatografi cecair berprestasi tinggi (HPLC). Untuk bahagian ketiga kajian, korelasi antara juzuk fitokimia dan bioaktiviti ditentukan menggunakan regresi PLS. Metabolit-metabolit yang telah dikenalpasti mempunyai kaitan yang kukuh dengan aktiviti antioksidan dan perencatan α -glukosidase, menunjukkan sumbangan besar mereka kepada aktiviti-aktiviti ini. Ekstrak *P. niruri* yang telah

dioptimumkan seterusnya dinilai bagi pengaruh antidiabetik secara *in vivo* pada dua kepekatan (250 dan 500 mg/kg bw) dan dibandingkan dengan metformin. Kesan ekstrak *P. niruri* kepada parameter biokimia tikus gemuk-diabetes juga disiasat. Pembentukan model gemuk diabetes dilakukan dengan memberi diet lemak yang tinggi sebagai makanan kepada tikus Sprague-Dawley dan mengaruh kencing manis dengan dos yang rendah streptozotocin (STZ). Keputusan *in vivo* menunjukkan ekstrak *P. niruri* sebanyak 500 mg/kg bw memaparkan pengurusan metabolit yang terganggu bagi tikus gemuk-diabetes ke arah kondisi normal. Fitokimia ekstrak *P. niruri* mungkin membawa kepada peningkatan tahap insulin, sekali gus merangsang β -sel pankreas sedia ada dalam tikus kencing manis. Ekstrak pada kepekatan 500 mg/kg bw juga mempamerkan kesan yang ketara dalam penurunan paras glukosa plasma dan meningkatkan profil lipid dalam tikus gemuk diabetes berbanding dengan 250 ekstrak mg/kg bw. Ekstrak *P. niruri* pada kedua-dua kepekatan menunjukkan pengurangan dalam paras kolesterol plasma dan lipoprotein berketumpatan rendah (LDL) dan peningkatan dalam lipoprotein berketumpatan tinggi (HDL). Kesan sinergi pelbagai metabolit dalam mengawal sindrom diabetes telah dibincangkan. Kajian ini mencadangkan *P. niruri* boleh menjadi tumbuhan ubatan yang penting dan berharga. Kesimpulannya, pendekatan metabolomik berasaskan ^1H NMR telah berjaya digunakan untuk mengoptimumkan parameter lepas tuai, memberi maklumat keberkesanan *P. niruri* sebagai ubat untuk kencing manis, dan menetapkan langkah awal ke arah membangunkan herba ini sebagai produk bertuntutan tinggi.

ACKNOWLEDGEMENTS

At the beginning, all praise due to ALLAH (SWT), lord of the world and peace be upon His messenger, prophet Mohammad (PBUH). Only by ALLAH Grace and Mercy this work has been completed and incha Allah, it will be successful.

Throughout my research period, there is so many people deserve appreciation and credit for their direct or indirect contribution in this work. First of all, the person who I respect most and deserve the most reverence expression is my supervisor Associate Professor Dr. Faridah Abas without her guidance; I cannot reach and achieve what I did. I would like to thank Associate Professor Dr. Faridah for her kind help, great support, beneficial advice and giving me freedom to pursue this project. I also highly appreciate her understanding, patience and motivating me through my research period.

I also want to give my warm appreciation to my committee members: Associate Professor Dr. Intan Safinar Ismail, Professor Dr. Amin Ismail, Professor Dr. Tan Chin Ping, and Associate Professor Dr. Alfi Khatib for their helpful suggestions, insights and great comments. I would like also to thank all the staffs of Faculty of Food Science and Technology and Laboratoy of Natural Products, Institute of Bioscience for sharing their valued knowledges, and rigorous attitudes toward scientific research.

I want to express my very sincere thankful and love to my wonderful family members: my parents, my grandmother, my brothers, my sisters, my uncles, my aunties and cousins for their support and prayers. Their encouragements and financial support push me to struggle for finishing and achieving the purpose of this project.

I would like to express my gratitude to my friendly lab mates and friends for their supports and encouragements. Sincere and special thanks are given to Mr Salahudin, Haji Azizul, Mrs Huda, and Dr Mulidiani for their guidance in NMR works.

Everyone whose named above or not listed, but whose amity is important to me. They deserve my intensive and earnest gratitude for helping me throughout this period. I would like to thank the various members of our Research group with whom I had the opportunity to work with to mention few Dr Leong, Dr Munirah, Ashikin, Khaleeda, Kayne, Soo Yee, Tahani, Azliana Abubakar, Umar Lawal, Khoo, Muhammad Abubakar Abo, Khaoula, and Zulaikha. They provided a friendly and cooperative atmosphere at work and also useful feedback and insightful comments on my work.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Faridah Abas, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Intan safinar binti Ismail, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Amin Ismail, PhD

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Tan Chin Ping, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice- Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Ahmed Mediani, GS34483

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____

Name of
Chairman of
Supervisory
Committee:

Associate Professor Dr. Faridah Abas

Signature: _____

Name of
Member of
Supervisory
Committee:

Associate Professor Dr. Intan Safinar Binti Ismail

Signature: _____

Name of
Member of
Supervisory
Committee:

Professor Dr. Tan Chin Ping

Signature: _____

Name of
Member of
Supervisory
Committee:

Professor Dr. Amin Ismail

TABLE OF CONTENTS

| | Page |
|--|-------------|
| ABSTRACT | i |
| ABSTRAK | iii |
| ACKNOWLEDGEMENT | v |
| APPROVAL | vii |
| DECLARATION | viii |
| LIST OF TABLES | xiv |
| LIST OF FIGURES | xv |
| LIST OF ABBREVIATIONS | xviii |
| | |
| CHAPTER | |
| | |
| 1 INTRODUCTION | 1 |
| 1.1 Background | 1 |
| 1.2 Scope and objectives | 3 |
| | |
| 2 LITERATURE REVIEW | 4 |
| 2.1 General description on genus <i>Phyllanthus</i> | 4 |
| 2.2 <i>Phyllanthus niruri</i> | 5 |
| 2.2.1 Traditional uses of <i>P. niruri</i> | 6 |
| 2.2.2 Medicinal uses of <i>P. niruri</i> | 7 |
| 2.2.3 Bioactive compound in <i>Phyllanthus niruri</i> | 8 |
| 2.3 <i>Phyllanthus urinaria</i> | 9 |
| 2.3.1 Bioactive compounds in <i>Phyllanthus urinaria</i> | 10 |
| 2.3.2 Medicinal uses of <i>Phyllanthus urinaria</i> | 11 |
| 2.4 Diabetes mellitus | 11 |
| 2.4.1 Relationship between obesity and diabetes | 12 |
| 2.4.2 Medicinal plants as anti-diabetic agents | 13 |
| 2.4.3 Natural compounds possessing antidiabetic properties | 14 |
| 2.4.4 α -Glucosidase inhibition | 14 |
| 2.4.5 Drugs altering insulin action | 15 |
| Enhancers of insulin effects: Biguanides (Metformin) | 15 |
| Alpha-Glucosidase Inhibitors: Acarbose and Miglitol | 15 |
| 2.5 Metabolomics | 16 |
| 2.5.1 Metabolomics in studying diseases | 17 |
| 2.5.2 Metabolomics in medicinal practices | 19 |
| 2.5.3 Metabolomics in diabetes research | 20 |
| 2.5.4 Metabolomics in studying metabolic disorders of DM | 21 |
| 2.5.5 The uses of NMR in studying antidiabetic compounds | 22 |
| 2.6 Data processing and statistical analysis in metabolomics | 23 |
| 2.6.1 Principal component analysis (PCA) | 23 |
| 2.6.2 Partial least square (PLS) | 24 |
| 2.6.3 Partial least square discriminate analysis (PLS-DA) | 24 |

| | | |
|----------|---|-----------|
| 3 | PHYTOCHEMICAL AND BIOLOGICAL FEATURES OF <i>P. niruri</i> AND <i>P. urinaria</i> HARVESTED AT DIFFERENT GROWTH STAGES REVEALED BY ¹H NMR-BASED METABOLOMICS | 25 |
| 3.1 | Introduction | 25 |
| 3.2 | Materials and Methods | 26 |
| 3.2.1 | Chemical and Reagent | 26 |
| 3.2.2 | Plant Material | 27 |
| 3.2.3 | Extraction of samples | 27 |
| 3.2.4 | NMR analysis | 27 |
| 3.2.5 | Bucketing of ¹ H NMR spectra | 28 |
| 3.2.6 | Total phenolic content assay | 28 |
| 3.2.7 | DPPH Free radical scavenging assay | 28 |
| 3.2.8 | α -Glucosidase inhibitory assay | 29 |
| 3.2.9 | Ultra performance liquid chromatography tandem mass spectrometry | 29 |
| 3.2.10 | Statistical analysis | 30 |
| 3.3 | Results and Discussion | 30 |
| 3.3.1 | Metabolite identification in <i>P. niruri</i> and <i>P. urinaria</i> | 30 |
| 3.3.2 | Principal component analysis (PCA) | 36 |
| 3.3.3 | Total phenolic content assay (TPC) and DPPH free radical scavenging assay and α -glucosidase inhibitory assay of <i>P. niruri</i> and <i>P. urinaria</i> . | 42 |
| 3.3.4 | Correlation between biological activities and phytochemicals using PLS | 44 |
| 3.3.5 | PLS model validation | 47 |
| 3.4 | Conclusions | 49 |
| 4 | RELATIONSHIP BETWEEN METABOLITES COMPOSITION AND BIOLOGICAL ACTIVITIES OF <i>P. niruri</i> EXTRACTS PREPARED BY DIFFERENT DRYING METHODS AND SOLVENTS EXTRACTION | 50 |
| 4.1 | Introduction | 50 |
| 4.2 | Materials and Methods | 51 |
| 4.2.1 | Chemicals and reagents | 51 |
| 4.2.2 | Plant material | 51 |
| 4.2.3 | Sample drying and extraction | 51 |
| 4.2.4 | Determination of total phenolic content (TPC) | 52 |
| 4.2.5 | Ferric reducing antioxidant potential (FRAP) | 52 |
| 4.2.6 | DPPH free radical scavenging assay | 52 |
| 4.2.7 | α -Glucosidase inhibitory activity | 52 |
| 4.2.8 | High performance liquid chromatography (HPLC) | 53 |
| 4.2.9 | Statistical analysis | 53 |
| 4.3 | Results and Discussion | 53 |
| 4.3.1 | Effect of drying method and solvent on the yield of extraction | 53 |
| 4.3.2 | Effects of drying methods and solvent on total phenolic content | 54 |
| 4.3.3 | Effect of drying methods and solvent on the antioxidant activity | 55 |

| | | |
|----------|--|-----------|
| 4.3.4 | Effect of drying methods and solvent on α -glucosidase inhibitory activity | 57 |
| 4.3.5 | Quantification of <i>P. niruri</i> metabolites using HPLC | 58 |
| 4.3.6 | PLS correlation | 62 |
| 4.4 | Conclusions | 64 |
| 5 | EFFECT OF POST-HARVEST TREATMENT ON PHYTOCHEMICAL COMPOSITION OF <i>P. niruri</i> USING NMR-BASED METABOLOMICS APPROACH | 65 |
| 5.1 | Introduction | 65 |
| 5.2 | Materials and Methods | 66 |
| 5.2.1 | Chemicals and reagent | 66 |
| 5.2.2 | Plant material and drying treatments | 66 |
| 5.2.3 | Extraction of samples | 67 |
| 5.2.4 | NMR analysis | 67 |
| 5.2.5 | Bucketing of ^1H NMR spectra and multivariate data analysis | 67 |
| 5.2.6 | Statistical analysis | 67 |
| 5.3 | Results and Discussion | 67 |
| 5.3.1 | ^1H NMR spectra of the samples and metabolites assignment | 67 |
| 5.3.2 | Discrimination of the dried samples extracted with different ethanol ratios by PCA | 69 |
| 5.3.3 | Correlation between bioactivities and metabolite changes among the samples | 75 |
| 5.4 | Conclusions | 80 |
| 6 | METABOLOMIC ANALYSIS AND BIOCHEMICAL CHANGES IN THE URINE AND SERUM OF STREPTOZOTOCIN INDUCED NORMAL- AND OBESE-DIABETIC RATS | 81 |
| 6.1 | Introduction | 81 |
| 6.2 | Materials and Methods | 83 |
| 6.2.1 | Chemicals and reagents | 83 |
| 6.2.2 | Animal disease model | 83 |
| 6.2.3 | Induction of obesity and diabetes in rats and metformin treatment | 83 |
| 6.2.4 | ^1H NMR preparation and analysis of urine | 85 |
| 6.2.5 | Blood collection and biochemical analysis of serum | 85 |
| 6.2.6 | Statistical analysis | 86 |
| 6.3 | Results and discussion | 86 |
| 6.3.1 | Body weight | 86 |
| 6.3.2 | Analysis of urine of high fat diet compared to normal diet rats | 88 |
| 6.3.3 | Biomarkers for normal, obese and obese-diabetic rats | 90 |
| 6.3.4 | Metabolic variation among normal, normal-diabetic and obese-diabetic rats | 95 |
| 6.3.5 | Metformin treatment | 101 |
| 6.3.6 | Biochemical analysis | 105 |
| 6.3.7 | Metabolic pathway analysis of diabetic models | 106 |
| 6.4 | Conclusions | 109 |

| | | |
|----------|--|------------|
| 7 | METABOLIC AND BIOCHEMICAL CHANGES IN STREPTOZOTOCIN INDUCED OBESE-DIABETIC RATS TREATED WITH <i>P. niruri</i> EXTRACT | 110 |
| 7.1 | Introduction | 110 |
| 7.2 | Materials and Methods | 111 |
| 7.2.1 | Chemicals and reagents | 111 |
| 7.2.2 | Characterization of <i>P. niruri</i> extract | 111 |
| 7.2.3 | Animal disease model | 112 |
| 7.2.4 | Experimental induction of diabetes in rats | 112 |
| 7.2.5 | Treatments | 113 |
| 7.2.6 | ¹ H NMR analysis and preparation of urine | 114 |
| 7.2.7 | Blood collection and biochemical analysis of serum | 114 |
| 7.2.8 | Statistical analysis | 114 |
| 7.3 | Results and discussion | 114 |
| 7.3.1 | Preliminary ¹ H NMR spectroscopic data of <i>P. niruri</i> extract | 114 |
| 7.3.2 | Identification of metabolites in urine samples | 116 |
| 7.3.3 | Metabolites analysis of urine samples and biomarkers of treated and non-treated obese-diabetic rats | 116 |
| 7.3.4 | Correlation among metabolites by Pearson's correlation analysis | 123 |
| 7.3.5 | Biochemical analysis | 125 |
| 7.3.6 | Metabolites association and pathway Analysis | 126 |
| | Amino acid metabolism | 129 |
| | Lipid metabolism | 129 |
| | Carbohydrate metabolism | 129 |
| | Tricarboxylic acid (TCA) | 130 |
| | Glyoxylate and dicarboxylate metabolism | 130 |
| 7.5 | Conclusions | 131 |
| 8 | CONCLUSIONS AND RECOMMENDATIONS | 132 |
| | BIBLIOGRAPHY | 134 |
| | BIODATA OF STUDENT | 150 |
| | LIST OF PUBLICATIONS | 151 |

LIST OF TABLES

| Table | | Page |
|-------|--|------|
| 2.1 | Botanical classification of <i>Phyllanthus niruri</i> | 5 |
| 2.2 | Botanical classification of <i>Phyllanthus urinaria</i> | 10 |
| 3.1 | The ¹ H NMR characteristics signals and UPLC-MS/MS data of both <i>P. niruri</i> and <i>P. urinaria</i> | 32 |
| 3.2 | VIP values of the major contributing compounds in the PLS | 46 |
| 4.1 | The crude extract (mg/100 mg) of dried sample after three different drying methods with different solvents | 54 |
| 4.2 | Total phenolic content (TPC) of <i>P. niruri</i> extracts | 55 |
| 4.3 | Free radical scavenging activity (DPPH) of <i>P. niruri</i> extracts | 56 |
| 4.4 | Ferric reducing ability power (FRAP) of <i>P. niruri</i> extracts | 57 |
| 4.5 | α -Glycosidase inhibitory activity of <i>P. niruri</i> extracts | 58 |
| 4.6 | The chemical constituents of three dried <i>P. niruri</i> extracted with different solvent ratios | 59 |
| 5.1 | ¹ H NMR characteristic signals of identified metabolites in <i>P. niruri</i> extracts | 68 |
| 5.2 | VIP values of the major contributing compounds in the PLS | 78 |
| 7.1 | The concentration of metabolites in <i>P. niruri</i> extract at 250 and 500 mg/Kg Bw | 115 |

LIST OF FIGURES

| Figure | | Page |
|--------|--|------|
| 2.1 | The plant of <i>Phyllanthus niruri</i> | 6 |
| 2.2 | Chemical structures of compounds isolated from <i>P. niruri</i> | 9 |
| 2.3 | The plant of <i>Phyllanthus urinaria</i> | 10 |
| 2.4 | Alpha-glucosidase and PNPG enzymatic reaction | 15 |
| 3.1 | ¹ H NMR spectra of the 80% ethanolic extract of <i>P. urinaria</i> (a) and <i>P. niruri</i> (b) with their expanded spectra for the range between δ 5.0 to 8.0 ppm. | 31 |
| 3.2 | The PCA score plot (PC1 vs. PC2, a) and PC1 loading column plot (b) for the discrimination between <i>P. niruri</i> (P N) and <i>P. urinaria</i> (P U) extracts. | 37 |
| 3.3 | The PCA score plot (PC1 vs. PC2, a) and the loading column plots of PC1 (b) and PC2 (c) of the ¹ H NMR data representing <i>P. niruri</i> extracts at the age of 8, 10 and 12 weeks old. | 39 |
| 3.4 | Relative quantification of the identified metabolites in both <i>P. niruri</i> and <i>P. urinaria</i> extracts (a) and their variations for <i>P. niruri</i> at different growth stages (b) based on the mean peak area of the ¹ H NMR signals. | 41 |
| 3.5 | Total phenolic content (a), DPPH free radical scavenging activity (b) and α -glucosidase inhibitory activities (c) of <i>P. niruri</i> and <i>P. urinaria</i> measured at different growth stages. | 43 |
| 3.6 | The biplot obtained from PLS describing the correlation among different ages of <i>P. niruri</i> . | 45 |
| 3.7 | The PLS models validation with 100 permutations of DPPH (a), TPC (b), and α -glucosidase (c) | 48 |
| 4.1 | A representative HPLC chromatogram of <i>Phyllanthus niruri</i> extracts | 58 |
| 4.2 | The PLS score plot (a) and loading plot (b) of PC1 vs. PC2, which represents the correlation among the dried <i>P. niruri</i> extracted with different solvent ratios and their chemical constituents. | 61 |
| 4.3 | The PLS models validation with 100 permutations of DPPH (a), FRAP (b), TPC (c), and α -glucosidase (d). | 63 |

| | | |
|------|---|-----|
| 5.1 | The PCA score plot (a) and loading score plot (b) of the discrimination of <i>P. niruri</i> extracts with various ethanol ratios (0, 50, 70, 80, 100%). | 70 |
| 5.2 | The PCA score plot (PC1 vs. PC2, a) and PC1 (a) and PC2 (b) loading column plots of the discrimination among FD, OD and AD <i>P. niruri</i> extracts. | 72 |
| 5.3 | Relative quantification of identified metabolites of different dried <i>P. niruri</i> based on the mean peak area of the ¹ H NMR signals. | 74 |
| 5.4 | The biplot obtained from PLS describing the correlation among the phytochemical constituents of different ethanol extracts (a) of various dried (b) <i>P. niruri</i> with TPC, FRAP, DPPH and α-glucosidase inhibitory activities | 76 |
| 5.5 | The regression coefficients of <i>Y</i> variables of PLS validation of DPPH, TPC and α-glucosidase inhibitory activity | 79 |
| 6.1 | Schematic diagram of the animal experimental design | 84 |
| 6.2 | The changes of body weight of the normal and obese rats (a) and the diabetic groups compared to controls (b). | 87 |
| 6.3 | The PCA score (a) and loading (b) plots (PC1 vs. PC2), comparing the urine of the rats receiving the high fat diet with the normal diet after three months. | 89 |
| 6.4 | The OPLSDA score (a) and S plots (b) of normal vs obese | 91 |
| 6.5 | The OPLSDA score (a) and S plots (b) of the normal vs obese diabetes | 92 |
| 6.6 | The OPLSDA SUS plot of normal vs obese and the normal vs obese diabetes | 94 |
| 6.7 | The PCA score plots (a, b) of obese- and normal-diabetic rats compared to ND and OB at week 13 (baseline, B) and week 17 (final, F) (a) and their respective loading column plots (c, d). | 96 |
| 6.8 | The OPLSDA score (a) and S plots (b) of normal vs obese-diabetic rats. | 97 |
| 6.9 | The OPLSDA score (a) and S plots (b) of normal vs normal-diabetic rats | 98 |
| 6.10 | The OPLSDA SUS plot of normal vs obese-diabetic and normal vs normal-diabetic rats | 99 |
| 6.11 | The PLS-DA score (a) and loading column (b) plots of obese- and normal-diabetic rats compared to ND at week 17 (final, F). | 100 |

| | | |
|------|--|-----|
| 6.12 | The VIP values derived from PLS-DA | 101 |
| 6.13 | The PCA score plot of the trajectories of metformin treatment of normal-diabetic (a) and obese-diabetic (b). I: initial, before injecting STZ. B: baseline after injecting STZ. F: final after 4 weeks of treatment. | 102 |
| 6.14 | The PCA score (a) and loading (b) plots of metformin treatment of obese diabetic rats | 104 |
| 6.15 | The biochemical parameters of rat serum for diabetic groups compared to controls. | 106 |
| 6.16 | Metabolic pathway analysis in normal and obese diabetes. | 107 |
| 7.1 | Schematic diagram of the animal experimental design | 113 |
| 7.2 | The PLS-DA score (a) and loading column (b) plots of the effects of low and high doses of <i>P. niruri</i> extract on the obese diabetic rats after 4 weeks of treatment. | 117 |
| 7.3 | The PCA score showing the trajectories of <i>P. niruri</i> treatment with HD (a) and metformin (b) and the loading plot of <i>P. niruri</i> treatment with HD (C). | 118 |
| 7.4 | The heatmap of the effects of low and high doses of <i>P. niruri</i> extract on the treated obese diabetic rats after 4 weeks of treatment. | 120 |
| 7.5 | The VIP values derived from PLS-DA | 121 |
| 7.6 | The box plots of the relative quantities of the most significant metabolites in the urine samples from the treated and untreated groups. | 122 |
| 7.7 | The overall Pearson's correlation of the most significant metabolites in the rats' urine sample | 124 |
| 7.8 | The biochemical parameters of the serum of rats treated with <i>P. niruri</i> extract compared to controls after 4 weeks of treatment. | 125 |
| 7.9 | Summary of the pathway analysis | 127 |
| 7.10 | Metabolic pathways and metabolite changes observed in the urine of obese diabetic rats treated with <i>P. niruri</i> extract compared to controls. | 128 |

LIST OF ABBREVIATIONS

| | |
|-------------------|---|
| $^1\text{H NMR}$ | Proton Nuclear Magnetic Resonance Spectroscopy |
| AA | Antioxidant Activity |
| AD | Air Drying |
| OD | Oven Drying |
| FD | Freeze Drying |
| CC | Column Chromatography |
| d | Doublet |
| DAD | Diode Array Detector |
| dd | Doublet of doublet |
| dH ₂ O | Deionized water |
| DPPH | Diphenylpicrylhydrazyl |
| ESI | Electrospray Ionization |
| g | Gram |
| GAE | Gallic Acid Equivalent |
| HMBC | Heteronuclear Multiple Bond Correlation |
| HCA | Hierarchical Cluster Analysis Hierarchical Cluster Analysis |
| HPLC | High Performance Liquid Chromatography |
| HREISMS | High Resolution Electron Impact Mass Spectroscopy |
| Hz | Hertz |
| IC ₅₀ | Inhibition Concentration at 50 percent |
| L | Litre |
| LC-MS | Liquid Chromatography–Mass Spectrometry |
| m | Multiplet |
| m/z | Mass to charge |
| MeOH | Methanol |
| EtOH | Ethanol |
| Aq | Water |
| MHz | MegaHertz |
| mL | Milliliter |
| MS | Mass Spectrometry |
| MVDA | multivariate data analysis |
| °C | Degree in Celsius |

| | |
|---|---|
| PC | Principal Component |
| PCA | Principal Component Analysis |
| PLS | Partial Least Squares |
| PLS-DA | Partial Least Squares–Discriminant Analysis |
| ppm | Part Per Million |
| RMSEE | Root Mean Square Error of Estimation |
| RMSEP | Root Mean Square Error of Prediction |
| ROS | Reactive Oxygen Species |
| RPLC | Reversed Phase Liquid Chromatography |
| s | Singlet |
| SIMCA | Soft Independent Modeling of Class Analogy |
| TPC | Total Phenolic Contents |
| UV | Ultraviolet |
| UV/VIS | Ultraviolet/visible |
| VIP | variable Importance in the Projection |
| δ | Chemical Shift in ppm |
| μg | Microgram |
| μL | Microliter |
| ^{13}C | Carbon-13 |
| LDL | Low-density Lipoprotein |
| HDL | High-density Lipoprotein |
| STZ | Streptozotocin |
| DM | Diabetes Mellitus |
| WHO | World Health Organization |
| PNPG | p-nitrophenyl- α -D-glucopyranose |
| ANOVA | Analysis of Variance |
| CID | Collision-induced Dissociation |
| NaOD | Sodium Deuterium Oxide |
| D ₂ O | Deuterium Oxide |
| CD ₃ OD | Methanol- <i>d</i> ₄ |
| KH ₂ PO ₄ | Potassium Dihydrogen Phosphate |
| CMC | Carboxymethyl Cellulose |
| CH ₃ OH- <i>d</i> ₄ | Deuterated Methanol- <i>d</i> ₄ |
| SD | Sprague-Dawley |

| | |
|------|---------------------------|
| HFD | High Fat Diet |
| ND | Normal Diet |
| 2D | Two Dimensional |
| HMDB | Human Metabolome Database |
| TG | Total Triglycerides |



CHAPTER 1

INTRODUCTION

1.1 Background

Diabetes mellitus (DM) is the dysregulation in metabolic function of an organism characterized by the disorder of the catabolic reaction of organic nutrients. This disorder is caused by a disturbance in the excretion and/or action of insulin, which is associated with chronic hyperglycemia. The DM is a chronic disease in which the pancreas does not produce enough insulin (Type 1) or the body is unable to use the insulin effectively (Type 2). Worldwide, 9% of adults aged 18 years and above were living with diabetes in 2014. In addition, DM caused 1.5 million deaths in 2013, with 80% of them occurring in low- and middle-income countries (WHO, 2014). One of the practical treatments of diabetes is the reduction of postprandial hyperglycemia by the retardation of glucose catabolism via the inhibition of hydrolysis enzymes (α -glucosidase and α -amylase). This inhibition reaction leads to the prolongation of the sugar digestion time and a decrease in the proportion of glucose absorption. These inhibitors can be provided by natural products.

Plant extracts as natural products that contain several phytochemicals are attracting more consideration for their potential usage in curing and preventing DM. Numerous natural biomolecules and plant extracts have been examined for their antidiabetic activities both *in vitro* and *in vivo*, and they showed very promising impacts, which indicates that phytochemical intake might be a key approach for diabetes prevention. Recent studies have indicated that plant substances possess different functions that aid to find the prominent candidate for innovative antidiabetic usage (Kordalewska & Markuszewski, 2015; Otunola & Afolayan, 2015; Patel et al., 2012).

The *Phyllanthus* genus (Euphorbiaceae) has significant medicinal properties used in the treatment of different diseases (Markom et al., 2007; Moreira et al., 2013). *Phyllanthus* genus is originally from India, and it spreads in tropical regions, including Malaysia (Ali et al., 2006). In Malaysia, it is locally known as 'dukung anak'. Specifically, the genus *Phyllanthus* has been used historically for the treatment of intestinal parasites, problems of the liver, kidney and bladder and the treatment of diseases such as infectious diseases (Calixto et al., 1998). This genus contains several species, and the most popular ones are *Phyllanthus niruri* and *Phyllanthus urinaria* (Khatoun et al., 2006; Kumaran & Karunakaran, 2007). *Phyllanthus niruri* has been utilized as a traditional medicine to cure kidney problems, diarrhea, fever, diabetes and colic (Kumaran & Karunakaran, 2007). *Phyllanthus niruri* also has the ability to reduce lipids peroxidation and exhibit an analgesic effect (Amin et al., 2012; Kumaran & Karunakaran, 2007). In China, this plant is used as a traditional folk medicine and is known as 'pearls under the leaves' (Murugaiyah and Chan, 2009). It has also been acclaimed for its effective remedy to treat diabetic sufferers in Nigeria and India (Okoli et al., 2010; Okoli et al., 2011). However, *P. urinaria* has several uses in traditional medicine including liver

protection, fever reduction, laxative, cure for inflammation, suppression of hyperactivity of the liver, improve eyesight, and urine flow as well as detoxification of poisons from the body (Amin et al., 2012; Kumaran & Karunakaran, 2007). Currently, *P. urinaria* is also widely used as an anti-inflammatory and antioxidant agent as well as a smooth muscle relaxant (Catapan et al., 2000; Kumaran & Karunakaran, 2007). Various bioactive and phytochemical compounds are found in *P. urinaria*, including flavonoids, tannins, lignans and other benzenoid compounds (Chang et al., 2003; Khatoon et al., 2006; Kumaran & Karunakaran, 2007). In addition, corilagin, gallic acid, caffeoylquinic acid, geraniin, and rutin have been isolated from *P. urinaria* (Khatoon et al., 2006; Kumar et al., 2015).

Metabolites can have significant variation at several growth stages and may be negatively affected by some processing methods, such as drying, in terms of their bioactivities and chemical structures. Thermal drying conditions were stated to possess wide negative influences on the functionality of plant materials due to a significant reduction of its metabolites (Lim & Murtijaya, 2007). A large variability was also marked in the biological activity of plants and their metabolite constituents as a result of extraction with different solvents (Siddhuraju & Becker, 2003). Therefore, it would be valuable to accomplish the conservation, efficiency and consistency values of plants and their derivative products against the negative effects and changes of these aspects using the appropriate approach.

Metabolomics is related to the metabolome, which represents the small molecules of the cells of organisms. It is a competent approach to study and evaluate the changes in the small molecule composition of the biological system under various circumstances. Furthermore, metabolomics is a global profiling tool in the detection of metabolites and metabolic pathways (Kaddurah-Daouk et al., 2008). Its central value is dramatically increased in biological regulation, including the control and management of diseases. Metabolomics sheds light on medicine with the help of technological advances, which bring it to the point that it can grow rapidly in the biological and biomedical areas. New insights and ideas regarding metabolites and their mechanisms of action provide the platform to be further used for studying many diseases. Proton nuclear magnetic resonance (^1H NMR) is emerging as a leading analytical tool in metabolomics studies due to the large number and groups of identified compounds. Thus, ^1H NMR-based metabolomics can be used as an approach to assess the variation in phytochemicals content and bioactivity of plant material and changes in biofluids (Cevallos-Cevallos et al., 2009; Deja et al., 2013).

Animal models play an important role in understanding metabolic disorders. Many studies have used animal models in simulating diabetic conditions to understand the molecular events associated with this disease and its complications (Bakar et al., 2015; Beckonert et al., 2007). In addition, the metabolic changes of body fluids associated with treatment with several herbal drugs can also be evaluated. Recently, herbal drugs, herbal drug preparations and herbal medicinal products are extensively practiced in most countries. There is thus a need for an appropriate approach for rapid screening, identification and preclinical testing. Metabolomics can provide the accuracy, reproducibility, selectivity and specificity that are needed by applying

different tools as well as coupling these tools (Bilia et al., 2002). Its mission is also to detect, identify, and correlate the metabolic variations in the biological system, regardless of the individual cell. It can still directly reflect the functionality of the cells. Therefore, it is preferred in many cases over genomic and proteomic methods for predicting the efficacy of drugs and extracts in the control and management of diseases.

1.2 Scope and objectives

The current study was initiated to solve several issues that can serve as platforms in obtaining valuable extracts for the treatment of DM and the improvement of its metabolic disturbances. The main aim of this study was to evaluate the antidiabetic activity of *Phyllanthus* extracts by ^1H NMR based metabolomics. To accomplish this goal, numerous specific objectives were proposed. The first part of the study aims to compare the antioxidant and α -glucosidase inhibitory activities of two *Phyllanthus* species and their phytochemicals content at different growth stages (Chapter 3). The most active species at the selected growth age was chosen to evaluate the effect of different solvents and drying methods on antioxidant and α -glucosidase inhibitory activities and quantifying the metabolites by high performance liquid chromatography (HPLC) (Chapter 4). Additionally profiling of the chemical constituents of the most desirable and potent extract and correlate the chemical profile with biological activities was further evaluated on the active species (Chapter 5). Then, the animal model of normal- and obese-diabetic rats was optimized and the metformin effect was tested (Chapter 6). Finally, the overall bio-markers (in animal bio-fluids) relating to the effect of *P. niruri* in streptozotocin induced diabetic rats were evaluated (Chapter 7).

The five specific objectives of the study are presented below:

1. To determine the antioxidant and α -glucosidase inhibitory activities of *Phyllanthus niruri* and *Phyllanthus urinaria* and their phytochemicals content at different growth ages (8, 10 and 12 weeks) using ^1H NMR-based metabolomics.
2. To profile the chemical constituents of the most desirable and potent extract and quantify the metabolites by HPLC and correlate the chemical profile with biological activities.
3. To determine the effect of different solvents and drying methods on antioxidant and α -glucosidase inhibitory activities of *P. niruri* and its phytochemical constituents using ^1H NMR-based metabolomics.
4. To evaluate the metabolic disorders associated with Type 1 and Type 2 DM on the urine composition of obese- and normal-diabetic rats as well as the effect of metformin treatment.
5. To identify the metabolic disorders in obese diabetic rats and evaluate the functionality and benefits of *P. niruri* extract on ameliorating the complications associated using a ^1H NMR-based metabolomics approach.



© COPYRIGHT UPM

BIBLIOGRAPHY

- Abas, F., Lajis, N. H., Israf, D., Khozirah, S., & Kalsom, Y. U. (2006). Antioxidant and nitric oxide inhibition activities of selected Malay traditional vegetables. *Food Chemistry*, 95(4), 566-573.
- Abas, F., Shaari, K., Israf, D., Syafri, S., Zainal, Z., & Lajis, N. H. (2010). LC–DAD–ESI-MS analysis of nitric oxide inhibitory fractions of tenggek burung (*Melicope ptelefolia* Champ. ex Benth.). *Journal of Food Composition and Analysis*, 23(1), 107-112.
- Abdel-Farid, I., Sheded, M., & Mohamed, E. (2014). Metabolomic profiling and antioxidant activity of some Acacia species. *Saudi Journal of Biological Sciences*, 21(5), 400-408.
- Ado, M. A., Abas, F., Leong, S. W., Shaari, K., Ismail, I. S., Ghazali, H. M., & Lajis, N. H. (2016). Chemical constituents and biological activities of *Callicarpa maingayi* leaves. *South African Journal of Botany*, 104, 98-104.
- Ahmed, O. M., Moneim, A. A., Yazid, I. A., & Mahmoud, A. M. (2010). Antihyperglycemic, antihyperlipidemic and antioxidant effects and the probable mechanisms of action of *Ruta graveolens* infusion and rutin in nicotinamide-streptozotocin-induced diabetic rats. *Diabetologia Croatica*, 39(1), 15-35.
- Alberti, Kurt George Matthew Mayer, Zimmet, P., & Shaw, J. (2006). Metabolic syndrome a new world wide definition. A consensus statement from the international diabetes federation. *Diabetic Medicine*, 23(5), 469-480.
- Albrecht, C., Stander, M., Grobbelaar, M., Colling, J., Kossmann, J., Hills, P., & Makunga, N. (2012). LC–MS-based metabolomics assists with quality assessment and traceability of wild and cultivated plants of *Sutherlandia frutescens* (Fabaceae). *South African Journal of Botany*, 82, 33-45.
- Ali, H., Houghton, P., & Soumyanath, A. (2006). α -Amylase inhibitory activity of some Malaysian plants used to treat diabetes; with particular reference to *Phyllanthus amarus*. *Journal of Ethnopharmacology*, 107(3), 449-455.
- Ali, Imen Ben El Hadj, Bahri, R., Chaouachi, M., Boussaïd, M., & Harzallah-Skhiri, F. (2014). Phenolic content, antioxidant and allelopathic activities of various extracts of *Thymus numidicus* Poir.organs. *Industrial Crops and Products*, 62, 188-195.
- Amin, Z. A., Abdulla, M. A., Ali, H. M., Alshawsh, M. A., & Qadir, S. W. (2012). Assessment of In vitro antioxidant, antibacterial and immune activation potentials of aqueous and ethanol extracts of *Phyllanthus niruri*. *Journal of the Science of Food and Agriculture*, 92(9), 1874-1877.
- Antuna-Puente, B., Feve, B., Fellahi, S., & Bastard, J. (2008). Adipokines: the missing link between insulin resistance and obesity. *Diabetes & Metabolism*, 34(1), 2-11.
- Bagalkotkar, G., Sagineedu, S. R., Saad, M. S., & Stanslas, J. (2006). Phytochemicals from *Phyllanthus niruri* Linn. and their pharmacological properties: a review. *Journal of Pharmacy and Pharmacology*, 58(12), 1559-1570.

- Bailey, N. J., Wang, Y., Sampson, J., Davis, W., Whitcombe, I., Hylands, P. J., Croft, S. L., & Holmes, E. (2004). Prediction of anti-plasmodial activity of *Artemisia annua* extracts: application of ^1H NMR spectroscopy and chemometrics. *Journal of Pharmaceutical and Biomedical Analysis*, 35(1), 117-126.
- Bailey, N. J., Sampson, J., Hylands, P. J., Nicholson, J. K., & Holmes, E. (2002). Multi-component metabolic classification of commercial feverfew preparations via high-field ^1H -NMR spectroscopy and chemometrics. *Planta Medica*, 68(8), 734-738.
- Bain, J. R., Stevens, R. D., Wenner, B. R., Ilkayeva, O., Muoio, D. M., & Newgard, C. B. (2009). Metabolomics applied to diabetes research: moving from information to knowledge. *Diabetes*, 58(11), 2429-2443.
- Bakar, M. H. A., Sarmidi, M. R., Cheng, K., Khan, A. A., Suan, C. L., Huri, H. Z., & Yaakob, H. (2015). Metabolomics—the complementary field in systems biology: a review on obesity and type 2 diabetes. *Molecular BioSystems*, 11(7), 1742-1774.
- Bansal, P., Paul, P., Mudgal, J., Nayak, P. G., Pannakal, S. T., Priyadarsini, K., & Unnikrishnan, M. (2012). Antidiabetic, antihyperlipidemic and antioxidant effects of the flavonoid rich fraction of *Pilea microphylla* (L.) in high fat diet/streptozotocin-induced diabetes in mice. *Experimental and Toxicologic Pathology*, 64(6), 651-658.
- Bao, Y., Zhao, T., Wang, X., Qiu, Y., Su, M., Jia, W., & Jia, W. (2009). Metabonomic variations in the drug-treated type 2 diabetes mellitus patients and healthy volunteers. *Journal of Proteome Research*, 8(4), 1623-1630.
- Bavarva, J. H. & Narasimhacharya, A. (2007). Comparative Antidiabetic, Hypolipidemic, and Antioxidant Properties of *Phyllanthus niruri*. in Normal and Diabetic Rats. *Pharmaceutical Biology*, 45(7), 569-574.
- Beckonert, O., Keun, H. C., Ebbels, T. M., Bundy, J., Holmes, E., Lindon, J. C., & Nicholson, J. K. (2007). Metabolic profiling, metabolomic and metabonomic procedures for NMR spectroscopy of urine, plasma, serum and tissue extracts. *Nature Protocols*, 2(11), 2692-2703.
- Bernhoft, A. (2010). A brief review on bioactive compounds in plants. *Bioactive Compounds in Plants-Benefits and Risks for Man and Animals*, 11-17.
- Berregi, I., Santos, J. I., del Campo, G., & Miranda, J. I. (2003a). Quantitative determination of (-)-epicatechin in cider apple juices by ^1H NMR. *Talanta*, 61(2), 139-145.
- Berregi, I., Santos, J. I., del Campo, G., Miranda, J. I., & Aizpurua, J. M. (2003b). Quantitation determination of chlorogenic acid in cider apple juices by ^1H NMR spectrometry. *Analytica Chimica Acta*, 486(2), 269-274.
- Bhattacharjee, R. & Sil, P. C. (2007). Protein isolate from the herb, *Phyllanthus niruri* L. (Euphorbiaceae), plays hepatoprotective role against carbon tetrachloride induced liver damage via its antioxidant properties. *Food and Chemical Toxicology*, 45(5), 817-826.
- Bilia, A. R., Bergonzi, M. C., Lazari, D., & Vincieri, F. F. (2002). Characterization of commercial kava-kava herbal drug and herbal drug preparations by means

- of nuclear magnetic resonance spectroscopy. *Journal of Agricultural and Food Chemistry*, 50(18), 5016-5025.
- Calixto, J. B., Santos, A. R., Filho, V. C., & Yunes, R. A. (1998). A review of the plants of the genus *Phyllanthus*: their chemistry, pharmacology, and therapeutic potential. *Medicinal Research Reviews*, 18(4), 225-258.
- Catapan, E., Otuki, M. F., Viana, A. M., Yunes, R. A., Bresciani, L. F., Ferreira, J., Santos, A. R., Calixto, J. B., & Cechinel-Filho, V. (2000). Pharmacological activity and chemical composition of callus culture extracts from selected species of *Phyllanthus*. *Die Pharmazie*, 55(12), 945-946.
- Cevallos-Cevallos, J. M., Reyes-De-Corcuera, J. I., Etxeberria, E., Danyluk, M. D., & Rodrick, G. E. (2009). Metabolomic analysis in food science: a review. *Trends in Food Science & Technology*, 20(11), 557-566.
- Chan, E., Lim, Y. Y., Wong, S. K., Lim, K., Tan, S., Lianto, F., & Yong, M. (2009). Effects of different drying methods on the antioxidant properties of leaves and tea of ginger species. *Food Chemistry*, 113(1), 166-172.
- Chang, C., Lien, Y., Liu, K. C. C., & Lee, S. (2003). Lignans from *Phyllanthus urinaria*. *Phytochemistry*, 63(7), 825-833.
- Chao, M., Zou, D., Zhang, Y., Chen, Y., Wang, M., Wu, H., Ning, G., & Wang, W. (2009). Improving insulin resistance with traditional Chinese medicine in type 2 diabetic patients. *Endocrine*, 36(2), 268-274.
- Chen, J., Zhao, X., Fritsche, J., Yin, P., Schmitt-Kopplin, P., Wang, W., Lu, X., Häring, H. U., Schleicher, E. D., & Lehmann, R. (2008). Practical approach for the identification and isomer elucidation of biomarkers detected in a metabolomic study for the discovery of individuals at risk for diabetes by integrating the chromatographic and mass spectrometric information. *Analytical Chemistry*, 80(4), 1280-1289.
- Chen, Y., Lo, Y., Chen, Y., Lai, N., Lin, N., Chung, T., Chen, W., & Tzen, J. T. (2015). Magnesium lithospermate B improves metabolic changes in high-fat diet-fed rats with metabolic syndrome. *Journal of Functional Foods*, 14, 163-173.
- Cheng, Q., Zhang, X., Wang, O., Liu, J., Cai, S., Wang, R., Zhou, F., & Ji, B. (2015). Anti-diabetic effects of the ethanol extract of a functional formula diet in mice fed with a fructose/fat-rich combination diet. *Journal of the Science of Food and Agriculture*, 95(2), 401-408.
- Cheng, Z., Su, L., Moore, J., Zhou, K., Luther, M., Yin, J. J., & Liangli (Lucy) Yu. (2006). Effects of postharvest treatment and heat stress on availability of wheat antioxidants. *Journal of Agricultural and Food Chemistry*, 54(15), 5623-5629.
- Cheng, H. Y., Yang, C. M., Lin, T. C., Lin, L. T., Chiang, L. C., & Lin, C. C. (2011). Excoecarianin, Isolated from *Phyllanthus urinaria* Linnaea, Inhibits Herpes Simplex Virus Type 2 Infection through Inactivation of Viral Particles. *Evidence-Based Complementary and Alternative Medicine : eCAM*, 2011, 259103.
- Chethan, S., Sreerama, Y., & Malleshi, N. (2008). Mode of inhibition of finger millet malt amylases by the millet phenolics. *Food Chemistry*, 111(1), 187-191.

- Choi, H., Choi, Y. H., Verberne, M., Lefeber, A. W., Erkelens, C., & Verpoorte, R. (2004). Metabolic fingerprinting of wild type and transgenic tobacco plants by ^1H NMR and multivariate analysis technique. *Phytochemistry*, 65(7), 857-864.
- Choi, Y. H., Sertic, S., Kim, H. K., Wilson, E. G., Michopoulos, F., Lefeber, A. W., Erkelens, C., Prat Kricun, S. D., & Verpoorte, R. (2005). Classification of *Ilex* species based on metabolomic fingerprinting using nuclear magnetic resonance and multivariate data analysis. *Journal of Agricultural and Food Chemistry*, 53(4), 1237-1245.
- Cnop, M., Welsh, N., Jonas, J. C., Jörns, A., Lenzen, S., & Eizirik, D. L. (2005). Mechanisms of pancreatic β -cell death in Type 1 and Type 2 diabetes many differences, few similarities. *Diabetes*, 54(suppl 2), S97-S107.
- Coman, C., Rugina, O. D., & Socaciu, C. (2012). Plants and natural compounds with antidiabetic action. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 40(1), 314.
- Connor, S. C., Hansen, M. K., Corner, A., Smith, R. F., & Ryan, T. E. (2010). Integration of metabolomics and transcriptomics data to aid biomarker discovery in type 2 diabetes. *Molecular Biosystems*, 6(5), 909-921.
- Cortés, M. E., Vigil, P., & Montenegro, G. (2011). The medicinal value of honey: A review on its benefits to human health, with a special focus on its effects on glycemic regulation. *Cien Inv Agr*, 38, 303-317.
- Dai, Y., Li, Z., Xue, L., Dou, C., Zhou, Y., Zhang, L., & Qin, X. (2010). Metabolomics study on the anti-depression effect of xiaoyaosan on rat model of chronic unpredictable mild stress. *Journal of Ethnopharmacology*, 128(2), 482-489.
- Dan, M., Su, M., Gao, X., Zhao, T., Zhao, A., Xie, G., Qiu, Y., Zhou, M., Liu, Z., & Jia, W. (2008). Metabolite profiling of *Panax notoginseng* using UPLC-ESI-MS. *Phytochemistry*, 69(11), 2237-2244.
- De Souza, T. P., Holzschuh, M. H., Lionço, M. I., González Ortega, G., & Petrovick, P. R. (2002). Validation of a LC method for the analysis of phenolic compounds from aqueous extract of *Phyllanthus niruri* aerial parts. *Journal of Pharmaceutical and Biomedical Analysis*, 30(2), 351-356.
- Deja, S., Barg, E., Młynarz, P., Basiak, A., & Willak-Janc, E. (2013). ^1H NMR-based metabolomics studies of urine reveal differences between type 1 diabetic patients with high and low HbA_{1c} values. *Journal of Pharmaceutical and Biomedical Analysis*, 83, 43-48.
- Deuschländer, M., Van de Venter, M., Roux, S., Louw, J., & Lall, N. (2009). Hypoglycaemic activity of four plant extracts traditionally used in South Africa for diabetes. *Journal of Ethnopharmacology*, 124(3), 619-624.
- Diao, C., Zhao, L., Guan, M., Zheng, Y., Chen, M., Yang, Y., Lin, L., Chen, W., & Gao, H. (2014). Systemic and characteristic metabolites in the serum of streptozotocin-induced diabetic rats at different stages as revealed by a ^1H -NMR based metabolomic approach. *Molecular BioSystems*, 10(3), 686-693.
- Dias, M. A., Campos, A. H., Yunes, R. A., & Calixto, J. B. (1995). Analysis of the mechanisms underlying the contractile response induced by the hydroalcoholic

extract of *Phyllanthus urinaria* in the guinea-pig urinary bladder in-vitro. *Journal of Pharmacy and Pharmacology*, 47(10), 846-851.

- Diaz, S. O., Pinto, J., Graça, G., Duarte, I. F., Barros, A. S., Galhano, E., Pita, C., Almeida, M. d. C., Goodfellow, B. J., & Carreira, I. M. (2011). Metabolic biomarkers of prenatal disorders: an exploratory NMR metabonomics study of second trimester maternal urine and blood plasma. *Journal of Proteome Research*, 10(8), 3732-3742.
- Dineshkumar, B., Mitra, A., & Manjunatha, M. (2010). A comparative study of alpha amylase inhibitory activities of common anti-diabetic plants at Kharagpur 1 block. *International Journal of Green Pharmacy*, 4(2), 115.
- Donath, M. Y., Størling, J., Maedler, K., & Mandrup-Poulsen, T. (2003). Inflammatory mediators and islet β -cell failure: a link between type 1 and type 2 diabetes. *Journal of molecular medicine*, 81(8), 455-470.
- DuPont, M. S., Bennett, R. N., Mellon, F. A., & Williamson, G. (2002). Polyphenols from alcoholic apple cider are absorbed, metabolized and excreted by humans. *The Journal of Nutrition*, 132(2), 172-175.
- Edmands, W. M., Beckonert, O. P., Stella, C., Campbell, A., Lake, B. G., Lindon, J. C., Holmes, E., & Gooderham, N. J. (2011). Identification of human urinary biomarkers of cruciferous vegetable consumption by metabonomic profiling. *Journal of Proteome Research*, 10(10), 4513-4521.
- Erdmann, J., Kallabis, B., Oppel, U., Sypchenko, O., Wagenpfeil, S., & Schusdziarra, V. (2008). Development of hyperinsulinemia and insulin resistance during the early stage of weight gain. *American Journal of Physiology. Endocrinology and Metabolism*, 294(3), E568-75.
- Eriksson, L., Antti, H., Gottfries, J., Holmes, E., Johansson, E., Lindgren, F., Long, I., Lundstedt, T., Trygg, J., & Wold, S. (2004). Using chemometrics for navigating in the large data sets of genomics, proteomics, and metabonomics (gpm). *Analytical and Bioanalytical Chemistry*, 380(3), 419-429.
- Eriksson, L., Kettaneh-Wold, N., Trygg, J., Wikström, C., & Wold, S. (2006). Multi- and megavariate data analysis: Part I: basic principles and applications.
- Fang, S., Rao, Y. K., & Tzeng, Y. (2008). Anti-oxidant and inflammatory mediator's growth inhibitory effects of compounds isolated from *Phyllanthus urinaria*. *Journal of Ethnopharmacology*, 116(2), 333-340.
- Felgines, C., Talavera, S., Texier, O., Gil-Izquierdo, A., Lamaison, J., & Remesy, C. (2005). Blackberry anthocyanins are mainly recovered from urine as methylated and glucuronidated conjugates in humans. *Journal of Agricultural and Food Chemistry*, 53(20), 7721-7727.
- Ferreira, A. S. (2009). Diagnostic accuracy of pattern differentiation algorithm based on Chinese medicine theory: a stochastic simulation study. *Chinese Medicine*, 4, 24-8546-4-24.
- Firuzi, O., Lacanna, A., Petrucci, R., Marrosu, G., & Saso, L. (2005). Evaluation of the antioxidant activity of flavonoids by "ferric reducing antioxidant power" assay and cyclic voltammetry. *Biochimica Et Biophysica Acta (BBA)-General Subjects*, 1721(1), 174-184.

- Fischer, U. A., Carle, R., & Kammerer, D. R. (2011). Identification and quantification of phenolic compounds from pomegranate (*Punica granatum* L.) peel, mesocarp, aril and differently produced juices by HPLC-DAD-ESI/MS n. *Food Chemistry*, 127(2), 807-821.
- Gami, B. (2011). Hemorrhoids—a common ailment among adults, causes & treatment: a review. *International Journal of Pharmacy and Pharmaceutical Sciences*, 3(Suppl 5), 5-13.
- Gandhi, G. R., Ignacimuthu, S., Paulraj, M. G., & Sasikumar, P. (2011). Antihyperglycemic activity and antidiabetic effect of methyl caffeate isolated from *Solanum torvum* Swartz. fruit in streptozotocin induced diabetic rats. *European Journal of Pharmacology*, 670(2), 623-631.
- Gao, P. (2008). The advantages of TCM syndrome type differentiation in diabetic retinopathy. *Beijing Journal of Traditional Chinese Medicine*, 27(5), 325-326.
- Gavaghan, C. L., Holmes, E., Lenz, E., Wilson, I. D., & Nicholson, J. K. (2000). An NMR-based metabonomic approach to investigate the biochemical consequences of genetic strain differences: application to the C57BL10J and Alpk: ApfCD mouse. *FEBS Letters*, 484(3), 169-174.
- Gedulin, B. R., Nikoulina, S. E., Smith, P. A., Gedulin, G., Nielsen, L. L., Baron, A. D., Parkes, D. G., & Young, A. A. (2005). Exenatide (exendin-4) improves insulin sensitivity and β -cell mass in insulin-resistant obese fa/fa Zucker rats independent of glycemia and body weight. *Endocrinology*, 146(4), 2069-2076.
- Giridharan, P., Somasundaram, S., Perumal, K., Vishwakarma, R., Karthikeyan, N., Velmurugan, R., & Balakrishnan, A. (2002). Novel substituted methylenedioxy lignan suppresses proliferation of cancer cells by inhibiting telomerase and activation of c-myc and caspases leading to apoptosis. *British Journal of Cancer*, 87(1), 98-105.
- Glauber, H. S., Rishe, N., & Karnieli, E. (2014). Introduction to personalized medicine in diabetes mellitus. *Rambam Maimonides Medical Journal*, 5(1), e0002.
- Gotoh, M., Li, C., Yatoh, M., Okabayashi, N., Habu, S., & Hirooka, Y. (2006). Hypothalamic monoamine metabolism is different between the diabetic GK (Goto-Kakizaki) rats and streptozotocin-induced diabetic rats. *Brain Research*, 1073, 497-501.
- Grebenstein, C., Choi, Y. H., Rong, J., de Jong, T. J., & Tamis, W. L. M. (2011). Metabolic fingerprinting reveals differences between shoots of wild and cultivated carrot (*Daucus carota* L.) and suggests maternal inheritance or wild trait dominance in hybrids. *Phytochemistry*, 72(11–12), 1341-1347.
- Heyman, H. & Meyer, J. (2012). NMR-based metabolomics as a quality control tool for herbal products. *South African Journal of Botany*, 82, 21-32.
- Hirst, J. A., Farmer, A. J., Ali, R., Roberts, N. W., & Stevens, R. J. (2012). Quantifying the effect of metformin treatment and dose on glycemic control. *Diabetes Care*, 35(2), 446-454.
- Holmes, E., Nicholls, A. W., Lindon, J. C., Connor, S. C., Connelly, J. C., Haselden, J. N., Damment, S. J., Spraul, M., Neidig, P., & Nicholson, J. K. (2000).

- Chemometric models for toxicity classification based on NMR spectra of biofluids. *Chemical Research in Toxicology*, 13(6), 471-478.
- Hsu, B., Coupar, I. M., & Ng, K. (2006). Antioxidant activity of hot water extract from the fruit of the Doum palm, *Hyphaene thebaica*. *Food Chemistry*, 98(2), 317-328.
- Ichiyanagi, T., Kashiwada, Y., Shida, Y., Sekiya, M., Hatano, Y., Takaishi, Y., & Ikeshiro, Y. (2013). Structural elucidation and biological fate of two glucuronyl metabolites of pelargonidin 3-O- β -D-glucopyranoside in rats. *Journal of Agricultural and Food Chemistry*, 61(3), 569-578.
- Ifeoma, O., Samuel, O., Itohan, A. M., & Adeola, S. O. (2013). Isolation, fractionation and evaluation of the antiplasmodial properties of *Phyllanthus niruri* resident in its chloroform fraction. *Asian Pacific Journal of Tropical Medicine*, 6(3), 169-175.
- Ilangkovan, M., Jantan, I., Mohamad, H. F., Husain, K., & Abdul Razak, A. F. (2013). Inhibitory effects of standardized extracts of *Phyllanthus amarus* and *Phyllanthus urinaria* and their marker compounds on phagocytic activity of human neutrophils. *Evidence-Based Complementary and Alternative Medicine*, 2013, 1-9.
- Iqbal, S. & Bhangar, M. (2006). Effect of season and production location on antioxidant activity of *Moringa oleifera* leaves grown in Pakistan. *Journal of Food Composition and Analysis*, 19(6), 544-551.
- Jalil, A. M. M., Ismail, A., Pei, C. P., Hamid, M., & Kamaruddin, S. H. S. (2008). Effects of cocoa extract on glucometabolism, oxidative stress, and antioxidant enzymes in obese-diabetic (Ob-db) rats. *Journal of Agricultural and Food Chemistry*, 56(17), 7877-7884.
- Javadi, N., Abas, F., Hamid, A. A., Simoh, S., Shaari, K., Ismail, I. S., Mediani, A., & Khatib, A. (2014). GC-MS-Based Metabolite Profiling of *Cosmos caudatus* Leaves Possessing Alpha-Glucosidase Inhibitory Activity. *Journal of Food Science*, 79(6), C1130-C1136.
- Jimenez, P., Cabrero, P., Basterrechea, J. E., Tejero, J., Cordoba-Diaz, D., Cordoba-Diaz, M., & Girbes, T. (2014). Effects of short-term heating on total polyphenols, anthocyanins, antioxidant activity and lectins of different parts of dwarf elder (*Sambucus ebulus* L.). *Plant Foods for Human Nutrition*, 69(2), 168-174.
- Kaddurah-Daouk, R., Kristal, B. S., & Weinshilboum, R. M. (2008). Metabolomics: a global biochemical approach to drug response and disease. *Annual Review of Pharmacology and Toxicology*, 48, 653-683.
- Kang, B., Racicot, K., Pilkenton, S. J., & Apostolidis, E. (2014). Evaluation of the In vitro Anti-hyperglycemic Effect of *Cinnamomum cassia* Derived Phenolic Phytochemicals, via Carbohydrate Hydrolyzing Enzyme Inhibition. *Plant Foods for Human Nutrition*, 69(2), 155-160.
- Kassuya, C. A., Leite, D. F., de Melo, L. V., Rehder, V. L. G., & Calixto, J. B. (2005). Anti-inflammatory properties of extracts, fractions and lignans isolated from *Phyllanthus amarus*. *Planta Medica*, 71(8), 721.

- Kemsley, E. K., Le Gall, G., Dainty, J. R., Watson, A. D., Harvey, L. J., Tapp, H. S., & Colquhoun, I. J. (2007). Multivariate techniques and their application in nutrition: a metabolomics case study. *British Journal of Nutrition*, 98(01), 1-14.
- Khanna, A., Rizvi, F., & Chander, R. (2002). Lipid lowering activity of *Phyllanthus niruri* in hyperlipemic rats. *Journal of Ethnopharmacology*, 82(1), 19-22.
- Khaton, S., Rai, V., Rawat, A. K. S., & Mehrotra, S. (2006). Comparative pharmacognostic studies of three *Phyllanthus* species. *Journal of Ethnopharmacology*, 104(1), 79-86.
- Kim, H. K., Choi, Y. H., Erkelens, C., Lefeber, A. W., & Verpoorte, R. (2005). Metabolic fingerprinting of *Ephedra* species using ¹H-NMR spectroscopy and principal component analysis. *Chemical and Pharmaceutical Bulletin*, 53(1), 105-109.
- Kim, J., Kwon, C., & Son, K. H. (2000). Inhibition of Alpha-glucosidase and Amylase by Luteolin, a Flavonoid. *Bioscience, Biotechnology, and Biochemistry*, 64(11), 2458-2461.
- Kim, K., Nam, K., Kurihara, H., & Kim, S. (2008). Potent α -glucosidase inhibitors purified from the red alga *Grateloupia elliptica*. *Phytochemistry*, 69(16), 2820-2825.
- Kim, S., Yang, S., Kim, H., Kim, Y., Park, T., & Choi, H. (2009). ¹H-nuclear magnetic resonance spectroscopy-based metabolic assessment in a rat model of obesity induced by a high-fat diet. *Analytical and Bioanalytical Chemistry*, 395(4), 1117-1124.
- Knowler, W. C., Barrett-Connor, E., Fowler, S. E., Hamman, R. F., Lachin, J. M., Walker, E. A., Nathan, D. M., & Diabetes Prevention Program Research Group. (2002). Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *The New England Journal of Medicine*, 346(6), 393-403.
- Kordalewska, M. & Markuszewski, M. J. (2015). Metabolomics in cardiovascular diseases. *Journal of Pharmaceutical and Biomedical Analysis*, 113, 121-136.
- Kumar, S., Chandra, P., Bajpai, V., Singh, A., Srivastava, M., Mishra, D., & Kumar, B. (2015). Rapid qualitative and quantitative analysis of bioactive compounds from *Phyllanthus amarus* using LC/MS/MS techniques. *Industrial Crops and Products*, 69, 143-152.
- Kumaran, A. & Karunakaran, R. J. (2007). In vitro antioxidant activities of methanol extracts of five *Phyllanthus* species from India. *LWT-Food Science and Technology*, 40(2), 344-352.
- Lanza, I. R., Zhang, S., Ward, L. E., Karakelides, H., Raftery, D., & Nair, K. S. (2010). Quantitative metabolomics by ¹H-NMR and LC-MS/MS confirms altered metabolic pathways in diabetes. *PLoS One*, 5(5), e10538.
- Lawal, U., Mediani, A., Maulidiani, H., Shaari, K., Ismail, I. S., Khatib, A., & Abas, F. (2015). Metabolite profiling of *Ipomoea aquatica* at different growth stages in correlation to the antioxidant and α -glucosidase inhibitory activities elucidated by ¹H NMR-based metabolomics. *Scientia Horticulturae*, 192, 400-408.

- Lee, S. Y., Mediani, A., Nur Ashikin, A. H., Azliana, A. B. S., & Abas, F. (2014). Antioxidant and α -glucosidase inhibitory activities of the leaf and stem of selected traditional medicinal plants. *International Food Research Journal*, 21(1), 165-172.
- Li, H., Xie, Z., Lin, J., Song, H., Wang, Q., Wang, K., Su, M., Qiu, Y., Zhao, T., & Song, K. (2008). Transcriptomic and metabonomic profiling of obesity-prone and obesity-resistant rats under high fat diet. *Journal of Proteome Research*, 7(11), 4775-4783.
- Lim, Y. Y. & Murtijaya, J. (2007). Antioxidant properties of *Phyllanthus amarus* extracts as affected by different drying methods. *LWT - Food Science and Technology*, 40(9), 1664-1669.
- Liu, J., He, T., Lu, Q., Shang, J., Sun, H., & Zhang, L. (2010). Asiatic acid preserves beta cell mass and mitigates hyperglycemia in streptozocin-induced diabetic rats. *Diabetes/Metabolism Research and Reviews*, 26(6), 448-454.
- Lordan, S., Smyth, T. J., Soler-Vila, A., Stanton, C., & Ross, R. P. (2013). The α -amylase and α -glucosidase inhibitory effects of Irish seaweed extracts. *Food Chemistry*, 141(3), 2170-2176.
- Luo, H. & Chen, J. (2007). Metabonomics and CM syndrome type research. *Chinese Journal of Information on TCM*, 14(5), 3-5.
- Luthria, D. L., Lin, L., Robbins, R. J., Finley, J. W., Banuelos, G. S., & Harnly, J. M. (2008). Discriminating between cultivars and treatments of broccoli using mass spectral fingerprinting and analysis of variance principal component analysis. *Journal of Agricultural and Food Chemistry*, 56(21), 9819-9827.
- Lv, H., Zhu, Y., Tan, J., Guo, L., Dai, W., & Lin, Z. (2015). Bioactive compounds from Pu-erh tea with therapy for hyperlipidaemia. *Journal of Functional Foods*, 19, Part A, 194-203.
- Määttä-Riihinen, K. R., Kamal-Eldin, A., & Törrönen, A. R. (2004). Identification and quantification of phenolic compounds in berries of *Fragaria* and *Rubus* species (family Rosaceae). *Journal of Agricultural and Food Chemistry*, 52(20), 6178-6187.
- Madsen, R., Lundstedt, T., & Trygg, J. (2010). Chemometrics in metabolomics—a review in human disease diagnosis. *Analytica Chimica Acta*, 659(1), 23-33.
- Mahdi, E. S., Noor, A. M., Sakeena, M. H., Abdullah, G. Z., Abdulkarim, M., & Sattar, M. A. (2011). Identification of phenolic compounds and assessment of in vitro antioxidants activity of 30% ethanolic extracts derived from two *Phyllanthus* species indigenous to Malaysia. *African Journal of Pharmacy and Pharmacology*, 5(17), 1967-1978.
- Maisuthisakul, P., Pasuk, S., & Ritthiruangdej, P. (2008). Relationship between antioxidant properties and chemical composition of some Thai plants. *Journal of Food Composition and Analysis*, 21(3), 229-240.
- Markom, M., Hasan, M., Daud, W. R. W., Singh, H., & Jahim, J. M. (2007). Extraction of hydrolysable tannins from *Phyllanthus niruri* Linn.: Effects of solvents and extraction methods. *Separation and Purification Technology*, 52(3), 487-496.

- Matsui, T., Ebuchi, S., Fujise, T., Abesundara, K. J., Doi, S., Yamada, H., & Matsumoto, K. (2004). Strong antihyperglycemic effects of water-soluble fraction of Brazilian propolis and its bioactive constituent, 3, 4, 5-tri-O-caffeoylquinic acid. *Biological and Pharmaceutical Bulletin*, 27(11), 1797-1803.
- Maulidiani, M., Sheikh, B. Y., Mediani, A., Wei, L. S., Ismail, I. S., Abas, F., & Lajis, N. H. (2015). Differentiation of *Nigella sativa* seeds from four different origins and their bioactivity correlations based on NMR-metabolomics approach. *Phytochemistry Letters*, 13, 308-318.
- Maulidiani, H., Khatib, A., Shaari, K., Abas, F., Shitan, M., Kneer, R., Neto, V., & Lajis, N. H. (2012). Discrimination of three pegaga (*Centella*) varieties and determination of growth-lighting effects on metabolites content based on the chemometry of ¹H nuclear magnetic resonance spectroscopy. *Journal of Agricultural and Food Chemistry*, 60(1), 410-417.
- Maulidiani, Abas, F., Khatib, A., Shaari, K., & Lajis, N. H. (2014). Chemical characterization and antioxidant activity of three medicinal Apiaceae species. *Industrial Crops and Products*, 55, 238-247.
- Maulidiani, Abas, F., Khatib, A., Shitan, M., Shaari, K., & Lajis, N. H. (2013). Comparison of Partial Least Squares and Artificial Neural Network for the prediction of antioxidant activity in extract of Pegaga (*Centella*) varieties from ¹H Nuclear Magnetic Resonance spectroscopy. *Food Research International*, 54(1), 852-860.
- Mediani, A., Abas, F., Khatib, A., Maulidiani, H., Shaari, K., Choi, Y. H., & Lajis, N. (2012a). ¹H-NMR-based metabolomics approach to understanding the drying effects on the phytochemicals in *Cosmos caudatus*. *Food Research International*, 49(2), 763-770.
- Mediani, A., Abas, F., Khatib, A., & Tan, C. P. (2013). *Cosmos Caudatus* as a potential source of polyphenolic compounds: Optimisation of oven drying conditions and characterisation of its functional properties. *Molecules*, 18(9), 10452-10464.
- Mediani, A., Abas, F., Ping, T. C., Khatib, A., & Lajis, N. H. (2012b). Influence of Growth Stage and Season on the Antioxidant Constituents of *Cosmos caudatus*. *Plant Foods for Human Nutrition*, 67(4), 344-350.
- Micali, S., Sighinolfi, M., Celia, A., De Stefani, S., Grande, M., Cicero, A., & Bianchi, G. (2006). Can *Phyllanthus niruri* affect the efficacy of extracorporeal shock wave lithotripsy for renal stones? A randomized, prospective, long-term study. *The Journal of Urology*, 176(3), 1020-1022.
- Mkele, G. (2013). A review of metformin and its place in the diabetes guidelines. *South African Family Practice*, 55(6), 504-506.
- Moreira, J., Klein-Júnior, L. C., Cechinel Filho, V., & de Campos Buzzi, F. (2013). Anti-hyperalgesic activity of corilagin, a tannin isolated from *Phyllanthus niruri* L.(Euphorbiaceae). *Journal of Ethnopharmacology*, 146(1), 318-323.
- Murugaiyah, V. & Chan, K. (2009). Mechanisms of antihyperuricemic effect of *Phyllanthus niruri* and its lignan constituents. *Journal of Ethnopharmacology*, 124(2), 233-239.

- Musabayane, C., Tufts, M., & Mapanga, R. (2010). Synergistic antihyperglycemic effects between plant-derived oleanolic acid and insulin in streptozotocin-induced diabetic rats. *Renal Failure*, 32(7), 832-839.
- Naik, A. D. & Juvekar, A. R. (2003). Effects of alkaloidal extract of *Phyllanthus niruri* on HIV replication. *Indian Journal of Medical Sciences*, 57(9), 387-393.
- Nardini, M., Cirillo, E., Natella, F., & Scaccini, C. (2002). Absorption of phenolic acids in humans after coffee consumption. *Journal of Agricultural and Food Chemistry*, 50(20), 5735-5741.
- Nemoto, T., Ando, I., Kataoka, T., Arifuku, K., Kanazawa, K., Natori, Y., & Fujiwara, M. (2007). NMR metabolic profiling combined with two-step principal component analysis for toxin-induced diabetes model rat using urine. *The Journal of Toxicological Sciences*, 32(4), 429-435.
- Nicholson, J. K. & Lindon, J. C. (2008). Systems biology: metabonomics. *Nature*, 455(7216), 1054-1056.
- Okoli, C., Ibiam, A., Ezike, A., Akah, P., & Okoye, T. (2010). Evaluation of antidiabetic potentials of *Phyllanthus niruri* in alloxan diabetic rats. *African Journal of Biotechnology*, 9(2), 248-259.
- Okoli, C., Obidike, I., Ezike, A., Akah, P., & Salawu, O. (2011). Studies on the possible mechanisms of antidiabetic activity of extract of aerial parts of *Phyllanthus niruri*. *Pharmaceutical Biology*, 49(3), 248-255.
- Otsuka, R., Tamakoshi, K., Yatsuya, H., Murata, C., Sekiya, A., Wada, K., Zhang, H. M., Matsushita, K., Sugiura, K., & Takefuji, S. (2006). Eating fast leads to obesity: findings based on self-administered questionnaires among middle-aged Japanese men and women. *Journal of Epidemiology*, 16(3), 117-124.
- Otunola, G. A. & Afolayan, A. J. (2015). Antidiabetic effect of combined spices of *Allium sativum*, *Zingiber officinale* and *Capsicum frutescens* in alloxan-induced diabetic rats. *Frontiers in Life Science*, 1-10.
- Pandey, K. B. & Rizvi, S. I. (2009). Plant polyphenols as dietary antioxidants in human health and disease. *Oxidative Medicine and Cellular Longevity*, 2(5), 270-278.
- Park, S., Sadanala, K. C., & Kim, E. K. (2015). A metabolomic approach to understanding the metabolic link between obesity and diabetes. *Molecules and Cells*, 38(7), 587-596.
- Patel, P., Harde, P., Pillai, J., Darji, N., & Patel, B. (2012). Antidiabetic herbal drugs a review. *Pharmacophore*, 3(1), 18-29.
- Patel, J. R., Tripathi, P., Sharma, V., Chauhan, N. S., & Dixit, V. K. (2011). *Phyllanthus amarus*: Ethnomedicinal uses, phytochemistry and pharmacology: A review. *Journal of Ethnopharmacology*, 138(2), 286-313.
- Phinney, K. W., Ballihaut, G., Bedner, M., Benford, B. S., Camara, J. E., Christopher, S. J., Davis, W. C., Dodder, N. G., Eppe, G., & Lang, B. E. (2013). Development of a Standard Reference Material for metabolomics research. *Analytical Chemistry*, 85(24), 11732-11738.
- Punithavathi, V. R., Prince, P. S. M., Kumar, R., & Selvakumari, J. (2011). Antihyperglycaemic, antilipid peroxidative and antioxidant effects of gallic

acid on streptozotocin induced diabetic Wistar rats. *European Journal of Pharmacology*, 650(1), 465-471.

- Rani, S. & Kumar, B. (2015). Glycemic control efficacy of *Phyllanthus niruri*. Linn extract in Diabetic Mice model. *Int.J.Curr.Microbiol.App.Sci*, 4(4), 599-609.
- Rastija, V. (2011). An overview of innovations in analysis and beneficial health effects of wine polyphenols. *Mini Reviews in Medicinal Chemistry*, 11(14), 1256-1267.
- Ribeiro, P. R., Willems, L. A., Mudde, E., Fernandez, L. G., de Castro, R. D., Ligterink, W., & Hilhorst, H. W. (2015). Metabolite profiling of the oilseed crop *Ricinus communis* during early seed imbibition reveals a specific metabolic signature in response to temperature. *Industrial Crops and Products*, 67, 305-309.
- Ribnicky, D. M., Kuhn, P., Poulev, A., Logendra, S., Zuberi, A., Cefalu, W. T., & Raskin, I. (2009). Improved absorption and bioactivity of active compounds from an anti-diabetic extract of *Artemisia dracunculus* L. *International Journal of Pharmaceutics*, 370(1), 87-92.
- Rochfort, S. (2005). Metabolomics reviewed: a new “omics” platform technology for systems biology and implications for natural products research. *Journal of Natural Products*, 68(12), 1813-1820.
- Rofiee, M., Yusof, M., Hisam, E. A., Bannur, Z., Zakaria, Z., Somchit, M., Teh, L., & Salleh, M. (2015). Isolating the metabolic pathways involved in the hepatoprotective effect of *Muntingia calabura* against CCl₄-induced liver injury using LC/MS Q-TOF. *Journal of Ethnopharmacology*, 166, 109-118.
- Roos, G., Roseler, C., Buter, K. B., & Simmen, U. (2004). Classification and correlation of St. John's wort extracts by nuclear magnetic resonance spectroscopy, multivariate data analysis and pharmacological activity. *Planta Medica*, 70(8), 771-777.
- Russell, C., Rahman, A., & Mohammed, A. R. (2013). Application of genomics, proteomics and metabolomics in drug discovery, development and clinic. *Therapeutic Delivery*, 4(3), 395-413.
- Sabir, S. & Rocha, J. (2008). Water-extractable phytochemicals from *Phyllanthus niruri* exhibit distinct in vitro antioxidant and in vivo hepatoprotective activity against paracetamol-induced liver damage in mice. *Food Chemistry*, 111(4), 845-851.
- Salek, R. M., Maguire, M. L., Bentley, E., Rubtsov, D. V., Hough, T., Cheeseman, M., Nunez, D., Sweatman, B. C., Haselden, J. N., Cox, R. D., Connor, S. C., & Griffin, J. L. (2007). A metabolomic comparison of urinary changes in type 2 diabetes in mouse, rat, and human. *Physiological Genomics*, 29(2), 99-108.
- Shaham, O., Wei, R., Wang, T. J., Ricciardi, C., Lewis, G. D., Vasan, R. S., Carr, S. A., Thadhani, R., Gerszten, R. E., & Mootha, V. K. (2008). Metabolic profiling of the human response to a glucose challenge reveals distinct axes of insulin sensitivity. *Molecular Systems Biology*, 4(1), 214-223.
- Sharma, P., Gujral, H. S., & Singh, B. (2012). Antioxidant activity of barley as affected by extrusion cooking. *Food Chemistry*, 131(4), 1406-1413.

- Sharma, V., Dhillon, P., Wambolt, R., Parsons, H., Brownsey, R., Allard, M. F., & McNeill, J. H. (2008). Metoprolol improves cardiac function and modulates cardiac metabolism in the streptozotocin-diabetic rat. *American Journal of Physiology. Heart and Circulatory Physiology*, 294(4), H1609-1620.
- Shetty, A., Rashmi, R., Rajan, M., Sambaiah, K., & Salimath, P. (2004). Antidiabetic influence of quercetin in streptozotocin-induced diabetic rats. *Nutrition Research*, 24(5), 373-381.
- Shuib, N. H., Shaari, K., Khatib, A., Maulidiani, Kneer, R., Zareen, S., Raof, S. M., Hj. Lajis, N., & Neto, V. (2011). Discrimination of young and mature leaves of *Melicope ptelefolia* using ¹H NMR and multivariate data analysis. *Food Chemistry*, 126(2), 640-645.
- Siddhuraju, P. & Becker, K. (2003). Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *Journal of Agricultural and Food Chemistry*, 51(8), 2144-2155.
- Song, L., Liu, H., Wang, Y., Wang, Y., Liu, J., Zhou, Z., Chu, H., Zhuang, P., & Zhang, Y. (2015). Application of GC/MS-based metabonomic profiling in studying the therapeutic effects of Huangbai–Zhimu herb-pair (HZ) extract on streptozotocin-induced type 2 diabetes in mice. *Journal of Chromatography B*, 997, 96-104.
- Soong, Y. Y. & Barlow, P. J. (2005). Isolation and structure elucidation of phenolic compounds from longan (*Dimocarpus longan* Lour.) seed by high-performance liquid chromatography–electrospray ionization mass spectrometry. *Journal of Chromatography A*, 1085(2), 270-277.
- Srinivasan, K. & Ramarao, P. (2007). Animal models in type 2 diabetes research: an overview. *Indian Journal of Medical Research*, 125(3), 451.
- Srinivasan, K., Viswanad, B., Asrat, L., Kaul, C., & Ramarao, P. (2005). Combination of high-fat diet-fed and low-dose streptozotocin-treated rat: a model for type 2 diabetes and pharmacological screening. *Pharmacological Research*, 52(4), 313-320.
- Sudjaroen, Y., Hull, W. E., Erben, G., Würtele, G., Changbumrung, S., Ulrich, C. M., & Owen, R. W. (2012). Isolation and characterization of ellagitannins as the major polyphenolic components of Longan (*Dimocarpus longan* Lour) seeds. *Phytochemistry*, 77, 226-237.
- Suhre, K. (2014). Metabolic profiling in diabetes. *The Journal of Endocrinology*, 221(3), R75-85.
- Sulaiman, S. F., Sajak, A. A. B., Ooi, K. L., Supriatno, & Seow, E. M. (2011). Effect of solvents in extracting polyphenols and antioxidants of selected raw vegetables. *Journal of Food Composition and Analysis*, 24(4–5), 506-515.
- Sultana, B., Anwar, F., & Przybylski, R. (2007). Antioxidant activity of phenolic components present in barks of *Azadirachta indica*, *Terminalia arjuna*, *Acacia nilotica*, and *Eugenia jambolana* Lam. trees. *Food Chemistry*, 104(3), 1106-1114.

- Sultana, B., Anwar, F., & Ashraf, M. (2009). Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. *Molecules*, 14(6), 2167-2180.
- TA, S. (2014). Diagnosis and classification of diabetes mellitus. *Diabetes Care*, 37, S81.
- Talavéra, S., Felgines, C., Texier, O., Besson, C., Mazur, A., Lamaison, J., & Rémésy, C. (2006). Bioavailability of a bilberry anthocyanin extract and its impact on plasma antioxidant capacity in rats. *Journal of the Science of Food and Agriculture*, 86(1), 90-97.
- Talaviya, P. A., Rao, S. K., Vyas, B. M., Indoria, S. P., Suman, R. K., & Suvagiya, V. P. (2014). A review on: Potential antidiabetic herbal medicines. *International Journal of Pharmaceutical Sciences and Research*, 5(2), 302.
- Tarachiwin, L., Masako, O., & Fukusaki, E. (2008). Quality evaluation and prediction of *Citrullus lanatus* by ^1H NMR-based metabolomics and multivariate analysis. *Journal of Agricultural and Food Chemistry*, 56(14), 5827-5835.
- Teul, J., Rupérez, F. J., Garcia, A., Vaysse, J., Balayssac, S., Gilard, V., Malet-Martino, M., Martin-Ventura, J. L., Blanco-Colio, L. M., & Tuñón, J. (2009). Improving metabolite knowledge in stable atherosclerosis patients by association and correlation of GC-MS and ^1H NMR fingerprints. *Journal of Proteome Research*, 8(12), 5580-5589.
- Tharakan, S. T. (2011). 1 Taxonomy of the Genus *Phyllanthus*. *Phyllanthus Species: Scientific Evaluation and Medicinal Applications*, 1.
- Trabelsi, N., Megdiche, W., Ksouri, R., Falleh, H., Oueslati, S., Soumaya, B., Hajlaoui, H., & Abdelly, C. (2010). Solvent effects on phenolic contents and biological activities of the halophyte *Limoniastrum monopetalum* leaves. *LWT-Food Science and Technology*, 43(4), 632-639.
- Ueno, H., Horie, S., Nishi, Y., Shogawa, H., Kawasaki, M., Suzuki, S., Hayashi, T., Arisawa, M., Shimizu, M., & Yoshizaki, M. (1988). Chemical and Pharmaceutical Studies on Medicinal Plants in Paraguay, Geraniin, an Angiotensin-Converting Enzyme Inhibitor from "Paraparai Mi," *Phyllanthus niruri*. *Journal of Natural Products*, 51(2), 357-359.
- Ursem, R., Tikunov, Y., Bovy, A., Van Berloo, R., & Van Eeuwijk, F. (2008). A correlation network approach to metabolic data analysis for tomato fruits. *Euphytica*, 161(1-2), 181-193.
- Van Der Werf, Mariet J, Jellema, R. H., & Hankemeier, T. (2005). Microbial metabolomics: replacing trial-and-error by the unbiased selection and ranking of targets. *Journal of Industrial Microbiology and Biotechnology*, 32(6), 234-252.
- Verma, S., Sharma, H., & Garg, M. (2014). *Phyllanthus amarus*: A Review. *Journal of Pharmacognosy and Phytochemistry*, 3(2), 18-22.
- Verpoorte, R., Choi, Y., & Kim, H. (2007). NMR-based metabolomics at work in phytochemistry. *Phytochemistry Reviews*, 6(1), 3-14.

- Vinayavekhin, N., Homan, E. A., & Saghatelian, A. (2009). Exploring disease through metabolomics. *ACS Chemical Biology*, 5(1), 91-103.
- Wang, P., Wang, J., Yang, M., & Kong, L. (2014). Neuroprotective effects of Huang-Lian-Jie-Du-Decoction on ischemic stroke rats revealed by ¹H NMR metabolomics approach. *Journal of Pharmaceutical and Biomedical Analysis*, 88, 106-116.
- Wang, W., Lee, P., Yeh, H., Smirnova, I. V., & Choi, I. (2012). Effects of acute and chronic hyperglycemia on the neurochemical profiles in the rat brain with streptozotocin-induced diabetes detected using in vivo ¹H NMR spectroscopy at 9.4 T. *Journal of Neurochemistry*, 121(3), 407-417.
- Wang, Y., Bollard, M. E., Keun, H., Antti, H., Beckonert, O., Ebbels, T. M., Lindon, J. C., Holmes, E., Tang, H., & Nicholson, J. K. (2003). Spectral editing and pattern recognition methods applied to high-resolution magic-angle spinning ¹H nuclear magnetic resonance spectroscopy of liver tissues. *Analytical Biochemistry*, 323(1), 26-32.
- Wansi, J. D., Lallemand, M., Chiozem, D. D., Toze, F. A. A., Mbaze, L. M., Naharkhan, S., Iqbal, M. C., Tillequin, F., Wandji, J., & Fomum, Z. T. (2007). α -Glucosidase inhibitory constituents from stem bark of *Terminalia superba* (Combretaceae). *Phytochemistry*, 68(15), 2096-2100.
- Wei, T., Zhao, L., Jia, J., Xia, H., Du, Y., Lin, Q., Lin, X., Ye, X., Yan, Z., & Gao, H. (2015). Metabonomic analysis of potential biomarkers and drug targets involved in diabetic nephropathy mice. *Scientific Reports*, 5.
- WHO (2014) (<http://www.who.int/mediacentre/factsheets/fs312/en/>).
- Wold, S., Sjöström, M., & Eriksson, L. (2001). PLS-regression: a basic tool of chemometrics. *Chemometrics and Intelligent Laboratory Systems*, 58(2), 109-130.
- Xia, J., Sinelnikov, I. V., Han, B., & Wishart, D. S. (2015). MetaboAnalyst 3.0--making metabolomics more meaningful. *Nucleic Acids Research*, 43(W1), W251-257.
- Xie, X., Wang, Y., Zuo, C., Fan, J., & Li, X. (2009). A case report of an effective treatment for diabetic foot ulcers with integration of traditional Chinese medicine and Western medicine. *Journal of Diabetes and its Complications*, 23(5), 360-364.
- Yan, J., Zhang, G., Pan, J., & Wang, Y. (2014). α -Glucosidase inhibition by luteolin: Kinetics, interaction and molecular docking. *International Journal of Biological Macromolecules*, 64, 213-223.
- Yang, C., Cheng, H., Lin, T., Chiang, L., & Lin, C. (2007). The in vitro activity of geraniin and 1, 3, 4, 6-tetra-O-galloyl- β -d-glucose isolated from *Phyllanthus urinaria* against herpes simplex virus type 1 and type 2 infection. *Journal of Ethnopharmacology*, 110(3), 555-558.
- Yang, C., Cheng, H., Lin, T., Chiang, L., & Lin, C. (2005). Acetone, ethanol and methanol extracts of *Phyllanthus urinaria* inhibit HSV-2 infection in vitro. *Antiviral Research*, 67(1), 24-30.

- Yuan, T., Ding, Y., Wan, C., Li, L., Xu, J., Liu, K., Slitt, A., Ferreira, D., Khan, I. A., & Seeram, N. P. (2012). Antidiabetic ellagitannins from pomegranate flowers: inhibition of α -glucosidase and lipogenic gene expression. *Organic Letters*, 14(20), 5358-5361.
- Zafrilla, P., Ferreres, F., & Tomás-Barberán, F. A. (2001). Effect of processing and storage on the antioxidant ellagic acid derivatives and flavonoids of red raspberry (*Rubus idaeus*) jams. *Journal of Agricultural and Food Chemistry*, 49(8), 3651-3655.
- Zhang, J., Liu, Q., Liu, Z., Li, L., & Yi, L. (2015). Antihyperglycemic activity of Anoctochilus roxburghii polysaccharose in diabetic mice induced by high-fat diet and streptozotocin. *Journal of Ethnopharmacology*, 164, 180-185.
- Zhang, S., Gowda, G. N., Asiago, V., Shanaiah, N., Barbas, C., & Raftery, D. (2008). Correlative and quantitative ^1H NMR-based metabolomics reveals specific metabolic pathway disturbances in diabetic rats. *Analytical Biochemistry*, 383(1), 76-84.
- Zhang, L. Z., Guo, Y. J., Tu, G. Z., Guo, W. B., & Miao, F. (2000). Studies on chemical constituents of *Phyllanthus urinaria* L. *Zhongguo Zhong Yao Za Zhi Zhongguo Zhongyao Zazhi China Journal of Chinese Materia Medica*, 25(10), 615-617.
- Zhou, S., Xu, C., Zhou, N., Huang, Y., Huang, L., Chen, X., Hu, Y., & Liao, Y. (1997). Mechanism of protective action of *Phyllanthus urinaria* L. against injuries injuries of liver cells. *China Journal of Chinese Materia Medica*, 22(2), 109-111.
- Zimmet, P., Alberti, K., & Shaw, J. (2001). Global and societal implications of the diabetes epidemic. *Nature*, 414(6865), 782-787.