



UNIVERSITI PUTRA MALAYSIA

***MECHANISM OF HEPATOPROTECTIVE ACTIVITY OF METHANOLIC
EXTRACT OF *Melastoma malabathricum* L. LEAVES***

SITI SYARIAH BINTI MAMAT

FPSK(M) 2017 15



**MECHANISM OF HEPATOPROTECTIVE ACTIVITY OF METHANOLIC
EXTRACT OF *Melastoma malabathricum* L. LEAVES**

By

SITI SYARIAH BINTI MAMAT

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Master of Science**

February 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

MECHANISM OF HEPATOPROTECTIVE ACTIVITY OF METHANOLIC EXTRACT OF *Melastoma malabathricum* L. LEAVES

By

SITI SYARIAH BINTI MAMAT

February 2017

Chairman : Associate Professor Zainul Amiruddin Zakaria, PhD
Faculty : Medicine and Health Sciences

The objective of this study was to determine the hepatoprotective activity of methanolic extracts of *Melastoma malabathricum* leaves (MEMM) and its partitions using rats' model, by evaluating the prophylactic effect of the plants extracts administered prior to the induction of liver toxicity using a hepatotoxic agent. The study were design as hepatoprotective potential of MEMM has never been reported. In an attempt to establish the pharmacological properties, the hepatoprotective potential of MEMM was investigated using carbon tetrachloride (CCl₄)- and paracetamol (PCM)- induced hepatotoxicity in rats. Throughout this study, the animals were divided into 22 groups containing 6 rats each group. In the first stage of *in vivo* study, rats were divided into groups and administered orally once daily with 10 % dimethyl sulfoxide (DMSO) as negative control, 200 mg/kg silymarin as positive control, or MEMM (50, 250, 500 mg/kg) for 7 days, followed by hepatotoxicity induction using CCl₄ or PCM. In the second stage, MEMM was partitioned into 3 fractions: petroleum ether extract (PEMM), ethyl acetate extract (EAMM), aqueous extract (AQMM). PEMM, EAMM and AQMM were then tested on PCM-induced hepatotoxicity in rats. Blood sample underwent biochemical analysis to evaluate alanine transferase (ALT), aspartate aminotransferase (AST) and alkaline phosphatase (ALP) levels; the livers were subjected to microscopic analysis. All extracts (MEMM, PEMM, EAMM AQMM) underwent antioxidant study using oxygen radical absorbance capacity (ORAC) test, Total phenolic compound (TPC), 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity and superoxide scavenging assay, and anti-inflammatory evaluation via lipoxygenase (LOX) and xanthine oxidase (XO) assays. Total phenolic content (TPC), phytochemical screening and high-performance liquid chromatography (HPLC) evaluation were also performed. From the histological observation, lymphocyte infiltration and

marked necrosis were observed in the DMSO-treated groups (negative control). MEMM showed encouraging activity for reducing the toxic effect of CCl₄ or PCM on the liver by reducing the weight of the liver in a dose independent manner; histological observation demonstrated normalization of the histopathological changes, preserving hepatocyte structure, causing a significant decline in AST and ALT levels ($p < 0.05$). PEMM, EAMM and AQMM which contains non-polar compounds, intermediate compounds and polar compounds, respectively, attenuated the liver enzyme levels in a dose-independent manner. Overall, EAMM had the best activity for attenuating the liver enzymes. MEMM had the highest TPC value, followed by AQMM, EAMM, and PEMM. All the extracts (MEMM, PEMM, EAMM, and AQMM) demonstrated potential free radical scavenging activity in SOD and DPPH assays. However, the different trend was showed by ORAC assay from that DPPH and SOD. EAMM and AQMM had high ORAC value, which determine the capacity of an extract to act as an antioxidant. In the anti-inflammatory assays, MEMM and EAMM showed the moderate inhibition in LOX activity, but weak anti-inflammatory activity in XO. Phytochemical screening showed that the extracts showed that MEMM, PEMM and EAMM contained flavonoids, triterpenes, tannins, saponins, polyphenolic compounds and steroids, but not alkaloids. In contrast, the AQMM extract contained fewer compounds. HPLC analysis demonstrated that four to eight peaks detected at different wavelengths of the chromatogram of MEMM, PEMM, EAMM and AQMM, which were suggested to be flavonoid-based compounds. In conclusions, MEMM exerted potential hepatoprotective activity that can be partly attributed to its antioxidant activity, and EAMM was considered to have the best activity among the fractions, which warrants further investigation.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**MEKANISMA AKTIVITI HEPATOPROTEKTIF OLEH EKTRAK
METANOL DARI DAUN *Melastoma malabathricum***

Oleh

SITI SYARIAH BINTI MAMAT

Februari 2017

Pengerusi : Profesor Madya Zainul Amiruddin Zakaria, PhD
Fakulti : Perubatan dan Sains Kesihatan

Objektif kajian ini adalah untuk menentukan aktiviti hepatoprotektif ekstrak metanol daripada daun *Melastoma malabathricum* dan pecahannya dengan menggunakan model tikus dengan menilai kesan profilaksi ekstrak tumbuhan yang diambil sebelum induksi ketoksikan hati menggunakan agen hepatotoksik. Kajian ini telah direka kerana potensi hepatoprotektif dari ekstrak MEMM tidak pernah dilaporkan. Dalam usaha untuk mewujudkan sifat-sifat farmakologi, potensi hepatoprotektif MEMM telah dikaji dengan menggunakan induksi karbon tetraklorida (CCl₄) - dan paracetamol (PCM) pada tikus. Sepanjang kajian ini, tikus telah dibahagikan kepada 22 kumpulan yang mempunyai 6 tikus bagi setiap kumpulan. Dalam peringkat pertama dalam kajian *in vivo*, tikus telah dibahagikan kepada kumpulan dan diberi makan sekali sehari dengan 10% dimetil sulfoxide (DMSO) sebagai kawalan negatif, 200 mg/kg silymarin sebagai kawalan positif, atau MEMM (50, 250, 500 mg / kg) selama 7 hari, diikuti dengan hepatoksiti induksi menggunakan CCl₄ atau PCM. Pada peringkat kedua, MEMM telah dibahagikan kepada 3 pecahan: ekstrak petroleum eter (PEMM), ekstrak etil asetat (EAMM), ekstrak akueus (AQMM). PEMM, EAMM dan AQMM kemudiannya diuji ke atas ransangan PCM pada tikus. Sampel darah telah diambil untuk menjalani analisis biokimia untuk menilai paras enzim seperti transferase alanine (ALT), aspartat aminotransferase (AST) dan tahap alkali phosphatase (ALP); sampel hati pula dinilai secara mikroskopik. Semua ekstrak (MEMM, PeMM, EAMM AQMM) menjalani kajian antioksidan menggunakan oksigen kapasiti penyerapan radikal (ORAC) ujian, Jumlah sebatian fenolik (TPC), 2,2-difenil-1-picrylhydrazyl (DPPH) aktiviti memerangkap radikal dan ujian perangkap aktiviti superoxide dismutase (SOD), dan penilaian anti-radang melalui ujian lipoxigenase (LOX) dan xanthine oxidase (XO). Jumlah kandungan fenolik (TPC), pemeriksaan fitokimia dan kromatografi cecair penilaian berprestasi

tinggi (HPLC) juga telah dijalankan. Dari pemerhatian histologi, penyusupan limfosit dan nekrosis diperhatikan dalam kumpulan yang diberikan DMSO (kawalan negatif). MEMM menunjukkan aktiviti yang menggalakkan bagi mengurangkan kesan toksik CCl₄ atau PCM pada hati dengan mengurangkan berat hati selari dengan jumlah dos yang diberi; pemerhatian histologi menunjukkan perubahan normal, memelihara struktur sel-sel hati, menyebabkan penurunan yang ketara dalam tahap AST dan ALT ($p < 0.05$). PEMM, EAMM dan AQMM yang mengandungi sebatian bukan kutub, sebatian perantaraan dan sebatian kutub, masing-masing, dapat mengurangkan tahap enzim hati tanpa bergantung pada dos. Secara keseluruhan, EAMM mempunyai aktiviti yang terbaik dalam penurunan enzim hati. MEMM mempunyai nilai TPC tertinggi, diikuti oleh AQMM, EAMM dan PEMM. Semua ekstrak (MEMM, PEMM, EAMM dan AQMM) menunjukkan potensi aktiviti memerangkap radikal bebas dalam ujian SOD dan DPPH. Walau bagaimanapun, ikutan yang berbeza telah ditunjukkan oleh ujian ORAC dari DPPH dan SOD. EAMM dan AQMM mempunyai nilai ORAC yang tinggi, yang menentukan kapasiti untuk bertindak sebagai antioksidan. Dalam ujian anti-radang, MEMM dan EAMM menunjukkan perencatan yang sederhana dalam aktiviti LOX, tetapi aktiviti anti-radang lemah dalam XO. Penyaringan fitokimia menunjukkan bahawa ekstrak menunjukkan bahawa MEMM, PEMM dan EAMM mengandungi flavonoid, triterpenes, tanin, saponin, sebatian polifenolik dan steroid, tetapi tidak alkaloid. Sebaliknya, ekstrak AQMM mengandungi hanya beberapa sebatian sahaja. Analisis HPLC menunjukkan bahawa empat hingga lapan puncak dikesan pada panjang gelombang yang berbeza daripada kromatogram daripada MEMM, PEMM, EAMM dan AQMM, yang telah dikategorikan sebagai jenis-jenis sebatian flavonoid. Sebagai kesimpulan, MEMM dikenakan berpotensi aktiviti hepatoprotective yang boleh sebahagiannya disebabkan oleh aktiviti antioksidan, dan EAMM dianggap mempunyai aktiviti yang terbaik dalam kalangan pecahan ekstrak, yang memerlukan siasatan lanjut.

ACKNOWLEDGEMENT

Alhamdulillah, All Praise to God Almighty for granting me grace and strength to persevere throughout my project and to overcome all the challenges that I had gone through during the project.

I wish to extend my deepest appreciation to my project supervisor, Associate Professor Dr. Zainul Amiruddin Zakaria, for his unrelenting guidance concern, valuable advice, patience, understanding, support and continuous supervision throughout this project. I truly thank him for giving me opportunity to be his postgraduate student.

I would like to express my gratitude and appreciation to my co-supervisor, Professor Teh Lay Kek for her kindness to teach and guide me in animal preparation and Dr. Norhafizah Mohtarrudin for her kindness to teach and guide me in histological study that is definitely important in order to complete this project.

My sincere appreciation dedicates to the Faculty of Medicine and Health Sciences, Universiti Putra Malaysia for giving me opportunity to carry out this project.

My special dedicates to the Pharmacology and Toxicology Research Laboratory science officer and assistants, Histopathology Research Laboratory science officer and assistants, and also Animal House Research Laboratory officer and assistants, for their assistance and patience.

My special also dedicate to all my fellow friends for their constant companion, inspiration, and assistance throughout this project. Last but not list, to my be-loved family, thanks for your unwavering love, encouragements and comfort that helped me to accomplish my project successfully.

I certify that a Thesis Examination Committee has met on 17 February 2017 to conduct the final examination of Siti Syariah binti Mamat on her thesis entitled "Mechanism of Hepatoprotective Activity of Methanolic Extract of *Melastoma malabathricum* L. Leaves" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Arifah binti Abdul Kadir, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Mohd Nasir bin Mohd Desa, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Internal Examiner)

Mohd Zaki Salleh, PhD

Professor
Universiti Teknologi MARA
Malaysia
(External Examiner)



NOR AINI AB. SHUKOR, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 April 2017

The thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Zainul Amiruddin Zakaria, Ph.D.

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Norhafizah bt Mohtarrudin, MBBS

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Member)

Teh Lay Kek, PhD

Professor
Faculty of Pharmacy
Universiti Teknologi MARA
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____

Date: _____

Name and Matric No: Siti Syariah binti Mamat / GS30434

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xviii
CHAPTER	
1 INTRODUCTION	
1.1 Problem statement	2
1.2 Research Hypothesis	3
1.2.1 Null Hypothesis	3
1.2.2 Alternative Hypothesis	3
1.3 Objectives of the Study	3
1.3.1 General Objectives	3
1.3.2 Specific Objectives	3
2 LITERATURE REVIEW	
2.1 Liver	
2.1.1 Gross morphology of the liver	4
2.1.2 Vascular system of the liver	4
2.1.3 Microanatomy of the liver	4
2.1.4 Functions of the liver	5
2.1.5 Role of drug metabolism and detoxification	6
2.1.6 Enzyme liver function	6
2.1.7 Disease of the liver	7
2.1.8 Mechanism of liver injury	7
2.2 Natural products	8
2.2.1 Plant-based products	9
2.2.2 Hepatoprotective-related natural products	9
2.2.3 <i>Melastoma malabathricum</i>	9
2.2.3.1 Geographical site	9
2.2.3.2 Botanical description	10
2.2.3.3 Traditional usages	11
2.2.3.4 Scientific findings	11
2.2.3.5 Phytochemical and chemical constituents	14
2.3 Hepatotoxicity model	16
2.3.1 Carbon tetrachloride	16
2.3.2 Paracetamol	17

3	EVALUATION OF HEPATOPROTECTIVE ACTIVITY OF METHANOL EXTRACT OF <i>MELASTOMA MALABATHRICUM</i> LEAVES AND ITS PARTITIONS	
3.1	Introduction	18
3.2	Methodology	19
3.2.1	Preparation of Methanol Extract of <i>Melastoma malabathricum</i> (MEMM)	19
3.2.1.1	Plant material collection	19
3.2.1.2	Preparation of crude methanol extract	19
3.2.1.3	Preparation of aqueous, ethyl acetate and petroleum ether plant extract	19
3.2.2	Preparation of normal, positive and negative control solution	19
3.2.3	Animals	20
3.2.4	Chemicals	20
3.2.5	Hepatotoxicity assays	20
3.2.6	Biochemical analysis	22
3.2.7	Histopathological investigation	22
3.2.8	Statistical analysis	23
3.3	Results	24
3.3.1	Production of methanolic extract of <i>m. malabathricum</i> and its partitions	24
3.3.2	Effect of MEMM on the body weight and liver weight after induction with CCl ₄	25
3.3.3	Macroscopic and microscopic study of the CCl ₄ -intoxicated liver with and without pre-treatment with MEMM	27
3.3.4	Biochemical parameters of the CCl ₄ -intoxicated liver with and without pre-treatment with MEMM	32
3.3.5	Effect of MEMM on the body weight and liver weight after induction with PCM	34
3.3.6	Macroscopic and microscopic study of the PCM-intoxicated liver with and without pre-treatment with MEMM	36
3.3.7	Biochemical parameters of the PCM-intoxicated liver with and without pre-treatment with MEMM	41
3.3.8	Effect of PEMM, EAMM and AQMM on the body weight and liver weight after induction with PCM	43
3.3.9	Microscopic study of the PCM-intoxicated liver with and without pre-treatment with PEMM, EAMM and AQMM	46
3.3.10	Biochemical parameters of the PCM-intoxicated liver with and without pre-treatment with PEMM, EAMM and AQMM	53
3.4	Discussion	55

4	EVALUATION OF THE ANTIOXIDANT ANTI- INFLAMMATORY ACTIVITIES OF THE METHANOLIC EXTRACT OF <i>MELASTOMA MALABATHRICUM</i> LEAVES AND ITS PARTITION	
4.1	Introduction	58
4.2	Methodology	60
4.2.1	Antioxidant assays	60
4.2.1.1	Oxygen Radical Absorbance Capacity (ORAC) test	60
4.2.1.2	Total Phenolic Content (TPC)	60
4.2.1.3	2,2-Diphenyl-2-Picryl-Hydrazyl (DDPH) Radical Scavenging Activity	61
4.2.1.4	Superoxide scavenging assay	61
4.2.2	Anti-inflammatory assays	61
4.2.2.1	Xanthine oxidase assay	61
4.2.2.2	Lipoxygenase assay	62
4.3	Results	62
4.3.1	2,2-Diphenyl-2-Picryl-Hydrazyl (DDPH) Radical Scavenging Activity	62
4.3.2	Superoxide scavenging assay	62
4.3.3	Total Phenolic Content (TPC)	62
4.3.4	Oxygen Radical Absorbance Capacity (ORAC) test	63
4.3.5	Xanthine oxidase assay and lipoxygenase activity	63
4.4	Discussion	68
5	PHYTOCHEMICAL SCREENING AND HIGH PERFORMANCE LIQUID CHROMATOGRAPHY PROFILING OF THE METHANOLIC EXTRACT OF <i>MELASTOMA MALABATHRICUM</i> AND ITS PARTITION	
5.1	Introduction	70
5.2	Methodology	71
5.2.1	Phytochemical screening	71
5.2.1.1	Saponins test	71
5.2.1.2	Alkaloids test	71
5.2.1.3	Flavonoids test	71
5.2.1.4	Tannins and Polyphenolic Compounds test	71
5.2.1.5	Triterpenes/Steroids test	71
5.2.2	HPLC profiling	72
5.2.3	Identification of flavonoids in MEMM via HPLC	72
5.3	Result	72
5.3.1	Phytochemical screening	72
5.3.2	HPLC profiling of MEMM	73
5.3.3	Identification of flavonoids in MEMM via HPLC HPLC and UV profiling of the partitions of	77
5.3.4	PEMM, EAMM and AQMM	77
5.4	Discussion	84

6	GENERAL DISCUSSION	86
7	SUMMARY, CONCLUSION AND RECOMMENDATION FOR FUTURE STUDY	90
	REFERENCES	91
	APPENDICES	106
	BIODATA OF STUDENT	108
	LIST OF PUBLICATIONS	109



LIST OF TABLES

Table		Page
1	Medicinal uses of <i>M. malabathricum</i> .	12
2	Pharmacological activities of <i>M. malabathricum</i>	13
3	Compounds of <i>M. malabathricum</i>	15
4	CCl ₄ -induced hepatotoxicity treatment groups	21
5	PCM-induced hepatotoxicity treatment groups	21
6	PCM-induced hepatotoxicity treatment groups of partition extracts	21
7	Liver histological scoring	23
8	Effect of MEMM on body and liver weights in CCl ₄ - treated rats	26
9	Histopathological scoring of the CCl ₄ -induced liver injury pre-treated with various doses of MEMM in rats.	26
10	Effect of CCl ₄ and protective treatments at ALT, AST and ALP (U/L)	33
11	Effect of MEMM on body and liver weights in PCM- treated rats	35
12	Histopathological scoring of the PCM-induced liver injury pre-treated with various doses of MEMM in rats.	35
13	Effect of PCM and protective treatments at ALT, AST and ALP (U/L)	42
14	Effect of PEMM, EAMM and AQMM on body and liver weights in PCM- treated rats	44
15	Histopathological scoring of the PCM-induced liver injury pre-treated with various doses of PEMM, EAMM and AQMM in rats	45
16	Effect of PEMM, EAMM and AQMM on liver function test, ALT, AST and ALP (U/L)	54
17	Antioxidant profile of MEMM, PEMM, EAMM and AQMM	64
18	Anti-inflammatory profile of MEMM, PEMM, EAMM and AQMM	67
19	Comparison of phytochemical constituents of the leaves in dried leaves powdered and in different extracts of <i>M. malabathricum</i> leaves	73

LIST OF FIGURES

Figure		Page
1	Location of the liver in human body	5
2	<i>M. malabathricum</i> leaves, flowers and fruits	10
3	The flowchart of method involved in the preparation of MEMM, PEMM, EAMM and AQMM and their yield of product	24
4	4A normal liver, 4B liver intoxicated with 1mL/kg CCl ₄ : gross image shows major color changes of liver lobes from dark marron to brownish with the coarse surface, 4C liver pre-treated with 200mg/kg Silymarin and induced with CCl ₄ : spot of color changes was noted, 4D liver pre-treated with 50mg/kg MEMM and induced by CCl ₄ , 4E liver pre-treated with 250mg/kg MEMM and induced by CCl ₄ , 4F liver pre-treated with 500mg/kg MEMM and induced by CCl ₄ .	28
5	5A Normal, 5B Section of liver tissue of 1mL/kg CCl ₄ -treated group (p.o) showing massive coagulative necrosis, haemorrhage, inflammation and steatosis. (H & E, 40x magnification). CV central vein. N necrosis. H haemorrhage. I inflammation. S steatosis.	29
5	5C Section of 200 mg/kg of Silymarin liver tissue pre-treated on the liver followed by CCl ₄ showing preservation of normal hepatocytes. 5D Section of pre-treated 50mg/kg MEMM liver tissue followed by CCl ₄ showing scattered massive steatosis, inflammation and necrosis. (H & E, 40x magnification). CV central vein. I inflammation. H haemorrhage. N necrosis. S steatosis.	30
5	5E Section of pre-treated 250mg/kg MEMM liver tissue followed by CCl ₄ showing moderate necrosis and inflammation with mild steatosis. 5F Section of pre-treated 500mg/kg MEMM followed by CCl ₄ showing normal histology with mild inflammation. (H & E, 40x magnification). CV central vein. I inflammation. H haemorrhage	31
6	6A normal liver, 6B liver intoxicated with 3g/kg PCM: gross image shows major colour changes of liver lobes, 6C liver pre-treated with 200mg/kg Silymarin and induced with PCM: spot of colour changes was noted, 6D liver pre-treated with 50mg/kg MEMM and induced by PCM, 6E liver pre-treated with 250mg/kg MEMM and induced by PCM, 6F liver pre-treated with 500mg/kg MEMM and induced by PCM	37
7	7A Normal, 7B Section of liver tissue of 3g/kg PCM-treated group (p.o) showing necrosis, haemorrhage and inflammation. (H & E, 40x magnification). CV central vein. N necrosis. H haemorrhage	38
7	7C Section of 200 mg/kg of Silymarin liver tissue pre-treated	39

	on the liver followed by PCM showing preservation of normal hepatocytes. 7D Section of pre-treated 50mg/kg MEMM liver tissue followed by PCM showing tissue necrosis, inflammation and haemorrhage. (H & E, 40x magnification). CV central vein. I inflammation. H haemorrhage.	
7	7E Section of pre-treated 250mg/kg MEMM liver tissue followed by PCM showing mild haemorrhage and inflammation. 7F Section of pre-treated 500mg/kg MEMM followed by PCM showing normal histology with mild inflammation. (H & E, 40x magnification). CV central vein. I inflammation. H haemorrhage	40
8	8A Normal, 8B Section of liver tissue of 3g/kg PCM-treated group showing massive coagulative necrosis, haemorrhage and inflammation. (H & E, 40x magnification). CV central vein. N necrosis. haemorrhage.	47
8	8C Section of 200 mg/kg of Silymarin liver tissue pre-treated on the liver followed by PCM showing preservation of normal hepatocytes. 8D Section of pre-treated 50 mg/kg PEMM liver tissue followed by PCM showing massive coagulative necrosis, inflammation and haemorrhage. (H & E, 40x magnification). CV central vein. I inflammation. H haemorrhage. N necrosis.	48
8	8E Section of pre-treated 250mg/kg PEMM liver tissue followed by PCM showing massive coagulative necrosis, inflammation and haemorrhage. 8F Section of pre-treated 500mg/kg PEMM followed by PCM showing spotty necrosis with mild inflammation. (H & E, 40x magnification). CV central vein. I inflammation. N necrosis.	49
8	8G Section of pre-treated 50mg/kg EAMM liver tissue followed by PCM showing mild spotty necrosis and inflammation. 8H Section of pre-treated 250mg/kg EAMM liver tissue followed by PCM showing mild haemorrhage and inflammation. (H & E, 40x magnification). CV central vein. N necrosis. H haemorrhage. I inflammation.	50
8	8I Section of pre-treated 500mg/kg EAMM liver tissue followed by PCM showing mild spotty necrosis and inflammation. 8J Section of pre-treated 50mg/kg AQMM liver tissue followed by PCM showing massive coagulative necrosis, inflammation and haemorrhage. (H & E, 40x magnification). CV central vein. N necrosis. H haemorrhage. I inflammation.	51
8	8K Section of pre-treated 250mg/kg AQMM liver tissue followed by PCM showing moderate inflammation with mild haemorrhage. 8L Section of pre-treated 500mg/kg AQMM liver tissue followed by PCM showing mild necrosis and inflammation. (H & E, 40x magnification). CV central vein. N necrosis. H haemorrhage. I inflammation.	52

9	9a) DPPH scavenging activity of MEMM 9b) DPPH scavenging activity of PEMM 9c) DPPH scavenging activity of EAMM 9d) DPPH scavenging activity of AQMM	65
10a	The HPLC profile of MEMM at the wavelengths of 254 and 366 nm	74
10b	The UV spectra analysis of MEMM demonstrated the presence of four major peak, namely peak 1 ($R_T = 3.65$ min), peak 2 ($R_T = 9.11$ min), peak 3 ($R_T = 15.33$ min), and peak 4 ($R_T = 31.72$ min), which were observed at their respective λ_{max} at the respective region of 234.9, 254.3-367.2, 204.9-348.2 and 255.5-369.4 nm, suggesting, in part, the presence of flavonoid-based compounds.	75
10c	Comparison between chromatogram of the standard compound quercitrin, rutin and quercetin with the chromatogram of MEMM at 366 nm showing all the peaks are parallel to each other, indicating the present of quercitrin, rutin and quercetin present in MEMM.	76
11a	The HPLC profile of PEMM at the wavelengths of 210, 254, 280, 300, 330 and 366 nm.	78
11b	The UV spectra analysis of PEMM demonstrated the presence of four major peak, namely peak 1 ($R_T = 2.77$ min), peak 2 ($R_T = 3.77$ min), peak 3 ($R_T = 18.91$ min), peak 4 ($R_T = 20.52$ min), and peak 5 ($R_T = 23.05$ min), which were observed at their respective λ_{max} at the respective region of 208.4, 207.2, 227.2, 253.1 and 254.3 nm.	79
12a	The HPLC profile of EAMM at the wavelengths of 210, 254, 280, 300, 330 and 366 nm.	80
12b	The UV spectra analysis of EAMM demonstrated the presence of four major peak, namely peak 1 ($R_T = 2.84$ min), peak 2 ($R_T = 3.90$ min), peak 3 ($R_T = 13.71$ min), peak 4 ($R_T = 18.92$ min), peak 5 ($R_T = 20.54$ min), peak 6 ($R_T = 23.15$ min), and peak 8 ($R_T = 24.83$ min), which were observed at their respective λ_{max} at the respective region of 214.3, 216.6-270.8, 214.3-272.0, 217.8-269.7, 254.3-366.6, 206.1-348.2 and 199.00-368.2 nm.	81
13a	The HPLC profile of AQMM at the wavelengths of 210, 254, 280, 300, 330 and 366 nm.	82
13b	The UV spectra analysis of AQMM demonstrated the presence of four major peak, namely peak 1 ($R_T = 2.81$ min), peak 2 ($R_T = 17.72$ min), peak 3 ($R_T = 18.47$ min), peak 4 ($R_T = 18.97$ min), peak 5 ($R_T = 19.71$ min), peak 6 ($R_T = 19.91$ min), peak 7 ($R_T = 20.56$ min), and peak 8 ($R_T = 23.12$ min), which were observed at their respective λ_{max} at the respective region of 226.0, 216.6-269.7, 217.8-263.8, 217.8-268.5, 217.8- 267.3, 221.3-267.3, 224.8-352.9 and 263.8-343.4 nm.	83

LIST OF ABBREVIATIONS

AAPH	2,2'-azobis-2-methyl-propanimide, dihydrochloride
ALP	Alkaline phosphatase
ALT	Alanine aminotransferase
AST	Aspartate aminotransferase
ANOVA	Analysis of variance
AQMM	Aqueous extract of <i>Melastoma malabathricum</i>
AUC	Area under curve
CCl ₃	Trichloromethyl free radical
CCl ₄	Carbon tetrachloride
Cl ₃ COO	Trichloromethyl peroxy
COX	Cyclooxygenase
CYP450	Cytochrome 450
DMSO	Dimethyl sulfoxide
DPPH	2,2-diphenyl-1-picrylhydrazyl
EAMM	Ethyl acetate extract of <i>Melastoma malabathricum</i>
H & E	Haematoxylin and eosin
H ₂ O ₂	Hydrogen peroxide
HO	Hydroxyl radical
HPLC	High-performance liquid chromatography
i.p	intraperitoneal
IC ₅₀	Median inhibitory concentration
LOX	Lipoxygenase
<i>M. malabathricum</i>	<i>Melastoma malabathricum</i>
MEMM	Methanol extract of <i>Melastoma malabathricum</i>
NAC	N-acetylcystein
NAPQI	<i>N</i> -acetyl <i>p</i> -benzoquinonimine
NIDDK	National Institute of Diabetes and Digestive and Kidney Diseases
o.p	Orally
O ₂	Oxygen
ORAC	Oxygen radical absorbance capacity
PCM	Paracetamol
PEMM	Petroleum ether extract of <i>Melastoma malabathricum</i>
PPAR-α	peroxisome proliferator-activated receptor-alpha
PUFA	Polyunsaturated fatty acid
ROS	Reactive oxygen species
SD	Sprague Dawley
SEM	Standard error of mean
SOD	Superoxide Dismutase
TPC	Total phenolic content
UPM	Universiti Putra Malaysia
WHO	World Health Organization
XO	Xanthine oxidase

CHAPTER 1

INTRODUCTION

Nowadays, liver diseases are a major worldwide health problem that occur throughout the world irrespective of sex, age, race and region. According to World Health Organization (WHO), about 46% of global diseases and 59% of the mortality is because of chronic diseases. Almost 35 million people in the world die of chronic diseases and liver disease rates are progressively increase over the years. According to National Statistics in the UK, liver diseases have been ranked as the fifth major cause of death. Deaths from cirrhosis have been estimated to increase and would make it as the 12th leading cause of death in 2020, as reported by Murray and Lopez, 1997.

Although liver damage is stereotypically linked to alcohol or drugs (e.g. paracetamol (PCM) or accidental exposure to certain chemicals (e.g. carbon tetrachloride (CCl₄)), the reality is that there are over 100 known forms of liver damage caused by a variety of factors and effecting everyone from infants to older adults such as autoimmune disorders, genetic inheritance, toxins, obesity and cryptogenic. Consequently, these caused chronic damage to the liver, in which scar tissue slowly replaces normal functioning tissue lost, nutrients, hormones, drugs and poisons are not processed effectively by the liver. In addition, protein production and other substances produced by the liver are inhibited.

Even though the modern drug technologies of high-throughput screening and synthetic chemistry of the 20th Century have been expanded upon greatly, nature, particularly plant-based therapies, has remained the most valued resource in the drug development arena (Helmstadter and Staiger, 2014). Drug discovery of current active agents has been discussed from a phytopharmaceutical viewpoint. An analysis from 1981 to 2010 by Newman and Cragg (2012) showed that more than two-thirds of the drug active compounds recently introduced are likely derived from natural sources, and only about 30% are of completely synthetic origin. For decades, therapeutic practices and roles of medicinal herbs for treating disease have been gathered through an array of trials and errors, and have been documented in the history of medicine. Regardless of the abundance of the number of modern drugs in the pharmaceutical market, traditional medicine has been favoured as the primary option for alternative medicines, considering its low cost and effectiveness, and cultural, historical and even religious inclinations (Priya *et al.*, 2010).

In retrospect, there has been tremendous drug discovery from natural products since World War I, but surprisingly, less than 10% of the 250,000 species from worldwide biodiversity has been studied for medicinal purposes (Ramasamy *et al.*, 2011), leaving many species awaiting therapeutic exploration. As a land rich with flora and faunal properties, Malaysia believed to be a reservoir of a large collection of potential medicinal plants. An increasing trend in Malaysia was the recent swing in

interest from synthetic allopathic drugs to herbal medicine. In 1999, the herbal and natural product domestic market was reported to be RM4.55 billion, and the current appraisal growth rate is estimated to be worth 15-20% annually (Nordin *et al.*, 2008). Alongside economic factors, the increased interest in the herbal industry in Malaysia has apparently been caused by changes in lifestyle, increased health consciousness, and the costliness of synthetic medicine (Aziz, 2003). An in depth report by the Ministry of Natural Resources and Environment on Biodiversity in Malaysia (2006) showed that Malaysia enjoys the advantage of genetic natural products, and indigenous knowledge (Biodiversity in Malaysia 2006).

Current drug used for treating liver injury, silymarin and N-acetylcysteine offer protection to the liver from damage or help to regenerate hepatic cells. However, current drugs mainly associated with have limited efficacy, causing adverse drug reactions and yet expensive (Stickle and Schuppan, 2007). Thus, there is an urgent need to find new hepatoprotective agents from natural sources to overcome those problems.

The plant kingdom is undoubtedly valuable as a source of new medicinal agents. Plants have always been a rich source of biochemical compounds. Many of these biochemical compounds are useful drugs in themselves and others have been the basis for synthetic drugs.

To explore these sources for hepatoprotective study, *Melastoma malabathricum* was selected to be investigated on a large scale. *M. malabathricum* is a primitive plant in Malaysia that has been widely tested and documented for its promising pharmacological properties, such as antinociceptive, anti-inflammatory, antipyretic, antiviral, antibacterial, antiparasitic, antioxidant, anticoagulant, antivenom, antiulcer, and antidiarrheal and wound healing activities. Nevertheless, its hepatoprotective properties in particular have not been explored properly. As such, further research on its hepatoprotective activity is significant for nominating another plant to the list of potential medicinal hepatoprotective plant-based products.

1.1 Problem statement

In spite of the advanced development of modern medicine, there are several obstacle faced by the publics, such as the high cost of available drugs, the presence of drug side effects that prevent patients with certain health conditions from consuming a certain drug, and lack of drug availability. Therefore it is highly recommended to search for alternative medicine treating liver ailments as a substitute for currently used drugs that have fewer or no side effects and are cheaper and widely available. Encouraging research on medicinal plants indicates that phytochemicals can be exploited for treating many health problems. Extensive studies have been conducted on plant natural products, and most of these products have shown potential as new promising hepatoprotective agents; thus, this study, which aimed to discover the potential of hepatoprotective activity of *M. malabathricum* leaves, might add another candidate to the list. Scientifically, *M. malabathricum* is not traditionally known to

have hepatoprotective properties. Nevertheless, the factors that might be involved in its cytoprotective effects can be evaluated and further studied for future plant-derived drug development. Previous studies on *M. malabathricum* reported the presence of antioxidant and anti-inflammatory activity that is relevant evidence for hepatoprotective activity. Therefore, this study is expected to discover the capacity of *M. malabathricum* for hepatoprotective activity.

1.2 Hypothesis

1.2.1 Null Hypothesis

Methanol extract of *M. malabathricum* (MEMM) leaves possess hepatoprotective activity in carbon tetrachloride (CCl₄) and PCM-induced liver toxicity assays, and one or more of its partition is expected to have hepatoprotective activity in PCM-induced liver toxicity.

1.2.2 Alternative Hypothesis

Methanol extract of *M. malabathricum* (MEMM) leaves and its partition do not possess hepatoprotective activity in PCM-induced liver toxicity.

1.3 Objectives of the Study

1.3.1 General Objective

The objectives of the proposal study are as follows:

1. To determine the hepatoprotective activity of *M. malabathricum* methanol leaf extract using various hepatotoxicity induced models in rats.

1.3.2 Specific Objectives

The specific objectives of the proposed study are:

1. To determine hepatoprotective effect of methanolic extract of *M. malabathricum* leaves against carbon tetrachloride and paracetamol-induced liver toxicity models in rats; and then find out the most effective partition of MEMM; petroleum ether, ethyl acetate and aqueous extracts on liver toxicity study,
2. To examine the involvement of antioxidant and anti-inflammatory activities of the extracts as part of the hepatoprotective pathway,
3. To screen for the bioactive constituents present in MEMM and its partition using high performance liquid chromatography (HPLC)

REFERENCES

- Adams, L. A., Angulo, P., & Lindor, K. D. (2005). Nonalcoholic fatty liver disease. *Canadian Medical Association Journal*, 172(7):899–905. <http://doi.org/10.1503/cmaj.045232>
- Adewusi, E. A. and Afolayan, A. J. (2010). A review of natural products with hepatoprotective activity. *Journal of Medicinal Plants Research*, 4(13), 1318-1334.
- Ahsan, M. R., Islam, K. M. and Afolayan, I. J. (2009). Hepatoprotective activity of Methanol Extract of some medicinal plants against carbon tetrachloride-induced hepatotoxicity in rats. *European Journal of Scientific Research*, 37:302-310.
- Ajay, K. G. and Neelam, M. (2006). Hepatoprotective activity of aqueous ethanolic extract of *Chamomile capitula* in paracetamol intoxicated albino rats. *American Journal of Pharmacology and Toxicology*, 1: 17-20
- Akah, P. A. and Odo, C. I. (2010). Hepatoprotective effect of the solvent fractions of the stem of *Hoslundia opposita* Vahl (Lamiaceae) against carbon tetrachloride- and paracetamol-induced liver damage in rats. *International Journal of Green Pharmacy*, 4: 54-58. 10.4103/0973-8258.62159.
- Akkol, E. K., Tatli, I. I. and Akdemir, Z. S. (2007). Antinociceptive and anti-inflammatory effects of saponin and iridoid glycosides from *Zeitschrift fur Naturforschung*. *Journal of Biosciences*. 68:813-820.
- Ali, D. M. H., Wong, K. C. and Boey, L. P. (2010). Triterpenoids, glycolipids and flavonoids of *Melastoma malabathricum*: Isolation, spectrometric characterization and antibacterial activity. Germany: VDM Verlag, pp. 1-228
- Amacher, D. E. (1998). Serum transaminase elevations as indicators of hepatic injury following the administration of drugs. *Regulatory Toxicology and Pharmacology*, 27(2):119-130.
- Amaral, J. D., Xavier, J. M. and Steer C. J. (2010). The role of p53 in apoptosis. *Discovery Medicine*, 9(45):145–152.
- Anderson, M. T., Staal, F. J. T., Gitler, C., Herzenberg, L. A. and Herzenberg, L. A. (1994). Separation of oxidant-initiated and redox-regulated steps in the NF- κ B signal transduction pathway. *Proceedings of the National Academy of Sciences of the United States of America*, 91(24): 11527–11531.
- Annegowda, H. V., Mordi, M.N., Ramanathan, S., Hamdan, M. R. and Mansor, S. (2012). Influence of sonication treatments and extraction on the phenolics and antioxidants in star fruits. *Journal of Food, Science and Technology*. 49(4):510-514.

- Arhoghro, E. M., Ekpo, K. E., Anosike, E. O., Ibeh, G. O. (2009). Effect of aqueous extract of bitter leaf (*Vernonia amygdalina* Del) on carbon tetrachloride (CCl₄) induced liver damage in albino Wistar rat's. *Eur J Sci Res*, 26: 122-130.
- Auvichayapat P, Prapochanung M, Tunkamnerdthai O, Sripanidkulchai B, Auyichayapat N, Thinkhamrop B, Kunhasura S, Wongpratoom S, Sinawat S and Hongprapa, P. (2008). Effectiveness of green tea on weight reduction in obese Thais: A randomized, controlled trial. *Physiol Behav*. 93: 486-491. 10.1016/j.physbeh.2007.10.009
- Aziz, R.A. (2003). Turning Malaysia Into A Global Herbal Producer: A personal Perspective. Universiti Teknologi Malaysia.
- Balan, T., Sani, M.H.M, Ahmad, S. H. M., Suppaiah, V., Mohtarrudin, N., Jamaludin, F., and Zakaria, Z.A. (2012). Antioxidant and antiinflammatory activities contribute to the prophylactic effect of semi-purified fractions obtained from the crude methanol extract of *Muntingia calabura* leaves against gastric ulceration in rats. *Journal of ethnopharmacology*. Doi: 10.1016/j.jep.2014.12.17
- Bharadwaj, S. and Gakhar, S. K. (2005). Ethnomedicinal plants used by tribals of Mizorum to use cuts and wounds. *Indian Journal of Traditional Knowledge*. 4:75–80.
- Biswas, S. K. and Lopes de Faria, J. B. (2006). Hypertension induces oxidative stress but not macrophage infiltration in the kidney in the early stage of experimental diabetes mellitus. *American Journal of Nephrology*, 26(5):415–422.
- Biswas, S. K. and Lopes de Faria, J. B. (2007). Which comes first: renal inflammation or oxidative stress in spontaneously hypertensive rats? *Free Radical Research*, 41(2):216–224.
- Blois, M.S (1958). Antioxidant determinations by the use of a stable free radical. *Nature*. 181 (4617): 1199-1200
- Bohm, B. (1998). Extraction, Purification and Identification of Flavonoids. In *Introduction to Flavonoids*; Harwood Academic Publishers: Amsterdam, The Netherlands, pp. 200–204.
- Bolling, B. W., Dolnikowski, G., Blumberg, J. B., and Chen, C.-Y. O. (2010). Polyphenol content and antioxidant activity of California almonds depend on cultivar and harvest year. *Food Chemistry*, 122(3):819–825.
- Burkill, I. H. (1966). *A Dictionary of Economic Products of Malay Peninsular*, Ministry of Agriculture and Cooperatives, Kuala Lumpur, Malaysia.

- Cao, G., Alessio, H.M. and Cutler, R.G. (1993). Oxygen-radical absorbance capacity assay for antioxidants. *Free Radical Biology and Medicine*. 14:303-311.
- Cao, G. and Prior, R.L. (1999). Measurement of oxygen radical absorbance capacity in biological samples. *Methods in Enzymology*, 299:50-62.
- Carolina Henríquez, Sergio Almonacid, Italo Chiffelle, Tania Valenzuela, Manuel Araya, Lorena Cabezas, Ricardo Simpson, and Hernán Speisky (2010). Determination of Antioxidant Capacity, Total Phenolic Content and Mineral Composition of Different Fruit Tissue of Five Apple Cultivars Grown in Chile. *Chilean Journal of Agricultural Research*, 70(4):523-536.
- Collins, T. (1999). "Acute and chronic inflammation," in Robbins Pathologic Basis of Disease, R. S. Cotran, V. Kumar, and T. Collins, Eds., pp. 50–88, W.B. Saunders, Philadelphia, Pa, USA, 1999.
- Dai, J. and Mumper, R. J. Plant Phenolics: Extraction, Analysis and Their Antioxidant and Anticancer Properties. *Molecules* 2010, 15, 7313-7352.
- Dangles, O.; Fargeix, G.; Dufour, C. (1999). One-electron oxidation of quercetin and quercetin derivatives in protic and non protic media. *Journal of the Chemical Society, Perkin Transactions 2*, 1999, 1387–1395.
- Daniel, L. (2006). Introduction. *Medicinal Plants*, Jan 2006:1-4.
- Dash, D. K., Yeligar, V. C., Nayak, S. S., Ghosh, T., Rajalingam, D., Sengupta, P., Maiti, B. C. and Maity, T. K. (2007). Evaluation of hepatoprotective and antioxidant activity of *Ichnocarpus frutescens* (Linn.) R.Br. on paracetamol-induced hepatotoxicity in rats. *Tropical Journal of Pharmaceutical Research*, 6(3):755-765.
- Dass, K. K. and Kotoky, J. J. (1988). A new aliphatic constituent of *Melastoma malabathricum* Linn. *Journal of the Indian Chemical Society*. 65:385–386.
- Davalos, D., Grutzendler, J., Yang, G., Kim, J. V., Zuo, Y., Jung, S., Littman, D. R., Dustin, M. L. and Gan, W. B. (2004). ATP mediates rapid microglial response to local brain injury *in vivo*. *Natural Neurosciences*, 8:752-758
- Devaraj, S., Ismail, S., Ramanathan, S., Marimuthu, S. and Fei, Y. M. (2010) Evaluation of the hepatoprotective activity of standardize ethanolic extract of *Curcuma xanthorrhiza* Roxb. *Journal of Medicinal Plants Research*, 4(23): 2512-2517.
- Dinda, B. and Saha, M. K. (1988). A flavonol-diglycoside from *Melastoma malabathricum*. *Journal of the Indian Chemical Society*. 65:209–211.
- DiRenzo A. B., Gandolfi A. J. and Sipes I. G. (1982). Microsomal bioactivation and covalent binding of aliphatic halides to DNA. *Toxicology Letter*, 11: 243–252.

- Doehner, W. and Landmesser, U. (2011). Xanthine oxidase and uric acid in cardiovascular disease: Clinical impact and therapeutic options. *Seminars in Nephrology*, 31(5):433-440.
- Dong, M. W. (2006). *Modern HPLC for practicing scientist*. John Wiley & Sons.
- El-Beshbishy, H. A., Mohamadin, A. M., Nagy, A. A. and Abdel-Naim, A. B. (2010). "Amelioration of tamoxifen-induced liver injury in rats by grape seed extract, black seed extract and curcumin", *Indian Journal of Experimental Biology*, 48(3):280–288.
- Faravani, M. (2009). The population biology of Straits Rhododendron (*Melastoma malabathricum* L. PhD thesis, University of Malaya, Kuala Lumpur, 2009.
- Ferreira, J. F. S., Luthria, D. L., Sasaki, T. and Heyerick, A. (2010). Flavonoids from *Artemisia annua*s antioxidants and their potential synergism with Artemisinin against malaria and cancer. *Molecules*, 15: 3135-3170. 10.3390/molecules15053135.
- Flohé, L. Brigelius-Flohé, R. Saliou, C., Traber, M.G. and Packer, L. (1997). Redox regulation of NF-κB activation. *Free Radical Biology and Medicine*, 22(6):1115–1126.
- Fromenty, B. and Pessayre, D. (1995). Inhibition of mitochondrial beta-oxidation as a mechanism of hepatotoxicity. *Pharmacol Ther* 67:101–154.
- Fuji, K. (1997). Preventive effect of isofluran on destruction of cytochrome P450 during dehalogenation of carbon tetrachloride in guinea pig microsomes. *Drug Metabol Drug Interact*, 14:99–107.
- Gawlik-Dziki, U. (2011). Dietary spices as a natural effectors of lipoxygenase, xanthine oxidase, peroxidase and antioxidant agents. *Food Science and Technology*, 47(1):138-146
- Golestan Jahromi, M., Nabavizadeh, F., Vahedian, J., Nahrevanian, H, Dehpour, A and Zare-Mehrjardi, A. (2010). Protective effect of ghrelin on acetaminophen-induced liver injury in rat. *Peptides*, 31(2010): 2114-2117.
- Guyton, A. C. and Hall, J. E. (2006). *Physiological Processes*. QT 104 G992t 2006]. QP34.5.G9 2006. 612-622.
- Gupta, A. K., Chitme, H., Dass S.K. and Misra, N. (2006) .Hepatoprotective activity of *Rauwolfia serpentina* rhizome in paracetamol intoxicated rats. *Journal of Pharmacology and Toxicology*, 1(1):82-88.
- Halliwell, B. and Gutteridge, J. M. C. (1999). *Free Radicals in Biology and Medicine*, Oxford University Press, London, UK, 3rd edition.

- Han, K. L., Jung, M. H., Sohn, J. H. and Hwang, J. K. (2006). Ginsenoside 20S-protopanaxatriol (PPT) activates peroxisome proliferator-activated receptor gamma (PPARgamma) in 3T3-L1 adipocytes. *Biological & Pharmaceutical Bulletin*, 29(1):110-113.
- Hassan, H. M. (2012). Hepatoprotective effect of red grape seed extracts against ethanol-induced cytotoxicity. *Global Journal of Biotechnology and Biochemistry*. 7:30-37.
- Haupt, S., Berger, M., Goldberg, Z., Haupt, Y. (2003). Apoptosis-the p53 network. *Journal of Cell Science*, 116: 4077-4085
- Helmstadter, A and Staiger, C. (2014). Traditional use of medicinal agents: a valid source of evidence. *Drug Discovery Today*.19:1
- Herman, B., Nieminen, A. L., Gores, G. J. and Lemasters, J. J. (1988). Irreversible injury in anoxic hepatocytes precipitated by an abrupt increase in plasma membrane permeability. *FASEB J* 2:146–151.
- Hernandez-Munoz R, Diaz-Munoz M, Suarez J, Chagoya de Sanches V. (1990). Adenosine partially prevents cirrhosis induced by carbon tetrachloride in rats. *Hepatology*, 12:242–8.
- <http://www.stanfordchildrens.org/content-public/topic/images/33/125533.gif>.
Accessed on 8 August 2016.
- Huang, D., Ou, B., Hampch-Woodill, M., Flanagan, J.A. and Prior, R.L. (2002). High throughput assay of oxygen radical absorbance capacity (ORAC) using a multichannel liquid handling system coupled with a microplate fluorescence reader in 96-well format. *Journal of Agricultural and Food Chemistry*. 5:4437-4444.
- Hussain F, Abdulla MA, Noor SM, Ismail S, Ali HM. (2008) Gastroprotective effects of *Melastoma malabathricum* aqueous leaf extract against ethanol-induced gastric ulcer in rats. *American Journal of Biochemistry and Biotechnology*. 4(4):438–441.
- Jain, S. K. and De Filippis, R. A. (1991). *Medicinal Plants of India*, Reference Publication Inc., Algonac, Mich, USA.
- Jaganath, I. B. and Ng L. T. *Herbs: The Green Pharmacy of Malaysia*. Kuala Lumpur, Malaysia: Vinpress Sdn. Bhd.; 2000
- Jantan I, Rafi IAA, Jalil J. (2005). Platelet-activating factor (PAF) receptor-binding antagonist activity of Malaysian medicinal plants. *Phytomedicine*. 12(1-2):88–92.

- Jodynis-Liebert J, Matlawska I, Bylka W, Murias M. (2005). Protective effect of *Aquilegia vulgaris* on APAP-induced oxidative stress in rats. *Journal of Ethnopharmacology*, 97: 351-358. 10.1016/j.jep.2004.11.027
- Johnny, L., Yusuf, U. K. and Nulit, R. (2010). The effect of herbal plant extracts on the growth and sporulation of *Colletotrichum gloeosporioides*. *Journal of Applied Biosciences*. 34:2218–2224.
- Jollow, D. J., Mitchell, J. R., Potter, W. Z., Davis, D. C., Gillette, J. R. and Brodie, B. B. (1973). Acetaminophen-induced hepatic necrosis. II. Role of covalent binding in vivo. *Journal of Pharmacology and Experimental Therapeutics*, 187: 195-202.
- Kamisan, F.H., Yahya, F., Ismail, N.A., Din, S.S., Mamat, S.S., Zabidi, Z., Zainuddin, Z., W.N., Mohtarrudin, N., Husain, H., Ahmad, Z. and Zakaria, Z.A. (2013). Hepatoprotective activity of methanol extract of *Melastoma malabathricum* leaf in rats. *Journal of Acupuncture and Meridian Studies*. 6:52-55.
- Karmen, A. (1955). A note on the spectrometric assay of glutamic-oxalacetic transaminase in human blood serum. *The Journal of Clinical Investigation*. 34 (1): 131–3. *PMC 438594. PMID 13221664*.
- Karmen, A., Wróblewski, F., & LaDue, J. S. (1955). Transaminase Activity In Human Blood. *Journal of Clinical Investigation*, 34(1), 126–133.
- Kaplowitz S. A., Campo S. , Chiu W. T. (2002) Cancer patients' desires for communication of prognosis information. *Health Communication*, 14(2):221-41.
- Koay, S. S. (2008). Establishment of cell suspension culture of *Melastoma malabathricum* L. for the production of anthocyanin,” PhD. Thesis, Universiti Sains Malaysia, Pulau Pinang, Malaysia.
- Kumar, G., Banu, G. S., Pappa, P. V., Sundararajan, M., and Pandian, M. R. (2004). Hepatoprotective activity of *Trianthema portulacastrum* L. against paracetamol and thioacetamide intoxication in albino rats. *Journal of Ethnopharmacology* 92(1):37–40.
- Kurt, C. K. and Kathleen, A. (2011). Chapter 12: Biochemical and Metabolic Principles, Goldfrank’s Toxicology Emergencies. *Annals of Internal Medicine*.
- Latiff, A. and Zakri, A. H. (2000). Protection of Traditional Knowledge, Innovations and Practices: The Malaysian Experience. in the *UNCTAD Expert Meeting on Systems and National Experiences for Protecting Traditional Knowledge, Innovations and Practices, Geneva, Switzerland, October-November 2000*

- Lee, H. O. and Ferguson, T. A. (2003). Biology of FasL. Cytokine Growth Factor Rev 14:325–335.
- Lee, H. L. and Halliwell, B. (2001). Antioxidant and prooxidant abilities of foods and beverages. *Methods in Enzymology*, 335:181-190.
- Lemasters, J. J. (1999). V. Necrapoptosis and the mitochondrial permeability transition: shared pathways to necrosis and apoptosis. *Am J Physiology*, 276:G1–G6.
- Liu, F., Ooi, V .E. C. and Chang, S. T. (1997). Free radical scavenging activities of mushroom polysaccharide extracts. *Life Sciences*, 60(10):763-771.
- Lohézic-Le Dévéhat, F., Bakhtiar, A., Bézivin, C., Amoros, M. and Boustie, J. (2002). Antiviral and cytotoxic activities of some Indonesian plants. *Fitoterapia*, 73(5):400–405.
- Lopez-Alarcon, C. and Denicola, A. (2013). Evaluating the antioxidant capacity of natural products: a review on chemical and cellular-based assays. *Analytica Chimica Acta*. 763:1-10.
- Luster, M. I., Simeonova, P. P., Gallucci, R. M., Matheson, J. M. and Yucesoy, B. (2000). Immunotoxicology: role of inflammation in chemical-induced hepatotoxicity. *International Journal of Immunopharmacology*, 22(12):1143-1147.
- Maji, S., Dandapat, P., Ojha, D., Maity, C., Halder, S. K., Das Mohapatra P. K., Pathak, T. K., Pati, B. R., Samanta, A. and Mondal, K. C. (2010). In vitro antimicrobial potentialities of different solvent extracts of ethnomedicinal plants against clinically isolated human pathogens. *Journal of Phytotherapy*. 2:57–64.
- Malik, A., Anis, I., Khan, S. B., Ahmed, E., Ahmed, Z., Nawas, S. A. and Choundry, M. I. (2004). Enzymes inhibiting lignants from *Vitex negundo*. *Chemical and Pharmaceutical bulletin*. 52: 1269-1272.
- Manautou, J. E., Emeigh, S.G., Khairallah, E. A and Cohen, S. D. (1996). Protection against acetaminophen hepatotoxicity by a single dose of clofibrate: Effects on selective protein arylation and glutathione depletion. *Fundamental and Applied Toxicology*, 29(2):229-237.
- Manicam, C., Abdullah, J. O., Mohd Tohit, E. R., Seman, Z., Sieo, C. C. and Hamid, M. (2010). In vitro anticoagulant activities of *Melastoma malabathricum* Linn. aqueous leaf extract: a preliminary novel finding. *Journal of Medicinal Plants Research*, 4:1464–1472.
- Manno, M., Ferrara, R., Cazzaro, S., Rigotti, P. and Ancona, E. (1992). Suicidal inactivation of human cytochrome P-450 by carbon tetrachloride and halothane in vitro. *Pharmacology and Toxicology*, 70:13–18.

- Manzoor-I-Khuda, M., Chowdhury, S. A., Reza, T., and Chowdhury, A. K. (1981). Chemical Investigation on *Melastoma Malabathricum*. Part 1: Isolation of Melastomic Acid and Beta-sitosterol from the Roots. *Journal of the Bangladesh Academy of Sciences*. 5:55–59
- Marzilawati, A. R., Ngau, Y. Y. and Mahadeva, S. (2012). Low rates of hepatotoxicity among Asian patients with paracetamol overdose: a review of 1024 cases. *BMC Pharmacology and Toxicology*. 13:8.
- Masih, N. G. and Singh, B. S. (2012). Phytochemical Screening of Some Plants Used in Herbal Based Cosmetic Preparations. *Chemistry of Phytopotentials: Health, Energy and Environment Perspectives* Springer Berlin Heidelberg 111-112.
- Mazura MP, Susanti D, Rasadah MA. (2007) Anti-inflammatory action of components from *Melastoma malabathricum*. *Pharmaceutical Biology*. 45(5):372–375.
- McCay P. B., Lay E. K., Poyer J. L., DuBose C. M. and Jansen E. G. (1984). Oxygen and carbon-centered free radical formation during carbon tetrachloride metabolism. *Journal of Biology and Chemistry*, 259: 2135–2143.
- Mehendale, H. M. (2000). PPAR- α : A key to the mechanism of hepatoprotection by clofibrate. *Toxicology Sciences*, vol. 57, no. 2, pp. 187-190.
- Mohandoss, S. and Ravindran, P. (1993). Flavonoids from *Melastoma malabathricum*. *Fitoterapia*, 64(3):277–278.
- Mohd Joffry, S., Yob, N. J., Rofiee, M. S., Affandi, M. M., Othman, F., Abdah, M. A. Mohd. Desa, M. N., Suhaili, Z., and Zakaria, Z. A. (2012). *Melastoma malabathricum* (L.) Smith: A review of its ethnomedicinal, chemical and pharmacological uses. *Evidence-Based Complementary and Alternative Medicine*, vol. 2012, Article ID 258434, 48 pages, 2012. doi:10.1155/2012/258434.
- Molyneux, P. (2004). The use of the stable free radical diphenylpicryl-hydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin Journal of Science and Technology*, 26(2): 211-219.
- Morrey, K. L. and Dalley, A. F. (2001). Clinically Oriented Anatomy. 4th ed. Baltimore: Lippincott Williams & Wilkins, Figure 5-25.
- Murray, C. J. L. and Lopez (1997). Alternative projections of mortality and disability by cause 1990–2020: Global Burden of Disease Study. *The Lancet*, Volume 349, Issue 9064, 1498 – 1504.

- Muthumani, M. and Prabu, S. M. (2014). Silibinin potentially attenuates arsenic-induced oxidative stress mediated cardiotoxicity and dyslipidemia in rats. *Cardiovascular Toxicology*, 14:83–97
- Nishimura, K., Tanaka, N., Kawano, T., Matsuura, K., Okamura, H. (1998). Changes in macrophage colony-stimulating factor concentration in serum and follicular fluid in in-vitro fertilization and embryo transfer cycles. *Fertil Steril.*, 9(1):53-7.
- Nazlina, I., Norha, S., Noor Zarina, A. W. and Ahmad, I. B. (2008). Cytotoxicity and antiviral activity of *Melastoma malabathricum* extracts. *Malaysian Journal of Applied Biology*, vol. 37, no. 2, pp. 53–55.
- Negi B. S., Dave B. P., Agarwal Y. K. (2012). Evaluation of Antimicrobial Activity of Bauhinia purpurea Leaves Under In Vitro Conditions. *Indian Journal of Microbiology*, 52(3):360-5.
- Newman, D. J. and Cragg, G.M. (2012). Natural products as sources of new drugs over 30 years from 1981 to 2010. *Journal of Natural Products*. 75:311-335.
- Nithianantham, K.; Shyamala, M.; Chen, Y.; Latha, L.Y.; Jothy, S.L.; Sasidharan, S. (2011). Hepatoprotective Potential of *Clitoria ternatea* Leaf Extract against Paracetamol Induced Damage in Mice. *Molecules*, 16:10134-10145.
- Nithianantham, K., Ping, K.Y., Latha, L.Y., Jothy, S.L., Darah, I., Chen, Y., Chew, A.-L., Sasidharan, S., (2013). Evaluation of hepatoprotective effect of methanolic extract of *Clitoria ternatea* (Linn.) flower against acetaminophen-induced liver damage. *Asian Pac. J. Trop. Dis.* 3, 314-319.
- Nordin, N., Othman, S.N., and Che Mat (2008). Technology implementation barriers in the Malaysian herbal industry: A case study. *Malaysian Management Journal*. 12:79-88.
- Noro, T., Miyase, T. and Kuroyanagi, M. (1983). Monoamine oxidase inhibitor from the rhizomes of *Kaempferia galangal* L. *Chemical and Pharmaceutical Bulletin*. 31: 2708-2711.
- Nuresti, S., Baek, S. H. and Asari, A. (2003). Chemical components of *Melastoma malabathricum*. *ACGC Chemical Research Communications*, vol. 16, pp. 28-33, 2003.
- Oboh, G. and Ademosun, A. O., (2012). Characterization of the antioxidant properties of phenolic extracts from some citrus peels. *Journal of Food, Science and Technology*, 49(6):729-36.
- Otsuka, H. (2005). Purification by solvent extraction using partition coefficient. *Natural Products Isolation*. Humana Press, 269-273

- Perez, M. B., Calderon, N. L. and Croci, C. A. (2007). Radiation-induced enhancement of antioxidant activity in extracts of rosemary *Rosmarinus officinalis* L. *Food Chemistry*. 104: 585-592.
- Pichiah PB, Moon HJ, Park JE, Moon YJ, Cha YS. 2012. Ethanolic extract of seabuckthorn (*Hippophae rhamnoides* L.) prevents high-fat diet-induced obesity in mice through down-regulation of adipogenic and lipogenic gene expression. *Nutrition Research*, 32: 856–864.
- Pierce, R. A., Glaug, M. R., Greco, R. S. (1987). Increased procollagen mRNA levels in carbon tetrachloride-induced liver fibrosis in rats. *Journal of Biology and Chemistry*, 262:1652–8
- Pithayanukul, P., Nithitanakool, S. and Bavovada, R. (2009). Hepatoprotective Potential of Extracts from Seeds of *Areca catechu* and Nutgalls of *Quercus infectoria*. *Molecules*, 14(12):4987-5000.
- Pramyothin, P., Ngamtin, C., Pongshompoo, S., Chaichantipyuth, C. (2007). Hepatoprotective activity of *Phyllanthus amarus* Schum. et. Thonn. extract in ethanol treated rats: In vitro and in vivo studies. *Journal of Ethnopharmacology*, 114: 169-173. 10.1016/j.jep.2007.07.037
- Priya, V., Niveda, S., Pratiksha, G. and Gayathri, R. (2010). A review of hepatoprotective natural products. *Recent Research in Science and Technology*. 26: 49-52.
- Pongprom N, Khanapan C, Tosert P. Chemical constituents of *Melastoma malabathricum* Linn. In: the 29th Congress on Science and Technology of Thailand; October 2003; Khon Kaen, Thailand. p. 130
- Porchezian, E. and Ansari, S. H. (2005). Hepatoprotective activity of *Abutilon indicum* on experimental liver damage in rats. *Phytomedicine*, vol. 12, no. 1-2, pp. 62–64.
- Potter WZ, Davis DC, Mitchell JR, Jollow DJ, Gillette JR, Brodie, B. B. (1973). A cetaminophen-induced hepatic necrosis. III. Cytochrome P-450-mediated covalent binding in vitro. *Journal of Pharmacology and Experimental Therapeutics*, 187: 203-210.
- Qader, S. W., Abdulla, M. A., Chua, L. S., Najim, N., Zain, M. M. and Hamdan, S. (2011). Antioxidant, Total Phenolic Content and Cytotoxicity Evaluation of Selected Malaysian Plants. *Molecules*. 16(4):3433-3443.
- Ramasamy, S., Wahab, N. A., Abidin, N. Z. and Manickam, S. (2011). Cytotoxicity evaluation of five selected Malaysian Phyllanthaceae species on various human cancer lines. *Journal of Medicinal Plant Research*. 5:2267-2273.

- Recknagel, R. O., Glende, E. A. Jr, Dolak, J. A. and Waller, R. L. (1989). Mechanism of carbon tetrachloride toxicity. *Pharmacology Therapeutic*, 43:139–154.
- Rolo A. P., Oliveira P. J., Moreno A. J., Palmeira C.M. Protection against post-ischemic mitochondrial injury in rat liver by silymarin or TUDC. *Hepatol. Res.* 2003(26):217–224.
- Rossi, A., Kapahi, P., Natoli, G., Takahashi, T., Chen, Y., Karin, M. (2000). Anti-inflammatory cyclopentenone prostaglandins are direct inhibitors of IkappaB kinase. *Nature*, 403:103–8.
- Sajem, A. L. and Gosai, K. (2006). Traditional use of medicinal plants by the Jaintia tribes in North Cachar Hills district of Assam, northeast India. *Journal of Ethnobiology and Ethnomedicine*, 2:33.
- Salama, S. M, Abdulla, M. A., Alrashid, A. S., Ismail, S., Alkiyumi, S. S and Golbabapour, S. (2013). Hepatoprotective effect of ethanolic extract of *Curcuma longa* on thioacetamide induced liver cirrhosis in rats. *BMC Complementary and Alternative Medicine*. 2013:13-56. DOI: 10.1186/1472-6882-13-56
- Sandhar, H. K., Kumar, B., Prasher, S., Tiwari, P., Salhan, M. and Sharma, P. (2011). A Review of Phytochemistry and Pharmacology of Flavonoids. *Internationale Pharmaceutica Scientia*, 1(1):24-41
- Sekine S. and Ichijo H. (2014). Mitochondrial proteolysis: Its emerging roles in stress responses. *Biochim. Biophys. Acta*. 2015(1850):274–280. doi: 10.1016/j.bbagen.2014.10.012.
- Sharma, H. K., Chhangte, L., and Dolui, A. K. (2001). Traditional medicinal plants in Mizoram, India. *Fitoterapia*. 72(2):146–161.
- Shimoda H, Seki E, Aitani M (2006). Inhibitory effect of green coffee bean extract on fat accumulation and body weight gain in mice. *BMC Complementary and Alternative Medicine*, 6: 9-10.1186/1472-6882-6-9. doi: 10.1186/1472-6882-6-9.
- Shnoy, A. K., Somayaji, S. N. and Bairy, K. L. (2002). Hepatoprotective activity of ethanolic extract of *Ginkgo biloba* against carbon tetrachloride induced hepatic injury in rats. *Indian Journal of Pharmacology*, vol. 46, pp. 167-74.
- Sies, H. (1985). Introductory remarks, in *Oxidative Stress*, H. Sies, Ed., pp. 1–7, Academic Press, Orlando, Fla, USA, 1985. Singleton, V. L and Rossi, J. A Jr. (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, vol. 16, no. 3, pp. 144-158.

- Singleton V. L. and Rossi, J. A Jr. (1965) Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, vol. 16, no. 3, pp. 144-158.
- Somchit, M. N., Zuraini, A., Bustamam, A. A., Sulaiman, M.R and Noratunlina, R. (2005). Protective activity of turmeric (*Curcuma longa*) in paracetamol-induced hepatotoxicity in rats. *International Journal of Pharmacology*, vol. 1(3):252-256.
- Starr, S, and Hand, H. (2002). Nursing care of chronic and acute liver failure. *Nursing Standard*, 16: 47-54.
- Stickel, F. and Schuppan, D. (2007). Herbal medicine in the treatment of liver diseases. *Dig Liver Dis*. 39(4):293-304.
- Sulaiman, M. R., Somchit, M. N., Israf, D. A., Ahmad, Z. and Moin, S. (2004). Antinociceptive effect of *Melastoma malabathricum* ethanolic extract in mice. *Fitoterapia*. 75(7-8):667-672.
- Sunilson, J. A. J, Anandarajagopal, K., Kumari, A.V. A. G. and Mohan, S. (2009). Antidiarrhoeal activity of leaves of *Melastoma malabathricum* linn. *Indian Journal of Pharmaceutical Sciences*. 71(6):691-695.
- Surai, P.F. (2015). Silymarin as a natural antioxidant; An overview of the current evidence and perspective. *Antioxidants (Basel)*. 4(1): 204-247.
- Susanti, D., Sirat, H. M., Ahmad, F. and Mat Ali, R. (2008). Bioactive constituents from the leaves of *Melastoma malabathricum* L. *Jurnal Ilmiah Farmasi*, 5(1):1-8.
- Tang, E. H. C., Libby, P., Vanhoutte, P. M. and Xu, A. (2012). Anti-inflammation therapy by activation of prostaglandin EP4 receptor in cardiovascular and other inflammatory diseases. *Journal of Cardiovascular Pharmacology*, 59:116-23.
- Tapas, A. R., Sakarkar, D. M. and Kakde, R. B. (2008). Flavonoids as Nutraceuticals: A Review. *Tropical Journal of Pharmaceutical Research*, 7(3):1089-1099.
- Thatoi, H. N., Panda, S. K., Rath, S. K., and Dutta, S. K. (2008) Antimicrobial activity and ethnomedicinal uses of some medicinal plants from Similipal Biosphere Reserve, Orissa. *Asian Journal of Plant Sciences*. 7(3):260-267.
- Tomasi, A., Albano, E., Bani, S., Botti, B., Corongiu, F., Dessi, M. A., Lannone, A., Vannini, V. and Dianzani, M. U. (1987). Free radical metabolism of carbon tetrachloride in rat liver mitochondria. *Biochemistry Journal*, 246:313-317.

- Tripathi D. and Hayes, P. C. (2001). A drug therapy for the prevention of variceal haemorrhage. *Alimentary Pharmacology & Therapeutics*, 15(3):291-310.
- Tsimogiannis, D. I. and Oreopoulou, V. (2006). The contribution of flavonoid C-ring on the DPPH free radical scavenging efficiency. A kinetic approach for the 3',4'-hydroxy substituted members. *Innovative Food Science and Emerging Technology*, 7:140–146.
- Tsimogiannis, D. and Oreopoulou, V. (2007). Defining the role of flavonoid structure on cottonseed oil stabilization: Study of A- and C-ring substitution. *The Journal of the American Oil Chemists' Society*, 84, 129–136.
- Tsimogiannis, D., Samiotaki, M., Panayotou, G. and Oreopoulou, V. (2007). Characterization of flavonoid subgroups and hydroxy substitution by HPLC-MS/MS. *Molecules*, 12: 593-606. 10.3390/12030593.
- Uawonggul N, Chaveerach A, Thammasirirak S, Arkaravichien T, Chuachan C, Daduang S. (2006) Screening of plants acting against *Heterometrus laoticus* scorpion venom activity on fibroblast cell lysis. *Journal of Ethnopharmacology*. 103(2):201–207.
- Uma, N. Jr., Fakurazi, S. and Hairuszah, I. (2010). *Moringa oleifera* Enhances Liver Antioxidant Status via Elevation of Antioxidant Enzymes Activity and Counteracts Paracetamol-induced Hepatotoxicity. *Malaysian Journal of Nutrition*, 16(2):293 – 307.
- Umali-Stuart, G. and Stiuart-Santiago, A. (2010). Philippine Medicinal Plants: Family Melastomaceae. February 2010.
- Van Gaal LF, Wauters MA and De Leeuw IH. (1997). The beneficial effects of modest weight loss on cardiovascular risk factors. *Int J Obes Relat Metab Disord*, 21: S5-S9
- Van Valkenberg, J. L. C. H and N. Bunyapraphatsara. 2001. *Melastoma malabathricum* L. In: van Valkenburg, J.L.C.H. and Bunyapraphatsara, N. (Eds). Plant Resources of South-East Asia No. 12(2): Medicinal and poisonous plants 2. Leiden, The Netherlands: *Backhuys Publisher*, pp. 365-366.
- Vermeulen, N.P.E., Bessems, J.G.M., Van De Straat, R., 1992. Molecular aspect of paracetamol-induced hepatotoxicity and its mechanism based prevention. *Drug Metab. Rev.* 24, 367.
- Vimala B, Nedunchezian M, Ramanathan S and Jayaprakas CA (2011). Evaluation of Orange-fleshed sweet potato for tuber yield and carotene content at different locations of Orissa. Abstracts in National Seminar on Climate Change and food Security: Challenges and Opportunities for Tuber Crops. CIG/P8. pp88

- Vishnu Priya, V., Jainu, M., Mohan, S. K. Saraswathi, P. and Chandra Sada Gopan, V. S. (2010). Antimicrobial Activity of Pericarp Extract of *Garcinia Mangostana* Linn. *International Journal of Pharma Sciences and Research*, 1(8):278-281.
- Weber L. W. D., Boll M. and Stampfl A. (2003). Hepatotoxicity and mechanism of action of haloalkanes: carbon tetrachloride as a toxicological model. *Critical Review in Toxicology*, 33:105–136.
- Wiar, C., Mogana, S., Khalifah, S., Mahan, S., Ismail, S., Buckle, M., Narayana, A. K. and Sulaiman, M. (2004). Antimicrobial screening of plants used for traditional medicine in the state of Perak, Peninsular Malaysia. *Fitoterapia*. 75(1):68–73
- Wong W. *Melastoma malabathricum: Too Beautiful to Be Called a Weed*. Singapore: Green Culture; 2008.
- World Health Organization. (2002). Traditional medicine strategy launched. *Bulletin of the world health organization*, 80:7
- Yahya F, Ismail NA, Kamisan FH, Mamat SS, Shamsahal Din S, Wan Zainulddin WN, Zabidi Z, Mohtarrudin N, Suhaili Z, Zakaria ZA: Effect of aqueous extract of *Melastoma malabathricum* leaves against paracetamol- and CCl₄-induced liver toxicity in rats. *Afr J Pharm Pharmacol* 2012, 6:2682–2685.
- Yang, J., Li, Y., Wang, F. and Wu, C. F. (2010) Hepatoprotective effects of apple polyphenols on CCl₄-induced acute liver damage in mice. *J Agric Food Chem.*, 58: 6525-6531. 10.1021/jf903070a.
- Yanpallewar, S. U., Sen, S., Tapas, S., Kumar, M., Raju, S. S. and Acharya, S. B. (2002). Effect of *Azadirachta indica* on paracetamol induced hepatic damage in albino rats. *Phytomedicine*, vol. 10, no. 5, pp. 391–396.
- Yob, N. J., Joffry, S. M., Affandi, M. M., Teh, L. K., Salleh M. Z. and Zakaria, Z. A. (2011). *Zingiber zerumbet* (L.) Smith: A Review of Its Ethnomedicinal, Chemical and Pharmacological Uses. *Evidence-based Complementary and Alternative Medicine*. 2011:543216
- Yoshida, T., Nakata, F., Hosotani, K., Nitta, A. and Okudat, T. (1992). Dimeric hydrolysable tannins from *Melastoma malabathricum*. *Phytochemistry*, vol. 31, no. 8, pp. 2829-2833.
- Zakaria, M. and Mohd, M. A. (1994). Traditional Malay Medicinal Plants, Fajar Bakti Sdn. Bhd., Kuala Lumpur, Malaysia.

- Zakaria, Z. A., Rofiee, M. S., Mohamed, A. M., Teh, L. K. and Salleh, M. Z. (2011). *In vitro* antiproliferative and antioxidant activities, and total phenolic contents of the extracts of *Melastoma malabathricum* leaves. *Journal of Acupuncture Meridian Studies*, 4(4):248-256.
- Zakaria, Z. A., Raden Mohd. Nor, R. N. S., Hanan Kumar, G., Abdul Ghani, Z. D. F., Sulaiman, M. R., Rathna Devi, G., Mat Jais, A. M., Somchit M. N. and Fatimah, C.A. (2006a). Antinociceptive, anti-inflammatory and anti-pyretic properties of *Melastoma malabathricum* leaves aqueous extract in experimental animals. *Canadian Journal of Physiology and Pharmacology*, 84(12):1291-1299.
- Zakaria ZA, Rofiee MS, Mohamed AM, Teh LK, Salleh M. Z. (2011). In vitro antiproliferative and antioxidant activities, and total phenolic contents of the extracts of *Melastoma malabathricum* leaves. *Journal of Acupuncture and Meridian Studies*, 4: 248-256. 10.1016/j.jams.2011.09.016.
- Zakaria, Z. A., Abdul Hisam, E. E., Norhafizah, M., Rofiee, M. S., Othman, F., Hasiah, A. H., and Vasudevan, M. (2012). Methanol Extract of *Bauhinia purpurea* Leaf Possesses Antiulcer Activity. *Medical Principles and Practice*. 21(5):476 – 482.