



UNIVERSITI PUTRA MALAYSIA

PHYTOCHEMICALS FROM THE STEM BARK OF *Mesua hexapetala* (HOOK. F.) P.S. ASHTON., *Mesua beccariana* (BAILL.) KOSTERM. AND *Garcinia mangostana* LINN. AND THEIR BIOLOGICAL ACTIVITIES

THIRUVENTHAN A/L KARUNAKARAN

FS 2017 78



**PHYTOCHEMICALS FROM THE STEM BARK OF *Mesua hexapetala*
(HOOK. F.) P.S. ASHTON., *Mesua beccariana* (BAILL.) KOSTERM. AND
Garcinia mangostana LINN. AND THEIR BIOLOGICAL ACTIVITIES**

By

THIRUVENTHAN A/L KARUNAKARAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

September 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This thesis is dedicated to my beloved family, Professor, supervisory committee and friends



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

PHYTOCHEMICALS FROM THE STEM BARK OF *Mesua hexapetala* (HOOK. F.) P.S. ASHTON., *Mesua beccariana* (BAILL.) KOSTERM. AND *Garcinia mangostana* LINN. AND THEIR BIOLOGICAL ACTIVITIES

By

THIRUVENTHAN A/L KARUNAKARAN

September 2017

Chairman : Professor Gwendoline Ee Cheng Lian, PhD
Faculty : Science

Phytochemical constituents isolated from the Calophyllaceae and Clusiaceae family of plants have gained impressive attention recently due to their amazing potencies in biological activities such as anti-inflammatory, cytotoxicity and anti-bacterial. Coumarins and xanthenes discovered from this family have exhibited potent biological activities and some of these compounds have been proposed as lead compounds for drug discovery processes. Nitric oxide plays a vital role in the key molecular signaling constituent involved in inflammatory processes and mass production of NO is one of the main factors that contribute towards chronic degenerative diseases such as cancer and arthritis. It is necessary to find an alternative sources from natural products such as plants as its offer greater hope in the identification of bioactive compounds, less adverse effects and their development into drugs for the treatment of inflammatory diseases. Phytochemical studies such as extraction, isolation and structure elucidation of the isolated compounds with the aid FTIR, GCMS, HRESIMS and NMR were performed on the stem bark of two selected species from the family of Calophyllaceae which are *Mesua hexapetala* and *Mesua beccariana* as well on the stem bark of species from the Clusiaceae family which is *Garcinia mangostana* afforded twenty secondary metabolites, three of which were identified as new compounds.

Phytochemical investigation on the stem bark of *Mesua hexapetala* yielded one new coumarin derivative, hexapetarin (**143**) together with 1,3,7-trihydroxy-2,4-di (3-methyl-2-butenyl)xanthone (**144**), trapezifolixanthone (**96**), cudraxanthone G (**19**), stigmasterol (**7**), β -sitosterol (**65**) and γ -sitosterol (**145**). These findings are considered novel as no previous reports nor research have been conducted on *Mesua hexapetala*. Meanwhile, from the stem bark of *Mesua beccariana*, a new xanthone derivative,

beccarixanthone T (**146**) and a new coumarin derivative, beccamarin T (**147**) were successfully isolated together with mesuarianone (**1**), mesuasinone (**2**), 1,5-dihydroxyxanthone (**57**), euxanthone (**20**) and four common triterpenoids, friedelin (**8**), **7**, **65** and **145**. The chemical constituents isolated from *Garcinia mangostana* were α -mangostin (**14**), β -mangostin (**88**), mangostenol (**137**), garcinone D (**132**) fuscaxanthone C (**148**), dulcisxanthone F (**94**) and **7**.

Moreover, the semi-synthesis reaction of β -mangostin which were acetylation and *O*-alkylation afforded six derivatives in which four are new derivatives, 6-monoacetate β -mangostin (**149**), 6-*O*-methyl β -mangostin (**148**), 6-*O*-benzyl β -mangostin (**150**), 6-*O*-*n*-hexyl β -mangostin (**151**), 6-*O*-*n*-butyl β -mangostin (**152**) and 6-*O*-*sec*-butyl β -mangostin (**153**). All extracted plant crude extracts, seven selected natural product compounds and four semi-synthetic derivatives of β -mangostin were evaluated for cell based cytotoxicity and nitric oxide inhibition assay. Nitric oxide inhibition activities indicated that four compounds **143**, **144**, **14** and **88** exhibited very significant activity towards inhibition of LPS/IFN- γ stimulated RAW 264.7 macrophages with IC₅₀ value of 30.79 ± 2.68 , 12.41 ± 0.89 , 29.81 ± 0.77 and 11.72 ± 1.16 μ M respectively. The ethyl acetate extract of *G.mangostana* and the *n*-hexane extract of *M.hexapetala* showed very potent activities. Structure-activity relationship studies of pure compounds and semisynthetic derivatives of β -mangostin towards anti-inflammatory activity demonstrated the importance of hydroxyl groups in their structure which influence bioactivity.

Anti-*Bacillus* tests were conducted on seven types of pure compounds - **145**, **146**, **147**, **144**, **14** and **88** as well as with all the plant extracts against four types *Bacillus* bacteria - *Bacillus subtilis*, *B.cereus*, *B.megaterium* and *B.pumilus*. Hexapetarin (**145**) showed potential anti-bacillus activities while the *n*-hexane and chloroform extract of *M.hexapetala* (MHH and MHC) gave good inhibitive activity towards these four bacteria. Thus, it can be summarized that the newly discovered coumarin derivative from *M.hexapetala*, hexapetarin (**145**), *n*-hexane crude extract of *M.hexapetala* and ethyl acetate crude extract of *G.mangostana* showed very significant nitric oxide inhibition and anti-*Bacillus* activity while compound **144**, **14** and **88** portrayed very potent nitric oxide inhibition activities. These compounds can be further developed into anti-inflammatory agents.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

SEBATIAN FITOKIMIA DARIPADA BATANG KULIT KAYU *Mesua hexapetala* (HOOK. F.) P.S. ASHTON., *Mesua beccariana* (BAILL.) KOSTERM. DAN *Garcinia mangostana* LINN., DAN AKTIVITI BIOLOGIKALNYA

Oleh

THIRUVENTHAN A/L KARUNAKARAN

September 2017

Pengerusi : Profesor Gwendoline Ee Cheng Lian, PhD
Fakulti : Sains

Pengasingan sebatian fitokimia daripada genus pokok keluarga Calophyllaceae dan Clusiaceae telah mendapat perhatian yang mengagumkan pada masa kini disebabkan potensinya terhadap aktiviti-aktiviti farmakologi seperti anti-radang, sitotoksik, anti-mikrobial, anti-diabetik dan anti-oksida. Kumarin dan xanton yang dijumpai daripada keluarga pokok ini didapati menunjukkan aktiviti biologikal yang sangat bagus, malah ada sesetengah sebatian semula jadi telah pun dicadangkan untuk dijadikan sampel ubat bagi proses penemuan ubat. Nitrik oksida memainkan peranan penting dalam menjadi kunci isyarat utama sebatian molekul terhadap proses keradangan dan produksi NO secara besar-besaran yang menjadi salah satu faktor yang menyumbang kearah penyakit degeneratif kronik seperti kanser dan arthritis. Adalah menjadi keperluan untuk mencari sumber alternatif daripada hasil semulajadi seperti pokok kerana ia memberikan harapan yang tinggi dalam mengenal pasti sebatian bioaktif, impak negatif yang kurang dan pembangunannya dalam menjadikan ubat untuk mengubati penyakit berkaitan anti-radang. Kajian fitokimia seperti pengekstrakan, pengasingan dan mengenal pasti bentuk sebatian hasil semulajadi dengan bantuan FTIR, GCMS, HRESIMS dan NMR telah dijalankan keatas batang kulit kayu, dua spesis keluarga pokok Calophyllaceae iaitu *Mesua hexapetala* dan *Mesua beccariana* serta satu spesis daripada keluarga pokok Clusiaceae iaitu *Garcinia mangostana* yang telah memberi dua puluh metabolit sekunder, tiga daripadanya telah dikenal pasti sebagai sebatian baru.

Kajian fitokimia terhadap batang kulit kayu *Mesua hexapetala* telah menghasilkan penemuan sebatian kumarin yang baru, hexapetarin (**143**) bersama-sama dengan 1,3,7-trihidroksi-2,4-di(3-metil-2-butenil)xanton (**144**), trapezifolixanton (**96**),

kudraxanton G (**19**), stigmasterol (**17**), β -sitosterol (**65**) dan γ -sitosterol (**145**). Tiada penyelidikan dan laporan kajian yang dilaporkan terhadap pokok ini. Di samping itu, melalui kajian terhadap batang kulit kayu, *Mesua beccariana*, xanton baru, bekarianton T (**146**) dan kumarin baru, bekamarin T (**147**) telah berjaya diasingkan bersama-sama dengan mesuarianon (**1**), mesuasanon (**2**), 1,5-dihidroksixanton (**57**), euxanton (**20**) dan empat triterpenoids biasa, fridelin (**8**), stigmasterol (**7**), β -sitosterol (**65**) dan γ -sitosterol (**145**). Daripada pokok *Garcinia mangostana*, sebatian kimia yang diasingkan ialah α -mangostin (**14**), β -mangostin (**88**), mangostenol (**137**), garcinon D (**132**), fuskaxanton C (**148**), dulsixanton F (**94**) dan **7**.

Malah, reaksi semi-sintesis terhadap β -mangostin iaitu asetilasi dan *O*-alkilasi yang memberikan hasil enam penerbitan β -mangostin dimana empat penerbitan merupakan penerbitan baru, 6-monoasetat β -mangostin (**149**), 6-*O*-metil β -mangostin (**148**), 6-*O*-benzil β -mangostin (**150**), 6-*O*-*n*-hexil β -mangostin (**151**), 6-*O*-*n*-butil β -mangostin (**152**) and 6-*O*-sek-butil β -mangostin (**153**). Kesemua ekstrak pokok dan juga tujuh sebatian semulajadi serta empat semi-sintetik β -mangostin telah dipilih untuk dinilai terhadap cerakin nitrik oksida dan sitotoksik berasaskan sel. Aktiviti anti-radang telah mengenal pasti empat sebatian iaitu **145**, **144**, **14** dan **88** telah menunjukkan aktiviti yang sangat ketara terhadap penyekatan RAW 264.7 makrofaj yang dirangsangkan oleh LPS/IFN- γ dengan nilai IC_{50} 30.79 ± 2.68 , 12.41 ± 0.89 , 29.81 ± 0.77 and 11.72 ± 1.16 μ M masing-masing. Ekstrak etil asetat *G.mangostana* dan ekstrak n-hexana *M.hexapetala* menunjukkan aktiviti yang sangat berpotensi. Kajian terhadap hubungan struktur-aktiviti sebatian tulen dan penerbitan semi-sintetik β -mangostin terhadap anti-radang menunjukkan kepentingan kewujudan kumpulan hidroksil dalam struktur sebatian yang mempengaruhi bioaktiviti sebatian tersebut.

Ujian anti-*Bacillus* yang dijalankan terhadap tujuh jenis sebatian tulen iaitu **145**, **146**, **147**, **144**, **2**, **14** dan **88** dengan semua ekstrak pokok daripada pokok-pokok yang dikaji terhadap empat jenis bakteria *Bacillus*. Hexapetarin (**143**) telah menunjukkan aktiviti anti-*Bacillus* yang sangat berpotensi terhadap empat jenis bakteria *Bacillus* - *Bacillus subtilis*, *B.cereus*, *B.megaterium* dan *B.pumilus*. Tuntasnya, boleh disimpulkan bahawa penemuan penerbitan baru kumarin, hexapetarin (**143**), ekstrak pokok n-hexana *M.hexapetala* dan ekstrak etil asetat pokok *G.mangostana* menunjukkan aktiviti anti-radang yang sangat signifikan terhadap anti-*Bacillus* sementara itu, **144**, **14** dan **88** mempamerkan aktiviti anti-radang yang sangat berpotensi dimana boleh diperkembangkan sebagai ejen anti-radang di masa hadapan.

ACKNOWLEDGEMENTS

First and foremost, I would like to express my deepest gratitude and appreciation to my beloved supervisor, YBhg. Prof. Dr. Gwendoline Ee Cheng Lian for giving me the chance to carry out my PhD studies under her guidance and supervision with great patience, as well as giving me many opportunities to gain knowledge in the field of organic chemistry, especially in phytochemistry, spectroscopic chemistry as well as in pharmacology. Her guidance has helped me throughout my research and writing of this thesis. I cannot imagine having a better advisor and mentor for my PhD study.

I would like to express my gratitude to my co-supervisor, Assoc. Prof Dr. Intan Safinar Ismail for giving me the chance to explore the world of biology and pharmacology under her guidance as well as for the motivation she has supplied me with along the way. I would also like to thank her for allowing me to use the facilities and equipment in her Laboratory of Natural Products (LHS), Institute of Bioscience. I would also like to thank my co-supervisor from the field of organic synthesis, Dr. Siti Mariam Mohd Nor, for her guidance and help in my semi-synthesis work. Special thanks goes to fellow researchers from LHS - Dr. Leong Sze Wei, Dr. Ramesh Kumar Santhanam, Assoc. Prof Dr. Yaya Rukayadi, Miss Kalaivani Palachandran and Miss Hong Xia - for their guidance and motivation in my biological work, which is a new field explored by me.

Not forgetting my friends who have always given me encouragement, motivation as well as helped me to face difficult situations during my PhD studies, Mr. Mahashanon Arumugam (PutraCAT, UPM), Miss Annushiah Vasanthakumar (USM), Mr. Kumar Arumgam (Jasin, Melaka), Dr. Shaari Daud (UiTM), Miss Naveena Reddy Kalidass (LHS, IBS, UPM), Miss Arthy Surendran (INTROP, UPM), Miss Pireya Arulu (Biotech, UPM), Mr. Yugesh Gannason (UCSI), Mrs. Guhan Uthayakumar (Ulsan, Korea), Mr. Fadzly Salleh (Vet, UPM), Dr. Raghunath Parayani (LHS, IBS, UPM), Dr. Karthivashan Govindasamy (Seoul, Korea), Dr. Peter Waziri (Nigeria) and others who were with me during these challenging times.

I would also like to express my gratitude to all my lab mates from the Chemistry Department, Faculty of Science, UPM, as well as lab mates from the Laboratory of Natural Products (LHS), Miss Nor Hisham Zamakshshari, Dr. Irene See, Mr. Lee Kar Wei, Dr. Teh Soek Sin, Mr. Ahmad Azri Fitri, Mr. Muniandy, Mr Wong Ka Woong, Mr. Lim Wui Zhuan, Mr. Taufiq Astifa, Miss Ashikin, Miss Kayne Lee, Miss Khaleeda, Mrs Ilya Iryani, Miss Amalina Azam, Miss Foong Yen Lum, Mrs Asila Mazlan, Mr. Abdel Ghani, Mr. Tee Keng Hong and Miss Ashkwinie Siramaloo, for their willingness to help and share their knowledge with me as well as being good listeners when I needed someone to talk to. All of them have made an awesome working environment for me which united us all as one team. I am really happy to have been blessed the opportunity to work with them all.

Special thanks goes to science officers Ms. Shareena Safiai (NMR) and Mr. Johadi Iskandar (NMR) in their guidance for using the NMR instrument. I am also grateful to Mr. Zainal Abidin Kassim (GCMS), Mrs. Rusnani Amirudin (FTIR), Mrs. Noriza Atan (UV-Vis), Mrs. Nurul Huda (Bioassay Lab, LHS) and Mr. Azizul Isha (Phyto Lab, LHS) for giving me the chance to analyze all my compounds for GCMS, IR, UV-Vis and biological analysis. I am highly indebted to Dr. Vivian Jong Yi Mian and Prof Dr. Jegak Uli for collection of plant samples.

My deepest appreciation goes to my beloved mother who has always given her moral support and wisdom in guiding me towards righteousness in the absence of my father in my life. Special thanks to my sister, Sharvina Karunakaran and relatives for their kindness and thoughtfulness. Moreover, I would like to express my appreciation to individuals around me for their beneficial advices and critics that have impacted my research methods and myself, which have indirectly led to the completion of my PhD studies successfully. Last but not least, my greatest and deepest gratitude to God Almighty who has always been beside me and answered all of my prayers. God Almighty has given me the courage to face all of the obstacles and difficulties that I have faced throughout my PhD journey and with His blessing and grace, I would be able to finish my PhD journey successfully.

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfillment of requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Gwendoline Ee Cheng Lian, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Intan Safinar Ismail, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Siti Mariam Mohd Nor, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: Professor Dr. Gwendoline Ee Cheng Lian

Signature: _____
Name of
Member of
Supervisory
Committee: Associate Professor Dr. Intan Safinar Ismail

Signature: _____
Name of
Member of
Supervisory
Committee: Dr. Siti Mariam Mohd Nor

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST APPENDICES	xx
LIST OF ABBREVIATIONS	xxii
 CHAPTER	
1 INTRODUCTION	1
1.1 General Introduction	1
1.2 Problem Statements	3
1.3 Objectives of Study	4
 2 LITERATURE REVIEW	 5
2.1 Family of Calophyllaceae and Clusiaceae	5
2.1.1 Taxonomy of Calophyllaceae	5
2.1.1.1 Genus <i>Mesua</i>	5
2.1.2 Taxonomy of Clusiaceae	9
2.1.2.1 Genus <i>Garcinia</i>	9
2.2 Phenolic compounds	11
2.2.1 Xanthones	12
2.2.2 Coumarins	14
2.2.3 Terpenoids	15
2.2.4 Triterpenoids	17
2.3 Phytochemical review on genus <i>Mesua</i>	17
2.3.1 Phytochemical constituent from genus <i>Mesua</i>	17
2.4 Phytochemical review on genus <i>Garcinia</i>	36
2.4.1 Phytochemical constituent of species <i>Garcinia mangostana</i>	36
2.5 Semisynthesis and structural modification of xanthones	48
2.6 Biological Activities of Family Calophyllaceae and Clusiaceae	50
2.6.1 Biological activities from <i>Mesua</i> species	50
2.6.2 Biological activities of species <i>Garcinia mangostana</i>	54
2.7 Inflammation	56
2.8 <i>Bacillus</i> type Bacteria as Foodborne Pathogens	58
 3 MATERIALS AND METHODS	 59
3.1 Materials	59
3.2 General Instruments	61

3.2.1	Melting points	61
3.2.2	Polarimeter	61
3.2.3	Ultra Violet Spectroscopy (UV)	61
3.2.4	Mass Spectrometer (MS)	61
3.2.5	Fourier Transform Infrared Spectroscopy (FTIR)	61
3.2.6	Nuclear Magnetic Resonance (NMR)	61
3.3	Extraction	62
3.4	Semi-Synthesis (Acetylation and <i>O</i> -Alkylation of β -mangostin (88)	62
3.4.1	Acetylation	62
3.4.2	<i>O</i> -Alkylation	62
3.5	Biological Activities	63
3.5.1	Sample preparation	63
3.5.2	Anti-Inflammatory Properties	63
3.5.2.1	Cell Culture	63
3.5.2.2	Cell Cytotoxic Test (MTT)	63
3.5.2.3	Nitrite Determination Assay (NO)	64
3.5.3	Anti- <i>Bacillus</i> Evaluation	65
3.5.3.1	Preparation of bacterial strains and inoculum	65
3.5.3.2	Disc Diffusion	66
3.5.3.3	Determination of Minimal Inhibitory Concentration (MIC) and Minimal Bactericidal Concentration (MBC)	66
3.6	Statistical analysis	67
4	RESULTS AND DISCUSSION	68
4.1	Isolation of Phytochemical Constituents from <i>Mesua hexapetala</i> , <i>Mesua beccariana</i> and <i>Garcinia mangostana</i>	68
4.1.1	<i>Mesua hexapetala</i>	68
4.1.2	<i>Mesua beccariana</i>	69
4.1.3	<i>Garcinia mangostana</i>	70
4.2	Phenolic Compounds from The Stem Bark of <i>Mesua hexapetala</i> , <i>Mesua beccariana</i> and <i>Garcinia mangostana</i>	70
4.2.1	New Phytochemical Constituents from Stem Bark of <i>Mesua hexapetala</i> and <i>Mesua beccariana</i> .	71
4.2.1.1	Isolation of Hexapetarin (143)	71
4.2.1.2	Characterization of Hexapetarin (143)	72
4.2.2	Isolation of Beccarixanthone T (146)	91
4.2.2.1	Characterization of Beccarixanthone T (146)	91
4.2.3	Isolation of Beccamarin T (147)	106
4.2.3.1	Characterization of Beccamarin T (147)	106
4.3	Known Phytochemical Constituents Isolated from Stem Bark <i>Mesua hexapetala</i> , <i>Mesua beccariana</i> and <i>Garcinia mangostana</i> .	123
4.3.1	Known Phytochemical Constituents from Stem Bark of <i>Mesua hexapetala</i>	123
4.3.1.1	Characterization of Trapezifolixanthone (96)	123
4.3.1.2	Characterization of Cudraxanthone G (19)	125

4.3.1.3	Characterization of 1,3,7-trihydroxy-2,4-di(3-methyl-2-enyl)xanthone (144)	128
4.3.2	Known Phytochemical Constituents from Stem Bark <i>Mesua beccariana</i>	130
4.3.2.1	Characterization of Mesuarianone (1)	130
4.3.2.2	Characterization of Mesuasinone (2)	133
4.3.2.3	Characterization of Euxanthone (20)	135
4.3.2.4	Characterization of 1, 5-dihydroxyxanthone (57)	137
4.3.3	Known Phytochemical Constituents from Stem Bark of <i>Garcinia mangostana</i>	139
4.3.3.1	Characterization of α -mangostin (14)	139
4.3.3.2	Characterization of β -mangostin (88)	141
4.3.3.3	Characterization of Dulcixanthone F (94)	144
4.3.3.4	Characterization of Garcinone D (132)	147
4.3.3.5	4.3.3.4 Characterization of Mangostenol (137)	150
4.3.3.6	Characterization of Fuscaxanthone C (148)	154
4.4	Semi-Synthesis (Acetylation and <i>O</i> -Alkylation of β -mangostin (88))	156
4.4.1	Acetylation of β -mangostin (88)	160
4.4.1.1	Characterization of 6-monoacetate β -mangostin (149)	163
4.4.2	<i>O</i> -Alkylation of β -mangostin (88)	164
4.4.2.1	Characterization of 6- <i>O</i> -alkylated β -mangostin (88)	166
4.5	Anti-Inflammatory Properties	173
4.5.1	Cytotoxicity Test (MTT)	174
4.5.2	Nitrite Determination (NO)	177
4.6	Anti- <i>Bacillus</i> Evaluation	183
4.6.1	Disc Diffusion Test	183
4.6.2	MIC and MBC Determination	185
5	SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	187
5.1	Summary	187
5.2	Conclusion	188
5.3	Recommendations for future research	189
	REFERENCES	190
	APPENDICES	197
	BIODATA OF STUDENT	291
	LIST OF PUBLICATIONS	292

LIST OF TABLES

Table		Page
1	Classification of Calophyllaceae	5
2	Classification of species <i>Mesua hexapetala</i>	6
3	Classification of species <i>Mesua beccariana</i>	7
4	Classification of Clusiaceae	9
5	Classification of species <i>Garcinia mangostana</i>	9
6	Phytochemical Constituents of <i>Mesua</i> species	18
7	Phytochemical constituent from <i>Garcinia mangostana</i>	37
8	Biological activities of <i>Mesua</i> species	52
9	Biological activities of the species <i>Garcinia mangostana</i>	54
10	¹ H-NMR (500 MHz, CDCl ₃) and ¹³ C-NMR (125 MHz, CDCl ₃) of hexapetarin (143)	75
11	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for	93
12	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for beccamarin T (147)	109
13	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃)	125
14	¹ H NMR (500 MHz, Acetone- <i>d</i> ₆), ¹³ C NMR (125 MHz, Acetone- <i>d</i> ₆) for	127
15	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for 1,3,7-trihydroxy-2,4-di(3-methyl-2-enyl)xanthone (144)	130
16	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for mesuarianone (1)	132
17	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for mesuasinone (2)	135
18	¹ H NMR (500 MHz, Acetone- <i>d</i> ₆), ¹³ C NMR (125 MHz, Acetone- <i>d</i> ₆) for euxanthone (20)	137

19	¹ H NMR (500 MHz, Acetone- <i>d</i> ₆), ¹³ C NMR (125 MHz, Acetone- <i>d</i> ₆) for 1,5-dihydroxyxanthone (57)	139
20	¹ H NMR (500 MHz, Acetone- <i>d</i> ₆), ¹³ C NMR (125 MHz, Acetone- <i>d</i> ₆)	141
21	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for β-mangostin (88)	144
22	¹ H NMR (500 MHz, Acetone- <i>d</i> ₆), ¹³ C NMR (125 MHz, Acetone- <i>d</i> ₆) for	147
23	¹ H NMR (500 MHz, Acetone- <i>d</i> ₆), ¹³ C NMR (125 MHz, Acetone- <i>d</i> ₆)	150
24	¹ H NMR (500 MHz, Acetone- <i>d</i> ₆), ¹³ C NMR (125 MHz, Acetone- <i>d</i> ₆) for mangostenol (137)	153
25	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for fuscaxanthone C (148)	156
26	Reactants, reaction conditions and product yield of semisynthetic β-mangostin derivatives.	158
27	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for 6-monoacetate β-mangostin (149)	164
28	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for	167
29	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for 6- <i>O</i> - <i>n</i> -hexyl β-mangostin (151)	169
30	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for 6- <i>O</i> - <i>n</i> -butyl β-mangostin (152)	171
31	¹ H NMR (500 MHz, CDCl ₃), ¹³ C NMR (125 MHz, CDCl ₃) for 6- <i>O</i> - <i>sec</i> -butyl β-mangostin (153)	173
32	Inhibitory activities of plant extracts and selected pure compounds on IFN-γ /LPS induced NO production in RAW 264.7 macrophages.	182
33	Inhibition zone of plant extracts and selected phytochemicals towards four type of <i>Bacillus</i> bacteria.	184
34	MIC and MBC activity of selected plant extracts and compound 143 towards four type of <i>Bacillus</i> bacteria.	186

LIST OF FIGURES

Figure		Page
1	Botanical features of <i>Mesua hexapetala</i> (Taken from Ashton, 1988)	7
2	The stem bark of <i>Mesua beccariana</i>	8
3	The leaves and twig of <i>Mesua beccariana</i>	8
4	The tree shrub of Mangosteen (<i>Garcinia mangostana.L</i>)	10
5	The view of <i>Garcinia mangostana.L</i> tree in its fruit season	10
6	Main streams of phenolic compounds	12
7	Basic Structure of Xanthone (Taken from Teh et al., 2010)	13
8	Biosynthesis for xanthenes (Taken from Negi et al. 2013)	14
9	The basic structure of coumarin (Taken from Venugopala et al., 2013)	15
10	Biosynthesis of coumarin	15
11	The main stream of terpenoids biosynthesis	16
12	Structure of squalene (Taken from Szakiel et al., 2012)	17
13	Biosynthesis of Nitric Oxide (NO)	57
14	Simplified overview of NO production and how it cause the physiological effects (taken from Ricciardolo et al, 2004.)	57
15	Simplified overview of NO production and how it cause the pathophysiological effects (taken from Ricciardolo et al, 2004.)	58
16	Flow Chart of Research Activities	60
17	Flow Chart of Cytotoxicity and Nitric Oxide Inhibition Assay	65
18	Phytochemicals Isolated from <i>Mesua hexapetala</i>	68
19	Phytochemicals Isolated from <i>Mesua beccariana</i>	69

20	Phytochemicals Isolated from <i>Garcinia mangostana</i>	70
21	FTIR Spectrum of Hexapetarin (143)	76
22	EIMS Spectrum of Hexapetarin (143)	77
23	HRESIMS Spectrum of Hexapetarin (143)	78
24	¹ H-NMR Spectrum of Hexapetarin (143) (500 MHz, CDCl ₃)	79
25	Expansion of ¹ H-NMR Spectrum of Hexapetarin (143) (500 MHz, CDCl ₃)	80
26	¹³ C-NMR Spectrum of Hexapetarin (143) (125 MHz, CDCl ₃)	81
27	DEPT NMR Spectrum of Hexapetarin (143) (125 MHz, CDCl ₃)	82
28	COSY NMR Spectrum of Hexapetarin (143)	83
29	Expansion of HMQC Spectrum of Hexapetarin (143)	84
30	Expansion of HMQC Spectrum Hexapetarin (143)	85
31	HMBC Spectrum of Hexapetarin (143)	86
32	Expansion of HMBC Spectrum of Hexapetarin (143)	87
33	Expansion of HMBC Spectrum of Hexapetarin (143)	88
34	Expansion of HMBC Spectrum of Hexapetarin (143)	89
35	Expansion of HMBC Spectrum of Hexapetarin (143)	90
36	FTIR Spectrum of Beccarixanthone T (146)	94
37	EIMS Spectrum of Beccarixanthone T (146)	95
38	HRESIMS Spectrum for Beccarixanthone T (146)	96
39	¹ H-NMR Spectrum of Beccarixanthone T (146) (500 MHz, CDCl ₃)	97
40	Expansion of ¹ H-NMR Spectrum of Beccarixanthone T (146) (500 MHz, CDCl ₃)	98
41	¹³ C-NMR Spectrum of Beccarixanthone T (146) (125 MHz, CDCl ₃)	99

42	DEPT NMR Spectrum of Beccarixanthone T (146) (125 MHz, CDCl ₃)	100
43	COSY NMR Spectrum of Beccarixanthone T (146)	101
44	HMQC Spectrum of Beccarixanthone T (146)	102
45	HMBC Spectrum of Beccarixanthone T (146)	103
46	Expansion of HMBC Spectrum of Beccarixanthone T (146)	104
47	Expansion of HMBC Spectrum Beccarixanthone T (146)	105
48	FTIR Spectrum of Beccamarin T (147)	110
49	EIMS Spectrum of Beccamarin T (147)	111
50	HRESIMS Spectrum of Beccamarin T (147)	112
51	¹ H-NMR Spectrum of Beccamarin T (147) (500 MHz, CDCl ₃)	113
52	Expansion of ¹ H-NMR Spectrum of Beccamarin T (147) (500 MHz, CDCl ₃)	114
53	¹³ C-NMR Spectrum of Beccamarin T (147) (125 MHz, CDCl ₃)	115
54	DEPT NMR Spectrum of Beccamarin T (147) (125 MHz, CDCl ₃)	116
55	COSY Spectrum of Beccamarin T (147)	117
56	HMQC Spectrum of Beccamarin T (147)	118
57	HMBC Spectrum of Beccamarin T (147)	119
58	Expansion of HMBC Spectrum of Beccamarin T (147)	120
59	Expansion of HMBC Spectrum of Beccamarin T (147)	121
60	Expansion of HMBC of Beccamarin T (147)	122
61	Reaction scheme of semi-synthetic derivatives of β-mangostin (88)	157
62	Mechanism reaction of acetylation for 6- <i>O</i> -monoacetate β-mangostin (149)	162
63	Mechanism of <i>O</i> -alkylation for β-mangostin derivatives	165

64	Cytotoxicity of Crude Extracts <i>M.hexapetala</i> against RAW 264.7 macrophages. All experiments were conducted in triplicate and results are expressed as mean \pm S.E.M.	175
65	Cytotoxicity of Selected Pure Compounds against RAW 264.7 macrophages. All experiments were conducted in triplicate and results are expressed as mean \pm S.E.M.	176
66	All experiments were conducted in triplicate and results are expressed as mean \pm S.E.M. P values are determined by one way analysis of variance followed by Dunnett test (### $p < 0.001$ vs control; *** $p < 0.001$, ** $p < 0.01$, and * $p < 0.05$ vs IFN- γ / LPS treatment).	179
67	All experiments were conducted in triplicate and results are expressed as mean \pm S.E.M. P values are determined by one way analysis of variance followed by Dunnett test (### $p < 0.001$ vs control; *** $p < 0.001$, ** $p < 0.01$, and * $p < 0.05$ vs IFN- γ / LPS treatment).	180
68	All experiments were conducted in triplicate and results are expressed as mean \pm S.E.M. P values are determined by one way analysis of variance followed by Dunnett test (### $p < 0.001$ vs control; *** $p < 0.001$, ** $p < 0.01$, and * $p < 0.05$ vs IFN- γ / LPS treatment).	181
69	Disc Diffusion Test of Selected Pure Compounds against Four Types of <i>Bacillus</i> Bacteria	184

LIST OF APPENDICES

Appendix	Page
A Characterization of Trapezifolixanthone (96)	197
B Characterization of Cudraxanthone G (19)	201
C Characterization of 1,3,7-trihydroxy-2,4-di(3-methyl-2-enyl)xanthone (144)	207
D Characterization of Mesuarianone (1)	211
E Characterization of Mesuasione (2)	215
F Characterization of Euxanthone (20)	220
G Characterization of 1,5-dihydroxyxanthone (57)	223
H Characterization of α -mangostin (14)	226
I Characterization of β -mangostin (88)	230
J Characterization of Dulcisxanthone F (94)	234
K Characterization of Garcinone D (132)	238
L Characterization of Mangostenol (137)	242
M Characterization of Fuscaxanthone C (148)	248
N Characterization of 6-monoacetate β -mangostin (149)	252
O Characterization of 6- <i>O</i> -benzyl β -mangostin (150)	257
P Characterization of 6- <i>O</i> - <i>n</i> -hexyl β -mangostin (151)	262
Q Characterization of 6- <i>O</i> - <i>n</i> -butyl β -mangostin (152)	267
R Appendix: Characterization of 6- <i>O</i> - <i>sec</i> -butyl β -mangostin (153)	272
S Cytotoxicity of Plant Extracts against RAW 264.7 macrophages.	277

T	Cytotoxicity of Selected Pure Compounds against RAW 264.7 macrophages.	279
U	Effects of various concentrations of plant extracts of <i>M. hexapetala</i> (MHC, MHE, MHA) on NO production by IFN- γ / LPS induced RAW 264.7 macrophages.	280
V	Effects of various concentrations of plant extracts of <i>M. beccariana</i> (MBH, MBC, MBE, MBA) on NO production by IFN- γ / LPS induced RAW 264.7 macrophages.	282
W	Effects of various concentrations of plant extracts of <i>G.mangostana</i> (GMH, GMC, GME, GMA) on NO production by IFN- γ / LPS induced RAW 264.7 macrophages.	284
X	Effects of various concentrations of selected phytochemicals isolated from <i>M.hexapetala</i> , <i>M.beccariana</i> and <i>G.mangostana</i> (14, 88, 146, 147 and 2) on NO production by IFN- γ / LPS induced RAW 264.7 macrophages.	286
Y	Effects of various concentration of selected semi-synthetic derivatives of β -mangostin (149, 148, 150, and 151) on NO production by IFN- γ / LPS induced RAW 264.7 macrophages.	289

LIST OF ABBREVIATIONS

δ	Chemical shift in ppm
λ_{\max}	Wavelength maxima in nm
$^{\circ}\text{C}$	Degree celcius
^1H	Proton
^{13}C	Carbon-13
CHX	Chlorhexidine
Cm	centimeter
D	Doublet
D	Deuterated proton
DMEM	Dulbecco's Modified Eagle's medium
DMSO	Dimethyl sulfoxide
ESI	Electrospray Ionization
FBS	Fetal Bovine Serum
FTIR	Fourier Transform Infra-Red
G	Gram
GC-MS	Gas chromatography-mass spectroscopy
HIV	Human immunodeficiency virus
HPLC	High performance liquid chromatography
Hz	Hertz
IC ₅₀	Initial Concentration to kill 50% of cells
IR	Infra-Red
IFN- γ	Interferon gamma
J	Coupling constant
LPS	Lipopolysaccharides
m	meter
[M] ⁺	Molecular ion
MBC	Minimal Bactericidal Concentration
MIC	Minimal Inhibitory Concentration

Mg	Milligram
MHA	Mueller Hinton Agar
MHB	Mueller Hinton Broth
MHz	Megahertz
m.p.	Melting point
MS	Mass spectrometry
MTT	3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide
m/z	Mass per charge
NMR	Nuclear Magnetic Resonance
NO	Nitric oxide
s	Singlet
d	Doublet
dd	Doublet of doublet
t	Triplet
q	Quartet
m	Multiplet
TLC	Thin layer chromatography
TMS	Tetramethylsilane
TSB	Tryptic Soybean Broth
µg	microgram
µM	Micro molar
UATR	Universal attenuated total reflection
UV	Ultraviolet
ν_{\max}	Wavenumber maxima in cm^{-1}
γ	gamma
α	alpha
β	beta

CHAPTER 1

INTRODUCTION

1.1 General Introduction

Natural products from flora and fauna have become the essential source for discovery of new drugs based on the new chemical compounds found through the studies conducted by researchers around the globe. Throughout the ages, since ancient times, natural products have played a vital role in the daily life of human beings especially in traditional medicine. It is known as the source of active ingredients of medicine used to treat various diseases. Historically, most of the new drugs discovered from natural products and compounds derived from it. Most bioactive natural products are discovered from sources like plants, bacteria and fungi. It has become the main source of active ingredients for medicines that been used to treat various diseases (Harvey, 2008).

The usage of natural products in the medicinal world can be traced from the ancient civilization such as Egyptian medicine which dates from 1500 BC. It was found in the Egyptian pharmaceutical record known as “Ebers Papyrus” that more than 700 drugs were mostly from plants, animal organs and natural minerals used during the civilization for medicinal purpose to treat various disease. In the eastern civilization like the Chinese and Indian traditional medicine, the Chinese *Materia Medica* and Indian *Ayurveda* scripts documented the usage of over 800 drugs from natural products especially from plants for various diseases dating from 1000 BC (Newman et al., 1999).

Bioactive natural products especially the secondary metabolites isolated from bioactive samples, including plants, microbes and animals and also semisynthetically derived compounds showed that a great diversity of chemical structures are used in a wide range of therapeutic medicinal applications such as anti-cancer, anti-inflammatory, anti-biotic, anti-diabetic and neurological disorders. Moreover, most of these natural products are found to obey the Lipinski's Rule of Five for orally available compounds indicating that natural product drugs have good absorbance properties compared to synthetic drugs (Harvey, 2008).

Plants have been used as drugs for thousands of years. Phytochemistry is the study of chemistry of plants which concentrates on the phytochemical constituents. Phytochemical constituents consist of primary and secondary metabolites produced by plants through biosynthesis. Phytochemical studies focused more on the isolation, structural identification and application of the secondary metabolites as these compounds have the potency to be developed as lead for various types of diseases and application. Phytochemistry is mainly focused on the secondary metabolites. Secondary metabolites are made up of molecules from reactions which are not

common to all life forms, but occur in certain species only. Secondary metabolism is of most interest to organic chemists. The structures of secondary metabolites are extraordinarily complex and their biosynthesis have been studied by organic chemists and biochemists to understand how plants system in produce phytochemicals. Secondary metabolism produces products that aid the growth and development of plants but are not required by the plants to survive. Secondary metabolism facilitates the primary metabolism in plants. Most of the phytochemicals discovered have been reported to be active towards human biology. Throughout the ages, plants have become the foundation for traditional medicine systems in eastern countries like China and India.

The field of phytochemistry has helped to explore and understand the world of traditional and herbal medicine. There is no proper documentation on the chemistry of drugs used in traditional and herbal medicine prescriptions since ancient times till the 1900s. Phytochemical studies have enabled researchers to identify bioactive lead compounds responsible for the medicinal properties with the aid from analytical chemistry and spectroscopic techniques. Moreover, the field of phytochemistry studies are required to expose the fourteen myths of traditional medicine which been argued by the synthetic chemists and pharmacist over the years. The fourteen myths are listed below (Cordell, 2014).

- Safety and effectiveness of traditional medicine eventhough it has been used for thousands of years.
- Using the “correct” plant is adequate.
- The biological effects are the same for different parts of plants.
- The biological effects are the same irrespective of the origin of the plants.
- The biological effects are the same irrespective of the plants preparation.
- Older plant material is less effective (or more toxic).
- “Wild”-collected plants are more active than cultivated plants.
- All of the materials in a complex prescription are necessary for effectiveness.
- Complex mixtures of plants cannot be standardized.
- This medicinal plant product is already well-regulated.
- Medicinal plants are safer than synthetic drugs.
- Medicinal plant will always be available.
- Traditional knowledge will always be there.
- Traditional medicines are not effective, and are therefore not worth considering as a part of a contemporary health care system.

Phytochemicals can be divided into six main groups which are phenolics, terpenoids, alkaloids, glycosides, tannins and saponins.

1.2 Problem Statements

Natural products or natural product derived compounds represent great structural diversity, which is not commonly seen in synthetic compounds. It can be concluded that natural products play a vital role in the discovery of leads for to be developed into drugs for treating human diseases. Natural products offer great hope in the identification of bioactive compounds and their development into drugs for the treatment many types of diseases. Besides that, plants have been the source for many traditional medicine systems throughout the world for thousands of years and continue to provide mankind with new remedies (Harvey, 2008). There is no proper documentation on the chemistry of drugs used in traditional and herbal medicine prescriptions since times till the 1900s (Newman et al., 1999).

Phytochemical studies have enabled researchers to identify bioactive hit compounds responsible for medicinal properties with the aid of analytical and spectroscopic chemistry techniques. Through phytochemical studies, researchers can prove the effectiveness of plant based medicines which have been used since ancient civilizations as well as the safety of plant based medicines compared to available synthetic drugs (Cordell, 2014). Moreover, it is necessity to explore alternative sources of medicinal plants rather than depend on available plants especially cultivated plants. Wild plants especially new species that have never been explored by any researchers should be starting point in the phytochemical and biological studies so as to provide new sources in the medicinal world hence enable medicinal supplies and stock which will continue to support the human needs.

There are several records in traditional medicine focusing on relief from pain and inflammation. Since ancient time, people suffering from inflammation were treated with phytochemicals, which is evident from the discovery of the first analgesic drug aspirin. The discovery of aspirin was based on the known analgesic and antipyretic properties of the bark of willow tree since 400 BC by the Greeks and Romans. Nitric oxide plays a vital role in the key molecular signaling constituent involved in inflammatory processes. Mass production of NO is one of the main factors that contribute towards chronic degenerative diseases such as cancer, arthritis, neurodegenerative and cardiovascular disorder (Leong et al. 2015; Leong et al. 2014). Biological interference of phytochemicals with NO production and bacterial inhibition are necessities in developing potent leads for anti-inflammatory and anti-bacterial related diseases.

From the previous researches on species such as *Mesua* and *Garcinia*, there are lot of potential secondary metabolites mostly xanthenes, coumarins and flavonoids that have been isolated recently with high bioactivities especially in anti-inflammatory especially in nitric oxide inhibition and anti-bacterial activities. In a nutshell, it is pivotal for chemists with the aid from the field of phytochemistry, analytical chemistry, spectroscopic chemistry and biological studies to continue the search for potential bioactive phytochemicals from plants in order to find an alternative source

of medicinal plants as well as do proper documentation of the chemistry of drugs present in the plants studied.

1.3 Objectives of Study

The research project is designed to isolate, purify and identify the structures of phytochemical compounds from *Mesua hexapetala*, *Mesua beccariana* and *Garcinia mangostana*. Structural modification will be carried out on the major constituents from the plants and the structural activity relationship study will be studied. The main purpose of this research is to study the phytochemistry of a new plant species *Mesua hexapetala* as well as the two other species *Mesua beccariana* and *Garcinia mangostana* and their potential bioactivities. Hence, the specific objectives of the research are:

1. To extract, isolate and elucidate the structures of the chemical constituents from *Mesua hexapetala*, *Mesua beccariana* and *Garcinia mangostana* using various chromatographic techniques and spectroscopic methods including IR, GCMS, UV-VIS, HRESIMS, and NMR.
2. To carry out structural modifications of β -mangostin isolated from *Garcinia mangostana* with simple synthesis method.
3. To determine cytotoxicity, nitric oxide inhibition and anti-bacillus test on the isolated compounds and plant extracts of *Mesua hexapetala*, *Mesua beccariana* and *Garcinia mangostana* as well as the semi-synthetic derivatives of β -mangostin and study the structure activity relationship on anti-inflammatory activities between natural and semi-synthesized compound.

REFERENCES

- Alam MS, Jain N, Kamil M and Liyas M. 1987. Mesuein : a novel flavanone glycoside from *Mesua Ferrea*. Chem. Ind. 16:565-566.
- Aoyama T, Katayama T, Yamamoto M, Tanaka H, and Kon K. 1992. 'A new antitumor antibiotic product, demethylchartreusin. Isolation and biological activities' J. Antibiot. 45(6): 875-878.
- Ashton PS. 1988. Manual of non-dipterocarp trees of Sarawak. Volume II. Dewan Bahasa dan Pustaka (Sarawak Branch) for Forest Department Sarawak.
- Asai F, Tosa H, Tanaka T and Inuma M. 1995. A xanthone from pericarps of *Garcinia mangostana*. Phytochemistry. 39: 943-944.
- Awang K, Chan G, Litaudon M, Ismail NH, Martin MT and Gueritte F. 2010. 4-Phenylcoumarins from *Mesua elegans* with acetylcholinesterase inhibitory activity. Bioorg. Med. Chem. 18:7873-7877.
- Bala KR and Seshadri TR .1971. Isolation and synthesis of some coumarin components of *Mesua Ferrea seed oil*. Phytochemistry. 10:1131-1134.
- Balasundram N, Sundram K and Saman S. 2006. Phenolic compounds in plants and agriindustrial by products: antioxidant activity, occurrence, and potential uses. Food. Chem. 99: 191-203.
- Bandaranayak WM, Selliah SS and Sultanbawa MUS. 1975. Xanthenes and 4-phenylcoumarins of *Mesua thwaitessi*. Phytochemistry, 14:265-269.
- Bhattacharyya P, Chakrabartty P and Chowdhury BK. 1988. Measuarin: a new 4-phenyl-coumarin from *Mesua Ferrea*. Chem. Ind. 7:239-240.
- Chan W-M, Liu DTL, Chan CKM, Chong KKL, and Lam DSC. 2003. Infective endophthalmitis caused by *Bacillus cereus* after cataract extraction surgery. Clin. Infect. Dis. 37, 31-34.
- Chen LG, Yang LL and Wang CC. 2008. Anti-inflammatory activity of mangostins from *Garcinia mangostana*. Food. Chem. Toxicol. 46: 688-693.
- Chin YW, Jung HA, Chai H, Keller WJ and Kinghorn AD. 2008. Xanthenes with quinone reductase-inducing activity from the fruits of *Garcinia mangostana* (Mangosteen). Phytochemistry. 69: 754-758.
- Choundhury S, Ahmed R, Barthel A and Leclercq PA. 1998. Volatile oils of *Mesua Ferrea* (L) from Assam, India. J. Essent. Oil. Res. 10:497-501.
- Chow YL & Quan HH. 1968. Chemical constituents of the heartwood of *Mesua Ferrea*. Phytochemistry. 7:1871-1874.

- Christenhusz JMM and Byng JW. 2016. The number of known plant species in the world and its annual increase. *Phytotaxa*, 261(3): 201-207.
- Clinical and Laboratory Standards Institute (CLSI). 2012. Reference method for performance standards for antimicrobial disk susceptibility tests; approved standard-eleventh edition; CLSI document M02-A11. Clinical and Laboratory Standards Institute, Wayne, PA, USA.
- Clinical and Laboratory Standards Institute (CLSI). 2012. Reference method for performance standards for antimicrobial disk susceptibility tests; approved standard-eleventh edition; CLSI document M02-A11. Clinical and Laboratory Standards Institute, Wayne, PA, USA.
- Cordell GA. 2014. Phytochemistry and traditional medicine -The revolution continues. *Phytochem Lett.* 10: 28-40
- Corner EJJ. 1988. Mangosteen family. In *Wayside Trees of Malaya*. Malayan Nature Society: Kuala Lumpur.
- Dai J and Mumper R. 2010. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*. 15: 7313-7352.
- Deachathai S, Mahabusarakam W, Phongpaichit S, Taylor WC, Zhang Y-J and Yang C-R. 2005. Phenolic compounds from the flower of *Garcinia dulcis*. *Phytochemistry*, 67: 464-469.
- Don M-J, Huang Y-J, Huang R-L and Lin Y-L. 2004. 'New phenolic principles from *Hypericum sampsonii*'. *Chem. Pharm. Bull.* 52(7):866-869.
- Ee GCL, Lim CK, Rahmat A and Lee HL. 2005. Cytotoxic activities of chemical constituents from *Mesua daphnifolia*. *Trop. Biomed.* 22:99-102.
- Ee GCL, Lim CK, Ong GP, Sukari MA and Lee HL. 2006. Daphnifolin, a new xanthone from *Mesua daphnifolia*. *J. Asian Nat. Prod. Res.* 8:567-570.
- Ee GCL, Teh SS, Kwong HC, Mah SH, Lim YM and Rahmani M. 2012. A new benzophenone from *Mesua congestiflora*, an inhibitor against human B lymphocyte cancer cell line. *Phytochem. Lett.* 5:545-548.
- Ee GCL, See I, Teh SS and Daud S. 2014. A new furanoxanthone from *Garcinia mangostana*. *J. Asian Nat. Prod. Res.* 16(7):790-794.
- Farnsworth RN and Bunyapraphatsara N. 1992. *Garcinia mangostana* Linn. In *Thai Medicinal Plants*. Prachachon Co., Ltd.: Bangkok, 160-162.
- Faudzi SMM, Leong SW, Abas F, Aluwi MFFM, Rullah K, Lam KW, Ahmad S, Tham CL, Shaari K and Lajis NH. 2015. Synthesis, biological evaluation and QSAR studies of diarylpentanoid analogues as potential nitric oxide inhibitors. *Med. Chem. Commun.* 6:1069-1080.

Friedli G-L. Retrieved from <http://www.friedli/herbs/phytochem/coumarins.html>

Gopalkrishnan C, Shankaranarayanan D, Nazimuddin SK, Vishwanathan S and Kameswaran L. 1980. Anti-inflammatory and C.N.S. depressant activities of xanthenes from *Calophyllum inophyllum* and *Mesua ferrea*. Indian. J. Pharmacol. 12:181-91.

Govindachari TR, Pai BR, Subramaniam PS, Rao UR and Muthukumaraswamy N. 1964. Constituents of *Mesua Ferrea* L-II: Ferruol A, a new 4-alkylcoumarin. Tetrahedron., 23:4161-4165.

Gunasekara SP and Sultanbawa MUS. 1977. Chemical investigation of Ceylonese plants. Part 20. Extractives of *Mesua myrtifolia*. Isolation and structure of a new triterpene hydroxy-acid, myrtifolic. J. Chem. Soc. 1(1):6-10.

Ha LD, Hansen PE, Vang O, Duus F, Pham HD and Nguyen LHD. 2009. Cytotoxic geranylated xanthenes and O-alkylated derivatives of α -Mangostin. Chem. Pharm. Bull. 57(8):830-834.

Harvey AL. 2008. Natural products in drug discovery. Drug. Discov. Today. 13(19/20):894-901.

Heywood VH. 1982. In Popular Encyclopedia of Plants, ed. S.R. Chant. England: Press Syndicate of the University of Cambridge.

Iranshahi M, Askari M, Sahebkar A and Hadjipavlou-Litina D. 2009. "Evaluation of antioxidant, anti-inflammatory and lipoxygenase inhibitory activities of the prenylated coumarin umbelliprenin," DARU. 17(2): 99–103.

Isakovic A, Jankovic T, Harhaji L, Kostic-Rajacic S, Nikolic Z, Vajs V and Trajkovic V. 2008. Antiglioma action of xanthenes from *Gentiana kochiana*: Mechanistic and structure–activity requirements. Bioorg. Med. Chem. 16(2008): 5683-5694.

Ito C, Itoigawa M, Takakura T, Ruangrunsi N, Enjo F, Tokuda H, Nishino and Furukawa H. 2003. Chemical Constituents of *Garcinia fusca*: Structure elucidation of eight new xanthenes and their cancer chemopreventive activity. J. Nat. Pro. 66:200-205.

Jung HA, Su BN, Keller WJ, Mehta RG and Kinghorn AD. 2006. Antioxidant xanthenes from the pericarp of *Garcinia mangostana* (Mangosteen). J. Agric. Food. Chem. 54: 2077–2082.

Kerrigan RA, Cowie ID and Dixon DJ. 2011. Clusiaceae. In Short, P.S. & Cowie, I.D. (eds), *Flora of the Darwin Region*. (Northern Territory Herbarium, Department of Natural Resources, Environment, the Arts and Sport). 1:1–7. Retrieved from: http://www.nt.gov.au/nreta/wildlife/plants_herbarium/index.html.

- Kimouli M, Vrioni G, Papadopoulou M, Koumali V, Petropoulou D, Gounaris A, Friedrich AW and Tsakris A. 2012. Two cases of severe sepsis caused by *Bacillus pumilus* in neonatal infants. *J. Med. Microbiol.* 61:596-599.
- Leong SW, Faudzi SMM, Abas F, Aluwi MFFM, Rullah K, Wai LK, Bahari MNA, Ahmad S, Tham CL, Shaari K and Lajis NH. 2014. Synthesis and sar study of diarylpentanoid analogues as new anti-inflammatory agents, *Molecules*, 19: 16058-16081.
- Leong SW, Faudzi SMM, Abas F, Aluwi MFFM, Rullah K, Wai LK, Bahari MNA, Ahmad S, Tham CL, Shaari K and Lajis NH. 2015. Nitric oxide inhibitory activity and antioxidant evaluations of 2-benzoyl-6-benzylidenecyclohexanone analogues, a novel series of curcuminoid and diarylpentanoid derivatives, *Bioorg. Med. Chem. Lett.* 25:3330-3337.
- Mohamed AG, Ibrahim RMS, Shabaan IAM and Ross AS. 2014. Mangostanaxanthones I and II new xanthones from the pericarp of *Garcinia mangostana*. *Fitoterapia*, 98:215-221.
- Morel C, Guilet D, Oger JM, Seraphin D, Sevenet T, Wiart C, Hadi AHA, Richomme P and Bruneton J. 1999. 6-Acylcoumarins from *Mesua racemosa*. *Phytochemistry*, 50:1243-1247.
- Morel C, Dartiguelongue C, Youhane T, Oger J.-M, Seraphin D, Duval O, Richomme P and Bruneton J. 1999. New coumarins from *Mesua racemosa*: isolation and synthesis. *Heterocycles*, 51(9):2183-2191.
- Nasir ND, Rahmani M, Shaari K, Kassim NK, Go R, Stanslas J and Jeyaraj EJ. 2013. Xanthones from *Calophyllum gracilipes* and their cytotoxic activity (Xanton daripada *Calofilum gracilipes* dan aktiviti ketoksikannya). *Sains Malays.* 42(9):1261-1266
- Negi JS, Bisht VK, Singh P, Rawat MSM and Joshi GP. 2013. Naturally occurring xanthones: chemistry and biology. *J. Appl. Chem.* 2013 (Article ID 621459): 1-10. DOI:10.1155/2013/621459.
- Newman DJ, Cragg GM and Snader KM. 1999. The influence of natural products upon drug discovery. *Nat. Pro. Rep.* 17:215-234.
- Nilar and Harrison LJ. 2002. Xanthones from the heartwood of *Garcinia mangostana*. *Phytochemistry*. 60:541–548
- Obolskiy D, Pischel I, Siriwatanametanon N and Heinrich M. 2009. *Garcinia mangostana* L.: a phytochemical and pharmacological review, *Phyther. Res.* 23(8):1047–1065.
- Oldenburg TBP, Wilkes H, Horsfield B, Van-Duin ACT, Stoddart D and Wilhelms A. 2002. "Xanthones—novel aromatic oxygen-containing compounds in crude oils," *Org Geochem*, 33(5): 595–609.

Pedraza-Chaverri J, Cardenas-Rodriguez N, Orozco-Ibarra M and Perez-Rojas JM. 2008

Medicinal properties of mangosteen (*Garcinia mangostana*). Food. Chem. Toxicol. 46(10):3227-3239.

Peres V, Nagem TJ and De Oliveira FF. 2000. Tetraoxygenated naturally occurring xanthenes. Phytochemistry, 55:683-710.

Piller NB. 1975. "A comparison of the effectiveness of some anti-inflammatory drugs on thermal oedema," Br. J. Exp. Pathol. 56(6): 554-560.

Rai LK, Pankaj, Prasad and Sharma E. 2000. Conservation threats to some important medicinal plants of the Sikkim Himalaya. Biol. Conservat, 93:27-33.

Ramage CM, Sando L, Peace CP, Carroll BJ and Drew RA. 2004. Genetic diversity in the apomictic fruit species *Garcinia mangostana* L. (mangosteen). Euphytica. 136: 1-10.

Ricciardolo FLM, Sterk PJ, Gaston B and Folkerts G. 2004. Nitric oxide in health and disease of the respiratory system. Physiol Rev. 84:731-765.

Rocha L, Marston A, Auxiliadora M, Kaplan C, Stoeckli-Evans H, Thull U, Testa B and Hostettmann K. 1994. An antifungal γ -pyrone and xanthenes with monoamine oxidase inhibitory activity from *Hypericum brasiliense*. Phytochemistry, 36:1381-138

Rouger C, Derbre S, Charreau B, Pabois A, Cauchy T, Litaudon M, Awang K and Richomme P. 2015. Lepidotol A from *Mesua lepidota* inhibits inflammatory and immune mediators in human endothelial cells. J. Nat. Prod. 78:2187-2197.

Rukayadi Y, Yong D and Hwang J-K. 2006. In vitro anticandidal activity of xanthorrhizol isolated from *Curcuma xanthorrhiza* Roxb. J. Antimicrob. Chemother. 57: 1231-1234

See I, Ee GCL, Teh SS, Kadir AA and Daud S. 2014. Two new chemical constituents from the stem bark of *Garcinia mangostana*. Molecules, 19:7308-7316.

Shivamurthy VM, Gantt S, Reilly C, Tilley P, Guzman J and Tucker L. 2016. *Bacillus pumilus* septic arthritis in a healthy child, Can. J. Infect. Dis. Med. Microbiol. 2016(Article ID 3265037):1-3. DOI:10.1155/2016/3265037

Suksamrarn S, Suwannapoch N, Ratananukul P, Aroonlerk N and Suksamrarn A. 2002. Xanthenes from the green fruit hulls of *Garcinia mangostana*. J. Nat. Prod. 65:761-763.

Suksamrarn S, Suwannapoch N, Phakhodee W, Thanuhiranlert J, Ratananukul P, Chimnoi N and Suksamrarn A. 2003. Antimycobacterial activity of prenylated xanthenes from the fruits of *Garcinia mangostana*. Chem. Pharm. Bull. 51: 857-859.

- Suksamrarn S, Komutiban O, Ratananukul P, Chinnoi N, Lartpornmatulee N and Suksamrarn A. 2006. Cytotoxic prenylated xanthenes from the young fruit of *Garcinia mangostana*. Chem. Pharm. Bull. 54: 301-305.
- Sultanbawa MUS. "Xanthonoids of tropical plants". 1980. Tetrahedron, 36(11):1465–1506.
- Syam S, Bustamam A, Abdullah R, Sukari MA, Hashim NM, Yahayu MA, Hassandarvish P, Mohan S and Abdelwahab SI. 2014. Cytotoxicity and oral acute toxicity studies of β -mangostin isolated from *Cratoxylum arborescens*. Phcog. J, 6(1):47–56.
- Szakiel A, Paczkowski C, Pensec F and Bertsch C. 2012. Fruit cuticular waxes as a source of biologically active triterpenoids. Phytochem Rev. 11(2-3):263-284.
- Tan WS, Palanisamy A, Govindarajan K and Fakurazi S. 2015. *Moringa oleifera* flower extract suppresses the activation of inflammatory mediators in lipopolysaccharide-stimulated RAW 264.7 macrophages via *NF-KB* pathway, Mediators Inflamm. 2015(Article ID 720171): 1-11. DOI:10.1155/2015/720171.
- Tanjung M, Saputri RD, Fitriati FF and Tjahjandarie TS. 2016. Antimalarial and antioxidant activities of isoprenylated coumarins from the stem bark of *Mesua borneensis* L. J. Biol. Act. Pro. Nat. 6(2):95-100.
- Teh SS, Ee GCL, Rahmani M, Sim WC, Mah SH and Teo SH. 2010. Two new pyranoxanthenes from *Mesua beccariana*. Molecules. 15:6733-6742.
- Teh SS, Ee GCL, Rahmani M, Yap YHT, Go R and Mah SH. 2011. Pyranoxanthenes from *Mesua ferrea*. Molecules, 16:5647-5654.
- Teh, SS, Ee GCL, Mah SH, Lim YM. 2013. Cytotoxic and structure-activity relationships of xanthone derivatives from *Mesua beccariana*, *Mesua ferrea*, and *Mesua congestiflora* towards nine cancer cell lines. Molecules, 18:1985-1994.
- Teh, SS, Ee GCL, Mah SH, Ahmad Z. 2016. Structure–activity relationship study of secondary metabolites from *Mesua beccariana*, *Mesua ferrea* and *Mesua congestiflora* for anti-cholinesterase activity. DOI: 10.1007/s00044-016-1531-0.
- Tena D, Torres JAM, Pomata MTP, Nieto JAS, Rubio V and Bisquert J. 2007. Cutaneous infection due to *Bacillus pumilus*: report of 3 cases. Clin. Infect. Dis. 44: 40-42.
- Venugopala KN, Rashmi V and Odhav B. 2013. Review on natural coumarin lead compounds for their pharmacological activity. 2013(Article ID 963248):1-14. DOI: 10.1155/2013/963248.

- Verotta L, Lovaglio E, Vidari G, Finzi PV, Neri MG, Raimondi A, Parapini S, Taramelli D, Riva A and Bombardelli E. 2004. 4-Alkyl- and 4-phenylcoumarins from *Mesua ferrea* as promising multidrug resistant antibacterials. *Phytochemistry*. 65(21):2867-2879.
- Vieira LMM and Kijjoa A. 2005. Naturally-occurring xanthenes: recent developments. *Curr Med Chem*. 12: 2413–2446.
- Walia S and Mukerjee SK. 1984. Ferrixanthone, A 1,3,5,6 - tetraoxygenated xanthone from *Mesua Ferrea*. *Phytochemistry*. 23(8):1816-1817.
- Wieble JEKC and Downton WJS. 1992. Mangosteen (*Garcinia mangostana* L.) – A potential crop for Northern Australia. *Acta Hort*. 321:132–137.
- Witaicenis A, Seito LN and Di-Stasi LC. 2010. “Intestinal anti-inflammatory activity of esculetin and 4-methylesculetin in the trinitrobenzenesulphonic acid model of rat colitis”. *Chem. Biol. Interact*. 186(2): 211–218.
- Walton NJ, Mayer MJ and Narbad A. 2003. Molecules of Interest: Vanillin. *Phytochemistry*, 63: 505-515
- Xu Z, Huang L, Chen X-H, Zhu X-F, Qian X-J, Feng GK, Lan WJ and Li H-J. 2014. Cytotoxic Prenylated Xanthenes from the Pericarps of *Garcinia mangostana*. *Molecules*. 19(2):1820-1827.
- Yahayu MA, Rahmani M, Hashim NM, Ee GCL, Sukari MA & Akim AM. 2013. Cytotoxic and anti-microbial xanthenes from *Cratoxylum arborescens* (Guttiferae). *Malay. J. Sci*. 32(1):53-60.
- Yousefi S, Bayat S, Rahman MBM, Ismail IS, Saki E and Abdulmalek E. 2015. Synthesis and in vitro bioactivity evaluation of new glucose and xylitol ester derivatives of 5-aminosalicylic acid, *RSC adv*. 5:97295-97370.
- Yu L, Zhao M, Yang B, Zhao Q and Jiang Y. 2007. Phenolics from hull of *Garcinia mangostana* fruit and their antioxidant activities. *Food. Chem*. 104: 176-181.
- Zhao Y, Liu JP, Lu D, Li PY and Zhang LX. 2010. A new antioxidant xanthone from the pericarp of *Garcinia mangostana* Linn. *Nat. Prod. Res*. 24:1664-1670.
- Zhou X, He L, Wu X, Zhong Y, Zhang Y, Wang Y, Wang B, Xu Z & Qiu Z. 2014. Two new xanthenes from the pericarp of *Garcinia mangostana*. *Nat. Prod. Res*. 29(1): 19-23.