



UNIVERSITI PUTRA MALAYSIA

***BIODEGRADATION OF PHENANTHRENE AND PYRENE USING
BACTERIA ISOLATED FROM USED VEHICLE
LUBRICANTCONTAMINATED
SOIL***

UMAR ZUBAIRU DARMA

FS 2017 79



**BIODEGRADATION OF PHENANTHRENE AND PYRENE USING
BACTERIA ISOLATED FROM USED VEHICLE LUBRICANT-
CONTAMINATED SOIL**

By

UMAR ZUBAIRU DARMA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillments of the Requirements for the Degree of Doctor of Philosophy**

July 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**BIODEGRADATION OF PHENANTHRENE AND PYRENE USING
BACTERIA ISOLATED FROM USED VEHICLE LUBRICANT-
CONTAMINATED SOIL**

By

UMAR ZUBAIRU DARMA

July 2017

Chairman : Associate Professor Muskhazli Mustafa, PhD
Faculty : Science

Petroleum industry contributes significantly towards global industrial civilization. Used vehicle lubricant oil being a petroleum product portrays dangerous impact on the environment as it generates hazardous polycyclic aromatic hydrocarbons (PAHs) through incomplete combustion. The rampant discharge of this oil in Malaysia causes serious environmental concern as PAHs are dangerous organic pollutants that cause mutation and cancer to living cells. The most commonly hazardous PAHs from used lubricating oil were phenanthrene and pyrene belonging to the low and high molecular weights PAHs. Biological removal of phenanthrene and pyrene provides an environmentally sound and state-of-the-art technique that employs natural biological processes for the complete elimination of such pollutants from the environment. However, slow degradation duration is a major PAHs biodegradation limitation. Therefore, the research objectives involved isolation of effective phenanthrene and pyrene degrading bacteria, optimizing the biodegradation conditions, analysing the biodegradation intermediates and evaluating the effects of heavy metals on the biodegradation effectiveness.

Phenanthrene and pyrene degrading bacteria were initially isolated from used lubricating oil contaminated soil using enrichment and plating techniques where both PAHs served as the sole bacteria carbon and energy sources. The effective phenanthrene and pyrene degrading bacteria were screened based on spray plate technique and colorimetric assay followed by 16S rRNA gene identification using universal primers. The biodegradation conditions of such identified bacteria were then optimized using single factor optimization before subjected to response surface methodology based on full factorial central composite design ($p \leq 0.05$). This was preceded with the analyses of phenanthrene and pyrene degrading metabolites using gas chromatography mass spectrophotometer (GC-MS). The identified bacteria were finally used for the formulation of bacteria consortium after the compatibility testing

based on cross spread technique and permutation assessments through colorimetric assay. The consortium was then used to degrade 500 mg/L phenanthrene and 250 mg/L pyrene in a complex culture that contained varying concentrations (2 mg/L to 12 mg/L) of Nickel, Lead, Vanadium and Cadmium separately.

Total of 93 different strains of bacteria were isolated from the enrichment technique. Among these bacteria, only 53 strains initiated the degradation of 5 g/L phenanthrene in 72 hours based on spray plate assessments ($n = 3$). Further screening by colorimetric assay indicated only two strains named MM045 and MM087 were able to degrade 75.2% and 80.2% of 500 mg/L phenanthrene in addition to 54.3% and 59.7% of 250 mg/L pyrene within 24 hours respectively ($n = 3$). These strains were then identified as *Cronobacter sakazakii* MM045 and *Enterobacter* sp. MM087 with accession numbers KT933253 and KT933254 respectively. The phenanthrene and pyrene degradation capabilities of *C. sakazakii* MM045 and *Enterobacter* sp. MM087 was statistically optimized through response surface methodology ($p \leq 0.05$). This optimization showed the combined efforts of the independent variables from each of *C. sakazakii* MM045 and *Enterobacter* sp. MM087 culture resulted in 100% degradations of both phenanthrene (500 mg/L) and pyrene (250 mg/L) in 24 hours. These were validated experimentally using the numerical optimization analyses ($n = 3$). The phenanthrene and pyrene biodegradation intermediates were then identified using the GC-MS analyses ($n = 3$). The pyrene identified metabolites from the *C. sakazakii* MM045 and *Enterobacter* sp. MM087 degradation cultures include pyrene *cis*-4,5-dihydrodiol, 3,4-dihydroxyphenanthrene, phthalic acid, pyruvic acid, acetic acid, lactic acid and formic acid. Additionally, phenanthrene identified metabolites from both cultures were 3,4-dihydroxyphenanthrene, phthalic acid, pyruvic acid, acetic acid and oxalic acid. These metabolites established the degradation pathways undergone by both bacteria. The bacteria were also found to tolerate more than 6 mg/L of Nickel, Cadmium, Lead and Vanadium which significantly exceeded the hazardous metals concentrations in the natural habitats were both bacteria survived prior to the isolation.

Considering the effective phenanthrene and pyrene degradation responses of *C. sakazakii* MM045 and that of *Enterobacter* sp. MM087 in 24 hours, both isolates can be used for commercial applications of degrading PAHs from contaminated environments. This commercial application could be possible as all the optimized independent variables are achievable within natural environments such as soil and water bodies. Furthermore, environmental availability of the *C. sakazakii* and *Enterobacter* sp. portrays another desirable character that made their choice as the best degradation alternative. Additionally, both bacteria can maintain their degradation effectiveness even in a complex environment that contained concurrent contamination of PAHs and heavy metals. Therefore, *C. sakazakii* MM045 and *Enterobacter* sp. MM087 can effectively and rapidly degrade phenanthrene and pyrene from used vehicle lubricant contaminated environment.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**BIODEGRADASI PHINANTREN DAN PIRIN MENGGUNAKAN
BAKTERIA TERASING DARIPADA MINYAK LUBRIKAN KENDERAAN
TERKONTAMINASI**

Oleh

UMAR ZUBAIRU DARMA

Julai 2017

Pengerusi : Profesor Madya Muskhazli Mustafa, PhD
Fakulti : Sains

Industri petroleum menyumbang secara signifikan ke arah tamadun perindustrian global. Minyak pelincir kenderaan yang digunakan sebagai produk petroleum memberikan gambaran kesan berbahaya terhadap alam sekitar kerana ia menghasilkan hidrokarbon aromatik polisiklik (PAHs) yang berbahaya melalui pembakaran tak lengkap. Pelepasan minyak ini secara meluas di Malaysia menyebabkan kebimbangan yang serius terhadap alam sekitar kerana PAHs adalah bahan pencemar organik berbahaya yang mampu menyebabkan mutasi dan kanser terhadap sel hidup. Jenis PAH yang paling kerap diperolehi dari minyak pelincir yang digunakan adalah phenantren dan pirin dari kumpulan PAHs dengan berat molekul yang tinggi dan rendah. Penyingkiran phenantren dan pirin secara biologi adalah teknik yang mesra alam dan terkini menggunakan proses biologi semula jadi untuk menghapuskan bahan pencemar sedemikian secara menyeluruh dari alam sekitar. Walau bagaimanapun, tempoh biodegradasi yang perlahan adalah pengehadan biodegradasi PAH yang utama. Oleh itu, objektif penyelidikan ini melibatkan pemisahan bakteria efektif yang merendahkan phenantren dan pirin, mengoptimumkan keadaan biodegradasi, menganalisis perantaraan biodegradasi dan menilai kesan logam berat terhadap keberkesanan biodegradasi.

Bakteria phenantren dan pirin yang terdegradasi pada mulanya diasingkan dari tanah yang tercemar oleh minyak pelincir yang digunakan menggunakan teknik pengayaan dan penyaduran di mana kedua-dua PAH berfungsi sebagai karbon tunggal dan sumber tenaga bakteria. Bakteria phenantren dan pirin yang merosakkan bakteria ditapis menggunakan teknik plat semburan dan cerakin kolorimeter diikuti pengenalan gen rRNA 16S menggunakan primer universal. Keadaan biodegradasi bakteria yang dikenal pasti kemudian dioptimumkan menggunakan pengoptimuman faktor tunggal sebelum diletakkan pada metodologi respon permukaan berdasarkan reka bentuk komposit pusat faktorial penuh ($p \leq 0.05$). Ini didahului dengan analisis metabolit yang

merendahkan phenantren dan pirin menggunakan spektrofotometer jisim kromatografi gas (GC-MS). Bakteria yang dikenal pasti akhirnya digunakan untuk perumusan konsortium bakteria selepas ujian keserasian berdasarkan teknik rebakan silang dan penilaian permutasi melalui ujian warna. Konsortium kemudiannya digunakan untuk menurunkan phenantrena 500 mg/L dan 250 mg/L pirin dalam kultur kompleks yang mengandungi Nikel, Plumbum, Vanadium dan Kadmium dengan kepekatan yang berbeza-beza (2 mg/L hingga 12 mg/L) secara berasingan.

Sejumlah 93 jenis bakteria yang berbeza telah diasingkan menggunakan teknik pengayaan. Antara bakteria ini, hanya 53 strain sahaja memulakan degradasi 5 g/L phenantrena dalam 72 jam berdasarkan penilaian plat semburan ($n = 3$). Penyaringan selanjutnya oleh cerakin kolorimeter menunjukkan hanya dua strain iaitu MM045 dan MM087 yang mampu menurunkan 75.2% dan 80.2% daripada 500 mg/L phenantrena serta 54.3% dan 59.7% daripada 250 mg/L pirin dalam tempoh 24 jam ($n = 3$). Strain ini kemudian dikenal pasti sebagai *Cronobacter sakazakii* MM045 dan *Enterobacter sp.* MM087 dengan nombor kemasukan KT933253 dan KT933254 masing-masing. Keupayaan degradasi phenantren dan piren oleh *C. sakazakii* MM045 dan *Enterobacter sp.* MM087 dioptimumkan secara statistik melalui metodologi permukaan tindak balas ($p \leq 0.05$). Pengoptimuman ini menunjukkan usaha gabungan pembolehkan bebas dari kultur *C. sakazakii* MM045 dan *Enterobacter sp.* MM087 dalam mendegradasi 100% phenantrena (500 mg/L) dan pirin (250 mg/L) dalam masa 24 jam. Ini telah disahkan secara eksperimen menggunakan analisis pengoptimuman berangka ($n = 3$). Perantara biodegradasi phenantren dan pirin kemudiannya dikenal pasti menggunakan analisis GC-MS ($n = 3$). Metabolit pirin yang dikenal pasti dari kultur degradasi *C. sakazakii* MM045 dan *Enterobacter sp.* MM087 termasuk pirin *cis*-4,5-dihydrodiol, 3,4-dihydroxyphenantren, asid ftalik, asid piruvik, asid asetik, asid laktik dan asid formik. Selain itu, metabolit phenantren yang dikenal pasti dari kedua-dua kultur adalah 3,4-dihydroxyphenantren, asid ftalik, asid piruvik, asid asetik dan asid osalik. Metabolit ini menubuhkan laluan degradasi yang dialami oleh kedua-dua bakteria. Bakteria juga didapati menoleransi lebih daripada 6 mg/L Nikel, Kadmium, Plumbum dan Vanadium yang melebihi kepekatan logam berbahaya di habitat semulajadi. Kedua-dua bakteria terselamat sebelum pengasingan.

Disebabkan keberkesanan tindak balas degradasi phenantren dan pirin oleh *C. sakazakii* MM045 dan *Enterobacter sp.* MM087 dalam masa 24 jam, kedua-dua pencilan boleh digunakan untuk aplikasi komersil dalam merendahkan PAH daripada persekitaran yang tercemar. Aplikasi komersil ini boleh dilakukan memandangkan semua pemboleh ubah bebas yang optimum dapat dicapai dalam persekitaran semula jadi seperti tanah dan hidupan air. Tambahan pula, ketersediaan alam sekitar *C. sakazakii* dan *Enterobacter sp.* menggambarkan ciri terpilih yang menjadikan mereka sebagai pilihan degradasi alternatif yang terbaik. Selain itu, kedua-dua bakteria boleh mengekalkan keberkesananannya walaupun dalam persekitaran yang kompleks yang mengandungi pencemaran serentak PAH dan logam berat. Oleh itu, *C. sakazakii* MM045 dan *Enterobacter sp.* MM087 dengan berkesan dan cepat boleh menghancurkan phenantren dan pirin daripada persekitaran tercemar oleh pelincir kenderaan yang digunakan.

ACKNOWLEDGEMENTS

All thanks are due to our Almighty Creator for the opportunity, life, guidance and protection to have witness this historical monument in my life. May He continue to guide and protect us in all our entire life.

Special gratitude to my late parents Alh Umar Darma and Hj Lubabatu Ibrahim, who coaches my life during their life time, may the Almighty Creator reward them with Aljannatul Firdaus. Life without you is indeed different and you have the credits of every single achievement in my life because I owe it all to you.

I thank the chairman of my supervisory committee Associate Professor Dr. Muskhazli Mustafa for your guidance, support and advice which helped me towards achieving better quality research. Indeed you are the best academician I have ever met who changes my academic thinking to better. I am also grateful to other committee members Associate Professor Dr. Nor Azwady Abd Aziz and Dr. Syaizwan Zahmir Zulkifli for your support, advice and patience towards overcoming several obstacles in the course of actions. May our Creator reward you all with Aljannatul Firdaus.

To my wife, Maryam Labaran Darma: indeed you are the best wife as you shared all my challenges, I thank you for your courageous advice, moral and emotional support; may the Almighty Creator reward you with Aljannatul Firdaus. To my only daughter Aminatu Zubairu Umar, I was exceptionally happy when you were born and I thanked my Creator for such special gift as you are a party to my PhD struggle. She is always keen to know what I was doing even though she never grasped what it was all about! I am always happy with your screams of joy. May our Almighty Creator guide your life, strengthen you, bless you and make you among the best Muslimah in the world.

I am also grateful to my family members, especially Abubakar Umar Darma (Bala) who supported my entire struggle financially and emotionally. Whenever significant momentous is achieved, he openly shows his happiness and kindness, thank you and may Aljannatul Firdaus be your reward.

My immense gratitude to the management of Umaru Musa Yar'adua University, Katsina, Nigeria for sponsoring my entire PhD programme financially: by the grace of our Creator I shall contribute my own quota towards uplifting the University to a greater height. To the Universiti Putra Malaysia, thank you for the Putra Grant (GP-IPS/2016/9489800) which helped immensely towards minimizing our financial stress during the study.

I thank my colleagues in our laboratory for your cooperation, feedback and of course friendship. Additionally, I would like to express my gratitude to the staff of ICP laboratory, faculty of Agriculture, UPM and GC-MS laboratory, faculty of Food Sciences, UPM for the analyses. Finally, all my friends at UPM were indeed kinds and courageous, thank you all.



I certify that a Thesis Examination Committee has met on 17 July 2017 to conduct the final examination of Umar Zubairu Darma on his thesis entitled "Biodegradation of Phenanthrene and Pyrene Using Bacteria Isolated from Used Vehicle Lubricant-Contaminated Soil" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Hishamuddin bin Omar, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Rosimah binti Nulit, PhD


Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Normala binti Halimoon, PhD

Senior Lecturer
Faculty of Environmental Studies
Universiti Putra Malaysia
(Internal Examiner)

Imran Hashmi, PhD

Professor
National University of Sciences and Technology
Pakistan
(External Examiner)



NOR AINI AB. SHUKOR, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 4 September 2017

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Muskhazli Mustafa, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Nor Azwady Abd Aziz, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Syaizwan Zahmir Zulkifli, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Umar Zubairu Darma, GS40787

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Associate Professor Muskhazli Mustafa

Signature: _____
Name of Member
of Supervisory
Committee: Associate Professor Dr. Nor Azwady Abd Aziz

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Syaizwan Zahmir Zulkifli

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xviii
CHAPTER	
1 INTRODUCTION	1
1.1 Background of the study	1
1.2 Problem statement	2
1.3 Significant of the study	3
1.4 Objective of the study	3
2 LITERATURE REVIEW	4
2.1 Used vehicle lubricant	4
2.2 Sources and toxicity of PAHs	5
2.2.1 Phenanthrene and pyrene	7
2.3 Remediation strategies of PAHs	8
2.3.1 Biodegradation of phenanthrene and pyrene	9
2.3.2 PAHs degrading bacteria and mechanisms of action	10
2.3.3 Phenanthrene and pyrene biodegradation metabolite	11
2.4 Factors influencing PAHs biodegradation	14
2.4.1 Temperature	14
2.4.2 Oxygenation	14
2.4.3 Hydrogen ion concentration (pH)	15
2.4.4 Salinity	15
2.4.5 Inoculums volume	15
2.5 Ways of improving PAHs biodegradation	16
2.5.1 Adaptation of PAHs degrading characters	16
2.5.2 Biodegradation consortium	16
2.5.3 Optimization of PAHs biodegradation conditions	17
2.6 Effects of hazardous heavy metals on PAHs biodegradation	18
2.7 PAHs biodegradation indicator	18
3 ISOLATION AND IDENTIFICATION OF PHENANTHRENE AND PYRENE DEGRADING BACTERIA	19
3.1 Introduction	19

3.2	Materials and methods	20
3.2.1	Sampling area	20
3.2.2	Soil sampling	23
3.2.3	Chemicals and Media	23
3.2.4	Preparation of solutions and reagents	23
3.2.4.1	Preparation of PAHs stock solutions	23
3.2.4.2	Preparation of DCPIP solution	23
3.2.4.3	Preparation of glycerol stock	24
3.2.4.4	Preparation of phosphate buffer saline	24
3.2.5	Culture media preparation	24
3.2.6	Isolation of PAHs degrading bacteria	24
3.2.6.1	Bacteria preservation	25
3.2.6.2	Bacteria resting cells preparation	25
3.2.7	Screening of effective phenanthrene and pyrene degrading bacteria	25
3.2.7.1	Spray plate assay	25
3.2.7.2	Colorimetric assay	26
3.2.8	Standard calibration curves	27
3.2.9	Bacteria identification	28
3.2.9.1	Morphology identification	28
3.2.9.2	16S rRNA gene sequencing	28
3.2.10	Data analyses	29
3.3	Results	29
3.3.1	Bacteria isolation and screening	29
3.3.2	Quantification of PAHs degradation	32
3.3.3	Bacteria identification	33
3.4	Discussion	36
3.5	Conclusion	40
4	OPTIMIZATION OF PHENANTHRENE AND PYRENE BIO-DEGRADATION CONDITIONS	41
4.1	Introduction	41
4.2	Materials and methods	42
4.2.1	Bacteria culture conditions	42
4.2.2	Single factor optimization (SFO)	42
4.2.3	Response surface methodology	43
4.2.4	Data analyses	45
4.3	Results	45
4.3.1	Single factor optimization	45
4.3.1.1	Effects of agitation	45
4.3.1.2	Effects of temperature	46
4.3.1.3	Effects of pH	47
4.3.1.4	Effects of inoculums volume	47
4.3.1.5	Effects of salinity	48
4.3.2	Response surface methodology	49
4.3.2.1	<i>Cronobacter sakazakii</i> MM045	49
4.3.2.2	<i>Enterobacter</i> sp. MM087	58
4.4	Discussions	63
4.5	Conclusion	70

5	IDENTIFICATION OF PHENANTHRENE AND PYRENE BIODEGRADATION METABOLITES	71
5.1	Introduction	71
5.2	Materials and methods	72
5.2.1	Bacteria growth conditions	72
5.2.2	Metabolites extraction	73
5.2.3	Metabolites purification	73
5.2.4	Metabolites identification	73
5.3	Results	74
5.3.1	Identification of pyrene metabolites	74
5.3.2	Identification of phenanthrene metabolites	77
5.3.3	Phenanthrene and pyrene biodegradation pathways	79
5.4	Discussions	82
5.5	Conclusion	84
6	IDENTIFICATION OF HEAVY METALS CO-CONTAMINANTS ON PAHs DEGRADING CONSORTIUM	85
6.1	Introduction	85
6.2	Materials and methods	86
6.2.1	Chemicals and reagents	86
6.2.2	Inoculums preparation	86
6.2.3	Formulation of PAHs degrading consortium	86
6.2.4	Analyses of heavy metals	87
6.2.5	Effects of selected metals on PAHs biodegradation	88
6.2.6	Data analyses	89
6.3	Results	89
6.3.1	Effective phenanthrene and pyrene degradation consortium	89
6.3.2	Toxic metals selection	90
6.3.3	Effects of selected metals on PAHs biodegradation	92
6.4	Discussions	95
6.5	Conclusion	97
7	CONCLUSION AND FUTURE RECOMMENDATIONS	98
7.1	Concluding Remarks	98
7.2	Future Recommendations	99
	REFERENCES	101
	APPENDICES	138
	BIODATA OF STUDENT	147
	LIST OF PUBLICATIONS	148

LIST OF TABLES

Table		Page
2.1	Comparisons between the physical, chemical and biological removal means of PAHs from contaminated environments	9
3.1	Soil sampling locations within Selangor, Malaysia	21
3.2	Bacteria screening for PAHs degradation by colorimetric assay (12 and 24h)	31
4.1	Coded and un-coded variables used for RSM optimisations of PAHs biodegradations by <i>C. sakazakii</i> MM045 and <i>Enterobacter</i> sp. MM087	44
4.2	RSM experimental settings of PAHs degradation responses by <i>C. sakazakii</i> MM045 and <i>Enterobacter</i> sp. MM087 in 24 hrs	50
4.3	Statistical summary of phenanthrene degradation by <i>C. sakazakii</i> MM045	53
4.4	Statistics summary of pyrene degradation by <i>C. sakazakii</i> MM045	54
4.5	Optimum phenanthrene and pyrene degradation response by <i>C. sakazakii</i> MM045.	57
4.6	Statistical summary of phenanthrene degradation by <i>Enterobacter</i> sp. MM087.	59
4.7	Statistical summary of pyrene degradation by <i>Enterobacter</i> sp. MM087.	60
4.8	Optimum phenanthrene and pyrene degradation values by <i>Enterobacter</i> sp. MM087.	63
6.1	PAHs degradation response from different consortium compositions made of <i>C. sakazakii</i> MM045 & <i>Enterobacter</i> sp. MM087.	89
6.2	ICP-OES analyses of MM045, MM087 and SRM-2587 soil samples.	91
6.3	Phenanthrene and pyrene degradation response of an effective consortium in different heavy metals concentrations	93

LIST OF FIGURES

Figure		Page
2.1	Sources, route of entrance and toxicity effects of environmental PAHs into the human body	7
2.2	Structural arrangements involving three and four aromatic compounds representing (a) phenanthrene and (b) pyrene	8
2.3	Phenanthrene degradation pathway of bacteria indicating metabolites involving (a) <i>cis</i> -3,4-dihydroxy-3,4-dihydrophenanthrene (b) 3,4-dihydroxyphenanthrene (c) 2-Hydroxy-2 <i>H</i> -benzo[<i>h</i>]chromene-2-carboxylate (d) <i>trans</i> -4-(1'-Hydroxynaphth-2'-yl)-2-oxobut-3-enoic acid (e) 1-Hydroxy-2-naphthaldehyde (f) 1-Hydroxy-2-naphthoate (g) <i>trans</i> -2'-Carboxybenzalpyruvate (h) phthalic acid	12
2.4	Pyrene degradation pathway undergone by bacteria where the specific metabolites involved (a) pyrene- <i>cis</i> -4,5-dihydrodiol (b) 4,5-Dihydroxypyrene (c) Phenanthrene-4,5-dicarboxylate (d) Phenanthrene-4-carboxylate (e) <i>cis</i> -3,4-Phenanthrenedihydrodiol-4-carboxylate (f) 3,4-Dihydroxyphenanthrene	13
3.1	Map of Peninsular Malaysia indicating the selected sampling locations used for the study.	22
3.2	Colorimetric assay indicating gradual colour changes from the PAHs degradation culture where (a) culture without DCPIP; (b) culture with DCPIP; (c) low degradation response (+); (d) moderate response (++); and (e) complete response (+++).	27
3.3	Spray plate screening indicating treatment plates (a) before and (b) after degradation based on phenanthrene crystals disappearance.	30
3.4	Phenanthrene and pyrene degradation responses by strain MM045 within 24 hours of incubation.	32
3.5	Phenanthrene and pyrene degradation response by strain MM087 within 24 hours of incubation.	33
3.6	Morphological appearance of PAHs degrading bacteria on LB agar plate indicating strains (a) MM045 and (b) MM087.	34
3.7	Gram reactions of PAHs degrading bacteria for strain (a) MM045 and (b) MM087	34

3.8	Agarose gel electrophoresis of the amplified 16S rRNA genes for strains MM045 and MM087	35
3.9	Neighbor Joining Tree indicating phylogenetic relationship among strain MM045 and MM087 with other closely related bacteria.	36
4.1	Effects of agitation speed on phenanthrene and pyrene degradations by (a) <i>C. sakazakii</i> MM045 and (b) <i>Enterobacter</i> sp. MM087	46
4.2	Effects of temperature phenanthrene and pyrene degradation by (a) <i>C. sakazakii</i> MM045 and (b) <i>Enterobacter</i> sp. MM087	46
4.3	Effects of pH on phenanthrene and pyrene degradation by (a) <i>C. sakazakii</i> MM045 and (b) <i>Enterobacter</i> sp. MM087.	47
4.4	Effects of inoculums volume on phenanthrene and pyrene degradation by (a) <i>C. sakazakii</i> MM045 and (b) <i>Enterobacter</i> sp. MM087.	48
4.5	Effects of salinity on phenanthrene and pyrene degradation by (a) <i>C. sakazakii</i> MM045 and (b) <i>Enterobacter</i> sp. MM087.	48
4.6	Phenanthrene degradation response by <i>C. sakazakii</i> MM045 showing interactions among (a) IV and rpm, (b) salinity and rpm, (c) pH and temperature, (d) IV and temperature, (e) salinity and temperature, (f) IV and pH, (g) salinity and pH, (h) salinity and IV.	55
4.7	Pyrene degradation response from <i>C. sakazakii</i> MM045 indicating interactions of (a) temperature and rpm, (b) pH and rpm, (c) salinity and rpm, (d) IV and temperature, (e) IV and pH, (f) salinity and IV.	56
4.8	Phenanthrene degradation response from <i>Enterobacter</i> sp. MM087 indicating interactions among (a) pH and rpm, (b) IV and rpm, (c) salinity and rpm, (d) pH and temperature, (e) IV and temperature, (f) salinity and temperature and (g) salinity and IV.	61
4.9	Pyrene degradation response from <i>Enterobacter</i> sp. MM087 indicating interactions among (a) temperature and rpm, (b) pH and temperature, (c) IV and temperature and (d) salinity and IV.	62
5.1	Mass spectra of pyrene intermediates where <i>C. sakazakii</i> MM045 generated (a) pyrene <i>cis</i> -4,5-dihydrodiol (b) 3,4-dihydroxyphenanthrene (c) phthalic acid (d) pyruvic acid (e) lactic acid (f) acetic acid (g) formic acid. Also <i>Enterobacter</i> sp. MM087 produced (a), (b), (c), (d), (f), and (g) intermediates	76

- 5.2 Mass spectra of phenanthrene intermediates where *C. sakazakii* MM045 produced (a) 3,4-dihydroxyphenathrene (b) phthalic acid (c) pyruvic acid (d) Oxalic acid and (e) acetic acid. The *Enterobacter* sp. MM087 produced intermediates were (a), (b), (c), and (e) 79
- 5.3 Phenanthrene and pyrene degradation pathway undergone by *C. sakazakii* MM045 and *Enterobacter* sp. MM087 indicating serial structural reduction of both PAHs 81
- 6.1 Mixed culture plate indicating the compatibility of *C. sakazakii* MM045 which appeared small milky and *Enterobacter* sp. MM087 being big white colonies. 90



LIST OF ABBREVIATIONS

ANOVA	Analysis of variance
As	Arsenic
ATSDR	Agency for Toxic Substances and Disease Registry
BHA	Bushnell Haas Agar
bp	Base pairs
CC	Column Chromatography
CCD	Central Composite Design
Cd	Cadmium
CFU	Colony forming unit
cm	Centimetre
Co	Cobalt
Cr	Chromium
CRMs	Certified reference materials
Cu	Copper
CV	Coefficient of variation
°C	Degree Celsius
DF	Degrees of freedom
DCPIP	Dichlorophenolindophenol
DNA	Deoxyribonucleic acid
dNTPs	Deoxyribonucleotides
EU	European Union
F-primer	Forward primer
FRLI	Federal Register of Legislative Instruments
g	Gram
GC-MS	Gas Chromatography- Mass Spectrometry

GDP	Gross domestic products
h	Hour
HMW	High Molecular Weight
HPLC	High performance liquid chromatography
HSD	Honest significant difference
ICP-OES	Inductively coupled plasma optical emission spectrophotometer
ISO	International standard organisation
IV	Inoculums volume
kDa	Kilo Dalton (unified atomic mass unit)
kg	Kilogram
L	Litre
LB	Luria-Bertani
LMW	Low Molecular Weight
LOQ	Limit of quantification
m Ω	Milliohm
mm ³	Millimetre cube
MgCl ₂	Magnesium Chloride
MEGA	Molecular Evolutionary Genetics Analysis
MEQR	Malaysian environmental quality report
mg	Milligram
min	Minute
mL	Millilitre
Mn	Manganese
MS	Mean square
MSM	Mineral Salts Medium
MT	Metric tonne

m/z	Mass per charge
NaCl	Sodium chloride
NADP	Nicotinamide adenine dinucleotide phosphate
NCBI	National Centre for Biotechnology Information
Ni	Nickel
nm	Nanometer
NIST	National institute of standards and technology
OD ₆₀₀	Optical Density at 600nm wavelength
PAHs	Polycyclic Aromatic Hydrocarbons
Pb	Lead
PBS	Phosphate buffer saline
PCR	Polymerase Chain Reaction
pH	Hydrogen ion concentration
PHN	Phenanthrene
PYR	Pyrene
R ²	Coefficient of determination
rpm	Revolution per minute
R-primer	Reverse primer
rRNA	Ribosomal ribonucleic acid
RSM	Response Surface Methodology
sec	Second
SFO	Single factor optimization
SRM	Standard reference material
SS	Sum of the square
TAE	Tri-acetic acid-Ethylenediaminetetraacetic acid
TRI	Texas Research Institute

U	Unit
US EPA	United States Environmental Protection Agency
US FDA	US Food and drugs administration
V	Vanadium
v/v	Volume per volume
WHO	World Health Organization.
wt	Weight
w/v	Weight per volume
Zn	Zinc
α	Alpha
β	Beta
3D	Three dimension
%	Percent
μL	Microlitre

CHAPTER 1

INTRODUCTION

1.1 Background of the study

Polycyclic aromatic hydrocarbons (PAHs) comprised large group of hazardous chemicals that resulted from incomplete combustion of organic products (Zhao *et al.*, 2008). Naturally, PAHs are sourced from forest fires, oil seepage and volcanoes while anthropogenic sources involved incineration, internal engine combustion, tobacco cigarette, carbon black, coal tar and asphalt (Feng *et al.*, 2012). These sources directly deposit toxic PAHs into the environment which persist for long duration due to chemical stability and low water solubility (Maliszewska, 1999). Such persistence has tremendously contributed in the bioaccumulation, adsorption, desorption and re-suspension of PAHs within different ecosystems (Arulazhagan and Vasudevan, 2011). However, very little proportion of such toxic PAHs undergo natural transformations while the larger part posed serious environmental concern as human health is the most critically affected (Lundstedt *et al.*, 2007; Bispo *et al.*, 1999). Hence, the current environmental challenges on PAHs need compliance with international best practices that eliminate lasting hazardous effects.

Environmental pollution involving used lubricating oil were reported in different parts of Malaysia where Selangor State recorded the highest reported cases (Mohd Radzi *et al.*, 2016; Keshavarzifard and Zakaria, 2015; Sany *et al.*, 2014). This created public health concerns due to the frequency of its occurrence and the severity it affects plants, animals, humans and the environments (Spinelli and Freitas, 2005; Akintunde *et al.*, 2015). Such oil damages animals' central nervous system and plants nutritional pathway (Todd *et al.*, 1999). Human being is also affected by used lubricating oil through the skin contact which absorbs PAHs into the bloodstream and eventually converts them into electrophilic derivatives that become cancer (Todd *et al.*, 1999). Additionally, humans are also affected on their central nervous systems which create disorderliness on the nerve, a situation termed as peripheral neuropathy (Todd *et al.*, 1999). Such situation causes feet and legs numbness which eventually becomes paralysis (Todd *et al.*, 1999). However, inadequate information on the toxicity of used lubricating oil and its post-disposal behaviour has further complicated issues as even sites not intended for its disposal are grossly affected (Oa and Lee, 2009; Park *et al.*, 2009; 2010).

Among the most hazardous PAHs commonly found in used lubricating oil are phenanthrene and pyrene which contained three and four aromatic structures arranged in angular and cluster forms respectively (Hossain and Salehuddin, 2012). These were classified as low and high molecular weights PAHs considering their number of aromatic rings (Wick *et al.*, 2011). Phenanthrene was reported to be a dangerous animal's skin photosensitizer which is characterized with mild allergen while pyrene has much toxicity effect on animal's body as it causes neurological disorder, lesions

on the kidney and reduces body weight (CCME, 2010; Patri *et al.*, 2010; Moody *et al.*, 2001).

The phenanthrene and pyrene conventional degradations were faced with high technological implication which generates more toxic intermediates (Park *et al.*, 2009; 2010). The biological removal is the scientifically proven technique due to its complete removal response and low cost implication when compared to conventional alternatives (Abdulsalam and Omale, 2009; Karamalidis *et al.*, 2010). In the biological removal, efficient microorganisms are utilized due to high abundance, species diversity, catabolic versatility and adaptation capability to adverse conditions (Moraes *et al.*, 2009). Bacteria were reported as the best microorganisms that that occupy suitable niche in used lubricating oil contaminated site due to broad enzymes possession that enable them utilize the PAHs as sole carbon sources (Madigan *et al.*, 1998). Previous studies indicates the PAHs removing capabilities of bacteria involving *Mycobacterium*, *Arthrobacter*, *Burkholderia*, *Sphingomonas*, and *Pseudomonas* (Baboshin *et al.*, 2008; Seo *et al.*, 2006; Kim *et al.*, 2003).

1.2 Problem statement

Despite the global acceptability of PAHs biodegradation, many degrading bacteria are faced with challenges that limit their rapid degradation responses (Seo *et al.*, 2009). Such limitations were caused by PAHs resistance to microbial attack, extended lag growth phase of the degrading isolates and unfavorable biodegradation conditions (Fankhauser, 2004). Among the most dangerous PAHs classified by USEPA were phenanthrene and pyrene which contained three and four benzene rings respectively (ASTDR, 2005). These phenanthrene and pyrene causes skin photosensitization, kidney and brain damages to human while destroying the soil and water ecosystems (Mesdhaghinia *et al.*, 2005; Patri *et al.*, 2010).

All the phenanthrene and pyrene degrading bacteria are very sensitive to physico-chemical conditions whose little alteration will negatively affect their metabolic activities (Leahy and Colwell, 1990). Such alteration will minimize degradation efficiency of the bacteria which subsequently will cause low enzymes production that may extend the isolates lag growth phase (Fankhauser, 2004; Richnow *et al.*, 1994). In this situation, PAHs intracellular localization is reduced which further complicates the entire biodegradation process as many bacteria survived at sole expense of PAHs (Chauhan *et al.*, 2008). However, optimizing the physico-chemical biodegradation conditions using statistically based approach is the best alternative to enhance degradation outcomes (Richnow *et al.*, 1994; Szabó *et al.*, 2007).

Another limitation that considerably affects phenanthrene and pyrene biodegradation is heavy metal co-contaminants whose complex interactions with the PAHs increase hazardous environmental risks (Markowicz, *et al.*, 2016; Sarma *et al.*, 2016). This negatively suppresses the bacterial metabolic activities due to combined effects from both PAHs and the heavy metals (Lu *et al.*, 2013; Thavamani *et al.*, 2012). Currently,

there are limited studies on the bacterial degradation response for PAHs in a mixed presence with heavy metals (Markowicz *et al.*, 2016; Sarma *et al.*, 2016).

Based on the prevailing slow phenanthrene and pyrene degradation limitation experienced by the bacteria, research questions that instigated the conduct of this study include; does used vehicle lubricant bioavailability perform effective phenanthrene and pyrene degradations at much faster rate? How do we determine the desirability of such microorganisms in a complex situation that involved PAHs and hazardous heavy metals?

1.3 Significant of the study

Effective biodegradation of phenanthrene and pyrene provides avenue of tackling used lubricating oil and its environmental challenges in Malaysia. This creates opportunity of using the locally isolated bacteria from the commonly natural inhabitants of used lubricating oil contaminated sites as fast effective biodegraders. The previously isolated bacteria were shown to degrade less than 120 mg/L phenanthrene and pyrene in more than 20 days of incubation (Abdul-Talib *et al.*, 2015; Sarma *et al.*, 2004). Therefore, isolating effective phenanthrene and pyrene degrading bacteria that can remove more than higher concentrations in 24 hours is much needed in Malaysia. This will improve the environmental condition which receives considerable amounts of used lubricating oil on daily basis. Additionally, opportunity will be provided to confirm the attainment of suitable niche by such bacteria in a complex system that is co-contaminated with heavy metals pollutants.

1.4 Objectives of the study

The main aim of the present study is to identify efficient bacteria that rapidly degrade phenanthrene and pyrene belonging to low and high molecular weight PAHs. The bacteria were enhanced by statistical based approach and tested for the biodegradation efficiency in complex mixture with heavy metals. The specific objectives designed to achieve this aim include:

1. To isolate and identify phenanthrene and pyrene degrading bacteria from used vehicle lubricant contaminated soil.
2. To optimize phenanthrene and pyrene biodegradation conditions.
3. To analyse phenanthrene and pyrene biodegradation intermediates.
4. To evaluate the effects of heavy metals co-contaminants on the phenanthrene and pyrene biodegradation.

REFERENCES

- Abbas, A., and Edwards, C. (1989). Effects of metals on a range of *Streptomyces* species. *Applied and Environmental Microbiology*, 55(8), 2030-2035.
- Abdulsalam, S., and Omale, A.B. (2009). Comparison of biostimulation and bioaugmentation techniques for the remediation of used motor oil contaminated soil. *Brazilian Archives of Biology and Technology* 52(3), 747-754.
- Abdul-Talib, S., Mohd-Kamil, N.A.F., Alias, S., and Othman, N. (2015). Degradation of phenanthrene in liquid culture and sand slurry by *Corynebacterium urealyticum*. *Civil and Environmental Research*, 7(3), 46-55.
- Achten, C., and Hofmann, T. (2009). Native polycyclic aromatic hydrocarbons in coals - a hardly recognized source of environmental contamination. *Science of the Total Environment*, 407(8), 2461-2473.
- Acinas, S.G., Rodríguez-Valera, F., and Pedrós-Alió, C. (1997). Spatial and temporal variation in marine bacterioplankton diversity as shown by RFLP fingerprinting of PCR amplified 16S rDNA. *FEMS Microbiology Ecology*, 24(1), 27-40.
- Adam, A.M., and Veenstra, J.N. (1993). Effects of selected environmental parameters on the biodegradation of 1,2-dichloropropane. *Proceedings of the 48th Industrial Waste Conference, Purdue University*, 48, 261.
- Adelowo, O.O., Alagbe, S.O., and Ayandele, A.A. (2006). Time-dependent stability of used engine oil degradation by cultures of *Pseudomonas fragi* and *Achromobacter aerogens*. *African Journal of Biotechnology*, 5(24), 2476-2479.
- Aelion, C.M., Dobbins, D.C., and Pfaender, F.K. (1989). Adaptation of aquifer microbial communities to the biodegradation of xenobiotic compounds: Influence of substrate concentration and pre-exposure. *Environmental Toxicology and Chemistry*, 8(1), 75-86.
- Aelion, C.M., Swindoll, C.M., and Pfaender, F.K. (1987). Adaptation to and biodegradation of xenobiotic compounds by microbial communities from a pristine aquifer. *Applied and Environmental Microbiology*, 53(9), 2212-2217.
- Ahn, Y., Sanseverino, J., and Sayler, G.S. (1999). Analyses of polycyclic aromatic hydrocarbon-degrading bacteria isolated from contaminated soils. *Biodegradation* 10(2), 49-157.
- Ahuatzi-Chacon, D., Ordorica-Morales, G., Ruiz-Ordaz, N., Cristiani-Urbina, E., Juárez-Ramírez, C., and Galíndez-Mayer, J. (2004). Kinetic study of phenol hydroxylase and catechol 1,2-dioxygenase biosynthesis by *Candida tropicalis* cells grown on different phenolic substrates. *World Journal of Microbiology and Biotechnology*, 20(7), 695-702.

- Aislabie, J., D.J. Saul, and Foght, J.M. (2006). Bioremediation of hydrocarbon-contaminated polar soils. *Extremophiles*, 10(3), 171-179.
- Akhnazarova, S. and Kafarov, V. (1982). *Experiment Optimization in Chemistry and Chemical Engineering*, Mir Publishers, Moscow.
- Akintunde, W. O., Olugbenga, O. A., and Olufemi, O. O. (2015). Some Adverse Effects of Used Engine Oil (Common Waste Pollutant) On Reproduction of Male Sprague Dawley Rats. *Open access Macedonian Journal of Medical Sciences*, 3(1), 46.
- Aksmann, A., and Tukaj, Z. (2008). Intact anthracene inhibits photosynthesis in algal cells; a fluorescence induction study on *Chlamydomonas reinhardtii* cw92 strain. *Chemosphere*, 74(1), 26-32.
- Alexander, M. (1999). *Biodegradation and Bioremediation*. Academic Press, San Deigo, USA.
- Alexander, M. (1978). *Advances in Microbial Ecology, Volume 2*. Plenum Press, New York.
- Ali, O.A., and Tarek, S.J. (2009). Removal of polycyclic aromatic hydrocarbons from Ismailia Canal water by chlorine, chlorine dioxide and ozone. *Desalination and Water Treatment*, 1(1-3), 289-298.
- Altaf, M., Naveena, B.J., and Reddy, G. (2007). Use of inexpensive nitrogen sources and starch for L(+) lactic acid production in anaerobic submerged fermentation. *Bioresource Technology*, 98(3), 498-503.
- Andrea, L., and Alleman, B.C. (1999). *Bioremediation Technologies for Polycyclic Aromatic Hydrocarbon Compounds*. Battelle Press Columbus, Ohio, USA
- Anfinsen, C.B. (1973). Studies on the principles that govern the folding of protein chains. *Science*, 181(4096), 223-30.
- Arulazhagan, P., and Vasudevan N. (2011). Biodegradation of polycyclic aromatic hydrocarbons by halotolerant bacterial strain *Ochrobactrum* sp. VA1. *Marine Pollution Bulletin*, 62(2), 388-394.
- Arulazhagan, P., Vasudevan, N., and Yeom, I.T. (2010). Biodegradation of polycyclic aromatic hydrocarbons by halotolerant bacterial consortium isolated from marine environment. *International Journal of Environmental Science and Technology*, 7(4), 639-652.
- Assinder, S.J., and Williams, P.A. (1990). The TOL plasmids: determinants of the catabolism of toluene and the xylenes. *Advances in Microbial Physiology*, 31, 1-69.

- Asturias, J.A., and Timmis, K.N. (1993). Three different 2,3-dihydroxybiphenyl-1,2-dioxygenase genes in the gram-positive polychlorobiphenyl-degrading bacterium *Rhodococcus globerulus* P6. *Journal of Bacteriology*, 175(15), 4631-4640.
- Atlas, R.M. (2010). *Handbook of microbiological media*. CRC press, Taylor and Francis Group, Boca Raton, Florida, USA.
- Atlas, R.M. (1995). Bioremediation of petroleum pollutants. *International Biodeterioration and Biodegradation*, 35(1-3), 317-327.
- Atlas, R.M. (1991). Microbial hydrocarbon degradation - bioremediation of oil spills. *Journal of Chemical Technology and Biotechnology*, 52(2), 149-156.
- Atlas, R.M. (1988). *Experimental Microbiology: Fundamentals and Applications*. Benjamin-Cummings Publishing Company, San Francisco, USA.
- Atlas, R.M. (1981). Microbial degradation of petroleum hydrocarbons; an environmental perspective. *Microbiological Reviews*, 45(1), 180.
- Atlas, R.M. (1978). Microorganisms and petroleum products. *BioScience*, 28(6), 387-390.
- Baboshin, M., Akimov, V., Baskunov, B., Born, T.L., Khan, S.U., and Golovleva, L. (2008). Conversion of polycyclic aromatic hydrocarbons by *Sphingomonas* sp. VKM B-2434. *Biodegradation*, 19(4), 567-576.
- Balabin, R.M. (2009). Polar (acyclic) isomer of formic acid dimer: Gas-phase Raman spectroscopy study and thermodynamic parameters. *The Journal of Physical Chemistry A*, 113(17), 4910-4918.
- Balashova, N.V., Koshelva, I.A., Golovchenko, N.P., and Boronin, A.M. (1999). Phenanthrene metabolism by *Pseudomonas* and *Burkholderia* strains. *Process of Biochemistry* 35(3), 291-296.
- Bamforth, S.M., and Singleton, I. (2005). Bioremediation of polycyclic aromatic hydrocarbons: current knowledge and future directions. *Journal of Chemical Technology and Biotechnology*, 80(7), 723-736.
- Banat, I.M., Satpute, S.K., Cameotra, S.S., Patil, R., and Nyayanit, N.V. (2014). Cost effective technologies and renewable substrates for biosurfactants' production. *Frontiers in Microbiology*, 5(697), 1-18.
- Bao, M.T., Wang, L.N., Sun, P.Y., Cao, L.X., Zou, J., and Li, Y.M. (2012). Biodegradation of crude oil using an efficient microbial consortium in a simulated marine environment. *Marine Pollution Bulletin*, 64(6), 1177-1185.
- Baranyi, J., and Roberts, T.A. (1995). Mathematics of predictive food microbiology. *International Journal of Food Microbiology*, 26(2), 199-218.

- Barea, J.M., Pozo, M.J., Azcon, R., and Azcon-Aguilar, C. (2005). Microbial cooperation in the rhizosphere. *Journal of Experimental Botany*, 56(417), 1761-1778.
- Bemgård, A., Colmsjö, A., and Lundmark, B.O. (1992). Gas chromatographic analysis of high-molecular-weight polynuclear aromatic hydrocarbons: Molecular weight 328. *Journal of Chromatography A*, 595(1-2), 247-258.
- Bentley, A., Atkinson, A., Jezek, J., and Rawson, D.M. (2001). Whole cell biosensors-electrochemical and optical approaches to ecotoxicity testing. *Toxicology in vitro*, 15(4), 469-475.
- Bhattacharyya, J.K., and Shekdar, A.V. (2003). Treatment and disposal of refinery sludges: Indian scenario. *Waste Management and Research*, 21(3), 249-261.
- Bidoia, E.D., Montagnolli, R.N., and Lopes, P.R.M. (2010). Microbial biodegradation potential of hydrocarbons evaluated by colorimetric technique. *Applied Microbiology and Microbial Biotechnology, Formatex*, 2, 1277-1288.
- Birgül, A., Tasdemir, Y., and Cindoruk, S.S. (2011). Atmospheric wet and dry deposition of polycyclic aromatic hydrocarbons determined using a modified sampler. *Atmospheric Research*, 101(1), 341-353.
- Bispo, A., Jourdain, M.J., and Jauzein, M. (1999). Toxicity and genotoxicity of industrial soils polluted by polycyclic aromatic hydrocarbons (PAHs). *Organic Geochemistry*, 30(8), 947-952.
- Bizzari, S.N. and Blagoev, M. (2010). *CEH Marketing Research Report: Formic acid*. Chemical Economics Handbook, SRI consulting, USA.
- Boehm, P.D. (2006). Polycyclic Aromatic Hydrocarbons (PAHs). In: Morrison, R.D. and B.L. Murphy. (Eds.). *Environmental Forensics Contaminant Specific Guide*. Academic Press, Burlington, USA.
- Boonchan, S., Britz, M.L., and Stanley, G.A. (2000). Degradation and mineralization of high-molecular weight polycyclic aromatic hydrocarbons by defined fungal-bacterial co-cultures, *Applied Environmental Microbiology*, 66(3), 1007-1019.
- Boone, D.R., Johnson, R.L., and Liu, Y. (1989). Diffusion of the interspecies electron carriers H₂ and formate in methanogenic ecosystems and its implications in the measurement of K_m for H₂ or formate uptake. *Applied and Environmental Microbiology*, 55(7), 1735-1741.
- Boopathy, R. (2000). Factors limiting bioremediation technologies. *Bioresource Technology*, 74(1), 63-67.

- Borden, R.C., Bedient, P.B., Lee, M.D., Ward, C.H., and Wilson, J.T. (1986). Transport of dissolved hydrocarbons influenced by oxygen limited biodegradation: Field application. *Water Resources Research*, 22(13), 1983-1990.
- Borden, R.C., and Bedient, P.B. (1986). Transport of dissolved hydrocarbons influenced by oxygen-limited biodegradation: Theoretical development. *Water Resources Research*, 22(13), 1973-1982.
- Bosma, T.N., Middeldorp, P.J., Schraa, G., and Zehnder, A.J. (1997). Mass transfer limitation of biotransformation: quantifying bioavailability. *Environmental Science and Technology*, 31(1), 248-252.
- Bossert I., and Bartha, R. (1984). The Fate of Petroleum in Soil Ecosystem: In Atlas, R.M. (Ed.) *Petroleum Microbiology*. Macmillan Publishing Company, New York, USA, 435-473.
- Boszczyk-Maleszak, H., Zabost, A., Wolicka, D., and Kacieszczenko, J. (2006). Effectiveness of biodegradation of petroleum products by mixed bacterial populations in liquid medium at different pH values. *Polish Journal of Microbiology*, 55(1), 69-73
- Box, G.E.P., Hunter W.G. and Hunter, J.S. (1978). *Statistics for Experimenters*. John Wiley and Sons, New York, USA, 291-334.
- Box, G.E.P. and Wilson, K.B. (1951). On the experimental attainment of optimum conditions. *Journal of the Royal Statistical Society*, B13, 1-45.
- Bradl H.E. (2005). Sources and origins of heavy metals. In Bradl H. E. (ed.), *Heavy Metals in the Environment: Origin, Interaction and Remediation*, Elsevier, Amsterdam
- Brady, N.C., and Weil, R.R. (2000). *Elements of the Nature and Properties of Soils* (No. 631.4 B733E.). Upper Saddle River, NJ, Prentice Hall, USA.
- Brandt, H.C.A., and Watson, W.P. (2003). Monitoring human occupational and environmental exposures to PAHs. *The Annals of Occupational Hygiene*, 47(5), 349-378.
- Brinda, L.M., Muthukumar, K., and Velan, M. (2013). Optimization of minimal salt medium for efficient phenanthrene biodegradation by *Mycoplana* sp. MVMB2 isolated from petroleum contaminated soil using factorial design experiments. *CLEAN–Soil, Air, Water*, 41(1), 51-59.
- Brown, L.D., and Ulrich, A.C. (2014). *Bioremediation of Oil Spills on Land*. Handbook of Oil Spill Science and Technology, 395-406.

- Cain, R.B. (1981). Regulation of aromatic and hydroaromatic catabolic pathways in *Nocardioform actinomycetes*. *Zentralbl. Bakteriolog. Mikrobiol. Hyg. I. Abt. Orig. Suppl.*, 11, 335-354.
- Cao, L., Shen, G., and Lu, Y. (2008). Combined effects of heavy metal and polycyclic aromatic hydrocarbon on soil microorganism communities. *Environmental Geology*, 54(7), 1531-1536.
- Calvo, C., Manzanera, M., Silva-Castro, G.A., Uad, I., and González-López, J. (2009). Application of bioemulsifiers in soil oil bioremediation processes: Future prospects. *Science of the Total Environment*, 407(12), 3634-3640.
- Carrington, B., Lowe, A., Shaw, L.E., and Williams, P.A. (1994). The lower pathway operon for benzoate catabolism in biphenyl-utilizing *Pseudomonas* sp. strain, IC and the nucleotide sequence of the bphE gene for catechol 2,3-dioxygenase. *Microbiology*, 140(3), 499-508.
- Cerniglia, C.E. (1992). Biodegradation of polycyclic aromatic hydrocarbons. *Biodegradation*, 3, 351-368.
- Cerniglia, C.E. (1984). Microbial metabolism of polycyclic aromatic hydrocarbons. *Advances in Applied Microbiology*, 30, 31-71.
- Chai, C.Y., Salanitro, J.P., Colthart, J.D., and Klein, C.L. (1987). Effects of dissolved oxygen on the biodegradation of BTX in a sandy aquifer. In *Proceedings of NWWA/API Conference on Petroleum Hydrocarbons and Organic Chemicals in Ground Water: Prevention, Detection, and Restoration*, Houston, 451- 461.
- Chaudhry, Q., Blom-Zandstra, M., Gupta, S.K., and Joner, E. (2005). Utilising the synergy between plants and rhizosphere microorganisms to enhance breakdown of organic pollutants in the environment. *Environmental Science and Pollution Research*, 12(1), 34-48.
- Chauhan, A., Oakeshott, J.G., and Jain, R.K. (2008). Bacterial metabolism of polycyclic aromatic hydrocarbons; strategies for bioremediation. *Indian Journal of Microbiology*, 48(1), 95-113.
- Chen, L., Li, G., and Zu-Liang, C. (2010). Biodegradation of naphthalene by strain *Bacillus fusiformis*. *Journal of Hazardous Materials*, 182(1), 771-777.
- Chen, B., Xuan, X., Zhu, L., Wang, J., Gao, Y., Yang, K., and Lou, B. (2004). Distributions of polycyclic aromatic hydrocarbons in surface waters, sediments and soils of Hangzhou City, China. *Water Research*, 38(16), 3558-3568.
- Cheung, P.Y., and Kinkle, B.K. (2001). *Mycobacterium* diversity and pyrene mineralization in petroleum contaminated soils, *Applied and Environmental Microbiology*, 67(5), 2222-2229.

- Cheung, H., Tanke, R.S., and Torrence, G.P. (2005). *Acetic Acid; Ullmann's Encyclopaedia of Industrial Chemistry*. Wiley-VCH, Weinheim, Germany.
- Chizhova, T., Hayakawa, K., Tishchenko, P., Nakase, H., and Koudryashova, Y. (2013). Distribution of polycyclic aromatic hydrocarbons in northwestern part of Japan Sea. *Deep Sea Research Part II: Topical Studies in Oceanography*, 86, 19-24.
- Chung, M.K., Hu, R., Cheung, K.C., and Wong, M.H. (2007). Pollutants in Hong Kong soils: Polycyclic aromatic hydrocarbons. *Chemosphere*, 67(3), 464–473
- Churchill, S.A., Harper, J.P., and Churchill, P.F. (1999). Isolation and characterization of a *Mycobacterium* species capable of degrading three and four ring aromatic and aliphatic hydrocarbons, *Applied Environmental Microbiology*, 65(2), 549-52.
- Clayden, J., Greeves, N., and Warren, S. (2002). *Organic Chemistry, Second Edition*. Oxford University Press, UK, 101–136.
- Claverys, J.P., and Håvarstein, L.S. (2007). Cannibalism and fratricide: mechanisms and raisons d'etre. *Nature Reviews Microbiology*, 5, 219-229.
- Cocaign-Bousquet, M., Garrigues, C., Loubiere, P., and Lindley, N.D. (1996). Physiology of pyruvate metabolism in *Lactococcus lactis*. *Antonie Van Leeuwenhoek*, 70(2), 253-267.
- Cody, G.D., Boctor, N.Z., Filley, T.R., Hazen, R.M., Scott, J.H., Sharma, A., and Yoder, H.S. Jr (2000). Primordial carbonylated iron-sulfur compounds and the synthesis of pyruvate. *Science*, 289(5483), 1337–1340.
- Connell, D.W. (2005). *Basic Concepts of Environmental Chemistry*. CRC Press, USA, 33487-2742.
- Cooper, C.S., Grover, P.L., and Sims, P. (1983). The metabolism and activation of benzo [a] pyrene. *Progress in Drug Metabolism*, 7, 295-396.
- Coral, G., and Karagoz, S. (2005). Isolation and characterization of phenanthrene-degrading bacteria from a petroleum refinery soil. *Annals of Microbiology*, 55(4), 255-259.
- Cosmetic Ingredient Review, CIR (2006). Annual review of cosmetic ingredient safety assessments (2004/2005). *International Journal of Toxicology*, 25, 1-89.
- Cousins, I.T., Beck, A.J., and Jones, K.C. (1999). A review of the processes involved in the exchange of semi-volatile organic compounds across the air-soil interface. *Science of the Total Environment*, 228(1), 5-24.

- Cowan, S.T., Barrow, G.I., Steel, K.J., and Feltham, R.K.A. (2004). *Cowan and Steel's Manual for the Identification of Medical Bacteria*. Cambridge University Press, UK.
- Cruz, J.A., Lantican, N.B., Delfin, E.F., and Paterno, E.S. (2015). Characterization and identification of growth-promoting *Actinomycetes*: A potential microbial inoculant. *Asia Life Sciences*, 24(1), 383-397.
- Cycoń, M., Żmijowska, A., Wójcik, M., and Piotrowska-Seget, Z. (2013). Biodegradation and bioremediation potential of diazinon-degrading *Serratia marcescens* to remove other organophosphorus pesticides from soils. *Journal of Environmental Management*, 117, 7-16.
- Czárán, T.L., Hoekstra, R.F., and Pagie, L. (2002). Chemical warfare between microbes promotes biodiversity. *Proceedings of the National Academy of Sciences*, 99(2), 786-790.
- Dabestani, R., and Ivanov, I. N. (1999). A compilation of physical, spectroscopic and photophysical properties of polycyclic aromatic hydrocarbons. *Photochemistry and Photobiology*, 70(1), 10-34.
- Dagley, S. (1986). *Biochemistry of aromatic hydrocarbon degradation*. In *The Biology of Pseudomonads*. The Bacteria: A Treatise on Structure and Function. Academic Press, USA, 527-555.
- Dagley, S. (1985). Microbial metabolism of aromatic compounds. In: M. Moo-Young, (Ed.), *Comprehensive Biotechnology, volume 1*. Pergamon Press, Elsevier, UK, 505.
- Dagley, S. (1978). *Pathways for the Utilization of Organic Growth Substrates*. Academic Press, Cambridge, Massachusetts, USA, 388.
- Dancer, G.I., Mah, J.H., Rhee, M.S., Hwang, I.G., and Kang, D.H. (2009). Resistance of *E. sakazakii* to environmental stresses. *Journal of Applied Microbiology*, 107(5), 1606-1614.
- Dawson, R.M.C., Elliott, D.C., Elliott, W.H., and Jones, K.M. (1986). Stability constants of metal complexes. *Data for Biochemical Research*, 399-415.
- de Carvalho, C.C., and da Fonseca, M.M.R. (2004). Solvent toxicity in organic-aqueous systems analysed by multivariate analysis. *Bioprocess and Biosystems Engineering*, 26(6), 361-375.
- Declercq, I., Cappuyns, V., and Duclos, Y. (2012). Monitored natural attenuation of contaminated soils: state of the art in Europe - a critical evaluation. *Science of the Total Environment*, 426, 393-405.

- Desai A., and Vyas, P. (2006). *Petroleum and Hydrocarbons Microbiology: Applied Microbiology*. Report from the Department of Microbiology, MS University of Baroda, Vadodara India, 1-22.
- Diaz, E. (2008). *Microbial Biodegradation: Genomics and Molecular Biology*. Caister Academic Press, Norfolk, U.K.
- Dibble, J.T., and Bartha, R. (1979). Effect of environmental parameters on the biodegradation of oil sludge. *Applied and Environmental Microbiology*, 37(4), 729-739.
- Djekrif-Dakhmouche, S., Gheribi-Aoulmi, Z., Meraihi, Z., and Bennamoun, L. (2006). Application of a statistical design to the optimization of culture medium for α -amylase production by *Aspergillus niger* ATCC 16404 grown on orange waste powder. *Journal of Food Engineering*, 73(2), 190–197.
- Doddamani, H.P., and Ninnekar, H.Z. (2000). Biodegradation of phenanthrene by a *Bacillus* sp. *Current Microbiology*, 41(1), 11-14.
- Donahue, R.L., Miller, R.W., and Shickluna, J.C. (1977). *Soils: An Introduction to Soils and Plant Growth*. Prentice-Hall International Inc, USA.
- Doyle, A., and Bryan, G.J. (1998). *Cell and tissue culture: laboratory procedure in biotechnology*. Chichester: John Willey and Sons Ltd, Baffins Lane Chichester, West Sussex P019 1UD, England
- Duffus, J. (2009). Heavy metals; a meaningless term? (IUPAC Technical Report). *Pure and Applied Chemistry*, 74(5), 793-807.
- Duetz, W.A., Rüedi, L., Hermann, R., O'Connor, K., Büchs, J., and Witholt, B. (2000). Methods for intense aeration, growth, storage, and replication of bacterial strains in microtiter plates. *Applied and Environmental Microbiology*, 66(6), 2641-2646.
- Duque, E., Haidour, A., Godoy, F., and Ramos, J. L. (1993). Construction of *Pseudomonas* hybrid strain that mineralizes 2,4,6-trinitrotoluene. *Journal of Bacteriology*, 175(8), 2278-2283.
- Duxbury, T., and Bicknell, B. (1983). Metal-tolerant bacterial populations from natural and metal-polluted soils. *Soil Biology and Biochemistry*, 15(3), 243-250.
- Dzombak, D.A., and Luthy, R.G. (1984). Estimating adsorption of polycyclic aromatic hydrocarbons on soils. *Soil Science*, 137(5), 292–308.
- Edewor, T.I., Adelowo, O.O., and Afolabi, T.J. (2004). Preliminary studies into the biological activities of a broad spectrum disinfectant formulated from used engine oil." *Pollution Research*, 23(4), 581-586.

- Eijsink, V.G., Axelsson, L., Diep, D.B., Håvarstein, L.S., Holo, H., and Nes, I.F. (2002). Production of class II bacteriocins by lactic acid bacteria; an example of biological warfare and communication. *Antonie Van Leeuwenhoek*, 81(1-4), 639-654.
- Eiland, F. (1981). The effects of application of sewage sludge on microorganisms in soil; microbial biomass, microbial activity, enzymatic activity, sewage sludge, heavy metals. *Tidsskrift for Planteavl, Denmark*.
- Elibol, M. (2004). Optimization of medium composition for actinorhodin production by *Streptomyces coelicolor* A3(2) with response surface methodology. *Process Biochemistry* 39(9), 1057–1062.
- Ellermeier, C.D., Hobbs, E.C., Gonzalez-Pastor, J.E., and Losick, R. (2006). A three-protein signaling pathway governing immunity to a bacterial cannibalism toxin. *Cell*, 124(3), 549-559.
- Erickson, B.D., and Mondello, F.J. (1993). Enhanced biodegradation of polychlorinated biphenyls after site-directed mutagenesis of a biphenyl dioxygenase gene. *Applied and Environmental Microbiology*, 59(11), 3858-3862.
- European Union (2011). Current EU approved additives and their E Numbers. *UK Food Standards Agency*, 10-27.
- Evans, W.C., Fernley, H.N., and Griffiths, E. (1965). Oxidative metabolism of phenanthrene and anthracene by soil *Pseudomonads*, the ring-fission mechanism. *Biochemical Journal*, 95(3), 819.
- Fahy, A., Ball, A.S., Lethbridge, G., Timmis, K.N., and McGenity, T.J. (2008). Isolation of alkali-tolerant benzene-degrading bacteria from a contaminated aquifer. *Letters in Applied Microbiology*, 47(1), 60-66.
- Fankhauser, D.B. (2004). *Bacteria Growth Curve*. University of Cincinnati Clermont College, 4200 Clermont College Drive, Batavia, Ohio 45103, USA
- Federal Register of Legislative Instruments, FRLI (2011). Australia and New Zealand food standard codes - Standard 1.2.4 - Labelling of Ingredients, F2011C00827, 125, 1-14
- Fellay, C., Dyson, P.J., and Laurency, G. (2008). A viable hydrogen storage system based on selective formic acid decomposition with a ruthenium Catalyst. *Angewandte Chemie*, 120(21), 4030-4032.
- Felsenstein, J. (1985). Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39(4), 783-791.

- Feng, T.C., Cui, C.Z., Dong, F., Feng, Y.Y., Liu, Y.D., and Yang, X.M., (2012). Phenanthrene biodegradation by halophilic *Marteella* sp. *Journal of Applied Microbiology*, 113(4), 779-789.
- Fetzer, J.C. (2000). *Large (C = 24) Polycyclic Aromatic Hydrocarbons: Chemistry and Analysis*, vol. 158. John Wiley and Sons.
- Fiehn, O. (2002). Metabolomics - the link between genotypes and phenotypes. *Plant Molecular Biology*, 48, 155-171.
- Freeze, R.A., and Cherry, J.A. (1979). *Groundwater, second edition*. Prentice-Hall International, Universiti Michigan, USA.
- Fusey, P., and Oudot, J. (1984). Relative influence of physical removal and biodegradation in the depuration of petroleum-contaminated seashore sediments. *Marine Pollution Bulletin*, 15, 136-141.
- Gabriel, J., Shah, V., Nesměrák, K., Baldrian, P., and Nerud, F. (2000). Degradation of polycyclic aromatic hydrocarbons by copper (II) hydrogen peroxide system. *Folia Microbiologica*, 45(6), 573-575.
- Ghanem, I., Orfi, M., and Shamma, M. (2007). Biodegradation of chlorpyrifos by *Klebsiella* sp. isolated from an activated sludge sample of waste water treatment plant in Damascus. *Folia Microbiologica*, 52(4), 423-427.
- Ghazali, F.M., Rahman, R.N.Z.A., Salleh, A.B., and Basri, M. (2004). Biodegradation of hydrocarbons in soil by microbial consortium. *International Biodeterioration and Biodegradation*, 54(1), 61-67.
- Ghosal, D., In-Soon, Y., and Gunsalus, I.C. (1987). Nucleotide sequence and expression of gene nahH of plasmid NAH7 and homology with gene xylE of TOL pWWO. *Gene*, 55(1), 19-28.
- Gibson, D.T., Zylstra, G.J., and Chauhan, S. (1990). Biotransformations catalyzed by toluene dioxygenase from *Pseudomonas putida* F 1, 132.
- Gibson, D.T. Koch, J.R. and Kallio, R.E. (1968). Oxidative degradation of aromatic hydrocarbons by microorganisms. In Enzymatic formation of catechol from benzene. *Biochemistry*, 7(7), 2653-2661.
- Gillespie, S.H., and Hawkey, P.M. (2006). *Principles and Practice of Clinical Bacteriology, second edition*, John Wiley Publisher, UK.
- Goodacre, R., Vaidyanathan, S., Dunn, W.B., Harrigan, G.G., and Kell, D.B. (2004). Metabolomics by numbers: acquiring and understanding global metabolite data. *Trends in Biotechnology*, 22(5), 245-252.
- Gogolev, A., and Wilke, B.M. (1997). Combination effects of heavy metals and fluoranthene on soil bacteria. *Biology and Fertility of Soils*, 25(3), 274-278.

- Gohel, V., Jiwan, D., Vyas, P., and Chatpar, H.S. (2005). Statistical optimization of chitinase production by *Pantoea dispersa* to enhance degradation of crustacean chitin waste. *Journal of Microbiology and Biotechnology*, 15(1), 197-201.
- Gold, V., Loening, K.L., McNaught, A.D., and Shemi, P. (1997). *IUPAC Compendium of Chemical Terminology*. Blackwell Science Ltd, Oxford, UK.
- Goyer, R., Golub, M., Choudhury, H., Hughes, M., Kenyon, E., and Stifelman, M. (2004). *Issue Paper on the Human Health Effects of Metals*. In USEPA risk assessment forum, volume 1200.
- González-Pastor, J.E., Hobbs, E.C., and Losick, R. (2003). Cannibalism by sporulating bacteria. *Science*, 301(5632), 510-513.
- Greenberg, H.J. (2004). *Mathematical programming glossary*. University of Colorado, USA.
- Grimberg, S.J., Stringfellow, W.T., and Aitken, M.D. (1996). Quantifying the biodegradation of phenanthrene by *Pseudomonas stutzeri* P16 in the presence of non-ionic surfactant. *Applied and Environmental Microbiology*, 62(7), 2387-2392.
- Grimmer, G., Dettbarn, G., Brune, H., Deutsch-Wenzel, R., and Misfeld, J. (1982). Quantification of the carcinogenic effect of polycyclic aromatic hydrocarbons in used engine oil by topical application onto the skin of mice. *International Archives of Occupational and Environmental Health*, 50(1), 95-100.
- Guo, W., He, M., Yang, Z., Lin, C., Quan, X., and Wang, H. (2007). Distribution of polycyclic aromatic hydrocarbons in water, suspended particulate matter and sediment from Daliao River watershed, China. *Chemosphere*, 68(1), 93-104.
- Gupta, S., Pathak, B., and Fulekar, M.H. (2015). Molecular approaches for biodegradation of polycyclic aromatic hydrocarbon compounds: a review. *Reviews in Environmental Science and Biotechnology*, 14(2), 241-269.
- Gurbiel, R.J., Batie, C.J., Sivaraja, M., True, A.E., Fee, J.A., Hoffman, B.M., and Ballou, D.P. (1989). Electron-nuclear double resonance spectroscopy of ¹⁵N-enriched phthalate dioxygenase from *Pseudomonas cepacia* proves that two histidines are coordinated to the [2Fe-2S] Rieske-type clusters. *Biochemistry*, 28(11), 4861-4871.
- Guzzo, A., and DuBow, M.S. (1994). Identification and characterization of genetically programmed responses to toxic metal exposure in *Escherichia coli*. *FEMS Microbiology Reviews*, 14(4), 369-374.
- Hampel, C.A. (1968). *The Encyclopedia of the Chemical Elements Reinhold*, New York, USA, 768.

- Hanks, R.J. (2012). *Applied soil physics: soil water and temperature applications, volume 8*. Springer, Germany.
- Hanson, K.G., Desai, J.D., and Desai, A.J. (1993). A rapid and simple screening technique for potential crude oil degrading microorganisms. *Biotechnology Techniques*, 7(10), 745-748
- Harayama, S., Kok, M., and Neidle, E.L. (1992). Functional and evolutionary relationships among diverse oxygenases. *Annual Reviews in Microbiology*, 46(1), 565-601.
- Harayama, S., and Rekik, M. (1989). Bacterial aromatic ring-cleavage enzymes are classified into two different gene families. *Journal of Biological Chemistry*, 264(26), 15328-15333.
- Harvey, R.G. (1996). Mechanisms of carcinogenesis of polycyclic aromatic hydrocarbons. *Polycyclic Aromatic Compounds*, 9(1-4), 1-23.
- Haynie, D.T. (2001). *Biological Thermodynamics*. Cambridge University Press, UK.
- Head, I.M., and Swannell, R.P. (1999). Bioremediation of petroleum hydrocarbon contaminants in marine habitats. *Current Opinion in Biotechnology*, 10, 234-239.
- Heiskanen, A., Yakovleva, J., Spégel, C., Taboryski, R., Koudelka-Hep, M., Emnéus, J., and Ruzgas, T. (2004). Amperometric monitoring of redox activity in living yeast cells: comparison of menadione and menadione sodium bisulfite as electron transfer mediators. *Electrochemistry Communications*, 6(2), 219-224.
- Heitkamp, M.A., and Cerniglia, C.E. (1988). Mineralization of polycyclic aromatic hydrocarbons by a bacterium isolated from sediment below an oil field. *Applied and Environmental Microbiology*, 54(6), 1612-1614.
- Herdan, J.M. (1997). Lubricating oil additives and the environment - an overview. *Lubrication Science*, 9(2), 161-172.
- Hilyard, E.J., Jones-Meehan, J.M., Spargo, B.J., and Hill, R.T. (2008). Enrichment, isolation, and phylogenetic identification of PAH-degrading bacteria from Elizabeth River sediments. *Applied and Environmental Microbiology*, 74(4), 1176-1182.
- Hiroki, M. (1994). Populations of cadmium tolerant microorganisms in soils polluted with heavy metals. *Soil Science and Plant Nutrition*, 40(3), 515-524.
- Hiroki, M. (1992). Effects of heavy metal contamination on soil microbial population. *Soil Science and Plant Nutrition*, 38(1), 141-147.

- Hirose, J., Kimura, N., Suyama, A., Kobayashi, A., Hayashida, S., and Furukawa, K. (1994). Functional and structural relationship of various extradiol aromatic ring-cleavage dioxygenases of *Pseudomonas* origin. *FEMS Microbiology Letters*, 118(3), 273-277.
- Hoover, R. (2014). Need to track organic nano-particles across the universe? NASA's got an app for that. NASA, USA. <http://www.nasa.gov/ames/need-to-track-organic-nano-particles-across-the-universenasas-got-an-app-for-that/#.VC2hdmeSzSs>
- Hossain, M.A., and Salehuddin, S.M. (2012). Polycyclic aromatic hydrocarbons in edible oils by gas chromatography coupled with mass spectroscopy. *Arabian Journal of Chemistry*, 5(3), 391-396.
- Hult, M.F., and Grabbe, R.R. (1985). Distribution of gases and hydrocarbon vapors in the unsaturated zone. In *US Geological Survey Toxic Substances Hydrology Program; Proceedings of the Technical Meeting, Cape Cod, Massachusetts*.
- Hussain, K., and Hoque, R.R. (2015). Seasonal attributes of urban soil polycyclic aromatic hydrocarbons of the Brahmaputra Valley. *Chemosphere*, 119, 794-802.
- Hutchins, S.R. (1991). Biodegradation of monoaromatic hydrocarbons by aquifer microorganisms using oxygen, nitrate, or nitrous oxide as the terminal electron acceptor. *Applied and Environmental Microbiology*, 57(8), 2403-2407.
- Ingraham, J.L., Low, K.B., Magasanik, B., Schaechter, M., and Umbarger, H.E. (1987). *Escherichia coli* and *Salmonella typhimurium*. *Cellular and Molecular Biology*, 2, 1334-1345.
- Ibarrolaza, A., Coppotelli, B.M., Del Panno, M.T., Donati, E.R., and Morelli, I.S. (2009). Dynamics of microbial community during bioremediation of phenanthrene and chromium (VI) contaminated soil microcosms. *Biodegradation*, 20(1), 95-107.
- Iftikhar, T., Niaz, M., Jabeen, R., and Haq, I.U. (2011). Purification and characterization of extracellular lipases. *Pakistan Journal of Botany*, 43(3), 1541-1545.
- Inouye, S., Kimoto, M., Nakazawa, A., and Nakazawa, T. (1990). Presence of flagella in *Pseudomonas putida* is dependent on the ntrA (rpoN) gene. *Molecular and General Genetics MGG*, 221(2), 295-298.
- International Agency for Research on Cancer, IARC (2010). Some non-heterocyclic polycyclic aromatic hydrocarbons and related exposures. *IARC Monographs on the Evolution of Carcinogenic Risks to Humans*, 92, 1-853.

- IARC (1989). Occupational exposures in petroleum refining; crude oil and major petroleum fuels. *IARC Monographs on the Evolution of Carcinogenic Risks to Humans*, 45, 1-322.
- Irha, N., Slet, J., and Petersell, V. (2003). Effect of heavy metals and PAH on soil assessed via dehydrogenase assay. *Environment International*, 28(8), 779-782.
- Isaac, P., Martínez, F.L., Bourguignon, N., Sánchez, L.A., and Ferrero, M.A. (2015). Improved PAHs removal performance by a defined bacterial consortium of indigenous *Pseudomonas* and *Actinobacteria* from Patagonia, Argentina. *International Biodeterioration and Biodegradation*, 101, 23-31.
- Iwabuchi, T., Yamauchi, Y.I., Katsuta, A., and Harayama, S. (1998). Isolation and characterization of marine *Nocardioides* capable of growing and degrading phenanthrene at 42 °C. *Journal of Marine Biotechnology*, 6, 86-90.
- Janbandhu, A., and Fulekar, M.H. (2011). Biodegradation of phenanthrene using adapted microbial consortium isolated from petrochemical contaminated environment. *Journal of Hazardous Materials*, 187(1), 333-340.
- Jaimés, 3rd, R., Kuzmiak-Glancy, S., Brooks, D.M., Swift, L.M., Posnack, N.G., and Kay, M.W. (2016). Functional response of the isolated, perfused normoxic heart to pyruvate dehydrogenase activation by dichloroacetate and pyruvate. *Pflügers Archiv-European Journal of Physiology*, 468(1), 131-142.
- Jiang, Y., Huang, H., Wu, M., Yu, X., Chen, Y., Liu, P., and Li, X. (2015). *Pseudomonas* sp. LZ-Q continuously degrades phenanthrene under hypersaline and hyperalkaline condition in a membrane bioreactor system. *Biophysics Reports*, 1(3), 156-167.
- Jonsson, P., Johansson, A.I., Gullberg, J., Trygg, J., Grung, B., Marklund, S., and Moritz, T. (2005). High-throughput data analysis for detecting and identifying differences between samples in GC/MS-based metabolomic analyses. *Analytical Chemistry*, 77(17), 5635-5642.
- Johnsen, A., and Karlson, U. (2005). PAH degradation capacity of soil microbial communities: Does it depend on PAH exposure? *Microbial Ecology*, 50(4), 488-495.
- Jones, S.H., and Alexander, M. (1988). Effect of inorganic nutrients on the acclimation period preceding mineralization of organic chemicals in lake water. *Applied and Environmental Microbiology*, 54(12), 3177-3179.
- Joutey, N.T., Bahafid, W., Sayel, H., and El Ghachtouli, N. (2013). Biodegradation: involved microorganisms and genetically engineered microorganisms. *Biodegradation - Life of Science. InTech, Rijeka*, 289-320.

- Juhasz, A.L., and Naidu, R. (2000). Bioremediation of high molecular weight polycyclic aromatic hydrocarbons: a review of the microbial degradation of benzo[a]pyrene. *International Biodeterioration and Biodegradation*, 45(1), 57-88.
- Jurak, E., Patyshakuliyeva, A., De Vries, R.P., Gruppen, H., and Kabel, M.A. (2015). Compost grown *Agaricus bisporus* lacks the ability to degrade and consume highly substituted xylan fragments. *PloS one*, 10(8), e0134169.
- Kanaly, R.A., and Harayama, S. (2000). Biodegradation of high-molecular-weight polycyclic aromatic hydrocarbons by bacteria. *Journal of Bacteriology*, 182(8), 2059-2067.
- Kafilzadeh, F., Khosrobak, A., and Jamali, H. (2016). Isolation and identification of phenanthrene degrading bacteria from the soil around oil company of Andimeshk and investigation of their growth kinetics. *Polycyclic Aromatic Compounds*, 36(1), 58-71.
- Kandhai, M.C., Reij, M.W., Gorris, L.G., Guillaume-Gentil, O., and Schothorst, M. (2004). Occurrence of *Enterobacter sakazakii* in food production environments and households. *Lancet*, 363(9402), 39-40.
- Kang, H., Hwang, S.Y., Kim, Y.M., Kim, E., Kim, Y.S., Kim, S.K., and Zylstra, G.J. (2003). Degradation of phenanthrene and naphthalene by a *Burkholderia* sp. strain. *Canadian Journal of Microbiology*, 49(2), 139-144.
- Karamalidis, A.K., Evangelou, A.C., Karabika, E., Koukkou, A.I., Drains, C., and Voudrias, E.A. (2010). Laboratory scale bioremediation of petroleum contaminated soil by indigenous microorganisms and added *Pseudomonas aeruginosa* strain Spet. *Bioresource Technology*, 101(16), 6545-6552.
- Kästner, M., Breuer-Jammali, M., and Mahro, B. (1998). Impact of inoculation protocols, salinity, and pH on the degradation of polycyclic aromatic hydrocarbons and survival of PAH-degrading bacteria introduced into soil. *Applied and Environmental Microbiology*, 64(1), 359-362.
- Katsoyiannis, A., and Samara, C. (2004). Persistent organic pollutants in the sewage treatment plant of Thessaloniki, northern Greece: occurrence and removal. *Water Research*, 38(11), 2685-2698.
- Keeler, J., and Wothers, P. (2008). *Chemical Structure and Reactivity, first edition*. Oxford University Press, UK, 27-46.
- Kelly, I., and Cerniglia, C.E. (1991). The metabolism of phenanthrene by species of *Mycobacterium*. *Journal of Industrial Microbiology*, 7(1), 19-26.

- Keshavarzifard, M., and Zakaria, M.P. (2015). Polycyclic aromatic hydrocarbons contamination of surface sediments from Port Dickson, Malaysia: Distribution, sources and ecological risk assessment. *Environmental Forensics*, 16(4), 322-332.
- Khanna, S., Lal, B., Chandra, R., Rajam, S., and Bajpai, U. (1998). *Enzymes and bioremediation*. In *Enzyme Chemistry: Impact and applications*. Suckling, C.J., Gibson, C.L., and Pitt, A.K., Eds., Blackie Academic and Professional, Chapman and Hall, London, 240.
- Khuri, A.I., and Mukhopadhyay, S. (2010). Response surface methodology. *Wiley Interdisciplinary Reviews: Computational Statistics*, 2(2), 128-149.
- Khuri, A.I. and Cornell, J.A. (1987). *Response Surfaces; Design and Analyses*. Marcel Dekker, Inc., New York, USA.
- Kim, Y.H., Freeman, J.P., Moody, J.D., Engesser, K.H., and Cerniglia, C.E. (2005). Effects of pH on the degradation of phenanthrene and pyrene by *Mycobacterium vanbaalenii* PYR-1. *Applied Microbiology and Biotechnology*, 67(2), 275-285.
- Kim, T.J., Lee, E.Y., Kim, Y.J., Cho, K.S., and Ryu, H.W. (2003). Degradation of polyaromatic hydrocarbons by *Burkholderia cepacia* 2A-12. *World Journal of Microbiology and Biotechnology*, 19(4), 411-417.
- Kim, S.J., Chun, J., Bae, K.S., and Kim, Y.C. (2000). Polyphasic assignment of an aromatic degrading *Pseudomonas* sp., strain DJ77, in the genus *Spingomonas* as *Spingomonas chungbukensis* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, 50(4), 1641-1647.
- Kimbara, K., Hashimoto, T., Fukuda, M., Koana, T., Takagi, M., Oishi, M., and Yano, K. (1989). Cloning and sequencing of two tandem genes involved in degradation of 2,3-dihydroxybiphenyl to benzoic acid in the polychlorinated biphenyl-degrading soil bacterium *Pseudomonas* sp. strain KKS102. *Journal of Bacteriology*, 171(5), 2740-2747.
- Kimura M. (1980). A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16(2), 111-120.
- Kitano, H. (2002). Systems biology; a brief overview. *Science*, 295(5560), 1662-1664.
- Kiyohara, H., Torigoe, S., Kaida, N., Asaki, T., Iida, T., Hayashi, H., and Takizawa, N. (1994). Cloning and characterization of a chromosomal gene cluster, pah, that encodes the upper pathway for phenanthrene and naphthalene utilization by *Pseudomonas putida* OUS82. *Journal of Bacteriology*, 176(8), 2439-2443.

- Kiyohara, H., Nagao, K., and Yana, K. (1982). Rapid screen for bacteria degrading water-insoluble, solid hydrocarbons on agar plates. *Applied and Environmental Microbiology*, 43(2), 454-457.
- Kiyohara, H., and Nagao, K. (1978). The catabolism of phenanthrene and naphthalene by bacteria. *Microbiology*, 105(1), 69-75.
- Kiyohara, H., and Nagao, K. (1977). Enzymatic conversion of 1-hydroxy-2-naphthoate in phenanthrene-grown *Aeromonas* sp. S45P1. *Agricultural and Biological Chemistry*, 41(4), 705-707.
- Kiyohara, H., Nagao, K., and Nomi, R. (1976). Degradation of phenanthrene through o-phthalate by an *Aeromonas* sp. *Agricultural and Biological Chemistry*, 40(6), 1075-1082.
- Knighton, W.B., and Grimsrud, E.P. (1983). Linearization of electron capture detector response to strongly responding compounds. *Analytical Chemistry*, 55(4), 713-718.
- Kröckel, L., and Focht, D.D. (1987). Construction of Chlorobenzene-utilizing recombinants by progenitive manifestation of a rare event. *Applied and Environmental Microbiology*, 53(10), 2470-2475.
- Kropp, K.G., and Fedorak, P.M. (1998). A review of the occurrence, toxicity, and biodegradation of condensed thiophenes found in petroleum. *Canadian Journal of Microbiology*, 44(7), 605-622.
- Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33(7), 1870-1874.
- Lakshmi, M.B, Muthukumar, K., and Velan, M. (2013). Optimization of minimal salt medium for efficient phenanthrene biodegradation by *Mycoplana* sp. MVMB2 isolated from petroleum contaminated soil using factorial design experiments. *CLEAN-Soil, Air, Water*, 41(1), 51-59.
- Lal, R. (1982). Accumulation, metabolism, and effects of organophosphorus insecticides on microorganisms. *Advances in Applied Microbiology*, 28, 149-200.
- Lal, R., and Saxena, D.M. (1982). Accumulation, metabolism, and effects of organochlorine insecticides on microorganisms. *Microbiological Reviews*, 46(1), 95.
- Lea, C. (2005). *Use of Crop-derived Raw Materials as Replacement for Mineral Oils in the Production of Metalworking Fluids: Challenges for the Lubricants Industry*. Fuchs Lubricants (UK) Pic. New Century Street, Hanley.

- Leadbetter, E.R., and Foster, J.W. (1960). Bacterial oxidation of gaseous alkanes. *Archives of Microbiology*, 35(1), 92-104.
- Leahy, J.G., and Colwell, R.R. (1990). Microbial degradation of hydrocarbons in the environment. *Microbiology Review*, 54(3), 305-315.
- Lee, J.Y., Jung, K.H., Choi, S.H., and Kim, H.S. (1995). Combination of the TOD and the TOL pathways in redesigning a metabolic route of *Pseudomonas putida* for the mineralization of a benzene, toluene, and p-xylene mixture. *Applied and Environmental Microbiology*, 61(6), 2211-2217.
- Leete, R. (2005). *Selangor's human development progress and challenges*. Malaysia UN resident coordinator, United Nations Development Programme, *Kuala Lumpur*.
- Leys, M.N., Bastiaens, L., Verstraete, W., and Springael, D. (2004). Influence of the carbon/nitrogen/phosphorus ratio on polycyclic aromatic hydrocarbons degradation by *Mycobacterium* and *Sphingomonas* in soil. *Applied Microbiology and Biotechnology*, 66(6), 726-736.
- Li, X., Li, P., Lin, X., Zhang, C., Li, Q., and Gong, Z. (2008). Biodegradation of aged polycyclic aromatic hydrocarbons (PAHs) by microbial consortia in soil and slurry phases. *Journal of Hazardous Materials*, 150(1), 21-26.
- Li, Q.Q., A. Loganath; Y.S. Chong and J.P. Obbard (2006). Persistent organic pollutants and adverse health effects in Humans. *Journal of Toxicology and Environmental Health*, 69(21), 1987-2005.
- Li, F., and Tan, T.C. (1994). Effect of heavy metal ions on the efficacy of a mixed *Bacilli* BOD sensor. *Biosensors and Bioelectronics*, 9(4-5), 315-324.
- Liang, D.W., Zhang, T., Fang, H.H., and He, J. (2008). Phthalates biodegradation in the environment. *Applied Microbiology and Biotechnology*, 80(2), 183-198.
- Lin, T.C., Pan, P.T., and Cheng, S.S. (2010). *Ex-situ* bioremediation of oil-contaminated soil. *Journal of Hazardous Materials*, 176(1), 27-34.
- Liu, J., Liu, S., Sun, K., Sheng, Y., Gu, Y., and Gao, Y. (2014). Colonization on root surface by a phenanthrene-degrading endophytic bacterium and its application for reducing plant phenanthrene contamination. *PloS one*, 9, e108249.
- Liu, K., Han, W., Pan, W.P., and Riley, J.T. (2001). Polycyclic aromatic hydrocarbon emissions from a coal fired pilot FBC system. *Journal of Hazardous Material*, 84(2), 175-188.
- Liu, B.L., and Tzeng, Y.M., (1998). Optimization of growth medium for production of spores from *Bacillus thuringiensis* using response surface methodology. *Bioprocess Engineering* 18(6), 413-418.

- Lloyd, C.A., and Cackette, T.A. (2001). Diesel Engines: Environmental Impact and Control. *Air and Waste Management Association*, 51, 805-847.
- Long, R.M. (2008). *The distribution and diversity of polycyclic aromatic compound-degrading bacteria and key degradative genes*. PhD dissertation, University of Exeter, UK.
- Lors, C., Ryngaert, A., Périé, F., Diels, L., and Damidot, D. (2010). Evolution of bacterial community during bioremediation of PAHs in coal tar contaminated soil. *Chemosphere*, 81(10), 1263-1271.
- Lu, J., Guo, C., Zhang, M., Lu, G., and Dang, Z. (2014). Biodegradation of single pyrene and mixtures of pyrene by a fusant bacterial strain F14. *International Biodeterioration and Biodegradation*, 87, 75-80.
- Lu, M., Xu, K., and Chen, J. (2013). Effect of pyrene and cadmium on microbial activity and community structure in soil. *Chemosphere*, 91(4), 491-497.
- Lundstedt, S., White, P.A., Lemieux, C.L., Lynes, K.D., Lambert, I.B., Öberg, L., and Tysklind, M. (2007). Sources, fate, and toxic hazards of oxygenated polycyclic aromatic hydrocarbons at PAH-contaminated sites. *AMBIO: A Journal of the Human Environment*, 36(6), 475-485.
- Luthy, R.G., Dzombak, D.A., Peters, C.A., Roy, S.B., Ramaswami, A., Nakles, D.V., and Nott, B.R. (1994). Remediating tar-contaminated soils at manufactured gas plant sites, *Environmental Science and Technology*, 28(6), 266-276.
- Madigan, M.T., J.M. Martinko and J. Parker (1998). *Brock: Biología de los Microorganismos*. Prentice Hall, España, 726.
- MacDougall, D., and Crummett, W.B. (1980). Guidelines for data acquisition and data quality evaluation in environmental chemistry. *Analytical Chemistry*, 52(14), 2242-2249.
- Maier, A., Schumann, B.L., Chang, X., Talaska, G., and Puga, A. (2002). Arsenic co-exposure potentiates benzo [a] pyrene genotoxicity. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 517(1), 101-111.
- Malaysia Environmental Quality Report, MEQR (2015). Ministry of Natural Resources and Environment Malaysia, Department of Environment.
- MEQR, (2014). Ministry of Natural Resources and Environment Malaysia, Department of Environment.
- MEQR, (2012). Ministry of Natural Resources and Environment Malaysia, Department of Environment.
- MEQR, (2010). Ministry of Natural Resources and Environment Malaysia, Department of Environment.

- Maliszewska-K.B., and Smreczak, B. (2003). Habitat functions of agricultural soils as affected by heavy metals and polycyclic aromatic hydrocarbons contamination. *Environment International*, 28(8), 719-728.
- Maliszewska, K.B. (1999). Sources, concentration, fate and effects of polycyclic aromatic hydrocarbons in the environment. Part A: PAHs in Air, *Polish Journal of Environmental Studies*, 8, 131-136.
- Malveda, M.P. (2015). *Chemical Economics Handbook Report on Plasticizers*. IHS Market Ltd, UK.
- Mandri, T., and Lin, J. (2007). Isolation and characterization of engine oil degrading indigenous microorganisms in Kwazulu-Natal, South Africa. *African Journal of Biotechnology*, 6(1), 23-27.
- Manoli, E., and Samara, C. (2008). The removal of polycyclic aromatic hydrocarbons in the wastewater treatment process: experimental calculations and model predictions. *Environmental Pollution*, 151(3), 477-485.
- Margesin, R. Labbe, D. Schninner, F. Greer, C.W. and Whyte, L.G. (2003). Characterisation of hydrocarbon degrading microbial populations in contaminated and pristine contaminated soils. *Applied and Environmental Microbiology*, 69(6), 3985-3092.
- Margesin, R., and Schinner, F. (2001). Biodegradation and bioremediation of hydrocarbons in extreme environments. *Applied Microbiology and Biotechnology*, 56(5), 650-663.
- Markowicz, A., Płaza, G., and Piotrowska-Seget, Z. (2016). Activity and functional diversity of microbial communities in long-term hydrocarbon and heavy metal contaminated soils. *Archives of Environmental Protection*, 42(4), 3-11.
- Martinek, R. (1969). Practical Clinical Enzymology: *Journal of American Medical Technologist*, 31, 162.
- Maslin, P., and Maier, R.M. (2000). Rhamnolipid enhanced mineralization of phenanthrene in organic-metal co-contaminated soils. *Bioremediation Journal*, 4(4), 295-308.
- McMeekin, T.A., Olley, J., and Ratkowsky, D.A. (1988). Temperature effects on bacterial growth rates. *Physiological Models in Microbiology*, 1, 75-89.
- Mesdhaghinia, A.R., Nasser, S., Arbabi, M., and Rezaie, S. (2005). Isolation of polycyclic aromatic hydrocarbon degrading bacteria associated with the petroleum contaminated soils in Iran. *Proceedings of the 9th International Conference on Environment Science and Technology*, 984-991.

- Megharaj, M., Ramakrishnan, B., Venkateswarlu, K., Sethunathan, N., and Naidu, R. (2011). Bioremediation approaches for organic pollutants: a critical perspective. *Environment International*, 37(8), 1362-1375.
- Michael, G.O. (2010). *Occupational Safety and Health Administration*. The Encyclopedia of Earth, Cutler Cleveland.
- Micó, C., Recatalá, L., Peris, M., and Sánchez, J. (2006). Assessing heavy metal sources in agricultural soils of European Mediterranean area by multivariate analysis. *Chemosphere*, 65(5), 863-872.
- Mielke, H.W., Gonzales, C.R., Smith, M.K., and Mielke, P.W. (2000). Quantities and associations of lead, zinc, cadmium, manganese, chromium, nickel, vanadium, and copper in fresh Mississippi delta alluvium and New Orleans alluvial soils. *Science of the Total Environment*, 246(2), 249-259.
- Mikkonen, A. (2012). *The Potential of Microbial Ecological Indicators to Guide Ecosophisticated Management of Hydrocarbon-Contaminated Soils*. PhD dissertation, University Helsinki, Finland.
- Miller, J.S., and Olejnik, D. (2004). Ozonation of polycyclic aromatic hydrocarbons in water solution. *Ozone: Science and Engineering*, 26(5), 453-464.
- Miller, C.D., Hall, K., Liang, Y.N., Nieman, K., Sorensen, D., and Issa, B. (2004). Isolation and characterization of polycyclic aromatic hydrocarbon degrading *Mycobacterium* isolates from soil. *Microbial Ecology*, 48(2), 230-238.
- Mishra, V., Lal, R., and Srinivasan. (2001). Enzymes and operons mediating xenobiotic degradation in bacteria. *Critical Reviews in Microbiology*, 27(2), 133-166.
- Mishra, V. (1997). *Degradation of Phenol and Benzoate by Bacteria; Cloning and Sequencing of a DNA Fragment of a Gene Encoding Hexachlorocyclohexane Dehydrochlorinase in Sphingomonas paucimobilis*. Ph.D. Thesis, University of Delhi, India.
- Moghadam, M.S., Ebrahimipour, G., Abtahi, B., and Ghassempour, A. (2013). Isolation, identification and optimization of phenanthrene degrading bacteria from the coastal sediments of nayband bay. *Jundishapur Journal of Microbiology*, 6(9), 1-8.
- Mohandass, R., Rout, P., Jiwal, S., and Sasikala, C. (2012). Biodegradation of Benzo[a]pyrene by the mixed culture of *Bacillus cereus* and *Bacillus vireti* isolated from the petrochemical industry. *Journal of Environmental Biology*, 33(6), 985-989.
- Mohd-Kamil, N.A.F., Hussain, N.H., Mizad, M.B., and Abdul-Talib, S. (2014). Enhancing performance of *Sphingobacterium spiritivorum* in bioremediation phenanthrene contaminated sand. *Remediation Journal*, 24(3), 119-128.

- Mohd Radzi, N.A.S., Abu Bakar, N.K., Emenike, C.U., and Abas, M.R. (2016). Polycyclic aromatic hydrocarbons: contamination level and risk assessment in urban areas, Kuala Lumpur, Malaysia. *Desalination and Water Treatment*, 57(1), 171-190.
- Mohn, W., and Stewart, R.G. (2000). Limiting factors for hydrocarbon biodegradation at low temperatures in Arctic soils. *Soil Biology and Biochemistry*, 32(8), 1161–1172.
- Montgomery, D.C. (2005). *Design and Analysis of Experiments: Response Surface Method and Designs*. John Wiley and Sons, Inc, New Jersey, USA.
- Montagnolli, R.N., Lopes, P.R.M., and Bidoia, E.D. (2015). Screening the toxicity and biodegradability of petroleum hydrocarbons by a rapid colorimetric method. *Archives of Environmental Contamination and Toxicology*, 68(2), 342-353.
- Mokross, H., Schmidt, E., and Reineke, W. (1990). Degradation of 3-chlorobiphenyl by *in vivo* constructed hybrid *Pseudomonads*. *FEMS Microbiology Letters*, 71(1-2), 179-185.
- Moody, J.D., Freeman, J.P., Doerge, D.R., and Cerniglia, C.E. (2001). Degradation of phenanthrene and anthracene by cell suspensions of *Mycobacterium* sp. strain PYR-1. *Applied and Environmental Microbiology*, 67(4), 1476-1483.
- Moore, J.W., and Stanitski, C.L. (2011). *Chemistry: The molecular science, 4th edition*. Cole Cengage Learning, USA.
- Moraes, B., Eduardo, T., and Sania, T.M (2009). Biodegradation of oil refinery residues using mixed-culture of microorganisms isolated from land farming. *Brazilian Archives of Biology and Technology*, 52(6), 1571-1578.
- Moret, S., Dyson, P.J., and Laurency, G. (2014). Direct synthesis of formic acid from carbon dioxide by hydrogenation in acidic media. *Nature Communications*, 5.
- Morgan, P., and Atlas, R.M. (1989). Hydrocarbon degradation in soils and methods for soil biotreatment. *Critical Reviews in Biotechnology*, 8(4), 305-333.
- Mueller, K.E., and Shann, J.R. (2006). Polycyclic aromatic hydrocarbons dissipation in spiked soil: Impacts of bioavailability, microbial activity, and trees. *Chemosphere*, 64(6), 1006-1014.
- Nagoba, B.S., Selkar, S.P., Wadher, B.J., and Gandhi, R.C. (2013). Acetic acid treatment of *Pseudomonas* wound infections - a review. *Journal of Infection and Public Health*, 6(6), 410-415.
- Nakai, C., Horiike, K., Kuramitsu, S., Kagamiyama, H., and Nozaki, M. (1990). Three isozymes of catechol 1,2-dioxygenase (pyrocatechase), alpha alpha, alpha beta, and beta beta, from *Pseudomonas arvilla* C-1. *Journal of Biological Chemistry*, 265(2), 660-665.

- Nasrollahzadeh, H.S., Najafpour, G.D., and Aghamohammadi, N. (2007). Biodegradation of phenanthrene by mixed culture consortia in batch bioreactor using central composite face-entered design. *International Journal of Environmental Research*, 1(2), 80-87.
- Nicholson, J.K., Holmes, E., Lindon, J.C., and Wilson, I.D. (2004). The challenges of modeling mammalian biocomplexity. *Nature Biotechnology*, 22, 1268-1274.
- NIST, (2017). Certificate of Analysis SRM 2587 trace elements in soil containing lead from paint (Nominal 3000 mg/kg Lead), January, 2017.
- Neddermeijer, H.G., Van Oortmarssen, G.J., Piersma, N., and Dekker, R. (2000). A framework for response surface methodology for simulation optimization. In *Proceedings of the 32nd conference on Winter simulation, Society for Computer Simulation International*, 129-136.
- Neelam, M., Nawaz, Z., and Riazuddin, S. (1987). Hydrocarbon biodegradation: Biochemical characterization of bacteria isolated from local soils. *Pakistan Journal of Scientific and Industrial Research*, 30(5), 382-385.
- Neidle, E.L., Hartnett, C., Ornston, L.N., Bairoch, A., Rekik, M., and Harayama, S. (1991). Nucleotide sequences of the *Acinetobacter calcoaceticus* benABC genes for benzoate 1,2-dioxygenase reveal evolutionary relationships among multicomponent oxygenases. *Journal of Bacteriology*, 173(17), 5385-5395.
- Nestler, C., Hansen, L., Ringelberg, D., and Talley, J. (2001). Remediation of soil PAH: comparison of biostimulation and bioaugmentation. In: *Sixth International in situ and on site Bioremediation Symposium*, 43-50.
- Nwinyi, O., Picardal, F.W., An, T.T., and Amund, O.O. (2013). Aerobic degradation of naphthalene, fluoranthene, pyrene and chrysene using indigenous strains of bacteria isolated from a former industrial site. *Canadian Journal of Pure and Applied Sciences*, 7, 2303-2314.
- Oa, S. W., and Lee, T. G. (2009). Investigation of soil pollution status for railroad depot. *Railway Bulletin*, 12(5), 788-792.
- Odu, C.T.I. (1977). *Oil Degradation and Microbiological Changes in Soils Deliberately Contaminated with Petroleum Hydrocarbons*. Institute of Petroleum (IP), UK.
- Oehlert, G.W. (2000). *Design and Analysis of Experiments: Response Surface Design*. W.H. Freeman and Company, New York, USA.
- Ogram, A.V., Jessup, R.E., Ou, L.T., and Rao, P.S.C. (1985). Effects of sorption on biological degradation rates of (2,4-dichlorophenoxy) acetic acid in soils. *Applied and Environmental Microbiology*, 49(3), 582-587.

- Okuda, T., Naoi, D., Tenmoku, M., Tanaka, S., He, K., Ma, Y. and Zhang, D. (2006). Polycyclic aromatic hydrocarbons in the aerosol in Beijing, China, measured by aminopropylsilane chemically-bonded stationary-phase column chromatography and HPLC/fluorescence detection. *Chemosphere*, 65(3), 427-435.
- Onakpoya, I., Hunt, K., Wider, B. and Ernst, E. (2014). Pyruvate supplementation for weight loss: a systematic review and meta-analysis of randomized clinical trials. *Critical Reviews in Food Science and Nutrition*, 51, 17-23.
- Pagnout, C., Frache, G., Poupin, P., Maunit, B., Muller, J.F., and Féraud, J.F. (2007). Isolation and characterization of a gene cluster involved in PAH degradation in *Mycobacterium* sp. strain SNP11: Expression in *Mycobacterium smegmatis* mc 2 155. *Research in Microbiology*, 158(2), 175-186.
- Park, S.W., Lee, J.Y., Kim, K.J., Yang, J.S., and Baek, K. (2010). Alkaline enhanced separation of waste lubricant oils from railway contaminated soil. *Separation Science and Technology*, 45(12-13), 1988-1993.
- Park, S.W., Lee, J.Y., Yang, J.S., Kim, K.J., and Baek, K. (2009). Electrokinetic remediation of contaminated soil with waste-lubricant oils and zinc. *Journal of Hazardous Materials*, 169(1), 1168-1172.
- Pasco, N., Baronian, K., Jeffries, C., and Hay, J. (2000). Biochemical mediator demand, a novel rapid alternative for measuring biochemical oxygen demand. *Applied Microbiology and Biotechnology*, 53(5), 613-618.
- Patri, M., Padmini, A., and Babu, P. P. (2010): Polycyclic aromatic hydrocarbons in air and their neurotoxic potency in association with oxidative stress: a brief perspective. *Annals of Neurosciences*, 16(1), 22-30.
- Paul, E.A. (2014). *Soil Microbiology, Ecology and Biochemistry*. Academic Press, an imprint of Elsevier USA.
- Pawar, R.M. (2015). The Effect of Soil pH on Bioremediation of PAHs. *Journal of Bioremediation and Biodegradation*, 6, 1-14.
- Pereira, P.A.D.P., Lopes, W.A., Carvalho, L.S., da Rocha, G.O., de Carvalho Bahia, N., Loyola, J., and de Andrade, J.B. (2007). Atmospheric concentrations and dry deposition fluxes of particulate trace metals in Salvador, Bahia, Brazil. *Atmospheric Environment*, 41(36), 7837-7850.
- Perez, J. (2012). *The Soil Remediation Industry in Europe: The Recent, Past and Future Perspectives*. Ernst and Young Et Associates, 1-2 Place des Saisons, Courbevoie, 92400, France, 2-22.
- Perry, J.J. (1979). Microbial cooxidations involving hydrocarbons. *Microbiological Reviews*, 43(1), 59.

- Peter Atkins, J.D.P. (2010). *Atkins' Physical Chemistry*. Oxford, 443.
- Peter, H.A. (2003). Petroleum and individual polycyclic aromatic hydrocarbon In D.J. Hoffman, B.A. Rattner, G.A. Buston, J. Cairns editors. *Handbook of Ecotoxicology*. CRC Press, Taylor and Francis USA, 342, 359.
- Pinyakong, O., Habe, H., Yoshida, T., Nojiri, H., and Omori, T. (2003). Identification of three novel salicylate 1-hydroxylases involved in the phenanthrene degradation of *Sphingobium* sp. strain P2. *Biochemical and Biophysical Research Communications*, 301(2), 350-357.
- Plaza, G.A., Lukasik, K., Wypych, J., Nalecz-Jawecki, G., Berry, C., and Brigmon, R.L. (2008). Biodegradation of crude oil and distillation products by biosurfactant-producing bacteria. *Polish Journal of Environmental Studies*, 17(1), 87.
- Prabhu, Y., and Phale, P.S. (2003). Biodegradation of phenanthrene by *Pseudomonas* sp. strain PP2: novel metabolic pathway, role of biosurfactant and cell surface hydrophobicity in hydrocarbon assimilation. *Applied Microbiology and Biotechnology*, 61(4), 342-351.
- Prasad, M.N.V., and Katiyar, S.C. (2010). Drill cuttings and fluids of fossil fuel exploration in north-eastern India: environmental concern and mitigation options. *Current Science*, 98(12), 1567.
- Rahman, K.S.M., Thahira-Rahman, J., Lakshmanaperumalsamy, P., and Banat, I.M. (2002). Towards efficient crude oil degradation by a mixed bacterial consortium. *Bioresource Technology*, 85(3), 257-261.
- Raja, P., Uma, S., Gopal, H., and Govindarajan, K. (2006). Impact of bio inoculants consortium on rice root exudates, biological nitrogen fixation and plant growth. *Journal Biological Science*, 6, 815-823.
- Ramadan, M.A., El-Tayeb, O.M., and Alexander, M. (1990). Inoculum size as a factor limiting success of inoculation in biodegradation. *Applied and Environmental Microbiology*, 56(5), 1392-1396.
- Raymond, R.L., Hudson, J.O., and Jamison, V.W. (1976). Oil degradation in soil. *Applied and Environmental Microbiology*, 31(4), 522-535.
- Ravindra, K., Sokhi, R., and Van Grieken, R. (2008). Atmospheric polycyclic aromatic hydrocarbons: source attribution, emission factors and regulation. *Atmospheric Environment*, 42(13), 2895-2921.
- Rebola, M., Silva, M.J., Louro, H., Antunes, A.M., José, S., Rebelo, M.H., and Cardoso, A.S. (2008). Chlorinated polycyclic aromatic hydrocarbons associated with drinking water chlorination - preparation, quantification and genotoxicity characterization. *Proceedings of ECOpole*, 2, 373-379.

- Renner, R. (1999). EPA to strengthen persistent, bioaccumulative, and toxic pollutant controls - mercury first to be targeted. *Environmental Science and Technology*, 33, 62A.
- Richnow, H.H., Seifert, R., Hefter, J., Kästner, M., Mahro, B., and Michaelis, W. (1994). Metabolites of xenobiotica and mineral oil constituents linked to macromolecular organic matter in polluted environments, *Organic Geochemistry*, 22(3-5), 671IN9-681IN10.
- Rijnaarts, H.H., Bachmann, A., Jumelet, J.C., and Zehnder, A.J. (1990). Effect of desorption and intra-particle mass transfer on the aerobic biomineralization of alpha-hexachlorocyclohexane in a contaminated calcareous soil. *Environmental Science and Technology*, 24(9), 1349-1354.
- Rodrigues, L., Teixeira, J., Oliveira, R., and Van Der Mei, H.C. (2006). Response surface optimization of the medium components for the production of biosurfactants by probiotic bacteria. *Process Biochemistry*, 41(1), 1-10.
- Rojo, F., Pieper, D.H., Engesser, K.H., Knackmuss, H.J., and Timmis, K.N. (1987). Assemblage of ortho cleavage route for simultaneous degradation of chloro- and methylaromatics. *Science*, 238(4832), 1395-1398.
- Rosen, K.H. (2007). Discrete mathematics and its applications. *Ars Mathematica Contemporanea (AMC)*, 10, 12.
- Rubio-Clemente, A., Torres-Palma, R.A., and Peñuela, G.A. (2014). Removal of polycyclic aromatic hydrocarbons in aqueous environment by chemical treatments: a review. *Science of the Total Environment*, 478, 201-225.
- Salmimies, R., Vehmaanperä, P., and Häkkinen, A. (2016). Acidic dissolution of magnetite in mixtures of oxalic and sulfuric acid. *Hydrometallurgy*, 163, 91-98.
- Samanta, S.K., Singh, O.V., and Jain, R.K. (2002). Polycyclic aromatic hydrocarbons: environmental pollution and bioremediation. *TRENDS in Biotechnology*, 20(6), 243-248.
- Samanta, S.K., Chakraborti, A.K., and Jain, R.K. (1999). Degradation of phenanthrene by different bacteria: evidence for novel transformation sequences involving the formation of 1-naphthol. *Applied Microbiology and Biotechnology*, 53(1), 98-107.
- Sandrin, T.R., and Maier, R.M. (2003). Impact of metals on the biodegradation of organic pollutants. *Environmental Health Perspectives*, 111(8), 1093.
- Sany, S.B.T., Hashim, R., Salleh, A., Rezayi, M., Mehdinia, A., and Safari, O. (2014). Polycyclic aromatic hydrocarbons in coastal sediment of Klang Strait, Malaysia: distribution pattern, risk assessment and sources. *PloS one*, 9, e94907.

- Sarkar, P., Meghvanshi, M., and Singh, R. (2011). Microbial consortium: A new approach in effective degradation of organic kitchen wastes. *International Journal of Environmental Science and Development*, 2(3), 170.
- Sarma, H., Islam, N.F., Borgohain, P., Sarma, A., and Prasad, M.N.V. (2016). Localization of polycyclic aromatic hydrocarbons and heavy metals in surface soil of Asia's oldest oil and gas drilling site in Assam, north-east India: implications for the bio-economy. *Emerging Contaminants*, 2(3), 119-127.
- Sarma, P.M., Bhattacharya, D., Krishnan, S., and Lal, B. (2004). Degradation of polycyclic aromatic hydrocarbons by a newly discovered enteric bacterium, *Leclercia adecarboxylata*. *Applied and Environmental Microbiology*, 70(5), 3163-3166.
- Sathishkumar, M., Binupriya, A.R., Baik, S.H., and Yun, S.E. (2008). Biodegradation of crude oil by individual bacterial strains and a mixed bacterial consortium isolated from hydrocarbon contaminated areas. *CLEAN–Soil, Air, Water*, 36(1), 92-96.
- Sayara, T.A., Adroguer, S., and Sánchez Ferrer, A. (2011). *Bioremediation of polycyclic aromatic hydrocarbons contaminated soil*. PhD Thesis, Universitat Autònoma de Barcelona, Brazil.
- Schwartz, E., Trinh, S.V., and Scow, K.M. (2000). Measuring growth of a phenanthrene degrading bacterial inoculum in soil with a quantitative competitive polymerase chain reaction method. *FEMS Microbiology Ecology*, 34(1), 1-7.
- Scow, K.M., and Hutson, J. (1992). Effect of diffusion and sorption on the kinetics of biodegradation: theoretical considerations. *Soil Science Society of America Journal*, 56(1), 119-127.
- Scow, K.M., and Alexander, M. (1992). Effect of diffusion on the kinetics of biodegradation: experimental results with synthetic aggregates. *Soil Science Society of America Journal*, 56(1), 128-134.
- Seo, J.S., Keum, Y.S., and Li, Q.X. (2009). Bacterial degradation of aromatic compounds. *International Journal of Environmental Research and Public Health*, 6(1), 278-309.
- Seo, J.S., Keum, Y.S., Hu, Y., Lee, S.E., and Li, Q.X. (2007). Degradation of phenanthrene by *Burkholderia* sp. C3: initial 1,2- and 3,4-dioxygenation and meta- and ortho-cleavage of naphthalene-1,2-diol. *Biodegradation*, 18(1), 123-131.

- Seo, J.S., Keum, Y.S., Hu, Y., Lee, S.E., and Li, Q.X. (2006). Phenanthrene degradation in *Arthrobacter* sp. P1-1: initial 1, 2, 3, 4 and 9, 10-dioxygenation, and meta-and ortho-cleavages of naphthalene-1,2-diol after its formation from naphthalene-1,2-dicarboxylic acid and hydroxyl naphthoic acid. *Chemosphere*, 65(11), 2388-2394.
- Seo, J.S. (2006). *Bacterial Proteomes and Metabolism of Aromatic Compounds*. Ph.D. dissertation, University of Hawaii Manoa, USA.
- Shaohua, C., Jianjun, L., Meiyang, H., Kaiping, L., Peng, G., and Huasheng, H. (2012). Enhancement of cypermethrin degradation by a co-culture of *Bacillus cereus* ZH-3 and *Streptomyces aureus* HP-S-01. *Bioresource Technology*, 110, 97-104.
- Shaohua, C., Qiongbo, H., Meiyang, H., Jianjun, L., Qunfang, W., and Kaiping, L. (2011). Isolation and characterization of a fungus able to degrade pyrethroids and 3 phenoxybenzaldehyde. *Bioresource Technology*, 102(17), 8110-8116.
- Shen, G., Lu, Y., and Hong, J. (2006). Combined effect of heavy metals and polycyclic aromatic hydrocarbons on urease activity in soil. *Ecotoxicology and Environmental Safety*, 63(3), 474-480.
- Shen, G., Cao, L., Lu, Y., and Hong, J. (2005). Influence of phenanthrene on cadmium toxicity to soil enzymes and microbial growth (5 pp). *Environmental Science and Pollution Research*, 12(5), 259-263.
- Sheng, X., Xiaobing C., and Linyan H. (2008). Characteristics of endophytic pyrene degrading bacterium of *Enterobacter* sp. 12J1 from *Allium macrostemon* Bunge. *International Biodeterioration and Biodegradation* 62(2), 88-95.
- Sherman, J.M., and Holm, G.E. (1922). Salt effects in bacterial growth: The growth on *Bacteria coli* in Relation to H⁺ ion concentration. *Journal of Bacteriology*, 7(5), 465.
- Shiaris, M.P. (1989). Seasonal biotransformation of naphthalene, phenanthrene, and Benzo[a]pyrene in surficial estuarine sediments. *Applied Environmental Microbiology*, 55(6), 1391-1399
- Sihag, S., Pathak, H., and Jaroli, D.P. (2014). Factors affecting the rate of biodegradation of polyaromatic hydrocarbons. *International Journal of Pure and Applied Biological Sciences*, 2(3), 185-202.
- Sikkema, J., De Bont, J.A., and Poolman, B. (1995). Mechanisms of membrane toxicity of hydrocarbons. *Microbiological Reviews*, 59(2), 201-222.
- Silva, C.J.S.M., and Roberto, I.C., (2001). Optimization of xylitol production by *Candida guilliermondii* FTI 20037 using response surface methodology. *Process Biochemistry*, 36(11), 1119-1124.

- Silva, A.S., Camargo, F.A.D.O., Andreatza, R., Jacques, R.J.S., Baldoni, D.B., and Bento, F.M. (2012). Enzymatic activity of catechol 1,2-dioxygenase and catechol 2, 3-dioxygenase produced by *Gordonia polyisoprenivorans*, *Quimica Nova*, 35(8), 1587-1592.
- Simarro, R., González, N., Bautista, L.F., Molina, M.C., and Schiavi, E. (2012). Evaluation of the influence of multiple environmental factors on the biodegradation of dibenzofuran, phenanthrene, and pyrene by a bacterial consortium using an orthogonal experimental design. *Water, Air and Soil Pollution*, 223(6), 3437-3444.
- Simarro, R., González, N., Bautista, L.F., Sanz, R., and Molina, M.C. (2011). Optimisation of key abiotic factors of PAH (naphthalene, phenanthrene and anthracene) biodegradation process by a bacterial consortium. *Water, Air and Soil Pollution*, 217(1-4), 365-374.
- Singh, A., and Ward, O.P. (2004). Biotechnology and bioremediation; an overview. In *Biodegradation and Bioremediation*, Springer Berlin Heidelberg, Germany, 1-17.
- Singh, S.N. and R.D. Tripathi (2007). *Environmental Bioremediation Technologies*. Springer-Verlag, Berlin Heidelberg, Germany.
- Skoog, D.A., Holler, F.J., and Crouch, S.R. (2007). *Principles of Instrumental Analysis*. Thomson Brooks/Cole, USA.
- Sogani, M., Mathur, N., Sharma, P., and Bhatnagar, P. (2012). Comparison of immobilized whole resting cells in different matrices vis-a-vis free cells of *Bacillus megaterium* for acyltransferase activity. *Journal of Environmental Research and Development*, 6(3A), 695-701.
- Sokhn, J., De Leij, F.A.A.M., Hart, T.D., and Lynch, J.M. (2001). Effect of copper on the degradation of phenanthrene by soil micro-organisms. *Letters in Applied Microbiology*, 33(2), 164-168.
- Song, H.G., and Bartha, R. (1990). Effects of jet fuel spills on the microbial community of soil. *Applied and Environmental Microbiology*, 56(3), 646-651.
- Spain, J.C., and van Veld, P.A. (1983). Adaptation of natural microbial communities to degradation of xenobiotic compounds: effects of concentration, exposure time, inoculum, and chemical structure. *Applied and Environmental Microbiology*, 45(2), 428-435.
- Speight, J.G. (2015). *Handbook of petroleum product analysis*. John Wiley and Sons, New Jersey, USA.
- Speight, J.G., and Arjoon, K.K. (2012). *Bioremediation of petroleum and petroleum products*. John Wiley & Sons, Hoboken, New Jersey USA, 443, 305-359.

- Spinelli, L., and Freitas, D.E. (2005). *Bioremediation, Toxicity and Cell Damage in gasoline spills* (Ph.D Thesis); Federal University of Rio Grande do Sul, Porto Alegre, Brazil.
- Springael, D., Kreps, S., and Mergeay, M. (1993). Identification of a catabolic transposon, Tn4371, carrying biphenyl and 4-chlorobiphenyl degradation genes in *Alcaligenes eutrophus* A5. *Journal of Bacteriology*, 175(6), 1674-1681.
- Srinivas, M.R.S., Chand, N., and Lonsane, B.K. (1994). Use of Plackett-Burman design for rapid screening of several nitrogen sources, growth/ product promoters, minerals, and enzyme inducers for the production of α -galactosidase by *Aspergillus niger* MRSS 234 in solid state fermentation. *Bioprocess Engineering*, 10(3), 139-144.
- Srogi, K. (2007). Monitoring of environmental exposure to polycyclic aromatic hydrocarbons: a review. *Environmental Chemistry Letters*, 5(4), 169-195.
- Stankovic S. and Stankovic A.R. (2013). Bioindicators of toxic metals. In E. Lichtfouse, J. Schwarzbauer, D. Robert 2013, *Green materials for energy, products and depollution*, Springer, Dordrecht, pp. 151–228.
- Starzecka, A., and Bednarz, T. (1993). Comparison of development and metabolic activity of algae and bacteria in soils under the influence of short-and long-term contamination with metallurgic industrial dusts. *Algological Studies/Archiv für Hydrobiologie, Supplement*, 71-88.
- Stingley, R.L., Khan, A.A., and Cerniglia, C.E. (2004). Molecular characterization of a phenanthrene degradation pathway in *Mycobacterium vanbaalenii* PYR-1. *Biochemical and Biophysical Research Communications*, 322(1), 133-146.
- Story, S.P., Parker, S.H., Hayasaka, S.S., Riley, M.B., and Kline, E.L. (2001). Convergent and divergent points in catabolic pathways involved in utilization of fluoranthene, naphthalene, anthracene, and phenanthrene by *Sphingomonas paucimobilis* var. EPA505. *Journal of Industrial Microbiology and Biotechnology*, 26(6), 369-382.
- Sudipt, K.S., Om, V.S., and Rakesh, K.J. (2002). Polycyclic aromatic hydrocarbons: environmental pollution and bioremediation. *Trends in Biotechnology*, 20(6), 243–248.
- Sun, G.D., Jin, J.H., Xu, Y., Zhong, Z.P., Liu, Y., and Liu, Z.P. (2014a). Isolation of high molecular weight PAH-degrading strain and enhancing removal of HMW-PAHs from heavily contaminated soil. *International Biodeterioration and Biodegradation*, 90, 23-28.
- Sun, K., Liu, J., Gao, Y., Jin, L., Gu, Y., and Wang, W. (2014b). Isolation, plant colonization potential, and phenanthrene degradation performance of the endophytic bacterium *Pseudomonas* sp. Ph6-gfp. *Scientific Reports*, 4, 10-1038.

- Sun, M., Luo, Y., Teng, Y., Jia, Z., Li, Z., and Deng, S. (2013). Remediation of polycyclic aromatic hydrocarbon and metal contaminated soil by successive methyl- β -cyclodextrin-enhanced soil washing–microbial augmentation: a laboratory evaluation. *Environmental Science and Pollution Research*, 20(2), 976-986.
- Swindoll, C.M., Aelion, C.M., and Pfaender, F.K. (1988). Influence of inorganic and organic nutrients on aerobic biodegradation and on the adaptation response of subsurface microbial communities. *Applied and Environmental Microbiology*, 54(1), 212-217.
- Szabó, K.É., Itor, P.O., Bertilsson, S., Tranvik, L., and Eiler, A. (2007). Importance of rare and abundant populations for the structure and functional potential of freshwater bacterial communities. *Aquatic Microbial Ecology*, 47(1), 1-10.
- Taira, K., Hirose, J., Hayashida, S., and Furukawa, K. (1992). Analysis of bph operon from the polychlorinated biphenyl-degrading strain of *Pseudomonas alcaligenes* KF707. *Journal of Biological Chemistry*, 267(7), 4844-4853.
- Tao, X.Q., Lu, G.N., Liu, J.P., Li, T., and Yang, L.N. (2009). Rapid degradation of phenanthrene by using *Sphingomonas* sp. GY2B immobilized in calcium alginate gel beads. *International Journal of Environmental Research and Public Health*, 6(9), 2470-2480.
- Tao, X.Q., Lu, G.N., Dang, Z., Yang, C., and Yi, X.Y. (2007). A phenanthrene-degrading strain *Sphingomonas* sp. GY2B isolated from contaminated soils. *Process Biochemistry*, 42(3), 401-408.
- Tatusova, T.A., and Madden, T.L. (1999). BLAST 2 Sequences, a new tool for comparing protein and nucleotide sequences. *FEMS Microbiology Letters*, 174(2), 247-250.
- Tenge, A.J., Kaihura, F.B.S., Lal, R., and Singh, B.R. (1998). Diurnal soil temperature fluctuations for different erosion classes of an oxisol at Mlingano, Tanzania. *Soil and Tillage Research*, 49, 211-217.
- Texas Research Institute, TRI (1982). *Enhancing the microbial degradation of underground gasoline by increasing available oxygen*. American Petroleum Institute, 4428, 1-25.
- Thavamani, P., Malik, S., Beer, M., Megharaj, M., and Naidu, R. (2012). Microbial activity and diversity in long-term mixed contaminated soils with respect to polyaromatic hydrocarbons and heavy metals. *Journal of Environmental Management*, 99, 10-17.
- Thavamani, P., Megharaj, M., Krishnamurti, G.S.R., McFarland, R., and Naidu, R. (2011). Finger printing of mixed contaminants from former manufactured gas plant (MGP) site soils: implications to bioremediation. *Environment International*, 37(1), 184-189.

- Thavasi, R., Jayalakshmi, S., Balasubramanian, T., and Banat, I.M. (2007). Effects of salinity, temperature, pH and crude oil concentration on biodegradation of crude oil by *Pseudomonas aeruginosa*. *Journal of Biological and Environmental Sciences*, 1(2), 51-57.
- Thomas, J., and Ward, C.H. (1989). *In situ* bioremediation of organic contaminants in the subsurface. *Environmental Science Technology*, 23(7), 760-765.
- Thomas, J.M., Lee, M.D., Bedient, P.B., Borden, R.C., and Canter, L.W. (1987). *Leaking underground storage tanks: remediation with emphasis on in situ bioremediation*. Rice University Houston, Texas, USA.
- Tian, L., Ma, P., and Zhong, J.J. (2002). Kinetics and key enzyme activities of phenanthrene degradation by *Pseudomonas mendocina*. *Process in Biochemistry*, 37(12), 1431-1437.
- Tiehm, A. (1994). Degradation of Polycyclic Aromatic-Hydrocarbons in the Presence of Synthetic Surfactants. *Applied and Environmental Microbiology* 60(1), 258-263
- Toccalino, P.L. (1992). Optimization of hydrocarbon biodegradation in a sandy soil. *Scholar Archive*, 289.
- Todd, G. D., Chessin, R. L., and Colman, J. (1999). Toxicological profile for total petroleum hydrocarbons (TPH). Research Triangle Institute, US Department of Health and Human Services, Agency for Toxic substances and Disease Registry, 205-93-0606.
- Toledo, F.L., Calvo, C., Rodelas, B., and González-López, J. (2006). Selection and identification of bacteria from waste crude oil with PAHs removal capacities. *Systematic and Applied Microbiology*, 29(3), 244-252.
- Tremolada, P., Villa, S., Bazzarin, P., Bizzotto, E., Comolli, R., and Vighi, M. (2008). POPs in mountain soils from the Alps and Andes: suggestions for a 'precipitation effect on altitudinal gradients. *Water, Air and Soil Pollution*, 188(1-4), 93-109.
- Ulrich, A.C., S.E. Guigard, J.M. Foght, K.M. Semple, and Pooley, K. (2009). Effect of salt on aerobic biodegradation of petroleum hydrocarbons in contaminated groundwater. *Biodegradation*, 20(1), 27-38.
- Upshall, C., Payne, J.F., and Hellou, J. (1992). Induction of MFO Enzymes and Production of Bile Metabolites in Rainbow Trout (*Oncorhynchus mykiss*) Exposed to Waste Crankcase Oil. *Environmental Toxicology and Chemistry*, 12(11), 2105-2112
- USEPA (2009). Ecotoxicological data; selected and distributed by technical database services (TDS), New York.

- USEPA (2008). National priority chemicals trends report, trends analyses for specific priority chemicals (2004–2006): Polycyclic aromatic compounds, hazardous waste minimization and management division, office of solid waste, USA.
- USEPA (2006). Standard operating procedure; determination of metals by inductively coupled plasma (ICP) methods, EPA/SW-846, methods 3015/3050B/6010B, SOP 1811. 3, 14-42.
- USEPA (1995). Manual bioventing principle and practice, volume I and II: Bioventing principles and design.
- USEPA (1988). Cleanup of releases from petroleum UST's: selected technologies. USEPA office of underground storage tanks, EPA/530/UST-88/00.
- US Food and Drug Administration (USFDA, 2013). Agency response letter GRAS notice no. GRN 000240.
- USFDA (2011). Listing of food additives status Part II: US Food and Drug Administration, 10-27.
- Vagi, M.C., Kostopoulou, M.N., Petsas, A.S., Lalousi, M.E., Rasouli, C., and Lekkas, T.D. (2005). Toxicity of organophosphorous pesticides to the green alga *Tetraselmis suecica*. *Global Nest, The International Journal*, 7(2), 222-227.
- Van Der Greef, J., Stroobant, P., and Van der Heijden, R. (2004). The role of analytical sciences in medical systems biology. *Current Opinion in Chemical Biology*, 8(5), 559-565.
- Van Der Meer, J.R., De Vos, W.M., Harayama, S., and Zehnder, A.J. (1992). Molecular mechanisms of genetic adaptation to xenobiotic compounds. *Microbiological Reviews*, 56(4), 677-694.
- Van Hamme, J.D., Singh, A., and Ward, O.P. (2003). Recent Advances in Petroleum Microbiology, *Microbiology and Molecular Biology Review*, 67(4), 649.
- Van Loosdrecht, M.C., Lyklema, J., Norde, W., and Zehnder, A.J. (1990). Influence of interfaces on microbial activity. *Microbiological Reviews*, 54(1), 75-87.
- Varjani, S.J., and Upasani, V.N. (2013). Comparative studies on bacterial consortia for hydrocarbon degradation. *International Journal of Innovative Research in Science, Engineering and Technology*, 2(10), 5377-5383.
- Van Rooij, J.G., Bodelier-Bade, M.M., and Jongeneelen, F.J. (1993). Estimation of individual dermal and respiratory uptake of polycyclic aromatic hydrocarbons in 12 coke oven workers. *British Journal of Industrial Medicine*, 50(7), 623-632.

- Veignie, E., Rafin, C., Landy, D., Fourmentin, S., and Surpateanu, G. (2009). Fenton degradation assisted by cyclodextrins of a high molecular weight polycyclic aromatic hydrocarbon benzo [a] pyrene. *Journal of Hazardous Materials*, 168(2), 1296-1301.
- Vela, N., Martínez-Menchón, M., Navarro, G., Pérez-Lucas, G., and Navarro, S. (2012). Removal of polycyclic aromatic hydrocarbons from groundwater by heterogeneous photocatalysis under natural sunlight. *Journal of Photochemistry and Photobiology A: Chemistry*, 232, 32-40.
- Vidali, M. (2001). Bioremediation; an overview. *Pure and Applied Chemistry*, 73(7), 1163-1172.
- Vig, K., Megharaj, M., Sethunathan, N., and Naidu, R. (2003). Bioavailability and toxicity of cadmium to microorganisms and their activities in soil: a review. *Advances in Environmental Research*, 8(1), 121-135.
- Wallace, J.M., and Hobbs, P.V. (2006). *Atmospheric Science: an Introductory Survey, volume 2*. Academic Press USA.
- Wang, Y.H., Lin, C., Lai, Y.C., and Chang-Chien, G.P. (2009). Characterization of PCDD/Fs, PAHs, and heavy metals in a secondary aluminum smelter. *Journal of Environmental Science and Health, Part A*, 44(13), 1335-1342.
- Weckwerth, W. (2003). Metabolomics in systems biology. *Annual Review of Plant Biology*, 54(1), 669-689.
- Weissenfels, W.D., Klewer, H.J., and Langhoff, J. (1992). Adsorption of PAHs by soil particles: influence on biodegradability and biotoxicity. *Applied Microbiology and Biotechnology*, 36(5), 689-696.
- Weissenfels, W.D., Beyer, M., and Klein, J. (1990). Degradation of phenanthrene, fluorene and fluoranthene by pure bacterial cultures. *Applied Microbiology and Biotechnology*, 32(4), 479-484.
- Wenster-Botz, D. (2000). Experimental design for fermentation media development: Statistical design or global random search? *Journal of Bioscience and Bioengineering*, 90(5), 473-483.
- WHO (1998). *Environmental Health Criteria 202: Selected Non-heterocyclic PAHs*. WHO, Geneva, Swisarland, 883.
- Wilkinson, S., Klar, J., and Applegarth, S. (2006). Optimizing biofuel cell performance using a targeted mixed mediator combination. *Electroanalysis*, 18(19-20), 2001-2007.
- William, M.W., and Gordon R.S. (2000). Limiting factors for hydrocarbon biodegradation at low temperature in Arctic soils. *Soil Biology and Biochemistry*, 32(8), 1161-1172.

- Willey, J. (2008). *Prescott, Harley, and Klein's Microbiology-7th International ed.*/Joanne M. Willey, Linda M. Sherwood, Christopher J. Woolverton. McGraw-Hill Higher Education, New York, USA.
- Wilson, S.C., and Jones, K.C. (1993). Bioremediation of soil contaminated with PAHs: a review. *Environmental Pollution*, 61(3), 229-249.
- Wick, A.F., Haus, N.W., Sukkariyah, B.F., Haering, K.C., and Daniels, W.L. (2011). *Remediation of PAH-contaminated soils and sediments: a literature review*. Virginia Polytechnic Institute and State University, USA, 102.
- Wong, K.W., Toh, B.A., Ting, Y.P., and Obbard, J.P. (2005). Biodegradation of phenanthrene by the indigenous microbial biomass in a zinc amended soil. *Letters in Applied Microbiology*, 40(1), 50-55.
- Wong, F., Harnerb, T., Liua, Q.T., and Diamonda, M.L. (2004). Using experimental and forest soils to investigate the uptake of PAHs along urban-rural gradient. *Environmental Pollution*, 129(3), 387-398.
- Wong, J.W.C., Lai, K.M., Wan, C.K., Ma, K.K., and Fang, M. (2002). Isolation and optimization of PAH-degradative bacteria from contaminated soil for PAHs bioremediation. *Water, Air and Soil Pollution*, 139(1-4), 1-13.
- Xia, Y., Min, H., Rao, G., Lv, Z.M., Liu, J., and Ye, Y.F. (2005). Isolation and characterization of phenanthrene-degrading *Spingomonas paucibilis* strain ZX4. *Biodegradation*, 16(5), 393-402.
- Yang, Y., Chen, R.F., and Shiaris, M.P. (1994). Metabolism of naphthalene, fluorene, and phenanthrene: preliminary characterization of a cloned gene cluster from *Pseudomonas putida* NCIB 9816. *Journal of Bacteriology*, 176(8), 2158-2164.
- Yoshida, N., Hoashi, J., Morita, T., McNiven, S.J., Nakamura, H., and Karube, I. (2001). Improvement of a mediator-type biochemical oxygen demand sensor for on-site measurement. *Journal of Biotechnology*, 88(3), 269-275.
- Young, M.E., Murray, M., and Cordiner, P. (1999). Stone consolidants and chemical treatments in Scotland. *Report to Historic Scotland*, 298.
- Yuan, S.Y., Chang, J.S., Yen, J.H., and Chang, B.V. (2001). Biodegradation of phenanthrene in river sediment. *Chemosphere*, 43(3), 273-278.
- Younas, R., Zhang, S., Zhang, L., Luo, G., Chen, K., Cao, L and Hao, S. (2016). Lactic acid production from rice straw in alkaline hydrothermal conditions in presence of NiO nanoplates. *Catalysis Today*, 274, 40-48.
- Zafar, M., Kumar, S., and Kumar, S. (2010). Optimization of naphthalene biodegradation by a genetic algorithm based response surface methodology. *Brazilian Journal of Chemical Engineering*, 27(1), 89-99.

- Zeng, L., Huang, J., Zhang, Y., Qiu, G., Tong, J., Chen, D., and Luo, X. (2008). An effective method of DNA extraction for bioleaching bacteria from acid mine drainage. *Applied Microbiology and Biotechnology*, 79(5), 881-888.
- Zhang, Y., and Tao S (2009). Global atmospheric emission inventory of PAHs 2004. *Atmospheric Environment*, 43(4), 812-819.
- Zhao, H., Zhang, Y., Xiao, X., Li, G., Zhao, Y., and Liang, Y. (2017). Different phenanthrene degrading bacteria cultured by *in situ* soil substrate membrane system and traditional cultivation. *International Biodeterioration and Biodegradation*, 117, 269-277.
- Zhao, H.P., Wang, L., Ren, J.R., Li, Z.Z., Li, M., and Gao, H.W. (2008). Isolation and characterization of phenanthrene-degrading strains *Sphingomonas* sp. ZP1 and *Tistrella* sp. ZP5, *Journal of Hazard Materials*, 152(3), 1293-1300.
- Zhao, J., Wang, M., Yang, Z., Wang, Z., Wang, H., and Yang, Z. (2007). The different behaviors of three oxidative mediators in probing the redox activities of the yeast *Saccharomyces cerevisiae*. *Analytica Chimica Acta*, 597(1), 67-74.
- Zhang, Z., Rengel, Z., Meney, K., Pantelic, L., and Tomanovic, R. (2011). Polynuclear aromatic hydrocarbons mediate cadmium toxicity to an emergent wetland species. *Journal of Hazardous Materials*, 189(1), 119-126.
- Zhang, W.X., Bouwer, E.J., and Ball, W.P. (1998). Bioavailability of hydrophobic organic contaminants: Effects and implications of sorption-related mass transfer on bioremediation. *Groundwater Monitoring and Remediation*, 18(1), 126-138.
- Zulkifli, S.Z., Ismail, A., Mohamat-Yusuff, F., Arai, T., and Miyazaki, N. (2010a). Johor Strait as a hotspot for trace elements contamination in Peninsular Malaysia. *Bulletin of Environmental Contamination and Toxicology*, 84(5), 568-573.
- Zulkifli, S.Z., Mohamat-Yusuff, F., Arai, T., Ismail, A., and Miyazaki, N. (2010b). An assessment of selected trace elements in intertidal surface sediments collected from the Peninsular Malaysia. *Environmental Monitoring and Assessment*, 169(1), 457-472.
- Zylstra, G.J., and Gibson, D.T. (1991). Aromatic hydrocarbon degradation: a molecular approach. *Genetic Engineering*, 183-203.