



UNIVERSITI PUTRA MALAYSIA

***OPTIMIZATION, KINETIC MODELING, AND CHARACTERIZATION OF
HYDRO-DISTILLATION AND SUBCRITICAL WATER EXTRACTION OF
AQUILARIA MALACCENSIS LAMK LEAVES AND WOOD ESSENTIAL
OIL***

MAHTAB SAMADI

FK 2017 17



**OPTIMIZATION, KINETIC MODELING, AND CHARACTERIZATION OF
HYDRO-DISTILLATION AND SUBCRITICAL WATER EXTRACTION OF
AQUILARIA MALACCENSIS LAMK LEAVES AND WOOD ESSENTIAL OIL**

By

MAHTAB SAMADI

**Thesis Submitted to the School of Graduate Studies,
Universiti Putra Malaysia, in Fulfilment of the
Requirements for the Degree of Doctor of Philosophy**

February 2017



© COPYRIGHT UPM

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

To my great family: my lovely parents, my unique husband, and my kind mother-in-law

Words cannot express how grateful I am for all their kindness and support. I am forever in their debt.

- ❖ My lovely parents (Ali Asghar Samadi and Maryam Kamali), whose smile means a world to me

My father is my kind-hearted hero. From the day that I remember, he has been trying to make the life better for all of his girls. He is the greatest and kindest dad of all. What can I say about my mother now that truly express my gratitude. She is my beautiful kind-hearted angel, whose biggest wish is the success and happiness of me and my sweet sisters. Her friendly, warm, and welcoming voice is the essential element of my daily life. My parents have always wanted the best for their children and have raised us with pure love. Their only dream is the success and happiness of their children and they have sacrificed everything to make that happen. Words cannot express their pure kindness. My dream has always been to make my parents happy. They're the best and I'm so lucky to be their daughter. Their endless love has always given me the strength no matter how far they were away from me.

- ❖ My unique, talented, kind, and supportive husband (Dr. Amir Masoud Tabatabaei Yazdi), whose love is the greatest gift of all and I'm so grateful for that

He is the love of my life. His kind support is beyond the words. I've tasted the sweet and precious love with him. He has been always beside me in sadness and happiness. He is the best of the bests, my best friend, and kindest of all. He has supported me during the saddest and hardest times of my life. His warm hug and kind voice is always soothing. I'm so grateful for his unconditional love and support. The truth is that without his love and support, I would have never been able to continue and finish my study. Even though our field of study is totally different, he never hesitated to help me. He owns my heart and his presence is the reason of my heart beat. Words cannot express my feeling and my appreciation. I only can say that I love him from the bottom of my heart, and I'm so thankful for all the things that he has done for me.

- ❖ My lovely mother in-law (Dr. Vida Samimi), who is not only like my mother, but also she is my kind, and supportive friend

She has always been supportive with her beautiful smile. Her flawless-pure love and support is priceless. She is the best mother in-law and I can't be luckier to have someone like her in my life. I love her like my own mother and her happiness is my happiness. She has always wanted the best for us and has sacrificed a lot for that. I want to express my gratitude for all the sacrifices that she has done. Appreciate all the love and support that she has been given us unconditionally. I hope that someday I can be able to make it up to her for all her kindness

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

OPTIMIZATION, KINETIC MODELING, AND CHARACTERIZATION OF HYDRO-DISTILLATION AND SUBCRITICAL WATER EXTRACTION OF *AQUILARIA MALACCENSIS* LAMK LEAVES AND WOOD ESSENTIAL OIL

By

MAHTAB SAMADI

February 2017

Chair: Associate Professor Zurina Zainal Abidin, PhD
Faculty: Engineering

Aquilaria malaccensis (gaharu), which is an Indo-Malaysian evergreen tree, is one of the most important higher-plants in south East Asia. Because of the high-demand for this valuable tree especially for its resinous wood oil, *Aquilaria malaccensis* is considered as an endangered species. On the other hand, the current extraction method (distillation) in gaharu industries not only requires lots of wood material for only small amount of essential oil, but also, it consumes lots of time and energy. Therefore, finding an efficient alternative to traditional distillation method such as Subcritical Water Extraction (SCWE; an extraction method that uses water at subcritical conditions) can make a significant difference in saving the materials, time, and energy. Thereby, the primary goal of this study was to assess whether SCWE can be used as an alternative to HD for extraction of *A. malaccensis* wood essential oil.

In addition, this study, for the first time, aimed to analyze *A. malaccensis* leaves oil since not only there had been some indications that the leaves of this evergreen tree may be as valuable as its wood, but also, the positive outcome would put a stop to the waste of *A. malaccensis* leaves.

For comparing both extraction methods (i.e. SCWE and HD), at first, the optimum conditions for both methods were identified by assessing the effect of different influential parameters (temperature/heating-power, sample-to-solvent ratio, and time). Hydro-distillation experiments conducted using cleverger apparatus, while SCWE was done using a batch subcritical reactor. Yield comparison between the essential oils extracted by HD and SCWE at optimum conditions showed that the yield of both leaves and wood essential oils extracted by SCWE (30 min) were almost 2.5 times the yield of HD (4-16 hrs). Additionally, the characterization tests were carried out on the essential oils and the leaves/wood samples. GC/MS results provided evidence that the quality of *A. malaccensis* wood and leave oils extracted by SCWE is significantly better compared to that extracted by HD as the amount of oxygenated and sesquiterpenoid compounds was found to be higher in oils extracted by SCWE. The essential oils extracted by SCWE also contained several value-added compounds useful in medicine such as furfural and guaiacol. FTIR, SEM, and BET/BJH on the wood and

leaves samples provided further evidence for better performance of SCWE, since the sample's pores, cell walls, cellulose, and hemicellulose were more damaged compared to those in HD sample.

Furthermore, in this study, kinetic modelling was conducted in order to provide a better understanding of both HD's and SCWE's mechanism. The result of kinetic modelling indicated that the unsteady-state diffusion model is the best model for describing HD, whereas, two-site kinetic desorption and second-order model were found to be the best models for explaining SCWE of wood and leaves essential oils respectively.

In short, the result of this study showed that SCWE is a better extraction method in terms of time, efficiency and quality. Both *A. malaccensis* leaves and wood oils were found to contain beneficial compounds useful for producing pesticides and medicine.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

PENGOPTIMUMAN, PEMODELAN KINETIC, DAN PENCIRIAN HYDRO-PENYULINGAN DAN AIR PERAHAN SUBGENTING DARIPADA AQUILARIA MALACCENSIS LAMK DEDAUNAN AND WOOD MINYAK ESSENTIAL

Oleh

MAHTAB SAMADI

Februari 2017

Pengerusi: Profesor Madya Zurina Zainal Abidin, PhD

Fakulti: kejuruteraan

Pada masa kini, kepentingan untuk produk semula jadi telah meningkat dengan ketara di seluruh dunia kerana produk semula jadi 'kurang kesan sampingan dan faedah yang lebih tinggi. Tinggi-tumbuhan, yang mengandungi pelbagai jenis metabolit sekunder dengan kepelbagaian struktur yang tinggi, telah digunakan sebagai salah satu sumber utama produk semula jadi untuk beberapa dekad. *Aquilaria malaccensis* (gaharu), yang merupakan pokok malar hijau Indo-Malaysia, adalah salah satu yang paling penting yang lebih tinggi-kilang di selatan Asia Timur. Oleh kerana permintaan tinggi untuk pokok ini berharga terutama bagi minyak kayu resin itu, *Aquilaria malaccensis* dianggap sebagai spesies terancam. Sebaliknya, kaedah semasa pengekstrakan (penyulingan) dalam industri gaharu bukan sahaja memerlukan banyak bahan kayu untuk jumlah hanya kecil minyak penting, tetapi juga, ia menggunakan banyak masa dan tenaga. Oleh itu, mencari alternatif yang cekap kepada kaedah penyulingan tradisional boleh membuat perbezaan yang signifikan dalam menyelamatkan bahan-bahan, masa, dan tenaga. pengambilan air subgenting (SCWE) adalah salah satu kaedah yang selamat, cekap dan mesra alam. Oleh itu, matlamat utama kajian ini adalah untuk menilai sama ada SCWE boleh digunakan sebagai alternatif kepada HD untuk pengekstrakan *A. malaccensis* minyak pati kayu. Di samping itu, kajian ini, buat kali pertama, bertujuan untuk menganalisis *A. malaccensis* meninggalkan minyak. Menganalisis *A. malaccensis* meninggalkan minyak pati adalah penting kerana bukan sahaja terdapat beberapa tanda-tanda bahawa daun pokok malar hijau ini mungkin yang berharga sama seperti kayu, tetapi juga, hasil positif akan menghentikan pembaziran *A. malaccensis* meninggalkan apabila pokok gaharu yang ditebang.

Untuk membandingkan kedua-dua kaedah pengekstrakan (iaitu SCWE dan HD), pada mulanya, keadaan optimum untuk kedua-dua kaedah telah dikenalpasti dengan menilai kesan parameter berpengaruh berbeza. Hasil perbandingan antara minyak pati diekstrak oleh HD dan SCWE pada keadaan optimum menunjukkan bahawa hasil daripada kedua-dua daun dan minyak pati kayu telah meningkat kepada hampir 2.5 kali oleh SCWE Masa perahan yang jauh lebih pendek (kurang dari 30 min). Selain itu, keputusan GC / MS memberikan bukti bahawa kualiti *A. malaccensis* kayu dan

meninggalkan minyak yang diekstrak oleh SCWE adalah jauh lebih baik berbanding yang diekstrak oleh HD kerana ia mengandungi sebatian nilai tambah. Hasil ujian pencirian (FTIR, SEM, dan BET / BJH) memberikan bukti lanjut untuk prestasi yang lebih baik daripada SCWE sejak liang sampel tersebut, dinding sel, selulosa, hemiselulosa dan selepas menjalani SCWE, lebih rosak berbanding sampel HD.

Tambahan pula, dalam kajian ini, model kinetik telah dijalankan untuk memberi pemahaman yang lebih baik daripada kedua-dua ini HD dan mekanisme SCWE ini. Hasil daripada model kinetik menunjukkan bahawa model penyebaran -State tak mantap adalah model yang terbaik untuk menerangkan HD, manakala, dua tapak kinetik desorption dan kedua untuk model didapati model yang terbaik untuk menerangkan SCWE kayu dan daun minyak pati masing-masing .

Pendek kata, hasil daripada kajian ini menunjukkan bahawa SCWE adalah kaedah pengekstrakan yang lebih baik dari segi masa, kecekapan dan kualiti. Kedua-dua *A. malaccensis* daun dan minyak kayu didapati mengandungi sebatian bermanfaat berguna untuk menghasilkan racun perosak dan perubatan.

ACKNOWLEDGEMENT

First and the foremost, I sincerely would like to express my utmost gratitude and respect to my great supervisor, Associate Prof. Dr. Zurina Zainal Abidin, for all the expert guidance, advices, support, and inspiration during these five years of my study. I am so lucky to have her as my supervisor and I shall never forget her kindness and support.

I would also like to express my respect and gratitude to my expert co-supervisors, Prof. Dr. Robiah Yunus, Dr. Dayang Radiah bt. Awang Biak, Prof. Dr. Hiroki Yoshida, and Dr. Eng Hai Lok, for their valuable-professional advices, time, and support during this project.

Moreover, I would like to thank my great friends and family. I'm very lucky to be surrounded by kind people like them. First, special thanks to my beautiful lovely sisters, Mahsa Samadi and Mahnoosh Samadi, for all the moral supports, encouragement, and kindness. They are the best sisters in the world and I'm the luckiest sister in the world to have such a kind sisters. Second, I would like to express my special appreciation to my great friends, who never cease to support me. I truly appreciate the expert helps, advices, and moral support of my kind friends, Dr. Soraya Hosseini and Dr. Farah Eghbali Babadi, regarding my thesis and cherish their true friendship. I also appreciate all the helps, inspirations, and moral support that Roya Esfahani gave me during my study. I shall never forget all those memorable days that I shared with all of them. Last but not least, I would like to express my special appreciation to my husband's family for all their supports and kindness.

APPROVAL

I certify that a Thesis Examination Committee has met on 28th February 2017 to conduct the final examination of Mahtab Samadi on her thesis entitled “Optimization, kinetic modeling, and characterization of hydro-distillation and subcritical water extraction of *Aquilaria malaccensis* lank leaves and wood essential oil” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Suraya bt. Abdul Rashid, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Thomas Choong Shean Yaw, PhD

Professor Ir.
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Mohd Halim Shah bin Ismail, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Elena Ibanez, PhD

Professor
Institute of Food Science Research
Spain
(External Examiner)

NOR AINI AB. SHUKOR, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 April 2017

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Zurina Zainal Abidin, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Robiah Yunus, PhD

Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Dayang Radiah bt. Awang Biak, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Hiroiyuki Yoshida, PhD

Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Eng Hai Lok, PhD

Senior Research Officer
Forest Research Institute Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by Graduate Student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Mahtab Samadi, GS32023

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee
Zurina Zainal Abidin

Signature: _____
Name of
Member of
Supervisory
Committee
Robiah Yunus

Signature: _____
Name of
Member of
Supervisory
Committee
Dayang Radiah Biak

Signature: _____
Name of
Member of
Supervisory
Committee
Hiroyuki Yoshida

Signature: _____
Name of
Member of
Supervisory
Committee
Eng Hai Lok

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENT	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xvi
CHAPTER	
1	
INTRODUCTION	1
1.1. Background	1
1.2. Problem statement	1
1.2.1. Why <i>Aquilaria malaccensis</i> ?	1
1.2.2. Why new extraction method is needed?	2
1.2.3. Gap	3
1.3. Significance of the study/Practical Contribution	3
1.4. Research Objectives	4
1.5. Scope	5
1.6. Structure of thesis	5
2	
LITERATURE REVIEW	7
2.1. Essential oil	7
2.1.1. Introduction	7
2.1.2. History/background	9
2.1.3. Definition	10
2.1.4. Chemical composition	10
2.1.5. Benefits and uses	12
2.2. Extraction	13
2.2.1. Sample preparation and pretreatment	14
2.2.1.1. Drying	14
2.2.1.2. Size Reduction	15
2.2.2. Hydro-distillation (HD)	15
2.2.2.1. Introduction	15
2.2.2.2. Background/ History	16
2.2.2.3. Principle	16
2.2.2.4. Equipment setup	17
2.2.2.5. Influential parameters	18
2.2.2.6. Application	19
2.2.2.7. Kinetic models	20
2.2.3. Subcritical water extraction (SCWE)	24
2.2.3.1. Introduction	24
2.2.3.2. Background/history	26
2.2.3.3. Principles	27
2.2.3.4. Extraction mechanism	31
2.2.3.5. Parameters affecting extraction	32

	2.2.3.6. Applications	35
	2.2.3.7. Kinetic models	37
2.3.	<i>Aquilaria malaccensis</i>	40
	2.3.1. Introduction	40
	2.3.2. <i>Aquilaria</i> genus	40
	2.3.3. Morphology, Ecology, and Distribution	44
	2.3.4. Resin formation	46
	2.3.5. Grading	46
	2.3.6. Trade	49
	2.3.7. Benefits and Usage	50
	2.3.7.1. Medicinal Application	50
	2.3.7.2. Other applications	51
	2.3.8. Extraction methods	51
	2.3.9. Identified chemical compounds	52
2.4.	Response Surface Methodology (RSM)	55
	2.4.1. Choice of the experimental design	56
	2.4.2. Codification of the levels of the variable	57
	2.4.3. Mathematical–statistical treatment of data	58
	2.4.4. RSM applications for extraction of essential oil	59
2.5.	Summary	60
3	MATERIALS AND METHODS	61
	3.1. Materials preparation	61
	3.1.1. Chemicals/Materials	62
	3.1.2. Sample preparation	62
	3.2. Extraction of essential oil	63
	3.2.1. Hydro-distillation of <i>A. malaccensis</i> leaves and wood	63
	3.2.1.1. Essential oil purification and yield calc.	65
	3.2.2. Subcritical Water Extraction of <i>A. malaccensis</i>	66
	3.2.2.1. Design of experiments and statistical	66
	3.2.2.2. Subcritical water extraction	71
	3.2.2.3. Essential oil purification and yield calc.	72
	3.3. Kinetic model assessment	73
	3.3.1. Hydro-distillation (HD)	73
	3.3.2. Subcritical water extraction (SCWE)	73
	3.4. Comparison of the extraction methods	75
	3.4.1. Gas chromatography Mass spectroscopy (GC/MS)	75
	3.4.2. Scanning electron microscopy	77
	3.4.3. Fourier transform-infrared (FT-IR) analysis	78
	3.4.4. BET surface area and BJH pore size distribution	79
	3.5. Summary	80
4	RESULTS AND DISCUSSION	82
	4.1. Extraction of <i>A. malaccensis</i> leaves essential oil	82
	4.1.1. Hydro-distillation extraction method	82
	4.1.1.1. Effect of Operational Conditions	82
	4.1.1.2. Optimization of operational condition	87
	4.1.1.3. Kinetic model	87
	4.1.2. Subcritical water extraction method	92
	4.1.2.1. Statistics/statistical Analysis of Expe.	92
	4.1.2.2. Analysis of variance (ANOVA)	95

4.1.2.3.	Regression analysis	99
4.1.2.4.	Interactive effect of operational condition	102
4.1.2.5.	Optimization of operational condition	104
4.1.2.6.	Kinetic modeling	105
4.1.3.	Qualitative assessment of the performance of the	112
4.1.3.1.	Gas Chromatography/ Mass Spectroscop	112
4.1.3.2.	Scanning electron microscope (SEM)	120
4.1.3.3.	Fourier transform infrared spectroscopy	122
4.1.3.4.	Brunauer-Emmett-Teller (BET) Surface	123
4.2.	Extraction of <i>A. malaccensis</i> wood essential oil	125
4.2.1.	Hydro-distillation extraction method	125
4.2.1.1.	Effect of Operational Conditions	125
4.2.1.2.	Optimization of operational condition	130
4.2.1.3.	Kinetic model	130
4.2.2.	Subcritical water extraction method	134
4.2.2.1.	Statistics/statistical Analysis of Expe.	134
4.2.2.2.	Analysis of variance (ANOVA)	137
4.2.2.3.	Regression analysis	141
4.2.2.4.	Interactive effect of variables	144
4.2.2.5.	Optimization of operational condition	146
4.2.2.6.	Kinetic modeling	148
4.2.3.	Qualitative assessment of the extraction methods	154
4.2.3.1.	Gas Chromatography/Mass Spectroscopy	154
4.2.3.2.	Scanning electron microscope (SEM)	162
4.2.3.3.	Fourier transform infrared spectroscopy	164
4.2.3.4.	Brunauer-Emmett-Teller (BET) Surface	165
5	CONCLUSION	168
5.1.	Overview of the study	168
5.2.	Summary of Study findings	168
5.3.	Conclusion and Implications	171
5.4.	Future research and recommendation	172
	REFERENCES	173
	APPENDICES	189
	BIODATA OF STUDENT	199
	PUBLICATION	200

LIST OF TABLES

Table	Page	
2.1	history of natural product medicine	8
2.2	Examples of secondary metabolites and their uses	9
2.3	some examples of chemical components of commercially important	11
2.4	Physiochemical properties of some common solvents used in natural	25
2.5	Examples of using SCWE for extraction or analysis of bioactive	26
2.6	various supercritical solvents and their critical conditions	29
2.7	SCWE of plant and food materials	36
2.8	Analyses of organic contaminants in environment by SCWE	36
2.9	Stocking of <i>Aquilaria</i> spp. in Malaysia during 200-2004	41
2.10	sical appearance of various grades of gaharu	47
2.11	Component based characteristic of agarwood oil	48
2.12	Examples of major chemical compounds of <i>A. malaccensis</i> wood	54
3.1	List of chemical materials	62
3.2	Important variables affecting subcritical water extraction	67
3.3	CCD design of extraction experiments for extraction of essential oils	67
3.4	applied kinetic models for hydro-distillation extraction	73
3.5	applied kinetic models for subcritical water extraction	74
3.6	GC/MS operational condition for identifying chemical compositions	76
4.1	Models used for studying kinetic of hydro-distillation	88
4.2	Values of the kinetic parameters	88
4.3	The yield of <i>A. malaccensis</i> based on Central Composite Design (CCD)	93
4.4	Sequential Model Sum of Squares	95
4.5	Lack of fit tests	96
4.6	Model summary statistics	97
4.7	The actual and predicted values of the response in the model	98
4.8	Analysis of variance of essential oil from <i>A. malaccensis</i> leaves	99
4.9	Analysis of variance and regression coefficients of extraction	101
4.10	Optimum condition derived by the response surface	105
4.11	Summary of kinetic models for extraction of <i>A. malaccensis</i>	107
4.12	Values of kinetic parameters of models for extraction	107
4.13	Chemical Compositions of Essential oil of <i>Aquilaria malaccensis</i> ' Leaves	114
4.14	Values of the kinetic parameters	132
4.15	The yield of <i>A. malaccensis</i> wood based on (CCD)	135
4.16	Sequential Model Sum of Squares	137
4.17	Lack of fit tests	138
4.18	Model summary statistics	139
4.19	The actual and predicted values of the yield in model	140
4.20	Analysis of variance and R-squared of extraction of essential	142
4.21	Analysis of variance and regression coefficients of extraction	143
4.22	Optimum condition derived by the response surface	147
4.23	Summary of kinetic models for extraction of <i>A. malaccensis</i>	149
4.24	Values of four kinetic models' parameters of <i>A. malaccensis</i>	150
4.25	Chemical Compositions of Essential oil of <i>Aquilaria malaccensis</i>	157

LIST OF FIGURES

Figure	Page	
2.1	Water phase temperature-pressure diagram	28
2.2	Some properties of water in the pressure range 21.8–30 MPa	30
2.3	graphical representation of dielectric constant of water vs. temperature	31
2.4	A proposed schematic presentation of the extraction steps	32
2.5	<i>Aquilaria</i> species	43
2.6	locations of <i>Aquilaria</i> populations in Malaysia	44
2.7	<i>Aquilaria malaccensis</i> Lam.	45
2.8	The geographical distribution of <i>Aquilaria malaccensis</i> in Malaysia	45
3.1	Summary of methods	61
3.2	(a) Clevenger-type apparatus hydro-distillation set up (b) clevenger	63
3.3	Extraction process by hydro-distillation method	64
3.4	HD's procedure	65
3.5	removing moisture of essential oil	66
3.6	Mathematical modeling steps in response surface methodology (RSM)	70
3.7	batch subcritical water extraction (SCWE) set up	71
3.8	stainless steel reactor	71
3.9	extraction set up by SCWE	72
3.10	liquid-liquid extraction process of essential oil	72
3.11	Characterization of essential oils and dry samples of <i>A. malaccensis</i>	75
4.1	Typical profile of <i>A. malaccensis</i> ' leaves essential oil during the hydro	83
4.2	The amount of extracted essential oil from <i>A. malaccensis</i> ' leaves	84
4.3	The amount of extracted essential oil from <i>A. malaccensis</i> ' leaves	85
4.4	Second order kinetic model of extraction of <i>A. malaccensis</i> ' leaves	89
4.5	Second order kinetic model of extraction of <i>A. malaccensis</i> ' leaves	90
4.6	Unsteady state diffusion kinetic model of extraction	90
4.7	Unsteady state diffusion kinetic model of extraction	91
4.8	Types of the model order	96
4.9	Scatter plot of predicted value versus actual value from central	98
4.10	Residual plot of runs from central composite design for extraction	99
4.11	Response surface plot of significant variables on extraction	103
4.12	profile of SCWE of <i>A. malaccensis</i> leaves essential oil at different	106
4.13	Second order kinetic model of extraction of <i>A. malaccensis</i>	108
4.14	Two-site desorption kinetic model of extraction of <i>A. malaccensis</i>	109
4.15	Thermodynamic partition with external mass transfer kinetic	110
4.16	GC/MS of the essential oil of <i>A. malaccensis</i> leaves by hydro	114
4.17	GC/MS chromatogram of essential oil of <i>A. malaccensis</i> leaves	114
4.18	Possible conversion reactions in essential oils	116
4.19	Hydrolysis of lignin to smaller compounds	117
4.20	Micrographs of (a) raw (i.e. untreated) <i>A. malaccensis</i> leaves	121
4.21	FTIR spectrum of <i>A. malaccensis</i> leaves before and after extraction	122
4.22	pore size plot of <i>A. malaccensis</i> leaves before extraction, after	123
4.23	Isotherm linear plot of <i>A. malaccensis</i> leaves before extraction	125
4.24	Typical profile of <i>A. malaccensis</i> wood essential oil extraction	126
4.25	The amount of extracted essential oil from <i>A. malaccensis</i>	127

4.26	The amount of extracted essential oil from <i>A. malaccensis</i>	128
4.27	Second order kinetic model of extraction of <i>A. malaccensis</i>	131
4.28	Second order kinetic model of extraction of <i>A. malaccensis</i>	131
4.29	Unsteady state diffusion kinetic model of extraction	133
4.30	Unsteady state diffusion kinetic model of extraction	133
4.31	Scatter plot of predicted value versus actual value from	140
4.32	Residual plot of runs from central composite design for yield	141
4.33	Response surface plot of significant variables on extraction	145
4.34	Profile of SCWE of <i>A. malaccensis</i> wood essential oil at different	149
4.35	Second order kinetic model of extraction of <i>A. malaccensis</i>	151
4.36	Two-site desorption kinetic model of extraction	152
4.37	Portioning coefficient with external mass transfer kinetic model	153
4.38	GC/MS chromatogram of <i>A. malaccensis</i> wood essential oil	156
4.39	GC/MS chromatogram of <i>A. malaccensis</i> wood essential oil	156
4.40	xylose and glucose pathway to furfural and HMF	159
4.41	Micrographs of <i>A. malaccensis</i> wood (a) before extraction	163
4.42	FTIR result of <i>A. malaccensis</i> wood before and after extraction	165
4.43	pore size of <i>A. malaccensis</i> wood before and after extraction	166
4.44	Isotherm linear plot of <i>A. malaccensis</i> wood before extraction	167

LIST OF ABBREVIATIONS

ANOVA	Analysis of Variance
BET	Brunauer Emmett Teller
BJH	Barrett-Joyner-Halenda
CCD	Central Composite Design
EO	Essential Oil
FT-IR	Fourier Transform-Infrared
GC	Gas Chromatography
GC/MS	Gas Chromatography/Mass Spectroscopy
HD	Hydro-Distillation
RSM	Response Surface Methodology
SCWE	Sub-Critical Water Extraction
SEM	Scanning Electron Microscopy
WHO	World Health Organization

CHAPTER 1

INTRODUCTION

1.1 Background

Aromatic herbs and plants are proven to be quite beneficial to human being as they have diverse range of applications in food, medicine, and etc. (Bonjar, 2004; Bakkali et al., 2008; Soran et al., 2015). For instance, the vast range of vegetables, herbs, fruits, and aromatic plants have been used in traditional medicine as they contain a rich source of biological active compounds (Nikolić, et al., 2014). It is not surprising that the interest for natural and organic products has increased significantly worldwide as they are considered to be less harmful in comparison with non-organic products (Jiao, et al., 2013). Nowadays, there are numerous natural products in the market for anti-parasitical, bactericidal, fungicidal, pesticides, cosmetic, perfumery, food additives, and medicinal purposes (Gechev et al., 2014; Sarker et al., 2006).

Lots of the above-mentioned organic products contain essential oils extracted from aromatic plants (Jiao et al., 2013). Essential oil, which is the key property of aromatic plants, is an intricate combination of volatile compounds usually exists in low amount within the plant (Reverchon, 1997; Hui, 1992; Lucchesi et al., 2004; Baser & Buchbauer, 2009). However, the amount of essential oil as well as chemical composition existing in essential oil varies significantly in each method of extraction (Jiao et al., 2013; Baser & Buchbauer, 2009). Therefore, given the aforementioned importance of essential oil for natural products, finding the highly efficient isolation method of essential oils as well as understanding and identifying the chemical composition of the essential oil extracted by each method are vitally important.

1.2 Problem statement

1.2.1 Why *Aquilaria malaccensis*?

Aquilaria malaccensis, also known as Agarwood or Gaharu, is one of the valuable *Aquilaria* species. *Aquilaria* species (23 species) is one of the important medicinal plant genres that have been used for centuries by both people and industries in south-east Asia. *Aquilaria* is a genus in the family *Thymeleaceae* and class *magnoliopsida* (Dashet al., 2008; Barden et al., 2000). For more than 2000 years, *Aquilaria malaccensis*/Agarwood has been considered precious as it has been used for perfumery, cosmetic, medicine and others (Barden et al., 2000). The importance of this evergreen tree is mostly because of its wood resin, which is produced as a natural immune response of the tree towards fungal attack. Despite the fact that 23 species of *Aquilaria* have been identified so far, only a few *Aquilaria* species- including *Aquilaria malaccensis*- produce this resin (Donovan & Puri, 2004; Barden et al., 2000). Thanks to its resin, *Aquilaria malaccensis*' wood has many useful medicinal applications and it has been used in traditional medicine to treat pain, fever, rheumatism, and asthma. Additionally, researches on non-traditional medicine revealed that the wood of *A. malaccensis* also has remarkable anticancer activity (Gunasekera et al., 1981). In the recent years, *Aquilaria malaccensis* has been widely used -resulting in over-harvesting- and hence it has been enlisted in red list of threatened trees (Barden et al., 2000).

1.2.2 Why new extraction method is needed?

Hydro-distillation (HD) and Steam distillation (SD) are two common methods used by industries for extracting essential oil from wood of *Aquilaria malaccensis*. Yet, these conventional methods are extremely time-consuming (Desai et al., 2014; Yoswathana et al., 2012; Sulaiman et al., 2015). Prior to extraction, these methods require one week for soaking (Fazila & Halim, 2012; Tajuddin & Yusoff et al., 2010). Additionally, they take about 16 hours to extract essential oil from the wood of *Aquilaria malaccensis*. Moreover, beside the disadvantages on the time consumption, HD and SD methods have other drawbacks as well. The loss of volatile compounds, low extraction efficiency, degradation of unsaturated compounds, and high energy consumption are the examples of these problems (Tam et al., 2007). To address these drawbacks, other extraction methods have been employed. Yet, other extraction methods were also found to be problematic or inefficient. For instance, Supercritical Fluids Extraction (SFE) method was developed to reduce the time consumption and also to improve the extraction efficiency (i.e. higher yield; Ibrahim et al., 2011). But, the carbon dioxide emitted during the extraction process is not environmental friendly as it has greenhouse effect (Jiménez-Carmona et al., 1999). Additionally, recently soxhlet extraction and Accelerated Solvent Extraction (ASE) methods have been used to improve the efficiency of essential oil extracted from *A. malaccensis* (Sulaiman et al., 2015). Despite the fact that both the soxhlet and ASE methods seem to improve the yield of essential oil, these methods still are not desirable since these methods use organic solvents (e.g. n-hexane), which are toxic, hazardous, and uneconomic. Thus, still a new extraction method for addressing aforementioned problems of HD and SD is required.

Subcritical Water Extraction (SCWE) is a new promising extraction method, which is safe, fast, economic, and environmental friendly in comparison with other new methods. Extraction of essential oil using SCWE uses water as solvent, which is cheap and environmental friendly (Ayala & Luquede Castro, 2001). Furthermore, SCWE has shown to require significantly less extraction time (around 2-3 times) and utilize lower amount of raw material, and it produces higher quality and quantity of essential oil (Herrero, Cifuentes, & Ibanez, 2006). The SCWE is rapidly emerging as an alternative for the extraction of essential oils compounds in the world (Luquede Castro et al., 1998).

Yoswathana et al. (2012) provided evidence for the suitability of SCWE method for extracting essential oil from *Aquilaria crassna* (i.e. one of *Aquilaria*/agarwood species). By using the SCWE for extracting essential oil from *A. crassna* (i.e. one of the species from *Aquilaria* family) in a very limited range of temperatures (i.e. 100, 125, and 150°C) and constant time, Yoswathana et al. (2012) concluded that in comparison with HD, Subcritical Water Extraction (SCWE) method resulted in higher yield of *A. crassna* oil in shorter time period. Hence, there are some indications that SCWE may be a better alternative for HD and SD methods in terms of time consumption, yield, and the quality *A. malaccensis*' oil.

1.2.3 Gap

As mentioned earlier, finding a better alternative for HD process can be quite useful for the industry. Additionally, assessing the extraction mechanism as well as the optimization is very beneficial for controlling and fine-tuning the extraction process. Yet, to the best of the author's knowledge, so far, no study on the extraction mechanism and optimization of subcritical water extraction (SCWE) for extraction of *A. malaccensis* oil has been carried out. Additionally, the study on the extraction mechanism and optimization of hydro-distillation (HD) of *A. malaccensis* is very limited (only one study exists; Sulaiman, et al., 2015). So, more investigation on the essential oil of *Aquilaria malaccensis*' wood is needed.

There are some indications that *Aquilaria malaccensis*' leaves can also be useful for variety of purposes. For instance, numerous studies (e.g. Suresh et al., 2008; Ibrahim et al., 2011; Morris et al., 2011; Dash et al., 2008; and Gurib Fakim, 2006 maintained that in higher plants (e.g. Agarwood), there is a good chance of finding rich amount of secondary metabolites, aromatic substances, microorganisms, and high structural diversity compounds in all of its parts. These materials have significant application in medicine, cosmetic, and pesticides (Gurib Fakim, 2006; Sarker et al., 2006). By taking into consideration that Agarwood has lots of green leaves throughout the year, finding secondary metabolites, aromatic substances, and microorganisms in Agarwood's leaves can be quite useful for providing necessary material for producing medicinal, cosmetic, and pesticide products.

Moreover, there are other indications that investigating on *Aquilaria malaccensis*' leaves are likely to be fruitful. For example, Huda et al. (2009), Khalil et al. (2013) and Wil et al. (2014) noted that solvent extraction of *A. malaccensis*' leaves shows significant antioxidant activity, and therefore, it can be considered as a natural antioxidant. Begum (2015), also provided evidences for both antibacterial and antioxidant activity of solvent extract of *A. malaccensis*' leaves. In spite of all the above-mentioned indications for the possible applications of *A. malaccensis*' leaves, surprisingly, no study on the essential oil of *A. malaccensis*' leaves or its chemical compounds has been conducted. So, by studying *A. malaccensis*' leave oil through characterization, this study aimed to fill this void.

1.3 Significance of the study/Practical Contribution

The significance of this research is threefold. First, by testing Subcritical Water Extraction (SCWE) method in terms of time consumption, yield, and the quality of product, this study tried to determine whether SCWE can be used as a better alternative for hydro-distillation (HD) and steam distillation (SD). By identifying SCWE as a better extraction method for *Aquilaria malaccensis*, this research can be quite beneficial for industries, in which *Aquilaria malaccensis*' essential oil is extracted, and help them to save a lot of time and money.

Second, by finding useful components from *Aquilaria malaccensis*' leaves such as secondary metabolites, aromatic substances, and microorganisms, which are useful for many industries (e.g. health care and agriculture), this study can take a big step in providing information for producing organic products. Finding useful components from *Aquilaria malaccensis*' leaves is importance since not only there are huge amount of leaves on *Aquilaria malaccensis* throughout the years that can be used, but also it will find alternative usage for the leaves of *Aquilaria malaccensis* which is already cut down (better usage of the whole tree).

Finally, by conducting a kinetic modelling on the extraction process using *Aquilaria malaccensis*, this study provides necessary information required for better controlling and optimizing the extraction process. This information can also help organizations to enhance the efficiency of extraction process and consequently improve their revenue (Paunović et al., 2014; Cassel et al., 2009). In addition, the result of kinetic modelling is essential for comparing traditional and modern extraction methods, scaling up purposes, and the design of unit operation (Chan et al., 2014; Silva et al., 2008; Minozzo, et al., 2012).

1.4 Research Objectives

This study attempts to fill the mentioned gaps and problem in three ways. First, by extracting and analyzing the essential oil of *Aquilaria malaccensis* from wood and (especially) leaves by both hydro-distillation and subcritical water extraction, this study attempts to identify its chemical compositions in order to provide a foundation for further investigations. Secondly, by conducting the kinetic-modeling for extraction method of the essential oil, this study tries to get fuller understanding of the mechanism of both hydro-distillation and subcritical water extraction processes, and consequently, to identify the optimum condition, in which higher yield of essential oil can be obtained. Finally, given the abundant benefits of finding better extraction methods for different industries, this study aimed to assess whether SCWE method is better than traditional methods (i.e. HD and SD) in terms of time consumption, yield efficiency and quality of the essential oil.

Overall, the objectives are divided into specific objectives as follows, which are the main contribution of the thesis to the body of knowledge:

- ❖ To optimize the hydro-distillation and subcritical water extraction of essential oils from *Aquilaria malaccensis*' leaves and wood.
- ❖ To assess kinetic modelling for both hydro-distillation and subcritical-water extraction process using the existing models and the experiment results in order to provide a better understanding of the extraction processes.
- ❖ To assess the effectiveness of hydro-distillation (HD) and subcritical-water extraction (SCWE) on extraction of the essential oil from *A. malaccensis* leaves and wood using GC/MS, SEM, FTIR, and BET results.

1.5 Scope

- ❖ In this study, extraction of the essential oils by hydro-distillation Clevenger apparatus method have been done as preliminary study (reference method) and main focus in on the extraction of the essential oils by subcritical water extraction (SCWE).
- ❖ A batch/static subcritical water extraction system, which has been imported from Japan, was employed in this study because not only this method is environmental friendly, also has the advantage of the short time of extraction.
- ❖ Response surface methodology (RSM) was used to design the extraction experiments by SCWE from both leaves and wood of *A. malaccensis*. A central composite design was used to optimize operational parameters of the extraction of essential oils by SCWE.
- ❖ The optimization of hydro-distillation method was assessed based on the yield of essential oil at different heating powers and sample-to-solvent ratios.
- ❖ The optimization of subcritical water extraction method was assessed based on the essential oil yield at different extraction temperatures.
- ❖ The effectiveness of HD and SCWE process on quality of the extracted essential oil was evaluated based on the chemical composition of *A. malaccensis* leave and wood essential oils identified by GC/MS
- ❖ The effectiveness of HD and SCWE in extraction of the essential oils was assessed based on morphological and structural changes of samples (using the results of SEM, FTIR, and BET on *A. malaccensis* leave and wood samples before and after extraction).

1.6 Structure of thesis

This thesis is divided into five chapters. Chapter one covers introduction, problem statements, objectives, scope and thesis structure. Chapter two includes descriptions on the essential oil, *Aquilaria malaccensis*, its benefits and its extraction methods. The advantages and disadvantages of hydro-distillation as current extraction method of *A. malaccensis* essential oil were explained. Also, Subcritical water extraction was introduced in full description as a possible replacement extraction method for gaharu industries. Furthermore, this chapter consists of the kinetic mechanism of extraction of agarwood oil by HD and SWE. Also, theory of multi-objective design optimization using response surface methodology was explained shortly. In chapter three, both the materials and methods for extracting essential oil from *A. malaccensis* wood and leave are elaborated. Moreover, the characterization methods (i.e. GC/MS, FTIR, BET, and SEM) for testing *A. malaccensis* leave and wood oil as well as for assessing leave and wood sample before and after extraction were explained. In chapter four, the results of experiments were thoroughly explained and discussed. Chapter four consists of preliminary study on extraction of essential oil from both leaves and wood of *A. malaccensis* by HD at optimum condition, which was found by studying the effect of key parameters and kinetic modeling of the process. An experimental design using

Central Composite Design (CCD) was done on the SCWE extraction of oils from wood and leaves of *A. malaccensis*. Then it continues to optimize operational parameters using response surface methodology. Finally, the kinetic study of the SCWE process for oil extraction from *A. malaccensis* was presented to have a better understanding of the process. This chapter also covers the results of several characterization tests reports of both essential oils and dry sample (before and after extraction) of HD and SCWE, in order to do a better comparison of the extraction methods. Finally, the overview of the study, conclusion, implication, and the direction for future studies were presented in Chapter 5.



REFERENCES

- Abd El-Aty, A., Mohamed, A., & Samhan, F. (2014). In vitro antioxidant and antibacterial activities of two fresh water Cyanobacterial species, *Oscillatoria agardhii* and *Anabaena sphaerica*. *Journal of Applied Pharmaceutical Science* Vol. , 4 (07): 069-075.
- Abdelmoez, W., Ashour, E., Naguib, S., Hilal, A., Al Mahdy, D., Mahrous, E., et al. (2016). Kinetic and Thermodynamics studies for Castor Oil Extraction Using Subcritical Water Technology. *Journal of Ole science*, 477-485.
- Adams, R. P. (2007). *Identification of Essential Oil Components By Gas Chromatography/Mass Spectrometry, 4th Edition*. Illinois: Allured Publishing Corporation.
- Ali, N. A., Ismail, N., & Nasir, T. M. (2012). Analysis of Agarwood Oil (*Aquilaria Malaccensis*) Based on GC-MS Data. *IEEE 8th International Colloquium on Signal Processing and its Applications* (pp. 470-473). Malacca: Advancing technology for humanity.
- Anderson, M. J., & Whitcomb, P. J. (2004). *RSM simplified : optimizing processes using response surface methods for design of experiments*. New York, N.Y.: Productivity Press .
- Aneupankul, T., Goto, M., Sasaki, M., Pavasant, P., & Shotipruk, A. (2007). Extraction of anti-cancer damnacanthol from roots of *Morinda citrifolia* by subcritical water. *Separation and Purification Technology*, 343-349.
- Annex2. (2003). *Review of Significant Trade: Aquilaria malaccensis*. Significant trade in plants: Implementation of Resolution Conf. 12.8: Progress with the Implementation of Species Reviews (CITES PC14 Doc.9.2.2).
- Anurukvorakun, O. (2013). Factorial Design Applied to Subcritical Water Extraction for the Investigation of Flavonoids and Antioxidant Capacity of *Gynura calciphila* Kerr. *Mahidol University Journal of Pharmaceutical Sciences*, 7-16.
- Aparna, V., Dileep, K., Mandal, P., Karthe, P., Sadasivan, C., & Haridas, M. (2012). Anti-Inflammatory Property of nHexadecanoic Acid: Structural Evidence and Kinetic Assessment. *Chemical Biology & Drug Design*, 434-439.
- Attokaran, M. (2011). *Natural Food Flavors and Colorants*. Iowa: Wiley blackwell.
- Ayala, R. S., & de Castro, M. L. (2001). Continuous subcritical water extraction as a useful tool for isolation of edible essential oils. *Food Chemistry*, 109-113.
- Azadfar, M., Gao, A. H., Bule, M. V., & Chen, S. (2015). Structural characterization of lignin: A potential source of antioxidants guaiacol and 4-vinylguaiacol. *International Journal of Biological Macromolecules*, 58-66.
- Bakkali, F., Averbeck, S., Averbeck, D., & Idaomar, M. (2008). Biological effects of essential oils – A review. *Food and Chemical Toxicology*, 446-475.
- Barden, A., Anak, N. A., Mulliken, T., & Song, M. (2000). *Heart of the Matter: Agarwood Use and Trade and CITES Implementation for Aquilaria malaccensis*. Cambridge: TRAFFIC International.
- Baş, D., & Boyacı, İ. H. (2007). Modeling and optimization I: Usability of response surface methodology. *Journal of Food Engineering*, 836-845.
- Baser, K. H., & Buchbauer, G. (2009). *Handbook of Essential Oils: Science, Technology, and Applications*. New York: CRC Press .
- Baser, K. H., & Demirci, F. (2007). Chemistry of Essential Oils. In R. G. Berger, *Flavours and Fragrances: Chemistry, Bioprocessing and Sustainability*. Berlin: Springer.

- Basile, A., Carmona, M. J., & Clifford, A. (1998). Extraction of rosemary by superheated water. *Journal of Agricultural and Food Chemistry*, 5204–5209.
- Bazaid, S., El-Amoudi, M., Ali, E., & Abdel-Hameed, E. (2013). Volatile oil studies of some aromatic plants in Taif region. *Journal of Medicinal Plants Studies*, 119-128.
- Begum, Y. (2015). Study on agarwood (*Aquilaria malaccensis*) to evaluate antibacterial and antioxidant activities of n – hexane, chloroform and ethyl acetate extracts. *PHARMATUTOR*, 2391.
- Beniwal, B. S. (1989). Silvical characteristics of *Aquilaria agallocha*. *Indian Forester*, 17–21.
- Bezerra, M. A., Santell, R. E., Oliveira, E. P., Villar, L. S., & Escalera, L. A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*, 965–977.
- Bonjar, G. H. (2004). Screening for antibacterial properties of some Iranian plants against two strains of *Escherichia coli*. *Asian Journal of Plant Sciences*. *Asian Journal of Plant Sciences*, 310-314.
- Boye, J. I., & Arcand, Y. (2011). *Green Technologies in Food Production and Processing*. New York: Springer.
- Brebu, M., & Vasile, C. (2010). Thermal degradation of lignin – a review. *Cellulose chemistry and technology*, 353-363.
- Brei, B., Edman, J. D., Gerade, B., & Clark, J. M. (2004). Relative abundance of two cuticular hydrocarbons indicates whether a mosquito is old enough to transmit malaria parasites. *Journal of Medical Entomology*, 807-809.
- Brereton, R. G. (2003). *Chemometrics: Data analysis for the laboratory and chemical plant (Vol. 8)*. John Wiley and Sons Ltd.
- Buang, F., Jantan, I., Amran, A. Z., & Dachyar, A. (2014). Optimization of Ginger (*Zingiber officinale*) Oil Yield from Malaysia in Different Hydrodistillation Physical Parameters via Central Composite Design of Response Surface Methodology (RSM). *Research Journal of Applied Sciences, Engineering and Technology*, 5098-5105.
- Burdock, G. A. (1997). *Encyclopedia of Food and Color Additives, volume 1*. New York: CRC Press.
- Cahyadi, C., Heng, P. W., & Chan, L. W. (2011). Optimization of process parameters for a quasi-continuous tablet coating system using design of experiments. *AAPS PharmSciTech*, 119–31.
- Carabias-Martinez, R., Rodriguez-Gonzalo, E., Revilla-Ruiz, P., & Hernandez-Mendez, J. (2005). Pressurized liquid extraction in the analysis of food and biological samples. *Journal of Chromatography A*, 1-17.
- Carmody, O., Frost, R., Xi, Y., & Kokot, S. (2007). Surface characterisation of selected sorbent materials for common hydrocarbon fuels. *Surface Science*, 2066-2076.
- Cassel, E., Vargas, R., Martinez, N., Lorenzo, D., & Dellacassa, E. (2009). Steam distillation modeling for essential oil extraction process. *Industrial Crops and Products*, 29: 171-76.
- Chai, W.-M., Liu, X., Hu, Y.-H., Feng, H.-L., Jia, Y.-L., Guo, Y.-J., et al. (2013). Antityrosinase and antimicrobial activities of furfuryl alcohol, furfural and furoic acid. *International Journal of Biological Macromolecules*, 151-155.
- Chakrabarty, K., Kumar, A., & Menon, V. (1994). *Trade in Agarwood*. New Delhi: TRAFFIC India and WWF-India.
- Chan, C.-H., Yusoff, R., & Ngoh, G.-C. (2014). Modeling and kinetics study of conventional and assisted batch solvent extraction. *Chemical Engineering Research and Design*, 1169-1186.

- Chemat, F., Vian, M. A., & Cravotto, G. (2012). Green Extraction of Natural Products: Concept and Principles. *International Journal of Molecular Sciences*, 8615-8627.
- Chen, H.-Q., Wei, J.-H., Yang, J.-S., Zhang, Z., Yang, Y., Gao, Z.-H., et al. (2012). Chemical Constituents of Agarwood Originating from the Endemic Genus *Aquilaria* Plants. *Chemistry & Biodiversity*, 236–250.
- Chen-Xing, Z., Mi, Z., Jing, H., & Bao-Cai, L. (2014). Chemical composition and antioxidant activity of the essential oil from the flowers of *Artemisia austro-yunnanensis*. *Journal of Chemical and Pharmaceutical Research*, 1583-1587.
- Chua, L. S. (2008). Agarwood (*Aquilaria malaccensis*) in Malaysia. *NDF WORKSHOP CASE STUDIES*, (p. 15). Malaysia.
- Chun-Pinga, X., Yuanshanga, L., Shanshana, Z., & Yinga, Z. (2015). Chemical Composition and In Vitro Antioxidant Activity of the Essential Oil of *Rehmannia glutinosa* Libosch Using Ultrasound as a Pretreatment. *Journal of Biologically Active Products from Nature*, 276-282.
- Cooperband, M., McElfresh, J., Millar, J., & Cardé, R. (2008). Attraction of female *Culex quinquefasciatus* Say (Diptera: Culicidae) to odors from chicken feces. *Journal of Insect Physiology*, 54(7):1184-92.
- Cortadella, J., & Reisig, W. (2004). *Applications and Theory of Petri Nets 2004: 25th International Conference*. Verlag Berlin Heidelberg: Springer.
- Council of Europe; European Pharmacopoeia Commission. (2008). *European pharmacopoeia*. Strasbourg: Council Of Europe.
- Critical point (thermodynamics)*. (2016, June 26). Retrieved June 2016, from Wikipedia, the free encyclopedia: [https://en.wikipedia.org/wiki/Critical_point_\(thermodynamics\)](https://en.wikipedia.org/wiki/Critical_point_(thermodynamics))
- Crozier, A., Clifford, M. N., & Ashihara, H. (2006). *Plant Secondary Metabolites: Occurrence, Structure and Role in the Human Diet*. oxford, UK: Blackwell Publishing Ltd.
- Dash, M., Patra, J. K., & Panda, P. P. (2008). Phytochemical and antimicrobial screening of extracts of *Aquilaria agallocha* Roxb. *African Journal of Biotechnology*, 3531-3534.
- Deng, C., Yao, N., Wang, A., & Zhang, X. (2005). Determination of essential oil in a traditional Chinese medicine, *Fructus amomi* by pressurized hot water extraction followed by liquid-phase microextraction and gas chromatography–mass spectrometry. *Analytica Chimica Acta*, 237–244.
- Desai, M., Parikh, J., & Kumar De, A. (2014). Modelling and optimization studies on extraction of lemongrass oil from *Cymbopogon flexuosus* (Steud.) Wats. *Chemical Engineering Research and Design*, 9 2: 793–803.
- Dias, D. A., Urban, S., & Roessner, U. (2012). A Historical Overview of Natural Products in Drug Discovery. *Metabolites*, 303-336.
- Donovan, D., & Puri, R. (2004). Learning from Traditional Knowledge of Non-timber Forest Products: Penan Benalui and the Autecology of *Aquilaria* in Indonesian Borneo. *Ecology and Society*, 3-26.
- Dubey, A., Boukouvala, F., Keyvan, G., Hsia, R., Saranteas, K., Brone, D., et al. (2012). Improvement of tablet coating uniformity using a quality by design approach. *AAPS PharmSciTech*, 231–246.
- Eikani, M. H., Golmohammad, F., & Rowshanzamir, S. (2007). Subcritical water extraction of essential oils from coriander seeds (*Coriandrum sativum* L.). *Journal of Food Engineering*, 735-740.

- Enriquez, M. M., Del, v. A., Chauvet, D., Lopez, V. A., Susano, P. M., & Campos, S. A. (2009). Acute Toxicity of Guaiacol Administered Subcutaneously in the Mouse. *Proceedings of the Western Pharmacology Society*, 92-3.
- Eurlings, M. C., & Gravendeel, B. (2005). TrnL-trnF sequence data imply paraphyly of *Aquilaria* and *Gyrinops* (Thymelaeaceae) and provide new perspectives for agarwood identification. *Plant Systematics and Evolution*, 1-12.
- Fang, Z., Sato, T., Smith Jr, R. L., Inomata, H., Arai, K., & Kozinski, J. A. (2008). Reaction chemistry and phase behavior of lignin in high-temperature and supercritical water. *Bioresource Technology*, 3424–3430.
- Faraone, N., Hillier, N. K., & Cutler, G. C. (2015). Plant Essential Oils Synergize and Antagonize Toxicity of Different Conventional Insecticides against *Myzus persicae* (Hemiptera: Aphididae). *PLoS ONE*, 1-12.
- Faridah, Q., Abdelmageed, A. H., Nor Hazirah, A. N., & Yaacob, M. (2010). Comparative study of essential oil composition of leaves and rhizomes of *Alpinia conchigera* Griff. At different post- harvest drying periods. *Journal of Medicinal Plants Research*, 2700-2705.
- Fazila, K. N., & Halim, K. K. (2012). Effects of soaking on Yield and Quality of Agarwood Oil. *Journal of Tropical Forest Science*, 557–564.
- Fernández-González, V., Concha-Graña, E., Muniategui-Lorenzo, S., López-Mahía, P., & Prada-Rodríguez, D. (2008). Pressurized hot water extraction coupled to solid-phase microextraction–gas chromatography–mass spectrometry for the analysis of polycyclic aromatic hydrocarbons in sediments. *Journal of Chromatography A*, 65–72.
- Figueiredo, C., Matsuo, A., Pereira, F., Rabaça, A., Farias, C., N., G., et al. (2014). *Pyrostegia venusta* heptane extract containing saturated aliphatic hydrocarbons induces apoptosis on B16F10-Nex2 melanoma cells and displays antitumor activity in vivo. *Pharmacognosy Magazine*, 363-376.
- Fujii, T., Khuwijtjaru, P., Kimura, Y., & Adachi, S. (2006). Decomposition kinetics of monoacyl glycerol and fatty acid in subcritical water under temperature-programmed heating conditions. *Food Chemistry*, 341-347.
- Gad, S. C. (2005). *Drug Discovery Handbook*. Hoboken: John Wiley and Sons Inc.
- Gámiz-Gracia, L., & Luque de Castro, M. (2000). Continuous subcritical water extraction of medicinal plant essential oil: comparison with conventional techniques. *Talanta*, 1179–1185.
- Gechev, T. S., Hille, J., Woerdenbag, H. J., Benina, M., Mehterov, N., Toneva, V., et al. (2014). Natural products from resurrection plants: Potential for medical applications. *Biotechnology Advances*, 1091-1101.
- Gilmour, S. G. (2006). Response surface designs for experiments in bioprocessing. *Biometrics*, 323-331.
- Giray, E. S., Kirici, S., Kaya, D. A., Turk, M., Sonmez, O., & Inan, M. (2008). Comparing the effect of sub-critical water extraction with conventional extraction methods on the chemical composition of *Lavandula stoechas*. *Talanta*, 930–935.
- Gogus, F., Ozel, M. Z., & Lewis, A. C. (2005). Superheated Water Extraction of Essential Oils of *Origanum micranthum*. *Journal of Chromatographic Science*, 87-91.
- Golmakani, M.-T., & Rezaei, K. (2008). Comparison of microwave-assisted hydrodistillation with the traditional hydrodistillation method in the extraction of essential oils from *Thymus vulgaris* L. *Food chemistry*, 925–930.
- Gong, Y., Zhang, X., He, L., Yan, Q., Yuan, F., & Gao, Y. (2013). Optimization of subcritical water extraction parameters of antioxidant polyphenols from sea

- buckthorn (*Hippophaë rhamnoides* L.) seed residue. *Journal of Food Science and Technology*, 1534-1542.
- Gumgumjee, N. M., & Hajar, A. S. (2015). . Antibacterial activities and gc-ms analysis of phytochemicals of *Ehretia abyssinica* r.br. ex fresen. *International Journal of Applied Biology and Pharmaceutical Technology*, 236-241.
- Gunasekera, S. P., Kinghorn, A. D., Cordell, G. A., & Farnsworth, N. R. (1981). Plant Anticancer Agents. XIX. Constituents of *Aquilaria malaccensis*. *Journal of natural products*, 569–572.
- Gunstone, F. D., Harwood, J. L., & Dijkstra, A. J. (2007). *The Lipid Handbook with Cd-Rom. 3rd edition* . Boca Raton: CRC Press.
- Gupta, S., & Dikshit, A. (2010). Biopesticides: An ecofriendly approach for pest control. *Journal of Biopesticides*, 186-188.
- Gurib Fakim, A. (2006). Medicinal Plants: Traditions of Yesterday and Drugs of Tomorrow. *Molecular Aspects of Medicine*, 1-93.
- Handa, S. S., Khanuja, S. P., Longo, G., & Rakesh, D. D. (2008). *Extraction Technologies for Medicinal and Aromatic Plants*. Trieste, Italy: ICS-UNIDO.
- Harada, H., Yamashita, U., Kurihara, H., Fukushi, E., Kawabata, J., & Kamei, Y. (2002). Antitumor activity of palmitic acid found as a selective cytotoxic substance in a marine red alga. . *Anticancer Research*, 2587–2590.
- Hashim, Y. Z.-Y., Ismail, N. I., & Abbas, P. (2014). Analysis of chemical compounds of agarwood oil from different species by gas chromatography mass spectrometry (gcms). *IIUM Engineering Journal*, 55-60.
- Hashim, Y. Z.-Y., Phirdaus, A., & Azura, A. (2014a). Screening of anticancer activity from agarwood essential oil. *Pharmacognosy Res.*, 191-194.
- Hawthorne, S. B., Yang, Y., & Miller, D. J. (1994). Extraction of Organic Pollutants from Environmental Solids with Sub- and Supercritical Water. *Analytical chemistry*, 2912–2920.
- Helmi, Z., Al Azzam, K., Tsymbalista, Y., Ghazleh, R., Shaibah, H., & Aboul-Enein, H. (2014). Analysis of Essential Oil in Jerusalem Artichoke (*Helianthus tuberosus* L.) Leaves and Tubers by Gas Chromatography-Mass Spectrometry. *Advanced Pharmaceutical Bulletin*, 521-526.
- Herrero, M., Cifuentes, A., & Ibanez, E. (2006). Sub- and supercritical fluid extraction of functional ingredients from different natural sources: Plants, food-by-products, algae and microalgae A review. *Food Chemistry* , 136–148.
- Hidayat, W., Shakaff, A. Y., Ahmad, M. N., & Adom, A. H. (2010). Classification of Agarwood Oil Using an Electronic Nose. *Sensors*, 4675-4685.
- Higher alkanes. (2016). Retrieved from Wikipedia: https://en.wikipedia.org/wiki/Higher_alkanes
- Ho, C. H., Cacace, J. E., & Mazza, G. (2008). Mass transfer during pressurized low polarity water extraction of lignans from flaxseed meal. *Journal of Food Engineering*, 64-71.
- Ho, G., Hao, V. L., HALL, D., LE, T., NGUYEN, D., CROSS, J., et al. (2014). (Z)-11-hexadecenyl acetate and (Z)-13-octadecenyl acetate improve the attractiveness of the standard sex pheromone of the yellow rice stem borer *Scirpophaga incertulas* (Lepidoptera: Pyralidae) in northern Vietnam. *J. Fac. Agr., Kyushu Univ.*, (2014), 59 (1): 85–89.
- Ho, Y., & McKay, G. (1999). Pseudo-second order model for sorption processes. *Process Biochemistry*, 451–465.
- Huda, A., Munira, M., Fitrya, S., & Salmah, M. (2009). Antioxidant activity of *Aquilaria malaccensis* (thymelaeaceae) leaves. *Pharmacognosy Research*, 1:5:270-273.

- Hui, Y. H. (1992). *Encyclopedia of food science and technology*. New York: Wiley.
- Ibáñez, E., Kubatova, A., Senorañs, F. J., Caverro, S., Reglero, G., & Hawthorne, S. B. (2003). Subcritical water extraction of antioxidant compounds from rosemary plants. *Journal of Agricultural and Food Chemistry*, 375–382.
- Ibrahim, A. H., Al-Rawi, S. S., Abdul Majid, A. M., Rahman, N. N., Abo- Salah, K. M., & Ab Kadir, M. O. (2011). Separation and fractionation of *Aquilaria Malaccensis* oil using supercritical fluid extraction and the cytotoxic properties of the extracted oil. *Procedia Food Science*, 1953-1959.
- Irchhaiya, R., Kumar, A., yadav, A., Gupta, N., Kumar, S., Gupta, N., et al. (2014). Metabolites in plants and its classification. *World Journal of Pharmacy and Pharmaceutical Sciences*, 287-305.
- Ishihara, M., Tsuneya, T., & Uneyama, K. (1993). Components of the Volatile Concentrate of Agarwood. *Journal of Essential Oil Research*, 283-289.
- Islam, M. N., Jo, Y.-t., & Park, J.-H. (2012). Determination of Partitioning Coefficient and Desorption Rate for Subcritical Water Extraction of PAHs. *IPCBE*, 45-48.
- Islam, M. N., Jo, Y.-T., Jung, S.-K., & Park, J.-H. (2013). Thermodynamic and kinetic study for subcritical water extraction of PAHs. *Journal of Industrial and Engineering Chemistry*, 129–136.
- Islam, M., Hussain, M., Kabir, E., & Hasan, M. (2014). Development of low cost agarwood essential oil extraction system for agar tree growers in Bangladesh. *Journal of the Bangladesh Agricultural University*, 167-170.
- Ismail, N., Ali, N. A., Jamil, M., Rahiman, M. H., Tajuddin, S. N., & Taib, M. N. (2014). A Review Study of Agarwood Oil and Its Quality Analysis. *Jurnal Teknologi (science and engineering)*, 37-42.
- Janey Alam, M., Mujahid, B., Azizur, R., Juber, A., Mohmmad, K., Yasmeen, J., et al. (2015). An insight of pharmacognostic study and phytopharmacology of *Aquilaria agallocha*. *Journal of Applied Pharmaceutical Science*, 173-181.
- Jayachandran, K., Sekar, I., Parthiban, K. T., Amirtham, D., & Suresh, K. K. (2014). Analysis of different grades of Agarwood (*Aquilaria malaccensis* Lamk.) oil through GC-MS. *Indian journal of natural products and resources*, 44-47.
- Jiao, J., Gai, Q.-Y., Fu, Y.-J., Zu, Y.-G., Luo, M., Zhao, C.-J., et al. (2013). Microwave-assisted ionic liquids treatment followed by hydro-distillation for the efficient isolation of essential oil from *Fructus forsythiae* seed. *Separation & Purification Technology*, 228-237.
- Jiménez-Carmona, M. M., & Castro, M. D. (1999). Isolation of eucalyptus essential oil for GC-MS analysis by extraction with subcritical water. *Chromatographia*, 578-582.
- Jong, P. L., Tsan, P., & Mohamed, R. (2014). Gas Chromatography-Mass Spectrometry Analysis of Agarwood Extracts from Mature and Juvenile *Aquilaria malaccensis*. *International Journal of Agriculture and Biology*, 644-648.
- Jouyban, A., Soltanpour, S., & Chan, H.-K. (2004). A simple relationship between dielectric constant of mixed solvents with solvent composition and temperature. *International Journal of Pharmaceutics*, 353-360.
- Jumepaeng, T., Jantayota, R., Hemmood, S., Komolwanich, S., Wuthisarn, S., Luthria, D., et al. (2013). Determination and Comparison of Volatile Aroma Compounds in Fresh and Dried Leaves Samples of Citronella Grass, Lemongrass, and Citronella Incense Products by Micro-Hydrodistillation. *Acta Chromatographica*, 177-190.

- Kamali, H., Aminimoghadamfarouj, N., & Nematollahi, A. (2014). Optimization of pressurized hot water extraction of Lavandin essential oils via central composite design. *International Journal of ChemTech Research*, 4853-4861.
- Kamiyama, M., Shibamoto, T., Horiuchi, M., Umamo, K., Kondo, K., & Otsuka, Y. (2013). Antioxidant/Anti-Inflammatory Activities and Chemical Composition of Extracts from the Mushroom *Trametes Versicolor*. *International Journal of Nutrition and Food Sciences*, 85-91.
- Kanmaz, E. O., & Ova, G. (2013). The effective parameters for subcritical water extraction of SDG lignan from flaxseed (*Linum usitatissimum* L.) using accelerated solvent extractor. *Eur Food Res Technol*, 159-166.
- Kapas, Á., Andras, C. D., Dobre, T. G., Vass, E., Szekely, G., Stroescu, M., et al. (2011). The kinetic of essential oil separation from fennel by microwave assisted hydrodistillation (MWHD). *U.P.B. Sci. Bull.*, 113-120.
- Karmakar, M., & Ray, R. R. (2011). Optimization of production conditions of extra cellular β -glucosidase in submerged fermentation of waterhyacinth using response surface methodology by *Rhizopus oryzae* MTCC 9642. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 299-308.
- Kasrati, A., Jamali, C. A., Fadli, M., Bekkouche, K., Hassani, L., Wohlmuth, H., et al. (2014). Antioxidative activity and synergistic effect of *Thymus saturejoides* Coss. essential oils with cefixime against selected food-borne bacteria. *Industrial Crops and Products*, 338-344.
- Keawsa-ard, S., Liawruangrath, B., Liawruangrath, S., Teerawutgulrag, A., & Pyne, S. G. (2012). Chemical constituents and antioxidant and biological activities of the essential oil from leaves of *Solanum spirale*. *Natural Product Communications*, 955-958.
- Khajavi, S. H., Kimura, Y., Oomori, T., Matsuno, R., & Adachi, S. (2005). Degradation kinetics of monosaccharides in subcritical water. *Journal of Food Engineering*, 309-313.
- Khajenoori, M., Asl, A. H., & Eikani, M. H. (2015). Subcritical Water Extraction of Essential Oils from *Trachyspermum ammi* Seeds. *Journal of essential oil bearing plants*, 1165-1173.
- Khajenoori, M., Haghghi Asl, A., & Hormozi, F. (2009). Proposed Models for Subcritical Water Extraction of Essential Oils. *Chinese Journal of Chemical Engineering*, 359-365.
- Khalil, A. S., Rahim, A. A., & Abdallah, K. B. (2013). Characterization of Methanolic Extracts of Agarwood Leaves. *Journal of Applied and Industrial Sciences*, 78-88.
- Konovalova, O., Gergel, E., & Herhel, V. (2013). GC-MS Analysis of Bioactive Components of *Shepherdia argentea* (Pursh.) Nutt. from Ukrainian Flora. *The Phama innovation journal*, 7-12.
- Kraber, S., Whitcomb, P. J., & Anderson, M. J. (2014). *Handbook for Experimenters-A concise collection of handy tips to help you set up and analyze your designed experiments*. Minneapolis: Stat-Ease, Inc.
- Kramer, P. J., & Kozlowski, T. T. (1979). *Physiology of woody plants*. New York: Academic Press.
- Kronholm, J., Hartonen, K., & Riekkola, M.-L. (2007). Analytical extractions with water at elevated temperatures and pressures. *Trends in Analytical Chemistry*, 396-412.
- Kubátová, A., Jansen, B., Vaudoisot, J.-F., & Hawthorne, S. B. (2002). Thermodynamic and kinetic models for the extraction of essential oil from

- savory and polycyclic aromatic hydrocarbons from soil with hot (subcritical) water and supercritical CO₂. *Journal of Chromatography A*, 175-188.
- Kumari, S., Pundhir, S., Priya, P., Jeena, G., Punetha, A., Chawla, K., et al. (2014). EssOilDB: a database of essential oils reflecting terpene composition and variability in the plant kingdom. *Database*, 1-12.
- Kumeta, Y., & Ito, M. (2010). Characterization of delta-guaiene synthases from cultured cells of *Aquilaria*, responsible for the formation of the sesquiterpenes in agarwood. *Plant Physiol.*, 1998-2007.
- Lafrankie, J. V. (1994). Population dynamics of some tropical trees that yield non-timber forest products. *Economic Botany*, 301-309.
- Lagadec, A. J., Miller, D. J., Lilke, A. V., & Hawthorne, S. B. (2000). Pilot-Scale Subcritical Water Remediation of Polycyclic Aromatic Hydrocarbon- and Pesticide-Contaminated Soil. *Environmental Science and Technology*, 1542-1548.
- Lee, S. Y., Weber, J., & Mohamed, R. (2011). Genetic Variation and Molecular Authentication of Selected *Aquilaria* Species from Natural Populations in Malaysia Using RAPD and SCAR Markers. *Asian Journal of Plant Sciences*, 202-211.
- Lee, S., Monnappa, A. K., & Mitchell, R. J. (2012). Biological activities of lignin hydrolysate-related compounds. *Society for Biochemistry and Molecular Biology*, 265-274.
- Li, Y., Fabiano-Tixier, A.-S., & Chemat, F. (2014). *essential oils as reagents in green chemistry*. Zug: Springer International Publishing.
- Liang, X., & Fan, Q. (2013). Application of Sub-Critical Water Extraction in Pharmaceutical Industry . *Journal of Materials Science and Chemical Engineering*, 1-6.
- Liu, T., Sui, X., Zhang, R., Yang, L., Zu, Y., Zhang, L., et al. (2011). Application of ionic liquids based microwave-assisted simultaneous extraction of carnosic acid, rosmarinic acid and essential oil from *Rosmarinus officinali*. . *Journal of Chromatography A*, 8480-8489.
- Liu, X., Jia, Y., Chen, J., Liang, G., Guo, H., Hu, Y., et al. (2015). Inhibition effects of benzylideneacetone, benzylacetone, and 4-phenyl-2-butanol on the activity of mushroom tyrosinase. *Journal of Bioscience and Bioengineering*, 275-279.
- Liu, Y., Huaiqiong, C., Yang, Y., Zhang, Z., Wei, J., Meng, H., et al. (2013). Whole-tree Agarwood-Inducing Technique: An Efficient Novel Technique for Producing High-Quality Agarwood in Cultivated *Aquilaria sinensis* Trees. *Molecules*, 3086-3106.
- Lou, X., Janssen, H.-G., & Cramers, C. A. (1997). Parameters Affecting the Accelerated Solvent Extraction of Polymeric Samples. *Analytical chemistry*, 1598-1603.
- Lowell, S., Thommes, M., Thomas, M. A., & Shields, J. E. (2004). *Characterization of Porous Solids and Powders: Surface Area, Pore Size and Density*. Netherlands: Kluwer Academic Publishers.
- Lucchesi, M. E., Chemat, F., & Smadja, J. (2004). Solvent-free microwave extraction of essential oil from aromatic herbs: comparison with conventional hydro-distillation. *Journal of Chromatography A* , 323-327.
- Lundstedt, T., Seifert, E., Abramo, L., Thelin, B., Nyström, Å., Pettersen, J., et al. (1998). Experimental design and optimization. *Chemometrics and Intelligent Laboratory Systems*, 3-40.

- Luque de Castro, M., & Jiménez-Carmona, M. (1998). Potential of water for continuous automated sample-leaching. *Trends in Analytical Chemistry*, 441–447.
- Ma, X., Yang, H., Yu, L., Chen, Y., & Li, Y. (2014). Preparation, Surface and Pore Structure of High Surface Area Activated Carbon Fibers from Bamboo by Steam Activation. *Materials*, 4431-4441.
- Maiti, R., Rodriguez, H. G., & Ivanova, N. S. (2016). *Autoecology and Ecophysiology of Woody Shrubs and Trees: Concepts and Applications*. Chichester: Wiley blackwell.
- Mallick, S. S., & Dighe, V. V. (2014). Detection and Estimation of alpha-Amyrin, beta-Sitosterol, Lupeol, and n-Triacontane in Two Medicinal Plants by High Performance Thin Layer Chromatography. *Advances in Chemistry*, 1-7.
- Mallikadvi, T., Paulsamy, S., Jamuna, S., & Karthika, K. (2012). Analysis for phytochemicals and bioinformatics approach for the evaluation of therapeutic properties of whole plant methanolic extract of mukia maderaspatana (L.) m.roem. (cucurbitaceae) – a traditional medicinal plant in western districts of tamil. *Asian Journal of Pharmaceutical & Clinical Research*, 163-168.
- Manilal, A., Sujith, S., Sabarathnam, B., Kiran, G., Selvin, J., Shakir, C., et al. (2011). Biological activity of the red alga *Laurencia brandenii*. *Acta Bot. Croat*, 70 : (1) 81–90 DOI: 10.2478/v10184-010-0001-x.
- Martos, J., Osinaga, P. W., Oliveira, E. d., & Castro, L. A. (2003). Hydrolytic Degradation of Composite Resins: Effects on the Microhardness. *Materials research*, 599-604.
- Masoumi, H. R., Basri, M., Kassim, A., Abdullah, D. K., Abdollahi, Y., Gani, S. S., et al. (2013). Statistical optimization of process parameters for lipase-catalyzed synthesis of triethanolamine-based esterquats using response surface methodology in 2-liter bioreactor. *The Scientific World Journal*, 1-9.
- Masoumi, H. R., Kassim, A., Basri, M., & Abdullah, D. K. (2011). Determining optimum conditions for lipase-catalyzed synthesis of triethanolamine (TEA)-based esterquat cationic surfactant by a taguchi robust design method. *Molecules*, 4672–4680.
- Mayr, G., Zeppetzaue, F., Zweckmair, T., Bauer, D., Hilad, S., Potthast, A., et al. (2015). The Reactions of Cellulose and Hemicellulose Degradation Products in the Viscose Fibre Spin Bath. *LENZINGER BERICHTS* 92 , 53-58.
- Miguel, M. G. (2010). Antioxidant and anti-inflammatory activities of essential oils: a short review. *Molecules*, 9252-9287.
- Miller, E. N., Jarboe, L. R., Yomano, L. P., York, S. W., & Shanmugam, K. T. (2009). Silencing of NADPH-dependent oxidoreductase genes (yqhD and dkgA) in furfural-resistant ethanologenic *Escherichia coli*. *Appl Environ Microbio*, 4315-4323.
- Milojević, S., Radosavljević, D., Pavićević, V., Pejanović, S., & Veljković, V. (2013). Modeling the kinetics of essential oil hydrodistillation from plant materials. 67 (5): 843–859 doi: 10.2298/HEMIND121026009M.
- Milojević, S., Stojanović, T., Palič, R., Lazič, M., & Veljković, V. (2008). Kinetic of distillation of essential oil from comminuted ripe juniper (*Juniperus communis* L.) berries. *Biochem Eng Journal*, 39: 547–53.
- Minitab, I. (2016). *Minitab 17 support*. Retrieved September 12, 2016, from Minitab.com: <http://www.http://support.minitab.com/en-us/minitab/17/topic-library/modeling-statistics/regression-and-correlation/regression-models/what-is-model-order/>

- Minozzo, M., Popielski, A., Prá, V. D., Treichel, H., Cansian, R. L., Oliveira, J. V., et al. (2012). Modeling of the overall kinetic extraction from *Maytenus aquifolia* using compressed CO₂. *Brazilian Journal of Chemical Engineering*, 835-843.
- Miraghaee, S. S., Karimi, I., & Becker, L. A. (2011). Psychobiological Assessment of Smoke of Agarwood (*Aquilaria* spp.) in Male Rats. *Journal of Applied Biological Sciences*, 45-53.
- Miyoshi, T., Ito, M., Kitayama, T., Isomori, S., & Yamashita, F. (2013). Sedative effects of inhaled benzylacetone and structural features contributing to its activity. *Biol Pharm Bull.*, 1474-1481.
- Mohamed, N. A., Morad, N. A., & Aziz, M. K. (2004). Process design of hydro-distillation in ginger oil production. *The 4th Annual Seminar of National Science Fellowship*, (pp. 527-531). Penang.
- Montané, D., S. J., Torras, C., & Farriol, X. (2002). High-temperature dilute-acid hydrolysis of olive stones for furfural production. *Biomass and Bioenergy* , 295-304 .
- Montgomery, D. C. (2013). *Design and analysis of experiments (8th edition)*. John Wiley and Sons, Inc.
- Moreno, E., Reza, J., & Trejo, A. (2007). Extraction of polycyclic aromatic hydrocarbons from soil using water under subcritical conditions. *Polycycl. Aromat. Compd.*, 239-260.
- Morris, P., Scragg, A. H., Stafford, A., & Fowler, M. W. (2011). *secondary metabolites in plant cell culture*. London: Cambridge University.
- Muhammad Hazwan, H., Hasfalina, C. M., Hishamuddin, J., & Zurina, Z. A. (2012). Optimization and Kinetics of Essential Oil Extraction from Citronella Grass by Ohmic Heated Hydro Distillation. *International Journal of Chemical Engineering and Applications*, 173-177.
- Muralidhar, R., Chirumamila, R., Marchant, R., & Nigam, P. (2001). A response surface approach for the comparison of lipase production by *Candida cylindracea* using two different carbon sources. *Biochemical Engineering Journal*, 17-23.
- Neustadt, J., & Pieczenik, S. (2009). *Foundations and Applications of Medical Biochemistry in Clinical Practice*. New York, Bloomington: iUniverse.
- Nikolić, M., Jovanović, K. K., Marković, T., Marković, D., Gligorijević, N., Radulović, S., et al. (2014). Chemical composition, antimicrobial, and cytotoxic properties of five Lamiaceae essential oils. *Industrial Crops and Products*, 225-232.
- Nishaa, S., Vishnupriya, M., Sasikumar, J., & Gopalakrishnan, V. (2013). Phytochemical Screening and GC-MS Analysis of *Madhuca longifolia* (L) Macbr. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 52-59.
- Noordin, M., Venkatesh, V., Sharif, S., Elting, S., & Abdullah, A. (2004). Application of response surface methodology in describing the performance of coated carbide tools when turning AISI 1045 steel. *Journal of Materials Processing Technology*, 46-58.
- Nor Azah, M., Chang, Y., Mailina, J., Abu Said, A., Husni S., A. M., Nor Hasnida, H., et al. (2008). Comparison of chemical profiles of selected gaharu oils from peninsular Malaysia. *The Malaysian Journal of Analytical Sciences*, 338 - 340.
- Nor Azah, M., Husni, S. S., Mailina, J., Sahrim, L., Abdul Majid, J., & Mohd Faridz, Z. (2013). Classification of agarwood (gaharu) by resin content. *Journal of Tropical Forest Science*, 213-219.

- Nor Azah, M., Ismail, N., Mailina, J., Taib, M., Rahiman, M., & Hafizi, Z. M. (2014). Chemometric study of selected agarwood oils by gas chromatography–mass spectrometry. *Journal of Tropical Forest Science*, 382–388.
- Okoh, O. O., Sadimenko, A., & Afolayan, A. J. (2010). Comparative evaluation of the antibacterial activities of the essential oils of *Rosmarinus officinalis* L. obtained by hydrodistillation and solvent free microwave extraction methods. *Food Chemistry*, 308–312.
- Okugawa, H., Ueda, R., Matsumoto, K., Kawanishi, K., & Kato, A. (1996). Effect of jinkoh-eremol and agarospirol from agarwood on the central nervous system in mice. *Planta Med*, 2-6.
- Okugawa, H., Ueda, R., Matsumoto, K., Kawanishi, K., & Kato, A. (2000). Effects of sesquiterpenoids from “Oriental incenses” on acetic acid-induced writhing and D2 and 5-HT2A receptors in rat brain. *Phytomedicine* , 417–422.
- Oldfield, S., Newton, A., Fragoso, G., Mathew, P., Miles, L., & Edwards, M. (1998). *Towards a Global Tree Conservation Atlas: Mapping the status and distribution of the world's threatened tree species*. Cambridge: UNEP-WCMC, FFI.
- Ong, E. S., & Len, S. M. (2003). Pressurized hot water extraction of berberine, baicalein and glycyrrhizin in medicinal plants. *Analytica Chimica Acta*, 81-89.
- Ong, E. S., Cheong, J. S., & Goh, D. (2006). Pressurized hot water extraction of bioactive or marker compounds in botanicals and medicinal plant materials. *Journal of Chromatography A*, 92–102.
- Ormeno, E., Goldstein, A., & Niinemets, U. (2011). Extracting and trapping biogenic volatile organic compounds stored in plant species. *Trends in Analytical Chemistry*, 978-989.
- Oyen, L., & Dung, N. X. (1999). *Plant Resources of South-East Asia No. 19 : Essential-oil plants*. Leiden: Backhuys Publishers.
- Ozel, M., & Kaymaz, H. (2004). Superheated water extraction, steam distillation and Soxhlet extraction of essential oils of *Origanum onites*. *Anal Bioanal Chem.*, 1127-1133.
- Palmitic acid*. (2016 , june). Retrieved from wikipedia: https://en.wikipedia.org/wiki/Palmitic_acid
- Paoli, G. D., Peart, D. R., Leighton, M., & Samsedin, I. (2001). An ecological and economic assessment of the non-timber forest product gaharu wood in Gunung Palung National Park, West Kalimantan, Indonesia. *Conservation Biology* , 1721–1732.
- Pathak, M. D., & Khan, Z. (1994). *Insect pests of rice*. Manila, Philippines: IRRI, ICIPE.
- Paunović, D. D., Mitić, S. S., Kostić, D. A., Mitić, M. N., Stojanović, B. T., & Pavlović, J. L. (2014). Kinetics and thermodynamics of the solid-liquid extraction process of total polyphenols from barley. *Advanced technologies*, 58-63.
- Perricone, M., Arace, E., Corbo, M. R., Sinigaglia, M., & Bevilacqua, A. (2015). Bioactivity of essential oils: a review on their interaction with food components. *Frontiers in microbiology*, 1-7.
- Plaza, M., & Turner, C. (2015). Pressurized hot water extraction of bioactives. *Trends in Analytical Chemistry*, 39-54.
- Plaza, M., Abrahamsson, V., & Turner, C. (2013). Extraction and Neof ormation of Antioxidant Compounds by Pressurized Hot Water Extraction from Apple Byproducts. *Journal of agricultural and food chemistry*, 5500–5510.

- Pornpunyapat, J., Chetpattananondh, P., & Tongurai, C. (2011). Mathematical modeling for extraction of essential oil from *Aquilaria crassna* by hydrodistillation and quality of agarwood oil. *A Journal of the Bangladesh Pharmacological Society*, 18-24.
- Pripdeevech, P., Khummueng, W., & Park, S.-K. (2011). Identification of Odor-active Components of Agarwood Essential Oils from Thailand by Solid Phase Microextraction-GC/MS and GC-O. *Journal of Essential Oil Research*.
- Quan, L. T., Tran, Q. K., Kouda, K., Nguyen, N. T., Maruyama, Y., Saiki, I., et al. (2003). A survey on agarwood in Vietnam. *Journal of Traditional Medicine*, 124-131.
- Rahimi, N., Shiva, M., Mortazavi, S. A., Elhamirad, A. H., Maskooki, A. M., & Rajabzadeh, G. (2015). Kinetic study of superheated water extraction of berberine from *Berberis vulgaris* root. *Bulgarian Chemical Communications*, 140 – 146.
- Rajeswari, G., Murugan, M., & Mohan, V. (2012). GC-MS analysis of bioactive components of *Hugonia mystax* L. (Linaceae). *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 301-308.
- Ramesh, B. (2013). Modern Techniques of Materials Characterisation. *Mallikarjunachari Gangapuram*.
- Ranitha, M., Abdurahman, H. N., Ziad, A. S., Azhari, H. N., Thana, & S., R. (2014). A Comparative Study of Lemongrass (*Cymbopogon Citratus*) Essential Oil Extracted by Microwave-Assisted Hydrodistillation (MAHD) and Conventional Hydrodistillation (HD) Method. *International Journal of Chemical Engineering and Applications*, 104-108.
- Ravindra, P., Bono, A., & Chu, C. (2013). Developments in sustainable chemical and bioprocess technology. *Springer Science & Business Media*.
- Reverchon, E. (1997). Supercritical fluid extraction and fractionation of essential oils and related products. *The Journal of Supercritical Fluids*, 1–37.
- Rivera, L. L., Vilarem, G., Gomez, R. S., Estrada, M. J., & Feijoo, J. V. (2010). Water Soluble Fractions of Caraway (*Carum carvi* L.) Essential Oil. *BOLETIN LATINOAMERICANO Y DEL CARIBE DE PLANTAS MEDICINALES Y AROMATICAS*, 495 - 500.
- Robeva, R., & Hodge, T. (2013). *Mathematical Concepts and Methods in Modern Biology: Using Modern Discrete Models*. San Diego, USA: Elsevier Inc. Academic Press.
- Rose, A. H. (1993). *Adv in Microbial Physiology APL, Volume 34*. NY: Academic Press.
- Rouquerol, F., Rouquerol, J., & Sing, K. (1999). *Adsorption by Powders & Porous Solids*. London: Academic Press.
- Rovio, S., Hartonen, K., Holm, Y., Hiltunen, R., & Riekkola, M. L. (1999). Extraction of Clove Using Pressurized Hot Water. *Flavour and Fragrance Journal*, 399-404.
- Rowe, D. J. (2005). *Chemistry and Technology of Flavors and Fragrances*. Oxford: Blackwell Publishing Ltd.
- Ruano, F., Hefetz, A., Lenoir, A., Francke, W., & Tinaut, A. (2005). Dufour's gland secretion as a repellent used during usurpation by the slave-maker ant *Rossomyrmex minuchae*. *Journal of Insect Physiology*, 1158-1164.
- Safaralie, A., Fatemi, S., & Sefidkon, F. (2008). Essential oil composition of *Valeriana officinalis* L. roots cultivated in Iran: Comparative analysis between supercritical CO₂ extraction and hydrodistillation. *Journal of Chromatography A*, 159–164.

- Salt, D. W., Ajmani, S., Crichton, R., & Livingstone, D. J. (2007). An improved approximation to the estimation of the critical F values in best subset regression. *Journal of Chemical Information and Modeling*, 143–149.
- Sangarunlert, W., Piumsomboon, P., & Ngamprasertsith, S. (2007). Furfural production by acid hydrolysis and supercritical carbon dioxide extraction from rice husk. *Korean Journal of Chemical Engineering*, 936-941.
- Sangwan, N., Farooqi, A., Shabih, F., & Sangwan, R. (2001). Regulation of essential oil production in plants. *Plant Growth Regulation*, 3-21.
- Sarip, M. S., Morad, N. A., Yamashita, Y., Tsuji, T., Yunus, M. A., Aziz, M. K., et al. (2016). Crude palm oil (CPO) extraction using hot compressed water (HCW). *Separation and Purification Technology*, 103-112.
- Sarker, S. D., Latif, Z., & Gray, A. I. (2006). *Methods in Biotechnology Natural Products Isolation*. New Jersey: Human press Inc.
- Savage, P. E., Gopalan, S., Mizan, T. I., Martino, C. J., & Brock, E. E. (1985). Reactions at Supercritical Conditions: Applications and Fundamentals. *Reactors, Kinetics and Catalysis*, 1723-1778.
- Sen, S., Talukdar, N. C., & Khan, M. (2015). A simple metabolite profiling approach reveals critical biomolecular linkages in fragrant agarwood oil production from *Aquilaria malaccensis* – a traditional agro-based industry in North East India. *Current Science*, 63-71.
- Sermakkani, M., & Thangapandian, V. (2012). Gc-ms analysis of *Cassia italica* leaf methanol extract. *Asian J Pharm Clin Res*, 90-94.
- Sharifi, A., Mortazavi, S., Maskooki, A., Niakousari, M., & Elhamirad, A. (2013). Optimization of Subcritical Water Extraction of Bioactive Compounds from Barberry Fruit (*Berberis vulgaris*) by Using Response Surface Methodology. *International Journal of Agriculture and Crop Sciences*, 89-96.
- Sharma, H. (2008). *Biotechnological Approaches for Pest Management and Ecological Sustainability*. USA: CRC Press.
- Shobha, R. P., & Agrawal, R. (2007). Volatile compounds of therapeutic importance produced by *leuconostoc paramesenteroides*, a native laboratory isolate. *Turkish Journal of Biology*, 35-40.
- Sihag, S., & Pathak, H. (2016). Biodegradation of 2T engine oil using soil microbe and gravimetric analysis. *International Journal of Scientific & Engineering Research*, 1286-1294.
- Silva, C. F., Mendes, M. F., Pessoa, F. L., & Queiroz, E. M. (2008). Supercritical carbon dioxide extraction of macadamia (*Macadamia integrifolia*) nut oil: experiments and modeling. *Brazilian Journal of Chemical Engineering*, 175-181.
- Silva, L., Nelson, D., Drummond, M., Dufossé, L., & Glória, M. (2005). Comparison of hydrodistillation methods for the deodorization of turmeric. *Food Research International*, 1087–1096.
- Simon, M. J. (2003). Concrete mixture optimization using statistical methods : Final report.
- Soran, M.-L., Stan, M., Niinemets, Ü., & Copolovici, L. (2015). Influence of microwave frequency electromagnetic radiation on terpene emission and content in aromatic plants. *J Plant Physiol.*, 1436-1443.
- Stanisavljevic, I. T., Lazic, M. L., & Veljkovic, V. B. (2010). Kinetics of Hydrodistillation and Chemical Composition of Essential Oil from Cherry Laurel (*Prunus laurocerasus* L. var. *serbica* Pancic) Leaves. *Journal of Essential Oil Research*, 564-567.

- Sulaiman, N., Idayu, M. I., Ramlan, A., Fashya, M. N., Farahiyah, A. N., Mailina, J., et al. (2015). Effects of extraction methods on yield and chemical compounds of gaharu (*aquilaria malaccensis*). *Journal of Tropical Forest Science*, 413-419.
- Suresh, K., Babu, S. S., & Harisaranraj, R. (2008). Studies on In Vitro Antimicrobial Activity of Ethanol Extract of *Rauvolfia tetraphylla*. *Ethnobotanical Leaflets*, 586-590.
- Tajuddin, S. N., & Yusoff, M. M. (2010). Chemical Composition of Volatile Oils of *Aquilaria malaccensis* (Thymelaeaceae) from Malaysia. *Natural Product Communications*, 1965 - 1968.
- Tajuddin, S. N., Muhamad, N. S., Yarmo, M. A., & Yusoff, M. M. (2013). Characterization of the chemical constituents of agarwood oils from Malaysia by comprehensive two-dimensional gas chromatography–time-of-flight mass spectrometry. *Mendeleev Communications*, 51–52.
- Takemoto, H., Ito, M., Shiraki, T., Yagura, T., & Honda, G. (2008). Sedative effects of vapor inhalation of agarwood oil and spikenard extract and identification of their active components. *Journal Of Natural Medicines*, 41–46.
- Tam, C., Yang, F., Zhang, Q., Guan, J., & Li, S. (2007). Optimization and comparison of three methods for extraction of volatile compounds from *Cyperus rotundus* evaluated by gas chromatography–mass spectrometry. *Journal of Pharmaceutical and Biomedical Analysis*, 444-449.
- Tamuli, P., Boruah, P., Nath, S. C., & Leclercq, P. (2005). Essential Oil of Eaglewood Tree: a Product of Pathogenesis. *Journal of essential oil research*, 601-604.
- Tan, Y. H., Davis, J. A., Fujikawa, K., Ganesh, N. V., Demchenko, A. V., & Stine, K. J. (2012). Surface area and pore size characteristics of nanoporous gold subjected to thermal, mechanical, or surface modification studied using gas adsorption isotherms, cyclic voltammetry, thermogravimetric analysis, and scanning electron microscopy. *Journal of materials chemistry*, 6733-6745.
- Tanko, H., Carrier, D. J., Duan, L., & Clause, E. (2005). Pre- and post-harvest processing of medicinal plants. *Plant Genetic Resources*, 304-313.
- Tatsuki, S. H. (1985). Identification of possible sex pheromone of the yellow stem borer moth *Scirpophaga incertulas*. *Appl. Entomol. Zool*, 20: 357–359.
- Teo, C. C., Tan, S. N., Yong, J. W., Hew, C. S., & Ong, E. S. (2008). Evaluation of the extraction efficiency of thermally labile bioactive compounds in *Gastrodia elata* Blume by pressurized hot water extraction and microwave-assisted extraction. *Journal of Chromatography A*, 34-40.
- Teo, C. C., Tan, S. N., Yong, J. W., Hew, C. S., & Ong, E. S. (2010). Pressurized hot water extraction (PHWE). *Journal of Chromatography A*, 2484–2494.
- Teófilo, R. F., & Ferreira, M. M. (2006). Quimiometria II: planilhas eletrônicas para cálculos de planejamentos experimentais, um tutorial. *Química Nova*, 338–350.
- Thakur, A., Panesar, P. S., & Saini, M. S. (2014). Response surface modeling of Lactic Acid extraction by emulsion liquid membrane: Box-Behnken experimental design. *International Journal of Biological, Veterinary, Agricultural and Food Engineering*, 8.
- Tiwari, R., & Rana, C. (2015). Plant Secondary Metabolites: A Review. *International Journal of Engineering Research and General Science*, 661-670.
- Tongnuanchan, P., & Benjakul, S. (2014). Essential Oils: Extraction, Bioactivities, and Their Uses for Food Preservation. *Journal of Food Science*, 1231-1249.
- Traub, R. E., & Rowley, G. L. (1991). An NCME instructional module on understanding reliability.

- Turek, C., & Stintzing, F. C. (2013). Stability of Essential Oils: A Review. *Comprehensive Reviews in Food Science and Food Safety*, 40-53.
- Ueda, J.-y., Imamura, L., Tezuka, Y., Tran, Q. L., Tsuda, M., & Kadota, S. (2006). New sesquiterpene from Vietnamese agarwood and its induction effect on brain-derived neurotrophic factor mRNA expression in vitro. *Bioorganic & Medicinal Chemistry*, 3571–3574.
- Ulitzur, S., & JW, H. (1979). Evidence for tetradecanal as the natural aldehyde in bacterial bioluminescence. *Proc Natl Acad Sci U S A*, 265-267.
- Vila, R., Santana, A. I., Pérez-Rosés, R., Valderrama, A., Castelli, M. V., Mendonca, S., et al. (2010). Composition and biological activity of the essential oil from leaves of *Plinia cerrocampaensis*, a new source of a-bisabolol. *Bioresource Technology*, 2510-2514.
- Wahyudiono, Machmudah, S., & Goto, M. (2013). Utilization of Sub and Supercritical Water Reactions in Resource Recovery of Biomass Wastes. *engineering journal*, 1-12.
- Wang, H., Liu, Y., Wei, S., & Yan, Z. (2012). Comparative seasonal variation and chemical composition of essential oils from the leaves and stems of *Schefflera heptaphylla* using microwave-assisted and conventional hydrodistillation. *Industrial Crops and Products*, 229–237.
- Wani, M. A., Farooqi, J. A., & Shah, W. A. (2016). Synthesis and Antimicrobial Activity of Some Furfural Substituted Bisdimedone Derivatives. *Journal of Pharmaceutical and Applied Chemistry*, 53-57 .
- Watanabe, H., Inaba, H., & Hastings, J. W. (1991). Effects of aldehyde and internal ions on bioluminescence expression of *Photobacterium phosphoreum*. *Archives of Microbiology*, 1-4.
- Weber, W., Schoenmakers, R., Keller, B., Gitzinger, M., Grau, T., Daoud-El Baba, M., et al. (2008). A synthetic mammalian gene circuit reveals antituberculosis compounds. *Proceedings of the National Academy of Sciences*, 9994-9998.
- Widiyarti, G., Abbas, J., & Anita, Y. (2014). Biotransformation and cytotoxic activity of guaiacol dimer. *Indonesian Journal of Chemistry*, 179-184.
- Wil, N. N., Omar, N. A., Ibrahim, N. A., & Tajuddin, S. N. (2014). In vitro antioxidant activity and phytochemical screening of *Aquilaria malaccensis* leaf extracts. *Journal of Chemical and Pharmaceutical Research*, 688-693 .
- Wink, M. (1988). Plant breeding: importance of plant secondary metabolites for protection against pathogens and herbivores. *Theor Appl. Genet*, 225-233.
- Wolff, K., & Nguyen, L. (2012). Strategies for potentiation of ethionamide and folate antagonists against *Mycobacterium tuberculosis*. *Expert Rev Anti Infect Ther*, 971-981.
- Yadav, D., Mudgal, V., Agrawal, J., Maurya, A., Bawankule, D., Chanotiya, C., et al. (2013). Molecular docking and ADME studies of natural compounds of Agarwood oil for topical anti-inflammatory activity. *Current Computer - Aided Drug Design*, 360-70.
- Yang, K., Zhou, Y., Wang, C., Du, S., Deng, Z., Liu, Q., et al. (2011). Toxicity of *Rhododendron anthopogonoides* Essential Oil and Its Constituent Compounds towards *Sitophilus zeamais*. *Molecules*, 7320-7330.
- Yang, L., Qu, H., Mao, G., Zhao, T., Li, F., Zhu, B., et al. (2013). Optimization of subcritical water extraction of polysaccharides from *Grifola frondosa* using response surface methodology. *Pharmacognosy Magazine*, 120-129.
- Yoswathana, N., Eshiaghi, M., & Jaturapornpanich, K. (2012). Enhancement of Essential Oil from Agarwood by Subcritical Water Extraction and A

- Pretreatments on Hydrodistillation. *International Journal of Chemical, Molecular, Nuclear, Materials and Metallurgical Engineering*, 453-459.
- Yusoff, N. A., Tajuddin, S. N., Hisyam, A., & Omar, N. A. (2015). Agarwood Essential Oil: Study on Optimum Parameter and Chemical Compounds of Hydrodistillation Extraction. *Journal of Applied Science and Agriculture*, 1-5.
- Zaridah, M. Z., Azah, M. A., & Rohani, A. (2006). MOSQUITOCIDAL ACTIVITIES OF MALAYSIAN PLANTS. *Journal of Tropical Forest Science*, 74-80.
- Zeitsch, K. (2000). *The Chemistry and Technology of Furfural and its Many By-Products*. Amsterdam: Elsevier.
- Zeković, Z., Vidović, S., Vladić, J., Radosavljević, R., Cvejic, A., Elgndi, M. A., et al. (2014). Optimization of subcritical water extraction of antioxidants from *Coriandrum sativum* seeds by response surface methodology. *The Journal of Supercritical Fluids*, 560-566.
- Zuzarte, M., & Salgueiro, L. (2015). Essential Oils Chemistry. In *Bioactive Essential Oils and Cancer* (pp. 19-28). Switzerland: Springer International Publishing.