



UNIVERSITI PUTRA MALAYSIA

***SOCIAL COGNITIVE PREDICTORS OF PHYSICAL ACTIVITY
AMONG HEALTH WORKERS***

TAN HOOI SHYUAN

FPSK(M) 2017 39



**SOCIAL COGNITIVE PREDICTORS OF PHYSICAL ACTIVITY
AMONG HEALTH WORKERS**

By

TAN HOOI SHYUAN

**Dissertation Submitted to the Department of Community Health,
Faculty of Medicine and Health Sciences, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Public Health**

August 2017

All material contained within the dissertation, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the dissertation for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of dissertation presented to the Department of Community Health,
Universiti Putra Malaysia in fulfilment of the requirement for the
Degree of Master of Public Health

SOCIAL COGNITIVE PREDICTORS OF PHYSICAL ACTIVITY AMONG HEALTH WORKERS

By

TAN HOOI SHYUAN

August 2017

Chairman : Dr. Ahmad Azuhairi bin Ariffin
Faculty : Medicine and Health Sciences

Background: In Malaysia, about 1 in every 3 adults are inactive. Health workers do not perform better in physical activity although they are seen as role models. Understanding of social cognitive predictors is important to motivate them to initiate and maintain regular physical activity.

Objective: The study aimed to determine social cognitive predictors of physical activity among health workers.

Methodology: An analytical cross sectional study was done among 310 health workers recruited by simple random sampling using validated self-administered questionnaires. Hierarchical multiple regression analysis was used to determine the predictors.

Results: The respondents spent 141 minutes (IQR=182.7) in total physical activity per day. Leisure time physical activity contributed to 16.6% of the total physical activity, with the remaining were non-leisure time physical activities during occupational, household chores and transportation. Leisure time physical activity had significant positive correlation with goal setting ($r=0.344$, $p<0.001$), self-efficacy ($r=0.283$, $p<0.001$), friends support ($r=0.305$, $p<0.001$), outcome expectation ($r=0.284$, $p<0.001$), family support ($r=0.193$, $p=0.001$) and male gender ($p=0.002$), while had significant negative correlation with barriers ($r=-0.256$, $p<0.001$) and present of family history of medical illness ($p=0.029$). Hierarchical multiple regression analysis found that 23.0% (adjusted $r^2 = 0.218$, $p=0.029$) of variance in leisure time physical activity was explained by social cognitive predictors with goal setting ($r^2=0.118$, $p<0.001$), barriers ($r^2=0.046$, $p<0.001$), self-efficacy ($r^2=0.033$, $p<0.001$), friend support ($r^2=0.021$, $p=0.004$) and outcome expectation ($r^2=0.012$, $p=0.029$). Gender added an additional of 1.8% of the variance ($r^2=0.018$, $p=0.007$).

Conclusion: Social cognitive predictors are useful to explain leisure time physical activity. The findings can be used in development of intervention to encourage behavioural change.

Keywords: physical activity, social cognitive predictors, health workers



Abstrak disertasi yang dikemukakan kepada Jabatan Kesihatan Komuniti,
Universiti Putra Malaysia sebagai memenuhi keperluan untuk
Ijazah Sarjana Kesihatan Awam

FAKTOR PERAMAL SOSIAL KOGNITIF YANG MEMPENGARUHI AKTIVITI FIZIKAL DI KALANGAN PEKERJA KESIHATAN

Oleh

TAN HOOI SHYUAN

Ogos 2017

Pengerusi: Dr Ahmad Azuhairi bin Ariffin
Fakulti: Perubatan dan Sains Kesihatan

Latar belakang: Di Malaysia, 1 dalam setiap 3 orang dewasa adalah tidak aktif. Pekerja kesihatan tidak lebih aktif dalam aktiviti fizikal walaupun selalu dianggap sebagai contoh. Memahami faktor-faktor peramal sosial kognitif adalah penting supaya dapat memberikan motivasi kepada mereka dalam memulakan dan meneruskan aktiviti fizikal.

Objektif: Objektif kajian ini adalah untuk menentukan faktor-faktor peramal sosial kognitif dalam aktiviti fizikal di kalangan pekerja kesihatan.

Metodologi: Kajian keratan rentas analitikal dijalankan di antara 310 orang pekerja kesihatan yang dipilih melalui pensampelan rawak ringkas dengan menggunakan soalan kaji selidik yang disahkan. Analisis pelbagai regresi secara hierarki telah digunakan untuk menentukan faktor-faktor peramal.

Keputusan: Responden menggunakan sebanyak 141 minit (IQR=182.7) dalam aktiviti fizikal seharian. Aktiviti fizikal waktu lapang hanya menyumbang kepada 16.6% daripada jumlah aktiviti fizikal, dan selebihnya daripada aktiviti fizikal bukan waktu lapang, iaitu semasa kerja bergaji, kerja rumah dan pengangkutan. Aktiviti fizikal waktu lapang mempunyai korelasi positif yang signifikan dengan penetapan matlamat ($r=0.334$, $p<0.001$), efikasi diri ($r=0.283$, $p<0.001$), sokongan kawan ($r=0.305$, $p<0.001$), kesan jangkaan ($r=0.283$, $p<0.001$), sokongan keluarga ($r=0.193$, $p=0.001$) dan kaum lelaki ($p=0.002$), manakala mempunyai korelasi negatif yang signifikan dengan halangan ($r=-0.256$, $p<0.001$) dan sejarah keluarga berpenyakit ($p=0.029$). Analisis pelbagai regresi secara hierarki menunjukkan sebanyak 23% (r^2 selaras=0.218, $p=0.029$) variasi dalam aktiviti fizikal waktu lapang dapat diterangkan oleh faktor peramal sosial kognitif, iaitu penetapan matlamat ($r^2=0.118$, $p<0.001$), halangan ($r^2=0.046$, $p<0.001$), efikasi diri ($r^2=0.033$, $p<0.001$), sokongan kawan ($r^2=0.021$,

$p=0.004$) dan kesan jangkaan ($r^2=0.012$, $p=0.029$). Pembolehubah jantina menyumbangkan 1.8% lagi dalam variasi aktiviti fizikal waktu lapang.

Kesimpulan: Faktor peramal sosial kognitif berguna untuk menerangkan variasi aktiviti fizikal waktu lapang. Hasil kajian ini boleh digunakan dalam perkembangan program intervensi untuk menggalakkan perubahan tingkah laku.

Kata kunci: Aktiviti fizikal, faktor peramal sosial kognitif, pekerja kesihatan



ACKNOWLEDGEMENTS

I would like to express my gratitude to my supervisor and co-supervisor, Dr Ahmad Azuhairi Ariffin and Associate Prof Dr Nor Afiah Mohd Zulkefli for guiding and advising me in this dissertation.

I am deeply thankful to my mentors from Ministry of Health, Dr Feisul Idzwan Mustapha and Dr Mohd Azahadi Omar for their guidance, encouragement and valuable feedback along this journey.

I truly appreciate the assistance from Dr Noraishah Jaafar and her team during the data collection period.

I am grateful to my dissertation examination committee including Associate Prof Dr Muhamad Hanafiah Juni, Dr Titi Rahmawati Hamedon and Professor Dr Krishna Gopal Rampal for their invaluable insights that helped me to improve the quality of this dissertation.

I would like to thank my colleagues who never fail to support and help each other whenever needed.

Most of all, I am blessed to have a wonderful husband and family. I would not be able to make this happen without their understanding, support and love.

I certify that a Dissertation Examination Committee has met on 3rd August 2017 to conduct the final examination of Tan Hooi Shyuan on her dissertation entitled “Social cognitive predictors of physical activity among health workers” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U. (A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Public Health.

Members of the Dissertation Examination Committee were as follows:

Dr. Muhamad Hanafiah Juni

MD (UKM), MPH (UM), MSc (London School of Economic)

Associate Professor (Medical)

Department of Community Health

Faculty of Medicine and Health Sciences

Universiti Putra Malaysia

(Chairman)

Dr. Titi Rahmawati Hamedon

MD (UKM), MPH (Occupational Health) (UKM)

Senior lecturer (Medical)

Department of Community Health

Faculty of Medicine and Health Sciences

Universiti Putra Malaysia

(Internal Examiner)

Dr. Krishna Gopal Rampal, PhD

MBBS (Agra), MPH (Mahidol), MPH (John Hopkins), PhD (Johns Hopkins),

FFOMI, FFOM, FAOEMM, FAMM

Professor (Community and Occupational)

Cyberjaya University College of Medical Sciences

(External Examiner)

Professor Dato' Dr. Abdul Jalil Nordin, DSIS

MD (UKM), M.Med. (Radiology) (UM)

Professor and Dean

Faculty of Medicine and Health Sciences

Universiti Putra Malaysia

Date:

This dissertation was submitted to Department of Community Health, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Public Health. The members of the Supervisory Committee were as follows:

Dr. Ahmad Azuhairi Ariffin
MBBS (KU), MCMed (OH) (UKM)
Senior lecturer (Medical)
Department of Community Health
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Dr. Nor Afiah Mohd Zulkefli, PhD
B.Med.Sc., MD (UKM), M. Community Health (Family Health) (UKM), PhD (UKM)
Associate Proffesor (Medical)
Department of Community Health
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Professor Dato' Dr. Abdul Jalil Nordin, DSIS
MD (UKM), M.Med. (Radiology) (UM)
Professor and Dean
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this dissertation is my original work;
- quotations, illustrations and citations have been duly referenced;
- this dissertation has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the dissertation and copyright of dissertation are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before dissertation is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the dissertation, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The dissertation has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No: Tan Hooi Shyuan, GS47196

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this dissertation was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Chairman of Supervisory Committee
Dr. Ahmad Azuhairi Ariffin
MBBS (KU), MCMed (OH) (UKM)

Signature: _____
Member of Supervisory Committee
Associate Professor Dr Nor Afiah Mohd Zulkefli
B.Med.Sc., MD (UKM), M. Community Health (Family Health) (UKM), PhD (UKM)

TABLE OF CONTENTS

ABSTRACT	Page
<i>ABSTRAK</i>	i
ACKNOWLEDGEMENTS	iii
APPROVAL	v
DECLARATION	vi
LIST OF TABLES	viii
LIST OF APPENDICES	xiv
LIST OF ABBREVIATIONS	xvi
CHAPTER	xvii
1 INTRODUCTION	1
1.1 Background	1
1.2 Problem Statement	2
1.3 Research Questions	2
1.4 Objectives of Study	3
1.4.1 General Objective	3
1.4.2 Specific Objectives	3
1.5 Hypothesis	3
1.6 Definition of Terms	4
1.6.1 Social Cognitive Predictors	4
1.6.2 Physical Activity	4
1.6.3 Health Workers	4
2 LITERATURE REVIEW	5
2.1 Burden of Non-Communicable Diseases and Its Risk Factors	5
2.2 Importance of Physical Activity	5
2.3 Prevalence of Physical Inactivity and Its Associated Sociodemographic Characteristics	6
2.4 Physical Activity among Health Workers	7
2.5 Social Cognitive Theory and Its Constructs in Physical Activity Behaviour	8
2.5.1 Self-efficacy	8
2.5.2 Outcome Expectation	9
2.5.3 Goal Setting	9
2.5.4 Barriers	10
2.5.5 Family and Friends Support	10
2.6 Other Correlates of Physical Activity	11
2.7 Conceptual Framework	11
Fig 2.1 The conceptual framework of the predictors of physical activity among health workers	12

3	METHODOLOGY	13
3.1	Study Location	13
3.2	Study Design	13
3.3	Study Duration	13
3.4	Study Population	13
3.5	Selection Criteria	13
	3.5.1 Inclusion Criteria	13
	3.5.2 Exclusion Criteria	13
3.6	Sampling Frame	14
3.7	Sampling Unit	14
3.8	Sampling Method	14
3.9	Sampling Size	14
3.10	Independent Variables	15
3.11	Dependent Variable	15
3.12	Operational Definition	15
	3.12.1 Gender	15
	3.12.2 Age	15
	3.12.3 Body Mass Index	15
	3.12.4 Ethnicity	16
	3.12.5 Marital Status	16
	3.12.6 Occupation	16
	3.12.7 Medical Illness	16
	3.12.8 Family History of Medical Illness	16
	3.12.9 Self-efficacy	16
	3.12.10 Outcome Expectation	16
	3.12.11 Goal Setting	17
	3.12.12 Barriers	17
	3.12.13 Family and Friends Support	17
	3.12.14 Physical Activity	17
3.13	Study Instrument	17
	3.13.1 Variables Measured by Instrument	17
	3.13.2 Validity of Instrument	19
	3.13.3 Reliability of Instrument	19
3.14	Data Collection	20
3.15	Data Analysis	20
3.16	Benefits and Risks for Participants	21
3.17	Informed Consent	21
3.18	Privacy and Confidentiality	21
3.19	Publication of the Study	21
3.20	Ethical Approval	21
4	RESULTS	22
4.1	Response Rate	22
4.2	Normality Tests of Variables	22
4.3	Sociodemographic and Health Status of Respondents	22
4.4	Social Cognitive Score of Physical Activity among respondents	24
4.5	Physical Activity of Respondents	24

4.6	Association between Physical Activity and Sociodemographic, Health Status	26
4.6.1	Gender	26
4.6.2	Age	27
4.6.3	Body Mass Index	27
4.6.4	Ethnicity	29
4.6.5	Marital Status	29
4.6.6	Occupation	29
4.6.7	Medical Illness	29
4.6.8	Family History of Medical Illness	29
4.7	Association between Physical Activity and Social Cognitive Constructs	31
4.7.1	Self-efficacy	31
4.7.2	Outcome Expectation	31
4.7.3	Goal Setting	31
4.7.4	Barriers	31
4.7.5	Family and Friends Support	31
4.8	Social Cognitive Predictors of Physical Activity	35
4.8.1	Social Cognitive Predictors of Total Physical Activity	35
4.8.2	Social Cognitive Predictors of Non-leisure Time Physical Activity	36
4.8.3	Social Cognitive Predictors of Leisure Time Physical Activity	38
5	DISCUSSION	41
5.1	Sociodemographic Characteristics of Respondents	41
5.2	Physical Activity of Respondents	41
5.3	Association between Physical Activity and Sociodemographic, Health Status	43
5.3.1	Gender	43
5.3.2	Age	44
5.3.3	Body Mass Index	44
5.3.4	Ethnicity	44
5.3.5	Marital Status	45
5.3.6	Occupation	45
5.3.7	Medical Illness	45
5.3.8	Family History of Medical Illness	45
5.4	Association between Physical Activity and Social Cognitive Constructs	46
5.4.1	Self-efficacy	46
5.4.2	Outcome Expectation	46
5.4.3	Goal Setting	46
5.4.4	Barriers	47
5.4.5	Family and Friends Support	48
5.5	Social Cognitive Predictors of Physical Activity	48

6	SUMMARY, CONCLUSION AND RECOMMENDATIONS	50
6.1	Strengths of the Study	50
6.2	Limitations of the Study	50
6.3	Recommendations	51
	REFERENCES	52
	APPENDICES	60
	BIODATA OF STUDENT	99



LIST OF TABLES

Table	Page
3.1 The internal consistency reliability tests	20
4.1. Sociodemographic and health status of respondents (N=310)	23
4.2. Social cognitive score among respondent (N=310)	24
4.3. Physical activity and sitting time of respondents per day in minutes (N=310)	25
4.4. Respondents without physical activity in different settings and intensity (N=310)	25
4.5. Composition (%) of physical activity	26
4.6. Physical activity of respondents in MET-minutes per week (N=310)	26
4.7. Physical activity per day in minutes by gender	27
4.8. Physical activity per day in minutes by age	28
4.9. Physical activity per day in minutes by body mass index	28
4.10. Physical activity per day in minutes by ethnicity, marital status, occupation, medical illness and family history of medical illness	30
4.11. Pearson Correlation and simple linear regression result analysis between total physical activity and sociodemographic, health status, social cognitive constructs	32
4.12. Pearson Correlation and simple linear regression result analysis between Non-leisure Time Physical Activity and sociodemographic, health status, social cognitive constructs	33
4.13. Pearson Correlation and simple linear regression result analysis between Leisure Time Physical Activity and sociodemographic, health status, social cognitive constructs	34

4.14.	Hierarchical multiple regression analysis summary for predictors of total physical activity (N=310)	35
4.15.	Model summary for predictors of total physical activity (N=310)	36
4.16.	Hierarchical multiple regression analysis summary for predictors of Non-leisure Time Physical Activity (N=310)	37
4.17.	Model summary for predictors of Non-leisure Time Physical Activity (N=310)	38
4.18.	Hierarchical multiple regression analysis summary for predictors of Leisure Time Physical Activity (N=310)	39
4.19.	Model summary for predictors of Leisure Time Physical Activity (N=310)	40

LIST OF APPENDICES

Appendix		Page
A1	Approval letter from the National Medical Research and Ethics Committee (NMREC) of National Institute of Health, Ministry of Health Malaysia	60
A2	Approval letter from the Ethics Committee for Research Involving Human Subjects of Universiti Putra Malaysia (JKEUPM)	62
B	Approval letter from Jabatan Kesihatan Negeri Pulau Pinang and Pejabat Kesihatan Daerah Seberang Perai Tengah	65
C1	Respondent's information sheet and consent-English version	68
C2	Respondent's information sheet and consent-Malay version	71
D1	Questionnaire-English version	74
D2	Questionnaire-Malay version	84
E1	Normality tests of variables	95
E2	Normality tests of variables after data transformation	96
E3	Itemised mean score for self-efficacy, outcome expectation, goal setting and barriers (N=310)	97
E4	Itemised mean score for family and friends support (N=310)	98

LIST OF ABBREVIATIONS

BMI	Body Mass Index
CI	Confidence Interval
HEPA	Health Enhancing Physical Activity
GPAQ	Global Physical Activity Questionnaire
IPAQ	International Physical Activity Questionnaire
LTPA	Leisure Time Physical Activity
M	Mean
MANS	Malaysian Adult Nutrition Survey
Mdn	Median
NCD	Non-Communicable Disease
NLTPA	Non-Leisure Time Physical Activity
NHMS	National Health and Morbidity Survey
PA	Physical Activity
SCT	Social Cognitive Theory
WHO	World Health Organisation



CHAPTER 1

INTRODUCTION

1.1 Background

Physical inactivity is the fourth leading risk factor for global mortality which contributes to 6% of deaths (World Health Organisation, 2009). Furthermore, physical inactivity increases the risk of non-communicable diseases (NCD), mainly coronary heart disease, type 2 diabetes, breast and colon cancers (Lee et al., 2012). In recognising the detrimental consequences of physical inactivity, World Health Organisation (WHO) aimed to have 10% relative reduction in physical inactivity prevalence by 2025 (World Health Organisation., 2013).

Physical inactivity is defined as failure to meet the minimum physical activity (PA) level recommended by World Health Organisation. Adult aged 18 to 64 years old are recommended to participate in moderate intensity aerobic PA for at least 150 minutes per week, or at least 75 minutes of vigorous intensity aerobic PA per week, or equivalent combination of both moderate and vigorous intensity activity (World Health Organisation, 2010). PA includes any activity that increases the energy expenditure, which could be done as part of occupation, household chores, transportation, or leisure time activity such as sport, game and planned exercise.

Despite various efforts in promoting PA, physical inactivity remains a public health concern. Worldwide, approaching one third or 31.1% of adults aged 15 years and above are physically inactive (Hallal et al., 2012). In Malaysia, the National Health and Morbidity Survey (NHMS) 2015 reported that 33.5% of adults aged 16 years and above were physically inactive (Institute for Public Health, 2015). With the rising prevalence of obesity and non-communicable diseases, physical inactivity as the main risk factor becomes even more relevant now. Therefore, prompt actions need to be taken to reduce physical inactivity in Malaysia.

Understanding the causes of PA behaviour or its predictors is essential for development and improvement of public health intervention in promoting regular physical activity. The complex PA behaviour is best to be studied by using a behavioural model. Bandura's social cognitive theory provides a useful framework to explain PA behaviour and has been tested across numerous populations (Mesters, Wahl, & Keulen, 2014; Young, Plotnikoff, Collins, Callister, & Morgan, 2014). The core constructs of social cognitive theory (SCT) are self-efficacy, outcome expectations, goals setting, barriers, family and friends support.

1.2 Problem Statement

In Penang state, the prevalence of physical inactivity increased from 19.9% in 2011 to 25.5% in 2015 as reported in NHMS (Institute for Public Health, 2011, 2015). The survey also revealed a worrying situation whereby 33.9% of government and semi-government employees were inactive (Institute for Public Health, 2015). Healthy workers are invaluable asset for an organisation. Poor health among health workers not only affected the individuals, but gave negative impact to the health system that safeguard population health (Blake & Harrison, 2013). Arvidson et al (2013) reported that higher level of leisure time physical activity was associated with improved work ability among workers in Sweden. Health workers are seen as role models in practicing regular PA. It was noted that physical activity habits of doctors influenced their counselling practice whereby physically active doctors were more likely to offer exercise counselling to patients (Lobelo, Duperly, & Frank, 2009). WHO also launched a healthy workplace initiative, "Walk the Talk", at WHO headquarters and regional offices in 2016 to promote healthy lifestyles includes PA while reflecting the commitment of WHO to set an example (World Health Organisation, 2017).

However, previous study done in a Malaysia university medical center showed that the prevalence of physical inactivity among the health worker respondents was 35.2% (Jamil et al., 2016). This was even higher than the prevalence of physical inactivity in Malaysia general population with 33.5% (Institute for Public Health, 2015). Another study among health workers working in a hospital in Malaysia showed high prevalence of obese (59%) and overweight (24%) which was significantly associated with poor physical activity (Ayiesah, Leonard, Vijaykumar, & Mohd Suhaimy, 2013).

In Malaysia, many researches have been conducted to study the sociodemographic characteristics of physical activity (Chan et al., 2014; Cheah & Poh, 2014; Lian, Bonn, Han, Choo, & Piau, 2016). However, little studies were done to address the social cognitive variables of PA, especially among health workers, which could explain the behaviour from another perspective.

1.3 Research Questions

- i. Is there any correlation between physical activity and sociodemographic, health status, self-efficacy, outcome expectation, goal setting, barriers, family and friends support among health workers?
- ii. What are the social cognitive predictors of physical activity among health workers?

1.4 Objectives of Study

1.4.1 General Objective

To determine social cognitive predictors of physical activity among health workers

1.4.2 Specific Objectives

- i. To describe the sociodemographic, health status, self-efficacy, outcome expectation, goal setting, barriers, family and friend support of physical activity among health workers.
- ii. To measure the total physical activity among health workers.
- iii. To measure the non-leisure time physical activity among health workers.
- iv. To measure the leisure time physical activity among health workers.
- v. To identify the association between total physical activity and sociodemographic, health status, and social cognitive constructs (self-efficacy, outcome expectation, goal setting, barriers, family and friend support).
- vi. To identify the association between non-leisure time physical activity and sociodemographic, health status, and social cognitive constructs (self-efficacy, outcome expectation, goal setting, barriers, family and friend support).
- vii. To identify the association between leisure time physical activity and sociodemographic, health status, and social cognitive constructs (self-efficacy, outcome expectation, goal setting, barriers, family and friend support).
- viii. To determine the social cognitive predictors of total physical activity among health workers.
- ix. To determine the social cognitive predictors of non-leisure physical activity among health workers.
- x. To determine the social cognitive predictors of leisure physical activity among health workers.

1.5 Hypothesis

- H₁: There is a significant association between total physical activity and sociodemographic, health status, self-efficacy, outcome expectation, goal setting, barriers, family and friends support among health workers.
- H₂: There is a significant association between non-leisure time physical activity and sociodemographic, health status, self-efficacy, outcome expectation, goal setting, barriers, family and friends support among health workers.
- H₃: There is a significant association between leisure time physical activity and sociodemographic, health status, self-efficacy, outcome expectation, goal setting, barriers, family and friends support among health workers.

1.6 Definition of Terms

1.6.1 Social Cognitive Predictors

Social cognitive predictors in this study referred to a set of constructs described in the Bandura's social cognitive theory of health behaviour (Bandura, 2004).

- a) self-efficacy that one can have control over PA
- b) outcome expectation or the expected costs and benefits in performing PA
- c) Goal setting in doing PA
- d) Barriers in carrying out PA
- e) family and friend support in involving PA

1.6.2 Physical Activity

According to Caspersen, Powell, & Christenson (1985), PA is defined as "any bodily movement produced by skeletal muscles that results in energy expenditure". PA includes activities of daily living, from occupational, household chores, transportation, or leisure time activity such as games, sports or exercises. Exercise is a subset of PA characterised by planned, structured and repetitive activity in order to improve or maintain health and physical fitness.

1.6.3 Health Workers

Health workers are defined as "all people whose job are to protect and improve the health of their communities" (World Health Organisation, 2006). There are two groups of health workers. The first group are health service providers who deliver health services. The second are health management and support workers who do not engage directly in the provision of service.

REFERENCES

- Anderson, E. S., Wojcik, J. R., Winett, R. a, & Williams, D. M. (2006). Social-cognitive determinants of physical activity: The influence of social support, self-efficacy, outcome expectations, and self-regulation among participants in a church-based health promotion study. *Health Psychology, 25*(4), 510–520.
- Anokye, N. K., Trueman, P., Green, C., Pavey, T. G., & Taylor, R. S. (2012). Physical activity and health related quality of life. *BMC Public Health, 12*(624), 1–8.
- Arem, H., Moore, S. C., Patel, A., Hartge, P., Berrington de Gonzalez, A., Visvanathan, K., ... Matthews, C. E. (2015). Leisure time physical activity and mortality. *JAMA Internal Medicine, 175*(6), 959–967.
- Armstrong, T., & Bull, F. (2006). Development of the World Health Organization Global Physical Activity Questionnaire (GPAQ). *Journal of Public Health, 14*(2), 66–70.
- Arvidson, E., Borjesson, M., Ahlborg, G., Lindegård, A., & Jonsdottir, I. H. (2013). The level of leisure time physical activity is associated with work ability-A cross sectional and prospective study of health care workers. *BMC Public Health, 13*(1), 855.
- Ayiesah, R., Leonard, J. H., Vijaykumar, P., & Mohd Suhaimy, R. (2013). Obesity and habitual physical activity level among staffs working in a military hospital in Malacca, Malaysia. *International Medical Journal Malaysia, 12*(1), 53–58.
- Bandura, A. (1977). Self-efficacy: Toward a unifying theory of behavioral change. *Psychological Review, 84*(2), 191–215.
- Bandura, A. (1986). *Social foundations of thought and action: A social cognitive theory*. Englewood Cliffs: Prentice-Hall.
- Bandura, A. (1994). Self-Efficacy. *Encyclopedia of human behavior* (Vol. 4). New York: Academic Press.
- Bandura, A. (1999). Social cognitive theory: An agentic perspective. *Asian Journal of Social Psychology, 2*(1), 21–41.
- Bandura, A. (2001). Social cognitive theory of mass communication. *Mediapsychology, 3*(3), 265–299.
- Bandura, A. (2004). Health promotion by social cognitive means. *Health Education & Behavior, 31*(2), 143–164.
- Bauman, A. E., Reis, R. S., Sallis, J. F., Wells, J. C., Loos, R. J. F., & Martin, B. W. (2012). Correlates of physical activity: Why are some people physically active and others not? *The Lancet, 380*(9838), 258–271.

- Bauman, A. E., Sallis, J. F., Dzewaltowski, D. A., & Owen, N. (2002). Toward a better understanding of the influences on physical activity: The role of determinants, correlates, causal variables, mediators, moderators and confounders. *American Journal of Preventive Medicine*, 23(2S), 5–14.
- Bauman, A., Ma, G., Cuevas, F., Omar, Z., Waqanivalu, T., Phongsavan, P., ... Bhushan, A. (2011). Cross-national comparisons of socioeconomic differences in the prevalence of leisure-time and occupational physical activity, and active commuting in six Asia-Pacific countries. *Journal of Epidemiology & Community Health*, 65(1), 35–43.
- Bellew, B., Bauman, A., Martin, B., Bull, F., & Matsudo, V. (2011). Public policy actions needed to promote physical activity. *Current Cardiovascular Risk Reports*, 5(4), 340–349.
- Biernat, E., Poznańska, A., & Gajewski, A. K. (2012). Is physical activity of medical personnel a role model for their patients. *Annals of Agricultural and Environmental Medicine*, 19(4), 707–710.
- Biswas, A., Oh, P. I., Faulkner, G. E., Bajaj, R. R., Silver, M. A., Mitchell, M. S., & Alter, D. A. (2015). Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults a systematic review and meta-analysis. *Annals of Internal Medicine*, 162(2), 123–132.
- Blake, H., & Harrison, C. (2013). Health behaviours and attitudes towards being role models. *British Journal of Nursing*, 22(2), 86–94.
- Bull, F. C., Maslin, T. S., & Armstrong, T. (2009). Global Physical Activity Questionnaire (GPAQ): Nine Country Reliability and Validity Study. *Journal of Physical Activity and Health*, 6(6), 790–804.
- Caspersen, C. J., Powell, K. E., & Christenson, G. M. (1985). Physical activity, exercise, and physical fitness: Definitions and distinctions for health-related research. *Public Health Reports*, 100(2), 126–131.
- Chan, Y. Y., Lim, K. K., Teh, C. H., Lim, K. H., Abd Hamid, H. A., Omar, M. A., ... Kee, C. C. (2014). Prevalence and factors associated with physical inactivity among Malaysian adults. *Southeast Asian Journal of Tropical Medicine and Public Health*, 45(2), 467–480.
- Chau, J. Y., Grunseit, A. C., Chey, T., Stamatakis, E., Brown, W. J., Matthews, C. E., ... van der Ploeg, H. P. (2013). Daily sitting time and all-cause mortality: A meta-analysis. *PLoS ONE*, 8(11), e80000.
- Cheah, Y. K. (2011). Influence of socio-demographic factors on physical activity participation in a sample of adults in Penang, Malaysia. *Malaysian Journal of Nutrition*, 17(3), 385–391.
- Cheah, Y. K., & Poh, B. K. (2014). The determinants of participation in physical activity in Malaysia. *Osong Public Health and Research Perspectives*, 5(1), 20–27.

- Cheah, Y. K., & Tan, A. K. G. (2014). Determinants of leisure-time physical activity: Evidence from Malaysia. *The Singapore Economic Review*, 59(2), 1450017.
- Choi, J. Y., Chang, A. K., & Choi, E.-J. (2015). Sex differences in social cognitive factors and physical activity in Korean college students. *Journal of Physical Therapy Science*, 27(6), 1659–1664.
- Church, T. S., Thomas, D. M., Tudor-Locke, C., Katzmarzyk, P. T., Earnest, C. P., Rodarte, R. Q., ... Bouchard, C. (2011). Trends over 5 decades in U.S. occupation-related physical activity and their associations with obesity. *PLoS ONE*, 6(5), e19657.
- Dans, A., Ng, N., Varghese, C., Tai, E. S., Firestone, R., & Bonita, R. (2011). The rise of chronic non-communicable diseases in southeast Asia: Time for action. *The Lancet*, 377, 680–689.
- Dixon, M. A. (2009). From their perspective: A qualitative examination of physical activity and sport programming for working mothers. *Sport Management Review*, 12(1), 34–48.
- Dlugonski, D., & Motl, R. W. (2014). Social cognitive correlates of physical activity among single mothers with young children. *Psychology of Sport and Exercise*, 15(6), 637–641.
- Dorough, A. E. (2005). *Thesis: The relationship of preferences and self-regulation among consistent exercisers*. Virginia Polytechnic Institute and State University.
- Eime, R. M., Young, J. A., Harvey, J. T., Charity, M. J., Payne, W. R., Brown, W., ... Deci, E. (2013). A systematic review of the psychological and social benefits of participation in sport for adults: Informing development of a conceptual model of health through sport. *International Journal of Behavioral Nutrition and Physical Activity*, 10(1), 135.
- Evans, L. K. (2011). Rural Black women's thoughts about exercise. *Applied Nursing Research*, 24(4), 200–206.
- Gebel, K., Ding, D., Chey, T., Stamatakis, E., Brown, W. J., & Bauman, A. E. (2015). Effect of moderate to vigorous physical activity on all-cause mortality in middle-aged and older Australians. *JAMA Internal Medicine*, 175(6), 970–977.
- Gordon, P. B., & Robert, S. B. (1994). A new sample size formula for regression. In *Annual meeting of the American Educational Research Association*. New Orleans, LA. <http://doi.org/ED419696>
- Greaves, C. J., Sheppard, K. E., Abraham, C., Hardeman, W., Roden, M., Evans, P. H., & Schwarz, P. (2011). Systematic review of reviews of intervention components associated with increased effectiveness in dietary and physical activity interventions. *BMC Public Health*, 11(1), 119.

- Hallal, P. C., Andersen, L. B., Bull, F. C., Guthold, R., Haskell, W., & Ekelund, U. (2012). Global physical activity levels: Surveillance progress, pitfalls, and prospects. *The Lancet*, 380(9838), 247–257.
- Holtermann, A., Marott, J. L., Gyntelberg, F., Søgaard, K., Suadicani, P., Mortensen, O. S., ... Schnohr, P. (2013). Does the benefit on survival from leisure time physical activity depend on physical activity at work? A prospective cohort study. *PLoS ONE*, 8(1), e54548.
- Ibrahim, S., Karim, N. A., Ng, L. O., & Wan Ngah, W. Z. (2013). Perceived physical activity barriers related to body weight status and sociodemographic factors among Malaysian men in Klang Valley. *BMC Public Health*, 13, 275.
- Institute for Public Health. (2011). *National Health and Morbidity Survey 2011. Vol. II: Non-communicable diseases*. Kuala Lumpur: Institute for Public Health, Ministry of Health Malaysia.
- Institute for Public Health. (2014). *National Health and Morbidity Survey 2014: Malaysian Adult Nutrition Survey (MANS). Vol II: Survey findings*. Kuala Lumpur: Institute for Public Health, Ministry of Health Malaysia.
- Institute for Public Health. (2015). *National Health and Morbidity Survey 2015. Vol. II: Non-communicable diseases, risk factors & other health problems*. Kuala Lumpur: Institute for Public Health, Ministry of Health Malaysia.
- IPAQ Research Committee. (2005). *Guidelines for data processing and analysis of the International Physical Activity Questionnaire (IPAQ) – Short and long forms*.
- Ismail, S., Mohd Zulkefli, N. A., Shaarin, M. T., Zainol A., N. F., & Tee, S. (2014). Predictors of physical activity among government servant in Malaysia. *International Journal of Public Health and Clinical Sciences*, 1(1), 80–87.
- Jamil, A. T., Ismail, A., Idris, I. B., Soo, K. C., Teng, A. J., Bahaman, N. A., & Fadzil, M. F. (2016). Levels of physical activity and its associated factors among health care workers. *Malaysian Journal of Public Health Medicine*, 16(3), 127–133.
- Jamil, A. T., Singh, R., Ismail, A., & Omar, A. (2015). Non-leisure time physical activity for adult Malaysian and determinant factors. *Malaysian Journal of Public Health Medicine*, 15(3), 84–93.
- Jekauc, D., Volkle, M., Wagner, M. O., Mess, F., Reiner, M., & Renner, B. (2015). Prediction of attendance at fitness center: A comparison between the theory of planned behavior, the social cognitive theory, and the physical activity maintenance theory. *Frontiers in Psychology*, 6(121), 1–10.
- Joseph, R. P., Ainsworth, B. E., Keller, C., & Dodgson, J. E. (2015). Barriers to physical activity among African American women: An integrative review of the literature. *Women Health*, 55(6), 679–699.

- Kandula, N. R., & Lauderdale, D. S. (2005). Leisure time, non-leisure time, and occupational physical activity in Asian Americans. *Annals of Epidemiology*, 15(4), 257–265.
- Kunene, S. H., & Taukobong, N. P. (2015). Level of physical activity of health professionals in a district hospital in KwaZulu-Natal , South Africa. *South African Journal of Physiotherapy*, 71(1), 1–6.
- Lee, I.-M., Shiroma, E. J., Lobelo, F., Puska, P., Blair, S. N., Katzmarzyk, P. T., ... Wells, J. C. (2012). Effect of physical inactivity on major non-communicable diseases worldwide: An analysis of burden of disease and life expectancy. *The Lancet*, 380(9838), 219–229.
- Lian, T. C., Bonn, G., Han, Y. S., Choo, Y. C., & Piau, W. C. (2016). Physical activity and its correlates among adults in Malaysia: A cross-sectional descriptive study. *PLoS ONE*, 11(6), e0157730.
- Lingesh, G., Khoo, S., Nahar, M., Mohamed, A., Taib, N. A., & Group, M. (2016). Comparing physical activity levels of Malay version of the IPAQ and GPAQ with accelerometer in nurses. *International Journal of Applied Exercise Physiology*, 5(3), 8–17.
- Lobelo, F., Duperly, J., & Frank, E. (2009). Physical activity habits of doctors and medical students influence their counselling practices. *British Journal of Sports Medicine*, 43(2), 89–92.
- Macniven, R., Bauman, A., & Abouzeid, M. (2012). A review of population-based prevalence studies of physical activity in adults in the Asia-Pacific region. *BMC Public Health*, 12(1), 41.
- Mailey, E. L., & McAuley, E. (2014). Impact of a brief intervention on physical activity and social cognitive determinants among working mothers: A randomized trial. *Journal of Behavioral Medicine*, 37(2), 343–355.
- McArthur, D., Dumas, A., Woodend, K., Beach, S., & Stacey, D. (2014). Factors influencing adherence to regular exercise in middle-aged women: a qualitative study to inform clinical practice. *BMC Women's Health*, 14(1), 49.
- Mesters, I., Wahl, S., & Keulen, V. H. M. (2014). Socio-demographic , medical and social-cognitive correlates of physical activity behavior among older adults (45 – 70 years): A cross-sectional study. *BMC Public Health*, 14(647), 1–12.
- Mo, P. K. H., Blake, H., & Batt, M. E. (2011). Getting healthcare staff more active: The mediating role of self-efficacy. *British Journal of Health Psychology*, 16(4), 690–706.
- Molanorouzi, K., Khoo, S., & Morris, T. (2015). Motives for adult participation in physical activity: Type of activity, age, and gender. *BMC Public Health*, 15(1), 66.

- Nocon, M., Hiemann, T., Muller-Riemenschneider, F., Thalau, F., Roll, S., & Williich, S. (2008). Association of physical activity with all-cause and cardiovascular mortality: A systematic review and meta-analysis. *European Journal of Cardiovascular Prevention & Rehabilitation*, 15(3), 239–46.
- Pan, S. Y., Cameron, C., DesMeules, M., Morrison, H., Craig, C. L., & Jiang, X. (2009). Individual, social, environmental, and physical environmental correlates with physical activity among Canadians: A cross-sectional study. *BMC Public Health*, 9(1), 21.
- Pekmezi, D., Marcus, B., Meneses, K., Baskin, M. L., Ard, J. D., Martin, M. Y., ... Demark-wahnefried, W. (2013). Developing an intervention to address physical activity barriers for African – American women in the deep south (USA). *Women's Health*, 9(3), 301–312.
- Physical Activity Guidelines Advisory Committee (PAGAC). (2008). *Physical Activity Guidelines Advisory Committee Report 2008*. Washington, DC: US Department of Health and Human Services.
- Poh, B. K., Safiah, M. Y., Tahir, A., Siti Haslinda, M. S., Siti Norazlin, N., Norimah, A. K., ... Fatimah, S. (2010). Physical activity pattern and energy expenditure of Malaysian adults: Findings from the Malaysian Adult Nutrition Survey (MANS). *Malaysian Journal of Nutrition*, 16(1), 13–37.
- Pucher, J., & Buehler, R. (2008). Making cycling irresistible: Lessons from the Netherlands, Denmark and Germany. *Transport Reviews*, 28(4), 495–528.
- Rhodes, R. E., & Nigg, C. R. (2011). Advancing physical activity theory: A review and future directions. *Exercise and Sport Sciences Reviews*, 39(3), 113–119.
- Rose, S., Spinks, N., & Canhoto, A. I. (2014). *Tests for the assumption that a variable is normally distributed. Management research: Applying the principles*. New York: Routledge.
- Rovniak, L. S., Anderson, E. S., & Winett, R. A. (2002). Social cognitive determinants of physical activity in young adults : A prospective structural equation analysis. *Annals of Behavioural Medicine*, 24(2), 149–156.
- Rubio, D. M., Berg-Weger, M., Tebb, S. S., Lee, E. S., & Rauch, S. (2003). Objectifying content validity: Conducting a content validity study in social work research. *Social Work Research*, 37(2), 94–104.
- Sallis, J. F., Cervero, R. B., Ascher, W., Henderson, K. A., Kraft, M. K., & Kerr, J. (2006). An ecological approach to creating active living communities. *Annual Review of Public Health*, 27(1), 297–322.
- Sallis, J. F., Grossman, R. M., Pinski, R. B., Patterson, T. L., & Nader, P. R. (1987). The development of scales to measure social support for diet and exercise behaviors. *Preventive Medicine*, 16(6), 825–836.

- Shores, K. A., & West, S. T. (2010). Pursuing leisure during leisure-time physical activity. *Journal of Physical Activity and Health*, 7(5), 685–694.
- Singh, A., & Purohit, B. (2011). Evaluation of Global Physical Activity Questionnaire (GPAQ) among healthy and obese health professionals in Central India. *Baltic Journal of Health and Physical Activity*, 3(1), 34–43.
- Soo, K. L., Manan, W. A., & Suriati, W. (2015). The Bahasa Melayu version of the Global Physical Activity Questionnaire: Reliability and validity study in Malaysia. *Asia Pacific Journal of Public Health*, 27(2), NP184–NP193.
- Stacey, F. G., James, E. L., Chapman, K., & Lubans, D. R. (2016). Social cognitive theory mediators of physical activity in a lifestyle program for cancer survivors and carers: Findings from the ENRICH randomized controlled trial. *International Journal of Behavioral Nutrition and Physical Activity*, 13(1), 49.
- Stensvold, D., Nauman, J., Nilsen, T. I. L., Wisloff, U., Slordahl, S. A., & Vatten, L. (2011). Even low level of physical activity is associated with reduced mortality among people with metabolic syndrome, a population based study (the HUNT 2 study, Norway). *BMC Medicine*, 9(1), 109.
- Tam, C. L., Bonn, G., Yeoh, S. H., & Wong, C. P. (2014). Investigating diet and physical activity in malaysia: Education and family history of diabetes relate to lower levels of physical activity. *Frontiers in Psychology*, 5(1328), 1–9.
- Tavakol, M., & Dennick, R. (2011). Making sense of Cronbach's alpha. *International Journal of Medical Education*, 2, 53–55.
- Taylor, D. (2014). Physical activity is medicine for older adults. *Postgraduate Medical Journal*, 90(1059), 26–32.
- Teh, C. H., Lim, K. K., Chan, Y. Y., Lim, K. H., Azahadi, O., Hamizatul Akmar, A. H., ... Fadhl, Y. (2014). The prevalence of physical activity and its associated factors among Malaysian adults: Findings from the National Health and Morbidity Survey 2011. *Public Health*, 128(5), 416–423.
- Wen, C. P., Wai, J. P. M., Tsai, M. K., Yang, Y. C., Cheng, T. Y. D., Lee, M. C., ... Wu, X. (2011). Minimum amount of physical activity for reduced mortality and extended life expectancy: A prospective cohort study. *The Lancet*, 378(9798), 1244–1253.
- White, S. M., Wojcicki, T. R., & McAuley, E. (2012). Social cognitive influences on physical activity behavior in middle-aged and older adults. *The Journals of Gerontology, Series B: Psychological Sciences and Social Sciences*, 67(1), 18–26.
- Wojcicki, T. R., White, S. M., & McAuley, E. (2009). Assessing outcome expectations in older adults: The multidimensional outcome expectations for exercise scale. *Journals of Gerontology - Series B Psychological Sciences and Social Sciences*, 64(1), 33–40.

- World Health Organisation. (2006). *The world health report 2006: Health workers*. Geneva: World Health Organisation.
- World Health Organisation. (2009). *Global health risks, mortality and burden of disease attributable to selected major risks*. Geneva: World Health Organisation.
- World Health Organisation. (2010). *Global recommendations on physical activity for health*. Geneva: World Health Organisation.
- World Health Organisation. (2011). *Global status report on noncommunicable diseases 2010*. Geneva: World Health Organisation.
- World Health Organisation. (2012). *Global Physical Activity Questionnaire (GPAQ) analysis guide*. Geneva: World Health Organisation.
- World Health Organisation. (2013). *Global action plan for the prevention and control of noncommunicable diseases 2013-2020*. Geneva: World Health Organisation.
- World Health Organisation. (2014a). *Global status report on noncommunicable diseases 2014*. Geneva: World Health Organisation.
- World Health Organisation. (2014b). *Noncommunicable diseases country profiles 2014*. Geneva: World Health Organisation.
- World Health Organisation. (2017). Walk the talk: WHO healthy work place initiative. Retrieved from <http://www.who.int/dietphysicalactivity/walk-the-talk-who-healthy-work-place-initiative/en/>
- Young, M. D., Plotnikoff, R. C., Collins, C. E., Callister, R., & Morgan, P. J. (2014). Social cognitive theory and physical activity: A systematic review and meta-analysis. *Obesity Reviews*, 15(12), 983–995.