



**UNIVERSITI PUTRA MALAYSIA**

***EFFECTS OF THERMAL PROCESSING AND THERMOSONICATION ON  
QUALITY OF HONEY FROM STINGLESS BEES***

**CHONG KAR YEEN**

**FK 2017 28**



**EFFECTS OF THERMAL PROCESSING AND THERMOSONICATION ON  
QUALITY OF HONEY FROM STINGLESS BEES**

By

**CHONG KAR YEEN**

**Thesis Submitted to the School of Graduate Studies,  
Universiti Putra Malaysia, in Fulfilment of the  
Requirements for the Degree of Master of Science**

**April 2017**

## **COPYRIGHT**

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment  
of the requirement for the degree of Master of Science

## EFFECTS OF THERMAL PROCESSING AND THERMOSONICATION ON QUALITY OF HONEY FROM STINGLESS BEES

By

CHONG KAR YEEN

April 2017

**Chairperson: Professor Ir. Chin Nyuk Ling, PhD**  
**Faculty: Engineering**

One of the distinctive characteristics of stingless bee honey is its higher moisture content than honeybee honey. Honey being a supersaturated sugar solution tends to crystallize and ferment easily. The objective of this research was to study the effects of thermal processing and thermosonication on the quality of a stingless bee honey in Malaysia, the *Kelulut*. A two-factor-five-level design was adopted and the factors were processing temperature ranging from 45 to 90 °C and processing time ranging from 30 to 120 minutes for both methods. Physicochemical properties including water activity, moisture content, colour intensity, viscosity, hydroxymethylfurfural (HMF) content, total phenolic content (TPC), and radical scavenging activity (RSA) were determined. Thermosonicated honey had its water activity and moisture content reduced by 7.9% and 16.6%, respectively compared to 3.5% and 6.9% by thermal processing. Thermosonicated honey had its colour intensity increased by 68.2 %, viscosity increased by 275.0%, TPC increased by 58.1%, and RSA increased by 63.0% when compared to its raw form. The increase in HMF to 62.46 mg/kg using thermosonication was within the limits of international standards. The second objective of this study was to optimise thermal processing and thermosonication conditions using response surface methodology (RSM) based on minimum water activity, moisture content, and HMF content while maximizing colour intensity, viscosity, TPC, and RSA. The optimum conditions for thermal processing were at 90 °C for 108 minutes while for thermosonication it was at 90 °C for 111 minutes. To examine potential anti-inflammatory effects of the honey samples, the ability of the optimized *Kelulut* honey to inhibit nitric oxide production in lipopolysaccharide (LPS)-stimulated murine macrophages, RAW 264.7 cells, was evaluated. Results showed that *Kelulut* honey was able to both inhibit and stimulate nitric oxide using honey concentrations of 10, 20, and 50 µg/mL. Additionally, *Kelulut* honey promotes cell growth of RAW 264.7 cells. Thermosonication was revealed to be an effective honey processing alternative to current practices and the use of *Kelulut* honey as a functional food is proposed.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

## **KESAN PEMPROSESAN TERMA DAN TERMOSONIKASI KE ATAS KUALITI MADU KELULUT**

Oleh

**CHONG KAR YEEN**

**April 2017**

**Pengerusi: Profesor Ir. Chin Nyuk Ling, PhD**  
**Fakulti: Kejuruteraan**

Salah satu ciri-ciri unik madu Kelulut adalah bahawa ianya mempunyai kandungan air yang lebih tinggi daripada madu lebah. Madu adalah sejenis larutan gula tepu yang mudah berubah menjadi hablur dan menjalani proses fermentasi. Objektif penyelidikan ini adalah untuk mengkaji kesan pemprosesan terma dan termosonikasi ke atas kualiti madu Kelulut. Eksperimen yang direka untuk kajian ini menggunakan dua faktor dan lima paras. Faktor-faktornya adalah suhu pemprosesan dari 45 ke 90 °C dan jangkamasa pemprosesan dari 30 ke 120 minit untuk kedua-dua jenis pemprosesan yang dinyatakan. Sifat-sifat fizikokimia seperti aktiviti air, kandungan air, intensiti warna, kelikatan, kandungan hydroxymethylfurfural (HMF), kandungan jumlah fenolik, dan aktiviti pemerangkapan radikal dikaji. Termosonikasi mampu mengurangkan aktiviti air dan kandungan air masing-masing sebanyak 7.9% dan 16.6% berbanding dengan 3.5% and 6.9% apabila pemprosesan terma digunakan. Termosonikasi juga meningkatkan intensiti warna sebanyak 68.2%, meningkatkan kelikatan sebanyak 275.0%, meningkatkan kandungan jumlah fenolik sebanyak 58.1%, dan meningkatkan aktiviti pemerangkapan radikal sebanyak 63.0% apabila dibandingkan dengan madu Kelulut yang tidak diproses. Peningkatan kandungan HMF kepada 62.46 mg/kg apabila termosonikasi digunakan masih berada di bawah had maksimum yang ditetapkan oleh piawaian antarabangsa. Objektif kedua kajian ini adalah untuk mengoptimakan suhu dan jangkamasa pemprosesan terma dan termosonikasi berdasarkan paras minimum aktiviti air, kandungan air, dan kandungan HMF manakala intensiti warna, kelikatan, kandungan jumlah fenolik, dan aktiviti pemerangkapan radikal dimaksimakan. Keadaan optima untuk pemprosesan terma adalah dengan menggunakan suhu 90 °C selama 108 minit manakala untuk termosonikasi, suhu 90 °C selama 111 minit disyorkan. Untuk menguji potensi sifat anti-keradangan dalam sampel madu Kelulut, keupayaan madu Kelulut yang telah dioptimakan untuk menyekat pengeluaran oksida nitrik dalam sel makrofaj murin, sel RAW 264.7 telah dikaji. Hasil kajian menunjukkan bahawa madu Kelulut mempunyai keupayaan untuk menyekat dan merangsang pengeluaran oksida nitrik apabila kepekatan madu sebanyak 10, 20, dan 50 µg/mL digunakan. Tambahan pula, madu

Kelulut menggalakkan pertumbuhan sel RAW 264.7. Termosonikasi terbukti sebagai satu kaedah pemprosesan madu alternatif kepada pemprosesan semasa manakala penggunaan madu Kelulut sebagai makanan fungsional disarankan.



## ACKNOWLEDGEMENTS

I would like to express my utmost gratitude and appreciation to my main supervisor, Professor Ir. Dr. Chin Nyuk Ling for her advice, guidance, motivation, and patience in assisting me throughout this research journey. I would also like to thank Associate Professor Dr. Yus Aniza binti Yusof for her moral support during my studies. Many thanks to Associate Professor Datin Dr. Hj. Sharida Fakurazi for her guidance in the field of immunology.

My sincere appreciation is also dedicated to the technicians and science officers, specifically Mr. Raman Morat, Madam Siti, and Mr. Zahir of the Process and Food Engineering Department laboratory, Madam Abby, Madam Mazni, and Madam Norhafiza of Laboratory for Vaccines & Immunotherapeutics, Institute of Bioscience, and Madam Asmawati of the Food Science and Technology Faculty laboratory. I would like to thank them for their guidance and assistance during my experiments. Without their permission to work in the laboratories during weekends or public holidays, it would be difficult to finish the experiments on time, particularly with cell culture experiments. Special thanks to Dr. Arulselvan Palanisamy for guidance in cell culture basics.

I would like to thank my comrade, Iswaibah Mustafa, who embarked on cell line research together. Thank you to Siok Peng, Poorani, Khaleel, Katya, Woan Sean, Jia Ning, Gothai, and Dr. Chee Wun for providing me with important tips, cells, and some urgent chemicals.

Last but not least, thank you to my family and friends for their support and understanding throughout my study.

## **REPLACE!**

I certify that a Thesis Examination Committee has met on 17<sup>th</sup> April 2017 to conduct the final examination of Chong Kar Yeen on her thesis entitled “Effects of Thermal Processing and Thermosonication on Quality of Honey from Stingless Bees” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Degree of Master of Science.

Members of the Thesis Examination Committee were as follows:

**Farah Saleena Taip, PhD**

Associate Professor

Faculty of Engineering

Universiti Putra Malaysia

(Chairman)

**Norashikin Abdul Aziz, PhD**

Associate Professor

Faculty of Engineering

Universiti Putra Malaysia

(Internal Examiner)

**Ida Idayu Muhammad, PhD**

Professor

Department of Bioprocess & Polymer Engineering

Faculty of Chemical & Energy Engineering

Universiti Teknologi Malaysia

(External Examiner)

---

**Nor Aini Ab. Shukor, PhD**

Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Ir. Chin Nyuk Ling, PhD**

Professor

Faculty of Engineering

Universiti Putra Malaysia

(Chairperson)

**Yus Aniza binti Yusof, PhD**

Associate Professor

Faculty of Engineering

Universiti Putra Malaysia

(Member)

**Sharida binti Fakurazi, PhD**

Associate Professor

Faculty of Medicine and Health Sciences

Universiti Putra Malaysia

(Member)

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

### **Declaration by Members of Supervisory Committee**

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_  
Name of  
Chairman of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

## TABLE OF CONTENTS

	Page
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	ii
<b>ACKNOWLEDGEMENTS</b>	iv
<b>APPROVAL</b>	v
<b>DECLARATION</b>	vii
<b>LIST OF TABLES</b>	xi
<b>LIST OF FIGURES</b>	xii
<b>LIST OF ABBREVIATIONS</b>	xiv
 <b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
1.1 Significance of Honey	1
1.2 Production of Stingless Bee Honey and Common Problems	2
1.3 Research Objectives	3
1.4 Thesis Scope and Organization	3
 <b>2 LITERATURE REVIEW</b>	<b>5</b>
2.1 Stingless Bee Honey	5
2.1.1 Honey Composition	5
2.1.2 Physical Properties of Honey	6
2.1.3 Chemical Properties of Honey	9
2.1.4 Common Problems Associated with Honey	11
2.1.5 Medicinal Benefits of Stingless Bee Honey and their Anti-inflammatory Activities	12
2.2 Processing Methods of Honey and their Effects	13
2.2.1 Thermal Processing	14
2.2.2 Thermosonication	15
2.3 Optimization using Response Surface Methodology	17
2.4 Summary	17
 <b>3 METHODOLOGY</b>	<b>18</b>
3.1 Honey Sample Preparation	18
3.2 Processing Methods of Honey	19
3.2.1 Thermal Processing of Honey	19
3.2.2 Thermosonication of Honey	20
3.3 Analysis of Physicochemical Properties	21
3.3.1 Physical Properties	21
3.3.2 Chemical Properties	24
3.3.3 Statistical Analysis	26
3.4 Experimental Design for Optimization of Honey Quality	26
3.5 Screening Anti-inflammatory Properties	28
3.5.1 Cell Line Information and its Medium	28
3.5.2 Basic Cell Culture	28
3.5.3 MTT Cell Viability Assay	30
3.5.4 Nitric Oxide Inhibition Assay	30
3.6 Summary	31

<b>4</b>	<b>STINGLESS BEE HONEY PROCESSING</b>	<b>32</b>
4.1	Physical Properties of Honey	32
4.1.1	Water Activity and Moisture Content	32
4.1.2	Colour Intensity	34
4.1.3	Flow Behaviour	35
4.2	Chemical Properties of Honey	36
4.2.1	Hydroxymethylfurfural Content	36
4.2.2	Total Phenolic Content	37
4.2.3	Radical Scavenging Activity	39
4.3	Summary	41
<b>5</b>	<b>OPTIMIZATION OF HONEY QUALITY</b>	<b>42</b>
5.1	Statistical Model for Honey Quality	42
5.2	Optimization of Physicochemical Properties	47
5.3	Summary	49
<b>6</b>	<b>ANTI-INFLAMMATORY PROPERTIES OF PROCESSED HONEY</b>	<b>50</b>
6.1	MTT Cell Viability Assay	50
6.2	Nitric Oxide Inhibition Assay	51
6.3	Summary	53
<b>7</b>	<b>SUMMARY, CONCLUSION, AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	<b>55</b>
7.1	Summary and Conclusion	55
7.2	Recommendations for Future Work	55
	<b>REFERENCES</b>	<b>56</b>
	<b>APPENDICES</b>	<b>68</b>
	<b>BIODATA OF STUDENT</b>	<b>73</b>

## LIST OF TABLES

Table		Page
2.1	Proximate composition and predominant sugars in raw Kelulut honey, processed Manuka honey, and commercial honey [adapted from Kek et al. (2016)]	5
2.2	Microorganisms growing in $a_w$ range of intermediate-moisture food (Beuchat, 1981)	7
2.3	Medicinal uses of stingless bee honey [adapted from Vit et al. (2004)]	12
3.1	Face-centred central composite design matrix used to evaluate effects of processing variables on honey quality	27
5.1	Regression coefficients, $R^2$ , adjusted $R^2$ , and P values for honey quality dependent variables for thermal processing of honey	43
5.2	Regression coefficients, $R^2$ , adjusted $R^2$ , and P values for honey quality dependent variables for honey thermosonication	43
5.3	Process variables, responses optimisation, desirability, and experimental value for the responses at optimum condition for thermally processed honey	48
5.4	Process variables, responses optimisation, desirability, and experimental value for the responses at optimum condition for thermosonicated honey	48
7.1	Tukey's test for multiple comparisons of means of responses of thermally processed honey	68
7.2	Tukey's test for multiple comparisons of means of responses of thermosonicated honey	79
7.3	P-values and F values for thermally processed honey and thermosonicated honey	70
7.4	Central composite design of independent variables and responses	71
7.5	RAW 264.7 cell viability studies	72
7.6	Nitric oxide concentrations ( $\mu\text{M}$ ) after treatment with various honey concentrations	72

## LIST OF FIGURES

Figure		Page
1.1	Honey pots made by stingless bees	2
2.1	Apparent viscosities of time-independent fluids [adapted from Sahin and Sumnu (2006)]	8
2.2	Time-dependent behaviour of fluids [adapted from Sahin and Sumnu (2006)]	9
3.1	Work sequence of this research	18
3.2	Fresh <i>Kelulut</i> honey in bottles	19
3.3	Thermostatic water bath (WNB 22, Memmert GmbH + Co. KG, Germany)	19
3.4	Ultrasonic bath tank (a) without cover and (b) with cover and suspended test tube rack	20
3.5	Test tube rack with honey samples suspended	20
3.6	Top view schematic diagram of the experimental set-up for thermosonication treatment: (1) side ultrasonic generator, (2) bottom ultrasonic generator, (3) steel rack containing test tubes, (4) heaters, and (5) insulation for ultrasonic tank	21
3.7	Water activity meter (Decagon Devices, Pullman, Washington, USA)	22
3.8	Digital refractometer (Digital ABBE Refractometer AR2008, A. Krüss, Germany)	23
3.9	UV/Visible spectrophotometer (Ultrospec 3100 pro, Amersham Biosciences, USA)	23
3.10	Rheometer (AR-G2, TA Instruments, New Castle, USA)	24
3.11	Benchtop centrifuge (Universal 320, Hettich, USA)	25
3.12	Mixture of honey and methanol	26
3.13	Table top centrifuge (Universal 32R, Hettich, USA)	29
4.1	Effect of thermal processing (a, c) and thermosonication (b, d) on water activity and moisture content	33
4.2	Effect of temperature and time on colour intensity using (a) thermal processing and (b) thermosonication	34
4.3	Effect of temperature and time on viscosity using (a) thermal processing and (b) thermosonication	35
4.4	Effect of temperature and time on hydroxymethylfurfural using (a) thermal processing and (b) thermosonication	37
4.5	Effect of temperature and time on total phenolic content using (a) thermal processing and (b) thermosonication	38
4.6	Effect of temperature and time on DPPH inhibition using (a) thermal processing and (b) thermosonication	40
5.1	Effects of temperature and time on (a) water activity, (b) moisture content, (c) colour intensity, (d) viscosity, (e) hydroxymethylfurfural content, (f) total phenolic content, and (g) DPPH radical scavenging activity for thermally processed honey	45

5.2	Effects of temperature and time on (a) water activity, (b) moisture content, (c) colour intensity, (d) viscosity, (e) hydroxymethylfurfural content, (f) total phenolic content, and (g) DPPH radical scavenging activity for thermosonicated honey	46
6.1	Effects of (a) raw honey, (b) thermally processed honey, and (c) thermosonicated honey on RAW 264.7 cell viability	51
6.2	Effects of (a) raw honey, (b) thermally processed honey, and (c) thermosonicated honey on nitric oxide production by LPS-stimulated RAW 264.7 cells	52



## LIST OF ABBREVIATIONS

ANOVA	Analysis of variance
DMSO	Dimethyl sulfoxide
DPPH	2,2-Diphenyl-1-picrylhydrazyl
HMF	Hydroxymethylfurfural
LPS	Lipopolysaccharide
MRP	Maillard reaction product
MTT	Methylthiazolyldiphenyl-tetrazolium bromide
NO	Nitric oxide
PBS	Phosphate buffered saline
RSA	Radical scavenging activity
RSM	Response surface methodology
TPC	Total phenolic content



## CHAPTER 1

### INTRODUCTION

#### 1.1 Significance of Honey

Honey is used for many purposes. It can be consumed as it is, be used as spread, or directly added to sweeten drinks. In bakery products such as *baklava*, a Turkish flaky pastry, gingerbreads, and fruit cakes, honey is used as it improves moisture retention and thus increases shelf life (Tong *et al.*, 2010). Apart from giving cakes a richer flavour, honey eliminates dryness and crumbliness in them. Honey is also used in confectionery such as caramel and nougat. One of the pull factors is the nutritional benefits of honey, as it is often used for medicinal purposes. Some of the medicinal uses of honey include treatment of digestive disorders, respiratory infections, and promotion of wound healing (Vit, Medina, & Eunice Enríquez, 2004). Home remedies also use honey to soothe sore throats and coughs. Honey is also integrated into cosmetics and skincare products as it has antioxidant and hygroscopic properties, thus providing a soothing effect on the skin.

In the wild, honeybees usually nest in hollow trees or rock crevices rather high above the ground. Over the years, honey hunting shifted to beekeeping in manmade hives where they can be conveniently accessible and safe from predators. This transition was important to increase honey production as its demand increased. From immovable combs, honey production advanced to movable-frame hive, and subsequently embossed beeswax foundation was invented to fit into the frames (Crane, 1980). A centrifugal extractor would spin the honey out and the frames are ready to be reused. This method is still presently used by small-scaled and artisanal honey producers.

Stingless bees which are much smaller in size, were prized for its cerumen and honey in native civilizations. The evolution from honey hunting to beekeeping is similar for stingless bees. From wild bees in hollow trees, beekeeping evolved to usage of a hollow log. The hollow log has a central flight entrance and closure and it was cut in such a way that it can be opened and resealed by the owner (Jones, 2013). At present, log hives are used along with boards.



**Figure 1.1: Honey pots made by stingless bees**

The tropical climate in Malaysia enables various plants to flourish which in turn becomes a suitable environment for foraging bees. Honey can be broadly categorized by its botanical origin and type of bees. Common honey found in Malaysia are the Acacia, Pineapple, *Tualang*, Borneo, *Gelam*, and *Kelulut* honey (Chua, Abdul-Rahaman, Sarmidi, & Aziz, 2012; Kek, Chin, Tan, Yusof, & Chua, 2016; Moniruzzaman, Khalil, Sulaiman, & Gan, 2013). The *Kelulut* honey is sourced from *Trigona* spp. stingless bees. It is a multifloral honey and some of the unique characteristics of stingless bee honey or 'pot honey' are its higher moisture content, higher electrical conductivity, higher acidity, and lower diastase activity (Chuttong, Chanbang, Sringarm, & Burgett, 2016).

## **1.2 Production of Stingless Bee Honey and Common Problems**

Similar to other bees, stingless bees collect nectar and pollen from flowers and carry them to the nest where the larvae are fed. The food goes into cerumen pots. At the hive, the bees ripen or dehydrate the nectar droplets by spinning them inside their mouthparts until honey is formed. Honey harvest was done by piercing or squeezing the pots of honey and pollen. Modern technology then introduced the use of a suction device to improve product quality. However, low productivity of stingless bee colonies is one of the factors deterring beekeepers and consumers (de Oliveira Alves, 2013).

Both stingless bee honey and honeybee honey have natural tendency to undergo crystallization and fermentation. Honey is a highly-saturated sugar solution and crystallization will occur spontaneously. When crystals are formed, two distinct phases which are crystallized phase at the bottom and liquid phase at the top can be seen. This undesirable appearance will deter consumers from purchasing the honey. Besides that, an increase in water activity becomes a favourable environment for naturally-present osmophilic yeasts in honey to multiply. As stingless bee honey itself has higher moisture content, fermentation degrades honey quality by altering its taste (Gleiter, Horn, & Isengard, 2006).

Current honey production practices by the beekeepers include refrigeration, dehydration, pasteurization, and maturation. Refrigeration involves keeping honey at approximately 4-8 °C just after harvesting and until consumption. Dehydration involves ventilating the honey in a dry room with a dehumidifier. Pasteurization is also performed at 72 °C for 15 seconds to eliminate pathogens. Another method is to develop maturity of the honey. Maturation is done by keeping honey inside closed bottles, and opened once a week to release gases produced by fermentation, and closed again. This process continues until no more gas is released. Then, the stabilized honey can be stored (Menezes, Vollet-Neto, Contrera, Venturieri, & Imperatriz-Fonseca, 2013).

Honey is well-known for its many benefits such as high antioxidants content, antimicrobial activity, wound-healing, and anti-inflammatory activities (Liu, Ye, Lin, Wang, & Peng, 2013). Further explanation can be found in Section 2.1.5. Hence, honeybee honey is often tested for its anti-inflammatory activities (Van den Berg et al., 2008). There are studies which used Malaysian honey such as Gelam honey to conduct *in vitro* and *in vivo* anti-inflammatory tests (Kassim, Achoui, Mansor, & Yusoff, 2010a; Kassim, Achoui, Mustafa, Mohd, & Yusoff, 2010b). However, there are insufficient studies on Kelulut honey and it is important to investigate the therapeutic properties to increase its consumption.

### **1.3 Research Objectives**

The objectives of this research were:

- i. to study the effects of temperature and time on the quality of *Kelulut* honey using thermal processing and thermosonication,
- ii. to optimize thermal processing and thermosonication by using response surface methodology, and
- iii. to evaluate the anti-inflammatory activity of selected treated honey by investigating inhibition of nitric oxide production in RAW 264.7 cells.

### **1.4 Thesis Scope and Organization**

Although there are many by-products of honey production such as propolis, bee pollen, royal jelly, beebread, and cerumen, this study focused only on honey. The processing methods used were conventional thermal processing and thermosonication. Honey quality was evaluated by its physicochemical properties covered by this research which were water activity, moisture content, colour intensity, viscosity, hydroxymethylfurfural content, total phenolic content, and radical scavenging activity. One of the ways to screen anti-inflammatory activities is by studying the inhibition of nitric oxide production in lipopolysaccharide-stimulated cells. For this purpose, mouse leukemic macrophages, also known as RAW 264.7 cells, were used.

The thesis comprises of seven main chapters. Chapter One gives an overview of the research, a brief introduction of objectives, and the problems encountered which led to initiation of this research. Chapter Two reviews literature of honey composition,

physicochemical properties of honey, anti-inflammatory activities of honey, detailed explanation on honey crystallization and fermentation, and effects of thermal processing and thermosonication on honey. Chapter Three focusses on the methodology including the experimental design and materials used for the entire research. Chapter Four presents the results and the effects of thermal processing and thermosonication on the quality of *Kelulut* honey. Chapter Five discusses optimization of honey quality. By using face-centred central composite design in the Minitab Statistical Software 16, the surface regression analysis, modelling response, surface plots, optimisation, and desirability of each response were obtained and discussed. Chapter Six concentrates on the investigation of anti-inflammatory properties of processed honey. It comprises of a cell viability assay and a nitric oxide assay. Chapter Seven summarizes the results and concludes the entire research.



## REFERENCES

- Abid, M., Jabbar, S., Wu, T., Hashim, M. M., Hu, B., Lei, S., ... Zeng, X. (2013). Effect of ultrasound on different quality parameters of apple juice. *Ultrasonics Sonochemistry*, 20(5), 1182–1187. <https://doi.org/10.1016/j.ultsonch.2013.02.010>
- Abramovič, H., Jamnik, M., Burkan, L., & Kač, M. (2008). Water activity and water content in Slovenian honeys. *Food Control*, 19(11), 1086–1090. <https://doi.org/10.1016/j.foodcont.2007.11.008>
- Abu-Jdayil, B., Al-Majeed Ghzawi, A., Al-Malah, K. I. M., & Zaitoun, S. (2002). Heat effect on rheology of light- and dark-colored honey. *Journal of Food Engineering*, 51(1), 33–38. [https://doi.org/10.1016/S0260-8774\(01\)00034-6](https://doi.org/10.1016/S0260-8774(01)00034-6)
- Ahmed, J., Prabhu, S. T., Raghavan, G. S. V., & Ngadi, M. (2007). Physico-chemical, rheological, calorimetric and dielectric behavior of selected Indian honey. *Journal of Food Engineering*, 79(4), 1207–1213. <https://doi.org/http://dx.doi.org/10.1016/j.jfoodeng.2006.04.048>
- Ahmed, J., Ramaswamy, H. S., Kasapis, S., & Boye, J. I. (2009). *Novel food processing: effects on rheological and functional properties*. CRC Press.
- Akhmazillah, M. F. N., Farid, M. M., & Silva, F. V. M. (2013). High pressure processing (HPP) of honey for the improvement of nutritional value. *Innovative Food Science & Emerging Technologies*, 20, 59–63. <https://doi.org/10.1016/j.ifset.2013.06.012>
- Al-Jadi, A.-M., Enchang, F. K., & Yusoff, K. M. (2014). The effect of Malaysian honey and its major components on the proliferation of cultured fibroblasts. *Turkish Journal of Medical Sciences*, 44(5), 733–740.
- Al-Waili, N. S. (2003). Identification of nitric oxide metabolites in various honeys: effects of intravenous honey on plasma and urinary nitric oxide metabolites concentrations. *Journal of Medicinal Food*, 6(4), 359–364. <https://doi.org/10.1089/109662003772519921>
- Al-Waili, N. S. (2005). Effects of Honey on the Urinary Total Nitrite and Prostaglandins Concentration. *International Urology and Nephrology*, 37(1), 107–111. <https://doi.org/10.1007/s11255-004-0871-8>
- Al-Waili, N. S., & Boni, N. S. (2004). Honey increased saliva, plasma, and urine content of total nitrite concentrations in normal individuals. *Journal of Medicinal Food*, 7(3), 377–380. <https://doi.org/10.1089/jmf.2004.7.377>
- Alimentarius, C. (2001). Codex standard for honey. *FAO, Rome*, 19–26.
- Aljadi, A. M., & Kamaruddin, M. Y. (2004). Evaluation of the phenolic contents and antioxidant capacities of two Malaysian floral honeys. *Food Chemistry*, 85(4), 513–518. [https://doi.org/10.1016/S0308-8146\(02\)00596-4](https://doi.org/10.1016/S0308-8146(02)00596-4)
- Almeida-Muradian, L. B. de, Matsuda, A. H., & Bastos, D. H. M. (2007). Physicochemical parameters of amazon Melipona honey. *Química Nova*, 30(3), 707–708.



- Alvarez-Suarez, J. M., Tulipani, S., Romandini, S., Bertoli, E., & Battino, M. (2010). Contribution of honey in nutrition and human health: a review. *Mediterranean Journal of Nutrition and Metabolism*, 3(1), 15–23. <https://doi.org/10.1007/s12349-009-0051-6>
- Amorati, R., & Valgimigli, L. (2015). Advantages and limitations of common testing methods for antioxidants. *Free Radical Research*, 49(5), 633–649. <https://doi.org/10.3109/10715762.2014.996146>
- Arnoldi, A. (2001). Thermal processing and food quality: analysis and control. *Thermal Technologies in Food Processing*, 138–159.
- Azeredo, L. d. C., Azeredo, M. A. A., de Souza, S. R., & Dutra, V. M. L. (2003). Protein contents and physicochemical properties in honey samples of *Apis mellifera* of different floral origins. *Food Chemistry*, 80(2), 249–254. [https://doi.org/10.1016/S0308-8146\(02\)00261-3](https://doi.org/10.1016/S0308-8146(02)00261-3)
- Baltrušaitytė, V., Venskutonis, P. R., & Čeksterytė, V. (2007). Radical scavenging activity of different floral origin honey and beebread phenolic extracts. *Food Chemistry*, 101(2), 502–514. <https://doi.org/http://dx.doi.org/10.1016/j.foodchem.2006.02.007>
- Benzing-Purdie, L. M., Ripmeester, J. A., & Ratcliffe, C. I. (1985). Effects of temperature on Maillard reaction products. *Journal of Agricultural and Food Chemistry*, 33(1), 31–33. <https://doi.org/10.1021/jf00061a009>
- Beretta, G., Gelmini, F., Lodi, V., Piazzalunga, A., & Maffei Facino, R. (2010). Profile of nitric oxide (NO) metabolites (nitrate, nitrite and N-nitroso groups) in honeys of different botanical origins: Nitrate accumulation as index of origin, quality and of therapeutic opportunities. *Journal of Pharmaceutical and Biomedical Analysis*, 53(3), 343–349. <https://doi.org/http://dx.doi.org/10.1016/j.jpba.2010.04.010>
- Beretta, G., Granata, P., Ferrero, M., Orioli, M., & Maffei Facino, R. (2005). Standardization of antioxidant properties of honey by a combination of spectrophotometric/fluorimetric assays and chemometrics. *Analytica Chimica Acta*, 533(2), 185–191. <https://doi.org/10.1016/j.aca.2004.11.010>
- Bermúdez-Aguirre, D., Mobbs, T., & Barbosa-Cánovas, G. (2011). Ultrasound Applications in Food Processing. In H. Feng, G. Barbosa-Canovas, & J. Weiss (Eds.), *Ultrasound Technologies for Food and Bioprocessing* (pp. 65–105). Springer New York. [https://doi.org/10.1007/978-1-4419-7472-3\\_3](https://doi.org/10.1007/978-1-4419-7472-3_3)
- Beuchat, L. R. (1981). Microbial stability as affected by water activity. *Cereal Foods World*, 26(7), 345–349.
- Biluca, F. C., Della Betta, F., de Oliveira, G. P., Pereira, L. M., Gonzaga, L. V., Costa, A. C. O., & Fett, R. (2014). 5-HMF and carbohydrates content in stingless bee honey by CE before and after thermal treatment. *Food Chemistry*, 159, 244–249. <https://doi.org/10.1016/j.foodchem.2014.03.016>
- Blois, M. S. (1958). Antioxidant Determinations by the Use of a Stable Free Radical. *Nature*, 181(4617), 1199–1200. Retrieved from <http://dx.doi.org/10.1038/1811199a0>
- Bogdan, C. (2001). Nitric oxide and the immune response. *Nat Immunol*, 2(10), 907–

916. Retrieved from <http://dx.doi.org/10.1038/ni1001-907>

- Bogdanov, S., Lüllmann, C., Martin, P., von der Ohe, W., Russmann, H., Vorwohl, G., ... Vit, P. (1999). Honey quality and international regulatory standards: review by the International Honey Commission. *Bee World*, 80(2), 61–69. <https://doi.org/10.1080/0005772X.1999.11099428>
- Bogdanov, S., Martin, P., & Lullmann, C. (2002). Harmonised methods of the international honey commission. *Swiss Bee Research Centre, FAM, Liebefeld*.
- Bogdanov, S., Ruoff, K., & Persano Oddo, L. (2004). Physico-chemical methods for the characterisation of unifloral honeys: a review. *Apidologie*, 35(Suppl 1), 4–17.
- Brand-Williams, W., Cuvelier, M. E., & Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT - Food Science and Technology*, 28(1), 25–30. [https://doi.org/10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5)
- Bulut, L., & Kilic, M. (2009). Kinetics of hydroxymethylfurfural accumulation and color change in honey during storage in relation to moisture content. *Journal of Food Processing and Preservation*, 33(1), 22–32. <https://doi.org/10.1111/j.1745-4549.2008.00233.x>
- Capuano, E., & Fogliano, V. (2011). Acrylamide and 5-hydroxymethylfurfural (HMF): A review on metabolism, toxicity, occurrence in food and mitigation strategies. *LWT - Food Science and Technology*, 44(4), 793–810. <https://doi.org/10.1016/j.lwt.2010.11.002>
- Cárcel, J. A., García-Pérez, J. V., Benedito, J., & Mulet, A. (2012). Food process innovation through new technologies: Use of ultrasound. *Journal of Food Engineering*, 110(2), 200–207. <https://doi.org/10.1016/j.jfoodeng.2011.05.038>
- Cavia, M. M., Fernáez-Muiño, M. A., Huidobro, J. F., & Sancho, M. T. (2004). Correlation between Moisture and Water Activity of Honeys Harvested in Different Years. *Journal of Food Science*, 69(5), C368–C370. <https://doi.org/10.1111/j.1365-2621.2004.tb10699.x>
- Chaikhram, P., Kemsawasd, V., & Apichartsrangkoon, A. (2016). Effects of conventional and ultrasound treatments on physicochemical properties and antioxidant capacity of floral honeys from Northern Thailand. *Food Bioscience*, 15, 19–26. <https://doi.org/10.1016/j.fbio.2016.04.002>
- Chaikhram, P., & Prangthip, P. (2015). Alteration of antioxidative properties of longan flower-Honey after High pressure, ultra-sonic and thermal processing. *Food Bioscience*.
- Chemat, F., Zill-e-Huma, & Khan, M. K. (2011). Applications of ultrasound in food technology: Processing, preservation and extraction. *Ultrasonics Sonochemistry*, 18(4), 813–835. <https://doi.org/10.1016/j.ultsonch.2010.11.023>
- Chin, N. L., Tan, M. C., Che Pa, N. F., & Yusof, Y. A. (2013). A method and apparatus for high intensity ultrasonic treatment of baking materials. United States: Universiti Putra Malaysia. Retrieved from <https://www.google.com/patents/US20130189407>
- Chirife, J., Zamora, M. C., & Motto, A. (2006). The correlation between water activity and % moisture in honey: Fundamental aspects and application to Argentine

- honeys. *Journal of Food Engineering*, 72(3), 287–292.  
<https://doi.org/10.1016/j.jfoodeng.2004.12.009>
- Chua, L. S., Abdul-Rahaman, N.-L., Sarmidi, M. R., & Aziz, R. (2012). Multi-elemental composition and physical properties of honey samples from Malaysia. *Food Chemistry*, 135(3), 880–887.  
<https://doi.org/10.1016/j.foodchem.2012.05.106>
- Chua, L. S., & Adnan, N. A. (2014). Biochemical and nutritional components of selected honey samples. *Acta Sci. Pol., Technol. Aliment*, 13(2), 169–179. Retrieved from [http://www.food.actapol.net/volume13/issue2/6\\_2\\_2014.pdf](http://www.food.actapol.net/volume13/issue2/6_2_2014.pdf)
- Chua, L. S., Adnan, N. A., Abdul-Rahaman, N. L., & Sarmidi, M. R. (2014). Effect of thermal treatment on the biochemical composition of tropical honey samples. *Int. Food Res. J*, 21(2), 773–778.
- Chuttong, B., Chanbang, Y., Sringarm, K., & Burgett, M. (2016). Physicochemical profiles of stingless bee (Apidae: Meliponini) honey from South East Asia (Thailand). *Food Chemistry*, 192, 149–155.  
<https://doi.org/10.1016/j.foodchem.2015.06.089>
- Clancy, J. (1998). *Basic concepts in immunology: a student's survival guide*. McGraw Hill Professional.
- Corrales, M., Toepfl, S., Butz, P., Knorr, D., & Tauscher, B. (2008). Extraction of anthocyanins from grape by-products assisted by ultrasonics, high hydrostatic pressure or pulsed electric fields: A comparison. *Innovative Food Science & Emerging Technologies*, 9(1), 85–91. <https://doi.org/10.1016/j.ifset.2007.06.002>
- Crane, E. (1975). *Honey. A comprehensive survey*. London: William Heinemann Ltd.
- Crane, E. (1980). *A book of honey*. Oxford: Oxford University Press.
- Cruz, R. M. S., Vieira, M. C., & Silva, C. L. M. (2007). Modelling kinetics of watercress (*Nasturtium officinale*) colour changes due to heat and thermosonication treatments. *Innovative Food Science & Emerging Technologies*, 8(2), 244–252. <https://doi.org/10.1016/j.ifset.2007.01.003>
- D'Arcy, B. (2007). *High-power ultrasound to control of honey crystallisation*. Rural Industries Research and Development Corporation.
- de Oliveira Alves, R. M. (2013). Production and Marketing of Pot-Honey. In P. Vit, S. R. M. Pedro, & D. Roubik (Eds.), *Pot-Honey: A legacy of stingless bees* (pp. 541–556). New York, NY: Springer New York. [https://doi.org/10.1007/978-1-4614-4960-7\\_40](https://doi.org/10.1007/978-1-4614-4960-7_40)
- Deliza, R., & Vit, P. (2013). Sensory evaluation of stingless bee pot-honey. In *Pot-Honey* (pp. 349–361). Springer.
- Efem, S. E. E. (1988). Clinical observations on the wound healing properties of honey. *British Journal of Surgery*, 75(7), 679–681.  
<https://doi.org/10.1002/bjs.1800750718>
- Escriche, I., Visquert, M., Juan-Borrás, M., & Fito, P. (2009). Influence of simulated industrial thermal treatments on the volatile fractions of different varieties of honey. *Food Chemistry*, 112(2), 329–338.



<https://doi.org/10.1016/j.foodchem.2008.05.068>

- Fallico, B., Arena, E., & Zappala, M. (2008). Degradation of 5-hydroxymethylfurfural in honey. *J Food Sci*, 73(9), C625-31. <https://doi.org/10.1111/j.1750-3841.2008.00946.x>
- Ferreira, E. L., Lencioni, C., Benassi, M. T., Barth, M. O., & Bastos, D. H. M. (2009). Descriptive sensory analysis and acceptance of stingless bee honey. *Food Science and Technology International*, 15(3), 251–258.
- Fontana Jr, A. J., Schmidt, S. J., & Labuza, T. P. (2008). *Water activity in foods: fundamentals and applications* (Vol. 13). John Wiley & Sons.
- Frankel, S., Robinson, G. E., & Berenbaum, M. R. (1998). Antioxidant capacity and correlated characteristics of 14 unifloral honeys. *Journal of Apicultural Research*, 37(1), 27–31. <https://doi.org/10.1080/00218839.1998.11100951>
- Fuenmayor, C. A., Díaz-Moreno, A. C., Zuluaga-Domínguez, C. M., & Quicazán, M. C. (2013). Honey of Colombian Stingless Bees: Nutritional Characteristics and Physicochemical Quality Indicators. In P. Vit, M. S. R. Pedro, & D. Roubik (Eds.), *Pot-Honey: A legacy of stingless bees* (pp. 383–394). New York, NY: Springer New York. [https://doi.org/10.1007/978-1-4614-4960-7\\_27](https://doi.org/10.1007/978-1-4614-4960-7_27)
- Gámbaro, A., Ares, G., Gimenez, A. N. A., & Pahor, S. (2007). Preference mapping of color of Uruguayan honeys. *Journal of Sensory Studies*, 22(5), 507–519.
- García Rodríguez, L., & Jick, H. (1994). Risk of upper gastrointestinal bleeding and perforation associated with Individual non-steroidal anti-inflammatory drugs. *The Lancet*, 343(8900), 769–772. [https://doi.org/http://dx.doi.org/10.1016/S0140-6736\(94\)91843-0](https://doi.org/http://dx.doi.org/10.1016/S0140-6736(94)91843-0)
- Garza, S., Ibarz, A., Pagán, J., & Giner, J. (1999). Non-enzymatic browning in peach puree during heating. *Food Research International*, 32(5), 335–343. [https://doi.org/http://dx.doi.org/10.1016/S0963-9969\(99\)00094-0](https://doi.org/http://dx.doi.org/10.1016/S0963-9969(99)00094-0)
- Gleiter, R. A., Horn, H., & Isengard, H. D. (2006). Influence of type and state of crystallisation on the water activity of honey. *Food Chemistry*, 96(3), 441–445. <https://doi.org/10.1016/j.foodchem.2005.03.051>
- Gonnet, M. (1977). Honey liquefaction, pasteurization and induced crystallization. *Apiacta; an International Technical Magazine of Apicultural and Economic Information*.
- Granato, D., Ribeiro, J. C. B., Castro, I. A., & Masson, M. L. (2010). Sensory evaluation and physicochemical optimisation of soy-based desserts using response surface methodology. *Food Chemistry*, 121(3), 899–906. <https://doi.org/10.1016/j.foodchem.2010.01.014>
- Guan, Y.-G., Zhang, B.-S., Yu, S.-J., Wang, X.-R., Xu, X.-B., Wang, J., ... Lin, H. (2011). Effects of ultrasound on a glycine–glucose model system—A means of promoting maillard reaction. *Food and Bioprocess Technology*, 4(8), 1391–1398.
- Guerrini, A., Bruni, R., Maietti, S., Poli, F., Rossi, D., Paganetto, G., ... Sacchetti, G. (2009). Ecuadorian stingless bee (Meliponinae) honey: A chemical and functional profile of an ancient health product. *Food Chemistry*, 114(4), 1413–1420. <https://doi.org/10.1016/j.foodchem.2008.11.023>

- Guler, A., Bakan, A., Nisbet, C., & Yavuz, O. (2007). Determination of important biochemical properties of honey to discriminate pure and adulterated honey with sucrose (*Saccharum officinarum* L.) syrup. *Food Chemistry*, 105(3), 1119–1125. <https://doi.org/10.1016/j.foodchem.2007.02.024>
- Hata, A. N., & Breyer, R. M. (2004). Pharmacology and signaling of prostaglandin receptors: Multiple roles in inflammation and immune modulation. *Pharmacology & Therapeutics*, 103(2), 147–166. <https://doi.org/10.1016/j.pharmthera.2004.06.003>
- Henson, P. M. (2005). Dampening inflammation. *Nat Immunol*, 6(12), 1179–1181. <https://doi.org/10.1038/ni1205-1179>
- Hermosín, I., Chicón, R. M., & Dolores Cabezudo, M. (2003). Free amino acid composition and botanical origin of honey. *Food Chemistry*, 83(2), 263–268. [https://doi.org/http://dx.doi.org/10.1016/S0308-8146\(03\)00089-X](https://doi.org/http://dx.doi.org/10.1016/S0308-8146(03)00089-X)
- Hoseney, R. C. (1984). Chemical changes in carbohydrates produced by thermal processing. *Journal of Chemical Education*, 61(4), 308. <https://doi.org/10.1021/ed061p308>
- Iglesias, M. T., Martín-Álvarez, P. J., Polo, M. C., de Lorenzo, C., González, M., & Pueyo, E. (2006). Changes in the Free Amino Acid Contents of Honeys During Storage at Ambient Temperature. *Journal of Agricultural and Food Chemistry*, 54(24), 9099–9104. <https://doi.org/10.1021/jf061712x>
- Jambrak, A. R. (2011). Experimental design and optimization of ultrasound treatment of food products. *Journal of Food Processing & Technology*.
- Jambrak, A. R., Mason, T. J., Paniwnyk, L., & Lelas, V. (2007). Accelerated drying of button mushrooms, Brussels sprouts and cauliflower by applying power ultrasound and its rehydration properties. *Journal of Food Engineering*, 81(1), 88–97. <https://doi.org/10.1016/j.jfoodeng.2006.10.009>
- Jones, R. (2013). Stingless Bees: A Historical Perspective. In P. Vit, S. R. M. Pedro, & D. Roubik (Eds.), *Pot-Honey: A legacy of stingless bees* (pp. 219–227). New York, NY: Springer New York. [https://doi.org/10.1007/978-1-4614-4960-7\\_14](https://doi.org/10.1007/978-1-4614-4960-7_14)
- Juszczak, L., & Fortuna, T. (2006). Rheology of selected Polish honeys. *Journal of Food Engineering*, 75(1), 43–49. <https://doi.org/10.1016/j.jfoodeng.2005.03.049>
- Kabbani, D., Sepulcre, F., & Wedekind, J. (2011). Ultrasound-assisted liquefaction of rosemary honey: influence on rheology and crystal content. *Journal of Food Engineering*, 107(2), 173–178.
- Karabournioti, S., & Zervaki, P. (2001). The effect of heating on honey HMF and invertase. *Apiacta*, 36(4), 177–181.
- Kassim, M., Achoui, M., Mansor, M., & Yusoff, K. M. (2010). The inhibitory effects of Gelam honey and its extracts on nitric oxide and prostaglandin E2 in inflammatory tissues. *Fitoterapia*, 81(8), 1196–1201. <https://doi.org/10.1016/j.fitote.2010.07.024>
- Kassim, M., Achoui, M., Mustafa, M. R., Mohd, M. A., & Yusoff, K. M. (2010). Ellagic acid, phenolic acids, and flavonoids in Malaysian honey extracts demonstrate in vitro anti-inflammatory activity. *Nutrition Research*, 30(9), 650–

659. <https://doi.org/10.1016/j.nutres.2010.08.008>

- Kek, S. P., Chin, N. L., Tan, S. W., Yusof, Y. A., & Chua, L. S. (2016). Classification of Honey from Its Bee Origin via Chemical Profiles and Mineral Content. *Food Analytical Methods*, 1–12. <https://doi.org/10.1007/s12161-016-0544-0>
- Kek, S. P., Chin, N. L., Yusof, Y. A., Tan, S. W., & Chua, L. S. (2014). Total Phenolic Contents and Colour Intensity of Malaysian Honeys from the *Apis* spp. and *Trigona* spp. Bees. *Agriculture and Agricultural Science Procedia*, 2, 150–155.
- Khalil, M. I., Alam, N., Moniruzzaman, M., Sulaiman, S. A., & Gan, S. H. (2011). Phenolic Acid Composition and Antioxidant Properties of Malaysian Honeys. *Journal of Food Science*, 76(6), C921–C928. <https://doi.org/10.1111/j.1750-3841.2011.02282.x>
- Khalil, M. I., Sulaiman, S. A., & Gan, S. H. (2010). High 5-hydroxymethylfurfural concentrations are found in Malaysian honey samples stored for more than one year. *Food and Chemical Toxicology*, 48(8–9), 2388–2392. <https://doi.org/10.1016/j.fct.2010.05.076>
- Korhonen, R., Lahti, A., Hämäläinen, M., Kankaanranta, H., & Moilanen, E. (2002). Dexamethasone Inhibits Inducible Nitric-Oxide Synthase Expression and Nitric Oxide Production by Destabilizing mRNA in Lipopolysaccharide-Treated Macrophages. *Molecular Pharmacology*, 62(3), 698 LP-704. Retrieved from <http://molpharm.aspetjournals.org/content/62/3/698.abstract>
- Kowalski, S. (2013). Changes of antioxidant activity and formation of 5-hydroxymethylfurfural in honey during thermal and microwave processing. *Food Chemistry*, 141(2), 1378–1382. <https://doi.org/10.1016/j.foodchem.2013.04.025>
- Lieu, L. N., & Le, V. V. M. (2010). Application of ultrasound in grape mash treatment in juice processing. *Ultrasonics Sonochemistry*, 17(1), 273–279. <https://doi.org/http://dx.doi.org/10.1016/j.ultsonch.2009.05.002>
- Liu, J.-R., Ye, Y.-L., Lin, T.-Y., Wang, Y.-W., & Peng, C.-C. (2013). Effect of floral sources on the antioxidant, antimicrobial, and anti-inflammatory activities of honeys in Taiwan. *Food Chemistry*, 139(1–4), 938–943. <https://doi.org/10.1016/j.foodchem.2013.02.015>
- Marcinkiewicz, E., Marcinkiewicz, J., & Chłopicki, S. (2004). Nitric oxide - a pro-inflammatory and anti-inflammatory mediator. *Central European Journal of Immunology*, 28(2), 74–78. Retrieved from <http://www.termedia.pl/Nitric-oxide-8211-a-pro-inflammatory-and-anti-inflammatory-mediator,10,1648,1,1.html>
- Marvin, G. E., Peterson, W. H., Fred, E. B., & Wilson, H. F. (1931). Some of the characteristics of yeasts found in fermenting honey. *J. Agr. Research*, 43(2), 121–131.
- Mason, T. J. (1991). Practical sonochemistry user's guide to application in chemistry and chemical engineering. UK: Ellis Horwood.
- Mason, T. J., & Lorimer, J. P. (2002). Applied sonochemistry. *The Uses of Power Ultrasound in Chemistry and Processing*, 1–48.
- Menezes, C., Vollet-Neto, A., Contrera, F. A. F. L., Venturieri, G. C., & Imperatriz-

- Fonseca, V. L. (2013). The Role of Useful Microorganisms to Stingless Bees and Stingless Beekeeping. In P. Vit, S. R. M. Pedro, & D. Roubik (Eds.), *Pot-Honey: A legacy of stingless bees* (pp. 153–171). New York, NY: Springer New York. [https://doi.org/10.1007/978-1-4614-4960-7\\_10](https://doi.org/10.1007/978-1-4614-4960-7_10)
- Milani, E. A., Ramsey, J. G., & Silva, F. V. M. (2016). High pressure processing and thermosonication of beer: Comparing the energy requirements and *Saccharomyces cerevisiae* ascospores inactivation with thermal processing and modeling. *Journal of Food Engineering*, 181, 35–41. <https://doi.org/10.1016/j.jfoodeng.2016.02.023>
- Molan, P. (2001). Why honey is effective as a medicine. *Bee World*, 82(1), 22–40. <https://doi.org/10.1080/0005772X.2001.11099498>
- Molyneux, P. (2004). The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakar J Sci Technol*, 26(2), 211–219.
- Moniruzzaman, M., Khalil, M. I., Sulaiman, S. A., & Gan, S. H. (2013). Physicochemical and antioxidant properties of Malaysian honeys produced by *Apis cerana*, *Apis dorsata* and *Apis mellifera*. *BMC Complementary and Alternative Medicine*, 13(1), 1–12. <https://doi.org/10.1186/1472-6882-13-43>
- Munro, J. A. (1943). The Viscosity and Thixotropy of Honey. *Journal of Economic Entomology*, 36(5), 769–777. <https://doi.org/10.1093/jee/36.5.769>
- Murphy, M., Cowan, C., Henchion, M., & O'Reilly, S. (2000). Irish consumer preferences for honey: a conjoint approach. *British Food Journal*, 102(8), 585–598. <https://doi.org/10.1108/00070700010348424>
- Musielak, G., Mierzwa, D., & Kroehnke, J. (2016). Food drying enhancement by ultrasound – A review. *Trends in Food Science & Technology*, 56, 126–141. <https://doi.org/10.1016/j.tifs.2016.08.003>
- Myers, R. H., Montgomery, D. C., & Anderson-Cook, C. M. (2009). *Response surface methodology: process and product optimization using designed experiments*. John Wiley & Sons.
- Naknean, P., Meenune, M., & Roudaut, G. (2009). Changes in physical and chemical properties during the production of palm sugar syrup by open pan and vacuum evaporator. *Asian Journal of Food and Agro-Industry*, 2(4), 448–456.
- Nava, E., Palmer, R. M. J., & Moncada, S. (1991). Inhibition of nitric oxide synthesis in septic shock: how much is beneficial? *The Lancet*, 338(8782), 1555–1557. [https://doi.org/http://dx.doi.org/10.1016/0140-6736\(91\)92375-C](https://doi.org/http://dx.doi.org/10.1016/0140-6736(91)92375-C)
- Nelson, J. M., & Cohn, D. J. (1924). Invertase in honey. *Journal of Biological Chemistry*, 61(1), 193–224. Retrieved from <http://www.jbc.org/content/61/1/193.short>
- Nicoli, M. C., Anese, M., Parpinel, M. T., Franceschi, S., & Lerici, C. R. (1997). Loss and/or formation of antioxidants during food processing and storage. *Cancer Lett*, 114(1–2), 71–74.
- Noci, F., Walkling-Ribeiro, M., Cronin, D. A., Morgan, D. J., & Lyng, J. G. (2009). Effect of thermosonication, pulsed electric field and their combination on inactivation of *Listeria innocua* in milk. *International Dairy Journal*, 19(1), 30–



35. <https://doi.org/10.1016/j.idairyj.2008.07.002>

- Oddo, L. P., Heard, T. A., Rodríguez-Malaver, A., Pérez, R. A., Fernández-Muiño, M., Sancho, M. T., ... Vit, P. (2008). Composition and Antioxidant Activity of Trigona carbonaria Honey from Australia. *Journal of Medicinal Food*, 11(4), 789–794. <https://doi.org/10.1089/jmf.2007.0724>
- Oroian, M., Amariei, S., Escriche, I., Leahu, A., Damian, C., & Gutt, G. (2014). Chemical Composition and Temperature Influence on the Rheological Behaviour of Honey. *International Journal of Food Properties*, 17(10), 2228–2240. <https://doi.org/10.1080/10942912.2013.791835>
- Örsi, F. (1973). Kinetic studies on the thermal decomposition of glucose and fructose. *Journal of Thermal Analysis and Calorimetry*, 5(2–3), 329–335.
- Page, J., & Henry, D. (2000). Consumption of nsoids and the development of congestive heart failure in elderly patients: An underrecognized public health problem. *Archives of Internal Medicine*, 160(6), 777–784. Retrieved from <http://dx.doi.org/10.1001/archinte.160.6.777>
- Pérez, R. A., Iglesias, M. T., Pueyo, E., González, M., & de Lorenzo, C. (2007). Amino Acid Composition and Antioxidant Capacity of Spanish Honey. *Journal of Agricultural and Food Chemistry*, 55(2), 360–365. <https://doi.org/10.1021/jf062055b>
- Piyasena, P., Mohareb, E., & McKellar, R. C. (2003). Inactivation of microbes using ultrasound: a review. *International Journal of Food Microbiology*, 87(3), 207–216. [https://doi.org/10.1016/S0168-1605\(03\)00075-8](https://doi.org/10.1016/S0168-1605(03)00075-8)
- Postmes, T. J., Bosch, M. M. C., Dutrieux, R., van Baare, J., & Hoekstra, M. J. (1997). Speeding Up the Healing of Burns with Honey. In A. Mizrahi & Y. Lensky (Eds.), *Bee Products: Properties, Applications, and Apitherapy* (pp. 57–63). Boston, MA: Springer US. [https://doi.org/10.1007/978-1-4757-9371-0\\_6](https://doi.org/10.1007/978-1-4757-9371-0_6)
- Pryce-Jones, J. (1952). ‘Stringiness’ in Honey and in Sugar Syrup Fed to Bees. *Bee World*, 33(9), 147–150. <https://doi.org/10.1080/0005772X.1952.11094748>
- Ratajski, A., Białobrzewski, I., Dajnowiec, F., & Bakier, S. (2010). The use of ultrasonic methods in the identification of honey types. *Technical Sciences/University of Warmia and Mazury in Olsztyn*, (13), 22–29.
- Raviyan, P., Zhang, Z., & Feng, H. (2005). Ultrasonication for tomato pectinmethylesterase inactivation: effect of cavitation intensity and temperature on inactivation. *Journal of Food Engineering*, 70(2), 189–196. <https://doi.org/10.1016/j.jfoodeng.2004.09.028>
- Rawson, A., Tiwari, B. K., Patras, A., Brunton, N., Brennan, C., Cullen, P. J., & O'Donnell, C. (2011). Effect of thermosonication on bioactive compounds in watermelon juice. *Food Research International*, 44(5), 1168–1173. <https://doi.org/10.1016/j.foodres.2010.07.005>
- Rodríguez-Malaver, A. J., Rasmussen, C., Gutiérrez, M. G., Gil, F., Nieves, B., & Vit, P. (2009). Properties of honey from ten species of Peruvian stingless bees. *Natural Product Communications*, 4(9), 1221–1226. Retrieved from <http://europepmc.org/abstract/MED/19831033>

- Rosales, G. R. O. (2013). Medicinal Uses of *Melipona beecheii* Honey, by the Ancient Maya. In P. Vit, S. R. M. Pedro, & D. Roubik (Eds.), *Pot-Honey: A legacy of stingless bees* (pp. 229–240). New York, NY: Springer New York. [https://doi.org/10.1007/978-1-4614-4960-7\\_15](https://doi.org/10.1007/978-1-4614-4960-7_15)
- Sahin, S., & Sumnu, S. G. (2006). Water Activity and Sorption Properties of Foods. In *Physical Properties of Foods* (pp. 193–228). New York, NY: Springer New York. [https://doi.org/10.1007/0-387-30808-3\\_5](https://doi.org/10.1007/0-387-30808-3_5)
- Sala, F. J., Burgos, J., Condón, S., Lopez, P., & Raso, J. (1995). Effect of heat and ultrasound on microorganisms and enzymes. In G. W. Gould (Ed.), *New Methods of Food Preservation* (pp. 176–204). Boston, MA: Springer US. [https://doi.org/10.1007/978-1-4615-2105-1\\_9](https://doi.org/10.1007/978-1-4615-2105-1_9)
- Šarić, G., Marković, K., Major, N., Krpan, M., Uršulin-Trstenjak, N., Hruškar, M., & Vahčić, N. (2012). Changes of antioxidant activity and phenolic content in acacia and multifloral honey during storage. *Food Technology and Biotechnology*, 50(4), 434–441.
- Saxena, S., Gautam, S., & Sharma, A. (2010). Physical, biochemical and antioxidant properties of some Indian honeys. *Food Chemistry*, 118(2), 391–397. <https://doi.org/http://dx.doi.org/10.1016/j.foodchem.2009.05.001>
- Saxena, S., Panicker, L., & Gautam, S. (2014). Rheology of Indian Honey: Effect of Temperature and Gamma Radiation. *International Journal of Food Science*, 2014, 6. <https://doi.org/10.1155/2014/935129>
- Serhan, C. N., Ward, P. A., & Gilroy, D. W. (2010). *Fundamentals of inflammation*. Cambridge University Press.
- Sesta, G., & Lusco, L. (2008). Refractometric determination of water content in royal jelly. *Apidologie*, 39(2), 225–232. <https://doi.org/10.1051/apido:2007053>
- Silva, I. A. A., Souza, A. L., Cordeiro, A. M. T. M., Soledade, L. E. B., Queiroz, N., & Souza, A. G. (2013). Thermal degradation of honeys and evaluation of physicochemical properties. *Journal of Thermal Analysis and Calorimetry*, 114(1), 353–358. <https://doi.org/10.1007/s10973-012-2926-x>
- Singh, V. R., & Dwivedi, S. (1995). Ultrasonic detection of adulteration in fluid foods. In *Engineering in Medicine and Biology Society, 1995 and 14th Conference of the Biomedical Engineering Society of India. An International Meeting, Proceedings of the First Regional Conference., IEEE* (p. 1/73-1/74). <https://doi.org/10.1109/RCEMBS.1995.508696>
- Singleton, V. L., Orthofer, R., & Lamuela-Raventos, R. M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*, (299C), 152–178.
- Smith, F. G. (1965). The Sucrose Content of Western Australian Honey. *Journal of Apicultural Research*, 4(3), 177–184. <https://doi.org/10.1080/00218839.1965.11100120>
- Strober, W. (2001). Trypan Blue Exclusion Test of Cell Viability. In *Current Protocols in Immunology*. John Wiley & Sons, Inc. <https://doi.org/10.1002/0471142735.ima03bs111>

- Subramanian, R., Umesh Hebbar, H., & Rastogi, N. K. (2007). Processing of Honey: A Review. *International Journal of Food Properties*, 10(1), 127–143. <https://doi.org/10.1080/10942910600981708>
- Suntiparapop, K., Prapaipong, P., & Chantawannakul, P. (2012). Chemical and biological properties of honey from Thai stingless bee (*Tetragonula leaviceps*). *Journal of Apicultural Research*, 51(1), 45–52. <https://doi.org/10.3896/IBRA.1.51.1.06>
- Suslick, K. S. (1989). The chemical effects of ultrasound. *Scientific American*, 260(2), 80–86.
- Sylvester, P. W. (2011). Optimization of the Tetrazolium Dye (MTT) Colorimetric Assay for Cellular Growth and Viability. In S. D. Satyanarayanajois (Ed.), *Drug Design and Discovery: Methods and Protocols* (pp. 157–168). Totowa, NJ: Humana Press. [https://doi.org/10.1007/978-1-61779-012-6\\_9](https://doi.org/10.1007/978-1-61779-012-6_9)
- Thrasyvoulou, A., Manikis, J., & Tselios, D. (1994). Liquefying crystallized honey with ultrasonic waves. *Apidologie*, 25, 297.
- Tiwari, B. K., Muthukumarappan, K., O'Donnell, C. P., & Cullen, P. J. (2008). Colour degradation and quality parameters of sonicated orange juice using response surface methodology. *LWT - Food Science and Technology*, 41(10), 1876–1883. <https://doi.org/10.1016/j.lwt.2007.11.016>
- Tiwari, B. K., Patras, A., Brunton, N., Cullen, P. J., & O'Donnell, C. P. (2010). Effect of ultrasound processing on anthocyanins and color of red grape juice. *Ultrasonics Sonochemistry*, 17(3), 598–604. <https://doi.org/10.1016/j.ultsonch.2009.10.009>
- Tong, Q., Zhang, X., Wu, F., Tong, J., Zhang, P., & Zhang, J. (2010). Effect of honey powder on dough rheology and bread quality. *Food Research International*, 43(9), 2284–2288. <https://doi.org/10.1016/j.foodres.2010.08.002>
- Tonks, A. J., Cooper, R. A., Jones, K. P., Blair, S., Parton, J., & Tonks, A. (2003). Honey stimulates inflammatory cytokine production from monocytes. *Cytokine*, 21(5), 242–247. [https://doi.org/10.1016/S1043-4666\(03\)00092-9](https://doi.org/10.1016/S1043-4666(03)00092-9)
- Tosi, E. A., Ré, E., Lucero, H., & Bulacio, L. (2004). Effect of honey high-temperature short-time heating on parameters related to quality, crystallisation phenomena and fungal inhibition. *LWT - Food Science and Technology*, 37(6), 669–678. <https://doi.org/10.1016/j.lwt.2004.02.005>
- Tsai, T.-H., Tsai, P.-J., & Ho, S.-C. (2005). Antioxidant and Anti-inflammatory Activities of Several Commonly Used Spices. *Journal of Food Science*, 70(1), C93–C97. <https://doi.org/10.1111/j.1365-2621.2005.tb09028.x>
- Turkmen, N., Sari, F., Poyrazoglu, E. S., & Velioglu, Y. S. (2006). Effects of prolonged heating on antioxidant activity and colour of honey. *Food Chemistry*, 95(4), 653–657. <https://doi.org/10.1016/j.foodchem.2005.02.004>
- Van den Berg, A. J. J., Van den Worm, E., Quarles van Ufford, H. C., Halkes, S. B. A., Hoekstra, M. J., & Beukelman, C. J. (2008). An in vitro examination of the antioxidant and anti-inflammatory properties of buckwheat honey. *Journal of Wound Care*, 17(4), 172–179.

- Vit, P., Bogdanov, S., & Kilchenmann, V. (1994). Composition of Venezuelan honeys from stingless bees (Apidae: Meliponinae) and *Apis mellifera* L. *Apidologie*, 25(3), 278–288.
- Vit, P., Medina, M., & Eunice Enríquez, M. (2004). Quality standards for medicinal uses of Meliponinae honey in Guatemala, Mexico and Venezuela. *Bee World*, 85(1), 2–5. <https://doi.org/10.1080/0005772X.2004.11099603>
- Wang, X. H., Gheldof, N., & Engeseth, N. J. (2004). Effect of Processing and Storage on Antioxidant Capacity of Honey. *Journal of Food Science*, 69(2), fct96-fct101. <https://doi.org/10.1111/j.1365-2621.2004.tb15509.x>
- White, J. W. (1979). Spectrophotometric method for hydroxymethylfurfural in honey. *Journal - Association of Official Analytical Chemists*, 62(3), 509–514. Retrieved from <http://europepmc.org/abstract/MED/479072>
- White Jr, J. W. (1978). Honey. In C. O. Chichester (Ed.), *Advances in Food Research* (Vol. Volume 24, pp. 287–374). Academic Press. [https://doi.org/10.1016/S0065-2628\(08\)60160-3](https://doi.org/10.1016/S0065-2628(08)60160-3)
- Wu, G., & Morris, S. M. (1998). Arginine metabolism: nitric oxide and beyond. *Biochemical Journal*, 336(1), 1 LP-17. Retrieved from <http://www.biochemj.org/content/336/1/1.abstract>
- Wu, J., Gamage, T. V., Vilku, K. S., Simons, L. K., & Mawson, R. (2008). Effect of thermosonication on quality improvement of tomato juice. *Innovative Food Science & Emerging Technologies*, 9(2), 186–195. <https://doi.org/10.1016/j.ifset.2007.07.007>
- Yanniotis, S., Skaltsi, S., & Karaburnioti, S. (2006). Effect of moisture content on the viscosity of honey at different temperatures. *Journal of Food Engineering*, 72(4), 372–377. <https://doi.org/10.1016/j.jfoodeng.2004.12.017>
- Zaitoun, S., Ghzawi, A. A.-M., Al-Malah, K. I. M., & Abu-Jdayil, B. (2001). Rheological Properties of Selected Light Coloured Jordanian Honey. *International Journal of Food Properties*, 4(1), 139–148. <https://doi.org/10.1081/JFP-100002192>
- Zamora, G. (2015). The antioxidant capacity and immunomodulatory activity of stingless bee honeys proceeding from Costa Rica. *Oxidants and Antioxidants in Medical Science*, 4(1), 49–55. <https://doi.org/10.5455/oams.180415.or.084>
- Zhao, L., Zhao, G., Chen, F., Wang, Z., Wu, J., & Hu, X. (2006). Different Effects of Microwave and Ultrasound on the Stability of (all-E)-Astaxanthin. *Journal of Agricultural and Food Chemistry*, 54(21), 8346–8351. <https://doi.org/10.1021/jf061876d>