



UNIVERSITI PUTRA MALAYSIA

***PHENOTYPIC AND MOLECULAR ANALYSES OF COAGULASE
NEGATIVE STAPHYLOCOCCI, PREDOMINANTLY COMPRISING
Staphylococcus epidermidis, ISOLATED FROM A STUDENT
POPULATION IN MALAYSIA***

PUNG HUI PING

FPSK(M) 2015 81



PHENOTYPIC AND MOLECULAR ANALYSES OF COAGULASE NEGATIVE STAPHYLOCOCCI, PREDOMINANTLY COMPRISING *Staphylococcus epidermidis*, ISOLATED FROM A STUDENT POPULATION IN MALAYSIA

By

PUNG HUI PING

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Master of Science

September 2015

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Science

**PHENOTYPIC AND MOLECULAR ANALYSES OF COAGULASE NEGATIVE
STAPHYLOCOCCI, PREDOMINANTLY COMPRISING *Staphylococcus
epidermidis*, ISOLATED FROM A STUDENT POPULATION IN MALAYSIA**

By

PUNG HUI PING

September 2015

Chair: Assoc. Prof. Mohd. Nasir bin Mohd. Desa, PhD
Faculty: Medicine and Health Science

Coagulase negative staphylococci (CoNS), predominantly *Staphylococcus epidermidis*, are commonly considered as commensal skin and nasal colonizing bacteria but have emerged as one of the leading causes of nosocomial infections. To date, limited studies have been conducted to investigate CoNS or methicillin resistant CoNS (MR-CoNS) in Malaysian community. Hence, this study was initiated to evaluate the status of CoNS in relation to their nasal carriage, risk factors, antimicrobial resistance, molecular characteristics and genetic background of the bacterial collection in a student population. This study was conducted at Universiti Putra Malaysia in November 2013 involving 192 health science students. A self-administrated questionnaire on socio-demographic and risk factors was distributed, followed by collection of nasal swab. Cultivation on mannitol salt agar (MSA) plates and basic phenotypic tests were used to preliminarily differentiate CoNS from *Staphylococcus aureus*. All CoNS isolates were subjected to antibiotic susceptibility test against 10 antibiotics and *mecA* gene screening. The *mecA* positive isolates were further subjected to staphylococcal chromosome cassette *mec* (SCC*mec*) typing and multilocus sequence typing (MLST) to identify the SCC*mec* types and sequence type (ST) of the isolates, respectively. CoNS isolates that were resistant to at least one antibiotic or were *mecA* positive were subjected to species identification based on *tuf* gene sequencing, screening of erythromycin resistance associated genes (*ermA*, *ermB*, *ermC*, and *msrA*), random amplified polymorphic DNA- (RAPD) and enterobacterial repetitive intergenic consensus-polymerase chain reaction (ERIC-PCR). In this study, a total of 120 isolates of CoNS (62.5%) were collected from 192 health sciences students. Chi-square

test showed significant association of nasal carriage of CoNS with gender ($P = 0.0455$), history of cold or fever ($P = 0.0147$) and presence of unhealed wound ($P = 0.0467$). Similarly, the P -value of the use of intravascular devices was close to significant ($P = 0.0589$). The other risk factors (ethnicity, health status and habit of touching nose) were deemed as statistically not significant. Resistance to penicillin was the highest among the isolates (58.3%) followed by tetracycline (9.2%), cefoxitin (8.3%), erythromycin (7.5%), oxacillin (6.7%), ceftazidime (5.0%), gentamicin (2.5%) and rifampin (1.7%). Two isolates of CoNS were identified as multidrug resistant strains (MDR) which are resistant to more than three antimicrobial families. Among 76 isolates that were subjected to *tuf* gene sequencing, 73 isolates were identified as *S. epidermidis* and three isolates as *S. haemolyticus*. Fifteen isolates of CoNS (12.5%) harbored *mecA* gene (13 *S. epidermidis*; 2 *S. haemolyticus*) whereas five isolates (4.2%) carried *msrA* gene (3 *S. epidermidis*; 2 *S. haemolyticus*). Twelve of the *mecA* positive isolates carried SCC*mec* type IV (11 *S. epidermidis*; 1 *S. haemolyticus*), two isolates carried SCC*mec* type II (1 *S. epidermidis*; 1 *S. haemolyticus*) and one isolate carried SCC*mec* type I (*S. epidermidis*). MLST analysis of MRSE isolates revealed four STs: ST57 (n=6), ST59 (n=1), ST72 (n=1) and ST193 (n=1). However, four of the isolates had non-typeable (new) ST. RAPD and ERIC analyses revealed that majority of the isolates were largely genetically distinct. We observed a potential dissemination of a few distinct MR-CoNS carrying predominantly SCC*mec* type IV in the population rather than a clonal spread in both analyses. Overall, the study showed a high prevalence of nasal carriage CoNS which had relatively low level of antibiotic resistance except for penicillin and had extensive genetic diversity.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Sarjana Sains

**ANALISIS FENOTIPIK DAN MOLEKUL UNTUK NEGATIF KOAGULASE
STAPHYLOCOCCI, TERUTAMANYA *Staphylococcus epidermidis*, YANG
DIKUMPUL DARIPADA POPULASI PELAJA DI MALAYSIA**

Oleh

PUNG HUI PING

September 2015

Pengerusi: Prof. Madya Mohd. Nasir bin Mohd. Desa, PhD
Fakulti: Perubatan dan Sains Kesihatan

Negative koagulase staphylococci (CoNS), terutamanya *Staphylococcus epidermidis* adalah mikrob yang biasanya hidup di permukaan kulit dan dalam rongga hidung tetapi telah menjadi penyebab utama bagi jangkitan mikrob di hospital. Sehingga kini, kajian mengenai CoNS atau ketahanan methicillin CoNS (MR-CoNS) terutamanya dalam kalangan komuniti di Malaysia kurang diberi perhatian. Maka, penyelidikan ini bertujuan untuk mengkaji taraf pembawa, factor-faktor risiko, ketahanan antimikrob, ciri-ciri molecular dan latar belakang genetik CoNS dalam kalangan populasi pelajar. Kajian ini dijalankan di Universiti Putra Malaysia pada November 2013 dengan melibatkan 192 pelajar Sains Kesihatan. Subjek kajian diberikan soal selidik yang mengandungi soalan mengenai sosio-demografik dan factor-faktor risiko sebelum pengambilan sampel swab hidung. Pertumbuhan bakteria di atas agar garam mannitol (MSA) dan ujian asas fenotipik digunakan sebagai ujian awal untuk membezakan CoNS daripada *Staphylococcus aureus*. Semua isolat bakteria telah menjalani ujian kepekaan antibiotic terhadap 10 jenis antibiotic dan penyaringan gen *mecA*. Selanjutnya, sampel yang mempunyai gen *mecA* telah menjalani ujian pengelasan staphylococcal kromosom kaset *mec* (SCC*mec*) dan pengelasan jujukan multilokus untuk menentukan jenis SCC*mec* dan penjenisan jujukan (ST). Ujian-ujian seperti penjujukan gen *tuf*, penyaringan gen-gen penentang erythromycin (*ermA*, *ermB*, *ermC*, dan *msrA*), Amplifikasi DNA Polimorf Rawak (RAPD) and *Enterobacterial Repetitive Intergenic Consensus*-PCR (ERIC-PCR) dijalankan untuk mengenalpastikan ciri-ciri CoNS isolat yang menentang minimum satu jenis antimikrob atau mengandungi gen *mecA*. Sebanyak 120 CoNS isolat (62.5%) telah dikumpulkan daripada 192 pelajar Sains Kesihatan. Ujian khi-kuasa dua menunjukkan hubungan penting antara taraf pembawa CoNS dengan tiga faktor, iaitu jantina ($P = 0.0455$), jangkitan selesema atau demam ($P = 0.0147$) dan luka yang belum sembuh ($P = 0.0467$). Factor-faktor lain seperti kaum, taraf kesihatan dan sebagainya dibuktikan tidak bererti tetapi factor

penggunaan alat suntikan ($P = 0.0589$) dibuktikan mempunyai hubungan yang mungkin berarti. Dalam penyelidikan ini, taraf ketahanan terhadap penisilin adalah tertinggi (58.3%), diikuti oleh tetracycline (9.2%), cefoxitin (8.3%), erythromycin (7.5%), oxacillin (6.7%), ceftazidime (5.0%), gentamicin (2.5%) and rifampin (1.7%). Hanya dua isolat telah dikesan mempunyai ketahanan terhadap pelbagai dadah (MDR) iaitu tahan terhadap sekurang-kurangnya tiga kelas antimikrob. Daripada 76 isolat CoNS yang menjalani ujian penyaringan jujukan gen *tuf*, 73 isolat adalah *S. epidermidis* dan tiga isolat adalah *S. haemolyticus*. Lima belas isolat (12.5%) mengandungi gen *mecA* (13 *S. epidermidis*; 2 *S. haemolyticus*) dan lima isolat (4.2%) mempunyai gen *msrA* (3 *S. epidermidis*; 2 *S. haemolyticus*). Dua belas isolat yang positif *mecA* membawa SCC*mec* jenis IV (11 *S. epidermidis*; 1 *S. haemolyticus*), dua isolat membawa SCC*mec* jenis II (1 *S. epidermidis*; 1 *S. haemolyticus*) and satu isolat SCC*mec* jenis I (*S. epidermidis*). Melalui analisis MLST untuk isolate MRSE, sebanyak empat STs ditemui iaitu ST57 (n=6), ST59 (n=1), ST72 (n=1) and ST193 (n=1). Walaubagaimanapun, ST bagi empat isolat MRSE tidak dapat dijeniskan. Analisis RAPD dan ERIC menunjukkan bahawa kebanyakan isolat berbeza dari segi genetik. Kami telah memerhati potensi pembawa yang terhad terhadap beberapa MR-CoNS, terutamanya SCC*mec* jenis IV dalam kalangan pelajar ini berbanding penyebaran klon dalam kedua-dua analisis. Kesimpulannya, penyelidikan ini menunjukkan kadar pembawa CoNS yang tinggi di dalam hidung di mana ia mempunyai taraf antimikrob yang rendah (kecuali penisilin) dan kepelbagaian genetic yang ekstrem.

ACKNOWLEDGEMENTS

First and foremost, I would like to thank God for giving me strength in completing this study. Furthermore, this study would not be completed without the endless help and support from the people mentioned below. It is my honor to express my gratitude towards these individuals.

I would like to express my deepest gratitude and appreciation to my supervisor, Assoc. Prof. Dr. Mohd. Nasir bin Mohd. Desa. He has been a great mentor throughout the course of this study by offering opinion and sharing your unsurpassed knowledge generously. Without his guidance, persistent help and patience, this study would not have been possible. I am really grateful with the encouragement that he has given me from all the time.

I would like to thank my co-supervisor, Assoc. Prof. Dr. Syafinaz Amin Nordin, for her suggestive comments and advices. I truly appreciate each of the helpful feedback that she had offered regarding the study.

My appreciation goes to the laboratory staff in the Applied Microbiology Laboratory, Mr. Sabri for his kindest help in providing the materials for my study and maintaining the equipment in the laboratory. On the other hand, I would like to thank my fellow coursemates who are always there to provide assistance, helpful opinions and knowledge. Moreover, we always motivate each other to stay strong and persevere through this journey together.

Last but not least, I am deeply indebted to my family. I am truly appreciated the sacrifices that they have made all the time. They are always there with me through ups and downs in the journey of learning by providing endless moral support. They are the main reason that keeps me going and persevere through all the obstacles.

I certify that a Thesis Examination Committee has met on 11 September 2015 to conduct the final examination of Pung Hui Ping on her thesis entitled "Phenotypic and Molecular Analyses of Coagulase Negative Staphylococci Predominantly Comprising *Staphylococcus epidermidis* Isolated from A Student Population in Malaysia" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Roslida binti Abd Hamid @ Abdul Razak, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Ngah Zasmy a/l Unyah, PhD

Senior Lecturer
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Internal Examiner)

Yeo Chew Chieng, PhD

Professor
Universiti Sultan Zainal Abidin
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 5 November 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Mohd. Nasir bin Mohd. Desa, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Syafinaz Amin Nordin, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Pung Hui Ping (GS37964)


Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: _____

Signature: _____
Name of Member of
Supervisory
Committee: _____



© COPYRIGHT UPM

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	ix
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF APPENDICES	xv
LIST OF ABBREVIATIONS	xvi
CHAPTER	
1 INTRODUCTION	1
1.1 Introduction	1
1.2 Problem statement	3
1.3 Research hypothesis	3
1.4 Objectives	3
1.4.1 General objectives	3
1.4.2 Specific objectives	3
2 LITERATURE REVIEW	4
2.1 Coagulase Negative Staphylococcus (CoNS)	4
2.1.1 <i>Staphylococcus epidermidis</i>	6
2.1.2 <i>Staphylococcus haemolyticus</i>	7
2.2 Pathogenesis and commensal lifestyle of CoNS	8
2.2.1 Antibiotic resistance	8
2.2.2 Toxins and virulence-associated factors	11
2.2.3 Commensal colonization of CoNS	12
2.3 Genotyping	13
2.3.1 Random amplified polymorphic DNA-PCR (RAPD) and enterobacterial repetitive intergenic consensus-PCR (ERIC)	13
2.3.2 Multilocus sequence typing (MLST)	14

3	METHODOLOGY	16
3.1	Study population	16
3.1.1	Sample size calculation	16
3.1.2	Sample collection	17
3.2	Questionnaires	17
3.2.1	Data collection	17
3.2.2	Statistical Analysis: Chi-square test	17
3.3	Identification of CoNS by phenotypic test	18
3.3.1	Gram stain	18
3.3.2	Coagulase test	18
3.3.3	Novobiocin susceptibility test	18
3.4	Antibiotic susceptibility test	19
3.4.1	Kirby Bauer disc diffusion method	19
3.4.2	Epsilometer test (E-test)	19
3.5	DNA extraction	20
3.6	Detection of antibiotic resistance associated determinants by PCR	20
3.6.1	Detection of <i>mecA</i> gene	20
3.6.2	SCC <i>mec</i> typing	21
3.6.3	Detection of erythromycin resistance associated genes	22
3.7	Identification of species by <i>tuf</i> -gene sequence	22
3.8	Genotyping	23
3.8.1	Multilocus sequence typing (MLST)	23
3.8.2	Random amplified polymorphic DNA-PCR (RAPD-PCR)	24
3.8.3	Enterobacterial repetitive intergenic consensus PCR (ERIC-PCR)	25
4	RESULTS	26
4.1	Identification and prevalence of CoNS	26
4.2	Data analysis	28
4.2.1	Association of nasal carriage CoNS with socio-demographic factors	28
4.2.2	Association of nasal carriage CoNS with various risk factors	29
4.3	Antibiotic susceptibility test of CoNS	31
4.3.1	Disc diffusion test	31
4.3.2	Determination of ceftazidime MIC by E-test	34
4.4	Identification of species by <i>tuf</i> -gene sequencing	34
4.5	Screening of <i>mecA</i> gene and SCC <i>mec</i> typing	35

4.6	Screening of erythromycin resistance associated genes	37
4.7	MLST	38
4.7.1	Identification of MRSE STs	38
4.7.2	e-BURST analysis of MLST data	39
4.7.3	RAPD analysis	41
4.7.4	ERIC analysis	44
5	DISCUSSION	47
5.1	Prevalence of CoNS and its association with socio-demographic and various risk factors	49
5.2	Correlation between antibiotic susceptibility patterns and antibiotic resistance associated genes	53
5.3	MLST and e-BURST analysis of MRSE	53
5.4	Characterization of genetic diversity of CoNS by RAPD and ERIC	55
6	CONCLUSION	57
6.1	Summary and conclusion	57
6.2	Limitations of study and recommendations for future studies	58
	REFERENCES	60
	APPENDICES	74
	BIODATA OF STUDENT	91
	LIST OF PUBLICATIONS	92

LIST OF TABLES

Table		Page
3.1	Nucleotide sequence of primers used in SCC <i>mec</i> typing	21
3.2	Nucleotide sequence of primers used in multiplex PCR of erythromycin resistance associated genes	22
3.3	Nucleotide sequence of primers used in PCR amplification of seven housekeeping genes	24
4.1	Chi-square test (P-value) showing the association between socio-demographic factors and nasal carriage CoNS	29
4.2a	Chi-square test (P-value) showing the association between various risk factors and nasal carriage CoNS	30
4.2b	Chi-square test (P-value) showing the association between various risk factors and nasal carriage of CoNS	31
4.3	Result of AST (Disc diffusion method)	32
4.4	MLST results of MRSE	39

LIST OF FIGURES

Figure		Page
2.1	Roles of <i>S. epidermidis</i> as commensal colonizing bacteria and pathogen	13
4.1	CoNS on MSA plate	26
4.2	Gram positive and cocci in grape-like clusters under 100X magnification	27
4.3	(a) Negative coagulase test of representative CoNS isolates. (b) Blank rabbit serum control; positive coagulase test of <i>S. aureus</i> ATCC 25923 and <i>S. aureus</i> ATCC 700699.	27
4.4	Novobiocin susceptibility test of representative CoNS isolates (E008)	28
4.5	Type of antibiotics with respect to percentage of resistant isolates	33
4.6	Antibiotics susceptibility tests by disc diffusion method for representative isolate (E2)	33
4.7	E-test (cefoxitin) of <i>S. epidermidis</i> ATCC 12228 and representative isolate (E111)	34
4.8	Agarose gel image of <i>tuf</i> gene (412 bp)	35
4.9	Agarose gel image of <i>mecA</i> gene (147 bp)	36
4.10	Agarose gel electrophoresis of PCR products for <i>SCCmec</i> types	37
4.11	Agarose gel images of multiplex PCR of erythromycin resistance associated genes	38
4.12a	E-BURST analysis of MLST data showing CC2	40
4.12b	Part of CC2 with focus on ST57 and ST59	40
4.13	E-BURST analysis of MLST data showing CC193	41
4.14a	Dendrogram of RAPD PCR profiles analysis showing genetic relationship between 76 isolates of CoNS	43
4.14b	Part of RAPD dendrogram (zoom in)	44
4.15	Genetic relationship between 76 isolates of CoNS	46

LIST OF APPENDICES

Appendix		Page
A1	Preparation of culture media	74
A1.1	Mannitol salt agar (MSA)	74
A1.2	Trypticase soy agar (TSA)	74
A1.3	Mueller Hinton agar (MHA)	74
A1.4	Glycerol stock	74
A2	Agarose gel electrophoresis	74
A2.1	1x TBE buffer	74
A2.2	1.5%, 1.8% or 2% agarose gel	74
B1	Representative gel electrophoresis images	75
B1.1	<i>mecA</i> gene	75
B1.2	<i>tuf</i> gene	75
B1.3	Multiplex PCR of <i>ermA</i> , <i>msrA</i> and 16S rRNA	76
B1.4	Multiplex PCR of <i>ermB</i> , <i>ermC</i> and 16S rRNA	76
B1.5	MLST – <i>arcC</i> gene	77
B1.6	MLST – <i>aroE</i> gene	77
B1.7	MLST – <i>gtr</i> gene	78
B1.8	MLST – <i>mutS</i> gene	78
B1.9	MLST – <i>pyrR</i> gene	79
B1.10	MLST – <i>tpi</i> gene	79
B1.11	MLST – <i>yqiL</i> gene	80
B2	Sequencing Data (BLAST results retrieved from GenBank at NCBI)	80
B2.1	Representative of <i>mecA</i> gene (E111)	80
B2.2	Representative of <i>msrA</i> gene (E23)	80
B2.3	Representatives of <i>tuf</i> gene	81
B2.4	Representative of seven housekeeping genes of MLST (<i>S. epidermidis</i> ATCC 12228)	82
B2.4.1	<i>arcC</i> gene	82
B2.4.2	<i>aroE</i> gene	83
B2.4.3	<i>gtr</i> gene	83
B2.4.4	<i>mutS</i> gene	84
B2.4.5	<i>pyrR</i> gene	84
B2.4.6	<i>tpi</i> gene	85
B2.4.7	<i>yqiL</i> gene	85
C1	Ethical approval	86
C2	Consent form	89
C3	Questionnaire	89

LIST OF ABBREVIATIONS

AMP	Antimicrobial peptide
AP-PCR	Arbitrarily primed polymerase chain reaction
ATCC	American Type Culture Collection
CNA	Colistin-nalidixic acid agar
CC	Clonal Complex
CLSI	Clinical and Laboratory Standards Institute
ERIC	Enterobacterial repetitive intergenic consensus
FAME	Fatty acid modifying enzyme
GlcNAc	<i>N</i> -acetylglucosamine
IS	Insertion sequence
MGE	Mobile genetic element
MHA	Mueller Hinton agar
MLST	Multilocus sequence typing
MR-CoNS	Methicillin resistant coagulase negative staphylococci
MRSA	Methicillin resistant <i>Staphylococcus aureus</i>
MRSE	Methicillin resistant <i>Staphylococcus epidermidis</i>
MSA	Mannitol salt agar
MSCRAMMs	Microbial surface components recognizing adhesive matrix molecules
PCR	Polymerase chain reaction
PFGE	Pulse field gel electrophoresis
PGA	Poly- γ -glutamic acid
PIA	Polysaccharide intercellular adhesion
PNAG	Poly- <i>N</i> -acetylglucosamine
PSM	Phenol-soluble modulins
RAPD	Random amplified polymorphic DNA
SCC	Staphylococcal Chromosome Cassette
SLV	Single locus variants
SPSS	Scientific Package of Social Science
ST	Sequence type
TSA	Trypticase soy agar
TSB	Trypticase soy broth
TSST-1	Toxic shock syndrome toxin 1

CHAPTER 1

INTRODUCTION

1.1 Introduction

Staphylococci are opportunistic pathogens that commonly colonize skin and mucous membrane of humans. The absence of enzyme coagulase distinguishes coagulase negative staphylococci (CoNS) from coagulase positive staphylococci, such as *Staphylococcus aureus*. *Staphylococcus epidermidis* is the most widespread CoNS species and is frequently isolated from human epithelia, axillae (armpits), head and nostril (Otto, 2009). The less frequently isolated species such as *Staphylococcus haemolyticus* and *Staphylococcus saprophyticus* are more commonly isolated from axillae and the female perineum area, respectively (Takeuchi et al., 2005). Generally, CoNS species distribute throughout the human body, indicating the importance of CoNS in maintaining healthy skin flora. The colonization of CoNS establishes on human skin almost immediately after birth and continues to be the dominant species during the first few months of life (Christensen & Brüggemann, 2014; Dong & Speer, 2014). CoNS usually maintain a mutual relationship with the human host by limiting the growth of pathogenic bacteria (Otto, 2014a). Despite the beneficial characteristics of CoNS, this bacterial species can cause infections in high risk individuals such as newborns, elder individuals and immunocompromised patients.

Two staphylococcal species, *S. aureus* and *S. epidermidis* have been associated with almost one million infections caused in hospitals every year around the world (Gill et al., 2005). Although CoNS have less pathogenic potential than *S. aureus*, the CoNS species particularly *S. epidermidis* and *S. haemolyticus* have emerged as the leading causative agents of nosocomial infections after *S. aureus* (Otto, 2009). CoNS infections are usually associated with the use of indwelling medical devices such as urinary tract catheters, pace maker, prosthetic joints and other medical implants (Ziebuhr et al., 2006). The lack of immunity barrier in high risk individuals eases the invasion and colonization of CoNS. The production of biofilm-associated proteins allows CoNS to adhere on inert surfaces and also host tissues. In addition to biofilm formation, clinical isolates of CoNS with multiple antibiotic resistances deteriorates the condition of nosocomial infections and increases the difficulty of treatments (Schoenfelder et al., 2010; Ziebuhr et al., 2006). Unlike methicillin resistant *S. aureus* (MRSA), eradication of antibiotic resistant CoNS particularly methicillin resistant strains is difficult because of the ubiquitous presence of CoNS and their role as commensal bacteria.

Multiple drug resistant (MDR) strains of CoNS which usually possess methicillin resistance, are not uncommon in clinical settings and are gradually emerging in

community settings. According to several studies, 60-85% of clinical strains are methicillin resistant CoNS (MR-CoNS) and 70-90% of hospital isolates of *S. epidermidis* are resistant to methicillin (MRSE), a rate which is actually higher than that of *S. aureus* (Otto, 2009; Jamaluddin et al., 2008). Moreover, the rate of MR-CoNS is relatively high in community settings (11-31%) (Lebeaux et al., 2012; Barbier et al., 2010; Ruppé et al., 2009; Jamaluddin et al., 2008; Silva, Mattos, Coimbra, Ferreira-Carvalho, & Figueiredo, 2001). Resistance to methicillin in staphylococci is due to the presence of the *mecA* gene which expresses mutated penicillin binding protein 2a (PBP2a), a transpeptidase with a low affinity for β - lactams (Zong, Peng, & Lu, 2011). The *mecA* gene is carried in mobile genetic elements (MGE) known as the staphylococcal cassette chromosome *mec* (SCC*mec*) (Zong et al., 2011). Due to the recombinases encoded by the cassette chromosome recombinase (*ccr*) gene, SCC*mec* complex can excise from and integrate into chromosomes among different staphylococcal species (Hanssen & Sollid, 2006). With flexible genomes, CoNS are believed to be the earlier reservoirs which transfer the methicillin resistance determinant into *S. aureus* to become MRSA (Otto, 2009; Hanssen, Kjeldsen, & Sollid, 2004).

In clinical settings, CoNS are usually considered as contaminants and are seldom identified to species level. However, with increased significance in infections, identification of CoNS species is important for better diagnosis and more efficient treatment. As compared to phenotypic identification methods, molecular methods involving polymerase chain reaction (PCR) and DNA sequencing are considered to produce more accurate results. The *tuf* gene sequencing which was found to have better discriminatory power than 16S rRNA, had been utilized in several studies to identify CoNS species (Bergeron et al., 2011; Hwang, Kim, Park, Song, & Kim, 2011; Heikens et al., 2005). Furthermore, genotyping methods such as random amplified polymorphic DNA- (RAPD) and enterobacterial repetitive intergenic consensus-PCR (ERIC) are useful in phylogenetic analysis of CoNS. These methods which are suitable for short term epidemiological studies can also be used to examine the occurrence of clonal dissemination in a community. On the other hand, multilocus sequence typing (MLST) is a relatively new genotyping method that analyzes phylogeny and evolution of bacterial species based on DNA sequencing of seven housekeeping genes. Currently, MLST is applicable for only *S. epidermidis* among CoNS species. However, with the advantages of standardized nomenclature and data comparison globally, this method has been utilized for many other bacterial species (Widerström, Wiström, Sjöstedt, & Monsen, 2012).

Community-acquired staphylococcal infections have been on the rise especially with increased antimicrobial resistance (Silva et al., 2001). Nonetheless, most research studies are concentrating on *S. aureus* or MRSA. Moreover, research on CoNS were usually focused on clinical isolates. There is limited information on the colonization of CoNS particularly antibiotic resistant strains in healthy individuals in community (Silva et al., 2001). Currently, there is only one report on clinical isolates of *S. epidermidis* in Malaysia and no research work on community acquired MR-CoNS have been reported. As a potential reservoir of

different antibiotic resistance determinants, CoNS can live as commensal microflora on human skin and can be potentially pathogenic to individuals with compromised immunity. Therefore, it is vital to investigate the dissemination of methicillin resistant strains among nasal isolates of CoNS from the community. Furthermore, genotyping analyzes can provide new insights and reveal the genetic relatedness between CoNS strains.

1.2 Problem statement

The prevalence rate of MR-CoNS in communities such as children, college students, public community and others from different countries was reported to be in the range of 11% to 31%. This raises the concern on the prevalent of nasal carriage CoNS especially potentially virulent strains in a student population. In addition, the antibiotic susceptibility patterns and the genetic background of CoNS collection from community were of great value for the literature as limited studies had been done on community isolates of CoNS.

1.3 Research hypothesis

It was hypothesized that there is high prevalence of nasal carriage CoNS in this student population. The prevalence rate of MR-CoNS was expected to be in range of 11 to 31% as reported in previous studies. The collection of nasal isolates CoNS was predicted to have a diverse genetic background.

1.4 Objectives

1.4.1 General objectives

In this study, our main objective was to investigate the phenotypic and molecular epidemiology of CoNS in a university student population. The status of CoNS was evaluated in relation to their nasal carriage and risk factors, antibiotic resistance patterns, molecular characteristics and genotypes.

1.4.2 Specific objectives

- To investigate the prevalence of CoNS nasal carriage within a student population and their association with socio-demographic and various risk factors.
- To examine the antibiotic susceptibility patterns of CoNS isolates through phenotypic methods (disc diffusion test).
- To identify the antibiotic resistance determinants such as *mecA* (SCC*mec*) and erythromycin resistance-associated genes (*ermA*, *ermB*, *ermC* and *msrA*) through PCR screening.
- To identify the representative CoNS isolates at the species level by *tuf* gene sequencing and to evaluate the genetic background of these isolates through different genotyping methods (RAPD-PCR, ERIC-PCR and MLST).

REFERENCES

- Amin Nordin, S., Za'im, N. A. N., Sahari, N. N., Jamaluddin, S. F., Ahmadi, S., & Mohd Desa, M. N. (2012). Staphylococcus aureus nasal carriers among medical students in a medical school. *The Medical Journal of Malaysia*, 67(6), 636–8.
- Agostino, M. (2012). Introduction to the BLAST Suite and BLASTN Key concepts. In *Practical Bioinformatics* (pp. 37–71). New York, USA: Garland Science.
- Ardic, N., Ozyurt, M., Sareyyupoglu, B., & Haznedaroglu, T. (2005). Investigation of erythromycin and tetracycline resistance genes in methicillin-resistant staphylococci. *International Journal of Antimicrobial Agents*, 26(3), 213–218. <http://doi.org/10.1016/j.ijantimicag.2005.06.013>
- Barbier, F., Ruppé, E., Hernandez, D., Lebeaux, D., Francois, P., Felix, B., ... Ruimy, R. (2010). Methicillin-resistant coagulase-negative staphylococci in the community: high homology of SCCmec IVa between *Staphylococcus epidermidis* and major clones of methicillin-resistant *Staphylococcus aureus*. *The Journal of Infectious Diseases*, 202(2), 270–281. <http://doi.org/10.1016/j.yder.2011.11.016>
- Barros, E. M., Lemos, M., Souto-Padrón, T., & Giambiagi-deMarval, M. (2015). Phenotypic and genotypic characterization of biofilm formation in *Staphylococcus haemolyticus*. *Current Microbiology*, 829–834. <http://doi.org/10.1007/s00284-015-0794-x>
- Berger-Bächli, B., & Rohrer, S. (2002). Factors influencing methicillin resistance in staphylococci. *Archives of Microbiology*, 178(3), 165–171. <http://doi.org/10.1007/s00203-002-0436-0>
- Bergeron, M., Dauwalder, O., Gouy, M., Freydiere, a. M., Bes, M., Meugnier, H., ... Boisset, S. (2011). Species identification of staphylococci by amplification and sequencing of the *tuf* gene compared to the gap gene and by matrix-assisted laser desorption ionization time-of-flight mass spectrometry. *European Journal of Clinical Microbiology and Infectious Diseases*, 30(3), 343–354. <http://doi.org/10.1007/s10096-010-1091-z>
- Biavasco, F., Vignaroli, C., & Varaldo, P. E. (2000). Glycopeptide Resistance in coagulase-negative staphylococci. *European Journal of Clinical Microbiology and Infectious Diseases*, 19, 403–417.
- Bilung, L. M., Fuh, Y. S., Linang, V., Benjamin, A., Vincent, M., Apun, K., ... Lin, C. S. (2014). Genomic diversity of cholera outbreak strains in east malaysia. *Malaysian Journal of Medicine and Health Sciences*, 10(June), 19–26.

- Boye, K., Bartels, M. D., Anderson, I. S., Moller, J. A., & Westh, H. (2007). A new multiplex PCR for easy screening of methicillin-resistant *Staphylococcus aureus* SCCmec types I-V. *Clinical Microbiology and Infection*, 13(7), 725–727. <http://doi.org/10.3109/10826087609056170>
- Burnie, J. P., Naderi-Nasab, M., Loudon, K. W., & Matthews, R. C. (1997). An epidemiological study of blood culture isolates of coagulase-negative staphylococci demonstrating hospital-acquired infection. *Journal of Clinical Microbiology*, 35(7), 1746–1750.
- Casey, A. L., Worthington, T., Caddick, J. M., Hilton, A. C., Lambert, P. A., & Elliott, T. S. J. (2006). RAPD for the typing of coagulase-negative staphylococci implicated in catheter-related bloodstream infection. *Journal of Infection*, 52, 282–289. <http://doi.org/10.1016/j.jinf.2005.06.005>
- Cavanagh, J. P., Hjerde, E., Holden, M. T. G., Kahlke, T., Klingenberg, C., Flægstad, T., ... Sollid, J. U. E. (2014). Whole-genome sequencing reveals clonal expansion of multiresistant *Staphylococcus haemolyticus* in European hospitals. *The Journal of Antimicrobial Chemotherapy*, (July), dku271–. <http://doi.org/10.1093/jac/dku271>
- Cetin, E. S., Gunes, H., Kaya, S., Aridogan, B. C., & Demirci, M. (2010). Distribution of genes encoding resistance to macrolides, lincosamides and streptogramins among clinical staphylococcal isolates in a Turkish university hospital. *Journal of Microbiology, Immunology and Infection*, 43(6), 524–529. [http://doi.org/10.1016/S1684-1182\(10\)60081-3](http://doi.org/10.1016/S1684-1182(10)60081-3)
- Chaieb, K., Abbassi, M. S., Touati, A., Hassen, A. B. E. N., Mahdouani, K., & Bakhrouf, A. (2005). Molecular characterization of *Staphylococcus epidermidis* isolated from biomaterials in a dialysis service. *Annals of Microbiology*, 55(4), 307–312.
- Chamberlain, N. R., & Brueggemann, S. a. (1997). Characterisation and expression of fatty acid modifying enzyme produced by *Staphylococcus epidermidis*. *Journal of Medical Microbiology*, 46(8), 693–697. <http://doi.org/10.1099/00222615-46-8-693>
- Chambers, H. F., Hartman, B. J., & Tomasz, A. (1985). Increased amounts of a novel penicillin binding protein in a strain of methicillin-resistant *Staphylococcus aureus*. *Journal of Clinical Investigation*, 76(July), 325–331.
- Cherifi, S., Byl, B., Deplano, A., Nagant, C., Nonhoff, C., Denis, O., & Hallin, M. (2014). Genetic characteristics and antimicrobial resistance of *Staphylococcus epidermidis* isolates from patients with catheter-related bloodstream infections and from colonized healthcare workers in a

Belgian hospital. *Annals of Clinical Microbiology and Antimicrobials*, 13(20), 1–8. <http://doi.org/10.1186/1476-0711-13-20>

Cherifi, S., Byl, B., Deplano, A., Nonhoff, C., Denis, O., & Hallin, M. (2013). Comparative epidemiology of *Staphylococcus epidermidis* isolates from patients with catheter-related bacteremia and from healthy volunteers. *Journal of Clinical Microbiology*, 51(5), 1541–1547. <http://doi.org/10.1128/JCM.03378-12>

Cheung, G. Y. C., Rigby, K., Wang, R., Queck, S. Y., Braughton, K. R., Whitney, A. R., ... Otto, M. (2010). *Staphylococcus epidermidis* strategies to avoid killing by human neutrophils. *PLoS Pathogens*, 6(10). <http://doi.org/10.1371/journal.ppat.1001133>

Christensen, G. J. M., & Brüggemann, H. (2014). Bacterial skin commensals and their role as host guardians. *Beneficial Microbes*, 5(2), 201–215. <http://doi.org/10.3920/BM2012.0062>

Cimiotti, J. P., Wu, F., Della-Latta, P., Nesin, M., & Larson, E. (2007). Emergence of resistant staphylococci on the hands of new graduate nurses. *Infection Control and Hospital Epidemiology*, 25(5), 431–435.

Clarridge III, J. E. (2004). Impact of 16S rRNA gene sequence analysis for identification of bacteria on clinical microbiology and infectious diseases. *Clinical Microbiology Reviews*, 17(4), 840–862. <http://doi.org/10.1128/CMR.17.4.840>

CLSI: *Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Third Informational Supplement*. (2012) (Vol. 31). Wayne, PA: Clinical and Laboratory Standards Institute.

Cuevas, O., Cercenado, E., Vindel, A., Sa, M., & Sa, M. (2004). Evolution of the antimicrobial resistance of *Staphylococcus* spp. in Spain: five Nationwide prevalence studies, 1986 to 2002. *Antimicrobial Agents and Chemotherapy*, 48(11), 4240–4245. <http://doi.org/10.1128/AAC.48.11.4240>

Delost, M. D. (1997). *Introduction to Diagnostic Microbiology*. Missouri: Mosby, Inc.

Didelot, X., & Maiden, M. C. J. (2010). Impact of recombination on bacterial evolution. *Trends in Microbiology*, 18(7), 315–322. <http://doi.org/10.1523/JNEUROSCI.2928-12.2013>. Gating

Dong, Y., & Speer, C. P. (2014). The role of *Staphylococcus epidermidis* in neonatal sepsis: Guarding angel or pathogenic devil? *International Journal of Medical Microbiology*, 304(5-6), 513–520. <http://doi.org/10.1016/j.ijmm.2014.04.013>

- Du, X., Zhu, Y., Song, Y., Li, T., Luo, T., Sun, G., ... Li, M. (2013). Molecular analysis of *Staphylococcus epidermidis* strains isolated from community and hospital environments in China. *PLoS ONE*, 8(5), 1–10. <http://doi.org/10.1371/journal.pone.0062742>
- Enright, M. C., & Spratt, B. G. (1999). Multi-locus sequence typing. *Methods in Molecular Medicine*, 7(12), 482–487.
- Falcone, M., Giannella, M., Raponi, G., Mancini, C., & Venditti, M. (2006). Teicoplanin use and emergence of *Staphylococcus haemolyticus*: Is there a link? *Clinical Microbiology and Infection*, 12(1), 96–97. <http://doi.org/10.1111/j.1469-0691.2005.01307.x>
- Fassler, J., & Cooper, P. (2011). *BLAST help*. Bethesda: National Center for Biotechnology Information (US). Retrieved from <http://www.ncbi.nlm.nih.gov/books/NBK1762/>
- Fatokun, O. (2014). Exploring antibiotic use and practices in a Malaysian community. *International Journal of Clinical Pharmacy*, 36(3), 564–569. <http://doi.org/10.1007/s11096-014-9937-6>
- Feil, E. J., Li, B. C., Aanensen, D. M., Hanage, W. P., & Spratt, B. G. (2004). eBURST: Inferring patterns of evolutionary descent among clusters of related bacterial genotypes from multilocus sequence typing data. *Journal of Bacteriology*, 186(5), 1518–1530. <http://doi.org/10.1128/JB.186.5.1518-1530.2004>
- Fredheim, E. G. A., Klingenberg, C., Rohde, H., Frankenberger, S., Gaustad, P., Flægstad, T., & Sollid, J. E. (2009). Biofilm formation by *Staphylococcus haemolyticus*. *Journal of Clinical Microbiology*, 47(4), 1172–1180. <http://doi.org/10.1128/JCM.01891-08>
- Frogatt, J. W., Johnston, J. L., Galetto, D. W., & Archer, G. L. (1989). Antimicrobial resistance in nosocomial isolates of *Staphylococcus haemolyticus*. *Antimicrobial Agents and Chemotherapy*, 33(4), 460–466. <http://doi.org/10.1128/AAC.33.4.460>
- Ghebremedhin, B., Layer, F., König, W., König, B., Ghebremedhin, B., Layer, F., & Ko, W. (2008). Genetic classification and distinguishing of staphylococcus species based on different partial *gap*, 16S rRNA, *hsp60*, *rpoB*, *sodA*, and *tuf* gene sequences. *Journal of Clinical Microbiology*, 46(3), 1019–1025. <http://doi.org/10.1128/JCM.02058-07>
- Gill, S. R., Fouts, D. E., Archer, G. L., Mongodin, E. F., Deboy, R. T., Ravel, J., ... Fraser, C. M. (2005). Insights on evolution of virulence and resistance from the complete genome analysis of an early methicillin-resistant *Staphylococcus aureus* strain and a biofilm-producing methicillin-resistant *Staphylococcus epidermidis* strain. *Journal of Bacteriology*, 187(7), 2426–2438. <http://doi.org/10.1128/JB.187.7.2426>

- Hanssen, A., Kjeldsen, G., & Sollid, J. U. E. (2004). Local variants of staphylococcal cassette chromosome mec in sporadic methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase negative staphylococci: Evidence of horizontal gene transfer? *Antimicrobial Agents and Chemotherapy*, 48(1), 285–296. <http://doi.org/10.1128/AAC.48.1.285>
- Hanssen, A., & Sollid, J. U. E. (2006). SCCmec in staphylococci : genes on the move. *FEMS Immunology and Medical Microbiology*, 46, 8–20. <http://doi.org/10.1111/j.1574-695X.2005.00009.x>
- Harder, J., & Schroder, J. M. (2005). Antimicrobial peptides in human skin. *Chemical Immunology and Allergy*, 86, 22–41. <http://doi.org/10.1159/000086651>
- Heikens, E., Fleer, A., Paauw, A., Florijn, A., & Fluit, A. C. (2005). Comparison of genotypic and phenotypic methods for species-level identification of clinical isolates of coagulase-negative staphylococci. *Journal of Clinical Microbiology*, 43(5), 2286–2290. <http://doi.org/10.1128/JCM.43.5.2286>
- Hellmark, B., Söderquist, B., Unemo, M., & Nilsson, Å. (2013). Comparison of *Staphylococcus epidermidis* isolated from prosthetic joint infections and commensal isolates in regard to antibiotic susceptibility , agr type , biofilm production , and epidemiology. *International Journal of Medical Microbiology*, 303(1), 32–39. <http://doi.org/10.1016/j.ijmm.2012.11.001>
- Hisata, K., Kuwahara-arai, K., Yamamoto, M., Ito, T., Nakatomi, Y., Cui, L., ... Hiramatsu, K. (2005). Dissemination of methicillin-resistant staphylococci among healthy japanese children. *Journal of Clinical Microbiology*, 43(7), 3364–3372. <http://doi.org/10.1128/JCM.43.7.3364>
- Huang, Y. C., Hwang, K. P., Chen, P. Y., Chen, C. J., & Lin, T. Y. (2007). Prevalence of Methicillin-Resistant *Staphylococcus aureus* Nasal Colonization among Taiwanese Children in 2005 and 2006. *Journal of Clinical Microbiology*, 45(12), 3992–3995. <http://doi.org/10.1128/JCM.01202-07>
- Hwang, S. M., Kim, M. S., Park, K. U., Song, J., & Kim, E. C. (2011). tuf gene sequence analysis has greater discriminatory power than 16S rRNA sequence analysis in identification of clinical isolates of coagulase-negative staphylococci. *Journal of Clinical Microbiology*, 49(12), 4142–4149. <http://doi.org/10.1128/JCM.05213-11>
- Ibrahim, S., Salmenlinna, S., Virolainen, A., Kerttula, A. M., Lyytikäinen, O., Jägerroos, H., ... Vuopio-Varkila, J. (2009). Carriage of methicillin-resistant staphylococci and their SCCmec types in a long-term-care facility. *Journal of Clinical Microbiology*, 47(1), 32–37. <http://doi.org/10.1128/JCM.01085-08>

- Ito, T., Katayama, Y., Asada, K., Mori, N., Tsutsumimoto, K., Tiensasitorn, C., & Hiramatsu, K. (2001). Structural comparison of three types of staphylococcal cassette chromosome *mec* integrated in the chromosome in methicillin-resistant *Staphylococcus aureus*. *Antimicrobial Agents and Chemotherapy*, 45(5), 1323–1336. <http://doi.org/10.1128/AAC.45.5.1323>
- Ito, T., Okuma, K., Ma, X. X., Yuzawa, H., & Hiramatsu, K. (2003). Insights on antibiotic resistance of *Staphylococcus aureus* from its whole genome: Genomic island SCC. *Drug Resistance Updates*, 6(1), 41–52. [http://doi.org/10.1016/S1368-7646\(03\)00003-7](http://doi.org/10.1016/S1368-7646(03)00003-7)
- Institute, C. and L. S. (2012). *Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Third Informational Supplement. Enterobacteriaceae* (Vol. 31)
- Iwase, T., Uehara, Y., Shinji, H., Tajima, A., Seo, H., Takada, K., ... Mizunoe, Y. (2010). *Staphylococcus epidermidis* Esp inhibits *Staphylococcus aureus* biofilm formation and nasal colonization. *Nature*, 465(7296), 346–349. <http://doi.org/10.1038/nature09074>
- Jamaluddin, T. Z. M., Tengku, M., Kuwahara-arai, K., Hisata, K., Terasawa, M., Cui, L., ... Hiramatsu, K. (2008). Extreme genetic diversity of methicillin-resistant *Staphylococcus epidermidis* strains disseminated among healthy Japanese children. *Journal of Clinical Microbiology*, 46(11), 3778–3783. <http://doi.org/10.1128/JCM.02262-07>
- Join-Lambert, O. F., Clauser, S., Guillet, C., Jais, J. P., Abachin, E., Quesnes, G., ... Ferroni, A. (2007). Comparison of cefoxitin and moxalactam 30 µg disc diffusion methods for detection of methicillin resistance in coagulase-negative staphylococci. *Journal of Antimicrobial Chemotherapy*, 59(4), 763–766. <http://doi.org/10.1093/jac/dkl548>
- Katayama, Y., Ito, T., & Hiramatsu, K. (2000). A New Class of Genetic Element , Staphylococcus Cassette Chromosome *mec* encodes methicillin resistance in *Staphylococcus aureus*. *Antimicrobial Agents and Chemotherapy*, 44(6), 1549–1555.
- Kim, M., Heo, S. R., Choi, S. H., Kwon, H., Park, J. S., Seong, M. W., ... Kim, E. C. (2008). Comparison of the MicroScan, VITEK 2, and Crystal GP with 16S rRNA sequencing and MicroSeq 500 v2.0 analysis for coagulase-negative staphylococci. *BMC Microbiology*, 8, 233. <http://doi.org/10.1186/1471-2180-8-233>
- Kloos, W. E., & Musselwhite, M. S. (1975). Distribution and persistence of *Staphylococcus* and *Micrococcus* species and other aerobic bacteria on human skin. *Applied Microbiology*, 30(3), 381–385.

- Kloos, W. E., & Schleifer, K. H. (1975). Simplified scheme for routine identification of human *Staphylococcus* species. *Journal of Clinical Microbiology*, 1(1), 82–88.
- Kocianova, S., Vuong, C., Yao, Y., Voyich, J. M., Fischer, E. R., DeLeo, F. R., & Otto, M. (2005). Key role of poly-gamma-DL-glutamic acid in immune evasion and virulence of *Staphylococcus epidermidis*. *The Journal of Clinical Investigation*, 115(3), 688–694. <http://doi.org/10.1172/JCI200523523>
- Lebeaux, D., Barbier, F., Angebault, C., Benmahdi, L., Ruppé, E., Felix, B., & Gaillard, K. (2012). Evolution of nasal carriage of methicillin-resistant coagulase-negative staphylococci in a remote population. *Antimicrobial Agents and Chemotherapy*, 56(1), 315–323. <http://doi.org/10.1128/AAC.00547-11>
- Leclercq, R. (2002). Mechanisms of resistance to macrolides and lincosamides: nature of the resistance elements and their clinical implications. *Clinical Infectious Diseases*, 34(4), 482–492. <http://doi.org/10.1086/324626>
- Li, M., Lai, Y., Villaruz, A. E., Cha, D. J., Sturdevant, D. E., & Otto, M. (2007). Gram-positive three-component antimicrobial peptide-sensing system. *Proceedings of the National Academy of Sciences of the United States of America*, 104(22), 9469–9474.
- Li, M., Wang, X., Gao, Q., Lu, Y., & Lu, Y. (2009). Molecular characterization of *Staphylococcus epidermidis* strains isolated from a teaching hospital in Shanghai, China. *Journal of Medical Microbiology*, 58, 456–461. <http://doi.org/10.1099/jmm.0.007567-0>
- Lim, J. A., Kwon, A. R., Kim, S. K., Chong, Y., Lee, K., & Choi, E. C. (2002). Prevalence of resistance to macrolide, lincosamide and streptogramin antibiotics in Gram-positive cocci isolated in a Korean hospital. *The Journal of Antimicrobial Chemotherapy*, 49(3), 489–495.
- Lim, K. K., & Teh, C. C. (2012). A cross sectional study of public knowledge and attitude towards antibiotics in Putrajaya, Malaysia. *Southern Medical Review*, 5(2), 26–33.
- Maiden, M. C., Bygraves, J. a, Feil, E., Morelli, G., Russell, J. E., Urwin, R., ... Spratt, B. G. (1998). Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proceedings of the National Academy of Sciences of the United States of America*, 95(6), 3140–3145. <http://doi.org/10.1073/pnas.95.6.3140>
- Mangold, K. A., Tajuddin, M., Huntington, A., Fritz, B., Thomson, R. B., & Kaul, K. L. (2005). Species-level identification of staphylococcal isolates by

real-time PCR and melt curve analysis. *Journal of Clinical Investigation*, 43(6), 2876–2880. <http://doi.org/10.1128/JCM.43.6.2876>

Martineau, F., Picard, F. J., Ke, D., Paradis, S., Roy, P. H., Ouellette, M., & Bergeron, M. G. (2001). Development of a PCR assay for identification of staphylococci at genus and species levels. *Journal of Clinical Microbiology*, 39(7), 2541–2547. <http://doi.org/10.1128/JCM.39.7.2541>

Martineau, F., Picard, F. J., Lansac, N., Ménard, C., Roy, P. H., Ouellette, M., & Michel, G. (2000). Correlation between the resistance genotype determined by multiplex PCR assays and the antibiotic susceptibility patterns of *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Antimicrobial Agents and Chemotherapy*, 44(2), 231–238. <http://doi.org/10.1128/AAC.44.2.231-238.2000>. Updated

Miragaia, M., Carriço, J. a., Thomas, J. C., Couto, I., Enright, M. C., & De Lencastre, H. (2008). Comparison of molecular typing methods for characterization of *Staphylococcus epidermidis*: Proposal for clone definition. *Journal of Clinical Microbiology*, 46(1), 118–129. <http://doi.org/10.1128/JCM.01685-07>

Miragaia, M., Couto, I., & De Lencastre, H. (2005). Genetic diversity among methicillin-resistant *Staphylococcus epidermidis* (MRSE). *Microbial Drug Resistance*, 11(2), 259–264.

Miragaia, M., Thomas, J. C., Couto, I., Enright, M. C., & Lencastre, H. De. (2007). Inferring a population structure for *Staphylococcus epidermidis* from multilocus sequence typing data □. *Journal of Bacteriology*, 189(6), 2540–2552. <http://doi.org/10.1128/JB.01484-06>

Oh, A. L., Hassali, M. A., Al-Haddad, M. S., Syed Sulaiman, S. A., Shafie, A. A., & Awaisu, A. (2011). Public knowledge and attitudes towards antibiotic usage: a cross-sectional study among the general public in the state of Penang, Malaysia. *The Journal of Infection in Developing Countries*, 5(05). <http://doi.org/10.3855/jidc.1502>

Otto, M. (2009). *Staphylococcus epidermidis* - the “accidental” pathogen. *Nature Reviews Microbiology*, 7(8), 555–567. <http://doi.org/10.1038/nrmicro.2182>. Staphylococcus

Otto, M. (2012a). Coagulase-negative staphylococci as reservoirs of genes facilitating MRSA infection. *Bioessays*, 35, 4–11. <http://doi.org/10.1002/bies.201200112>

Otto, M. (2012b). Molecular basis of *Staphylococcus epidermidis* infections. *Seminars in Immunopathology*, 34, 201–214. <http://doi.org/10.1007/s00281-011-0296-2>

- Otto, M. (2014a). Phenol-soluble modulins. *International Journal of Medical Microbiology*, 304(2), 164–169. <http://doi.org/10.1016/j.biotechadv.2011.08.021>. Secreted
- Otto, M. (2014b). *Staphylococcus epidermidis* pathogenesis. In *Staphylococcus Epidermidis: Methods and Protocols* (Vol. 1106, pp. 17–31). <http://doi.org/10.1007/978-1-62703-736-5>
- Otto, M., Echner, H., & Voelter, W. (2001). Pheromone cross-inhibition between *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Infection and Immunity*, 69(3), 1957–1960. <http://doi.org/10.1128/IAI.69.3.1957>
- Pereira, M. S. V., Leal, N. C., Leal, T. C. a., Sobreira, M., De Almeida, a. M. P., Siqueira-Júnior, J. P., & Campos-Takaki, G. M. (2002). Typing of human and bovine *Staphylococcus aureus* by RAPD-PCR and ribotyping-PCR. *Letters in Applied Microbiology*, 35(1), 32–36. <http://doi.org/10.1046/j.1472-765X.2002.01127.x>
- Piette, A., & Verschraegen, G. (2009). Role of coagulase-negative staphylococci in human disease. *Veterinary Microbiology*, 134(1-2), 45–54. <http://doi.org/10.1016/j.vetmic.2008.09.009>
- Podkowik, M., Park, J. Y., Seo, K. S., Bystro, J., & Bania, J. (2013). Enterotoxigenic potential of coagulase-negative staphylococci. *International Journal of Food Microbiology*, 163, 34–40. <http://doi.org/10.1016/j.ijfoodmicro.2013.02.005>
- Preacher, K. J., & Briggs, N. E. (2001). Calculation for the chi-square test: An interactive calculation tool for chi-square tests of goodness of fit and independence [Computer software]. Retrieved from <http://quantpsy.org>.
- Putt, S. N. H., Shaw, A. P. M., Woods, A. J., Tyler, L., & James, A. D. L. (1987). *Veterinary epidemiology and economics in Africa - A manual for use in the design and appraisal of livestock health policy*. Retrieved from <http://www.fao.org/wairdocs/ilri/x5436e/x5436e07.htm>
- Queck, S. Y., Khan, B. a., Wang, R., Bach, T. H. L., Kretschmer, D., Chen, L., ... Otto, M. (2009). Mobile genetic element-encoded cytolysin connects virulence to methicillin resistance in MRSA. *PLoS Pathogens*, 5(7), 1–12. <http://doi.org/10.1371/journal.ppat.1000533>
- Raimundo, O., Heussler, H., Bruhn, J. B., Suntrarachun, S., Kelly, N., Deighton, M. A., & Garland, S. M. (2002). Molecular epidemiology of coagulase-negative staphylococcal bacteraemia in a newborn intensive care unit. *Journal of Hospital Infection*, 51, 33–42. <http://doi.org/10.1053/jhin.2002.1203>

- Rolo, J., Lencastre, D., & Miragaia, M. (2012). Strategies of adaptation of *Staphylococcus epidermidis* to hospital and community : amplification and diversification of SCCmec. *Journal of Antimicrobial Chemotherapy*, 67, 1333–1341. <http://doi.org/10.1093/jac/dks068>
- Rubinstein, E., & Keynan, Y. (2014). Vancomycin revisited – 60 years later. *Frontiers in Public Health*, 2(October), 1–7. <http://doi.org/10.3389/fpubh.2014.00217>
- Ruppé, E., Barbier, F., Mesli, Y., Cojocar, R., Benkhalfat, M., Hassaine, H., ... Soumare, S. (2009). Diversity of staphylococcal cassette chromosome *mec* structures in methicillin-resistant *Staphylococcus epidermidis* and *Staphylococcus haemolyticus* strains among outpatients from four countries. *Antimicrobial Agents and Chemotherapy*, 53(2), 442–449. <http://doi.org/10.1128/AAC.00724-08>
- Sahal, G., & Bilkay, I. S. (2014). Multi drug resistance in strong biofilm forming clinical isolates of *Staphylococcus epidermidis*. *Brazilian Journal of Microbiology*, 45(2), 539–544.
- Sakoulas, G., & Moellering, R. C. (2008). Increasing antibiotic resistance among methicillin-resistant *Staphylococcus aureus* strains. *Clinical Infectious Diseases*, 46(5), S360–S367. <http://doi.org/10.1086/533592>
- Sani, N. A. M., Sapri, H. F., Neoh, H., & Hussin, S. (2014). First report on the molecular epidemiology of Malaysian *Staphylococcus epidermidis* isolated from a university teaching hospital. *BMC Research Notes*, 7(1), 1–4. <http://doi.org/10.1186/1756-0500-7-597>
- Schoenfelder, S. M. K., Lange, C., Eckart, M., Hennig, S., Kozytska, S., & Ziebuhr, W. (2010). Success through diversity - How *Staphylococcus epidermidis* establishes as a nosocomial pathogen. *International Journal of Medical Microbiology*, 300(6), 380–386. <http://doi.org/10.1016/j.ijmm.2010.04.011>
- Shah, M. M., Iihara, H., Noda, M., Song, S. X., Nhung, P. H., Ohkusu, K., ... Ezaki, T. (2007). dnaJ gene sequence-based assay for species identification and phylogenetic grouping in the genus *Staphylococcus*. *International Journal of Systematic and Evolutionary Microbiology*, 57(1), 25–30. <http://doi.org/10.1099/ijms.0.64205-0>
- Shin, J. H., Kim, S. H., Jeong, H. S., Oh, S. H., Kim, H. R., Lee, J. N., ... Kim, Y. H. (2011). Identification of coagulase-negative staphylococci isolated from continuous ambulatory peritoneal dialysis fluid using 16S ribosomal RNA, *tuf*, and *SodA* gene sequencing. *Peritoneal Dialysis International*, 31(3), 340–346. <http://doi.org/10.3747/pdi.2010.00073>
- Shore, A. C., & Coleman, D. C. (2013). Staphylococcal cassette chromosome *mec* : Recent advances and new insights. *International Journal of Medical*

Microbiology, 303(6-7), 350–359. <http://doi.org/10.1016/j.ijmm.2013.02.002>

Silva, F. R., Mattos, E. M., Coimbra, M. V. S., Ferreira-Carvalho, B. T., & Figueiredo, A. M. S. (2001). Isolation and molecular characterization of methicillin-resistant coagulase-negative staphylococci from nasal flora of healthy humans at three community institutions in Rio de Janeiro City. *Epidemiology & Infection*, 127, 57–62.

Stepanović, S., Hauschild, T., Dakić, I., Al-Doori, Z., Švabić-Vlahović, M., Ranin, L., & Morrison, D. (2006). Evaluation of phenotypic and molecular methods for detection of oxacillin resistance in members of the *Staphylococcus sciuri* group. *Journal of Clinical Microbiology*, 44(3), 934–937. <http://doi.org/10.1128/JCM.44.3.934-937.2006>

Stojanov, M., Sakwinska, O., & Moreillon, P. (2013). Expression of *scmec* cassette chromosome recombinases in methicillin-resistant *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Journal of Antimicrobial Chemotherapy*, 68(4), 749–757. <http://doi.org/10.1093/jac/dks494>

Suhaili, Z., Johari, S. A., Sajili, M. H., Yahya, A., Zakaria, Z. A., Nasir, M., ... Ali, A. M. (2013). In silico PCR verification and simplex real-time PCR detection of methicillin-resistant *Staphylococcus aureus* (MRSA) from east coast Malaysian clinical isolates. *Walailak Journal*, 10(3), 237–246.

Swenson, J. M., Tenover, F. C., Addison, R., D'Souza, H., O'Connor, J., Rothberg, J., ... Willey, B. (2005). Results of disk diffusion testing with cefoxitin correlate with presence of *mecA* in *Staphylococcus* spp. *Journal of Clinical Microbiology*, 43(8), 3818–3823. <http://doi.org/10.1128/JCM.43.8.3818-3823.2005>

Takeuchi, F., Watanabe, S., Baba, T., Yuzawa, H., Morimoto, Y., Kuroda, M., ... Hiramatsu, K. (2005). Whole-genome sequencing of *Staphylococcus haemolyticus* uncovers the extreme plasticity of its genome and the evolution of human-colonizing staphylococcal species. *Journal of Bacteriology*, 187(21), 7292–7308. <http://doi.org/10.1128/JB.187.21.7292>

Thomas, J. C., Vargas, M. R., Miragaia, M., Peacock, S. J., Archer, G. L., & Enright, M. C. (2007). Improved multilocus sequence typing scheme for *Staphylococcus epidermidis*. *Journal of Clinical Microbiology*, 45(2), 616–619. <http://doi.org/10.1128/JCM.01934-06>

Trzcinski, K., Cooper, B. S., Hryniewicz, W., & Dowson, C. G. (2000). Expression of resistance to tetracyclines in strains of methicillin-resistant *Staphylococcus aureus*. *Journal of Antimicrobial Chemotherapy*, 45(6), 763–770. <http://doi.org/10.1093/jac/45.6.763>

- Tsubakishita, S., Kuwahara-Arai, K., Sasaki, T., & Hiramatsu, K. (2010). Origin and molecular evolution of the determinant of methicillin resistance in staphylococci. *Antimicrobial Agents and Chemotherapy*, *54*(10), 4352–4359. <http://doi.org/10.1128/AAC.00356-10>
- Turner, K. M. E., Hanage, W. P., Fraser, C., Connor, T. R., & Spratt, B. G. (2007). Assessing the reliability of eBURST using simulated populations with known ancestry. *BMC Microbiology*, *7*, 30. <http://doi.org/10.1186/1471-2180-7-30>
- Van Belkum, A., Bax, R., Peerbooms, P., Goessens, W. H., van Leeuwen, N., & Quint, W. G. (1993). Comparison of phage typing and DNA fingerprinting by polymerase chain reaction for discrimination of methicillin-resistant *Staphylococcus aureus* strains. *Journal of Clinical Microbiology*, *31*(4), 798–803.
- Vasanthakumari, N., Alshrari, A. S. D., Rad, E. G., Moghaddam, H. G., van Belkum, A., Alreshidi, M. A., ... Nor Shamsudin, M. (2009). Highly dynamic transient colonization by *Staphylococcus aureus* in healthy Malaysian students. *Journal of Medical Microbiology*, *58*(11), 1531–1532. <http://doi.org/10.1099/jmm.0.011692-0>
- Vitali, L. A., Petrelli, D., Lamikanra, A., Prenna, M., & Akinkunmi, E. O. (2014). Diversity of antibiotic resistance genes and staphylococcal cassette chromosome *mec* elements in faecal isolates of coagulase-negative staphylococci from Nigeria. *BMC Microbiology*, *14*(106), 1–8. <http://doi.org/10.1186/1471-2180-14-106>
- Von Eiff, C., Heilmann, C., Herrmann, M., & Peters, G. (1999). Basic aspects of the pathogenesis of staphylococcal polymer-associated infections. *Infection*, *27*(1), 7–10. <http://doi.org/10.1007/BF02561610>
- Vuong, C., & Otto, M. (2002). *Staphylococcus epidermidis* infections. *Microbes and Infection*, *4*, 481–489.
- Wang, R., Khan, B. a., Cheung, G. Y. C., Bach, T. H. L., Jameson-Lee, M., Kong, K. F., ... Otto, M. (2011). *Staphylococcus epidermidis* surfactant peptides promote biofilm maturation and dissemination of biofilm-associated infection in mice. *Journal of Clinical Investigation*, *121*(1), 238–248. <http://doi.org/10.1172/JCI42520>
- Wang, X. M., Noble, L., Kreiswirth, B. N., Eisner, W., McClements, W., Jansen, K. U., & Anderson, A. S. (2003). Evaluation of a multilocus sequence typing system for *Staphylococcus epidermidis*. *Journal of Medical Microbiology*, *52*(Pt 11), 989–998. <http://doi.org/10.1099/jmm.0.05360-0>
- Wei, W., Cao, Z., Zhu, Y., Wang, X., Ding, G., Xu, H., ... Li, Y. (2006). Conserved genes in a path from commensalism to pathogenicity: Comparative phylogenetic profiles of *Staphylococcus epidermidis* RP62A

and ATCC12228. *BMC Genomics*, 12(112), 1–12.
<http://doi.org/10.1186/1471-2164-7-112>

- Weller, T. M. . (1999). The distribution of *mecA*, *mecR1* and *mecI* and sequence analysis of *mecI* and the *mec* promoter region in staphylococci expressing resistance to methicillin. *Journal of Antimicrobial Chemotherapy*, 43, 15–22.
- Widerström, M., Wiström, J., Ek, E., Edebro, H., & Monsen, T. (2011). Near absence of methicillin-resistance and pronounced genetic diversity among *Staphylococcus epidermidis* isolated from healthy persons in northern Sweden. *Apmis*, 119(8), 505–512. <http://doi.org/10.1111/j.1600-0463.2011.02757.x>
- Widerström, M., Wiström, J., Sjöstedt, a., & Monsen, T. (2012). Coagulase-negative staphylococci: Update on the molecular epidemiology and clinical presentation, with a focus on *Staphylococcus epidermidis* and *Staphylococcus saprophyticus*. *European Journal of Clinical Microbiology and Infectious Diseases*, 31(1), 7–20. <http://doi.org/10.1007/s10096-011-1270-6>
- Wieser, M., & Busse, H.-J. (2000). Rapid identification of *Staphylococcus epidermidis*. *International Journal of Systematic and Evolutionary Microbiology*, 50, 1087–1093.
- Wilson, L. A., & Sharp, P. M. (2006). Enterobacterial repetitive intergenic consensus (ERIC) sequences in *Escherichia coli*: Evolution and implications for ERIC-PCR. *Molecular Biology and Evolution*, 23(6), 1156–1168. <http://doi.org/10.1093/molbev/msj125>
- Wu, S. W., De Lencastre, H., & Tomasz, A. (2001). Recruitment of the *mecA* gene homologue of *Staphylococcus sciuri* into a resistance determinant and expression of the resistant phenotype in *Staphylococcus aureus*. *Journal of Bacteriology*, 183(8), 2417–2424. <http://doi.org/10.1128/JB.183.8.2417>
- Ye, Y., Jiang, Q., Wu, Q., Zhang, J., Lu, J., & Lin, L. (2012). The characterization and comparison of *Staphylococcus aureus* by antibiotic susceptibility testing, enterobacterial repetitive intergenic consensus–polymerase chain reaction, and random amplified polymorphic DNA–polymerase chain reaction. *Foodborne Pathogens and Disease*, 9(2), 168–171. <http://doi.org/10.1089/fpd.2011.0927>
- Zhang, Y., Agidi, S., & Lejeune, J. T. (2009). Diversity of staphylococcal cassette chromosome in coagulase-negative staphylococci from animal sources. *Journal of Applied Microbiology*, 107, 1375–1383. <http://doi.org/10.1111/j.1365-2672.2009.04322.x>

- Zhang, Y. Q., Ren, S. X., Li, H. L., Wang, Y. X., Fu, G., Yang, J., ... Wen, Y. M. (2003). Genome-based analysis of virulence genes in a non-biofilm-forming *Staphylococcus epidermidis* strain (ATCC 12228). *Moleucular Microbiology*, 49(6), 1577–1593. <http://doi.org/10.1046/j.1365-2958.2003.03671.x>
- Ziebuhr, W., Hennig, S., Eckart, M., Kranzler, H., Batzilla, C., & Kozitskaya, S. (2006). Nosocomial infections by *Staphylococcus epidermidis*: how a commensal bacterium turns into a pathogen. *International Journal of Antimicrobial Agents*, 28S, 14–20. <http://doi.org/10.1016/j.ijantimicag.2006.05.012>
- Zmantar, T., Kouidhi, B., Miladi, H., & Bakhrouf, A. (2011). Detection of macrolide and disinfectant resistance genes in clinical *Staphylococcus aureus* and coagulase-negative staphylococci. *BMC Research Notes*, 4(1), 453. <http://doi.org/10.1186/1756-0500-4-453>
- Zong, Z. (2013). Characterization of a complex context containing *mecA* but lacking genes encoding cassette chromosome recombinases in *Staphylococcus haemolyticus*. *BMC Microbiology*, 13(1), 64. <http://doi.org/10.1186/1471-2180-13-64>
- Zong, Z., Peng, C., & Lü, X. (2011). Diversity of SCC*mec* elements in methicillin-resistant coagulase-negative staphylococci clinical isolates. *PLoS ONE*, 6(5), 1–6. <http://doi.org/10.1371/journal.pone.0020191>