



UNIVERSITI PUTRA MALAYSIA

***EVALUATION OF BANANA AND PEAR FRUIT MATURITY STAGES
USING LASER BACKSCATTERING IMAGES, ARTIFICIAL NEURAL
NETWORK AND SUPPORT VECTOR MACHINE TECHNIQUES***

ADEBAYO SEGUN EMMANUEL

FK 2017 65



**EVALUATION OF BANANA AND PEAR FRUIT MATURITY STAGES
USING LASER BACKSCATTERING IMAGES, ARTIFICIAL NEURAL
NETWORK AND SUPPORT VECTOR MACHINE TECHNIQUES**

By

ADEBAYO SEGUN EMMANUEL

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

June 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This work is dedicated to my wife and children:

Adebayo Olanike Oluwabusayomi (Mrs), Iyanuoluwa Aduke and Oluwabunmi Ajike



© COPYRIGHT UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

EVALUATION OF BANANA AND PEAR FRUIT MATURITY STAGES USING LASER BACKSCATTERING IMAGES, ARTIFICIAL NEURAL NETWORK AND SUPPORT VECTOR MACHINE TECHNIQUES

By

ADEBAYO SEGUN EMMANUEL

June 2017

Chairman : Norhashila Hashim, PhD
Faculty : Engineering

Consumers considered ripeness of fruit as a very important factor in making choices of purchase. Ripeness in fruit generally affects their eating quality and market price. Quality attributes of fruit determined the extent of its acceptability and satisfaction by the consumers. Many quality attributes have been used for the determination of fruit quality, among them are colour, firmness and soluble solid contents (SSC). Majority of the techniques used to determine the maturity stages are destructive involving the removal of little quantity of fruit tissue especially for the measurement of SSC, total acidity and nutritional content. These techniques resulted in large amount of postharvest losses, inability to measure the whole batch and laborious. However, over the last decades several attempts have been made to develop optical techniques for monitoring quality indices of fruits via non-destructive approach. Laser light backscattering imaging (LLBI) system is one of the emerging optical techniques which is inexpensive and easy to use.

Bananas samples at six ripening stages i.e. from ripening stage 2 to 7 and pear samples at different days after full bloom (dafb) were obtained from a ripening facility at Potsdam Bornim and Sachsenobst orchard Germany respectively. The samples were kept at 14 °C with 79 to 89 % relative humidity (RH). Laser light backscattering imaging (LLBI) with 5 laser diodes wavelengths in the visible and near infrared region i.e. 532, 660, 785, 830 and 1060 nm were employed to acquire the backscattering images of the samples and features were extracted from the backscattering images of both fruit using transform-based textural techniques viz: Wavelet transform, Gabor transform, Tamura texture and optical properties i.e. absorption and reduced scattering coefficients with Farrell's diffusion theory. The reference measurements of index of chlorophyll, elasticity, firmness and SSC were measured with ΔA meter, texture analyzer, penetrometer and refractometer respectively immediately after backscattering images acquisition.

The extracted features and optical properties at individual and combined wavelengths were used as an input into the prediction and classification models in the predictions and classification of quality attributes and maturity stages of both fruit. Two computational intelligence techniques, artificial neural network (ANN) and support vector machines (SVM) were used to build the prediction and classification models. Root mean square error of calibration (RMSEC), root mean square error of cross-validation (RMSECV), coefficient of determination (R^2) and bias were used to evaluate the performance of the prediction models while overall classification accuracy was used to evaluate the classification models.

The results showed that there was a very strong correlation between the absorption and reduced scattering coefficients with ripening stage of banana and pear development. The range values of absorption coefficient and reduced scattering coefficient of bananas at 532 nm were between 0.312 and 0.963, and 2.637 and 4.893 for ripening stages 2 to 7 respectively. The result shown a decreasing trend of optical properties with increasing wavelength. For pear at 532 nm, the range of absorption coefficient was between 0.033 and 0.308 while reduced scattering coefficient was between 3.160 and 6.741. For banana, analysis using ANN with visible wavelength region of 532, 660 and 785nm resulted in high R^2 values ranging from 0.977 to 0.981 for the prediction of index of chlorophyll and 0.955 to 0.976 for elasticity; while near infrared region of 830 and 1060nm resulted in R^2 range between 0.964 and 0.980 for SSC prediction when absorption and reduced scattering coefficients at individual and combined wavelengths were used. For the classification of banana into ripening stages 2 to 7, visible wavelength region using ANN gave the highest classification accuracy of 98.77% with combined wavelengths while the highest overall classification accuracy of 96.30 % was recorded at 830nm with SVM. For pear the highest R^2 of 0.947 and 0.818 were obtained for firmness and SSC respectively using ANN while R^2 values of 0.890 and 0.808 for firmness and SSC using SVM. For pear classification into different maturity stages, the highest classification accuracy of 90.42 % was achieved with both ANN and SVM. Similar results though lower were obtained when transform-based textural techniques were used for the prediction of banana and pear quality parameters and classification of banana and pear into different ripening and maturity stages. This study has shown that transform-based textural techniques and optical properties of banana and pears with ANN and SVM as prediction and classification models can be employed to predict the quality parameters and classify banana and pears into different ripening and maturity stages non-destructively.

Abstrak tesis dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENILAIAN TAHAP KEMATANGAN BUAH PISANG DAN PEAR
MENGUNAKAN KAEDAH PENGIMEJAN PENGHAMBURAN LASER,
RANGKAIAN NEURAL BUATAN DAN MESIN SOKONGAN VEKTOR.**

Oleh

ADEBAYO SEGUN EMMANUEL

Jun 2017

Pengerusi : Norhashila Hashim, PhD
Fakulti : Kejuruteraan

Pengguna menganggap kematangan buah sebagai satu faktor yang sangat penting dalam membuat pilihan pembelian. Kematangan buah secara amnya memberi kesan kepada kualiti pemakanan dan harga pasaran. Ciri-ciri kualiti buah menentukan tahap penerimaan dan kepuasan pengguna. Pelbagai ciri-ciri kualiti telah digunakan bagi menentukan kualiti buah-buahan, antaranya ialah warna, ketegasan dan kandungan pepejal larut (SSC). Kebanyakan teknik-teknik yang digunakan untuk menentukan tahap kematangan adalah bersifat musnah melibatkan pembuangan sedikit kuantiti tisu buah terutama untuk mengukur SSC, jumlah keasidan dan kandungan nutrien. Teknik-teknik ini menghasilkan kerugian lepas-tuai dalam jumlah yang besar, ketidakbolehan mengukur seluruh kumpulan dan menggunakan tenaga buruh yang tinggi. Walau bagaimanapun, sejak beberapa dekad yang lalu, beberapa percubaan telah dijalankan untuk membangunkan teknik optik bagi pemantauan indeks kualiti buah-buahan melalui pendekatan tanpa-musnah. Sistem cahaya laser penghamburan adalah salah satu teknik optik yang muncul, yang mempunyai kos yang murah dan mudah untuk digunakan.

Sampel buah pisang pada enam tahap kematangan iaitu daripada tahap kematangan 2 hingga 7 dan sampel buah pear pada hari-selepas-mekar (dafb) yang berlainan telah diperolehi masing-masing daripada pusat pemeraman di Potsdam Bornim dan dusun Sachsenobst, Jerman. Sampel kajian telah disimpan pada suhu 14 °C dengan kelembapan bandingan 79 hingga 89% (RH). Cahaya laser pengimejan penghamburan (LLBI) dengan 5 gelombang diod laser dalam rantau nampak dan berhampiran inframerah iaitu 532, 660, 785, 830 dan 1060 nm telah digunakan untuk mendapatkan imej-imej penghamburan sampel dan ciri-ciri imej diekstrak daripada imej-imej tersebut menggunakan teknik berasaskan-ubahan tekstur iaitu: ubahan Wavelet, ubahan Gabor, tekstur Tamura dan sifat-sifat optik iaitu pekali penyerapan dan penyebaran-berkurang dengan teori penyebaran Farrell's. Pengukuran rujukan indeks klorofil, keanjalan, ketegasan dan SSC diukur masing-masing dengan ΔA

meter, penganalisa tekstur, *penetrometer* dan *refractometer* sebaik selepas pengukuran pengimejan penghamburan.

Ciri-ciri yang diekstrak dan sifat-sifat optik pada gelombang individu dan gabungan telah digunakan sebagai input kepada model-model ramalan dan pengelasan ciri-ciri kualiti dan tahap kematangan kedua-dua buah. Dua teknik kecerdasan pengkomputeran, teknik rangkaian neural buatan (ANN) dan mesin sokongan vektor (SVM) telah digunakan untuk membina model-model ramalan dan pengelasan. Punca kuasa dua ralat penentuan (RMSEC), punca kuasa dua ralat pengesahan-silang (RMSECV), pekali penentuan (R^2) dan kecenderungan telah digunakan untuk menilai prestasi model ramalan manakala ketepatan keseluruhan pengelasan digunakan untuk menilai model klasifikasi. Hasil kajian menunjukkan bahawa wujud hubungan yang sangat kuat antara pekali penyerapan dan penyebaran-berkurang dengan tahap kematangan buah pisang dan pembentukan buah pear.

Nilai julat pekali penyerapan dan penyebaran-berkurang buah pisang pada 532 nm adalah masing-masing antara 0.312 dan 0.963, dan 2.637 dan 4.893 bagi tahap kematangan 2 hingga 7. Keputusan menunjukkan gaya menurun oleh sifat optik dengan peningkatan gelombang. Bagi buah pear pada 532 nm, julat pekali penyerapan adalah antara 0.033 dan 0.308 manakala pekali penyebaran-berkurang adalah antara 3.160 dan 6.741. Bagi buah pisang, analisis menggunakan ANN dengan rantau gelombang nampak iaitu 532, 660 dan 785nm menghasilkan nilai R^2 yang tinggi antara 0.977 hingga 0.981 bagi ramalan indeks klorofil dan 0.955 hingga 0.976 bagi keanjalan; sementara gelombang berhampiran inframerah iaitu 830 dan 1060nm menghasilkan R^2 berjulat antara 0.964 dan 0.980 untuk ramalan SSC apabila pekali penyerapan dan penyebaran-berkurang pada gelombang individu dan gabungan digunakan. Bagi pengelasan buah pisang ke tahap kematangan 2 hingga 7, rantau gelombang nampak menggunakan ANN memberikan ketepatan klasifikasi tertinggi iaitu 98.77% dengan gelombang gabungan manakala ketepatan klasifikasi keseluruhan tertinggi sebanyak 96,30% dicatatkan pada 830nm dengan SVM. Bagi buah pear, nilai R^2 tertinggi masing-masing 0.947 dan 0.818 telah diperolehi untuk keanjalan dan SSC menggunakan ANN manakala nilai R^2 0.890 dan 0.808 diperolehi untuk keanjalan dan SSC menggunakan SVM. Bagi pengelasan buah pear ke tahap kematangan berbeza, ketepatan klasifikasi tertinggi sebanyak 90.42% telah dicapai dengan ANN dan SVM. Keputusan yang sama walaupun rendah diperolehi apabila teknik berasaskan-ubahan tekstur digunakan untuk ramalan parameter kualiti buah pisang dan pear dan pengelasan buah pisang dan pear ke tahap kemasakan dan kematangan berbeza. Kajian ini telah menunjukkan bahawa teknik berasaskan-ubahan tekstur dan ciri-ciri optik pisang dan buah pear dengan ANN dan SVM sebagai model ramalan dan pengelasan boleh digunakan untuk meramalkan parameter kualiti dan mengelaskan buah pisang dan pear ke tahap kemasakan dan kematangan yang berbeza secara tanpa musnah.

ACKNOWLEDGEMENTS

To God Almighty be glory, honour, power and adorations for making this journey a success.

To the Chairman and members of my supervisory committee Dr. Norhashila Hashim, Dr. Marsyita Hanafi and Assoc. Prof. Dr. Khalina Abdan I am grateful for all your unalloyed efforts and assistance throughout the duration of this programme. Thanks for bearing with my inadequacies.

Of very significant is the contribution of Professor Manuela Zude-Sasse and other members of staff of the Department of Horticultural Engineering of the Leibniz-Institut für Agrartechnik and Bioökonomie Potsdam-Bornim e.V (ATB) Germany for providing the opportunity for data collection and other technical input. Prof., I am grateful for granting to me five months of mentoring. Almighty God will bless your labour of love. I also wish to acknowledge the support and assistance received from Dr. Kaveh Mollazade of the Department of Biosystems Engineering, University of Kurdistan, Sanandaj, Iran.

Similarly, I am grateful to the Management of Federal University of Technology, Minna for the award of study fellowship to undertake this programme. Of note is the scholarship provided by Tertiary Education Trust fund (TETFUND) for the programme. Equally important is the support received from colleagues and members of staff of the Department of Agricultural and Bioresources Engineering, Federal University of Technology, Minna. In addition, I like to appreciate the spiritual supports from Pastors Noah Otene and Prince Dania. Thank you all and God bless you real good. I am equally grateful for unmeasurable support received from my extended family. Thanks for been there for me.

To my lovely daughters Iyanuoluwa Aduke and Oluwabunmi Ajike, thanks for bearing with my absence. Thinking about you always gave me inspiration and motivation to get down to work. To my wife, Olanike Oluwabusayomi, indeed you have proven time and time again to be reliable, dependable and supportive. Thanks for holding forth the home front while I am absent from home.

Thank you all!

I certify that a Thesis Examination Committee has met on 5 June 2017 to conduct the final examination of Adebayo Segun Emmanuel on his thesis entitled "Evaluation of Banana and Pear Fruit Maturity Stages using Laser Backscattering Images, Artificial Neural Network and Support Vector Machine Techniques" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Aimrun Wayayok, PhD
Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Abd. Rahman bin Ramli, PhD
Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Rosnah binti Shamsudin, PhD
Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Mahmoud Omid, PhD
Professor
University of Tehran
Iran
(External Examiner)



NOR AINI AB. SHUKOR, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 8 August 2017

This thesis was submitted to the senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Norhashila Hashim, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Marsyita Hanafi, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Khalina Abdan, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date :

Declaration by the student

I hereby confirm that:

- this thesis is my original work
- quotations, illustrations and citations have been duly referenced
- the thesis has not been submitted previously or concurrently for any other degree at any institutions
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be owned from supervisor and deputy vice –chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____

Date: _____

Name and Matric No: Adebayo, Segun Emmanuel, GS39837

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Dr. Norhashila Hashim

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Marsyita Hanafi

Signature: _____
Name of Member
of Supervisory
Committee: Associate Professor Dr. Khalina Abdan

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xix
CHAPTER	1
1 INTRODUCTION	1
1.1 Background of study	1
1.2 Statement of problem	1
1.3 Objectives of the study	2
1.4 Scope and limitation of the study	3
1.5 Significance of the study	3
2 LITERATURE REVIEW	5
2.1 Physiological background of fruit maturity and ripening	5
2.2 Fruit quality	6
2.2.1 External appearance	6
2.2.2 Chemical changes	7
2.2.3 Mechanical Properties	9
2.2.3.1 Fruit firmness	9
2.2.3.2 Elasticity or elastic modulus	10
2.3 Optical properties of fruit	10
2.3.1 Electromagnetic Radiation	12
2.3.2 Light interaction in biological materials	14
2.4 Non-destructive optical techniques for monitoring fruit maturity and ripening	16
2.4.1 Optical methods for analyzing fruit quality	16
2.4.2 Sensor system for backscattering imaging in the visible and shortwave near infrared wavelength range	18
2.4.3 Optical geometry	19
2.4.4 Light backscattering imaging concept	21
2.5 Backscattering Image Analysis	22
2.5.1 Image pre-processing	22
2.5.2 Image segmentation	23
2.5.3 Region of interest	25
2.5.4 Prediction and classification	25
2.6 Extraction of optical properties	26
2.7 Summary of the review	27

3	MATERIALS AND METHODS	29
3.1	Study flowchart	29
3.2	Sample collection and preparation	30
3.3	Data collection	31
3.3.1	Reference Measurements	31
3.3.2	Backscattering Imaging	32
3.3.3	Features extraction	33
3.4	Data analysis	40
3.5	Prediction and classification models	41
3.5.1	Artificial neural network	41
3.5.1.1	Types of activation function	44
3.5.1.2	Multilayer perceptron (MLP)	45
3.5.1.3	Back-propagation Algorithm	46
3.5.2	Support vector machine	50
3.5.3	Performance evaluation of the classifiers and prediction Models	54
3.5.3.1	Correlation coefficient	55
3.5.3.2	Specificity	55
3.5.3.3	Sensitivity	55
3.5.3.4	Accuracy	55
3.5.3.5	Other performance criteria for prediction models	56
4	RESULTS AND DISCUSSION	57
4.1	Reference measurements data	57
4.1.1	Index of chlorophyll, elasticity and soluble solid content (SSC) of banana	57
4.1.2	Firmness and soluble solid content (SSC) of pear	61
4.2	Backscattered images of banana and pear	64
4.3	Prediction of quality parameters of banana and classification into maturity stages	65
4.3.1	Farrell optical properties of bananas	66
4.3.2	Tamura features of bananas	75
4.3.3	Gabor features of bananas	83
4.3.4	Wavelets features of bananas	91
4.4	Prediction of quality parameters of pear and classification into maturity stages	99
4.4.1	Farrell optical properties of pears	99
4.4.2	Tamura features of pears	107
4.4.3	Gabor features of pears	115
4.4.4	Wavelets features of pears	122
4.5	Comparison of the capability of transform-based processing techniques and diffusion theory for prediction and classification of quality attributes of banana and pear fruits	129
5	CONCLUSIONS AND RECOMMENDATIONS	131
5.1	Conclusion	131
5.2	Recommendation	132

REFERENCES	133
APPENDICES	151
BIODATA OF STUDENT	171
LIST OF PUBLICATIONS	172



© COPYRIGHT UPM

LIST OF TABLES

Table		Page
2.1	External appearance of colour for fruit maturity determination	7
4.1	Descriptive statistics of banana reference data and ripening stage	60
4.2	Correlation of reference measurements and the ripening stage of banana	60
4.3	Descriptive statistics of Pear reference measurements and maturity	63
4.4	Correlation of reference measurements and pear development	63
4.5	Optical properties of banana from Farrell diffusion theory	66
4.6	Correlation of Optical properties of bananas from Farrell diffusion theory and reference measurements with ripening stages at different wavelengths	68
4.7	Correlation of optical properties of bananas extracted with laser wavelength used	69
4.8	Performance evaluation of the prediction models using μ_a and μ_s' coefficients of bananas from Farrell diffusion theory at individual wavelength	70
4.9	Performance evaluation of the prediction models using μ_a and μ_s' coefficients of bananas from Farrell diffusion theory at combined wavelengths	71
4.10	Overall classification accuracy using μ_a and μ_s' coefficients of bananas from Farrell diffusion theory at individual and combined wavelengths	75
4.11	The mean values and standard error of Tamura features extracted from the backscattering images of banana at different ripening stages	76
4.12	Correlation of Tamura features of bananas and ripening stage	77
4.13	Performance evaluation of the prediction models using Tamura features of bananas at individual wavelength	78
4.14	Performance evaluation of the prediction models using Tamura features of bananas at combined wavelengths	79
4.15	Overall classification Accuracy using Tamura features of bananas at individual and combined wavelengths	83

4.16	The mean values and standard errors of Gabor features of bananas at 0.1768 Hz and 0 radian	84
4.17	Correlation of Gabor features of bananas with ripening stages	85
4.18	Performance evaluation of the prediction models using Gabor features of bananas at individual wavelength	86
4.19	Performance evaluation of the prediction models using Gabor features of bananas with combined wavelengths	87
4.20	Overall classification accuracy using Gabor features of bananas at individual and combined wavelengths	91
4.21	The mean and standard error of Wavelets features of bananas at first level of decomposition and horizontal coefficient	92
4.22	Correlation of Wavelets features of banana with ripening stage	93
4.23	Performance evaluation of the prediction models using Wavelets features of bananas at individual wavelength	94
4.24	Performance evaluation of the prediction models using Wavelet features of bananas with combined wavelengths	95
4.25	Overall classification accuracy of banana ripening stages using Wavelets features of bananas at individual and combined wavelengths	99
4.26	Optical properties of pears from Farrell diffusion theory	100
4.27	Correlation of μ_a and μ_s' coefficients of pears from Farrell diffusion theory and reference measurements with days after full bloom at different wavelengths	101
4.28	Correlation of μ_a and μ_s' coefficients of pears extracted from Farrell diffusion theory with laser wavelength used	102
4.29	Performance evaluation of the prediction models using μ_a and μ_s' coefficients of pears from Farrell diffusion	103
4.30	Performance evaluation of the prediction models using μ_a and μ_s' coefficients of pears at combined wavelengths	104
4.31	Overall classification accuracy using μ_a and μ_s' coefficients of pears from Farrell diffusion theory at individual and combined wavelengths	107
4.32	The mean values and standard error of Tamura features extracted from the backscattering images of pear at different days after full bloom	108

4.33	Correlation of Tamura features of pears and reference measurements with time (days after full bloom) at different wavelengths	110
4.34	Performance evaluation of the prediction models using Tamura features of pears at individual wavelength	111
4.35	Performance evaluation of the prediction models using Tamura features of pear at combined wavelengths	112
4.36	Overall classification accuracy using Tamura features of pears at individual and combined wavelengths	115
4.37	The mean values and standard error of Gabor features of pears at 0.1768 Hz and 0 radian	116
4.38	Correlation of Gabor features of pears with pear development	117
4.39	Performance evaluation of the prediction models using Gabor features of pear at individual wavelength	118
4.40	Performance evaluation of the prediction models using Gabor features of pears with combined wavelengths	119
4.41	Overall classification accuracy using Gabor features of pears at individual and combined wavelengths	122
4.42	Wavelets features of pears at first level of decomposition and horizontal coefficient	123
4.43	Correlation of Wavelet features of pears with pear maturity	124
4.44	Performance evaluation of the prediction models using Wavelets features of pears at individual wavelength	125
4.45	Performance evaluation of the prediction models using Wavelets features of pears with combined wavelengths	126
4.46	Overall classification accuracy using Wavelets features of pears at individual and combined wavelengths	129
4.47	Comparison of overall classification accuracy of the feature extraction Methods for banana and pear maturity classification	129
4.48	Comparison of feature extraction methods for banana and pear quality parameters predictions (coefficient of determination)	130

LIST OF FIGURES

Figure	Page
2.1 Electromagnetic spectrum	13
2.2 Incident light on a fruit or vegetable results in specular reflectance (gloss), diffuse reflectance from features at depths to about 5 mm (body reflectance or interreflectance), diffuse transmittance, or absorbance	14
2.3 Three approaches to generate spectral image	20
2.4 Backscattered images of banana and pear	22
3.1 Flowchart of Research work	29
3.2 Different ripening stages of banana	30
3.3 Pear Samples used for the experiment	31
3.4 (a) Laser light Backscattering Imaging System (b) Schematic diagram of LLBI a-the CCD Camera b-the laser diode arrangement c-the fruit platform d-the laptop.	33
3.5 Implementation of Gabor features for prediction and classification	34
3.6 Implementation of Wavelets features for prediction and classification	35
3.7 Implementation of Tamura features for prediction and classification	39
3.8 Implementation of optical properties for prediction and classification	40
3.9 Nonlinear model of a neuron	43
4.1 Reference data and banana ripening stage (a) Index of Chlorophyll (b) Elasticity (c) SSC with ripening stages of banana. The error bars and Tukey's test are shown. Same letter shows no significant difference ($p \leq .05$)	59
4.2 Reference data and pear development (a) SSC (b) Firmness with pear development. The error bars and Tukey's test are shown. Same letter shows no significant difference ($p \leq .05$)	62
4.3 Backscattering images of banana at different wavelengths and ripening stage	64

4.4	Pear backscattering image at different wavelength and days after full bloom	65
4.5	Index of chlorophyll prediction at different wavelength with μ_a and μ_s' coefficients of bananas from Farrell diffusion theory (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	72
4.6	Elasticity prediction at different wavelength with μ_a and μ_s' coefficients of bananas from Farrell diffusion theory (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	73
4.7	SSC prediction at different wavelength with μ_a and μ_s' coefficients of bananas from Farrell diffusion theory (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	74
4.8	Elasticity prediction at different wavelength with Tamura features of bananas (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	80
4.9	Index of chlorophyll prediction at different wavelength with Tamura features of bananas (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	81
4.10	SSC prediction at different wavelength with Tamura features of bananas (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	82
4.11	Elasticity prediction at different wavelength with Gabor features of bananas (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	88
4.12	Index of chlorophyll prediction at different wavelength with Gabor features of bananas (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	89
4.13	SSC prediction at different wavelength with Gabor features of bananas (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	90
4.14	Elasticity prediction at different wavelength with Wavelets features of bananas (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	96
4.15	Index of chlorophyll prediction at different wavelength with Wavelets features of bananas (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	97
4.16	SSC prediction at different wavelength with Wavelets features of bananas (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	98

4.17	SSC prediction at different wavelength with μa and $\mu s'$ coefficients of pears from Farrell diffusion theory (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	105
4.18	Firmness prediction at different wavelength with μa and $\mu s'$ coefficients of pears from Farrell diffusion (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	106
4.19	SSC prediction at different wavelength with Tamura features of pears (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	113
4.20	Firmness prediction at different wavelength with Tamura features of pears (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	114
4.21	Firmness prediction at different wavelength with Gabor features of pears (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	120
4.22	Firmness prediction at different wavelength with Gabor features of pears (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	121
4.23	Firmness prediction at different wavelength with Wavelets features of pears (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	127
4.24	SSC prediction at different wavelength with Wavelets features of pears (a) 532 nm (b) 660 nm (c) 785 nm (d) 830 nm (e) 1060 nm using ANN model	128

LIST OF ABBREVIATIONS

ABR	Acid-Brix Ratio
ANN	Artificial neural network
AOTF	Acousto-optic tunable filter
AU	Auxilliary unit
BIL	Band-interleaved-by-line
CCD	Charge-coupled device
CMOS	Complementary metal oxide semiconductor
CMY	Cyan, magenta and yellow
CT	Computed Tomography
DA	Discriminant analysis
DAFB	Days after full bloom
DM	Dry Matter
EM	Electromagnetic spectrum
EN	Electronic Nose
FF	Flesh firmness
FN	False negative
FP	False positive
GLCM	Gray level co-occurrence matrix
GLRLM	Gray level run length matrix
GUI	Graphical user interface
HSI	Hue, saturation and intensity
LBP	Local binary pattern
LED	Laser emission diode
LCTF	Liquid crystal tunable filter
LDA	Linear discriminant analysis
LLBI	Laser Light Backscattering Imaging
LR	Linear regression

LS-SVM	Least square support vector machine
MC	Monte Carlo
MLR	Multiple linear regression
MLP	Multilayer perceptron
MSI	Multispectral Imaging
MRI	Magnetic resonance imaging
NB	Naïve bayes
NIR	Near Infrared
NIRS	Near infrared reflectance spectroscopy
NMR	Nuclear Magnetic Resonance
QDA	Quadratic discriminant analysis
RBF	Radial-basis function
RGB	Red, green, blue
RMSE	Root mean square error
RP	Ripening stage
SIMCA	Soft independence modelling of class analogy
SSC	Soluble solids content
SVM	Support vector machine
SVLA	Support vector learning algorithm
SWNIR	Short wave near infrared
TA	Titrateable acidity
TN	True negative
TP	True positive
TSS	Total soluble solids
UV	Ultra Violet
VC	Vapnik-Chervonenkis
μ_a	Absorption coefficient
μ'_s	Reduced scattering coefficient

CHAPTER 1

INTRODUCTION

1.1 Background of study

The quality, shelf life, and storage are principles that determine the time which fruit and vegetables are harvested. According to Barbosa-Cánovas (2003), there are three distinct stages in the life span of fruit and vegetable and these are maturation, ripening, and senescence. At maturity, the fruit is fully developed to be harvested. This is due to an edible portion of the fruit that has been completely developed in size, though it may not be available for consumption immediately. The stage following or sometimes overlapping with maturation is ripening. At this stage, the fruit has undergone biochemical processes marked by taste and aroma and it is ready for immediate consumption. The final stage is senescence, where this stage is marked by an essential decline in the quality parameters such as texture, flavour, etc of the fruit and vegetable. Senescence is often described as the death of fruit tissue.

Fresh fruit quality has been defined as the sum total of attributes, properties, or characteristics that enhance or attract them for consumption (Achilleas and Anastasios, 2008; Giusti et al., 2008; Schreiner et al., 2013). For instance, the concept of fruit quality changes along the supply chain. To producers or farmers, quality is defined in terms of high yield, ease of harvest, and long transportation without/with minimal damage, storability, ratio of soluble solids content (SSC) to acidity, as well as attractive appearance (Narrod et al., 2009; Van der Vorst et al., 2009). Subsequently to the wholesalers and retail marketers, fruit quality is defined in terms of good appearance, firmness, and extended shelf life, while the consumer of fruit looks at quality from the perspective of freshness, firmness, flavour attributes, nutritional quality, and appearance in terms of absence of defects or decay (Punan et al., 2000; Zúñiga-Arias, 2007).

1.2 Statement of problem

Discrimination of fruit into maturity stages with deployment of non-destructive assessment is critical because it assist in proper grading of fruit for handling and storage. As such various non-destructive techniques such as multispectral, hyperspectral, spectrophotometers as well as machine vision systems have been deployed for variety of fruit.

In a number of researches carried out on the non-destructive fruit quality assessments, light intensity-based features in the space domain has been utilized to build calibration models between the acquired images from the various systems and the reference measurements to be studied. Features such as the area of the captured image regions (Qing et al., 2007), radial averaging (Lu, 2004) wherein the photon scattering region is partitioned into several circular rings and the average value of the pixels within the

ring is used as features to determine the quality of interest. Employing the radial averaging, various mathematical functions such as Lorentzian, Modified Lorentzian and Gompertz functions were fit into one-dimensional scattering profile as a function of distance. Other approaches used to process backscattering images is using one dimensional features of pixel brightness using colour models such as RGB, HSI, $L^*a^*b^*$ and others and using the colour features to assess or predict fruit quality of interest.

Consequent utilizing the one dimensional features focus solely on the pixel values neglecting pixel intensity pattern or location from the overall two-dimensional captured scattering images. Therefore, employing the two-dimensional scattering images i.e using the pixel values and location could help in improvement of the performance of backscattering images processing to study or assess the quality parameters of fruit and discriminating them into maturity stages.

Image texture is a two-dimensional processing approach which takes into account the pixels values and locations (Zheng et al., 2006). Though, a number of image texture-based analysis techniques has been used in image processing problems, majority has been done individually with different fruits, very few studies has reported comparative studies of these techniques. Therefore, this study investigates the comparative study of transform-based textural techniques to analyze backscattering images of banana and pear for prediction of the fruit quality parameters and discrimination into maturity stages.

Furthermore, processing of backscattering images using light propagation in tissues by extracting absorption and reduced scattering coefficients otherwise known as optical properties which is guided by the radiative transport theory has been employed sparingly. Few studies has reported the use of optical properties extracted from backscattering images for assessment of quality parameters of fruit and discrimination into maturity stages. Therefore, this study investigate the use of optical properties of banana and pear to assess/predict their quality parameters and discriminate them into maturity stages.

1.3 Objectives of the study

The main objective of this study is to evaluate quality parameters of banana and pear fruit related to the development and ripening by means of backscattering imaging coupled with computational intelligence techniques. The specific objectives are:

1. To evaluate banana and pear quality parameters using backscattering imaging and standard reference measurements i.e. firmness, elasticity, SSC, and index of chlorophyll.
2. To analyse the capability of transform-based processing techniques (wavelet, Gabor and Tamura) and diffusion theory (Farrell) features extraction methods for prediction of quality attributes of banana (index of chlorophyll, elasticity and SSC) and pear (Firmness and SSC) and classification of the fruit into maturity stages.

3. To develop intelligent models for bananas and pears quality parameters prediction and classification into maturity stages by artificial neural network and support vector machine.

1.4 Scope and limitation of the study

This study focuses on the evaluation of quality parameters of two climacteric fruits which are banana (Cavendish) and pear (*Pyrus communis* 'Conference'). The optical technique employed is LLBI system with five laser diodes 532, 660, 785, 830, and 1060 nm to capture the images of the fruit. Measurements were carried out on samples of banana at six ripening stages and on pears at different days after full bloom once a week for eight weeks. The quality parameters of interest were elasticity, firmness, index of chlorophyll and SSC and these were determined destructively using standard methods.

Optical properties and textural features were extracted from the backscattered images of both fruit using Farrell diffusion approximation and transform-based techniques (Wavelet, Gabor and Tamura). Labview and Matlab were used to develop a program to extract optical properties and textural features respectively.

The optical properties and textural features serve as input for neural networks and support vector machines models to predict the quality parameters of banana and pear, and also to classify banana into six ripening stages. Statistica 12, WEKA and Unscrambler x were used in the data analysis. This work is limited to two climacteric fruit that are banana and pear and one variety each for the two fruit, with three quality parameters for banana and two for pear. Pear sample collection is from one orchard.

1.5 Significance of the study.

Feasible applications of the method include packinghouse sorting of bananas into different ripening stages for effective postharvest handling to prolong shelf-life and pears for similar postharvest handling and effective sorting into ripe and unripe pears and banana. Using this technique banana and pears may be sorted into six ripening stages and pears into two classes of ripe and unripe fruits. The technique may also potentially be extended to other climacteric fruits.

The industrial relevance of the method presented in this research is immediate and high. The methods are simple to implement therefore its applicability in the industry can be undertaken with minimal expertise. Likewise, the system set-up is less cumbersome so it can be easily assembled for use in the industry. Due to low processing time per object with laser backscattering imaging system, time for image processing will be greatly reduced thereby enhancing the sorting and grading of fruit in the industry.

Since the techniques are non-destructive, samples which usually would have been destroyed with destructive technique will be saved thereby reducing waste and subsequently maximum numbers of samples can be assessed to have a fair representation of the entire lots. Less man power will be required as the technique can be automated thereby few hands will be needed to run the system with good accuracy for maturity stages discrimination of fruit.



REFERENCES

- Abubaker, H.M., Tománek, P. and Grmela, L., 2011. Measurement of dynamic variations of polarized light in processed meat due to aging, SPIE Optics+ Optoelectronics. International Society for Optics and Photonics, pp. 80730U-80730U-6.
- Achilleas, K. and Anastasios, S., 2008. Marketing aspects of quality assurance systems: The organic food sector case. *British Food Journal*, 110(8): 829-839.
- Agati, G., Meyer, S., Matteini, P. and Cerovic, Z.G., 2007. Assessment of anthocyanins in grape (*Vitis vinifera* L.) berries using a noninvasive chlorophyll fluorescence method. *Journal of Agricultural and Food Chemistry*, 55(4): 1053-1061.
- Alexander, L. and Grierson, D., 2002. Ethylene biosynthesis and action in tomato: a model for climacteric fruit ripening. *Journal of Experimental Botany*, 53(377): 2039-2055.
- Ali, M.M., Hashim, N., Bejo, S.K. and Shamsudin, R., 2017. Quality evaluation of watermelon using laser-induced backscattering imaging during storage. *Postharvest Biology and Technology*, 123: 51-59.
- Amann, M.-C., Bosch, T., Lescure, M., Myllyla, R. and Rioux, M., 2001. Laser ranging: a critical review of usual techniques for distance measurement. *Optical Engineering*, 40(1): 10-19.
- Arridge, S., Schweiger, M., Hiraoka, M. and Delpy, D., 1993. A finite element approach for modeling photon transport in tissue. *Medical Physics*, 20(2): 299-309.
- ASAE, A., 1997. Standards—Standards, Engineering Practices, Data. American Society of Agricultural Engineers, St. Joseph, MI, 556, pp 571-575.
- Ayub, R., Gris, M., Amor, M.B., Gillot, L., Roustan, J-P., Latché, A., Bouzayen, M. and Pech, J-C., 1996. Expression of ACC oxidase antisense gene inhibits ripening of cantaloupe melon fruits. *Nature Biotechnology*, 14(7): 862-866.
- Babar, M.A., Reynolds, M.P., Van Ginkel, M., Klatt, A.R., Raun, W.R. and Stone, M.L., 2006. Spectral reflectance to estimate genetic variation for in-season biomass, leaf chlorophyll, and canopy temperature in wheat. *Crop Science*, 46(3): 1046-1057.
- Bain, J.M. and Robertson, R., 1951. The physiology of growth in apple fruits I. Cell size, cell number, and fruit development. *Australian Journal of Biological Sciences*, 4(2): 75-91.
- Bajcsy, R., 1973. Computer description of textured surfaces, Proceedings of the 3rd international joint conference on Artificial intelligence. Morgan Kaufmann

Publishers Inc., pp. 572-579.

- Balibrea, M.E., Martínez-Andújar, C., Cuartero, J., Bolarín, M.C. and Pérez-Alfocea, F., 2006. The high fruit soluble sugar content in wild *Lycopersicon* species and their hybrids with cultivars depends on sucrose import during ripening rather than on sucrose metabolism. *Functional Plant Biology*, 33(3): 279-288.
- Baranyai, L. and Zude, M., 2009. Analysis of laser light propagation in kiwifruit using backscattering imaging and Monte Carlo simulation. *Computers and Electronics in Agriculture*, 69(1): 33-39.
- Barbosa-Cánovas, G.V., 2003. Handling and preservation of fruits and vegetables by combined methods for rural areas: Technical manual, 149. Food & Agriculture Org, pp 508-520.
- Bartley, I.M. and Knee, M., 1982. The chemistry of textural changes in fruit during storage. *Food Chemistry*, 9(1): 47-58.
- Bashir, H.A. and Abu-Goukh, A.-B.A., 2003. Compositional changes during guava fruit ripening. *Food Chemistry*, 80(4): 557-563.
- Beckles, D.M., 2012. Factors affecting the postharvest soluble solids and sugar content of tomato (*Solanum lycopersicum* L.) fruit. *Postharvest Biology and Technology*, 63(1): 129-140.
- Biale, J.B., 1964. Growth, maturation, and senescence in fruits. *Science*, 146: 880-888.
- Biale, J.B., Young, R.E. and Olmstead, A.J., 1954. Fruit respiration and ethylene production. *Plant physiology*, 29(2): 168-175.
- Birth, G., 1976a. How light interacts with foods. In "Quality Detection in Foods" Ed. Gaffney, American Society Agricultural Engineers St. Joseph, MI: 6, pp 20-31.
- Birth, G.S., 1976b. How light interacts with foods. In "Quality Detection in Foods" Ed. Gaffney, Journal of American Society Agricultural Engineers St. Joseph, MI: 7, pp 45-54.
- Blasco, J., Aleixos, N. and Moltó, E., 2003. Machine vision system for automatic quality grading of fruit. *Biosystems Engineering*, 85(4): 415-423.
- Blasco, J., Aleixos, N. and Moltó, E., 2007. Computer vision detection of peel defects in citrus by means of a region oriented segmentation algorithm. *Journal of Food Engineering*, 81(3): 535-543.
- Brady, C., 1987. Fruit ripening. *Annual Review of Plant Physiology*, 38(1): 155-178.
- Brummell, D.A., 2006. Cell wall disassembly in ripening fruit. *Functional Plant Biology*, 33(2): 103-119.

- Brummell, D.A. and Harpster, M.H., 2001. Cell wall metabolism in fruit softening and quality and its manipulation in transgenic plants, *Plant Cell Walls*. Springer, pp. 311-340.
- Bureau, S., Ruiz, D., Reich, M., Gouble, B., Bertrand, D., Audergon, J-M. and Renard, C. MGC., 2009. Rapid and non-destructive analysis of apricot fruit quality using FT-near-infrared spectroscopy. *Food Chemistry*, 113(4): 1323-1328.
- Bushberg, J.T. and Boone, J.M., 2011. *The essential physics of medical imaging*. Lippincott Williams & Wilkins, pp 40-52.
- Butz, P., Hofmann, C. and Tauscher, B., 2005. Recent developments in noninvasive techniques for fresh fruit and vegetable internal quality analysis. *Journal of Food Science*, 70(9): 131-141.
- Bylesjö, M., Rantalainen, M., Cloarec, O., Nicholson, J.K., Holmes, E. and Trygg, J., 2006. OPLS discriminant analysis: combining the strengths of PLS-DA and SIMCA classification. *Journal of Chemometrics*, 20(8-10): 341-351.
- Camps, C. and Christen, D., 2009. Non-destructive assessment of apricot fruit quality by portable visible-near infrared spectroscopy. *LWT-Food Science and Technology*, 42(6): 1125-1131.
- Cao, F., Wu, D. and He, Y., 2010. Soluble solids content and pH prediction and varieties discrimination of grapes based on visible–near infrared spectroscopy. *Computers and Electronics in Agriculture*, 71: S15-S18.
- Carlini, P., Massantini, R. and Mencarelli, F., 2000. Vis-NIR measurement of soluble solids in cherry and apricot by PLS regression and wavelength selection. *Journal of Agricultural and Food Chemistry*, 48(11): 5236-5242.
- Chang, C.-L. and Chang, K.-P., 2014. The growth response of leaf lettuce at different stages to multiple wavelength-band light-emitting diode lighting. *Scientia Horticulturae*, 179: 78-84.
- Chang, T. and Kuo, C.-C., 1993. Texture analysis and classification with tree-structured wavelet transform. *Image Processing, IEEE Transactions*, 2(4): 429-441.
- Chapman, S.J., 2015. *MATLAB programming for engineers*. Nelson Education.
- Chen, Y.-R., Chao, K. and Kim, M.S., 2002. Machine vision technology for agricultural applications. *Computers and electronics in Agriculture*, 36(2): 173-191.
- Cheniclet, C., Rong, W.Y., Causse, M., Frangne, N., Bolling, L., Carde, J-P. and Renaudin, J-P., 2005. Cell expansion and endoreduplication show a large genetic variability in pericarp and contribute strongly to tomato fruit growth. *Plant Physiology*, 139(4): 1984-1994.

- Chin, L.-H., Ali, Z.M. and Lazan, H., 1999. Cell wall modifications, degrading enzymes and softening of carambola fruit during ripening. *Journal of Experimental Botany*, 50(335): 767-775.
- Choi, C., Lee, K. and Park, B., 1997. Prediction of soluble solid and firmness in apple by visible/near-infrared spectroscopy. *Journal of the Korean Society for Agricultural Machinery (Korea Republic)*, 8: 23-35.
- Choi, K., Lee, G., Han, Y. and Bunn, J., 1995. Tomato maturity evaluation using color image analysis. *Transactions of the ASAE*, 38(1): 171-176.
- Chu, K., 1999. An introduction to sensitivity, specificity, predictive values and likelihood ratios. *Emergency Medicine*, 11(3): 175-181.
- Cluff, K., Naganathan, G.K., Subbiah, J., Lu, R., Calkins, C.R. and Samal, A., 2008. Optical scattering in beef steak to predict tenderness using hyperspectral imaging in the VIS-NIR region. *Sensing and Instrumentation for Food Quality and Safety*, 2(3): 189-196.
- Commission, J.F.W.C.A., 1994. *Codex Alimentarius: Processed and Quick Frozen Fruits and Vegetables*. Food & Agriculture Org.
- CooMbe, B.G. and McCarthy, M., 2000. Dynamics of grape berry growth and physiology of ripening. *Australian Journal of Grape and Wine Research*, 6(2): 131-135.
- Cox, K.A., McGhie, T.K., White, A. and Woolf, A.B., 2004. Skin colour and pigment changes during ripening of 'Hass' avocado fruit. *Postharvest Biology and Technology*, 31(3): 287-294.
- Cozzolino, D., Esler, M.B., Damberg, R.G., Cynkar, W.U., Boehm, D.R., Francis, I.L. and Gishen, M., 2004. Prediction of colour and pH in grapes using a diode array spectrophotometer (400-1100 nm). *Journal of Near Infrared Spectroscopy*, 12: 105-112.
- Crisosto, C.H., Day, K.R., Crisosto, G.M. and Garner, D., 2001. Quality attributes of white flesh peaches and nectarines grown under California conditions. *Journal-American Pomological Society*, 55: 45-51.
- Cubeddu, R., D'Andrea, C., Pifferi, A., Taroni, P., Torricelli, A., Valentini, G., Ruiz-Altisent, M., Valero, C., Ortiz, C. and Dover, C., 2001. Time-resolved reflectance spectroscopy applied to the nondestructive monitoring of the internal optical properties in apples. *Applied Spectroscopy*, 55(10): 1368-1374.
- Cubeddu, R., Pifferi, A., Taroni, P. and Torricelli, A., 2002. New perspectives for quality assessment: time resolved optical methods. *Fruit and Vegetable Processing: Maximizing Quality*. CRC Press–Woodhead Publishing, Cambridge, UK: 150-169.

- Cubero, S., Aleixos, N., Moltó, E., Gómez-Sanchis, J. and Blasco, J., 2011. Advances in machine vision applications for automatic inspection and quality evaluation of fruits and vegetables. *Food and Bioprocess Technology*, 4(4): 487-504.
- Daubechies, I., 1988. Orthonormal bases of compactly supported wavelets. *Communications on Pure and Applied Mathematics*, 41(7): 909-996.
- Deikman, J., 1997. Molecular mechanisms of ethylene regulation of gene transcription. *Physiologia Plantarum*, 100(3): 561-566.
- Deulin, X. and L'Huillier, J., 2006. Finite element approach to photon propagation modeling in semi-infinite homogeneous and multilayered tissue structures. *The European Physical Journal Applied Physics*, 33(02): 133-146.
- Du, C.-J. and Sun, D.-W., 2004. Recent developments in the applications of image processing techniques for food quality evaluation. *Trends in Food Science & Technology*, 15(5): 230-249.
- Elmasry, G., Kamruzzaman, M., Sun, D.-W. and Allen, P., 2012a. Principles and applications of hyperspectral imaging in quality evaluation of agro-food products: a review. *Critical Reviews in Food Science and Nutrition*, 52(11): 999-1023.
- ElMasry, G. and Sun, D.-W., 2010. Principles of hyperspectral imaging technology. *Hyperspectral Imaging for Food Quality Analysis and Control*: pp 3-43.
- ElMasry, G., Sun, D.-W. and Allen, P., 2012b. Near-infrared hyperspectral imaging for predicting colour, pH and tenderness of fresh beef. *Journal of Food Engineering*, 110(1): 127-140.
- ElMasry, G., Wang, N., ElSayed, A. and Ngadi, M., 2007. Hyperspectral imaging for nondestructive determination of some quality attributes for strawberry. *Journal of Food Engineering*, 81(1): 98-107.
- ElMasry, G., Wang, N. and Vigneault, C., 2009. Detecting chilling injury in Red Delicious apple using hyperspectral imaging and neural networks. *Postharvest Biology and Technology*, 52(1): 1-8.
- ElMasry, G., Wang, N., Vigneault, C., Qiao, J. and ElSayed, A., 2008. Early detection of apple bruises on different background colors using hyperspectral imaging. *LWT-Food Science and Technology*, 41(2): 337-345.
- Everingham, M., Van Gool, L., Williams, C.K., Winn, J. and Zisserman, A., 2010. The pascal visual object classes (voc) challenge. *International Journal of Computer Vision*, 88(2): 303-338.
- Fang, Z.-H., Fu, X.-P. and He, X.-M., 2015. Investigation of absorption and scattering characteristics of kiwifruit tissue using a single integrating sphere system. *Journal of Zhejiang University*, 4(1): 98-107.
- Farrell, T.J., Patterson, M.S. and Wilson, B., 1992. A diffusion theory model of

spatially resolved, steady-state diffuse reflectance for the noninvasive determination of tissue optical properties *in vivo*. *Medical Physics*, 19(4): 879-888.

Fekete, A., 1993a. Elasticity characteristics of fruits, *International Symposium on Postharvest Treatment of Horticultural Crops* 368, pp. 199-205.

Fekete, A., 1993b. Non-destructive method of fruit elasticity determination, *Proc. 4th Int. Symposium on Fruit, Nut, and Vegetable Production Engineering*, pp. 309-315.

Fischer, R.L. and Bennett, A.B., 1991. Role of cell wall hydrolases in fruit ripening. *Annual Review of Plant Biology*, 42(1): 675-703.

Fulponi, L., 2006. Private voluntary standards in the food system: The perspective of major food retailers in OECD countries. *Food Policy*, 31(1): 1-13.

García-Ramos, F.J., Valero, C., Homer, I., Ortiz-Cañavate, J. and Ruiz-Altisent, M., 2005. Non-destructive fruit firmness sensors: a review. *Spanish Journal of Agricultural Research*, 3(1): 61-73.

Garcia, J.M. and Yousfi, K., 2005. Non-destructive and objective methods for the evaluation of the maturation level of olive fruit. *European Food Research and Technology*, 221(3-4): 538-541.

Giangiaco, R., 2006. Study of water–sugar interactions at increasing sugar concentration by NIR spectroscopy. *Food Chemistry*, 96(3): 371-379.

Gierson, D. and Kader, A., 1986. Fruit ripening and quality, *The tomato crop*. Springer, pp. 241-280.

Gillaspy, G., Ben-David, H. and Gruissem, W., 1993. Fruits: a developmental perspective. *The Plant Cell*, 5(10): 1439-1444.

Giovannoni, J., 2001. Molecular biology of fruit maturation and ripening. *Annual Review of Plant Biology*, 52(1): 725-749.

Giusti, A.M., Bignetti, E. and Cannella, C., 2008. Exploring new frontiers in total food quality definition and assessment: From chemical to neurochemical properties. *Food and Bioprocess Technology*, 1(2): 130-142.

Goetz, A.F., Vane, G., Solomon, J.E. and Rock, B.N., 1985. *Imaging Spectrometry for Earth Remote Sensing*. *Science*, 228(4704): 1147-1153.

Golparvar-Fard, M., Bohn, J., Teizer, J., Savarese, S. and Peña-Mora, F., 2011. Evaluation of image-based modeling and laser scanning accuracy for emerging automated performance monitoring techniques. *Automation in Construction*, 20(8): 1143-1155.

Gomez, A.H., He, Y. and Pereira, A.G., 2006. Non-destructive measurement of

acidity, soluble solids and firmness of Satsuma mandarin using Vis/NIR-spectroscopy techniques. *Journal of Food Engineering*, 77(2): 313-319.

Gorton, H.L., Brodersen, C.R., Williams, W.E. and Vogelmann, T.C., 2010. Measurement of the optical properties of leaves under diffuse light. *Photochemistry and photobiology*, 86(5): 1076-1083.

Gray, J., Picton, S., Giovannoni, J. and Grierson, D., 1994. The use of transgenic and naturally occurring mutants to understand and manipulate tomato fruit ripening. *Plant, Cell & Environment*, 17(5): 557-571.

Gunasekaran, S., 1996. Computer vision technology for food quality assurance. *Trends in Food Science & Technology*, 7(8): 245-256.

Gunasekaran, S., Paulsen, M. and Shove, G., 1985. Optical methods for nondestructive quality evaluation of agricultural and biological materials. *Journal of Agricultural Engineering Research*, 32(3): 209-241.

Hadfield, R.H., 2009. Single-photon detectors for optical quantum information applications. *Nature photonics*, 3(12): 696-705.

Haralick, R.M. and Shanmugam, K., 1973. Textural features for image classification. *IEEE Transactions on Systems, Man, and Cybernetics*(6): 610-621.

Haralick, R.M., Shanmugam, K. and Dinstein, I.H., 1973. Textural features for image classification. *IEEE Transactions on Systems, Man and Cybernetics*, (6): 610-621.

Harker, F.R., Marsh, K.B., Young, H., Murray, S.H., Gunson, F.A. and Walker, S.B., 2002. Sensory interpretation of instrumental measurements 2: sweet and acid taste of apple fruit. *Postharvest Biology and Technology*, 24(3): 241-250.

Harker, F.R., Redgwell, R.J., Hallett, I.C., Murray, S.H. and Carter, G., 2010. Texture of fresh fruit. *Horticultural Reviews*, Volume 20: 121-224.

Hashim, N., Pflanz, M., Regen, C., Janius, R.B., Abdul Rahman, R., Osman, A., Shitan, M. and Zude, M., 2013. An approach for monitoring the chilling injury appearance in bananas by means of backscattering imaging. *Journal of Food Engineering*, 116(1): 28-36.

Haskell, R.C., Svaasand, L.O., Tsay, T-T., Feng, T.-C., Tromberg, B.J. and McAdams, M.S., 1994. Boundary conditions for the diffusion equation in radiative transfer. *JOSA A*, 11(10): 2727-2741.

Heney, L.G. and Greenstein, J.L., 1941. Diffuse radiation in the galaxy. *The Astrophysical Journal*, 93: 70-83.

Hiraoka, Y., Sedat, J.W. and Agard, D.A., 1987. The use of a charge-coupled device for quantitative optical microscopy of biological structures. *Science*, 238(4823): 36-41.

- Hörtensteiner, S., 2006. Chlorophyll degradation during senescence*. *Annual Review of Plant Biology*, 57: 55-77.
- Huber, D., 1984. Strawberry fruit softening: the potential roles of polyuronides and hemicelluloses. *Journal of Food Science*, 49(5): 1310-1315.
- Hulme, A.C., 1971. The biochemistry of fruits and their products. Vol. 2. *The Biochemistry of Fruits and their Products*. Vol. 2, pp 50-62.
- Ishimaru, A., 1978. Wave propagation and scattering in random media, 2. Academic press New York, pp 23-32.
- Jacques, S.L., 1998. Light distributions from point, line and plane sources for photochemical reactions and fluorescence in turbid biological tissues. *Photochemistry and Photobiology*, 67(1): 23-32.
- Jaiswal, P., Jha, S.N., Kaur, P.P., Bhardwaj, R., Singh, A.K. and Wadhawan, V., 2014. Prediction of textural attributes using color values of banana (*Musa sapientum*) during ripening. *Journal of Food Science and Technology*, 51(6): 1179-1184.
- Jha, S., Chopra, S. and Kingsly, A., 2007. Modeling of color values for nondestructive evaluation of maturity of mango. *Journal of Food Engineering*, 78(1): 22-26.
- Jha, S.N., 2010. Colour measurements and modeling, *Nondestructive Evaluation of Food Quality*. Springer, pp. 17-40.
- Jha, S.N., Jaiswal, P., Narsaiah, K., Gupta, M., Bhardwaj, R. and Singh, A.K., 2012. Non-destructive prediction of sweetness of intact mango using near infrared spectroscopy. *Scientia Horticulturae*, 138: 171-175.
- Kader, A.A., 2002. Postharvest technology of horticultural crops, 3311. UCANR Publications, pp 1-20.
- Kader, A.A., 2008. Flavor quality of fruits and vegetables. *Journal of the Science of Food and Agriculture*, 88(11): 1863-1868.
- Kamruzzaman, M., Barbin, D., ElMasry, G., Sun, D.-W. and Allen, P., 2012. Potential of hyperspectral imaging and pattern recognition for categorization and authentication of red meat. *Innovative Food Science & Emerging Technologies*, 16: 316-325.
- Kienle, A., Lilge, L., Patterson, M.S., Hibst, R., Steiner, R. and Wilson, B.C., 1996. Spatially resolved absolute diffuse reflectance measurements for noninvasive determination of the optical scattering and absorption coefficients of biological tissue. *Applied Optics*, 35(13): 2304-2314.
- Kondo, N., Ahmad, U., Monta, M. and Murase, H., 2000. Machine vision based quality evaluation of Iyokan orange fruit using neural networks. *Computers and Electronics in Agriculture*, 29(1): 135-147.

- Laine, A. and Fan, J., 1996. Frame representations for texture segmentation. *IEEE Transactions on Image Processing*, 5(5): 771-780.
- Lammertyn, J., Nicolai, B., Ooms, K., De Smedt, V. and De Baerdemaeker, J., 1998. Non-destructive measurement of acidity, soluble solids, and firmness of Jonagold apples using NIR-spectroscopy. *Transactions of the ASAE*, 41(4): 1089.
- Lelièvre, J.M., Latchè, A., Jones, B., Bouzayen, M. and Pech, J.C., 1997. Ethylene and fruit ripening. *Physiologia Plantarum*, 101(4): 727-739.
- Liew, C. and Lau, C., 2012. Determination of quality parameters in Cavendish banana during ripening by NIR spectroscopy. *International Food Resources Journal*, 19(2): 751-758.
- Liu, C., Liu, W., Lu, X., Ma, F., Chen, W., Yang, J. and Zheng, L., 2014. Application of multispectral imaging to determine quality attributes and ripeness stage in strawberry fruit. *PloS One*, 9(2): e87818:1-8.
- Liu, Y., Chen, Y.-R., Kim, M.S., Chan, D.E. and Lefcourt, A.M., 2007. Development of simple algorithms for the detection of fecal contaminants on apples from visible/near infrared hyperspectral reflectance imaging. *Journal of Food Engineering*, 81(2): 412-418.
- Lizada, C., 1993. *Mango, Biochemistry of fruit ripening*. Springer, pp. 255-271.
- Lobit, P., Soing, P., Génard, M. and Habib, R., 2002. Theoretical analysis of relationships between composition, pH, and titratable acidity of peach fruit. *Journal of Plant Nutrition*, 25(12): 2775-2792.
- Lorente, D., Zude, M., Idler, C., Gómez-Sanchis, J. and Blasco, J., 2015. Laser-light backscattering imaging for early decay detection in citrus fruit using both a statistical and a physical model. *Journal of Food Engineering*, 154: 76-85.
- Lorente, D., Zude, M., Regen, C., Palou, L., Gómez-Sanchis, J. and Blasco, J., 2013. Early decay detection in citrus fruit using laser-light backscattering imaging. *Postharvest Biology and Technology*, 86: 424-430.
- Lu, R., 2001. Predicting firmness and sugar content of sweet cherries using near-infrared diffuse reflectance spectroscopy. *Transactions of the ASAE*, 44(5): 1265-1275.
- Lu, R., 2004a. Multispectral imaging for predicting firmness and soluble solids content of apple fruit. *Postharvest Biology and Technology*, 31(2): 147-157.
- Lu, R., 2004b. Prediction of apple fruit firmness by near-infrared multispectral scattering. *Journal of texture studies*, 35(3): 263-276.
- Lu, R., 2007. Nondestructive measurement of firmness and soluble solids content for apple fruit using hyperspectral scattering images. *Sensing and Instrumentation for Food Quality and Safety*, 1(1): 19-27.

- Lu, R. and Ariana, D., 2002. A near-infrared sensing technique for measuring internal quality of apple fruit. *Applied Engineering in Agriculture*, 18(5): 585-597.
- Lu, R., Cen, H., Huang, M. and Ariana, D.P., 2010. Spectral absorption and scattering properties of normal and bruised apple tissue. *Transactions of the ASABE*, 53(1): 263-269.
- Lu, R., Guyer, D.E. and Beaudry, R.M., 2000. Determination of firmness and sugar content of apples using near-infrared diffuse reflectance. *Journal of Texture Studies*, 31(6): 615-630.
- Lu, R. and Peng, Y., 2006. Hyperspectral scattering for assessing peach fruit firmness. *Biosystems Engineering*, 93(2): 161-171.
- Ludford, P.M., 1995. Postharvest hormone changes in vegetables and fruit, *Plant Hormones*. Springer, pp. 725-750.
- Manickavasagan, A. and Jayasuriya, H., 2014. *Imaging with Electromagnetic Spectrum. Applications in Food and Agriculture*, Springer, pp 232-240.
- Manley, M., Williams, P., Nilsson, D. and Geladi, P., 2009. Near infrared hyperspectral imaging for the evaluation of endosperm texture in whole yellow maize (*Zea mays* L.) kernels. *Journal of Agricultural and Food Chemistry*, 57(19): 8761-8769.
- Marsh, K., Attanayake, S., Walker, S., Gunson, A., Boldingh, H. and MacRae, E., 2004. Acidity and taste in kiwifruit. *Postharvest Biology and Technology*, 32(2): 159-168.
- Mattheis, J.P. and Fellman, J.K., 1999. Preharvest factors influencing flavor of fresh fruit and vegetables. *Postharvest Biology and Technology*, 15(3): 227-232.
- McGlone, V., Abe, H. and Kawano, S., 1997. Kiwifruit firmness by near infrared light scattering. *Journal of Near Infrared Spectroscopy*, 5: 83-90.
- McGlone, V., Kawano, A. and Kawano, S., 1998. Kiwifruit firmness by near infrared light scattering. *Journal of Near Infrared Spectroscopy*, 5(2): 83-89.
- Medlicott, A.P. and Thompson, A.K., 1985. Analysis of sugars and organic acids in ripening mango fruits (*Mangifera indica* L. var Keitt) by high performance liquid chromatography. *Journal of the Science of Food and Agriculture*, 36(7): 561-566.
- Mehl, P.M., Chen, Y.-R., Kim, M.S. and Chan, D.E., 2004. Development of hyperspectral imaging technique for the detection of apple surface defects and contaminations. *Journal of Food Engineering*, 61(1): 67-81.
- Mendoza, F. and Aguilera, J., 2004. Application of image analysis for classification of ripening bananas. *Journal of Food Science*, 69(9): E471-E477.
- Mendoza, F., Aguilera, J. and Dejmek, P., 2004. Predicting ripening stages of bananas

(Musa cavendish) by computer vision, V International Postharvest Symposium 682, pp. 1363-1370.

Mendoza, F., Lu, R., Ariana, D., Cen, H. and Bailey, B., 2011. Integrated spectral and image analysis of hyperspectral scattering data for prediction of apple fruit firmness and soluble solids content. *Postharvest Biology and Technology*, 62(2): 149-160.

Merry, C.J., 2010. 5 Basic Electromagnetic Radiation. *Manual of Geospatial Science and Technology*: pp 55-66.

Miller, B.K. and Delwiche, M.J., 1989. A color vision system for peach grading. *Transactions of the ASAE*, 32(4): 1484-1490.

Mireei, S., 2010. Nondestructive determination of effective parameters on maturity of mozafati & shahani date fruits by NIR spectroscopy technique, PhD thesis, Department of Mechanical Engineering of Agricultural Machinery, University of Tehran, Iran, pp 120-200.

Mireei, S., Mohtasebi, S., Massudi, R., Refiee, S. and Arabanian, A., 2010. Feasibility of near infrared spectroscopy for analysis of date fruits. *International Agrophysics*, 24: 351-356.

Moghimi, A., Aghkhani, M.H., Sazgarnia, A. and Sarmad, M., 2010. Vis/NIR spectroscopy and chemometrics for the prediction of soluble solids content and acidity (pH) of kiwifruit. *Biosystems Engineering*, 106(3): 295-302.

Mohsenin, N.N., 1970. Physical properties of plant and animal materials. Vol. 1. Structure, physical characteristics and mechanical properties. *Physical properties of plant and animal materials. Vol. 1. Structure, Physical Characteristics and Mechanical Properties.*, pp 1-50.

Mollazade, K., Omid, M., Akhlaghian Tab, F., Kalaj, Y.R., Mohtasebi, S.S. and Zude, M., 2013. Analysis of texture-based features for predicting mechanical properties of horticultural products by laser light backscattering imaging. *Computers and Electronics in Agriculture*, 98: 34-45.

Mollazade, K., Omid, M., Tab, F.A. and Mohtasebi, S.S., 2012. Principles and applications of light backscattering imaging in quality evaluation of agro-food products: a review. *Food and Bioprocess Technology*, 5(5): 1465-1485.

Monje, O.A. and Bugbee, B., 1992. Inherent limitations of nondestructive chlorophyll meters: a comparison of two types of meters. *HortScience*, 27(1): 69-71.

Moons, E., Sinnaeve, G. and Dardenne, P., 1998. Non destructive visible and NIR spectroscopy measurement for the determination of apple internal quality, XXV International Horticultural Congress, Part 7: Quality of Horticultural Products 517, pp. 441-448.

Mori, S., Monden, Y. and Mori, T., 1973. Edge representation in gradient space.

Computer Graphics and Image Processing, 2(3-4): 321-325.

- Mounet, F., Moing, A., Garcia, V., Petit, J., Maucourt, M., Deborde, C., Bernillon, S., Le Gall, G., Colquhoun, I. and Defernez, M., 2009. Gene and metabolite regulatory network analysis of early developing fruit tissues highlights new candidate genes for the control of tomato fruit composition and development. *Plant Physiology*, 149(3): 1505-1528.
- Narro, C., Roy, D., Okello, J., Avendaño, B., Rich, K. and Thorat, A., 2009. Public-private partnerships and collective action in high value fruit and vegetable supply chains. *Food Policy*, 34(1): 8-15.
- Nicolai, B.M., Beullens, K., Bobelyn, E., Peirs, A., Saeys, W., Theron, K.I. and Lammertyn, J., 2007. Nondestructive measurement of fruit and vegetable quality by means of NIR spectroscopy: A review. *Postharvest Biology and Technology*, 46(2): 99-118.
- Nicolai, B.M., Verlinden, B.E., Desmet, M., Saevels, S., Saeys, W., Theron, K., Cubeddu, R., Pifferi, A. and Torricelli, A., 2008. Time-resolved and continuous wave NIR reflectance spectroscopy to predict soluble solids content and firmness of pear. *Postharvest Biology and Technology*, 47(1): 68-74.
- Nogata, Y., Ohta, H. and Voragen, A., 1993. Polygalacturonase in strawberry fruit. *Phytochemistry*, 34(3): 617-620.
- Oeller, P.W. and Min-Wong, L., 1991. Reversible inhibition of tomato fruit senescence by antisense RNA. *Science*, 254(5030): 437.
- Okada, E., Firbank, M., Schweiger, M., Arridge, S.R., Cope, M. and Delpy, D.T., 1997. Theoretical and experimental investigation of near-infrared light propagation in a model of the adult head. *Applied Optics*, 36(1): 21-31.
- Ollinger, S., 2011. Sources of variability in canopy reflectance and the convergent properties of plants. *New Phytologist*, 189(2): 375-394.
- Otsu, N., 1975. A threshold selection method from gray-level histograms. *Automatica*, 11(285-296): 23-27.
- Özdemir, A.E., Candir, E.E., Toplu, C., Kaplankiran, M., Demirkeser, T.H. and Yildiz, E., 2009. The effects of physical and chemical changes on the optimum harvest maturity in some avocado cultivars. *African Journal of Biotechnology*, 8(9).
- Paliyath, G. and Murr, D.P., 2008. Biochemistry of fruits. *Food Biochemistry and Food Processing*: 487-514.
- Palmer, G.M. and Ramanujam, N., 2006. Monte Carlo-based inverse model for calculating tissue optical properties. Part I: Theory and validation on synthetic phantoms. *Applied Optics*, 45(5): 1062-1071.

- Park, B., Abbott, J.A., Lee, K., Choi, C. and Choi, K., 2003. Near-infrared diffuse reflectance for quantitative and qualitative measurement of soluble solids and firmness of Delicious and Gala apples. *Transactions of the ASAE*, 46(6): 1721.
- Paull, R.E. and Chen, N.J., 1983. Postharvest variation in cell wall-degrading enzymes of papaya (*Carica papaya* L.) during fruit ripening. *Plant Physiology*, 72(2): 382-385.
- Paz, P., Sánchez, M.-T., Pérez-Marín, D., Guerrero, J.-E. and Garrido-Varo, A., 2008. Nondestructive determination of total soluble solid content and firmness in plums using near-infrared reflectance spectroscopy. *Journal of Agricultural and Food Chemistry*, 56(8): 2565-2570.
- Paz, P., Sánchez, M.-T., Pérez-Marín, D., Guerrero, J.E. and Garrido-Varo, A., 2009. Instantaneous quantitative and qualitative assessment of pear quality using near infrared spectroscopy. *Computers and Electronics in Agriculture*, 69(1): 24-32.
- Pedrycz, W., Bonissone, P.P. and Ruspini, E.H., 1998. *Handbook of fuzzy computation*. Institute of Physics Pub., pp 67-80.
- Peng, Y. and Lu, R., 2006. Improving apple fruit firmness predictions by effective correction of multispectral scattering images. *Postharvest Biology and Technology*, 41(3): 266-274.
- Peng, Y. and Lu, R., 2007. Prediction of apple fruit firmness and soluble solids content using characteristics of multispectral scattering images. *Journal of Food Engineering*, 82(2): 142-152.
- Peng, Y. and Lu, R., 2008. Analysis of spatially resolved hyperspectral scattering images for assessing apple fruit firmness and soluble solids content. *Postharvest Biology and Technology*, 48(1): 52-62.
- Phillips, T.A., 2006. Spectral Reflectance Imagery and Baseline Analysis of Anthocyanin Concentration in *Gossypium Hirsutum* L, Dakota Wesleyan University, pp 80-87.
- Picton, S., Barton, S.L., Bouzayen, M., Hamilton, A.J. and Grierson, D., 1993. Altered fruit ripening and leaf senescence in tomatoes expressing an antisense ethylene-forming enzyme transgene. *The Plant Journal*, 3(3): 469-481.
- Prasanna, V., Prabha, T. and Tharanathan, R., 2007. Fruit ripening phenomena—an overview. *Critical Reviews in Food Science and Nutrition*, 47(1): 1-19.
- Punan, M.S., Rahman, A.S.A., Nor, L.M., Muda, P., Sapii, A.T., Yon, R.M. and Som, F.M., 2000. Establishment of a quality assurance system for minimally processed jackfruit, *ACIAR Proceedings*. ACIAR; 1998, pp. 115-122.
- Qin, J., 2007. Measurement of the optical properties of horticultural and food products

by hyperspectral imaging, *Applied Spectroscopy*, 61(3): 23-30.

- Qin, J. and Lu, R., 2007. Measurement of the absorption and scattering properties of turbid liquid foods using hyperspectral imaging. *Applied Spectroscopy*, 61(4): 388-396.
- Qin, J. and Lu, R., 2008. Measurement of the optical properties of fruits and vegetables using spatially resolved hyperspectral diffuse reflectance imaging technique. *Postharvest Biology and Technology*, 49(3): 355-365.
- Qin, J. and Lu, R., 2009. Monte Carlo simulation for quantification of light transport features in apples. *Computers and Electronics in Agriculture*, 68(1): 44-51.
- Qin, J., Lu, R. and Peng, Y., 2009. Prediction of apple internal quality using spectral absorption and scattering properties. *Transactions of the ASABE*, 52(2): 499-486.
- Qing, Z., Ji, B. and Zude, M., 2007a. Predicting soluble solid content and firmness in apple fruit by means of laser light backscattering image analysis. *Journal of Food Engineering*, 82(1): 58-67.
- Qing, Z., Ji, B. and Zude, M., 2007b. Wavelength selection for predicting physicochemical properties of apple fruit based on near-infrared spectroscopy. *Journal of Food Quality*, 30(4): 511-526.
- Qing, Z., Ji, B. and Zude, M., 2008. Non-destructive analyses of apple quality parameters by means of laser-induced light backscattering imaging. *Postharvest Biology and Technology*, 48(2): 215-222.
- Rady, A., Guyer, D. and Lu, R., 2015. Evaluation of Sugar Content of Potatoes using Hyperspectral Imaging. *Food and Bioprocess Technology*: 1-16.
- Rajkumar, P., Wang, N., Elmasry, G., Raghavan, G. and Garipey, Y., 2012. Studies on banana fruit quality and maturity stages using hyperspectral imaging. *Journal of Food Engineering*, 108(1): 194-200.
- Redgwell, R.J., MacRae, E., Hallett, I., Fischer, M., Perry, J. and Harker, R., 1997. In vivo and in vitro swelling of cell walls during fruit ripening. *Planta*, 203(2): 162-173.
- Robertson, G.L., 2012. *Food packaging: principles and practice*. CRC press, pp 15-25.
- Romano, G., Baranyai, L., Gottschalk, K. and Zude, M., 2008. An approach for monitoring the moisture content changes of drying banana slices with laser light backscattering imaging. *Food and Bioprocess Technology*, 1(4): 410-414.
- Romano, G., Nagle, M., Argyropoulos, D. and Müller, J., 2011. Laser light backscattering to monitor moisture content, soluble solid content and hardness of apple tissue during drying. *Journal of Food Engineering*, 104(4): 657-662.

- Sadler, G.D. and Murphy, P.A., 2010. pH and titratable acidity, Food analysis. Springer, pp. 219-238.
- Saeys, W., Velazco-Roa, M.A., Thennadil, S.N., Ramon, H. and Nicolai, B.M., 2008. Optical properties of apple skin and flesh in the wavelength range from 350 to 2200 nm. *Applied Optics*, 47(7): 908-919.
- Sakai, T., Sakamoto, T., Hallaert, J. and Vandamme, E.J., 1993. [Pectin, Pectinase, and Protopectinase: Production,] Properties, and Applications. *Advances in Applied Microbiology*, 39: 213-294.
- Sampsel, J.B., 1996. Spatial light modulator scanning system. Google Patents, pp 1-5.
- Sanaeifar, A., Bakhshipour, A. and de la Guardia, M., 2016a. Prediction of banana quality indices from color features using support vector regression. *Talanta*, 148: 54-61.
- Sanaeifar, A., Mohtasebi, S.S., Ghasemi-Varnamkhasti, M. and Ahmadi, H., 2016b. Application of MOS based electronic nose for the prediction of banana quality properties. *Measurement*, 82: 105-114.
- Saranwong, S., Sornsrivichai, J. and Kawano, S., 2004. Prediction of ripe-stage eating quality of mango fruit from its harvest quality measured nondestructively by near infrared spectroscopy. *Postharvest Biology and Technology*, 31(2): 137-145.
- Schaare, P. and Fraser, D., 2000. Comparison of reflectance, interactance and transmission modes of visible-near infrared spectroscopy for measuring internal properties of kiwifruit (*Actinidia chinensis*). *Postharvest Biology and Technology*, 20(2): 175-184.
- Schreiner, M., Korn, M., Stenger, M., Holzgreve, L. and Altmann, M., 2013. Current understanding and use of quality characteristics of horticulture products. *Scientia Horticulturae*, 163: 63-69.
- Selvaraj, Y., Kumar, R. and Pal, D., 1989. Changes in sugars, organic acids, amino acids, lipid constituents and aroma characteristics of ripening mango (*Mangifera indica* L) fruit. *Journal of Food Science and Technology*, 26(6): 308-313.
- Serrano, M., Zapata, P.J., Guillén, F., Martínez-Romero, D., Castillo, S. and Valero, D., 2008. Post-harvest ripening of tomato. *Tomatoes and Tomato Products*. Edited by Preedy VR and Watson RR. Enfield: Science Publishers: 67-84.
- Shaw, G. and Manolakis, D., 2002. Signal processing for hyperspectral image exploitation. *IEEE Signal Processing Magazine*, 19(1): 12-16.
- Shewfelt, R.L., 1999. What is quality? *Postharvest Biology and Technology*, 15(3): 197-200.
- Simpson, C.R., Kohl, M., Essenpreis, M. and Cope, M., 1998. Near-infrared optical

- properties of ex vivo human skin and subcutaneous tissues measured using the Monte Carlo inversion technique. *Physics in Medicine and Biology*, 43(9): 2465.
- Smith, W.H., 1950. Cell-multiplication and cell-enlargement in the development of the flesh of the apple fruit. *Annals of Botany*, 14(53): 23-38.
- Soft, S., 2013. *Statistica 12*. Tulsa, OK: Stat Soft Inc.
- Soltani, M., Alimardani, R. and Omid, M., 2010. Prediction of banana quality during ripening stage using capacitance sensing system. *Australian Journal of Crop Science*, 4(6): 443.
- Soltani, M., Alimardani, R. and Omid, M., 2011. Evaluating banana ripening status from measuring dielectric properties. *Journal of Food Engineering*, 105(4): 625-631.
- Steiner, R., 2011. Laser-tissue interactions, *Laser and IPL Technology in Dermatology and Aesthetic Medicine*. Springer, pp. 23-36.
- Sun, D.-W., 2000. Inspecting pizza topping percentage and distribution by a computer vision method. *Journal of Food Engineering*, 44(4): 245-249.
- Tamura, H., Mori, S. and Yamawaki, T., 1978. Textural features corresponding to visual perception. *IEEE Transactions on Systems, Man and Cybernetics*, 8(6): 460-473.
- Theologis, A., Oeller, P.W., Wong, L.m., Rottmann, W.H. and Gantz, D.M., 1993. Use of a tomato mutant constructed with reverse genetics to study fruit ripening, a complex developmental process. *Developmental Genetics*, 14(4): 282-295.
- Tsai, C.-L., Yang, Y.-F., Han, C.-C., Hsieh, J.-H. and Chang, M., 2001. Measurement and simulation of light distribution in biological tissues. *Applied Optics*, 40(31): 5770-5777.
- Tsuchikawa, S. and Hamada, T., 2004. Application of time-of-flight near infrared spectroscopy for detecting sugar and acid contents in apples. *Journal of Agricultural and Food Chemistry*, 52(9): 2434-2439.
- Tucker, G. and Grierson, D., 2013. Fruit ripening. *The Biochemistry of Plants*, 12: 265-381.
- Unser, M., 1995. Texture classification and segmentation using wavelet frames. *IEEE Transactions on Image Processing*, 4(11): 1549-1560.
- Van der Vorst, J.G., Tromp, S.-O. and Zee, D.-J.v.d., 2009. Simulation modelling for food supply chain redesign; integrated decision making on product quality, sustainability and logistics. *International Journal of Production Research*, 47(23): 6611-6631.

- Vangdal, E., 1985. Quality criteria for fruit for fresh consumption. *Acta Agriculturae Scandinavica*, 35(1): 41-47.
- Ventura, M., de Jager, A., de Putter, H. and Roelofs, F.P., 1998. Non-destructive determination of soluble solids in apple fruit by near infrared spectroscopy (NIRS). *Postharvest Biology and Technology*, 14(1): 21-27.
- Villa-Rodríguez, J.A., Molina-Corral, F.J., Ayala-Zavala, J.F., Olivas, G.I. and González-Aguilar, G.A., 2011. Effect of maturity stage on the content of fatty acids and antioxidant activity of 'Hass' avocado. *Food Research International*, 44(5): 1231-1237.
- Vinha, A.F., Moreira, J. and Barreira, S.V., 2013. Physicochemical Parameters, Phytochemical Composition and Antioxidant Activity of the Algarvian Avocado (*Persea americana* Mill.). *Journal of Agricultural Science*, 5(12): 100-110.
- Voragen, A.G., Coenen, G.-J., Verhoef, R.P. and Schols, H.A., 2009. Pectin, a versatile polysaccharide present in plant cell walls. *Structural Chemistry*, 20(2): 263-275.
- Waikato, U., 2009. Weka. Verzija.
- Wang, H., Schauer, N., Usadel, B., Frasse, P., Zouine, M., Hernould, M., Latché, A., Pech, J.-C., Fernie, A.R. and Bouzayen, M., 2009. Regulatory features underlying pollination-dependent and-independent tomato fruit set revealed by transcript and primary metabolite profiling. *The Plant Cell*, 21(5): 1428-1452.
- Wang, L.V. and Liang, G., 1999. Absorption distribution of an optical beam focused into a turbid medium. *Applied Optics*, 38(22): 4951-4958.
- Wei, X., Liu, F., Qiu, Z., Shao, Y. and He, Y., 2014. Ripeness classification of astringent persimmon using hyperspectral imaging technique. *Food and Bioprocess Technology*, 7(5): 1371-1380.
- Weszka, J.S., Dyer, C.R. and Rosenfeld, A., 1976. A comparative study of texture measures for terrain classification. *IEEE Transactions on Systems, Man, and Cybernetics*, 4(SMC-6): 269-285.
- Wills, R.B.H., McGlasson, W.B., Graham, D. and Joyce, D., 2007. *Postharvest: an introduction to the physiology and handling of fruit, vegetables and ornamentals*. CABI, pp 54-60.
- Wu, D. and Sun, D.-W., 2013. Colour measurements by computer vision for food quality control—A review. *Trends in Food Science & Technology*, 29(1): 5-20.
- Yam, K.L. and Papadakis, S.E., 2004. A simple digital imaging method for measuring and analyzing color of food surfaces. *Journal of Food Engineering*, 61(1): 137-142.

- Yamaki, S., Machida, Y. and Kakiuchi, N., 1979. Changes in cell wall polysaccharides and monosaccharides during development and ripening of Japanese pear fruit. *Plant and Cell Physiology*, 20(2): 311-321.
- Yang, Q., 1994. An approach to apple surface feature detection by machine vision. *Computers and Electronics in Agriculture*, 11(2): 249-264.
- Zhang, D., Lee, D.-J., Tippetts, B.J. and Lillywhite, K.D., 2014. Date maturity and quality evaluation using color distribution analysis and back projection. *Journal of Food Engineering*, 131: 161-169.
- Zheng, C., Sun, D.-W. and Zheng, L., 2006. Recent applications of image texture for evaluation of food qualities—a review. *Trends in Food Science & Technology*, 17(3): 113-128.
- Zhu, Q., He, C., Lu, R., Mendoza, F. and Cen, H., 2015. Ripeness evaluation of ‘Sun Bright’ tomato using optical absorption and scattering properties. *Postharvest Biology and Technology*, 103: 27-34.
- Zude, M., 2003. Non-destructive prediction of banana fruit quality using VIS/NIR spectroscopy. *Fruits*, 58(03): 135-142.
- Zude, M., Herold, B., Roger, J.-M., Bellon-Maurel, V. and Landahl, S., 2006. Non-destructive tests on the prediction of apple fruit flesh firmness and soluble solids content on tree and in shelf life. *Journal of Food Engineering*, 77(2): 254-260.
- Zúñiga-Arias, G.E., 2007. Quality management and strategic alliances in the mango supply chain from Costa Rica: an interdisciplinary approach for analysing coordination, incentives and governance, 3. Wageningen Academic Pub, pp 102-110.