



**UNIVERSITI PUTRA MALAYSIA**

***OPTIMIZED TECHNIQUES FOR LANDSLIDE DETECTION AND  
CHARACTERISTICS USING LiDAR DATA***

**MUSTAFA RIDHA MEZAAL**

**FK 2018 79**



**OPTIMIZED TECHNIQUES FOR LANDSLIDE DETECTION AND  
CHARACTERISTICS USING LiDAR DATA**

By

**MUSTAFA RIDHA MEZAAL**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of  
Doctor of Philosophy**

**May 2018**

## **COPYRIGHT**

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment  
of the requirement for the degree of Doctor of Philosophy

## **OPTIMIZED TECHNIQUES FOR LANDSLIDE DETECTION AND CHARACTERISTICS USING LiDAR DATA**

By

**MUSTAFA RIDHA MEZAAL**

**May 2018**

**Chairman : Professor Biswajeet Pradhan, PhD**  
**Faculty : Engineering**

Landslides are one of the major natural disasters that cause severe damage to lives and properties worldwide. Historically, landslide occurrences are usually mapped by taking inventory of location and magnitude of the landslide in a region. This information is used to examine and manage slope failures and distributions effectively. A good landslide inventory map is a prerequisite for analyzing landslide susceptibility, hazard, and risk. Field survey, optical remote sensing, and synthetic aperture radar techniques are traditional techniques use for landslide detection in tropical regions. However, such techniques are time consuming and costly. In addition, the dense vegetation in tropical forests affects the accuracy of the landslide inventory maps. Furthermore, it is difficult to distinguish different types of landslides due to geomorphological development along hillsides using the traditional approach. so, it necessary to develop more innovative approach that can resolve the aforementioned challenges.

Therefore, in line with the objectives of this research, very- high-resolution LiDAR point cloud data and orthophotos image, have been utilized to map the landslide events in Cameron Highlands, Malaysia. The segmentation process was optimized using Fuzzy-based Segmentation Parameter. Also, six techniques: Ant Colony Optimization (ACO), Gain Ratio (GR), Particle Swarm Optimization (PSO) and Genetic Algorithm (GA), Random forest (RF), and Correlation-based Feature Selection (CFS) were used for the feature selection. The locations of landslides were detected accurately by employing two Machine learning classifiers, namely, SVM and RF, decision rule and hierarchal rules sets were developed by applying decision tree (DT) algorithm to provide improved landslide inventory. In this task, two neural network algorithms, Recurrent Neural Networks (RNN) and Multi-Layer Perceptron Neural Networks

(MLP-NN) were used and the hyper-parameters of the network architecture was optimized based on a systematic grid search.

The performance of the outcome was validated based on the receiver operating characteristic (ROC) area under the curve (AUC) values, confusion matrix and Cross Validation method. Transferability of each of the models was verified by testing in another site for consistency. The overall accuracy of the Support Vector Machine SVM and Random Forest RF classifiers revealed that three of the six algorithms exhibited higher ranks in the landslide detection. The classification accuracy of the RF classifier is observed to be higher than that of SVM using either all features or only the optimal features. The proposed techniques performed well in detecting landslides in tropical area in Malaysia. Furthermore, the transferability indicates that the techniques can easily be extended to any region with similar characteristics.

The result show that the accuracy of shallow and deep-seated landslides were 0.80 and 0.83, respectively. The intensity derived from the LiDAR data, geometric and texture features significantly affects the accuracy of differentiating shallow from deep-seated landslides. While, the results of shallow and deep using hierarchal rules set were observed to be 87.2%, and 90% respectively, for site A (Analysis area). More so, the hierarchal rules set were evaluated using another site named site B (Test area), and the accuracies of shallow and deep seated were found to be 86.4% and 80.8% respectively. This indicates that LiDAR data are highly efficient in detecting landslide characteristics in tropical forested areas.

Furthermore, RNN and MLP-NN models in the test area showed 81.11%, and 74.56%, accuracy level, respectively. These results indicated that the proposed models with optimized hyper-parameters produced the accurate classification results. The LiDAR-derived data, orthophotos and textural features significantly affected the classification results. The results indicated that the proposed methods have the potential to produce accurate and appropriate landslide inventory in tropical regions such as Malaysia. Hopefully, this innovative method can be deployed in detecting landslide and distinguish between different types of landslides (shallow and deep-seated landslides) in the near future for landslide management due to its transferability capabilities to different environments.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PEMBANGUNAN TEKNIK YANG DIKENAKAN UNTUK PENGESANAN  
TANAH RUNTUH DAN SIFAT DARI PEMANTAUAN DATA DARI LASER  
UDARA DAN ORTHOFOTO**

Oleh

**MUSTAFA RIDHA MEZAAL**

**Mei 2018**

**Pengerusi : Profesor Biswajeet Pradhan, PhD**  
**Fakulti : Kejuruteraan**

Tanah runtuh merupakan salah satu bencana alam yang boleh mengakibatkan kemalangan jiwa dan kerosakan harta benda. Menurut sejarah, kejadian tanah runtuh kebiasaannya dikesan dengan mengenal pasti lokasi dan magnitud kejadian tanah runtuh tersebut di sesuatu kawasan dan maklumat ini digunakan untuk mengkaji corak tebaran tanah runtuh bagi menguruskan hakisan tanah dengan berkesan. Peta kajian tanah runtuh yang baik merupakan satu keperluan penting bagi mengenal pasti kawasan yang terdedah dan berisiko untuk berlakunya tanah runtuh selain daripada untuk mengkaji perubahan landskap yang berlaku akibat daripada kejadian tanah runtuh. Pengukuran tanah, sensor penderiaan jauh dan teknik sintetik bukaan radar adalah antara teknik tradisional untuk mengenal pasti tanah runtuh di kawasan tropika. Walaubagaimanapun, teknik-teknik tradisional ini adalah lambat dan menelan kos yang sangat besar. Tambahan pula, ketebalan tumbuhan hutan tropika merumitkan penghasilan peta kajian tanah runtuh di kawasan ini dengan lebih berkesan. Teknik-teknik tradisional ini bukan sahaja lambat bahkan sukar untuk membezakan jenis-jenis kejadian tanah runtuh kerana pembangunan geomorfologi di kawasan persisiran bukit dan pergerakan tektonik dalam tanah itu sendiri.

Hasil kajian ini membentangkan beberapa kaedah untuk mengesan tanah runtuh dan berupaya bagi membezakan antara tanah runtuh yang cetek dan dalam dengan menggunakan pengesan cahaya dan berbagai-bagai set data-LiDar dan kepenggunaannya mengikut keadaan persekitaran yang berbeza. Antara objektif khusus kajian ini adalah: 1) mengenal pasti algoritma yang sesuai untuk pemilihan ciri bagi menambah baik proses mengesan tanah runtuh, 2) untuk mengoptimumkan ketetapan peraturan keputusan data set dan hierarki peraturan untuk membezakan ciri-

ciri kegelongsoran yang berlainan, 3) untuk membangunkan prosedur pembelajaran (Seni Bina Saraf) bagi pemetaan tanah runtuh. Bagi mencapai matlamat ini, pertama, proses segmentasi dioptimumkan dengan menggunakan selia logik kabur. Selepas itu, enam teknik: Pengoptimuman Koloni Semut (ACO), Nisbah Perolehan (GR), Pengoptimuman Kawan Zarah (PSO) dan Algoritma Genetik (GA), Hutan Rawak (RF), dan Hubungan Pemilihan Ciri (CFS) telah digunakan untuk pemilihan ciri. Di sini, lokasi tanah runtuh telah dikesan dengan tepat dengan menggunakan dua teknik mesin pengasingan pembelajaran, iaitu, MSV dan HR. Di samping itu, peraturan keputusan dan peraturan set hierarki telah dibangunkan dengan menggunakan Pohon Keputusan (DT) algoritma untuk menyediakan inventori tanah runtuh yang lebih baik. Dalam tugas ini, dua rangkaian neural algoritma telah digunakan iaitu Rangkaian Saraf Berulang (RNN) dan Rangkaian Saraf Pelbagai Lapisan Perceptron (MLP-NN). Dalam proses ini juga, lebih parameter dari rangkaian seni bina telah dioptimumkan berdasarkan carian grid yang sistematik.

Kesemua prestasi tersebut telah disahkan berdasarkan nilai di bawah kurva (AUC) dalam ciri-ciri operasi penerima (ROC), matriks kekeliruan dan pengesanan silang. Setiap pemindahan model telah disahkan dengan diguna pakai di kawasan lain. Untuk objektif pertama, keseluruhan ketepatan daripada Mesin Sokongan Vektor (SVM) dan pengelas Hutan Rawak (RF) mendedahkan bahawa tiga daripada enam algoritma berjaya mempamerkan ketepatan yang lebih tinggi dalam mengesan kejadian tanah runtuh. Ketepatan pengelasan bagi pengelas (RF) adalah lebih tinggi daripada pengelas (SVM) walau sama ada dengan menggunakan kesemua ciri atau ciri optimum sahaja. Teknik yang dicadangkan ini adalah yang terbaik dalam mengesan tanah runtuh di kawasan tropika seperti Malaysia. Oleh yang demikian, teknik-teknik ini mudah untuk diguna pakai di mana-mana kawasan yang mempunyai ciri-ciri yang hampir serupa.

Objektif kedua menunjukkan bahawa ketepatan bagi mengenal pasti jenis-jenis tanah runtuh sama ada cetek dan mendalam adalah 0.80 dan 0.83. Keamatan yang diperolehi daripada tekstur dan data LiDAR dengan ketara sangat mempengaruhi ketepatan dalam membezakan tanah runtuh cetek dengan mendalam. Walau bagaimanapun, didapati bahawa keputusan data bagi tanah runtuh cetek dan mendalam dengan menggunakan aturan hierarki adalah 87.2% dan 90.0%., untuk kawasan A (iaitu kawasan yang dianalisis). Sementara itu, set aturan hierarki yang sama diguna pakai di kawasan B (iaitu kawasan yang dikaji), dan ketepatan dalam mengenal pasti tanah runtuh cetek dan mendalam adalah 86.4% dan 80.8%. Oleh yang demikian, hasil kajian ini menunjukkan bahawa data LiDAR adalah sangat berkesan dalam mengesan dan mengenal pasti jenis-jenis tanah runtuh di kawasan hutan hujan tropika.

Selain itu, hasil daripada objektif ketiga pula menunjukkan bahawa ketepatan bagi model RNN dan MLP-NN yang dijalankan di kawasan uji kaji masing-masing adalah 81.11% dan 74.56%. Oleh itu, keputusan ini menunjukkan bahawa model yang dicadangkan dalam kajian ini menghasilkan hasil klasifikasi yang sangat tepat dengan

mengoptimumkan beberapa parameter-hiper. Di sini, data daripada LiDAR, orthofotos dan ciri-ciri tekstur amat mempengaruhi hasil klasifikasi kajian.

Keputusan dari hasil kajian ini menunjukkan bahawa kaedah yang dicadangkan berpotensi untuk menghasilkan inventori jenis tanah runtuh dengan tepat dan sesuai diguna pakai di kawasan hutan hujan tropika seperti Malaysia. Algoritma dan maklumat yang diperolehi daripada kajian ini dapat menyumbang kepada pengurusan tanah runtuh di negara-negara hutan hujan tropika.





## ACKNOWLEDGEMENTS

I praise ALLAH for his great loving kindness, which has brought all of us to tell and encourage each other and who has pulled us from the darkness to the light. All respect for our holy prophet (Peace be upon him), who guided us to identify our creator. I also thank all my brothers and sister who answered ALLAH's call and have made their choice to be in the straight path of ALLAH.

As always it is impossible to mention everybody who had an impact to this work however there are those whose spiritual support is even more important. I feel a deep sense of gratitude for my mother and father, who formed part of my vision and taught me good things that really matter in life. Their infallible love and support has always been my strength. Their patience and sacrifice will remain my inspiration throughout my life. I am also very much grateful to all my family members for their constant inspiration and encouragement specially my mother in law.

My heartfelt thanks to my love and my wife Doaa Talib Hashim for her guidance and moral support. She always helped me out when I got any difficulties or queries regarding all the aspect of life. Again I thank her for standing by my side and sharing a great relationship as compassionate wife. I will always remember the warmth shown by her.

I take this opportunity to express my profound gratitude and deep regards to my guide Prof. Dr. Biswajeet Pradhan for his exemplary guidance, monitoring and constant encouragement throughout the course of this thesis. The blessing, help and guidance given by him time to time shall carry me a long way in the journey of life on which I am about to embark. He created an atmosphere that encouraged innovation and shared his extraordinary experiences throughout the work. Without his unflinching encouragement, it would have been impossible for me to finish this research.

I am so thankful to my committee Assoc. Prof. Dr. Helmi Zulhaidi bin Mohd Shafri and Dr. Zainuddin bin Md Yusoff, for the valuable information provided by them in their respective fields. I am obliged for their cooperation during the period of my study.

I certify that a Thesis Examination Committee has met on 17 May 2018 to conduct the final examination of Mustafa Ridha Mezaal on his thesis entitled "Optimized Techniques for Landslide Detection and Characteristics Using LiDAR Data" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

**Farzad Hejazi, PhD**

Associate Professor  
Faculty of Engineering  
Universiti Putra Malaysia  
(Chairman)

**Dato' Shattri bin Mansor, PhD**

Professor  
Faculty of Engineering  
Universiti Putra Malaysia  
(Internal Examiner)

**Abdul Rashid bin Mohamed Sharif, PhD**

Professor Sr.Gs.  
Faculty of Engineering  
Universiti Putra Malaysia  
(Internal Examiner)

**Hyung-Sup Jung, PhD**

Professor  
University of Seoul  
South Korea  
(External Examiner)



**RUSLI HAJI ABDULLAH, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 30 July 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Biswajeet Pradhan, PhD**

Professor  
Faculty of Engineering  
Universiti Putra Malaysia  
(Chairman)

**Helmi Zulhaidi Mohd Shafri, PhD**

Associate Professor  
Faculty of Engineering  
Universiti Putra Malaysia  
(Member)

**Zainuddin Bin Md Yusoff, PhD**

Senior Lecturer  
Faculty of Engineering  
Universiti Putra Malaysia  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Mustafa Ridha Mezaal, GS45396

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: \_\_\_\_\_  
Name of  
Chairman of  
Supervisory  
Committee: Professor Dr. Biswajeet Pradhan

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: Associate Professor Dr. Helmi Zulhaidi Mohd Shafri

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: Dr. Zainuddin Bin Md Yusoff

## TABLE OF CONTENTS

	Page
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	vi
<b>APPROVAL</b>	vii
<b>DECLARATION</b>	ix
<b>LIST OF TABLES</b>	xv
<b>LIST OF FIGURES</b>	xvii
<b>LIST OF ABBREVIATIONS</b>	xx
 <b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
1.1 Introduction	1
1.2 Problem Statement	2
1.3 Motivation Behind this Research	2
1.4 Research Objectives	3
1.5 Research Questions	3
1.6 Scope of this Thesis	4
1.7 Research Hypothesis	4
1.8 Thesis Organization	5
 <b>2 LITERATURE REVIEW</b>	<b>6</b>
2.1 Introduction	6
2.2 Landslide detection	6
2.2.1 Types of Landslides	6
2.2.1.1 Deep-seated Landslide	7
2.2.1.2 Shallow Landslide	9
2.3 Traditional Based Techniques	10
2.3.1 Geomorphological Field Mapping	10
2.3.2 Visual Interpretation	11
2.4 Innovative and Emerging Techniques	12
2.4.1 Image Classification	12
2.4.1.1 Pixel Based Techniques	13
2.4.1.2 Object Based Techniques	14
2.5 Analysis of Very-High Resolution DEMs	17
2.6 LiDAR data and Geo-Morphometric Features	18
2.7 Machine Learning Algorithms	22
2.7.1 Support Vector Machine (SVM)	23
2.7.2 Random Forest (RF)	24
2.7.3 Decision Trees (DT)	24
2.7.4 Deep Neural Networks (DNN)	25
2.7.4.1 Multi-Layer Perceptrons (MLP)	26
2.7.4.2 Recurrent Neural Networks (RNN)	26

2.8	Hierarchal Approach	27
2.9	Feature Selection	28
2.9.1	Ant Colony Optimization (ACO)	29
2.9.2	Gain ratio (GR)	29
2.9.3	Particle swarm optimization (PSO)	30
2.9.4	Random Forest (RF)	30
2.9.5	Genetic Algorithms (GA)	30
2.9.6	Correlation Feature Selection (CFS)	31
2.10	Summary	31
<b>3</b>	<b>MATERIALS AND METHODS</b>	<b>33</b>
3.1	Introduction	33
3.2	Overall Methodology	33
3.3	Study Areas	35
3.3.1	Geological, and Morphological Characteristics of the Study Area	36
3.4	Data Used	38
3.4.1	Landslide Inventory	38
3.4.2	LiDAR Data	40
3.4.2.1	LiDAR Point Cloud	40
3.4.2.2	Orthophoto	40
3.5	Pre-Analysis	40
3.5.1	Pre-processing	40
3.5.2	Object Based Approach	44
3.5.2.1	Image Segmentation	44
3.6	Investigate the Suitable Selection Features Algorithm	47
3.6.1	Overall Methodology	47
3.6.2	Feature Selection Using Six Algorithms (ACO, GRA, PSO, GA, RF and CFS)	48
3.6.2.1	ACO	50
3.6.2.2	GR	51
3.6.2.3	PSO	52
3.6.2.4	RF	53
3.6.2.5	GA	54
3.6.2.6	CFS	54
3.7	Machine Learning Algorithms	55
3.7.1	SVM Classifier	55
3.7.2	RF Classifier	56
3.8	Optimize Rule Sets to Differentiate Various Landslide Characteristics and Compare its Performance with Hierarchical Rule-Set Classification Approach.	58
3.8.1	Selecting the most Suitable Algorithm for Feature Selection	58
3.8.1.1	Overall Methodology	58
3.8.1.2	Feature selection using Three Algorithms (ACO, RF and CFS)	59
3.8.1.3	Evaluating the Feature Selection Algorithms	60



3.8.2	Develop Decision Rule Sets and Hierarchical Rule Sets for Distinguishing Landslide Types (i.e. Shallow and Deep seated)	60
3.8.2.1	Overall Methodology	60
3.8.2.2	Decision Tree	61
3.8.2.3	Implementations of Hierarchical Rules Set	63
3.9	Applied Deep Learning (Neural Architecture) Procedure for Landslide Mapping	64
3.9.1	MLP-NN Model	64
3.9.1.1	Architecture of the MLP-NN Model	65
3.9.2	RNN Model	66
3.9.2.1	Architecture of RNN Model	66
3.9.3	Training data for MLP-NN and RNN Models	67
3.10	Accuracy Assessment	67
3.11	Summary	69
<b>4</b>	<b>RESULTS AND DISCUSSION</b>	<b>70</b>
4.1	The Results of Selected Feature Techniques	70
4.1.1	Optimized Segmentation	70
4.1.2	Selection of the Relevant Features by Various Algorithms	71
4.1.3	Results of the SVM and RF Classifiers	77
4.1.4	Accuracy Assessment for Landslide Detection	78
4.1.5	Discussion	80
4.1.6	Field Investigation	81
4.2	Distinguish two Types of Landslide Namely Shallow and Deep seated	82
4.2.1	Optimizing the Boundary of the Types of Landslide	82
4.2.2	Selection Relevant Feature through three Algorithms (CFS, ACO and RF) for Different types of Landslide	83
4.2.3	RF Classifier for Differentiation between Shallow and Deep Seated Landslide in the Analysis Area	88
4.2.4	Evaluation of CFS based Feature Selection to other Site (i.e. Test site)	90
4.2.5	Accuracy Assessment	91
4.2.6	Discussion	93
4.2.7	Field Investigation	94
4.3	Optimized Rule Sets for Detecting Landslide and Differentiation	95
4.3.1	Segmentation Parameters Selected Using a Fuzzy Logic Supervised Approach	95
4.3.2	Features Selected Using the CFS Method	96
4.3.3	Rule Sets Developed for Landslide Detection and Characterization	97
4.3.4	Model Transferability	98
4.3.5	Discussion	101
4.3.6	Accuracy Assessment	103
4.3.7	Field Investigation	105



4.4	Hierarchical Rule-based Classification for differentiating Shallow and Deep Seated Landslide	105
4.4.1	Optimized Segmentation based on FbSP Optimizer	105
4.4.2	Feature Selection Using CFS method	106
4.4.3	Developed Rules Set based on Hierarchal Classification	107
4.4.4	Evaluation of the Developed Rules Sets	112
4.5	Effect of Using Intensity on the Image Segmentation	113
4.5.1	Accuracy Assessment	114
4.5.2	Discussion	116
4.5.3	Filed Investigation	117
4.6	Optimized Neural Architecture for Landslide Detection	117
4.6.1	Supervised Approach for Optimizing Segmentation	117
4.6.2	Relevant Feature Subset based on CFS Algorithm	118
4.6.3	Results of Landslide Detection	119
4.6.4	Performance of the MLP-NN and RNN Models	121
4.6.5	Sensitivity Analysis	122
4.6.6	Field Investigation	124
4.7	Comprehensive Comparison Assessment	125
4.8	Summary	125
<b>5</b>	<b>CONCLUSION AND RECOMMENDATIONS</b>	127
5.1	Overview	127
5.2	Contributions	127
5.3	Overall Findings	128
5.4	Conclusion	128
5.5	Recommendations for Future Work	131
	<b>REFERENCES</b>	132
	<b>APPENDICES</b>	158
	<b>BIODATA OF STUDENT</b>	163
	<b>LIST OF PUBLICATIONS</b>	164

## LIST OF TABLES

Table		Page
2.1	Types of Landslide with brief version	7
2.2	Overview of Techniques for the Collection of Landslide Information Obtained from (Van Westen et al., 2008)	20
3.1	The Feature Selection Using in the Current Research	50
3.2	Optimizing the Parameters of Support Vector Machine	56
3.3	Object Features used in the Current this Study	60
4.1	Multi- resolution Segmentation Parameters	70
4.2	Optimal Feature Selection for Detection Landslide using Various Algorithms	73
4.3	Correlation Coefficient for the Best Feature Selection in Landslide Detection	76
4.4	Results Comparison Based on Overall Accuracy and Kappa Coefficient for Important Features and Full Features using RF and SVM Algorithms	79
4.5	Results Comparison Based on User's Accuracy and Producer's Accuracy for Important Features and Full Features using RF and SVM Algorithms	79
4.6	Multi-resolution Segmentation Parameters	83
4.7	The Important Features Selection through three Algorithms for Detecting Types of Landslide (i.e. Shallow and Deep Seated landslides)	85
4.8	Correlation Coefficient for the Best Feature Selection in Distinguishing Landslide Types	87
4.9	Results Comparison based on Overall Accuracy for Important and Full Features using RF Classifier	92
4.10	Results Comparison based on User's Accuracy and Producer's Accuracy for Important and Full Features using RF Classifier	93
4.11	Optimized Segmentation Parameters	96
4.12	Results of the Features Selection using CFS Method	97
4.13	Rules Defied for Identification Landslide from Non-Landslide	97
4.14	Rules Defined for Distinguishing between Types of Landslide	100
4.15	Relevant Features Selected based on CFS Algorithm	107

4.16	Rule sets Developed by the DT Algorithm using the Important Feature Subset	109
4.17	Shows the Results of Overall Accuracy, Kappa Coefficient, user's Accuracy and Producer's accuracy for Analysis Area	115
4.18	Shows the Results of Overall Accuracy, Kappa Coefficient, User's Accuracy and Producer's Accuracy for test site "B"	115
4.19	Multi-resolution Segmentation Parameters	118
4.20	CFS Results for the most Relevant Feature Subset	119
4.21	Cross-Validation Accuracy Results of the Proposed Models	122



## LIST OF FIGURES

Figure	Page
3.1 Overall Methodology Flowchart	34
3.2 Shows the Location of Different Study Areas in Cameron Highland Malaysia	35
3.3 The Characteristic Maps of (A) LULC, (B) Geological, and, (C) Soil Maps of the entire Study Area	36
3.4 The Landslide Inventory Maps	39
3.5 Shows LiDAR derived data (A) Orthophotos, (B) DTM, (C) DSM, (D) Intensity, (E) Height, (F) Slope, (G) Aspect, and (H) Hillshade	42
3.6 Workflow of the Proposed FbSP Optimizer	46
3.7 Flowchart of the Proposed Methodology	48
3.8 ACO-Based Feature Selection Workflow	51
3.9 The Flowchart Designed for this research	59
3.10 An Overview of the Method Adopted in the Present Study	61
3.11 The Flowchart Illustrates the Overview of the Proposed Method	63
3.12 Architecture of the MLR-NN Model	65
3.13 Architecture of the RNN Model	66
4.1 Shows the Result of the Segmentation using Optimized Parameters (when the Scale is 75.52) for the Analysis Area. It can be Seen that the Landslide Objects were Accurately Delineated Highlighted by Red Color	71
4.2 Shows the Results of Support Vector Machine (A) Analysis area (B) Test site	77
4.3 Shows the Results of Random Forest, (A) Analysis Area, (B) Test Sit	78
4.4 Landslides Locations in the Study Area	82
4.5 The Segmentation Process and Optimal Segmentation using FbSP Optimizer for the Location of Landslides. (A) And (B) Represent the Initial Segmentation, while (C) Represent the Optimized Segmentation	83

4.6	[A] Result of RF Classifier in the Analysis Area, While [B] Presented The Inventory Map And [C] Showed how the Proposed Method Differentiated between two classes (Shallow And Deep Seated)	88
4.7	Illustrates Sketch of Shallow Landslide in (Yellow Polygon) And Deep Seated Landslide in (White Polygon) in Cameron Highland	89
4.8	Result of Transferability Model Showed the Locations of Shallow and Deep Seated Landslide in the Test Site	91
4.9	Field Photos Showing the Location of Landslide During field Investigation	95
4.10	Shows the process of segmentation, (A) Initial Segmentation (left) and, (B) Optimal Segmentation (right)	96
4.11	Analysis Area for Detection the Location of the Landslide	98
4.12	Show the Results of Transferability to whole Study Area	99
4.13	Show the Location of the Shallow Landslides Represent in (Red polygon), Deep-Seated Landslides Represent in (Yellow polygon)	101
4.14	ROC Curve for, (A) Landslide Detection and their Characteristics, (B) Shallow Landslide, (C) Deep Seated Landslide	104
4.15	Landslide Photos Taken in Study Area During Field Investigation	105
4.16	Shows the Process of Optimization Segmentation, (A) Initial Segmentation and, (B) Optimized Segmentation	106
4.17	Illustrates the Structure of Different Types of Soil	108
4.18	Results of Hierarchal Rules Set Classification at Site “A”	110
4.19	Show the Digital Number (DN) Values of, (A) Texture, (B) Intensity and, (C) Average of Visible Bands, which they Contributed in Distinguish between the Classes (Shallow, Deep Seated, Cut-Slope and Bare Soil)	112
4.20	Results of Hierarchal Rules Set Classification at Site “B”	113
4.21	Shows that the Amount of Intensity Value Involved in Landslide	114
4.22	Shows the Field Photos in Some Locations of Landslide at (A) Taman Desa Corina, Kampung Raja (B) Tanah-Rata	117

4.23	Parameter Optimization of the Multiresolution Segmentation Algorithm: (A) Initial Segmentation and (B) Optimized Segmentation	118
4.24	Results of the Qualitative Assessment of (A) RNN and (B) MLP-NN For the Analysis Area	120
4.25	Results of the Qualitative Assessment of (A) RNN And (B) MLP-NN for the Test site	120
4.26	Impact of the Optimization Algorithm on the Performance of MLP-NN and RNN Models	122
4.27	Impact of Batch Size on the Performance of The MLP-NN and RNN Models	123
4.28	Influence of Dropout Rate on the Performance of the RNN Model	124
4.29	Field Photographs Showing Landslide Locations during Field Investigation in (A) Tanah Rata (101°23'24.89"E and 4°26'24.33"N), and (B) Ringlet (101°22'55.37"E and 4°24'45.47"N)	124

## LIST OF ABBREVIATIONS

ACO	Ant Colony Optimization
AUC	The area under curve
CFS	Correlation-based Feature Selection
DEM	Digital Elevation Model
DNN	Deep Neural Network
CWT	Continuous Wavelet Transform
DSM	Digital surface model
DN	Digital Number
EC	Evolutionary Computation
DT	Decision Tree
RBF	Radial Basis Function
SAR	Synthetic Aperture Radar
GA	Genetic Algorithm
GIS	Geographic information system
GR	Gain ratio
LiDAR	Light Detection and Ranging
MLP	Multi-Layer Perceptron
nDSM	Normalized Digital Surface Model
OBIA	Object-based Image Analysis
PSO	Particle Swarm Optimization
RF	Random Forest
RNN	Recurrent Neural Networks
ROC	Relative Operating Characteristic
RS	Remote Sensing
SPOT	Satellite Pour l'Observation de la Terre
SVM	Support Vector Machine
DM	Data Mining
VHR	Very High Resolution
OOB	“Out-of-Bag
LULC	Land Use /Cover
BPNN	Back-Propagation neural network

PNN	Probability Neural Network
UA	User Accuracy
PA	Producer Accuracy
OA	Overall Accuracy
Cor	Correlation
Stdev	Standard Deviation
GLCM	Gray-Level Co-occurrence Matrix
GLDV	Grey level difference vector
TP	True Positives,
TN	True Negatives
FP	False Positives
FN	False negatives



## CHAPTER 1

### INTRODUCTION

#### 1.1 Introduction

The rapid expansion of cities and the continuously increasing population in urban areas lead to the establishment of settlements in mountainous areas. This phenomenon has increased the impact of natural disasters, particularly landslides, in these mountainous areas. Landslides result in severe property losses, human casualties, and environmental damage. Landslides are considered one of the geological phenomena – along with earthquakes, subsidence, volcanic activity and glacial rebound – to contribute to ground deformation (Pradhan and Lee, 2010). In the near future, climate change induced by anthropogenic actions such as urban expansion and deforestation, will lead to higher landslide occurrences. Considering the wide coverage of landslide damages, planners and decision makers have to identify location of landslide to mitigate the event (Pradhan et al., 2011). Two major aspects should be considered to produce an efficient landslide inventory map. One aspect is related to the data used, whereas the other is related to the methods employed (Pradhan et al., 2016).

Landslides are geological disasters with catastrophic effects on human lives and properties. Landslide incidence is related to a cluster of triggering factors, such as intense rainfall, volcanic eruptions, rapid snowmelt, elevated water levels, and earthquakes (Tehrany et al., 2014a). Landslide inventory maps are required for various purposes such as: (i) recording the magnitude of landslide in a region, (ii) performing the initial steps in analysing the susceptibility, hazard, and risk of the landslide (Van Westen et al., 2008; Guzzetti et al., 2012; Althuwaynee et al., 2014a; Althuwaynee et al., 2014b; Pradhan et al., 2016), and (iii) examining the patterns of landslide distributions and studying the evolution of landscape affected by landslides (Parker et al., 2011; Pradhan et al., 2016). Mapping of landslide susceptibility, hazards, and risks is imperative to achieve the mitigation planning (Pradhan and Buchroithner, 2010). The identification and mitigation of landslide occurrences are also crucial within the field of hazard management and mitigation research (Guzzetti et al., 2012).

According to Metternicht et al. (2005), remote sensing (RS) technology provides detail information about landslides for policy-makers and emergency managers. Selecting relevant features is important for distinguishing between landslides and non-landslides as well as classifying landslides (Van Westen et al., 2008). Mapping a landslide inventory in tropical areas is challenging because the dense vegetation cover in these regions obscures underlying landforms (Chen et al., 2015). The precision of derived maps depends not only on the methodology adopted but also on the quality of the features selection. The use of a large number of features can decrease the accuracy of landslide detection because of the presence of irrelevant features (Stumpf and Kerle, 2011). Moreover, the dense vegetation of tropical forests complicates the generation

of an accurate landslide inventory map in these regions. Therefore, the present study aims to utilize light detection and ranging (LiDAR) data and orthophoto image along with recent optimized techniques, in detecting landslides and differentiating between their characteristics.

## **1.2 Problem Statement**

Field survey, optical remote sensing, and synthetic aperture radar techniques are traditional techniques use for landslide detection in Remote Sensing and Geographic Information System (GIS) (Pradhan et al., 2016). However, there exist some challenges that affect its accuracy and performance. In a dense vegetation cover, it is difficult to use the traditional technique due to the obstruction from mountains and shadows (Chen et al., 2014). Many researches have attempted to develop rule-set based on time trial-error method. However, the rule-set development with the trial-and-error optimization method is time consuming, and the optimum rulesets are difficult to be determined (Sameen et al., 2017). Also, many investigations have been carried out using hierarchical approach with multiresolution sets of images. However, this approach has not been adopted in differentiating between types of landslide (shallow and deep seated) using LiDAR data. Despite the viability of Neural Architecture techniques in many areas of endeavour such as GIS, the technique has not been deployed with LiDAR data in detecting landslide.

## **1.3 Motivation Behind this Research**

Nowadays, natural hazards are common in today's life. Increasing amounts of natural catastrophes have proved to the human the vital importance of the natural hazards issues for the safety of the environment, and the populations. Rapid urbanization and climate change are expected to raise the amount of landslide. Landslide results in sever property losses, human casualties and environmental damage. The dramatic landslides which occur in tropical countries, especially Malaysia, emphasize the extremity in climatic variations. Since large-scale landslides often cause severe damage to human lives and properties, it is desirable to identify and map them for hazard mitigation purposes (Lin et al., 2013a). This phenomenon triggers due to the unexpected variations of the state of features due to natural forces, such as rainfall, and human activities. In most cases, human beings are not capable of controlling and predicting these disasters precisely. When major natural catastrophes such as landslides, earthquakes, floods and land subsidence occur, they have a definite impact upon human lives, belongings, infrastructure, farming and environment. The influence of natural hazards varies based on the intensity and coverage of region.

Landslides are of the most commonly occurring natural catastrophes that influence human beings and their surrounding environment. Inhabitants of Asia and the Pacific regions are affected in terms of social and economic stability. According to Pradhan (2010), high percentage of the destructions related to natural catastrophes in Malaysia are produced by landslide. Furthermore, average annual landslide damage is as high

as \$10 million USD. The attention towards providing proper landslide management has risen over the last few centuries. The recent reasons for recurrent landslides within some regions are mostly due to unplanned urbanization, construction and deforestation. In spite of these factors, it is possible to predict and mitigate effects of landslides with the use of various technology. The use of technology can facilitate landslide prevention actions to detect the landslide areas and to have an early warning for this catastrophe.

This thesis attempts to propose novel techniques for mapping the landslide in vulnerable locations using state-of-the-art machine learning methods. The key motivation of this research is to establish high-quality landslide inventory maps by applying semi-automatic techniques. To recognize those susceptible regions, landslide inventory maps should be generated as a basis of landslide susceptibility mapping in order to avoid more urbanization in hazardous areas. To reduce the damage and victims in case of a landslide occurrence, it is critical to locate the susceptible areas. Additionally, optimized techniques vastly improve the accuracy of classifications. Governments and planners can utilize the generated results of this study to identify safe regions for inhabitants, support the first ones to respond to emergencies, and update the urban planning strategies. Such data can decrease the requirements for performing field surveys by agencies such as departments of surveying.

#### **1.4 Research Objectives**

The present thesis proposes and applies various new methods that clearly contribute to the gap in the literature. These methods are simple, repeatable, and comprehensive. The following are the main objectives of the thesis:

1. To investigate the suitable selection features algorithms in landslide detection.
2. To optimize rule sets to distinguish various landslide characteristics and compare its performance with hierarchical rule-set classification approach.
3. To apply deep learning (Neural Architecture) classifier for landslide mapping.

#### **1.5 Research Questions**

This thesis comprehensively addresses the following research questions:

1. What are the results of the selected feature techniques using high resolution LiDAR data? How do the selected relevant features contribute to detection of landslides and differentiation of two landslide types in a highly dense, vegetated area?

2. What is the result of the optimize rule data with high density LiDAR data? How can this be helpful in detecting landslides and differentiating between two types of landslide?
3. How effective is the supervised approach classification method in defining the landslide prone areas?
4. How do the LiDAR derived data aid in differentiation of deep seated and shallow landslides?
5. How does the optimized segmentation aid in delineating the boundary of landslides and improve in its exploitation?
6. How do the decision rule sets aid to distinguish between two types of landslides?
7. How do the hierarchal rules set contribute to differentiation of two types of landslides?
8. How do deep neural networks help in detecting landslides within tropical areas?

## **1.6 Scope of this Thesis**

This study is carried out to develop a method for detecting and distinguish landslide areas using remote sensing and GIS techniques in a tropical environment like Cameron Highland, Malaysia. Also, transferability capability to similar geological and geomorphological areas are investigated and tested in other study areas. In this research, airborne laser scanning techniques is used to detect landslides and to distinguish between shallow and deep-seated landslides. Six (6) algorithms; Ant Colony Optimization (ACO), Gain Ratio (GR), Particle Swarm Optimization (PSO) and Genetic Algorithm (GA), Random forest (RF), and Correlation-based Feature Selection (CFS) are used for the feature selection. The performance of the feature selection algorithm is evaluated using two Machine learning classifiers, namely support vector machine (SVM) and random forest (RF). Decision rule and hierarchal rules set are developed by applying decision tree (DT) to provide improved landslide inventory map. In this task, two neural network algorithms, Recurrent Neural Networks (RNN) and Multi-Layer Perceptron Neural Networks (MLP-NN) are used and the hyper-parameters of the network architecture was optimized based on a systematic grid search.

## **1.7 Research Hypothesis**

1. The optimized techniques for both segmentation parameters and feature selection may improve and facilitate the generation of landslide inventory map with high density LiDAR data.
2. Decision rule and hierarchal rule sets along with relevant features may improve significantly in differentiating between the landslide types.
3. Deep neural network may contribute in improving accuracy of a landslide inventory map.

## **1.8 Thesis Organization**

This thesis is organized into five chapters. The summary of each chapter is as follows:

### **(i) CHAPTER 1: INTRODUCTION**

This chapter introduces the problem statement of the study, aim, objectives and scope of the study. Furthermore, this chapter highlights the research questions proposed for the thesis as well as the significant contribution of new knowledge and the overall structure of the thesis.

### **(ii) CHAPTER 2: LITERATURE REVIEW**

This chapter provides an overview of landslide status in various regions and previous work of using remote sensing and GIS to detect landslides. The two types of landslide detection techniques were outlined followed by the traditional and innovative and emerging techniques for detecting landslide prone areas. Next, the methodology used to detect and distinguish landslides using different techniques were discussed. Finally, validation methods used to assess the accuracy of maps produced are summarized.

### **(iii) CHAPTER 3: MATERIALS AND METHODOLOGY**

This chapter details the characteristics of the study area followed by the following details in order: materials, data, methodology, optimized segmentation, evaluation of the training sample, features selection algorithms, detection, differentiation of the types of landslide (i.e. deep seated and shallow), and various remote sensing techniques.

### **(iv) CHAPTER 4: RESULTS AND DISCUSSION**

This chapter focuses on the results of the study, which include the analysis results, optimized segmentation, and selected feature algorithms using LiDAR data – supported by photos, tables, diagram and charts. This chapter also discusses the results of optimized segmentation. Afterwards, decision rule and hierarchal rules were used in differentiating between two types landslides with regards to their characteristics. Lastly, the results of the deep neural network – using only the LiDAR derived data – were employed for detecting landslides and differentiating between the two types of landslides are discussed.

### **(v) CHAPTER 5: CONCLUSION AND FUTURE WORK RECOMMENDATIONS**

This chapter provides the overall conclusion from this study, recommendation and further research for the study area.



## REFERENCES

- Abbaspour, K. C., Schulin, R., Van Genuchten, M. T. (2001). Estimating unsaturated soil hydraulic parameters using ant colony optimization. *Advances in Water Resources* 24(8), 827-841.
- Abe, S. (2005) Support vector machines for pattern classification (Vol. 53) London: Springer.
- Agliardi, F., Crosta, G. B., Zanchi, A., Ravazzi, C. (2009). Onset and timing of deep-seated gravitational slope deformations in the eastern Alps, Italy. *Geomorphology*, 103(1), 113-129.
- Aiello, M., Gianinetto, M. (2017). A combined use of multispectral and SAR images for ship detection and characterization through object based image analysis. In *Earth Resources and Environmental Remote Sensing/GIS Applications VIII* (Vol. 10428, p. 104280A). International Society for Optics and Photonics.
- Althuwaynee, O. F., Pradhan, B., Park, H. J., Lee, J. H. (2014a). A novel ensemble bivariate statistical evidential belief function with knowledge-based analytical hierarchy process and multivariate statistical logistic regression for landslide susceptibility mapping. *Catena*, 114, 21-36.
- Althuwaynee, O. F., Pradhan, B., Park, H. J., Lee, J. H. (2014b). A novel ensemble decision tree-based CHi-squared Automatic Interaction Detection (CHAID) and multivariate logistic regression models in landslide susceptibility mapping. *Landslides*, 11(6), 1063-1078.
- Alwan, H. B., Ku-Mahamud, K. R. (2012, December). Optimizing support vector machine parameters using continuous ant colony optimization. In *Computing and Convergence Technology (ICCCT) 2012 7th International Conference on* (pp. 164-169).
- Amodei, D., Ananthanarayanan, S., Anubhai, R., Bai, J., Battenberg, E., Case, C., Chen, J. (2016). Deep speech 2: End-to-end speech recognition in english and mandarin. In *International Conference on Machine Learning*, (pp. 173-182).
- Antonini, G., Ardizzone, F., Cardinali, M., Galli, M., Guzzetti, F., Reichenbach, P. (2002). Surface deposits and landslide inventory map of the area affected by the 1997 Umbria-Marche earthquakes. *Bollettino Della Società Geologica Italiana*, 121(1), 843-853.
- Ardizzone, F., Cardinali, M., Galli, M., Guzzetti, F., Reichenbach, P. (2007) Identification and mapping of recent rainfall-induced landslides using elevation data collected by airborne Lidar. *Natural Hazards and Earth System Science*, 7(6), 637– 650.

- Baatz, M., Schäpe, A. (1999). Object based and multi-scale image analysis in semantic networks. Paper presented at the 2nd international symposium: operationalization of remote sensing.
- Baczyński, D., Parol, M. (2004). Influence of artificial neural network structure on quality of short-term electric energy consumption forecast. *IEE Proceedings-Generation, Transmission and Distributio*, 151(2), 241-245.
- Bai, S., Wang, J., Zhang, Z., Cheng, C. (2012). Combined landslide susceptibility mapping after Wenchuan earthquake at the Zhouqu segment in the Bailongjiang Basin, China. *Catena* 99, 18-25.
- Bajracharya, B., Bajracharya, S. R. (2008, January). Landslide mapping of the Everest region using high resolution satellite images and 3D visualization. In *Proceedings of the Mountain GIS e-Conference, Kathmandu, Nepal* (pp. 14-25).
- Barbarella, M., Fiani, M., Lugli, A. (2013). Application of LiDAR-derived DEM for detection of mass movements on a landslide. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences* (5/W3), 89 –98.
- Barlow, J., Martin, Y., Franklin, S. E. (2003). Detecting translational landslide scars using segmentation of Landsat ETM+ and DEM data in the northern Cascade Mountains, British Columbia. *Canadian journal of remote sensing* 29(4), 510-517.
- Bartels, M., Wei, H. (2010). Threshold-free object and ground point separation in LIDAR data. *Pattern Recognition Letters* 31(10), 1089-1099.
- Belgiu, M., Drăguț, L. (2014). Comparing supervised and unsupervised multiresolution segmentation approaches for extracting buildings from very high resolution imagery. *ISPRS Journal of Photogrammetry and Remote Sensing* 96, 67–75.
- Benediktsson, J. A., Sveinsson, J. R. (1997). Feature extraction for multisource data classification with artificial neural networks. *International Journal of Remote Sensing* 18(4), 727-740.
- Benz, U.C., Hofmann, P., Willhauck, G., Lingenfelder, I., Heynen, M. (2004). Multi-resolution, object based fuzzy analysis of remote sensing data for GIS-ready information. *ISPRS Journal of photogrammetry and Remote Sensing* 58(3), 239–258.
- Bhaduri, K., Wolff, R., Giannella, C., Kargupta, H. (2008). Distributed decision-tree induction in peer-to-peer systems. *Statistical Analysis and Data Mining: The ASA Data Science Journal* 1(2), 85-103.

- Bibi, T., Razak, K. A., Rahman, A. A., Latif, A. (2017). Spatio temporal detection and virtual mapping of landslide using high-resolution airborne laser altimetry (lidar) in densely vegetated areas of tropics. *International Archives of the Photogrammetry, Remote Sensing & Spatial Information Sciences*, 42.
- Bignel, F., Snelling, G. (1977). The geochronology of the main range Batholith: Cameron highlands road and Gunong Bujang Melaka. *Overseas Geol Miner Resour*, 47, 3–35.
- Bishop, C.M. (1995). Neural networks for pattern recognition. Oxford university press.
- Blaschke, T., Hay, G.J., Kelly, M., Lang, S., Hofmann, P., Addink, E., Tiede, D. (2014). Geographic object-based image analysis—towards a new paradigm. *ISPRS Journal of Photogrammetry and Remote Sensing* 87, 180–191.
- Blaschke, T. (2010). Object based image analysis for remote sensing. *ISPRS Journal of Photogrammetry and Remote Sensing* 65(1), 2–16.
- Blonda, P., Bennardo, A., Satalino, G., Pasquariello, G. (1996). Fuzzy logic and neural techniques integration: an application to remotely sensed data. *Pattern Recognition letters* 17(13), 1343–1348.
- Boiman, O., Shechtman, E., Irani, M. (2008, June). In defense of nearest-neighbor based image classification. In *Computer Vision and Pattern Recognition, 2008. CVPR 2008. IEEE Conference on*, (pp. 1–8). IEEE.
- Booth, A.M., Roering, J. J., Perron, J. T. (2009). Automated landslide mapping using spectral analysis and high-resolution topographic data: Puget Sound lowlands, Washington, and Portland Hills, Oregon. *Geomorphology* 109(3), 132–147.
- Borghuis, A., Chang, K., Lee, H. (2007). Comparison between automated and manual mapping of typhoon-triggered landslides from SPOT-5 imagery. *International Journal of Remote Sensing* 28(8), 1843–1856.
- Breiman, L.F., Olshen, J H RA, Stone, C.J. (1984). Classification and Regression Trees, Belmont, CA: Wadsworth.
- Breiman, L. (2001). Random forests. *Machine learning* 45(1), 5–32.
- Brunetti, M. T., Guzzetti, F., Rossi, M. (2009). Probability distributions of landslide volumes. *Nonlinear Processes in Geophysics* 16(2), 179.
- Bugnion, L., Volkwein, A., Denk, M. (2009). Artificial full scale shallow landslides. In *EGU General Assembly Conference Abstracts* (Vol. 11, p. 13064).



- Bui, D.T., Pradhan, B., Lofman, O., Revhaug, I., Dick, O.B. (2012). Spatial prediction of landslide hazards in Hoa Binh province (Vietnam): a comparative assessment of the efficacy of evidential belief functions and fuzzy logic models. *Catena* 96, 28-40.
- Bui, D. T., Tuan, T. A., Klempe, H., Pradhan, B., Revhaug, I. (2016). Spatial prediction models for shallow landslide hazards: a comparative assessment of the efficacy of support vector machines, artificial neural networks, kernel logistic regression, and logistic model tree. *Landslides*, 13(2), 361-378.
- Cao, L.Y., Xia, L. Z. (2007). An ant colony optimization approach for SAR image segmentation. In *Wavelet Analysis and Pattern Recognition, 2007. ICWAPR'07. International Conference on* (Vol. 1, pp. 296-300). IEEE.
- Chang, K. T., Hwang, J. T., Liu, J. K., Wang, E. H., Wang, C. I. (2011). Apply two hybrid methods on the rainfall-induced landslides interpretation. In *Geoinformatics, 2011 19th International Conference on* (pp. 1-5).
- Chang, K. T., Liu, J. K., Chang, Y.M., Kao, C. S. (2010). An accuracy comparison for the landslide inventory with the BPNN and SVM methods. In *Proc. Gi4DM 2010*.
- Chang, K. T., Liu, J. K., Wang, C. I. (2012). An object based analysis for characterizing the rainfall-induced shallow landslide. *Journal of Marine Science and Technology* 20(6), 647-656.
- Chaplot, S., Patnaik, L. M., Jagannathan, N. R. (2006). Classification of magnetic resonance brain images using wavelets as input to support vector machine and neural network. *Biomedical Signal Processing and Control* 1(1), 86-92.
- Chen, C.H. (2012). *Signal and image processing for remote sensing*. CRC press.
- Chen, H., Zeng, Z., Tang, H. (2015). Landslide deformation prediction based on recurrent neural network. *Neural Processing Letters* 41(2), 169-178.
- Chen, Q., Chen, Y., Jiang, W. (2016). Genetic Particle Swarm Optimization–Based Feature Selection for Very-High-Resolution Remotely Sensed Imagery Object Change Detection. *Sensors* 16(8), 1204.
- Chen, R. F., Lin, C.W., Chen, Y. H., He, T. C., Fei, L.Y. (2015). Detecting and Characterizing Active Thrust Fault and Deep-Seated Landslides in Dense Forest Areas of Southern Taiwan Using Airborne LiDAR DEM. *Remote Sensing* 7(11), 15443-15466.
- Chen, T., Trinder, J. C., Niu, R. (2017). Object based Landslide Mapping Using ZY-3 Satellite Imagery, Random Forest and Mathematical Morphology, for the Three-Gorges Reservoir, China. *Remote Sensing* 9(4), 333.

- Chen, W., Li, X., Wang, Y., Chen, G., Liu, S. (2014). Forested landslide detection using LiDAR data and the random forest algorithm: A case study of the Three Gorges, China. *Remote Sensing of Environment*, 152, 291-301.
- Chen, Y., Su, W., Li, J., Sun, Z. (2009). Hierarchical object based classification using very high resolution imagery and LIDAR data over urban areas. *Advances in Space Research*, 43(7), 1101-1110.
- Chow, W. S., Zakaria, M., Ferdaus, A., Nurzaidi, A. (2003). Geological terrain mapping. *JMG unpublished report, JMG. SWP. GS*, 1-42.
- Clague, J. J., Stead, D. (2012). Landslides: types, mechanisms and modeling. Cambridge University Press.
- Cobbing, E., Pitfield, P., Darbyshire, D., Mallick, D. (1992). The granites of the SE Asian tin belt. British Geological Survey, Overseas Memoir No. 10: HMSO, London.
- Congalton, R.G. (1991). A review of assessing the accuracy of classifications of remotely sensed data. *Remote sensing of environment*, 37(1), 35-46.
- Cortes, C., Vapnik, V. (1995). Support-vector networks. *Machine learning*, 20(3), 273-297.
- Costa, H., Foody, G. M., Boyd, D. S. (2018). Supervised methods of image segmentation accuracy assessment in land cover mapping. *Remote Sensing of Environment*, 205, 338-351.
- Cruden, D. M., Varnes, D. J. (1996). Landslides: investigation and mitigation. Chapter 3-Landslide types and processes. *Transportation Research Board Special Report*, (247).
- Czuchlewski, K. R., Weissel, J. K., Kim, Y. (2003). Polarimetric synthetic aperture radar study of the Tsaoling landslide generated by the 1999 Chi-Chi earthquake, Taiwan. *Journal of Geophysical Research: Earth Surface*, 108(F1).
- Daniel, S. (2014). Predictive modeling of trust to Social Media content (Master's thesis, University of Stavanger, Norway).
- Danneels, G., Pirard, E., Havenith, H. B. (2007). Automatic landslide detection from remote sensing images using supervised classification methods. *Geoscience and Remote Sensing Symposium* 3014–3017.
- Darwish, A., Leukert, K., Reinhardt, W. (2003). Image segmentation for the purpose of object-based classification. In *Geoscience and Remote Sensing Symposium, 2003. IGARSS'03. Proceedings. 2003 IEEE International* (Vol. 3, pp. 2039-2041). IEEE.

- De Blasio, F.V. (2011). Landslides in Valles Marineris (Mars): a possible role of basal lubrication by sub-surface ice. *Planetary and Space Science* 59(13), 1384–1392.
- Delgado, J., Vicente, F., García-Tortosa, F., Alfaro, P., Estevez, A., Lopez-Sanchez, J. M., Mallorquí, J.J. (2011). A deep seated compound rotational rock slide and rock spread in SE Spain: structural control and DInSAR monitoring. *Geomorphology*, 129(3), 252-262.
- Deng, L. (2014). A tutorial survey of architectures, algorithms, and applications for deep learning. *APSIPA Transactions on Signal and Information Processing*, 3.
- Derron, M. H., Jaboyedoff, M. (2010). Preface" LIDAR and DEM techniques for landslides monitoring and characterization". *Natural Hazards and Earth System Science* 10(9), 1877-1879.
- Dey, V., Zhang, Y., Zhong, M. (2010). A review on image segmentation techniques with remote sensing perspective (pp. 31-42). na.
- Díaz-Uriarte, R., De Andres, S.A. (2006). Gene selection and classification of microarray data using random forest. *BMC bioinformatics*, 7(1), 3.
- Dibs, H., Idrees, M. O., Alsalhin, G. B. A. (2017). Hierarchical classification approach for mapping rubber tree growth using per-pixel and object based classifiers with SPOT-5 imagery. *The Egyptian Journal of Remote Sensing and Space Science* 20(1), 21-30.
- Domlija, P., Bernat, S., Mihalić, S.A., Benac, Č. (2014). Landslide inventory in the area of Dubračina River Basin (Croatia). In *Landslide science for a safer geoenvironment*(pp. 837-842). Springer, Cham.
- Donati, L., Turrini, M.C. (2002). An objective method to rank the importance of the factors predisposing to landslides with the GIS methodology: application to an area of the Apennines (Valnerina; Perugia, Italy). *Engineering Geology*, 63(3), 277-289.
- Dópido, I., Li, J., Marpu, P. R., Plaza, A., Dias, J. M. B., Benediktsson, J.A. (2013). Semisupervised self-learning for hyperspectral image classification. *IEEE Transactions on Geoscience and Remote Sensing*, 51(7), 4032-4044.
- Dorigo, M., Birattari, M., Stutzle, T. (2006). Ant colony optimization. *IEEE Computational Intelligence Magazine*, 1(4), 28-39.
- Dorigo, M., Stutzle, T. (2003). The ant colony optimization metaheuristic: Algorithms, applications, and advances. *International Series in Operations Research and Management Science*, 251-286.

- Dou, J., Chang, K.-T., Chen, S., Yunus, A.P., Liu, J. K., Xia, H., Zhu, Z. (2015). Automatic case-based reasoning approach for landslide detection: integration of object based image analysis and a genetic algorithm. *Remote Sensing* 7(4), 4318–4342.
- Drăguț, L., Tiede, D., Levick, S. R. (2010). a tool to estimate scale parameter for multiresolution image segmentation of remotely sensed data. *International Journal of Geographical Information Science* 24(6), 859–871.
- Duro, D. C., Franklin, S. E., Dubé, M. G. (2012). Multi-scale object-based image analysis and feature selection of multi-sensor earth observation imagery using random forests. *International Journal of Remote Sensing* 33(14), 4502–4526.
- Eeckhaut, M., Poesen, J., Verstraeten, G., Vanacker, V., Nyssen, J., Moeyersons, J., Vandekerckhove, L. (2007). Use of LIDAR-derived images for mapping old landslides under forest. *Earth Surface Processes and Landforms*, 32(5), 754–769.
- Evans, J. S., Hudak, A. T. (2007). A multiscale curvature algorithm for classifying discrete return LiDAR in forested environments. *IEEE Transactions on Geoscience and Remote Sensing*, 45(4), 1029–1038.
- Fawcett, T. (2006). An introduction to ROC analysis. *Pattern recognition letters*, 27(8), 861–874.
- Feizizadeh, B., Blaschke, T., Tiede, D., Moghaddam, M. H. R. (2017). Evaluating fuzzy operators of an object-based image analysis for detecting landslides and their changes. *Geomorphology*, 293, 240–254.
- Fiorucci, F., Cardinali, M., Carlà, R., Rossi, M., Mondini, A., Santurri, L., Guzzetti, F. (2011). Seasonal landslide mapping and estimation of landslide mobilization rates using aerial and satellite images. *Geomorphology* 129(1), 59–70.
- Foody, G. M., Mathur, A. (2004). A relative evaluation of multiclass image classification by support vector machines. *IEEE Transactions on Geoscience and Remote Sensing* 42(6), 1335–1343.
- Fu, G., Liu, C., Zhou, R., Sun, T., Zhang, Q. (2017). Classification for High Resolution Remote Sensing Imagery Using a Fully Convolutional Network. *Remote Sensing* 9(5), 498.
- Galli, M., Ardizzone, F., Cardinali, M., Guzzetti, F., Reichenbach, P. (2008). Comparing landslide inventory maps. *Geomorphology* 94(3), 268–289.
- Gamanya, R., De Maeyer, P., De Dapper, M. (2009). Object based change detection for the city of Harare, Zimbabwe. *Expert Systems with Applications*, 36(1), 571–588.

- Gao, J., Maro, J. (2010). Topographic controls on evolution of shallow landslides in pastoral Wairarapa, New Zealand, 1979–2003. *Geomorphology* 114(3), 373-381.
- Gao, T., Zhu, J., Zheng, X., Shang, G., Huang, L., Wu, S. (2015). Mapping spatial distribution of larch plantations from multi-seasonal Landsat-8 OLI imagery and multi-scale textures using random forests. *Remote Sensing* 7(2), 1702-1720.
- Geertsema, M., Blais-Stevens, A., Kwoil, E., Menounos, B., Venditti, J., Grenier, A., Wiebe, K. (2017). Sensitive clay landslide detection and characterization in and around Lakelse Lake, British Columbia, Canada. *Sedimentary Geology*.
- Georganos, S., Grippa, T., Vanhuysse, S., Lennert, M., Shimoni, M., Kalogirou, S., Wolff, E. (2018). Less is more: optimizing classification performance through feature selection in a very-high-resolution remote sensing object-based urban application. *GIScience & Remote Sensing*, 55(2), 221-242.
- Gardner, M.W., Dorling, S.R. (1998). Artificial neural networks (the multilayer perceptron)—a review of applications in the atmospheric sciences. *Atmospheric Environment* 32(14), 2627-2636.
- Gislason, P.O., Benediktsson, J.A., Sveinsson, J. R. (2006). Random forests for land cover classification. *Pattern Recognition Letters*, 27(4), 294-300.
- Goldberg, D. (1989). Genetic algorithms in search, optimization and learning. Addison-Wesley Pub. Co.
- Gorsevski, P. V., Brown, M. K., Panter, K., Onasch, C. M., Simic, A., Snyder, J. (2016). Landslide detection and susceptibility mapping using LiDAR and an artificial neural network approach: a case study in the Cuyahoga Valley National Park, Ohio. *Landslides*, 13(3), 467-484.
- Gottlieb, J., Puchta, M., Solnon, C. (2003). A study of greedy, local search, and ant colony optimization approaches for car sequencing problems. In *EvoWorkshops* (Vol. 2611, pp. 246-257).
- Goudie, A. (Ed.). (2004). Encyclopedia of geomorphology (Vol. 2). Psychology Press.
- Grimm, R., Behrens, T., Märker, M., Elsenbeer, H. (2008). Soil organic carbon concentrations and stocks on Barro Colorado Island digital soil mapping using Random Forests analysis. *Geoderma* 146(1), 102–113.
- Guo, Z., Zhang, L., Zhang, D. (2010). A completed modeling of local binary pattern operator for texture classification. *IEEE Transactions on Image Processing*, 19(6), 1657-1663.

- Guzzetti, F., Cardinali, M., Reichenbach, P., Carrara, A. (2000). Comparing landslide maps: A case study in the upper Tiber River Basin, central Italy. *Environmental Management*, 25(3), 247-263.
- Guzzetti, F., Galli, M., Reichenbach, P., Ardizzone, F., Cardinali, M. (2006). Landslide hazard assessment in the Collazzone area, Umbria, Central Italy. *Natural Hazards and Earth System Science*, 6(1), 115-131.
- Guzzetti, F., Mondini, A.C., Cardinali, M., Fiorucci, F., Santangelo, M., Chang, K.-T. (2012). Landslide inventory maps: New tools for an old problem. *Earth-Science Reviews* 112(1), 42-66.
- Hall, M. A. (1999). Correlation-based feature selection for machine learning.
- Hall, M. A., Holmes, G. (2003). Benchmarking attribute selection techniques for discrete class data mining. *IEEE transactions on knowledge and data engineering*, 15(6), 1437-1447.
- Hamedianfar, A., Shafri, H. Z. M. (2016). Integrated approach using data mining-based decision tree and object-based image analysis for high-resolution urban mapping of WorldView-2 satellite sensor data. *Journal of Applied Remote Sensing*, 10(2), 025001-025001.
- Han, J., Pei, J., Kamber, M. Data Mining: Concepts and Techniques; Elsevier: Amsterdam, The Netherlands, 2011.
- Haneberg, W.C. (2005). The ins and outs of airborne LIDAR: An introduction for practicing engineering geologists. *AEG News*, 48, 16-19.
- Haneberg, W. C., Cole, W. F., Kasali, G. (2009). High-resolution lidar-based landslide hazard mapping and modeling, UCSF Parnassus Campus, San Francisco, USA. *Bulletin of engineering geology and the environment*, 68(2), 263-276.
- Hay, G. J., Castilla, G. (2008). Geographic Object-Based Image Analysis (GEOBIA): A new name for a new discipline. In *Object-based image analysis*, (pp. 75-89). Springer, Berlin, Heidelberg.
- Heleno, S., Lousada, M., Pereira, M.J., Pina, P. (2013, December). Automated Detection Of Small And Shallow Landslides After The 2010 Madeira Island Flash-Floods In VHR Imagery. In *ESA Living Planet Symposium* (Vol. 722, p. 285).
- Heleno, S., Matias, M., Pina, P., Sousa, A.J. (2016). Semiautomated object-based classification of rain-induced landslides with VHR multispectral images on Madeira Island. *Natural Hazards and Earth System Sciences*, 16, 1035-1048.



- Heleno, S., Matias, M., Pina, P., Sousa, A. J. (2015). Automated object-based classification of rain-induced landslides with VHR multispectral images in Madeira Island. *Natural Hazards and Earth System Sciences Discussions*, 3(9).
- Hinton, G., Osindero, S., Welling, M., Teh, Y. W. (2006). Unsupervised discovery of nonlinear structure using contrastive backpropagation. *Cognitive science*, 30(4), 725-731.
- Hochreiter, S., Schmidhuber, J. (1997). Long short-term memory. *Neural computation* 9(8), 1735-1780.
- Hofmann, P., Blaschke, T., Strobl, J. (2011). Quantifying the robustness of fuzzy rule sets in object-based image analysis. *International Journal of Remote Sensing*, 32(22), 7359-7381.
- Hölbling, D., Friedl, B., Eisank, C. (2015). An object-based approach for semi-automated landslide change detection and attribution of changes to landslide classes in northern Taiwan. *Earth Science Informatics*, 8(2), 327-335.
- Hong, H., Pradhan, B., Xu, C., Bui, D. T. (2015). Spatial prediction of landslide hazard at the Yihuang area (China) using two-class kernel logistic regression, alternating decision tree and support vector machines. *Catena*, 133, 266-281.
- Huang, S. S., Hormati, A., Bacon, D. F., Rabbah, R. M. (2008). Liquid Metal: Object based Programming Across the Hardware/Software Boundary. In *ECOOP* (Vol. 8, pp. 76-103).
- Im, J., Jensen, J. R., Tullis, J. A. (2008). Object-based change detection using correlation image analysis and image segmentation. *International Journal of Remote Sensing*, 29(2), 399-423.
- Iovine, G., D'Ambrosio, D., Di Gregorio, S. (2005). Applying genetic algorithms for calibrating a hexagonal cellular automata model for the simulation of debris flows characterised by strong inertial effects. *Geomorphology*, 66(1-4), 287-303.
- Jabari, S., Zhang, Y. (2013). Very high resolution satellite image classification using fuzzy rule-based systems. *Algorithms*, 6(4), 762-781.
- Jebur, M. N., Pradhan, B., Tehrany, M. S. (2014). Detection of vertical slope movement in highly vegetated tropical area of Gunung pass landslide, Malaysia, using L-band InSAR technique. *Geosciences Journal*, 18(1), 61-68.
- Johansen, K., Coops, N. C., Gergel, S. E., Stange, Y. (2007). Application of high spatial resolution satellite imagery for riparian and forest ecosystem classification. *Remote sensing of Environment*, 110(1), 29-44.

- Jozefowicz, R., Vinyals, O., Schuster, M., Shazeer, N., Wu, Y. (2016). Exploring the limits of language modeling. *arXiv preprint arXiv:1602.02410*.
- Kamps, M., Bouten, W., Seijmonsbergen, A. (2017). LiDAR and Orthophoto Synergy to optimize Object-Based Landscape Change: Analysis of an Active Landslide. *Remote Sensing*, 9(8), 805.
- Kanellopoulos, I., Wilkinson, G.G. (1997). Strategies and best practice for neural network image classification. *International Journal of Remote Sensing*, 18(4), 711-725.
- Karegowda, A. G., Manjunath, A. S., Jayaram, M. A. (2010). Comparative study of attribute selection using gain ratio and correlation based feature selection. *International Journal of Information Technology and Knowledge Management*, 2(2), 271-277.
- Kasai, M., Ikeda, M., Asahina, T., Fujisawa, K. (2009). LiDAR-derived DEM evaluation of deep-seated landslides in a steep and rocky region of Japan. *Geomorphology* 113(1), 57-69.
- Kellerer-Pirklbauer, A., Proske, H., Strasser, V. (2010). Paraglacial slope adjustment since the end of the Last Glacial Maximum and its long-lasting effects on secondary mass wasting processes: Hauser Kaibling,
- Kennedy, J., Eberhart, R. (1995). PSO optimization. In *Proc. IEEE Int. Conf. Neural Networks* (Vol. 4, pp. 1941-1948). IEEE Service Center, Piscataway, NJ.
- Keyport, R. N., Oommen, T., Martha, T. R., Sajinkumar, K. S., Gierke, J. S. (2018). A comparative analysis of pixel-and object-based detection of landslides from very high-resolution images. *International Journal of Applied Earth Observation and Geoinformation*, 64, 1-11.
- Korup, O. (2006). Effects of large deep-seated landslides on hillslope morphology, western Southern Alps, New Zealand. *Journal of Geophysical Research: Earth Surface*, 111(F1).
- Krause, G., Bock, M., Weiers, S., Braun, G. (2004). Mapping land-cover and mangrove structures with remote sensing techniques: A contribution to a synoptic GIS in support of coastal management in North Brazil. *Environmental Management*, 34(3), 429-440.
- Kumar, D. N., Reddy, M. J. (2006). Ant colony optimization for multi-purpose reservoir operation. *Water Resources Management* 20(6), 879-898.
- Kuo, B. C., Li, C. H., Yang, J. M. (2009). Kernel nonparametric weighted feature extraction for hyperspectral image classification. *IEEE Transactions on Geoscience and Remote Sensing*, 47(4), 1139-1155.



- Kursa, M. B., Rudnicki, W. R. (2010). Feature selection with the Boruta package. *Journal*.
- Kurtz, C., Passat, N., Gancarski, P., Puissant, A. (2012). Extraction of complex patterns from multiresolution remote sensing images: A hierarchical top-down methodology. *Pattern Recognition*, 45(2), 685-706.
- Kurtz, C., Stumpf, A., Malet, J. P., Gancarski, P., Puissant, A., Passat, N. (2014). Hierarchical extraction of landslides from multiresolution remotely sensed optical images. *ISPRS Journal of Photogrammetry and Remote Sensing*, 87, 122-136.
- Ladha, L., Deepa, T. (2011). Feature selection methods and algorithms. *International Journal on Computer Science and Engineering* 1(3), 1787–1797.
- Lakany, H., Conway, B. A. (2007). Understanding intention of movement from electroencephalograms. *Expert Systems*, 24(5), 295-304.
- Laliberte, A. S., Rango, A. (2009). Texture and scale in object-based analysis of subdecimeter resolution unmanned aerial vehicle (UAV) imagery. *IEEE Transactions on Geoscience and Remote Sensing*, 47(3), 761-770.
- Lawrence, R. L., Wood, S.D., Sheley, R. L. (2006). Mapping invasive plants using hyperspectral imagery and Breiman Cutler classifications (RandomForest). *Remote Sensing of Environment* 100(3), 356-362.
- Lee, S., Song, K. Y., Oh, H. J., Choi, J. (2012). Detection of landslides using web-based aerial photographs and landslide susceptibility mapping using geospatial analysis. *International Journal of Remote Sensing* 33(16), 4937–4966.
- Lefsky, M. A., Cohen, W. B., Spies, T. A. (2001). An evaluation of alternate remote sensing products for forest inventory, monitoring, and mapping of Douglas-fir forests in western Oregon. *Canadian Journal of Forest Research*, 31(1), 78-87.
- Lemp, D., Weidner, U. (2005, July). Segment-Based characterization of roof surfaces using hyperspectral and laser scanning data. In *Geoscience and Remote Sensing Symposium, 2005. IGARSS'05. Proceedings. 2005 IEEE International* (Vol. 7, pp. 4942-4945). IEEE.
- Li, L. L., Wang, J. K. (2012). SAR image ship detection based on ant colony optimization. In *Image and Signal Processing (CISP), 2012 5th International Congress on* (pp. 1100-1103). IEEE.
- Li, M., Ma, L., Blaschke, T., Cheng, L., Tiede, D. (2016). A systematic comparison of different object-based classification techniques using high spatial resolution imagery in agricultural environments. *International Journal of Applied Earth Observation and Geoinformation* 49, 87–98.

- Li, X., Cheng, X., Chen, W., Chen, G., Liu, S. (2015). Identification of forested landslides using LiDAR data, object-based image analysis, and machine learning algorithms. *Remote Sensing* 7(8), 9705–9726.
- Liaw, A., Wiener, M. (2002). Classification and regression by random Forest. *R news*, 2(3), 18–22.
- Lin, C. W., Tseng, C. M., Tseng, Y. H., Fei, L.Y., Hsieh, Y. C., Tarolli, P. (2013a). Recognition of large scale deep-seated landslides in forest areas of Taiwan using high resolution topography. *Journal of Asian Earth Sciences* 62, 389-400.
- Lin, M. L., Chen, T. W., Lin, C. W., Ho, D. J., Cheng, K. P., Yin, H. Y., Chen, M. C. (2013b). Detecting large-scale landslides using LiDAR data and aerial photos in the Namasha-Liuogwey area, Taiwan. *Remote Sensing*, 6(1), 42-63.
- Liu, Y., Zhang, H. H., Wu, Y. (2011). Hard or soft classification? Large-margin unified machines. *Journal of the American statistical association* 106(493), 166-177.
- Lu, J., Xiong, C., Parikh, D., Socher, R. (2016). Knowing when to look: Adaptive attention via A visual sentinel for image captioning. *arXiv preprint arXiv:1612.01887*.
- Lu, P., Stumpf, A., Kerle, N., Casagli, N. (2011). Object based change detection for landslide rapid mapping. *IEEE Geoscience and remote sensing letters*, 8(4), 701-705.
- Lyu, H., Lu, H., Mou, L. (2016). Learning a transferable change rule from a recurrent neural network for land cover change detection. *Remote Sensing*, 8(6), 506.
- Ma, H.R., Cheng, X., Chen, L., Zhang, H., Xiong, H. (2016). Automatic identification of shallow landslides based on Worldview2 remote sensing images. *Journal of Applied Remote Sensing* 10(1), 016008-016008.
- Ma, L., Cheng, L., Li, M., Liu, Y., Ma, X. (2015). Training set size, scale, and features in Geographic Object-Based Image Analysis of very high resolution unmanned aerial vehicle imagery. *ISPRS Journal of Photogrammetry and Remote Sensing*, 102, 14-27.
- Ma, L., Fu, T., Blaschke, T., Li, M., Tiede, D., Zhou, Z., Chen, D. (2017). Evaluation of Feature Selection Methods for Object-Based Land Cover Mapping of Unmanned Aerial Vehicle Imagery Using Random Forest and Support Vector Machine Classifiers. *ISPRS International Journal of Geo-Information* 6(2), 51.
- Magee, K.S. (2011). Segmentation, object based applications for remote sensing land cover and land use classification. University of Cincinnati.

- Malamud, B. D., Turcotte, D. L., Guzzetti, F., Reichenbach, P. (2004). Landslide inventories and their statistical properties. *Earth Surface Processes and Landforms* 29(6), 687-711.
- Mantas, C.J., Abellán, J. (2014). Credal-C4. 5: Decision tree based on imprecise probabilities to classify noisy data. *Expert Systems with Applications* 41(10), 4625-4637.
- Marc, O., Hovius, N. (2015). Amalgamation in landslide maps: effects and automatic detection. *Natural Hazards and Earth System Sciences* 15(4), 723.
- Martha, T. R., Kerle, N., Jetten, V., van Westen, C. J., Kumar, K. V. (2010). Characterising spectral, spatial and morphometric properties of landslides for semi-automatic detection using object based methods. *Geomorphology* 116(1), 24-36.
- Martha, T.R., Kerle, N., Van Westen, C.J., Jetten, V., Kumar, K.V. (2011). Segment optimization and data-driven thresholding for knowledge-based landslide detection by object-based image analysis. *IEEE Transactions on Geoscience and Remote Sensing* 49(12), 4928–4943.
- Matinfar, H. R., Panah, S. K. A., Zand, F., Khodaei, K. (2013). Detection of soil salinity changes and mapping land cover types based upon remotely sensed data. *Arabian Journal of Geosciences* 6(3), 913-919.
- Maxwell, T.L., Zhang, Y. (2005, August). A fuzzy approach to supervised segmentation parameter selection for object-based classification. In *Proceedings of SPIE, the International Society for Optical Engineering* (pp. 59091O-1). Society of Photo-Optical Instrumentation Engineers.
- McCalpin, J. (1984). Preliminary age classification of landslides for inventory mapping. In *Proceedings 21st annual Engineering Geology and Soils Engineering Symposium, Moscow, Idaho* (pp. 99-111).
- Mehrotra, A., Singh, K. K., Nigam, M. J., Pal, K. (2015). Detection of tsunami-induced changes using generalized improved fuzzy radial basis function neural network. *Natural Hazards* 77(1), 367-381.
- Metternicht, G., Hurni, L., Gogu, R. (2005). Remote sensing of landslides: An analysis of the potential contribution to geo-spatial systems for hazard assessment in mountainous environments. *Remote sensing of Environment*, 98(2), 284-303.
- Mia, M. M. A., Biswas, S. K., Urmi, M. C., Siddique, A. (2015). An algorithm for training multilayer perceptron (MLP) for Image reconstruction using neural network without overfitting. *Int J Sci Techno Res* 4(2), 271-275.

- Miner, A., Flentje, P., Mazengarb, C., Windle, D. (2010). Landslide Recognition using LiDAR derived Digital Elevation Models-Lessons learnt from selected Australian examples.
- Mahrooghy, M., Aanstoos, J. V., Nobrega, R. A., Hasan, K., Prasad, S., Younan, N. H. (2015). A machine learning framework for detecting landslides on earthen levees using spaceborne SAR imagery. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 8(8), 3791-3801.
- Mohammed, A. W., Zhang, M., Johnston, M. (2009). Particle swarm optimization based adaboost for face detection. In *Evolutionary Computation, 2009. CEC'09. IEEE Congress on* (pp. 2494-2501). IEEE.
- Moine, M., Puissant, A., Malet, J. P. (2009). Detection of landslides from aerial and satellite images with a semi-automatic method. Application to the Barcelonnette basin (Alpes-de-Hautes-Provence, France). In *Landslide processes-from geomorphologic mapping to dynamic modelling* 63–68.
- Mokhtarzade, M., Zoj, M. V. (2007). Road detection from high-resolution satellite images using artificial neural networks. *International Journal of Applied Earth Observation and Geoinformation* 9(1), 32-40.
- Mondini, A. C., Guzzetti, F., Reichenbach, P., Rossi, M., Cardinali, M., Ardizzone, F. (2011). Semi-automatic recognition and mapping of rainfall induced shallow landslides using optical satellite images. *Remote Sensing of Environment* 115(7), 1743-1757.
- Mondini, A. C., Marchesini, I., Rossi, M., Chang, K. T., Pasquariello, G., Guzzetti, F. (2013). Bayesian framework for mapping and classifying shallow landslides exploiting remote sensing and topographic data. *Geomorphology*, 201, 135-147.
- Moosavi, V., Talebi, A., Shirmohammadi, B. (2014). Producing a landslide inventory map using pixel-based and object based approaches optimized by Taguchi method. *Geomorphology*, 204, 646-656.
- Mora, O. E., Lenzano, M. G., Toth, C. K., Grejner-Brzezinska, D. A., Fayne, J. V. (2018). Landslide Change Detection Based on Multi-Temporal Airborne LiDAR-Derived DEMs. *Geosciences*, 8(1), 23.
- Moradi, P., Gholampour, M. (2016). A hybrid particle swarm optimization for feature subset selection by integrating a novel local search strategy. *Applied Soft Computing*, 43, 117-130.
- Mountrakis, G., Luo, L. (2011). Enhancing and replacing spectral information with intermediate structural inputs: A case study on impervious surface detection. *Remote Sensing of Environment* 115(5), 1162-1170.

- Mountrakis, G., Im, J., Ogole, C. (2011). Support vector machines in remote sensing: A review. *ISPRS Journal of Photogrammetry and Remote Sensing*, 66(3), 247-259.
- Murakmi, S., Nishigaya, T., Tien, T. L., Sakai, N., Lateh, H. H., Azizat, N. (2014). Development of historical landslide database in Peninsular Malaysia. In *Telecommunication Technologies (ISTT), 2014 IEEE 2nd International Symposium on* (pp. 149-153). IEEE.
- Mutanga, O., Adam, E., Cho, M. A., 2012. High density biomass estimation for wetland vegetation using WorldView-2 imagery and random forest regression algorithm. *International Journal of Applied Earth Observation and Geoinformation* 18, 399–406.
- Navarro-Sanchez, V. D., Lopez-Sanchez, J. M., Vicente-Guijalba, F. (2010). A contribution of polarimetry to satellite differential SAR interferometry: Increasing the number of pixel candidates. *IEEE Geoscience and Remote Sensing Letters*, 7(2), 276-280.
- Nefeslioglu, H. A., Sezer, E., Gokceoglu, C., Bozkir, A. S., Duman, T. Y. (2010). Assessment of landslide susceptibility by decision trees in the metropolitan area of Istanbul, Turkey. *Mathematical Problems in Engineering*, 2010.
- Nichol, J. E., Shaker, A., Wong, M. S., 2006. Application of high-resolution stereo satellite images to detailed landslide hazard assessment. *Geomorphology* 76(1), 68–75.
- O'Connell, J., Bradter, U., Benton, T. G. (2015). Wide-area mapping of small-scale features in agricultural landscapes using airborne remote sensing. *ISPRS Journal of Photogrammetry and Remote Sensing* 109, 165-177.
- Olaya, V. (2009). Basic land-surface parameters. *Developments in Soil Science* 33, 141–169.
- Omran, M. G., Salman, A., Engelbrecht, A. P. (2006). Dynamic clustering using particle swarm optimization with application in image segmentation. *Pattern Analysis and Applications*, 8(4), 332.
- Oppikofer, T., Bunkholt, H. S. S., Fischer, L., Saintot, A., Hermanns, R. L., Carrea, D., Jaboyedoff, M. (2012, June). Investigation and monitoring of rock slope instabilities in Norway by terrestrial laser scanning. In *Landslides and Engineered Slopes. Protecting Society through Improved Understanding: Proceedings of the 11th International & 2nd North American Symposium on Landslides, Banff, Canada*, (pp. 3-8).
- Ordóñez, F. J., Roggen, D. (2016). Deep convolutional and lstm recurrent neural networks for multimodal wearable activity recognition. *Sensors*, 16(1), 115.

- Osei-Bryson, K. M. (2014). Overview on decision tree induction. In *Advances in research methods for information systems research* (pp. 15-22). Springer US.
- Otukei, J. R., Blaschke, T. (2010). Land cover change assessment using decision trees, support vector machines and maximum likelihood classification algorithms. *International Journal of Applied Earth Observation and Geoinformation*, 12, S27-S31.
- Pal, M., Foody, G. M. (2010). Feature selection for classification of hyperspectral data by SVM. *IEEE Transactions on Geoscience and Remote Sensing* 48(5), 2297-2307.
- Palenzuela, J. A., Marsella, M., Nardinocchi, C., Pérez, J. L., Fernández, T., Chacón, J., Irigaray, C. (2015). Landslide detection and inventory by integrating LiDAR data in a GIS environment. *Landslides*, 12(6), 1035-1050.
- Parker, R. N., Densmore, A. L., Rosser, N. J., De Michele, M., Li, Y., Huang, R., Petley, D. N. (2011). Mass wasting triggered by the 2008 Wenchuan earthquake is greater than orogenic growth. *Nature Geoscience* 4(7), 449-452.
- Parpinelli, R. S., Lopes, H. S., Freitas, A. A. (2002). Data mining with an ant colony optimization algorithm. *IEEE Transactions on Evolutionary Computation* 6(4), 321-332.
- Pawłuszek, K., Borkowski, A. (2016). Landslides identification using airborne laser scanning data derived topographic terrain attributes and support vector machine classification. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 8, 145-149.
- Pawłuszek, K., Borkowski, A. (2017). Impact of DEM-derived factors and analytical hierarchy process on landslide susceptibility mapping in the region of Rożnów Lake, Poland. *Natural Hazards*, 86(2), 919-952.
- Pawłuszek, K., Borkowski, A., Tarolli, P. (2018). Sensitivity analysis of automatic landslide mapping: numerical experiments towards the best solution. *Landslides*, 1-15.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Vanderplas, J. (2011). Scikit-learn: Machine learning in Python. *Journal of Machine Learning Research* 12(Oct), 2825-2830.
- Pei, Y., Wang, W., Zhang, S. (2012). Basic ant colony optimization. In *Computer Science and Electronics Engineering (ICCSEE), 2012 International Conference on* (Vol. 1, pp. 665-667). IEEE.
- Permuter, H., Francos, J., Jermyn, I. (2006). A study of Gaussian mixture models of color and texture features for image classification and segmentation. *Pattern Recognition*, 39(4), 695-706.



- Perumal, K., Bhaskaran, R. (2010). Supervised classification performance of multispectral images. *arXiv preprint arXiv:1002.4046*.
- Pesaresi, M., Gerhardinger, A., Kayitakire, F. (2008). A robust built-up area presence index by anisotropic rotation-invariant textural measure. *IEEE Journal of selected topics in applied earth observations and remote sensing*, 1(3), 180-192.
- Pontius Jr, R. G., Millones, M. (2011). Death to Kappa: birth of quantity disagreement and allocation disagreement for accuracy assessment. *International Journal of Remote Sensing* 32(15), 4407-4429.
- Pradhan, B. (2010). Flood susceptible mapping and risk area delineation using logistic regression, GIS and remote sensing. *Journal of Spatial Hydrology* 9(2).
- Pradhan, B. (2011). Use of GIS-based fuzzy logic relations and its cross application to produce landslide susceptibility maps in three test areas in Malaysia. *Environmental Earth Sciences*, 63(2), 329-349.
- Pradhan, B. (2013). A comparative study on the predictive ability of the decision tree, support vector machine and neuro-fuzzy models in landslide susceptibility mapping using GIS. *Computers and Geosciences*, 51, 350-365.
- Pradhan, B., Alsaleh, A. (2017). A Supervised Object-Based Detection of Landslides and Man-Made Slopes Using Airborne Laser Scanning Data. In *Laser Scanning Applications in Landslide Assessment*, (pp. 23-50). Springer, Cham.
- Pradhan, B., Buchroithner, M. F. (2010). Comparison and validation of landslide susceptibility maps using an artificial neural network model for three test areas in Malaysia. *Environmental and Engineering Geoscience*, 16(2), 107-126.
- Pradhan, B., Jebur, M. N., Shafri, H. Z. M., Tehrany, M. S. (2016). Data Fusion Technique Using Wavelet Transform and Taguchi Methods for Automatic Landslide Detection From Airborne Laser Scanning Data and QuickBird Satellite Imagery. *IEEE Transactions on Geoscience and Remote Sensing* 54(3), 1610 –1622.
- Pradhan, B., Lee, S. (2010). Regional landslide susceptibility analysis using back-propagation neural network model at Cameron Highland, Malaysia. *Landslides* 7(1), 13–30.
- Pradhan, B., Mansor, S., Pirasteh, S., Buchroithner, M. F. (2011). Landslide hazard and risk analyses at a landslide prone catchment area using statistical based geospatial model. *International Journal of Remote Sensing*, 32(14), 4075-4087.



- Pradhan, B., Sameen, M. I., Nampak, H. (2017). Integration of LiDAR and QuickBird Data for Automatic Landslide Detection Using Object-Based Analysis and Random Forests. In *Laser Scanning Applications in Landslide Assessment* (pp. 69-81). Springer, Cham.
- Pratola, C., Del Frate, F., Schiavon, G., Solimini, D., Licciardi, G. (2011). Characterizing land cover from X-band COSMO-SkyMed images by neural networks. In *Urban Remote Sensing Event (JURSE), 2011 Joint* (pp. 49-52). IEEE.
- Puissant, A., Rougier, S., Stumpf, A. (2014). Object based mapping of urban trees using Random Forest classifiers. *International Journal of Applied Earth Observation and Geoinformation* 26, 235-245.
- Quinlan, J. R. (1993). C4. 5: Programming for machine learning. *Morgan Kaufmann*, 38.
- Radoux, J., Bogaert, P. (2014). Accounting for the area of polygon sampling units for the prediction of primary accuracy assessment indices. *Remote sensing of environment*, 142, 9-19.
- Rau, J. Y., Chang, K. T., Shao, Y. C., Lau, C.C. (2012). Semi-automatic shallow landslide detection by the integration of airborne imagery and laser scanning data. *Natural Hazards* 61(2), 469–480.
- Rau, J. Y., Jhan, J. P., Rau, R. J. (2014). Semiautomatic object based landslide recognition scheme from multisensor optical imagery and DEM. *IEEE Transactions on Geoscience and Remote Sensing*, 52(2), 1336-1349.
- Razak, K. A., Straatsma, M. W., Van Westen, C. J., Malet, J. P., De Jong, S. M. (2011). Airborne laser scanning of forested landslides characterization: Terrain model quality and visualization. *Geomorphology*, 126(1-2), 186-200.
- Rejer, I., Lorenz, K. (2013). Genetic algorithm and forward method for feature selection in EEG feature space. *Journal of Theoretical and Applied Computer Science*, 7(2), 72-82.
- Rodriguez-Galiano, V.F., Ghimire, B., Rogan, J., Chica-Olmo, M., Rigol-Sanchez, J.P.(2012). An assessment of the effectiveness of a random forest classifier for land-cover classification. *ISPRS Journal of Photogrammetry and Remote Sensing* 67, 93–104.
- Rosin, P.L., Hervás, J. (2005). Remote sensing image thresholding methods for determining landslide activity. *International Journal of Remote Sensing* 26(6), 1075-1092.
- Rozenstein, O., Karnieli, A. (2011). Comparison of methods for land-use classification incorporating remote sensing and GIS inputs. *Applied Geography*, 31(2), 533-544.

- Ruiz, L. Á., Recio, J. A., Crespo-Peremarch, P., Sapena, M. (2018). An object-based approach for mapping forest structural types based on low-density LiDAR and multispectral imagery. *Geocarto International*, 33(5), 443-457.
- Saeys, Y., Inza, I., Larrañaga, P. (2007). A review of feature selection techniques in bioinformatics. *Bioinformatics* 23(19), 2507–2517.
- Salehi, B., Zhang, Y., Zhong, M., Dey, V. (2012). Object-based classification of urban areas using VHR imagery and height points ancillary data. *Remote Sensing*, 4(8), 2256-2276.
- Sameen, M. I., Pradhan, B., Shafri, H. Z., Mezaal, M. R., bin Hamid, H. (2017). Integration of Ant Colony Optimization and Object-Based Analysis for LiDAR Data Classification. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 10(5), 2055–2066.
- Sameen, M.I., Pradhan, B. (2017). A Two-Stage Optimization Strategy for Fuzzy Object-Based Analysis Using Airborne LiDAR and High-Resolution Orthophotos for Urban Road Extraction. *Journal of Sensors*, 2017.
- Sameen, M.I., Pradhan, B. (2017). Severity Prediction of Traffic Accidents with Recurrent Neural Networks. *Applied Sciences* 7(6), 476.
- Samui, P. (2008). Slope stability analysis: a support vector machine approach. *Environmental Geology*, 56(2), 255.
- Samy, I. E., Marghany, M. M., Mohamed, M. M. (2014). Landslide modelling and analysis using remote sensing and GIS: A case study of Cameron highland, Malaysia. *Journal of Geomatics*, 8(2).
- Santangelo, á., Cardinali, á., Rossi, á., Mondini, A., Guzzetti, F. (2010). Remote landslide mapping using a laser rangefinder binocular and GPS. *Natural Hazards and Earth System Sciences* 10(12), 2539-2546.
- Schmidhuber, J. (2015). Deep learning in neural networks: An overview. *Neural networks*, 61, 85-117.
- Schulz, W.H. (2007). Landslide susceptibility revealed by LIDAR imagery and historical records, Seattle, Washington. *Engineering Geology* 89(1), 67-87.
- Segoni, S., Leoni, L., Benedetti, A. I., Catani, F., Righini, G., Falorni, G., Rebor, N. (2009). Towards a definition of a real-time forecasting network for rainfall induced shallow landslides. *Natural Hazards and Earth System Sciences*, 9(6), 2119.
- Shackelford, A. K., Davis, C. H. (2003). A hierarchical fuzzy classification approach for high-resolution multispectral data over urban areas. *IEEE transactions on geoscience and remote sensing*, 41(9), 1920-1932.

- Shahabi, H., Hashim, M. (2015). Landslide susceptibility mapping using GIS-based statistical models and Remote sensing data in tropical environment. *Scientific reports*, 5, 9899.
- Sheikhpour, R., Sarram, M. A., Sheikhpour, R. (2016). Particle swarm optimization for bandwidth determination and feature selection of kernel density estimation based classifiers in diagnosis of breast cancer. *Applied Soft Computing* 40, 113-131.
- Shi, W., Deng, S., Xu, W. (2018). Extraction of multi-scale landslide morphological features based on local Gi\* using airborne LiDAR-derived DEM. *Geomorphology*, 303, 229-242.
- Shi, L., Wan, Y., Gao, X., Wang, M. (2018). Feature Selection for Object-Based Classification of High-Resolution Remote Sensing Images Based on the Combination of a Genetic Algorithm and Tabu Search. *Computational Intelligence and Neuroscience*, 2018.
- Singh, A., Singh, K. K. (2017). Satellite image classification using Genetic Algorithm trained radial basis function neural network, application to the detection of flooded areas. *Journal of Visual Communication and Image Representation* 42, 173-182.
- Singh, D. A. A. G., Leavline, E. J., Valliyappan, K., Srinivasan, M. (2015). Enhancing the performance of classifier using particle swarm optimization (PSO)-based dimensionality reduction. *International Journal of Information and Communication Sciences*, 6, 19-26.
- Singh, K. K., Nigam, M. J., Pal, K., Mehrotra, A. (2014). A fuzzy kohonen local information C-means clustering for remote sensing imagery. *IETE Technical Review*, 31(1), 75-81.
- Singh, K.K., Singh, A. (2016). Detection of 2011 Sikkim earthquake-induced landslides using neuro-fuzzy classifier and digital elevation model. *Natural Hazards* 83(2), 1027-1044.
- Siregar, V. P., Agus, S. B., Subarno, T., Prabowo, N. W. (2018). Mapping shallow waters habitats using OBIA by applying several approaches of depth invariant index in North Kepulauan Seribu. In *IOP Conference Series: Earth and Environmental Science* (Vol. 149, No. 1, p. 012052). IOP Publishing.
- Siyahghalati, S., Saraf, A. K., Pradhan, B., Jebur, M. N., Tehrany, M. S. (2016). Rule-based semi-automated approach for the detection of landslides induced by 18 September 2011 Sikkim, Himalaya, earthquake using IRS LISS3 satellite images. *Geomatics, Natural Hazards and Risk*, 7(1), 326-344.
- Slatton, K. C., Carter, W. E., Shrestha, R. L., Dietrich, W. (2007). Airborne laser swath mapping: achieving the resolution and accuracy required for geosurficial research. *Geophysical Research Letters*, 34(23).

- Srivastava, N., Hinton, G. E., Krizhevsky, A., Sutskever, I., Salakhutdinov, R. (2014). Dropout: a simple way to prevent neural networks from overfitting. *Journal of Machine Learning Research* 15(1), 1929–1958.
- Statnikov, A., Wang, L., Aliferis, C. F. (2008). A comprehensive comparison of random forests and support vector machines for microarray-based cancer classification. *BMC bioinformatics*, 9(1), 319.
- Stumpf, A., Kerle, N.(2011). Object based mapping of landslides using Random Forests. *Remote Sensing of Environment* 115(10), 2564–2577.
- Stumpf, A., Lachiche, N., Malet, J.P., Kerle, N., Puissant, A. (2014). Active learning in the spatial domain for remote sensing image classification. *IEEE Transactions on Geoscience and Remote Sensing*, 52(5), 2492-2507.
- Su, W., Zhang, C., Zhu, X., Li, D. (2009). A hierarchical object based method for land cover classification of SPOT 5 imagery. *WSEAS Transactions on Information Science and Applications*, 6(3), 437-446.
- Suliman, A., Zhang, Y. (2016). Segment-Based Terrain Filtering Technique for Elevation-Based Building Detection in VHR Remote Sensing Images. *Advances in Remote Sensing*, 5(03), 192.
- Swets, J.A. (1988). Measuring the accuracy of diagnostic systems. *Science*, 240(4857), 1285-1293.
- Tamimi, E., Ebadi, H., Kiani, A. (2017). Hybrid Optimization of Object-Based Classification in Highresolution Images Using Continous Ant Colony Algorithm With Emphasis On Building Detection. *International Archives of the Photogrammetry, Remote Sensing & Spatial Information Sciences*, 42.
- Tang, A. M., Quek, C., Ng, G. S. (2005). GA-TSKfnn: Parameters tuning of fuzzy neural network using genetic algorithms. *Expert Systems with Applications* 29(4), 769-781.
- Team, R. D.C. (2011). R Development Core Team: R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2011.
- Tehrany, M.S., Pradhan, B., Jebur, M.N. (2014a). Flood susceptibility mapping using a novel ensemble weights-of-evidence and support vector machine models in GIS. *Journal of Hydrology* 512, 332-343.
- Tehrany, M.S., Pradhan, B., Jebur, M.N. (2014b). A comparative assessment between object and pixel-based classification approaches for land use/land cover mapping using SPOT 5 imagery. *Geocarto International* 29(4), 351-369.

- Tehrany, M.S., Pradhan, B., Jebur, M. N. (2013). Spatial prediction of flood susceptible areas using rule based decision tree (DT) and a novel ensemble bivariate and multivariate statistical models in GIS. *Journal of Hydrology* 504, 69-79.
- Tong, H., Maxwell, T., Zhang, Y., Dey, V. (2012). A supervised and fuzzy-based approach to determine optimal multi-resolution image segmentation parameters. *Photogrammetric Engineering and Remote Sensing* 78(10), 1029-1044.
- Travelletti, J., Delacourt, C., Allemand, P., Malet, J. P., Schmittbuhl, J., Toussaint, R., Bastard, M. (2012). Correlation of multi-temporal ground-based optical images for landslide monitoring: Application, potential and limitations. *ISPRS Journal of Photogrammetry and Remote Sensing* 70, 39-55.
- Turcotte, D.L., Malamud, B.D., Guzzetti, F., Reichenbach, P. (2002). Self-organization, the cascade model, and natural hazards. *Proceedings of the National Academy of Sciences* 99 (suppl 1), 2530-2537.
- Unler, A., Murat, A. (2010). A discrete particle swarm optimization method for feature selection in binary classification problems. *European Journal of Operational Research* 206(3), 528-539.
- van Asselen, S., Seijmonsbergen, A.C. (2006). Expert-driven semi-automated geomorphological mapping for a mountainous area using a laser DTM. *Geomorphology* 78(3), 309-320.
- Van Coillie, F. M., Verbeke, L. P., De Wulf, R. R. (2007). Feature selection by genetic algorithms in object-based classification of IKONOS imagery for forest mapping in Flanders, Belgium. *Remote Sensing of Environment*, 110(4), 476-487.
- Van Den Eeckhaut, M., Hervás, J. (2012). State of the art of national landslide databases in Europe and their potential for assessing landslide susceptibility, hazard and risk. *Geomorphology* 139, 545-558.
- Van Den Eeckhaut, M., Kerle, N., Poesen, J., Hervás, J. (2012). Object based identification of forested landslides with derivatives of single pulse LiDAR data. *Geomorphology* 173, 30-42.
- Van Westen, C. J., Van Asch, T. W., Soeters, R. (2006). Landslide hazard and risk zonation—why is it still so difficult?. *Bulletin of Engineering geology and the Environment*, 65(2), 167-184.
- Van Westen, C. J., Castellanos, E., Kuriakose, S.L. (2008). Spatial data for landslide susceptibility, hazard, and vulnerability assessment: an overview. *Engineering Geology* 102(3), 112-131.

- Vapnik, V. (1999). The Nature of Statistical Learning Theory. *Springer-Verlag* 344 p.
- Vennari, C., Gariano, S. L., Antronico, L., Brunetti, M. T., Iovine, G., Peruccacci, S., ... Guzzetti, F. (2014). Rainfall thresholds for shallow landslide occurrence in Calabria, southern Italy. *Natural hazards and earth system sciences*, 14(2), 317-330.
- Verikas, A., Gelzinis, A., Bacauskiene, M. (2011). Mining data with random forests: A survey and results of new tests. *Pattern Recognition*, 44(2), 330-349.
- Vieira, M. A., Formaggio, A. R., Rennó, C. D., Atzberger, C., Aguiar, D. A., Mello, M. P. (2012). Object based image analysis and data mining applied to a remotely sensed Landsat time-series to map sugarcane over large areas. *Remote Sensing of Environment*, 123, 553-562.
- Vohora, V. K., Donoghue, S. L. (2004). Application of remote sensing data to landslide mapping in Hong Kong. *International Archives of Photogrammetry, Remote Sensing and Spatial Information Sciences*.
- Vosselman, G. (2000). Slope based filtering of laser altimetry data. *International Archives of Photogrammetry and Remote Sensing*, 33(B3/2; PART 3), 935-942.
- Wang, S., Yao, X. (2013). Relationships between diversity of classification ensembles and single-class performance measures. *IEEE Transactions on Knowledge and Data Engineering*, 25(1), 206-219.
- Watts, J. D., Lawrence, R. L., Miller, P. R., Montagne, C. (2009). Monitoring of cropland practices for carbon sequestration purposes in north central Montana by Landsat remote sensing. *Remote Sensing of Environment* 113(9), 1843-1852.
- Wei, L., Qi, J. J. (2010). Relation between concept lattice reduction and rough set reduction. *Knowledge-Based Systems* 23(8), 934-938.
- Wiegand, C., Rutzinger, M., Heinrich, K., Geitner, C. (2013). Automated extraction of shallow erosion areas based on multi-temporal ortho-imagery. *Remote Sensing*, 5(5), 2292-2307.
- Wilson, J. P., Gallant, J. C. (Eds.). (2000). *Terrain analysis: principles and applications*. John Wiley & Sons.
- Witten, I. H., Frank, E. (2005). Data mining: Practical machine learning tools and techniques, (morgan kaufmann series in data management systems). *Morgan Kaufmann, June*, 104, 113.
- Wu, H., Prasad, S. (2017). Convolutional recurrent neural networks for hyperspectral data classification. *Remote Sensing*, 9(3), 298.



- Wu, X., Ren, F., Niu, R. (2014). Landslide susceptibility assessment using object mapping units, decision tree, and support vector machine models in the Three Gorges of China. *Environmental earth sciences*, 71(11), 4725-4738.
- Wu, Y., Schuster, M., Chen, Z., Le, Q. V., Norouzi, M., Macherey, W., Klingner, J. (2016). Google's neural machine translation system: Bridging the gap between human and machine translation. *arXiv preprint arXiv:1609.08144*.
- Xiang, S., Nie, F., Zhang, C. (2008). Learning a Mahalanobis distance metric for data clustering and classification. *Pattern Recognition* 41(12), 3600-3612.
- Xiaoliang, Z., Hongcan, Y., Jian, W., Shangzhuo, W. (2009, December). Research and application of the improved algorithm C4. 5 on decision tree. In *Test and Measurement, 2009. ICTM'09. International Conference on* (Vol. 2, pp. 184-187). IEEE.
- Xue, B., Zhang, M., Browne, W. N. (2013). Particle swarm optimization for feature selection in classification: A multi-objective approach. *IEEE transactions on cybernetics* 43(6), 1656-1671.
- Yan, G., Mas, J. F., Maathuis, B. H. P., Xiangmin, Z., Van Dijk, P. M. (2006). Comparison of pixel-based and object-oriented image classification approaches—a case study in a coal fire area, Wuda, Inner Mongolia, China. *International Journal of Remote Sensing*, 27(18), 4039-4055.
- Yang, G. Y. C. (1995). Geological mapping from multi-source data using neural networks. *Geomatics Engineering*
- Yang, M. D. (2007). A genetic algorithm (GA) based automated classifier for remote sensing imagery. *Canadian Journal of Remote Sensing*, 33(3), 203-213.
- Yang, X., Chen, L. (2010). Using multi-temporal remote sensor imagery to detect earthquake-triggered landslides. *International Journal of Applied Earth Observation and Geoinformation* 12(6), 487-495.
- Yu, T.T., Wang, T. S., Cheng, Y. S. (2015). Analysis of factors triggering shallow failure and deep-seated landslides induced by single rainfall events. *Journal of Disaster Research*, 10(5), 966-972.
- Yu, W., Wu, B., Huang, T., Li, X., Williams, K., Zhao, H. (2006). Statistical methods in proteomics. *Springer Handbook of Engineering Statistics* 623–638.
- Yuan, H., Van Der Wiele, C. F., Khorram, S. (2009). An automated artificial neural network system for land use/land cover classification from Landsat TM imagery. *Remote Sensing*, 1(3), 243-265.
- Zaremba, W., Sutskever, I., Vinyals, O. (2014). Recurrent neural network regularization. *arXiv preprint arXiv:1409.2329*.



- Zêzere, J. L., Trigo, R. M., Trigo, I. F. (2005). Shallow and deep landslides induced by rainfall in the Lisbon region (Portugal): assessment of relationships with the North Atlantic Oscillation. *Natural Hazards and Earth System Science* 5(3), 331-344.
- Zhang, Y., Maxwell, T., Tong, H., Dey, V. (2010). Development of a supervised software tool for automated determination of optimal segmentation parameters for ecognition 5–7.
- Zhen, Z., Quackenbush, L.J., Stehman, S.V., Zhang, L. (2013). Impact of training and validation sample selection on classification accuracy and accuracy assessment when using reference polygons in object-based classification. *International Journal of Remote Sensing* 34(19), 6914–6930.
- Zizioli, D., Meisina, C., Bordoni, M., Zucca, F. (2014). Rainfall-triggered shallow landslides mapping through Pleiades images. In *Landslide science for a Safer Geoenvironment* (pp. 325-329). Springer, Cham.
- Zurada, J. M. (1992). Introduction to artificial neural systems (Vol. 8). St. Paul: West.