



UNIVERSITI PUTRA MALAYSIA

***ANTIMICROBIAL ACTIVITY, PHYTOCHEMICAL ANALYSIS AND
TOXICITY TEST OF *Cosmos caudatus* Kunth EXTRACT***

NOR ASMA HUSNA MOHAMED YUSOFF

IB 2015 34



**ANTIMICROBIAL ACTIVITY, PHYTOCHEMICAL ANALYSIS AND
TOXICITY TEST OF *Cosmos caudatus* Kunth EXTRACT**

By

NOR ASMA HUSNA MOHAMED YUSOFF

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

December 2015

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Doctor of Philosophy

**ANTIMICROBIAL ACTIVITY, PHYTOCHEMICAL ANALYSIS AND
TOXICITY TEST OF *Cosmos caudatus* Kunth EXTRACT**

By

NOR ASMA HUSNA YUSOFF

December 2015

Chairman : Associate Professor Yaya Rukayadi, PhD
Institute : Bioscience

The uses of natural products in the elimination of microorganism contaminations in food have increased nowadays. In this study, air-dried leaf methanolic extract of *Cosmos caudatus* Kunth was used. The antimicrobial activity of *C. caudatus* extract and time-kill curve analysis was evaluated against foodborne pathogens. The stability of extract at different pH and temperature also were evaluated. The identification and quantification of bioactive compounds were conducted using LC-MS/MS and HPLC. Three representative microorganisms were observed under transmission electron microscope (TEM). The effect of *C. caudatus* extract on natural microflora in selected food materials and its sensory acceptability were tested at different concentrations of extract for different exposure time. The toxicity level of extract was assessed using *Artemia salina* spp. Results showed that all tested foodborne pathogens were susceptible to *C. caudatus* extract at different range of MIC value from 6.25 to 50.00 mg/ml. Time-kill curve study shows that all tested foodborne pathogens can be killed by the extract with different time-killing curve. Generally, the antimicrobial activities of *C. caudatus* extract were not significantly affected on tested pH and temperatures. LC-MS/MS analysis identified the presence of rutin, quercetin 3-*O*-glucoside, quercetin 3-*O*-arabinoside, quercetin 3-*O*-rhamnoside and quercetin. HPLC analysis showed the highest concentration was quercetin rhamnoside (29.66 mg/g). All the identified active compounds proved possess antimicrobial activities. TEM observation showed changing on the pathogen cells's structure. Generally the significant reduction of natural microflora in raw food materials were started at 0.05% of *C. caudatus* extract for 10 min of exposure time, and the sensory acceptability at this concentration and time were accepted by the panellists. The toxicity study demonstrated that *C. caudatus* extract was not toxic to *A. salina* ($LC_{50} = 3.54$ mg/ml). In conclusion, *C. caudatus* extract exhibits antimicrobial activities, thus can be developed as natural sanitizer for washing raw food materials.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**AKTIVITI ANTIMIKROB, ANALISIS FITOKIMIA DAN UJIAN TOKSISITI
OLEH EKSTRAK *Cosmos caudatus* Kunth**

Oleh

NOR ASMA HUSNA YUSOFF

Disember 2015

Pengerusi : Profesor Madya Yaya Rukayadi, PhD
Institut : Biosains

Sekarang ini, penggunaan produk semula jadi dalam penghapusan pencemaran mikroorganisma dalam bahan makanan meningkat. Dalam kajian ini, sampel daun pengeringan-angin metanolik ekstrak *Cosmos caudatus* Kunth digunakan. Aktiviti antimikrob ekstrak *C. caudatus* dan analisis keluk masa-pembunuhan dinilai terhadap patogen bawaan makanan. Kestabilan ekstrak pada pH dan suhu yang berbeza juga dinilai. Mengenalpasti dan mengukur kandungan sebatian bioaktif dalam ekstrak dilakukan menggunakan LC-MS/MS dan HPLC. Tiga wakil mikroorganisma diperhatikan di bawah mikroskop elektron transmisi (TEM). Kesan ekstrak *C. caudatus* terhadap mikroflora semulajadi dalam bahan makanan dan penerimaan deria diuji pada kepekatan ekstrak berbeza untuk masa berbeza. Tahap ketoksikan ekstrak dinilai menggunakan *Artemia salina* spp. Keputusan menunjukkan semua patogen yang diuji dipengaruhi oleh ekstrak *C. caudatus* pada nilai MIC yang berbeza bermula dari 6.25 hingga 50.00 mg/ml. Kajian keluk masa-pembunuhan menunjukkan bahawa semua patogen bawaan makanan boleh dibunuh oleh ekstrak pada keluk masa-pembunuhan yang berlainan. Secara umumnya, aktiviti antimikrobial ekstrak *C. caudatus* terhadap semua patogen bawaan makanan yang diuji tidak terjejas dengan ketara oleh pH dan suhu yang diuji. Analisis LC-MS/MS menentukan kehadiran rutin, kuersetin 3-*O*-glukosida, kuersetin 3-*O*-arabinosida, kuersetin 3-*O*-ramnosida dan kuersetin. Analisis HPLC menunjukkan kepekatan tertinggi adalah kuersetin ramnosida (29.66 mg/g). Semua sebatian ini telah dibuktikan mempunyai aktiviti antimikrob. Pemerhatian TEM menunjukkan perubahan kepada struktur sel. Secara keseluruhannya, jumlah pengurangan mikroflora semulajadi dalam bahan makanan mentah yang dirawat adalah ketara bermula pada kepekatan ekstrak *C. caudatus* 0.05% selama 10 min masa pendedahan, dan penerimaan deria pada kepekatan dan pendedahan masa ini telah diterima oleh ahli panel. Kajian ketoksikan menunjukkan bahawa ekstrak *C. caudatus* tidak toksik kepada *A. salina* ($LC_{50} = 3.54$ mg/ml). Kesimpulannya, ekstrak *C. caudatus* mempamerkan aktiviti antimikrob, dengan itu boleh dimajukan sebagai pembersih semulajadi untuk mencuci bahan makanan mentah.

ACKNOWLEDGEMENTS

In the name of Allah, most Gracious and most Merciful. All praise to Allah for His blessing that allows me to complete this piece of work.

I would like to express my greatest thanks and appreciation to my research supervisor Assoc. Prof. Dr. Yaya Rukayadi for his invaluable guidance, understanding, patience and constant encouragement throughout the course of my study. Without his guidance this thesis would not have been possible to be finished.

My thanks extended to my committee members; Assoc. Prof. Dr. Faridah Abas for her guidance especially on the HPLC and LC-MS problem. Special thanks also to Assoc. Prof. Dr. Alfi Khatib for his early guidance before he had to transfer to International Islamic University of Malaysia (IIUM). Many thanks also go to all staffs in IBS, Mrs Mazina, Mrs Zurina, Mr Salahuddin and Mr Azizul for their valuable help. Very special thanks to all my friends; Muhammad Safwan Bustamam, Lew Kok Fang, Ahmed Mediani, Syahidah Ahmad, Nurul Syazwani Zainin, Lau Kah Yan, Maya Zakaria, Nurul Husna and others who are so many which is impossible to mention all of them here.

Last but not least, I am greatly indebted to my dear husband, Mohd Lokmal Md Hussein for his understanding, practical and emotional support throughout my study. My lovely daughter Sofie Nur Hanna, thank you for being a good girl whenever I bring you to lab. Not to be forgotten my beloved family, my mother Asiah Mat Noor, my father Mohamed Yusoff Yaacob and siblings who have always prayed for my success.

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Yaya Rukayadi, PhD

Associate Professor
Institute of Bioscience
Universiti Putra Malaysia
(Chairman)

Faridah Abas, PhD

Associate Professor
Institute of Bioscience
Universiti Putra Malaysia
(Member)

Alfi Khatib, PhD

Associate Professor
Department of Food Science and Technology
International Islamic Universiti Malaysia
(External member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by Members of Supervisory Committee

This is to confirm that:

- The research conducted and the writing of this thesis was under our supervision;
- Supervisions responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012 - 2013) are adhered to.

Signature: _____

Name of

Chairman of

Supervisory

Committee: Assoc. Prof. Dr. Yaya Rukayadi

Signature: _____

Name of

Member of

Supervisory

Committee: Assoc. Prof. Dr. Faridah Abas

Signature: _____

Name of

Member of

Supervisory

Committee: Assoc. Prof. Dr. Alfi Khatib

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	ii
ACKNOWLEDGEMENTS	iii
APPROVAL	iv
DECLARATION	vi
LIST OF TABLES	xii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xvii
 CHAPTER	
 1 INTRODUCTION	
1.1 General Introduction	1
1.2 Objectives	3
 2 LITERATURE REVIEW	
2.1 Foodborne Pathogens	5
2.2 Sources of Foodborne Pathogens	7
2.2.1 Milk and dairy product	7
2.2.2 Meat-based product	9
2.2.3 Fruits and vegetables	9
2.3 Natural Plant Antimicrobials	11
2.3.1 Phenolic compounds	12
2.3.2 Essential oils	14
2.4 <i>Cosmos caudatus</i> Kunth.	16
2.4.1 Biology of <i>C. caudatus</i> Kunth.	16
2.4.1.1 Classification of <i>C. caudatus</i> Kunth.	16
2.4.1.2 Morphology of <i>C. caudatus</i> Kunth.	16
2.4.1.3 Distribution of <i>C. caudatus</i> Kunth.	19
2.4.1.4 Traditional use of <i>C. caudatus</i> Kunth.	19
2.4.2 Metabolites identified in <i>C. caudatus</i> Kunth.	19
2.4.3 Pharmacological reports of <i>C. caudatus</i> Kunth.	22
2.4.3.1 Antioxidant activity	23
2.4.3.2 Anticancer and antimutagen activity	25
2.4.3.3 Antimicrobial activity	26
2.5 Major compounds in <i>C. caudatus</i> extract	28
2.5.1 Quercetin	28
2.5.2 Catechin	29
2.5.3 Rutin	31
2.5.4 Chlorogenic acid	32
2.6 Food Sanitizations	33
2.6.1 Chemical-based control sanitizer	33
2.6.2 Natural-based control sanitizer	34

3	MATERIALS AND METHODS	
3.1	Sample Collection	36
3.1.1	<i>Cosmos caudatus</i> Kunth.	36
3.1.2	Raw food materials	37
3.2	Extraction of <i>C. caudatus</i>	37
3.3	Antimicrobial Activity of <i>C. caudatus</i> Extract	37
3.3.1	Microbial strains and maintenance	37
3.3.2	Preparation of media and chemicals	39
3.3.2.1	Preparation of microbiological agar/broth	39
3.3.2.2	Preparation of phosphate buffer saline	40
3.3.2.3	Preparation of chlorhexidine	40
3.3.2.4	Preparation of dimethyl sulfoxide(DMSO)	41
3.3.3	Preparation of antimicrobial agents	41
3.3.4	Preparation of inoculums	41
3.3.5	Disc-diffusion test	41
3.3.6	Minimum inhibitory concentration (MIC) and minimum bactericidal/fungicidal concentration (MBC/MFC)	42
3.3.7	Time-kill curves analysis	42
3.3.8	Stability of <i>C. caudatus</i> extract at different pH and Temperature	43
3.4	Light Sheet Electron Microscope (LSEM)	43
3.5	Transmission Electron Microscope (TEM)	43
3.6	Identification and Quantification of Active Compounds in <i>C. caudatus</i> Extract	44
3.6.1	High performance liquid chromatography (HPLC)	44
3.6.2	Liquid chromatography mass spectrometry (LC-MS/MS)	45
3.7	Antimicrobial Activity of <i>C. caudatus</i> Single Compounds	46
3.8	Application of <i>C. caudatus</i> Extract on Raw Food Materials	46
3.8.1	Preparation of <i>C. caudatus</i> extract for treatment Solution	46
3.8.2	Preparation of selective media agar	46
3.8.3	Treatment of raw food materials with <i>C. caudatus</i> extract solution	47
3.9	Sensory Evaluation	48
3.9.1	Evaluation of sensory acceptability of treated raw food materials	48
3.9.2	Sensory statistical analysis	48
3.10	Toxicity Evaluation of <i>C. caudatus</i> Extract	48
3.10.1	Hatching shrimp	48
3.10.2	Brine shrimp (<i>Artemia salina</i> spp.) lethality assay	48
3.11	Statistical Analysis	48

4	RESULTS AND DISCUSSIONS	
4.1	Yield of <i>C. caudatus</i> Crude Extract	49
4.2	Susceptibility of <i>C. caudatus</i> Extract on Foodborne Pathogens	51
4.2.1	Disc-diffusion, minimum inhibitory concentration (MIC), minimum bactericidal/fungicidal concentration (MBC/MFC)	51
4.3	Time-kill Curve Analysis of <i>C. caudatus</i> Extract on Foodborne Pathogens	62
4.3.1	Time-kill curve of <i>B. cereus</i> (ATCC33019)	63
4.3.2	Time-kill curve of <i>B. subtilis</i> (ATCC6633)	64
4.3.3	Time-kill curve of <i>E.coli</i> O157:H7 (ATCC43895)	65
4.3.4	Time-kill curve of <i>K. pneumoniae</i> (ATCC13773)	66
4.3.5	Time-kill curve of <i>L.monocytogenes</i> (ATCC19112)	67
4.3.6	Time-kill curve of <i>P. mirabilis</i> (ATCC21100)	68
4.3.7	Time-kill curve of <i>P. aeruginosa</i> (ATCC9027)	69
4.3.8	Time-kill curve of <i>S. Typhimurium</i> (ATCC14028)	70
4.3.9	Time-kill curve of <i>S. aureus</i> (ATCC29737)	71
4.3.10	Time-kill curve of <i>C. albicans</i> (ATCC10231)	72
4.4	Stability of <i>C. caudatus</i> Extract at Different pH and Temperature	73
4.5	The Effect of <i>C. caudatus</i> Extract on the Viability and Cellular Injury to the Structures of <i>B. cereus</i> , <i>K. pneumoniae</i> and <i>C. albicans</i>	76
4.5.1	Light sheet electron microscope (LSEM)	76
4.5.2	Transmission electron microscope (TEM)	79
4.6	Identification of Bioactive Compounds in <i>C. caudatus</i> Extract using Liquid Chromatography Mass Spectrometry (LC-MS/MS)	84
4.7	Quantification of Quercetin rhamnoside, Rutin and Quercetin in <i>C.caudatus</i> Extract using High Performance Liquid Chromatography (HPLC)	88
4.8	Antimicrobial Activity of Single Compounds in <i>C.caudatus</i> Extract using Agar Disc-Diffuison Test	88
4.9	Application of <i>C. caudatus</i> Extract on Raw Food Materials	90
4.9.1	Detection of natural microbial populations in raw food materials using selective media agar	90
4.9.2	Natural microbial populations detected in beef meat, chicken meat, shrimp and oyster mushroom	90
4.9.3	Effect of different concentration and exposure time of <i>C. caudatus</i> extract on the number of natural microbial populations in raw food materials	93
4.9.3.1	Beef meat	93
4.9.3.2	Chicken meat	95
4.9.3.3	Shrimp	97
4.9.3.4	Oyster mushroom	99
4.10	Sensory Evaluation of Treated Beef Meat, Chicken Meat, Shrimp and Oyster Mushroom with <i>C. caudatus</i> Extract at Different Concentration and Exposure Time	102
4.10.1	Sensory evaluation of treated beef meat with <i>C.</i>	102

	<i>caudatus</i> extract at different concentration and exposure time	
4.10.2	Sensory evaluation of treated chicken meat with <i>C. caudatus</i> extract at different concentration and exposure time	105
4.10.3	Sensory evaluation of treated shrimp with <i>C. caudatus</i> extract at different concentration and exposure time	107
4.10.4	Sensory evaluation of treated oyster mushroom with <i>C. caudatus</i> extract at different concentration and exposure time	109
4.11	Toxicity of <i>C. caudatus</i> Extract on <i>A. salina</i> spp.	112
5	SUMMARY, CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCH	115
5.1	Summary	115
5.2	Conclusion	116
5.3	Recommendations for Future Research	117
	REFERENCES	118
	APPENDICES	153
	BIODATA OF STUDENT	160
	LIST OF PUBLICATIONS	161

LIST OF TABLES

Table		Page
2.1	Classification of <i>C. caudatus</i> Kunth.	16
2.2	Metabolite compounds identified in <i>C. caudatus</i> Kunth.	20
2.3	Previous studies on pharmacological reports of <i>C. caudatus</i> Kunth.	23
3.1	Microbial species, source and their cultivation medium	38
3.2	Type of media and preparations	39
3.3	HPLC gradient elution for determination of flavonoids in <i>C. caudatus</i> extract	44
3.4	LC-MS/MS gradient elution for compounds identification in <i>C. caudatus</i> extract	45
4.1	Total yield of <i>C. caudatus</i> extract	49
4.2	The yield of <i>C. caudatus</i> extract compared with other previous studies	50
4.3	Inhibition zone, minimum inhibitory concentration (MIC) and minimum bactericidal/fungicidal concentration (MBC/MFC) of <i>C. caudatus</i> extract against foodborne pathogens	52
4.4	Previous studies on the antimicrobial activity of <i>C. caudatus</i> extract	55
4.5	The susceptibility of <i>C. caudatus</i> extract against foodborne pathogens compared with other plant extracts	59
4.6	Stability of <i>C. caudatus</i> extract at different pH on the MIC and MBC/MFC of foodborne pathogens	74
4.7	Stability of <i>C. caudatus</i> extract at different temperature on the MIC and MBC/MFC of foodborne pathogens	75
4.8	List of compounds identified in <i>C. caudatus</i> extract using LC-MS/MS analysis	87
4.9	Antimicrobial activity of single compounds from <i>C. caudatus</i> extract on representative food-borne pathogens	89

4.10	Total plate count and number of natural microbial detected in raw food samples using different selective media agar	92
4.11	Effects of different concentration and exposure time of <i>C. caudatus</i> extract on natural microbial in beef meat	94
4.12	Effects of different concentration and exposure time of <i>C. caudatus</i> extract on natural microbial in chicken meat	96
4.13	Effects of different concentration and exposure time of <i>C. caudatus</i> extract on natural microbial in shrimp	98
4.14	Effects of different concentration and exposure time of <i>C. caudatus</i> extract on natural microbial in oyster mushroom	100
4.15	Sensory acceptability of treated beef meat with <i>C. caudatus</i> extract at different concentration and exposure time	104
4.16	Sensory acceptability of treated chicken meat with <i>C. caudatus</i> extract at different concentration and exposure time	106
4.17	Sensory acceptability of treated shrimp with <i>C. caudatus</i> extract at different concentration and exposure time	108
4.18	Sensory acceptability of treated oyster mushroom with <i>C. caudatus</i> extract at different concentration and exposure time	110
4.19	Toxicity of <i>C. caudatus</i> extract using brine shrimp lethality assay	114

LIST OF FIGURES

Figure		Page
2.1	Incidence rates of food poisoning cases in Malaysia	6
2.2	Mortality rates of food poisoning cases in Malaysia	6
2.3	Whole plant parts of <i>C. caudatus</i> Kunth.	17
2.4	Leaf parts of <i>C. caudatus</i> Kunth.	17
2.5	Flower parts of <i>C. caudatus</i> Kunth.	18
2.6	Fruit parts of <i>C. caudatus</i> Kunth.	18
2.7	Quercetin structure	29
2.8	Catechin structure	30
2.9	Rutin structure	31
2.10	Chlorogenic acid structure	32
3.1	Fresh <i>C. caudatus</i> sample	36
4.1	Time-kill curve plots for <i>B. cereus</i> (0, 3.13, 6.25, 12.5, 25, and 50 mg/ml) following exposure to <i>C. caudatus</i> extract. Values given in the brackets after species are 0× MIC, 1/2× MIC, 1× MIC, 2× MIC, 4× MIC and 8× MIC, as respectively	63
4.2	Time-kill curve plots for <i>B. subtilis</i> (0, 6.25, 12.5, 25 and 50 mg/ml) following exposure to <i>C. caudatus</i> extract. Values given in the brackets after species are 0× MIC, 1/2× MIC, 1× MIC, 2× MIC and 4× MIC, as respectively	64
4.3	Time-kill curve plots for <i>E. coli</i> O157: H7 (0, 6.25, 12.5, 25 and 50 mg/ml) following exposure to <i>C. caudatus</i> extract. Values given in the brackets after species are 0× MIC, 1/2× MIC, 1× MIC, 2× MIC and 4× MIC, as respectively	65
4.4	Time-kill curve plots for <i>K. pneumoniae</i> (0, 6.25, 12.5, 25 and 50 mg/ml) following exposure to <i>C. caudatus</i> extract. Values given in the brackets after species are 0× MIC, 1/2× MIC, 1× MIC, 2× MIC and 4× MIC, as respectively	66

4.5	Time-kill curve plots for <i>L. monocytogenes</i> (0, 6.25, 12.5, 25 and 50 mg/ml) following exposure to <i>C. caudatus</i> extract. Values given in the brackets after species are 0× MIC, 1/2× MIC, 1× MIC, 2× MIC and 4× MIC, as respectively	67
4.6	Time-kill curve plots for <i>P. mirabilis</i> (0, 3.13, 6.25, 12.5, 25 and 50 mg/ml) following exposure to <i>C. caudatus</i> extract. Values given in the brackets after species are 0× MIC, 1/2× MIC, 1× MIC, 2× MIC, 4× MIC and 8× MIC, as respectively	68
4.7	Time-kill curve plots for <i>P. aeruginosa</i> (0, 6.25, 12.5, 25 and 50 mg/ml) following exposure to <i>C. caudatus</i> extract. Values given in the brackets after species are 0× MIC, 1/2× MIC, 1× MIC, 2× MIC and 4× MIC, as respectively	69
4.8	Time-kill curve plots for <i>S. Typhimurium</i> (0, 6.25, 12.5, 25 and 50 mg/ml) following exposure to <i>C. caudatus</i> extract. Values given in the brackets after species are 0× MIC, 1/2× MIC, 1× MIC, 2× MIC and 4× MIC, as respectively	70
4.9	Time-kill curve plots for <i>S. aureus</i> (0, 3.13, 6.25, 12.5, 25 and 50 mg/ml) following exposure to <i>C. caudatus</i> extract. Values given in the brackets after species are 0× MIC, 1/2× MIC, 1× MIC, 2× MIC, 4× MIC and 8× MIC, as respectively	71
4.10	Time-kill curve plots for <i>C. albicans</i> (0, 6.25, 12.5, 25 and 50 mg/ml) following exposure to <i>C. caudatus</i> extract. Values given in the brackets after species are 0× MIC, 0.5× MIC, 1× MIC, 2× MIC and 4× MIC, as respectively	72
4.11	The stained <i>B. cereus</i> viewed under light-sheet electron microscope [(a) control; (b) after treated with <i>C. caudatus</i> extract for 24 h]	78
4.12	Transmission electron micrograph of <i>B. cereus</i> [(c) non-treated; (d) after treated with <i>C. caudatus</i> extract for 24 h]	80
4.13	Transmission electron micrograph of <i>K. pneumoniae</i> [(c) non-treated; (d) after treated with <i>C. caudatus</i> extract for 24 h]	82
4.14	Transmission electron micrograph of <i>C. albicans</i> [(c) non-treated; (d) after treated with <i>C. caudatus</i> extract for 24 h]	83
4.15	Total ion chromatogram of identified compounds in <i>C. caudatus</i> extract	85
4.16	The standard curve of brine shrimp lethality assay after treated with <i>C. caudatus</i> extract for 24 h	113



LIST OF ABBREVIATIONS

amu	-	atomic mass unit
cfu/g	-	colony forming unit/gram
DCD	-	developmental coordination order
EDTA	-	ethylene diamine tetra acetic acid
FRAP	-	ferric reducing ability of plasma
HPLC	-	high performance liquid chromatography
H ₂ O ₂	-	hydrogen peroxide
LC	-	lethality concentration
LC-MS	-	liquid chromatography - mass spectrometry
MAP	-	modified atmosphere packaging
Mg ²⁺	-	magnesium ion
MRSA	-	methicillin resistant <i>Staphylococcus aureus</i>
OH	-	hydroxyl radical
O ₂	-	Oxygen
O ₂ ⁻	-	superoxide anion radicals
Ppm	-	parts per million
ROS	-	reactive oxygen species
USFDA	-	united state of food and drug administration
WHO	-	world health organization

CHAPTER 1

INTRODUCTION

1.1 General Introduction

An illness caused by the consumption of food contaminated with pathogens has a wide economic and public health impact worldwide (Gandhi & Chikindas, 2007). The spoiled foods include lipid oxidation, color changes, off-flavours and off-odours are mainly caused by the growth of microbes and excess of their metabolisms in the food products (Gram et al., 2002). In England and Wales, it has been estimated that there are around 1.3 million reported cases of foodborne illnesses which have caused up to 21,000 people got hospitalized and 500 deaths annually (Adak et al., 2002). Meanwhile, in Australia, about 5.4 million cases were reported with 15,000 people got hospitalized and 120 deaths annually (Soon et al., 2011). In the United States, the reported foodborne cases has shown an increase of approximately 48 million illnesses with 128,000 hospitalizations and 3,000 deaths each year (Scallan et al., 2011). The same trend has been shown in Malaysia where the reported cases of foodborne illnesses had increased from 17.76 cases in 2005 (MOH, 2006) to 47.79 per 100,000 populations in 2013 (MOH, 2014).

In general, foodborne illnesses occur are due to improper food handling and poor hygienic practices, especially during food preparation and storage period (Scott, 2003). Common examples of widely reported foodborne pathogens include *Bacillus cereus*, *Campylobacter*, *Clostridium*, *Escherichia coli* O157:H7, *Listeria monocytogenes*, *Salmonella* and *Staphylococcus aureus* (Singh et al., 2003). Food contamination can occur during post-harvest processing or even earlier which is during pre-harvesting period due to the contamination from a farm environment, water irrigation systems, manure and so on (Doyle & Erickson, 2006). Without proper decontamination techniques, pathogens will keep on growing especially during food storage period and during food preparation caused by cross contamination (Scott, 2003).

In fresh produce foods such as fruits and vegetables, washing with tap or chlorinated water is the common decontamination practice used. The washing treatment is to ensure the removal of soil and other plant debris to improve the appearance of the product (Delaquis et al., 2004) and reduce the microbial proliferation during subsequent storage (Simons & Sanguansri, 1997). As recommended by United States of Food and Drug Administration (USFDA) (1998), the total chlorine concentration allowed for the washing treatment was about 50 - 200 mg/l (pH 6.0 - 7.5) with contact time of 1 - 2 minutes. Chlorinated water is reported to have disinfectant effect on fresh produce (Delaquis et al., 2004; Baur et al., 2004). However, the practice of using chlorinated water is being questioned due to the formation of carcinogenic by-products (trihalomethanes) when chlorine reacts with organic matter such as decayed leaves (Chang et al., 2000). In addition, water treated with high chlorine residues will also produce some odour which might affect the food flavour and smell.

Moreover, food sanitization on meat and poultry products such as cattle, pork, beefs and chicken carcasses have also become a concern. Some bacteria associated with poultry products such as *B. cereus*, *Campylobacter*, *Clostridium*, *L. monocytogenes* and *Salmonella* have been widely reported to cause food spoilage (Hinton Jr et al., 2004). All these bacteria usually come from the farm or during slaughtering process (Solomon et al., 2002). Even though the bacteria's population is at a safe level earlier, they will rapidly multiply especially during storage period if no proper decontamination treatment is applied. Some people believe by chilling these poultry can stop and kill the foodborne pathogens attached, without realizing that the pathogens have a special ability to adjust their intracellular cells in a stress environment. As reported by Capita et al. (2002), even though these poultry foods are refrigerated or frozen prior to cooking, the survival of psychrotrophic pathogens such as *L. monocytogenes* is a food safety concern. A study reported by Al-Nehlawi et al. (2014) revealed the ability of *L. monocytogenes* to spoil a poultry sausage during refrigeration due to their ability to grow anaerobically during refrigeration, thus contaminating the food products.

There are many food decontamination treatments that have been applied to eliminate or substantially decrease bacterial populations both in fresh produce and raw poultries. A number of antimicrobial treatments have been studied including chlorine, organic acids, bacteriocins, hydrogen peroxide, ozonation, irradiation, UV light and many more. All these techniques were applied alone or sometimes through the combination called as hurdle technology. However, those treatments seemed to be ineffective since there were many drawbacks reported such as the deterioration in food quality, health risks, unattractive cost implication and consumer preference (Leistner, 2000). In addition, the use of advanced methods for food treatment may also cause loss of organoleptic properties of food which in turns reduce consumer acceptability (Negi, 2012). Others have also reported that by applying too much techniques will somehow affect the food quality in terms of their loss of nutrient contents, poor appearance and at the same time increase the cost of food processing (Fisher & Phillips, 2006). The application of hurdle technologies in food preservation also initiate food safety issues even though the technique is sometimes successful in controlling the growth of foodborne pathogens (Leistner, 2000). Moreover, reportedly that the long-term use of chlorine sanitizer may cause several carcinogenic effects (Neo et al., 2013).

Hence, an alternative method to eliminate and reduce microbial populations needs to be investigated. With a great consumer concern regarding on synthetic antimicrobials has lead the researchers and food processors to look for more on natural decontamination agents with a broad spectrum of antimicrobial activity (Marino et al., 2001). In this context, plant antimicrobials are gaining wide interest as most of them are classified as GRAS (Generally Recognized as Safe) status which has been discovered to have higher levels of food safety (Alzarkey & Nakahara, 2003). Plants or leafy green vegetables are widely consumed by people worldwide, either eaten fresh or cooked using various methods. These vegetables contain a large amount of phytochemicals such as phenolic acids, flavonoids and aromatic compounds and have been proven to possess a wide range of biological functions including as antioxidant and having antimicrobial activity (Tzortzakis & Economakis, 2007; Gutierrez et al., 2008; Moon et al., 2013).

Cosmos caudatus Kunth. are widely distributed in tropical countries and have long been used in folk medicines. Several bioactive compounds have been isolated from leaves of *C. caudatus* including flavonoids, phenols and alkaloids with reference to several biological properties such as antioxidant, antifungal, antibacterial and anticancer activities (Abas et al., 2006; Pebriana et al., 2008; Salehan et al., 2013). However, in terms of antimicrobial activities, only a few of antifungal compounds were isolated from roots of *C. caudatus* extract have been reported. One of them was hydroxyeugenol, which was proved to inhibit *C. albicans* (Fuzzati et al., 1995). Thus, study was undertaken to discover the antimicrobial activity of *Cosmos caudatus* Kunth. extract against several foodborne pathogens namely *Bacillus cereus*, *Bacillus subtilis*, *Candida albicans*, *Escherichia coli* O157:H7, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Salmonella Typhimurium* and *Staphylococcus aureus*. This study is important to assess the ability of plant extract to decrease, inhibit or eliminate microbial activity which can provide basic knowledge on the development of *C. caudatus* extract as a natural food sanitizer. Moreover, this research was also believed to significantly contribute to the increase in application of other natural plant extracts as food sanitizer or perhaps, in combination with other food preservation methods to ensure the safety of preserved foods.

1.2 Objectives

Therefore, the objectives of this study are:

1. To determine the susceptibility of *Bacillus cereus*, *Bacillus subtilis*, *Escherichia coli* O157:H7, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Salmonella Typhimurium*, *Staphylococcus aureus* and *Candida albicans* by *C. caudatus* extract in term of disc diffusion test, MIC, MBC/MFC and killing time curves.
2. To evaluate the stability of *C. caudatus* extract at different pH and temperature against *Bacillus cereus*, *Bacillus subtilis*, *Escherichia coli* O157:H7, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Salmonella Typhimurium*, *Staphylococcus aureus* and *Candida albicans* in term of MIC and MBC/MFC.
3. To analyse the mode of actions of antimicrobial activity of *C. caudatus* extract on representative foodborne pathogens using light sheet electron microscope (LSEM) and transmission electron microscope (TEM).
4. To identify and quantify the bioactive compounds in *C. caudatus* extract which are responsible for antimicrobial activity using Liquid Chromatography Mass-Spectrometry (LC-MS/MS) and High Performance Liquid Chromatography (HPLC), respectively.
5. To analyse the effect of *C. caudatus* extract on foodborne pathogens in raw food materials including beef meat, chicken meat, shrimp and oyster mushroom.

6. To evaluate the sensory acceptability of raw food materials; beef meat, chicken meat, shrimp and oyster mushroom after treated with *C. caudatus* extract at different concentration and exposure time.
7. To evaluate the toxicity effect of *C. caudatus* extract on eukaryotic cells using brine shrimp (*Artemia salina* spp.) lethality assay.



REFERENCES

- Ababutain, I.M. (2011). Antimicrobial activity of ethanolic extract from some medicinal plant. *Australian Journal of Basic and Applied Sciences*. 5(11): 678-683.
- Abadias, M., Cañamás, T.P., Asensio, A., Anguera, M. & Viñas, I. (2006). Microbial quality of commercial 'Golden Delicious' apples throughout production and shelf-life and Lleda (Catalonia, Spain). *International Journal of Food Microbiology*. 108: 404-409.
- Abadias, M., Alegre, I., Usall, J., Torres, R. & Viñas, I. (2011). Evaluation of alternative sanitizers to chlorine disinfection for reducing foodborne pathogens in fresh-cut apple. *Postharvest Biology and Technology*. 59: 289-297.
- Abas, F., Shaari, K., Lajis, N.H., Israf, D.A. & Kalsom, Y.U. (2003). Antioxidative and radical scavenging properties of the constituents isolated from *Cosmos caudatus* Kunth. *Natural Product Sciences*. 9(4): 245-248.
- Abas, F., Lajis, N.H., Israf, D.A., Khozirah, S. & Umi Kalsom, Y. (2006). Antioxidant and nitric oxide inhibition activities of selected Malay traditional vegetables. *Food Chemistry*. 95: 566-573.
- Abdullah, E., Raus, R.A. & Jamal, P. (2012). Extraction and evaluation of antibacterial activity from selected flowering plants. *American Medical Journal*. 3(1): 27-32.
- Abdullah, A., Dhaliwal, K.K., Roslan, N.N.F., Lee, C.H., Kalaiselvam, M., Radman, M., Saad, Q.H.M., Yusof, K. & Jaarin, K. (2015). The effects of *Cosmos caudatus* (Ulam raja) on detoxifying enzymes in extrahepatic organs in mice. *Journal of Applied Pharmaceutical Science*. 5(01): 082-088.
- Adak, G.K., Long, S.M. & O'Brien, S.J. (2002). Trends in indigenous foodborne disease and deaths, England and Wales: 1992-2000. *GUT*. 51: 832-841.
- Addgene. (2014). Creating bacterial glycerol stocks for long-term storage of plasmids. Retrieved 27 May 2015 from <https://www.addgene.org/plasmids-protocols/create-glycerol-stocks/>
- Agata, N., Ohta, M. & Yokoyama, K. (2002). Production of *Bacillus cereus* emetic toxin (cereulide) in various foods. *International Journal of Food Microbiology*. 73: 23-27.
- Ajaykumar, T.V., Anandarajagopal, K., Sunilson, J.A.J., Arshad, A., Jainaf, R.A.M. & Venkateshan, N. (2012). Anti-inflammatory activity of *Cosmos caudatus*. *International Journal of Universal Pharmacy and Bio Sciences*. ISSN: 2319-8141.

- Akbar, A. & Anal, A.K. (2013). Prevalence and antibiogram study of *Salmonella* and *Staphylococcus aureus* in poultry meat. *Asian Pasific Journal of Tropical Biomedicine*. 3(2): 163-168.
- Alcaraz, L.E., Blanco, S.E., Puig, O.N., Tomas, F. & Ferretti, F.H. (2000). Antibacterial activity of flavonoids against methicillin-resistant *Staphylococcus aureus* strains. *Journal of Theorological Biology*. 205: 23-240.
- Ali, M.A. & Takwa, H.I. (2010). Improve of the quality and shelf-life of minced beef mixed with soy protein by sage (*Saliva officinalis*). *International Food Research Journal*. 17: 1125-1130.
- Allende, A., Selma, M.V., López-Gálvez, F., Villaescusa, R. & Gil, M.I. (2008). Role of commercial sanitizers and washing systems on epiphytic microorganisms and sensory quality of fresh-cut escarole and lettuce. *Postharvest Biology and Technology*. 49: 155-163.
- Alma, M.H., Mavi, A., Yildirim, A., Digrak, M. & Hirata, T. (2003). Screening chemical compositions and *in vitro* antioxidant and antimicrobial activities of the essential oils from *Origanum syriacum* L. growing in Turkey. *Biological and Pharmaceutical Bulletin*. 26: 1725-1729.
- Alwash, M.S., Ibrahim, N. & Ahmad, W.Y. (2013). Identification and mode of action of antibacterial components from *Melastoma malabathricum* Linn. leaves. *American Journal of Infectious Diseases*. 9(2): 46-58.
- Alzarokay, N.S. & Nakahara, K. (2003). Antibacterial activity of extracts from some edible plants commonly consumed in Asia. *International Journal of Food Microbiology*. 80: 223-230.
- Al-Nehlawi, A., Guri, S., Guamis, B. & Saldo, J. (2014). Synergistic effect of carbon dioxide atmosphere and high hydrostatic pressure to reduce spoilage bacteria on poultry sausage. *LWT – Food Science and Technology*. 58: 404-411.
- Amalia, L., Anggadiredja, K., Sukrasno, Fidrianny, I. & Inggriani, R. (2012). Anti-hypertensive potency of wild cosmos (*Cosmos caudatus* Kunth, Asteraceae) leaf extract. *Journal of Pharmacology and Toxicology*. 7(8): 359-368.
- Amna, O.F., Nooraain, H., Noriham, A., Azizah, A.H. & Husna, N. (2013). Acute and oral subacute toxicity study of ethanolic extract of *Cosmos caudatus* leaf in 119neumon dawley rats. *International Journal of Bioscience, Biochemistry and Bioinformatics*. 3(4).
- Andarwulan, N., Batari, R., Sandrasari, D.A., Bolling, B. & Wijaya, H. (2010). Flavonoid content and antioxidant activity of vegetables from Indonesia. *Food Chemistry*. 121: 1231-1235.

- Andrews, J.M. (2001). Determination of minimum inhibitory concentration. *Journal of Antimicrobial Chemotherapy*. 1: 5-16.
- Annegowda, H.V., Anwar, L.N., Mordi, M.N., Ramanathan, S. & Mansor, S.M. (2010). Influence of sonication on the phenolic content and antioxidant activity of *Terminalia catappa* L. leaves. *Pharmacognosy Research*. 2(6): 368-373.
- Apostolidis, E., Karayannakidis, P.D., Kwon, Y.I., Lee, C.M. & Seeram, N.P. (2011). Seasonal variation of phenolic antioxidant mediated α -glucosidase inhibition of *Ascophyllum nodosum*. *Plant Foods and Human Nutrition*. 66: 313-319.
- Arabshahi-D. S., Vishalakshi Devi, D. & Urooj, A. (2007). Evaluation of antioxidant activity of some plant extracts and their heat, pH and storage stability. *Food Chemistry*. 100: 1100-1105.
- Arakawa, H., Maeda, M., Okubo, S. & Shimamura, T. (2004). Role of hydrogen peroxide in bactericidal action of catechin. *Europe Pubmed Central*. 27(3): 277-281.
- Arbianti, R., Utami, T.S., Kurmana, A. & Sinaga, A. (2007). Comparison of antioxidant activity and total phenolic content of *Dillenia indica* leaves extracts obtained using various techniques. *Proceedings of the 14th Regional Symposium on Chemical Engineering*. December 4-5, Yogyakarta, Indonesia.
- Asao, T., Kumeda, Y & Kawai (2003). An extensive outbreak of staphylococcal food poisoning due to low-fat milk in Japan: estimation of enterotoxin A in the incriminated milk and powdered skim milk. *Epidemiology and Infection*. 130: 33-40.
- Atanda, O.O., Akpan, I. & Oluwafemi, F. (2007). The potential of some spice essential oils in the control of *Aspergillus parasiticus* CFR 223 and aflatoxin production. *Food Control*. 18: 601-607.
- Atlas, R.M. (2010). Handbook of microbiological media. CRC Press. Taylor & Francis Group. Boca Raton, Florida, US.
- Bacteriological Analytical Manual (BAM). (1998). 8th Edition, Revision A.
- Balasundram, N., Sundram, K. & Samman, S. (2006). Phenolic compounds in plants and agri-industrial by-products: antioxidant activity, occurrence and potential uses. *Food Chemistry*. 99: 191-203.
- Barlow, S.M. (1990). Toxicological aspects of antioxidants used as food additives. *Food Antioxidants*. 253-308.

- Basma, A.A., Zuraini, Z. & Sasidharan, S. (2011). A transmission electron microscopy study of the diversity of *Candida albicans* induced by *Euphorbia hirta* L. leaf extract *in vitro*. *Asian Pacific Journal of Tropical Biomedicine*. 1(1): 20-22.
- Baur, S., Klaiber, R., Hammes, W.P. & Carle, R. (2004). Sensory and microbiological quality of shredded, packaged iceberg lettuce as affected by pre-washing procedures with chlorinated and ozonated water. *Innovative Food Science and Emerging Technologies*. 5: 45-55.
- Belewu, M.A., Ahmed El-Imam, A.M., Adeyemi, K.D. & Oladunjoye, S.A. (2012). Eucalyptus oil and lemon grass oil: effect on chemical composition and shelf-life of soft cheese. *Environment and Natural Resources Research*. 2(1): 114-118.
- Beltran, D., Selma, M.V., Tudela, J.A. & Gil, M.I. (2005). Effect of different sanitizers on microbial and sensory quality of fresh-cut potato strips stored under modified atmosphere or vacuum packaging. *Postharvest Biology and Technology*. 37: 37-46.
- Bennett, S.D., Walsh, K.A. & Gould, L.H. (2013). Foodborne disease outbreak caused by *Bacillus cereus*, *Clostridium perfringens* and *Staphylococcus aureus* – United States, 1998 – 2008. *Clinical Infectious Diseases*. 57(3): 425-33.
- Berne, S., Pohleven, F., Turk, T. & Sepcic, K. (2008). Induction of fruiting in oyster mushroom (*Pleurotus ostreatus*) by polymeric 3-alkylpyridinium salts. *Mycological Research*. 112: 1085-1087.
- Beuchat, L.R., Nail, B.V., Adler, B.B. & Clavero, M.R.S. (1998). Efficacy of spray application of chlorinated water in killing pathogenic bacteria on raw apples, tomatoes and lettuce. *Journal of Food Protection*. 61: 1305-1311.
- Beuchat, L.R. (2002). Ecological factors influencing survival and growth of human pathogens on raw fruits and vegetables. *Microbes and Infection*. 4: 413-423
- Beuchat, L.R., Adler & Lang. (2004). Efficacy of chlorine and peroxyacetic acid sanitizer in killing *Listeria monocytogenes* on iceberg and romaine lettuce using simulated commercial processing conditions. *Journal of Food Protection*. 67: 1238-1242.
- Bingol, E.B., Cetin, O. & Muratoglu, K. (2011). Effect of lemon juice on the survival of *Salmonella* Enteritidis and *Escherichia coli* in cig kofte (raw meatball). *British Food Journal*. 113(9): 1183-1194.
- Bisha, B. & Brehm-Stecher, B.F. (2010). Combination of adhesive-tape-based sampling and fluorescence *in situ* hybridization for rapid detection of *Salmonella* on fresh produce. *Journal of Vis Exp*. 44: 2308.

- Bolling, B.W., Dolnikowski, G., Blumberg, J.B. & Chen, C.Y. (2010). Polyphenol content and antioxidant activity of California almonds depend on cultivar and harvest year. *Food Chemistry*. 122(3): 819-825.
- Boughalleb, N., Debbabi, N., Jannet, H.B., Mighri, Z. & Mahjoub, M.E. (2005). Antifungal activity of volatile components extracted from leaves, stems and flowers of four plants growing in Tunisia. *Phytopathology Mediterranean*. 44: 307-312.
- Brackett, R.E. (1992). Shelf stability and safety of fresh produce as influenced by sanitation and disinfection. *Journal of Food Protection*. 55: 808-814.
- Brasil, I.M., Gomes, C., Puerta-Gomez, A., Castell-Perez, M.E. & Moreira, R.G. (2012). Polysaccharide-based multi-layered antimicrobial edible coating enhances quality of fresh-cut papaya. *LWT-Food Science and Technology*. 47: 39-45.
- Bunawan, H., Baharum, S.N., Bunawan, S.N., Amin, N.M. & Noor, N.M. (2014). *Cosmos caudatus* Kunth: a traditional medicinal herb. *Global Journal of Pharmacology*. 8(3): 420-426.
- Burt, S. (2004). Essential oils: their antibacterial properties and potential applications in foods-a review. *International Journal of Food Microbiology*. 94: 223-253.
- Busatta, C., Mossi, A.J., Rodrigues, M.R.A., Cansian, R.L. & de Oliveira, J.V. (2007). Evaluation of *Origanum vulgare* essential oils as antimicrobial agent in sausage. *Brazilian Journal of Microbiology*. 38: 610-616.
- Candiracci, M., Citterio, B., Diamantini, G., Blasa, M., Accorsi, A. & Piatti, E. (2011). Honey flavonoids, natural antifungal agents against *Candida albicans*. *International Journal of Food Properties*. 14(4): 799-808.
- Capita, R., Allonso-Calleja, C., del Camino García-Fernández, M. & Moreno, B. (2002). Activity of trisodium phosphate compared with sodium hydroxide wash solutions against *Listeria monocytogenes* attached to chicken skin during refrigerated storage. *Food Microbiology*. 19(1): 57-63.
- Careaga, M., Fernández, E., Dorantes, L., Mota, L., Jaramillo, M.E. & Hernandez-Sanchez, H. (2003). Antibacterial activity of *Capsicum* extract against *Salmonella* Typhimurium and *Pseudomonas aeruginosa* inoculated in raw beef meat. *International Journal of Food Microbiology*. 83: 331-335.
- Carlin, F., Brillard, J., Broussolle, V., Clavel, T., Duport, C., Jobin, M., Guinebretiere, M.H., Auger, S., Sorokine, A. & Nguyen-The, C. (2010). Adaptation of *Bacillus cereus*, an ubiquitous worldwide-distributed foodborne pathogen, to a changing environment. *Food Research International*. 43: 1885-1894.

- Castenmiller, J.J.M., Linssen, J.P.H., Heinonen, I.M., Hopia, A.I., Schwarz, K. & Hollmann, P.C.H. (2002). Antioxidant properties of differently processed spinach products. *Nahrung Journal of Food*. 46: 290-293.
- Caunii, A., Pribac, G., Grozea, I., Gaitin, D. & Samfira, I. (2012). Design of optimal solvents for extraction of bio-active ingredients from six varieties of *Medicagosativa*. *Chemistry Central Journal*. 6: 13-131.
- CDC (Centers for Disease Control and Prevention). (2011). Investigation update: outbreak of *Shiga toxin-producing E. coli* O104 (STEC O104:H4) infections associated with travel to Germany.
- Ceylan, E. & Fung, D.Y.C. (2004). Antimicrobial activity of spices. Food Science Institute, Kansas State University, Manhattan.
- Chahardehi, A.M., Ibrahim, D., Sulaiman, S.F. & Mousavi, L. (2015). Time-kill study of ethyl acetate extract of stinging nettle on *Bacillus subtilis* subsp. *spizizenii* ATCC CRM-6633 strain NRS 231. *Annual Research and Review in Biology*. 6(1): 33-40.
- Chan, S.C., Yau, W.L., Wang, W., Smith, D.K., Sheu, F.S. & Chen, H.M. (1998). Microscopic observations of the different morphological changes caused by anti-bacterial peptides on *Klebsiella penumoniae* and HL-60 leukemia cells. *Journal of Peptide Sciences*. 4: 413-425.
- Chang, C.Y., Hsieh, Y.H., Hsu, S.S., Hu, P.Y. & Wang, K.H. (2000). The formation of disinfection by-products in water treated with chlorine dioxide. *Journal of Hazardous Materials B79*. 89-102.
- Chang, J-M. & Fang, T.J. (2007). Survival of *Escherichia coli* O157:H7 and *Salmonella enterica* serovars Typhimurium in iceberg lettuce and the antimicrobial effect of rice vinegar against *E. coli* O157:H7. *Food Microbiology*. 24(7-8): 745-751.
- Chen, C.Z. & Cooper, S.L. (2002). Interactions between dendrimer biocides and bacterial membranes. *Biomaterials*. 23(16): 3359-68.
- Chen, J.H. & Hotchkiss, J.H. (2003). *Listeria monocytogenes*: low levels equals low risk. *Journal of Food Protection*. 66: 570-7.
- Clifford, M.N. (2000). Chlorogenic acids and other cinnamates-nature, occurrence, dietary burden, absorption and metabolism. *Journal of Science Food Agriculture*. 80: 1033-43.
- Clinical and Laboratory Standrads Institute (CLSI). (2008). Reference method for broth dilution antifungal susceptibility testing of yeasts; approved standard-third edition; CLSI document M27 – A3. Clinical and Laboratory Standards Institue, Wayne, PA, USA.

- Clinical and Laboratory Standards Institute (CLSI). (2012). Reference method for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard-ninth edition; CLSI document M07-A9. Clinical and Laboratory Standards Institute, Wayne, PA, USA.
- Clinical and Laboratory Standards Institute (CLSI). (2012). Reference method for performance standards for antimicrobial disk susceptibility tests; approved standard-eleventh edition; CLSI document M02-A11. Clinical and Laboratory Standards Institute, Wayne, PA, USA.
- Cloete, T.E. (2003). Resistance mechanisms of bacteria to antimicrobial compounds. *Intenational Biodeterioration and Biodegradation*. 51: 277-282.
- Cock, I. (2007). Antibacterial activity of selected Australian native plant extracts. *The Internet Journal of Microbiology*. 4:2.
- Cook, N.C. & Samman, S. (1996). Flavonoids-chemistry, metabolism, cardioprotective effects and dietary sources. *Journal of Nutrients Biochemistry*. 7: 66-76.
- Corbo, M.R., Bevilacqua, A., Campaniello, D., D'Amato, D., Speranza, B. & Sinigaglia, M. (2009). Prolonging microbial shelf life of foods through the use of natural compounds and non-thermal approaches-a review. *International Journal of Food Science and Technology*. 44: 223-241.
- Cowan, M.M. (1999). Plant products as antimicrobial agents. *Clinical Microbiology Revision*. 12: 564-582.
- Cushnie, T.P.M. & Lamb, A.J. (2005). Antimicrobial activity of flavonoids. *International Journal of Antimicrobial Agents*. 26: 343-356.
- Cushnie, T.P.T. & Lamb, A.J. (2011). Recent advances in understanding the antibacterial properties of flavonoids. *International Journal of Antimicrobial Agents*. 38: 99-107.
- Dahiya, P. & Purkayastha, S. (2012). Phytochemical screening and antimicrobial activity of some medicinal plants against multi-drug resistant bacteria from clinical isolates. *Indian Journal of Pharmaceutical Sciences*. 74(5): 443-450.
- Darah, I., Lim, S.H. & Nithianantham, K. (2013). Effects of methanol extract of *Wedelia chinensis* Osbeck (Asteraceae) leaves against pathogenic bacteria with emphasise on *Bacillus cereus*. *Indian Journal of Pharmaceutical Sciences*. 75(5): 533-539.
- Delaquis, P.J., Fukumoto, L.R., Toivonen, P.M.A. & Cliff, M.A. (2004). Implications of wash water chlorination and temperature for the microbiological and sensory properties of fresh-cut iceberg lettuce. *Postharvest Biology and Technology*. 31: 81-91.

- Derakhshan, S., Sattari, M. & Bigdeli, M. (2008). Effect of subinhibitory concentrations of cumin (*Cuminum cyminum* L.) seed essential oil and alcoholic extract on the morphology, capsule expression and urease activity of *Klebsiella 125neumonia*. *International Journal of Antimicrobial Agents*. 32(5): 432-436.
- Derrickson-Tharrington, E., Kendall, P.A. & Sofos, J.N. (2005). Inactivation of *Escherichia coli* O157:H7 during storage or drying of apple slices pretreated with acid solutions. *International Journal of Food Microbiology*. 99: 79-89.
- Deri, A.S. & Rajkumar, J. (2013). *In vitro* antibacterial activity and stability of *Avicennia marina* against urinary tract infection pathogens at different parameters. *Pakistan Journal of Biological Sciences*. 16(19): 1034-1039.
- Deza, M.A., Araujo, M. & Garrido, M.J. (2005). Inactivation of *Escherichia coli*, *Listeria monocytogenes*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* on stainless steel and glass surfaces by neutral electrolyzed water. *Letters in Applied Microbiology*. 40: 341-346.
- Dian-Nashiela, F., Noriham, A., Nooraain, H. & Azizah, A.H. (2014). Antioxidant activity of herbal tea prepared from *Cosmos caudatus* leaves at different maturity stages. *International Food and Research Journal*. 22(3): 1189-1194.
- Dickens, J.A. & Ingram, K.D. (2001). Efficacy of an herbal extract, at various concentrations, on the microbiological quality of broiler carcasses after simulated chilling. *Poultry Science Association*. 10: 194-198.
- Ding, T., Rahman, S.M.E. & Oh, D. (2011). Inhibitory effects of low concentration electrolyzed water and other sanitizers against foodborne pathogens on oyster mushroom. *Food Control*. 22: 318-322.
- Dos Santos, E.C.G., Donnici, C.L., Camargos, E.R.S., de Rezende, A.A., Andrada, E.H.A., Soares, L.A.L., Farias, L.M., Carvalho, M.A.R & Almeida, M.G. (2013). Effects of *Copaifera duckei* Dwyer oleoresin on the cell wall and cell division of *Bacillus cereus*. *Journal of Medical Microbiology*. 62: 1032-1037.
- Doyle, M.E. (2005). Food antimicrobials, cleaners and sanitizers. *Food Research International Briefing*. University of Wisconsin-Madison.
- Doyle, M.P. & Erickson, M.C. (2006). Reducing the carriage of foodborne pathogens in livestock and poultry. *Poultry Science*. 85(6): 960-973.
- Du, H. & Li, H. (2008). Antioxidant effect of Cassia essential oil on deep-fried beef during the frying process. *Meat Science*. 78: 461-468.
- Dubal, Z.B., Paturkar, A.M., Waskar, V.S., Zende, R.J., Latha, C., Rawool, D.B. & Kadam, M.M. (2004). Effect of food grade organic acids on inoculated *S. aureus*, *L. monocytogenes*, *E. coli* and *S. Typhimurium* in sheep/goat meat stored at refrigeration temperature. *Meat Science*. 66: 817-821.

- Durairaj, S., Srinivasan, S. & Lakshmanaperumalsamy, P. (2009). *In vitro* antibacterial activity and stability of garlic extract at different pH and temperature. *Electronic Journal of Biology*. 5(1): 5-10.
- Elgayyar, M., Droughon, F.A., Golden, D.A. & Mount, J.R. (2001). Antimicrobial activity of essential oils from plants against selected pathogenic and saprophytic microorganisms. *Journal of Food Protection*. 64: 1019-1024.
- Erlund, I., Meririnne, E., Alfthan, G. & Aro, A. (2001). Plasma kinetics and urinary excretion of the flavanones naringenin and hesperetin in humans after ingestion of orange juice and grapefruit juice. *The Journal of Nutrition*. 0022-3166/01.
- Ezat, S.W.P., Netty, D. & Sangaran, G. (2013). Paper review of factors, surveillance and burden of foodborne disease outbreak in Malaysia. *Malaysian Journal of Public Health Medicine*. 13(2): 98-105.
- Farber, J.M., Daley, E., Coates, F., Beausoleil, N. & Fournier, J. (1991). Feeding trials of *Listeria monocytogenes* with a nonhuman primate model. *Journal of Clinical Microbiology*. 29(11): 2606-2608.
- Felgines, C., Texier, O., Morand, C., Manach, C., Scalbert, A., Regeat, F. & Remesy, C. (2000). Bioavailability of the flavanone naringenin and its glycoside in rats. *American Journal of Physiol Gastrointestine Liver Physiology*. 279: G1148-G1154.
- Ferguson, D.D., Scheftel, J., Cronquist, A., Smith, K., Woo-Ming, A. & Anderson, A. (2005). Temporally distinct *Escherichia coli* O157 outbreaks associated with alfalfa sprouts linked to a common seed source – Colorado and Minnesota. *Epidemiology Infections*. 133: 439-447.
- Firouzi, R., Shekarforoush, S.S., Nazer, A.H., Borumand, Z. & Jooyandeh, A.R. (2007). Effects of essential oils of oregano and nutmeg on growth and survival of *Yersinia enterocolitica* and *Listeria monocytogenes* in barbecued chicken. *Journal of Food Protection*. 70(11): 2626-2630.
- Fisher, K. & Phillips, C.A. (2006). The effect of lemon, orange and bergamot essential oils and their components on the survival of *Campylobacter jejuni*, *Escherichia coli* O157:H7, *Listeria monocytogenes*, *Bacillus cereus* and *Staphylococcus aureus* *in vitro* and in food system. *Journal of Applied Microbiology*. 101: 1232-1240.
- Foster, J.W. (1995). Low pH adaptation and the acid tolerance response of *Salmonella* Typhimurium. *Criticals Review in Microbiology*. 21(4): 215-37.
- Freidovich, I. (1999). Fundamental aspects of reactive oxygen species or what's the matter with oxygen?. *Annual New York Academy Science*. 893: 13.

- Fukai, T., Marumo, A., Kaitou, K., Kanda, T., Tarada, S. & Nomura, T. (2002). Antimicrobial activity of licorice flavonoids against methicillin-resistant *Staphylococcus aureus*. *Fitoterapia*. 73: 536-539.
- Fuzzati, N., Dyatmiko, W., Rahman, A. & Hostettmann, K. (1995). Phenylpropane derivatives from roots of *Cosmos caudatus*. *Phytochemistry*. 39(2): 409-412.
- Gamboa-Angulo, M.M., Cristobal-Alejo, J., Medina-Baizabal, I.L., Chi-Romero, F., Mendez-Gonzalez, R., Sima-Polanco, P. & May-Pat, F. (2008). Antifungal properties of selected plants from the Yucatan peninsula, Mexico. *World Journal Microbial Biotechnology*. 24: 1955-1959.
- Gandhi, M. & Chikindas, M.L. (2007). *Listeria*: A foodborne pathogen that knows how to survive. *International Journal of Food Microbiology*. 113: 1-15.
- Gangoué-Piéboji, J., Eze, N., Djintchui, A.N., Ngameni, B., Tsbang, N., Pegnyemb, D.E., Biyiti, L., Ngassam, P., Koulla-Shiro, S. & Galleni, M. (2009). The *in vitro* antimicrobial activity of some traditionally used medicinal plants against beta-lactam-resistant bacteria. *The Journal of Infection in Developing Countries*. 3(9): 671-680.
- Ghoshal, S., Krishna Prasad, B.N. & Lakshmi, V. (1996). Antimoebic activity of *Piper longum* fruits against *Entamoeba histolytica* *in vitro* and *in vivo*. *Journal of Ethnopharmacology*. 50: 167-170.
- Gil, M.I., Selma, M.V., Lopez-Galvez, F. & Allende, A. (2009). Fresh-cut product sanitation and wash disinfection: problems and solutions. *International Journal of Food Microbiology*. 134(1-2): 37-45.
- Glass, R.I., Parashar, U.D. & Estes, M.K. (2009). Norovirus gastroenteritis. *The New England Journal of Medicine*. 361: 1776-85.
- Gordon, N.C. & Wareham, D.W. (2010). Multidrug-resistant *Acinetobacter baumannii*: mechanisms of virulence and resistance. *International Journal of Antimicrobial Agents*. 35: 219-226.
- Goyal, A., Singh, J., Jain, S. & Pathak, D.P. (2009). Phytochemicals as potential antimutagens. *International Journal of Pharmaceutical Research and Development*. ISSN 0974-9446.
- Gram, L., Ravn, L., Rasch, M., Bruhn, J.B., Christensen, A.B. & Givskov, M. (2002). Food spoilage-interactions between food spoilage bacteria. *International Journal of Food Microbiology*. 78: 79-97.
- Greene, S.K., Daly, E.R., Talbot, E.A., Demma, L.J., Holzbauer, S. & Patel, N.J. (2008). Recurrent multistate outbreak of *Salmonella* Newport associated with tomatoes from contaminated fields. *Epidemiology Infection*. 136: 157-165.

- Greig, J.D. & Ravel, A. (2009). Analysis of foodborne outbreak data reported internationally for source attribution. *International Journal of Food Microbiology*. 130: 77-87.
- Griffith, C.J., Cooper, R.A., Gilmore, J., Davies, C. & Lewis, M. (2000). An evaluation of hospital cleaning regimes and standards. *Journal of Hospital Infection*. 45: 19-28.
- Guardia, T., Rotelli, A.E., Juarez, A.O. & Pelzer, L.E. (2001). Anti-inflammatory properties of plant flavonoids. Effects of rutin, quercetin and hesperidin on adjuvant arthritis in rat. *II Farmaco*. 53: 421-424.
- Gündüz, G.T. (2013). Antimicrobial activity of sloe berry purees on *Salmonella* spp. *Food Control*. 32: 354-358.
- Gurib-Fakim, A. (2006). Medicinal plants: Traditions of yesterday and drugs of tomorrow. *Molecular Aspects of Medicine*. 27: 1-93.
- Gutierrez, J., Barry-Ryan, C. & Bourke, P. (2008). The antimicrobial efficacy of plant essential oil combinations and interactions with food ingredients. *International Journal of Food Microbiology*. 124: 91-97.
- Gutierrez, J., Bourke, P., Lonchamp, J. & Barry-Ryan, C. (2009). Impact of plant essential oils on microbiological, organoleptic and quality markers of minimally processed vegetables. *Innovative Food Science and Emerging Technologies*. 10: 195-202.
- Harborne, J.B. & Williams, C.A. (2000). Advances in flavonoid research since 1992. *Phytochemistry*. 55: 481-504.
- Hassan, K.I. & Ali, A.A. (2014). Rapid detection system of *Salmonella Typhi* in some commercial brands of milk in Sulamani of Iraq. *International Journal of Advanced Research*. 2(5): 1063-1067.
- Hazwani, Z. (2012). Optimization of extraction parameters of total phenolic compound from *Cosmos caudatus*. Universiti Malaysia Pahang.
- Hendra, R., Ahmad, S., Sukari, A., Yunus Shukor, M. & Oskoueian, E. (2011). Flavonoid analyses and antimicrobial activity of various parts of *Phaleria macrocarpa* (Scheff.) Boerl Fruit. *International Journal of Molecular Sciences*. 12: 3422-3431.
- Higginbotham, K., Burris, K.P., Zivanovic, S., Davidson, P.M. & Stewart Jr., C.N. (2014). Aqueous extracts of *Hibiscus sabdariffa* calyces as an antimicrobial rinse on hot dogs against *Listeria monocytogenes* and methicillin-resistant *Staphylococcus aureus*. *Food Control*. 40: 274-277.

- Hil, P.J.R., Dalmas, E., Nout, M.J.R. & Abee, T. (2009). Soy bean tempe extracts show antibacterial activity against *Bacillus cereus* cells and spores. *Journal of Applied Microbiology*. 109: 137-145.
- Hinton Jr., Cason, J.A. and Ingram, K.D. (2004). Tracking spoilage bacteria in commercial poultry processing and refrigerated storage of poultry carcasses. *International Journal of Food Microbiology*. 91: 155-165.
- Hirasawa, M. & Takada, K. (2004). Multiple effects of green tea catechin on the antifungal activity of antimycotics against *Candida albicans*. *Journal of Antimicrobial Chemotherapy*. 53: 225-229.
- Hirun, N., Dokmaisrijan, S. & Tantishaiyakul, V. (2012). Experimental FTIR and theoretical studies of gallic acid-acetonitrile clusters. *Spectrochim Acta A Molecular Biomolecule Spectroscopy*. 86: 93-100.
- Hodzic, Z., Pasalic, H., Memisevic, A., Srabovic, M. & Poljakovic, M. (2009). The influence of total phenols content on antioxidant capacity in the whole grain extracts. *European Journal of Scientific Research*. 28: 471-477.
- Cosmos caudatus*. (n.d). in *Wikipedia*. Retrieved August 5, 2015, from https://en.wikipedia.org/wiki/Cosmos_caudatus.
- Huda-Faujan, N., Noriham, N., Norrakiah, A.S. & Babji, A.S. (2006). Antioxidant effects of *Cosmos caudatus*, *Polygonum minus* and *Murraya Koenigii* in spent hen burger. *Journal of Animal Science*. 1: 72-77.
- Huda-Faujan, N., Noriham, A., Norrakiah, A.S. & Babji, A.S. (2007). Antioxidative activities of water extracts of some Malaysian herbs. *Asian Food Journal*. 14(1): 61- 68.
- Huda-Faujan, N., Noriham, N., Norrakiah, A. & Babji, A.S. (2009). Antioxidant activity of plant methanolic extracts containing phenolic compounds. *African Journal of Biotechnology*. 8: 484-489.
- Hurst, J.K., Barrette Jr., W.C., Michel, B.R. & Rosen, H. (1991). Hypochlorous acid and myeloperoxidase-catalyzed oxidation of iron-sulfur clusters in bacterial respiratory dehydrogenase. *European Journal of Biochemistry*. 202: 1275-1282.
- Ikhlas, B., Huda, N. & Ismail, N. (2012). Effect of *Cosmos caudatus*, *Polygonum minus* and BHT on physical properties, oxidative process and microbiology growth of quail meatball during refrigeration storages. *Journal of Food Processing and Preservation*. 36: 55-66.
- Ikigai, H., Nakae, T. & Hara, Y. (1993). Bactericidal catechins damage the lipid bilayer. *Biochimica et Biophysica Acta*. 1147: 132-6.

- Ilic, S.B., Konstantinovic, S.S. & Todoravic, Z.B. (2004). Antimicrobial activity of bioactive component from flower of *Linum capitatum* Kit. *Facta Universitatis, series Physics and Chemistry Technology*. 3: 73-77.
- Inatsu, Y., Bari, M.L., Kawasaki, S., Isshiki, K. & Kawamoto, S. (2005). Efficacy of acidified sodium chloride treatments in reducing *Escherichia coli* O157:H7 on Chinese cabbage. *Journal of Food Protection*. 68: 251-255.
- Indira, G. (2014). *In vitro* antifungal susceptibility testing of five antifungal agents against *Dermatophytic species* by CLSI (M38-A) microdilution method. *Clinical Microbiology*. 3: 3.
- Inouye, S., Takizawa, T. & Yamaguchi, H. (2001). Antibacterial activity of essential oils and their major constituents against respiratory tract pathogens by gaseous contact. *Journal of Antimicrobial Chemotherapy*. 47: 565-573.
- Iqbal, S. & Bhanger, M.I. (2006). Effect of season and production location on the antioxidant activity of *Moringa oleifera* leaves grown in Pakistan. *Journal of Food Composition and Analysis*. 102: 544-551.
- Ismail, S. (2000). Sayuran tradisional ulam dan penyedap rasa. Malaysia: Universiti Kebangsaan Malaysia.
- Issa-Zaccharia, A., Kamitani, Y., Morita, K. & Iwasaki, K. (2009). Decontamination of ready-to-eat Japanese mustard green (*Brassica japonica*) from *Escherichia coli* using slightly acidic electrolyzed water. *African Journal of Microbiology Research*. 3(10): 649-653.
- Jami, M., Ghanbari, M., Zunabovic, M., Domig, K.J. & Kneifel, W. (2014). *Listeria monocytogenes* in aquatic food products-a review. *Comprehensive Reviews in Food Science and Food Safety*. 13.
- Janisiewicz, W.J., Conway, W.S., Brown, M.W., Sapers, G.M., Fratamico, P. & Buchanan, R.L. (1999). Fate of *Escherichia coli* O157:H7 on fresh-cut apple tissue and its potential for transmission by fruit flies. *Applied and Environmental Microbiology*. 65(1): 1-5.
- Javadi, N., Abas, F., Hamid, A., Simoh, S., Shaari, K., Ismail, I.S., Mediani, A. & Khatib, A. (2014). GC-MS-based metabolite profiling of *Cosmos caudatus* leaves possessing alpha-glucosidase inhibitory activity. *Food Chemistry*. 79(6): 1130-1136.
- Jayaprakasha, G.K., Jagan Mohan Rao, L. & Sakariah, K.K. (2005). Chemistry and biological activities of *Curcuma longa*. *Trends Food Science and Technology*. 16(12): 533-548.
- Jay-Russell, M.T. (2010). Raw (unpasteurized) milk: are health-conscious consumers making an unhealthy choice?. *Clinical Infectious Disease*. 51(12): 1418-1419.

- Jobim, M.L., Santos, R.C.V., Alves, C.F.D.S., Oliveira, R.M., Mostardeiro, C.P., Sagrillo, M.R., Filho, O.C.S., Garcia, L.F.M., Manica-Cattani, M.F., Ribeiro, E.E. & Manica da Cruz, I.B. (2014). Antimicrobial activity of Amazon *Astrocaryum aculeatum* extracts and its association to oxidative metabolism. *Microbiological Research*. 169: 314-323.
- Johari, N.H. (2006). Kajian aktiviti antimikrob ekstrak tumbuhan *Cosmos caudatus* terhadap *Staphylococcus aureus*, *Bacillus cereus* and *Candida albicans*. Univeristi Malaysia Sabah.
- Jun, H., Kim, J., Bang, J., Kim, H., Beuchat, L.R. & Ryu, J.H. (2013). Combined effects of plant extracts in inhibiting the growth of *Bacillus cereus* in reconstituted infant rice meal. *International Journal of Food Microbiology*. 160(3): 260-266.
- Kajiya, K., Hojo, H., Suzuki, M., Nanjo, F., Kumazawa, S. & Nakayama, T. (2004). Relationship between antibacterial activity of (+) – catechin derivatives and their interaction with a model membrane. *Journal of Agricultural and Food Chemistry*. 52(6): 1514-1519.
- Kanatt, S.R., Chander, R. & Sharma, A. (2010). Antioxidant and antimicrobial activity of pomegranate peel extract improves the shelf-life of chicken products. *International Journal of Food Science and Technology*. 45: 216-222.
- Karimy, M.F., Julendra, H., Hayati, S.N., Sofyan, A., Damayanti, E. & Priyowidodo, D. (2013). Efektifitas ekstrak daun kenikir (*Cosmos caudatus*), daun mengkudu (*Morinda citrifolia*) dan tepung cacing tanah (*Lumbricus rubellus*) dalam sediaan granul larut air sebagai koksidiostat alami terhadap infeksi *Eimeria tanella* pada ayam broiler. *Indonesian Journal of Animal and Veterinary Science*. 18(2): 88-98.
- Keraita, B., Konradsen, F., Drechsel, P. & Abaido, R. (2007). Reducing microbial contamination on wastewater-irrigated lettuce by cessation of irrigation before harvesting. *Tropical Medicinal International Health*. 12: 8-14.
- Khalil, O.A.K., de Faria Oliveira, O.M.M., Velloso, J.C.R., de Quadros, A.U., Dalposso, L.M., Karam, T.K., Mainardes, R.M. & Khalil, N.M. (2012). Curcumin antifungal and antioxidant activities are increased in the presence of ascorbic acid. *Food Chemistry*. 133: 1001-1005.
- Kim, J.B., Kim, J.M., Cho, S.H., Oh, H.S., Choi, N.J. & Oh, D.H. (2011). Toxin genes profiles and toxin production ability of *Bacillus cereus* isolated from clinical and food samples. *Journal of Food Science*. 76(1): 26-29.
- Klaiber, R.G., Baur, S., Wolf, G., Hammes, W.P. & Carle, R. (2005). Quality of minimally processed carrots as affected by warm water washing and chlorination. *Innovative Food Science and Emerging Technologies*. 6: 351-362.

- Klausmeyer, P., Chmurny, G.N., McCloud, T.G., Tucker, K.D. & Shoemaker, R.H. (2004). A novel antimicrobial indolizinium alkaloid from *Aniba panurensis*. *Journal of Natural Product*. 67: 1732-1735.
- Knekt, P., Jarvinen, R. & Seppanen, R. (1997). Dietary flavonoids and the risk of lung cancer and other malignant neoplasms. *American Journal of Epidemiology*. 146: 223-230.
- Kofidis, G., Bosabalidis, A.M. & Moustakas, M. (2003). Contemporary seasonal and altitudinal variations of leaf structural features in *oregano* (*Origanum vulgare* L.). *Annals of Botany*. 92: 635-645.
- Konate, K. & Souza, A. (2010). Polyphenol contents, antioxidant and anti-inflammatory activities of six malvaceae species traditionally used to treat hepatitis B in Burkina Faso. *European Journal Science Research*. 44(4): 570-580.
- Konate, K., Hilou, A., Mavoungou, J.F., Lepengue, A.N., Souza, A., Barro, N., Datte, J.Y., M'Batchi, B. & Nacoulma, O.G. (2012). Antimicrobial activity of polyphenol-rich fractions from *Sida alba* L. (Malvaceae) against co-trimoxazol-resistant bacteria strains. *Annals of Clinical Microbiology and Antimicrobials*. 11:5.
- Koseki, S., Yoshida, K., Isobe, S. & Itoh, K. (2004). Efficacy of acidic electrolyzed water for microbial decontamination of cucumbers and strawberries. *Journal of Food Protection*. 67: 1247-1251.
- Kramer, J.M. & Gilbert, R.J. (1989). *Bacillus cereus* and other *Bacillus* species. Foodborne Bacterial Patogens (Doyle M.P. ed), pp: 21-70. Marcel Dekker, New York.
- Krishnaraju, A.V., Rao, T.V.N., Sundaraju, D., Vanisree, M., Tsay, H.S. & Subbaraja, G.V. (2005). Assessment of bioactivity of Indian medicinal plants using brine shrimp (*Artemia salina*) lethality assay. *International Journal of Applied Science and Engineering*. 3(2): 125-134.
- Kroupitski, Y., Pinto, R., Brandl, M.T., Belausov, E. & Sela, S. (2009). Interactions of *Salmonella enterica* with lettuce leaves. *Journal of Applied Microbiology*. 106: 1876-1885.
- Kumudavally, K.V., Tabassum, A., Radhakrishna, K. & Bawa, A.S. (2011). Effect of ethanolic extract of clove on the keeping quality of fresh mutton during storage at ambient temperature ($25 \pm 2^\circ\text{C}$). *Journal of Food Science and Technology*. 48(4): 466-471.
- Lachumy, S.J.T., Sasidharan, S., Sumathy, V. & Zuraini, Z. (2010). Pharmacological activity, phytochemical analysis and toxicity of methanol extract of *Etlingera elatior* (torch ginger) flowers. *Asian Pasific Journal of Tropical Medicine*. 769-774.

- Lakhanpal, P. & Rai, D.K. (2007). Quercetin: A versatile flavonoid. *Internet Journal of Medical Update*. 2(2): July-Dec.
- Lambert, R.J.N., Skandamis, P.N., Coote, P.J. & Nychas, G.J.E. (2001). A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *Journal of Applied Microbiology*. 91: 453-462.
- Lambert, P.A. (2002). Cellular impermeability and uptake of biocides and antibiotics in Gram-positive bacteria and mycobacteria. *Journal of Applied Microbiology Symposium Supplement*. 92: 46S-54S.
- Lawrie, R.A. & Ledward, D.A. (2006). *Lawrie's Meat Science: Seventh English*, edition ed. Cambridge England: Woodhead Publishing Limited.
- Lazaroaie, M. (2007). The tolerance of gram-negative bacterial strain to saturated and aromatic hydrocarbons. *Romanian Academy of Science*. 59: 2006-2007.
- Lazarova, R. & Krystallis, A. (2010). Consumer's perception of food quality and its relation to the choice of food. Department of Marketing and Statistics. Aarhus Universitet.
- Lee, K.W., Everts, H. & Beynen, A.C. (2004). Essential oils in broiler nutrition. *International Journal of Poultry Science*. 3(12): 738-752.
- Lee, M. (2009). Basic skills in interpreting laboratory data: fourth edition. American Society of Health-System Pharmacist, Inc. US.
- Lee, T.K. & Vairappan, C.S. (2011). Antioxidant, antibacterial and cytotoxic activities of essential oils and ethanol extracts of selected South East Asian herbs. *Journal of Medicinal Plant Research*. 5(21): 5284-5290.
- Lee, K.A., Moon, S., Lee, J., Kim, K., Park, Y. & Paik, H. (2013). Antibacterial activity of a novel flavonoid, 7-O-butyl naringenin, against methicillin-resistant *Staphylococcus aureus* (MRSA). *Food and Science Biotechnology*. 22(6): 1725-1728.
- Leifert, C., Ball, K., Volakakis, N. & Cooper, J.M. (2008). Control of enteric pathogens in ready-to-eat vegetable crops in organic in 'low input' production system: a HACCP-based approach. *Journal of Applied Microbiology*. 105(4): 931-50.
- Leistner, L. (2000). Basic aspects of food preservation by hurdle technology. *International Journal of Food Microbiology*. 55: 181-186.
- Levison, M.E. & Levison J.H. (2009). Pharmacokinetics and pharmacodynamics of antibacterial agents. *Infectious Disease Clinics of North America*. 23(4): 791-vii.

- Leyer, G.J. & Johnson, E.A. (1997). Acid adaptation sensitizes *Salmonella* Typhimurium to hypochlorous acid. *Applied Environmental Microbiology*. 64: 461-467.
- Li, M. & Xu, Z. (2008). Quercetin in a lotus leaves extract may be responsible for antibacterial activity. *Arch of Pharmaceutical Research*. 31(5): 640-644.
- Li, J., Han, Q., Chen, W. & Ye, L. (2012). Antimicrobial activity of Chinese bayberry extract for the preservation of surimi. *Journal of Science and Food Agricultural*. 92(11): 2358-65.
- Lin, Y.T., Labbe, R.G. & Shetty, K. (2004). Inhibition of *Listeria monocytogenes* in fish and meat systems by use of oregano and cranberry phytochemical synergies. *Applied and Environmental Microbiology*. 70: 5672-5678.
- Lis-Balchin, M., Steyrl, H. & Krenn, E. (2003). The comparative effect of novel *Pelargonium* essential oils and their corresponding hydrosols as antimicrobial agents in a model food system. *Phytotherapy Research*. 17(1): 60-65.
- Loh, S.P. & Hadira, O. (2011). *In vitro* inhibitory potential of selected Malaysian plants against key enzymes involved in hyperglycemia and hypertension. *Malaysian Journal of Nutrition*. 17(1): 77-86.
- Lorian, V. (2005). Antimicrobial susceptibility on solid media. In *Antibiotic in Laboratory Medicine*, fifth ed. USA: Lippincott William and Wilkins. Pp. 8-61.
- Lou, Z., Wang, H., Zhu, S., Zhang, M., Gao, Y., Ma, C. & Wang, Z. (2010). Improved extraction and identified by ultra performance liquid chromatography tandem mass spectrometry of phenolic compounds in burdock leaves. *Journal of Chromatography A*. 1217: 2441-2446.
- Lou, Z., Wang, H., Zhu, S., Ma, C. & Wang, Z. (2011). Antibacterial activity and mechanism of action of chlorogenic acid. *Journal of Food Science*. 76(6).
- Lucera, A., Costa, C., Conte, A. & Del Nobile, M.A. (2012). Food applications of natural antimicrobial compounds. *Frontiers in Microbiology*. 3: 287.
- Luend, B.M. & O'Brien, S.J. (2011). The occurrence and prevention of foodborne disease in vulnerable people. *Foodborne Pathogens and Disease*. 8(9).
- Lungu, B., Ricke, S.C. & Johnson, M.G. (2009). Growth, survival, proliferation and pathogenesis of *Listeria monocytogenes* under low oxygen or anaerobic conditions: a review. *Anaerobe*. 15: 7-17,
- Lynch, M.F., Tauxe, R.V. & Hedberg, C.W. (2009). The growing burden of foodborne outbreaks due to contaminated fresh produce: risks and opportunities. *Epidemiology of Infections*. 137: 307-315.

- Mahmood, A.K., Khan, M.S., Khan, M.A. & Bilal, M. (2014). Prevalence of *Salmonella* in diarrheic adult goats in field conditions. *The Journal of Animal and Plant Sciences*. 24(1): 98-102.
- Maldonado, V.A., Glode, M.P. & Bhatia, J. (2014). Consumption of raw and unpasteurized milk and milk products by pregnant women and children. *Pediatrics*. 133(1).
- Mamman, P.H., Mshelia, W.P., Susbatrus, S.C. & Sambo, K.W. (2013). Antibacterial effects of crude extract of *Azadirachta indica* against *Escherichia coli*, *Salmonella* spp. and *Staphylococcus aureus*. *International Journal of Medicine and Medical Sciences*. 5: 14-18.
- Manach, C., Scalbert, A., Morand, C., Remesy, C. & Jimenez, L. (2004). Polyphenols: food sources and bioavailability. *American Journal of Clinical Nutrition*. 79: 727-747.
- Mandalari, G., Bennett, R.N., Bisignano, G., Trombetta, D., Saija, A., Faulds, C.B. & Gasson, M.J. (2007). Antimicrobial activity of flavonoids extracted from bergamot (*Citrus bergamia* Risoo) peel, a byproduct of the essential oil industry. *Journal of Applied Microbiology*. 2056-2064.
- Mandal, S., DebMandal, M., Pal, N.K. & Saha, K. (2011). Inhibitory and killing activities of black tea (*Camellia sinensis*) extract against *Salmonella enterica* serovar Typhi and *Vibrio cholera* O1 biotype E1 Tor serotype Ogawa isolates. *Jundishapur Journal of Microbiology*. 4: 115-121.
- Marino, M., Bersani, C. & Comi, G. (2001). Impedance measurements to study the antimicrobial activity of essential oils from *Lamiaceae* and *Compositae*. *International Journal of Food Microbiology*. 67: 187-195.
- Martins, C.V.B., da Silva, A.A., Neres, A.T.M., Magalhaes, T.F.F., Watanabe, G.A., Modolo, L.V., Sabino, A.A., de Fatima, A. & de Resende, M.A. (2009). Curcumin as a promising antifungal of clinical interest. *Journal of Antimicrobial Chemotherapy*. 63: 337-339.
- Maulidiani, H., Alfi, K., Khozirah, S., Faridah, A., Mahendran, S., Ralf, K., Victor, N. & Nordin, H.L. (2012). Discrimination of three Pegaga (*Cantella*) varieties and determination of growth-lighting effects on metabolite contents based on the chemometry of ¹H nuclear magnetic resonance spectroscopy. *Journal of Agricultural and Food Chemistry*. 60:410-417.
- McCabe-Sellers, B.J. and Beattie, S.E. (2004). Food safety: emerging trends in foodborne illness surveillance and prevention. *Journal of The American Dietetic Association*. 104: 1708-1717.
- McDonnell, G. & Russell, A.D. (1999). Antiseptics and disinfectants: activity, action and resistance. *Clinical Microbiology Reviews*. 12(1): 147-79.

- Mead, P.S., Slutsker, L., Dietz, V., McCaig, L.F., Breese, J.S., Shapiro, C., Griffin, P.M. & Tauxe, R.V. (1999). Food related illness and death in the United States. *Emergence of Infectious Diseases*. 5: 607-625.
- Mediani, A., Abas, F., Ping, T.C., Khatib, A. & Lajis, N.H. (2012). Influence of growth stage and season on the antioxidant constituents of *Cosmos caudatus*. *Plant Foods and Human Nutritions*. 67: 344-350.
- Mediani, A., Abas, F., Khatib, A. and Tan, C.P. (2013). *Cosmos caudatus* as a potential source of polyphenolic compounds: optimisation of oven drying condition and characterisation of its functional properties. *Molecules*. 18(9): 19452-10464.
- Meyer, B.N., Ferrigni, N.R., Putnam, J.E., Jascobsen, L.B., Nichols, D.E. & McLaughlin, J.L. (1982). Brine shrimp: a convenient general bioassay for active plant constituents. *Plant Medica*. 45(1): 31-34.
- Mhone, T.A., Matope, G. & Saidi, P.T. (2012). Detection of *Salmonella* spp. *Candida albicans*, *Aspergillus* spp. and antimicrobial residues in raw and processed cow milk from selected smallholder farms of Zimbabwe. *Veterinary Medicine International*. Doi: 10.1155/2012/301902.
- Michael, A.S., Thompson, C.G. & Abramovitz, M. (1956). *Artemia salina* as a test organism for a bioassay. *Science*. 123: 464.
- Mikusanti, M., Jenie, B.S.L., Syarief, R., Pontjo, B. & Mulyadi, G.T. (2009). Antibacterial activity of *temukunci* tuber (*Kaempferia pandurata*) essential oil against *B. cereus*. *Medicine Journal of Indones*. 18: 10-17.
- Miller, N.J. & Rice-Evans, C.A. (1997). The relative contributions of ascorbic acid and phenolic antioxidants to the total antioxidants to the activity of orange and apple fruit juices and black currant drink. *Food Chemistry*. 60: 331-337.
- Mimarian, M., Toolabi, A., Javadi, A., Narges, K. & Jalalkamali, B. (2012). Evaluating the anti microbial effects of the extract from 3 plants (thyme, pennyroyal, savory) on spore growth, spore elimination and growth of *Bacillus cereus*. *International Journal of Research in Chemistry and Environment*. 2(3): 186-190.
- Ministry of Health (MOH). (2005). *Annual report 2005*. Malaysia.
- Ministry of Health (MOH). (2006). *Annual report 2006*. Malaysia.
- Ministry of Health (MOH). (2007). *Annual report 2007*. Malaysia.
- Ministry of Health (MOH). (2008). *Annual report 2008*. Malaysia.
- Ministry of Health (MOH). (2009). *Annual report 2009*. Malaysia.

- Ministry of Health (MOH). (2010). *Annual report 2010*. Malaysia.
- Ministry of Health (MOH). (2011). *Annual report 2011*. Malaysia.
- Ministry of Health (MOH). (2012). *Annual report 2012*. Malaysia.
- Ministry of Health (MOH). (2013). *Annual report 2013*. Malaysia.
- Moghadamtousi, S.Z., Abdul Kadir, H., Paydar, M., Rouhollahi, E. & Karimian, H. (2014). *Annona muricata* leaves induced apoptosis in A549 cell through mitochondrial-mediated pathway and involvement of NF- κ B. *Complementary and Alternative Medicine*. 14: 299.
- Mohamed, N., Sahlugi, Z., Ramli, E.S.M. & Muhammad, N. (2013). The effects of *Cosmos caudatus* (*Ulam raja*) on dynamic and cellular bone histomorphometry in ovariectomized rats. *BMC Research Notes*. 6: 239.
- Mohanraj, R. (2014). Plant based antibacterial agents. *Plant Based Medicines*. 1-19.
- Molan, P.C. (1992). The antibacterial activity of honey-the nature of antibacterial activity. *Bee World*. 73(1): 5-28.
- Moon, S., Lee, H., Kim, K., Park, Y., Nah, S., Ahn, D. & Paik, H. (2013). Antimicrobial effect of 7-O-butylnaringenin, a novel flavonoid and various natural flavonoids against *Helicobacter pylori* strains. *International Journal of Environmental Research and Public Health*. 10: 5459-5469.
- Moreira, M.R., Ponce, A.G., Del Valle, C.E. & Roura, S.I. (2005). Inhibitory parameters of essential oils to reduce a foodborne pathogen. *LWT-Food Science and Technology*. 38: 565-570.
- Mueller, M., de la Pena, A. & Derendorf, H. (2004). Issues in pharmacokinetics and pharmacodynamics of anti-infective agents: kill curves versus MIC. *Antimicrobial Agents and Chemotherapy*. 48(2): 369-377.
- Muniandy, K., Hassan, Z. & Isa, M.H.M. (2014). Antimicrobial activity of ethanolic extract of *Coleus aromaticus* against common wound pathogens. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 5(3): 1286.
- Murakami, A., Ali, A.M., Mat-Salleh, K., Koshimizu, K. & Ohigashi, H. (2000). Screening for the *in-vitro* anti-tumor-promoting activities of edible plants from Malaysia. *Bioscience, Biotechnology and Biochemistry*. 64(1): 9-16.
- Murcia, A.M., Martinez-Tome, M., Jimenez, A.M., Vera, A.M., Honrubia, M. & Parras, P. (2002). Antioxidant activity of edible fungi (truffles and mushrooms): losses during industrial processing. *Journal of Food Protection*. 65: 1614-1622.

- Murinda, S.E., Nguyen, L.T., Ivey, S.J., Gillespie, B.E., Almeida, R.A., Draughon, F.A. & Oliver, S.P. (2002). Molecular characterization of *Salmonella* spp. Isolated from bulk tank milk and cull dairy cow fecal samples. *Journal of Food Protection*. 65: 1100-1105.
- Mustafa, R.A., Abdul Hamid, A., Mohamed, S. & Abu Bakar, F. (2010). Total phenolic compounds, flavonoids and radical scavenging activity of 21 selected tropical plants. *Journal of Food Science*. 75(1).
- Mytle, N., Anderson, G.L., Doyle, M.P. & Smith, M.A. (2006). Antimicrobial activity of clove (*Syzygium aromaticum*) oil in inhibiting *Listeria monocytogenes* on chicken frankfurters. *Food Control*. 17: 102-107.
- Naczki, M. & Shahidi, F. (2006). Phenolics in cereals, fruits and vegetables: occurrence, extraction and analysis. *Journal of Pharmaceutical and Biomedical Analysis*. 41: 1523-1542.
- Naidu, A.S. (2000). Phyto-phenols. *Natural Food Antimicrobial Systems*. Pp. 265-295. Boca Raton, Florida: CRC Press.
- Nascimento, G.G.F., Locatelli, J., Freitas, P.C. & Silva, G.L. (2000). Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. *Brazilian Journal of Microbiology*. 31(4): 247-256.
- NCCLS (2004). Reference method for antifungal disk diffusion susceptibility testing of yeasts; approved guideline. NCCLS document M44-A. National Committee for Clinical and Laboratory Standards, Wayne, PA, USA.
- Nebahat, O., Gulmez, M., Vatansever, L. & Guven, A. (2008). Application of antimicrobial ice for extending shelf life of fish. *Journal of Food Protection*. 71(1): 218-222.
- Nedorostova, L., Kloucek, P., Kokoska, L., Stolcova, M. & Pulkrabek, J. (2009). Antimicrobial properties of selected essential oils in vapour phase against foodborne bacteria. *Food Control*. 20: 157-160.
- Negi, P.S. (2012). Plant extracts for the control of bacterial growth: Efficacy, stability and safety issues for food application. *International Journal of Food Microbiology*. 156: 7-17.
- Neo, S.Y., Lim, P.Y., Phua, L.K., Khoo, G.H., Kim, S.J., Lee, S.C. & Yuk, H.G. (2013). Efficacy of chlorine and peroxyacetic acid on reduction of natural microflora, *Escherichia coli* O157: H7, *Listeria monocytogenes* and *Salmonella* spp. on mung bean sprouts. *Food Microbiology*. 36: 475-480.
- Newkirk, R., Hedberg, C. & Bender, J. (2011). Establishing a milkborne disease outbreak profile: potential food defense implications. *Foodborne Pathogens Diseases*. 8(3): 433-437.

- Newman, S.E. (2004). Disinfecting irrigation water for disease management. In 'Proceedings of the 20th Annual Conference on Pest Management on Ornamentals'. Society of America, Florists, San Jose, Canada.
- Nitiema, L.W., Savadogo, A., Simpore, J., Dianou, D. & Traore, A.S. (2012). *In vitro* antimicrobial activity of some phenolic compounds (coumarin and quercetin) against gastroenteritis bacterial strains. *International Journal of Microbiological Research*. 3(3): 183-187.
- Norhanom, A.W., Ashril, Y. & Mustafa, A.M. (1990). Phytochemicals and Biopharmaceutics from the Malaysian Rainforest. FRIM, Kuala Lumpur, Malaysia.
- Norlaili, A.H., Roselina, K.T. & Muhammad, A.S. (2014). Effect of *Cosmos caudatus* Kunth. (*Ulam raja*) aqueous and dry extracts on the physiochemical and functional properties and sensory acceptability of herbal yellow alkaline noodles. *Malaysian Journal of Nutritions*. 20(3): 403-415.
- Nuengchamngong, N., Krittasilp, K. & Ingkaninan, K. (2009). Rapid screening and identification of antioxidants in aqueous extracts of *Houttuynia cordata* using LC-ESI-MS coupled with DPPH assay. *Food Chemistry*. 117: 750-756.
- Nurul, H., Ruzita, A. & Aronal, A.P. (2010). The antioxidant effect of *Cosmos caudatus* and *Polygonum minus* in refrigerated meatballs. *International Food Research Journal*. 17: 893-904.
- Nyenje, M.E., Odjadjare, C.E., Tanih, N.F., Green, E. & Ndip, R.N. (2012). Foodborne pathogens recovered from ready-to-eat foods from roadside cafeterias and retail outlets in Alice, Eastern Cape Province, South Africa: public health implications. *International Journal of Environmental Research Public Health*. 9: 2608-2619.
- Okigbo, R.N. & Mmeka, E.C. (2008). Antimicrobial effects of three tropical plant extracts on *Staphylococcus aureus*, *Escherichia coli* and *Candida albicans*. *African Journal Traditional, Complementary and Alternative Medicines*. 5(3): 226-229.
- Oliver, S.P., Jayarao, B.M. & Almeida, R.A. (2005). Foodborne pathogens in milk and the dairy farms environment: food safety and public health implications. *Foodborne Pathogens and Disease*. 2 (2).
- Olthof, M.R., Hollman, P.C.H. & Katan, M.B. (2001). Chlorogenic acid and caffeic acid are absorbed in human. *Journal of Nutrition*. 131(1): 66-71.
- Olugu, S.V., Nyegue, M., Kamga, P.B., Bayoi, J.R. & Etoa, F.X. (2014). Activity of conessine at various temperature and pH on inhibition of germination of *Bacillus cereus* and *Bacillus stearothermophilus* spores. *African Journal of Microbiology Research*. 8(4): 320-326.

- Omidbeygi, M., Barzegar, M., Hamidi, Z. & Naghdibadi, H. (2007). Antifungal activity of thyme, summer savory and clove essential oils against *Aspergillus flavus* in liquid medium and tomato paste. *Food Control*. 18: 1518-1523.
- Ong, H.C. & Norzalina, J. (1999). Malay herbal medicine in Gemencheh, Negeri Sembilan, Malaysia. *Fitoterapia*. 70: 10-14.
- Orhan, D.D., Ozcelik, B., Ozgen, S. & Ergun, F. (2010). Antibacterial, antifungal and antiviral activities of some flavonoids. *Microbiological Research*. 165: 496-504.
- Othman, M., Loh, H.S., Wiart, C., Khoo, T.J., Lim, K.H. & Ting, K.N. (2011). Optimal methods for evaluating antimicrobial activities from plant extracts. *Journal of Microbiological Methods*. 84: 161-166.
- Oussalah, M., Caillet, S., Saucier, L. & Lacroix, M. (2006). Antimicrobial effects of selected plant essential oils on the growth of four pathogenic bacteria: *E. coli* O157:H7, *Salmonella* Typhimurium, *Staphylococcus aureus* and *Listeria monocytogenes*. *Food Control*. 18(5): 414-420.
- Palumbo, M. & Harris, L.J. (2011). Microbiological food safety of olive oil: a review of the literature. *Olive Center*.
- Pankey, G.A. & Ashcraft, D.S. (2009). *In vitro* antibacterial activity of tigecycline against resistant Gram-negative bacilli and enterococci by time-kill assay. *Diagnostic Microbiology and Infectious Disease*. 64: 300-304.
- Paolillo, R., Carratelli, C.R. & Rizzo, A. (2011). Effect of resveratrol and quercetin in experimental infection by *Salmonella enterica* serovar Typhimurium. *International Immunopharmacology*. 11: 149-156.
- Parekh, J., Jadeja, D. & Chanda, S. (2005). Efficacy of aqueous and methanol extracts of some medicinal plants for potential antibacterial activity. *Turkish Journal of Biology*. 29: 203-210.
- Parekh, J. & Chanda, S. (2008). Phytochemicals screening of some plants from western region of India. *Plant Archives*. 8: 657-662.
- Park, C.M., Taormina, P.J. & Beuchat, L.R. (2000). Efficacy of allyl isothiocyanate in killing enterohemorrhagic *Escherichia coli* O157:H7 on alfalfa seeds. *International Journal of Food Microbiology*. 56: 13-20.
- Park, H., Hung, Y. & Chung, D. (2004). Effects of chlorine and pH on efficacy of electrolyzed water for inactivating *Escherichia coli* O157:H7 and *Listeria monocytogenes*. *International Journal of Food Microbiology*. 91: 13-18.
- Parnell, T.L., Harris, L.J. & Suslow, T.V. (2005). Reducing *Salmonella* on cantaloupes and honeydew melons using wash practices applicable to postharvest handling,

- foodservice and consumer preparation. *International Journal of Food Microbiology*. 99: 59-70.
- Pasvolsky, R., Zakin, V., Ostrova, I. & Shemesh, M. (2014). Butyric acid released during milk lipolysis triggers biofilm formation of *Bacillus* species. *International Journal of Food Microbiology*. 181: 19-27.
- Patil, D.G. & Nikam, S.V. (2013). *In vitro* antimicrobial, antioxidant activity and phytochemical analysis of *Cosmos caudatus* (wild cosmos). *Universal Journal of Pharmacy*. 2(6): 64-70.
- Pebriana, R.B., Wardhani, B.W.K., Widayanti, E., Wijayanti, N.L.S., Wijayanti, T.R., Riyanto, S. & Meiyanto, E. (2008). Apoptotic effect of Kenikir leaves (*Cosmos caudatus* Kunth.) methanolic extract on breast cancer cell line. Cancer Chemoprevention Research Center, Universitas Gadjah Mada Yogyakarta.
- Pei, R., Zhou, F., Ji, B. & Xu, J. (2009). Evaluation of combined antibacterial effects of eugenol, cinnamaldehyde, thymol and carvacrol against *E. coli* with an improved method. *Journal of Food Science*. 74(7).
- Pereira, V., Dias, C., Vasconcelos, M.C., Rosa, E. & Saavedra, M.J. (2014). Antibacterial activity and synergistic effects between *Eucalyptus globulus* leaf residues (essential oils and extracts) and antibiotics against several isolates of respiratory tract infections (*Pseudomonas aeruginosa*). *Industrial Crops and Products*. 52: 1-7.
- Perumal, V., Hamid, A.A., Ismail, A., Saari, K., Abas, F., Ismail, I.S., Maulidiani, Lajis, N.H. & Khatib, A. (2014). Effects of *Cosmos caudatus* Kunth leaves on the lipid profile of a hyperlipidemia-induced animal model. *Journal of Food Chemistry and Nutrition*. 2(1): 43-51.
- Perumalla, A.V.S. & Hettiarachchy, N.S. (2011). Green tea and grape seed extracts. Potential application in food safety and quality. *Food Research International*. 44(4): 827-839.
- Phillips, C.A., Laird, K. & Allen, S.C. (2012). The use of Citri-VTM – An antimicrobial citrus essential oil vapour for the control of *Penicillium chrysogenum*, *Aspergillus niger* and *Alternaria alternate* *in vitro* and on food. *Food Research International*. 47: 310-314.
- Pimentel, R.B.Q., Costa, C.A., Albuquerque, P.M. & Junior, S.D. (2013). Antimicrobial activity and rutin identification in honey produced by the stingless bee *Melipona compressipes manausensis* and commercial honey. *BMC Complementary and Alternative Medicine*. 13:151.
- Pitella, F., Dutra, R.C., Junior, D.D., Lopes, M.T.P. & Barbosa, N.R. (2009). Antioxidant and cytotoxic activities of *Cantella asiatica* (L.) urban. *International Journal of Molecular Sciences*. 10(9): 3713-3721.

Plant Resources of South East Asia (PROSEA). (1993). 5: 1-3.

Plazonic, A., Bucar, F., Males, Z., Mornar, A., Nigovic, B. & Kujundzic, N. (2009). Identification and quantification of flavonoids and phenolic acids in Burr Parsley (*Caucalis platycarpos* L.), using high performance liquid chromatography with diode array detection and electrospray ionization mass spectrometry. *Molecules*. 14: 2466-2490.

Prakash, B., Shukla, R., Singh, P., Kumar, A., Mishra, P.K. & Dubey, N.K. (2010). Efficacy of chemically characterized *Piper betle* L. essential oil against fungal and aflatoxin contamination of some edible commodities and its antioxidant activity. *International Journal of Food Microbiology*. 142: 114-119.

Proestos, C., Boziaris, I.S., Nychas, G-J.E. & Komaitis, M. (2006). Analysis of flavonoids and phenolic acids in Greek aromatic plants: investigation of their antioxidant capacity and antimicrobial activity. *Food Chemistry*. 95: 664-671.

Pui, C.F., Wong, W.C., Chai, L.C., Lee, H.Y., Noorlis, A., Zainazor, T.C.T., Tang, J.Y.H., Ghazali, F.M., Cheah, Y.K., Nakaguchi, Y., Nishibuchi, M. & Radu, S. (2011). Multiplex PCR for the concurrent detection and differentiation of *Salmonella* spp., *Salmonella* Typhi and *Salmonella* Typhimurium. *Tropical Medicine and Health*. 39(1): 9-15.

Puupponen-Pimia, R., Nohynek, L., Meier, C., Kahkonen, M., Heinonen, M. & Hopia, A. (2001). Antimicrobial properties of phenolic compounds from berries. *Journal of Applied Microbiology*. 90: 494-507.

Radman, H.M., Yusof, K., Mohd Saad, Q., Wan Ngah, W.Z. & Abdullah, A. (2014). The effect of *Ulam raja* (*Cosmos caudatus*) on drug-metabolizing enzymes, lipid peroxidation and antioxidant status in mice liver. *International Journal of Pharm Tech Research*. 6(4): 1213-1225.

Ragaert, P., Devlieghree, F. & Debevere, F. (2007). Role of microbiological and physiological spoilage mechanisms during storage of minimally processed vegetables. *Postharvest Biology and Technology*. 44: 185-194.

Ragasa, C.Y., Nacpil, Z.D., Penalosa, B.A., Coll, J.C. & Rideout, J.A. (1999). Antimutagen and antifungal compounds from *Cosmos caudatus*. *Philippines Journal Science*. 126: 199.

Ragasa, C.Y., Ha, H.K.P., Hasika, M., Maridable, J.B., Gaspillo, P.D. & Rideout, J.A. (2008). Antimicrobial and cytotoxic terpenoids from *Cymbopogon citratus* Stapf. *Phillipines Scientists*. 45: 111-122.

Rajabi, S., Ramazani, A., Hamidi, M. & Naji, T. (2015). *Artemia salina* as a model organism in toxicity assessment of nanoparticles. *DARU Journal of Pharmaceutical Sciences*. 23:20.

- Rajeh, M.A.B., Zuraini, Z., Sasidharan, S., Latha, L.Y. & Amutha, S. (2010). Assessment of *Euphorbia hirta* L. leaf, flower, stem and root extracts for their antibacterial and antifungal activity and brine shrimp lethality. *Molecules*. 15: 6008-6018.
- Ramachandran, S., Vamsikrishna, M., Gowthami, K.V., Heera, B. & Dhanaraju, M.D. (2011). Assessment of cytotoxic activity of *Agave cantula* using brine shrimp (*Artemia salina*) lethality bioassay. *Asian Journal of Scientific Research*. 4(1): 90-94.
- Ramos, F.A., Takaishi, Y., Shirotori, M., Kawaguchi, Y., Tsuchiya, K., Shibata, H., Higuti, T., Tadokoro, T. & Takeuchi, M. (2006). Antibacterial and antioxidant activities of quercetin oxidation products from Yellow Onion (*Allium cepa*) skin. *Journal of Agricultural and Food Chemistry*. 54: 3551-3557.
- Ranasinghe, L., Jayawardena, B. & Abeywickrama, K. (2002). Fungicidal activity of essential oils of *Cinnamomum zeylanicum* (L.) and *Syzygium aromaticum* (L.) Merr et L.M. Perry against crown rot and anthracnose pathogens isolated from banana. *Letters in Microbiology*. 35: 205-211.
- Rasdi, N.H.M., Samah, O.A., Sule, A. & Ahmed, Q.U. (2010). Antimicrobial studies of *Cosmos caudatus* Kunth. (Compositae). *Journal of Medicinal Plants Research*. 4: 669-673.
- Rauha, J.P., Remes, S., Heinonen, M., Hopia, A., Kahkonen, M., Kujala, T., Pihlaja, K., Vuorela, H. & Vuorela, P. (2000). Antimicrobial effects of Finnish plant extracts containing flavonoids and other phenolic compounds. *International Journal of Food Microbiology*. 56: 3-12.
- Reihani, S.F.S. & Azhar, M.E. (2012). Antioxidant activity and total phenolic content in aqueous extracts of selected traditional Malay salads (*Ulam*). *International Food Research Journal*. 19(4): 1439-1444.
- Reihani, S.F.S., Tan, T.C., Huda, N. & Easa, A.M. (2014). Frozen storage stability of beef patties incorporated with extracts from *Ulam raja* leaves (*Cosmos caudatus*). *Food Chemistry*. 155: 17-23.
- Ren, W., Qiao, Z., Wang, H., Zhu, L. & Zhang, L. (2003). Flavonoids: Promising anticancer agents. *Medicinal Plant Research Reviews*. 23(4): 519-534.
- Rice-Evans, C.A., Miller, N.J. & Paganga, G. (1996). Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radical Biology and Medicine*. 20: 933-56.
- Rios, J.L. & Recio, M.C. (2005). Medicinal plants and antimicrobial activity. *Journal of Ethnopharmacol*. 100: 80-84.

- Rosyid, T.A., Karim, R., Adzahan, N.M. & Ghazali, F.M. (2011). Antibacterial activity of several Malaysian leaves extracts on the spoilage bacteria of yellow alkaline noodles. *African Journal of Microbiology Research*. 5(8): 898-903.
- Rufus, P.G., Mohamed, N. & Shuid, A.N. (2015). *Cosmos caudatus* enhances fracture healing in ovariectomised rats: a preliminary biochemical evaluation. *International Journal of Applied Research in Natural Products*. 8(1): 12-19.
- Rukayadi, Y. & Hwang, J.K. (2006). *In vitro* activity of xanthorrhizol against *Streptococcus mutans* biofilms. *Letters in Applied Microbiology*. 42: 400-404.
- Rukayadi, Y., Shim, J.S. & Hwang, J.K. (2008). Screening of Thai medicinal plants for anticandidal activity. *Mycoses*. 51: 308-312.
- Rukayadi, Y., Lau, K.Y., Zainin, N.S., Zakaria, M. & Abas, F. (2013). Screening of antimicrobial activity of tropical edible medicinal plant extracts against five standard microorganisms for natural food preservatives. *International Food Research Journal*. 20(5): 2905-2910.
- Saftner, R.A., Bai, J.H., Abbott, J.A. & Lee, Y.S. (2003). Sanitary dips with calcium propionate, calcium chloride or a calcium amino acid chelate maintain quality and shelf stability of fresh-cut honeydew chunks. *Postharvest Biology and Technology*. 29: 257-269.
- Sahgal, G., Ramanathan, S., Sasidharan, S., Mordi, M.N., Ismail, S. & Mansor, S.M. (2010). Brine shrimp lethality assay and acute oral toxicity studies on *Swietenia mahagoni* (Linn.) Jacq. seed methanolic extract. *Pharmacognosy Research*. 2(4): 215-220.
- Salehan, N.M., Meon, S. & Ismail, I.S. (2013). Antifungal activity of *Cosmos caudatus* extracts against seven economically important plant pathogens. *International Journal of Agriculture and Biology*. 15: 864-870.
- Samy, R.P. (2005). Antimicrobial activity of some medicinal plants from India. *Fitoterapia*. 76: 697-699.
- Sánchez-Rabaneda, F., Jáuregui, O., Lamuela-Raventós, R. M., Viladomat, F., Bastida, J. & Codina, C. (2004). Qualitative analysis of phenolic compounds in apple pomace using liquid chromatography coupled to mass spectrometry in tandem mode. *Rapid Communications in Mass Spectrometry : RCM*. 18(5): 553-563.
- Sanmee, R., Dell, B., Lumyong, P., Izumori, K. & Lumyong, S. (2003). Nutritive value of popular wild edible mushrooms from northern Thailand. *Food Chemistry*. 84: 527-532.
- Sarker, S.D. & Nahar, L. (2012). Natural products isolation: methods in molecular biology. *Springer Science + Business Media*. Vol. 864. Pp. 33-34.

- Satta, G., Cornaglia, G., Mazzariol, A., Golini, G., Valisena, S. & Fontana, R. (1995). Target for bacteriostatic and bactericidal activities of β -lactam antibiotics against *Escherichia coli* resides in different penicillin-binding proteins. *Antimicrobial Agents Chemotherapy*. 39: 812-818.
- Scallan, E., Hoekstra, R.M., Angulo, F.J., Tauxe, R.V., Widdowson, M., Roy, S.L., Jones, J.L. & Griffin, P.M. (2011). Foodborne illness acquired in the United States – major pathogens. *Emerging Infectious Diseases*. 17(1): 7-15.
- Schlesser, J.E., Gerdes, R., Ravishankar, S., Madsen, K., Mowbray, J. & Teo, A.Y. (2006). Survival of a five-strain cocktail of *Escherichia coli* O157:H7 during the 60-day aging period of cheddar cheese made from unpasteurized milk. *Journal of Food Protection*. 69(5): 990-998.
- Scott, E. (2003). Food safety and foodborne disease in 21st century homes. *The Canadian Journal of Infectious Diseases*. 14(5): 277-280.
- Selma, M.V., Allende, A., Lopez-Galvez, F., Conesa, M.A. & Gil, M.I. (2008). Disinfection potential of ozone, ultraviolet-c and their combination in wash water for the fresh-cut vegetable industry. *Food Microbiology*. 25: 809-814.
- Sengun, I.Y. & Karapinar, M. (2004). Effectiveness of lemon juice, vinegar and their mixture in the elimination of *Salmonella* Typhimurium on carrots (*Daucus carota* L.). *International Journal of Food Microbiology*. 96: 301-305.
- Seow, Y.X., Yeo, C.R., Chung, H.L. & Yuk, H.G. (2014). Plant essential oils as active antimicrobial agents. *Critical Reviews in Food Science and Nutrition*. 54: 625-644.
- Shan, B., Cai, Y., Brooks, J.D. & Corke, H. (2007). The *in vitro* antibacterial activity of dietary spice and medicinal herb extracts. *International Journal of Food Microbiology*. 117: 112-119.
- Shimamura, T., Zhao, W.H. & Hu, Z.Q. (2007). Mechanism of action and potential for use of tea catechin as an anti-infective agent. *Anti-infective Agents in Medicinal Chemistry*. 6: 57-62.
- Shu, Y., Liu, Y., Li, L., Feng, J., Lou, B., Zhou, X. & Wu, H. (2011). Antibacterial activity of quercetin on oral infectious pathogens. *African Journal of Microbiology Research*. 5(30): 5358-5361.
- Shui, G., Leong, L.P. & Wong, S.P. (2005). Rapid screening and characterization of antioxidants of *Cosmos caudatus* using liquid chromatography coupled with mass spectrometry. *Journal of Chromatography B*. 827: 127-138.
- Shuib, N.H., Shaari, K., Khatib, A., Maulidiani, Kneer, R., Zareen, S., Raof, S.M., Lajis, N.H. & Neto, V. (2011). Discrimination of young and mature leaves of

- Melicope ptelefolia* using ¹HNMR and multivariate data analysis. *Food Chemistry*. 126: 640-645.
- Silva Jr., I.F., Oliveira, R.G., Soares, I.M., Alvim, T.D.C., Ascencio, S.D. & Oliveira, D.T. (2014). Evaluation of acute toxicity, antibacterial activity and mode of action of the hydroethanolic extract of *Piper umbellatum* L. *Journal of Ethnopharmacology*. 151: 137-143.
- Simons, L.K. & Sanguansri, P. (1997). Advances in the washing of minimally processed vegetables. *Food Australia*. 49: 75-80.
- Singh, N., Singh, R.K., Bhunia, A.K. & Stroshine, R.L. (2002). Efficacy of chlorine dioxide, ozone and thyme essential oil or sequential washing in killing *Escherichia coli* O157:H7 on lettuce and baby carrots. *Lebensmittel-Wissenschaft and Technology*. 35: 720-729.
- Singh, A., Singh, R.K., Bhunia, A.K. & Singh, N. (2003). Efficacy of plant essential oils as antimicrobial agents against *Listeria monocytogenes* in hotdogs. *Food Science and Technology*. 36: 787-794.
- Singleton, V.L., Orthofer, R. & Lamuela-Raventos, R.M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*. 299: 152-178.
- Škerget, M., Majhenić, L., Bezjak, M. & Knez, Z. (2009). Antioxidant, radical scavenging and antimicrobial activities of red onion (*Allium cepa* L.) skin and edible part extracts. *Chemical and Biochemistry of England Q*. 23(4): 435-444.
- Soares, C.M., Kabuki, D.Y. & Kuaye, A.Y. (2012). Growth of enterotoxin producing *Bacillus cereus* in meat substrates at 10°C and 30°C. *Brazilian Journal of Microbiology*. 1401-1405.
- Soderstrom, A., Osterberg, P., Lindqvist, A., Jonsson, B., Lindberg, A. & Blide Ulander, S. (2008). A large *Escherichia coli* O157 outbreak in Sweden associated with locally produced lettuce. *Foodborne Pathogens Disease*. 5: 339-349.
- Sohn, H.Y., Son, K.H., Kwon, C.S., Kwon, G.S. & Kang, S.S. (2004). Antimicrobial and cytotoxic activity of 18 prenylated flavonoids isolated from medicinal plants: *Morus alba* L., *Morus mongolica* Schneider, *Broussonetia papyrifera* (L.) Vent., *Sophora flavescens* Ait and *Echinosophora koreensis* Nakai. *Phytomedicine*. 11: 666-672.
- Solomon, E.B., Yaron, S. & Matthews, K.R. (2002). Transmission of *Escherichia coli* O157:H7 from contaminated manure and irrigation water to lettuce plant tissue and its subsequent internalization. *Applied of Environmental Microbiology*. 68: 397-400.

- Solomon, O.O., Olufemi, O.O. & Toyosi, A.T. (2014). Organoleptic and chemical evaluations of suya (an intermediate moisture meat) cured with *Ocimum gratissimum* extract. *Nova Journal of Engineering and Applied Sciences*. 2(2): 1-7.
- Soon, J.M., Singh, H. & Baines, R. (2011). Foodborne diseases in Malaysia: A review. *Food Control*. 22(6): 823-830.
- Soylu, S. (2006). Accumulation of cell-wall bound phenolic compounds and phytoalexin in *Arabidopsis thaliana* leaves following inoculation with pathovores of *Pseudomonas syringae*. *Plant Sciences*. 170: 942-952.
- Srinu, B., Kumar, A.V., Kumar, M.S., Narayana, B.V.L. & Rao, T.M. (2012). Assessment of microbiological quality and associated health risks of raw milk sold in and around Hyderabad city. *International Journal of Pharma and Bio Sciences*. 3(4): 609-614.
- Stapleton, J.T. Williams, C.F. & Xiang, J. (2004). Minireview: GB virus type C: a beneficial infection?. *Journal of Clinical Microbiology*. 4(2): 3915-3919.
- Stefan, M., Zamfirache, M., Padurariu, C., Truta, E. & Gostin, I. (2013). The composition and antibacterial activity of essential oils in three *Oscimum* species growing in Romania. *Central European Journal of Biology*. 8(6): 600-608.
- Stepanovic, S., Antic, N., Dakic, I. & Svabic-Vlahovic, M. (2003). *In vitro* antimicrobial activity of propolis and synergism between propolis and antimicrobial drugs. *Microbiological Research*. 158: 353-357.
- Stivarius, M.R., Pohlman, F.W., McElyea, K.S. & Apple, J.K. (2002). Microbial, instrumental color and sensory color and odor characteristics of ground beef produced from beef trimmings treated with ozone or chlorine dioxide. *Meat Science*. 60: 299-305.
- Stockinger Lab. (2001). Glycerol stocks. Retrieved 27 May 2015 from www.oardc.ohio-state/stockingerlab/
- Stopforth, J.D., Ikeda, J.S., Kendall, P.A. & Sofos, J.N. (2004). Survival of acid-adapted or nonadapted *Escherichia coli* O157:H7 in apple wounds and surrounding tissue following chemical treatments and storage. *International Journal Food Microbiology*. 90: 51-61.
- Suchandra, C., Zareena, N., Gautam, S., Sournyakanti, A., Prasad, S.V. & Arun, S. (2007). Antioxidant activity of some phenolic constituents from green pepper (*Piper nigrum* L.) and fresh nutmeg mace (*Myristica fragrans*). *Food Chemistry*. 101: 515-523.

- Sukrasno, Fidriany, I., Anggadiredja, K., Handayani, W.A. & Anam, K. (2011). Influence of drying method on flavonoid content of *Cosmos caudatus* (Kunth) leaves. *Research Journal of Medicinal Plant*. 5(2): 189-195.
- Sulaiman, S.F., Sajak, A.A.B., Ooi, K.L., Suprianto & Seow, E.M. (2011). Effects of solvents in extracting polyphenols and antioxidants of selected raw vegetables. *Journal of Food Composition and Analysis*. 24: 506-515.
- Sultana, B., Anwar, F. & Ashraf, M. (2009). Effect of extraction solvent / technique on the antioxidant activity of selected medicinal plant extracts. *Molecules*. 14: 2167-2180.
- Sultanbawa, Y. (2011). Plant antimicrobials in food applications: Minireview. *FORMATX*.
- Sumazian, Y., Syahida, A., Hakiman, M. & Maziah, M. (2010). Antioxidant activities, flavonoids, ascorbic acid and phenolic contents of Malaysian vegetables. *Journal of Medicinal Plant Research*. 4(10): 881-890.
- Supardy, Afifah, N., Ibrahim, D., Sulaiman, S.F. & Zakaria, N.A. (2012). Inhibition of *Klebsiella 148neumonia* ATCC 13883 by hexane extract of *Halimeda discoidea* (Decaisne) and the identification of its potential bioactive compounds. *Journal of Microbiology and Biotechnology*. 22(6): 872-881.
- Svoboda, K., Brooker, J.D. & Zrustova, J. (2006). Antibacterial and antioxidant properties of essential oils: Their potential applications in food industries. *Acta Horticulturae*. 709: 35-44.
- Swamy, T.A., Obey, J. & Mutuku, N.C. (2013). Phytochemical analysis of *Vernonia adoensis* leaves and roots used as a traditional medicinal plant in Kenya. *International Journal of Pharmaceutical and Biological*. 3: 46-52.
- Syahmi, A.R.M., Vijayarathna, S., Sasidharan, S., Latha, L.Y., Kwan, Y.P., Lau, Y.L., Shin, L.N. & Chen, Y. (2010). Acute oral toxicity and brine shrimp lethality of *Elaeis guineensis* Jacq., (oil palm leaf) methanol extract. *Molecules*. 15: 8111-8121.
- Tajkarimi, M.M., Ibrahim, S.A. & Cliver, D.O. (2010). Antimicrobial herb and spice compounds in food. *Food Control*. 21(9): 1199-1218.
- Talcott, S.T., Howard, L.R. & Bernes, C.H. (2000). Antioxidant changes and sensory properties of carrot puree processed with and without periderm tissue. *Journal of Agricultural and Food Chemistry*. 48: 1315-1321.
- Talcott, S.T., Percival, S.S., Pittet-Moore, J. & Celoria, C. (2003). Phytochemical composition and antioxidant stability of fortified yellow passion fruit (*Passiflora edulis*). *Journal of Agricultural and Food Chemistry*. 51: 935-941.

- Tanaka, T. (1997). Chemoprevention of human cancer: biology and therapy. *Critical Reviews in Oncology/Hematology*. 25: 139-174.
- Teixeira, S., Siquet, C., Alves, C., Boal, L., Marques, P.M., Borges, F., Lima, J.L.F.C. & Reis, S. (2005). Structure-property studies on the antioxidant activity of flavonoids present in diet. *Free Radical Biology and Medicine*. 39: 1099-1108.
- Tian, N., Liu, Z., Huang, J., Luo, G., Liu, S. & Liu, X. (2007). Isolation and preparation of flavonoids from the leaves of *Nelumbo nucifera* Gaertn by preparative reversed-phase high-performance liquid chromatography. *Sepu*. 25: 88-92.
- Tirado, C. & Schmidt, K. (2001). WHO surveillance program for control of foodborne infections and intoxications in Europe. Preliminary results and trends across greater Europe. *Journal of Infection*. 43: 80-84.
- Tiwari, B.K., Vald Ramidi, V.P., O'Donnell, C.P., Muthukumarappan, K., Bourke, P. & Cullen, P.J. (2009). Application of antimicrobials for food preservation. *Journal of Agricultural and Food Chemistry*. 57: 5987-6000.
- Tombola, F., Campello, S., De Luca, L., Ruggiero, P., Del Giudice, G., Papini, E. & Zoratti, M. (2003). Plant polyphenols inhibit VacA, a toxin secreted by the gastric pathogen *Helicobacter Pylori*. *Federation of European Biochemical Societies Letters*. 543: 184-189.
- Tornuk, F., Cankurt, H., Ozturk, I., Sagdic, O., Bayram, O. & Yetim, H. (2011). Efficacy of various plant hydrosols as natural food sanitizers in reducing *Escherichia coli* O157:H7 and *Salmonella* Typhimurium on fresh cut carrots and apples. *International Journal of Food Microbiology*. 148: 30-35.
- Tserennadmid, R., Takó, M., Galgóczy, L., Papp, T., Pesti, M., Vágvölgyi, C., Almassy, K. & Krisch, J. (2011). Anti-yeast activities of some essential oils in growth medium, fruit juices and milk. *International Journal of Food Microbiology*. 144: 480-486.
- Tsuchiya, H., Sato, M., Miyazaki, T., Fujiwara, S. & Tanigaki, S. (1996). Comparative study on the antibacterial activity of phytochemical flavanones against methicillin-resistant *Staphylococcus aureus*. *Journal of Ethnopharmacology*. 50: 27-34.
- Tsuchiya, H. & Linuma, M. (2000). Reduction of membrane fluidity by antibacterial sophoraflavanone G isolated from *Sophora exigua*. *Phytomedicine*. 7: 161-165.
- Tsuchiya, H. (2001). Stereospecificity in membrane effects of catechins. *Chemico-Biological Interactions*. 134(1): 41-54.

- Tzortzakis, N.G. & Economakis, C.D. (2007). Antifungal activity of lemongrass (*Cymbopogon citratus* L.) essential oil against key postharvest pathogens. *Innovative Food Science and Emerging Technologies*. 8: 253-258.
- Ukuku, D.O. & Fett, W.E. (2004). Method of applying sanitizers and sample preparation affects recovery of native microflora and *Salmonella* on whole cantaloupe surfaces. *Journal of Food Protection*. 67: 999-1004.
- Ukuku, D.O., Fett, W.F. & Sapers, G.M. (2004). Inhibition of *Listeria monocytogenes* by native microflora of whole cantaloupe. *Journal of Food Safety*. 24: 129-146.
- Upadhyay, A., Upadhyaya, I., Kollanoor-Johny, A., Baskaran, S.A., Mooyottu, S., Karumathil, D. & Venkitanarayanan, K. (2013). Inactivation of *Listeria monocytogenes* on frankfurters by plant-derived antimicrobials alone or in combination with hydrogen peroxide. *International Journal of Food Microbiology*. 163(2-3): 114-118.
- US Food and Drug Administration (USFDA). (1998). Guideline for industry. Guide to minimize microbial safety hazards for fresh fruits and vegetables. US Food and Drug Administration, Washington, DC.
- Valero, A., Perez-Rodriguez, F., Carrasco, E., Fuentes-Alventosa, J.M., Garcia-Gimeno, R.M. & Zurera, G. (2009). Modelling the growth boundaries of *Staphylococcus aureus*. Efficacy of temperature, pH and water activity. *International Journal of Food Microbiology*. 133: 186-194.
- Van Acker, F.A.A., Schouten, O., Haenen, G.R.M.M., Va der Vijgh, W.J.F. & Bast, A. (2000). Flavonoids can replace α -tocopherol as an antioxidant. *Federal of European Biochemical Societies Letters*. 473: 145-148.
- Van Zeebroeck, M., Ramon, H., DeBaerdmaeker, J., Nicolai, B. & Tijskens, E. (2007). Impact damage of apples during transport and handling. *Postharvest Biology and Technology*. 45: 157-167.
- Vikram, A. (2010). Grapefruit bioactive limonoids modulate *Escherichia coli* O157:H7 TTSS and biofilm. *International Journal of Food Microbiology*. 140(2-3): 109-116.
- Vilgis, T.A. (2013). Texture, taste and aroma: multi-scale materials and the gastrphysics of food. *Flavour*. 2: 12.
- Vital, P.G. & Rivera, W.L. (2009). Antimicrobial activity and cytotoxicity of *Chromolaena odorata* (Lf) King and Robinson and *Uncaria perrottetii* (A. Rich) Merr. extracts. *Journal of Medicinal Plant Research*. 3(7): 511-518.
- Wachtel, M.R. & Charkowski, A.O. (2002). Cross-contamination of lettuce with *Escherichia coli* O157: H7. *Journal of Food Protection*. 65(3): 465-470.

- Wan Norhana, M.N., Poole, S.E., Deeth, H.C. & Dykes, G.A. (2010). Prevalence, persistence and control of *Salmonella* and *Listeria* in shrimp and shrimp products: a review. *Food Control*. 21: 343-361.
- Wang, G., Zhao, T. & Doyle, M.P. (1997). Survival and growth of *Escherichia coli* O157:H7 in unpasteurized and pasteurized milk. *Journal of Food Protection*. 60(6): 610-613.
- Wani, N.S., Bhalerao, A.K., Ranaware, V.P. & Zanje, R. (2013). Formulation and evaluation of herbal sanitizer. *International Journal of PharmTech Research*. 5: 40-43.
- Weerakkody, N.S., Caffin, N., Turner, M.S. & Dykes, G.A. (2010). *In vitro* antimicrobial activity of less-utilized spice and herb extracts against selected food-borne bacteria. *Food Control*. 21: 1408-1414.
- Westrell, T., Ciampa, N., Boelaert, F., Helwigh, B., Korsgaard, H., & Chriel, M. (2009). Zoonotic infections in Europe in 2007: a summary of the EFSA-ECDC annual report. *Euro Surveill*. 14: pii=19100.
- White, D.G., Zhao, S., Sudler, R., Ayers, S., Friedman, S., Chen, S., McDermott, P.F., McDermott, S., Wagner, D.D. & Meng, J. (2001). The isolation of antibiotic-resistant *Salmonella* from retail ground meats. *The New England Journal of Medicine*. 345(16): 1147-1154.
- Wiedmann, M. & Roberts, A.J. (2003). Pathogen, host and environmental factors contributing to the pathogenesis of listeriosis. *Cellular Molecular Life Science* 60: 904-918.
- Williams, D.B. & Carter, C.B. (1996). Transmission electron microscopy. 2nd edition. Springer Science+Business Media, New York, USA.
- Witkowska, A.M., Hickey, D.K., Alonso-Gomez, M. & Wilkinson, M. (2013). Antimicrobial activity of commercial herb and spices extracts against selected foodborne bacteria. *Journal of Food Research*. 2(4): 37-54.
- World Health Organization (WHO). (1996). International stability testing: guidelines for stability of pharmaceutical products containing well established drug substances in conventional dosage forms. Switzerland: WHO.
- World Health Organization (WHO). (2010). Foodborne diseases – a growing risk. Initiative to estimate the global burden of foodborne diseases. Geneva: World Health Organization.
- Wong, S.P., Leong, L.P. & Koh, J.H.W. (2006). Antioxidant activities of aqueous extracts of selected plants. *Food Chemistry*. 99: 775-783.

- Wu, L., Si, J., Yuan, X. & Shi, X. (2008). Quantitative variation of flavonoids in *Houttuynia cordata* from different geographic origins in China. *Chinese Journal of Natural Medicines*. 7(1): 40-46.
- Yang, S.E., Yu, R.C. & Chou, C.C. (2001). Influence of holding temperature on the growth and survival of *Salmonella* spp. and *Staphylococcus aureus* and the production of staphylococcal enterotoxin in egg products. *International Journal of Food Microbiology*. 63: 99-107.
- Yazdani, D., Tan, Y.H., Zainal Abidin, M.A. & Jaganath, I.B. (2011). A review on bioactive compounds isolated from plants against plant pathogenic fungi. *Journal of Medicinal Plant Research*. 5(30): 6584-6589.
- Yusoff, N.A.H., Bustamam, M.S., Abas, F., Khatib, A. & Rukayadi, Y. (2014). Antimicrobial activity of *Cosmos caudatus* extract against foodborne pathogens. *Journal of Pure and Applied Microbiology*. 8(5): 3681-3688.
- Yuste, J. & Fung, D.Y.C. (2002). Inactivation of *Listeria monocytogenes* Scott A 49594 in apple juice supplemented with cinnamon. *Journal of Food Protection*. 65: 1663-1666.
- Zabka, M., Pavela, R. & Slezakova, L. (2009). Antifungal effect of *Pimenta dioica* essential oil against dangerous pathogenic and toxinogenic fungi. *Industrial Crops and Products*. 30: 250-253.
- Zainin, N.S., Lau, K.Y., Zakaria, M., Son, R., Abdull Razis, A.F. & Rukayadi, Y. (2013). Antibacterial activity of *Boesenbergia rotunda* (L.) Mansf. A. extract against *Escherichia coli*. *International Food Research Journal*. 2016: 3319-3323.
- Zheng, W. & Wang, S. (2001). Antioxidant activity and phenolic compounds in selected herbs. *Journal of Agricultural and Food Chemistry*. 49: 5165-5170.
- Zhou, G.H., Xu, X.L. & Liu, Y. (2010). Preservation technologies for fresh meat-a review. *Meat Science*. 86: 119-128.
- Zhu, M.Z., Wu, W., Jiao, L.L., Yang, P.F. & Guo, M.Q. (2015). Analysis of flavonoids in lotus (*Nelumbo nucifera*) leaves and their antioxidant activity using macroporous resin chromatography couples with LC-MS/MS and antioxidant biochemical assays. *Molecules*. 20: 10553-10565.
- Zongo, C., Savadogo, A., Somda, K.M., Koudou, J. & Traore, A.S. (2011). *In vitro* evaluation of the antimicrobial and antioxidant properties of extracts from whole plant of *Alternanthera pungens* H.B. & K. and leaves of *Combretum sericeum* G. Don. *International Journal of Phytomedicine*. 3: 182-191.