



**UNIVERSITI PUTRA MALAYSIA**

***PHENOLOGICAL, POSTHARVEST PHYSICOCHEMICAL AND  
ULTRASTRUCTURAL CHARACTERISTICS OF TORCH GINGER  
(*Etilingera elatior* (Jack) R.M. Smith) INFLORESCENCE AS A CUT  
FLOWER***

**CHOON SEA YEAT**

**FP 2016 55**



**PHENOLOGICAL, POSTHARVEST PHYSICOCHEMICAL AND  
ULTRASTRUCTURAL CHARACTERISTICS OF TORCH GINGER  
(*Etilingera elatior* (Jack) R.M. Smith) INFLORESCENCE AS A CUT FLOWER**

By

**CHOON SEA YEAT**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

**June 2016**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**PHENOLOGICAL, POSTHARVEST PHYSICOCHEMICAL AND  
ULTRASTRUCTURAL CHARACTERISTICS OF TORCH GINGER  
(*Etilingera elatior* (Jack) R.M. Smith) INFLORESCENCE AS A CUT FLOWER**

By

**CHOON SEA YEAT**

**June 2016**

**Chairman : Associate Professor Phebe Ding, PhD**  
**Faculty : Agriculture**

The extravagant and showy inflorescence of torch ginger (*Etilingera elatior*) with bright colour is suitable to be used as a cut flower. The major weakness of this inflorescence is bract browning that shortens its vase life and reduces marketability. The causes of the bract browning disorder and the timing of its occurrence have not been investigated in detail. Understanding the characteristics of torch ginger and the mechanisms that regulate bract browning is needed for the production of high quality inflorescence. The objective of this work was to assess the potential of torch ginger inflorescence as a cut flower. In study one, the growth and development cycle of torch ginger plant was determined. Results showed that the growth and development cycle of torch ginger plants was divided into vegetative and reproductive phases. The entire inflorescence development period took about 60 days from the inflorescence shoot emergence until blooming stage. In study two, the mechanisms of bract opening and peduncle strength of inflorescence was elucidated. The experiment was conducted in a nested design. Data were analysed using general linear model. The means comparison was performed using Tukey's test. Results showed that fresh weight and respiration rate of inflorescence increased gradually from tight bud to bloom stage. The significant higher respiration rate of inflorescence at bloom stage was coincided with a drastic decrease in soluble sugar content in involucre. This result indicates that soluble sugars depletion is the cause of bract browning. As the inflorescence developed, the cellulose content in involucre and floral bracts decreased significantly. In involucre, the cellulase activity showed a significant increase from tight bud to 6-tip opens stage and decreased thereafter. A significant increase in pectin methylesterase activity was recorded from 6-tip opens to torch shows stage in involucre. In peduncle, the ethanol insoluble residue and cellulose content increased significantly from tight bud to bloom stage. In study three, postharvest performance of inflorescences supplemented with sucrose was evaluated. The experiment was conducted in a randomized complete block design with factorial arrangement. Data were analysed using analysis of variance. There was a significant quadratic relationship ( $P \leq 0.05$ ) between vase life and sucrose concentrations on inflorescences at torch shows stage.

The vase life of inflorescences increased quadratically from 0 to 4% sucrose solutions and decreased thereafter. The changes of fresh weight in inflorescences were positively correlated with vase life ( $r=0.67$ ). The excessive water loss via transpiration in bract cells is the main factor in reducing vase life. Further study should be conducted to reduce the transpiration rate of inflorescence in order to prolong its longevity.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**CIRI-CIRI FENOLOGI, FIZIKOKIMIA LEPAS TUAI DAN  
ULTRASTRUKTUR PERBUNGAAN KANTAN (*Etilingera elatior* (Jack) R.M.  
Smith) SEBAGAI BUNGA KERATAN**

Oleh

**CHOON SEA YEAT**

**Jun 2016**

**Pengerusi : Profesor Madya Phebe Ding, PhD**  
**Fakulti : Pertanian**

Ciri-ciri perbungaan kantan (*Etilingera elatior*) yang menonjolkan dan warna yang menarik adalah sesuai untuk digunakan sebagai bunga keratan. Pemerangan pada brakta merupakan kelemahan pada perbungaan kantan yang memendekkan jangka hayat dan mengurangkan nilai pasarnya. Sebeb berlaku pemerangan pada brakta masih belum disiasat secara terperinci. Pemahaman pada ciri-ciri pokok kantan dan mekanisme yang mengawal pemerangan pada brakta adalah diperlukan untuk pengeluaran perbungaan kantan yang berkualiti tinggi. Objektif kajian ini adalah untuk menilai potensi perbungaan kantan sebagai bunga keratan. Kajian pertama adalah menentukan kitaran tumbesaran dan perkembangan kantan. Keputusan menunjukkan tumbesaran dan perkembangan kantan terbahagi kepada fasa vegetatif dan reproduktif. Tempoh perkembangan perbungaan bermula daripada pertumbuhan pucuk bunga sehingga perbungaan mekar mengambil masa 60 hari. Dalam kajian kedua, mekanisme pengembangan brakta dan kekukuhan tangkai bunga dikaji. Eksperimen ini dijalankan dalam rekabentuk tersarang. Data dianalisa dengan menggunakan model linear am. Perbandingan antara rawatan dengan menggunakan uji Tukey. Keputusan menunjukkan bahawa berat basah dan kadar respirasi meningkat secara beransur-ansur daripada tahap kematangan kudup ketat sehingga mekar. Kadar respirasi perbungaan yang menunjukkan tahap signifikan tinggi pada tahap kematangan mekar diikuti oleh penurunan kandungan gula terlarut dalam brakta. Pengurangan kandungan gula terlarut merupakan punca bagi pemerangan brakta. Apabila tahap kematangan perbungaan melanjut, selulosa dalam brakta menurun secara signifikan. Aktiviti selulase dalam brakta menunjukkan peningkatan secara signifikan daripada kematangan kudup ketat ke 6-hujung brakta terbuka dan menurun selepas itu. Aktiviti pektin metilesterase dalam brakta pula meningkat secara signifikan daripada kematangan 6-hujung brakta terbuka ke bentuk obor kelihatan. Dinding sel bagi brakta dihidrolisis oleh selulase dan pektin metilesterase untuk pengembangan brakta. Di tangkai bunga, sisa etanol tidak terlarut dan kandungan selulosa meningkat secara signifikan daripada tahap kematangan kudup ketat sehingga mekar. Dalam kajian ketiga, penilaian prestasi lepas tuai perbungaan yang dirawat dengan sukrosa telah dijalankan. Eksperimen ini dijalankan dengan

rekabentuk rawak blok lengkap dengan susunan faktorial. Data dianalisa dengan analisis varians. Perbandingan min antara rawatan dijalankan menggunakan uji berjarak ganda Duncan. Keputusan analisis regresi menunjukkan hubungan kuadratik secara signifikan ( $P \leq 0.05$ ) antara jangka hayat dan kepekatan sukrosa pada perbungaan di kematangan kudup ketat, 6-hujung brakta terbuka, bentuk obor kelihatan dan mekar. Jangka hayat perbungaan kantan meningkat secara kuadratik daripada larutan sukrosa pada kepekatan 0-4% dan menurun selepas itu. Analisis korelasi menunjukkan perubahan berat basah perbungaan adalah secara korelasi positif ( $r=0.67$ ) dengan jangka hayatnya. Kehilangan air yang berlebihan melalui transpirasi dalam sel-sel brakta merupakan faktor utama dalam pemendekkan jangka hayat perbungaan. Kajian selanjutnya perlu dijalankan untuk mengurangkan kadar transpirasi untuk memanjangkan jangka hayat perbungaan.




## ACKNOWLEDGEMENTS

I would like to extend my gratitude to Associate Professor Dr. Phebe Ding, the chairman of my supervisory committee; Professor Dr. Mahmud Tengku Muda Mohamed and Professor Dr. Khozirah Shaari, members of my supervisory committee, for their invaluable advice, guidance, encouragement and patience throughout the research and the preparation of this thesis.

Thanks to the staffs and friends of the Postharvest Laboratory who had helped me in completing the experiments successfully.

Last but not least, my sincere appreciation to my dearest parents, sisters and my best friend for their understanding, patience and moral support throughout my study.





I certify that a Thesis Examination Committee has met on 29 June 2016 to conduct the final examination of Choon Sea Yeat on her thesis entitled “Phenological, Postharvest Physicochemical and Ultrastructural Characteristics of Torch Ginger (*Etilingera elatior* (Jack) R.M. Smith) Inflorescence As A Cut Flower” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the degree of Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

**Siti Aishah binti Hassan, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Siti Hajar binti Ahmad, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Nur Ashikin Psyquay binti Abdullah, PhD**

Associate Professor  
Faculty of Agriculture and Food Sciences  
Universiti Putra Malaysia (Bintulu Campus)  
(Internal Examiner)

**Hazem Mohamed Kalaji, PhD**

Professor  
Warsaw University of Life Sciences  
Poland  
(External Examiner)



---

**NOR AINI AB. SHUKOR, PhD**  
Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 3 November 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Phebe Ding, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Mahmud Tengku Muda Mohamed, PhD**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Khozirah Shaari, PhD**

Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Member)

---

**BUJANG KIM HUAT, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Choon Sea Yeat (GS 28551)

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_  
Name of Chairman  
of Supervisory  
Committee: Associate Professor Dr. Phebe Ding

Signature: \_\_\_\_\_  
Name of Member of  
Supervisory  
Committee: Professor Dr. Mahmud Tengku Muda Mohamed

Signature: \_\_\_\_\_  
Name of Member of  
Supervisory  
Committee: Professor Dr. Khozirah Shaari

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xiii
<b>LIST OF FIGURES</b>	xiv
<b>LIST OF ABBREVIATIONS</b>	xvii
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	
2.1 Torch ginger	2
2.2 Flower/inflorescence development and opening	2
2.3 Criteria for selection as a cut flower	3
2.4 Causes of vase life termination	3
2.4.1 Soluble sugars depletion in cut flowers	4
2.4.2 Water relations of cut flowers	5
2.4.3 Ethylene effects on cut flowers	6
2.5 Postharvest treatment for cut flowers	6
<b>3 GROWTH AND DEVELOPMENT OF TORCH GINGER (<i>Etilingera elatior</i> (Jack) R.M. Smith) PLANT</b>	
3.1 Introduction	9
3.2 Materials and methods	9
3.2.1 Plant materials	9
3.2.2 Determination of morphological characteristics of torch ginger plant	9
3.2.3 Statistical analysis	10
3.3 Results	10
3.4 Discussion	12
3.5 Conclusions	13
<b>4 PHYSICAL, PHYSIOLOGICAL, BIOCHEMICAL AND STRUCTURAL CHARACTERISTICS OF TORCH GINGER (<i>Etilingera elatior</i> (Jack) R.M. Smith) INFLORESCENCE IN RELATION TO INFLORESCENCE OPENING AND PEDUNCLE STRENGTH</b>	
4.1 Introduction	14
4.2 Materials and methods	14
4.2.1 Plant materials	14

4.2.2	Determination of physical characteristics	15
4.2.3	Determination of physiological characteristics – respiration and ethylene production rates	15
4.2.4	Determination of biochemical characteristics	16
4.2.5	Histochemical and structural characteristics	19
4.2.6	Experimental design and statistical analysis	19
4.3	Results	19
4.3.1	Fresh weight of torch ginger inflorescence	19
4.3.2	Dry weight of torch ginger inflorescence	21
4.3.3	Water content of torch ginger inflorescence	21
4.3.4	Respiration and ethylene production rates	21
4.3.5	Mucilage content of peduncle	21
4.3.6	Ethanol insoluble residue	23
4.3.7	Cellulose content	23
4.3.8	Cellulase activity	24
4.3.9	Polygalacturonase activity	24
4.3.10	Pectin methylesterase activity	26
4.3.11	Soluble sugars content of involucre	26
4.3.12	Structural studies	27
4.4	Discussion	29
4.4.1	The characteristic of inflorescences during growth and development in relation to its longevity	29
4.4.2	Mechanisms of inflorescence opening	30
4.4.3	Structural characteristics of peduncle in relation to peduncle strength	31
4.5	Conclusions	32

## **5 SUCROSE SUPPLEMENTATION ON POSTHARVEST CHARACTERISTICS OF TORCH GINGER (*Etilingera elatior* (Jack) R.M. Smith) INFLORESCENCE**

5.1	Introduction	33
5.2	Materials and methods	33
5.2.1	Plant materials	33
5.2.2	Determination of postharvest characteristics	34
5.2.3	Determination of soluble sugar content	35
5.2.4	Tissue preparation for scanning electron microscopy	35

5.2.5	Experimental design and statistical analysis	35
5.3	Results	36
5.3.1	Vase life and bract opening of inflorescence	36
5.3.2	Solution uptake rate	39
5.3.3	Relative fresh weight and dry weight	40
5.3.4	Respiration and ethylene production rates	41
5.3.5	Soluble sugar content of inflorescence bracts	42
5.3.6	Scanning electron microscopy	43
5.4	Discussion	45
5.4.1	Sucrose solution on postharvest performance of torch ginger inflorescence	45
5.4.2	Cellular structure of torch ginger inflorescence in relation to its vase life	45
5.5	Conclusions	46
<b>6</b>	<b>SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	47
	<b>BIBLIOGRAPHY</b>	48
	<b>APPENDICES</b>	61
	<b>BIODATA OF STUDENT</b>	63
	<b>LIST OF PUBLICATIONS</b>	64

## LIST OF TABLES

Table		Page
1	Main effect of developmental stages on ethanol insoluble residue of bract of torch ginger inflorescence	23
2	Main effect of developmental stages on polygalacturonase activity of bract of torch ginger inflorescence	26
3	Main and interaction effects between developmental stages and sucrose concentrations on vase life, solution uptake rate, relative fresh weight, dry weight, respiration, ethylene production rate and soluble sugar content of cut torch ginger inflorescences	37
4	Effects of sucrose concentration on bract opening of torch ginger inflorescence at tight bud and 6-tip opens stages during vase life observation	38
5	Correlation coefficients ( $r$ ) between vase life (VL), solution uptake rate (ML), relative fresh weight (RFW), dry weight (DW) and respiration rate ( $\text{CO}_2$ ) of torch ginger inflorescences	41



## LIST OF FIGURES

Figure		Page
1	Growth chart of torch ginger plant. (A) Emergence of leafy shoot from ground. Leafy shoot was formed by layers of leaf sheath. (B) The mature leafy shoot in vegetative phase may reach a height of 3-4 m. (C) Emergence of inflorescence shoot from rhizome connected to leafy shoot signalled the start of reproductive phase. (D) Torch ginger inflorescence at the flowering stage. (E) Opening of true flowers from the fridge towards the torch of inflorescence. (F) Senescence of torch ginger inflorescence without fruit forming	11
2	Growth rate of peduncle length (A) and inflorescence diameter (B) of torch ginger inflorescence	12
3	Changes in fresh weight (A), dry weight (B), water content (C) and respiration rate (D) of torch ginger inflorescences at the four developmental stages. Means followed by the different letters are significantly different by Tukey's test at $P \leq 0.05$ . Each value is the mean of 3 replications $\pm$ SE	20
4	Mucilage content (A), ethanol insoluble residue (EIR) (B) and cellulose content (C) of torch ginger inflorescence peduncle at the four developmental stages. Means of developmental stage within the same peduncle part followed by the different letters are significantly different by Tukey's test at $P \leq 0.05$ . Each value is the mean of 3 replications	22
5	Cellulose content (A), cellulase activity (B) and pectin methylesterase activity (C) of bracts of torch ginger inflorescence at the four developmental stages. Means of developmental stage within the same bract type followed by the different letters are significantly different by Tukey's test at $P \leq 0.05$ . Each value is the mean of 3 replications	25
6	Soluble sugar content of involucre of torch ginger inflorescence at the four developmental stages. Means followed by the different letters are significantly different by Tukey's test at $P \leq 0.05$ . Each value is the mean of 3 replications $\pm$ SE	27
7	(A) Transverse section of the basal part of torch ginger peduncle. (B) Scanning electron micrograph of basal part of bloom stage peduncle (20x). (C) Scanning electron micrograph of upper part of tight bud stage peduncle (20x). OS = outer sheath, IS = inner sheath, OZ = outer heart zone, IZ = inner heart zone, VB = vascular bundles, P = pericycle ring	28

- 8 Light microscopy images of basal part of torch ginger inflorescence peduncle (10x). The transverse section revealed that oleoresin canal (OC), vascular bundles (VB) with its elements of xylem (X), phloem (PH) and fibers (F), and mucilage content (M) were found in the peduncle 28
- 9 Transverse section of the basal part of torch ginger inflorescence peduncle. (A) Light micrograph of the heart of peduncle (4x). (B) Scanning electron micrographs of the heart of peduncle in outer heart zone (OZ) (350x) and (C) inner heart zone (IZ) (350x). VB = vascular bundles, P = pericycle ring 29
- 10 Vase life of torch ginger inflorescences at three developmental stages treated with different concentrations of sucrose. Means of different vase solution concentration within the same developmental stage followed by the different letters are significantly different by DMRT at  $P \leq 0.05$  36
- 11 Relationship between vase life and sucrose concentration of torch ginger inflorescence at torch shows stage. ( $\blacktriangle$ ),  $y = -0.030x^2 + 0.034x + 4.365$ ,  $R^2 = 0.49$  38
- 12 Solution uptake rate of torch ginger inflorescences at four developmental stages treated with different concentrations of sucrose. Means of different vase solution concentration within the same developmental stage followed by the different letters are significantly different by DMRT at  $P \leq 0.05$  39
- 13 Relationships between relative fresh weight and sucrose concentration at different developmental stages of torch ginger inflorescence. Tight bud ( $\blacklozenge$ ),  $y = -0.363x^2 + 3.448x - 0.155$ ,  $R^2 = 0.73$ ; 6-tip opens ( $\blacksquare$ ),  $y = -0.093x^2 + 0.620x + 0.487$ ,  $R^2 = 0.68$ ; torch shows ( $\blacktriangle$ ),  $y = -0.039x^2 - 0.066x - 1.125$ ,  $R^2 = 0.73$ ; and bloom ( $\bullet$ ),  $y = -0.003x^2 - 0.211x - 1.526$ ,  $R^2 = 0.40$  40
- 14 Respiration rate of torch ginger inflorescences at four developmental stages treated with different concentrations of sucrose. Means of different vase solution concentration within the same developmental stage followed by the different letters are significantly different by DMRT at  $P \leq 0.05$  42
- 15 Relationships between soluble sugar content and sucrose concentration at different developmental stages of torch ginger inflorescence. Tight bud ( $\blacklozenge$ ),  $y = 2.5x^2 + 46.50x + 27.267$ ,  $R^2 = 0.94$ ; 6-tip opens ( $\blacksquare$ ),  $y = 8.106x^2 + 8.597x + 27.444$ ,  $R^2 = 0.88$ ; torch shows ( $\blacktriangle$ ),  $y = 1.063x^2 + 55.832x + 8.723$ ,  $R^2 = 0.97$ ; and bloom ( $\bullet$ ),  $y = 6.984x^2 + 1.143x + 24.772$ ,  $R^2 = 0.96$  43

- 16 Scanning electron micrographs of peduncle end of torch ginger inflorescences. (A) Longitudinal section of peduncle end harvested on day 0 (x200); (B) longitudinal section of peduncle end from inflorescence placed in 0% sucrose solution at the end of vase life (x400) and (C) longitudinal section of peduncle end from inflorescence placed in 10% sucrose solution at the end of vase life (x400). XV = xylem vessel 44
- 17 Scanning electron micrographs of involucre and floral bracts of torch ginger inflorescence. (A) abaxial surface of involucre bract (x250); (B) adaxial surface of involucre bract (x100); (C) abaxial surface of floral bract (x230) and (D) adaxial surface of floral bract (x200). S = opened-stoma 44



## LIST OF ABBREVIATIONS

1-MCP	1-methylcyclopropene
8-HQC	8-hydroxyquinoline citrate
8-HQS	8-hydroxyquinoline sulphate
AgNO <sub>3</sub>	Silver nitrate
ANOVA	Analysis of variance
AOA	Aminooxyacetic acid
AVG	Aminoethoxyvinylglycine
ClO <sub>2</sub>	Chlorine dioxide
CO <sub>2</sub>	Carbon dioxide
C <sub>2</sub> H <sub>4</sub>	Ethylene
DACP	Diazocyclopentadiene
DICA	Dichloroisocyanuric acid
DMRT	Duncan's multiple range test
DNS	3,5-dinitrosalicylic acid
DW	Dry weight
EIR	Ethanol insoluble residue
FAA	Formalin – acetic acid – ethanol – water
FW	Fresh weight
MVG	Methoxyvinylglycine
PME	Pectin methylesterase
PG	Polygalacturonase activity
SAS	Statistical Analysis Software
SE	Standard error
STS	Silver thiosulphate

## CHAPTER 1

### INTRODUCTION

Torch ginger (*Etilingera elatior* (Jack) R.M. Smith) is a perennial herbaceous clumping plant that belongs to the family Zingiberaceae. It is native to Malaysia, Indonesia and southern Thailand (Burt & Smith, 1986; Ong, 2008). The current commercial production of torch ginger in Malaysia is focused in supplying inflorescence buds for culinary purposes. However, the torch ginger inflorescence bud sold in the market is poor in quality. There is no proper postharvest handling to extend their shelf life or to maintain the eating quality. Moreover, the wholesale and retail prices of inflorescence bud are fluctuated according to market demand. Usually, high market demand only occurs during festive celebrations or when the production is low in dry season. When the market demand is low, the wholesale and retail prices are low. The inflorescence bud tend to be negligent and left to bloom, and finally dying on plant. Consequently, studies should be conducted to evaluate its diversity for other usage in order to increase its economic importance.

In recent years, torch ginger has been gaining recognition and planted as an ornamental and landscaping plant in urban areas. The extravagant and showy inflorescence with brilliant colour has demonstrated its potential as a cut flower. The torch ginger inflorescence has been used as floral arrangements in hotels and restaurants in Australia, Brazil, Costa Rica, Hong Kong, Thailand and United States of America (Larsen et al., 1999; Prasongchan et al., 2009; Loges et al., 2011; Carneiro et al., 2014). The use of torch ginger inflorescence as a cut flower is relatively new in Malaysia due to limited supply and the unawareness of growers on the potential usage. There is only small number of florists using the cut inflorescence as bouquet fillers and small scale exportation to Hong Kong during Chinese New Year. However, at farm price, the margin of cut inflorescence is five times higher than young inflorescence bud.

From the observation, bract browning is the major problem in torch ginger inflorescence to be used as cut flower. The bract browning symptom reduces the postharvest quality of the inflorescence and shortens its vase life. This reduces the marketability and export potential of the inflorescence. However, the causes of the bract browning still remain unknown. Consumers demand for high quality flowers. Therefore, reducing bract browning in improving the inflorescence quality is a high priority. There is a need to understand the characteristics of the crop and the mechanisms that regulate bract browning. Thus, the general objective of this work was to assess the potential of torch ginger inflorescence as a cut flower. Three studies were conducted with the aims (i) to determine the growth and development cycle of torch ginger plant; (ii) to elucidate the inflorescence opening and peduncle strength during inflorescence development and (iii) to reveal the postharvest performance of inflorescences at different developmental stages supplemented with sucrose as vase solution. Results from this study would enhance the understanding of this crop as a cut flower.

## BIBLIOGRAPHY

- AbdelKader, H. H. (2012). Postharvest physiology of cut hippeastrum (*Hippeastrum hybridum* Herb.) inflorescences. *World Applied Sciences Journal*, 19(7), 943-950.
- Abdelmageed, A. H. A., Faridah, Q. Z., Nur Amalina, A., & Muhamad, Y. (2011). The influence of organ and post-harvest drying period on yield and chemical composition of the essential oils of *Etilingera elatior* (Zingiberaceae). *Journal of Medicinal Plants Research*, 5(15), 3432-3439.
- Abdelwahab, S. I., Zaman, F. Q., Mariod, A. A., Yaacob, M., Abdelmageed, A. H. A., & Khamis, S. (2010). Chemical composition, antioxidant and antibacterial properties of the essential oils of *Etilingera elatior* and *Cinnamomum pubescens* Kochummen. *Journal of the Science of Food and Agriculture*, 90, 2682-2688.
- Abeles, F. B., Morgan, P. W., & Saltveit Jr, M. E. (1992). Regulation of ethylene production by internal, environmental, and stress factors. In *Ethylene in plant biology* (2nd ed., pp. 56-119). San Diego: Academic Press.
- Akoumianaki-Ioannidou, A., Darras, A. I., & Diamantaki, A. (2010). Post-harvest vase solutions and storage effects on cut *Nerium oleander* inflorescences. *The Journal of Horticultural Science and Biotechnology*, 85, 1-6.
- Asrar, A. A. (2012). Effects of some preservative solutions on vase life and keeping quality of snapdragon (*Antirrhinum majus* L.) cut flowers. *Journal of the Saudi Society of Agricultural Sciences*, 11, 29-35.
- Baille, A., Gutierrez Colomer R. P., & Gonzalez-Real, M. M. (2006). Analysis of intercepted radiation and dry matter accumulation in rose flower shoots. *Agricultural and Forest Meteorology*, 137, 68-80.
- Baker, J. E., Anderson, J. D., Adams, D. O., Apelbaum, A., & Lieberman, M. (1982). Biosynthesis of ethylene from methionine in aminoethoxyvinylglycine-resistant avocado tissue. *Plant Physiology*, 69, 93-97.
- Baker, J. E., Wang, C. Y., Lieberman, M., & Hardenburg, R. (1977). Delay of senescence in carnations by rhizobitoxine analog and sodium benzoate. *HortScience*, 12, 38-39.
- Bayogan, E. R. V., & Gratuito, M. B. B. (2015). Vase life of torch ginger (*Etilingera elatior*) inflorescences as influenced by harvest maturity and 1-methylcyclopropene. *Acta Horticulturae*, 1088, 313-317.
- Beauzamy, L., Nakayama, N., & Boudaoud, A. (2014). Flowers under pressure: ins and outs of turgor regulation in development. *Annals of Botany*, 114(7), 1517-1533.



- Blankenship, S. M., & Dole, J. M. (2003). 1-Methylcyclopropene: A review. *Postharvest Biology and Technology*, 28, 1-25.
- Blankenship, S. M., & Sisler, E. C. (1993). Response of apples to diazocyclopentadiene inhibition of ethylene binding. *Postharvest Biology and Technology*, 3, 95-101.
- Bleeksma, H. C., & van Doorn, W. G. (2003). Embolism in rose stems as a result of vascular occlusion by bacteria. *Postharvest Biology and Technology*, 29(3), 334-340.
- Borochoy, A., & Woodson, W. R. (1989). Physiology and biochemistry of flower petal senescence. In J. Janick (Ed.), *Horticultural reviews*. (Vol. 11, pp. 15-43). Hoboken, NJ: John Wiley & Sons.
- Bratcher, C. B. (1992). *Methods of producing field-grown specialty cut flowers* (Unpublished Masters thesis). Retrieved from <https://shareok.org/bitstream/handle/11244/13428/Thesis-1992-B824m.pdf?sequence=1>
- Brummell, D. A. (2006). Cell wall disassembly in ripening fruit. *Functional Plant Biology*, 33, 103-119.
- Bunya-atichart, K., Ketsa, S., & van Doorn, W. G. (2004). Postharvest physiology of *Curcuma alismatifolia* flowers. *Postharvest Biology and Technology*, 34(2), 219-226.
- Burge, G. K., Morgan, E. R., Eason, J. R., Clark, G. E., Catley, J. L., & Seelye, J. F. (2008). *Sandersonia aurantiaca*: Domestication of a new ornamental crop. *Scientia Horticulturae*, 118, 87-99.
- Burt, B. L., & Smith, R. M. (1986). *Etilingera*: the inclusive name for *Achasma*, *Geanthus* and *Nicolaia* (Zingiberaceae). *Notes from the Royal Botanic Garden Edinburgh*, 43, 235-241.
- Carneiro, D. N. M., Paiva, P. D. D. O., Carneiro, L. F., Rodrigues, R. D. S., Lima, L. C. D. O., Dias, G. D. M. G., & Pedroso, R. G. A. V. (2014). Developmental stage and pulsing in inflorescences of torch ginger. *Revista Brasileira de Horticultura Ornamental*, 20(2), 163-170.
- Çelikel, F. G., Cevallos, J. C., & Reid, M. S. (2010). Temperature, ethylene and the postharvest performance of cut snapdragons (*Antirrhinum majus*). *Scientia Horticulturae*, 125, 429-433.
- Chan, E. W. C., Lim, Y. Y., & Wong, S. K. (2011). Phytochemistry and pharmacological properties of *Etilingera elatior*: A review. *Pharmacognosy Journal*, 3(22), 6-10.

- Chutichudet, P., Chutichudet, B., & Boontiang, K. (2010). Effect of 1-MCP fumigation on vase life and other postharvest qualities of Siam Tulip (*Curcuma aeruginosa* Roxb) cv. Laddawan. *International Journal of Agricultural Research*, 5(1), 1-10.
- Criley, R. A., & Paull, R. E. (1993). Review: Postharvest handling of bold tropical cut flowers *Anthurium*, *Alpinia purpurata*, *Heliconia*, and *Strelitzia*. *Acta Horticulturae*, 337, 201-211.
- da Silva J. A. T. 2003. The cut flower: postharvest considerations. *Journal of Biological Sciences*, 3(4), 406-442.
- Dai, J., & Paull, R. E. (1995). Source-sink relationship and *Protea* postharvest leaf blackening. *Journal of the American Society for Horticultural Science*, 120(3), 475-480.
- Damunupola, J. W., & Joyce, D. C. (2008). When is a vase solution biocide not, or not only, antimicrobial? *Journal of the Japanese Society for Horticultural Science*, 77(3), 211-228.
- Damunupola, J. W., Qian, T., Muusers, R., Joyce, D. C., Irving, D. E., van Meeteren, U. (2010). Effect of S-carvone on vase life parameters of selected cut flower and foliage species. *Postharvest Biology and Technology*, 55, 66-69.
- Darras, A. I., Akoumianaki-Ioannidou, A., & Pompodakis, N. E. (2010). Evaluation and improvement of post-harvest performance of cut *Viburnum tinus* inflorescence. *Scientia Horticulturae*, 124(3), 376-380.
- de Witte, Y., Harkema, H., & van Doorn, W. G. (2014). Effect of antimicrobial compounds on cut *Gerbera* flowers: Poor relation between stem bending and numbers of bacteria in the vase water. *Postharvest Biology and Technology*, 91, 78-83.
- Dole, J. M., Vilorio, Z., Fanelli, F. L., & Fonteno, W. (2009). Postharvest evaluation of cut dahlia, linaria, lupine, poppy, rudbeckia, trachelium, and zinnia. *HortTechnology*, 19(3), 593-600.
- Eason, J. R., Morgan, E. R., Mullan, A. C., & Burge, G. K. (2001). Postharvest characteristics of Santonia 'Golden Lights' a new hybrid cut flower from *Sandersonia aurantiaca* x *Littonia modesta*. *Postharvest Biology and Technology*, 22, 93-97.
- Emongor, V. E. (2004). Effects of gibberellic acid on postharvest quality and vase life of gerbera cut flowers (*Gerbera jamesonii*). *Journal of Agronomy*, 3(3), 191-195.
- Evans, R. Y., & Reid, M. S. (1988). Changes in carbohydrates and osmotic potential during rhythmic expansion of rose petals. *Journal of the American Society for Horticultural Science*, 113(6), 884-888.



- Gago, C. M. L., & Monteiro, J. A. (2011). NAA and STS effects on bract survival time, carbohydrate content, respiration rate and carbohydrate balance of potted *Bougainvillea spectabilis* Willd. *Postharvest Biology and Technology*, *60*, 235-243.
- Galen, C., Dawson, I. E., & Stanton, M. L. (1993). Carpels as leaves: Meeting the carbon cost of reproduction in an alpine buttercup. *Oecologia*, *95*, 187-193.
- Gendy, A. S. H., & Mahmoud, A. A. (2012). Effect of some preservative solution treatments on characters of *Strelitzia reginae* cut flowers. *Australian Journal of Basic and Applied Sciences*, *6*(5), 260-267.
- Ghose, T. K. (1987). Measurement of cellulase activities. *Pure and Applied Chemistry*, *59*(2), 257-268.
- Gookin, T. E., Hunter, D. A., & Reid, M. S. (2003). Temporal analysis of alpha and beta-expansin expression during floral opening and senescence. *Plant Science*, *164*(5), 769-781.
- Hagerman, A. E., & Austin, P. J. (1986). Continuous spectrophotometric assay for plant pectin methyl esterase. *Journal of Agricultural and Food Chemistry* *34*, 440-444.
- Halevy, A. H., & Mayak, S. (1979). Senescence and postharvest physiology of cut flowers: Part 1. In J. Janick (Ed.), *Horticultural reviews* (Vol. 1, pp. 204-236). Hoboken, NJ: John Wiley & Sons.
- Harada, T., Torii, Y., Morita, S., Onodera, R., Hara, Y., Yokoyama, R., Nishitani, K., & Satoh, S. (2011). Cloning, characterization, and expression of xyloglucan endotransglucosylase/hydrolase and expansin genes associated with petal growth and development during carnation flower opening. *Journal of Experimental Botany*, *62*(2), 815-823.
- He, S. G., Joyce, D. C., & Irving, D. E. (2006a). Competition for water between inflorescences and leaves in cut flowering stems of *Grevillea* 'Crimson Yul-lo'. *The Journal of Horticultural Science and Biotechnology*, *81*(5), 891-897.
- He, S. G., Joyce, D. C., Irving, D. E., & Faragher, D. E. (2006b). Stem end blockage in cut *Grevillea* 'Crimson Yul-lo' inflorescences. *Postharvest Biology and Technology*, *41*, 78-84.
- Ho, L. C., & Nichols, R. (1977). Translocation of <sup>14</sup>C-sucrose in relation to changes in carbohydrate content in rose corollas cut at different stages of development. *Annals of Botany*, *41*(1), 227-242.
- Hu, Y., Doi, M., & Imanishi, H. (1998). Competitive water relations between leaves and flower bud during transport of cut roses. *Journal of the Japanese Society for Horticultural Science*, *67*, 532-536.

- Ichimura, K., Kawabata, Y., Kishimoto, M., Goto, R., & Yamada, K. (2002). Variation with the cultivar in the vase life of cut rose flowers. *Bulletin of the National Institute of Floricultural Science*, 2, 9-20.
- Imsabai, W., Ketsa, S., & van Doorn, W. G. (2010). Role of ethylene in the lack of floral opening and in petal blackening of cut lotus (*Nelumbo nucifera*) flowers. *Postharvest Biology and Technology*, 58, 57-64.
- Imsabai, W., Leethiti, P., Netlak, P., & van Doorn, W. G. (2013). Petal blackening and lack of bud opening in cut lotus flowers (*Nelumbo nucifera*): Role of adverse water relations. *Postharvest Biology and Technology*, 79, 32-38.
- Ito, A., Hayama, H., & Kashimura, Y. (2004). Possible roles of sugar concentration and its metabolism in the regulation of flower bud formation in Japanese pear (*Pyrus pyrifolia*). *Acta Horticulturae*, 363, 365-373.
- Jackie, T., Haleagrahara, N., & Chakravarthi, S. (2011). Antioxidant effects of *Etlingera elatior* flower extract against lead acetate – induced perturbations in free radical scavenging enzymes and lipid peroxidation in rats. *BMC Research Notes*, 4, 67-74.
- Jang, J., Leon, P., Zhou, L., & Sheen, J. (1997). Hexokinase as a sugar sensor in higher plants. *Plant Cell*, 9, 5-19.
- Jaroenkit, T., & Paull, R. E. (2003). Postharvest handling of heliconia, red ginger, and bird-of-paradise. *HortTechnology*, 13(2), 259-266.
- Javad, N. D. M., Mahmood, P. Y., Roya, K., & Hamideh, J. M. (2012). Effect of cultivar on water relations and postharvest quality of gerbera (*Gerbera jamesonii* Bolus ex. Hook f.) cut flower. *World Applied Sciences Journal*, 18(5), 698-703.
- Jones, M. L., Kim, E. S., & Newman, S. E. (2001). Role of ethylene and 1-MCP in flower development and petal abscission in zonal geraniums. *HortScience*, 36(7), 1305-1309.
- Jones, R. B., & Truett, J. K. (1992). Postharvest handling of cut *Gloriosa rothschildiana* O'Brien (Liliaceae) flowers. *Journal of the American Society for Horticultural Science*, 117(3), 442-445.
- Joyce, D. C., & Jones, P. N. (1992). Water balance of the foliage of cut Geraldton waxflower. *Postharvest Biology and Technology*, 2, 31-39.
- Kazemi, M., & Ameri, A. (2012). Postharvest life of cut gerbera flowers as affected by nano-silver and acetylsalicylic acid. *Asian Journal of Biochemistry*, 7, 106-111.
- Ketsa, S., & Chinprayoon, S. (2007). Effect of holding solutions on vase life of cut miniature roses. *Acta Horticulturae*, 751, 459-463.

- Knee, M. (2000). Selection of biocides for use in floral preservatives. *Postharvest Biology and Technology*, 18(3), 227-234.
- Lachumy, S. J. T., Sasidharan, S., Sumathy, V., & Zuraini, Z. (2010). Pharmacological activity, phytochemical analysis and toxicity of methanol extract of *Etilingera elatior* (torch ginger) flowers. *Asian Pacific Journal of Tropical Medicine*, 3, 769-774.
- Larsen, K., Ibrahim, H., Khaw, S. H., & Saw, L. G. (1999). *Gingers of Peninsular Malaysia and Singapore*. Kota Kinabalu: Natural History Publications (Borneo).
- Lau, O. L., & Yang, S. F. (1976). Inhibition of ethylene production by cobaltous ion. *Plant Physiology*, 58, 114-117.
- Loges, V., da Costa, A. S., Guimaraes, W. N. R., & Teixeira, M. D. C. F. (2011). Market potential of torch ginger and beehive: Part 2. *Bulletin Heliconia Society International*, 17(4), 1-4.
- Loubaud, M., & van Doorn, W. G. (2004). Wound-induced and bacteria-induced xylem blockage in roses, *Astilbe*, and *Viburnum*. *Postharvest Biology and Technology*, 32, 281-288.
- Macnish, A. J., Leonard, R. T., & Nell, T. A. (2008). Treatment with chlorine dioxide extends the vase life of selected cut flowers. *Postharvest Biology and Technology*, 50, 197-207.
- Mansouri, H. (2012). Salicylic acid and sodium nitroprusside improve postharvest life of chrysanthemums. *Scientia Horticulturae*, 145, 29-33.
- McConchie, R., Lang, N. S., & Gross, K. C. (1991). Carbohydrate depletion and leaf blackening in *Protea neriljiolia*. *Journal of the American Society for Horticultural Science*, 116(6), 1019-1024.
- McCully, M. E., & Boyer, J. S. (1997). The expansion of maize root-cap mucilage during hydration: 3. Changes in water potential and water content. *Physiologia Plantarum*, 99, 169-177.
- Miller, G. L. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry*, 31(3), 426-428.
- Mohamad, H., Lajis, N. H., Abas, F., Ali, A. M., Sukari, M. A., Kikuzaki, H., & Nakatani, N. (2005). Antioxidative constituents of *Etilingera elatior*. *Journal of Natural Products*, 68, 285-288.
- Mujaffar, S., & Sankat, C. K. (2003). Effect of waxing on the water balance and keeping qualities of cut anthuriums. *International Agrophysics*, 17, 77-84.

- Nishikawa, F., Iwama, T., Masaya, K., Hyodo, H., Ikoma, Y., & Yano, M. (2005). Effect of sugars on ethylene synthesis and responsiveness in harvested broccoli florets. *Postharvest Biology and Technology*, 36, 157-165.
- O'Donoghue, E. M., Somerfield, S. D., & Heyes, J. A. (2002). Organization of cell walls in *Sandersonia aurantiaca* floral tissue. *Journal of Experimental Botany*, 53(368), 513-523.
- OIV. (2008). Cellulases. *International Oenological Codex*. Paris: International Organisation of Vine and Wine.
- OIV. (2012a). Polygalacturonase. *International Oenological Codex*. Paris: International Organisation of Vine and Wine.
- OIV. (2012b). Pectin Methylsterase. *International Oenological Codex*. Paris: International Organisation of Vine and Wine.
- Ong, H. C. (2008). *Rempah-ratus: Khasiat makanan & ubatan*. Kuala Lumpur: Utusan Publications & Distributors.
- Owusu, S. K. (2013). *The effects of different levels of sugar and sodium benzoate on the extension of Heliconia psittacorum cut flowers* (Unpublished Master thesis). Retrieved from <http://dspace.knust.edu.gh:8080/jspui/handle/123456789/5313>
- Perik, R. R. J., Razé D., Harkema, H., Zhong, Y., & van Doorn, W. G. (2012). Bending in cut *Gerbera jamesonii* flowers relates to adverse water relations and lack of stem sclerenchyma development, not to expansion of the stem central cavity or stem elongation. *Postharvest Biology and Technology*, 74, 11-18.
- Philosoph-Hadas, S., Meir, S., Rosenberger, I., & Halevy, A. H. (1996). Regulation of the gravitropic response and ethylene biosynthesis in gravistimulated snapdragon spikes by calcium chelators and ethylene inhibitors. *Plant Physiology*, 110, 301-310.
- Piromruen, B., Buanong, M., & Kanlayanarat, S. (2008). Effect of thidiazuron on quality and vase life of heliconia (*Heliconia* spp. cv. Bigbug). *Acta Horticulturae*, 804, 283-286.
- Porat, R., Shlomo, E., Serek, M., Sisler, E. C., & Borochoy, A. (1995). 1-methylcyclopropene inhibits ethylene action in cut phlox flowers. *Postharvest Biology and Technology*, 6(3), 313-319.
- Prasongchan, N., Jitphusa, A., & Suksawat, M. (2009). Evaluation of different chemicals treatments on vase life of torch ginger [*Etilingera elatior* (Jack) R.M. Smith]. In I. Warrington, M. Grossman, W. Schaefer, K. Schaller, J. L. Silva, T. Kamimoto, S. Frusher, R. Zawadzki, T. Kiatsiroj, P. Toedtoon et al. (Eds.). Paper presented at the International Conference on the Role of Universities in Hands-on Education 2009, Rajamangala University of

Technology Lanna (pp. 304-311). Chiang-Mai: Rajamangala University of Technology Lanna Press.

- Price, A. M., Orellana, A. D. F., Salleh, F. M., Stevens, R., Acock, R., Buchanan-Wollaston, V., Stead, A. D., & Rogers, H. J. (2008). A comparison of leaf and petal senescence in wallflower reveals common and distinct patterns of gene expression and physiology. *Plant Physiology*, *147*, 1898-1912.
- Pun, U. K., & Ichimura, K. (2003). Role of sugars in senescence and biosynthesis of ethylene in cut flowers. *Japan Agricultural Research Quarterly*, *37*, 219-224.
- Ranwala, A. P., & Miller, W. B. (2009). Comparison of the dynamic of non-structural carbohydrate pools in cut tulip stems supplied with sucrose or trehalose. *Postharvest Biology and Technology*, *52*, 91-96.
- Razali, Z., Boyce, A. N., Nair, H., & Chandran, S. (2013). Cell wall enzymes and changes in sugar content in pollinated orchids treated with ethylene inhibitors and sugars. *Scientia Horticulturae*, *159*, 143-151.
- Reid, M. S. (1985). Ethylene and abscission. *HortScience*, *20*(1), 45-50.
- Reid, M. S. (2005). Flower development: from bud to bloom. *Acta Horticulturae*, *669*, 105-109.
- Reid, M. S., & Evans, R. Y. (1986). Control of cut flower opening. *Acta Horticulturae*, *181*, 45-54.
- Reid, M. S., Evans, R. Y., Dodge, L. L., & Mor, Y. (1989). Ethylene and silver thiosulfate influence opening of cut rose flowers. *Journal of the American Society for Horticultural Science*, *114*, 436-440.
- Reid, M. S., & Wu, M. J. (1992). Ethylene and flower senescence. *Plant growth regulation*, *11*, 37-43.
- Rogers, H. J. (2006). Programmed cell death in floral organs: How and why do flowers die? *Annals of Botany*, *97*, 309-315.
- Sakai, S., Kato, M., & Inoue, T. (1999). Three pollination guilds and variation in floral characteristics of Bornean gingers (Zingiberaceae and Costaceae). *American Journal of Botany*, *86*(5), 646-658.
- Sass, J. E. (1958). *Botanical microtechnique* (3rd ed.). Ames, Iowa: Iowa State College Press.
- Satoh, S., Nukui, H., & Inokuma, T. (2005). A method for determining the vase life of cut spray carnation flowers. *Journal of Applied Horticulture*, *7*(1), 8-10.



- Scariot, V., Paradiso, R., Rogers, H., & De Pascale, S. (2014). Ethylene control in cut flowers: Classical and innovative approaches. *Postharvest Biology and Technology*, 97, 83-92.
- Schroeder, K. R., & Stimart, D. P. (2005). Comparison of stomatal density and postharvest transpiration between long- and short-lived cut flower genotypes of *Antirrhinum majus* L. *Journal of the American Society for Horticultural Science*, 130(5), 742-746.
- Scofield, G. N., Ruuska, S. A., Aoki, N., Lewis, D. C., Tabe, L. M., & Jenkins, C. L. D. (2009). Starch storage in the stem of wheat plants: Localization and temporal changes. *Annals of Botany*, 103, 859-868.
- Scott Jr, T. A., & Melvin, E. H. (1953). Determination of dextran with anthrone. *Analytical Chemistry*, 25(11), 1656-1661.
- Sengupta, S., & Majumder, A. L. (2014). Physiological and genomic basis of mechanical-functional trade-off in plant vasculature. *Frontiers in Plant Science*, 5, 224.
- Serek, M., Sisler, E. C., & Reid, M. S. (1995a). Effects of 1-MCP on the vase life and ethylene response of cut flowers. *Plant Growth Regulation*, 16, 93-97.
- Serek, M., Sisler, E. C., & Reid, M. S. (1995b). 1-Methylcyclopropene, a novel gaseous inhibitor of ethylene action, improves the vase life of fruits, cut flowers and potted plants. *Acta Horticulturae*, 394, 337-346.
- Serek, M., Woltering, E. J., Sisler, E. C., Frello, S., & Sriskandarajah, S. (2006a). Controlling ethylene responses in flowers at the receptor level. *Biotechnology Advances*, 24(4), 368-381.
- Serek, M., Sisler, E. C., Frello, S., Sriskandarajah, S. (2006b). Postharvest technologies for extending the shelf life of ornamental crops. *International Journal of Postharvest Technology and Innovation*, 1, 69-75.
- Shahri, W., & Tahir, I. (2011). Flower development and senescence in *Ranunculus asiaticus* L. *Journal of Fruit and Ornamental Plant Research*, 19(2), 123-131.
- Shanan, T. N., Emara, K. S., & Barakat, S. O. (2010). Prolonging vase life of carnation flowers using natural essential oils and its impact on microbial profile of vase solutions. *Australian Journal of Basic and Applied Sciences*, 4(8), 3559-3574.
- Shimizu-Yumoto, H., & Ichimura, K. (2010). Postharvest physiology and technology of cut *Eustoma* flowers. *Journal of the Japanese Society for Horticultural Science*, 79(3), 227-238.

- Shtein, I., Meir, S., Riov, J., & Philosoph-Hadas, S. (2011). Interconnection of seasonal temperature, vascular traits, leaf anatomy and hydraulic performance in cut *Dodonaea* 'Dana' branches. *Postharvest Biology and Technology*, *61*, 184-192.
- Shu, Z., Shi, Y., Qian, H., Tao, Y., & Tang, D. (2010). Distinct respiration and physiological changes during flower development and senescence in two *Freesia* cultivars. *HortScience*, *45*(7), 1088-1092.
- Sisler, E. C., & Serek, M. (2003). Compounds interacting with the ethylene receptor in plants. *Plant Biology*, *5*(5), 473-480.
- Sood, S., Vyas, D., & Nagar, P. K. (2006). Physiological and biochemical studies during flower development in two rose species. *Scientia Horticulturae*, *108*, 390-396.
- Spikman, G. (1989). Development and ethylene production of buds and florets of cut freesia inflorescences as influenced by silver thiosulphate, aminoethoxyvinylglycine and sucrose. *Scientia Horticulturae*, *39*, 73-81.
- Sun, Y., Christensen, B., Liu, F., Wang, H., & Müller, R. (2009). Effects of ethylene and 1-MCP (1-methylcyclopropene) on bud and flower drop in mini *Phalaenopsis* cultivars. *Plant Growth Regulation*, *59*, 83-91.
- Takahashi, R., Fujitani, C., Yamaki, S., & Yamada, K. (2007). Analysis of the cell wall loosening proteins during rose flower opening. *Acta Horticulturae*, *755*, 483-488.
- Thanatcha, R., & Pranee, A. (2011). Extraction and characterization of mucilage in *Ziziphus mauritiana* Lam. *International Food Research Journal*, *18*, 201-212.
- Tripathi, S. K., & Tuteja, N. (2007). Integrated signaling in flower senescence. *Plant Signaling and Behavior*, *2*(6), 437-445.
- Trusty, S. E., & Miller, W. B. (1991). Postproduction carbohydrate levels in pot chrysanthemums. *Journal of the American Society for Horticultural Science*, *116*(6), 1013-1018.
- Unemoto, L. K., de Faria, R. T., Takahashi, L. S. A., de Assis, A. M., & Lone, A. B. (2011). Longevity of torch ginger inflorescences with 1-methylcyclopropene and preservative solutions. *Acta Scientiarum: Agronomy*, *33*(4), 649-653.
- Updegraff, D. M. (1969). Semimicro determination of cellulose in biological materials. *Analytical Biochemistry*, *32*, 420-424.
- van der Meulen-Muisers, J. J. M., van Oeveren, J. C., van der Plas, L. H. W., & van Tuyl, J. M. (2001). Postharvest flower development in Asiatic hybrid lilies as related to tepal carbohydrate status. *Postharvest Biology and Technology*, *21*, 201-211.

- van Doorn, W. G. (1996). Water relations of cut flowers. In J. Janick (Ed.), *Horticultural reviews* (Vol. 18, pp. 1-85). Oxford: John Wiley & Sons.
- van Doorn, W. G. (1999). Vascular occlusion in cut flowers. I. General principles and recent advances. *Acta Horticulturae*, 482, 59-63.
- van Doorn, W. G. (2004). Is petal senescence due to sugar starvation? *Plant Physiology*, 134, 35-42.
- van Doorn, W. G. (2012). Water relations of cut flowers: An update. In J. Janick (Ed.), *Horticultural reviews* (Vol. 40, pp. 55-106). Hoboken, NJ: John Wiley & Sons.
- van Doorn, W. G., Balk, P. A., van Houwelingen, A. M., Hoerberichts, F. A., Hall, R. D., Vorst, O., van der Schoot, C., & van Wordragen, M. F. (2003). Gene expression during anthesis and senescence in *Iris* flowers. *Plant Molecular Biology*, 53, 845-863.
- van Doorn, W. G., Groenewegen, G., van de Pol, P., & Berkholst, E. M. (1991). Effects of carbohydrate and water status on flower opening of cut Madelon roses. *Postharvest Biology and Technology*, 1(1), 47-57.
- van Doorn, W. G., de Witte, Y., & Perik, R. R. J. (1990). Effect of antimicrobial compounds on the number of bacteria in stems of cut rose flowers. *Journal of Applied Microbiology*, 68(2), 117-122.
- van Doorn, W. G., & Han, S. S. (2011). Postharvest quality of cut lily flowers. *Postharvest Biology and Technology*, 62, 1-6.
- van Doorn, W. G., & Kamdee, C. (2014). Flower opening and closure: An update. *Journal of Experimental Botany*, 65(20), 5749-5757.
- van Doorn, W. G., Perik, R. R. J., Abadie, P., & Harkema, H. (2011). A treatment to improve the vase life of cut tulips: Effects on tepal senescence, tepal abscission, leaf yellowing and stem elongation. *Postharvest Biology and Technology*, 61, 56-63.
- van Doorn, W. G., & van Meeteren, U. (2003). Flower opening and closure: A review. *Journal of Experimental Botany*, 54(389), 1801-1812.
- van Doorn, W. G., & Woltering, E. J. (2008). Physiology and molecular biology of petal senescence. *Journal of Experimental Botany*, 59(3), 453-480.
- van Meeteren, U., van Ieperen, W., Scheenen, T., & van As, H. (2001). Processes and xylem anatomical properties involved in rehydration dynamics of cut flowers. *Acta Horticulturae*, 543, 207-215.
- Veen, H. (1979). Effects of silver on ethylene synthesis and action in cut carnations. *Planta*, 145, 467-470.



- Walton, E. F., Boldingh, H. L., McLaren, G. F., Williams, M. H., & Jackman, R. (2010). The dynamics of starch and sugar utilisation in cut peony (*Paeonia lactiflora* Pall.) stems during storage and vase life. *Postharvest Biology and Technology*, 58(2), 142-146.
- Wang, C. Y., Baker, J. E., Hardenburg, R., & Lieberman, M. (1977). Effects of two analogs of rhizobitoxine sodium benzoate on senescence of snapdragons. *Journal of the American Society for Horticultural Science*, 102, 517-520.
- Wang, R., Zheng, X., & Xu, X. (2014). Evidence for physiological vascular occlusion in stems of cut *Gerbera* cv. Hongyan. *Journal of Agricultural Science and Technology*, 16, 365-372.
- Waseem, S., & Inayatullah, T. (2011). Physiological and biochemical changes associated with flower development and senescence in *Consolida ajacis* Nieuwl cv. Violet blue. *Frontiers of Agriculture in China*, 5(2), 201-208.
- Wernett, H. C., Wilfret, G. J., Sheehan, T. J., Marousky, F. J., Lyrene, P. M., & Knauft, D. A. (1996). Postharvest longevity of cut-flower *Gerbera*. I. Response to selection for vase life components. *Journal of the American Society for Horticultural Science*, 121(2), 216-221.
- Woltering, E. J., & van Doorn, W. G. (1988). Role of ethylene in senescence of petals-morphological and taxonomical relationships. *Journal of Experimental Botany*, 208, 1605-1616.
- Woolf, A. B., Combes, S., Petley, M., Olsson, S. R., Wohlers, M., Jackman, R. C. (2012). Hot water treatments reduce leaf yellowing and extend vase life of Asiatic hybrid lilies. *Postharvest Biology and Technology*, 64, 9-18.
- Yamada, K., Ito, M., Oyama, T., Nakada, M., Maesaka, M., & Yamaki, S. (2007). Analysis of sucrose metabolism during petal growth of cut roses. *Postharvest Biology and Technology*, 43, 174-177.
- Yamada, K., Norikoshi, R., Suzuki, K., Imanishi, H., & Ichimura, K. (2009a). Determination of subcellular concentrations of soluble carbohydrates in rose petals during opening by nonaqueous fractionation method combined with infiltration-centrifugation method. *Planta*, 230, 1115-1127.
- Yamada, K., Norikoshi, R., Suzuki, K., Nishijima, T., Imanishi, H., & Ichimura, K. (2009b). Cell division and expansion growth during rose petal development. *Journal of the Japanese Society for Horticultural Science*, 78(3), 356-362.
- Yamada, T., Takatsu, Y., Manabe, T., Kasumi, M., & Marubashi, W. (2003). Suppressive effect of trehalose on apoptotic cell death leading to petal senescence in ethylene-insensitive flowers of gladiolus. *Plant Science*, 164, 213-221.

- Yap, Y. M., Loh, C. S., & Ong, B. L. (2008). Regulation of flower development in *Dendrobium crumenatum* by changes in carbohydrate contents, water status and cell wall metabolism. *Scientia Horticulturae*, 119, 59-66.
- Yeat, C. S., Szydlik, M., & Łukaszewska, A. J. (2012). The effect of postharvest treatments on flower quality and vase life of cut alstroemeria 'Dancing Queen'. *Journal of Fruit and Ornamental Plant Research*, 20(2), 147-160.
- Yemm, E. W., & Willis, A. J. (1954). The estimation of carbohydrates in plant extracts by anthrone. *Biochemical Journal*, 57(3), 508-514.
- Zhao, D., Han, C., Tao, J., Wang, J., Hao, Z., Geng, Q., & Du, B. (2012). Effects of inflorescence stem structure and cell wall components on the mechanical strength of inflorescence stem in herbaceous peony. *International Journal of Molecular Sciences*, 13, 4993-5009.

